

RIGIDITY OF THE BRYANT RICCI SOLITON

ZIYI ZHAO[†] AND XIAOHUA ZHU[‡]

ABSTRACT. We introduce a new curvature-pinching condition, which is weaker than the positive sectional curvature or the positive isotropic curvature, and then we prove several rigidity results for the rotational solutions of steady Ricci solitons, i.e., the Bryant Ricci solitons.

CONTENTS

0.	Introduction	1
1.	Preliminary	4
2.	Fine estimates of curvature behavior	6
2.1.	$\frac{1}{2}$ -order improvement	7
2.2.	Fine roundness of level sets	14
3.	Solutions of (0.6)	15
4.	Almost Killing VFs with fast decay of Lie derivative	18
5.	Evolution solution of Lichnerowicz equation	24
6.	Proof of main theorem and corollaries	28
	References	29

0. INTRODUCTION

In [4], Brendle solved the following Perelman’s conjecture [32]:

Any 3-dimensional (complete) non-flat κ -noncollapsed steady gradient soliton is rotationally. Namely, it is isometric to the Bryant Ricci soliton up to scaling.

For general dimensional steady gradient solitons, Brendle also proved [5]

Theorem 0.1. *Any steady gradient Ricci soliton (M^n, g) ($n \geq 3$) with positive sectional curvature must be isometric to the Bryant Ricci soliton up to scaling if it is asymptotically cylindrical.*

Since any 3-dimensional non-flat steady gradient soliton has always positive sectional curvature [10] and it is also asymptotically cylindrical by the κ -noncollapsed property [32], Theorem 0.1 in particular also confirms the Perelman’s conjecture.

The asymptotically cylindrical behavior is an asymptotic property for a Riemannian manifold (M, g) [5] (also see Definition 1.2). In this paper, we purpose to weaken the global condition of positive sectional curvature to generalize Theorem 0.1.

Let Rm_g be the curvature operator tensor and $R = R(x)$ the scalar curvature of (M, g) . We introduce the following curvature-pinching condition,

$$(0.1) \quad 2P < R,$$

where the function $P = P(x)$ is defined on M by

$$(0.2) \quad P(\cdot) = \sup_h \frac{\text{Rm}_g(\cdot)(h, h)}{|h|^2} = \sup_h \frac{\mathbf{R}_{ijkl}(\cdot)h_{ik}h_{jl}}{h_{ik}h_{jk}}, \quad \forall \text{ symmetric } 2\text{-tensor } h \neq 0.$$

The following result is our main result in this paper.

2000 *Mathematics Subject Classification.* Primary: 53C25; Secondary: 53C55, 58J05, 32Q20.

Key words and phrases. Bryant Ricci soliton, Ricci flow, curvature pinching, positive isotropic curvature.

[†] [‡] partially supported by NSFC 12271009.

Theorem 0.2. *Let (M^n, g) ($n \geq 4$) be a steady gradient Ricci soliton which satisfies (0.1). Then (M, g) must be isometric to the Bryant Ricci soliton up to scaling if it is asymptotically cylindrical.*

It can be checked that there are two special cases of (M^n, g) with curvatures: positive sectional curvature ($n \geq 4$) or positive isotropic curvature (PIC, $n \geq 5$) on which (0.1) holds (cf. Proposition 1.6). Thus, in particular, Theorem 0.2 is a generalization of Theorem 0.1 for the steady gradient solitons with positive sectional curvature

As an application of Theorem 0.2, we prove the following rigidity of the Bryant soliton with respect to the linear decay curvature .

Corollary 0.3. *Let (M^n, g) ($n \geq 4$) be a κ -noncollapsed steady gradient Ricci soliton with positive Ricci curvature. Suppose that there exists C_0 and r_0 such that for any $x \in M$ with $\rho(x) \geq r_0$ it holds*

$$(0.3) \quad \text{Rm}_g(x) \geq 0 \text{ and } R(x) \leq \frac{C_0}{\rho(x)}.$$

Then (M, g) is isometric to the Bryant Ricci soliton if it satisfies (0.1). In particular, any n -dimensional κ -noncollapsed steady gradient Ricci soliton with either the positive Ricci curvature and PIC ($n \geq 5$) or the positive sectional curvature ($n \geq 4$) must be isometric to the Bryant Ricci soliton up to scaling under the curvature decay (0.3).

The above corollary generalizes a result of Deng-Zhu [18] instead of the positive sectional curvature by the pinching-curvature (0.1). We note that the global condition of positive Ricci curvature and the condition (0.3) near the infinity are both necessary according to examples of κ -noncollapsed steady gradient Ricci soliton constructed in [23, 2], which are not isometric to the Bryant Ricci soliton up to scaling. But we guess that the positive Ricci curvature condition can be removed under (0.1).

The notion of PIC was first introduced by Micallef and Moore [30] in their work on the index problem of minimal two-spheres. The PIC condition is preserved under the Ricci flow, proved by Hamilton [27] for 4-dimension, and Nguyen [31] and Brendle-Schoen [6] independently for higher dimensions. Two other closely related PIC1 and PIC2, a little more strong curvatures than PIC, which were introduced by Brendle-Schoen play important roles in their proof of the differentiable sphere theorem [6]. There are also important classification results for differential structures of compact manifolds with PIC via surgery method, we refer the reader to [27, 11, 12, 7], etc.

For the steady solitons with PIC, Brendle proved that any non-flat κ -noncollapsed, uniformly PIC steady gradient Ricci soliton of dimension 4 must be the Bryant soliton up to scaling [5]. Recently, Cho-Li generalized the Brendle's result for the steady solitons with $n \geq 12$ [13]. Thus our result, Corollary 0.3 gives a character for the steady solitons with PIC in particular for the dimension between 5 and 11. We would like to point out that just assuming the PIC instead of the uniformly PIC is not sufficient to prove the rigidity of steady gradient Ricci solitons, thanks for the new examples of Lai about κ -noncollapsed steady gradient Ricci solitons of dimension $n \geq 4$ with the positive curvature operator which are not rotationally symmetric [34].

Definition 0.4. *(M^n, g) is called $C^{2,\tau}$ -asymptotic to the n -dimensional Bryant Ricci soliton (\mathbb{R}^n, g_0) if there are a ball $B_R(0) \subset \mathbb{R}^n$ and a diffeomorphism*

$$F : (\mathbb{R}^n \setminus B_R, g_0) \rightarrow (M, g)$$

*such that $\hat{g} = F^*g$ satisfies*

$$(0.4) \quad |\hat{g} - g_0|_{g_0}(x) \leq O(\rho(x)^{-\tau-\frac{1}{2}}), \quad |\nabla^k \hat{g}|_{g_0} \leq O(\rho(x)^{-\tau-k}),$$

where $k = 1, 2, \tau \in (0, 1)$.

By Corollary 0.3, we can get the following rigidity theorem for the steady gradient Ricci solitons.

Corollary 0.5. *Any steady gradient Ricci soliton with the curvature condition (0.1), which is $C^{2,\tau}$ -asymptotic to the n -dimensional Bryant Ricci soliton (\mathbb{R}^n, g_0) , must be isometric to (\mathbb{R}^n, g_0) .*

In general, there is no rigidity for the steady soliton under the metric $C^{l,\tau}$ -asymptotic behavior if there is no curvature restriction. For examples, Conlon and Deruelle have constructed a complete steady gradient Kähler-Ricci soliton in every Kähler class of a crepant resolution over a Calabi-Yau cone, which unique up to the flow of the soliton vector field converges at a polynomial rate to the Cao's steady gradient Kähler-Ricci soliton on the cone [17].

It is interesting to mention that a similar version of Corollary 0.5 has been established for the shrinking gradient Ricci soliton [28] and the expanding gradient Ricci soliton [21, 22], respectively. We hope that the condition (0.1) can be weakened by the positive Ricci curvature. Actually, we have the following conjecture.

Conjecture 0.6. *Any steady gradient Ricci soliton with positive Ricci curvature, which is $C^{2,\tau}$ -asymptotic to the n -dimensional Bryant Ricci soliton (\mathbb{R}^n, g_0) , must be isometric to (\mathbb{R}^n, g_0) .*

To prove Theorem 0.2, we will follow an approach of Brendle to construct $\frac{n(n-1)}{2}$ Killing vector fields (VFs) on the steady Ricci soliton (M^n, g) as in the proof of Theorem 0.1 [5], where the positivity of sectional curvature plays a crucial role in his decay estimate for the solutions of Lichnerowicz equation via the maximal principle (also see Remark 4.5). However, in our case with the curvature condition (0.1), we use the heat flow method to do the decay estimate by solving the parabolic version of Lichnerowicz equation (cf. Section 5). Thus we need to show that the Lie derivatives of metric g associated to the constructed $\frac{n(n-1)}{2}$ approximate Killing VFs have fast polynomial decay (cf. Section 4).

We now outline the main steps in the proof of Theorem 0.2.

Step 1 (cf. Section 2, 3). As in [4, 5], we construct a collection of approximate Killing VFs U_a , $a \in \{1, \dots, \frac{n(n-1)}{2}\}$, near the infinity of (M, g) . By several improvements of asymptotic estimates for curvatures (cf. Subsection 2.2), we show that those U_a satisfy (cf. Proposition 3.1),

$$(0.5) \quad |\mathcal{L}_{U_a}(g)| \leq O(\rho^{-\frac{1}{2}-\frac{1}{4}\delta}) \text{ and } |\Delta U_a + D_X U_a| \leq O(\rho^{-1-\frac{1}{4}\delta}).$$

The above estimate allows us to get global solutions of the following elliptic equation associated to U_a on (M, g) ,

$$(0.6) \quad \Delta W_a + D_X W_a = 0.$$

The method is to solve the Dirichlet problem via perturbation and exhausting as in [5, Section 3].

Step 2 (cf. Section 4). We shall do the decay estimate of $h_a = \mathcal{L}_{W_a} g$ for solutions of Lichnerowicz-typed equation associated to the steady Ricci soliton (M, g) ,

$$(0.7) \quad \Delta_L h_a + \mathcal{L}_X(h_a) = 0.$$

We prove that there exists a constant λ such that $|\tilde{h}_a|$, where $\tilde{h}_a = h_a - \lambda \text{Ric}(g)$, decays to zero at any polynomial rate (cf. Theorem 4.3).

Step 3 (cf. Section 5). We prove that the uniqueness of solutions of (0.7) and \tilde{h}_a is in fact zero then we will finish the proof of Theorem 0.2. Our method is to consider the following parabolic Lichnerowicz equation for the initial value \tilde{h}_a constructed in Step 2,

$$(0.8) \quad \frac{\partial}{\partial t} h = \Delta_{L, g(t)} h,$$

where $g(t)$ is the Ricci flow solution induced by g via transformations group Φ_t^* generated by the soliton VF. By modifying the pinching estimate of Anderson-Chow for the solution of (0.8) in 3-dimension [1] (also see [35]), we introduce the following heat equation for any dimension $n \geq 3$,

$$(0.9) \quad \frac{\partial}{\partial t} H - \Delta_{g(t)} H - 2P(\cdot, t)H = 0,$$

where $P(\cdot, t) = \Phi_t^* P(\cdot)$ and $P(\cdot)$ is the function defined by (0.2). We will show that there exists a bounded positive super-solution of (0.9) (cf. Lemma 5.1), and then use the heat kernel method to prove that the limit solution of (0.8) is just zero (cf. Proposition 5.4).

Theorem 0.2, Corollary 0.3 and Corollary 0.5 will be finally proved in Section 6.

1. PRELIMINARY

A complete Riemannian metric g on M is called a gradient Ricci soliton if there exists a smooth function f (which is called a defining function) on M such that

$$(1.1) \quad R_{ij}(g) + \rho g_{ij} = \nabla_i \nabla_j f,$$

where $\rho \in \mathbb{R}$ is a constant. The gradient Ricci soliton is called expanding, steady and shrinking according to the sign $\rho >, =, < 0$, respectively. These three types of Ricci solitons correspond to three different blow-up solutions of Ricci flow [26].

In case of steady Ricci solitons, we can rewrite (1.1) as

$$(1.2) \quad 2 \operatorname{Ric}(g) = \mathcal{L}_X g,$$

where \mathcal{L}_X is the Lie operator along the gradient VF $X = \nabla f$ generalized by f . Let $\{\Phi_t^*\}_{t \in (-\infty, \infty)}$ be a 1-ps of transformations generated by $-X$. Then $g(t) = \Phi_t^*(g)$ ($t \in (-\infty, \infty)$) is a solution of Ricci flow. Namely, $g(t)$ satisfies

$$(1.3) \quad \frac{\partial g}{\partial t} = -2 \operatorname{Ric}(g), \quad g(0) = g.$$

For simplicity, we call $g(t)$ the soliton Ricci flow of (M, g) .

By (1.2), we have

$$\langle \nabla R, \nabla f \rangle = -2 \operatorname{Ric}(\nabla f, \nabla f),$$

where R is the scalar curvature of g . It follows

$$R + |\nabla f|^2 = \operatorname{Const}.$$

Since R is always positive ([36, 9]), the above equation can be normalized by

$$(1.4) \quad R + |\nabla f|^2 = 1.$$

In dimension $n = 2$, it is classified that the cigar is only solution [27, 16]. Brendle prove that the Bryant soliton is only 3-dimensional non-flat κ -noncollapsed steady gradient soliton up to scaling [4]. Here we call a steady gradient soliton as the Bryant Ricci soliton if it is rotational and is normalized by (1.4). In this paper, we always assume $n \geq 4$.

The following result has been proved in [9, 14].

Lemma 1.1. *Let (M, g) be a steady gradient soliton. Suppose that the scalar curvature decays uniformly and have positive Ricci curvature outside of a compact set $K \subset M$. Then there exists positive constants r_0, C_1, C_2 such that for all $x \in M$ with $\rho(x) \geq r_0$ it holds*

$$(1.5) \quad C_1 \rho(x) \leq f(x) \leq C_2 \rho(x),$$

where $\rho(x) = d(p, x)$.

Recall (cf. [5])

Definition 1.2. *We say (M, g) is asymptotically cylindrical if the following holds:*

(i) *The scalar curvature satisfies $\frac{C_1}{\rho(x)} \leq R(x) \leq \frac{C_2}{\rho(x)}$ as $\rho(x) \gg 1$, where C_1, C_2 are two positive constants.*

(ii) *Let p_m be an arbitrary sequence of marked points going to infinity. Consider the rescaled metrics*

$$\hat{g}^{(m)}(t) = r_m^{-1} \Phi_{r_m t}^*(g),$$

where $r_m R(p_m) = \frac{n-1}{2} + o(1)$ as $m \rightarrow \infty$, the flow $(M, \hat{g}^{(m)}(t), p_m)$ converges in the Cheeger-Gromov sense to a family of shrinking cylinders $(\mathbb{S}^{n-1} \times \mathbb{R}, \bar{g}(t))$, $t \in (0, 1)$. The metric $\bar{g}(t)$ is given by

$$(1.6) \quad \bar{g}(t) = (n-2)(2-2t)g_{\mathbb{S}^{n-1}(1)} + dr^2,$$

where $\mathbb{S}^{n-1}(1)$ is the unit sphere in the Euclidean space.

Under the condition (ii) in Definition 1.2, the following curvature behavior estimate for the steady gradient solitons has been proved in [19, Lemma 6.5, 6.6].

Lemma 1.3. *Let (M, g) be a noncompact steady Ricci soliton, which satisfies the condition (ii) in Definition 1.2. For $p \in M$, let $\{e_1, e_2, \dots, e_n\}$ be an orthonormal basis of $T_p M$ with respect to metric g , and $e_n = \frac{\nabla f}{|\nabla f|}$. Then*

$$\begin{aligned} \frac{|\text{Hess } R| + |\Delta \text{Ric}(g)|}{R^2} &\rightarrow 0, \quad \text{as } \rho(x) \rightarrow \infty, \\ \frac{\text{Ric}(e_i, e_j)}{R} &\rightarrow \frac{\delta_{ij}}{(n-1)}, \quad \text{as } \rho(x) \rightarrow \infty, \\ \frac{\text{Ric}(e_i, e_n)}{R} &\rightarrow 0, \quad \text{as } \rho(x) \rightarrow \infty, \\ \frac{\text{Rm}(e_j, e_k, e_k, e_l)}{R} &\rightarrow \frac{(1 - \delta_{jk})\delta_{jl}}{(n-1)(n-2)}, \quad \text{as } \rho(x) \rightarrow \infty, \\ \frac{\text{Rm}(e_n, e_i, e_j, e_k)}{R} &\rightarrow 0, \quad \text{as } \rho(x) \rightarrow \infty. \end{aligned}$$

In particular, there exists a compact set $K \subset M$ such that (M, g) has positive sectional curvature outside of K .

By the above lemma (also see [14, Section 3]), the first condition in Definition 1.2 can be deduced from the second one. Hence, we may say that (M, g) is asymptotically cylindrical with only assuming the second condition.

Lemma 1.3 also implies that (M, g) has positive sectional curvature outside of a compact set of M . Then by Lemma 1.1, the defining function f has a linear growth. Thus we may assume $f \geq 2$ on M .

Definition 1.4. *A Riemannian manifold (M^n, g) ($n \geq 4$) is said to have PIC, if for any $p \in M$, and any orthonormal 4-frame $\{e_1, e_2, e_3, e_4\}$ of $T_p M$, we have*

$$R_{1313} + R_{1414} + R_{2323} + R_{2424} - 2R_{1234} > 0.$$

Lemma 1.5. *If (M^n, g) , $n \geq 5$ has PIC, then the largest eigenvalue of Ricci tensor is bounded above by $\frac{1}{2}R$.*

Proof. By Definition 1.4, for any orthonormal 4-frame $\{e_i, e_j, e_k, e_l\}$, and $\{-e_i, e_j, e_k, e_l\}$, we have

$$R_{ikik} + R_{ilil} + R_{jkjk} + R_{jljl} - 2R_{ijkl} > 0$$

and

$$R_{ikik} + R_{ilil} + R_{jkjk} + R_{jljl} + 2R_{ijkl} > 0.$$

Combining these two relations together, we get

$$(1.7) \quad R_{ikik} + R_{ilil} + R_{jkjk} + R_{jljl} > 0.$$

By choosing an orthonormal n -frame $\{e_1, e_2, \dots, e_n\}$, we may assume that the largest eigenvalue of Ricci tensor is equal to Ric_{nn} . Thus by (1.7) it is easy to see that as $n \geq 5$ it holds

$$R - 2\text{Ric}_{nn} = \sum_{k,l=1}^{n-1} R_{klkl} > 0.$$

□

Proposition 1.6. *If (M^n, g) is an n -dimensional Riemannian manifold. Then the PIC implies (0.1) when $n \geq 5$, and the positive sectional curvature implies (0.1) when $n \geq 4$.*

Proof. Recall that $P = \sup_h \frac{Rm(h,h)}{|h|^2}$ for any symmetric 2-tensor $h \neq 0$. Then at each point in M , we can choose local coordinates such that h is diagonal and have

$$P = \sup_h \frac{\sum_{i,j=1}^n R_{ijij} h_{ii} h_{jj}}{\sum_{i=1}^n h_{ii}^2}.$$

Denote $(M_{ij}) = (R_{ijij})$ as a matrix, which is symmetric and whose diagonal elements are all 0. Thus P is bounded by the norm of the largest eigenvalue of (M_{ij}) . By the Gerschgorin circle theorem,

$$(1.8) \quad P \leq \max_i \sum_{j \neq i} M_{ij} = \max_i \text{Ric}_{ii}.$$

If (M, g) has PIC, by Lemma 1.5, we see

$$(1.9) \quad R - 2P \geq R - 2 \text{Ric}_{ii} = \sum_{j,l \neq i} R_{jllj} > 0,$$

when $n \geq 5$. If (M, g) has positive sectional curvature, (1.9) obviously holds for $n \geq 4$. Hence the proposition is true. \square

2. FINE ESTIMATES OF CURVATURE BEHAVIOR

In this section, we modify the argument in [5, Section 2] to improve the estimate for the curvature behavior around level sets of the steady Ricci soliton with asymptotically cylindrical property. The result will be used to construct the solutions of elliptic equation (0.6) for VFs in Section 3.

For a fixed point $p \in M$, we choose an orthonormal frame $\{e_1, e_2, \dots, e_n\}$ with $e_n = \frac{\nabla f}{|\nabla f|}$. By [5, Proposition 2.1] (also see Lemma 1.3), we have

$$(2.1) \quad \text{Ric}(e_i, e_j) = \frac{1}{n-1} R \delta_{ij} + o(r^{-1}), \quad i, j \in \{1, 2, \dots, n-1\}.$$

and

$$(2.2) \quad 2 \text{Ric}(e_i, X) = -\langle e_i, \nabla R \rangle = o(r^{-\frac{3}{2}}).$$

Moreover, we have

$$(2.3) \quad |\text{Ric}|^2 - \frac{1}{n-1} R^2 = O(r^{-\frac{5}{2}})$$

and

$$(2.4) \quad 2 \text{Ric}(X, X) = -\langle X, \nabla R \rangle = \Delta R + 2|\text{Ric}|^2 = \frac{2}{n-1} R^2 + o(r^{-2}).$$

Let

$$(2.5) \quad T = (n-1) \text{Ric}(g) - Rg + Rdf \otimes df$$

be defined as in [5, Section 2]. It has been prove by Brendle

Lemma 2.1. *If (M, g) is asymptotically cylindrical, it holds*

$$(2.6) \quad |T| \leq O(r^{-\frac{3}{2}}), \quad |\nabla T| \leq O(r^{-\frac{3}{2}-16\delta}).$$

As a consequence,

$$(2.7) \quad |\nabla R| \leq O(r^{-\frac{3}{2}-16\delta}), \quad |\Delta R| \leq O(r^{-2-8\delta})$$

and

$$(2.8) \quad |\nabla \text{Ric}(g)| \leq O(r^{-\frac{3}{2}-16\delta}), \quad |D^2 \text{Ric}(g)| \leq O(r^{-2-8\delta}), \quad |D^3 \text{Ric}(g)| \leq O(r^{-\frac{5}{2}-4\delta}).$$

The following roundness estimate has been also proved by Brendle.

Lemma 2.2.

$$(2.9) \quad fR = \frac{n-1}{2} + O(r^{-8\delta}),$$

$$(2.10) \quad f \operatorname{Ric}(g) \leq \left(\frac{1}{2} + o(1)\right)g, \quad f^2 \operatorname{Ric}(g) \geq cg, \quad c > 0,$$

and

$$(2.11) \quad \begin{aligned} fR_{ijkl} &= \frac{1}{2(n-2)} (g_{ik} - \partial_i f \partial_k f) (g_{jk} - \partial_j f \partial_k f) \\ &\quad - \frac{1}{2(n-2)} (g_{il} - \partial_i f \partial_l f) (g_{jk} - \partial_j f \partial_k f) \\ &\quad + O(r^{-8\delta}). \end{aligned}$$

2.1. $\frac{1}{2}$ -order improvement. In this subsection, we will improve the estimates in Lemma 2.1 and Lemma 2.2.

First we prove

Lemma 2.3.

$$(2.12) \quad |D_X T| \leq O(r^{-2-8\delta}) \text{ and } |D_{X,X}^2 T| \leq O(r^{-\frac{5}{2}-8\delta}).$$

Proof. By (2.8), we have

$$(2.13) \quad R_{ijkl} \partial^l f = D_i \operatorname{Ric}_{jk} - D_j \operatorname{Ric}_{ik} = O\left(r^{-\frac{3}{2}-16\delta}\right).$$

Then by (2.7) and (2.8), we get

$$\begin{aligned} &-D_X T_{ik} \\ &= (n-1)\Delta \operatorname{Ric}_{ik} - \Delta R(g_{ik} - \partial_i f \partial_k f) + 2(n-1)R_{ijkl} \operatorname{Ric}^{jl} \\ &\quad - 2|\operatorname{Ric}|^2(g_{ik} - \partial_i f \partial_k f) - R \operatorname{Ric}(X, e_i) \partial_k f - R \operatorname{Ric}(X, e_k) \partial_i f \\ &= 2(n-1)R_{ijkl} \operatorname{Ric}^{jl} - 2|\operatorname{Ric}|^2(g_{ik} - \partial_i f \partial_k f) + O(r^{-2-8\delta}). \end{aligned}$$

By (1.2), it follows

$$\begin{aligned} &-D_X T_{ik} \\ &= \frac{2}{(n-1)(n-2)} R^2(n-2|\nabla f|^2 + |\nabla f|^4)(g_{ik} - \partial_i f \partial_k f) \\ &\quad - \frac{2}{(n-1)(n-2)} R^2(g_{ik} - \partial_i f \partial_k f) + \frac{2}{(n-1)(n-2)} R^2(2-3|\nabla f|^2 + |\nabla f|^4) \partial_i f \partial_k f \\ &\quad - \frac{2}{(n-1)^2} R^2(n-2|\nabla f|^2 + |\nabla f|^4)(g_{ik} - \partial_i f \partial_k f) + O(r^{-2-8\delta}) \\ &= \frac{2}{(n-1)(n-2)} R^2((n-2) + O(r^{-1}))(g_{ik} - \partial_i f \partial_k f) \\ &\quad - \frac{2}{(n-1)^2} R^2((n-1) + O(r^{-1}))(g_{ik} - \partial_i f \partial_k f) + O(r^{-2-8\delta}) \\ &= O(r^{-2-8\delta}). \end{aligned}$$

Thus the first relation of (2.12) is true.

For the second relation, we use the fact

$$-D_{X,X}^2 T_{ik} - \langle D_X X, \nabla T_{ik} \rangle = \langle X, \nabla(-D_X T_{ik}) \rangle.$$

By (2.6), we see

$$(2.14) \quad \langle D_X X, \nabla T_{ik} \rangle = \operatorname{Ric}(X, e_j) \nabla_j T_{ik} = O(r^{-3}).$$

On the other hand,

$$\begin{aligned}
& \langle X, \nabla(-D_X T_{ik}) \rangle \\
&= (n-1)D_X \Delta \text{Ric}_{ik} - D_X \Delta R(g_{ik} - \partial_i f \partial_k f) \\
&+ (\Delta R + 2|\text{Ric}|^2) \text{Ric}(X, e_i) \partial_k f + (\Delta R + 2|\text{Ric}|^2) \text{Ric}(X, e_k) \partial_i f \\
&+ 2(n-1)D_X R_{ijkl} \text{Ric}^{jl} + 2(n-1)R_{ijkl} D_X \text{Ric}^{jl} \\
&- 4D_X \text{Ric}_{jl} \text{Ric}^{jl}(g_{ik} - \partial_i f \partial_k f) \\
&- D_X R(\text{Ric}(X, e_i) \partial_k f + \text{Ric}(X, e_k) \partial_i f) \\
&- R(D_X \text{Ric}(X, e_i) \partial_k f + D_X \text{Ric}(X, e_k) \partial_i f) \\
&- 2R \text{Ric}(X, e_i) \text{Ric}(X, e_k).
\end{aligned}$$

Then by (2.7) and (2.8), it follows

$$\begin{aligned}
& \langle X, \nabla(-D_X T_{ik}) \rangle \\
&= (n-1)D_X \Delta \text{Ric}_{ik} - D_X \Delta R(g_{ik} - \partial_i f \partial_k f) + O(r^{-\frac{5}{2}-8\delta}) \\
&= (n-1)\Delta D_X \text{Ric}_{ik} - \Delta D_X R(g_{ik} - \partial_i f \partial_k f) + Rm * (\nabla \text{Ric} + \nabla R) \\
&+ (n-1)\nabla_j (R_{lji}^m \text{Ric}_{mk} \partial^l f + R_{ljk}^m \text{Ric}_{im} \partial^l f) + O(r^{-\frac{5}{2}-8\delta}) \\
&= O(r^{-\frac{5}{2}-8\delta}).
\end{aligned}$$

Thus combining this with (2.14), we get the second relation of (2.12). □

Next we improve the C^0 -norm of T -tensor by the above lemma.

Proposition 2.4.

$$(2.15) \quad |T| \leq O(r^{-\frac{3}{2}-4\delta}).$$

Proof. We use the parabolic maximum principle to get the estimate (2.15) of T as in [5, Section 2]. We note that by the Ricci equation (1.3) the Ricci tensor satisfies the equation

$$\Delta \text{Ric}_{ik} + D_X \text{Ric}_{ik} = -2 \sum_{j,l=1}^n R_{ijkl} \text{Ric}^{jl}$$

Then by the identity

$$\Delta X + D_X X = 0,$$

we obtain

$$\begin{aligned}
& \Delta (Rg_{ik} - R\partial_i f \partial_k f) + D_X (Rg_{ik} - R\partial_i f \partial_k f) \\
&= (\Delta R + \langle X, \nabla R \rangle) (g_{ik} - \partial_i f \partial_k f) - 2 \text{Ric}_{ij} \partial_j R \partial_k f - 2 \text{Ric}_{jk} \partial_j R \partial_i f \\
(2.16) \quad & - 2R \text{Ric}_{ij} \text{Ric}_{jk} - R\Delta \partial_i f \partial_k f - R\Delta \partial_k f \partial_i f - RD_X \partial_i f \partial_k f - RD_X \partial_k f \partial_i f \\
&= -2|\text{Ric}|^2 (g_{ik} - \partial_i f \partial_k f) + O\left(r^{-\frac{5}{2}-16\delta}\right).
\end{aligned}$$

In the last relation we used (2.13).

By (2.1) and (2.2) together with (2.13), we have

$$\Delta \partial_i f \partial_k f = \partial_i \Delta f + R_{jijl} \partial^l f = \partial_i R + R_{jijl} \partial^l f \leq O(r^{-\frac{3}{2}-16\delta}).$$

Then

$$\begin{aligned}
& \Delta T_{ik} + D_X T_{ik} \\
&= -2 \sum_{j,l=1}^{n-1} R_{ijkl} T^{jl} - 2 \sum_{j=1}^n R_{ijkn} \text{Ric}^{jn} - 2 \sum_{l=1}^n R_{inlk} \text{Ric}^{nl} + 2R_{inkn} \text{Ric}^{nn} \\
&- 2R \text{Ric}_{ik} + 2RR_{inkn} + 2 \sum_{j,l=1}^n R_{ijkl} R \partial^j f \partial^l f + 2|\text{Ric}|^2 (g_{ik} - \partial_i f \partial_k f) + O\left(r^{-\frac{5}{2}-16\delta}\right).
\end{aligned}$$

By (2.13), it follows

$$\begin{aligned} & \Delta T_{ik} + D_X T_{ik} \\ &= -2 \sum_{j,l=1}^{n-1} R_{ijkl} T^{jl} - 2R \text{Ric}_{ik} \\ &+ 2|\text{Ric}|^2 (g_{ik} - \partial_i f \partial_k f) + O\left(r^{-\frac{5}{2}-16\delta}\right). \end{aligned}$$

Hence,

$$\begin{aligned} & \Delta (|T|^2) + \langle X, \nabla (|T|^2) \rangle \\ &= 2|DT|^2 - 4 \sum_{j,l=1}^{n-1} R_{ijkl} T^{ik} T^{jl} - 4R \sum_{i,k=1}^n \text{Ric}_{ik} T^{ik} \\ &+ 4|\text{Ric}|^2 \sum_{i,k=1}^n (g_{ik} - \partial_i f \partial_k f) T^{ik} + O\left(r^{-\frac{5}{2}-16\delta}\right) |T| \\ &= 2|DT|^2 - 4 \sum_{j,l=1}^{n-1} R_{ijkl} T^{ik} T^{jl} - \frac{4}{n-1} R |T|^2 \\ &+ 4 \left(|\text{Ric}|^2 - \frac{1}{n-1} R^2 \right) \sum_{i,k=1}^n (g_{ik} - \partial_i f \partial_k f) T^{ik} + O\left(r^{-\frac{5}{2}-16\delta}\right) |T|. \end{aligned}$$

Note

$$|X|^2 = |\nabla f|^2 = 1 - R = 1 + O(r^{-1}).$$

Then

$$\begin{aligned} & \sum_{i,k=1}^n (g_{ik} - \partial_i f \partial_k f) T^{ik} \\ &= -R + 2R|\nabla f|^2 - R|\nabla f|^4 - (n-1) \text{Ric}(X, X) \\ &= -R^3 - (n-1) \text{Ric}(X, X) = O(r^{-2}). \end{aligned}$$

Therefore, by (2.3), we get

$$(2.17) \quad \begin{aligned} & \Delta (|T|^2) + \langle X, \nabla (|T|^2) \rangle \\ & \geq -4 \sum_{j,l=1}^{n-1} R_{ijkl} T^{ik} T^{jl} - \frac{4}{n-1} R |T|^2 - O\left(r^{-\frac{5}{2}-16\delta}\right) |T| - O\left(r^{-\frac{9}{2}}\right). \end{aligned}$$

On the other hand, by (2.11), we have

$$(2.18) \quad \begin{aligned} R_{ijkl} &= \frac{1}{(n-1)(n-2)} R (g_{ik} - \partial_i f \partial_k f) (g_{jl} - \partial_j f \partial_l f) \\ &- \frac{1}{(n-1)(n-2)} R (g_{il} - \partial_i f \partial_l f) (g_{jk} - \partial_j f \partial_k f) \\ &+ O\left(r^{-1-8\delta}\right). \end{aligned}$$

Moreover, we can estimate

$$\begin{aligned} \text{tr}(T) &= -R^2 = O\left(r^{-2}\right), \\ T(\nabla f, \cdot) &= (n-1) \text{Ric}(\nabla f, \cdot) - R^2 \nabla f = O\left(r^{-\frac{3}{2}}\right), \\ T(\nabla f, \nabla f) &= (n-1) \text{Ric}(\nabla f, \nabla f) - R^2 |\nabla f|^2 = O\left(r^{-2}\right). \end{aligned}$$

Thus

$$\begin{aligned}
& \sum_{j,l=1}^{n-1} R_{ijkl} T^{ik} T^{jl} \\
&= \frac{1}{(n-1)(n-2)} R(-R^3 - (n-1) \operatorname{Ric}(X, X)) (g_{jl} - \partial_j f \partial_l f) T^{jl} \\
&\quad - \frac{1}{(n-1)(n-2)} R |T|^2 + \frac{1}{(n-1)(n-2)} R((n-1) \operatorname{Ric}(X, X) - R + R|\nabla f|^4)^2 \\
&\quad + \frac{1}{(n-1)(n-2)} R((n-1)(\operatorname{Ric}(X, e_j) \partial_l f + \operatorname{Ric}(X, e_l) \partial_j f) - 2R^2 \partial_j f \partial_l f) T^{jl} \\
&\quad - \frac{1}{(n-1)(n-2)} R((n-1) \operatorname{Ric}(X, X) \partial_j f \partial_l f - R^2 |\nabla f|^2 \partial_j f \partial_l f) T^{jl} \\
&\quad + o(r^{-1}) |T|^2 \\
(2.19) \quad &= -\frac{1}{(n-1)(n-2)} R |T|^2 + O\left(r^{-\frac{5}{2}-8\delta}\right) |T| + o(r^{-1}) |T|^2 + O(r^{-5}).
\end{aligned}$$

Hence, inserting (2.19) into (2.17), we obtain

$$\begin{aligned}
& \Delta(|T|^2) + \langle X, \nabla(|T|^2) \rangle \\
(2.20) \quad & \geq -\frac{4(n-3)}{(n-1)(n-2)} R |T|^2 - o(r^{-1}) |T|^2 - O\left(r^{-\frac{5}{2}-8\delta}\right) |T| - O(r^{-4-8\delta})
\end{aligned}$$

Let Σ be the level surface of $\{f = r\}$ on (M, g) . Since the mean curvature H with respect to the metric g on Σ satisfies

$$H = \frac{1}{|\nabla f|} R - \frac{1}{|\nabla f|^3} \operatorname{Ric}(X, X) = O(r^{-1}),$$

by Lemma 2.3, we can rewrite (2.20) as

$$\begin{aligned}
& \Delta_{\Sigma}(|T|^2) + \langle X, \nabla(|T|^2) \rangle \\
&= \Delta(|T|^2) + \langle X, \nabla(|T|^2) \rangle - D_{X,X}^2(|T|^2) - H \langle X, \nabla(|T|^2) \rangle \\
&\geq -\frac{2(n-3)}{n-2} f^{-1} |T|^2 - o(r^{-1}) |T|^2 - O\left(r^{-\frac{5}{2}-8\delta}\right) |T| - O(r^{-4-8\delta}),
\end{aligned}$$

where Δ_{Σ} denotes the Laplacian operator with respect to the induced metric on Σ . Furthermore, by the fact $f = \text{const.}$ on Σ , we can estimate

$$\begin{aligned}
& \Delta_{\Sigma}(f^2 |T|^2) + \langle X, \nabla(f^2 |T|^2) \rangle \\
&= f^2 \Delta_{\Sigma}(|T|^2) + f^2 \langle X, \nabla(|T|^2) \rangle + 2f |\nabla f|^2 |T|^2 \\
&\geq -\frac{2(n-3)}{n-2} f |T|^2 + 2f(1 - O(r)) |T|^2 - o(r) |T|^2 - O\left(r^{-\frac{1}{2}-8\delta}\right) |T| - O(r^{-2-8\delta}) \\
&\geq \frac{2}{n-2} f |T|^2 - o(r) |T|^2 - O\left(r^{-\frac{1}{2}-8\delta}\right) |T| - O(r^{-2-8\delta}) \geq -O(r^{-2-8\delta}).
\end{aligned}$$

Note that $f^2 |T|^2 \rightarrow 0$ at infinity. Hence by the parabolic maximum principle, we conclude

$$f^2 |T|^2 \leq O(r^{-1-8\delta}),$$

which implies (2.4). □

By Proposition 2.4, we see

$$|T| \leq O(r^{-\frac{3}{2}-4\delta}).$$

By Shi's estimates for higher order derivatives of curvature [33], it follows

$$(2.21) \quad |D^m T| \leq O(r^{\frac{m+2}{2}}), \quad \forall m \geq 1.$$

Thus by the standard interpolation inequalities, we get

$$(2.22) \quad |DT| \leq O(r^{-2-2\delta}) \text{ and } |D^2T| \leq O(r^{-\frac{5}{2}-\delta}).$$

On the other hand, by the relation

$$\begin{aligned} D^i D^k T_{ik} &= \frac{n-3}{2} \Delta R + D_X D_X R + \frac{7}{2} R D_X R + R^3 \\ &\quad + \partial_i R \text{Ric}_i^k \partial_k f + R |\text{Ric}(g)|^2 \\ &= \frac{n-3}{2} \Delta R + D_X D_X R + O\left(r^{-\frac{5}{2}-16\delta}\right) \\ &= \frac{n-3}{2} \Delta R - D_X (\Delta R + 2 |\text{Ric}(g)|^2) + O\left(r^{-\frac{5}{2}-16\delta}\right) \\ &= \frac{n-3}{2} \Delta R - \Delta(D_X R) - R_{mii}^l \partial_l R \partial^m f + O\left(r^{-\frac{5}{2}-16\delta}\right) \end{aligned}$$

and

$$D_X R = -\Delta R + |\text{Ric}(g)|^2,$$

we know from (2.13),

$$D^i D^k T_{ik} = \frac{n-3}{2} \Delta R + O\left(r^{-\frac{5}{2}-16\delta}\right).$$

Hence by (2.22), we improve the Lapalace estimate of scalar curvature in (2.7) by

$$(2.23) \quad |\Delta R| \leq O(r^{-\frac{5}{2}-\delta}).$$

By (2.23), (2.9) in Lemma 2.2 can be improved as follows.

Proposition 2.5.

$$(2.24) \quad fR = \frac{n-1}{2} + O(r^{-\frac{1}{2}-\delta}).$$

Proof. By Proposition 2.4, we have

$$|\text{Ric}(g)| = \frac{1}{n-1} R |g - df \otimes df| + O\left(r^{-\frac{3}{2}-4\delta}\right) = \frac{1}{\sqrt{n-1}} R + O\left(r^{-\frac{3}{2}-4\delta}\right),$$

and so

$$|\text{Ric}(g)|^2 = \frac{1}{n-1} R^2 + O(r^{-\frac{5}{2}-4\delta}).$$

By (2.23), it follows

$$-\langle X, \nabla R \rangle = \Delta R + 2 |\text{Ric}(g)|^2 = \frac{2}{n-1} R^2 + O\left(r^{-\frac{5}{2}-\delta}\right).$$

Thus

$$\left\langle X, \nabla \left(\frac{1}{R} - \frac{2}{n-1} f \right) \right\rangle = O\left(r^{-\frac{1}{2}-\delta}\right).$$

Integrating the above relation along the integral curves of X , we obtain

$$\frac{1}{R} - \frac{2}{n-1} f = O\left(r^{\frac{1}{2}-\delta}\right).$$

This implies (2.24). □

Also we prove

Lemma 2.6.

$$(2.25) \quad \begin{aligned} i). & \quad |\nabla_{e_i} e_j| \leq O(r^{-\frac{1}{2}-8\delta}); \\ ii). & \quad |D_{i,j}^2 R| \leq O(r^{-\frac{5}{2}-\delta}); \\ iii). & \quad |D_{i,j}^2 \text{Ric}(g)| \leq O(r^{-\frac{5}{2}-\delta}). \end{aligned}$$

Proof. i). By (2.8) and (2.13), we have

$$\begin{aligned}
D_X \nabla_{e_i} e_j &= D_i D_j \nabla f + R_{kij}^l \partial^k f \\
&= D_i \operatorname{Ric}(e_j, \cdot) + R_{kij}^l \partial^k f \\
(2.26) \qquad &= O(r^{-\frac{3}{2}-8\delta}).
\end{aligned}$$

Then

$$|D_X \nabla_{e_i} e_j| \leq O(r^{-\frac{3}{2}-8\delta}).$$

Since $|\nabla_{e_i} e_j| \rightarrow 0$ at infinity, by integrating (2.26) along the integral curves of X , we get the first relation in (2.25) immediately.

ii). By definition, we have

$$(2.27) \qquad D_{i,j}^2 R = D_i D_j R - D_{\nabla_{e_i} e_j} R.$$

Since

$$\begin{aligned}
D^k T_{ik} &= \frac{n-3}{2} \partial_i R + \langle \nabla f, \nabla R \rangle \partial_i f + R^2 \partial_i f + R \operatorname{Ric}_i^k \partial_k f \\
&= \frac{n-3}{2} \partial_i R + (-\Delta R + |\operatorname{Ric}(g)|^2) \partial_i f + O(r^{-2}) \\
&= \frac{n-3}{2} \partial_i R + O(r^{-2}),
\end{aligned}$$

By (2.22), we get

$$(2.28) \qquad |DR| = O(r^{-2}),$$

which is also an improvement of the first relation in (2.7). Thus by the proved relation in i), we obtain

$$(2.29) \qquad |D_{\nabla_{e_i} e_j} R| \leq O(r^{-\frac{5}{2}-8\delta}).$$

On the other hand, we also have

$$(2.30) \qquad D^k T_{ik} = \frac{n-3}{2} \partial_i R - \Delta R \partial_i f - 2|\operatorname{Ric}|^2 \partial_i f + R^2 \partial_i f + R \operatorname{Ric}_i^k \partial_k f.$$

Then

$$\begin{aligned}
D_j D^k T_{ik} &= \frac{n-3}{2} \partial_j \partial_i R - \partial_j \Delta R \partial_i f - \Delta R \operatorname{Ric}_{ij} - 4D_j \operatorname{Ric}_{kl} \operatorname{Ric}^{kl} \partial_i f - 2|\operatorname{Ric}|^2 \operatorname{Ric}_{ij} \\
(2.31) \qquad &+ 2R \partial_j R \partial_i f + R^2 \operatorname{Ric}_{ij} + \partial_j R \operatorname{Ric}_i^k \partial_k f + R \partial_j \operatorname{Ric}_i^k \partial_k f + R \operatorname{Ric}_i^k \operatorname{Ric}_{jk}.
\end{aligned}$$

Thus by (2.22), we get

$$|D_i D_j R| \leq O(r^{-\frac{5}{2}-\delta}).$$

Hence, inserting this relation together with (2.29) into (2.27), we prove the second relation in (2.25).

iii). By (2.5), we have

$$(n-1) D_{i,j}^2 \operatorname{Ric}_{kl} = D_{i,j}^2 (T_{kl} + R(g_{kl} - \partial_k f \partial_l f)).$$

Note

$$\begin{aligned}
&D_{i,j}^2 (R(g_{kl} - \partial_k f \partial_l f)) \\
&= D_i D_j (R(g_{kl} - \partial_k f \partial_l f)) - D_{\nabla_{e_i} e_j} (R(g_{kl} - \partial_k f \partial_l f)) \\
&= D_{i,j}^2 R(g_{kl} - \partial_k f \partial_l f) - \partial_j R \operatorname{Ric}_{ik} \partial_l f - \partial_j R \operatorname{Ric}_{il} \partial_k f \\
&\quad - \partial_i R \operatorname{Ric}_{jk} \partial_l f - \partial_i R \operatorname{Ric}_{jl} \partial_k f - R \partial_i \operatorname{Ric}_{jk} \partial_l f - R \partial_i \operatorname{Ric}_{jl} \partial_k f \\
&\quad - R \operatorname{Ric}_{jk} \operatorname{Ric}_{il} - R \operatorname{Ric}_{jl} \operatorname{Ric}_{ik} \\
&\quad + R \operatorname{Ric}(\nabla_{e_i} e_j, e_k) \partial_l f + R \operatorname{Ric}(\nabla_{e_i} e_j, e_l) \partial_k f.
\end{aligned}$$

Then by the estimate proved in ii), we see

$$D_{i,j}^2(R(g_{kl} - \partial_k f \partial_l f)) \leq O(r^{-\frac{5}{2}-\delta}).$$

Thus combining (2.22), we also get the third relation in (2.25). \square

At last, we improve (2.11) in Lemma 2.2 as follows.

Proposition 2.7.

$$(2.32) \quad \begin{aligned} fR_{ijkl} &= \frac{1}{2(n-2)} (g_{ik} - \partial_i f \partial_k f) (g_{ik} - \partial_i f \partial_k f) \\ &\quad - \frac{1}{2(n-2)} (g_{il} - \partial_i f \partial_l f) (g_{jk} - \partial_j f \partial_k f) \\ &\quad + O(r^{-\frac{1}{2}-\delta}). \end{aligned}$$

Proof. By [8, Proposition 2.10], we have

$$\begin{aligned} -D_X R_{ijkl} &= D_{i,k}^2 \text{Ric}_{jl} - D_{i,l}^2 \text{Ric}_{jk} - D_{j,k}^2 \text{Ric}_{il} + D_{j,l}^2 \text{Ric}_{ik} \\ &\quad + \sum_{m=1}^n \text{Ric}_i^m R_{mjkl} + \sum_{m=1}^n \text{Ric}_j^m R_{imkl}. \end{aligned}$$

On the other hand, by (2.11),

$$\begin{aligned} \sum_{m=1}^n \text{Ric}_i^m R_{mjkl} &= \frac{1}{n-1} R \sum_{m=1}^n (\delta_i^m - \partial_i f \partial^m f) R_{mjkl} + O(r^{-\frac{5}{2}-4\delta}) \\ &= \frac{1}{n-1} R R_{ijkl} + O(r^{-\frac{5}{2}-4\delta}). \end{aligned}$$

Then, by Lemma 2.6, we get

$$\begin{aligned} -D_X R_{ijkl} &= \frac{2}{n-1} R R_{ijkl} + O(r^{-\frac{5}{2}-4\delta}) \\ &= f^{-1} R_{ijkl} + O(r^{-\frac{5}{2}-\delta}). \end{aligned}$$

It follows

$$(2.33) \quad |D_X(fR_{ijkl})| \leq O(r^{-\frac{3}{2}-\delta}).$$

Let

$$\begin{aligned} S_{ijkl} &= \frac{1}{2(n-2)} (g_{ik} - \partial_i f \partial_k f) (g_{jl} - \partial_j f \partial_l f) \\ &\quad - \frac{1}{2(n-2)} (g_{il} - \partial_i f \partial_l f) (g_{jk} - \partial_j f \partial_k f). \end{aligned}$$

It is easy to see

$$|D_X S_{ijkl}| \leq C |\text{Ric}(X, \cdot)|.$$

Note that by (2.2) and (2.7) it holds

$$\text{Ric}(X, \cdot) \leq O(r^{-\frac{3}{2}-8\delta}).$$

Then

$$(2.34) \quad |D_X S_{ijkl}| \leq O(r^{-\frac{3}{2}-8\delta}).$$

Hence, combining (2.33) and (2.34), we obtain

$$(2.35) \quad |D_X(fR_{ijkl} - S_{ijkl})| \leq O(r^{-\frac{3}{2}-\delta}).$$

Since $|fR_{ijkl} - S_{ijkl}| \rightarrow 0$ at infinity, by the integration along the integral curves of X , we deduce from (2.35),

$$|fR_{ijkl} - S_{ijkl}| \leq O(r^{-\frac{1}{2}-\delta}),$$

which implies (2.32). \square

2.2. Fine roundness of level sets. By the asymptotically cylindrical property, each level surface $\Sigma_r \subset M$ of f is diffeomorphic to S^{n-1} . Then there is a family of diffeomorphisms

$$F_r : S^{n-1} \rightarrow \Sigma_r$$

such that $\frac{\partial}{\partial r} F_r = \frac{X}{|X|^2}$. Thus

$$\gamma_r = \frac{1}{2(n-2)r} F_r^*(g)$$

defines a family of metrics on S^{n-1} .

By Proposition 2.7, we have

Proposition 2.8. *For each $l \geq 0$, it holds*

$$(2.36) \quad \left\| \frac{d}{dr} \gamma_r \right\|_{C^l(S^{n-1}, \gamma_r)} \leq O\left(r^{-\frac{3}{2}-\frac{1}{2}\delta}\right).$$

Proof. Let

$$\bar{R} = \bar{R}_{ijkl} dx^i \otimes dx^j \otimes dx^k \otimes dx^l$$

be the curvature tensor of induced metric $g|_{\Sigma_r}$ on Σ_r . Then by the Gauss formula,

$$\bar{R}_{ijkl} - R_{ijkl} = \frac{1}{|\nabla f|^2} (\text{Ric}_{ik} \text{Ric}_{jl} - \text{Ric}_{il} \text{Ric}_{jk}).$$

By Proposition 2.7, it follows

$$(2.37) \quad \begin{aligned} \bar{R}_{ijkl} &= \frac{1}{2(n-2)r} (g_{ik} - \partial_i f \partial_k f) (g_{jk} - \partial_j f \partial_k f) \\ &\quad - \frac{1}{2(n-2)r} (g_{il} - \partial_i f \partial_l f) (g_{jk} - \partial_j f \partial_k f) \\ &\quad + O(r^{-\frac{3}{2}-\delta}). \end{aligned}$$

Since the normal velocity of the flow is given by

$$\frac{1}{|X|} = 1 + O(r^{-1}),$$

we have (cf. [18, Section 3]),

$$\frac{d}{dr} F_r^*(g) = F_r^* \left(\frac{2}{|\nabla f|^2} \text{Ric}(g) \right).$$

Hence by (2.37), we get

$$\sup_{S^{n-1}} \left| \frac{d}{dr} F_r^*(g) - \frac{1}{r} F_r^*(g) \right|_{F_r^*(g)} \leq O\left(r^{-\frac{3}{2}-\delta}\right).$$

Note

$$\frac{d}{dr} \gamma_r = \frac{1}{r} \left(\frac{d}{dr} F_r^*(g) - \frac{1}{r} F_r^*(g) \right).$$

Therefore, we obtain

$$(2.38) \quad \sup_{S^{n-1}} \left| \frac{d}{dr} \gamma_r \right|_{\gamma_r} \leq O\left(r^{-\frac{3}{2}-\delta}\right).$$

By (2.37) we see that the manifold (S^{n-1}, γ_r) has uniformly bounded curvature and all the derivatives of the curvature are also uniformly bounded. Thus

$$\left\| \frac{1}{r} \gamma_r \right\|_{C^l(S^{n-1}, \gamma_r)} \leq O(r^{-1}).$$

On the other hand, by Shi's estimate

$$\sup_{\{f=r\}} |D^l \text{Ric}| \leq O\left(r^{-\frac{l+2}{2}}\right),$$

which is equal to

$$\sup_{\{r-\sqrt{r} \leq f \leq r+\sqrt{r}\}} \left| D^l \left(\mathcal{L}_{\frac{X}{|X|^2}}(g) \right) \right| \leq O\left(r^{-\frac{l+2}{2}}\right),$$

we have

$$\left\| F_r^* \left(\frac{\mathcal{L}_{\frac{X}{|X|^2}}(g)}{|X|^2} \right) \right\|_{C^l(S^{n-1}, \gamma_r)} \leq O(1).$$

Since

$$\frac{d}{dr} \gamma_r + \frac{1}{r} \gamma_r = \frac{1}{2(n-2)r} F_r^* \left(\frac{\mathcal{L}_{\frac{X}{|X|^2}}(g)}{|X|^2} \right),$$

we get

$$(2.39) \quad \left\| \frac{d}{dr} \gamma_r \right\|_{C^l(S^{n-1}, \gamma_r)} \leq O(r^{-1})$$

for each $l \geq 0$. Now we can obtain (2.36) by (2.38) and (2.39) via the standard interpolation inequalities. \square

Since the metrics γ_r converge in C^∞ to a smooth metric $\bar{\gamma}$ as $r \rightarrow \infty$. By (2.37), the limit metric $\bar{\gamma}$ must be the standard sphere metric with constant sectional curvature 1. Moreover, by Proposition 2.8, we get

$$(2.40) \quad \|\gamma_r - \bar{\gamma}\|_{C^l(S^{n-1}, \bar{\gamma})} \leq O\left(r^{-\frac{1}{2} - \frac{1}{2}\delta}\right), \text{ for each } l \geq 0.$$

3. SOLUTIONS OF (0.6)

In this section, we construct a collection of VFs W_a , $a \in \{1, \dots, \frac{n(n-1)}{2}\}$, which solve (0.6) with property

$$(3.1) \quad |W_a| = O(r^{\frac{1}{2}}) \text{ and } |DW_a| \leq O(r^{-\frac{1}{2}}).$$

As in [5], we let $\bar{U}_a, a \in \{1, \dots, \frac{n(n-1)}{2}\}$, be the Killing VFs on $(S^{n-1}, \bar{\gamma})$ such that

$$(3.2) \quad \sum_{\alpha=1, \dots, \frac{n(n-1)}{2}} \bar{U}_\alpha \otimes \bar{U}_\alpha = \frac{1}{2} \sum_{i=1}^{n-1} \bar{e}_i \otimes \bar{e}_i,$$

where $\{\bar{e}_1, \dots, \bar{e}_{n-1}\}$ is a local orthonormal frame on $(S^{n-1}, \bar{\gamma})$. Then for the each \bar{U}_a above, we define a VF U_a on M such that outside a compact set of M it holds

$$(3.3) \quad F_r^* U_a = \bar{U}_a,$$

which is tangent to the level sets Σ_r for all r sufficiently large. By (2.40), we see

$$(3.4) \quad \sum_{a=1, \dots, \frac{n(n-1)}{2}} U_a \otimes U_a = r \left(\sum_{i=1}^{n-1} e_i \otimes e_i + O\left(r^{-\frac{1}{2} - \frac{1}{4}\delta}\right) \right),$$

where $\{e_1, \dots, e_{n-1}\}$ is a local orthonormal frame on Σ_r . Moreover,

$$(3.5) \quad \|\mathcal{L}_{\bar{U}_a}(\gamma_r)\|_{C^l(S^{n-1}, \gamma_r)} = \|\mathcal{L}_{\bar{U}_a}(\gamma_r - \bar{\gamma})\|_{C^l(S^{n-1}, \gamma_r)} \leq O\left(r^{-\frac{1}{2} - \frac{1}{2}\delta}\right).$$

Proposition 3.1. *Each U_a above satisfies (0.5) on (M, g) . Namely,*

$$(3.6) \quad \sup_{\Sigma_r} |\mathcal{L}_{U_a}(g)| \leq O\left(r^{-\frac{1}{2} - \frac{1}{2}\delta}\right)$$

and

$$(3.7) \quad \sup_{\Sigma_r} |\Delta U_a + D_X U_a| \leq O\left(r^{-1 - \frac{1}{4}\delta}\right).$$

Proof. By

$$(3.8) \quad \left[U_a, \frac{X}{|X|^2} \right] = 0,$$

we have

$$(3.9) \quad [U_a, X] = U_a(|X|^2) \frac{X}{|X|^2}.$$

Since by (2.7),

$$U_a(|X|^2) = -\langle U_a, \nabla R \rangle = O(r^{-1-8\delta}),$$

we get

$$(3.10) \quad |[U_a, X]| \leq O(r^{-1-8\delta}).$$

By (3.5) and (3.10), it is easy to see

$$\begin{aligned} & \mathcal{L}_{U_a}(g)(e_i, e_j) \\ &= \langle D_{e_i} U_a, e_j \rangle + \langle D_{e_j} U_a, e_i \rangle = O\left(r^{-\frac{1}{2}-\frac{1}{2}\delta}\right), \end{aligned}$$

$$\begin{aligned} & \frac{1}{2} \mathcal{L}_{U_a}(g)(X, X) \\ &= \langle D_X U_a, X \rangle = \langle D_{U_a} X, X \rangle - \langle [U_a, X], X \rangle = -\frac{1}{2} U_a(|X|^2) = O(r^{-1-8\delta}). \end{aligned}$$

Moreover,

$$\begin{aligned} & \mathcal{L}_{U_a}(g)(X, e_j) \\ &= \langle D_X U_a, e_j \rangle + \langle D_{e_j} U_a, X \rangle = \langle D_{U_a} X, e_j \rangle - \langle U_a, D_{e_j} X \rangle - \langle [U_a, X], e_j \rangle = 0. \end{aligned}$$

Thus we prove (3.6) by the above three relations.

Since

$$\|\bar{U}_a\|_{C^l(S^{n-1}, \gamma_r)} \leq O(1),$$

we have

$$\sup_{\Sigma_r} |D_\Sigma^l U_a| \leq O\left(r^{-\frac{l-1}{2}}\right), \quad \forall l \geq 1.$$

Together with (3.8), we derive

$$(3.11) \quad \sup_{\Sigma_r} |D^l U_a| \leq O\left(r^{-\frac{l-1}{2}}\right), \quad \forall l \geq 1.$$

As a consequence,

$$\sup_{\Sigma_r} |D^l h_a| \leq O(r^{-\frac{l}{2}}), \quad \forall l \geq 1,$$

where $h_a = \mathcal{L}_{U_a}(g)$. Hence by the standard interpolation inequalities, we obtain

$$(3.12) \quad \sup_{\{f=r\}} |D h_a| \leq O(r^{-1-\frac{1}{4}\delta}).$$

On the other hand, we have

$$\operatorname{div}(h_a) - \frac{1}{2} \nabla(\operatorname{tr}(h_a)) = \Delta U_a + \operatorname{Ric}(U_a, \cdot).$$

Therefore,

$$\sup_{\Sigma_r} (\Delta U_a + \operatorname{Ric}(U_a, \cdot)) \leq O\left(r^{-1-\frac{1}{4}\delta}\right).$$

Since

$$|\operatorname{Ric}(U_a, \cdot) - D_X U_a| = |[U_a, X]|,$$

by (3.10),

$$|\operatorname{Ric}(U_a, \cdot) - D_X U_a| \leq O(r^{-1-8\delta}).$$

Combining the above two relations, we prove (3.7) immediately. \square

By Proposition 3.1, there are $\frac{n(n-1)}{2}$ VFs U which satisfy the elliptic equation,

$$(3.13) \quad \Delta U + D_X U = Q,$$

where Q satisfies

$$(3.14) \quad |Q| = O\left(r^{-1-\frac{1}{4}\delta}\right).$$

We need to perturb each U to get a solution of (0.6). First we have the following decay estimate.

Lemma 3.2. *Let U be a solution of (3.13) in the region $\{f \leq \rho\}$, which satisfies (3.14). Then there exist a uniform constant $B \geq 1$ such that*

$$(3.15) \quad \sup_{\{f \leq \rho\}} (|U| - Bf^{-\epsilon}) \leq \sup_{f=\rho} |U| - B\rho^{-\epsilon}.$$

Proof. By the identity (1.4), we have

$$(3.16) \quad \begin{aligned} & -\Delta(f^{-\epsilon}) - \langle X, \nabla(f^{-\epsilon}) \rangle \\ &= \epsilon f^{-1-\epsilon} (\Delta f + |\nabla f|^2) - \epsilon(\epsilon+1)f^{-2-\epsilon} |\nabla f|^2 \\ &> \frac{\epsilon}{2} f^{-1-\epsilon}. \end{aligned}$$

Then by (3.14) with $\epsilon = \frac{1}{4}\delta$, there is a constant $B > 1$ such that

$$|Q| < \frac{\epsilon B}{2} f^{-1-\epsilon}.$$

On the other hand, by the Kato's inequality,

$$\begin{aligned} \Delta(|U|^2) + \langle X, \nabla(|U|^2) \rangle &= 2|DU|^2 + 2\langle U, Q \rangle \\ &\geq 2|\nabla|U||^2 - 2|Q||U|, \end{aligned}$$

we have

$$(3.17) \quad \Delta(|U|) + \langle X, \nabla|U| \rangle \geq -|Q|.$$

Thus we get

$$\Delta(|U| - Bf^{-\epsilon}) + \langle X, \nabla(|U| - Bf^{-\epsilon}) \rangle > 0.$$

Hence, by the maximum principle, we prove (3.15). \square

By Lemma 3.2, we are able to prove

Theorem 3.3. *Let U be a smooth VF on (M, g) which satisfies (3.13) and (3.14). Then there exists a smooth bounded solution V of (0.6) on (M, g) , which satisfies*

$$(3.18) \quad |DV| \leq O(r^{-\frac{1}{2}}).$$

Proof. Let ρ_m be a sequence to the infinity and $V^{(m)}$ a sequence of solutions of (3.13) in the region $\{f \leq \rho\}$ with zero boundary values on the boundary Σ_{ρ_m} . Then by Lemma 3.2, we see

$$(3.19) \quad \sup_{\{f \leq \rho_m\}} |V^{(m)}| \leq B2^{-\epsilon} - B\rho_m^{-\epsilon} \leq C,$$

where C is independent of m . By the regularity, we also get

$$\sup_{\{f \leq \rho_m\}} |D^k V^{(m)}| \leq C_k.$$

Thus by taking a subsequence of $V^{(m)}$, we will obtain a smooth bounded solution V of (0.6) on (M, g)

It remain to prove (3.18). Let $r_m \rightarrow 0$ be a sequence and

$$\hat{g}^{(m)}(t) = r_m^{-1} \Phi_{r_m t}^*(g),$$

where $t \in [-\frac{1}{2}, 0]$. We define

$$\hat{V}^{(m)}(t) = \Phi_{r_m t}^*(V)$$

and

$$\hat{Q}^{(m)}(t) = r_m \Phi_{r_m t}^*(Q).$$

Then $\hat{V}^{(m)}(t)$ satisfies the parabolic equation,

$$\frac{\partial}{\partial t} \hat{V}^{(m)}(t) = \Delta_{\hat{g}^{(m)}(t)} \hat{V}^{(m)}(t) + \text{Ric}_{\hat{g}^{(m)}(t)} \left(\hat{V}^{(m)}(t) \right) - \hat{Q}^{(m)}(t).$$

By (3.14), we see

$$\sup_{t \in [-\frac{1}{2}, 0]} \sup_{\{r_m - \sqrt{r_m} \leq f \leq r_m + \sqrt{r_m}\}} \left| \hat{Q}^{(m)}(t) \right|_{\hat{g}^{(m)}(t)} \leq O \left(r_m^{-\frac{1}{2} - \epsilon} \right)$$

and

$$\sup_{t \in [-\frac{1}{2}, 0]} \sup_{\{r_m - \sqrt{r_m} \leq f \leq r_m + \sqrt{r_m}\}} \left| \hat{V}^{(m)}(t) \right|_{\hat{g}^{(m)}(t)} \leq O \left(r_m^{-\frac{1}{2}} \right).$$

Thus by the standard interior estimates for parabolic equations (cf. [29, Theorem 7.22]), we obtain

$$\begin{aligned} & \sup_{\{f=r_m\}} \left| D\hat{V}^{(m)}(0) \right|_{\hat{g}^{(m)}(0)} \\ & \leq C \sup_{t \in [-\frac{1}{2}, 0]} \sup_{\{r_m - \sqrt{r_m} \leq f \leq r_m + \sqrt{r_m}\}} \left| \hat{V}^{(m)}(t) \right|_{\hat{g}^{(m)}(t)} \\ & + C \sup_{t \in [-\frac{1}{2}, 0]} \sup_{\{r_m - \sqrt{r_m} \leq f \leq r_m + \sqrt{r_m}\}} \left| \hat{Q}^{(m)}(t) \right|_{\hat{g}^{(m)}(t)} \\ & \leq O(r_m^{-\frac{1}{2}}). \end{aligned}$$

Hence by taking a limit, we prove (3.18). □

Corollary 3.4. *There exists a collection of VFs W_a , $a \in \{1, \dots, \frac{n(n-1)}{2}\}$, each of which solves (0.6),*

$$\Delta W_a + D_X W_a = 0, \text{ on } M.$$

Moreover, W_a satisfies (3.1).

Proof. Let $Q = \Delta U_a + D_X U_a$ for each U_a constructed in (3.3). Then by Lemma 3.3, we obtain a solution V_a of (3.13). Thus $W_a = U_a - V_a$ solves (0.6) which satisfies (3.1) by (3.4), (3.11) and Theorem 3.3. □

Remark 3.5. *In [5, Section 3], Brendle used a delicate blow-down analysis for the equation (3.13) to get a locally uniformly bounded estimate for the sequence of $V^{(m)}$. We can prove that $V^{(m)}$ are uniformly bounded by the decay estimate in Lemma (3.2) (also see (3.19), so in particular simple his proof to get a limit solution of (0.6).*

4. ALMOST KILLING VFS WITH FAST DECAY OF LIE DERIVATIVE

In this section, we prove that there exists a collection of approximate Killing VFs Y_a , $a \in \{1, \dots, \frac{n(n-1)}{2}\}$, each of which has fast decay of the Lie derivative for the soliton metric (M, g) near the infinity.

Recall the Lichnerowicz Laplacian Δ_L for $(0, 2)$ -symmetric tensors associated to (M, g) defined by

$$(\Delta_L h)_{ij} \doteq \Delta h_{ij} + 2R_{kijl} h_{kl} - \text{Ric}_{ik} h_{kj} - \text{Ric}_{jk} h_{ki}.$$

Then the Ricci curvature tensor is a special solution of (0.7) (cf. [3, 4, 5]). Namely, we have

$$(4.1) \quad \Delta_L(\text{Ric}(g)) + \mathcal{L}_X(\text{Ric}(g)) = 0.$$

In [5], Brendle found a way to construct the solution of (0.7) by solving the elliptic equation (0.6) for VFs. In fact, if W is a solution of (0.6), then $h = \mathcal{L}_W g$ will satisfy (0.7), i.e.,

$$\Delta_L h + \mathcal{L}_X h = 0.$$

The following lemma shows that the solution of (0.7) has any polynomial decay near the infinity associated to an asymptotically cylindrical steady Ricci soliton.

Lemma 4.1. *Let (M, g) be an asymptotically cylindrical steady Ricci soliton. Let h be a solution of the Lichnerowicz-type equation (0.7) on the region $\{\rho'_0 \leq f \leq \rho\}$ associated to (M, g) . Then for any large l , there exists a $\rho_0 \geq \rho'_0$ such that*

$$(4.2) \quad \sup_{\{\rho_0 \leq f \leq \rho\}} f^l |h| \leq \max\{\rho^l \sup_{\{f=\rho\}} |h|, \rho_0^l \sup_{\{f=\rho_0\}} |h|\}.$$

Proof. Since (M, g) is asymptotically cylindrical, there exists a set $K \subset M$ such that (M, g) admits positive curvature operator outside of K by Lemma 1.3. Moreover, by (2.10) in Lemma 2.1, for any large $l > 0$ there is another compact set K' such that

$$(4.3) \quad f \operatorname{Ric} < \left(\frac{l}{2} - \frac{l(l+1)}{2} f^{-1} |\nabla f|^2\right) g, \quad M \setminus K'.$$

Thus we can a ρ_0 large enough such that $K \cup K' \subset \{f \leq \rho_0\}$.

It suffices to show that

$$(4.4) \quad f^l h \leq \max\{\rho^l \sup_{\{f=\rho\}} |h|, \rho_0^l \sup_{\{f=\rho_0\}} |h|\} g.$$

Then $-h$ will be also true since $-h$ is also a solution of (0.7). Thus (4.4) implies (4.2).

Following the argument in [5, Section 4], we choose the smallest real number θ such that $\theta f^{-l} g - h$ is positive semi-definite at each point in the region $\{\rho_0 \leq f \leq \rho\}$. Then there exists a point $p_0 \in \{\rho_0 \leq f \leq \rho\}$ and an orthonormal basis $\{e_1, \dots, e_n\}$ of $T_{p_0} M$ such that at the point p_0 it holds

$$(4.5) \quad \theta f^{-l} - h(e_1, e_1) = 0.$$

Without loss of generality, we may assume that $p_0 \in \{\rho_0 < f < \rho\}$. Thus we get at p_0 ,

$$(4.6) \quad \theta \Delta(f^{-l}) - (\Delta h)(e_1, e_1) \geq 0$$

and

$$(4.7) \quad \theta \langle X, \nabla(f^{-l}) \rangle - (D_X h)(e_1, e_1) = 0.$$

Note

$$\Delta(f^{-l}) = l(l+1)f^{-l-2} |\nabla f|^2 - lf^{-l-1} \Delta f$$

and

$$\langle X, \nabla(f^{-l}) \rangle = -lf^{-l-1} |\nabla f|^2.$$

Then by (0.7) with help of (4.5)-(4.7), we have at p_0 ,

$$\begin{aligned} 0 &= (\Delta h)(e_1, e_1) + (D_X h)(e_1, e_1) + 2 \sum_{i,k=1}^n R(e_1, e_i, e_1, e_k) h(e_i, e_k) \\ &\leq \theta \Delta(f^{-l}) + \theta \langle X, \nabla(f^{-l}) \rangle + 2 \sum_{i,k=1}^n R(e_1, e_i, e_1, e_k) h(e_i, e_k) \\ &= -l\theta f^{-l-1} \left(1 - (l+1)f^{-1} |\nabla f|^2 - \frac{2}{l} f \operatorname{Ric}(e_1, e_1)\right) \\ &\quad - 2 \sum_{i,k=1}^n R(e_1, e_i, e_1, e_k) (\theta f^{-l} g(e_i, e_k) - h(e_i, e_k)). \end{aligned}$$

Since (M, g) has positive sectional curvature on $M \setminus K$,

$$\sum_{i,k=1}^n R(e_1, e_i, e_1, e_k) (\theta f^{-l} g(e_i, e_k) - h(e_i, e_k)) \geq 0.$$

Thus we obtain

$$-l\theta f^{-l-1} \left(1 - (l+1)f^{-1}|\nabla f|^2 - \frac{2}{l}f \operatorname{Ric}(e_1, e_1) \right) \geq 0, \text{ on } M \setminus K \cup K'.$$

By (4.3), it follows that $\theta \leq 0$. Since $h \leq \theta f^{-l}g$, we conclude that $h \leq 0$ at each point in the region $\{\rho_0 \leq f \leq \rho\}$. Hence, (4.4) is automatically satisfied. The lemma is proved. \square

The following lemma is due to [5, Lemma 4.1].

Lemma 4.2. *Let $(S^{n-1} \times \mathbb{R}, \bar{g}(t))$ ($t \in (0, 1)$) be the shrinking cylinders, namely, $\bar{g}(t)$ is a form of metric on $S^{n-1} \times \mathbb{R}$,*

$$(4.8) \quad \bar{g}(t) = (n-2)(2-2t)g_{S^{n-1}} + dr^2.$$

Suppose that $\bar{h}(t)$ ($t \in (0, 1)$) is a solution of (0.8) associated to $\bar{g}(t)$, which is invariant under translations along the axis of the cylinder, and satisfies

$$(4.9) \quad |\bar{h}(t)|_{\bar{g}(t)} \leq (1-t)^{-l}$$

for some $l > 0$ and all $t \in (0, \frac{1}{2}]$. Then

$$\inf_{\lambda \in \mathbb{R}} \sup_{S^{n-1} \times \mathbb{R}} |\bar{h}(t) - \lambda \operatorname{Ric}_{\bar{g}(t)}|_{\bar{g}(t)} \leq N(1-t)^{\frac{1}{2(n-2)} - \frac{1}{2}}, \quad \forall t \in [\frac{1}{2}, 1).$$

Here N is a positive constant.

By Lemma 4.1 and Lemma 4.2, we prove the following decay estimate.

Theorem 4.3. *For all large l , there exists a collection of approximate Killing VFs Y_a , $a \in \{1, \dots, \frac{n(n-1)}{2}\}$, such that*

$$(4.10) \quad |\mathcal{L}_{Y_a}g| \leq O(r^{-l}),$$

outside a compact set K of M . Moreover, $\mathcal{L}_{Y_a}g$ are solutions of Lichnerowicz type equation (0.7) on M .

Proof. Let W_a , $a \in \{1, \dots, \frac{n(n-1)}{2}\}$, be a collection of VFs on M constructed in Corollary 3.4. Then $h_a = \mathcal{L}_{W_a}g$ satisfy (0.7). Moreover, by (3.6),

$$(4.11) \quad |h_a| \leq O(r^{-\frac{1}{2}}).$$

As in [5, Section 4], we define a function on $[r_0, \infty)$ by

$$A_a(r) = \inf_{\lambda \in \mathbb{R}} \sup_{\{f=r\}} |h_a - \lambda \operatorname{Ric}(g)|.$$

Clearly

$$(4.12) \quad A_a(r) \leq \sup_{\{f=r\}} |h_a| \leq O(r^{-\frac{1}{2}}) \leq O(r^{\frac{1}{2(n-2)} - \frac{1}{2} - \epsilon'}),$$

where $\epsilon' = \frac{1}{1000n}$ is small number. Fix a number $\tau \in (0, \frac{1}{2})$ such that $\tau^{-\epsilon'} > 2N$, where N is the constant in Lemma 4.2. Then by (4.12) there exists a sequence of $r_m \rightarrow \infty$ such that

$$A_a(r_m) \leq 2\tau^{\frac{1}{2} - \frac{1}{2(n-2)} + \epsilon'} A_a(\tau r_m).$$

For the above sequence $\{r_m\}$, there are two cases as follows.

Case 1: There exists a subsequence of $r_m \rightarrow \infty$, we still denote them by r_m , such that

$$(4.13) \quad \rho_0^l \sup_{\{f=\rho_0\}} |h_a - \lambda_m \operatorname{Ric}(g)| \leq r_m^l \sup_{\{f=r_m\}} |h_a - \lambda_m \operatorname{Ric}(g)|.$$

We claim that there is some $\lambda \in \mathbb{R}$ such that

$$(4.14) \quad h_a = \lambda \operatorname{Ric}(g), \text{ on } \{f \geq \rho_0\}.$$

Then we let $Y_a = W_a - \frac{1}{2}\lambda X$, which satisfies

$$\mathcal{L}_{Y_a}g = 0, \quad \{f \geq \rho_0\},$$

and so (4.10) is obviously true. By Corollary 3.4 and the fact $|X| \leq 1$, we notice that Y_a is a non-zero VF and it is globally well-defined on M .

To prove (4.14), we need to show that there exists a sequence of numbers $r_m \rightarrow \infty$ such that $A_a(r_m) = 0$. Namely,

$$(4.15) \quad h_a - \lambda_m \operatorname{Ric}(g) = 0, \quad \text{on } \{f = r_m\}.$$

In fact, by applying Lemma 4.1 to $h_a - \lambda_m \operatorname{Ric}(g)$ on the region $\{\rho_0 \leq f \leq r_m\}$, it follows by (4.13), we have

$$\begin{aligned} & \sup_{\{\rho_0 \leq f \leq r_m\}} f^l |h_a - \lambda_m \operatorname{Ric}(g)| \\ & \leq \max\{r_m^l \sup_{\{f=r_m\}} |h_a - \lambda_m \operatorname{Ric}(g)|, \rho_0^l \sup_{\{f=\rho_0\}} |h_a - \lambda_m \operatorname{Ric}(g)|\} \\ & \leq r_m^l \sup_{\{f=r_m\}} |h_a - \lambda_m \operatorname{Ric}(g)| = 0. \end{aligned}$$

Thus $h_a - \lambda_m \operatorname{Ric}(g) = 0$ in the region $\{\rho_0 \leq f \leq r_m\}$ for any λ_m . As a consequence, all λ_m are same and so (4.14) is true.

Suppose that (4.15) is not true. Then there is a sufficiently large r such that

$$(4.16) \quad A_a(r) > 0.$$

By choosing a λ_m such that $r \in [\rho_0, r_m]$ and then applying Lemma 4.1 to the tensor

$$\tilde{h}_a^{(m)} = \frac{1}{A_a(r_m)} (h_a - \lambda_m \operatorname{Ric}(g)),$$

it follows by (4.13),

$$(4.17) \quad \begin{aligned} & \sup_{\{f=r\}} |\tilde{h}_a^{(m)}| \\ & \leq \frac{r_m^l}{r^l} \sup_{\{f=r_m\}} |\tilde{h}_a^{(m)}| = \frac{r_m^l}{r^l A_a(r_m)} \sup_{\{f=r_m\}} |h_a - \lambda_m \operatorname{Ric}(g)| \\ & = \frac{r_m^l}{r^l}. \end{aligned}$$

Define

$$\hat{g}^{(m)}(t) = r_m^{-1} \Phi_{r_m t}^*(g)$$

and

$$\hat{h}_a^{(m)}(t) = r_m^{-1} \Phi_{r_m t}^*(\tilde{h}_a^{(m)}).$$

Then the metrics $\hat{g}^{(m)}(t)$ evolve by the Ricci flow for each m and the tensors $\hat{h}_a^{(m)}(t)$ satisfy the parabolic Lichnerowicz equation associated to the flow $\hat{g}^{(m)}(t)$,

$$\frac{\partial}{\partial t} \hat{h}_a^{(m)}(t) = \Delta_{L, \hat{g}^{(m)}(t)} \hat{h}_a^{(m)}(t).$$

By (4.17), it is easy to see

$$\limsup_{m \rightarrow \infty} \sup_{t \in [\delta', 1 - \delta']} \sup_{\{r_m - \delta'^{-1} \sqrt{r_m} \leq f \leq r_m + \delta'^{-1} \sqrt{r_m}\}} \left| \hat{h}_a^{(m)}(t) \right|_{\hat{g}^{(m)}(t)} < \infty,$$

where $\delta' \in (0, \frac{1}{2})$ is any fixed number.

Choose a sequence of marked points $p_m \in M$ such that $f(p_m) = r_m$. Then manifolds $(M, \hat{g}^{(m)}(t), p_m)$ converge to a family of shrinking cylinders $(S^{n-1} \times \mathbb{R}, \bar{g}(t))$, $t \in (0, 1)$ in the sense of Cheeger-Gromov, where $\bar{g}(t)$ is given by (4.8). Moreover, the vector fields $r_m^{\frac{1}{2}} X$ converge to the axial vector field $\frac{\partial}{\partial z}$ on $S^{n-1} \times \mathbb{R}$, and the sequence $\hat{h}_a^{(m)}(t)$ converges to a family of tensors $\bar{h}_a(t)$, $t \in (0, 1)$, which solve the parabolic Lichnerowicz equation (0.8) associated to $\bar{g}(t)$.

Note that

$$\Phi_{\sqrt{r_m}s}^* \left(\hat{h}_a^{(m)}(t) \right) = \hat{h}_a^{(m)} \left(t + \frac{s}{\sqrt{r_m}} \right).$$

Then

$$\Psi_s^*(\bar{h}_a(t)) = \bar{h}_a(t),$$

where $\Psi_s : S^{n-1} \times \mathbb{R} \rightarrow S^{n-1} \times \mathbb{R}$ denotes the flow generated by the axial vector field $-\frac{\partial}{\partial z}$. Namely, $\bar{h}_a(t)$ is invariant under translations along the axis of the cylinder. Moreover, by (4.17), we have

$$|\bar{h}_a(t)|_{\bar{g}(t)} \leq (1-t)^{-l}, \quad \forall t \in (0, \frac{1}{2}].$$

Thus by Lemma 4.2, we get

$$(4.18) \quad \inf_{\lambda \in \mathbb{R}} \sup_{S^{n-1} \times \mathbb{R}} |\bar{h}_a(t) - \lambda \text{Ric}_{\bar{g}(t)}|_{\bar{g}(t)} \leq N(1-t)^{\frac{1}{2(n-2)} - \frac{1}{2}}, \quad \forall t \in [\frac{1}{2}, 1).$$

On the other hand, we have

$$\begin{aligned} & \inf_{\lambda \in \mathbb{R}} \sup_{\Phi_{r_m(\tau-1)}(\{f=\tau r_m\})} \left| \hat{h}_a^{(m)}(1-\tau) - \lambda \text{Ric}_{\hat{g}^{(m)}(1-\tau)} \right|_{\hat{g}^{(m)}(1-\tau)} \\ &= \inf_{\lambda \in \mathbb{R}} \sup_{\{f=\tau r_m\}} \left| \tilde{h}_a^{(m)} - \lambda \text{Ric}_g \right|_g \\ &= \frac{1}{A_a(r_m)} \inf_{\lambda \in \mathbb{R}} \sup_{\{f=\tau r_m\}} |h_a - \lambda \text{Ric}_g|_g \\ &= \frac{A_a(\tau r_m)}{A_a(r_m)} \\ &\geq \frac{1}{2} \tau^{\frac{1}{2(n-2)} - \frac{1}{2} - \epsilon'}. \end{aligned}$$

Then as $m \rightarrow \infty$ we obtain

$$(4.19) \quad \inf_{\lambda \in \mathbb{R}} \sup_{S^{n-1} \times \mathbb{R}} |\bar{h}_a(1-\tau) - \lambda \text{Ric}_{\bar{g}(1-\tau)}|_{\bar{g}(1-\tau)} \geq \frac{1}{2} \tau^{\frac{1}{2(n-2)} - \frac{1}{2} - \epsilon'}.$$

Hence, by choosing a small τ so that $\tau^{-\epsilon'} > 2N$, we see that (4.18) is a contradiction with (4.19). Therefore we prove (4.14).

Case 2: There is a ρ_1 large enough, such that

$$(4.20) \quad \rho_0^l \sup_{\{f=\rho_0\}} |h_a - \lambda_m \text{Ric}| > r_m^l \sup_{\{f=r_m\}} |h_a - \lambda_m \text{Ric}|,$$

for all $r_m > \rho_1$. Then applying Lemma 4.1 to $h_a - \lambda_m \text{Ric}(g)$ on region $\{\rho_0 \leq f \leq r_m\}$, for all $r_m > \rho_1$, we have

$$\begin{aligned} & \sup_{\{f=r\}} f^l |h_a - \lambda_m \text{Ric}| \\ & \leq \max \left\{ r_m^l \sup_{\{f=r_m\}} |h_a - \lambda_m \text{Ric}|, \rho_0^l \sup_{\{f=\rho_0\}} |h_a - \lambda_m \text{Ric}| \right\} \\ (4.21) \quad & \leq \rho_0^l \sup_{\{f=\rho_0\}} |h_a - \lambda_m \text{Ric}|, \quad r \in [\rho_0, r_m]. \end{aligned}$$

Claim 4.4. *The sequence λ_m are uniformly bounded.*

By the above claim, we can take a subsequence of λ_m which converges to a bounded number λ , and the VF $W_a - \frac{1}{2}\lambda_m X$ converges to a vector field $Y_a = W_a - \frac{1}{2}\lambda X$ on M smoothly. Thus by (4.21), we obtain

$$(4.22) \quad \sup_{\{f=r\}} |h_a - \lambda \text{Ric}(g)| \leq \frac{\rho_0^l}{r^l} \sup_{\{f=\rho_0\}} |h_a - \lambda \text{Ric}| \leq O(r^{-l}).$$

Hence, (4.10) holds for the above Y_a . The proof of Theorem 4.3 is finished.

On the contrary, there exists a subsequence $\lambda_m \rightarrow \infty$ such that $\lambda_{m+1} - \lambda_m > 1$. Since $|h_a| \leq O(r^{-\frac{1}{2}})$, we have $|\text{Ric}(g)| \geq O(r^{-1})$ by (2.1). Note that λ_m is chosen by the relation

$$A_a(r_m) = \sup_{\{f=r_m\}} |h_a - \lambda_m \text{Ric}(g)| = \inf_{\lambda} \sup_{\{f=r_m\}} |h_a - \lambda \text{Ric}(g)|.$$

Then it is easy to see that

$$(4.23) \quad \lambda_m \leq O(r_m^{\frac{1}{2}}).$$

By (4.21), we have

$$\sup_{\{f=r\}} |h_a - \lambda_m \text{Ric}(g)| \leq \frac{C'}{r^l} \lambda_m, \quad r \in [\rho_0, r_m].$$

In particular,

$$(4.24) \quad \sup_{\{f=r_m\}} |h_a - \lambda_m \text{Ric}(g)| \leq \frac{C'}{r_m^l} \lambda_m.$$

It follows that

$$(4.25) \quad \sup_{\{f=r_m\}} |h_a - \lambda_{m+1} \text{Ric}(g)| \leq \frac{C'}{r_m^l} \lambda_{m+1}.$$

Note that

$$\begin{aligned} & \sup_{\{f=r_m\}} (\lambda_{m+1} - \lambda_m) |\text{Ric}(g)| \\ & \leq \sup_{\{f=r_m\}} |h_a - \lambda_{m+1} \text{Ric}(g)| + \sup_{\{f=r_m\}} |h_a - \lambda_m \text{Ric}(g)|. \end{aligned}$$

Thus combining (4.23)-(4.25), we get

$$(4.26) \quad \begin{aligned} & (\lambda_{m+1} - \lambda_m) \frac{C'}{r_m} \\ & \leq \sup_{\{f=r_m\}} (\lambda_{m+1} - \lambda_m) |\text{Ric}(g)| \leq \frac{C'}{r_m^l} (\lambda_m + \lambda_{m+1}) \\ & \leq \frac{C'}{r_m^l} (\lambda_{m+1} - \lambda_m) + \frac{C''}{r_m^{l-\frac{1}{2}}}. \end{aligned}$$

As a consequence, by choosing $l \geq 2$, we derive

$$(\lambda_{m+1} - \lambda_m) \leq O(r_m^{-l+\frac{3}{2}}),$$

and so by (4.23),

$$(4.27) \quad \lambda_{m+1} \leq O(r_m^{\frac{1}{2}}).$$

Therefore, by (4.26), we conclude

$$(4.28) \quad \begin{aligned} & \sup_{\{f=r_m\}} (\lambda_{m+1} - \lambda_m) |\text{Ric}(g)| \leq \frac{C'}{r_m^l} (\lambda_m + \lambda_{m+1}) \\ & \leq O(r_m^{-l+\frac{1}{2}}). \end{aligned}$$

However, by the fact $\lambda_{m+1} - \lambda_m > 1$, we see

$$\sup_{\{f=r_m\}} (\lambda_{m+1} - \lambda_m) |\text{Ric}(g)| \geq O(r_m^{-1}),$$

which is a contradiction with (4.28)! Claim 4.4 is proved. \square

Remark 4.5. *Because of lack of the positivity of sectional curvature of g , we only do the decay estimates for the tensors h_a and $\mathcal{L}_{Y_a}g$ away from a compact set of M in Lemma 4.1 and Theorem 4.3, respectively. Moreover, we need to do the fast decay estimate for those tensors in order to use the heat kernel method in the latter Section 5.*

5. EVOLUTION SOLUTION OF LICHNEROWICZ EQUATION

In this section, we study the evolution equation of Lichnerowicz (0.8) by using the heat kernel method under the curvature-pinching condition (0.1).

Let $g(t)$ ($t \in (0, T]$) be a solution of Ricci flow and G the heat kernel of following heat-typed equation associated to $g(t)$,

$$(5.1) \quad \frac{\partial}{\partial t} H = \Delta_{g(t)} H + 2PH,$$

where $P = P(\cdot, t)$ is a smooth function on $M \times (0, T]$. Namely, $G > 0$ satisfies

$$(5.2) \quad \begin{aligned} \partial_t G(x, t; y, s) &= \Delta_{x,t} G(x, t; y, s) + 2PG(x, t; y, s) \\ \lim_{t \searrow s} G(\cdot, t; y, s) &= \delta_y. \end{aligned}$$

The existence of heat kernel G on a complete manifold can be found in [15, Chapter 24]. Moreover, G satisfies the following estimate,

$$(5.3) \quad G(x, t; y, 0) \leq \frac{C_3 e^{C_1 t} \exp\left(-\frac{d_{g(t)}^2(x, y)}{C_4 t}\right)}{\text{Vol}_{g(t)}^{1/2} B_{g(t)}\left(x, \sqrt{\frac{t}{2}}\right) \cdot \text{Vol}_{g(t)}^{1/2} B_{g(t)}\left(y, \sqrt{\frac{t}{2}}\right)}.$$

We first prove the existence of positive bounded super-solution of (5.1).

Lemma 5.1. *Let (M^n, g) ($n \geq 4$) be a steady gradient Ricci soliton and $g(t)$ the soliton Ricci flow of g . Suppose that g has an asymptotically cylindrical property and satisfies the following curvature-pinching condition,*

$$(5.4) \quad 2P < \frac{m}{m+1} R,$$

where P is the function defined by (0.2). Then for any large $m \geq 2$ there exists a smooth positive uniformly bounded function $s(x, t)$ such that

$$(5.5) \quad \left(\frac{\partial}{\partial t} - \Delta_{g(t)} - 2P(\cdot, t)\right)s(x, t) \geq 0,$$

where $P(\cdot, t) = \Phi_t^* P(\cdot)$. Moreover $s(x, 0)$ satisfies

$$(5.6) \quad \lim_{\rho} \frac{|s(x, 0)|}{\rho(x)^m} = A \neq 0,$$

where $\rho(x) = d(p, x)$.

Proof. We reduce to find a smooth positive uniformly bounded function s such that

$$(5.7) \quad \Delta s + D_X s + 2Ps \leq 0.$$

Then $s(x, t) = \Phi_t^*(s)$ will satisfy (5.5), where Φ_t^* is the transformations group generated by $-X$.

By a direct calculation,

$$(5.8) \quad \begin{aligned} \Delta e^{-f} + D_X e^{-f} + 2P e^{-f} \\ = -\Delta f e^{-f} + |\nabla f|^2 e^{-f} - |\nabla f|^2 e^{-f} + 2P e^{-f} \\ = -(R - 2P)e^{-f} \leq -\frac{1}{m+1} R e^{-f}. \end{aligned}$$

Then

$$(5.9) \quad \begin{aligned} \Delta f^{-m} + D_X f^{-m} + 2P f^{-m} \\ = -m f^{-m-1} \Delta f + m(m+1) f^{-m-2} |\nabla f|^2 - m f^{-m-1} |\nabla f|^2 + 2P f^{-m} \\ \leq -m f^{-m-1} (1 - (m+1) |\nabla f|^2 f^{-1} - \frac{1}{m+1} R f). \end{aligned}$$

Note that $f \geq 2$ and by (2.9) we may choose m large enough, such that $\frac{1}{m+1} \frac{n}{2} < \frac{1}{4}$. Thus there exists a compact set K_1 such that $\frac{1}{m+1} R f < \frac{1}{4}$ outside of K_1 . Moreover, we can choose another

compact set K_2 such that $(m+1)f^{-1} < \frac{1}{4}$ outside of K_2 . Let $K' = K_1 \cup K_2$. Hence, by (5.9) we get

$$(5.10) \quad \Delta f^{-m} + D_X f^{-m} + 2P f^{-m} \leq 0, \quad M \setminus K'.$$

By the compactness of K' , it is easy to see

$$(5.11) \quad |m f^{-m-1} (1 - (m+1)|\nabla f|^2 f^{-1} - \frac{1}{m+1} R f)| \leq C_1,$$

and

$$(5.12) \quad \frac{1}{m+1} R e^{-f} > C_2,$$

where C_1 and C_2 are two constants. Let

$$s(\cdot) = \frac{C_1 + 1}{C_2} e^{-f} + f^{-m}.$$

Then by (5.8) and (5.10), we get

$$\Delta s + D_X s + 2P s \leq 0, \quad M \setminus K'.$$

Moreover, by (5.8), (5.9), (5.11) and (5.12), we also obtain

$$\Delta s + D_X s + 2P s \leq 0, \quad K'.$$

Hence, the function $s(x)$ satisfies (5.7) and

$$|s(x)| \leq O(\rho^{-m}).$$

The lemma is proved. □

Let $G(x, t; y, s)$ be the heat kernel in (5.2) for the soliton Ricci flow $g(t)$ and the function $P(\cdot, t) = \Phi_t^* P(\cdot)$ in Lemma 5.1. For any fixed $D > 0$, we set

$$(5.13) \quad u_D(x, t) = \int_{B_0(p, D)} G(x, t; y, 0) dv_{g(0)} y,$$

where p is the maximal point of R and $B_0(p, D)$ denotes the D -geodesic ball centered at p with respect to the Ricci soliton g . We have the following estimate for $u_D(x, t)$.

Lemma 5.2. *Let $(M, g(t))$ be the soliton Ricci flow of g in Lemma 5.1. Suppose that (0.1) holds for g . Then*

$$(5.14) \quad \sup_{B_t(p, D)} u_D(\cdot, t) \rightarrow 0, \quad \text{as } t \rightarrow \infty,$$

where $B_t(p, D)$ denotes the D -geodesic ball of $g(t)$, which exhausts M as $t \rightarrow \infty$.

Proof. By the definition of G , we have

$$s(x, 0) = \int_M G(x, 0; y, 0) s(y, 0) d_0 y.$$

Since $s(x, t)$ is a super-solution of (5.1) by Lemma 5.1, by the maximum principle we see

$$(5.15) \quad s(x, t) \geq \int_M G(x, t; y, 0) s(y, 0) d_0 y.$$

Note that there is some $c > 0$ such that

$$s(y, 0) \geq c, \quad \text{in } B_0(p, D).$$

Thus we obtain

$$\begin{aligned} & c \int_{B_0(p,D)} G(x,t;y,0) dv_{g(0)} y \\ & \leq c \int_M G(x,t;y,0) dv_{g(0)} y \leq \int_M G(x,t;y,0) s(y,0) dv_{g(0)} y \\ & \leq s(x,t) \leq C. \end{aligned}$$

It follows

$$(5.16) \quad u_D(x,t) \leq c^{-1}C = C_1.$$

Since $u_D(x,t)$ satisfies the equation (5.1), by a standard parabolic estimate, we obtain

$$(5.17) \quad |\partial_t u_D(x,t)| + |\nabla u_D(x,t)| \leq C_2.$$

Now we use the contradiction argument as in [35, Section 7] to prove (5.14). Suppose that there exist $\epsilon > 0$ and sequences of $t_i \rightarrow \infty$ and $x_i \in B_{t_i}(p,D)$ such that $u_D(x_i, t_i) > \epsilon$. Without loss of generality we may assume that $t_{i+1} \geq t_i + 1$. Thus by (5.17), we have

$$(5.18) \quad \begin{aligned} & \int_{B_{t_i}(p,D)} \int_{B_0(p,D)} G(x,t;y,0) dv_{g(0)} y dv_{g(t)} x \\ & = \int_{B_{t_i}(p,D)} u_D(x,t) dv_{g(t)} x \geq \delta_1 > 0, \quad \forall t \in [t_i, t_i + \delta_1], \end{aligned}$$

where δ_1 is a small constant.

By the Gaussian upper bound of G in (5.3), we use the divergence theorem to see

$$\int_M \Delta_{x,t} G(x,t;y,0) dv_{g(t)} x = 0.$$

It follows

$$(5.19) \quad \begin{aligned} & \partial_t \int_M G(x,t;y,0) dv_{g(t)} x = \int_M [\partial_t G(x,t;y,0) - R(x,t)G(x,t;y,0)] dv_{g(t)} x \\ & = \int_M [\Delta_{x,t} G(x,t;y,0) + 2P(x,t)G(x,t;y,0) - R(x,t)G(x,t;y,0)] dv_{g(t)} x \\ & = \int_M [2P(x,t) - R(x,t)] G(x,t;y,0) dv_{g(t)} x, \end{aligned}$$

where the function $P(\cdot, t) = \Phi_t^* P(\cdot)$. We notice that by (0.1) there is a $\delta_2 > 0$ such that

$$(5.20) \quad \sup_{B_{t_i}(p,D)} [2P(\cdot, t) - R(\cdot, t)] = \sup_{B_0(p,D)} [2P(\cdot) - R(\cdot)] \leq -\delta_2.$$

Let

$$G_D(t) = \int_{B_0(p,D)} \int_M G(x,t;y,0) dv_{g(t)} x dv_{g(0)} y.$$

Then by (5.19), we get

$$(5.21) \quad \partial_t G_D(t) = \int_{B_0(p,D)} \int_M [2P(x,t) - R(x,t)] G(x,t;y,0) dv_{g(t)} x dv_{g(0)} y < 0.$$

More precisely, by (5.18) and (5.20) we obtain

$$\partial_t G_D(t) \leq -\delta_1 \delta_2, \quad t \in [t_i, t_i + \delta_1].$$

Since G is everywhere positive and so as $G_D(t) > 0$, we derive

$$\begin{aligned} -G_D(0) & \leq \lim_{t \rightarrow \infty} G_D(t) - G_D(0) \\ & \leq -\sum_{i=1}^{\infty} \int_{t_i}^{t_i + \delta_1} \delta_1 \delta_2 dt = -\sum_{i=1}^{\infty} \delta_1^2 \delta_2. \end{aligned}$$

The above relation implies that $G_D(0) = \infty$, which is impossible!. Therefore (5.14) is true and the lemma is proved.

□

The following is a generalization of pinching estimate of Anderson-Chow for the solution of (0.8) in higher dimensions [1].

Lemma 5.3. *Let $(M^n, g(t))$ ($t \in (0, T]$) be an $n \geq 4$ -dimension Ricci flow. Let H be a positive solution of (5.1) associated to the flow $g(t)$ with the function $P(\cdot, t)$ given by*

$$(5.22) \quad P(\cdot, t) = \sup_h \frac{\text{Rm}_{g(t)}(\cdot)(h, h)}{|h|^2} = \sup_h \frac{\text{R}_{t,ijkl}(\cdot)h_{ik}h_{jl}}{h_{ik}h_{ik}}, \quad \forall t \in (0, T].$$

Then any solution h of (0.8) satisfies the following pinching estimate,

$$(5.23) \quad \partial_t \left(\frac{|h|^2}{H^2} \right) \leq \Delta \left(\frac{|h|^2}{H^2} \right) + \frac{2}{H} \nabla H \cdot \nabla \left(\frac{|h|^2}{H^2} \right),$$

where $\Delta = \Delta_{g(t)}$ and $|\cdot| = |\cdot|_{g(t)}$.

Proof. By a direct computation with help of (0.8) and (5.1), we have the following identities

$$\begin{aligned} & \partial_t \left(\frac{|h|^2}{H^2} \right) \\ &= -\frac{2}{H^2} \frac{\partial}{\partial t} g_{ij} h_{ik} h_{kj} + \frac{2}{H^2} \left(\frac{\partial}{\partial t} h \right) \cdot h - 2 \frac{|h|^2}{H^3} \frac{\partial}{\partial t} H \\ &= \frac{4 \text{Ric}(h, h)}{H^2} + \frac{2}{H^2} (\Delta h_{ij} + 2R_{kijl} h_{kl} - R_{ik} h_{kj} - R_{jk} h_{ki}) h_{ij} - 2 \frac{|h|^2}{H^3} (\Delta H + 2PH), \end{aligned}$$

and

$$\begin{aligned} & \Delta \left(\frac{|h|^2}{H^2} \right) \\ &= 2 \frac{\Delta h \cdot h}{H^2} + 2 \frac{|\nabla h|^2}{H^2} - 8 \frac{h}{H^3} \langle \nabla h, \nabla H \rangle - 2 \frac{|h|^2 \Delta H}{H^3} + 6 \frac{|h|^2 |\nabla H|^2}{H^4}. \end{aligned}$$

Note

$$\frac{1}{H} \nabla H \cdot \nabla \left(\frac{|h|^2}{H^2} \right) = 2 \frac{h}{H^3} \langle \nabla h, \nabla H \rangle - 2 \frac{|h|^2 |\nabla H|^2}{H^4}.$$

Then combining the above three relations, we get

$$\begin{aligned} & \partial_t \left(\frac{|h|^2}{H^2} \right) \\ &= \Delta \left(\frac{|h|^2}{H^2} \right) + \frac{2}{H} \nabla H \cdot \nabla \left(\frac{|h|^2}{H^2} \right) + \frac{4}{H^2} (R_{ijkl} h_{ik} h_{jl} - P|h|^2) \\ &\quad - 2 \frac{|H \nabla_i h_{jk} - (\nabla_i H) h_{jk}|^2}{H^4} \\ &\leq \Delta \left(\frac{|h|^2}{H^2} \right) + \frac{2}{H} \nabla H \cdot \nabla \left(\frac{|h|^2}{H^2} \right). \end{aligned}$$

The last inequality follows from the condition (5.22). □

By the above three lemmas, we are able to get an asymptotic behavior estimate for the solution of (0.8).

Proposition 5.4. *Let $(M^n, g(t))$ be the soliton Ricci flow of g with dimension $n \geq 4$ in Lemma 5.1. Let $h(t)$ ($t \in [0, \infty)$) be a solution of (0.8) with an initial value $h(x, 0)$ satisfying*

$$(5.24) \quad |h|(x, 0) \leq O(\rho^{-l}),$$

for some $l > m$, where m is the large number chosen as in (5.4). Then

$$(5.25) \quad \lim_t \sup_{B_t(p, D)} |h|(\cdot, t) = 0.$$

Proof. Let $G(x, t; y, s)$ be the heat kernel in (5.13). Set

$$H(x, t) = \int_M G(x, t; y, 0) |h|(y, 0) d_{g(0)} y.$$

Then $H(\cdot, t)$ is a solution of (5.1) with $P(\cdot, t) = \Phi_t^* P(\cdot)$ for any $t > 0$. Thus by Lemma 5.3, (5.23) holds for $\frac{|h|^2}{H^2}$. By the maximum principle, we see that $|h| \leq H$, that is,

$$(5.26) \quad |h(x, t)| \leq \int_M G(x, t; y, 0) |h|(y, 0) d_{g(0)} y.$$

On the other hand, by (5.24) and (5.6) in Lemma 5.1, for any $\epsilon > 0$ we can find some $D > 0$ such that

$$|h|(y, 0) \leq \epsilon s(y, 0), \quad \forall y \in M \setminus B_0(p, D).$$

We may assume $D > \frac{1}{\epsilon}$. Hence, by (5.26) and (5.15) together with Lemma 5.2, we see that for sufficiently large t and $x \in B_t(p, D)$,

$$\begin{aligned} |h(x, t)| &\leq \int_{B_0(p, D)} G(x, t; y, 0) |h|(y, 0) d_{g(0)} y + \int_{M \setminus B_0(p, D)} G(x, t; y, 0) |h|(y, 0) d_{g(0)} y \\ &\leq \epsilon + \epsilon \int_M G(x, t; y, 0) |s|(y, 0) d_{g(0)} y \\ &\leq \epsilon + \epsilon s(x, t) \leq \epsilon(1 + C). \end{aligned}$$

Since ϵ can be taken arbitrary small, we will get (5.25). The proposition is proved. \square

6. PROOF OF MAIN THEOREM AND COROLLARIES

In this section, we complete the proofs of Theorem 0.2, Corollary 0.3 and Corollary 0.5.

Proof of Theorem 0.2. By Lemma 1.3, it is easy to see

$$\lim_{x \rightarrow \infty} \frac{P(x)}{R(x)} = \frac{2}{n-1}.$$

On the other hand, by the condition (0.1), for any compact set K there exists a small positive number μ such that

$$2P < (1 - \mu)R \leq \frac{m}{m+1}R, \quad \text{on } K,$$

where m can be chosen sufficiently large. Thus the condition (5.4) holds on the total space (M, g) .

By Theorem 4.3, we already know that there is a collection of approximate Killing VFs Y_a ($a \in \{1, \dots, \frac{n(n-1)}{2}\}$) such that

$$|\mathcal{L}_{Y_a} g| \leq O(r^{-l}),$$

outside a compact set, where l can be chosen so that $l > m$. Let $g(t) = \Phi_t^*(g)$ and $Y_a(t) = \Phi_t^*(Y_a)$. Then $h_a(t) = \mathcal{L}_{Y_a(t)} g(t) = \Phi_t^* h_a$ is a solution of (0.8) for any $t > 0$. This is because $h_a = \mathcal{L}_{Y_a} g$ is a solution of (0.7). Thus applying Proposition 5.4 to each solution $h_a(t)$ of (0.7) with the initial value h_a , we see

$$\lim_t \sup_{B_t(p, D)} |h_a(\cdot, t)| = 0.$$

Note that $h_a(\cdot) = \Phi_{-t}^* h_a(t)$. Hence, $|h_a| \equiv 0$ on any compact D -geodesic ball $B_0(p, D) \subset M$. This implies that $|h_a| \equiv 0$ on M . Therefore we prove that each Y_a is a Killing VF on (M, g) .

By [5, Proposition 4.1], Y_a satisfies

$$\Delta Y_a + \text{Ric}(g)(Y_a, \cdot) = 0.$$

Since Y_a also satisfies by the construction of Y_a ,

$$\Delta Y_a + D_X Y_a = 0,$$

we get

$$[Y_a, X] = \text{Ric}(g)(Y_a, \cdot) - D_X Y_a = 0.$$

It follows

$$D^2(\mathcal{L}_{Y_a}(f)) = \mathcal{L}_{Y_a}(D^2 f) = \frac{1}{2}\mathcal{L}_{Y_a}(\mathcal{L}_X g) = \frac{1}{2}\mathcal{L}_X(\mathcal{L}_{Y_a} g) = 0.$$

Hence, the function $\mathcal{L}_{Y_a}(f) = \langle Y_a, X \rangle$ is constant. Since X vanishes at the point where f attains its minimum, the function $\langle Y_a, X \rangle$ must vanish identically. This implies that (M, g) is rotationally symmetric and Theorem 0.2 is proved. \square

The first part of Corollary 0.3 is a direct application of Theorem 0.2 since (M, g) has an asymptotically cylindrical property under the linear decay condition of curvature in (0.3) by [18, Theorem 1.3]. Then by Proposition 1.6, the second part is also true.

Remark 6.1. *In 4d, the condition $\text{Rm}_g(x) \geq 0$ ($\rho(x) \geq r_0$) in (0.3) can be weakened as the non-negativity of sectional curvature. This is because all level sets of f are diffeomorphic to \mathbb{S}^3 and so any limit of Ricci flow from a blow-down sequence of Ricci flow obtained in [18, Theorem 1.4] satisfies Definition 1.2-(ii) by Hamilton's result for closed 3d-Ricci flow with positive Ricci curvature [24] (also see [19, Theorem 1.5]).*

Proof of Corollary 0.5. We need to check that (M, g) has an asymptotically cylindrical property under the condition (0.4). In fact, for a fixed $p \in (\mathbb{R}^n, g_0)$, we can choose an orthonormal basis $\{e_1, e_2, \dots, e_n\}$ of $T_p M$ with respect to metric g_0 with $e_n = \frac{\nabla f}{|\nabla f|}$. Then it is easy to see

$$(6.1) \quad \sup_{\partial B_p(r)} |\hat{R}_{ijkl} - R_{ijkl}|_{g_0} \leq O(r^{-1-\tau}), \quad 1 \leq i, j, k, l \leq n-1,$$

$$(6.2) \quad \sup_{\partial B_p(r)} |\hat{R}_{ijkn} - R_{ijkn}|_{g_0} \leq O(r^{-\frac{3}{2}-\tau}), \quad 1 \leq i, j, k \leq n-1,$$

and

$$(6.3) \quad \sup_{\partial B_p(r)} |\hat{R}_{inkn} - R_{inkn}|_{g_0} \leq O(r^{-2-\tau}), \quad 1 \leq i, k \leq n-1.$$

Here we use the sectional curvature decay of Bryant soliton and (2.13). By the above three relations, we see that (M, g) has positive curvature operator outside of a compact set K . Moreover, its scalar curvature has exactly linear decay, i.e. there exists positive numbers r_0, C_1 and C_2 such that

$$(6.4) \quad \frac{C_1}{\rho(x)} \leq R(x) \leq \frac{C_2}{\rho(x)}, \quad \forall \rho(x) \geq r_0.$$

Also one can check that (M, g) is κ -noncollapsed for some $\kappa > 0$.

According to the proof of [18, Theorem 1.4], it suffices to prove that the Gauss sphere Σ_r of geodesic ball of $B_r(p)$ in (M, g) is diffeomorphic to the unit sphere in the Euclidean space and

$$\text{Diam}(\Sigma_r, g) \leq C\sqrt{r}$$

as long as r is sufficiently large. But these two facts can be guaranteed by (0.4). Thus the corollary is proved. \square

REFERENCES

- [1] Anderson, G. and Chow, B., *A pinching estimate for solutions of the linearized Ricci flow system on 3-manifolds*, *Calc. Var. Partial Differential Equations*, **23** (2005), 1–12.
- [2] Appleton, A., *A family of non-collapsed steady Ricci solitons in even dimensions greater or equal to four*, arXiv:1708.00161.
- [3] Besse, A., *Einstein manifolds*, volume 10 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*, Springer-Verlag, Berlin, 1987.
- [4] Brendle, S., *Rotational symmetry of self-similar solutions to the Ricci flow*, *Invent. Math.*, **194** (2013), 731–764.

- [5] Brendle, S., *Rotational symmetry of Ricci solitons in higher dimensions*, *J. Differential Geom.*, **97** (2014), 191–214.
- [6] Brendle, S. and Schoen, R., *Manifolds with 1/4-pinched curvature are space forms*, *J. Amer. Math. Soc.*, **22** (2009), 287–307.
- [7] Brendle, S., *Ricci flow with surgery on manifolds with positive isotropic curvature*, *Ann. of Math.*, **190** (2019), 465–559.
- [8] Brendle, S., *Ricci flow and the sphere theorem*, American Mathematical Soc., 2010.
- [9] Cao, H. D. and Chen, Q., *On locally conformally flat gradient steady Ricci solitons*, *Trans. Amer. Math. Soc.*, **364** (2012), 2377–2391.
- [10] Chen, B. L., *Strong uniqueness of the Ricci flow*, *J. Differential Geom.*, **82** (2009), 363–382.
- [11] Chen, B. L., Tang, S. and Zhu, X. P., *Complete classification of compact four-manifolds with positive isotropic curvature*, *J. Differential Geom.*, **91** (2012), 41–80.
- [12] Chen, B. L. and Zhu, X. P., *Ricci flow with surgery on four-manifolds with positive isotropic curvature*, *J. Differential Geom.*, **74** (2006), 177–264.
- [13] Cho, J. H. and Li, Y., *Ancient solutions to the Ricci flow with isotropic curvature conditions*, *Mathematische Annalen*, (2022), 1–33.
- [14] Chow, B., Deng, Y. and Ma, Z., *On four-dimensional steady gradient Ricci solitons that dimension reduce*, *Adv. Math.*, **403** (2022), 61 pp.
- [15] Chow, B., Chu, S., Glickenstein, D., Guenther, C., Isenberg, J., Ivey, T., Knopf, D., Lu, P., Luo, F. and Ni, L., *The Ricci flow: techniques and applications. Part III: geometric-analytic aspects.*, American Mathematical Society, 163, 2010.
- [16] Chow, B., Lu, P. and Ni, L., *Hamilton’s Ricci flow*, American Mathematical Soc., 2006.
- [17] Conlon, R. and Deruelle, A., *Steady gradient Kähler-Ricci solitons on crepant resolutions of Calabi-Yau cones*, arXiv:2006.03100.
- [18] Deng, Y. and Zhu, X. H., *Higher dimensional steady Ricci solitons with linear curvature decay*, *J. Eur. Math. Soc. (JEMS)*, **22** (2020), 4097–4120.
- [19] Deng, Y. and Zhu, X. H., *Classification of gradient steady Ricci solitons with linear curvature decay*, *Sci. China Math.*, **63** (2020), 135–154.
- [20] Deng, Y. and Zhu, X. H., *Steady Ricci solitons with horizontally ϵ -pinched Ricci curvature*, *Sci. China Math.*, **64** (2021), 1411–1428.
- [21] Deruelle, A., *Asymptotic estimates and compactness of expanding gradient Ricci solitons*, *Ann. Sc. Norm. Super. Pisa Cl. Sci.*, **17** (2017), 485–530.
- [22] Deruelle, A., *Smoothing out positively curved metric cones by Ricci expanders*, *Geom. Funct. Anal.*, **26** (2016), 188–249.
- [23] Dancer, A. and Wang, M., *Some New Examples of Non-Kähler Ricci Solitons*, *Math. Res. Lett.*, **16** (2009), 349–363.
- [24] Hamilton, R. S., *Three-manifolds with positive Ricci curvature*, *J. Differential Geometry*, **17** (1982), 255–306.
- [25] Hamilton, R. S., *The Ricci flow on surfaces*, *Contemporary Mathematics*, **71** (1988), 237–261.
- [26] Hamilton, R. S., *Formation of singularities in the Ricci flow*, *Surveys in Diff. Geom.*, **2** (1995), 7–136.
- [27] Hamilton, R. S., *Four-manifolds with positive isotropic curvature*, *Comm. Anal. Geom.*, **5** (1997), 1–92.
- [28] Kotschwar, B. and Wang, L., *Rigidity of asymptotically conical shrinking gradient Ricci solitons*, *J. Differential Geom.*, **100** (2015), 55–108.
- [29] Lieberman, G., *Second order parabolic differential equations*, World scientific, 1996.
- [30] Micallef, M. and Moore, J., *Minimal two-spheres and the topology of manifolds with positive curvature on totally isotropic two-planes*, *Ann. of Math.*, **127** (1988), 199–227.
- [31] Nguyen, H., *Isotropic curvature and the Ricci flow*, *Int. Math. Res. Not.*, (2010), 536–558.
- [32] Perelman, G., *The entropy formula for the Ricci flow and its geometric applications*, arXiv:0211159, 2002.
- [33] Shi, W. X., *Ricci deformation of the metric on complete noncompact Riemannian manifolds*, *J. Differential Geom.*, **30** (1989), 303–394.
- [34] Lai, Y., *A family of 3d steady gradient solitons that are flying wings*, arXiv:2010.07272, 2020.
- [35] Lai, Y., *O (2)-symmetry of 3D steady gradient Ricci solitons*, arXiv:2205.01146, 2022.
- [36] Zhang, Z. H., *On the completeness of gradient Ricci solitons*, *Proc. Amer. Math. Soc.*, **137** (2009), 2755–2759.

BICMR AND SMS, PEKING UNIVERSITY, BEIJING 100871, CHINA.

Email address: 1901110027@pku.edu.cn, xhzhu@math.pku.edu.cn