

ATTAINABLE MEASURES FOR CERTAIN TYPES OF P-ADIC DUFFIN-SCHAEFFER SETS

MATHIAS L. LAURSEN

Abstract

This paper settles recent conjectures concerning the p -adic Haar measure applied to a family of sets defined in terms of Diophantine approximation. This is done by determining the spectrum of measure values for each family and seeing that this contradicts the corresponding conjectures.

1. Introduction

In Diophantine approximation, we are often interested in determining when a given number $\alpha \in [0, 1]$ has infinitely many solutions to equations of the form

$$\left| x - \frac{a}{n} \right| < \frac{\psi(n)}{n}, \quad (1.1)$$

for some chosen function $\psi : \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$. In 1924, Khintchine [7] gave one of the first metric results on regarding this equation, as stated below.

THEOREM (KHINTCHINE). *Let $\psi : \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$ and write*

$$\mathcal{K}(\psi) = \{ \alpha \in [0, 1] : (1.1) \text{ holds for infinitely many } (a, n) \in \mathbb{Z} \times \mathbb{N} \}.$$

If $\psi(n)$ is monotonic, then $\mathcal{K}(\psi)$ has Lebesgue measure 1 if $\sum_{n=1}^{\infty} \psi(n) = \infty$ and has Lebesgue measure 0 otherwise.

Later, Duffin and Schaeffer [3] tried to remove the monotonicity condition and found a counterexample, which led them to suppose that it would be more natural to consider the set

$$\mathcal{A}(\psi) = \left\{ x \in [0, 1] : (1.1) \text{ holds for infinitely many } \frac{a}{n} \in \mathbb{Q} \right\}, \quad (1.2)$$

where the fractions a/n are assumed to be reduced, i.e. $\gcd(a, n) = 1$. They then famously conjectured the below theorem, which was only proven a couple of years ago by Koukoulopoulos and Maynard [8].

DUFFIN–SCHAEFFER THEOREM. *Let $\psi : \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$. Then*

$$|\mathcal{A}(\psi)| = \begin{cases} 0 & \text{if } \sum_{n=1}^{\infty} \frac{\psi(n)\phi(n)}{n} < \infty, \\ 1 & \text{if } \sum_{n=1}^{\infty} \frac{\psi(n)\phi(n)}{n} = \infty, \end{cases}$$

Here and throughout this paper, ϕ denotes the Euler totient function. Named after the conjecture, we will refer to $\mathcal{A}(\psi)$ as the (real) Duffin–Schaeffer set. In handling the conjecture, one usually writes it as a lim-sup set $\mathcal{A}(\psi) = \limsup_{n \rightarrow \infty} \mathcal{A}_n(\psi)$ where

$$\mathcal{A}_n(\psi) = \bigcup_{\substack{1 \leq a \leq n \\ \gcd(a,n)=1}} B_{\mathbb{R}}\left(\frac{a}{n}, \frac{\psi(n)}{n}\right) \cap [0, 1].$$

We use $B_X(x, r)$ to denote the closed ball of radius r around the point x in metric space X when $r > 0$, and the singleton $\{x\}$ when $r = 0$. The convergence case is then a simple application of the Borel–Cantelli lemma, but the divergence case is much deeper and was settled in the affirmative only a couple of years ago by Koukoulopoulos and Maynard [8].

Before the conjecture was settled, Haynes [5] proposed a p -adic variant of the conjecture in terms of the p -adic Haar measure μ_p with $\mu_p(\mathbb{Z}_p) = 1$ and a p -adic Duffin–Schaeffer set, \mathcal{A}^p . He here chose to first translate \mathcal{A}_n into

$$\mathcal{A}_n^p(\psi) := \bigcup_{\substack{|a| \leq n \\ \gcd(a,n)=1}} B_{\mathbb{Q}_p}\left(\frac{a}{n}, \frac{\psi(n)}{n}\right) \cap \mathbb{Z}_p$$

and then define \mathcal{A}^p as the limsup set of $\mathcal{A}_n^p(\psi)$. From this, Haynes phrased a p -adic Duffin–Schaeffer conjecture for the set $\mathcal{A}^p(\psi)$. One of the main results of Haynes’ paper was that if a certain ‘quasi-independentness on average’ criterion, closely related to the real Duffin–Schaeffer conjecture, were to hold, then his conjecture would follow. In [9], Kristensen and Laursen modified arguments from [8] to prove this ‘quasi-independentness on average’ for relevant ψ , thus settling the conjecture in the affirmative, so we may now state it as a theorem.

p -ADIC DUFFIN–SCHAEFFER THEOREM. *We have*

$$\mu_p(\mathcal{A}^p(\psi)) = \begin{cases} 0 & \text{if } \sum_{n=1}^{\infty} \mu_p(\mathcal{A}_n^p(\psi)) < \infty \\ 1 & \text{if } \sum_{n=1}^{\infty} \mu_p(\mathcal{A}_n^p(\psi)) = \infty. \end{cases}$$

However, the set $\mathcal{A}^p(\psi)$ is not the only natural choice for a p -adic variant of $\mathcal{A}(\psi)$, as one might instead start from equation (1.2) when translating the

question to a p -adic context. For this, we first need a translation of equation (1.1). Following Jarník [6] and Lutz [10], this should be

$$\left| x - \frac{a}{n} \right|_p \leq \frac{\psi(\max\{|a|, |n|\})}{\max\{|a|, |n|\}}. \quad (1.3)$$

By fixing $\max\{|a|, |n|\}$, we limit both the numerator and the denominator. If we, as is done in the real case, merely compared based on n , the equation would trivially have infinitely many answers $a \in \mathbb{Z}$ for any $x \in \mathbb{Z}_p$ whenever $\psi(n) > 0$, as the fractions a/n would be dense in the ball of radius $|n|_p^{-1}$ around the origin, by virtue of \mathbb{Z} being dense in \mathbb{Z}_p . If we impose the condition that a/n be a reduced fraction, the argument becomes slightly more obscure, but the fractions would still be dense in \mathbb{Z}^p when $\gcd(a, n) = 1$ – just add the proper multiple of a sufficiently large power of p to a when $\gcd(a, n) > 1$. We then define the alternative p -adic Duffin–Schaeffer set as

$$\mathcal{B}^p(\psi) = \left\{ x \in \mathbb{Z}_p : (1.3) \text{ holds for infinitely many } \frac{a}{n} \in \mathbb{Q} \right\}.$$

This set was also briefly considered by Haynes [5], who showed that this set allowed for $\mu_p(\mathcal{B}^p(\psi)) \neq 0, 1$, by explicitly constructing a ψ such that the measure was p^{-1} . Kristensen and Laursen [9] revisited the set and found an uncountable collection of possible values for $\mu_p(\mathcal{B}^p(\psi))$ when ψ varies. For $p = 2$, they found that the possible values were exactly those in the set $[0, 1/2] \cup \{0\}$ and conjectured that the set of possible values would be $[0, 1/p] \cup \{1\}$ for $p > 2$. The first main result of this paper is a rejection of that conjecture by proving that the spectrum of possible values of $\mu_p(\mathcal{B}^p(\psi))$ is in fact limited to those already found in [9], which make up a Lebesgue-null set of Hausdorff dimension $\log 2 / \log p$.

Furthermore, a new paper by Badziahin and Bugeaud [1] introduces another Duffin–Schaeffer-like set of p -adic numbers, which we will denote by $\mathfrak{A}^p(\psi)$. The definition of this set is related to that of $\mathcal{B}^p(\psi)$ where equation (1.3) is replaced by

$$\left| x - \frac{a}{b} \right|_p < \frac{\psi(|ab|)}{|ab|}, \quad (1.4)$$

so that $\mathfrak{A}^p(\psi)$ becomes

$$\mathfrak{A}^p(\psi) = \left\{ x \in \mathbb{Z}_p : (1.4) \text{ holds for infinitely many } \frac{a}{b} \in \mathbb{Q} \right\}.$$

This set is not quite a generalisation of the original Duffin–Schaeffer set, but it appears to be of a similar nature. In their paper, Badziahin and Bugeaud prove a theorem inspired by the above theorem due to Khintchine and by the Duffin–Schaeffer Theorem. Akin to Khintchine, they achieve their result by imposing certain restrictions on the approximation function ψ , including

some weak growth restrictions. After stating the theorem, they suggest that it may hold with these restrictions weakened or perhaps even removed. We will in this paper show that said restrictions cannot be removed entirely as the spectrum of values attainable by $\mu_p(\mathfrak{A}^p(\psi))$ is the entire unit interval when ψ is allowed to be any function onto $\mathbb{R}_{\geq 0}$.

2. Main Results

For ease of notation, we will suppress ψ when talking about the various Duffin–Schaeffer-inspired sets, in the cases where there is no ambiguity towards the underlying ψ .

The most obvious reason for the difference between \mathcal{A}^p and \mathcal{B}^p is that \mathcal{A}^p only allows fractions where the numerator is bounded by the denominator while \mathcal{B}^p allows numerators of any size relative to the denominator. As such, it appears natural to consider what happens with the remaining fractions, where the numerator is greater than the denominator. This leads to a third p -adic Duffin–Schaeffer set, \mathcal{C}^p , which we define as the limsup set of sets \mathcal{C}_n^p given by

$$\mathcal{C}_n^p = \bigcup_{\substack{|a| < n \\ \gcd(a,n)=1}} B_{\mathbb{Q}_p} \left(\frac{n}{a}, \frac{\psi(n)}{n} \right) \cap \mathbb{Z}_p,$$

for $n > 1$, and $\mathcal{C}_1^p = \emptyset$ for $n = 1$. It is then easy to see that \mathcal{B}^p is the limsup set of $\mathcal{B}_n^p := \mathcal{A}_n^p \cup \mathcal{C}_n^p$. By Dirichlet’s pigeonhole principle, we then get

$$\mathcal{B}^p = \mathcal{A}^p \cup \mathcal{C}^p. \quad (2.1)$$

If $p \nmid n$ and $p \mid a$, then $|n/a|_p > 1$, so that $B(n/a, \psi(n)/n) \cap \mathbb{Z}_p = \emptyset$ unless $\psi(n)/n \geq |a|_p^{-1} > 1$, in which case $\mathcal{C}_n^p = \mathbb{Z}_p = B(n/1, \psi(n)/n) \cap \mathbb{Z}_p$. By this realization, we may instead write

$$\mathcal{C}_n^p(\psi) = \bigcup_{\substack{|a| < n \\ \gcd(a, pn)=1}} B_{\mathbb{Q}_p} \left(\frac{n}{a}, \frac{\psi(n)}{n} \right) \cap \mathbb{Z}_p,$$

Note that this is also applicable for $n = 1$, as the union is empty in that case. We will therefore use this as the *de facto* definition of $\mathcal{C}_n^p(\psi)$ going forward.

We are now ready to state the first main result of this paper, which determines the spectra of values for $\mu_p(\mathcal{C}^p)$ and $\mu_p(\mathcal{B}^p)$, respectively.

THEOREM 2.1. *Let $x \in [0, 1]$. There exists a function $\psi : \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$ such that $\mu_p(\mathcal{C}^p(\psi)) = x$ if, and only if, x is of the form*

$$x = \sum_{k=0}^{\infty} x_k (p-1)p^{-k-1}, \quad x_k \in \{0, 1\} \quad \forall k \in \mathbb{N}_0.$$

There exists a function $\psi : \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$ such that $\mu_p(\mathcal{B}^p(\psi)) = x$ if, and only if, $x = 1$ or x is of the form

$$x = \sum_{k=1}^{\infty} x_k (p-1)p^{-k-1}, \quad x_k \in \{0, 1\} \quad \forall k \in \mathbb{N}.$$

We now turn our attention towards the set \mathfrak{A}^p from [1]. In order to have notation more in line with the Duffin–Schaeffer sets \mathcal{A} and \mathcal{B}^p , we alter equation (1.4) slightly,

$$\left| x - \frac{a}{b} \right|_p \leq \frac{\psi(|ab|)}{|ab|}, \quad (2.2)$$

which leads to a set

$$\mathfrak{A}^p(\psi) = \left\{ x \in \mathbb{Z}_p : (1.4) \text{ holds for infinitely many } \frac{a}{b} \in \mathbb{Q} \right\}.$$

To see that this set will have the same spectrum of measures as \mathfrak{A}^p , define ψ' by

$$\psi'(n) = \begin{cases} p\psi(n) & \text{if } \psi(n) = n/p^k \text{ for some } k \in \mathbb{Z}, \\ \psi(n) & \text{otherwise.} \end{cases}$$

Note that

$$\left| x - \frac{a}{n} \right|_p \leq 0$$

has at most 1 solution $a/n \in \mathbb{Q}$ for any given x , and so the collection of n with $\psi(n) = 0$ contributes to neither $\mathfrak{A}^p(\psi)$ nor $\mathfrak{A}^p(\psi')$. For $\psi(n) > 0$, equation (1.4) with ψ' is equivalent to equation (2.2) with ψ . It thus follows that $\mathfrak{A}^p(\psi) = \mathfrak{A}^p(\psi')$. Since there is an obvious bijection between ψ and ψ' , \mathfrak{A}^p and \mathfrak{A}^p have the same spectrum of measures.

We will also consider a Khintchine-like variant of the set, defined as

$$\mathfrak{K}^p(\psi) = \{ x \in \mathbb{Z}_p : (2.2) \text{ holds for infinitely many } (a, b) \in \mathbb{Z} \times \mathbb{Z} \setminus 0 \}.$$

By taking \mathbb{Z}_{∞} , \mathbb{Q}_{∞} , $|\cdot|_{\infty}$, and μ_{∞} to mean $[0, 1]$, \mathbb{R} , $|\cdot|$, and the Lebesgue measure, respectively, we define related sets over the real numbers when allowing $p = \infty$.

As is the case for the ‘proper’ Duffin–Schaeffer sets, we will want to write \mathfrak{A}^p and \mathfrak{K}^p as the limsup set of sets \mathfrak{A}_n^p and \mathfrak{K}_n^p , respectively. Write $B_{\mathbb{Q}_p}(\pm a, r) = B_{\mathbb{Q}_p}(a, r) \cup B_{\mathbb{Q}_p}(-a, r)$ and $n = \prod_{i=1}^{\omega(n)} p_i^{\nu_{p_i}(n)}$, where $\omega(n)$ denotes the number

of distinct prime divisors of n . By defining

$$\mathfrak{R}_n^p(\psi) = \bigcup_{a|n} B_{\mathbb{Q}_p} \left(\pm \frac{a}{n/a}, \frac{\psi(n)}{n} \right) \cap \mathbb{Z}_p,$$

$$\mathfrak{A}_n^p(\psi) = \bigcup_{a_1, \dots, a_{\omega(n)} \in \{0,1\}} B_{\mathbb{Q}_p} \left(\pm \frac{\prod_{a_i=1} p_i^{\nu_{p_i}(n)}}{\prod_{a_i=0} p_i^{\nu_{p_i}(n)}}, \frac{\psi(n)}{n} \right) \cap \mathbb{Z}_p,$$

we immediately achieve

$$\mathfrak{R}^p = \limsup_{n \rightarrow \infty} \mathfrak{R}_n^p, \quad \mathfrak{A}^p = \limsup_{n \rightarrow \infty} \mathfrak{A}_n^p.$$

It so happens that the spectra of values for $|\mathfrak{A}^\infty|$ and $|\mathfrak{R}^\infty|$ are surprisingly easy to settle while the spectra of values for $\mu_p(\mathfrak{A}^p)$ and $\mu_p(\mathfrak{R}^p)$ when $p < \infty$ require significantly more care. However, the spectra are nonetheless independent of p as they always take up the entire unit interval as seen by the below theorem. We will prove the easy case of $p = \infty$ immediately and save $p < \infty$ for Section 5. That section will be independent of Sections 3 and 4.

THEOREM 2.2. *Let p be a prime or $p = \infty$, and let $x \in [0, 1]$. Then there exists a $\psi : \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$ such that $\mu_p(\mathfrak{A}^p(\psi)) = \mu_p(\mathfrak{R}^p(\psi)) = x$.*

PROOF FOR $p = \infty$. Let $x \in [0, 1]$ and define $\psi : \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$ by

$$\psi(n) = \begin{cases} nx & \text{if } n \text{ is a prime} \\ 0 & \text{otherwise} \end{cases}$$

Since ψ is supported on the primes, we immediately have that $\mathfrak{A}^\infty = \mathfrak{R}^\infty$. We are thus left with showing that $|\mathfrak{A}^\infty| = x$. Upon applying the Borel-Cantelli Lemma, this follows by a brief calculation, where we use q to denote primes:

$$\begin{aligned} |\mathfrak{A}^\infty| &= \left| \limsup_{q \rightarrow \infty} \bigcup_{e_1, e_2 \in \{-1, 1\}} B_{\mathbb{R}}(e_1 q^{e_2}, x) \cap [0, 1] \right| \\ &= \left| \limsup_{q \rightarrow \infty} [0, x + q^{-1}] \right| = x. \end{aligned}$$

This completes the proof

Note that $\mathfrak{A}_n^p(\psi) \subseteq \mathfrak{R}_n^p(\psi)$ will hold with strict inclusion for all small enough ψ , so we should not expect equality between $\mu_p(\mathfrak{A}^p)$ and $\mu_p(\mathfrak{R}^p)$ in general. In fact, if we in the above proof had taken

$$\psi(n) = \begin{cases} nx & \text{if } n \text{ is the square of a prime,} \\ 0 & \text{otherwise,} \end{cases}$$

we would still find $|\mathfrak{A}^\infty| = x$ while $|\mathfrak{R}^p| = \min\{2x, 1\}$ as \mathfrak{R}^p now also accepts infinitely many copies of $1 = q/q$ as approximants and thereby includes the interval $[1 - x, 1]$ in \mathfrak{R}^p .

3. Some p -adic measure theory

We will in this section present a series of general measure theoretical results that will be used for the proof of Theorem 2.1. Except for Lemma 3.5, these results all appear to have been applied to some extent in [5], even though only Lemma 3.2 was formally stated. All results are to be applied in proving Proposition 4.2, which will be introduced in Section 4 and plays a central role in proving the ‘only if’ parts of Theorem 2.1.

Note that each non-negative real number $x \in \mathbb{R}_{\geq 0}$ has a canonical base p expansion of the form

$$x = \sum_{n=N}^{\infty} a_n p^{-n},$$

where $a_n \in \{0, \dots, p-1\}$, $\liminf a_n < p-1$, and $N \in \mathbb{Z}_{\leq 0}$ is maximal. Throughout this paper, the function $\iota_p : \mathbb{R}_{\geq 0} \rightarrow \mathbb{Q}_p$ denotes the associated map

$$x = \sum_{n=N}^{\infty} a_n p^{-n} \mapsto \iota_p(x) = \sum_{n=N}^{\infty} a_n p^n.$$

This function is measure preserving in that $\lambda(A) = p\mu_p(\iota_p(A))$ for any measurable subset $A \subseteq \mathbb{R}$.

LEMMA 3.1. *The preimage map ι_p^{-1} maps balls $B \subseteq \mathbb{Q}_p$ to half-open interval in \mathbb{R} of length $p\mu_p(B)$. In particular, ι_p is measurable, and the associated push-forward measure on the Lebesgue measure, $\iota_{p\#}$, is equal to $p\mu_p$.*

PROOF. Notice that

$$\begin{aligned} \iota_p^{-1}(\mathbb{Z}_p) &= \left\{ \sum_{m=0}^{\infty} b_m p^{-m} \in \mathbb{R} : b_m \in \{0, \dots, p-1\}, \liminf_{m \rightarrow \infty} b_m < p-1 \right\} \\ &= [0, p). \end{aligned}$$

Let B be a ball in \mathbb{Z}_p and write $B = \sum_{m=0}^{M-1} b_m p^m + p^M \mathbb{Z}_p$ for some $M \in \mathbb{N}_0$ and $b_0, \dots, b_{M-1} \in \{0, \dots, p-1\}$. We then have

$$\iota_p^{-1}(B) = \iota_p^{-1} \left(\sum_{m=0}^{M-1} b_m p^m + p^{-M} \iota_p^{-1}(\mathbb{Z}_p) \right) = \sum_{m=0}^{M-1} b_m p^{-m} + [0, p^{1-M}),$$

which proves the first part of the lemma. Since the Borel algebra of \mathbb{Z}_p is generated by the collection of balls in \mathbb{Z}_p , and the intervals are contained

in the Borel algebra of $\mathbb{R}_{\geq 0}$, ι_p is a measurable function, and the the push-forward $\iota_{p\#}$ is a Borel-measure on \mathbb{Q}_p . From the above equation, we notice in particular that for all balls B in \mathbb{Q}_p ,

$$\iota_{p\#}(B) = p\mu_p(B).$$

Since all balls have finite measure, and the collection of the empty set and all balls in \mathbb{Q}_p is preserved under pairwise intersection and generates the Borel algebra of \mathbb{Q}_p , this implies that $\iota_{p\#} = p\mu_p$, by the Uniqueness of Measures Theorem, and the proof is complete.

The alternative definition of μ_p provided by the above lemma gives a tool to translate measure theoretic results regarding \mathbb{R} into measure theoretic results regarding \mathbb{Q}_p . It appears that Haynes may have used this alternative definition in [5], though it was not explicitly stated. One result that very easily translates using ι_p is the below lemma from [4], which is a modification of a lemma from [2].

LEMMA 3.2. *Let $\{I_n\}_{n \in \mathbb{N}}$ be a sequence of real intervals with $\lambda(I_n) \xrightarrow[n \rightarrow \infty]{} 0$, and let $\{U_n\}_{n \in \mathbb{N}}$ be a sequence of measurable sets such that*

$$U_n \subseteq I_n, \quad |U_n| \geq \epsilon |I_n| \quad \forall n \in \mathbb{N},$$

for some fixed $0 < \epsilon < 1$. Then $|\limsup_{n \rightarrow \infty} U_n| = |\limsup_{n \rightarrow \infty} I_n|$.

COROLLARY 3.3. *Let $\{B_n\}_{n \in \mathbb{N}}$ be a sequence of p -adic balls with $\mu_p(B_n) \xrightarrow[n \rightarrow \infty]{} 0$. Suppose $\{U_n\}_{n \in \mathbb{N}}$ is a sequence of measurable sets such that, for some*

$$U_n \subseteq B_n, \quad \mu_p(U_n) \geq \epsilon \mu_p(B_n) \quad \forall n \in \mathbb{N},$$

for some fixed $0 < \epsilon < 1$. Then $\mu_p(\limsup_{n \rightarrow \infty} U_n) = \mu_p(\limsup_{n \rightarrow \infty} B_n)$.

PROOF. Applying $p\mu_p = \iota_{p\#}$ by Lemma 3.1, we have

$$\begin{aligned} p\mu_p\left(\limsup_{n \rightarrow \infty} U_n\right) &= \left| \iota_p^{-1} \left(\bigcap_{N=1}^{\infty} \bigcup_{n \geq N} U_n \right) \right| = \lim_{N \rightarrow \infty} \left| \iota_p^{-1} \left(\bigcup_{n \geq N} U_n \right) \right| \\ &= \left| \limsup_{n \rightarrow \infty} \iota_p^{-1}(U_n) \right|. \end{aligned}$$

By repeating the process for B_n , the statement follows by Lemma 3.2, and the proof is complete.

Another relevant result that may be derived using ι_p is a p -adic version of the Lebesgue Density Theorem, as presented below.

LEMMA 3.4 (Lebesgue Density Theorem for \mathbb{Q}_p). *Let A be a measurable subset of \mathbb{Q}_p such that $\mu_p(A) > 0$. Then A contains a point of density 1, which is to say that there exists $a \in A$ such that if $\{B_M\}_{M \in \mathbb{N}}$ is a sequence of balls $B_M \ni a$ of radius $r(B_M) \xrightarrow{M \rightarrow \infty} 0$, then*

$$\frac{\mu_p(A \cap B_M)}{\mu_p(B_M)} \xrightarrow{M \rightarrow \infty} 1.$$

PROOF. By Lemma 3.1, we have $p\mu_p = \iota_{p\#}$. Thus $\tilde{A} = \iota_p^{-1}(A)$ has positive Lebesgue measure, which means that it must contain a point $x \in \tilde{A}$ of density 1, by the Lebesgue Density Theorem. Put $a = \iota_p(x)$, let $\{B_M\}$ be a collection of balls around a in \mathbb{Q}_p of radius $r(B_M) \xrightarrow{M \rightarrow \infty} 0$. Let $\varepsilon > 0$, and put $\tilde{B}_M = \iota_p^{-1}(B_M)$. Since ι_p^{-1} maps balls to intervals, and $\tilde{B}_M \ni x$, we must have that $|\tilde{A} \cap \tilde{B}_M| > 0$. We may thus pick open intervals $I_M \supseteq \tilde{B}_M$ such that $|\tilde{A} \cap I_M| \leq p^\varepsilon |\tilde{A} \cap \tilde{B}_M|$ and $|I_M| \leq p^\varepsilon |\tilde{B}_M|$. Then $I_M \ni x$ and $|I_M| \leq p^{1+\varepsilon} \mu_p(B_M) \xrightarrow{\varepsilon \rightarrow 0} 0$. Since x is of density 1, we then have

$$\frac{\mu_p(A \cap B_M)}{\mu_p(B_M)} = \frac{\lambda(\tilde{A} \cap \tilde{B}_M)}{\lambda(\tilde{B}_M)} \geq p^{-\varepsilon} \frac{\lambda(\tilde{A} \cap I_M)}{\lambda(I_M)} \xrightarrow{M \rightarrow \infty} p^{-\varepsilon},$$

and the lemma follows by letting ε tend to 0.

We will also use the below lemma, which relies on elementary algebra and then the Uniqueness of Measures Theorem. This does not use ι_p .

LEMMA 3.5. *Let $f : \mathbb{Z}_p^\times \rightarrow \mathbb{Z}_p^\times$ denote the map $x \mapsto 1/x$. Then f maps balls in \mathbb{Z}_p^\times to balls of the same measure in \mathbb{Z}_p^\times . In particular, f preserves μ_p restricted to \mathbb{Z}_p^\times .*

PROOF. This is a simple calculation.

4. Proof of Theorem 2.1

To prove Theorem 2.1, we will use two propositions. The first one, below, puts $\mu_p(\mathcal{B}^p)$ equal to $\mu_p(\mathcal{A}^p)$ or $\mu_p(\mathcal{C}^p)$, depending on a divergence criterion, and splits \mathcal{C}^p into smaller pieces to be dealt with individually.

In the proof and for the rest of this paper, we will use \sqcup to denote the disjoint union of sets.

PROPOSITION 4.1. *Let l be the minimal $k \in \mathbb{N}_0$ such that $\psi(n)/n \geq p^{-k}$ for infinitely many $n \in p^k \mathbb{Z}_p$, if such a k exists. Otherwise, put $l = \infty$ and*

write $p^{-\infty} = 0$. Then

$$\mu_p(\mathcal{B}^p) = \begin{cases} \mu_p(\mathcal{A}^p) = 1 & \text{if } \sum_{p \nmid n} \frac{\phi(n)\psi(n)}{n} = \infty, \\ \mu_p(\mathcal{C}^p) & \text{if } \sum_{p \nmid n} \frac{\phi(n)\psi(n)}{n} < \infty, \end{cases} \quad (4.1)$$

$$\mu_p(\mathcal{C}^p) = p^{-l} + \sum_{0 \leq k < l} \mu_p(\mathcal{C}^p \cap p^k \mathbb{Z}_p^\times). \quad (4.2)$$

Furthermore, if $k < l$, then $\mathcal{C}^p \cap p^k \mathbb{Z}_p^\times = \limsup_{\nu_p(n)=k} \mathcal{C}_n^p$.

PROOF. We start by proving equation (4.1). If $\sum_{p \nmid n} \frac{\phi(n)\psi(n)}{n} = \infty$, this is [9, Theorem 2], so suppose not. Then $\sum_{p \nmid n} \mu_p(\mathcal{A}_n^p) < \infty$. If $\sum_{p \mid n} \mu_p(\mathcal{A}_n^p) = \infty$, then we must have $\psi(n) \geq pn$ infinitely often, and so $\mathcal{B}^p = \mathcal{C}^p = \mathbb{Z}_p$. If $\sum_{p \mid n} \mu_p(\mathcal{A}_n^p) < \infty$, then the Borel-Cantelli Lemma implies $\mu_p(\mathcal{A}^p) = 0$, and the statement follows by equation (2.1).

Moving on to equation (4.2), notice that

$$\mathcal{C}^p = (\mathcal{C}^p \cap p^l \mathbb{Z}_p) \sqcup \bigsqcup_{0 \leq k < l} \mathcal{C}^p \cap p^k \mathbb{Z}_p^\times,$$

with the convention of $p^\infty = 0$ (as an element of \mathbb{Z}_p) in the case of $l = \infty$. Equation (4.2) is then equivalent to $\mu_p(\mathcal{C}^p \cap p^l \mathbb{Z}_p) = p^l$. If $l = \infty$, this is trivial, so suppose $l < \infty$. Let $\alpha \in p^l \mathbb{Z}_p$ and $n \in p^l \mathbb{Z}_p$. Then clearly $|\alpha - n/1|_p \leq p^{-l}$. Since $\psi(n)/n \geq p^{-l}$ for infinitely many such n , we must have $\mathcal{C}^p \supseteq p^l \mathbb{Z}_p$, and so $\mu_p(\mathcal{C}^p \cap p^l \mathbb{Z}_p) = p^l$.

As for the final part of the statement, this is vacuous if $l = 0$, so suppose $l > 0$ and let $0 \leq j \leq k < l$. Then $\psi(n)/n < p^{-j}$ for all but finitely many $n \in p^j \mathbb{N}$. If $\nu_p(n) = j$, then $\alpha \in \mathcal{C}_n^p$ would imply $|\alpha|_p = |\alpha - \frac{n}{a} + \frac{n}{a}|_p = p^{-j}$ for some $\gcd(a, pn) = 1$ when n is sufficiently large, so that

$$\limsup_{\nu_p(n)=j} \mathcal{C}_n^p \subseteq p^j \mathbb{Z}_p^\times.$$

If $\nu_p(n) > k$, we would instead find $|\alpha|_p \leq \max\{|\alpha - \frac{n}{a}|, |\frac{n}{a}|_p\} < p^{-k}$, i.e.,

$$\limsup_{\nu_p(n)>k} \mathcal{C}_n^p \subseteq p^{k+1} \mathbb{Z}_p.$$

By the pigeon-hole principle, the proposition is proven upon calculating

$$\begin{aligned} \mathcal{C}^p \cap p^k \mathbb{Z}_p^\times &= \left(\limsup_{\nu_p(n)>k} \mathcal{C}_n^p \cap p^k \mathbb{Z}_p^\times \right) \cup \bigcup_{j=0}^k \limsup_{\nu_p(n)=j} \mathcal{C}_n^p \cap p^k \mathbb{Z}_p^\times \\ &= \limsup_{\nu_p(n)=k} \mathcal{C}_n^p. \end{aligned}$$

The other proposition leading to Theorem 2.1 is a bit more involved and may be thought of as a shell-wise zero-full law for \mathcal{C}^p . This relates to, and is inspired by, the zero-one laws $|\mathcal{A}| \in \{0, 1\}$ [4] and $\mu_p(\mathcal{A}^p) \in \{0, 1\}$ [5].

PROPOSITION 4.2. *Let $k \in \mathbb{N}_0$. Then $\mu_p(\mathcal{C}^p \cap p^k \mathbb{Z}_p^\times) \in \{0, (p-1)/p^{k+1}\}$.*

The proof of this proposition follows the same overall structure as the proof of the 0-1 law in [5], with some modifications. In that light, it is perhaps not surprising that we will need the below lemma, which corresponds to the less trivial part of [5, Lemma 2]. We will here continue to use $\omega(n)$ to denote the number of prime divisors of n . The proof of the lemma will apply the Möbius μ function, which is defined as $\mu(d) = (-1)^{\omega(d)}$ when d is square free, and $\mu(d) = 0$ otherwise. In this context, the following three facts, which can be found in [11], will be applied without proof.

$$\sum_{d|n} \mu(d) = \begin{cases} 1 & \text{if } d = 1, \\ 0 & \text{if } d \neq 1, \end{cases} \quad (4.3)$$

$$\sum_{d|n} \mu(d) \frac{n}{d} = \phi(n), \quad (4.4)$$

$$\frac{\phi(n)}{n} = \prod_{q|n} (1 - q^{-1}). \quad (4.5)$$

LEMMA 4.3. *If $n > 1$ and $\psi(n) > 4^{\omega(n)}$, then $\mathcal{C}_n^p \supseteq p^{\nu_p(n)} \mathbb{Z}_p^\times$.*

PROOF. Put $k = \nu_p(n)$, and let $\alpha \in p^k \mathbb{Z}_p^\times$. Then $n/\alpha \in \mathbb{Z}_p^\times$, and we write

$$\frac{n}{\alpha} = \sum_{m=0}^{\infty} b_m p^m, \quad b_m \in \{0, \dots, p-1\} \forall m \in \mathbb{N}_0, \quad b_0 \neq 0.$$

If $\psi(n) \geq n/p^k$, then clearly $\alpha \in \mathcal{C}_n^p$ since $|n|_p = |\alpha|_p = p^{-k}$ implies $|\alpha - \frac{n}{1}|_p \leq p^{-k}$, so suppose not. Pick $N \in \mathbb{Z}$ such that $\psi(n)/n \in [p^{-N}, p^{-N+1})$. Note that $N > k$. Our job of proving $\alpha \in \mathcal{C}_n^p$ then reduces to finding an a with $|a| < n$ and $\gcd(a, pn) = 1$ such that $\nu_p(\alpha - n/a) \geq N$. As $\nu_p(\alpha) = k$ and $p \nmid a$, the latter part is equivalent to $\nu_p(a - n/\alpha) \geq N - k$. The proof is then complete if we find an a with $|a| < n$ and $\gcd(a, pn) = 1$ such that

$$a = \frac{n}{\alpha} + \sum_{m=N-k}^{\infty} c_m p^m = \sum_{m=0}^{N-k-1} b_m p^m + \sum_{m=N-k}^{\infty} (b_m + c_m) p^m,$$

with $c_m \in \{0, \dots, p-1\}$ for all m . Write $b = \sum_{m=0}^{N-k-1} b_m p^m$. Since $p \nmid b$ as $b_0 \neq 0$ and $N > k$, all elements of the set

$$A = \{a \in \mathbb{Z} : |a| < n, \gcd(a, n) = 1, a \equiv b \pmod{p^{N-k}}\}$$

satisfy these criteria, and so we are done if we can show that $\#A > 0$. By equation (4.3), we have

$$\begin{aligned} \#A &= \sum_{\substack{|a| < n \\ a \equiv b \pmod{p^{N-k}}} } \sum_{d | \gcd(a, n)} \mu(d) = \sum_{\substack{d | n \\ p \nmid d}} \sum_{\substack{|a| < n, d | a \\ a \equiv b \pmod{p^{N-k}}} } \mu(d) \\ &= \sum_{d | n/p^k} \mu(d) \sum_{\substack{|l| < n/d \\ l \equiv bd^{-1} \pmod{p^{N-k}}} } 1. \end{aligned}$$

To simplify notation, let $\tilde{n} = n/p^k$. For each $d | \tilde{n}$, pick $x_d \in \mathbb{Z}$ such that $x_d \equiv bd^{-1} \pmod{p^{N-k}}$, and let k_d denote the difference $\#((-n/d, n/d) \cap (x_d + p^{N-k}\mathbb{Z})) - 2n/(p^{N-k}d)$. Then

$$\begin{aligned} \#A &= \sum_{d | \tilde{n}} \mu(d) \sum_{\substack{|l| < n/d \\ l \in x_d + p^{N-k}\mathbb{Z}}} 1 = \sum_{d | \tilde{n}} \mu(d) \left(\frac{2n}{p^{N-k}d} + k_d \right) \\ &= \frac{2p^k}{p^{N-k}} \sum_{d | \tilde{n}} \mu(d) \frac{\tilde{n}}{d} + \sum_{d | \tilde{n}} \mu(d) k_d = 2p^{-N} p^{2k} \phi(\tilde{n}) + \sum_{d | \tilde{n}} \mu(d) k_d \\ &\geq 2p^k \frac{\psi(n)}{n} \phi(n) + \sum_{d | \tilde{n}} \mu(d) k_d, \end{aligned}$$

where the final equality and the inequality follow from equation (4.4) and the choices of N and k , respectively. Notice that $|k_d| \leq 1$, which combined with assumption $\psi(n) > 4^{\omega(n)}/n$ implies

$$\#A > 2p^k \frac{4^{\omega(n)}}{n} \phi(n) - \sum_{d | \tilde{n}} |\mu(d)| = 2p^k \frac{\phi(n)}{n} 4^{\omega(n)} - 2^{\omega(\tilde{n})}.$$

By equation (4.5), we have

$$2 \frac{\phi(n)}{n} 4^{\omega(n)} = 2 \prod_{\substack{q | n \\ q \text{ prime}}} 4(1 - q^{-1}) \geq 2 \prod_{\substack{q | n \\ q \text{ prime}}} 2 = 2^{\omega(n)+1},$$

and so $\#A > 2^{\omega(n)} > 0$. We conclude $\alpha \in \mathcal{C}_n^p$ and thus $\mathcal{C}_n^p \supseteq p^k \mathbb{Z}_p^\times$, and the proof is complete.

We will also need the below lemma, which is essentially a specialisation of Corollary 3.3 in the context of \mathcal{C}^p .

LEMMA 4.4. *Let $k \in \mathbb{N}_0$, and suppose $\text{supp}(\psi) \subseteq p^k \mathbb{N} \setminus p^{k+1} \mathbb{N}$. Then $\mu_p(\mathcal{C}^p(x\psi) \cap p^k \mathbb{Z}_p^\times) = \mu_p(\mathcal{C}^p(\psi) \cap p^k \mathbb{Z}_p^\times)$ for all $x > 0$.*

PROOF. There is nothing to prove if $x = 1$ or if $\psi(n) > 0$ for only finitely many n , so suppose neither is the case. By replacing ψ with $\psi' = x\psi$ and x by $x' = 1/x$ if necessary, we may assume without loss of generality that $x \in (0, 1)$. If $x\psi(n) \geq 4^{\omega(n)}$ infinitely often, then Lemma 4.3 implies that $\mathcal{C}^p(\psi), \mathcal{C}^p(x\psi) \supseteq p^k \mathbb{Z}_p^\times$, so suppose not. Then, for n sufficiently large,

$$\frac{\psi(n)}{n} \leq x^{-1} \frac{4^{\omega(n)}}{n} \leq 16x^{-1} \frac{4^{\log_5(n)}}{n} \xrightarrow{n \rightarrow \infty} 0.$$

Since $\mu_p(\mathcal{C}_n^p(\psi)) = 0$ when $\psi(n) = 0$, we have by the Borel–Cantelli Lemma that

$$\begin{aligned} \mu_p(\mathcal{C}^p(\psi)) &= \mu_p\left(\limsup_{\psi(n) > 0} \mathcal{C}_n^p(\psi)\right), \\ \mu_p(\mathcal{C}^p(x\psi)) &= \mu_p\left(\limsup_{\psi(n) > 0} \mathcal{C}_n^p(x\psi)\right). \end{aligned}$$

Notice, for $\psi(n) > 0$, that $\mathcal{C}_n^p(\psi)$ is a finite union of proper balls $B_{i_n}, \dots, B_{i_{n+1}-1}$ of radius $\psi(n)/n \xrightarrow{n \rightarrow \infty} 0$, and that each ball B_{i_n+j} is matched one-to-one by a ball U_{i_n+j} from $\mathcal{C}_n^p(x\psi)$ satisfying

$$U_{i_n+j} \subseteq B_{i_n+j}, \quad \mu_p(U_{i_n+j}) \geq \frac{x}{p} \mu_p(B_{i_n+j}).$$

From Corollary 3.3, it follows that $\mu_p(\mathcal{C}^p(x\psi)) = \mu_p(\mathcal{C}^p(\psi))$. This completes the proof since clearly $\mathcal{C}^p(x\psi) \subseteq \mathcal{C}^p(\psi)$.

We are now ready to complete the proof of the shell-wise p -adic zero-full law. The remaining part of the proof is where it differs the most from the proof of the 0-1 law in [5], though it still follows the same overall idea.

PROOF OF PROPOSITION 4.2. Let l be defined as in Proposition 4.1. If $k \geq l$, it follows from the proof of that proposition that $\mathcal{C}_n^p(\psi) \supseteq p^l \mathbb{Z}_p^\times$, and we are done, so suppose $k > l$. We then have that $\mathcal{C}^p(\psi) \cap p^k \mathbb{Z}_p^\times = \limsup_{\nu_p(n)=k} \mathcal{C}_n^p(\psi)$, and we may hence assume, without loss of generality, that $\text{supp}(\psi) \subseteq p^k \mathbb{N} \setminus p^{k+1} \mathbb{N}$. Following the arguments in the proof of Lemma 4.4, $\limsup_{n \rightarrow \infty} \psi(n)/n > 0$ would likewise imply $\mathcal{C}_n^p(\psi) \supseteq p^k \mathbb{Z}_p^\times$, so suppose $\limsup_{n \rightarrow \infty} \psi(n)/n = 0$. We then have, for any fixed j , that

$$\frac{\psi(n)}{n} < p^{-j} \quad \text{for all } n \geq N_j, \text{ for some } N_j \in \mathbb{N}. \quad (4.6)$$

Let $\tau_1 : \mathbb{Z}_p^\times \rightarrow \mathbb{Z}_p^\times$ and $\tau_2 : p^k \mathbb{Z}_p^\times \rightarrow p^k \mathbb{Z}_p^\times$ be given by

$$\tau_1(b) = \begin{cases} \sum_{m=0}^{\infty} b_{m+1} p^m & \text{if } b_1 \neq 0, \\ 1 + \sum_{m=0}^{\infty} b_{m+1} p^m & \text{if } b_1 = 0, \end{cases}$$

$$\tau_2(p^k b) = p^k / \tau_1(b),$$

for $b = \sum_{m=0}^{\infty} b_m p^m \in \mathbb{Z}_p^\times$. For $K \geq 2$ and $b = \sum_{m=0}^{K-1} b_m p^m \in \mathbb{Z}_p^\times$, note that

$$\tau_1(b + p^K \mathbb{Z}_p) = \tau_1(b_1) + \sum_{m=1}^{K-2} b_{m+1} p^m + p^{K-1} \mathbb{Z}_p. \quad (4.7)$$

Thus, τ_1 maps balls of centre b and radius p^{-K} to balls of centre b and radius p^{1-K} when $K \geq 2$. This makes any restriction of τ_1 to a ball in \mathbb{Z}_p^\times of radius at most p^{-2} into a homeomorphism onto its image as it is clearly a bijection under such a restriction. By Lemma 3.5, these properties extend to a restriction of τ_2 when replacing K by $M \geq k+2$. Let B be a ball of radius at most $k+2$. The inverse of τ_2 restricted to B , $\tau_2|_B^{-1}$ is thus measurable and has a push-forward measure, $(\tau_2|_B^{-1})_{\#} \mu$, satisfying

$$(\tau_2|_B^{-1})_{\#} \mu(\tilde{B}) = \mu_p(\tau_2(\tilde{B})) = p \mu_p(\tilde{B}),$$

for all balls $\tilde{B} \subseteq \tau_2(B)$. By the proof of Lemma 3.1, this means that $(\tau_2|_B^{-1})_{\#} \mu = p \mu_p$, i.e.,

$$\mu_p(\tau_2(A)) = p \mu_p(A), \quad (4.8)$$

for all Borel subsets $A \subseteq B$.

Let $\alpha \in \mathcal{C}^p(\psi)$ and write

$$\alpha' := p^k / \alpha = \sum_{m=0}^{\infty} a_m p^m \in \mathbb{Z}_p^\times, \quad a_i \in \{0, \dots, p-1\}.$$

Let $n \in \mathbb{N}$ such that $\alpha \in \mathcal{C}_n^p(\psi)$. Since $\lim_{n \rightarrow \infty} \psi(n)/n = 0$, we have $\psi(n)/n \leq p^{-k-2}$ for n large enough. By the assumption on the support of ψ , we may write $n = p^k n'$ for some $n' \in \mathbb{N} \setminus p\mathbb{N}$ and pick some $|a| < |n|$ with $\gcd(a, pn) = 1$ such that

$$|n' \alpha' - a|_p = \left| \frac{n}{\alpha} - a \right|_p = p^k \left| \alpha - \frac{n}{a} \right|_p \leq p^k \frac{\psi(n)}{n}.$$

If $a_1 \neq 0$, put $a' = (a - a_0 n')/p$. Then

$$\begin{aligned} \left| \frac{n}{\tau_2(\alpha)} - a' \right|_p &= \left| n' \tau_1(\alpha') - \frac{a - a_0 n'}{p} \right|_p = \left| \frac{n'(\alpha' - a_0) - (a - a_0 n')}{p} \right|_p \\ &= p |n' \alpha' - a|_p \leq p^{k+1} \frac{\psi(n)}{n}, \\ |a'| &= \left| \frac{a - a_0 n'}{p} \right| \leq \frac{|a| + (p-1)n}{p} < n, \\ a' &= \frac{a - a_0 n'}{p} = \frac{a - (a_0 + p a_1) n'}{p} + a_1 n'. \end{aligned}$$

Since $p^2 \mid a - \alpha' n'$, we have $p \mid (a - (a_0 + p a_1) n')/p \in \mathbb{Z}$. From the last equation, we can thus deduce that $\gcd(a', p n') = \gcd(a', p n) = 1$, as $p \nmid a_1 n'$ and $\gcd(a, n') = 1$. Combined with the other two equations, we conclude that $\tau_2(\alpha) \in \mathcal{C}_n^p(p\psi)$. If $a_1 = 0$, we put $a' = (a + (p - a_0) n')/p$ and reach the same conclusion, based on similar calculations. Hence, $\tau_2(\mathcal{C}^p(\psi)) \subseteq \mathcal{C}^p(p\psi)$. By induction, we then have

$$\tau_2^j(\mathcal{C}^p(\psi)) \subseteq \mathcal{C}^p(p^j \psi), \quad (4.9)$$

for all $j \in \mathbb{N}$, where τ_2^j denotes the composition of j copies of τ_2 .

If $\mu_p(\mathcal{C}^p(\psi)) = 0$, we are done, so suppose that $\mu_p(\mathcal{C}^p(\psi)) > 0$. Then Lemma 3.4 implies that $\mathcal{C}^p(\psi)$ contains a point α of density 1, i.e.,

$$\frac{\mu_p(\mathcal{C}^p(\psi) \cap B_M(\alpha))}{\mu_p(B_M(\alpha))} \xrightarrow{M \rightarrow \infty} 1,$$

where we use $B_M(\alpha)$ as short-hand notation of the ball $B_{\mathbb{Q}_p}(\alpha, p^{-M}) = \alpha + p^M \mathbb{Z}_p$. Let $\varepsilon > 0$ and pick $M \geq k + 2$ such that

$$\mu_p(\mathcal{C}^p(\psi) \cap B_M(\alpha)) \geq (1 - \varepsilon) \mu_p(B_M(\alpha)) = (1 - \varepsilon) p^{-M}.$$

Pick $x = p^k \sum_{m=0}^{M-k-1} x_m p^m \in \mathbb{N}$ such that $B_M(x) = B_M(\alpha)$. Put $A = \tau_2^{M-k-1}(\mathcal{C}^p \cap B_M(x))$. By Lemma 4.4 and equations (4.7) and (4.9), we have

$$\begin{aligned} \mu_p(\mathcal{C}^p(\psi) \cap p^k \mathbb{Z}_p^\times) &= \mu_p(\mathcal{C}^p(p^{M-k} \psi) \cap p^k \mathbb{Z}_p^\times) \\ &\geq \mu_p(\tau_2^{M-k}(\mathcal{C}^p(\psi) \cap B_M(x))) = \mu_p(\tau_2(A)) \end{aligned} \quad (4.10)$$

Let $\tilde{x} = \tau_2^{M-k-1}(x)$ and note that $\tilde{x} = p^k \tilde{x}_0$ for some $\tilde{x}_0 \in \{1, \dots, p-1\}$. Then

$$A \subseteq B_{k+1}(\tilde{x}) = p^k \bigsqcup_{i=0}^{p-1} (\tilde{x}_0 + ip + p^2 \mathbb{Z}_p).$$

By applying equation (4.8) and then later the above inclusion, we find

$$\begin{aligned}
\mu_p(\tau_2(A)) &\geq \sum_{i=1}^{p-1} \mu_p(\tau_2(A \cap B_{k+2}(\tilde{x} + ip))) \\
&= p \sum_{i=1}^{p-1} \mu_p(A \cap B_{k+2}(\tilde{x} + ip)) \\
&= p\mu_p(A \setminus B_{k+2}(\tilde{x})) \geq p(\mu_p(A) - p^{-k-2}) \\
&\geq (p-1)\mu_p(A)
\end{aligned}$$

Meanwhile, an iterative application of equations (4.7) and (4.8) yield

$$\mu_p(A) = p^{M-k-1} \mu_p(\mathcal{C}^p \cap B_M(x)) \geq \frac{1-\epsilon}{p^{k+1}},$$

so that $\mu_p(\mathcal{C}^p(\psi) \cap p^k \mathbb{Z}_p^\times) \geq \frac{p-1}{p^{k+1}}(1-\epsilon)$, by equation (4.10). This completes the proof.

PROOF OF THEOREM 2.1. We start by the ‘only if’ parts. For \mathcal{C}^p , it follows by Proposition 4.2 upon noting

$$\mathcal{C}^p = (\mathcal{C}^p \cap \{0\}) \sqcup \bigsqcup_{k=0}^{\infty} \mathcal{C}^p \cap p^k \mathbb{Z}_p^\times.$$

As for \mathcal{B}^p , suppose $\mu_p(\mathcal{B}^p) < 1$. By Proposition 4.1, $\sum_{p \nmid n} \phi(n)\psi(n)/n < \infty$, $\mu_p(\mathcal{B}^p) = \mu_p(\mathcal{C}^p)$, and $\mathcal{C}^p \cap \mathbb{Z}_p^\times = \limsup_{p \nmid n} \mathcal{C}_n^p$. Since

$$\sum_{p \nmid n} \mu_p(\mathcal{C}_n^p) \leq \sum_{p \nmid n} \frac{2\phi(n)\psi(n)}{n} < \infty,$$

the Borel-Cantelli Lemma implies $\mu_p(\mathcal{C}^p \cap \mathbb{Z}_p^\times) = 0$, and we are done by the above consideration of \mathcal{C}^p .

The ‘if’ parts of the theorem are already dealt with in the proof of Theorem 3 of [9] (at least for \mathcal{B}^p), but we will repeat the argument here for clarity, shortened by means of Proposition 4.1. Let $x_k \in \{0, 1\}$ for $k \in \mathbb{N}_0$ and define $\psi : \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$ by

$$\psi(n) = x_{\nu_p(n)} \frac{n}{p^{\nu_p(n)+1}}.$$

Let $k \in \mathbb{N}_0$, and let $q > p$ be some large prime. If $x_k = 0$, we clearly have $\limsup_{\nu_p(n)=k} \mathcal{C}_n^p = \emptyset$. If, on the other hand, $x_k = 1$, then $\psi(p^k q)/(p^k q) =$

p^{-k-1} , and so

$$\begin{aligned} \mathcal{C}_{p^k q}^p &= \bigcup_{\substack{|a| < p^k q \\ \gcd(a, p^{k+1}q) = 1}} B_{\mathbb{Q}_p} \left(\frac{p^k q}{a}, p^{-k-1} \right) \\ &\supseteq \bigcup_{a=1}^{p-1} p^k \frac{q}{a} + p^{k+1} \mathbb{Z}_p = p^k \bigcup_{a=1}^{p-1} \frac{q}{a} + p \mathbb{Z}_p \end{aligned}$$

Since q is a unit modulo p , and since inversion and multiplication by units only permute the set of units modulo p , this implies that $\mathcal{C}_{p^k q}^p \supseteq p^k \mathbb{Z}_p^\times$. As there are infinitely many such q , we get $\mathcal{C}^p \supseteq p^k \mathbb{Z}_p^\times$. By Proposition 4.1, we then have

$$\begin{aligned} \mu_p(\mathcal{C}^p) &= \sum_{x_k=1} \mu_p(p^k \mathbb{Z}_p^\times) = \sum_{k=0}^{\infty} x_k (p-1) p^{-k-1}, \\ \mu_p(\mathcal{B}^p) &= \begin{cases} 1 & \text{if } x_0 = 1, \\ \sum_{k=1}^{\infty} x_k (p-1) p^{-k-1} & \text{if } x_0 = 0, \end{cases} \end{aligned}$$

as the value l from the proposition is clearly infinite. This completes the proof of Theorem 2.1.

5. Proof of Theorem 2.2 for $p < \infty$

The proof of Theorem 2.2 follows the same main idea as the ‘if’ part of Theorem 2.1, though some details are different. The main difference is that additional care is needed for the choice of the support. This will rely heavily on the below theorem due to Dirichlet, which may be found in [11].

DIRICHLET’S THEOREM ON PRIMES. *Let $a, b \in \mathbb{N}$ such that $\gcd(a, b) = 1$. Then there are infinitely many primes $q \equiv a \pmod{b}$.*

To simplify the notation, the symbol \pm will be used to implicitly denote the union of the cases of $+$ and $-$, respectively, in place of \pm . We thus write

$$\begin{aligned} B(\pm a^{\pm 1}, r) &= B(a^{\pm 1}, r) \cup B(-a^{\pm 1}, r) \\ &= B(a, r) \cup B(a^{-1}, r) \cup B(-a, r) \cup B(-a^{-1}, r). \end{aligned}$$

Furthermore, we continue using \sqcup to denote the disjoint union of sets. Recall that we already proved Theorem 2.2 for $p = \infty$ in Section 2.

PROOF OF THEOREM 2.2, FOR $p < \infty$. Let $x \in [0, 1]$. If $x = 1$, the statement is trivial, so suppose not. We then write $x = \sum_{k=0}^{\infty} x_k p^{-k-1}$ where the x_k are chosen such that $\liminf_{k \rightarrow \infty} x_k < p - 1$. Pick $K = \min\{k \in \mathbb{N}_0 : x_k < p - 1\}$, and let $g \in \mathbb{N}$ be such that $g + p\mathbb{Z}_p$ generates the multiplicative group

$\mathbb{Z}_p/p\mathbb{Z}_p$. As the rest of the construction will depend on the prime p , we split into four cases. In the first two cases, which construct ψ for primes $p > 5$ according to their congruency classes modulo 4, we do not put any further restrictions on the choice of g . In the other two cases, which deal with $p = 2$ and $p = 3, 5$, respectively, we will need some further restrictions on g and therefore fix a specific value, depending on p .

Case 1: $5 < p \equiv 1 \pmod{4}$. For $a \in \{0, \dots, p-1\}$ and $k \in \mathbb{N}_0$, define

$$I_a := \begin{cases} \emptyset & \text{if } a < 4, \\ \{1, g^{\frac{p-1}{4}}\} \cup \{g^i : 2 \leq i \leq a/4\} & \text{if } 4 \leq a < p-1, \\ \{1, \dots, p-1\} & \text{if } a = p-1, \end{cases}$$

$$r_k := \begin{cases} x_k - 4 \lfloor \frac{x_k}{4} \rfloor + \sum_{l=k+1}^{\infty} x_l p^{k-l} & \text{if } x_k < p-1, \\ 0 & \text{if } x_k = p-1. \end{cases}$$

Since clearly $r_k \in [0, 4]$, we may write

$$\frac{r_k}{4} = \sum_{i=1}^{\infty} b_{k,i} p^{-i}, \quad b_{k,i} \in \{0, 1, \dots, p-1\}.$$

Based on this, we construct ψ as

$$\psi(n) := \begin{cases} f_k(q) & \text{if } n = p^k q, \text{ where } k \leq K, \\ 0 & \text{otherwise,} \end{cases}$$

where we use q to denote primes other than p , and

$$f_k(q) = \begin{cases} q/p & \text{if } q \equiv m \pmod{p}, \text{ where } m \in I_{x_k}, \\ q/p^{i+1} & \text{if } q \equiv g + b'p^i \pmod{p^{i+1}}, \\ & \text{where } 1 \leq b' \leq b_{k,i}, i \in \mathbb{N}, \\ 0 & \text{otherwise.} \end{cases}$$

Let $k \leq K$. Then

$$\begin{aligned} \mathfrak{A}_{p^k q}^p &= \begin{cases} B_{\mathbb{Q}_p}(\pm q^{\pm 1} p^k, p^{-k-1}) & \text{if } q \equiv m \pmod{p}, \text{ where } m \in I_{x_k}, \\ B_{\mathbb{Q}_p}(\pm q^{\pm 1} p^k, p^{-k-i-1}) & \text{if } q \equiv g + b'p^i \pmod{p^{i+1}}, \\ & \text{where } 1 \leq b' \leq b_{k,i}, i \in \mathbb{N}, \\ B_{\mathbb{Q}_p}(\pm q^{\pm 1} p^k, 0) & \text{otherwise} \end{cases} \\ &= p^k \begin{cases} \pm q^{\pm 1} + p\mathbb{Z}_p & \text{if } q \equiv g^i \pmod{p}, \text{ where } g^i \in I_{x_k}, \\ \pm q^{\pm 1} + p^{i+1}\mathbb{Z}_p & \text{if } q \equiv g + b'p^i \pmod{p^{i+1}}, \\ & \text{where } 1 \leq b' \leq b_{k,i}, i \in \mathbb{N}, \\ \{\pm q^{\pm 1}\} & \text{otherwise} \end{cases} \quad (5.1) \\ &\subseteq p^k \mathbb{Z}_p^\times. \end{aligned}$$

If $k < K$, then $I_a = \{1, \dots, p-1\}$, and so we are in the first case for all q , implying that $\limsup_{q \rightarrow \infty} \mathfrak{A}_{p^k q}^p = p^k \mathbb{Z}_p^\times$, by Dirichlet's Theorem on primes. By Dirichlet's pigeonhole principle, this means that

$$\mathfrak{R}^p \supseteq \mathfrak{A}^p \supseteq \limsup_{\substack{n \rightarrow \infty \\ n=p^k q, k < K}} \mathfrak{A}_n^p = \bigcup_{k=0}^{K-1} \limsup_{q \rightarrow \infty} \mathfrak{A}_{p^k q}^p = \mathbb{Z}_p \setminus p^K \mathbb{Z}_p. \quad (5.2)$$

We are thus left to consider $\mathfrak{R}^p \cap p^K \mathbb{Z}_p$ and $\mathfrak{A}^p \cap p^K \mathbb{Z}_p$. Since q is prime, and $\psi(p^k q)/(p^k q) \leq p^{-k-1}$, we note, for arbitrary $k \in \mathbb{N}_0$,

$$\mathfrak{R}_{p^k q}^p \subseteq \mathfrak{A}_{p^k q}^p \cup \bigcup_{0 < j \leq k/2} p^{k-2j} \mathbb{Z}_p^\times. \quad (5.3)$$

Since \mathfrak{R}_n^p is finite when $\psi(n) = 0$, and $\mathfrak{R}_{p^k}^p \subseteq p^k \mathbb{Z}_p$ by equations (5.1) and (5.3), we have

$$\begin{aligned} \mu_p(\mathfrak{R}^p \cap p^K \mathbb{Z}_p) &= \mu_p \left(\limsup_{n \rightarrow \infty} \mathfrak{R}_n^p \cap p^K \mathbb{Z}_p \right) \\ &= \mu_p \left(\limsup_{q \rightarrow \infty} \mathfrak{R}_{p^K q}^p \cap p^K \mathbb{Z}_p \right) \\ &= \mu_p \left(\limsup_{q \rightarrow \infty} \mathfrak{A}_{p^K q}^p \right) = \mu_p(\mathfrak{A}^p \cap p^K \mathbb{Z}_p). \end{aligned}$$

Since $x_k = p-1$ for $0 \leq k < K$, equation (5.2) implies

$$\begin{aligned} \mu_p(\mathfrak{R}^p) &= \mu_p(\mathfrak{A}^p) = \mu_p(\mathbb{Z}_p \setminus p^K \mathbb{Z}_p) + \mu_p \left(\limsup_{q \rightarrow \infty} \mathfrak{A}_{p^K q}^p \right) \\ &= \sum_{k=0}^{K-1} x_k p^{-k-1} + \mu_p \left(\limsup_{q \rightarrow \infty} \mathfrak{A}_{p^K q}^p \right). \end{aligned}$$

The theorem hence follows for the current case if we can show that

$$\mu_p \left(\limsup_{q \rightarrow \infty} \mathfrak{A}_{p^K q}^p \right) = \sum_{k=K}^{\infty} x_k p^{-k-1}. \quad (5.4)$$

Applying Dirichlet's Theorem on primes on equation (5.1) for $k = K$, we find

$$\begin{aligned}
\limsup_{q \rightarrow \infty} \mathfrak{A}_{p^{Kq}}^p &= \bigcup_{m \in I_{x_K}} (\pm m^{\pm 1} p^K + p^{K+1} \mathbb{Z}_p) \\
&\cup \bigcup_{i=1}^{\infty} \bigcup_{b'=1}^{b_{K,i}} (\pm (g + b' p^i)^{\pm 1} p^K + p^{K+i+1} \mathbb{Z}_p) \\
&= \bigcup_{m \in I_{x_K}} (\pm m^{\pm 1} p^K + p^{K+1} \mathbb{Z}_p) \\
&\sqcup \bigcup_{i=1}^{\infty} \bigcup_{b'=1}^{b_{K,i}} (\pm (g + b' p^i)^{\pm 1} p^K + p^{K+i+1} \mathbb{Z}_p),
\end{aligned} \tag{5.5}$$

using that $(\pm m_1^{\pm 1} p^K + p^{K+1} \mathbb{Z}_p) \cap (\pm m_2^{\pm 1} p^K + p^{K+1} \mathbb{Z}_p) = \emptyset$ for distinct $m_1, m_2 \in \{g^j : 0 \leq j \leq (p-1)/4\}$. We clearly also have

$$(\pm (g + b'_1 p^i)^{\pm 1} p^K + p^{K+i+1} \mathbb{Z}_p) \cap (\pm (g + b'_2 p^j)^{\pm 1} p^K + p^{K+j+1} \mathbb{Z}_p) = \emptyset,$$

for $i \neq j$ or $b'_1 \neq b'_2$. Applying this to the above calculation, we find

$$\begin{aligned}
\mu_p \left(\limsup_{q \rightarrow \infty} \mathfrak{A}_{p^{Kq}}^p \right) &= \sum_{m \in I_{x_K}} \mu_p(\pm m^{\pm 1} p^K + p^{K+1} \mathbb{Z}_p) \\
&+ \sum_{l=1}^{\infty} \sum_{b'=1}^{b_{K,l}} \mu_p(\pm (g + b' p^l)^{\pm 1} p^K + p^{K+l+1} \mathbb{Z}_p) \\
&= p^{-K} \left(\sum_{m \in I_{x_K}} \mu_p(\pm m^{\pm 1} + p \mathbb{Z}_p) \right. \\
&\quad \left. + \sum_{l=1}^{\infty} \sum_{b'=1}^{b_{K,l}} \mu_p(\pm (g + b' p^l)^{\pm 1} + p^{l+1} \mathbb{Z}_p) \right).
\end{aligned} \tag{5.6}$$

In order to determine the first sum in equation (5.6), note that

$$-g^i \equiv g^{\frac{p-1}{2}+i}, \quad g^{-i} \equiv g^{p-1-i}, \quad -g^{-i} \equiv g^{\frac{p-1}{2}-i}$$

modulo p . For $1 \leq i < (p-1)/4$, we have

$$\begin{aligned}
\frac{p-1}{4} &< \frac{p-1}{2} - i < \frac{p-1}{2} < \frac{p-1}{2} + i < 3\frac{p-1}{4} \\
&< p-1-i < p-1,
\end{aligned}$$

implying that $g^i, g^{-i}, -g, -g^{-i}$ are not congruent modulo p^l for $l \geq 1$. A similar argument yields $1 = 1^{-1} \not\equiv -1$ and $g^{(p-1)/4} \equiv -g^{-(p-1)/4} \not\equiv -g^{(p-1)/4}$

modulo p . Hence, for integers $1 \leq i < (p-1)/4$, $0 \leq b' \leq p-1$, $l \geq 1$, and $j \in \{0, (p-1)/4\}$,

$$\begin{aligned} \mu_p(\pm(g^i + b'p^l)^{\pm 1} + p^{l+1}\mathbb{Z}_p) &= 4/p^{l+1}, \\ \mu_p(\pm g^{\pm j} + p\mathbb{Z}_p) &= 2/p. \end{aligned} \quad (5.7)$$

We are now ready to handle the first sum of (5.6). For $x_K \geq 4$, we find

$$\begin{aligned} \sum_{m \in I_{x_K}} \mu_p(\pm m^{\pm 1} + p\mathbb{Z}_p) &= \sum_{i=2}^{\lfloor x_K/4 \rfloor} \mu_p(\pm g^{\pm(p-1)/4} + p\mathbb{Z}_p) \\ &\quad + \mu_p(\pm 1^{\pm 1} + p\mathbb{Z}_p) + \mu_p(\pm g^{\pm(p-1)/4} + p\mathbb{Z}_p) \\ &= \left(\sum_{i=2}^{\lfloor x_K/4 \rfloor} \frac{4}{p} \right) + \frac{2}{p} + \frac{2}{p} = \frac{4}{p} \lfloor \frac{x_K}{4} \rfloor. \end{aligned}$$

For $x_K < 4$, $I_{x_K} = \emptyset$, and so we reach the same conclusion in that case.

Equation (5.6) and the definitions of $b_{a,i}$, r_a , and x_k then allow us to conclude equation (5.4) as we see

$$\begin{aligned} \mu_p(\limsup_{q \rightarrow \infty} \mathfrak{A}_{p^k q}^p) &= p^{-K} \left(\frac{4}{p} \lfloor \frac{x_K}{4} \rfloor + \sum_{i=1}^{\infty} \sum_{b'=1}^{b_{K,i}} \frac{4}{p^{i+1}} \right) \\ &= p^{-K-1} 4 \left(\lfloor \frac{x_K}{4} \rfloor + \sum_{i=1}^{\infty} b_{K,i} p^{-i} \right) \\ &= p^{-K-1} 4 \left(\lfloor \frac{x_K}{4} \rfloor + r_K \right) \\ &= p^{-K-1} \sum_{l=K}^{\infty} p^{l-K} x_l = \sum_{l=K}^{\infty} x_l p^{l-1}. \end{aligned}$$

Case 2: $5 < p \equiv 3 \pmod{4}$. This case closely follows the structure of case 1. The main difference is a modification to the definitions of I_a and r_k , so that

$$\begin{aligned} I_a &:= \begin{cases} \emptyset & \text{if } a < 2, \\ \{1\} \cup \{g^i : 2 \leq i \leq (a+2)/4\} & \text{if } 2 \leq a < p-1, \\ \{1, \dots, p-1\} & \text{if } a = p-1, \end{cases} \\ r_k &:= \begin{cases} \sum_{l=k}^{\infty} x_l p^{k-l} & \text{if } x_k < 2, \\ \sum_{l=k}^{\infty} x_l p^{k-l} - 4 \lfloor \frac{x_k-2}{4} \rfloor & \text{if } 2 \leq x_k < p-1, \\ 0 & \text{if } x_k = p-1. \end{cases} \end{aligned}$$

Note that all arguments until and including equation (5.7) remain valid, except that we no longer have an integer $j = (p-1)/4$. For $x_K < 2$, the remaining arguments are unchanged, so suppose $x_K \geq 2$. Then

$$\begin{aligned} \sum_{m \in I_{x_K}} \mu_p(\pm m^{\pm 1} + p\mathbb{Z}_p) &= \sum_{i=2}^{\lfloor (x_K+2)/4 \rfloor} \mu_p(\pm g^{\pm(p-1)/4} + p\mathbb{Z}_p) \\ &\quad + \mu_p(\pm 1^{\pm 1} + p\mathbb{Z}_p) \\ &= \left(\sum_{i=2}^{\lfloor (x_K+2)/4 \rfloor} \frac{4}{p} \right) + \frac{2}{p} = \frac{4}{p} \left\lfloor \frac{x_K - 2}{4} \right\rfloor. \end{aligned}$$

Applying this and equation (5.7) to equation (5.4), we conclude

$$\begin{aligned} \mu_p(\limsup_{q \rightarrow \infty} \mathfrak{A}_{p^k q}^p) &= p^{-K} \left(\frac{4}{p} \left\lfloor \frac{x_k - 2}{4} \right\rfloor + \sum_{i=1}^{\infty} \sum_{b'=1}^{b_{K,i}} \frac{4}{p^{i+1}} \right) \\ &= p^{-K-1} 4 \left(\left\lfloor \frac{x_k - 2}{4} \right\rfloor + \sum_{i=1}^{\infty} b_{K,i} p^{-i} \right) \\ &= p^{-K-1} 4 \left(\left\lfloor \frac{x_k - 2}{4} \right\rfloor + r_K \right) \\ &= p^{-K-1} \sum_{l=K}^{\infty} p^{l-K} x_l = \sum_{l=K}^{\infty} x_l p^{l-1}. \end{aligned}$$

Case 3: $p = 2$. In this case, we will use the same construction as in case 1, except that we fix $g = 1$ and change $b_{K,i}$ such that

$$\frac{r_K}{2} = \sum_{i=1}^{\infty} b_{K,i} 2^{-i}.$$

In terms of I_a , the case $a = p - 1$ is more important than the case $a < 4$, so we read the construction as $I_1 = \{1\}$ for $p = 2$. Note that all arguments of the proof for case 1 until equation (5.6) remain valid. Since $(1 + 2^i)^{-1} \equiv 1 + 2^i$

mod 2^{i+1} , $b_{K,i} \in \{0, 1\}$, and $x_K = 0$, we find

$$\begin{aligned} \mu_2 \left(\limsup_{q \rightarrow \infty} \mathfrak{A}_{2^k q}^2 \right) &= 2^{-K} \sum_{i=1}^{\infty} b_{K,i} \mu_2(\pm(1+2^i) + 2^{i+1}\mathbb{Z}_2) \\ &= 2^{-K} \sum_{i=1}^{\infty} b_{K,i} 2 \cdot 2^{-i-1} = 2^{-K} \frac{r_K}{2} \\ &= 2^{-K-1} \sum_{l=K+1}^{\infty} x_l 2^{K-l} = \sum_{l=K}^{\infty} x_l 2^{-l-1}, \end{aligned}$$

which completes the proof in this case.

Case 4: $p = 3, 5$. We use the same construction as in case 2, except that we further change r_k and ψ so that

$$r_k = \begin{cases} x_K - 2 \lfloor \frac{x_K}{2} \rfloor + \sum_{l=K+2}^{\infty} x_l p^{x_K-l} & \text{if } k = K, \\ x_k - 2 \lfloor \frac{x_k}{2} \rfloor & \text{if } k \neq K \end{cases}$$

$$\psi(n) := \begin{cases} f_k(q) & \text{if } n = p^k q, \quad k \leq K+1, \\ 0 & \text{otherwise.} \end{cases}$$

The value of g will also be fixed, depending on p . Note that the arguments of case 2 (which follows the arguments of case 1) remain valid all the way to equation (5.3) and that equation (5.1) now also holds for $k = K+1$. We then find

$$\begin{aligned} \mu_p(\mathfrak{R}^p \cap p^K \mathbb{Z}_p) &= \mu_p \left(\limsup_{q \rightarrow \infty} \mathfrak{A}_{p^K q}^p \right) + \mu_p \left(\limsup_{q \rightarrow \infty} \mathfrak{A}_{p^{K+1} q}^p \right) \\ &= \mu_p(\mathfrak{A}^p \cap p^K \mathbb{Z}_p), \end{aligned}$$

by the same arguments as for the corresponding equation in case 1. Recalling $x_k = p-1$ for $k < K$, equation (5.2) implies

$$\begin{aligned} \mu_p(\mathfrak{R}^p) = \mu_p(\mathfrak{A}^p) &= \sum_{k=0}^{K-1} x_k p^{-k-1} + \mu_p \left(\limsup_{q \rightarrow \infty} \mathfrak{A}_{p^K q}^p \right) \\ &\quad + \mu_p \left(\limsup_{q \rightarrow \infty} \mathfrak{A}_{p^{K+1} q}^p \right), \end{aligned}$$

so that we are left to prove

$$\mu_p \left(\limsup_{q \rightarrow \infty} \mathfrak{A}_{p^K q}^p \right) + \mu_p \left(\limsup_{q \rightarrow \infty} \mathfrak{A}_{p^{K+1} q}^p \right) = \sum_{k=K}^{\infty} x_k p^{-k-1}.$$

Note that equation (5.5) is preserved and that it remains valid for K replaced by $K + 1$. Let $k \in \{K, K + 1\}$. Then

$$\bigcup_{m \in I_{x_k}} \pm m p^k + p^{k+1} \mathbb{Z}_p = p^k \begin{cases} \mathbb{Z}_p^\times & \text{if } x_k = p - 1, \\ (\pm 1 + p \mathbb{Z}_p) & \text{if } 2 \leq x_k < p - 1, \\ \emptyset & \text{otherwise.} \end{cases}$$

From this follows

$$\begin{aligned} \mu_p(\limsup_{q \rightarrow \infty} \mathfrak{A}_{p^k q}^p) &= p^{-k-1} 2 \left\lfloor \frac{x_k}{2} \right\rfloor \\ &\quad + p^{-k} \sum_{i=1}^{\infty} \mu_p \left(\bigcup_{b'=1}^{b_{k,i}} (\pm (g + b' p^i)^{\pm 1} + p^{i+1} \mathbb{Z}_p) \right). \end{aligned}$$

We are now done if we, for each of $p = 3, 5$, can show

$$\sum_{i=1}^{\infty} \mu_p \left(\bigcup_{b'=1}^{b_{K,i}} (\pm (2 + b' p^i)^{\pm 1} + p^{i+1} \mathbb{Z}_p) \right) = r_k, \quad (5.8)$$

as this would imply

$$\begin{aligned} \mu_p(\limsup_{q \rightarrow \infty} \mathfrak{A}_{p^k q}^p) &= p^{-k-1} 2 \left\lfloor \frac{x_k}{2} \right\rfloor + p^{-k} r_k \\ &= \begin{cases} x_k p^{-K} + \sum_{l=K+2}^{\infty} x_l p^{-l} & \text{if } k = K, \\ x_{K+1} p^{-(K+1)} & \text{if } k = K + 1. \end{cases} \end{aligned}$$

For $p = 3$, fix $g = 2$. Then $\liminf_{l \rightarrow \infty} x_l < 2$, and we have

$$\frac{r_k}{4} \leq \frac{1}{4} \left(1 + \sum_{l=k+2}^{\infty} x_l 2^{k-l} \right) < \frac{1}{4} \left(1 + \frac{1}{3} \right) = \frac{1}{3},$$

i.e., $b_{1,k} = 0$. For $i \geq 2$ and $1 \leq b' \leq b_{i,k}$, notice that

$$\begin{aligned} (2 + b' 3^i) + 3^{i+1} \mathbb{Z}_3 &\subseteq 2 + 3^2 \mathbb{Z}_3, \\ -(2 + b' 3^i) + 3^{i+1} \mathbb{Z}_3 &\subseteq 7 + 3^2 \mathbb{Z}_3, \\ (2 + b' 3^i)^{-1} + 3^{i+1} \mathbb{Z}_3 &\subseteq 5 + 3^2 \mathbb{Z}_3, \\ -(2 + b' 3^i)^{-1} + 3^{i+1} \mathbb{Z}_3 &\subseteq 4 + 3^2 \mathbb{Z}_3. \end{aligned}$$

Since all balls on the right-hand-side of the above four inclusions are disjoint, the same holds for the four balls defining $(\pm 2^{\pm 1} + b' 3^i + 3^{i+1} \mathbb{Z}_3)$. When we then vary i , we note that the sets

$$(\pm 2 + b' 3^i)^{\pm 1} + 3^{i+1} \mathbb{Z}_3 \subseteq \pm 2^{\pm 1} + 3^i \mathbb{Z}_3^\times$$

are also disjoint. We conclude equation (5.8) by calculating

$$\begin{aligned} \sum_{i=1}^{\infty} \mu_3 \left(\bigcup_{b'=1}^{b_{k,i}} (\pm(2+b'3^i)^{\pm 1} + 3^{i+1}\mathbb{Z}_3) \right) &= \sum_{i=1}^{\infty} \sum_{b'=1}^{b_{k,i}} 4\mu_3(3^{i+1}\mathbb{Z}_3) \\ &= 4 \sum_{i=1}^{\infty} b_{k,i} 3^{-i-1} = r_k. \end{aligned}$$

For $p = 5$, fix $g = 3$ and estimate

$$\frac{r_k}{4} \leq \frac{1}{4} \left(1 + \sum_{l=K+2}^{\infty} x_l 2^{x_K-l} \right) < \frac{1}{4} \left(1 + \frac{1}{5} \right) < \frac{2}{5},$$

so that $b_{k,1} \in \{0, 1\}$. For $i \geq 1$ and $1 \leq b' \leq b_{k,i}$, let $a_i \in \{0, 1\}$ such that $a_1 = b'$ and $a_i = 0$ for $i > 1$. We then find

$$\begin{aligned} (3 + b'5^i) + 5^{i+1}\mathbb{Z}_5 &\subseteq 3 + a_i 5 + 5^2\mathbb{Z}_5, \\ -(3 + b'5^i) + 5^{i+1}\mathbb{Z}_5 &\subseteq 2 + (4 - a_i)5 + 5^2\mathbb{Z}_5, \\ (3 + b'5^i)^{-1} + 5^{i+1}\mathbb{Z}_5 &\subseteq 2 + (3 + a_i)5 + 5^2\mathbb{Z}_5, \\ -(3 + b'5^i)^{-1} + 5^{i+1}\mathbb{Z}_5 &\subseteq 3 + (1 - a_i)5 + 5^2\mathbb{Z}_5. \end{aligned}$$

We are again left with four disjoint balls on the right-hand-side, for i fixed. We then apply arguments in parallel to those for $p = 3$ and conclude equation (5.8). This completes the proof.

In cases 3 and 4 of the above proof, note that the choice of generator g matters; for $p = 2$, $g > 1$ would lead to $\pm g^{\pm 1}$ representing four unique values modulo 2^i for sufficiently large i , where we want exactly 2 unique values. For $p = 3$, $g \equiv -1 \pmod{9}$ would on the other hand lead to $\pm g^{\pm 1}$ representing only two unique values modulo 9, where we want exactly 4. The same issue arises modulo 25 for $p = 5$ when $g + a_1 5 = g + 5 \equiv \pm 7 \pmod{25}$.

Note also that the alterations introduced in case 4 would not be sufficient for $p = 5$ if they were to be applied to case 1, even though $5 \equiv 1 \pmod{4}$, since it would allow any $b_{k,1}$ between 0 and 4, which would lead to $g + a_1 5 \equiv \pm 7 \pmod{25}$ for some a_1 , regardless of g .

Finally, note that the construction in case 4 would also work for $p = 2$ by putting $g = 3$ and $r_k = 0$ for $k \neq K$, though that would not actually shorten the proof as we would then have to give $p = 2$ the same amount of special attention as we gave each of $p = 3$ and $p = 5$. This suggests that we are in a peculiar case of $p = 2$ *not* being the most troublesome prime, as that title appears to go to $p = 5$, with $p = 3$ as a close second.

6. Concluding Remarks

Considering how Proposition 4.2 acts as a shell-wise \mathcal{C}^p -variant of the 0-1 law on \mathcal{A}^p from [5], it appears rather plausible that the p -adic Duffin–Schaeffer theorem should also have a shell-wise \mathcal{C}^p -variant, as formally stated below.

CONJECTURE. *Let $\psi : \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$, and let p be a prime. Suppose $\text{supp } \psi \subseteq p^k \mathbb{N} \setminus p^{k+1} \mathbb{N}$ for some $k \in \mathbb{N}_0$. Then*

$$\mu_p(\mathcal{C}^p \cap p^k \mathbb{Z}_p^\times) = \begin{cases} (p-1)/p^{k+1} & \text{if } \sum_{n=1}^{\infty} \mu_p(\mathcal{C}_n^p) = \infty, \\ 0 & \text{if } \sum_{n=1}^{\infty} \mu_p(\mathcal{C}_n^p) < \infty. \end{cases}$$

As with the original and p -adic Duffin–Schaeffer Theorems, the Borel–Cantelli Lemma directly implies $\mu_p(\mathcal{C}^p) = 0$ when the series converges. If the conjecture holds true, it combines with Proposition 4.1 to provide an explicit formula for determining the p -adic measures of \mathcal{B}^p and \mathcal{C}^p . It is expected that the conjecture will follow from a modification of the proof of [5, Theorem 2] combined with [8, Proposition 5.4], following the structure presented in [9]. In the light of [9, Theorem 2], it appears only natural if the measures $\mu_p(\mathcal{C}_n^p)$ in the divergence criterion may be replaced by the fractions $\phi(n)\psi(n)/n$ from the real Duffin–Schaeffer Theorem, by following a similar argument.

As for the set \mathfrak{A}^p , one might try to modify the construction with the aim of decreasing the spectrum of possible measure values, similarly to what Haynes [5] achieved in constructing \mathcal{A}^p instead of \mathcal{B}^p . In his construction, Haynes effectively removed the sets \mathcal{B}_n^p that were restricted to specific shells $p^k \mathbb{Z}_p^\times$ from consideration as he indirectly forced the sets \mathcal{A}_n^p with $p \mid n$ to be either empty or full [5, Lemma 2]. Trying to get a similar modification of \mathfrak{A}^p , we might consider the set

$$\limsup_{n \rightarrow \infty} \bigcup_{\substack{d|n \\ \gcd(d, pn/d)=1}} B_{\mathbb{Q}_p} \left(\frac{d}{n/d}, \frac{\psi(n)}{n} \right).$$

However, for $p \neq 3, 5$ and $x \in [0, (p-1)/p] \cup \{1\}$, the ψ constructed in the proof of Theorem 2.2 will still produce measure x . For $p = 3, 5$, ψ only works for $x \in \{1, (p-1)/p\} \cup \bigcup_{x_0=0}^{p-2} \frac{x_0}{p} + [0, p^{-2})$, but it seems reasonable that there should also exist a ψ for the remaining $x \in [0, (p-1)/p] \cup \{1\}$; perhaps a hybrid between the constructions from cases 3 and 4 would do the trick. Note that this attack would work identically if the above modification were carried out on \mathfrak{R}^p instead. As such, there does not seem to be any immediate ‘correction’ to the set \mathfrak{A}^p that would make it satisfy a 0-1 law in general.

Acknowledgements. I thank my supervisor Simon Kristensen for aiding me in writing this paper, and the Independent Research Fund Denmark for funding my research (Grant ref. 1026-00081B).

REFERENCES

1. Dzmitry Badziahin and Yann Bugeaud. “Multiplicative p-adic Approximation”. *Michigan Mathematical Journal* 71.1 (2022), pp. 121–143. DOI: 10.1307/mmj/20195785. URL: <https://doi.org/10.1307/mmj/20195785>.
2. J. W. S. Cassels. “Some metrical theorems in Diophantine approximation. I”. *Mathematical Proceedings of the Cambridge Philosophical Society* 46.2 (1950), pp. 209–218. DOI: 10.1017/S0305004100025676.
3. R. J. Duffin and A. C. Schaeffer. “Khintchine’s problem in metric Diophantine approximation”. *Duke Mathematical Journal* 8.2 (1941), pp. 243–255. DOI: 10.1215/S0012-7094-41-00818-9. URL: <https://doi.org/10.1215/S0012-7094-41-00818-9>.
4. P. Gallagher. “Approximation by reduced fractions”. *Journal of The Mathematical Society of Japan* 13.4 (1961), pp. 342–345. ISSN: 0025-5645.
5. A. K. Haynes. “The Metric Theory of p-Adic Approximation”. *International mathematics research notices IMRN*. 1 (2009), pp. 18–52. ISSN: 1073-7928.
6. V. Jarník. “Sur les approximations diophantiques des nombres p-adiques”. *Revista de Ciencias* 47 (1945), pp. 489–505. ISSN: 0034-7760.
7. A. Khintchine. “Einige Sätze über Kettenbrüche, mit Anwendungen auf die Theorie der Diophantischen Approximationen”. *Mathematische Annalen*. 92.1-2 (1924), pp. 115–125. ISSN: 0025-5831.
8. D. Koukoulopoulos and J. Maynard. “On the Duffin-Schaeffer conjecture”. *Annals of mathematics*. 192.1 (2020), pp. 251–307. ISSN: 0003-486X.
9. Simon Kristensen and Mathias Løkkegaard Laursen. *The p-adic Duffin-Schaeffer conjecture*. 2021. DOI: 10.48550/ARXIV.2110.02598. URL: <https://arxiv.org/abs/2110.02598>.
10. É. Lutz. *Sur les approximations diophantiennes linéaires p-adiques*. fre. Actualités Scientifiques et Industrielles ; 1224. Paris: Hermann, 1955.
11. H. L. Montgomery and R. C. Vaughan. *Multiplicative Number Theory I. Classic Theory*. Cambridge University Press, 2007. Chap. 2.2. ISBN: 978-1-107-40582-0.

DEPARTMENT OF MATHEMATICS
 AARHUS UNIVERSITY
 8000 AARHUS C,
 DENMARK
 E-mail: mll@math.au.dk