

# DE RHAM - WITT KZ EQUATIONS

*Crystalline bozonization*

Vadim Schechtman and Alexander Varchenko

December 5, 2022

*Те, кому приходилось столкнуться с векторами Витта  
в собственной работе, часто относятся к ним  
с теплотой*

*Д. Каледин<sup>1</sup>*

## Introduction

In this note we propose a de Rham - Witt version of the derived KZ equations, and of their hypergeometric realizations, cf. [SV2]. In particular we describe a de Rham - Witt version of Theorems 7.2.5', 7.2.5'' from [SV1].

We also propose de Rham - Witt versions of some classical theorems related to arbitrary hyperplane arrangements.

### *Contents of the paper*

In §1 we recall some background material about the de Rham - Witt complex. In §2 we present a de Rham - Witt version of the first part of [SV1] concerning arbitrary hyperplane arrangements over  $\mathbb{F}_p$ . In 2.3.3 - 2.3.7 we discuss also a motivic version as suggested by A.Beilinson. In §3 we present a de Rham - Witt version of [SV2] which in its turn is a development of [SV1], Part 2. For simplicity of exposition we discuss only the case  $\mathfrak{g} = \mathfrak{sl}_2$  but the generalization to arbitrary Kac - Moody algebras is straightforward.

---

<sup>1</sup>See [K]

## Acknowledgements

We are grateful to Alexander Beilinson for an interesting discussion and consultations; the idea of 2.3.3 - 2.3.7 belongs to him, and the proofs are obtained jointly with him. We express our gratitude to H el ene Esnault for important corrections.

Our reflections on the topics of this paper have been triggered by a talk of Rub en Mu noz-Bertrand; we are thankful to him.

##  1. The de Rham - Witt complex

### 1.1. De Rham - Witt complex of a polynomial ring. Cf. [B],  2.

$p$  - a prime number.  $W(\mathbb{F}_p) = \mathbb{Z}_p$ ,  $F = \text{Id}$ ,  $V = p\text{Id}$ . Thus

$$W_n(\mathbb{F}_p) := W(\mathbb{F}_p)/V^n W(\mathbb{F}_p) = \mathbb{Z}/p^n \mathbb{Z}.$$

(a) *Witt ring*

Let  $A := \mathbb{F}_p[t_1, \dots, t_n]$ .

We consider, together with S.Lubkin et al., cf. [L, I1, I2, B], a (nonnoetherian) algebra

$$C = \cup_{i \geq 0} \mathbb{Q}_p[t_1^{p^{-i}}, \dots, t_n^{p^{-i}}]$$

**1.1.1. Remark.**  $C$  is filtered by subalgebras

$$C_m = \cup_{0 \leq i \leq m} \mathbb{Q}_p[t_1^{p^{-i}}, \dots, t_n^{p^{-i}}],$$

and each

$$C_m = \mathbb{Q}_p[t_1^{p^{-m}}, \dots, t_n^{p^{-m}}].$$

We may also consider a bigger ring

$$\hat{C} = \left\{ \sum_{(i_1, \dots, i_n) \in \mathbb{N}[p^{-1}]} a_{i_1, \dots, i_n} t_1^{i_1} \dots t_n^{i_n} \right\}$$

(formal power series).

Both  $C$  and  $\hat{C}$  are equipped with a  $\mathbb{Q}_p$ -linear automorphism  $F$ ,

$$F(t_i) = t_i^p,$$

and we define  $V := pF^{-1}$ .

*Notations.* For  $i \in \mathbb{Z}[p^{-1}] \subset \mathbb{Q} \subset \mathbb{Q}_p$ ,

$$d(i) := -v_p(i);$$

for  $I = (i_1, \dots, i_n) \in \mathbb{Z}[p^{-1}]^n$ ,

$$\begin{aligned} |I| &= \sum i_k, \\ d(I) &= \max\{d(i_k)\}. \end{aligned}$$

The Witt ring  $W(\mathbb{F}_p[t_1, \dots, t_n])$  is a subring of  $\hat{C}$ , namely

$$W(\mathbb{F}_p[t_1, \dots, t_n]) = \left\{ \sum_{(i_1, \dots, i_n) \in \mathbb{N}[p^{-1}]^n} a_{i_1, \dots, i_n} t_1^{i_1} \dots t_n^{i_n} \mid \forall I \ v_p(a_I) \geq d(I), \lim_{|I| \rightarrow \infty} a_I = 0 \right\},$$

cf. [L], VI (a). It is stable with respect to  $F$  and  $V$ .

(b) *De Rham complex*  $\Omega_{C/\mathbb{Q}_p}^\bullet$

Consider the de Rham complex  $\Omega_{C/\mathbb{Q}_p}^\bullet$ .

Each  $m$ -form  $\omega \in \Omega_{C/\mathbb{Q}_p}^m$  may be uniquely written as

$$\omega = \sum_{1 \leq i_1 < \dots < i_m \leq n} a_{i_1, \dots, i_m}(t) d \log t_{i_1} \wedge \dots \wedge d \log t_{i_m}$$

where  $a_I(t) \in C$ .

Define automorphisms  $F : \Omega_{C/\mathbb{Q}_p}^m \xrightarrow{\sim} \Omega_{C/\mathbb{Q}_p}^m$  by

$$\begin{aligned} F\left(\sum_I a_{i_1, \dots, i_m}(t_1, \dots, t_n) d \log t_{i_1} \wedge \dots \wedge d \log t_{i_m}\right) = \\ \sum_I a_{i_1, \dots, i_m}(t_1^p, \dots, t_n^p) d \log t_{i_1} \wedge \dots \wedge d \log t_{i_m}. \end{aligned}$$

*Compatibility with d:*

We have

$$dF = pFd$$

As before, we put  $V := pF^{-1}$ .

Thus we have two morphisms  $F, W : \Omega_{C/\mathbb{Q}_p}^\bullet \rightarrow \Omega_{C/\mathbb{Q}_p}^\bullet$  which satisfy:

$$FV = VF = p, FdV = d, dF = pFd, Vd = pdV,$$

$$V(x \cdot Fy) = Vx \cdot y$$

(c) *De Rham - Witt complex*

Let us say that  $\omega \in \Omega_{C/\mathbb{Q}_p}^m$  is *integral* if all coefficients of all  $a_I(t)$  belong to  $\mathbb{Z}_p$ . Define subspaces

$$E^m = \{\omega \in \Omega_{C/\mathbb{Q}_p}^m \mid \omega, d\omega \text{ are integral}\},$$

for example

$$E^0 = \cup_{i \geq 0} p^i \mathbb{Z}_p[t_1^{p^{-i}}, \dots, t_n^{p^{-i}}].$$

They form a subcomplex  $E^\bullet \subset \Omega_{C/\mathbb{Q}_p}^\bullet$  stable with respect to  $F$  and  $V$ .

Define

$$\text{Fil}^a E^m = V^a E^m + d(V^a E^{m-1}) \subset E^m;$$

these subspaces form a subcomplex  $\text{Fil}^a E^\bullet \subset E^\bullet$ , and we may consider the quotient complex

$$W_a \Omega_A^\bullet = E^\bullet / \text{Fil}^a E^\bullet.$$

When  $a \geq 1$  varies, these complexes form a projective system.

$$W \Omega_A^\bullet = \varprojlim W_a \Omega_A^\bullet.$$

**1.2. De Rham - Witt complex of Laurent polynomials.** The construction is similar, cf. [I2] II.2.

Consider the case of one variable.

$$A = \mathbb{F}_p[t, t^{-1}],$$

$$C = \cup_{r \geq 0} \mathbb{Q}_p[t^{1/p^r}, t^{-1/p^r}]$$

$$F : C \xrightarrow{\sim} C, F(a(t, t^{-1})) = a(t^p, t^{-p}), V = pF^{-1}.$$

The de Rham complex  $\Omega_{C/\mathbb{Q}_p}^\bullet$  has length 1:

$$0 \longrightarrow C = \Omega_{C/\mathbb{Q}_p}^0 \longrightarrow \Omega_{C/\mathbb{Q}_p}^1 \longrightarrow 0.$$

Each form  $\omega \in \Omega_{C/\mathbb{Q}_p}^1$  can be uniquely written as

$$\omega = \sum_{i \in \mathbb{N}[p-1]} a_i(t, t^{-1}) d \log t, \quad a_i \in C,$$

$$F : \Omega_{C/\mathbb{Q}_p}^1 \xrightarrow{\sim} \Omega_{C/\mathbb{Q}_p}^1,$$

$$F\left(\sum_{i \in \mathbb{N}[p-1]} a_i(t, t^{-1}) d \log t\right) = \sum_{i \in \mathbb{N}[p-1]} a_i(t^p, t^{-p}) d \log t,$$

$$V = pF^{-1}.$$

Let us say that  $\omega \in \Omega_{C/\mathbb{Q}_p}^m$  is *integral* if all coefficients of all  $a_I(t)$  belong to  $\mathbb{Z}_p$ . Define subspaces

$$E^m = \{\omega \in \Omega_{C/\mathbb{Q}_p}^m \mid \omega, d\omega \text{ are integral}\},$$

for example

$$E^0 = \cup_{i \geq 0} p^i \mathbb{Z}_p[t^{p^{-i}}, t^{-p^{-i}}].$$

They form a subcomplex  $E^\bullet \subset \Omega_{C/\mathbb{Q}_p}^\bullet$  stable with respect to  $F$  and  $V$ .

Define

$$\text{Fil}^a E^m = V^a E^m + d(V^a E^{m-1}) \subset E^m;$$

these subspaces form a subcomplex  $\text{Fil}^a E^\bullet \subset E^\bullet$ , and we may consider the quotient complex

$$W_a \Omega_A^\bullet = E^\bullet / \text{Fil}^a E^\bullet.$$

When  $a \geq 1$  varies, these complexes form a projective system.

$$W \Omega_A^\bullet = \varprojlim W_a \Omega_A^\bullet.$$

**1.3. Shift.** Given  $z \in \mathbb{F}_p$ , let

$$A = \mathbb{F}_p[t, (t-z)^{-1}] = \mathbb{F}_p[t-z, (t-z)^{-1}].$$

We introduce an algebra

$$C = \cup_{r \geq 0} \mathbb{Q}_p[(t-z)^{1/p^r}, (t-z)^{-1/p^r}],$$

an isomorphism  $F : C \xrightarrow{\sim} C$ ,

$$F(a(t-z, (t-z)^{-1})) = a((t-z)^p, (t-z)^{-p}),$$

and an endomorphism  $V : C \rightarrow C$ ,

$$V = pF^{-1}.$$

Then we proceed as before: 1-forms will have the form

$$\omega = \sum_{i \in \mathbb{N}[p^{-1}]} a_i(t-z, (t-z)^{-1}) d \log(t-z),$$

etc.

**1.4. Laurent polynomials of several variables.** Let  $I \subset [n] = \{1, \dots, n\}$ , and

$$A = \mathbb{F}_p[t_1, \dots, t_n, t_i^{-1}, i \in I]$$

We proceed as before, starting with an algebra

$$C = \cup_{r \geq 0} \mathbb{Q}_p[t_1^{1/p^r}, \dots, t_n^{1/p^r}, t_i^{\pm 1/p^r}, i \in I],$$

etc.

The multiplicative group  $A^*$  is a direct sum of  $\mathbb{F}_p^*$  and of a free abelian group generated by  $t_i$ ,  $i \in I$ . The Teichmüller homomorphism

$$T : A^* \rightarrow W(A)^* \subset \hat{C}^*$$

sends  $t_i$  to  $t_i \in \hat{C}^*$ .

**1.5.** More generally, we will be interested in the de Rham - Witt complex of the ring

$$A = \mathbb{F}_p[t, z_1, \dots, z_n, (t-z_1)^{-1}, \dots, (t-z_n)^{-1}].$$

Similarly to the previous considerations, we introduce a ring

$$C = \cup_{r \geq 0} \mathbb{Q}_p[t^{1/p^r}, (t-z_1)^{\pm 1/p^r}, \dots, (t-z_n)^{\pm 1/p^r}]$$

and its completion  $\hat{C}$ .

The complex  $W\Omega_A^\bullet$  will be a subcomplex of  $\Omega_{\hat{C}/\mathbb{Q}_p}^\bullet$ .

## §2. Du côté de chez Orlik - Solomon

### 2.1. Orlik - Solomon subalgebra, aka fixed points of Frobenius.

**Notation.** Given a whatever object  $?$  acted upon by a Frobenius  $F$ , the superscript  $?^{F^{-1}}$  will denote  $\{x \in ? \mid Fx = x\}$ .

**2.1.1.** Let  $A = \mathbb{F}_p[t, t^{-1}]$ . Consider the fixed submodule of Frobenius

$$W\Omega_A^{\bullet F^{-1}} \hookrightarrow W\Omega_A^{\bullet}. \quad (2.1.1)$$

We have

$$W\Omega_A^{0F^{-1}} = \mathbb{Z}_p, \quad W\Omega_A^{1F^{-1}} = \mathbb{Z}_p d \log t,$$

the differential in  $\Omega_A^{\bullet F^{-1}}$  being zero.

The embedding (2.1.1) is a quasiisomorphism.

**2.1.2.** A more general case of 1.4 is similar: for

$$A = \mathbb{F}_p[t_1, \dots, t_n, t_i^{-1}, i \in I]$$

the subcomplex  $W\Omega_A^{\bullet F^{-1}}$  is a free exterior  $\mathbb{Z}_p$ -algebra with generators  $d \log t_i, i \in I$  and zero differential,

$$W\Omega_A^{\bullet F^{-1}} = \Lambda_{\mathbb{Z}_p}^{\bullet}(\oplus_{i \in I} \mathbb{Z}_p d \log t_i).$$

It will be called the *Orlik - Solomon subalgebra*, and denoted also  $W\Omega_A^{\bullet OS}$ .

The embedding

$$W\Omega_A^{\bullet OS} \hookrightarrow W\Omega_A^{\bullet}. \quad (2.1.2)$$

is a quasiisomorphism.

**2.2.** This is a general phenomenon, valid for any hyperplane arrangement. Namely, consider an affine space

$$\mathbb{A}_{\mathbb{F}_p}^n = \text{Spec } \mathbb{F}_p[t_1, \dots, t_n]$$

and a finite collection  $\mathcal{H} = \{H_i\}_{i \in I}$  of affine hyperplanes in it,  $H_i$  being given by an equation  $f_i(t) = 0$ .

Let

$$U = \mathbb{A}_{\mathbb{F}_p}^n \setminus \bigcup_{i \in I} H_i = \text{Spec } A$$

where

$$A = \mathbb{F}_p[t_1, \dots, t_n, f_i^{-1}, i \in I].$$

We will use both notations  $W\Omega_A^\bullet = W\Omega^\bullet(U)$ .

**2.2.1. Fixed points of Frobenius are closed.** If  $x \in W\Omega_A^i$ ,  $x = Fx$  then

$$dx = dFx = pFdx,$$

i.e.

$$(1 - pF)dx = 0.$$

But the operator  $1 - pF$  is invertible:

$$(1 - pF)^{-1} = \sum_{i \geq 0} p^i F^i,$$

and the series in the RHS converges. Whence  $dx = 0$ .

We've got an embedding of complexes

$$os : W\Omega_A^{\bullet OS} := W\Omega_A^{\bullet F-1} \hookrightarrow W\Omega_A^\bullet. \quad (2.2.1)$$

where the first one is equipped with the zero differential.

**2.2.2. Logarithmic forms.** Let  $H \subset \mathbb{A} = \mathbb{A}_{\mathbb{F}_p}^n$  be a hyperplane given by an equation  $f(t) = 0$ . Let us choose new coordinates  $u_1, \dots, u_n$  in  $\mathbb{A}$  such that in new coordinates  $H$  will be given by the equation  $u_n = 0$ . Thus we get an isomorphism

$$\phi : \mathbb{A} \setminus H \xrightarrow{\sim} \text{Spec } \mathbb{F}_p[u_1, \dots, u_{n-1}, u_n^{\pm 1}]$$

We may apply 1.4 to the target and get closed 1-forms

$$d \log u_n \in W\Omega^1(\mathbb{F}_p[u_1, \dots, u_{n-1}, u_n^{\pm 1}]),$$

whence

$$d \log f := \phi^* d \log u_n \in W\Omega^1(\mathbb{A} \setminus H).$$

Otherwise one can define  $d \log f$  as in [I2] 3.23, as

$$d \log f = d\tilde{f}/\tilde{f}$$

where  $\tilde{f} \in W\mathcal{O}(\mathbb{A} \setminus H)$  is the Teichmüller representative of  $f$ .

**2.2.3.** Returning to our arrangement, we obtain closed 1-forms  $d \log f_i \in W\Omega_A^1$ ; by construction they belong to  $W\Omega_A^{1F-1}$ .



Compare with [I2] CH. I, (3.23).

According to *op. cit.* I, Thm. 5.7.2

$$W\Omega_U^{\bullet F^{-1}} \subset W\Omega_U^{\bullet}$$

is the same as the subsheaf generated by logarithmic forms.

**2.3. Orlik - Solomon.** Recall that the Orlik - Solomon algebra  $\mathcal{A}(\mathcal{H})$ , cf. [OS], §2, is an  $\mathbb{N}$ -graded skew commutative ring with  $\mathcal{A}(\mathcal{H})_0 = \mathbb{Z}$ , the quotient of the exterior algebra with generators  $(H_i), i \in I$ , of degree 1, modulo the ideal generated by the elements

$$(H_1, \dots, H_p) := H_1 \wedge \dots \wedge H_p$$

if  $H_i$  are not in general position, and

$$\sum_{i=1}^{p+1} (-1)^i (H_1, \dots, \hat{H}_i, \dots, H_{p+1}) = 0$$

if  $H_1, \dots, H_{p+1}$  are not in general position and  $H_1 \cap \dots \cap H_{p+1} \neq \emptyset$ , cf. [SV1], 1.2.

Multiplication is given by concatenation.

Recall that all  $\mathcal{A}(\mathcal{H})_p$  are free abelian groups.

**2.3.1. Proposition.** *There exists a unique algebra map*

$$\psi : \mathcal{A}(\mathcal{H}) \longrightarrow W\Omega_A^{\bullet OS} = W\Omega^{\bullet OS}(U)$$

such that  $\psi(H_i) = d \log f_i$ .  $\square$

*Integer structure*

Let

$$W\Omega^{\bullet OS}(U)_{\mathbb{Z}} = \text{Im}(\psi) \subset W\Omega^{\bullet OS}(U)$$

denote the  $\mathbb{Z}$ -subalgebra generated by  $d \log f_i$ .

**2.3.2. Proposition.** *The map  $\psi$  induces an isomorphism*

$$\psi : \mathcal{A}(\mathcal{H}) \xrightarrow{\sim} W\Omega^{\bullet OS}(U)_{\mathbb{Z}},$$

whence an isomorphism

$$\psi_{\mathbb{Z}_p} : \mathcal{A}(\mathcal{H}) \otimes \mathbb{Z}_p \xrightarrow{\sim} W\Omega^{\bullet OS}(U).$$

**Proof** the same as in [OS], Thm. 5.2.  $\square$

*Motivic variation: Milnor K-theory and Orlik - Solomon*

**2.3.3.** As was suggested by Beilinson, we may consider a multiplicative version of the above.

At the beginning we may replace  $\mathbb{F}_p$  by an arbitrary base field  $k$ .

Let

$$k(A) = k(U) = k(\mathbb{A}_k^n) = k(t_1, \dots, t_n)$$

be the field of rational functions on  $\mathbb{A}_k^n$ . Let  $K_{\bullet}^M(k(U))$  be the algebra of Milnor  $K$ -functors. Thus

$$K_0^M(k(U)) = \mathbb{Z}, \quad K_1^M(k(U)) = k(t_1, \dots, t_n)^*,$$

and  $K_{\bullet}^M(k(U))$  is a an  $\mathbb{N}$ -graded skew commutative algebra generated by  $K_1^M(k(U))$  subject to the Steinberg relation

$$\{x, 1 - x\} = 0, \quad x, 1 - x \in K_1^M(k(U)), \quad (2.3.1)$$

cf. [M].

Let

$$\tilde{K}_1^M(U) \subset K_1^M(k(U))$$

be the subgroup generated by  $f_i$ ; it is a free abelian group with a base  $\{f_i, i \in I\}$ . Let

$$\tilde{K}_{\bullet}^M(U) \subset K_{\bullet}^M(k(U))$$

be the  $\mathbb{Z}$ -subalgebra generated by  $\tilde{K}_1^M(U)$ .

We have a map

$$\chi_{\mathbb{Z}1} : \tilde{K}_1^M(U) \longrightarrow \mathcal{A}_1(\mathcal{H})$$

sending  $f_i$  to  $(H_i)$ .

**2.3.3.1. Claim.** *The map  $\chi_{\mathbb{Z}1}$  extends to an algebra map*

$$\chi_{\mathbb{Z}} : \tilde{K}_{\bullet}^M(U) \longrightarrow \mathcal{A}(\mathcal{H}) \quad (2.3.2)$$

**2.3.3.2. Example. Identity "chère à Gelfand": magic of the Steinberg relation.** Consider an arrangement  $\mathcal{H}$  in  $\mathbb{A}_k^2 = \text{Spec } k[x, y]$  consisting of 3 lines

$$\ell_1 : x = 0, \ell_2 : y = 0, \ell_3 : x = y,$$

so

$$U = \mathbb{A}_k^2 \setminus \bigcup_{i=1}^3 \ell_i.$$

We have a familiar relation

$$\frac{1}{x(x-y)} - \frac{1}{y(x-y)} = -\frac{1}{xy},$$

and its counterpart in  $\mathcal{A}(\mathcal{H})$ . Let us write down its counterpart in  $K_2^M(U)$ .

We have the following equalities in  $K_2^M(k(U)) \otimes \mathbb{Z}[1/2]$ :

$$\begin{aligned} \{x, x-y\} &= \{x, x\} + \{x, 1-x^{-1}y\} = \{x, 1-x^{-1}y\}; \\ \{y, x-y\} &= \{y, xy^{-1}-1\} = \{y, xy^{-1}\} + \{y, 1-x^{-1}y\} = \\ &= \{y, x\} + \{y, y^{-1}\} + \{y, 1-x^{-1}y\} = \{y, x\} + \{y, 1-x^{-1}y\}. \end{aligned}$$

Here we have used the equality  $\{y, y\} = 0$ ; by that reason we had to tensorise by  $\mathbb{Z}[1/2]$ .

On the other hand

$$\{x, 1-x^{-1}y\} = \{x, 1-x^{-1}y\} + \{x^{-1}y, 1-x^{-1}y\} = \{xx^{-1}y, 1-x^{-1}y\} = \{y, 1-x^{-1}y\},$$

whence

$$\{x, x-y\} - \{y, x-y\} = -\{y, x\} = \{x, y\}.$$

This is the Gelfand identity in  $K_2^M$ .  $\square$

**2.3.4. Conjecture (A.Beilinson) .** *The map  $\chi_{\mathbb{Z}}$  (2.3.2) is an isomorphism.*

The map  $\chi_{\mathbb{Z}}$  composed with  $\psi$  (see Prop. 2.3.2) gives rise to a map

$$d \log : \tilde{K}_{\bullet}^M(U) \longrightarrow W\Omega^{\bullet OS}(U)_{\mathbb{Z}} \quad (2.3.4)$$

sending  $f_i$  to  $d \log f_i$ .

**2.3.5. Motivic Orlik - Solomon.** *A canonical map*

$$H_{Mot}^\bullet(U; \mathbb{Q})^{\text{diag}} \longrightarrow H_{Mot}^\bullet(U; \mathbb{Q})$$

*is an isomorphism.*

Here  $H_{Mot}^\bullet(U; \mathbb{Q})$  denotes the motivic cohomology realized as appropriate eigenspaces of Adams operations in the Quillen  $K$ -theory, see [Be], 2.2, and  $H_{Mot}^\bullet(U; \mathbb{Q})^{\text{diag}}$  is its part consisting of all elements of type  $(i, i)$ .

**Proof.** Our assertion says that all elements of  $H_{Mot}^\bullet(U; \mathbb{Q})$  are of type  $(i, i)$ . This may be proven by induction on  $n$ .  $\square$

**2.3.6.** Instead of rational motivic cohomology we could use more refined version, the Voevodsky's motivic cohomology, [MVW].

It is also probable that a reduced version from [ES] works for  $U$ .

Let us return to the de Rham - Witt realization.

**2.4. Theorem.** *The embedding os (2.2.1) is a quasiisomorphism.*

**Proof** the same as in [OS], Thm. 5.2 which uses [Br] Lemme 5. Brieskorn uses the Lefschetz theorems on hyperplane sections which hold true for the crystalline cohomology.  $\square$

**2.5. Turning on a connection, "Aomoto conjecture".** Given a collection of exponents  $a_i \in \mathbb{Z}_p$ ,  $i \in I$ , we get a closed one-form

$$\omega = \sum_i a_i d \log f_i \in W\Omega_A^{1OS},$$

and a connection

$$\nabla = d + \omega \wedge \bullet : W\Omega_A^i \longrightarrow W\Omega_A^{i+1}.$$

which restricted to  $W\Omega_A^{\bullet OS}$  is simply the multiplication by  $\omega$ .

**2.6. Theorem** (Aomoto conjecture, a weak form). *For sufficiently small  $a_i$  (in the  $p$ -adic metric) the embedding*

$$(W\Omega_A^{\bullet OS}, \omega) \hookrightarrow (W\Omega_A^\bullet, d + \omega)$$

is a quasiisomorphism.

**Proof.** The same as in [SV1] Thm 4.6.  $\square$

**2.7.** It is probable that a strong form of the Aomoto conjecture similar to [ESV] is valid as well.

### §3. De Rham - Witt Coulomb - KZ - Chevalley cocycle

**3.1.** Consider a simple example. Suppose that  $p > 2$ . Suppose we are given  $n$  numbers  $a_i \in \mathbb{Z}_p$ ,  $1 \leq i \leq n$ . Consider an affine space

$$\mathbb{A} = \text{Spec } \mathbb{F}_p[t, z_1, \dots, z_n]$$

and a hyperplane arrangement in it consisting of hyperplanes  $H_i : t = z_i$ ,  $H_{ij} : z_i = z_j$ .

Consider the complement

$$U = \mathbb{A} \setminus \left( \left( \bigcup_i H_i \right) \cup \left( \bigcup_{i,j} H_{ij} \right) \right) = \text{Spec } \mathbb{F}_p[t, z_1, \dots, z_n, (t - z_i)^{-1}, (z_i - z_j)^{-1}]$$

We have a Coulomb 1-form

$$\omega = - \sum_i 2a_i d \log(t - z_i) + \sum_{i,j} a_i a_j d \log(z_i - z_j) \in W\Omega^1(U)$$

*Construction of a hypergeometric cocycle*

We follow the pattern of [SV2], Section 2.

**3.2. Coulomb connection.** Suppose we are given natural  $n, N \geq 1$ , and an  $n$ -tuple

$$\mathbf{m} = (m_1, \dots, m_n) \in \mathbb{Z}_p^n.$$

Let

$$U_n = \text{Spec } \mathbb{F}_p[t_1, \dots, t_N, z_1, \dots, z_n, (z_b - z_c)^{-1}],$$

$$U_{n,N} = \text{Spec } A, \quad A = \mathbb{F}_p[t_1, \dots, t_N, z_1, \dots, z_n, (t_i - t_j)^{-1}, (t_i - z_b)^{-1}, (z_b - z_c)^{-1}].$$

We define the de Rham - Witt Coulomb closed 1-form

$$\omega_{\mathbf{m}} = \sum_{1 \leq b < c \leq n} \frac{m_b m_c}{2} d \log(z_b - z_c) + \sum_{1 \leq i < j \leq N} 2d \log(t_i - t_j) -$$

$$- \sum_{1 \leq i \leq N, 1 \leq b \leq n} m_b d \log(t_i - z_b) \in W\Omega^1(A) = W\Omega^1(U_{n,N}).$$

Given an invertible  $\kappa \in \mathbb{Z}_p$ , we define a Coulomb connection

$$\nabla_{Coul,KZ} = d + \frac{1}{\kappa} \omega_{\mathfrak{m}} : W\Omega^\bullet(U_{n,N}) \longrightarrow W\Omega^{\bullet+1}(U_{n,N}).$$

**3.3. Chevalley complex.** Let  $\mathfrak{g}$  be the Lie algebra  $\mathfrak{sl}_2$  over  $\mathbb{Z}_p$  with generators  $e, f, g$ ,  $\mathfrak{n} = \mathbb{Z}_p f \subset \mathfrak{g}$ .

For  $m \in \mathbb{Z}_p$  let

$$M(m) = \oplus_{i \geq 0} \mathbb{Z}_p f^i v$$

be the Verma module,  $ev = 0$ ; let

$$M(\mathfrak{m}) = M(m_1) \otimes_{\mathbb{Z}_p} \dots \otimes_{\mathbb{Z}_p} M(m_n) = \oplus_{i \geq 0} M(\mathfrak{m})_i$$

We consider a Chevalley complex

$$C^\bullet(M(\mathfrak{m})) : 0 \longrightarrow M(\mathfrak{m}) \xrightarrow{d} \mathfrak{n} \otimes_{\mathbb{Z}_p} M(\mathfrak{m}), \quad d(x) = f \otimes ex$$

living in degrees 0, 1,

$$C^\bullet(M(\mathfrak{m})) = \oplus_{i \geq 0} C^\bullet(M(\mathfrak{m}))_i$$

Let

$$\Omega = \frac{1}{2} h \otimes h + e \otimes e + f \otimes f \in \mathfrak{g} \otimes \mathfrak{g}$$

be the Casimir element. For  $1 \leq i < j \leq n$  we will denote by

$$\Omega_{ij} : M(\mathfrak{m}) \longrightarrow M(\mathfrak{m})$$

the action of  $\Omega$  through  $i$ -th and  $j$ -th factors.

**3.4. Coulomb - KZ - Chevalley complex.** Let

We define a bicomplex

$$C_{\mathfrak{m},N}^{\bullet\bullet} = \{W\Omega^p(U_{n,N}) \otimes_{\mathbb{Z}_p} C^q(M(\mathfrak{m}))_N\}_{p \geq 0, 0 \leq q \leq 1}.$$

The horizontal differential

$$d' = \nabla_{Coul,KZ} = d + \frac{1}{\kappa} \omega_{\mathfrak{m}} - \frac{1}{\kappa} \omega_{KZ} :$$

$$W\Omega^\bullet(U_{n,N}) \otimes_{\mathbb{Z}_p} C^\bullet(M(\mathfrak{m}))_N \longrightarrow W\Omega^{\bullet+1}(U_{n,N}) \otimes_{\mathbb{Z}_p} C^\bullet(M(\mathfrak{m}))_N$$

acts on index  $p$ , where

$$\omega_{KZ} := \sum_{1 \leq b < c \leq n} \Omega_{bc} d \log(z_b - z_c) : C_{\mathfrak{m}, N}^{pq} \longrightarrow C_{\mathfrak{m}, N}^{p+1, q},$$

the Casimirs  $\Omega_{bc}$  acting through  $M(\mathfrak{m})$  factor.

The vertical differential  $d''$  acts on index  $q$  and is induced by the differential  $d_{Ch}$  in the Chevalley complex  $C^\bullet(M(\mathfrak{m}))_N$ .

**3.5. Hypergeometric forms.** They are defined by the same formulas as in [SV2] 2.2 but now they are considered as elements of the de Rham - Witt complex.

For  $b = (b_1, \dots, b_n) \in \mathbb{N}^n$ ,  $|b| := \sum b_i = N$ , define

$$\begin{aligned} u_b = & d \log(t_1 - z_1) \wedge \dots \wedge d \log(t_{b_1} - z_1) + \\ & + d \log(t_{b_1+1} - z_2) \wedge \dots \wedge d \log(t_{b_1+b_2} - z_2) + \\ & \dots + d \log(t_{b_1+\dots+b_{n-1}+1} - z_n) \wedge \dots \wedge d \log(t_N - z_n) \in W\Omega^N(U_{n, N}), \end{aligned}$$

and for  $c = (c_1, \dots, c_n) \in \mathbb{N}^n$ ,  $|c| = N - 1$ , define

$$\begin{aligned} u_c = & -\kappa \left( d \log(t_2 - z_1) \wedge \dots \wedge d \log(t_{c_1+1} - z_1) + \right. \\ & + d \log(t_{c_1+2} - z_2) \wedge \dots \wedge d \log(t_{c_1+c_2+1} - z_2) + \\ & \left. \dots + d \log(t_{c_1+\dots+c_{n-1}+2} - z_n) \wedge \dots \wedge d \log(t_1 - z_n) \right) \in W\Omega^{N-1}(U_{n, N}), \end{aligned}$$

### *Symmetrization*

Starting from here we suppose that  $p > N$ .

The symmetric group acts on  $W\Omega^\bullet(U_{n, N})$  by permuting  $t_i$ . For  $w(t_1, \dots, t_N) \in W\Omega^\bullet(U_{n, N})$  we denote

$$\text{Alt } w(t_1, \dots, t_N) = \sum_{\sigma \in \Sigma_N} (-1)^{|\sigma|} w(t_{\sigma(1)}, \dots, t_{\sigma(N)}).$$

We define

$$w_b = \frac{1}{b_1! \dots b_n!} \text{Alt } u_b, \quad w_c = \frac{1}{c_1! \dots c_n!} \text{Alt } u_c.$$

**3.6.** Out of these forms we define a cochain in our bicomplex of total degree  $N$ : set

$$\mathcal{J}_0 = \sum_{|b|=N} w_b \otimes f^b v \in C_{\mathfrak{m},N}^{N,0}$$

where

$$f^b v = f^{b_1} \dots f^{b_n} v,$$

and

$$\mathcal{J}_1 = \sum_{|c|=N-1} w_c \otimes f \otimes f^c v \in C_{\mathfrak{m},N}^{N-1,1}.$$

**3.7. Theorem.**  $\mathcal{J} = (\mathcal{J}_0, \mathcal{J}_1)$  is a cocycle, i.e.

$$\nabla_{KZ, Coul} \mathcal{J}_0 = 0,$$

$$d_{Ch} \mathcal{J}_0 + \nabla_{KZ, Coul} \mathcal{J}_1 = 0.$$

**Proof** the same as in [SV2], 2.5.  $\square$

**3.8. Bozonization map.** Similarly to [SV2], 2.6 we can interpret our cocycle as a map of (filtered) complexes.

Namely, our bicomplex may be rewritten as a tensor product

$$C_{\mathfrak{m},N}^{\bullet\bullet} = W\Omega^\bullet(C^\bullet(M(\mathfrak{m}))_N) \otimes_{W\Omega^\bullet(U_n)} W\Omega_{\mathfrak{m}}^\bullet(U_{n,N})$$

where

$$W\Omega^\bullet(C^\bullet(M(\mathfrak{m}))_N) = W\Omega^\bullet(U_n) \otimes_{\mathbb{Z}_p} C^\bullet(M(\mathfrak{m}))_N$$

equipped with the KZ connection

$$\nabla_{KZ} = d_z - \frac{1}{\kappa} \omega_{KZ},$$

and

$$W\Omega_{\mathfrak{m}}^\bullet(U_{n,N}) = W\Omega^\bullet(U_{n,N})$$

equipped with the Coulomb connection

$$\nabla_{Coul} = d_z + \frac{1}{\kappa} \omega_{Coul}.$$



Define the dual Chevalley complex

$$C_{\bullet}(M(\mathfrak{m})^*)_N = (C^{\bullet}(M(\mathfrak{m}))_N)^*$$

living in degrees  $-1, 0$ . The  $N$ -cocycle  $\mathcal{J}$  gives rise to a map of complexes

$$\eta(\mathcal{J}) : W\Omega^{\bullet}(C_{\bullet}(M(\mathfrak{m})^*)_N) \longrightarrow W\Omega_{\mathfrak{m}}^{\bullet}(U_{n,N})^{\text{Alt}}[N]. \quad (3.8.1)$$

Moreover, it is a map of filtered complexes, where

$$F^i W\Omega^{\bullet}(C_{\bullet}(M(\mathfrak{m})^*)_N) \subset W\Omega^{\bullet}(C_{\bullet}(M(\mathfrak{m})^*)_N)$$

is the subcomplex of degree  $\geq i$  forms, and

$$F^i W\Omega_{\mathfrak{m}}^{\bullet}(U_{n,N}) \subset W\Omega_{\mathfrak{m}}^{\bullet}(U_{n,N})$$

is the subcomplex of forms having  $\geq i$  of differentials  $dz_a$ .

Let  $\mathcal{L}(\mathfrak{m}, N)$  denote the structure sheaf  $\mathcal{O}_{U_{n,N}}$  equipped with the Coulomb de Rham - Witt connection

$$\nabla_{\mathfrak{m}} = d + \frac{1}{\kappa}\omega_{\mathfrak{m}}.$$

Let  $p : U_{n,N} \longrightarrow U_n$  be the projection. The direct image  $Rp_*\mathcal{L}(\mathfrak{m}, N)$  carries the Gauss - Manin connection  $\nabla_{GM}$ , the de Rham - Witt version of [KO].

The induced by (3.8.1) map of  $E_1$ -pages of spectral sequences will be

$$\begin{aligned} \eta^i : (W\Omega^{\bullet}(H_i(M(\mathfrak{m})^*)_N)(U_n), \nabla_{KZ}) &\longrightarrow \\ \longrightarrow (W\Omega^{\bullet}(Rp_*^{N-i}(\mathcal{L}(\mathfrak{m}, N))^{\text{Alt}})(U_n), \nabla_{GM}), & i = 0, 1. \end{aligned}$$

**3.9. Theorem.** *The maps  $\eta^i$  are isomorphisms for  $1/\kappa$  close to zero in the  $p$ -adic metrics.*

**Proof.** This is a corollary of Thm. 2.6.  $\square$

**3.10.** All the above generalizes to the case of an arbitrary symmetrizable Kac - Moody algebra  $\mathfrak{g}$ . The details are left to an interested reader.

## References

- [Be] A.Beilinson, Higher regulators and values of  $L$ -functions, *Current problems in mathematics* **24**, 181–238, *Itogi Nauki i Tekhniki, VINITI*, Moscow, 1984.
- [B] S.Bloch, Crystals and de Rham - Witt connections, *J. Inst. Math. Jussieu* **3** (2004), 315 - 326.
- [Br] E.Brieskorn, Sur les groupes de tresses (d'après V.I.Arnold), Séminaire Bourbaki **401**, 1973.
- [ES] J.N.Eberhardt, J.Scholbach, Integral motivic sheaves and geometric representation theory
- [ESV] H.Esnault, V.Schechtman, E.Viehweg, Cohomology of local systems on the complement of hyperplanes, *Inv. Math.* **109** (1992), 557 - 561.
- [I1] L.Illusie, Complexe de de Rham - Witt, *Journées de Géométrie Algébrique de Rennes*, 83–112, *Astérisque* **63**, 1979.
- [I2] L.Illusie, Complexe de de Rham - Witt et cohomologie cristalline, *Ann. Sci. ENS* **12** (1979), 501 - 661.
- [K] D.Kaledin, Witt vectors, commutative and noncommutative, *Russian Math. Surveys*, **73** (2018), 3 - 34.
- [KO] N.Katz, T.Oda, On the differentiation of De Rham cohomology classes with respect to parameters, *J. Math. Kyoto Univ.* **8** (1968), 199 - 213.
- [L] S.Lubkin, Generalization of  $p$ -adic cohomology; bounded Witt vectors. A canonical lifting of a variety in characteristic  $p \neq 0$  back to characteristic zero, *Compos. Math.* **34** (1977), 225 - 277.
- [MVW] C.Mazza, V.Voevodsky, Ch.Weibel, Lecture notes on motivic cohomology
- [M] J.Milnor, Introduction to algebraic  $K$ -theory, *Annals of Math. Studies* **72**, Princeton University Press, Princeton, NJ.; University of Tokyo Press, Tokyo, 1971.
- [OS] P.Orlik, L.Solomon, Combinatorics and topology of complements of hyperplanes, *Inv. Math.* **56** (1980), 167 - 189.
- [SV1] V.Schechtman, A.Varchenko, Arrangements of hyperplanes and Lie algebra homology, *Inv. Math.* **106** (1991), 139 - 194.
- [SV2] V.Schechtman, A.Varchenko, Derived KZ equations, *J. Singul.* **25** (2022), 422 - 442.

V.S.: Institut de Mathématiques de Toulouse, Université Paul Sabatier, 118 route de Narbonne, 31062 Toulouse, France; Kavli IPMU, 5-1-5 Kashiwa-no-ha, Kashiwa-shi, Chiba, 277-8583 Japan

A.V.: Department of Mathematics, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599-3250, USA; Faculty of Mathematics and Mechanics, Lomonosov Moscow State University, Leninskiye Gory 1, 119991 Moscow GSP-1, Russia