Optimal transport map estimation in general function spaces

Vincent Divol^{1,2}, Jonathan Niles-Weed^{1,2}, Aram-Alexandre Pooladian¹

¹Center for Data Science, New York University ²Courant Institute of Mathematical Sciences, New York University vincent.divol@nyu.edu, jnw@cims.nyu.edu, aram-alexandre.pooladian@nyu.edu

December 8, 2022

Abstract

We consider the problem of estimating the optimal transport map between a (fixed) source distribution P and an unknown target distribution Q, based on samples from Q. The estimation of such optimal transport maps has become increasingly relevant in modern statistical applications, such as generative modeling. At present, estimation rates are only known in a few settings (e.g. when P and Q have densities bounded above and below and when the transport map lies in a Hölder class), which are often not reflected in practice. We present a unified methodology for obtaining rates of estimation of optimal transport maps in general function spaces. Our assumptions are significantly weaker than those appearing in the literature: we require only that the source measure P satisfies a Poincaré inequality and that the optimal map be the gradient of a smooth convex function that lies in a space whose metric entropy can be controlled. As a special case, we recover known estimation rates for bounded densities and Hölder transport maps, but also obtain nearly sharp results in many settings not covered by prior work. For example, we provide the first statistical rates of estimation when P is the normal distribution and the transport map is given by an infinite-width shallow neural network.

1 Introduction

The estimation of optimal transport maps is an increasingly relevant task in machine learning [Gra+18; Hua+21; Fin+20; ACB17; GPC18; Sal+18], with applications in computational biology [Sch+19; Mor+21; Dem+21; Yan+20a], computer graphics [Sol+16; Sol+15; Fey+17], and economics [CCG16; Che+17; TGR21; GX21]. These developments in the applied sciences have been accompanied by several recent works in the area of *statistical* estimation of optimal transport maps: we suppose that our source and target probability measures, P and Q respectively, exhibit densities on \mathbb{R}^d . Then, in accordance with Brenier's theorem (see Section 2), there exists a convex function φ_0 , called the Brenier potential, such that the optimal transport map from P to Q is given by $\nabla \varphi_0$.

Our statistical setup is the following: we have full access to P, but Q can only be accessed through i.i.d. samples $Y_1, \ldots, Y_n \sim Q$. The task of a statistician is to propose an estimator $\nabla \hat{\varphi}$ that can approximate $\nabla \varphi_0$ on the basis of these samples alone.

A modern use-case of the aforementioned setup lies in generative modeling (see e.g. the reviews [Gui+21; KPB20; YZH22]), where practitioners are given samples from a data distribution Q (e.g. a distribution of images), with the goal of generating new samples. To achieve this goal, a map T is learned so that the pushforward of P, the standard Gaussian, by T is approximately equal to Q. A new sample of law close to Q can then be generated by first sampling $X \sim P$, and then computing T(X).

Statistical transport map estimation has received considerable attention in recent years, pioneered by [HR21]. They consider the case where φ_0 lies in the class of (s + 1)-times differentiable, β -smooth, α -strongly convex functions, with P having a density bounded away from 0 and ∞ on a bounded convex domain Ω . They show that their wavelet-based estimator, $\nabla \hat{\varphi}_{(n,W)}$, achieves the following estimation rate

$$\mathbb{E} \|\nabla \hat{\varphi}_{(n,\mathrm{W})} - \nabla \varphi_0\|_{L^2(P)}^2 \lesssim n^{-\frac{2s}{2s-2+d}} \log^2(n) \,,$$

which is minimax optimal up to logarithmic factors. Subsequent work by [Man+21; DGS21; PN21; Muz+21] proposed alternative estimators for φ_0 lying in the same function class, with similar assumptions on P and Q. While seemingly natural, these assumptions are often too restrictive for practical applications. Moreover, the estimators analyzed in prior works largely do not correspond to those employed in practice. Indeed, for large-scale tasks, a number of works have proposed optimizing the empirical dual problem (see Eq. (1) below) over a class of input convex neural networks [Mak+20; BKC22]; however, this approach currently comes with no statistical guarantees.

In this work, we bridge this gap in the literature by presenting a unified perspective on estimating optimal transport maps. To do this, we leverage the fact that the Brenier potential φ_0 minimizes the function $\varphi \mapsto S(\varphi) = \int \varphi \, \mathrm{d}P + \int \varphi^* \, \mathrm{d}Q$, where φ^* is the convex conjugate of the function φ . Writing the empirical target measure as $Q_n \coloneqq n^{-1} \sum_{i=1}^n \delta_{Y_i}$, we study convergence properties of the estimator $\nabla \hat{\varphi}_{\mathcal{F}}$, where

$$\hat{\varphi}_{\mathcal{F}} \in \operatorname*{argmin}_{\varphi \in \mathcal{F}} S_n(\varphi) \coloneqq \int \varphi \, \mathrm{d}P + \int \varphi^* \, \mathrm{d}Q_n \,, \tag{1}$$

and \mathcal{F} is some function class that φ_0 either lies in or is close to. We give a general decomposition of the risk of $\hat{\varphi}_{\mathcal{F}}$ (Theorem 2 and Theorem 3) that is given in terms of the covering numbers of the class \mathcal{F} . Such structural assumptions allow us to obtain near-optimal rates for a large number of cases with a single result, such as when φ_0 belongs to:

- 1. the set of quadratics $x \mapsto \frac{1}{2}x^{\top}Ax + b^{\top}x$,
- 2. a finite set,
- 3. a parametric set,
- 4. a Reproducting Kernel Hilbert Space (RKHS),
- 5. $\mathcal{C}_L^s(\Omega)$, the Hölder ball of radius L for functions of regularity s,

- 6. a class of "spiked" potential functions (borrowing terminology from [NR22]),
- 7. a class of shallow neural networks (i.e., a Barron space [Bac17; EMW22; Bar93]),
- 8. or the space of input convex neural networks.

Additionally, unlike previous results in the literature, we are able to give general decomposition results for the risk of $\nabla \hat{\varphi}_{\mathcal{F}}$ under minimal regularity assumptions on the source measure (Theorem 2 and Theorem 3.1): we only require that P satisfies a Poincaré inequality. In particular, P can have a density (bounded away from 0 and ∞) on any domain with Lipschitz boundary, while densities proportional to $e^{-\|x\|^s}$ for $s \ge 1$ are also admissible. We therefore cover the important case where P is a Gaussian, which is novel to the best of our knowledge.

As an appetizer to our main theorems (stated in full generality in Section 3), let us consider two very different situations. In the first one (Section 4.3), we assume that φ_0 belongs to a parametric class \mathcal{F} , indexed by a finite dimensional set Θ . Then, under mild conditions on the parametrization, we show that

$$\mathbb{E} \|\nabla \hat{\varphi}_{\mathcal{F}} - \nabla \varphi_0\|_{L_2(P)}^2 \lesssim_{\log n} n^{-1},$$
(2)

that is a parametric rate of convergence holds. Although simple and expected, such a result is not present in the literature so far.

A second, non-parametric example covered by our results is when $\nabla \varphi_0$ is a shallow neural network with ReLu activation function (Section 4.7). Under such an assumption, for an appropriate class of candidate neural networks \mathcal{F} , we are able to show that

$$\mathbb{E} \|\nabla \hat{\varphi}_{\mathcal{F}} - \nabla \varphi_0\|_{L_2(P)}^2 \lesssim_{\log n} n^{-\frac{1}{2} - \frac{1}{d}},\tag{3}$$

a rate which we show is close to being minimax optimal.

Organization

We begin by providing some background on optimal transport maps in Section 2. Our main results are presented in Section 3, where we state rates of estimation in two settings: (i) when P is compactly supported, but φ_0 is not necessarily strongly convex, and (ii) when Pis unbounded, but φ_0 is strongly convex. In both cases, we require φ_0 to be (locally) smooth, so that the optimal transport map $\nabla \varphi_0$ is at least (locally) Lipschitz. We present our proofs in Section 5, omitting technical lemmas to the Appendix. We provide several examples that verify the (log-)covering condition on \mathcal{F} required to obtain our proposed convergence rates in Section 4.

Notation

Let \mathbb{R}^d be the Euclidean space for $d \geq 3$. The ball of radius R > 0 centered at $x \in \mathbb{R}^d$ is denoted by B(x; R). We assume that we have access to a collection of i.i.d. random variables $(Y_i)_{i\geq 1}$ from law Q, that are all defined on the same probability space. Depending on the context, we write either P(f), $\int f \, dP$ or $\mathbb{E}_P[f(X)]$ for the integral of a function f against a (probability) measure P, with $P(||x||^m)$ denoting the m^{th} moment of P. The pushforward of P by a measurable function $T : \mathbb{R}^d \to \mathbb{R}^m$ is written as $T_{\sharp}P$. The $L_2(\rho)$ -norm of a function $f : \mathbb{R}^d \to \mathbb{R}^m$ with respect to a σ-finite measure ρ is written as $||f||_{L_2(\rho)} = (\int ||f||^2 d\rho)^{1/2}$. For $\alpha > 1$, we write $\mathcal{C}^{\alpha}(\Omega)$ to be the space of $\lfloor \alpha \rfloor$ -times differentiable functions defined on a domain Ω whose $\lfloor \alpha \rfloor$ th derivative is $\alpha - \lfloor \alpha \rfloor$ Hölder smooth; see Equation (24) for more details. The gradient of f is written as ∇f , whereas its Hessian is written as $\nabla^2 f$. If $w : \mathbb{R}^d \to \mathbb{R}$ is a nonnegative function, we let $L_{\infty}(w)$ be the space of functions $f : \mathbb{R}^d \to \mathbb{R}$ endowed with the norm $\|f\|_{L_{\infty}(w)} = \sup_{x \in \mathbb{R}^d} |f(x)|w(x)$. For $a \in \mathbb{R}$, we write $\langle x \rangle^a$ for $(1 + \|x\|)^a$. We also let $\log_+(x)$ denote the function max $\{1, \log(x)\}$. For $d \in \mathbb{N}$, we write \mathbb{S}^d_+ (resp. \mathbb{S}^d_{++}) to be the space of symmetric positive semi-definite (resp. positive definite) matrices. The smallest eigenvalue of a symmetric matrix A is written as $\lambda_{\min}(A)$, whereas its operator norm is $||A||_{\text{op}}$. Finally, we use $a \leq b$ to indicate that there exists a constant C > 0 such that $a \leq Cb$, and will often use e.g. $\leq_{\log(n)}$ to omit logarithmic factors in n.

Throughout this work, we will repeatedly consider suprema of collections of random variables, and such a supremum may not necessarily be measurable. If this is the case, the symbols \mathbb{E} and \mathbb{P} have to be considered as representing the outer expectation of the supremum. All relevant results used to bound the expectation of such suprema hold for outer expectations, so that we will not make the distinction between expectation and outer expectation. We will also always assume implicitly that $\hat{\varphi}_{\mathcal{F}}$ is measurable: this can always be ensured by replacing \mathcal{F} by a countable dense subset (with respect to the ∞ -norm).

2 Background on optimal transport under the quadratic cost

For a domain $\Omega \subseteq \mathbb{R}^d$, let $\mathcal{P}(\Omega)$ be the space of probability measures on Ω , and $\mathcal{P}_{ac}(\Omega)$ be those with densities. Our analysis primarily hinges on existence results and stability bounds for transport maps, which we will later relate to empirical processes. We refer the reader to standard texts on optimal transport for more details e.g. [Vil09; San15].

A fundamental result concerning the existence and uniqueness of optimal transport maps is due to Brenier [Bre91]. Essentially, Theorem 1 below states that, provided the source measure has a density, a unique optimal transport map exists and is the gradient of a convex function, see [San15, Theorem 1.22, Theorem 1.40].

Theorem 1 (Brenier's Theorem). Let $P \in \mathcal{P}_{ac}(\mathbb{R}^d)$ and $Q \in \mathcal{P}(\mathbb{R}^d)$ be measures with finite second moments. Then, there exists a solution T_0 to

$$\inf_{T \in \mathcal{T}(P,Q)} \int \|x - T(x)\|^2 \,\mathrm{d}P(x)\,,\tag{4}$$

where $\mathcal{T}(P,Q) := \{T : \mathbb{R}^d \to \mathbb{R}^d \mid T_{\sharp}P = Q\}$ is the set of transport maps from P to Q. Moreover, $T_0 = \nabla \varphi_0$, for a convex function φ_0 solving

$$\inf_{\varphi \in L^1(P)} S(\varphi) := \inf_{\varphi \in L^1(P)} \int \varphi \, \mathrm{d}P + \int \varphi^* \, \mathrm{d}Q,\tag{5}$$

where φ_0^* is the convex conjugate to φ_0 . The squared 2-Wasserstein distance is written as

$$\frac{1}{2}W_2^2(P,Q) = \frac{1}{2}P(||x||^2) + \frac{1}{2}Q(||x||^2) - S(\varphi_0).$$

Moreover, as Brenier potentials are defined up to constants (indeed, shifting $\tilde{\varphi}(x) = \varphi(x) + c$ does not change the objective function in Eq. (5)), we will assume that $\varphi_0(0) = 0$, and also extend this assumption later to our general function classes of interest. Particular attention will be paid to functions φ which have a polynomial growth. The function φ is said to be (β, a) -smooth (for $\beta, a \ge 0$) if it is twice differentiable and if

$$\forall x \in \mathbb{R}^d, \ \|\nabla^2 \varphi(x)\|_{\text{op}} \le \beta \langle x \rangle^a.$$
(6)

When a = 0, this implies that the function φ is β -smooth in the classical sense. Similarly, we say that φ is (α, a) -convex (for $\alpha, a \ge 0$) if

$$\forall x \in \mathbb{R}^d, \ \lambda_{\min}(\nabla^2 \varphi(x)) \ge \alpha \langle x \rangle^a.$$
(7)

Once again, $(\alpha, 0)$ -convex functions correspond to α -strongly convex functions in the usual sense. We gather properties of such functions, that we will repeatedly use in our proofs, in Appendix B. Note that for a = 0, twice-differentiability is not required, and the usual definitions of smoothness and strong convexity can be used.

This leads us to our first proposition which states that the functional S grows at least quadratically around its minimizer φ_0 (up to logarithmic factors). We allow the potentials to either have bounded domains (where we denote by dom(φ) the domain of a function φ) or to be smooth of some exponent $a \ge 0$, while we require P to have subexponential tails: there exists a number c > 0 with $\mathbb{E}_{X \sim P}[e^{\|X\|/c}] \le 2$, where we define $\|P\|_{\exp}$ as the smallest number c > 0 satisfying this property. Equivalently, a distribution is subexponential if it has tails of the form $P(\|X\| > ct) \le Ce^{-t}$, see Appendix A for more details on subexponential distributions. A proof of Proposition 1 is found in Appendix B.

Proposition 1 (Map stability). Let P be a probability distribution with subexponential tails. Consider one of the two following settings:

- 1. the potentials φ_0 and φ_1 are (β, a) -smooth, and let b = a(a+1).
- 2. P is supported in B(0; R) and the potentials φ_0 and φ_1 are twice differentiable, with $\|\nabla \varphi_i\| \leq R$ on the support of P and $B(0; 2R) \subseteq \operatorname{dom}(\varphi_i)^1$ (i = 0, 1). In this case, let b = 0.

Assume that $\|\nabla \varphi_1(0) - \nabla \varphi_0(0)\| \leq M$ and that φ_0 is convex. Let $Q \coloneqq (\nabla \varphi_0)_{\sharp} P$ and $S(\varphi_1) \coloneqq P(\varphi_1) + Q(\varphi_1^*)$. Denoting $\ell \coloneqq S(\varphi_1) - S(\varphi_0)$, we have

$$\|\nabla\varphi_1 - \nabla\varphi_0\|_{L^2(P)}^2 \lesssim \log_+(1/\ell)^b \ell,\tag{8}$$

where the suppressed constant does not depend on d.

Remark 1. Proposition 1 is a variant of [HR21, Proposition 10], where they consider the case where P has a bounded support, resulting in the bound

$$\|\nabla \varphi - \nabla \varphi_0\|_{L^2(P)}^2 \lesssim \ell$$

Remark 2. If needed, one can weaken the assumptions on the regularity of the potentials: it is enough to assume that they are differentiable, with locally Lipschitz continuous gradients, and corresponding (local) Lipschitz norm that grows at most polynomially.

¹The constant 2 does not play any special role here, and can be replaced by any C > 1.

3 Main results

Recall the setup: we suppose that there exists a convex function φ_0 such that $Q = (\nabla \varphi_0)_{\sharp} P$. We have full access to P and to i.i.d. samples $Y_1, \ldots, Y_n \sim Q$. We are interested in studying the convergence rate of the plugin estimator $\nabla \hat{\varphi}_{\mathcal{F}}$ to $\nabla \varphi_0$, where

$$\hat{\varphi}_{\mathcal{F}} \in \operatorname*{argmin}_{\varphi \in \mathcal{F}} \int \varphi \, \mathrm{d}P + \int \varphi^* \, \mathrm{d}Q_n \, ,$$

with \mathcal{F} being a class of twice differentiable (non-necessarily convex) functions that gives a good approximation of φ_0 . This is similar to the setup proposed in [HR21], though we consider *any* class of candidate potentials \mathcal{F} , whereas they only consider a class of strongly convex functions with truncated wavelet expansions. For a subset A of a metric space (E, d), the covering number $\mathcal{N}(h, A, E)$ is defined as the smallest number of balls of radius h in Ethat are needed to cover A. We make the following assumptions throughout this paper.

- (A1) There exists $\beta, a \ge 0$ such that every $\varphi \in \mathcal{F}$ is (β, a) -smooth;
- (A2) There exist $\gamma \in [0, 2), \gamma' \ge 0$ and $D_{\mathcal{F}} \ge 1$ such that for every h > 0 and $\delta > 0$,

$$\log \mathcal{N}(h, \mathcal{F}, L_{\infty}(e^{-\delta \|\cdot\|})) \le C_{\delta} D_{\mathcal{F}} \log_{+}(1/h)^{\gamma'} h^{-\gamma};$$
(9)

(A3) The probability measure P satisfies a Poincaré inequality: for every differentiable function $f : \mathbb{R}^d \to \mathbb{R}$, it holds that

$$\operatorname{Var}_{P}(f) \leq C_{\operatorname{PI}} \int \|\nabla f(x)\|^{2} \,\mathrm{d}P(x), \tag{10}$$

where $0 \leq C_{\text{PI}} < \infty$.

Remark 3. The estimator $\hat{\varphi}_{\mathcal{F}}$ is nothing but an *M*-estimator. Therefore, it should not come as a surprise that the excess risk $S(\hat{\varphi}_{\mathcal{F}}) - S(\varphi_0)$ is related to the suprema of empirical processes of the form $\sup_{\varphi \in \mathcal{F}} \int \varphi^* d(Q - Q_n)$. Assumption (A1) enables us, through the use of the stability property Proposition 1, to relate $\|\nabla \hat{\varphi}_{\mathcal{F}} - \nabla \varphi_0\|^2$ to $S(\hat{\varphi}_{\mathcal{F}}) - S(\varphi_0)$. The latter supremum can then be controlled thanks to standard results by the covering numbers of the class of dual potentials $\{\varphi^* : \varphi \in \mathcal{F}\}$, which has to be related to the covering number of the class \mathcal{F} in order to use Assumption (A2). However, such an approach would lead to suboptimal rates of convergence. Our improved rates require to use an additional localization scheme, to replace the empirical process indexed by \mathcal{F} by a process indexed by a small ball in \mathcal{F} for the metric $(\varphi, \varphi') \mapsto \|\nabla \varphi - \nabla \varphi'\|_{L_2(P)}$. Assumption (A3) comes into play to relate the L_2 -metric at the gradient level to the L_2 -metric on the potentials.

Remark 4. Assumption (A2) allows us to cover two different scenarios. In one case, we think of the class \mathcal{F} as being a non-parametric class, fixed once and for all, that contains the true potential φ_0 . In this case, we will typically have a control of the form $\log \mathcal{N}(h, \mathcal{F}, L_{\infty}(e^{-\delta \|\cdot\|})) \leq D_{\mathcal{F}}h^{-\gamma}$ for some $0 < \gamma < 2$, so that \mathcal{F} is a Donsker class and $D_{\mathcal{F}}$ is a fixed constant. In the other scenario, we think of $\mathcal{F} = \mathcal{F}_D$ as being a "large" parametric class of dimension D (for instance, one can think of \mathcal{F} as being the class of functions

with Fourier expansions up to a finite level D). Then, the log-covering number will behave logarithmically ($\gamma' = 1$ and $\gamma = 0$), but with a prefactor $D_{\mathcal{F}}$ proportional to the dimension D of the parametric class. We will then have a standard bias-variance trade-off, with a bias representing how efficiently the potential φ_0 can be represented by the class \mathcal{F}_D . Typically, the dimension D will be chosen as n^s for some s > 0 to balance this trade-off. In particular, one should not think of $D_{\mathcal{F}}$ as being a fixed constant in this scenario, but as playing the role of the increasing dimension of an approximating parameter class.

Remark 5. The Poincaré inequality is ubiquitous in probability and analysis, being for instance connected to the phenomenon of concentration of measure [BL97; Goz10]. In general, the main obstruction to obtain a Poincaré inequality is when the measure has disconnected support. Examples of conditions that give rise to a probability measure satisfying (A3) are:

- any measure having a density bounded away from zero and infinity on a bounded Lipschitz domain (or more generally with a support satisfying the cone condition, see [AF03, Section 6]),
- $P \propto e^{-V}$ with V strongly convex (via the Brascamp-Lieb inequality [BL02]),
- $P \propto e^{-V}$, with positive constants c and a such that (i) $\langle x, \nabla V(x) \rangle \ge c ||x||$ for x large enough, or (ii) $a ||\nabla V(x)||^2 \Delta V(x) \ge c$ for x large enough (see [Bak+08]).

Special cases include the standard normal distribution, and log-concave measures with $V(x) = ||x||^s$ for $s \ge 1$. As a last comment, let us mention that the Poincaré inequality implies that P has subexponential tails [BL97]:

$$\forall t > 0, \ P(||X|| > ct) \le Ce^{-t},$$
(11)

a fact that we will repeatedly use.

3.1 The bounded case

Besides conditions (A1), (A2) and (A3), very few requirements are needed to obtain rates of convergence in the bounded setting.

(B) *P* is supported in B(0; R). Moreover, for every $\varphi \in \mathcal{F}$, φ is convex, dom(φ) = B(0; 2R) and $\|\nabla \varphi\| \leq R$ on the support of *P*.

To put it another way, we require that the candidate transport maps be uniformly bounded on the support of P, while the associated potentials are finite on a ball strictly containing the support. Such a requirement is equivalent to having an *a priori* bound R on the size of the supports of P and Q.

Theorem 2. Assume that (A1), (A2), (A3) and (B) are satisfied. Defining $\tilde{n} = n/D_F$ (that we assume is at least 2), we have

$$\mathbb{E} \|\nabla \hat{\varphi}_{\mathcal{F}} - \nabla \varphi_0\|_{L^2(P)}^2 \lesssim \inf_{\varphi \in \mathcal{F}} (S(\varphi) - S(\varphi_0)) + (\log \tilde{n})^q \tilde{n}^{-\frac{2}{2+\gamma}}, \tag{12}$$

where $q = 3 \max\{1, \gamma'\}$ and the suppressed constant does not depend on the dimension d.

Remark 6. In the above theorem, the parameter $\tilde{n} = n/D_{\mathcal{F}}$ plays the role of an effective number of samples. This is similar to what we observe for the risk of the sample mean in the Gaussian location model; the squared loss for estimating the mean θ of $N(\theta, I_{D_{\mathcal{F}}})$ is precisely \tilde{n}^{-1} . In our context, this observation is reasonable if we think about \mathcal{F} as being a set of dimension $D_{\mathcal{F}}$.

3.2 The strongly convex case

In the second setting, we do not assume that P has a bounded support anymore, but assume instead that φ_0 is strongly convex of exponent $a \ge 0$.

(C1) The potential φ_0 is (β, a) -smooth and (α, a) -convex $\alpha, \beta > 0$ for some $a \ge 0$.

To obtain improved rates of convergence, we also require mild regularity assumptions on P and its support. We say that a set A is a L-Lipschitz basic set if it is the image of the unit cube $[0, 1]^d$ by a Lipschitz diffeormorphism Φ with both Φ and Φ^{-1} that are L-Lipschitz continuous. A domain Ω is a L-Lipschitz domain if there exists a partition $(A_k)_k$ of Ω into L-Lipschitz basic sets.

(C2) The measure P has a density p and its support is a Lipschitz domain Ω . Furthermore, there exist constants $B, \theta \geq 0$ such that for every $R \geq 1$, the function $\log p$ is Lipschitz continuous on B(0; R) with Lipschitz constant BR^{θ} .

In the bounded case, assumption (C2) covers any probability distribution P having a Lipschitz density bounded away from 0 and ∞ on a Lipschitz domain. When P has full domain, assumption (C2) covers all the potentials of the form $V(x) = ||x||^s$ for some $s \ge 1$, including in particular the Gaussian case.

For $\alpha > 0$, let $\mathcal{F}_{\alpha} \subseteq \mathcal{F}$ be the set of potentials in \mathcal{F} that are (α, a) -convex for the same parameter a as in conditions (A1) and (C1).

Theorem 3. Consider P and \mathcal{F} that satisfy (A1), (A2), (A3) and (C1).

1. Define $\tilde{n} = n/D_{\mathcal{F}}$ (that we assume is at least 2). Then,

$$\mathbb{E} \|\nabla \hat{\varphi}_{\mathcal{F}} - \nabla \varphi_0\|_{L^2(P)}^2 \lesssim_{\log n} \inf_{\varphi \in \mathcal{F}_\alpha} (S(\varphi) - S(\varphi_0)) + \tilde{n}^{-\frac{2}{2+\gamma}}.$$
 (13)

2. If P also satisfies (C2), then

$$\mathbb{E} \|\nabla \hat{\varphi}_{\mathcal{F}} - \nabla \varphi_0\|_{L^2(P)}^2 \lesssim_{\log n, d} \inf_{\varphi \in \mathcal{F}_\alpha} (S(\varphi) - S(\varphi_0)) + D_{\mathcal{F}}^c n^{-\frac{2(d-\gamma)}{2d+\gamma(d-4)}},$$
(14)

where $c = 2(d-2)/(2d + \gamma(d-4))$.

The second inequality (14) does not follow directly from the entropy bound outlined in Remark 3, and requires a more delicate decomposition strategy, reminiscent of the wavelet decomposition used in [HR21]. This strategy consists in considering piecewise linear approximations of the dual potentials φ^* on a fine enough grid of the ambient space \mathbb{R}^d . The supremum of the empirical process indexed by the piecewise linear approximations is bounded through the Rademacher complexity of the process, while the remaining term is bounded by the metric entropy of the set. The key point of this decomposition is to leverage the regularity of P stated in Assumption (C2) to show that there exists a small envelope function for the remainder terms in the decomposition. As this approach is based on a decomposition of the ambient space \mathbb{R}^d , the corresponding rates of convergence now depend explicitly on d, while being faster than the rates in (13).

Remark 7. Recall our two main scenarios: either \mathcal{F} is a Donsker class, and the approximation error is zero, or \mathcal{F} is a large parametric class, so that $\gamma = 0$ and $D_{\mathcal{F}}$ plays the role of a dimension. In the first case, we obtain a rate of estimation of order $n^{-\frac{2(d-\gamma)}{2d+\gamma(d-4)}}$, that ranges from $n^{-1/2}$ (when γ approaches 2) to n^{-1} (when γ approaches 0). We will see that such a rate is close to being tight in the presence of shallow neural networks, see Section 4.7. In the other scenario (with $\gamma = 0$), the upper bound becomes

$$\inf_{\varphi \in \mathcal{F}_{\alpha}} (S(\varphi) - S(\varphi_0)) + D_{\mathcal{F}}^{\frac{d-2}{d}} n^{-1}.$$
(15)

We will see in Section 4.5 that such a decomposition leads to minimax rates when the potential φ_0 belongs to a Hölder class.

Remark 8. We do not impose that candidate potentials $\varphi \in \mathcal{F}$ are convex in Theorem 3. However, one can always obtain a convex potential by considering the biconjugate $\hat{\varphi}_{\mathcal{F}}^{**}$. As $\hat{\varphi}_{\mathcal{F}}^{**} \leq \hat{\varphi}_{\mathcal{F}}$, it always holds that $S(\hat{\varphi}_{\mathcal{F}}^{**}) \leq S(\hat{\varphi}_{\mathcal{F}})$. Griewank and Rabier [GR90] show that the biconjugate φ^{**} is locally $\mathcal{C}^{1,1}$ if the potential φ is. One can check in the proofs of [GR90] that, if φ satisfies the growth condition (A1), then the local $\mathcal{C}^{1,1}$ norm of the biconjugate also grows polynomially with the distance from the origin. This is enough to apply the stability property (Remark 2), which then shows that the risk of $\nabla \hat{\varphi}_{\mathcal{F}}^{**}$ will be of the same order as the risk of $\nabla \hat{\varphi}_{\mathcal{F}}$, while being the gradient of a convex function.

4 Examples

4.1 Transport between normal distributions

We first verify our general theorems on a simple example: the Gaussian-to-Gaussian case. It holds that for $P = N(0, I_d)$ and Q = N(b, A), the optimal Brenier potential is given by

$$\varphi_0: x \mapsto \frac{1}{2} x^\top A^{1/2} x + b^\top x \,. \tag{16}$$

Therefore, φ_0 belongs to the set of candidate potentials

$$\mathcal{F}_{\text{quad}} = \{ x \mapsto \frac{1}{2} x^{\top} A^{1/2} x + b^{\top} x : A \in \mathbb{S}^d_+, \ b \in \mathbb{R}^d \}.$$
(17)

As per our setup, we assume to have access to n samples from the distribution Q and our goal is to estimate T_0 . The natural estimator in this setting is a plug-in estimator based on the empirical covariance matrix and empirical mean of the samples from Q. It is easy to show by a direct argument that this estimator achieves the parametric rate [FLF19]. As a proof of concept, we show that this same result can be obtained by our general techniques.

One can check that the plug-in estimator is precisely the estimator that arises when optimizing the (constrained) semidual problem

$$\min_{\varphi \in \mathcal{F}_{\text{quad}}} \int \varphi \, \mathrm{d}P + \int \varphi^* \, \mathrm{d}Q_n$$

This estimator does not fall into our framework directly since the set \mathcal{F}_{quad} is not precompact, and therefore its covering numbers are infinite. However, we show in Appendix D that it suffices to control the covering number of the smaller set

$$\mathcal{F}_{\text{quad},0} = \{ x \mapsto \frac{1}{2} x^{\top} A^{1/2} x : A \in \mathbb{S}^d_+, \|A\|_{\text{op}} = 1 \}$$

which grows at most logarithmically. Theorem 3 then immediately yields the following bound.

Proposition 2. It holds that

$$\mathbb{E} \|\nabla \hat{\varphi}_{\mathcal{F}_{\text{quad}}} - \nabla \varphi_0\|_{L_2(P)}^2 \lesssim_{\log(n)} n^{-1}.$$
 (18)

Full details appear in Appendix D.

4.2 Finite set

We now consider the setup from [VV21], where Vacher and Vialard consider the problem of estimating an optimal Brenier potential over a finite class of functions $\mathcal{F} = \{\varphi_1, \ldots, \varphi_K\}$. In addition, the authors focus on the case where (i) the measure P has a bounded support, (ii) the potentials φ_k are strongly convex and smooth, and (iii) the "true" potential φ_0 does not necessarily belong to \mathcal{F} . Their main result reads

$$\mathbb{E} \|\nabla \hat{\varphi}_{\mathcal{F}} - \nabla \varphi_0\|_{L_2(P)}^2 \lesssim \min_{k=1\dots K} \|\nabla \varphi_k - \nabla \varphi_0\|_{L_2(P)}^2 + \sqrt{\frac{\log n}{n}}.$$
(19)

Theorem 2 allows us to strengthen this result if we assume a Poincaré inequality. Indeed, the log-covering number of \mathcal{F} is 0 for h small enough, so that condition (A2) holds with $\gamma = \gamma' = 0$. Therefore, if P satisfies a Poincaré inequality, a direct application of Theorem 2 yields

$$\mathbb{E} \|\nabla \hat{\varphi}_{\mathcal{F}} - \nabla \varphi_0\|_{L_2(P)}^2 \lesssim \min_{k=1\dots K} (S(\varphi_k) - S(\varphi_0)) + n^{-1} (\log n)^3.$$
⁽²⁰⁾

As in [VV21], if we also assume that the potentials in \mathcal{F} are strongly convex, we can apply a *reverse stability bound* (see [HR21, Proposition 10]), resulting in

$$\mathbb{E} \|\nabla \hat{\varphi}_{\mathcal{F}} - \nabla \varphi_0\|_{L_2(P)}^2 \lesssim \min_{k \in [K]} \|\nabla \varphi_k - \nabla \varphi_0\|_{L_2(P)}^2 + n^{-1} (\log n)^3.$$
(21)

To put it another way, assuming a Poincaré inequality is enough to improve the excess of risk from $n^{-1/2}$ to n^{-1} (up to logarithmic factors). By applying Theorem 3, the same control holds when P is not bounded and the candidate potentials $\{\varphi_k\}_{k=1}^K$ are (β, a) -smooth and (α, a) -convex for some $a \ge 0$.

4.3 Parametric set

More generally, one can consider a set of potentials parametrized by a bounded set $\Theta \subseteq \mathbb{R}^m$. Consider a set $\mathcal{F} = \{\varphi_\theta : \theta \in \Theta\}$ and assume that there exist constants $L, p \ge 0$ such that

$$\forall \theta, \theta' \in \Theta, \ \forall x \in \mathbb{R}^d, \ |\varphi_{\theta}(x) - \varphi_{\theta'}(x)| \le L \|\theta - \theta'\| \langle x \rangle^p.$$
(22)

Then, the log-covering number at scale h of \mathcal{F} for the norm $L_{\infty}(e^{-\delta \|\cdot\|})$ is controlled by the log-covering number of the set Θ , which scales as $m \log_+(1/h)$. Therefore, condition (A2) is satisfied for $\gamma = 0$, $\gamma' = 1$. Let P be a probability measure satisfying a Poincaré inequality. Theorem 3 implies that whenever $\varphi_0 \in \mathcal{F}$ is strongly convex of some exponent $a \geq 0$, it holds that

$$\mathbb{E} \|\nabla \hat{\varphi}_{\mathcal{F}} - \nabla \varphi_0\|_{L^2(P)}^2 \lesssim_{\log n} n^{-1}, \tag{23}$$

that is we obtain a parametric rate of convergence.

4.4 Reproducing Kernel Hilbert Spaces

We turn our attention to a class of transport maps given by smooth potential functions that lie in a Reproducing Kernel Hilbert Space (RKHS). We briefly recall that a RKHS is a Hilbert space \mathcal{H} of functions over a domain \mathcal{X} where there exists a *kernel* $\mathcal{K} : \mathcal{X} \times \mathcal{X} \to \mathbb{R}$ such that for all $f \in \mathcal{H}$, for all $x \in \mathcal{X}$,

$$f(x) = \langle f, \mathcal{K}(\cdot, x) \rangle_{\mathcal{H}}.$$

We refer to [Wai19; SSB+02] for more details on the properties of such spaces.

For our application of interest, we let \mathcal{F} be the unit ball in some RKHS over $\mathcal{X} = \mathbb{R}^d$, pertaining to some fixed kernel \mathcal{K} . We will assume that \mathcal{K} is of class \mathcal{C}^4 , which is enough to ensure that all potentials $\varphi \in \mathcal{F}$ are β -smooth for some $\beta > 0$ according to [Zho08]. To complete our analysis, we use existing bounds on the L_{∞} -covering numbers over such a set (see e.g. [Yan+20b]), which are expressed in terms of the spectral properties of the kernel operator. For simplicity, we assume that either the spectrum is finite or has eigenvalues that decay exponentially fast, and that the kernel is smooth enough to ensure that $f \in \mathcal{H}$ are smooth; see Appendix H for details on these definitions and assumptions. For example, when the kernel is associated to a finite-dimensional feature mapping, the RKHS is finitedimensional and thus exhibits a finite spectrum. The well-known Gaussian kernel

$$\mathcal{K}^2_{\sigma}(x,y) = \exp(-\|x-y\|^2/\sigma^2),$$

is an example of a kernel with exponentially fast decaying spectrum over the sphere, with $\sigma^2 > 0$. For such smooth kernels, one can show that the log-covering numbers exhibit the following bound [Yan+20b, Lemma D.2]

$$\log \mathcal{N}(h, \mathcal{F}, L_{\infty}(e^{-\delta \|\cdot\|})) \lesssim \log_{+}(1/h)^{\gamma'}$$

for some $\gamma' \geq 1$. Thus, if P satisfies (A3) and $\varphi_0 \in \mathcal{F}$ is strongly convex, then by Theorem 3, we again obtain parametric convergence i.e.

$$\mathbb{E} \|\nabla \hat{\varphi}_{\mathcal{F}} - \nabla \varphi_0\|_{L^2(P)}^2 \lesssim_{\log(n)} n^{-1}.$$

4.5 Hölder potentials

The first optimal transport map estimator was proposed by Hütter and Rigollet in [HR21]. For a k-times differentiable function φ , we let $d^k \varphi$ be its k^{th} order differential. Hütter and Rigollet consider the setting where the potential φ_0 belongs to $\mathcal{F} = \mathcal{C}_L^s(\Omega)$, the set of functions φ defined on some bounded domain Ω being $\lfloor s \rfloor$ -times differentiable, with $\lfloor s \rfloor$ derivatives being $(s - \lfloor s \rfloor)$ -Hölder continuous, and norm

$$\|\varphi\|_{C^{s}(\Omega)} = \max_{0 \le i \le \lfloor s \rfloor} \sup_{x \in \Omega} \|d^{i}\varphi(x)\|_{\mathrm{op}} + \sup_{x,y \in \Omega} \frac{\|d^{\lfloor s \rfloor}\varphi(x) - d^{\lfloor s \rfloor}\varphi(y)\|_{\mathrm{op}}}{\|x - y\|^{s - \lfloor s \rfloor}}$$
(24)

smaller than L. When Ω is a bounded convex domain, the authors study the estimator

$$\hat{\varphi}_{(n,\mathrm{W})} = \hat{\varphi}_{\mathcal{F}_J(\mathrm{W}^{\alpha,\beta})} \coloneqq \operatorname*{argmin}_{\varphi \in \mathcal{F}_J(\mathrm{W}^{\alpha,\beta})} S_n(\varphi), \tag{25}$$

where $\mathcal{F}_J(W^{\alpha,\beta})$ is the space of functions having a finite wavelet expansions up to some depth J, while also being α -strongly convex and β -smooth (details on wavelet expansions are given in Appendix J). Hütter and Rigollet show that, if the source measure P has a density bounded away from zero and infinity on Ω , then this estimator benefits from the rate of convergence (which is minimax optimal up to logarithmic factors), for $s \geq 2$,

$$\mathbb{E} \|\nabla \hat{\varphi}_{(n,\mathrm{W})} - \nabla \varphi_0\|_{L_2(P)}^2 \lesssim_{\log n} n^{-\frac{2(s-1)}{2s+d-4}}.$$
(26)

We are able to improve the results of Hütter and Rigollet in several directions. First, we relax the assumptions on P, which does not need to have a bounded support but is only required to satisfy a Poincaré inequality (assumption **(A3)**) and some mild regularity conditions on its density (assumption **(C2)**). The second improvement we make is related to the computation of the estimator. Indeed, in their numerical implementations, Hütter and Rigollet did not compute the estimator $\hat{\varphi}_{(n,W)}$, but removed the α -strong convexity assumptions (for otherwise, the estimator $\hat{\varphi}_{(n,W)}$ is not computable). More precisely, for J > 0 an integer and R > 0, let $\mathcal{F}_J(R)$ be the set of potentials having a finite wavelet expansion of depth J on the cube of side-length R, with bounded coefficients, and extended by an arbitrary fixed convex function outside the cube (see Appendix J for a precise definition). Then, Hütter and Rigollet use the estimator $\hat{\varphi}_{\mathcal{F}_J(R)}^{**}$ for their numerical simulations, although they do not give any theoretical guarantees for doing so. Our general theory, together with Remark 8, allows us to bound the risk of this estimator, showing the soundness of Hütter and Rigollet's numerical implementation. The proof of Proposition 3 can be found in Appendix J.

Proposition 3. Let $s \geq 2$. Assume that $\varphi_0 \in \mathcal{C}_L^s(\mathbb{R}^d)$ is α -strongly convex and β -smooth. Assume that P satisfies (A3) and (C2). Then, for some choice of R and J,

$$\mathbb{E} \|\nabla \hat{\varphi}_{\mathcal{F}_J(R)} - \nabla \varphi_0\|^2 \lesssim_{\log n} n^{-\frac{2(s-1)}{2s+d-4}}.$$
(27)

Remark 8 implies that the same rate of convergence holds for the biconjugate $\nabla \hat{\varphi}_{\mathcal{F}_J(R)}^{**}$ considered by Hütter and Rigollet, although this biconjugation step is not necessary to obtain a minimax rate of convergence.

A seminal result by Caffarelli [Caf00] states that the assumptions of Proposition 3 holds for s = 2 when the source and target measures are appropriately log-smooth and log-strongly concave: $P = \exp(-V)$ and $Q = \exp(-W)$ with $0 \leq \alpha_V I \leq \nabla^2 V(x) \leq \beta_V I$ and similarly $0 \leq \alpha_W I \leq \nabla^2 W(y) \leq \beta_W I$, with P and Q having full support on \mathbb{R}^d . In this setting, the optimal Brenier potential is $\sqrt{\alpha_V/\beta_W}$ -strongly convex and $\sqrt{\beta_V/\alpha_W}$ -smooth, thus verifying the smoothness and strong convexity requirements of Proposition 3. This principle is known as *Caffarelli's contraction theorem*; see e.g. [CP22] for a short proof based on entropic optimal transport. As a corollary, we obtain the following new result.

Corollary 1. Let $d \ge 3$, and let P and Q be log-smooth and log-strongly convex, with $Q = (\nabla \varphi_0)_{\sharp} P$. Then, for some choice of R and J,

$$\mathbb{E} \|\nabla \hat{\varphi}_{\mathcal{F}_J(R)} - \nabla \varphi_0\|^2 \lesssim_{\log n} n^{-\frac{2}{d}}.$$
(28)

4.6 "Spiked" potential functions

The spiked transport model was recently proposed by Niles-Weed and Rigollet to model situations where two probability measures of interest living in a high-dimensional space \mathbb{R}^d only differ on a low-dimensional subspace; see [NR22]. Let $\mathcal{V}_{k\times d}$ be the Stiefel manifold, consisting of all $k \times d$ matrices with orthonormal rows. We assume that there exists a matrix $U \in \mathcal{V}_{k\times d}$ such that $Y \sim Q$ can be obtained from $X \sim P$ through a map T that can be decomposed into $T(x) = U^{\top}T'(Ux) + (I - U^{\top}U)x$, while we will make structural assumption on T'. On the level of the potentials, this is equivalent to assuming that the Brenier potential φ_0 belongs to the function space

$$\mathcal{F}_{\text{spiked}} \coloneqq \left\{ x \mapsto \varphi'(Ux) + \frac{1}{2} \| x - U^{\top} Ux \|^2 : \ \varphi' \in \mathcal{F}, \ U \in \mathcal{V}_{k \times d} \right\},\tag{29}$$

where \mathcal{F} is a class of potentials defined on \mathbb{R}^k . The covering numbers of the class $\mathcal{F}_{\text{spiked}}$ are controlled by the covering numbers of the class \mathcal{F} up to logarithmic factors.

Proposition 4. It holds that

$$\log \mathcal{N}(h, \mathcal{F}_{\text{spiked}}, L_{\infty}(e^{-\delta \|\cdot\|})) \le \log \mathcal{N}(h/2, \mathcal{F}, L_{\infty}(e^{-\delta \|\cdot\|})) + C_{\delta} dk \log_{+}(1/h).$$
(30)

A proof of Proposition 4 can be found in Appendix I. This property is enough to show how one can obtain rates of convergence with exponents depending only on the effective dimension k of the problem, the dependency on d only existing through prefactors. As an example, if φ_0 is assumed to be of Hölder regularity s on the vector space spanned by U, one can use a wavelet estimator in dimension k (as in Section 4.5) to obtain a rate of convergence of order $n^{-\frac{2(s-1)}{2s+k-4}} \ll n^{-\frac{2(s-1)}{2s+d-4}}$ (with prefactors depending on d). Finding the optimal function in \mathcal{F}_{spiked} requires to solve an optimization problem defined over the Stiefel manifold. Such a problem is non trivial, and, as in [NR22], we conjecture that a computational-statistical gap exists.

4.7 Barron spaces

For large-scale tasks, neural networks are often used to parametrize mappings or deformations from a set of data to another. These deformations can be interpreted as transport maps between a source distribution (often a standard Gaussian measure) and a target distribution, which is to be learned from samples. Several works have proposed to estimate such maps using optimal transport principles [Mak+20; Hua+21; BKC22], but with few accompanying guarantees.

Our general techniques allow us to obtain statistical results in this setting, namely when the Brenier potential lies in a class of shallow neural networks with unbounded width, referred to as a *Barron space* [EMW22].

Let \mathcal{M} be a *m*-dimensional smooth compact manifold (possibly with smooth boundary), and consider an activation function $\sigma : (x, v) \in \mathbb{R}^d \times \mathcal{M} \to \mathbb{R}$. We define the $\mathcal{C}^s(\mathcal{M})$ -norm of a function $f : \mathcal{M} \to \mathbb{R}$ as in the Euclidean case (24), by using an arbitrary systems of charts on \mathcal{M} . We assume that

- (D1) for every $v \in \mathcal{M}$, $x \mapsto \sigma(x, v)$ is convex with $\sigma(0, v) = 0$, and is (β, a) -smooth,
- (D2) for every $x \in \mathbb{R}^d$, the function $v \mapsto \sigma(x, v)$ belongs to $\mathcal{C}^s(\mathcal{M})$ for some s > 0, with a uniformly bounded $\mathcal{C}^s(\mathcal{M})$ -norm.

For a given activation function, we define the Barron space \mathcal{F}_{σ} as the space of functions $\varphi : \mathbb{R}^d \to \mathbb{R}$ for which there exists a measure θ on \mathcal{M} with finite total variation such that

$$\forall x \in \mathbb{R}^d, \ \varphi(x) = \int \sigma(x, v) \,\mathrm{d}\theta(v), \tag{31}$$

with the Barron norm of φ is defined as the infimum of the total variations of measures θ such that the representation (31) holds. We let \mathcal{F}_{σ}^{1} be the unit ball in \mathcal{F}_{σ} .

Remark 9. Note that the optimal Brenier potential, φ_0 , is therefore associated with an optimal measure θ_0 on the manifold \mathcal{M} . In other words,

$$\varphi_0(x) = \int \sigma(x, v) \,\mathrm{d} heta_0(v) \,\mathrm{d} heta_0(v)$$

From a practitioner's perspective, it therefore suffices to find the optimal weights associated to the neural network by traditional means (e.g. stochastic gradient descent on the objective function of interest).

Theorem 4. Let P be a probability measure satisfying (A3) and (C2), and let σ be an activation function satisfying (D1) and (D2). Let $\varphi_0 \in \mathcal{F}_{\sigma}^1$ be (α, a) -strongly convex. Then,

$$\mathbb{E}[\|\nabla \hat{\varphi}_{\mathcal{F}_{\sigma}^{1}} - \nabla \varphi_{0}\|_{L_{2}(P)}^{2}] \lesssim_{\log n, d} n^{-\frac{2s+m(1-2/d)}{2s+2m(1-2/d)}}.$$
(32)

The exponent in Theorem 4 always lies between -1 and -1/2, in particular not depending on the ambient dimension d. Note however that the hidden prefactor does depend exponentially on d, so that finer bounds would be needed to completely break the curse of dimensionality. A case of particular interest for applications is when $\sigma(x, v) = \sigma(\langle x, v \rangle)$ for

some convex function $\sigma : \mathbb{R} \to \mathbb{R}$, where v is in the unit sphere $\mathcal{M} = \mathcal{S}^{d-1}$ (of dimension m = d - 1). We assume that $\sigma = \sigma_s$ is given by $\sigma_s(u) = u_+^s$. In this situation, we may use Theorem 4 to get the following rate, that we show is almost minimax when the dimension is large.

Proposition 5. Let P be a probability measure satisfying (A3) and (C2), Let $\sigma = \sigma_s$ for some $s \geq 2$. Let $\varphi_0 \in \mathcal{F}_{\sigma}^1$ be (α, a) -strongly convex. Then,

$$\mathbb{E}[\|\nabla \hat{\varphi}_{\mathcal{F}_{\sigma_s}^1} - \nabla \varphi_0\|_{L_2(P)}^2] \lesssim_{\log n, d} n^{-\frac{2s+d-3+2/d}{2s+2(d-3+2/d)}}.$$
(33)

Furthermore, if d is odd, it holds that

$$\inf_{\hat{T}} \sup_{Q \in \mathcal{Q}_{\sigma_s,P}} \mathbb{E}[\|\hat{T} - \nabla \varphi_0\|_{L_2(P)}^2] \gtrsim n^{-\frac{2s+d-1}{2s+2d-3}}.$$
(34)

For large d, the exponent in both the upper and lower bounds is $-\frac{2s+d}{2s+2d}(1+o_d(1))$. Theorem 4 and Proposition 5 are proven in Appendix K.

As a particular case of the above theorem, when s = 2, we have $\sigma_s(u) = \frac{u_+^2}{2}$, so that $\sigma'_s(u) = u_+$ is the ReLu activation function. The transport maps arising from the set $\mathcal{F}_{\sigma_s}^1$ are of the form $x \mapsto \int \langle x, v \rangle_+ v \cdot d\theta(v)$, which are shallow neural networks with a ReLu activation function. Proposition 5 gives the upper bound $n^{-\frac{1}{2}-\frac{1}{d-1+1/d}}$, with the near-matching lower bound $n^{-\frac{1}{2}-\frac{5}{4d+2}}$. For instance, for d = 11, the exponent in the upper bound is 0.599, whereas the exponent in the lower bound is approximately equal to 0.612. We conjecture that the minimax rate for this problem is $n^{-\frac{1}{2}-\frac{1}{d-1}}$.

4.8 Input convex neural networks

As a last example, Makkuva & al. [Mak+20] propose to use input convex neural networks to estimate Brenier potentials. In our language, their estimator is exactly $\hat{\varphi}_{\mathcal{F}}$, where \mathcal{F} is a class of input convex neural networks that can be parametrized using m weights (for some large m). Assuming that all the weights of the neural network are uniformly bounded, one can check that (22) is satisfied, so that the log-covering number at scale h is of order $m \log_+(1/h)$. One can then use Theorem 3 to obtain a rate of convergence of order

$$\mathbb{E} \|\nabla \hat{\varphi}_{\mathcal{F}} - \nabla \varphi_0\|_{L^2(P)}^2 \lesssim_{\log n} \inf_{\varphi \in \mathcal{F}_\alpha} (S(\varphi) - S(\varphi_0)) + (n/m)^{-1}.$$
(35)

To obtain a more satisfying rate of convergence, one needs to understand the approximation properties of input convex neural networks, in order to be able to control the bias term in the above upper bound. Quantifying this bias is an important question in the theory of the quantitative approximation properties of input convex neural networks, which is an attractive topic for future work.

5 Proofs

The key technique to proving the stated rates is van de Geer's "one-shot" localization technique, which was successfully used by Hütter and Rigollet for dealing with Hölder potentials [HR21]. To ease notation, we will write $\hat{\varphi}$ instead of $\hat{\varphi}_{\mathcal{F}}$.

Let $\overline{\varphi} \in \mathcal{F}$ be any potential. Let $\tau > 0$ be a parameter to fix and consider

$$\hat{\varphi}_t = t\hat{\varphi} + (1-t)\overline{\varphi}, \text{ where } t = \frac{\tau}{\tau + \|\nabla\hat{\varphi} - \nabla\overline{\varphi}\|_{L_2(P)}}.$$
 (36)

The potential $\hat{\varphi}_t$ is localized in the sense that

$$\|\nabla\hat{\varphi}_t - \nabla\overline{\varphi}\|_{L_2(P)} = \frac{\tau \|\nabla\hat{\varphi} - \nabla\overline{\varphi}\|_{L_2(P)}}{\tau + \|\nabla\hat{\varphi} - \nabla\overline{\varphi}\|_{L_2(P)}} \le \tau.$$
(37)

Let $\ell_t = S(\hat{\varphi}_t) - S(\varphi_0)$ and $\ell = S(\overline{\varphi}) - S(\varphi_0)$. As both $\hat{\varphi}$ and φ_0 satisfy the smoothness condition **(A1)**, so does $\hat{\varphi}_t$. Therefore, the stability property (Proposition 1) implies that (with b = a(a+1) or b = 0 depending on whether P has a bounded support or not)

$$\left(\frac{\tau \|\nabla \hat{\varphi} - \nabla \overline{\varphi}\|_{L_2(P)}}{\tau + \|\nabla \hat{\varphi} - \nabla \overline{\varphi}\|_{L_2(P)}}\right)^2 = \|\nabla \hat{\varphi}_t - \nabla \overline{\varphi}\|_{L_2(P)}^2 \le 2\|\nabla \hat{\varphi}_t - \nabla \varphi_0\|_{L_2(P)}^2 + 2\|\nabla \overline{\varphi} - \nabla \varphi_0\|_{L_2(P)}^2 \\ \le \kappa \log_+ (1/\ell_t)^b \ell_t + \kappa \log_+ (1/\ell)^b \ell, \tag{38}$$

where κ depends on an upper bound on $\|\nabla \hat{\varphi}_t(0) - \nabla \varphi_0(0)\|$ and $\|\nabla \overline{\varphi}(0) - \nabla \varphi_0(0)\|$. Let us first bound ℓ_t . The convexity of S_n and the optimality of $\hat{\varphi}$ for S_n implies that

$$S_n(\hat{\varphi}_t) \le t S_n(\hat{\varphi}) + (1-t) S_n(\overline{\varphi}) \le S_n(\overline{\varphi}).$$
(39)

Remark that the functionals S and S_n are invariant under translations: it holds that $S(\varphi) = S(\varphi + c)$ for every $c \in \mathbb{R}$ and every potential φ . Let $\varphi_c = \varphi - P(\varphi)$ be the centered version of φ . We obtain

$$\ell_{t} = S(\hat{\varphi}_{t}) - S(\varphi_{0}) \leq S(\hat{\varphi}_{t,c}) - S_{n}(\hat{\varphi}_{t,c}) + S_{n}(\overline{\varphi}_{c}) - S(\varphi_{0})$$

$$= \left\{ S(\hat{\varphi}_{t,c}) - S_{n}(\hat{\varphi}_{t,c}) + S_{n}(\overline{\varphi}_{c}) - S(\overline{\varphi}_{c}) \right\} + S(\overline{\varphi}) - S(\varphi_{0})$$

$$= (Q - Q_{n})(\hat{\varphi}_{t,c}^{*} - \overline{\varphi}_{c}^{*}) + \ell$$

$$\leq Z_{n} + \ell,$$
(40)

where $Z_n = |(Q - Q_n)(\hat{\varphi}_{t,c}^* - \overline{\varphi}_c^*)|$. Therefore,

$$\left(\frac{\tau \|\nabla \hat{\varphi} - \nabla \overline{\varphi}\|_{L_2(P)}}{\tau + \|\nabla \hat{\varphi} - \nabla \overline{\varphi}\|_{L_2(P)}}\right)^2 \le 2\kappa \log_+ \left(\frac{1}{\ell + Z_n}\right)^b (\ell + Z_n) = v_n.$$
(41)

The following implication holds:

$$v_n \le \frac{\tau^2}{4} \implies \|\nabla\hat{\varphi} - \nabla\overline{\varphi}\|_{L_2(P)}^2 \le \tau^2.$$
 (42)

Thus, the bulk of the proof consists in finding a good bound on Z_n that holds with high probability. This is where the different assumptions of Theorem 2 and Theorem 3 will come into play.

Let A be a subset of a metric space (E, d) consisting of functions from \mathbb{R}^d to \mathbb{R} . Given two functions $f_1, f_2 : \mathbb{R}^d \to \mathbb{R}$, the *bracket* $[f_1, f_2]$ is the set of functions $g : \mathbb{R}^d \to \mathbb{R}$ such that $f_1(x) \leq g(x) \leq f_2(x)$ for every $x \in \mathbb{R}^d$. The *h*-bracketing number $\mathcal{N}_{[]}(h, A, E)$ is the smallest number of brackets $[f_1, f_2]$ needed to cover A, with $d(f_1, f_2) \leq h$. For $\delta > 0$, we define the δ -bracketing integral of A as the number

$$\mathcal{J}_{[\]}(\delta, A, E) \coloneqq \int_0^\delta \sqrt{\log(2\mathcal{N}_{[\]}(h, A, E))} \,\mathrm{d}h \in (0, +\infty].$$
(43)

We say that a class of functions \mathcal{F} is *E-Donsker* if the integral $\mathcal{J}_{[]}(1, \mathcal{F}, E)$ is finite, where E is some metric space containing \mathcal{F} . A function $G : \mathbb{R}^d \to \mathbb{R}$ is an *envelope* for a class of functions \mathcal{G} if $|g(x)| \leq G(x)$ for every $x \in \mathbb{R}^d$ and every $g \in \mathcal{G}$.

Proposition 6 (Remark 3.5.14 in [GN21]). Let $\sigma > 0$ be a number such that $\sigma^2 \ge \sup_{g \in \mathcal{G}} Qg^2$ and let G be an envelope for \mathcal{G} . Assume that $\mathcal{J}_{[\]}(2||G||_{L_2(Q)}, \mathcal{G}, L_2(Q))$ is finite and let

$$\omega = \sqrt{\frac{n\sigma^2}{32\log(2\mathcal{N}_{[]}(\sigma/2,\mathcal{G},L_2(Q)))}}.$$
(44)

Then,

$$\mathbb{E}\left[\left|\sup_{g\in\mathcal{G}}(Q_n-Q)(g)\right|\right] \leq \frac{116}{\sqrt{n}}\mathcal{J}_{[\]}(2\sigma,\mathcal{G},L_2(Q)) + 8\int G(y)\mathbf{1}\{G(y)>\omega\}\,\mathrm{d}Q(y).$$
(45)

In both cases (bounded support or φ_0 strongly convex), we require the following threestep procedure: (i) find an envelope function of \mathcal{G} , (ii) find a bound on $\sigma^2 = \sup_{g \in \mathcal{G}} ||g||_{L_2(Q)}$, and (iii) find a bound on the bracketing numbers $\mathcal{N}_{[]}(h, \mathcal{G}, L_2(Q))$ of \mathcal{G} for the $L_2(Q)$ -norm.

In what follows, we will always choose the parameter τ so that $\tau^2 \geq \ell$.

5.1 Proof of Theorem 2

We first consider the case where P is bounded (assumption **(B)**). In this case, the parameter b is equal to zero, and the logarithmic factor in (41) disappears. Furthermore, as all transport maps involved are bounded by R, the constant κ in (38) only depends on R. We also let $\overline{\varphi}$ be the potential minimizing $S(\varphi) - S(\varphi_0)$ over $\varphi \in \mathcal{F}$ (if the infimum is not attained, we use a standard approximation scheme).

Introduce the set $\operatorname{Li}(\mathcal{F}) = \{t\varphi + (1-t)\varphi' : \varphi, \varphi' \in \mathcal{F}, 0 \leq t \leq 1\}$ of lines between elements of \mathcal{F} . Remark that the potential $\hat{\varphi}_t$ is in $\operatorname{Li}(\mathcal{F})$ and satisfies $\|\nabla \hat{\varphi}_t - \nabla \overline{\varphi}\|_{L_2(P)} \leq \tau$ (recall (37)); this encourages the definition

$$\operatorname{Li}(\mathcal{F})_{\tau} = \left\{ \varphi \in \operatorname{Li}(\mathcal{F}) : \|\nabla \varphi - \nabla \overline{\varphi}\|_{L_2(P)} \le \tau \right\}.$$
(46)

Therefore, we bound Z_n by $|\sup_{q\in\mathcal{G}}(Q-Q_n)(g)|$, where

$$\mathcal{G} = \{ \varphi_c^* - \overline{\varphi}_c^* : \varphi \in \operatorname{Li}(\mathcal{F})_\tau \}.$$
(47)

As per our three-step procedure, we require the three following lemmas, whose proofs are deferred to Appendix E.

Lemma 1 (Envelope for \mathcal{G} , bounded case). The constant function equal to $G \equiv 5R^2$ is an envelope function for \mathcal{G} .

Lemma 2 (Uniform second moment on \mathcal{G} , bounded case). It holds that

$$\sup_{g \in \mathcal{G}} \|g\|_{L_2(Q)} \le \sigma \coloneqq C\tau.$$
(48)

Lemma 3 (Bracketing numbers of \mathcal{G} , bounded case). Pick $\delta = 1/(2R)$. For h > 0, it holds that

$$\log \mathcal{N}_{[]}(h,\mathcal{G},L_2(Q)) \le \log \mathcal{N}(ch,\mathcal{F},L_{\infty}(e^{-\delta \|\cdot\|})) + C\log_+(1/h).$$
(49)

Equipped with these lemmas, we are ready to apply Proposition 6. By assumption (A2), and Lemma 3, we obtain that for h > 0

$$\log \mathcal{N}_{[]}(h, \mathcal{G}, L_2(Q)) \lesssim D_{\mathcal{F}} \log_+(1/h)^{\gamma''} h^{-\gamma},$$
(50)

where

$$\gamma'' = \begin{cases} \gamma' & \text{if } \gamma > 0, \\ \max\{1, \gamma'\} & \text{if } \gamma = 0. \end{cases}$$

Therefore, the bracketing integral is smaller than

$$\mathcal{J}(2\sigma, \mathcal{G}, L_2(Q)) \lesssim \int_0^{2\sigma} \sqrt{D_{\mathcal{F}} \log_+(1/h)^{\gamma''} h^{-\gamma}} \,\mathrm{d}h \lesssim \left\{ D_{\mathcal{F}} \log_+(1/\tau)^{\gamma''} \tau^{2-\gamma} \right\}^{\frac{1}{2}}.$$

Let $\tilde{n} = n/D_{\mathcal{F}}$ and

$$\tau = \eta \max\left\{\tilde{n}^{-\frac{1}{2+\gamma}} (\log \tilde{n})^{\frac{q}{2}}, \ell^{\frac{1}{2}}\right\},\tag{51}$$

where $q = 3 \max\{1, \gamma'\}$ and $\eta > 0$ is a parameter to fix, and we recall that $\ell = S(\overline{\varphi}) - S(\varphi_0) = \inf_{\varphi \in \mathcal{F}} (S(\varphi) - S(\varphi_0))$. Then, the estimate (50) yields that

$$\omega = \sqrt{\frac{n\sigma^2}{32\log(2\mathcal{N}_{[]}(\sigma/2,\mathcal{G},L_2(Q)))}} \ge G$$
(52)

if the constant η is chosen large enough. By Proposition 6, it holds that

$$\mathbb{E}[Z_n] \le \mathbb{E}[|\sup_{g \in \mathcal{G}} (Q - Q_n)(g)|] \le C \left\{ \frac{\log_+(1/\tau)^{\gamma''} \tau^{2-\gamma}}{\tilde{n}} \right\}^{\frac{1}{2}} =: m.$$
(53)

Although the expectation of Z_n is controlled, we are not done yet: to be able to leverage Eq. (42), the random variable Z_n needs to be controlled with high probability. To do this, we incorporate the following lemma, which is a special case of Lemma 11 (see Appendix A).

Lemma 4. Let \mathcal{G} be a class of functions such that for every $g \in \mathcal{G}$, $|g(y)| \leq L\langle y \rangle^s$ for some $s \geq 1$, $L \geq 0$. Assume that $Q = (\nabla \varphi_0)_{\#} P$ with φ_0 that is (β, a) -smooth for some $\beta, a \geq 0$. Let m be such that $m \geq \mathbb{E}[|\sup_{g \in \mathcal{G}} (Q_n - Q)(g)|]$ and $m \geq n^{-1} (\log n)^{s(a+1)}$. Then, for every u > 0,

$$\mathbb{P}\left\{\left|\sup_{g\in\mathcal{G}}(Q_n-Q)(g)\right| > cu^{s(a+1)}m\right\} \le Ce^{-u}.$$
(54)

Let us now conclude. First, if $|Z_n| \leq \ell$, by choosing η large enough, we obtain that

$$v_n = 2\kappa(Z_n + \ell) \le 4\kappa\ell \le \frac{\tau^2}{2},$$

so that we can use (42) to control $\|\nabla \hat{\varphi} - \nabla \overline{\varphi}\|_{L_2(P)}$. In the case $\ell \leq |Z_n|$, we apply Lemma 4 with s = 1, a = 0 and $u = \log \tilde{n}$. Indeed, the condition $m \geq (\log n)n^{-1}$ is satisfied for m appearing in (53) (for η large enough), so that, with probability $1 - C\tilde{n}^{-1}$,

$$v_n \le 4\kappa |Z_n| \lesssim (\log \tilde{n})m_s$$

a quantity that is smaller than $\frac{\tau^2}{2}$ (once again, for η large enough).

In total, we have shown that with probability $1 - C\tilde{n}^{-1}$, $\|\nabla\hat{\varphi} - \nabla\overline{\varphi}\|_{L_2(P)} \leq \tau$. As we also have the crude bound $\|\nabla\hat{\varphi} - \nabla\overline{\varphi}\| \leq 2R$, we obtain the final decomposition

$$\mathbb{E}[\|\nabla\hat{\varphi} - \nabla\overline{\varphi}\|_{L_2(P)}^2] \le \tau^2 + 4R^2 \mathbb{P}\left\{\|\nabla\hat{\varphi} - \overline{\varphi}\|_{L_2(P)} > \tau\right\} \le \tau^2 + 4CR^2\tilde{n}^{-1}.$$
 (55)

We may also bound

$$\mathbb{E}[\|\nabla\hat{\varphi} - \nabla\varphi_0\|_{L_2(P)}^2] \le 2\|\nabla\overline{\varphi} - \nabla\varphi_0\|_{L_2(P)}^2 + 2\mathbb{E}[\|\nabla\hat{\varphi} - \nabla\overline{\varphi}\|_{L_2(P)}^2] \\ \lesssim \ell + \tau^2 + \tilde{n}^{-1}$$

(where we apply Proposition 1). This gives us the desired bound.

5.2 Proof of Theorem 3

The proof of Theorem 3, where we do not assume that P has a bounded support, is more delicate. As the exponents in the different logarithmic factors start to have complicated expressions (that are polynomial in a and linear in d), we keep them implicit in the proof.

First, we show in Appendix F that we can assume without loss of generality that there exists a constant r such that $\|\nabla\varphi(0)\| \leq r$ for every $\varphi \in \mathcal{F}$, a property that will be useful. In particular, the constant κ appearing in (38) is controlled. We assume that the set \mathcal{F}_{α} is nonempty, for otherwise there is nothing to prove. Let $\overline{\varphi}$ be a potential that attains the infimum of $S(\varphi) - S(\varphi_0)$ over \mathcal{F}_{α} . Let $\lambda > 1$. We write

$$\mathbb{E}[\|\nabla\hat{\varphi}_{\mathcal{F}} - \nabla\overline{\varphi}\|_{L_{2}(P)}^{2}] \leq (\lambda\tau)^{2} + \mathbb{E}[\|\nabla\hat{\varphi}_{\mathcal{F}} - \nabla\overline{\varphi}\|_{L_{2}(P)}^{2}\mathbf{1}\{\|\nabla\hat{\varphi}_{\mathcal{F}} - \nabla\overline{\varphi}\|_{L_{2}(P)}^{2} \geq \lambda\tau\}].$$
(56)

As $\hat{\varphi}_{\mathcal{F}}$ and $\overline{\varphi}$ are (β, a) -smooth, their gradients grow at most as $2\beta \langle x \rangle^{a+1}$ (see Lemma 14). Therefore, the norm $\|\nabla \hat{\varphi}_{\mathcal{F}} - \nabla \overline{\varphi}\|_{L_2(P)}^2$ is bounded by a constant *C* depending on the moment of *P* of order 2a + 1 (which is finite: recall that the Poincaré inequality implies a subexponential tail), and we have the bound

$$\mathbb{E}[\|\nabla\hat{\varphi}_{\mathcal{F}} - \nabla\overline{\varphi}\|_{L_2(P)}^2] \le (\lambda\tau)^2 + C \cdot \mathbb{P}\left\{\|\nabla\hat{\varphi}_{\mathcal{F}} - \nabla\overline{\varphi}\|_{L_2(P)}^2 \ge \lambda\tau\right\}.$$
(57)

Assume that $\|\nabla \hat{\varphi}_{\mathcal{F}} - \nabla \overline{\varphi}\|_{L_2(P)} \ge \lambda \tau$. Then, the parameter t defined in (36) is smaller than $1/(1+\lambda)$. Choose $\lambda = 2(1+\beta/\alpha)$, so that $-t\beta + (1-t)\alpha \ge \frac{\alpha}{2}$. As the potential $\overline{\varphi}$ is (α, a) -strongly convex and the potential $\hat{\varphi}$ is (β, a) -smooth, the previous inequality shows that the potential $\hat{\varphi}_t$ is $(\alpha/2, a)$ -strongly convex. Therefore, the function $g = \hat{\varphi}_{t,c}^* - \overline{\varphi}_c^*$ belongs to the set

$$\mathcal{G} = \{\varphi_c^* - \overline{\varphi}_c^* : \varphi \in \operatorname{Li}(\mathcal{F})_\tau, \varphi \text{ is } (\alpha/2, a) \text{-convex}\},$$
(58)

As in the previous proof, we require three lemmas, whose proofs are deferred to Appendix F.

Lemma 5 (Envelope for \mathcal{G} , strongly convex case). The function $G: y \mapsto C(r^2 + ||y||^2) + \tau \sqrt{C_{PI}}$ is an envelope function for \mathcal{G} .

Lemma 6 (Uniform second moment on \mathcal{G} , strongly convex case). It holds that

$$\sup_{g \in \mathcal{G}} \|g\|_{L_2(Q)} \le \sigma \coloneqq C \log_+(1/\tau)^c \tau.$$
(59)

Lemma 7 (Bracketing numbers of \mathcal{G} , strongly convex case). For δ a small enough constant, and for h > 0, it holds that

$$\log \mathcal{N}_{[]}(h, \mathcal{G}, L_2(Q)) \le \log \mathcal{N}(ch, \mathcal{F}, L_{\infty}(e^{-\delta \|\cdot\|})) + C \log_+(1/h).$$
(60)

Having these three lemmas at our disposal, we can conclude as in the proof of Theorem 2: this yields an excess of risk of order $\tilde{n}^{-\frac{2}{2+\gamma}}$ (up to polylogarithmic factors). The first inequality in Theorem 3 is obtained in this fashion, and we leave the details to the reader. Such a rate is however suboptimal when P satisfies condition (C2). In this situation, a better bound on $Z_n \leq \mathbb{E}[|\sup_{g \in \mathcal{G}} (Q - Q_n)(g)|]$ is possible. First, we show that we can safely assume that Q has a bounded support, of radius of polylogarithmic order.

Lemma 8. Let $g_{\leq R}$ be the function g restricted to B(0; R), and Q^R the probability Q conditioned on being in B(0; R), with Q_n^R being defined likewise. For $R = c(\log n)^{a+1}$ and some constant c large enough, it holds that

$$\mathbb{E}[|\sup_{g\in\mathcal{G}}(Q-Q_n)(g)|] \le \mathbb{E}[|\sup_{g\in\mathcal{G}}(Q^R-Q_n^R)(g_{\le R})|] + n^{-1}.$$
(61)

Furthermore, the measure Q^R has a L-Lipschitz domain for some $L \leq_{\log n} 1$ and the density q^R satisfies for every ball U of radius r

$$\frac{\sup_U q^R}{\inf_U q^R} \lesssim_{\log n} e^{cr(\log n)^{\kappa}} \tag{62}$$

for some constants c and κ .

Lemma 8 is proven in Appendix F. We will now write Q, Q_n, g instead of $Q^R, Q_n^R, g_{\leq R}$ to ease notation. We consider a partition \mathcal{A}_J of the support Ω of Q into cube-like shapes of size of order 2^{-J} (see Appendix G for details). Any function $g \in \mathcal{G}$ is approximated by $\Pi_J g$, which is given by the $L_2(Q)$ -projection of g on the sets of piecewise linear functions, linear on each block of the partition. We then split the expectation

$$\mathbb{E}[|\sup_{g\in\mathcal{G}}(Q-Q_n)(g)|] \le \mathbb{E}[|\sup_{g\in\mathcal{G}}(Q-Q_n)(\Pi_J g)|] + \mathbb{E}[|\sup_{g\in\mathcal{G}}(Q-Q_n)(g-\Pi_J g)|].$$
(63)

As the functions $\Pi_J g$ depend only on a finite number of parameters, the first supremum can be shown to be of order $n^{-1/2}$. We now assume that τ and 2^{-J} are at least of order n^{-q} for some parameter q > 0, so that $\log_+(1/\tau)$ and J are all of order at most $\log n$. This condition will be satisfied for our final choices of τ and J. Lemma 9. It holds that

$$\mathbb{E}[|\sup_{g\in\mathcal{G}}(Q-Q_n)(\Pi_J g)|] \lesssim_{\log n,d} \frac{2^{\frac{d-2}{2}J}\tau}{\sqrt{n}}.$$
(64)

The remainder term $\mathbb{E}[|\sup_{g\in\mathcal{G}}(Q-Q_n)(g-\Pi_J g)|]$ is bounded using Proposition 6. However, unlike in the proof of Theorem 2, we may exploit to our advantage that, by a Taylor expansion, the functions $g - \Pi_J g$ are bounded by a small constant with respect to the L_{∞} norm, of order 2^{-2J} , while they have a $L_2(Q)$ -diameter of order $2^{-J}\tau$. These considerations are enough to obtain the following bound.

Lemma 10. Let $\mathcal{G}_{>J} = \{g - \prod_J g : g \in \mathcal{G}\}$. Assume that $2^{-J(2-\gamma)} \ge c_1(\log n)^{c_2} \frac{n}{D_F} \tau^{2+\gamma}$ for some constants c_1 and c_2 large enough. Then, it holds that

$$\mathbb{E}[|\sup_{g\in\mathcal{G}}(Q-Q_n)(g-\Pi_J g)|] \lesssim_{\log n} \left(\frac{D_{\mathcal{F}}(2^{-J}\tau)^{2-\gamma}}{n}\right)^{1/2}.$$
(65)

The proofs of Lemma 9 and Lemma 10 are given in Appendix G. Let J be such that $2^{-J} \simeq (D_F \tau^{-\gamma})^{-\frac{1}{d-\gamma}}$ and let

$$\tau = \eta \max\{(\log n)^q \left(\frac{D_F^{d-2}}{n^{d-\gamma}}\right)^{\frac{1}{2d+\gamma(d-4)}}, \ell^{1/2}\},$$

where we choose η , q large enough. By Lemma 9 and Lemma 10, we obtain

$$\mathbb{E}[|Z_n|] \lesssim_{\log n} \tau^{\frac{d(2-\gamma)}{2(d-\gamma)}} \frac{D_{\mathcal{F}}^{\frac{d-2}{2(d-\gamma)}}}{\sqrt{n}}.$$
(66)

One obtains using Lemma 4 that the random variable $Z_n + \ell$ is controlled by (66) (up to polylogarithmic factors) with probability $1 - Cn^{-1}$. We check that this upper bound is smaller than $\frac{(\lambda \tau)^2}{2}$, should we choose η and q large enough. We therefore have proven using (42) that $\mathbb{P}\left\{ \|\nabla \hat{\varphi}_{\mathcal{F}} - \nabla \overline{\varphi}\|_{L_2(P)} > \lambda \tau \right\} \leq Cn^{-1}$, which concludes the proof using (57).

Remark 10. The proofs of Theorem 2 and Theorem 3.1 never use that P has a density; only the Poincaré inequality is required. Hence, uniform probability measures P on compact submanifolds of \mathbb{R}^d , such as the sphere are also admissible. The proof of Theorem 3.2 can also be adapted to the manifold setting, by covering a manifold by a locally finite number of charts, and using a partition of each chart by small cube-like shapes. The rates obtained would then depend on the dimension on the manifold instead of the ambient dimension d. We leave the details to the interested reader.

Acknowledgements

JNW is partially supported by National Science Foundation grant DMS-2210583. AAP is partially supported by Natural Sciences and Engineering Research Council of Canada (NSERC), the National Science Foundation through NSF Award 1922658, and Meta AI Research.

A Properties of subexponential distributions

We gather in this section results on subexponential distributions relevant to us.

Lemma 11. Let \mathcal{G} be a class of measurable functions with measurable envelope $G(y) = L\langle y \rangle^s$ for some $s \geq 1$. Assume that $Q = T_{\sharp}P$ where P is subexponential and and $||T(x)|| \leq L\langle x \rangle^r$ for some $r \geq 1$. Let $Z = |\sup_{a \in \mathcal{G}} (Q_n - Q)(g)|$. Then, for u > 0,

$$\mathbb{P}\left\{Z > u^{rs}c(\mathbb{E}[Z] + n^{-1}(\log n)^{rs})\right\} \le Ce^{-u}.$$
(67)

Proof. We apply Theorem 2.14.5 in [VW96]. Indeed, the random variable $Y \sim Q$ belongs to the Orlicz space of exponent 1/r, while the measurable envelope has a finite Orlicz norm of exponent 1/(rs).

Lemma 4 follows from this inequality. Indeed, $\|\nabla \varphi_0(x)\| \leq \langle x \rangle^r$ for r = a + 1 (see Lemma 14 below), so that one can apply Lemma 11. Subexponential variables also have controlled moments.

Lemma 12. Let P be a subexponential probability distribution. Let $a \ge 1$ and $m \ge ||P||_{exp}$. Let $r \ge r_0 = 2m(a-1)$. Then,

$$\mathbb{E}_{P}[\|X\|^{a}] \leq C_{a}m^{a}$$

$$\mathbb{E}_{P}[\|X\|^{a}\mathbf{1}\{\|X\| \geq r\}] \leq C_{a}r^{a}e^{-r/m}.$$
(68)

Proof. The first inequality is standard, see e.g. [Ver18, Section 2.7]. The second is obtained by integrating the bound $P(||X|| \ge mt) \le 2e^{-t}$ (that is obtained through a Chernoff's bound).

We conclude this section by a technical lemma, that shows that the integral of a function f against a subexponential distribution P is stable under polynomial perturbations of f.

Lemma 13. Let $f : \mathbb{R}^d \to \mathbb{R}$ be a function with $0 \le f(x) \le L\langle x \rangle^a$ for every $x \in \mathbb{R}^d$ and some $L, a \ge 0$. Let P be a subexponential distribution and let $I = \mathbb{E}_P[f(X)]$. Then, for every $b \ge 0$,

$$\mathbb{E}_{P}[f(X)\langle X\rangle^{-b}] \ge CI \log_{+}(1/I)^{-b} \\
\mathbb{E}_{P}[f(X)\langle X\rangle^{b}] \le CI \log_{+}(1/I)^{b}.$$
(69)

Proof. We only prove the first inequality, the second one being proven similarly. Fix a threshold $r \ge 1$ and write

$$\int f(x) \langle x \rangle^{-b} \, \mathrm{d}P(x) \ge \frac{1}{(2r)^b} \int_{\|x\| \le r} f(x) \, \mathrm{d}P(x).$$

According to Lemma 12 with $m = ||P||_{exp}$, if $r \ge r_0$,

$$\int_{\|x\| \ge r} |f(x)| \, \mathrm{d}P(x) \le LC_a r^a e^{-r/m}.$$

Pick $r = m\lambda \log_+(1/I)$. For λ large enough with respect to the different constants involved, the above display is smaller than I/2. Therefore,

$$\int_{\|x\| \le r} |f(x)| \, \mathrm{d}P(x) = I - \int_{\|x\| \ge r} |f(x)| \, \mathrm{d}P(x) \ge I/2,$$

concluding the proof.

B Properties of strongly convex and smooth potentials

We gather in this section the different growth properties of (α, a) -strongly convex and (β, a) smooth potentials (as well as their convex conjugates). We start with a simple lemma that we give without proof, and that follows easily from writing Taylor expansions.

Lemma 14. Let φ be a (β, a) -smooth function for some $\beta, a \ge 0$. Let $x, y \in B(0; r)$ for some $r \ge 0$. Then,

$$\|\nabla\varphi(x) - \nabla\varphi(y)\| \le \beta \langle r \rangle^a \|x - y\| \le 2\beta \langle r \rangle^{a+1},\tag{70}$$

$$|\varphi(x) - \varphi(y)| \le (2\beta \langle r \rangle^{a+1} + \|\nabla\varphi(0)\|)\|x - y\| \le 2\beta' \langle r \rangle^{a+2}, \tag{71}$$

where $\beta' = 2\beta + \|\nabla\varphi(0)\|$. Furthermore, if $\nabla\varphi$ is invertible with $\|\nabla\varphi^{-1}(x)\|, \|\nabla\varphi^{-1}(y)\| \leq t$, then

$$\|\nabla\varphi^{-1}(x) - \nabla\varphi^{-1}(y)\| \ge \frac{\|x - y\|}{\beta} \langle t \rangle^{-a}.$$
(72)

Lemma 15. Let φ be a (α, a) -strongly convex function for some $\alpha, a \geq 0$. Let $x, y \in \mathbb{R}^d$ with $r = \max\{\|x\|, \|y\|\}$. Then, $\nabla \varphi$ is bijection on \mathbb{R}^d , and

$$\|\nabla\varphi(x) - \nabla\varphi(y)\| \ge \frac{\alpha}{2^{a+2}} \langle r \rangle^a \|x - y\|,$$
(73)

$$\|\nabla\varphi^{-1}(x)\| \le 4\alpha^{-\frac{1}{a+1}} (\|x\| + \|\nabla\varphi(0)\|)^{\frac{1}{a+1}},\tag{74}$$

$$\|\nabla\varphi^{-1}(x) - \nabla\varphi^{-1}(y)\| \le \frac{2^{a+2}}{\alpha} \|x - y\|$$
 (75)

Proof. First, note that φ is in particular α -strongly convex of exponent 0 (i.e. strongly convex in the classical sense), so that $\nabla \varphi$ is indeed a bijection. We prove the first inequality. Assume without loss of generality that ||y|| = r. It holds that

$$\|\nabla\varphi(x) - \nabla\varphi(y)\| \ge \int_0^1 \lambda_{\min}(\nabla^2\varphi(tx + (1-t)y))\|x - y\| \,\mathrm{d}t.$$

For $0 \le t \le 1/4$, we have $||tx + (1-t)y|| \ge r - 2tr \ge r/2$. Therefore,

$$\|\nabla\varphi(x) - \nabla\varphi(y)\| \ge \frac{\alpha}{4} \langle \frac{r}{2} \rangle^a \|x - y\| \ge \frac{\alpha}{2^{a+2}} \langle r \rangle^a \|x - y\|$$

The second inequality is obtained by inverting the first one and using that $\langle r \rangle \geq r$. The third one is also obtained by inverting the first one, and using that $\langle r \rangle \geq 1$.

We now turn to bounds on the convex conjugates.

Lemma 16. Let φ be a (α, a) -strongly convex function for some $\alpha, a \ge 0$, with $\varphi(0) = 0$. Then, $\nabla \varphi$ is invertible, and, for $y \in \mathbb{R}^d$,

$$\varphi^*(y) \le \alpha^{-\frac{1}{1+a}} \|\nabla\varphi(0) - y\|^{\frac{2+a}{1+a}}.$$
 (76)

Proof. We write a Taylor expansion around 0. For every $z \in \mathbb{R}^d$,

$$\begin{split} \varphi(z) &\geq \varphi(0) + \langle \nabla \varphi(0), z \rangle + \int_0^1 \lambda_{\min}(\nabla^2 \varphi(tz)) \|z\|^2 t \, \mathrm{d}t \\ &\geq \langle \nabla \varphi(0), z \rangle + \int_0^1 \alpha \langle t \|z\| \rangle^a \|z\|^2 t \, \mathrm{d}t \geq \langle \nabla \varphi(0), z \rangle + \int_0^1 \alpha(t\|z\|)^a \|z\|^2 \, \mathrm{d}t \\ &\geq \langle \nabla \varphi(0), z \rangle + \frac{\alpha}{a+2} \|z\|^{a+2}. \end{split}$$

Therefore,

$$\varphi^*(y) \le \sup_{z \in \mathbb{R}^d} \langle z, y - \nabla \varphi(0) \rangle - \frac{\alpha}{a+2} \|z\|^{a+2}.$$

We conclude thanks to the formula for the convex conjugate of the function $x \mapsto ||x||^{a+2}$. \Box

The following lemma will be useful. It shows that, for points y that stays at a bounded distance from $\nabla \varphi(x)$ the convex conjugate $\varphi^*(y)$ is lower bounded by a quadratic approximation of φ at x.

Lemma 17 (Bounds of convex conjugates: quadratic behavior). Let φ be a (β, a) -smooth function for some $\beta, a \ge 0$ Then, for y satisfying $||y - \nabla \varphi(x)|| \le \ell$,

$$\varphi^*(y) \ge -\varphi(x) + \langle y, x \rangle + \frac{1}{2D_x} \|y - \nabla\varphi(x)\|^2, \tag{77}$$

where $D_x = \beta \langle ||x|| + \frac{\ell}{\beta} \rangle^a$.

Proof. Let $x \in \mathbb{R}^d$ and $z \in \mathbb{R}^d$, with $||x||, ||z|| \leq r$. A Taylor expansion yields

$$\varphi(z) \le \varphi(x) + \langle \nabla \varphi(x), z - x \rangle + \frac{\beta}{2} \langle r \rangle^a ||z - x||^2 = q_x(z).$$
(78)

Let $z = x + \frac{y - \nabla \varphi(x)}{D_x}$, so that we may pick $r = ||x|| + \frac{\ell}{D_x}$. We obtain

$$\varphi^{*}(y) \geq \langle z, y \rangle - \varphi(z) \geq \langle z, y \rangle - q_{x}(z)$$

$$\geq -\varphi(x) + \langle y, x \rangle + \frac{1}{D_{x}} \|y - \nabla \varphi(x)\|^{2} \left(1 - \frac{\beta}{2D_{x}} \langle r \rangle^{a}\right).$$
(79)

A little bit of algebra yields $1 - \frac{\beta}{2D_x} \langle r \rangle^a \ge \frac{1}{2}$, giving the result.

We are now in position to prove Proposition 1

Proof of Proposition 1. We write the proof when the potentials are smooth and defined on \mathbb{R}^d , the case where the support of P is contained in B(0; R) being a straightforward adaptation of the proof of [HR21, Proposition 10]. We use Lemma 14 to get

$$\|\nabla\varphi_1(x) - \nabla\varphi_0(x)\| \le M + 4\beta \langle x \rangle^{a+1}.$$
(80)

We apply Lemma 17 to $\varphi = \varphi_1$, with $y = \nabla \varphi_0(x)$ and ℓ given by the upper bound in (80). Note that $S(\varphi_0) = \int \langle \nabla \varphi_0(x), x \rangle$. Integrating (77) against P yields

$$S(\varphi_1) \ge \int \langle \nabla \varphi_0(x), x \rangle \, \mathrm{d}P(x) + \int \frac{1}{2D_x} \| \nabla \varphi_0(x) - \nabla \varphi_1(x) \|^2 \, \mathrm{d}P(x)$$
$$= S(\varphi_0) + \int \frac{1}{2D_x} \| \nabla \varphi_0(x) - \nabla \varphi_1(x) \|^2 \, \mathrm{d}P(x)$$

Note that D_x grows at most polynomially with ||x||, with degree a(a + 1). The inequality then follows from Lemma 13: we obtain that, for $I = ||\nabla \varphi_0 - \nabla \varphi_1||^2_{L_2(P)}$,

$$I\log_+(1/I)^{-b} \le C\ell. \tag{81}$$

One can invert this inequality to obtain an inequality of the form $I \leq C' \ell \log_+(1/\ell)^b$.

When y is far away from $\nabla \varphi(x)$, as φ grows polynomially (of order 2 + a), we expect the convex conjugate to behave like the convex conjugate of the function $z \mapsto ||z - \nabla \varphi(x)||^a$, which scales like $||y - \nabla \varphi(x)||^{\frac{2+a}{1+a}}$, where $\frac{2+a}{1+a}$ is the conjugate exponent of 2 + a.

Lemma 18 (Lower bound of convex conjugates: polynomial behavior). Under the same assumptions as Lemma 17, if $||y - \nabla \varphi(x)|| \ge \beta \max\{1, ||x||^{a+1}\}$, then

$$\varphi^*(y) \ge -\varphi(x) + \langle y, x \rangle + 2^{-2a-1} \beta^{-\frac{1}{a+1}} \|y - \nabla\varphi(x)\|^{\frac{2+a}{1+a}}.$$
(82)

Proof. Let $v = \|y - \nabla \varphi(x)\|$. We start again from (78). Let $e = \frac{y - \nabla \varphi(x)}{\|y - \nabla \varphi(x)\|}$ and choose $z = x + e \cdot t \cdot (v/\beta)^{\frac{1}{1+a}}$ for some parameter t to fix. Thanks to the condition on v, we may choose the upper bound r on the norm of x and z as $r = (t+1)(v/\beta)^{\frac{1}{1+a}}$. As $v \ge \beta$, it holds that $\langle r \rangle^a \le (t+2)^a (v/\beta)^{\frac{a}{1+a}}$. We obtain

$$\begin{split} \varphi^*(y) &\geq \langle z, y \rangle - q_x(z) \\ &\geq -\varphi(x) + \langle x, \nabla\varphi(x) \rangle + \langle z, y - \nabla\varphi(x) \rangle - \frac{\beta}{2} \langle r \rangle^a t^2 (v/\beta)^{\frac{2}{1+a}} \\ &\geq -\varphi(x) + \langle y, x \rangle + t v^{1+\frac{1}{1+a}} \beta^{-\frac{1}{1+a}} - \frac{\beta}{2} t^2 (2+t)^a (v/\beta)^{\frac{2+a}{1+a}} \\ &\geq -\varphi(x) + \langle y, x \rangle + \beta^{-\frac{1}{1+a}} v^{\frac{2+a}{1+a}} (t - \frac{t^2 (2+t)^a}{2}). \end{split}$$

We let $t = 2^{-2a}$ to obtain that $t - \frac{t^2(2+t)^a}{2} \ge \frac{t}{2}$, concluding the proof.

If one does not care about the tight exponent $\frac{2+a}{1+a}$, it is always possible to lower bound the last term of (82) by a term proportional to $||y - \nabla \varphi(x)||$ (recall that we assume that $||y - \nabla \varphi(x)|| \ge \beta$ in this lemma).

We can also gather Lemma 17 and Lemma 18 together to obtain, with $\ell = \beta \max\{1, \|x\|^{a+1}\}$ and $v = \|y - \nabla \varphi(x)\|$,

$$\varphi^*(y) \ge -\varphi(x) + \langle y, x \rangle + \min\{\frac{1}{2D_x}v^2, C_{a,\beta}v\},\tag{83}$$

where D_x is bounded by a polynomial expression of degree a(a+1) in ||x||.

C Properties of covering and bracketing numbers

We state four simple properties of covering and bracketing numbers that we will repeatedly use.

Lemma 19 (Covering of lines). Let \mathcal{F} be a subset of a normed space $(E, \|\cdot\|)$, bounded by R, and consider the set of lines $\text{Li}(\mathcal{F}) = \{tf + (1-t)g : f, g \in \mathcal{F}, 0 \le t \le 1\}$. Then, there exists a constant c depending on R such that for h > 0,

$$\log \mathcal{N}(h, \mathcal{F}, E) \le 2\log \mathcal{N}(h/3, \mathcal{F}, E) + c\log_+(1/h).$$
(84)

Proof. Let A be a minimal (h/3)-covering of \mathcal{F} and let T be a (h/(6R))-covering of [0, 1]. For $tf + (1-t)g \in \mathcal{F}$, we let f_0 be the closest element to f in the covering A, and define g_0 and t_0 likewise. Then,

$$||tf + (1-t)g - (t_0f_0 + (1-t_0)g_0)|| \le ||f - f_0|| + ||g - g_0|| + 2R|t - t_0| \le h.$$

Therefore, the set $\{t_0f_0 + (1 - t_0)g_0 : f_0, g_0 \in A, t_0 \in T\}$ is a *h*-covering with logarithm of the size at most the right-hand side of (84).

Lemma 20 (Subadditivity of brackets). Let $\mathcal{F}_1, \ldots, \mathcal{F}_k$ be classes of functions and

$$\mathcal{F} \coloneqq \{f_1 + \dots + f_k : f_i \in \mathcal{F}_i \text{ for } i = 1, \dots, k\}.$$

Let h > 0 and let $h_1, \ldots, h_k > 0$ with $h_1 + \cdots + h_k \leq h$. Then, for every metric space E,

$$\log \mathcal{N}_{[]}(h, \mathcal{F}, E) \le \sum_{i=1}^{k} \log \mathcal{N}_{[]}(h_i, \mathcal{F}_i, E)$$
(85)

Proof. A sum of u_i -brackets is a u-bracket.

Lemma 21 (Brackets and translations). Let \mathcal{F} be a class of functions, and consider the class $\tilde{\mathcal{F}} = \{\varphi - H(\varphi) : \varphi \in \mathcal{F}\}$, where $H : \mathcal{F} \to [-M, M]$ is an arbitrary function. Let (E, d) be a metric space such that d(a, b) = |a - b| for any constant functions a, b. Then, there exists a constant c depending on M such that for h > 0,

$$\log \mathcal{N}_{[]}(h, \mathcal{F}, E) \le \log \mathcal{N}_{[]}(h/2, \mathcal{F}, E) + c \log_+(1/h).$$
(86)

Proof. We use the subadditivity of the brackets, so that

$$\log \mathcal{N}_{[]}(h, \tilde{\mathcal{F}}, E) \leq \log \mathcal{N}_{[]}(h/2, \mathcal{F}, E) + \log \mathcal{N}_{[]}(h/2, \{H(\varphi): \varphi \in \mathcal{F}\}, E).$$

The second term is of order $\log_+(1/h)$.

Lemma 22 (Bracketing and dilations). Let $1 \leq p \leq \infty$ and let ρ be a σ -finite measure on \mathbb{R}^d . Let \mathcal{F} be a class of functions with envelope function G and $\|G\|_{L_p(\rho)} \leq \sigma$. Consider a class

$$\tilde{\mathcal{F}} \subseteq \{ x \mapsto a\varphi(x) : \varphi \in \mathcal{F}, \ 0 \le a \le R \},$$
(87)

for some $R \geq 1$. Then, for h > 0,

$$\log \mathcal{N}_{[]}(h, \tilde{\mathcal{F}}, L_p(\rho)) \le \log \mathcal{N}_{[]}(h/(2R), \mathcal{F}, L_p(\rho)) + c \log_+(1/h),$$
(88)

where c depends on σ and R.

Proof. Let h > 0. Let $[f_1, g_1], \ldots, [f_K, g_K]$ be a minimal covering of \mathcal{F} by brackets of size u = h/(2R). Note that we can always pick the functions f_k and g_k so that $|f_k|, |g_k| \leq G$. Let $[a_1, b_1], \ldots, [a_L, b_L]$ be a covering of [0, R] by intervals of length $v = h/(4\sigma)$. The brackets [f, g], where

$$f = a_l(f_k)_+ - b_l(f_k)_-$$
 and $g = b_l(g_k)_+ - a_l(g_k)_-$ (89)

cover $\tilde{\mathcal{F}}$. Furthermore, the size of each bracket is bounded by

$$\|f - g\|_{L_p(\rho)} \le 2\sigma(b_l - a_l) + b_l \|f_k - g_k\|_{L_p(\rho)} < 2\sigma v + Ru = h.$$

The logarithmic of the number of such brackets is $\log K + \log L$, while L scales like 1/h.

D Affine transformations of potentials

In Section 4.1, we claimed that as a particular case of our general theorems, we cover the case where P is the normal distribution and

$$\mathcal{F}_{\text{quad}} = \{ x \mapsto \frac{1}{2} x^{\top} A^{1/2} x + b^{\top} x : A \in \mathbb{S}^d_+, b \in \mathbb{R}^d \}$$
(90)

is the set of quadratics. However, as the set \mathcal{F}_{quad} is not precompact, the covering numbers of \mathcal{F}_{quad} are infinite. Thus, assumption (A2) is not satisfied and Theorem 3 cannot be readily applied. Despite this, all potentials in \mathcal{F}_{quad} can be obtained by scaling and translating potentials from the set

$$\mathcal{F}_{\text{quad},0} = \{ x \mapsto \frac{1}{2} x^{\top} A^{1/2} x : A \in \mathbb{S}^d_+, \|A\|_{\text{op}} = 1 \},\$$

whose bracketing number can easily be controlled, thus satisfying (A2). The next proposition asserts that if one performs such transformations on an arbitrary "base set" \mathcal{F}_0 satisfying (A2), then the fast rates of convergence also holds on the larger class \mathcal{F} that is induced by

scaling and translating. To this end, given a class of potentials \mathcal{F} and $0 \leq r_1, r_2 \leq +\infty$, we introduce the modified class

$$\mathcal{F}_{\mathrm{aff}}^{r_1, r_2} := \{ x \mapsto b\varphi(x) + \langle t, x \rangle : \varphi \in \mathcal{F}, \ 0 \le b \le r_1, \ \|t\| \le r_2 \}$$
(91)

and define the uniform Jensen constant

$$C_J(\mathcal{F}, P) \coloneqq \inf\{\mathbb{E}_P[\varphi(X)] - \varphi(\mathbb{E}_P[X]) : \varphi \in \mathcal{F}\}.$$
(92)

The constant is nonnegative if all the functions in \mathcal{F} are convex, and is positive as long as every potential $\varphi \in \mathcal{F}$ is α -strongly convex and P is not degenerate, with $C_J(\mathcal{F}, P) \geq \alpha \operatorname{Var}_P(X)$.

Proposition 7. Let P be a subexponential distribution and let $Q = (\nabla \varphi_0)_{\sharp} P$ for some (β, a) smooth convex potential φ_0 with $\beta, a \ge 0$. Let \mathcal{F} be a class of potentials satisfying **(A1)** and $C_J(\mathcal{F}, P) > -\infty$, with $\nabla \varphi(0) = 0$ for every $\varphi \in \mathcal{F}$. Let $\tilde{\mathcal{F}} \subseteq \mathcal{F}_{aff}^{r_1, +\infty}$. Assume that either $r_1 < +\infty$ or that $C_J(\mathcal{F}, P) > 0$. Further assume that there exists an α -strongly convex
potential in \mathcal{F} . Then, there exists r > 0 such that

$$\mathbb{E}[\|\nabla \hat{\varphi}_{\tilde{\mathcal{F}}} - \nabla \varphi_0\|_{L_2(P)}^2 \mathbf{1}\{\hat{\varphi}_{\tilde{\mathcal{F}}} \neq \hat{\varphi}_{\tilde{\mathcal{F}}^r}\}] \le c \exp\left(-Cn\right),\tag{93}$$

where $\tilde{\mathcal{F}}^r = \tilde{\mathcal{F}} \cap \mathcal{F}_{aff}^{r,r}$, with constants r, c and C depending on the different parameters involved, and on P through $||P||_{exp}$.

Proof. Let us write the sample $Y_1, \ldots, Y_n \sim Q$ as $Y_i = \nabla \varphi_0(X_i)$, where $X_i \sim P$. We can write $\hat{\varphi}_{\tilde{\mathcal{F}}}(x) = b\varphi(x) + \langle t, x \rangle$ for $x \in \mathbb{R}^d$, where $\varphi \in \mathcal{F}$, and b and t are two parameters. Let $\overline{\varphi}$ be an α -strongly convex potential in \mathcal{F} . Note that $S_n(\hat{\varphi}_{\tilde{\mathcal{F}}}) - S_n(\overline{\varphi}) \leq 0$. We have for $y \in \mathbb{R}^d$

$$\hat{\varphi}_{\tilde{\mathcal{F}}}^*(y) = b\varphi^*\left(\frac{y-t}{b}\right)$$

so that

$$S_n(\hat{\varphi}_{\tilde{\mathcal{F}}}) = P(\hat{\varphi}_{\tilde{\mathcal{F}}}) + Q_n(\hat{\varphi}_{\tilde{\mathcal{F}}}^*) = b\mathbb{E}_P[\varphi(X)] + \langle t, \mathbb{E}_P[X] \rangle + \frac{b}{n} \sum_{i=1}^n \varphi^*\left(\frac{\nabla \varphi_0(X_i) - t}{b}\right)$$

Our goal is to show that the condition $S_n(\hat{\varphi}_{\tilde{\mathcal{F}}}) - S_n(\overline{\varphi}) \leq 0$ implies that both b and ||t|| are upper bounded by some parameter r, which implies that $\hat{\varphi}_{\tilde{\mathcal{F}}} \in \tilde{\mathcal{F}}^r$. In particular, we then have $\hat{\varphi}_{\tilde{\mathcal{F}}} = \hat{\varphi}_{\tilde{\mathcal{F}}^r}$.

Bound on *b*: By definition of the convex conjugate, it holds that

$$b\varphi^*\left(\frac{\nabla\varphi_0(X_i)-t}{b}\right) \ge -b\varphi(\mathbb{E}_P[X]) + \langle \mathbb{E}_P[X], \nabla\varphi_0(X_i)-t\rangle.$$
(94)

We obtain

$$S_n(\hat{\varphi}_{\tilde{\mathcal{F}}}) \ge b(\mathbb{E}_P[\varphi(X)] - \varphi(\mathbb{E}_P[X])) + \frac{1}{n} \sum_{i=1}^n \langle \mathbb{E}_P[X], \nabla \varphi_0(X_i) \rangle$$

Therefore,

$$S_{n}(\hat{\varphi}_{\tilde{\mathcal{F}}}) - S_{n}(\overline{\varphi}) = P(\hat{\varphi}_{\tilde{\mathcal{F}}}) + Q_{n}(\hat{\varphi}_{\tilde{\mathcal{F}}}^{*}) - P(\overline{\varphi}) - Q_{n}(\overline{\varphi}^{*})$$

$$\geq bC_{J}(\mathcal{F}, P) - A_{n}, \qquad (95)$$

where $A_n = P(\overline{\varphi}) + Q_n(\overline{\varphi}^*) - \langle \mathbb{E}_P[X], P_n(\nabla \varphi_0) \rangle$. Assume that $C_J(\mathcal{F}, P) > 0$. Then, as $S_n(\hat{\varphi}_{\tilde{\mathcal{F}}}) - S_n(\overline{\varphi}) \leq 0$, it holds that

$$0 \le b \le b_{\max} \coloneqq \max\{1, \frac{A_n}{C_J(\mathcal{F}, P)}\}.$$
(96)

If $r_1 < +\infty$, we instead define $b_{\max} = \max\{1, r_1\}$, so that the inequality $b \le b_{\max}$ also holds. Note also that we have $b_{\max} \ge 1$, a property which will be used.

Bound on ||t||: Finding a bound on the norm of the parameter t proves to be more delicate, and relies on lower bounds on the convex conjugate given in Appendix B. Apply (83) to $x = \mathbb{E}_P[X]$ and $y = \frac{\nabla \varphi_0(X_i) - t}{b}$. The parameter D_x appearing in (83) can be upper bounded by a constant depending on β and the norm of $\mathbb{E}_P[X]$ (which is in turn bounded in term of $||P||_{\exp}$). In total, we obtain a lower bound of the form

$$b\varphi^*\left(\frac{\nabla\varphi_0(X_i)-t}{b}\right) \ge -b\varphi(\mathbb{E}_P[X]) + \langle \mathbb{E}_P[X], \nabla\varphi_0(X_i)-t \rangle + C\min\{v_i^2, v_i\},$$

where $v_i = \|\frac{\nabla \varphi_0(X_i) - t}{b} - z\|$, and we define $z = \nabla \varphi(\mathbb{E}_P[X])$. Assuming without loss of generality that $\|t\| \ge 1$, we lower bound v_i^2 ,

$$v_i^2 \ge \frac{\|t\|^2 - 2\|t\|(\|\nabla\varphi_0(X_i)\| + b_{\max}\|z\|)}{b_{\max}^2} \ge \frac{\|t\| - 2\|\nabla\varphi_0(X_i)\| - 2b_{\max}\|z\|}{b_{\max}^2} \eqqcolon U_i,$$

as well as v_i ,

$$v_i \ge \frac{\|t\| - \|\nabla \varphi_0(X_i)\| - b_{\max}\|z\|}{b_{\max}} \ge U_i,$$

where we use that $b_{\text{max}} \geq 1$. As for (95), we obtain

$$S_{n}(\hat{\varphi}_{\tilde{\mathcal{F}}}) - S_{n}(\overline{\varphi}) \geq bC_{J}(\mathcal{F}, P) - A_{n} + \frac{C}{n} \sum_{i=1}^{n} U_{i}$$

$$\geq -b_{\max}|C_{J}(\mathcal{F}, P)| - A_{n} + C\frac{\|t\|}{b_{\max}^{2}} - 2C\frac{\|z\|}{b_{\max}} - 2C\frac{P_{n}(\|\nabla\varphi_{0}\|)}{b_{\max}^{2}}.$$
(97)

Therefore,

$$C||t|| \le 2CP_n(||\nabla\varphi_0||) + 2Cb_{\max}||z|| + b_{\max}^2 A_n + b_{\max}^3 |C_J(\mathcal{F}, P)|.$$

As $b_{\max}^2 A_n \leq \frac{1}{2}(b_{\max}^4 + A_n^2)$, ||t|| is bounded by a polynomial expression T_{\max} of degree 4 in the variables $P_n(||\nabla \varphi_0||)$ and $Q_n(\overline{\varphi}^*)$.

Conclusion: Let $r \ge 1$. One can make the key observation that if $\hat{\varphi}_{\tilde{\mathcal{F}}} \neq \hat{\varphi}_{\tilde{\mathcal{F}}^r}$, then either $b_{\max} > r$ or $T_{\max} > r$. Therefore,

$$\mathbb{E}[\|\nabla \hat{\varphi}_{\tilde{\mathcal{F}}} - \nabla \varphi_0\|_{L_2(P)}^2 \mathbf{1}\{\hat{\varphi}_{\tilde{\mathcal{F}}} \neq \hat{\varphi}_{\tilde{\mathcal{F}}^r}\}] \leq 2\mathbb{E}[\|\nabla \hat{\varphi}_{\tilde{\mathcal{F}}}\|_{L^2(P)}^2 \mathbf{1}\{\hat{\varphi}_{\tilde{\mathcal{F}}} \neq \hat{\varphi}_{\tilde{\mathcal{F}}^r}\}] \\ + 2\|\nabla \varphi_0\|_{L_2(P)}^2 \mathbb{P}(\hat{\varphi}_{\tilde{\mathcal{F}}} \neq \hat{\varphi}_{\tilde{\mathcal{F}}^r}).$$

By Lemma 14, it holds that

$$\|\nabla \hat{\varphi}_{\tilde{\mathcal{F}}}(x)\| \le b_{\max} \|\nabla \hat{\varphi}(x)\| + T_{\max} \|x\| \le (b_{\max}\beta \langle x \rangle^a + T_{\max}) \|x\|$$

Therefore,

$$\mathbb{E}[\|\nabla\hat{\varphi}_{\tilde{\mathcal{F}}} - \nabla\varphi_0\|_{L_2(P)}^2 \mathbf{1}\{\hat{\varphi}_{\tilde{\mathcal{F}}} \neq \hat{\varphi}_{\tilde{\mathcal{F}}^r}\}] \le c\mathbb{E}[b_{\max}^2 \mathbf{1}\{b_{\max} > r\}] + c\mathbb{E}[T_{\max}^2 \mathbf{1}\{T_{\max} > r\}]$$

for some constant c depending on $\overline{\varphi}$, $\|P\|_{\exp}$, β and a.

Both b_{\max} and T_{\max} are bounded by polynomial expressions in $P_n(\|\nabla\varphi_0\|)$ and $Q_n(\overline{\varphi}^*)$. As $\overline{\varphi}$ is strongly convex, $\overline{\varphi}^*(\nabla\varphi_0(X_i))$ grows at most quadratically by Lemma 16. In any case, both $\|\nabla\varphi_0(X_i)\|$ and $\overline{\varphi}^*(\nabla\varphi_0(X_i))$ grow polynomially with respect to $\|X_i\|$. We conclude thanks to the following lemma.

Lemma 23. Let P be a subexponential distribution and let \overline{X}_n be the sample mean of n i.i.d. observations of law P. Then, there exists $r_0 > 0$ such that for $a \ge 1$ and $r \ge r_0$,

$$\mathbb{E}[\overline{X}_n^a \mathbf{1}\{|\overline{X}_n| > r\}] \le c \exp\left(-Cn\right).$$
(98)

Proof. We choose $r \ge r_0 = (2|\mathbb{E}_P[X]|)^a$, so that

$$\mathbb{E}[|\overline{X}_n|^a \mathbf{1}\{|\overline{X}_n| > r\}] = \int_r^{+\infty} \mathbb{P}\left\{|\overline{X}_n|^a > t\right\} \, \mathrm{d}t \le \int_r^{+\infty} \mathbb{P}\left\{|\overline{X}_n - \mathbb{E}_P[X]| > t^{1/a}/2\right\} \, \mathrm{d}t.$$

We then use Bernstein's inequality to conclude, see e.g. [Ver18, Corollary 2.8.3]. \Box

As a consequence of Proposition 7, we can show that Theorem 3 holds as well for affine transformations of a base set \mathcal{F} .

Corollary 2. Consider a probability distribution P and a class of potentials \mathcal{F} with $C_J(\mathcal{F}, P) > -\infty$. Let $\tilde{\mathcal{F}} \subseteq \mathcal{F}_{aff}^{r_1,+\infty}$, and further assume that φ_0 , P and \mathcal{F} satisfy the assumptions of Theorem 3.1 (resp. Theorem 3.2). Assume that either $r_1 < +\infty$ or that $C_J(\mathcal{F}, P) > 0$. Then,

$$\mathbb{E} \|\nabla \hat{\varphi}_{\tilde{\mathcal{F}}} - \nabla \varphi_0\|_{L^2(P)}^2 \lesssim v_n, \tag{99}$$

where v_n is the rate appearing in of Theorem 3.1 (resp. Theorem 3.2).

Proof. Let $\mathcal{F}_0 = \{x \mapsto \varphi(x) - \langle \nabla \varphi(0), x \rangle : \varphi \in \mathcal{F}\}$. Then, $\tilde{\mathcal{F}} \subseteq (\mathcal{F}_0)_{\text{aff}}^{1,+\infty}$ and $C_J(\mathcal{F}, P) = C_J(\mathcal{F}_0, P) > -\infty$. Note that thanks to the assumptions of Theorem 3, \mathcal{F} contains a strongly convex potential. Proposition 7 implies that for some r > 0,

$$\mathbb{E}[\|\nabla\hat{\varphi}_{\tilde{\mathcal{F}}} - \nabla\varphi_0\|_{L_2(P)}^2 \mathbf{1}\{\hat{\varphi}_{\tilde{\mathcal{F}}} \neq \hat{\varphi}_{\tilde{\mathcal{F}}^r}\}] \le \frac{1}{n}.$$
(100)

To conclude, it suffices to check that the set $\tilde{\mathcal{F}}^r$ satisfy the assumptions of Theorem 2 (resp. Theorem 3). Assumption (A2) is satisfied for $\tilde{\mathcal{F}}^r$ by Lemma 22 and Lemma 21. It is easy to check that if \mathcal{F} satisfies one of the other assumptions, then $\tilde{\mathcal{F}}^r$ will also satisfy it. \Box

Let us come back to the Gaussian case. Recall that we introduced the "base" set

$$\mathcal{F}_{\text{quad},0} = \{ x \mapsto \frac{1}{2} x^{\top} A^{1/2} x : A \in \mathbb{S}^d_+, \|A\|_{\text{op}} = 1 \},\$$

whereas $(\mathcal{F}_{quad,0})_{aff}^{+\infty,+\infty} = \mathcal{F}_{quad}$. We are therefore in position to apply Corollary 2. Note that every $\varphi \in \mathcal{F}_{quad,0}$ is 1-smooth and convex, while $P = N(0, I_d)$ satisfies both the Poincaré inequality (A3) and condition (C2). Although not all potentials in $\mathcal{F}_{quad,0}$ are strongly convex, the uniform Jensen constant $C_J(\mathcal{F}_{quad,0}, P)$ is at least 1/2:

$$\mathbb{E}_P[\varphi_A(X)] - \varphi_A(\mathbb{E}_P[X]) = \frac{1}{2} \operatorname{Tr}(A) \ge \frac{1}{2} > 0,$$

since $||A||_{\text{op}} = 1$. Assumption (A2) is verified in Lemma 24, with log-covering numbers scaling logarithmically. All in all, with the absence of a bias term, Corollary 2 implies that the estimator $\nabla \hat{\varphi}_{\mathcal{F}_{\text{rund}}}$ attains the rate n^{-1} .

Lemma 24. Let $\delta > 0$. It holds that $\log \mathcal{N}(h, \mathcal{F}_{quad,0}, L_{\infty}(e^{-\delta |\cdot||})) \leq c \log_{+}(1/h)$.

Proof. If $\Sigma_1, \Sigma_2 \in \mathbb{S}^d_+$, then $\sup_x x^\top (\Sigma_2 - \Sigma_1) x \cdot e^{-\delta \|x\|} \leq C_{\delta} \|\Sigma_2 - \Sigma_1\|_{\text{op.}}$ As the set of matrices is finite dimensional, the covering number scales polynomially with h. \Box

E Lemmas: bounded case

Proof of Lemma 1. Let us first bound $\varphi \in \mathcal{F}$. Let $x \in B(0; R)$. Then

$$|\varphi(x)| \le \int_0^1 |\langle \nabla \varphi(tx), x \rangle| \, \mathrm{d}t \le R^2.$$

Remark that Q is supported in B(0; R). Let $y \in B(0; R)$. We have

$$\varphi^*(y) = \sup_{\|x\| \le 2R} \langle x, y \rangle - \varphi(x) \le 2R^2 + R^2 = 3R^2.$$
(101)

Also, choosing x = 0, we obtain $\varphi^*(y) \ge 0$. Therefore, $|\varphi^*(y) - \overline{\varphi}^*(y)| \le 3R^2$. Furthermore, $|P\varphi - P\overline{\varphi}| \le 2R^2$. Therefore, $5R^2$ is an envelope of \mathcal{G} .

Proof of Lemma 2. Let φ be any potential in \mathcal{F} . As $Q = (\nabla \varphi_0)_{\sharp} P$, we may write y in the support of Q as $y = \nabla \varphi_0(x)$ for x in the support of P. Therefore, $\varphi_0^*(y) = \langle y, x \rangle - \varphi_0(x)$. Let x' be a point attaining the supremum in the definition of $\varphi^*(y)$. Note that $||x'|| \leq 2R$ and that $\nabla \varphi(x') = y$. Then,

$$\varphi_c^*(y) - \varphi_{0,c}^*(y) = \langle y, x' - x \rangle - \varphi_c(x') + \varphi_{0,c}(x)$$
$$= \langle y, x' - x \rangle + (\varphi_c(x) - \varphi_c(x')) + (\varphi_{0,c}(x) - \varphi_c(x)).$$

As φ_c is convex, it holds that $\varphi_c(x') \ge \varphi_c(x) + \langle \nabla \varphi(x), x' - x \rangle$, so that

$$\varphi_c^*(y) - \varphi_{0,c}^*(y) \le \langle \nabla \varphi_0(x) - \nabla \varphi(x), x' - x \rangle + (\varphi_{0,c}(x) - \varphi_c(x)).$$

Likewise, $\varphi_c(x) \ge \varphi_c(x') + \langle y, x - x' \rangle$. Therefore,

$$\varphi_c^*(y) - \varphi_{0,c}^*(y) \ge \varphi_{0,c}(x) - \varphi_c(x).$$

In total, as $||x - x'|| \le R + 2R \le 3R$,

$$|\varphi_{c}^{*}(y) - \varphi_{0,c}^{*}(y)| \leq 3R \|\nabla\varphi_{0}(x) - \nabla\varphi(x)\| + |\varphi_{0,c}(x) - \varphi_{c}(x)|.$$
(102)

By the Poincaré inequality, $\|\varphi_{0,c} - \varphi_c\|_{L_2(P)} \leq C_{PI}^{1/2} \|\nabla \varphi_0 - \nabla \varphi\|_{L_2(P)}$. In total, we obtain that

$$\|\varphi_{c}^{*} - \varphi_{0,c}^{*}\|_{L_{2}(Q)} \leq (3R + C_{\mathtt{PI}}^{1/2}) \|\nabla\varphi_{0} - \nabla\varphi\|_{L_{2}(P)}.$$
(103)

Now, let $g = \varphi_c^* - \overline{\varphi}_c^* \in \mathcal{G}$. Then, applying the stability property (Proposition 1) leads to

$$\begin{aligned} \|g\|_{L_{2}(Q)} &\leq \|\varphi_{c}^{*} - \varphi_{0,c}^{*}\|_{L_{2}(Q)} + \|\overline{\varphi}_{c}^{*} - \varphi_{0,c}^{*}\|_{L_{2}(Q)} \\ &\leq (3R + C_{\mathrm{PI}}^{1/2})(\|\nabla\varphi_{0} - \nabla\varphi\|_{L_{2}(P)} + \|\nabla\varphi_{0} - \nabla\overline{\varphi}\|_{L_{2}(P)}) \\ &\leq (3R + C_{\mathrm{PI}}^{1/2})(\|\nabla\overline{\varphi} - \nabla\varphi\|_{L_{2}(P)} + 2\|\nabla\varphi_{0} - \nabla\overline{\varphi}\|_{L_{2}(P)}) \\ &\leq (3R + C_{\mathrm{PI}}^{1/2})(\tau + 2\sqrt{C_{0}\ell}), \end{aligned}$$

As $\ell \leq \tau^2$ by assumption, we obtain the conclusion.

Proof of Lemma 3. First, as $|P(\varphi)| \leq R^2$ for any $\varphi \in \mathcal{F}$, it holds by Lemma 21 that for h > 0,

$$\log \mathcal{N}_{[]}(h,\mathcal{G},L_2(Q)) \le \log \mathcal{N}_{[]}(h/2,\tilde{\mathcal{G}},L_2(Q)) + c \log_+(1/h),$$
(104)

where $\tilde{\mathcal{G}} = \{\varphi^* - \overline{\varphi}^* : \varphi \in \operatorname{Li}(\mathcal{F})_{\tau}\}$. Furthermore, The application $\varphi \mapsto \varphi^*$ is order-reversing and is an isometry for the ∞ -norm. Remark also that the bracketing number is unchanged when all the functions in a class are shifted by the same function. Those two remarks imply that

$$\mathcal{N}_{[]}(h,\mathcal{G},L_2(Q)) \le \mathcal{N}_{[]}(h,\mathcal{G},L_\infty(\rho)) \le \mathcal{N}_{[]}(h,\operatorname{Li}(\mathcal{F})_{\tau},L_\infty(\rho)),$$
(105)

where ρ is the Lebesgue measure on B(0; 2R). The bracketing numbers for the L_{∞} -norm are controlled by the covering numbers for the same norm, see [VW96, Theorem 2.7.11]. Threfore, as, for any function f, it holds that $||f||_{L_{\infty}(\rho)} \leq ||f||_{L_{\infty}(e^{-||\cdot||/(2R)})}e^{1}$, we have the conclusion using Lemma 19.

F Lemmas: strongly convex case

Our first lemma states that we may without loss of generality assume that every potential φ in \mathcal{F} satisfies $\|\nabla \varphi(0)\| \leq r$ for some constant r.

Lemma 25. Let $\mathcal{F}^r = \{ \varphi \in \mathcal{F} : \|\nabla \varphi(0)\| \leq r \}$. Then, for r large enough,

$$\mathbb{E}[\|\nabla \hat{\varphi}_{\mathcal{F}} - \nabla \varphi_0\|_{L_2(P)}^2 \mathbf{1}\{\hat{\varphi}_{\mathcal{F}} \neq \hat{\varphi}_{\mathcal{F}^r}\}] \le \frac{1}{n}.$$
(106)

Proof. Let $\mathcal{F}_0 = \{x \mapsto \varphi(x) - \langle \nabla \varphi(0), x \rangle : \varphi \in \mathcal{F}\}$. Then, $\mathcal{F} \subseteq (\mathcal{F}_0)^{1,+\infty}_{aff}$. Condition (A1) and Lemma 14 imply that

$$C_J(\mathcal{F}_0, P) \ge -2\beta \int \langle x \rangle^{a+2} \, \mathrm{d}P(x) > -\infty.$$
(107)

We may therefore apply Proposition 7, which implies that there exists r > 0 such that the statement of the lemma holds.

This preliminary remark shows that we can safely assume that $\|\nabla\varphi(0)\|$ is uniformly bounded over $\varphi \in \mathcal{F}$: we will now simply write \mathcal{F} instead of \mathcal{F}^r and keep this assumption in mind. To ease notation, we will now write T instead of $\nabla\varphi$ for a general potential $\varphi \in \mathcal{F}$, with $T_0 = \nabla\varphi_0$ and $\overline{T} = \nabla\overline{\varphi}$.

Proof of Lemma 5. As φ is $(\alpha/2, a)$ -strongly convex, we have by Lemma 16 that $\varphi^*(y) \leq (\alpha/2)^{-\frac{1}{1+a}} ||T(0) - y||^{\frac{2+a}{1+a}} \leq C(r^2 + ||y||^2)$, where we used the fact that $||T(0)|| \leq r$. As $\overline{\varphi}$ is also $(\alpha/2)$ -strongly convex, the same inequality holds for $\overline{\varphi}$ (where we pick r larger than $||\overline{T}(0)||$). Together, with the Poincaré inequality for P, this yields,

$$\begin{aligned} |\varphi_c^*(y) - \overline{\varphi}_c^*(y)| &\leq 2C(r^2 + \|y\|^2) + |\mathbb{E}[\overline{\varphi}(X) - \varphi(X)]| \leq 2C(r^2 + \|y\|^2) + \|\overline{\varphi} - \varphi\|_{L^2(P)} \\ &\leq 2C(r^2 + \|y\|^2) + \tau \sqrt{C_{\text{PI}}} \,. \end{aligned}$$

Proof of Lemma 6. Let $g = \varphi_c^* - \overline{\varphi}_c^* \in \mathcal{G}$. By definition of a convex conjugate, it holds that

$$\varphi_c^*(y) = \langle y, T^{-1}(y) \rangle - \varphi(T^{-1}(y)) + P(\varphi) = \langle y, T^{-1}(y) \rangle - \varphi_c(T^{-1}(y))$$

Omitting the argument y for notational brevity, we obtain

$$\begin{split} \varphi_c^* - \varphi_{0,c}^* &= \langle \mathrm{id}, T^{-1} - T_0^{-1} \rangle + \varphi_{0,c} \circ T_0^{-1} - \varphi_c \circ T^{-1} \\ &= \langle \mathrm{id}, T^{-1} - T_0^{-1} \rangle + (\varphi_{0,c} \circ T_0^{-1} - \varphi_c \circ T_0^{-1}) + (\varphi_c \circ T_0^{-1} - \varphi_c \circ T^{-1}) \\ &= A_1 + A_2 + A_3. \end{split}$$

We bound the norms of each of the three terms separately:

• According to Lemma 15, $||T^{-1}(y) - T^{-1}(y')|| \le \frac{2^{a+3}}{\alpha} ||y' - y||$ for every y, y'. Therefore

$$\|A_1\|_{L_2(Q)}^2 = \int \langle \mathrm{id}, T^{-1} - T_0^{-1} \rangle^2 \, \mathrm{d}Q = \int \langle T_0, T^{-1} \circ T_0 - \mathrm{id} \rangle^2 \, \mathrm{d}P$$

$$\leq c \int \|T_0\|^2 \|T_0 - T\|^2 \, \mathrm{d}P \leq C \int \|T_0(x) - T(x)\|^2 \langle x \rangle^{2a+2} \, \mathrm{d}P(x),$$

where we use Lemma 14 at the last line. Let $I = \int ||T_0(x) - T(x)||^2 dP(x)$. We may apply Lemma 13 to obtain $||A_1||_{L_2(Q)}^2 \leq CI \log_+(1/I)^{2a+2}$.

• By the Poincaré inequality, $||A_2||_{L_2(Q)} = ||\varphi_{0,c} \circ T_0^{-1} - \varphi_c \circ T_0^{-1}||_{L_2(Q)} = ||\varphi_{0,c} - \varphi_c||_{L_2(P)} \le \sqrt{C_{\mathtt{PI}}} ||T - T_0||_{L_2(P)}.$

• By Lemma 14, it holds that

$$\begin{split} \|A_3\|_{L_2(Q)}^2 &= \int \|\varphi \circ T_0^{-1} - \varphi \circ T^{-1}\|^2 \,\mathrm{d}Q \le c \int \langle \|T_0^{-1}\| + \|T^{-1}\| \rangle^{2a+2} \|T_0^{-1} - T^{-1}\|^2 \,\mathrm{d}Q \\ &\le c \int \langle \|\operatorname{id}\| + \|T^{-1} \circ T_0\| \rangle^{2a+2} \|\operatorname{id} - T^{-1} \circ T_0\|^2 \,\mathrm{d}P \\ &\le C \int \langle x \rangle^{2a+2} \|T(x) - T_0(x)\|^2 \,\mathrm{d}P(x), \end{split}$$

where we use Lemma 15 at the last line. As for the bound on A_1 , we can use Lemma 13 to obtain that this term is controlled by $CI \log_+(1/I)^{2a+2}$.

In total, $\|\varphi_c^* - \varphi_{0,c}^*\|_{L_2(Q)}^2 \leq C \log_+(1/I)^{2a+2}I$ for some constant C. By Proposition 1, the integral I is bounded by

$$2\|T - \overline{T}\|_{L_2(P)}^2 + 2\|T_0 - \overline{T}\|_{L_2(P)}^2 \le 2\tau^2 + C\ell \log_+(1/\ell)^{a(a+1)} \le C\tau^2 \log_+(1/\tau)^{a(a+1)},$$

where we use that $\tau^2 \ge \ell$. We obtain

$$\|\varphi_c^* - \varphi_{0,c}^*\|_{L_2(Q)}^2 \le C\tau^2 \log_+(1/\tau)^{a^2 + 3a + 2}.$$

The same inequality holds for $\overline{\varphi}$ (as $\overline{\varphi}$ is also strongly convex and in $\operatorname{Li}(\mathcal{F})_{\tau}$), so that we get our final bound on $\|\varphi_c^* - \overline{\varphi}_c^*\|_{L_2(Q)}$ by using the triangle inequality.

Eventually, the bracketing numbers are controlled thanks to the next lemma.

Lemma 26. Let φ_1, φ_2 be two functions that are (α, a) -strongly convex and (β, a) -smooth, with $\varphi_i(0) = 0$ and $\|\nabla \varphi_i(0)\| \leq r$ (for i = 1, 2). Let $x \in \mathbb{R}^d$. Then, for every $\delta > 0$,

$$|\varphi_1^*(T_0(x)) - \varphi_2^*(T_0(x))| \le C_{\delta} \|\varphi_1 - \varphi_2\|_{L_{\infty}(e^{-\delta}\|\cdot\|)} e^{C\delta\|x\|}.$$
(108)

Proof. Let $y \in \mathbb{R}^d$. It holds that $\varphi_1^*(y) = \langle x_1, y \rangle - \varphi_1(x_1)$ for $x_1 = T_1^{-1}(y)$. Therefore,

$$\varphi_1^*(y) - \varphi_2^*(y) = (\langle x_1, y \rangle - \varphi_1(x_1)) - \varphi_2^*(y) \le (\langle x_1, y \rangle - \varphi_1(x_1)) - (\langle x_1, y \rangle - \varphi_2(x_1))$$
$$\le |\varphi_1(x_1) - \varphi_2(x_1)|.$$

By symmetry, it holds that $|\varphi_1^*(y) - \varphi_2^*(y)| \le |\varphi_1(x_1) - \varphi_2(x_1)| + |\varphi_1(x_2) - \varphi_2(x_2)|$, with $x_2 = T_2^{-1}(y)$. Let $y = T_0(x)$. Then,

$$\begin{aligned} |\varphi_1(x_1) - \varphi_2(x_1)| &\leq |\varphi_1(T_1^{-1} \circ T_0(x)) - \varphi_2(T_1^{-1} \circ T_0(x))|e^{-\delta \|T_1^{-1} \circ T_0(x)\|} e^{\delta \|T_1^{-1} \circ T_0(x)\|} \\ &\leq \|\varphi_1 - \varphi_2\|_{L_{\infty}(e^{-\delta \|\cdot\|})} e^{\delta \|T_1^{-1} \circ T_0(x)\|}. \end{aligned}$$

By Lemma 15, $||T_1^{-1}(T_0(x))|| \leq 4\alpha^{-\frac{1}{a+1}}(||T_0(x)|| + r)^{\frac{1}{a+1}}$. By Lemma 14, $||T_0(x)|| \leq r + 2\beta\langle x\rangle^{a+1}$. In total, we obtain that $||T_1^{-1}(T_0(x))|| \leq C\langle x\rangle$ for a large enough constant C. As a similar inequality holds for x_2 , we obtain the conclusion.

Proof of Lemma 7. Lemma 14 implies that the potentials φ in \mathcal{F} are such that $|P(\varphi)| \leq M$ for some constant M. Therefore, by Lemma 21, we have for h > 0

$$\log \mathcal{N}_{[]}(h,\mathcal{G},L_2(Q)) \le \log \mathcal{N}_{[]}(h/2,\mathcal{G},L_2(Q)) + c\log_+(1/h), \tag{109}$$

where $\tilde{\mathcal{G}} = \{\varphi^* - \overline{\varphi}^* : \varphi \in \operatorname{Li}(\mathcal{F})_{\tau}, \varphi \text{ is } (\alpha/2)\text{-strongly convex}\}$. The conclusion of Lemma 7 is then obtained through Lemma 26. Indeed, for $\varphi_1, \varphi_2 \in \operatorname{Li}(\mathcal{F})_{\tau}$ that are $(\alpha/2)$ -strongly convex,

$$\int |\varphi_1^*(y) - \varphi_2^*(y)|^2 \, \mathrm{d}Q(y) = \int |\varphi_1^*(T_0(x)) - \varphi_2^*(T_0(x))|^2 \, \mathrm{d}P(x)$$
$$\leq \|\varphi_1 - \varphi_2\|_{L_{\infty}(e^{-\delta \|\cdot\|})}^2 \int e^{2C\delta \|x\|} \, \mathrm{d}P(x).$$

As bracketing numbers are controlled by covering numbers for the L_{∞} -norm (see [VW96, Theorem 2.7.11]), we may choose δ small enough so that the latter integral is finite, and then conclude using Lemma 19.

Proof of Lemma 8. For R > 0, let $g_{>R} = g - g_{\leq R}$. Let G be the envelope function given by Lemma 5. Then,

$$\mathbb{E}[|\sup_{g \in \mathcal{G}} (Q - Q_n)(g_{>R})| \le 2Q(G_{>R}) = 2 \int_{\|\nabla \varphi_0(x)\| > R} G(\nabla \varphi_0(x)) \, \mathrm{d}P(x).$$
(110)

If R is large enough and $\|\nabla \varphi_0(x)\| > R$, then $\|x\| > cR^{1/(a+1)}$ according to Lemma 14. By Lemma 12, as P is subexponential (it satisfies the Poincaré inequality), the right-hand side of Equation (110) is smaller than n^{-2} if $R = \kappa (\log n)^{a+1}$ for some κ large enough.

The probability that there exists one sample point not in B(0; R) is bounded by $nQ(||Y|| > R) \le n^{-2}$ for κ large enough. If this event is satisfied, then $Q_n^R = Q_n$, so that

$$\begin{split} \mathbb{E}[|\sup_{g \in \mathcal{G}} (Q - Q_n)(g_{\leq R})|] &\leq \mathbb{E}[|\sup_{g \in \mathcal{G}} (Q^R - Q_n^R)(g_{\leq R})|] + \mathbb{E}[|\sup_{g \in \mathcal{G}} ((Q - Q^R) - (Q_n - Q_n^R))(g_{\leq R})|] \\ &\leq \mathbb{E}[|\sup_{g \in \mathcal{G}} (Q^R - Q_n^R)(g_{\leq R})|] + (\sup_{B(0;R)} G)(n^{-2} + \frac{1}{Q(B(0;R))} - 1) \\ &\leq \mathbb{E}[|\sup_{g \in \mathcal{G}} (Q^R - Q_n^R)(g_{\leq R})|] + n^{-1}, \end{split}$$

where we use that $(\sup_{B(0;R)} G) \leq C(1 + \log n)^{2a+2}$ according to Lemma 5, and that $1 - Q(B(0;R)) = \int_{\|\nabla \varphi_0(x)\| > R} dP(x) \leq n^{-2}$.

The probability distribution Q is obtained as the pushforward of P by the map $T_0 = \nabla \varphi_0$, where φ_0 is (α, a) -strongly convex and (β, a) -smooth. Let Ω^R be the support of Q^R . By Lemma 15, every $y \in \Omega^R$ is written as $T_0(x)$ for some $x \in B(0; C_0 R^{1/(a+1)})$ for some constant C_0 . By definition of smoothness, on such a ball, the map T_0 is a Lipschitz diffeomorphism with operator norm of the derivative bounded by $C_1 R^{a/(a+1)} \leq C_1 R$. Furthermore, by definition of strong convexity, the operator norm of the differential of the inverse map T_0^{-1} at $y \in B(0; R)$ is bounded by C_2 . The set Ω^R is partitioned into images the $T_0(A_k) \cap B(0; R)$, where the A_k s form a partition of the support of Ω into basic Lipschitz sets. To conclude, we

use that (i) the intersection a Lipschitz domain with a ball is still a Lipschitz domain, and (ii) T_0^{-1} is a Lipschitz diffeomorphism with polylogarithmic distortion on B(0; R). Therefore, $\Omega^{\hat{R}}$ is a *L*-Lipschitz domain with $L \leq R \leq_{\log n} 1$. Eventually, the density $q^{\hat{R}}$ of $Q^{\hat{R}}$ is obtained by a change of variables: for $y \in B(0; R)$,

we have

$$q^{R}(y) = \frac{p(T_{0}^{-1}(y))|\det(DT_{0}^{-1}(y))|}{Q(B(0;R))}.$$
(111)

By strong convexity and smoothness, we have $1 \leq_{\log n,d} |\det(DT_0^{-1}(y))| \leq_{\log n,d} 1$. Furthermore, using condition (C2) and strong convexity of φ_0 , $\log p \circ T_0^{-1}$ is Lipschitz continuous of with constant of order $R^{\theta} \leq_{\log n} 1$. These two properties are enough to prove the last statement of Lemma 8.

G Partitions of Lipschitz domains

We prove in this section Lemma 9 and Lemma 10. To do so, we first give a decomposition of a Lipschitz domain into small cube-like shapes. Let Ω be the support of Q. By Lemma 8, we can assume Ω to be a L-Lipschitz domain included in B(0; R) for $L, R \leq_{\log n} 1$, and q to satisfy condition (62). By definition of a Lipschitz domain, Ω is partitioned (up to a negligible set) into K basic Lipschitz sets $A_k = \Phi_k([0, 1]^d)$ for some Lipschitz diffeomorphisms Φ_k whose differentials and inverse differentials are bounded by L. Note that the number K of such sets is at most of order $(R/L)^d \lesssim_{\log n,d} 1$.

Let $J \geq 1$. We consider a partition $\mathcal{A}_{J,d}$ of $[0,1]^d$ by a regular grid of cubes of side-length 2^{-J} . A partition of A_k is obtained by considering the sets $\Phi_k(\Box)$ for $\Box \in \mathcal{A}_{J,d}$.

The union of these partitions for the different A_k s gives a partition \mathcal{A}_J of Ω . For $\Box \in \mathcal{A}_J$, let Q_{\Box} be the measure Q conditioned on \Box . Let m_{\Box} be the expected value of Q_{\Box} and Σ_{\Box} be its covariance matrix. We choose J such that $2^{-J}(\log n)^{\kappa} \leq 1$, where κ is the constant in (62).

Lemma 27. Let $\Box \in \mathcal{A}_J$.

- 1. The diameter of $\Box \in \mathcal{A}_J$ is smaller than $cL2^{-J}$.
- 2. The volume of \Box is between $cL^{-d}2^{-Jd}$ and $CL^{d}2^{-Jd}$.
- 3. The Poincaré constant of Q_{\Box} satisfies $C_{PI}(Q_{\Box}) \lesssim_{\log n,d} 2^{-2J}$.
- 4. We have $\lambda_{\min}(\Sigma_{\Box}) \gtrsim_{\log n.d} 2^{-2J}$.

Proof. Let $\Box \in \mathcal{A}_J$, written as $\Phi_k(\widetilde{\Box})$ for some square $\widetilde{\Box}$ of side-length 2^{-J} . Points 1. and 2. follow immediately from Φ_k and Φ_k^{-1} being *L*-Lipschitz. [VV12, Proposition 2.1] implies that the Poincaré constant of the uniform measure on \Box is at most $c2^{-2J}L^2$. The same control holds for the Poincaré constant of Q_{\Box} , with a Poincaré constant multiplied by $\frac{\sup_{\Box} q}{\inf_{\Box} q} \lesssim_{\log n,d} e^{c2^{-J}(\log n)^{\kappa}} \lesssim_{\log n,d} 1 \text{ (where we use (62))}.$

It remains to bound the smallest eigenvalue of Σ_{\Box} . The Lipschitz continuity of Φ_k^{-1} implies that $\Box = \Phi_k(\tilde{\Box})$ contains a ball B of radius $2^{-J}/L$ centered at a certain x_0 . For u a unit vector, writing λ for the Lebesgue measure,

$$u^{\top} \Sigma_{\Box} u = \frac{1}{Q(\Box)} \int_{\Box} \langle x - m_{\Box}, u \rangle^{2} dQ(x)$$

$$\geq \frac{\inf_{\Box} q}{\sup_{\Box} q} \frac{1}{\lambda(\Box)} \int_{\Box} \langle x - m_{\Box}, u \rangle^{2} dx$$

$$\gtrsim_{\log n, d} \frac{\lambda(B)}{\lambda(\Box)} \frac{1}{\lambda(B)} \int_{B} \langle x - m_{\Box}, u \rangle^{2} dx$$

$$\gtrsim_{\log n, d} \frac{\lambda(B)}{\lambda(\Box)} \frac{1}{\lambda(B)} \int_{B} \langle x - x_{0}, u \rangle^{2} dx,$$

where in the last step, we used that the covariance over B is centered at x_0 . The average integral is larger than the smallest eigenvalue of the covariance matrix uniform measure on a ball of radius $2^{-J}/L$, which is of order $2^{-2J}/L^2$. Also, as

$$\frac{\lambda(B)}{\lambda(\Box)}\gtrsim \frac{2^{-Jd}L^{-d}}{2^{-Jd}L^d}\gtrsim L^{-2d},$$

we obtain the result.

For $g \neq L_2$ -function, recall that $\prod_J g$ is the piecewise linear function given by the $L_2(Q)$ projection of g on the sets of piecewise linear functions on each part of the partition \mathcal{A}_J . We
write g_{\Box} for the Q-average of g on a given element $\Box \in \mathcal{A}_J$.

Lemma 28 (Control of the L_2 -norm of the projection). Every $g \in \mathcal{G}$ satisfies $\|\Pi_J g - g\|_{L_2(Q)} \lesssim_{\log n,d} 2^{-J} \tau$.

Proof. Let $\Pi_J g$ be the piecewise constant function, equal to g_{\Box} on $\Box \in \mathcal{A}_J$. By definition of the projection,

$$\|\Pi_J g - g\|_{L_2(Q)}^2 \le \|\tilde{\Pi}_J g - g\|_{L_2(Q)}^2 = \sum_{\Box \in \mathcal{A}_J} \int_{\Box} (g - g_{\Box})^2 \, \mathrm{d}Q.$$

By applying the Poincaré inequality on each part of the partition, we obtain

$$\|\Pi_J g - g\|_{L_2(Q)}^2 \lesssim_{\log n,d} \sum_{\Box \in \mathcal{A}_J} 2^{-2J} \int_{\Box} \|\nabla g\|_{L_2(Q)}^2 \lesssim_{\log n,d} 2^{-2J} \|\nabla g\|_{L_2(Q)}^2.$$

We use Lemma 6 to conclude.

Lemma 29 (Control of the L_{∞} -norm of the projection). Let $M = \sup_{g \in \mathcal{G}} ||g||_{\mathcal{C}^{2}(\Omega)}$. Every $g \in \mathcal{G}$ satisfies $||\Pi_{J}g - g||_{\infty} \lesssim_{\log n,d} 2^{-2J}M$.

Proof. Fix $\Box \in \mathcal{A}_J$ and let $x \in \Box$. One can check (by writing out explicitly an orthonormal basis) that

$$\Pi_{J}g(x) = g_{\Box} + \frac{1}{Q(\Box)} \int_{\Box} (x - m_{\Box})^{\top} \Sigma_{\Box}^{-1} (y - m_{\Box}) g(y) \, \mathrm{d}Q(y).$$
(112)

Therefore, as we control the diameter of \Box and the smallest eigenvalue of Σ ,

$$|\Pi_J g(x)| \lesssim_{\log n, d} \left(1 + \frac{1}{\lambda_{\min}(\Sigma_{\Box})} 2^{-2J}\right) ||g||_{L_{\infty}(\Box)} \lesssim_{\log n, d} ||g||_{L_{\infty}(\Box)}.$$
 (113)

For $y \in \Box$, we can write by a Taylor expansion $g(y) = g(x) + \langle \nabla g(x), y - x \rangle + \varepsilon_x(y) = g_x(y) + \varepsilon_x(y)$, with $|\varepsilon_x(y)| \leq_{\log n,d} M2^{-2J}$. Note that g_x is a linear function, so that $\prod_J g_x = g_x$. Therefore, by (113)

$$|\Pi_J g(x) - g(x)| = |\Pi_J (g - g_x)(x) - (g - g_x)(x)| \lesssim_{\log n, d} |\Pi_J (\varepsilon_x)(x)| + M2^{-2J} \lesssim_{\log n, d} M2^{-2J}. \square$$

We are now in position to prove Lemma 9 and Lemma 10.

Proof of Lemma 9. Recalling the expression of the projection (112), we write $\Pi_J g = \tilde{\Pi}_J g + R_J g$, where $\tilde{\Pi}_J g$ is given by local averages g_{\Box} on each $\Box \in \mathcal{A}_J$ whereas $R_J g = \Pi_J g - \tilde{\Pi}_J g$ is a remainder term. We have

$$\mathbb{E}[|\sup_{g\in\mathcal{G}}(Q-Q_n)(R_Jg)|] = \mathbb{E}[|\sup_{g\in\mathcal{G}}\sum_{\Box}(Q-Q_n)(R_Jg\mathbf{1}\{\Box\})|].$$

Let $\Box \in \mathcal{A}_J$. Let x_{\Box} be the identity function on \Box . As $\int_{\Box} (y - m_{\Box}) dQ(y) = 0$, we have

$$(Q-Q_n)R_Jg\mathbf{1}\{\Box\} = \frac{1}{Q(\Box)} \int_{\Box} ((Q-Q_n)(x_{\Box}))^{\top} \Sigma_{\Box}^{-1}(y-m_{\Box})g(y) \, \mathrm{d}Q(y)$$

$$= \frac{1}{Q(\Box)} \int_{\Box} ((Q-Q_n)(x_{\Box}))^{\top} \Sigma_{\Box}^{-1}(y-m_{\Box})(g(y)-g_{\Box}) \, \mathrm{d}Q(y)$$

$$\lesssim_{\log n,d} 2^J ||(Q-Q_n)(x_{\Box})|| ||g-g_{\Box}||_{L_2(Q_{\Box})} \lesssim_{\log n,d} ||(Q-Q_n)(x_{\Box})|| ||\nabla g\mathbf{1}\{\Box\}||_{L_2(Q_{\Box})},$$

where we use the Poincaré inequality and Lemma 27 at the last line. By Cauchy-Schwartz inequality,

$$\mathbb{E}[|\sup_{g\in\mathcal{G}}\sum_{\Box}(Q-Q_n)(R_Jg\mathbf{1}\{\Box\})|]$$

$$\lesssim_{\log n,d}\left(\sum_{\Box}\|\nabla g\mathbf{1}\{\Box\}\|_{L_2(Q)}^2\right)^{1/2}\left(\sum_{\Box}\mathbb{E}[\|(Q-Q_n)(x_{\Box})\|^2]\right)^{1/2}$$

$$\lesssim_{\log n,d}\tau\left(\sum_{\Box}\frac{1}{n}\operatorname{Var}_Q(x_{\Box})\right)^{1/2}\lesssim_{\log n,d}\frac{\tau}{\sqrt{n}}\left(\sum_{\Box}Q(\Box)2^{-2J}\right)^{1/2}\lesssim_{\log n,d}\frac{\tau 2^{-J}}{\sqrt{n}},$$

where we use the Poincaré inequality to bound the variance, and the control on the Poincaré constant given in Lemma 27.

The second term to control is $\mathbb{E}[|\sup_{g\in\mathcal{G}}(Q-Q_n)(\tilde{\Pi}_J g)|]$. This quantity is left unchanged if we replace g by g + c for some constant c, so that we assume without loss of generality that $\int g \, \mathrm{d}Q = 0$. The partitions \mathcal{A}_J are nested, so that we may write

$$\tilde{\Pi}_J g = \sum_{j=1}^J (\tilde{\Pi}_j g - \tilde{\Pi}_{j-1} g)$$

The function $\Delta_j g = \tilde{\Pi}_j g - \tilde{\Pi}_{j-1} g$ is piecewise constant on each subcube of $\Box \in \mathcal{A}_{j-1}$. More precisely, let $\mathcal{C}(\Box)$ be the set of subcubes of \Box in \mathcal{A}_j . Then, on $E \in \mathcal{C}(\Box)$, $\Delta_j g$ is equal to $b_{j,\Box,E} = \frac{1}{Q(E)} \int_E g - \frac{1}{Q(\Box)} \int_{\Box} g$.

Lemma 30. We have

$$\sum_{\Box \in \mathcal{A}_{j-1}} \sum_{E \in \mathcal{C}(\Box)} Q(E) b_{j,E}(g)^2 \lesssim_{\log n,d} 2^{-2j} \|\nabla g\|_{L_2(Q)}^2.$$
(114)

Proof. By Jensen's inequality and Poincaré inequality,

$$b_{j,E}(g)^2 = \left(\frac{1}{Q(E)}\int_E (g - g_{\Box})\right)^2 \le \frac{1}{Q(E)}\int_E (g - g_{\Box})^2 \lesssim_{\log n,d} \frac{1}{Q(E)}2^{-2j}\int_{\Box} \|\nabla g\|^2$$

As there are 2^d subcubes for each part of the partition, we obtain

$$\sum_{\Box \in \mathcal{A}_{j-1}} \sum_{E \in \mathcal{C}(\Box)} Q(E) b_{j,E}(g)^2 \lesssim_{\log n,d} 2^{-2j} \sum_{\Box \in \mathcal{A}_{j-1}} \int_{\Box} \|\nabla g\|^2 \lesssim_{\log n,d} 2^{-2j} \|\nabla g\|_{L_2(Q)}^2.$$

By Cauchy-Schwartz inequality and Lemma 30,

$$\begin{aligned} \mathbb{E}[|\sup_{g\in\mathcal{G}}(Q-Q_n)(\tilde{\Pi}_Jg)| &\leq \sum_{j=1}^J \mathbb{E}[|\sup_{g\in\mathcal{G}}(Q-Q_n)(\Delta_Jg)|] \\ &\leq \sum_{j=1}^J \mathbb{E}[|\sup_{g\in\mathcal{G}}\sum_{\square\in\mathcal{A}_{j-1},E\in\mathcal{C}(\square)} b_{j,E}(g)(Q-Q_n)(E)|] \\ &\leq \sum_{j=1}^J \sup_{g\in\mathcal{G}} \left(\sum_{\square,E}Q(E)b_{j,E}(g)^2\right)^{1/2} \left(\sum_{\square,E}\frac{1}{Q(E)}\mathbb{E}[(Q_n-Q)(E)^2]\right)^{1/2} \\ &\lesssim_{\log n,d} \sum_{j=1}^J 2^{-j}\tau \left(\sum_{\square,E}\frac{1}{n}\right)^{1/2} \lesssim_{\log n,d} n^{-1/2}2^{\frac{d-2}{2}J}\tau, \end{aligned}$$

where we use at the last line that there the number of elements in the partition \mathcal{A}_j is of order 2^{jd} (up to polylogarithmic factors in n).

Proof of Lemma 10. Let $g = \varphi_c^* - \overline{\varphi}_c^* \in \mathcal{G}$, with φ and $\overline{\varphi}$ that are $(\alpha/2, a)$ -strongly convex. Then, the \mathcal{C}^2 -norm of φ_c^* restricted to B(0; R) is controlled by $\sup_{\|y\| \leq R} \lambda_{\min}^{-1}(\nabla^2 \varphi(\nabla \varphi^{-1}(y)))$, which grows at most polynomially in R by Lemma 15 and the definition of strongly convex. Let $\mathcal{G}_{>J} = \{\Pi_J g - g : g \in \mathcal{G}\}$. Then, by Lemma 29, $\mathcal{G}_{>J}$ has an envelope function G_J with $G_J \leq_{\log n,d} 2^{-2J}$. Furthermore, by Lemma 28, we have $\sup_{f \in \mathcal{G}_{>J}} \|f\|_{L_2(Q)} \leq \sigma$ for $\sigma \asymp_{\log n,d} 2^{-J}\tau$. Let us use Proposition 6 to bound $\mathbb{E}[|\sup_{g \in \mathcal{G}} (Q - Q_n)(g - \Pi_J g)|]$. We use the following bound on bracketing numbers, proven below.

Lemma 31 (Control of the bracketing numbers). For δ a small enough constant, and for h > 0, it holds that

$$\log \mathcal{N}_{[]}(h, \mathcal{G}_{>J}, L_2(Q)) \le \log \mathcal{N}(ch, \mathcal{F}, L_{\infty}(e^{-\delta \|\cdot\|})) + C \log_+(1/h).$$
(115)

Using this lemma and condition (A2), one can check that the condition $2^{-J(2-\gamma)} \geq c_1(\log n)^{c_2} \frac{n}{D_F} \tau^{2+\gamma}$ ensures that the parameter

$$\omega = \sqrt{\frac{n\sigma^2}{32\log(2\mathcal{N}_{[]}(\sigma/2,\mathcal{G}_{>J},L_2(Q)))}}$$

is smaller than G_J , so that the second term in (45) is null. We therefore obtain

$$\mathbb{E}\left[|\sup_{g\in\mathcal{G}}(Q-Q_n)(g-\Pi_J g)|\right] \le \frac{116}{\sqrt{n}}\mathcal{J}_{[]}(2\sigma,\mathcal{G}_{>J},L_2(Q)).$$
(116)

Lemma 31 and condition (A2) then gives the desired bound on the entropic integral. \Box *Proof of Lemma 31.* Let $\varphi_1, \varphi_2 \in \text{Li}(\mathcal{F})_{\tau}$ that are $(\alpha/2)$ -strongly convex. By (113)

$$\int |\Pi_J \varphi_1^*(y) - \Pi_J \varphi_2^*(y)|^2 \, \mathrm{d}Q(y) = \sum_{\square \in \mathcal{A}_J} \int_{\square} |\Pi_J \varphi_1^*(y) - \Pi_J \varphi_2^*(y)|^2 \, \mathrm{d}Q(y)$$
$$\lesssim_{\log n, d} \sum_{\square \in \mathcal{A}_J} Q(\square) \|\varphi_1^* - \varphi_2^*\|_{L_{\infty}(\square)}^2.$$

Let $y_{\Box} = T_0(x_{\Box})$ be the point of \Box with the largest norm. By Lemma 15, every point $y \in \Box$ is equal to $T_0(x)$ for some x satisfying $||x|| \leq ||T(x)||^{1/(a+1)} \leq ||y_{\Box}||^{1/(a+1)}$. By Lemma 26, we therefore have for every $\delta > 0$,

$$\|\varphi_1^* - \varphi_2^*\|_{L_{\infty}(\Box)} \lesssim \|\varphi_1 - \varphi_2\|_{L_{\infty}(e^{-\delta \|\cdot\|})} e^{C\delta \|y_{\Box}\|^{1/(a+1)}} \lesssim \|\varphi_1 - \varphi_2\|_{L_{\infty}(e^{-\delta \|\cdot\|})} e^{C'\delta \|x_{\Box}\|}, \quad (117)$$

where we also use Lemma 14. Therefore,

$$\int |\Pi_J \varphi_1^*(y) - \Pi_J \varphi_2^*(y)|^2 \,\mathrm{d}Q(y) \lesssim_{\log n, d} \|\varphi_1 - \varphi_2\|_{L_\infty(e^{-\delta \|\cdot\|})}^2 \sum_{\Box \in \mathcal{A}_J} Q(\Box) e^{C'\delta \|x_{\Box}\|}$$

For every $y \in \Box$, we have by Lemma 15

$$||x_{\Box}|| = ||T_0^{-1}(y_{\Box})|| \le ||T_0^{-1}(y_{\Box}) - T_0^{-1}(y)|| + ||T_0^{-1}(y)|| \le cL2^{-J} + ||T_0^{-1}(y)||.$$

We may therefore bound the sum by a corresponding integral:

$$\sum_{\square \in \mathcal{A}_J} Q(\square) e^{C'\delta \|x_\square\|} \le \int e^{C'\delta(cL2^{-J} + \|T_0^{-1}(y)\|)} \,\mathrm{d}Q(y) \lesssim \int e^{C'\delta \|x\|} \,\mathrm{d}P(x) \lesssim 1,$$

where we pick δ small enough so that the last integral is finite. In total, as bracketing numbers are controlled by covering numbers for the L_{∞} -norm (see [VW96, Theorem 2.7.11]), we have shown that

$$\log \mathcal{N}_{[]}(h, \mathcal{G}_{>J}, L_2(Q)) \le \log \mathcal{N}(ch, \mathcal{F}, L_{\infty}(e^{-\delta \|\cdot\|})) + C \log_+(1/h)$$

where we used Lemma 19 and Lemma 22 to respectively account for the fact that \mathcal{G} is indexed by a subset of $\text{Li}(\mathcal{F})$ and for the centering of the functions in \mathcal{G} .

H On the spectrum of kernels for RKHS

Here we provide some details about the spectrum of kernels, following the discussion of Section 4.4. To this end, recall that a kernel $\mathcal{K} : \mathcal{X} \times \mathcal{X} \to \mathbb{R}$ defines a Reproducing Kernel Hilbert Space (RKHS) if for all $f \in \mathcal{H}$ (the Hilbert space over \mathcal{X}) and all $x \in \mathcal{X}$, it holds that

$$f(x) = \langle f, \mathcal{K}(\cdot, x) \rangle_{\mathcal{H}}.$$

In Section 4.4, we provided log-covering bounds from [Yan+20b] that required that the kernel have either finite spectrum, or exponentially decaying eigenvalues; the following is borrowed from [Yan+20b, Section 2.2 and Appendix D]. To this end, we note that \mathcal{K} induces an integral operator $T_{\mathcal{K}}: L^2(\mathcal{X}) \to L^2(\mathcal{X})$ defined as

$$T_{\mathcal{K}}f(y) = \int \mathcal{K}(y, x)f(x) \,\mathrm{d}x,$$

for all $f \in L^2(\mathcal{X})$. Mercer's theorem (see e.g. [SSB+02]) states that this operator has countably many, positive eigenvalues with corresponding eigenfunctions $\{(\sigma_i, \vartheta_i)\}_{i=1}^{\infty}$ which form an orthonormal basis of $L^2(\mathcal{X})$, whence the kernel admits the following spectral decomposition

$$\mathcal{K}(x,y) = \sum_{i=1}^{n} \sigma_i \vartheta_i(x) \vartheta_i(y) \,. \tag{118}$$

The two conditions we presented are the following:

- 1. γ -finite spectrum: $\sigma_j = 0$ for all $j \geq \gamma$,
- 2. γ -exponential decay: there exist absolute constants C_1 and C_2 such that for all $j \ge 1$, $\sigma_j \le C_1 \exp(-C_2 j^{\gamma})$,

where for the latter, we also require that there exist constants $\tau \in [0, \frac{1}{2}), C_{\vartheta} > 0$ such that $\sup_{x \in \mathcal{X}} \sigma_j^{\tau} |\vartheta_j(x)| \leq C_{\vartheta}$ for all $j \geq 1$. In fact, over the sphere in \mathbb{R}^d , it holds that the Gaussian kernel exhibits γ -exponential decay for any $\tau > 0$ [Yan+20b, Appendix B.3], wherein several more examples of kernels are presented. The following is a proposition from [Yan+20b], which is an upper bound on the covering numbers we employ in Section 4.4.

Proposition 8. Let $\mathcal{F}_{\mathcal{K}}$ be the unit ball in an RKHS corresponding to some kernel \mathcal{K} i.e. $\mathcal{F}_{\mathcal{K}} := \{f : ||f||_{\mathcal{H}} \leq 1\}$, and let \mathcal{K} be a kernel with γ -finite spectrum, or γ -exponentially decaying spectrum, as described above, such that $\sup_{z} \mathcal{K}(z, z) \leq 1$. Then,

$$\log \mathcal{N}(h, \mathcal{F}_{\mathcal{K}}, \|\cdot\|_{\infty}) \leq \begin{cases} C\gamma(\log(1/h) + c) & \gamma\text{-finite spectrum}, \\ C(\log(1/h) + c)^{1+1/\gamma} & \gamma\text{-exponentially decaying spectrum}, \end{cases}$$

where the absolute constants c, C > 0 depend on $C_{\vartheta}, C_1, C_2, \gamma$ and τ .

I Covering number for the spiked transport case

Proof of Proposition **4**. Recall that

$$\mathcal{F}_{\text{spiked}} = \left\{ x \mapsto \varphi(Ux) + \frac{1}{2} x^{\top} (I - U^{\top} U) x : \varphi \in \mathcal{F}, U \in \mathcal{V}_{k \times d} \right\} ,$$

where $\mathcal{V}_{k \times d}$ is the Stiefel manifold, and \mathcal{F} is some function class. Log-covering number bounds for Stiefel manifold are already known (see [NR22, Lemma 4]):

$$\log \mathcal{N}(h, \mathcal{V}_{k \times d}, \|\cdot\|_{\mathrm{op}}) \le dk \log(c\sqrt{k}/h), \qquad (119)$$

where c > 0 is some absolute constant. The same bound holds for the space $L_{\infty}(e^{-\delta \|\cdot\|})$ up to a constant depending on δ , as in our previous calculations. To complete the proof, we can decompose a function $\varphi' \in \mathcal{F}_{\text{spiked}}$ into two parts; $\varphi \circ U$ and a quadratic. The quadratic part contributes the same degree of complexity as a parametric class (see Section 4.3): indeed we have that

$$\|x^{\top}(I - U^{\top}U)x - x^{\top}(I - V^{\top}V)x\| = \|x^{\top}(U + V)^{\top}(U - V)x\|$$

$$\leq \|x\|^{2}\|U + V\|_{F}\|U - V\|_{F}$$

$$\leq C_{d}\|x\|^{2}\|U - V\|_{\text{op}},$$

using that $||U||_{op}$, $||V||_{op} \leq 1$ and that the Frobenius and operator norms are equivalent. Thus, we can apply known bounds on the covering numbers for the Stiefel manifold to determine the size of the (log-)covering numbers that the quadratic contributes to the class \mathcal{F}_{spiked} . Finally, log-covering numbers for the composition $\varphi \circ U$ can simply be bounded by considering the sum of the log-covering numbers of the two classes. Thus, applying Eq. (119), completes the proof.

J Besov spaces

We now prove the results in Section 4.5 by relevant recalling facts about Besov spaces, which can be found in many textbooks (see e.g. [GN21]). The Besov spaces $B_{p,q}^s$ are a family of functional spaces, with s representing a regularity index, that can be defined using a wavelet basis. A wavelet basis is an orthonormal basis $(\Gamma_{j,k})_{j\in\mathbb{Z}, k\in\mathbb{Z}^d}$ of $L_2(\mathbb{R}^d)$ that satisfy some desirable properties. Let s > 0. We will always assume that the regularity of the wavelet basis is large enough with respect to s and d. Let $1 \leq p, q \leq \infty$. If a function φ admits a wavelet representation

$$\varphi = \sum_{j \in \mathbb{Z}} \sum_{k \in \mathbb{Z}^d} \gamma_{k,j} \Gamma_{k,j}, \qquad (120)$$

then, the Besov norm $B_{p,q}^s$ of φ is defined as

$$\|\varphi\|_{B^{s}_{p,q}} = \|\gamma\|_{b^{s}_{p,q}} = \left(\sum_{j \in \mathbb{Z}} 2^{jq(s+\frac{d}{2}-\frac{d}{p})} \left(\sum_{k \in \mathbb{Z}^{d}} |\gamma_{j,k}|^{p}\right)^{\frac{q}{p}}\right)^{\frac{1}{q}},$$
(121)

with generalizations when p or q equal to ∞ .

Let R > 0. Each wavelet $\Gamma_{k,j}$ has compact support of diameter of order 2^{-j} , and, for a fixed j, the number of wavelets $\Gamma_{k,j}$ that intersect the ball B(0; R) is of order $R^d 2^{jd}$. Let $I_j(R)$ be the set of indexes k such that this intersection is nonempty. Let $J_0 \in \mathbb{Z}$ be such that B(0; R) is included in the support of a single wavelet Γ_{k,J_0} . We can pick J_0 such that $J_0 \gtrsim -\log R$. Let L > 0 and define the function $\eta_R(x)$ to be a smooth convex function equal to $||x||^2$ for $||x|| \ge R/2$, and equal to 0 for $||x|| \le R/4$. We define the set $\mathcal{F}_J(R)$ as the set of functions of the form

$$\eta_R(x) + \sum_{j=J_0}^J \sum_{k \in I_j(R)} \gamma_{k,j} \Gamma_{k,j},$$

where, $\|\gamma\|_{b_{1,\infty}^s} \leq L$.

To put it another way, functions $\varphi \in \mathcal{F}_J(R)$ are functions with finite wavelet expansions up to depth J and bounded $B_{1,\infty}^s$ norm on a set of size roughly R, whereas we add a quadratic term to ensure that they are strongly convex at infinity. Note however that functions in $\mathcal{F}_J(R)$ are not necessarily convex.

Let us show that, under the assumptions of Proposition 3, we can apply Theorem 3 to obtain a rate of convergence. By hypothesis, assumptions (A3), (C1) and (C2) are satisfied. Furthermore, as the C^2 -norm is controlled by the $B^2_{1,\infty}$ -norm and that $s \ge 2$, all potentials in $\mathcal{F}_J(R)$ are $(\beta, 0)$ -smooth. It remains to check the bracketing assumption (A2). Let $\delta > 0$ and let $\varphi_1, \varphi_2 \in \mathcal{F}_J(R)$, with associated coordinates γ^1 and γ^2 . We have

$$\begin{aligned} |\varphi_1(x) - \varphi_2(x)| e^{-\delta ||x||} &= |\sum_{j=J_0}^J \sum_{k \in I_j(R)} (\gamma_{j,k}^1 - \gamma_{j,k}^2) \Gamma_{k,j}(x)| e^{-\delta ||x||} \\ &\lesssim R^{d/2} 2^{Jd/2} ||\gamma^1 - \gamma^2||_{\infty}, \end{aligned}$$

where we bound $e^{-\delta \|\cdot\|}$ by 1, and use a standard bound on the L_{∞} -norm in terms of wavelet coefficients (see e.g. [HR21, Lemma 24]). Note that only the coefficients $\gamma_{k,j}^1 - \gamma_{k,j}^2$ with $k \in I_j(R)$ are non zero. Therefore, the bracketing number of $\mathcal{F}_J(R)$ at scale h with respect to $L_{\infty}(e^{-\delta \|\cdot\|})$ is controlled by the covering number at scale of order $hR^{-d/2}2^{-Jd/2}$ of the ℓ_{∞} -ball in dimension $\sum_{j=-J_0}^{J} |I_j(R)| \leq R^d 2^{Jd}$. By [Sch84], the logarithm of such a covering number is of order

$$R^d 2^{Jd} \log(R^{d/2} 2^{Jd/2} / h) \lesssim R^d 2^{Jd} (\log(R) + J) \log_+(1/h)$$

that is assumption (A2) holds with $\gamma = 0$, $\gamma' = 1$ and $D_{\mathcal{F}} = R^d 2^{Jd} (\log(R) + J)$.

We are in position to apply Theorem 3, which gives a control up to logarithmic term

$$\inf_{\varphi \in \mathcal{F}_{\alpha}} (S(\varphi) - S(\varphi_0)) + D_{\mathcal{F}}^{\frac{d-2}{d}} n^{-1}.$$

To conclude, it remains to bound the bias term $\inf_{\varphi \in \mathcal{F}_{\alpha}}(S(\varphi) - S(\varphi_0))$. Let γ^0 be the vector of coefficients of φ_0 in the wavelet basis, and let $\varphi_{0,J}(x) = \eta_R(x) + \psi_{0,J}(x)$, where

$$\psi_{0,J}(x) = \sum_{j=J_0}^{J} \sum_{k \in I_j(R)} \gamma_{j,k}^0 \Gamma_{jk}(x).$$
(122)

As $s \geq 2$, the norm of the Hessian of $\psi_{0,J} - \varphi_0$ at $x \in B(0; R)$, is bounded by $2^{(J_0 - J)2} \|\varphi_0\|_{\mathcal{C}^s} \leq \alpha/2$ for J large enough. Therefore, as φ_0 is α -strongly convex, we obtain that $\psi_{0,J}$ is $(\alpha/2)$ -strongly convex. As $\varphi_{0,J}$ is obtained by adding the convex function η_R , $\varphi_{0,J}$ is also $(\alpha/2)$ -strongly convex. We apply a reverse stability bound for strongly convex potentials (that would be proven in the same fashion as [HR21, Proposition 10]) to obtain that

$$S(\varphi_{0,J}) - S(\varphi_{0}) \lesssim \|\nabla\varphi_{0} - \nabla\varphi_{0,J}\|_{L_{2}(P)}^{2} = \int \|\nabla\varphi_{0}(x) - \nabla\varphi_{0,J}(x)\|^{2} \,\mathrm{d}P(x)$$

$$\leq \|p\|_{\infty} \int_{\|x\| \le R/4} \|\nabla\varphi_{0}(x) - \nabla\varphi_{0,J}(x)\|^{2} \,\mathrm{d}x + \int_{\|x\| > R/4} \|\nabla\varphi_{0}(x) - \nabla\varphi_{0,J}(x)\|^{2} \,\mathrm{d}P(x).$$

The first integral is bounded by a quantity of order $2^{2(J_0-J)(s-1)} \|\varphi_0\|_{\mathcal{C}^s}^2$ (see e.g. [HR21, Lemma 13]). The second integral is of order e^{-cR} as P is subexponential. We pick $R \approx 2^{J_0} = \log n$. Up to logarithmic terms, we obtain the final bias-variance decomposition

$$2^{-2J(s-1)} + 2^{J(d-2)}n^{-1}.$$

Eventually, we let $2^{-J} \simeq n^{-\frac{1}{2s+d-4}}$ to conclude.

K Barron spaces

Proof of Theorem 4. By construction, every $\varphi \in \mathcal{F}_{\sigma}^{1}$ is (β, a) -smooth, so that assumptions (A1) and (C1) are satisfied. To conclude, it remains to bound the covering numbers. \Box

Lemma 32. It holds that, for $h, \delta > 0$,

$$\log \mathcal{N}(h, \mathcal{F}_{\sigma}^{1}, L_{\infty}(e^{-\delta \|\cdot\|})) \lesssim_{\log_{+}(1/h)} h^{-\frac{2m}{2s+m}}.$$
(123)

Let B be a Banach space. We say that B is of type 2 if there exists a constant $T_B > 0$ such that for all $x_1, \ldots, x_n \in B$,

$$\mathbb{E}\left[\left\|\sum_{i=1}^{n}\varepsilon_{i}x_{i}\right\|_{B}^{2}\right] \leq T_{B}^{2}\sum_{i=1}^{n}\|x_{i}\|_{B}^{2},\qquad(124)$$

where $\varepsilon_1, \ldots, \varepsilon_n$ are i.i.d. Rademacher variables. Siegel and Xu show in [SX21] that, if $\sigma : \mathcal{M} \to B$ is of regularity *s* for some Banach space *B* of type 2 (where regularity is defined through dual maps, see [SX21, Definition 1] for details), then the covering numbers of \mathcal{F}_{σ}^1 satisfy for h > 0

$$\log \mathcal{N}(h, \mathcal{F}_{\sigma}^{1}, B) \leq C \left(\frac{h}{T_{B}}\right)^{-\frac{2m}{2s+m}}.$$
(125)

Such a result is almost enough to conclude. Indeed, the Banach space $L_{\infty}(e^{-\delta \|\cdot\|})$ is not of type 2. We use an approximation scheme, that shows that the covering numbers for the ∞ -norm cannot be too different from the covering numbers for the *p*-norm, where *p* is large. As spaces L_p are of type 2, this gives us the conclusion.

Lemma 33. Let \mathcal{F} be a set of (β, a) -smooth functions, with $\sup_{f \in \mathcal{F}} \|\nabla f(0)\| \leq M$ and f(0) = 0 for all $f \in \mathcal{F}$. Assume that \mathcal{F} satisfies that, for all finite measures ρ and $2 \leq p < \infty$,

$$\log \mathcal{N}(h, \mathcal{F}, L_p(\rho)) \le C \left(\frac{h}{T_{L_p(\rho)}}\right)^{-\gamma}.$$
(126)

Then, for $\delta, h > 0$,

$$\log \mathcal{N}(h, \mathcal{F}, L_{\infty}(e^{-\delta \|\cdot\|})) \lesssim_{\log_{+}(1/h)} h^{-\gamma}.$$
(127)

Before writing a proof, let us remark that one can use this lemma with (125) to obtain that (A2) holds with $\gamma = \frac{2m}{2s+m}$, which gives Theorem 4 through Theorem 3.

Proof. Let us first recall Khintchine's inequality. It states that for any finite measure ρ and $2 \leq p < \infty$, the space $L_p(\rho)$ is of type 2 with $T_{L_p(\rho)} \leq C\rho(\mathbb{R}^d)^{1/p}\sqrt{p}$, for some absolute constant C (see e.g. [Ver18, Section 2.6]). Let ρ_R be the Lebesgue measure on B(0; R) and let $\{f_1, \ldots, f_K\}$ be an h'-covering of \mathcal{F} for the norm $L_p(\rho_R)$ for some parameter h' to fix. By assumption, we can pick K such that

$$\log K \le C \left(\frac{h'}{\rho_R(\mathbb{R}^d)^{1/p} \sqrt{p}}\right)^{-\gamma}$$

Let $f \in \mathcal{F}$, and let f_k be such that $||f - f_k||_{L_p(\rho_R)} \leq h'$. By Lemma 14, we have

$$\|f - f_k\|_{L_{\infty}(e^{-\delta \|\cdot\|})} \le \|f - f_k\|_{L_{\infty}(\rho_R)} + 4(2\beta + M)\langle R \rangle^{a+2} e^{-\delta R}.$$
 (128)

We pick $R = c \log_+(1/h)$ for c large enough, so that the second term is smaller than h/2. Let us bound the first term. According to Lemma 14, the function $f - f_k$ is L-Lipschitz continuous on B(0; R), with $L = 4\beta \langle R \rangle^{a+1} + 2M$. Let $x_0 \in B(0; R)$ be such that $m = \sup_{\|x\| \leq R} |f(x) - f_k(x)|$ is attained at x_0 . Therefore, $|f(x) - f_k(x)| \geq m - Lr$ for $x \in B(x_0; r)$. Then, for $r \leq R$

$$(h')^p \ge \int_{B(0;R)} |f(x) - f_k(x)|^p \, \mathrm{d}x \ge \int_{B(x_0;r) \cap B(0;R)} (m - Lr)^p \, \mathrm{d}x \ge c_d r^d (m - Lr)^p.$$

We pick r = m/(2L) (that is smaller than R for h small enough) to obtain that $h' \ge c_{d,L}^{\frac{1}{p}} m^{1+\frac{d}{p}}$. Pick $h' = c_{d,L}^{\frac{1}{p}} (h/2)^{1+\frac{d}{p}}$ to obtain that $m = ||f - f_k||_{L_{\infty}(\rho_R)} \le h/2$. Therefore, the set $\{f_1, \ldots, f_K\}$ is a h-covering of \mathcal{F} for the norm $L_{\infty}(e^{-\delta||x||})$ and we have shown that

$$\log \mathcal{N}(h, \mathcal{F}, L_{\infty}(e^{-\delta \|\cdot\|})) \leq \log K \leq C \left(\frac{h'}{\rho_R(\mathbb{R}^d)^{1/p}\sqrt{p}}\right)^{-\gamma}$$
$$\leq C \left(\frac{c_{d,L}^{\frac{1}{p}}(h/2)^{1+\frac{d}{p}}}{(c_d R^d)^{1/p}\sqrt{p}}\right)^{-\gamma}.$$

We pick $p = \log_+(1/h)$ to conclude.

Eventually, we prove Proposition 5.

Proof of Proposition 5. The upper bound follows from Theorem 4. Consider $C^{s+\frac{d+1}{2}}(B(0;1))$ to be the space of functions of regularity $s + \frac{d+1}{2}$ on the unit ball B(0;1) in \mathbb{R}^d . According to [HR21, Remark 3], the minimax rate of estimation for potentials $\varphi \in C^{s+\frac{d+1}{2}}(B(0;1))$ is of order at least $n^{-\frac{2s+d-1}{2s+2d-3}}$. According to Proposition 5 and the homogeneous reformulation in [Bac17], the space $C^{s+\frac{d+1}{2}}(B(0;1))$ is a subset of \mathcal{F}^1_{σ} . Therefore, the minimax lower bound on \mathcal{F}^1_{σ} is larger than the one on $C^{s+\frac{d+1}{2}}(B(0;1))$, giving the minimax lower bound.

References

[ACB17] M. Arjovsky, S. Chintala, and L. Bottou. "Wasserstein generative adversarial networks". In: Proceedings of the 34th International Conference on Machine Learning. Proceedings of Machine Learning Research 70 (2017). Ed. by D. Precup and Y. W. Teh, pp. 214–223. [AF03] R. A. Adams and J. J. Fournier. *Sobolev spaces*. Elsevier, 2003. [Bac17] F. Bach. "Breaking the curse of dimensionality with convex neural networks". In: The Journal of Machine Learning Research 18.1 (2017), pp. 629–681. [Bak+08]D. Bakry et al. "A simple proof of the Poincaré inequality for a large class of probability measures". In: Electronic Communications in Probability 13 (2008), pp. 60–66. [Bar93] A. R. Barron. "Universal approximation bounds for superpositions of a sigmoidal function". In: IEEE Trans. Inform. Theory 39.3 (1993), pp. 930–945. ISSN: 0018-9448. C. Bunne, A. Krause, and M. Cuturi. "Supervised Training of Conditional [BKC22] Monge Maps". In: arXiv preprint arXiv:2206.14262 (2022). H. J. Brascamp and E. H. Lieb. "On extensions of the Brunn-Minkowski and [BL02] Prékopa-Leindler theorems, including inequalities for log concave functions, and with an application to the diffusion equation". In: Inequalities. Springer, 2002, pp. 441–464. [BL97] S. Bobkov and M. Ledoux. "Poincaré's inequalities and Talagrand's concentration phenomenon for the exponential distribution". In: Probability Theory and *Related Fields* 107.3 (1997), pp. 383–400. [Bre91] Y. Brenier. "Polar factorization and monotone rearrangement of vector-valued functions". In: Communications on pure and applied mathematics 44.4 (1991), pp. 375–417. L. A. Caffarelli. "Monotonicity properties of optimal transportation and the [Caf00]fkg and related inequalities". In: Communications in Mathematical Physics 214.3 (2000), pp. 547–563. [CCG16] G. Carlier, V. Chernozhukov, and A. Galichon. "Vector quantile regression: an optimal transport approach". In: The Annals of Statistics 44.3 (2016), pp. 1165–1192.

- [Che+17] V. Chernozhukov et al. "Monge–Kantorovich depth, quantiles, ranks and signs". In: *The Annals of Statistics* 45.1 (2017), pp. 223–256.
- [CP22] S. Chewi and A.-A. Pooladian. "An entropic generalization of Caffarelli's contraction theorem via covariance inequalities". In: *arXiv preprint arXiv:2203.04954* (2022).
- [Dem+21] P. Demetci et al. "Unsupervised integration of single-cell multi-omics datasets with disparities in cell-type representation". In: *bioRxiv* (2021).
- [DGS21] N. Deb, P. Ghosal, and B. Sen. "Rates of Estimation of Optimal Transport Maps using Plug-in Estimators via Barycentric Projections". In: *arXiv preprint arXiv:2107.01718* (2021).
- [EMW22] W. E, C. Ma, and L. Wu. "The Barron space and the flow-induced function spaces for neural network models". In: Constr. Approx. 55.1 (2022), pp. 369– 406. ISSN: 0176-4276.
- [Fey+17] J. Feydy et al. "Optimal transport for diffeomorphic registration". In: International Conference on Medical Image Computing and Computer-Assisted Intervention. Springer. 2017, pp. 291–299.
- [Fin+20] C. Finlay et al. "Learning normalizing flows from Entropy-Kantorovich potentials". In: *arXiv preprint arXiv:2006.06033* (2020).
- [FLF19] R. Flamary, K. Lounici, and A. Ferrari. "Concentration bounds for linear monge mapping estimation and optimal transport domain adaptation". In: arXiv preprint arXiv:1905.10155 (2019).
- [GN21] E. Giné and R. Nickl. *Mathematical foundations of infinite-dimensional statistical models*. Cambridge University Press, 2021.
- [Goz10] N. Gozlan. "Poincaré inequalities and dimension free concentration of measure". In: Annales de l'IHP Probabilités et statistiques. Vol. 46. 3. 2010, pp. 708– 739.
- [GPC18] A. Genevay, G. Peyré, and M. Cuturi. "Learning generative models with Sinkhorn divergences". In: Proceedings of the 21st International Conference on Artificial Intelligence and Statistics. 2018, pp. 1608–1617.
- [GR90] A. Griewank and P. Rabier. "On the smoothness of convex envelopes". In: Transactions of the American Mathematical Society 322.2 (1990), pp. 691–709.
- [Gra+18] W. Grathwohl et al. "Ffjord: Free-form continuous dynamics for scalable reversible generative models". In: *arXiv preprint arXiv:1810.01367* (2018).
- [Gui+21] J. Gui et al. "A review on generative adversarial networks: Algorithms, theory, and applications". In: *IEEE Transactions on Knowledge and Data Engineering* (2021).
- [GX21] F. Gunsilius and Y. Xu. "Matching for causal effects via multimarginal optimal transport". In: *arXiv preprint arXiv:2112.04398* (2021).
- [HR21] J.-C. Hütter and P. Rigollet. "Minimax estimation of smooth optimal transport maps". In: *The Annals of Statistics* 49.2 (2021), pp. 1166–1194.

- [Hua+21] C.-W. Huang et al. "Convex Potential Flows: Universal Probability Distributions with Optimal Transport and Convex Optimization". In: International Conference on Learning Representations. 2021.
- [KPB20] I. Kobyzev, S. J. Prince, and M. A. Brubaker. "Normalizing flows: An introduction and review of current methods". In: *IEEE transactions on pattern* analysis and machine intelligence 43.11 (2020), pp. 3964–3979.
- [Mak+20] A. Makkuva et al. "Optimal transport mapping via input convex neural networks". In: *International Conference on Machine Learning*. PMLR. 2020, pp. 6672–6681.
- [Man+21] T. Manole et al. "Plugin estimation of smooth optimal transport maps". In: arXiv preprint arXiv:2107.12364 (2021).
- [Mor+21] N. Moriel et al. "NovoSpaRc: flexible spatial reconstruction of single-cell gene expression with optimal transport". In: *Nature Protocols* 16.9 (2021), pp. 4177–4200.
- [Muz+21] B. Muzellec et al. "Near-optimal estimation of smooth transport maps with kernel sums-of-squares". In: *arXiv preprint arXiv:2112.01907* (2021).
- [NR22] J. Niles-Weed and P. Rigollet. "Estimation of wasserstein distances in the spiked transport model". In: *Bernoulli* 28.4 (2022), pp. 2663–2688.
- [PN21] A.-A. Pooladian and J. Niles-Weed. "Entropic estimation of optimal transport maps". In: *arXiv preprint arXiv:2109.12004* (2021).
- [Sal+18] T. Salimans et al. "Improving GANs Using Optimal Transport". In: International Conference on Learning Representations. 2018.
- [San15] F. Santambrogio. "Optimal transport for applied mathematicians". In: *Birkäuser*, NY 55.58-63 (2015), p. 94.
- [Sch+19] G. Schiebinger et al. "Optimal-transport analysis of single-cell gene expression identifies developmental trajectories in reprogramming". In: Cell 176.4 (2019), pp. 928–943.
- [Sch84] C. Schütt. "Entropy numbers of diagonal operators between symmetric Banach spaces". In: *Journal of approximation theory* 40.2 (1984), pp. 121–128.
- [Sol+15] J. Solomon et al. "Convolutional Wasserstein distances: Efficient optimal transportation on geometric domains". In: ACM Transactions on Graphics (TOG) 34.4 (2015), p. 66.
- [Sol+16] J. Solomon et al. "Entropic metric alignment for correspondence problems". In: ACM Trans. Graph. 35.4 (2016), 72:1–72:13.
- [SSB+02] B. Schölkopf, A. J. Smola, F. Bach, et al. *Learning with kernels: support vec*tor machines, regularization, optimization, and beyond. MIT press, 2002.
- [SX21] J. W. Siegel and J. Xu. "Sharp Bounds on the Approximation Rates, Metric Entropy, and *n*-widths of Shallow Neural Networks". In: *arXiv preprint arXiv:2101.12365* (2021).

- [TGR21] W. Torous, F. Gunsilius, and P. Rigollet. "An Optimal Transport Approach to Causal Inference". In: *arXiv preprint arXiv:2108.05858* (2021).
- [Ver18] R. Vershynin. *High-dimensional probability: An introduction with applications in data science.* Vol. 47. Cambridge university press, 2018.
- [Vil09] C. Villani. Optimal transport: old and new. Vol. 338. Springer, 2009.
- [VV12] A. Veeser and R. Verfürth. "Poincaré constants for finite element stars". In: IMA Journal of Numerical Analysis 32.1 (2012), pp. 30–47.
- [VV21] A. Vacher and F.-X. Vialard. "Convex transport potential selection with semidual criterion". In: *arXiv preprint arXiv:2112.07275* (2021).
- [VW96] A. W. Vaart and J. A. Wellner. "Weak convergence and empirical processes with applications to statistics". In: *Weak convergence and empirical processes*. Springer, 1996, pp. 16–28.
- [Wai19] M. J. Wainwright. *High-dimensional statistics: A non-asymptotic viewpoint*. Vol. 48. Cambridge University Press, 2019.
- [Yan+20a] K. D. Yang et al. "Predicting cell lineages using autoencoders and optimal transport". In: *PLoS computational biology* 16.4 (2020), e1007828.
- [Yan+20b] Z. Yang et al. "On function approximation in reinforcement learning: optimism in the face of large state spaces". In: Proceedings of the 34th International Conference on Neural Information Processing Systems. 2020, pp. 13903– 13916.
- [YZH22] L. Yang, Z. Zhang, and S. Hong. "Diffusion models: A comprehensive survey of methods and applications". In: *arXiv preprint arXiv:2209.00796* (2022).
- [Zho08] D.-X. Zhou. "Derivative reproducing properties for kernel methods in learning theory". In: Journal of computational and Applied Mathematics 220.1-2 (2008), pp. 456–463.