

A CLT FOR THE LSS OF LARGE DIMENSIONAL SAMPLE COVARIANCE MATRICES WITH DIVERGING SPIKES

BY ZHIJUN LIU^{*}, JIANG HU[†], ZHIDONG BAI[‡], AND HAIYAN SONG[§]

KLASMOE and School of Mathematics and Statistics, Northeast Normal University, China. ^{*}liuzj037@nenu.edu.cn; [†]huj156@nenu.edu.cn; [‡]baizd@nenu.edu.cn; [§]songhy716@nenu.edu.cn

In this paper, we establish the central limit theorem (CLT) for linear spectral statistics (LSS) of large-dimensional sample covariance matrix when the population covariance matrices are not uniformly bounded, which is a non-trivial extension of the Bai-Silverstein theorem (BST) (2004). The latter has strongly influenced the development of high-dimensional statistics, especially in applications of random matrix theory to statistics. However, the assumption of uniform boundedness of the population covariance matrices has seriously limited the applications of the BST. The aim of this paper is to remove the barriers for the applications of the BST. The new CLT, allows spiked eigenvalues to exist, which may be bounded or tend to infinity. An important feature of our result is that the roles of either spiked eigenvalues or the bulk eigenvalues predominate in the CLT, depending on which variance is nonnegligible in the summation of the variances. The CLT for LSS is then applied to compare four linear hypothesis tests: The Wilk's likelihood ratio test, the Lawly-Hotelling trace test, the Bartlett-Nanda-Pillai trace test, and Roy's largest root test. We also derive and analyze their power function under particular alternatives.

1. Introduction. We consider the general sample covariance matrix $\mathbf{B}_n = \frac{1}{n} \mathbf{T}_p \mathbf{X}_n \mathbf{X}_n^* \mathbf{T}_p^*$, where \mathbf{X}_n is a $p \times n$ matrix with independent and identically distributed (i.i.d.) standardized entries, \mathbf{T}_p is a $p \times p$ deterministic matrix, $\mathbf{T}_p \mathbf{X}_n$ is considered a random sample from the population with the population covariance matrix $\mathbf{T}_p \mathbf{T}_p^* = \mathbf{\Sigma}$, and $*$ represents the complex conjugate transpose. In the sequel, we simply write $\mathbf{B} \equiv \mathbf{B}_n$, $\mathbf{T} \equiv \mathbf{T}_p$ and $\mathbf{X} \equiv \mathbf{X}_n$ when there is no confusion. $\lambda_1, \lambda_2, \dots, \lambda_p$ denotes the eigenvalues of \mathbf{B} . For a known test function f , we call $\sum_{j=1}^p f(\lambda_j)$ the linear spectral statistic (LSS) of \mathbf{B} . As most of the classical test statistics in multivariate statistical analysis are associated with the eigenvalues of sample covariance matrices, LSSs are remarkable tools in many statistical problems (see [Anderson \(2003\)](#); [Yao et al. \(2015\)](#) for details). By extensively studying high-dimensional data, it was found that the performance of LSSs is significantly different between low dimensions and high-dimensional data. For example, under the low-dimensional setting, Wilks' theorem (see [Wilks \(1938\)](#)) provides the χ^2 approximation for the likelihood ratio test (LRT) statistics, which is a kind of LSS. However, when p is large compared with the sample size n , the LRTs have Gaussian fluctuations (see [Bai et al. \(2009\)](#); [Jiang and Yang \(2013\)](#)). More generally, [Bai and Silverstein \(2004\)](#) proved the central limit theorem for the LSSs of high-dimensional \mathbf{B} under very mild conditions by using random matrix theory (RMT). We refer to this result as the Bai-Silverstein theorem (BST) for brevity. Following the development of [Bai and Silverstein \(2004\)](#), there have been many extensions under different settings. [Pan and Zhou \(2008\)](#) generalized the BST by removing the constraint on the fourth moment of the underlying random variables. [Zheng \(2012\)](#) and [Yang and Pan \(2015\)](#) extended the BST to multivariate F matrices and canonical correlation matrices, respectively. [Pan \(2014\)](#) showed the CLT of the LSS for noncentered sample covariance matrices, and [Zheng et al.](#)

MSC2020 subject classifications: Primary 60B20; secondary 60F05.

Keywords and phrases: Empirical spectral distribution, linear spectral statistic, random matrix, Stieltjes transform.

(2015) studied the unbiased sample covariance matrix when the population mean is unknown. Chen and Pan (2015) focused on the ultrahigh dimensional case in which the dimension p is much larger than the sample size n . Gao et al. (2017) and Li et al. (2021) studied the CLT for the LSSs of the high-dimensional Spearman correlation and Kendall's rank correlation matrices, respectively. Without attempting to be comprehensive, we refer readers to other extensions (Bai et al., 2007, 2015, 2019; Zheng et al., 2019; Banna et al., 2020; Najim and Yao, 2016; Baik et al., 2018; Hu et al., 2019; Jiang and Bai, 2021).

Almost all the theories mentioned above have traditionally assumed that the spectral norms of Σ are bounded in n . This assumption severely limits their applications in data analysis because in many fields, such as economics and wireless communication networks, the leading eigenvalues may tend to infinity. We present two examples here.

- **Signal detection**(Johnstone and Nadler (2017)): We consider a single signal model

$$\mathbf{x} = \chi_s^{1/2} u \mathbf{h} + \sigma \mathbf{v},$$

where \mathbf{h} is an unknown p -dimensional vector, u is a random variable distributed as $N(0, 1)$, χ_s is the signal strength, σ is the noise level, and \mathbf{v} is a random noise vector that is independent of u and distributed as a multivariate Gaussian $N_p(0, \Sigma_v)$. It is easy to check that the covariance matrix of \mathbf{x} is $\Sigma_x = \sigma^2 \Sigma_v + \chi_s \mathbf{h} \mathbf{h}'$. When the noise level is low, while the signal strength is large and sometimes tends to infinity, it is illogical to assume the boundedness of Σ_x .

- **Factor model** (Bai and Ng (2002)): Many economic analyses such as arbitrage pricing theory and the rank of a demand system, fit naturally into the framework of factor model:

$$\underset{(N \times 1)}{\mathbf{x}_t} = \underset{(N \times r)}{\mathbf{\Lambda}} \underset{(r \times 1)}{\mathbf{f}_t} + \underset{(N \times 1)}{\mathbf{e}_t}$$

where \mathbf{x}_t is the observed data, and \mathbf{f}_t , $\mathbf{\Lambda}$ and \mathbf{e}_t represent the true common factors, the factor loadings and the idiosyncratic error term, respectively. Normalizing the covariance matrix of \mathbf{f}_t to be an identity matrix, we find that $\Sigma_x = \mathbf{\Lambda} \mathbf{\Lambda}' + \Sigma_e$, where Σ_x and Σ_e are the covariance matrices of \mathbf{x}_t and \mathbf{e}_t , respectively. A commonly used estimator of Σ_x is $\hat{\Sigma}_x = (1/T) \sum_{t=1}^T (\mathbf{x}_t - \bar{\mathbf{x}}_t)(\mathbf{x}_t - \bar{\mathbf{x}}_t)'$, where T is a large time dimension. Inference on the true common factors and the factor loadings can be based on the eigenvalues of $\hat{\Sigma}_x$. When N is large, the first r large eigenvalues of Σ_x can diverge, but the $(r + 1)$ th eigenvalue can be bounded; see Chamberlain and Rothschild (1983).

For these reasons, it is of practical value to obtain the asymptotic properties of the LSS when Σ is unbounded. Therefore, in this paper, we focus on the CLT for the LSS of a general covariance matrix structure

$$(1.1) \quad \Sigma = \mathbf{V} \begin{pmatrix} \mathbf{D}_1 & 0 \\ 0 & \mathbf{D}_2 \end{pmatrix} \mathbf{V}^*,$$

where \mathbf{V} is a unitary matrix, \mathbf{D}_1 is a diagonal matrix consisting of the descending unbounded eigenvalues, and \mathbf{D}_2 is the diagonal matrix of the bounded eigenvalues. The setting (1.1) is attributed to the famous spiked model in which a few large eigenvalues of the population covariance matrix are assumed to be well separated from the rest of eigenvalues (Johnstone, 2001). The spiked model has provided the foundations for a rich theory of principal component analysis through the performance of extreme eigenvalues, as discussed in Baik and Silverstein (2006); Paul (2007); Bai and Yao (2008); Nadler (2008); Jung and Marron (2009); Bai and Yao (2012); Onatski et al. (2014); Bloemendal et al. (2016); Wang and Yao (2017); Donoho et al. (2018); Perry et al. (2018); Johnstone and Paul (2018); Yang and Johnstone (2018); Yao et al. (2018); Dobriban (2020); Johnstone and Onatski

(2020); Cai et al. (2020); Jiang and Bai (2021). There are also several works in which the asymptotics of various quantities are considered as the spike strengths tend to infinity. Specifically, Zhou and Marron (2015) focused on the consistency of the sample eigenvector, corresponding to the largest eigenvalue of the sample covariance matrix, under high dimension and low sample size settings, when Σ is unbounded and the data set is Gaussian. Wang and Fan (2017) derived the asymptotic distributions of the spiked eigenvalues and eigenvectors under a generalized regime, which assumes $p > n$, and $p/(n\alpha_i)$ is bounded for all the spikes α_i as p, n tend to infinity. Recently, Li et al. (2020), Yin (2021) and Zhang et al. (2022) investigated the trace of the large sample covariance matrix with the spiked model assumption.

We highlight the main contributions of the present paper. First, we prove a nontrivial extension of the BST to the case in which the population covariance matrices are unbounded in the spectral norm. In particular, we show how the test functions and the divergence rate of the population spectral norm affect the new CLT. Second, it is known that Gaussian-like moments, i.e., the first four moments, coincide with a standard Gaussian distribution, or the diagonality of the population covariance matrix is necessary for the CLT of the LSSs (e.g., Zheng et al. (2015)). However, we prove that these restrictions can be completely removed by normalization. More importantly, even if the limit of the LSS variance does not exist, the normalized CLT can still hold. Third, technically, the entire paper is built on the decomposition of the LSS, which is split into two parts, an unbounded part and a bounded part. The bounded part contains bounded spikes. The bounded part of the LSS cannot be studied using the result in Bai and Silverstein (2004) directly since it is not an LSS of some sample covariance matrix. It approximates the LSS of a bounded sample covariance matrix but we need to estimate the difference between them. To address this challenge, we make full use of RMT and estimate the limit of that difference. Moreover, we also estimate the difference between the corresponding center term of LSS based on the sample covariance matrix and the bounded covariance matrix. In addition, we prove that the LSSs of the unbounded and bounded parts are asymptotically independent.

As an application, the established CLT is employed to study the asymptotic behavior of four linear hypothesis tests: The Wilk's likelihood ratio test, the Lawly-Hotelling trace test, the Bartlett-Nanda-Pillai trace test, and the Roy's largest root test, under the hypothesis

$$H_0 : \Sigma = \mathbf{I}_p \text{ v.s. } H_1 : \Sigma \neq \mathbf{I}_p.$$

The Wilk's likelihood ratio, the Lawly-Hotelling trace, and the Bartlett-Nanda-Pillai trace tests are typical examples of LSSs with test functions $f(x) = \log(x+1)$, x and $\frac{x}{1+x}$. In this paper, we start from a different perspective by studying the asymptotic distribution of four linear hypothesis tests under certain alternatives. Then we establish and compare their power functions under these alternatives. Roughly speaking, the Lawly-Hotelling trace test and Roy's largest root test are relatively better than the other two tests.

The remaining sections are organized as follows: Section 2 contains a detailed description of our notation and assumptions. The main results for the CLT for the LSS of the sample covariance matrix are stated in Section 3. In Section 4, we explore an application of our main results. Technical proofs of the theorems are presented in Section 5. One extra section of supplementary material completes this paper. It contains some postponed proofs and derivations of results found in Section 3 and Section 4.

2. Notation and assumptions. Throughout the paper, we use bold capital letters and bold lowercase letters to represent matrices and vectors, respectively. Scalars are often in regular letters. e_i denotes a standard basis vector whose components are all zero, except the i -th component, which equals 1. We use $\text{tr}(\mathbf{A})$, \mathbf{A}' and \mathbf{A}^* to denote the trace, transpose and conjugate transpose of matrix \mathbf{A} , respectively. We also use f' to denote the derivative of

function f , and we use $\frac{\partial}{\partial z_1} f(z_1, z_2)$ to denote the partial derivative of function f with respect to z_1 ; Let $[\mathbf{A}]_{ij}$ denote the (i, j) -th entry of the matrix \mathbf{A} and $\oint_{\mathcal{C}} f(z) dz$ denote the contour integral of $f(z)$ on the contour \mathcal{C} . Let $\lambda_i^{\mathbf{A}}$ denote the i th largest eigenvalue of matrix \mathbf{A} . Weak convergence is denoted by \xrightarrow{d} . Throughout this paper, we use $o(1)$ (resp. $o_p(1)$) to denote a negligible scalar (resp. in probability), and the notation C represents some generic constants that may vary from line to line.

Let $\mathbf{X} = (\mathbf{x}_1, \dots, \mathbf{x}_n) = (x_{ij})$, $1 \leq i \leq p$, $1 \leq j \leq n$; let \mathbf{T} be a $p \times p$ deterministic matrix and $\mathbf{\Sigma} = \mathbf{T}\mathbf{T}^*$. The spectrum of $\mathbf{\Sigma}$ is formed as $\rho_1 \geq \dots \geq \rho_p$. We define the singular value decomposition of \mathbf{T} as

$$(2.2) \quad \mathbf{T} = \mathbf{V}\mathbf{D}^{1/2}\mathbf{U}^* = \mathbf{V} \begin{pmatrix} \mathbf{D}_1^{\frac{1}{2}} & 0 \\ 0 & \mathbf{D}_2^{\frac{1}{2}} \end{pmatrix} \mathbf{U}^*$$

where \mathbf{U} and \mathbf{V} are unitary matrices, and $\mathbf{D}_1 = \text{diag}(\alpha_1, \dots, \alpha_1, \alpha_2, \dots, \alpha_2, \dots, \alpha_K, \dots, \alpha_K)$ is a diagonal matrix of the spiked eigenvalues for which the components tend to infinity. To avoid confusion, we call them 'remote spikes'. \mathbf{D}_2 is the diagonal matrix of the eigenvalues with the bounded components, including bounded spiked eigenvalues and bulk eigenvalues. $\alpha_1 > \dots > \alpha_K$ denotes the remote spiked eigenvalues of $\mathbf{\Sigma}$ with the multiplicity $d_k, k = 1, \dots, K$, and $d_1 + \dots + d_K = M$. Moreover, let $\rho_i = \alpha_k$ if $i \in J_k$, where $J_k = \{j_k + 1, \dots, j_k + d_k\}$ is the set of ranks of the eigenvalue α_k with multiplicities d_k . Then, the corresponding sample covariance matrix

$$\mathbf{B} = \frac{1}{n} \mathbf{T}\mathbf{X}\mathbf{X}^*\mathbf{T}^*$$

is the so-called generalized spiked sample covariance matrix. $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_p$ denotes the eigenvalues of \mathbf{B} . Corresponding to the decomposition of \mathbf{D} , we decompose $\mathbf{V} = (\mathbf{V}_1, \mathbf{V}_2)$, $\mathbf{U} = (\mathbf{U}_1, \mathbf{U}_2)$, and $\mathbf{\Gamma} = \mathbf{V}_2\mathbf{D}_2^{1/2}\mathbf{U}_2^*$, $\mathbf{r}_j = \frac{1}{\sqrt{n}}\mathbf{\Gamma}\mathbf{x}_j$, $\mathbf{A}_j = \mathbf{B} - z\mathbf{I} - \mathbf{r}_j\mathbf{r}_j^*$. Let \mathbb{E}_j be the conditional expectation with respect to the σ -field generated by $\mathbf{r}_1, \dots, \mathbf{r}_j$. For any matrix \mathbf{A} with real eigenvalues, the empirical spectral distribution of \mathbf{A} is denoted by

$$F^{\mathbf{A}}(x) = \frac{1}{p} (\text{number of eigenvalues of } \mathbf{A} \leq x).$$

For any function of bounded variation F on the real line, its Stieltjes transform is defined by

$$m_F(z) = \int \frac{1}{\lambda - z} dF(\lambda), \quad z \in \mathbb{C}^+ := \{z \in \mathbb{C} : \Im z > 0\}.$$

The assumptions used in the results of this paper are as follows:

ASSUMPTION 1. $\{x_{ij}, 1 \leq i \leq p, 1 \leq j \leq n\}$ are independent random variables with common moments

$$\mathbb{E}x_{ij} = 0, \quad \mathbb{E}|x_{ij}|^2 = 1, \quad \beta_x = \mathbb{E}|x_{ij}|^4 - |\mathbb{E}x_{ij}^2|^2 - 2, \quad \alpha_x = |\mathbb{E}x_{ij}^2|^2,$$

and satisfy the following Lindeberg-type condition:

$$\frac{1}{np} \sum_{i=1}^p \sum_{j=1}^n \mathbb{E} \left\{ |x_{ij}|^4 \mathbb{1}_{\{|x_{ij}| \geq \eta\sqrt{n}\}} \right\} \rightarrow 0, \quad \text{for any constant } \eta > 0.$$

ASSUMPTION 2. As $\min\{p, n\} \rightarrow \infty$, the ratio of the dimension-to-sample size (RDS) $c_n := p/n \rightarrow c > 0$.

REMARK 2.1. *Assumptions 1-2 are standard in RMT. If $\mathbb{E}x_{ij} \neq 0$, we can use the centralized sample covariance matrices and $n - 1$ instead of \mathbf{B}_n and n , respectively, and the following results also hold. Details can be found in (Zheng et al., 2015). Therefore, in this sequel, we assume $\mathbb{E}x_{ij} = 0$ without loss of generality.*

ASSUMPTION 3. \mathbf{T} is nonrandom. As $\min\{p, n\} \rightarrow \infty$, $\alpha_K \rightarrow \infty$ and $H_n := F^{\mathbf{T}\Gamma^*} \xrightarrow{d} H$, where H is a distribution function on the real line. M is fixed.

REMARK 2.2. *It was shown by Silverstein (1995) that under Assumptions 1-3, $F^{\mathbf{B}} \xrightarrow{d} F^{c,H}$ almost surely, where $F^{c,H}$ is a nonrandom distribution function whose Stieltjes transform $m := m_{F^{c,H}}(z)$ satisfies equation*

$$(2.3) \quad m = \int \frac{1}{t(1 - c - czm) - z} dH(t).$$

In this sequel, we call $F^{c,H}$ the limiting spectral distribution (LSD) of \mathbf{B} . Moreover, as the matrix $\underline{\mathbf{B}} = \frac{1}{n} \mathbf{X}^ \mathbf{T}^* \mathbf{T} \mathbf{X}$ shares the same nonzero eigenvalues with \mathbf{B} , equation (2.3) can be rewritten as*

$$\underline{m} = - \left(z - c \int \frac{t}{1 + t\underline{m}} dH(t) \right)^{-1},$$

where $\underline{m} := m_{\underline{F}^{c,H}}(z)$ represents the Stieltjes transform of the LSD of $\underline{\mathbf{B}}$.

ASSUMPTION 4. Test functions f_1, \dots, f_h are analytic on a connected open region of the complex plane containing the support of F^{c_n, H_n} for almost all n . Moreover, we suppose that for any $l = 1, \dots, h$,

$$\lim_{\substack{\{x_n, y_n\} \rightarrow \infty \\ x_n/y_n \rightarrow 1}} \frac{f'_l(x_n)}{f'_l(y_n)} = 1.$$

REMARK 2.3. *In fact, Assumption 4 is not too restrictive for application, many common functions such as logarithmic and polynomial functions satisfy it. However, it is worth noting that the exponential function may not satisfy this assumption.*

ASSUMPTION 5. \mathbf{T} is real or the variables x_{ij} are complex satisfying $\alpha_x = 0$.

ASSUMPTION 6. $\mathbf{T}^* \mathbf{T}$ is diagonal or $\beta_x = 0$.

REMARK 2.4. *When x_{ij} are from a complex Gaussian distribution, then $\alpha_x = 0$ and $\beta_x = 0$, which are referred to as Gaussian-like moment conditions. Assumption 5 and Assumption 6 are used as a replacement for Gaussian moment conditions. It is proved by Zheng et al. (2015) that these two structural assumptions on population matrices are necessary for their results when the Gaussian-like moment conditions in the BST do not hold. However, we will show in the next section that these two assumptions are not needed by normalizing the LSS.*

3. Main results. Now, we are in a position to present our main theorems, and their proofs are provided in Section 5 and the supplementary material. Note that

$$\sum_{j=1}^p f(\lambda_j) = p \int f(x) dF^{\mathbf{B}}(x).$$

Thus, for brevity, we define the normalized LSSs as

$$Y_l = \int f_l(x) dG_n(x) - \sum_{k=1}^K d_k f_l(\phi_n(\alpha_k)) - \frac{M}{2\pi i} \oint_{\mathcal{C}} f_l(z) \frac{\underline{m}'(z)}{\underline{m}(z)} dz, \quad l = 1, 2, \dots, h,$$

where

$$G_n(x) = p[F^{\mathbf{B}}(x) - F^{c_n, H_n}(x)], \quad \phi_n(x) = x \left(1 + c_n \int \frac{t}{x-t} dH_n(t) \right),$$

and F^{c_n, H_n} is the LSD $F^{c, H}$ with c, H replaced by c_n, H_n . $\underline{F}^{c, H}$ denotes the LSD of matrix $n^{-1} \mathbf{X}^* \mathbf{U}_2 \mathbf{D}_2 \mathbf{U}_2^* \mathbf{X}$, $\mathbf{U} = (\mathbf{U}_1, \mathbf{U}_2)$, $\mathbf{U}_1 = (u_{ij})_{i=1, \dots, p; j=1, \dots, M}$, $\phi_k = \phi(x) |_{x=\alpha_k} = \alpha_k \left(1 + c \int \frac{t}{\alpha_k - t} dH(t) \right)$,

$$b_n(z) = \frac{1}{1 + n^{-1} \mathbb{E} \text{tr} \mathbf{\Gamma} \mathbf{\Gamma}^* \mathbf{A}_j^{-1}(z)}, \quad \mathcal{U}_{i_1 j_1 i_2 j_2} = \sum_{t=1}^p \bar{u}_{t i_1} u_{t j_1} u_{t i_2} \bar{u}_{t j_2},$$

$$\theta_k = \phi_k^2 \underline{m}_2(\phi_k), \quad \nu_k = \phi_k^2 \underline{m}^2(\phi_k),$$

$$\underline{m}(\lambda) = \int \frac{1}{x - \lambda} d\underline{F}^{c, H}(x), \quad \underline{m}_2(\lambda) = \int \frac{1}{(\lambda - x)^2} d\underline{F}^{c, H}(x),$$

$$c_{nM} = \frac{p - M}{n}, \quad H_{2n} = F^{\mathbf{D}_2},$$

$$\mathbf{P}_n(z) = ((1 - c_{nM}) \mathbf{\Gamma} \mathbf{\Gamma}^* - z c_{nM} m_{2n0}(z) \mathbf{\Gamma} \mathbf{\Gamma}^* - z \mathbf{I}_p)^{-1}.$$

Here, $m_{2n0}(z)$ is the Stieltjes transform of $F^{c_{nM}, H_{2n}}$ and $\underline{m}_{2n0}(z) = -\frac{1 - c_{nM}}{z} + c_{nM} m_{2n0}(z)$. For clarification purposes, $m_{1n0}(z)$ denotes the Stieltjes transform of F^{c_n, H_n} , $m_n = \frac{1}{p} \text{tr}(\mathbf{B} - z \mathbf{I}_p)^{-1}$, and $m_{2n} = \frac{1}{p - M} \text{tr}(\mathbf{S}_{22} - z \mathbf{I}_{p - M})^{-1}$, which is used later in the proof.

We first establish a CLT of the LSS without any restrictions imposed on the Gaussian moments or on the structures of the population covariance matrix by normalizing the LSS.

THEOREM 3.1. *Under Assumptions 1–4, we have*

$$\frac{Y_1 - \mu_1}{\sqrt{\varsigma_1^2}} \xrightarrow{d} N(0, 1),$$

where the mean function is

$$\mu_1 = -\frac{\alpha_x}{2\pi i} \cdot \oint_{\mathcal{C}} \frac{c_{nM} f_1(z) \int \underline{m}_{2n0}^3(z) t^2 (1 + t \underline{m}_{2n0}(z))^{-3} dH_{2n}(t)}{\left(1 - c_{nM} \int \frac{\underline{m}_{2n0}^2(z) t^2}{(1 + t \underline{m}_{2n0}(z))^2} dH_{2n}(t) \right) \left(1 - \alpha_x c_{nM} \int \frac{\underline{m}_{2n0}^2(z) t^2}{(1 + t \underline{m}_{2n0}(z))^2} dH_{2n}(t) \right)} dz$$

(3.4)

$$- \frac{\beta_x}{2\pi i} \cdot \oint_{\mathcal{C}} \frac{c_{nM} f_1(z) \int \underline{m}_{2n0}^3(z) t^2 (1 + t \underline{m}_{2n0}(z))^{-3} dH_{2n}(t)}{1 - c_{nM} \int \underline{m}_{2n0}^2(z) t^2 (1 + t \underline{m}_{2n0}(z))^{-2} dH_{2n}(t)} dz,$$

and the covariance function is

$$\varsigma_1^2 = \sum_{k=1}^K \frac{\phi_n^2(\alpha_k)}{n} (f_1'(\phi_n(\alpha_k)))^2 s_k^2 - \frac{1}{4\pi^2} \oint_{\mathcal{C}_1} \oint_{\mathcal{C}_2} f_1(z_1) f_1(z_2) \vartheta_n^2 dz_1 dz_2.$$

Here, $s_k^2 = \frac{\sum_{j \in J_k} ((\alpha_x + 1)\theta_k + \beta_x \mathcal{U}_{jjj} \nu_k) + \sum_{j_1 \neq j_2} \beta_x \mathcal{U}_{j_1 j_1 j_2 j_2} \nu_k}{\theta_k^2}$, $\vartheta_n^2 = \Theta_{0,n}(z_1, z_2) + \alpha_x \Theta_{1,n}(z_1, z_2) + \beta_x \Theta_{2,n}(z_1, z_2)$, where

$$\begin{aligned}\Theta_{0,n}(z_1, z_2) &= \frac{\underline{m}'_{2n0}(z_1) \underline{m}'_{2n0}(z_2)}{(\underline{m}_{2n0}(z_1) - \underline{m}_{2n0}(z_2))^2} - \frac{1}{(z_1 - z_2)^2}, \\ \Theta_{1,n}(z_1, z_2) &= \frac{\partial}{\partial z_2} \left\{ \frac{\partial \mathcal{A}_n(z_1, z_2)}{\partial z_1} \frac{1}{1 - \alpha_x \mathcal{A}_n(z_1, z_2)} \right\}, \\ \mathcal{A}_n(z_1, z_2) &= \frac{z_1 z_2}{n} \underline{m}'_{2n0}(z_1) \underline{m}'_{2n0}(z_2) \text{tr} \mathbf{\Gamma}^* \mathbf{P}_n(z_1) \mathbf{\Gamma} \mathbf{\Gamma}' \mathbf{P}_n(z_2)' \bar{\mathbf{\Gamma}}, \\ \Theta_{2,n}(z_1, z_2) &= \frac{z_1^2 z_2^2 \underline{m}'_{2n0}(z_1) \underline{m}'_{2n0}(z_2)}{n} \sum_{i=1}^p [\mathbf{\Gamma}^* \mathbf{P}_n^2(z_1) \mathbf{\Gamma}]_{ii} [\mathbf{\Gamma}^* \mathbf{P}_n^2(z_2) \mathbf{\Gamma}]_{ii}.\end{aligned}$$

In the equation above, $\mathcal{C}, \mathcal{C}_1$ and \mathcal{C}_2 are closed contours in the complex plane enclosing the support of $F^{c,H}$, and \mathcal{C}_1 and \mathcal{C}_2 are nonoverlapping. It is worth noting that $\Theta_{1,n}(z_1, z_2)$ and $\Theta_{2,n}(z_1, z_2)$ may not converge.

As a minor price for the removal of the bounded spectrum condition, the new CLT described above only applies to a single LSS. To guarantee that the new CLT applies to multiple normalized LSSs, structural assumptions about the population covariance matrices are needed (Assumptions 5 and 6). The following theorem is a nontrivial extension of the BST:

THEOREM 3.2. *Under Assumptions 1–6, the random vector*

$$\left(\frac{Y_1 - \mu_1}{\sqrt{\sigma_1^2}}, \dots, \frac{Y_h - \mu_h}{\sqrt{\sigma_h^2}} \right)' \xrightarrow{d} N_h(0, \mathbf{\Psi}),$$

with mean function

$$\mu_l = -\frac{\alpha_x}{2\pi i} \cdot \oint_{\mathcal{C}} f_l(z) \frac{c_{nM} \int \underline{m}_{2n0}^3(z) t^2 (1 + t \underline{m}_{2n0}(z))^{-3} dH_{2n}(t)}{\left(1 - c_{nM} \int \frac{\underline{m}_{2n0}^2(z) t^2}{(1 + t \underline{m}_{2n0}(z))^2} dH_{2n}(t)\right) \left(1 - \alpha_x c_{nM} \int \frac{\underline{m}_{2n0}^2(z) t^2}{(1 + t \underline{m}_{2n0}(z))^2} dH_{2n}(t)\right)} dz$$

(3.5)

$$-\frac{\beta_x}{2\pi i} \cdot \oint_{\mathcal{C}} f_l(z) \frac{c_{nM} \int \underline{m}_{2n0}^3(z) t^2 (1 + t \underline{m}_{2n0}(z))^{-3} dH_{2n}(t)}{1 - c_{nM} \int \underline{m}_{2n0}^2(z) t^2 (1 + t \underline{m}_{2n0}(z))^{-2} dH_{2n}(t)} dz, \quad l = 1, \dots, h,$$

variance function

$$\sigma_l^2 = \sum_{k=1}^K \frac{\phi_n^2(\alpha_k)}{n} (f'_l(\phi_n(\alpha_k)))^2 s_k^2 + \kappa_{ll}, \quad l = 1, \dots, h,$$

and covariance matrix $\mathbf{\Psi} = (\psi_{st})_{h \times h}$,

$$\psi_{st} = \frac{\sum_{k=1}^K \varpi_s^k \varpi_t^k s_k^2 + \kappa_{st}}{\sqrt{\sum_{k=1}^K (\varpi_s^k)^2 s_k^2 + \kappa_{ss}} \sqrt{\sum_{k=1}^K (\varpi_t^k)^2 s_k^2 + \kappa_{tt}}},$$

where

$$\kappa_{st} = -\frac{1}{4\pi^2} \oint_{\mathcal{C}_1} \oint_{\mathcal{C}_2} \frac{f_s(z_1) f_t(z_2)}{(\underline{m}(z_1) - \underline{m}(z_2))^2} d\underline{m}(z_1) d\underline{m}(z_2) - \frac{c\beta_x}{4\pi^2} \oint_{\mathcal{C}_1} \oint_{\mathcal{C}_2} f_s(z_1) f_t(z_2)$$

$$\begin{aligned} & \times \left[\int \frac{t^2}{(\underline{m}(z_1)t+1)^2(\underline{m}(z_2)t+1)^2} dH(t) \right] d\underline{m}(z_1) d\underline{m}(z_2) \\ & - \frac{1}{4\pi^2} \oint_{\mathcal{C}_1} \oint_{\mathcal{C}_2} f_s(z_1) f_t(z_2) \left[\frac{\partial^2}{\partial z_1 \partial z_2} \log(1-a(z_1, z_2)) \right] dz_1 dz_2, \\ a(z_1, z_2) &= \alpha_x \left(1 + \frac{\underline{m}(z_1)\underline{m}(z_2)(z_1-z_2)}{\underline{m}(z_2)-\underline{m}(z_1)} \right). \end{aligned}$$

Note that s_k^2 is defined in Theorem 3.1, and $\varpi_l^k = \frac{\phi_n(\alpha_k)}{\sqrt{n}} f_l'(\phi_n(\alpha_k))$ is allowed to be infinity.

REMARK 3.1. If we rewrite the covariance matrix Ψ in Theorem 3.2 in expressions with n , such as ς_1^2 in Theorem 3.1, then Assumptions 5-6 should be removed analogously. However, in this case, we need to verify the nonsingularity of these covariance matrices. It is obvious that Ψ is singular if the set of test functions are linearly dependent. However, it is not easy to see whether or not Ψ is invertible when the test functions are totally linearly independent. For not lengthening the paper too much, we decide not to pursue that direction in this paper.

REMARK 3.2. From the theorem, it is not difficult to find that the asymptotic distributions of the LSSs depend on the divergence rates of $\phi_n(\alpha_k)$ and $f_l'(\phi_n(\alpha_k))$. In particular, when $\frac{\phi_n(\alpha_1)}{\sqrt{n}} f_l'(\phi_n(\alpha_1)) \rightarrow 0$ ($n \rightarrow \infty$), Theorem 3.2 reduces to Theorem 2.1 in Zheng et al. (2015); when $\frac{\phi_n(\alpha_1)}{\sqrt{n}} f_l'(\phi_n(\alpha_1)) \rightarrow \infty$ ($n \rightarrow \infty$), Theorem 3.2 is a nontrivial extension of the CLT derived by Jiang and Bai (2021). Furthermore, Yin (2021) recently obtained a CLT when the test functions were polynomial.

REMARK 3.3. If $\mathbf{D}_2 = \mathbf{I}_{p-M}$, then the mean function μ_1 and κ_{st} in the covariance function of Theorem 3.2 can be simplified from the results in Wang and Yao (2013) and Zheng et al. (2015), i.e.,

$$\begin{aligned} \phi_n(x) &= x + \frac{x(p-M)}{n(x-1)}, \quad \mu_1 = \alpha_x I_1(f_1) + \beta_x I_2(f_1), \\ \kappa_{st} &= (\alpha_x + 1) J_1(f_s, f_t) + \beta_x J_2(f_s, f_t), \\ I_1(f_1) &= \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} f_1 \left(|1 + \sqrt{c}z|^2 \right) \left[\frac{z}{z^2 - r^{-2}} - \frac{1}{z} \right] dz, \\ I_2(f_1) &= \frac{1}{2\pi i} \oint_{|z|=1} f_1 \left(|1 + \sqrt{c}z|^2 \right) \frac{1}{z^3} dz, \\ J_1(f_s, f_t) &= \lim_{r \downarrow 1} \frac{-1}{4\pi^2} \oint_{|z_1|=1} \oint_{|z_2|=1} \frac{f_s \left(|1 + \sqrt{c}z_1|^2 \right) f_t \left(|1 + \sqrt{c}z_2|^2 \right)}{(z_1 - rz_2)^2} dz_1 dz_2, \\ J_2(f_s, f_t) &= -\frac{1}{4\pi^2} \oint_{|z_1|=1} \frac{f_s \left(|1 + \sqrt{c}z_1|^2 \right)}{z_1^2} dz_1 \oint_{|z_2|=1} \frac{f_t \left(|1 + \sqrt{c}z_2|^2 \right)}{z_2^2} dz_2. \end{aligned}$$

4. Application. Linear hypothesis testing plays an important role in the analysis of multivariate data. Four criteria are used to test linear hypotheses: the likelihood ratio criterion, the Lawley-Hotelling trace criterion, the Bartlett-Nanda-Pillai trace criterion, and Roy's largest root criterion, which are also referred to as: the Wilk's U , the Lawley-Hotelling W , the Bartlett-Nanda-Pillai V and Roy's largest root R , respectively. They are defined as:

- Wilk's likelihood ratio $U = \sum_{i=1}^p \log(1 + \lambda_i)$
- Lawley-Hotelling trace $W = \sum_{i=1}^p \lambda_i$
- Bartlett-Nanda-Pillai trace $V = \sum_{i=1}^p \frac{\lambda_i}{1 + \lambda_i}$
- Roy's largest root $R = \lambda_1$

where $\lambda_i, i = 1, \dots, p$ are the eigenvalues of an F matrix. The four statistics are all functions of the eigenvalues of an F matrix under a certain transformation. The four statistics can be classified into two categories. The first category can be expressed as linear spectral statistics and the second category is based on extreme eigenvalues, such as the largest or smallest eigenvalues. Obviously, the first category includes U, W, V , corresponding to test function $f_U(x) = \log(x + 1), f_W(x) = x, f_V(x) = \frac{x}{1+x}$. The second category includes R . For more details, we refer readers to read [Anderson \(2003\)](#).

In practical applications, people are most concerned about which test has the best effect. To select the best one among the four tests above, one needs to know their asymptotic power under alternatives. In this section, we apply Theorem 3.1 to obtain the asymptotic distribution of U, W, V under different alternative settings, and we also provide their approximated power functions.

The settings for the population covariance matrices under alternatives are as follows:

$$H_1: \Sigma = \text{diag}(\alpha_1, \underbrace{1, 1, \dots, 1}_{p-1}).$$

$$H_2: \Sigma = \text{diag}(\alpha_1, \alpha_2, \underbrace{1, \dots, 1}_{p-2}).$$

$$H_3: \Sigma = \text{diag}(1 + a/n, \dots, 1 + a/n).$$

4.1. *Testing the hypothesis when the alternative is a spiked model: H_1 and H_2 .* In this subsection, we consider the hypothesis $H_0 = \mathbf{I}_p$ versus the alternative population covariance as a spiked model, such as H_1 and H_2 above. Theorems 4.1–4.3 derive the asymptotic distribution of U, W, V under H_1 and H_2 , respectively. We also obtain and analyze the asymptotic power of U, W, V under certain alternatives. Moreover, the asymptotic distribution and power of R under the spiked model is established in [Hou et al. \(2019\)](#). Without loss of generality, we consider the trivial case $c < 1$.

For brevity, we introduce some notations. We define

$$\varrho(c) = \frac{c + \sqrt{c^2 + 4}}{2}, \quad \tilde{c} = \frac{4c}{(2 + c + \sqrt{c^2 + 4})^2}.$$

$$CT(x, c, \tilde{c}) = \log(1 + x) + \frac{-\left(\sqrt{c} - \frac{1}{\sqrt{c}}\right)^2 (\log(1 - \sqrt{\tilde{c}c}) + \sqrt{\tilde{c}c}) - \sqrt{\tilde{c}} (\sqrt{c} - (\sqrt{c})^3)}{1 - c}.$$

To avoid misunderstandings, we define the values of $\varrho(c_n), \varrho(c_{nM})$ to be the same as $\varrho(c)$ above with the substitution of c_n and c_{nM} for c in these quantities, respectively. The same substitution also holds for \tilde{c}_n .

THEOREM 4.1 (U statistics). *Under Assumptions 1–4 with $c_n = p/n \rightarrow c \in (0, 1)$, we have under H_1 or H_2 ,*

$$\frac{U - (p - M) \int f_U(x) dF^{c_{nM}, H_{2n}}(x) - \mu_U}{\sqrt{s_U^2}} \xrightarrow{d} N(0, 1),$$

where

$$\begin{aligned} \int f_U(x) dF^{c_{nM}, H_{2n}}(x) &= CT(\varrho(c_{nM}), c_{nM}, \tilde{c}_{nM}), \\ \mu_U &= \alpha_x I_1(f_U) + \beta_x I_2(f_U) + \sum_{k=1}^K d_k \log(1 + \phi_n(\alpha_k)) + M \log\left(1 - \sqrt{\tilde{c}_n c_n}\right), \\ \varsigma_U^2 &= \sum_{k=1}^K \frac{\phi_n^2(\alpha_k)}{n(1 + \phi_n(\alpha_k))^2} s_k^2 + (\alpha_x + 1) J_1(f_U, f_U) + \beta_x J_2(f_U, f_U), \\ I_1(f_U) &= \log(1 + \varrho(c_{nM})) - \log(2 + c_{nM}) + \sum_{k=1}^{\infty} \left(\frac{\sqrt{c_{nM}}}{2 + c_{nM}}\right)^{2k} \frac{(2k-1)!}{k!k!}, \\ I_2(f_U) &= -\sum_{k=1}^{\infty} \left(\frac{\sqrt{c_{nM}}}{2 + c_{nM}}\right)^{2k} \frac{(2k-1)!}{(k-1)!(k+1)!}, \\ J_1(f_U, f_U) &= J_2(f_U, f_U) = \left(\sum_{k=1}^{\infty} \left(\frac{\sqrt{c_{nM}}}{2 + c_{nM}}\right)^{2k-1} \frac{(2k-2)!}{k!(k-1)!}\right)^2. \end{aligned}$$

COROLLARY 4.1. *Under the same assumptions as in Theorem 4.1, we have*

$$P_U - \Phi\left(\frac{(p-M)CT(\varrho(c_{nM}), c_{nM}, \tilde{c}_{nM}) - pCT(\varrho(c_n), c_n, \tilde{c}_n) + A_1}{\varsigma_U} - z_\xi \frac{A_3}{\varsigma_U}\right) \rightarrow 0,$$

where P_U is the power function of Wilk's likelihood ratio test, Φ is the distribution of a standard normal variable, and z_ξ is the upper quantile of the of the standard normal distribution Φ at level ξ , and

$$\begin{aligned} A_1 &= \sum_{k=1}^K d_k \log(1 + \phi_n(\alpha_k)) + M \log\left(1 - \sqrt{\tilde{c}_n c_n}\right), \\ A_3 &= \sqrt{\alpha_x + \beta_x + 1} \sum_{k=1}^{\infty} \left(\frac{\sqrt{c_n}}{2 + c_n}\right)^{2k-1} \frac{(2k-2)!}{k!(k-1)!}. \end{aligned}$$

THEOREM 4.2 (W statistics). *Under Assumptions 1–4 with $c_n = p/n \rightarrow c \in (0, 1)$, we have under H_1 or H_2 ,*

$$\frac{W - (p-M) \int f_W(x) dF^{c_{nM}, H_{2n}}(x) - \mu_W}{\sqrt{\varsigma_W^2}} \xrightarrow{d} N(0, 1),$$

where

$$\begin{aligned} \int f_W(x) dF^{c_{nM}, H_{2n}}(x) &= 1, \quad \mu_W = \sum_{k=1}^K d_k \phi_n(\alpha_k) - M c_n, \\ \varsigma_W^2 &= \sum_{k=1}^K \frac{\phi_n^2(\alpha_k)}{n} s_k^2 + \alpha_x c_{nM} + \beta_x c_{nM} + c_{nM}. \end{aligned}$$

COROLLARY 4.2. *Under the same assumptions as in Theorem 4.2, we have*

$$P_W - \Phi \left(\frac{\sum_{k=1}^K d_k \phi_n(\alpha_k) - M c_n - M}{\varsigma_W} - z_\xi \frac{\sqrt{\alpha_x c_n + \beta_x c_n + c_n}}{\varsigma_W} \right) \rightarrow 0,$$

where P_W is the power function of Lawley-Hotelling trace test, Φ is the distribution of a standard normal variable, and z_ξ is the upper quantile of the of the standard normal distribution Φ at level ξ .

THEOREM 4.3 (V statistic). *Under Assumptions 1–4 with $c_n = p/n \rightarrow c \in (0, 1)$, we have under H_1 or H_2 ,*

$$\frac{V - (p - M) \int f_V(x) dF^{c_{nM}, H_{2n}}(x) - \mu_V}{\sqrt{\varsigma_V^2}} \xrightarrow{d} N(0, 1),$$

where

$$\begin{aligned} \int f_V(x) dF^{c_{nM}, H_{2n}}(x) &= \frac{1}{1 + \varrho(c_{nM})}, \\ \mu_V &= \alpha_x I_1(f_V) + \beta_x I_2(f_V) + \sum_{k=1}^K d_k \frac{\phi_n(\alpha_k)}{1 + \phi_n(\alpha_k)} - \frac{M(c_n - 2)}{(2 + c_n + \sqrt{c_n^2 + 4})(1 - \tilde{c}_n)} - \frac{M}{2}, \\ \varsigma_V^2 &= \sum_{k=1}^K \frac{\phi_n^2(\alpha_k)}{n(1 + \phi_n(\alpha_k))^4} s_k^2 + (\alpha_x + 1) J_1(f_V, f_V) + \beta_x J_2(f_V, f_V), \\ I_1(f_V) &= \frac{1}{2 + c_{nM}} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c_{nM}}}{2 + c_{nM}} \right)^{2k} \frac{(2k)!}{k!k!} - \\ &\quad \frac{1}{1 + \varrho(c_{nM})} \left(\frac{\tilde{c}_{nM}}{-(1 - \tilde{c}_{nM})^2} + \frac{1}{2(\sqrt{\tilde{c}_{nM}} - 1)^2} + \frac{1}{2(\sqrt{\tilde{c}_{nM}} + 1)^2} \right), \\ I_2(f_V) &= -\frac{1}{2 + c_{nM}} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c_{nM}}}{2 + c_{nM}} \right)^{2k} \frac{(2k)!}{(k-1)!(k+1)!}, \\ J_1(f_V, f_V) = J_2(f_V, f_V) &= \left(\frac{1}{2 + c_{nM}} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c_{nM}}}{2 + c_{nM}} \right)^{2k+1} \frac{(2k+1)!}{k!(k+1)!} \right)^2. \end{aligned}$$

COROLLARY 4.3. *Under the same assumptions as in Theorem 4.3, we have*

$$P_V - \Phi \left(\frac{\frac{(p-M)}{1+\varrho(c_{nM})} - \frac{p}{1+\varrho(c_n)} + \sum_{k=1}^K d_k \frac{\phi(\alpha_k)}{1+\phi(\alpha_k)} - \frac{M}{2} - \frac{(c_n-2)M}{2(1+\varrho(c_n))(1-\tilde{c}_n)}}{\varsigma_V} - z_\xi \frac{A_4}{\varsigma_V} \right) \rightarrow 0,$$

where P_V is the power function of Bartlett-Nanda-Pillai trace test, Φ is the distribution of a standard normal variable, and z_ξ is the upper quantile of the of the standard normal distribution Φ at level ξ , and

$$A_4 = \frac{\sqrt{\alpha_x + \beta_x + 1}}{2 + c_n} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c_n}}{2 + c_n} \right)^{2k+1} \frac{(2k+1)!}{k!(k+1)!}.$$

REMARK 4.1. For P_U , $(p - M)CT(\varrho(c_{nM}), c_{nM}, \tilde{c}_{nM}) - pCT(\varrho(c_n), c_n, \tilde{c}_n)$, $M \log(1 - \sqrt{\tilde{c}_n c_n})$, A_3 , and ς_U are all of constant order $O(1)$. When spiked eigenvalues α_k are of constant order $O(1)$, then A_1 is a constant, hence we can assert that P_U does not tend to 1 as n tends to infinity. When α_k tends to infinity, then A_1 tends to infinity, therefore we have P_U tends to 1.

For P_W , $\sqrt{\alpha_x c_n + \beta_x c_n + c_n}$ is of constant order $O(1)$. Hence when α_k is of constant order $O(1)$, then ς_W and $\sum_{k=1}^K d_k \phi_n(\alpha_k)$ are both constants, thus we can assert that P_W does not tend to 1 as n tends to infinity. When α_k tends to infinity, P_W tends to 1 as n tends to infinity. Notably, different orders of α_k lead to different speed of P_W . When α_k tends to infinity and the order of α_k is lower than or equal to $n^{1/2}$, then ς_W is a constant and $\sum_{k=1}^K d_k \phi_n(\alpha_k)$ tends to infinity. Therefore P_W tends to 1 of order $\phi(\alpha_k)$. When the order of α_k is higher than $n^{1/2}$, both $\sum_{k=1}^K d_k \phi_n(\alpha_k)$ and ς_W tend to infinity, of order $\phi(\alpha_k)$ and $\frac{\phi(\alpha_k)}{\sqrt{n}}$, respectively. Then P_W tends to 1 of order $n^{1/2}$. As a comparison, when the order of α_k is lower than or equal to $n^{1/2}$, P_W is of order $\phi(\alpha_k)$, and P_U is of order $\log(\phi(\alpha_k))$, then we can assert that P_W tends to 1 faster than P_U . But when the order of α_k is higher than $n^{1/2}$, P_W is of order $n^{1/2}$, and P_U is of order $\log(\phi(\alpha_k))$, then the convergence rate depends on the value of α_k .

For P_V , $\frac{2(p-M)}{2+c_{nM}+\sqrt{c_{nM}^2+4}} - \frac{2p}{2+c_n+\sqrt{c_n^2+4}}$, A_4 and ς_V are all of order $O(1)$. When $\alpha_k > 1 + \sqrt{c}$, whether it tends to infinity or not, we can assert that $\sum_{k=1}^K d_k \frac{\phi(\alpha_k)}{1+\phi(\alpha_k)}$ is of constant order $O(1)$. Then P_V does not tend to 1 when n tends to infinity.

For the power of R , according to [Hou et al. \(2019\)](#),

$$P_R - 1 + \Xi_{d_1} \left(\tau \sqrt{n} \left(\frac{\mu_{n,p}}{n} + t_\xi \frac{\sigma_{n,p}}{n} - \phi(\alpha_1) \right) \right) \rightarrow 0,$$

where Ξ_{d_1} is the distribution of the largest eigenvalue of a Gaussian orthogonal ensemble (GOE) matrix, $\mu_{n,p} = (\sqrt{n-1} + \sqrt{p})^2$, $\sigma_{n,p} = (\sqrt{n-1} + \sqrt{p}) \left(\frac{1}{\sqrt{n-1}} + \frac{1}{\sqrt{p}} \right)^{1/3}$, and τ is a constant. Note that $\frac{\mu_{n,p}}{n}$ is of constant order $O(1)$, and the order of $\frac{\sigma_{n,p}}{n}$ is $n^{-2/3}$. The term $\tau \sqrt{n} \left(\frac{\mu_{n,p}}{n} - \phi(\alpha_1) \right)$ tends to negative infinity. Then we find P_R tends to 1 whether the spiked eigenvalues α_1 are bounded or unbounded, and the convergence rate is $\alpha_1 \sqrt{n}$. When the largest population eigenvalue α_1 is simple, that is, $d_1 = 1$, therefore, the largest eigenvalue turns out to be the only element of the GOE, a normal random variable. Then comparing P_W with P_R , we can imagine that when the number of spikes increases to infinity, test W may have better performance than test R . But it is not the focus of this paper, therefore, we decide to leave it in our further research.

We conclude the subsection by reporting a small Monte-Carlo experiment that demonstrates our analysis of the power of U, W, V, R under the spiked model. We assume samples are drawn from Gaussian, $p = 150, n = 300$. We set spiked eigenvalues α_1 with five different orders. Under H_2 there are two spiked eigenvalues and we set $\alpha_1 = \alpha_2$. The simulation results are listed in Table 1. It is worth noting that when the spiked eigenvalue is of constant order $O(1)$, R is significantly better than others. R has the best effect whether the spiked eigenvalues are of constant order $O(1)$ or tend to infinity. Moreover, when α_1 tends to infinity, the powers of U and W also tend to 1 and W is faster than U , which is consistent with Remark 4.1. V has the worst performance under the spiked model. When comparing H_2 with H_1 , we find that as the number of spikes increases, although R has the best effect, there is an obvious improvement in the performance of U, W, V . We guess that R is not the only top-performing test when the number of spikes is large.

TABLE 1
Empirical power from 10000 repeated simulations based on H_1 and H_2
when $p = 150, n = 300$

Model	Test	$\alpha_1 = 3$	$\alpha_1 = n^{1/4}$	$\alpha_1 = n^{1/3}$	$\alpha_1 = n^{1/2}$	$\alpha_1 = n^{2/3}$
H_1	U	0.3803	0.5807	0.8551	0.9985	1
	W	0.6386	0.9279	0.9999	1	1
	V	0.2999	0.3823	0.4966	0.6276	0.6786
	R	1	1	1	1	1
H_2	U	0.8580	0.9842	0.9997	1	1
	W	0.9883	0.9999	1	1	1
	V	0.6677	0.8166	0.9284	0.9831	0.9931
	R	1	1	1	1	1

4.2. *Testing the hypothesis when eigenvalues are equal: H_3 .* The second test problem we consider concerns $H_0 = \mathbf{I}_p$ versus H_3 ; that is, the eigenvalues of the population covariance matrix are equal and close to 1. In this subsection, we present the asymptotic distribution of the four statistics U, W, V, R under H_3 in Theorems 4.4–4.7, including the approximated power functions of the four tests. Here, we consider a trivial case when $c < 1$.

For brevity, we introduce some notations. We define

$$\varepsilon(c) = \frac{l(1+c) - 1 + \sqrt{(1-l(1+c))^2 - 4(l^2 - l(1+c))}}{2}, \quad \check{c} = \frac{l^2 c}{(1 + \varepsilon(c))^2}, \quad l = 1 + \frac{a}{n}.$$

To avoid misunderstandings, we define the values of $\varepsilon(c_n)$ to be the same as $\varepsilon(c)$ with the substitution of c_n for c . The same substitution also holds for \check{c}_n .

THEOREM 4.4 (U statistic). *Under Assumptions 1–3 with $c_n = p/n \rightarrow c \in (0, 1)$, we have under H_3 ,*

$$\frac{U - p \int f_U(x) dF^{c_n, H_n}(x) - \mu_U}{\sqrt{\varsigma_U^2}} \xrightarrow{d} N(0, 1),$$

where

$$\begin{aligned} \int f_U(x) dF^{c_n, H_n}(x) &= CT(\varepsilon(c_n), c_n, \check{c}_n), \\ \mu_U &= \alpha_x I_1(f_U) + \beta_x I_2(f_U), \quad \varsigma_U^2 = (\alpha_x + 1) J_1(f_U, f_U) + \beta_x J_2(f_U, f_U), \\ I_1(f_U) &= \log(1 + \varepsilon(c_n)) - \log(1 + l + lc_n) + \sum_{k=1}^{\infty} \left(\frac{l\sqrt{c_n}}{1 + l + lc_n} \right)^{2k} \frac{(2k-1)!}{k!k!}, \\ I_2(f_U) &= - \sum_{k=1}^{\infty} \left(\frac{l\sqrt{c_n}}{1 + l + lc_n} \right)^{2k} \frac{(2k-1)!}{(k-1)!(k+1)!}, \\ J_1(f_U, f_U) &= J_2(f_U, f_U) = \left(\sum_{k=1}^{\infty} \left(\frac{l\sqrt{c_n}}{1 + l + lc_n} \right)^{2k-1} \frac{(2k-2)!}{k!(k-1)!} \right)^2. \end{aligned}$$

COROLLARY 4.4. *Under the same assumptions as in Theorem 4.4, we have*

$$P_U - \Phi \left(\frac{pCT(\varepsilon(c_n), c_n, \check{c}_n) - pCT(\varrho(c_n), c_n, \check{c}_n) + A_5}{\varsigma_U} - z_\xi \frac{A_3}{\varsigma_U} \right) \rightarrow 0,$$

where P_U is the power function of Wilk's likelihood ratio test, Φ is the distribution of a standard normal variable, and z_ξ is the upper quantile of the of the standard normal distribution Φ at level ξ . A_3 is defined in Corollary 4.1, and

$$\begin{aligned} A_5 = & \alpha_x \left[\log \left(\frac{1 + \varepsilon(c_n)}{1 + \varrho(c_n)} \right) + \log \left(\frac{2 + c_n}{1 + l + lc_n} \right) + \right. \\ & \left. \frac{(2k-1)!}{k!k!} \left(\sum_{k=1}^{\infty} \left(\frac{l\sqrt{c_n}}{1 + l + lc_n} \right)^{2k} - \sum_{k=1}^{\infty} \left(\frac{\sqrt{c_n}}{2 + c_n} \right)^{2k} \right) \right] + \\ & \beta_x \left[\frac{(2k-1)!}{(k+1)!(k-1)!} \left(\sum_{k=1}^{\infty} \left(\frac{\sqrt{c_n}}{2 + c_n} \right)^{2k} - \sum_{k=1}^{\infty} \left(\frac{\sqrt{lc_n}}{1 + l + lc_n} \right)^{2k} \right) \right]. \end{aligned}$$

THEOREM 4.5 (W statistic). Under Assumptions 1–3 with $c_n = p/n \rightarrow c \in (0, 1)$, we have under H_3 ,

$$\frac{W - lp}{l\sqrt{(\alpha_x + \beta_x + 1)c_n}} \xrightarrow{d} N(0, 1).$$

COROLLARY 4.5. Under the same assumptions as in Theorem 4.5, we have

$$P_W - \Phi \left(\frac{(l-1)p}{l\sqrt{(\alpha_x + \beta_x + 1)c_n}} - \frac{z_\xi}{l} \right) \rightarrow 0,$$

where P_W is the power function of Lawley-Hotelling trace test, Φ is the distribution of a standard normal variable, and z_ξ is the upper quantile of the standard normal distribution Φ at level ξ .

THEOREM 4.6 (V statistic). Under Assumptions 1–3 with $c_n = p/n \rightarrow c \in (0, 1)$, we have under H_3 ,

$$\frac{V - p \int f_V(x) dF^{c_n, H_n}(x) - \mu_V}{\sqrt{\varsigma_V^2}} \xrightarrow{d} N(0, 1),$$

where

$$\begin{aligned} \int f_V(x) dF^{c_n, H_n}(x) &= \frac{l}{1 + \varepsilon(c_n)}, \\ \mu_V &= \alpha_x I_1(f_V) + \beta_x I_2(f_V), \quad \varsigma_V^2 = (\alpha_x + 1) J_1(f_V, f_V) + \beta_x J_2(f_V, f_V), \\ I_1(f_V) &= \frac{1}{1 + l + lc_n} \sum_{k=0}^{\infty} \left(\frac{l\sqrt{c_n}}{1 + l + lc_n} \right)^{2k} \frac{(2k)!}{k!k!} - \\ & \frac{1}{1 + \varepsilon(c_n)} \left(\frac{\check{c}_n}{-(1 - \check{c}_n)^2} + \frac{1}{2(\sqrt{\check{c}_n} - 1)^2} + \frac{1}{2(\sqrt{\check{c}_n} + 1)^2} \right), \\ I_2(f_V) &= -\frac{1}{1 + l + lc_n} \sum_{k=0}^{\infty} \left(\frac{l\sqrt{c_n}}{1 + l + lc_n} \right)^{2k} \frac{(2k)!}{(k-1)!(k+1)!}, \\ J_1(f_V, f_V) = J_2(f_V, f_V) &= \left(\frac{1}{1 + l + lc_n} \sum_{k=0}^{\infty} \left(\frac{l\sqrt{c_n}}{1 + l + lc_n} \right)^{2k+1} \frac{(2k+1)!}{k!(k+1)!} \right)^2. \end{aligned}$$

COROLLARY 4.6. *Under the same assumptions as in Theorem 4.6, we have*

$$P_V - \Phi \left(\frac{\frac{pl}{1+\varepsilon(c_n)} - \frac{p}{1+\varrho(c_n)} + A_6}{\varsigma_V} - z_\xi \frac{A_4}{\varsigma_V} \right) \rightarrow 0$$

where P_V is the power function of Bartlett-Nanda-Pillai trace test, Φ is the distribution of a standard normal variable, and z_ξ is the upper quantile of the of the standard normal distribution Φ at level ξ , and A_4 is defined in Corollary 4.3,

$$\begin{aligned} A_6 = & \alpha_x \left[\frac{(2k)!}{k!k!} \left(\frac{1}{1+l+lc_n} \sum_{k=0}^{\infty} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k} - \frac{1}{2+c_n} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c_n}}{2+c_n} \right)^{2k} \right) + \right. \\ & \frac{1}{1+\varrho(c_n)} \left(\frac{\tilde{c}_n}{-(1-\tilde{c}_n)^2} + \frac{1}{2(\sqrt{\tilde{c}_n}-1)^2} + \frac{1}{2(\sqrt{\tilde{c}_n}+1)^2} \right) - \\ & \left. \frac{1}{1+\varepsilon(c_n)} \left(\frac{\check{c}_n}{-(1-\check{c}_n)^2} + \frac{1}{2(\sqrt{\check{c}_n}-1)^2} + \frac{1}{2(\sqrt{\check{c}_n}+1)^2} \right) \right] + \beta_x \frac{2k!}{(k-1)!(k+1)!} \\ & \left[\frac{1}{2+c_n} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c_n}}{2+c_n} \right)^{2k} - \frac{1}{1+l+lc_n} \sum_{k=0}^{\infty} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k} \right]. \end{aligned}$$

THEOREM 4.7 (R statistic). *Under Assumptions 1–3 with $c_n = p/n \rightarrow c \in (0, 1)$, we have under H_3 ,*

$$\frac{R - l \frac{\mu_{n,p}}{n}}{l \frac{\sigma_{n,p}}{n}} \xrightarrow{d} TW_1,$$

where

$$\begin{aligned} \mu_{n,p} &= (\sqrt{n-1} + \sqrt{p})^2, \\ \sigma_{n,p} &= (\sqrt{n-1} + \sqrt{p}) \left(\frac{1}{\sqrt{n-1}} + \frac{1}{\sqrt{p}} \right)^{1/3}, \end{aligned}$$

and TW_1 denotes the type-1 Tracy-Widom distribution.

COROLLARY 4.7. *Under the same assumptions as in Theorem 4.7, we have*

$$P_R - 1 + F_1 \left(\frac{t_\xi}{l} + \frac{(1-l) \frac{\mu_{n,p}}{n}}{l \frac{\sigma_{n,p}}{n}} \right) \rightarrow 0,$$

where P_R is the power function of Roy's largest root test, F_1 is the cumulative distribution function of a type-1 Tracy-Widom distribution variable, and t_ξ is the upper quantile of the type-1 Tracy-Widom distribution F_1 at level ξ .

REMARK 4.2. *For P_U , A_3 , A_5 , and ς_U are all of constant order $O(1)$. As a in H_3 increases, the value of l becomes larger. When $n(l-1)$ tends to a constant, $pCT(\varepsilon(c_n), c_n, \check{c}_n) - pCT(\varrho(c_n), c_n, \tilde{c}_n)$ tends to a constant, then we can assert that P_U does not tend to 1. When $n(l-1)$ tends to infinity as a increases, $pCT(\varepsilon(c_n), c_n, \check{c}_n) - pCT(\varrho(c_n), c_n, \tilde{c}_n)$ tends to infinity, then P_U tends to 1.*

For P_W , $l\sqrt{(\alpha_x + \beta_x + 1)c_n}$ is of constant order $O(1)$. Similar to the U statistic, when $n(l-1)$ tends to a constant, then $(l-1)p$ is also of constant order; therefore, we can assert

TABLE 2
Empirical power from 10000 repeated simulations based on H_3 : eigenvalues are equal
when $p = 100, n = 300$

Test	$a = 1$	$a = 3$	$a = 5$	$a = 10$	$a = 18$	$a = 50$
U	0.0898	0.3085	0.6215	0.9871	1	1
W	0.1108	0.3449	0.6504	0.9903	1	1
V	0.1261	0.3515	0.6353	0.9849	1	1
R	0.0594	0.0980	0.1497	0.3233	0.7190	1

that P_W does not tend to 1. When l becomes larger and $n(l-1)$ tends to infinity, then $(l-1)p$ also tends to infinity; therefore, P_W tends to 1.

For P_V , A_4 , A_6 , and ς_V are all of constant order $O(1)$. When $n(l-1)$ tends to a constant, $\frac{pl}{1+\varepsilon(c_n)} - \frac{p}{1+\varrho(c_n)} = p \left(\frac{l}{1+\varepsilon(c_n)} - \frac{1}{1+\varrho(c_n)} \right)$ tends to a constant, then we assert that P_V does not tend to 1. When l becomes larger and $n(l-1)$ tends to infinity, $\frac{pl}{1+\varepsilon(c_n)} - \frac{p}{1+\varrho(c_n)}$ tends to infinity. Therefore, in this case, P_V tends to 1.

For P_R , $\frac{\mu_{n,p}}{n}$ is of constant order $O(1)$, and the order of $\frac{\sigma_{n,p}}{n}$ is $n^{-2/3}$. When $n^s(l-1)$ tends to a constant, and $\frac{2}{3} \leq s \leq 1$, since t_ξ and l are two constants, then $\frac{t_\xi}{l} + \frac{(1-l)\frac{\mu_{n,p}}{n}}{l\frac{\sigma_{n,p}}{n}}$ tends to a constant as n tends to infinity. Therefore, we conclude that $F_1 \left(\frac{t_\xi}{l} + \frac{(1-l)\frac{\mu_{n,p}}{n}}{l\frac{\sigma_{n,p}}{n}} \right)$ does not tend to 0. In other words, P_R does not tend to 1. When $s < \frac{2}{3}$, then $\frac{t_\xi}{l} + \frac{(1-l)\frac{\mu_{n,p}}{n}}{l\frac{\sigma_{n,p}}{n}}$ tends to negative infinity, which yields $F_1 \left(\frac{t_\xi}{l} + \frac{(1-l)\frac{\mu_{n,p}}{n}}{l\frac{\sigma_{n,p}}{n}} \right)$ tends to 0. In other words, P_R tends to 1 when $s < \frac{2}{3}$, but the convergence rate is slower than $n^{2/3}$. For example, when $s = \frac{1}{2}$, then the convergence rate is $n^{1/6}$, which is a slow rate. As a comparison, the convergence rates of P_W and P_V are both $p(l-1)$, while the convergence rate of P_U is $p \log(l)$, and that of P_R is $(1-l)n^{2/3}$. Then, we assert that P_R has the slowest convergence rate, and P_W and P_V converge the fastest.

We conclude this subsection by reporting a Monte-Carlo experiment. It is similar to the one in the previous subsection and designed to examine the analysis of the power of U, W, V, R under H_3 . We assume Gaussian samples, with $p = 100, n = 300$. We adopt six different settings for a . The simulation results are listed in Table 2. The differences are significant for $a = 5$ and $a = 10$, which is consistent with Remark 4.2. For R , when $a = 18$, it also tends to 1 but with a slow rate.

5. Technical proofs. In this section, we present some lemmas that are needed in the proofs of main results. Proofs of Theorems 3.1–4.7, and Corollaries 4.1–4.7 are postponed to the supplementary materials. First, we truncate and renormalize the random variables to ensure the existence of their higher-order moments.

5.1. *Truncation and renormalization.* Let $\hat{x}_{ij} = x_{ij} \mathbb{1}_{\{|x_{ij}| < \eta_n \sqrt{n}\}}$ and $\tilde{x}_{ij} = \frac{\hat{x}_{ij} - \mathbb{E}\hat{x}_{ij}}{\hat{\sigma}_n}$, where $\hat{\sigma}_n^2 = \mathbb{E}|\hat{x}_{ij} - \mathbb{E}\hat{x}_{ij}|^2$ and $\eta_n \rightarrow 0$ with a slow rate. Correspondingly, define $\hat{\mathbf{B}} = \frac{1}{n} \mathbf{T} \hat{\mathbf{X}} \hat{\mathbf{X}}^* \mathbf{T}^*$ and $\tilde{\mathbf{B}} = \frac{1}{n} \mathbf{T} \tilde{\mathbf{X}} \tilde{\mathbf{X}}^* \mathbf{T}^*$, where $\hat{\mathbf{X}} = (\hat{x}_{ij})$ and $\tilde{\mathbf{X}} = (\tilde{x}_{ij})$. \hat{G}_n and \tilde{G}_n denote the analogs of G_n with the matrix \mathbf{B} replaced by $\hat{\mathbf{B}}$ and $\tilde{\mathbf{B}}$, respectively. Next, we demonstrate that the entries of \mathbf{X} in the LSSs are equivalent to and can be replaced by the truncated and renormalized entries.

According to the Lindeberg-type condition in Assumption 1, we find that as $\min\{n, p\} \rightarrow \infty$,

$$\mathbb{P}(\mathbf{B} \neq \hat{\mathbf{B}}) \leq \sum_{i,j} \mathbb{P}(|x_{ij}| \geq \eta_n \sqrt{n}) \rightarrow 0.$$

It follows from the definition of LSSs that for any $l = 1, \dots, h$,

$$\begin{aligned} & \left| \int f_l(x) d\hat{G}_n - \int f_l(x) d\tilde{G}_n(x) \right| = \sum_{i=1}^p \left| f_l(\lambda_i^{\hat{\mathbf{B}}}) - f_l(\lambda_i^{\tilde{\mathbf{B}}}) \right| \\ (5.6) \quad & \leq \sum_{i=1}^M \left| f_l(\lambda_i^{\hat{\mathbf{B}}}) - f_l(\lambda_i^{\tilde{\mathbf{B}}}) \right| + \sum_{i=M+1}^p \left| f_l(\lambda_i^{\hat{\mathbf{B}}}) - f_l(\lambda_i^{\tilde{\mathbf{B}}}) \right|. \end{aligned}$$

Using the same discussion in Bai and Silverstein (2004), we can easily find that the second term of (5.6) tends to 0 in probability. For the first term of (5.6), from the arguments in Supplement B of Jiang and Bai (2021), we know that

$$(5.7) \quad \left| \lambda_i^{\hat{\mathbf{B}}} - \lambda_i^{\tilde{\mathbf{B}}} \right| = o_p(n^{-\frac{1}{2}} \rho_i).$$

We recall that $\rho_i = \alpha_k$ if $i \in J_k$, where $J_k = \{j_k + 1, \dots, j_k + d_k\}$ is the set of ranks of the eigenvalue α_k with multiplicities d_k . Then, for brevity, we denote $\beta_i = (\lambda_i^{\hat{\mathbf{B}}} - \lambda_i^{\tilde{\mathbf{B}}})/\rho_i$ and obtain that

$$\begin{aligned} & f_l(\lambda_i^{\hat{\mathbf{B}}}) - f_l(\lambda_i^{\tilde{\mathbf{B}}}) = \int_0^{\beta_i \rho_i} f_l'(t + \lambda_i^{\tilde{\mathbf{B}}}) dt \\ (5.8) \quad & = \int_0^1 \beta_i \rho_i f_l'(\beta_i \rho_i s + \lambda_i^{\tilde{\mathbf{B}}}) ds = \beta_i \rho_i f_l'(\rho_i) \int_0^1 \frac{f_l'(\rho_i(\beta_i s + \frac{\lambda_i^{\tilde{\mathbf{B}}}}{\rho_i}))}{f_l'(\rho_i)} ds, \end{aligned}$$

since $\beta_i = o_p(n^{-\frac{1}{2}})$, $\frac{\lambda_i^{\tilde{\mathbf{B}}}}{\rho_i}$ tends to 1, then from Assumption 4 we obtain

$$(5.9) \quad \sqrt{n} \sum_{i=1}^M \frac{\left| f_l(\lambda_i^{\hat{\mathbf{B}}}) - f_l(\lambda_i^{\tilde{\mathbf{B}}}) \right|}{f_l'(\rho_i) \rho_i} = o_p(1).$$

Since $\frac{Y_1 - \mu_1}{\sqrt{\varsigma_1^2}} = \frac{Y_1 - \mu_1}{\sqrt{\sum_{k=1}^K \frac{\phi_n^2(\alpha_k)}{n} (f_1'(\phi_n^2(\alpha_k)))^2 s_k^2 + A}}$ where $A = -\frac{1}{4\pi^2} \oint_{\mathcal{C}_1} \oint_{\mathcal{C}_2} f_1(z_1) f_1(z_2) \vartheta_n^2 dz_1 dz_2$ and (5.9), we obtain

$$\begin{aligned} & \left| \frac{\hat{Y}_1 - \mu_1}{\sqrt{\varsigma_1^2}} - \frac{\tilde{Y}_1 - \mu_1}{\sqrt{\varsigma_1^2}} \right| = \left| \frac{\hat{Y}_1 - \tilde{Y}_1}{\sqrt{\varsigma_1^2}} \right| = \left| \frac{\hat{Y}_1 - \tilde{Y}_1}{\sqrt{\sum_{k=1}^K \frac{\phi_n^2(\alpha_k)}{n} (f_1'(\phi_n^2(\alpha_k)))^2 s_k^2 + A}} \right| \\ & \leq \left| \frac{\hat{Y}_1 - \tilde{Y}_1}{\sqrt{\sum_{k=1}^K \frac{\phi_n^2(\alpha_k)}{n} (f_1'(\phi_n^2(\alpha_k)))^2 s_k^2}} \right| = o_p(1), \end{aligned}$$

where the last equality is due to $\left| \frac{\hat{Y}_1 - \tilde{Y}_1}{\sqrt{\sum_{k=1}^K \frac{\phi_n^2(\alpha_k)}{n} (f_1'(\phi_n^2(\alpha_k)))^2 s_k^2}} \right|$ has the same order as (5.9).

Thus, it is concluded that the procedure of truncation does not affect the limiting distribution of LSS. Therefore, in the following proofs, we can safely assume that $|x_{ij}| < \eta_n \sqrt{n}$.

5.2. *Some primary definitions and lemmas.* In this section, we provide some useful results that are used later in the proofs of Theorems 3.1 and 3.2. For the population covariance matrix $\Sigma = \mathbf{T}\mathbf{T}^*$, we consider the corresponding sample covariance matrix $\mathbf{B} = \mathbf{T}\mathbf{S}_x\mathbf{T}^*$, where $\mathbf{S}_x = \frac{1}{n}\mathbf{X}\mathbf{X}^*$. By singular value decomposition of \mathbf{T} (see (2.2)),

$$\mathbf{B} = \mathbf{V} \begin{pmatrix} \mathbf{D}_1^{\frac{1}{2}} \mathbf{U}_1^* \mathbf{S}_x \mathbf{U}_1 \mathbf{D}_1^{\frac{1}{2}} & \mathbf{D}_1^{\frac{1}{2}} \mathbf{U}_1^* \mathbf{S}_x \mathbf{U}_2 \mathbf{D}_2^{\frac{1}{2}} \\ \mathbf{D}_2^{\frac{1}{2}} \mathbf{U}_2^* \mathbf{S}_x \mathbf{U}_1 \mathbf{D}_1^{\frac{1}{2}} & \mathbf{D}_2^{\frac{1}{2}} \mathbf{U}_2^* \mathbf{S}_x \mathbf{U}_2 \mathbf{D}_2^{\frac{1}{2}} \end{pmatrix} \mathbf{V}^*.$$

Note that

$$\mathbf{S} = \begin{pmatrix} \mathbf{D}_1^{\frac{1}{2}} \mathbf{U}_1^* \mathbf{S}_x \mathbf{U}_1 \mathbf{D}_1^{\frac{1}{2}} & \mathbf{D}_1^{\frac{1}{2}} \mathbf{U}_1^* \mathbf{S}_x \mathbf{U}_2 \mathbf{D}_2^{\frac{1}{2}} \\ \mathbf{D}_2^{\frac{1}{2}} \mathbf{U}_2^* \mathbf{S}_x \mathbf{U}_1 \mathbf{D}_1^{\frac{1}{2}} & \mathbf{D}_2^{\frac{1}{2}} \mathbf{U}_2^* \mathbf{S}_x \mathbf{U}_2 \mathbf{D}_2^{\frac{1}{2}} \end{pmatrix} \triangleq \begin{pmatrix} \mathbf{S}_{11} & \mathbf{S}_{12} \\ \mathbf{S}_{21} & \mathbf{S}_{22} \end{pmatrix}.$$

Moreover, \mathbf{B} and \mathbf{S} have the same eigenvalues.

Recall that $\underline{\mathbf{B}} = \frac{1}{n}\mathbf{X}^*\mathbf{T}^*\mathbf{T}\mathbf{X}$ (the spectrum of which differs from that of \mathbf{B} by $|n-p|$ zeros). Its limiting spectral distribution is $\underline{F}^{c,H}$, $\underline{F}^{c,H} \equiv (1-c)\mathbb{1}_{[0,\infty)} + cF^{c,H}$, and its Stieltjes transform is $\underline{m}(z)$. Let $\tilde{\lambda}_j$ be the eigenvalues of \mathbf{S}_{22} so that the LSS of \mathbf{S}_{22} is $\sum_{j=1}^{p-M} f(\tilde{\lambda}_j)$. Correspondingly, recall that $c_{nM} := \frac{p-M}{n}$, $H_{2n} := F^{\mathbf{D}^2}$ and $m_{2n0} := m_{2n0}(z)$ are the Stieltjes transforms of $F^{c_{nM}, H_{2n}}$. First, in Lemma 5.1 we derive that the difference between the two centers is 0.

LEMMA 5.1. *Under Assumptions 1–4,*

$$(p-M) \int f(x) dF^{c_{nM}, H_{2n}} = p \int f(x) dF^{c_n, H_n}(x).$$

PROOF. By the Cauchy integral formula,

$$p \int f(x) dF^{c_n, H_n} = -\frac{p}{2\pi i} \oint_{\mathcal{C}} f(z) m_{1n0} dz = -\frac{n}{2\pi i} \oint_{\mathcal{C}} f(z) \underline{m}_{1n0} dz,$$

$$(p-M) \int f(x) dF^{c_{nM}, H_{2n}} = -\frac{p-M}{2\pi i} \oint_{\mathcal{C}} f(z) m_{2n0} dz = -\frac{n}{2\pi i} \oint_{\mathcal{C}} f(z) \underline{m}_{2n0} dz,$$

where \underline{m}_{1n0} and \underline{m}_{2n0} are the Stieltjes transforms of \underline{F}^{c_n, H_n} and $\underline{F}^{c_{nM}, H_{2n}}$, respectively. Then,

$$(p-M) \int f(x) dF^{c_{nM}, H_{2n}} - p \int f(x) dF^{c_n, H_n} = \frac{n}{2\pi i} \oint_{\mathcal{C}} f(z) (\underline{m}_{1n0} - \underline{m}_{2n0}) dz.$$

Next, we prove that $\underline{m}_{1n0} = \underline{m}_{2n0}$.

Note that m_{1n0} and m_{2n0} are the unique solutions to

$$(5.10) \quad z = -\frac{1}{\underline{m}_{1n0}} + c_n \int \frac{tdH_n(t)}{1 + t\underline{m}_{1n0}}$$

$$(5.11) \quad z = -\frac{1}{\underline{m}_{2n0}} + c_{nM} \int \frac{tdH_{2n}(t)}{1 + t\underline{m}_{2n0}},$$

respectively, where $\underline{m}_{1n0} = -\frac{1-c_n}{z} + c_n m_{1n0}$ and $\underline{m}_{2n0} = -\frac{1-c_{nM}}{z} + c_{nM} m_{2n0}$. Since

$$H_n(t) = \frac{1}{p} \left[\sum_{i=1}^M \mathbb{1}_{\{0 \leq t\}} + \sum_{i=M+1}^p \mathbb{1}_{\{\alpha_i \leq t\}} \right] = \frac{M}{p} + \frac{1}{p} \sum_{i=M+1}^p \mathbb{1}_{\{\alpha_i \leq t\}}$$

and $H_{2n}(t) = \frac{1}{p-M} \sum_{i=M+1}^p \mathbb{1}_{\{\alpha_i \leq t\}}$, (5.10) can be written as

$$\begin{aligned}
z &= -\frac{1}{\underline{m}_{1n0}} + \frac{p}{n} \int \frac{td \left(\frac{M}{p} + \frac{1}{p} \sum_{i=M+1}^p \mathbb{1}_{\{\alpha_i \leq t\}}(t) \right)}{1 + t\underline{m}_{1n0}} \\
&= -\frac{1}{\underline{m}_{1n0}} + \frac{p}{n} \int \frac{td \left(\frac{1}{p} \sum_{i=M+1}^p \mathbb{1}_{\{\alpha_i \leq t\}}(t) \right)}{1 + t\underline{m}_{1n0}} \\
(5.12) \quad &= -\frac{1}{\underline{m}_{1n0}} + \frac{1}{n} \sum_{i=M+1}^p \frac{\alpha_i}{1 + \alpha_i \underline{m}_{1n0}}.
\end{aligned}$$

Similarly, equation (5.11) can be written as

$$(5.13) \quad z = -\frac{1}{\underline{m}_{2n0}} + \frac{1}{n} \sum_{i=M+1}^p \frac{\alpha_i}{1 + \alpha_i \underline{m}_{2n0}}.$$

Thus, according to the fact that m_{1n0} and m_{2n0} are the unique solutions of (5.12) and (5.13), respectively, we have $m_{1n0} = m_{2n0}$, which completes the proof of this lemma. \square

Recall that in the introduction we mentioned that the bounded part of the LSS does not equal the LSS of \mathbf{S}_{22} since the off-diagonal sample covariance matrix blocks are not 0. The following lemma measures the difference between the two parts mentioned above.

LEMMA 5.2. *Under Assumptions 1–4,*

$$\sum_{j=M+1}^p f(\lambda_j) - \sum_{j=1}^{p-M} f(\tilde{\lambda}_j) - \frac{M}{2\pi i} \oint_{\mathcal{C}} f(z) \frac{\underline{m}'(z)}{\underline{m}(z)} dz = o_p(1).$$

PROOF. Note that

$$L_1 := \sum_{j=M+1}^p f(\lambda_j).$$

By the Cauchy integral formula, we have

$$L_1 = -\frac{p}{2\pi i} \oint_{\mathcal{C}} f(z) m_n(z) dz,$$

where $m_n = \frac{1}{p} \text{tr}(\mathbf{S} - z\mathbf{I}_p)^{-1} = \frac{1}{p} \text{tr}(\mathbf{B} - z\mathbf{I}_p)^{-1}$. Analogously, we have

$$L_2 := \sum_{j=1}^{p-M} f(\tilde{\lambda}_j) = -\frac{p-M}{2\pi i} \oint_{\mathcal{C}} f(z) m_{2n}(z) dz,$$

where $m_{2n} = \frac{1}{p-M} \text{tr}(\mathbf{S}_{22} - z\mathbf{I}_{p-M})^{-1}$. By applying the block matrix inversion formula to m_n , we can obtain

$$(5.14) \quad L_1 - L_2 = -\frac{1}{2\pi i} \oint_{\mathcal{C}} f(z) (T_1 - T_2) dz,$$

where

$$\begin{aligned}
T_1 &= \text{tr} \left(\mathbf{S}_{11} - z\mathbf{I}_M - \mathbf{S}_{12} (\mathbf{S}_{22} - z\mathbf{I}_{p-M})^{-1} \mathbf{S}_{21} \right)^{-1}, \\
T_2 &= -\text{tr} \left[\left(\mathbf{S}_{11} - z\mathbf{I}_M - \mathbf{S}_{12} (\mathbf{S}_{22} - z\mathbf{I}_{p-M})^{-1} \mathbf{S}_{21} \right)^{-1} \mathbf{S}_{12} (\mathbf{S}_{22} - z\mathbf{I}_{p-M})^{-2} \mathbf{S}_{21} \right].
\end{aligned}$$

Note that for any matrix \mathbf{Z} ,

$$\mathbf{Z}(\mathbf{Z}^*\mathbf{Z} - \lambda\mathbf{I})^{-1}\mathbf{Z}^* = \mathbf{I} + \lambda(\mathbf{Z}\mathbf{Z}^* - \lambda\mathbf{I})^{-1},$$

which, together with the notation $\Upsilon_n := \frac{1}{n}\mathbf{D}_1^{\frac{1}{2}}\mathbf{U}_1^*\mathbf{X}\left(\frac{1}{n}\mathbf{X}^*\mathbf{U}_2\mathbf{D}_2\mathbf{U}_2^*\mathbf{X} - z\mathbf{I}_n\right)^{-1}\mathbf{X}^*\mathbf{U}_1\mathbf{D}_1^{\frac{1}{2}}$, implies that

$$\begin{aligned} T_1 &= -z^{-1}\text{tr}(\mathbf{I}_M + \Upsilon_n)^{-1}, \\ T_2 &= z^{-1}\text{tr}\left[(\mathbf{I}_M + \Upsilon_n)^{-1}\mathbf{S}_{12}(\mathbf{S}_{22} - z\mathbf{I}_{p-M})^{-2}\mathbf{S}_{21}\right], \end{aligned}$$

$\underline{m}_{2n} = \underline{m}_{2n}(z)$ denotes the Stieltjes transform of $F_n^{\frac{1}{n}\mathbf{X}^*\mathbf{U}_2\mathbf{D}_2\mathbf{U}_2^*\mathbf{X}}$. Thus, we have that $\underline{m}_{2n}(z) - \underline{m}(z) = o_p(1)$ for any $z \in \mathcal{C}$. From Theorem 3.1 of (Jiang and Bai, 2021), we know that

$$(5.15) \quad \frac{1}{n}\mathbf{U}_1^*\mathbf{X}\left(\frac{1}{n}\mathbf{X}^*\mathbf{U}_2\mathbf{D}_2\mathbf{U}_2^*\mathbf{X} - z\mathbf{I}_n\right)^{-1}\mathbf{X}^*\mathbf{U}_1 = \underline{m}_{2n}(z)\mathbf{I}_M + O_p(n^{-\frac{1}{2}}).$$

Thus, under Assumption 3, we find that

$$(5.16) \quad \mathbf{D}_1^{1/2}(\mathbf{I}_M + \Upsilon_n)^{-1}\mathbf{D}_1^{1/2} = \frac{1}{\underline{m}(z)}\mathbf{I}_M + o_p(1),$$

which yields

$$(5.17) \quad T_1 = o_p(1).$$

It follows that

$$\begin{aligned} & \mathbf{D}_1^{-1/2}\mathbf{S}_{12}(\mathbf{S}_{22} - z\mathbf{I}_{p-M})^{-2}\mathbf{S}_{21}\mathbf{D}_1^{-1/2} \\ &= \frac{1}{n}\text{tr}\left[(\mathbf{S}_{22} - z\mathbf{I}_{p-M})^{-2}\mathbf{S}_{22}\right]\mathbf{I}_M + O_p(n^{-\frac{1}{2}}) \\ &= \frac{1}{n}\text{tr}(\mathbf{S}_{22} - z\mathbf{I}_{p-M})^{-1}\mathbf{I}_M + \frac{z}{n}\text{tr}(\mathbf{S}_{22} - z\mathbf{I}_{p-M})^{-2}\mathbf{I}_M + O_p(n^{-\frac{1}{2}}) \\ &= cm(z)\mathbf{I}_M + zcm'(z)\mathbf{I}_M + o_p(1) \\ (5.18) \quad &= \underline{m}(z)\mathbf{I}_M + z\underline{m}'(z)\mathbf{I}_M + o_p(1), \end{aligned}$$

where the last equality is derived from $\underline{m} = -\frac{1-c}{z} + cm(z)$. Therefore, according to (5.16) and (5.18), we obtain

$$T_2 = M\frac{\underline{m}(z) + z\underline{m}'(z)}{z\underline{m}(z)} + o_p(1),$$

which, together with (5.14) and (5.17), implies that

$$L_1 - L_2 = \frac{M}{2\pi i} \oint_{\mathcal{C}} f(z) \frac{\underline{m}(z) + z\underline{m}'(z)}{z\underline{m}(z)} dz + o_p(1) = \frac{M}{2\pi i} \oint_{\mathcal{C}} f(z) \frac{\underline{m}'(z)}{\underline{m}(z)} dz + o_p(1).$$

Therefore, the proof of this lemma is complete. \square

Define random vector $\gamma_k = (\gamma_{kj})' = \left(\sqrt{n}\frac{\lambda_j - \phi_n(\alpha_k)}{\phi_n(\alpha_k)}, j \in J_k\right)'$, where J_k is the indicator set of a packet of d_k consecutive sample eigenvalues. Then, we present the following lemma, which is borrowed from Jiang and Bai (2021) and characterizes the limiting distribution of the spiked eigenvalues of the sample covariance matrix.

LEMMA 5.3. (*Jiang and Bai (2021)*) Under Assumptions 1–4, random vector γ_k converges weakly to the joint distribution of d_k eigenvalues of a Gaussian random matrix

$$-\frac{1}{\theta_k} [\mathbf{\Omega}_{\phi_k}]_{kk},$$

where

$$\theta_k = \phi_k^2 \underline{m}_2(\phi_k), \quad \underline{m}_2(\lambda) = \int \frac{1}{(\lambda - x)^2} d\underline{F}^{c,H}(x),$$

with $\underline{F}^{c,H}$ being the LSD of matrix $n^{-1} \mathbf{X}^* \mathbf{U}_2 \mathbf{D}_2 \mathbf{U}_2^* \mathbf{X}$, $\phi_k = \alpha_k \left(1 + c \int \frac{t}{\alpha_k - t} dH(t)\right)$. $[\mathbf{\Omega}_{\phi_k}]_{kk}$ is the k th diagonal block of matrix $\mathbf{\Omega}_{\phi_k}$. The variances and covariances of the elements ω_{ij} of $\mathbf{\Omega}_{\phi_k}$ are:

$$\text{Cov}(\omega_{i_1, j_1}, \omega_{i_2, j_2}) = \begin{cases} (\alpha_x + 1)\theta_k + \beta_x \mathcal{U}_{iiii} \nu_k, & i_1 = j_1 = i_2 = j_2 = i \\ \theta_k + \beta_x \mathcal{U}_{ijij} \nu_k, & i_1 = i_2 = i \neq j_1 = j_2 = j \\ \beta_x \mathcal{U}_{i_1 j_1 i_2 j_2} \nu_k, & \text{other cases} \end{cases}$$

where $\beta_x \mathcal{U}_{i_1 j_1 i_2 j_2} = \sum_{t=1}^p \bar{u}_{ti_1} u_{tj_1} u_{ti_2} \bar{u}_{tj_2} \beta_x$, $\mathbf{u}_i = (u_{1i}, \dots, u_{pi})'$ are the i th column of the matrix \mathbf{U}_1 , $\nu_k = \phi_k^2 \underline{m}_2^2(\phi_k)$.

Recall that λ_j is the eigenvalue of \mathbf{B} , and $\tilde{\lambda}_j$ is the eigenvalue of \mathbf{S}_{22} . The following lemma shows the asymptotic independence between $\sum_{j=1}^M f(\lambda_j)$ and $\sum_{j=1}^{p-M} f(\tilde{\lambda}_j)$.

LEMMA 5.4. Under Assumptions 1–4, $\sum_{j=1}^M f(\lambda_j)$ and $\sum_{j=1}^{p-M} f(\tilde{\lambda}_j)$ are asymptotically independent.

PROOF. It is sufficient to prove that for a given $\sum_{j=1}^{p-M} f(\tilde{\lambda}_j)$, the asymptotic limiting distribution of $\sum_{j=1}^M f(\lambda_j)$ does not depend on the random part of $\sum_{j=1}^{p-M} f(\tilde{\lambda}_j)$, that is, it only depends on its limit.

First, we consider $f(x) = x$. From the proof of Theorem 3.1 in *Jiang and Bai (2021)*, we have the following determinant equation that

$$0 = |[\mathbf{\Omega}_M(\phi_k)]_{kk} + \lim \gamma_{kj} \{ \phi_k^2 \underline{m}_2(\phi_k) \} \mathbf{I}_{d_k}|,$$

where $\mathbf{\Omega}_M(\phi_k)$

$$= \frac{\phi_k}{\sqrt{n}} \left[\text{tr} \left\{ \left(\phi_k \mathbf{I} - \frac{1}{n} \mathbf{X}^* \mathbf{U}_2 \mathbf{D}_2 \mathbf{U}_2^* \mathbf{X} \right)^{-1} \right\} \mathbf{I} - \mathbf{U}_1^* \mathbf{X} \left(\phi_k \mathbf{I} - \frac{1}{n} \mathbf{X}^* \mathbf{U}_2 \mathbf{D}_2 \mathbf{U}_2^* \mathbf{X} \right)^{-1} \mathbf{X}^* \mathbf{U}_1 \right],$$

and $\underline{m}_2(\phi_k)$ is the limit of $\text{tr} \left(\phi_k \mathbf{I} - \frac{1}{n} \mathbf{X}^* \mathbf{U}_2 \mathbf{D}_2 \mathbf{U}_2^* \mathbf{X} \right)^{-2}$. Then, we know that γ_{kj} has the same asymptotic distribution with eigenvalues of $-\frac{[\mathbf{\Omega}_M(\phi_k)]_{kk}}{\phi_k^2 \underline{m}_2(\phi_k)}$ in order. From *Jiang and Bai (2021)*, we can obtain that the limiting distribution of $\mathbf{\Omega}_M(\phi_k)$ is not changed if \mathbf{X} is replaced by \mathbf{Y} , which is independent of \mathbf{X} but has the same distribution as it. Therefore $\mathbf{U}_1^* \mathbf{X}$ and $\mathbf{U}_2^* \mathbf{X}$ can be assumed to be independent since $\mathbf{U}_2^* \mathbf{X}$ can be replaced by $\mathbf{U}_2^* \mathbf{Y}$. Then, given $\mathbf{U}_2^* \mathbf{X}$, the limiting distribution of γ_{kj} only depends on the limit of $\text{tr} \left(\phi_k \mathbf{I} - \frac{1}{n} \mathbf{X}^* \mathbf{U}_2 \mathbf{D}_2 \mathbf{U}_2^* \mathbf{X} \right)^{-1}$, that is, $\underline{m}_2(\phi_k)$, and has nothing to do with the random part. Therefore, the independence between $\sum_{j=1}^M f(\lambda_j)$ and $\sum_{j=1}^{p-M} f(\tilde{\lambda}_j)$ is obtained when $f(x) = x$.

When $f(x) \neq x$, by using the Newton-Leibniz formula, we have

$$\begin{aligned}
& \sum_{j=1}^M f(\lambda_j) - \sum_{k=1}^K d_k f(\phi_n(\alpha_k)) = \sum_{k=1}^K \sum_{j \in J_k} (f(\lambda_j) - f(\phi_n(\alpha_k))) \\
&= \sum_{k=1}^K \sum_{j \in J_k} \int_0^{\frac{\phi_n(\alpha_k)}{\sqrt{n}} \gamma_{kj}} f'(t + \phi_n(\alpha_k)) dt \\
&= \sum_{k=1}^K \sum_{j \in J_k} \int_0^1 \frac{\phi_n(\alpha_k)}{\sqrt{n}} \gamma_{kj} \frac{f'(\phi_n(\alpha_k) (1 + \frac{\gamma_{kj}}{\sqrt{n}} s))}{f'(\phi_n(\alpha_k))} f'(\phi_n(\alpha_k)) ds \\
&= \sum_{k=1}^K \sum_{j \in J_k} \int_0^1 \varpi^k \gamma_{kj} \frac{f'(\phi_n(\alpha_k) (1 + \frac{\gamma_{kj}}{\sqrt{n}} s))}{f'(\phi_n(\alpha_k))} ds \\
(5.19) \quad & \rightarrow \sum_{k=1}^K \sum_{j \in J_k} \int_0^1 \gamma_{kj} \varpi^k ds = \sum_{k=1}^K \sum_{j \in J_k} \varpi^k \gamma_{kj},
\end{aligned}$$

where (5.19) is true due to Assumption 4, and we denote $\varpi^k = \frac{\phi_n(\alpha_k)}{\sqrt{n}} f'(\phi_n(\alpha_k))$. Thus, we turn it into a function of γ_{kj} . Since we have proven above that given $\sum_{j=1}^{p-M} f(\tilde{\lambda}_j)$, the limiting distribution of γ_{kj} is concerned only with the limit of $\sum_{j=1}^{p-M} f(\tilde{\lambda}_j)$, as is $\sum_{k=1}^K \sum_{j \in J_k} \varpi^k \gamma_{kj}$, accordingly, we can conclude that $\sum_{j=1}^M f(\lambda_j)$ and $\sum_{j=1}^{p-M} f(\tilde{\lambda}_j)$ are asymptotically independent. The proof is complete. \square

In the following lemma, we derive the asymptotic distribution of the LSS generated from submatrix \mathbf{S}_{22} .

LEMMA 5.5. Define $Q_1 = \sum_{j=1}^{p-M} f_1(\tilde{\lambda}_j) - (p-M) \int f_1(x) dF^{c_{nM}, H_{2n}}$; then, under Assumptions 1–4, we have

$$\kappa_1^{-1} (Q_1 - \mu_1) \xrightarrow{d} N(0, 1)$$

with mean function

$$\begin{aligned}
\mu_1 &= -\frac{\alpha_x}{2\pi i} \cdot \oint_{\mathcal{C}} f_1(z) \frac{c_{nM} \int \underline{m}_{2n0}^3(z) t^2 (1 + t \underline{m}_{2n0}(z))^{-3} dH_{2n}(t)}{\left(1 - c_{nM} \int \frac{\underline{m}_{2n0}^2(z) t^2}{(1 + t \underline{m}_{2n0}(z))^2} dH_{2n}(t)\right) \left(1 - \alpha_x c_{nM} \int \frac{\underline{m}_{2n0}^2(z) t^2}{(1 + t \underline{m}_{2n0}(z))^2} dH_{2n}(t)\right)} dz \\
&\quad - \frac{\beta_x}{2\pi i} \cdot \oint_{\mathcal{C}} f_1(z) \frac{c_{nM} \int \underline{m}_{2n0}^3(z) t^2 (1 + t \underline{m}_{2n0}(z))^{-3} dH_{2n}(t)}{1 - c_{nM} \int \underline{m}_{2n0}^2(z) t^2 (1 + t \underline{m}_{2n0}(z))^{-2} dH_{2n}(t)} dz,
\end{aligned}$$

and the covariance function is

$$\kappa_1^2 = -\frac{1}{4\pi^2} \oint_{\mathcal{C}_1} \oint_{\mathcal{C}_2} f_1(z_1) f_1(z_2) \vartheta_n^2 dz_1 dz_2,$$

where $\vartheta_n^2 = \Theta_{0,n}(z_1, z_2) + \alpha_x \Theta_{1,n}(z_1, z_2) + \beta_x \Theta_{2,n}(z_1, z_2)$,

$$\Theta_{0,n}(z_1, z_2) = \frac{\underline{m}'_{2n0}(z_1) \underline{m}'_{2n0}(z_2)}{(\underline{m}_{2n0}(z_1) - \underline{m}_{2n0}(z_2))^2} - \frac{1}{(z_1 - z_2)^2},$$

$$\begin{aligned}\Theta_{1,n}(z_1, z_2) &= \frac{\partial}{\partial z_2} \left\{ \frac{\partial \mathcal{A}_n(z_1, z_2)}{\partial z_1} \frac{1}{1 - \alpha_x \mathcal{A}_n(z_1, z_2)} \right\}, \\ \mathcal{A}_n(z_1, z_2) &= \frac{z_1 z_2}{n} \underline{m}'_{2n0}(z_1) \underline{m}'_{2n0}(z_2) \text{tr} \mathbf{\Gamma}^* \mathbf{P}_n(z_1) \mathbf{\Gamma} \mathbf{\Gamma}' \mathbf{P}_n(z_2)' \bar{\mathbf{\Gamma}}, \\ \Theta_{2,n}(z_1, z_2) &= \frac{z_1^2 z_2^2 \underline{m}'_{2n0}(z_1) \underline{m}'_{2n0}(z_2)}{n} \sum_{i=1}^p [\mathbf{\Gamma}^* \mathbf{P}_n^2(z_1) \mathbf{\Gamma}]_{ii} [\mathbf{\Gamma}^* \mathbf{P}_n^2(z_2) \mathbf{\Gamma}]_{ii},\end{aligned}$$

and the definitions of \mathbf{P}_n , $\mathbf{\Gamma}$, and \underline{m}_{2n0} are defined in Section 3.

PROOF. From [Zheng et al. \(2015\)](#), we have that under Assumptions 1–4, the random variable $(\kappa_1^0)^{-1} (Q_1 - \mu_1) \xrightarrow{d} N(0, 1)$, with mean function

$$\begin{aligned}\mu_1 &= -\frac{\alpha_x}{2\pi i} \cdot \oint_{\mathcal{C}} \frac{f_1(z) c_{nM} \int \underline{m}_{2n0}^3(z) t^2 (1 + t \underline{m}_{2n0}(z))^{-3} dH_{2n}(t)}{\left(1 - c_{nM} \int \frac{\underline{m}_{2n0}^2(z) t^2}{(1 + t \underline{m}_{2n0}(z))^2} dH_{2n}(t)\right) \left(1 - \alpha_x c_{nM} \int \frac{\underline{m}_{2n0}^2(z) t^2}{(1 + t \underline{m}_{2n0}(z))^2} dH_{2n}(t)\right)} dz \\ &\quad - \frac{\beta_x}{2\pi i} \cdot \oint_{\mathcal{C}} \frac{f_1(z) c_{nM} \int \underline{m}_{2n0}^3(z) t^2 (1 + t \underline{m}_{2n0}(z))^{-3} dH_{2n}(t)}{1 - c_{nM} \int \underline{m}_{2n0}^2(z) t^2 (1 + t \underline{m}_{2n0}(z))^{-2} dH_{2n}(t)} dz,\end{aligned}$$

and the covariance function is

$$(\kappa_1^0)^2 = -\frac{1}{4\pi^2} \oint_{\mathcal{C}_1} \oint_{\mathcal{C}_2} f_1(z_1) f_1(z_2) (\vartheta_n^0)^2 dz_1 dz_2,$$

where

$$\begin{aligned}(\vartheta_n^0)^2 &= \frac{b_n(z_1) b_n(z_2)}{n^2} \sum_{j=1}^n \text{tr} \mathbb{E}_j \mathbf{\Gamma} \mathbf{\Gamma}^* \mathbf{A}_j^{-1}(z_1) \mathbb{E}_j \left(\mathbf{\Gamma} \mathbf{\Gamma}^* \mathbf{A}_j^{-1}(z_2) \right) \\ &\quad + \frac{\alpha_x b_n(z_1) b_n(z_2)}{n^2} \sum_{j=1}^n \text{tr} \mathbb{E}_j \mathbf{\Gamma}^* \mathbf{A}_j^{-1}(z_1) \mathbf{\Gamma} \mathbb{E}_j \left(\mathbf{\Gamma}' (\mathbf{A}_j')^{-1}(z_2) \bar{\mathbf{\Gamma}} \right) \\ &\quad + \frac{\beta_x b_n(z_1) b_n(z_2)}{n^2} \sum_{j=1}^n \sum_{i=1}^p \mathbf{e}_i' \mathbf{\Gamma}^* \mathbf{A}_j^{-1}(z_1) \mathbf{\Gamma} \mathbf{e}_i \cdot \mathbf{e}_i' \mathbf{\Gamma}^* \mathbf{A}_j^{-1}(z_2) \mathbf{\Gamma} \mathbf{e}_i,\end{aligned}$$

where $b_n(z) = \frac{1}{1 + n^{-1} \mathbb{E} \text{tr} \mathbf{\Gamma} \mathbf{\Gamma}^* \mathbf{A}_j^{-1}(z)}$. The notation $\mathbf{A}_j, \mathbf{e}_i$ is defined in Section 2. Moreover [Najim and Yao \(2016\)](#) provided an estimation ϑ_n^2 for $(\vartheta_n^0)^2$ and proved that $(\vartheta_n^0)^2$ is close to ϑ_n^2 in the Lévy–Prohorov distance, where $\vartheta_n^2 = \Theta_{0,n}(z_1, z_2) + \alpha_x \Theta_{1,n}(z_1, z_2) + \beta_x \Theta_{2,n}(z_1, z_2)$,

$$\begin{aligned}\Theta_{0,n}(z_1, z_2) &= \frac{\underline{m}'_{2n0}(z_1) \underline{m}'_{2n0}(z_2)}{(\underline{m}_{2n0}(z_1) - \underline{m}_{2n0}(z_2))^2} - \frac{1}{(z_1 - z_2)^2}, \\ \Theta_{1,n}(z_1, z_2) &= \frac{\partial}{\partial z_2} \left\{ \frac{\partial \mathcal{A}_n(z_1, z_2)}{\partial z_1} \frac{1}{1 - \alpha_x \mathcal{A}_n(z_1, z_2)} \right\}, \\ \mathcal{A}_n(z_1, z_2) &= \frac{z_1 z_2}{n} \underline{m}'_{2n0}(z_1) \underline{m}'_{2n0}(z_2) \text{tr} \mathbf{\Gamma}^* \mathbf{P}_n(z_1) \mathbf{\Gamma} \mathbf{\Gamma}' \mathbf{P}_n(z_2)' \bar{\mathbf{\Gamma}}, \\ \Theta_{2,n}(z_1, z_2) &= \frac{z_1^2 z_2^2 \underline{m}'_{2n0}(z_1) \underline{m}'_{2n0}(z_2)}{n} \sum_{i=1}^p [\mathbf{\Gamma}^* \mathbf{P}_n^2(z_1) \mathbf{\Gamma}]_{ii} [\mathbf{\Gamma}^* \mathbf{P}_n^2(z_2) \mathbf{\Gamma}]_{ii}.\end{aligned}$$

The definitions of \mathbf{P}_n , $\mathbf{\Gamma}$, and \underline{m}_{2n0} are defined in Section 3. Notably, if $\mathbf{\Gamma}$ is not real, the convergence of $\Theta_{1,n}(z_1, z_2)$ is not guaranteed. However, if $\mathbf{\Gamma}$ and entries x_{ij} are real, that is,

$\alpha_x = 1$, then it can be easily proven that $\Theta_{0,n}(z_1, z_2) = \Theta_{1,n}(z_1, z_2)$. Similarly, the convergence of $\Theta_{2,n}(z_1, z_2)$ depends on the assumption that $\mathbf{\Gamma}^*\mathbf{\Gamma}$ is diagonal; thus, under Assumptions 1–4, $\Theta_{1,n}(z_1, z_2)$ and $\Theta_{2,n}(z_1, z_2)$ may have no limits.

Thus, the covariance term $(\kappa_1^0)^2$ is estimable, and the estimation is κ_1^2 , with

$$\kappa_1^2 = -\frac{1}{4\pi^2} \oint_{\mathcal{C}_1} \oint_{\mathcal{C}_2} f_1(z_1) f_1(z_2) \vartheta_n^2 dz_1 dz_2.$$

Therefore, the proof is finished. \square

Acknowledgments. The authors would like to thank Editor, one Associate Editor and two referees for their constructive comments. Zhidong Bai was partially supported by NSFC Grant NO.12171198 and Team Project of Jilin Provincial Department of Science and Technology NO.20210101147JC. Jiang Hu was partially supported by NSFC Grant NO.12171078.

REFERENCES

- Theodore Wilbur Anderson. *An Introduction to Multivariate Statistical Analysis. Third Edition.* Wiley New York, 2003.
- Jushan Bai and Serena Ng. Determining the number of factors in approximate factor models. *Econometrica*, 70(1):191–221, 2002.
- Zhidong Bai, Baiqi Miao, and Guangming Pan. On asymptotics of eigenvectors of large sample covariance matrix. *The Annals of Probability*, 35(4):1532–1572, 2007.
- Zhidong Bai, Jiang Hu, Guangming Pan, and Wang Zhou. Convergence of the empirical spectral distribution function of beta matrices. *Bernoulli*, 21(3):1538–1574, 2015.
- Zhidong Bai, Huiqin Li, and Guangming Pan. Central limit theorem for linear spectral statistics of large dimensional separable sample covariance matrices. *Bernoulli*, 25(3):1838–1869, 2019.
- Zhidong Bai and Xue Ding. Estimation of spiked eigenvalues in spiked models. *Random Matrices: Theory and Applications*, 1(02):1150011, 2012.
- Zhidong Bai and Jack W. Silverstein. No eigenvalues outside the support of the limiting spectral distribution of large-dimensional sample covariance matrices. *The Annals of Probability*, 26(1):316–345, 1998.
- Zhidong Bai and Jack W. Silverstein. CLT for linear spectral statistics of large-dimensional sample covariance matrices. *The Annals of Probability*, 32(1A):553 – 605, 2004.
- Zhidong Bai and Jianfeng Yao. Central limit theorems for eigenvalues in a spiked population model. *Annales de l’Institut Henri Poincaré, Probabilités et Statistiques*, 44(3):447 – 474, 2008.
- Zhidong Bai and Jianfeng Yao. On sample eigenvalues in a generalized spiked population model. *Journal of Multivariate Analysis*, 106:167–177, 2012.
- Zhidong Bai, Dandan Jiang, Jianfeng Yao, and Shurong Zheng. Corrections to LRT on large-dimensional covariance matrix by RMT. *The Annals of Statistics*, 37(6B):3822 – 3840, 2009.
- Jinho Baik and Jack W. Silverstein. Eigenvalues of large sample covariance matrices of spiked population models. *Journal of Multivariate Analysis*, 97(6):1382–1408, 2006.
- Jinho Baik, Ji Oon Lee, and Hao Wu. Ferromagnetic to paramagnetic transition in spherical spin glass. *Journal of Statistical Physics*, 173(5):1484–1522, 2018.
- Badi H. Baltagi, Chihwa Kao, and Fa Wang. Asymptotic power of the sphericity test under weak and strong factors in a fixed effects panel data model. *Econometric Reviews*, 36(6-9):853–882, 2017.
- Marwa Banna, Jamal Najim, and Jianfeng Yao. A clt for linear spectral statistics of large random information-plus-noise matrices. *Stochastic Processes and their Applications*, 130(4):2250–2281, 2020.
- Alex Bloemendal, Antti Knowles, Horng-Tzer Yau, and Jun Yin. On the principal components of sample covariance matrices. *Probability Theory and Related Fields*, 164(1-2):459–552, 2016.
- T. Tony Cai, Xiao Han, and Guangming Pan. Limiting laws for divergent spiked eigenvalues and largest nonspiked eigenvalue of sample covariance matrices. *The Annals of Statistics*, 48(3):1255–1280, 2020.
- Gary Chamberlain and Michael Rothschild. Arbitrage, factor structure, and mean-variance analysis in large asset markets. *Econometrica*, 51:1305–1324, 1983.
- Binbin Chen and Guangming Pan. Clt for linear spectral statistics of normalized sample covariance matrices with the dimension much larger than the sample size. *Bernoulli*, 21(2):1089–1133, 2015.
- Edgar Dobriban. Permutation methods for factor analysis and pca. *The Annals of Statistics*, 48(5):2824–2847, 2020.

- David Donoho, Matan Gavish, and Iain Johnstone. Optimal shrinkage of eigenvalues in the spiked covariance model. *The Annals of Statistics*, 46(4):1742–1778, 2018.
- Jiti Gao, Xiao Han, Guangming Pan, and Yanrong Yang. High dimensional correlation matrices: The central limit theorem and its applications. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 79(3):677–693, 2017.
- Jiang Hu, Weiming Li, Zhi Liu, and Wang Zhou. High-dimensional covariance matrices in elliptical distributions with application to spherical test. *The Annals of Statistics*, 47(1):527–555, 2019.
- Zhiqiang Hou, Yan Liu, Zhidong Bai and Jiang Hu. Approximation of the power functions of Roy’s largest root test under general spiked alternatives. *Random Matrices: Theory and Applications*, 10(1), 2019.
- Dandan Jiang and Zhidong Bai. Generalized four moment theorem and an application to clt for spiked eigenvalues of high-dimensional covariance matrices. *Bernoulli*, 27(1):274–294, 2021.
- Tiefeng Jiang and Fan Yang. Central limit theorems for classical likelihood ratio tests for high-dimensional normal distributions. *The Annals of Statistics*, 41(4):2029–2074, 2013.
- Iain M. Johnstone. On the distribution of the largest eigenvalue in principal components analysis. *The Annals of Statistics*, 29(2):295 – 327, 2001.
- Iain M. Johnstone and Boaz Nadler. Roy’s largest root test under rank-one alternatives. *Biometrika*, 104(1): 181–193, 2017.
- Iain M. Johnstone and Alexei Onatski. Testing in high-dimensional spiked models. *The Annals of Statistics*, 48(3):1231–1254, 2020.
- Iain M. Johnstone and Debashis Paul. PCA in high dimensions: An orientation. *Proceedings of the IEEE*, 106(8): 1277–1292, 2018.
- Dag Jonsson. Some limit theorems for the eigenvalues of a sample covariance matrix. *Journal of Multivariate Analysis*, 12(1):1–38, 1982.
- Sungkyu Jung and J. S. Marron. PCA consistency in high dimension, low sample size context. *The Annals of Statistics*, 37(6B):4104–4130, 2009.
- Olivier Ledoit and Michael Wolf. Some hypothesis tests for the covariance matrix when the dimension is large compared to the sample size. *The Annals of Statistics*, 30(4):1081–1102, 2002.
- Zeng Li, Fang Han, and Jianfeng Yao. Asymptotic joint distribution of extreme eigenvalues and trace of large sample covariance matrix in a generalized spiked population model. *The Annals of Statistics*, 48(6):3138–3160, 2020.
- Zeng Li, Qinwen Wang, and Runze Li. Central limit theorem for linear spectral statistics of large dimensional kendall’s rank correlation matrices and its applications. *The Annals of Statistics*, 49(3):1569–1593, 2021.
- Boaz Nadler. Finite sample approximation results for principal component analysis: A matrix perturbation approach. *The Annals of Statistics*, 36(6):2791–2817, 2008.
- Hisao Nagao. On Some Test Criteria for Covariance Matrix *The Annals of Statistics* 1(4):700–709, 1973.
- Jamal Najim and Jianfeng Yao. Gaussian fluctuations for linear spectral statistics of large random covariance matrices. *The Annals of Applied Probability*, 26(3):1837–1887, 2016.
- Alexei Onatski, Marcelo J. Moreira, and Marc Hallin. Asymptotic power of sphericity tests for high-dimensional data. *The Annals of Statistics*, 41(3):1204–1231, 2013.
- Alexei Onatski, Marcelo J. Moreira, and Marc Hallin. Signal detection in high dimension: The multispiked case. *The Annals of Statistics*, 42(1):225–254, 2014.
- Guangming Pan. Comparison between two types of large sample covariance matrices. *Annales de l’Institut Henri Poincaré, Probabilités et Statistiques*, 50(2):655–677, 2014.
- Guangming Pan and Wang Zhou. Central limit theorem for signal-to-interference ratio of reduced rank linear receiver. *The Annals of Applied Probability*, 18(3):1232–1270, 2008.
- Debashis Paul. Asymptotics of sample eigenstructure for a large dimensional spiked covariance model. *Statistica Sinica*, 17(4):1617–1642, 2007.
- Amelia Perry, Alexander S. Wein, Afonso S. Bandeira, and Ankur Moitra. Optimality and sub-optimality of pca i: Spiked random matrix models. *The Annals of Statistics*, 46(5):2416–2451, 2018.
- Jack W. Silverstein. Strong convergence of the empirical distribution of eigenvalues of large dimensional random matrices. *Journal of Multivariate Analysis*, 55(2):331–339, 1995.
- Qinwen Wang and Jianfeng Yao. On the sphericity test with large-dimensional observations. *Electronic Journal of Statistics*, 7:2164–2192, 2013.
- Qinwen Wang and Jianfeng Yao. Extreme eigenvalues of large-dimensional spiked fisher matrices with application. *The Annals of Statistics*, 45(1):415–460, 2017.
- Weichen Wang and Jianqing Fan. Asymptotics of empirical eigenstructure for high dimensional spiked covariance. *The Annals of Statistics*, 45(3):1342–1374, 2017.
- S. S. Wilks. The Large-Sample Distribution of the Likelihood Ratio for Testing Composite Hypotheses. *The Annals of Mathematical Statistics*, 9(1):60–62, 1938.

- Jeha Yang and Iain M. Johnstone. Edgeworth correction for the largest eigenvalue in a spiked pca model. *Statistica Sinica*, 2018.
- Yanrong Yang and Guangming Pan. Independence test for high dimensional data based on regularized canonical correlation coefficients. *The Annals of Statistics*, 43(2):467–500, 2015.
- Jianfeng Yao, Shurong Zheng, and Zhidong Bai. *Large Sample Covariance Matrices and High-Dimensional Data Analysis*. Cambridge University Press, 2015.
- Zhigang Yao, Ye Zhang, Zhidong Bai, and William F. Eddy. Estimating the number of sources in magnetoencephalography using spiked population eigenvalues. *Journal of the American Statistical Association*, 113(522):505–518, 2018.
- Yanqing Yin. Spectral statistics of high dimensional sample covariance matrix with unbounded population spectral norm. *accepted by Bernoulli*, 2021.
- Zhixiang Zhang, Shurong Zheng, Guangming Pan, and Pingshou Zhong. Asymptotic independence of spiked eigenvalues and linear spectral statistics for large sample covariance matrices. *accepted by The Annals of Statistics*, 2022.
- Shurong Zheng. Central limit theorems for linear spectral statistics of large dimensional \mathbf{F} -matrices. *Annales de l’Institut Henri Poincaré, Probabilités et Statistiques*, 48(2):444–476, 2012.
- Shurong Zheng, Zhidong Bai, and Jianfeng Yao. Substitution principle for CLT of linear spectral statistics of high-dimensional sample covariance matrices with applications to hypothesis testing. *The Annals of Statistics*, 43(2):546 – 591, 2015.
- Shurong Zheng, Zhidong Bai, and Jianfeng Yao. CLT for eigenvalue statistics of large-dimensional general fisher matrices with applications. *Bernoulli*, 23(2):1130–1178, 2017.
- Shurong Zheng, Guanghui Cheng, Jianhua Guo, and Hongtu Zhu. Test for high-dimensional correlation matrices. *The Annals of Statistics*, 47(5):2887–2921, 2019.
- Yihui Zhou and James Stephen Marron. High dimension low sample size asymptotics of robust PCA. *Electronic Journal of Statistics*, 9(1):204–218, 2015.

SUPPLEMENTARY MATERIAL FOR: A CLT FOR THE LSS OF LARGE DIMENSIONAL SAMPLE COVARIANCE MATRICES WITH DIVERGING SPIKES

BY ZHIJUN LIU^{*}, JIANG HU[†], ZHIDONG BAI[‡], AND HAIYAN SONG[§]

KLASMOE and School of Mathematics and Statistics, Northeast Normal University, China. ^{*}liuzj037@nenu.edu.cn;
[†]huj156@nenu.edu.cn; [‡]baizd@nenu.edu.cn; [§]songhy716@nenu.edu.cn

In this document we present many of the technical details from the contribution [Liu et al. \(2022\)](#). The number of scheme(equations, theorems, lemmas, etc.) is shared with the main document so that there are no misunderstandings with the use of references.

6. Proof of Theorems and Corollaries.

6.1. *Proof of Theorem 3.1.* The proof of Theorem 3.1 builds on the decomposition analysis of the LSS and we split the LSS into part (I) $\sum_{j=1}^M f(\lambda_j)$ and part (II) $\sum_{j=M+1}^p f(\lambda_j)$. Enlightened by the BST in [Bai and Silverstein \(2004\)](#), we have

$$\begin{aligned} & \sum_{j=1}^p f(\lambda_j) - p \int f(x) dF^{c_n, H_n} \\ &= \sum_{j=1}^M f(\lambda_j) + \sum_{j=M+1}^p f(\lambda_j) - p \int f(x) dF^{c_n, H_n} \\ &= \sum_{j=1}^M f(\lambda_j) + \sum_{j=1}^{p-M} f(\tilde{\lambda}_j) - (p-M) \int f(x) dF^{c_{nM}, H_{2n}} + \sum_{j=M+1}^p f(\lambda_j) - \sum_{j=1}^{p-M} f(\tilde{\lambda}_j) \\ &+ (p-M) \int f(x) dF^{c_{nM}, H_{2n}} - p \int f(x) dF^{c_n, H_n}. \end{aligned}$$

Since Lemma 5.1 has shown the difference between $(p-M) \int f(x) dF^{c_{nM}, H_{2n}}$ and $p \int f(x) dF^{c_n, H_n}$ is 0. Moreover, in Lemma 5.2 we have proved

$$\sum_{j=M+1}^p f(\lambda_j) - \sum_{j=1}^{p-M} f(\tilde{\lambda}_j) = \frac{M}{2\pi i} \oint_C f(z) \frac{\underline{m}'(z)}{\underline{m}(z)} dz + o_p(1).$$

It follows that

$$\begin{aligned} & \sum_{j=1}^p f(\lambda_j) - p \int f(x) dF^{c_n, H_n} \\ &= \sum_{j=1}^M f(\lambda_j) + \sum_{j=1}^{p-M} f(\tilde{\lambda}_j) - (p-M) \int f(x) dF^{c_{nM}, H_{2n}} + \frac{M}{2\pi i} \oint_C f(z) \frac{\underline{m}'(z)}{\underline{m}(z)} dz + o_p(1), \end{aligned}$$

MSC2020 subject classifications: Primary 60B20; secondary 60F05.

Keywords and phrases: Empirical spectral distribution, linear spectral statistic, random matrix, Stieltjes transform.

which yields

(6.20)

$$\sum_{j=1}^p f(\lambda_j) - p \int f(x) dF^{c_n, H_n} - \sum_{k=1}^K d_k f(\phi_n(\alpha_k)) - \frac{M}{2\pi i} \oint_C f(z) \frac{m'(z)}{\underline{m}(z)} dz$$

(6.21)

$$= \sum_{j=1}^M f(\lambda_j) - \sum_{k=1}^K d_k f(\phi_n(\alpha_k)) + \sum_{j=1}^{p-M} f(\tilde{\lambda}_j) - (p-M) \int f(x) dF^{c_{nM}, H_{2n}} + o_p(1).$$

The analysis below is executed by dividing (6.21) into two parts: (I) $\sum_{j=1}^M f(\lambda_j) - \sum_{k=1}^K d_k f(\phi_n(\alpha_k))$ and (II) $\sum_{j=1}^{p-M} f(\tilde{\lambda}_j) - (p-M) \int f(x) dF^{c_{nM}, H_{2n}}$, where we ignore the impact of $o_p(1)$ on the asymptotic distribution. Since we have derived the asymptotic distribution of part (II) in Lemma 5.5, we only need to consider the asymptotic distribution of part (I) $\sum_{j=1}^M f(\lambda_j) - \sum_{k=1}^K d_k f(\phi_n(\alpha_k))$. From the proof of Lemma 5.4, $\sum_{j=1}^M f(\lambda_j) - \sum_{k=1}^K d_k f(\phi_n(\alpha_k))$ has the same limiting distribution as $\sum_{k=1}^K \frac{\phi_n(\alpha_k)}{\sqrt{n}} f'(\phi_n(\alpha_k)) \sum_{j \in J_k} \gamma_{kj}$. From Lemma 5.3, we have $(\gamma_{kj}, j \in J_k)' \xrightarrow{d} -\frac{1}{\theta_k} [\mathbf{\Omega}_{\phi_k}]_{kk}$, so $\sum_{j \in J_k} \gamma_{kj} \xrightarrow{d} -\frac{1}{\theta_k} \text{tr}[\mathbf{\Omega}_{\phi_k}]_{kk}$. Recall that ω_{ij} is the element of $\mathbf{\Omega}_{\phi_k}$, and $\text{tr}[\mathbf{\Omega}_{\phi_k}]_{kk}$ is the summation of the diagonal element, that is, $\sum_{j \in J_k} \omega_{jj}$. Because the diagonal elements are i.i.d., $\mathbb{E}(\sum_{j \in J_k} \omega_{jj}) = 0$, $\text{Var}(\sum_{j \in J_k} \omega_{jj}) = \sum_{j \in J_k} \text{Var}(\omega_{jj}) + \sum_{j_1 \neq j_2} \text{cov}(\omega_{j_1 j_1}, \omega_{j_2 j_2}) = \sum_{j \in J_k} ((\alpha_x + 1)\theta_k + \beta_x \mathcal{U}_{jjjj} \nu_k) + \sum_{j_1 \neq j_2} \beta_x \mathcal{U}_{j_1 j_1 j_2 j_2} \nu_k$.

Therefore, from Lemma 5.3, we obtain that the asymptotic distribution of $\sum_{j \in J_k} \gamma_{kj}$ is a Gaussian distribution with

$$\mathbb{E} \sum_{j \in J_k} \gamma_{kj} = 0,$$

$$s_k^2 \triangleq \text{Var} \left(\sum_{j \in J_k} \gamma_{kj} \right) = \frac{\sum_{j \in J_k} ((\alpha_x + 1)\theta_k + \beta_x \mathcal{U}_{jjjj} \nu_k) + \sum_{j_1 \neq j_2} \beta_x \mathcal{U}_{j_1 j_1 j_2 j_2} \nu_k}{\theta_k^2},$$

and then, we directly derive that the mean function of $\sum_{k=1}^K \frac{\phi_n(\alpha_k)}{\sqrt{n}} f'(\phi_n(\alpha_k)) \sum_{j \in J_k} \gamma_{kj}$ is 0 and that its covariance function is

$$\text{Var}(Y_{f_1}) = \sum_{k=1}^K \frac{\phi_n^2(\alpha_k)}{n} (f_1'(\phi_n(\alpha_k)))^2 s_k^2.$$

Finally, we focus on the asymptotic distribution of equation (6.21). Because of Lemma 5.4, the two LSSs are asymptotically independent; thus, the random variable

$$\varsigma_1^{-1} (Y_1 - \mathbb{E}Y_1) \xrightarrow{d} N(0, 1),$$

with mean function $\mathbb{E}Y_1 = \mu_1$ being

$$\begin{aligned} & -\frac{\alpha_x}{2\pi i} \cdot \oint_C f_1(z) \frac{c_{nM} \int \underline{m}_{2n0}^3(z) t^2 (1 + t \underline{m}_{2n0}(z))^{-3} dH_{2n}(t)}{\left(1 - c_{nM} \int \frac{\underline{m}_{2n0}^2(z) t^2}{(1 + t \underline{m}_{2n0}(z))^2} dH_{2n}(t)\right) \left(1 - \alpha_x c_{nM} \int \frac{\underline{m}_{2n0}^2(z) t^2}{(1 + t \underline{m}_{2n0}(z))^2} dH_{2n}(t)\right)} dz \\ & -\frac{\beta_x}{2\pi i} \cdot \oint_C f_1(z) \frac{c_{nM} \int \underline{m}_{2n0}^3(z) t^2 (1 + t \underline{m}_{2n0}(z))^{-3} dH_{2n}(t)}{1 - c_{nM} \int \underline{m}_{2n0}^2(z) t^2 (1 + t \underline{m}_{2n0}(z))^{-2} dH_{2n}(t)} dz, \end{aligned}$$

and covariance function ς_1^2 being

$$\sum_{k=1}^K \frac{\phi_n^2(\alpha_k)}{n} (f_1'(\phi_n(\alpha_k)))^2 s_k^2 - \frac{1}{4\pi^2} \oint_{\mathcal{C}_1} \oint_{\mathcal{C}_2} f_1(z_1) f_1(z_2) (\vartheta_n)^2 dz_1 dz_2,$$

where ϑ_n^2 is defined in Lemma 5.5. Therefore, the proof is finished.

6.2. *Proof of Theorem 3.2.* Similar to the proof of Theorem 3.1, we divide the LSSs into two parts. Different from the above analysis, in this section, we focus on the multidimensional case under Assumptions 1–6. Recall that we defined

$$G_n(x) = p [F^{\mathbf{B}}(x) - F^{c_n, H_n}(x)],$$

$$Y_l = \int f_l(x) dG_n(x) - \sum_{k=1}^K d_k(\phi_n(\alpha_k)) - \frac{M}{2\pi i} \oint_{\mathcal{C}} f_l(z) \frac{\underline{m}'(z)}{\underline{m}(z)} dz.$$

Because of equation (6.21), the random vector (Y_1, \dots, Y_h) shares the same asymptotic distribution with the summation of two random vectors

$$\left(\sum_{k=1}^K \frac{\phi_n(\alpha_k)}{\sqrt{n}} f_1'(\phi_n(\alpha_k)) \sum_{j \in J_k} \gamma_{kj}, \dots, \sum_{k=1}^K \frac{\phi_n(\alpha_k)}{\sqrt{n}} f_h'(\phi_n(\alpha_k)) \sum_{j \in J_k} \gamma_{kj} \right)$$

and

$$\left(\sum_{j=1}^{p-M} f_1(\tilde{\lambda}_j) - (p-M) \int f_1(x) dF^{c_{nM}, H_{2n}}, \dots, \sum_{j=1}^{p-M} f_h(\tilde{\lambda}_j) - (p-M) \int f_h(x) dF^{c_{nM}, H_{2n}} \right).$$

First, we focus on the first random vector. Similar to the proof of Theorem 3.1, we derive that the mean function of the first random vector is 0 and that the covariance function is

$$\text{Cov}(Y_{f_s}, Y_{f_t}) = \sum_{k=1}^K \frac{\phi_n^2(\alpha_k)}{n} f_s'(\phi_n(\alpha_k)) f_t'(\phi_n(\alpha_k)) s_k^2 = \sum_{k=1}^K \varpi_s^k \varpi_t^k s_k^2,$$

where $\varpi_t^k = \frac{\phi_n(\alpha_k)}{\sqrt{n}} f_t'(\phi_n(\alpha_k))$. Moreover, the asymptotic distribution of the second random vector is derived in Zheng et al. (2015). Because of Lemma 5.4, two random vectors are asymptotically independent; thus, the random vector

$$(Y_1 - \mathbb{E}Y_1, \dots, Y_h - \mathbb{E}Y_h)' \xrightarrow{d} N_h(0, \mathbf{\Omega}),$$

with mean function $\mathbb{E}Y_l$ is the same as μ_l , and the covariance matrix is $\mathbf{\Omega}$ with its entries

$$\omega_{st} = \sum_{k=1}^K \varpi_s^k \varpi_t^k s_k^2 + \kappa_{st},$$

where

$$\begin{aligned} \kappa_{st} = & -\frac{1}{4\pi^2} \oint_{\mathcal{C}_1} \oint_{\mathcal{C}_2} \frac{f_s(z_1) f_t(z_2)}{(\underline{m}(z_1) - \underline{m}(z_2))^2} d\underline{m}(z_1) d\underline{m}(z_2) - \frac{c\beta_x}{4\pi^2} \oint_{\mathcal{C}_1} \oint_{\mathcal{C}_2} f_s(z_1) f_t(z_2) \\ & \left[\int \frac{t}{(\underline{m}(z_1)t + 1)^2} \times \frac{t}{(\underline{m}(z_2)t + 1)^2} dH(t) \right] d\underline{m}(z_1) d\underline{m}(z_2) \end{aligned}$$

$$-\frac{1}{4\pi^2} \oint_{C_1} \oint_{C_2} f_s(z_1) f_t(z_2) \left[\frac{\partial^2}{\partial z_1 \partial z_2} \log(1 - a(z_1, z_2)) \right] dz_1 dz_2,$$

$$a(z_1, z_2) = \alpha_x \left(1 + \frac{\underline{m}(z_1) \underline{m}(z_2) (z_1 - z_2)}{\underline{m}(z_2) - \underline{m}(z_1)} \right).$$

Then, we obtain the random vector

$$\left(\frac{Y_1 - \mathbb{E}Y_1}{\sqrt{\sigma_1^2}}, \dots, \frac{Y_h - \mathbb{E}Y_h}{\sqrt{\sigma_h^2}} \right)' \xrightarrow{d} N_h(0, \Psi),$$

which has a mean function that is the same as that in Theorem 3.1, and variance function

$$\sigma_l^2 = \sum_{k=1}^K \frac{\phi_n^2(\alpha_k)}{n} (f'_l(\phi_n(\alpha_k)))^2 s_k^2 + \kappa_{ll}, \quad l = 1, \dots, h,$$

and the covariance matrix $\Psi = (\psi_{st})_{h \times h}$ is the correlation coefficient matrix of random vector $(Y_1, \dots, Y_h)'$ with its entries

$$\psi_{st} = \frac{\sum_{k=1}^K \varpi_s^k \varpi_t^k s_k^2 + \kappa_{st}}{\sqrt{\sum_{k=1}^K (\varpi_s^k)^2 s_k^2 + \kappa_{ss}} \sqrt{\sum_{k=1}^K (\varpi_t^k)^2 s_k^2 + \kappa_{tt}}}.$$

Note that renormalization is necessary to guarantee that elements in the correlation coefficient matrix Ψ are limited. Therefore, the proof is finished.

6.3. *Proof of Theorem 4.1.* Now we prove the Theorem 4.1. Recall that

$$G_n(x) = p [F^{\mathbf{B}}(x) - F^{c_n, H_n}(x)],$$

$$Y = \int f_U(x) dG_n(x) - \sum_{k=1}^K d_k f_U(\phi_n(\alpha_k)) - \frac{M}{2\pi i} \oint_{\mathcal{C}} f_U(z) \frac{\underline{m}'(z)}{\underline{m}(z)} dz.$$

When $f_U(x) = \log(1+x)$, after some calculations, we obtain

$$\int f_U(x) dG_n(x) = U - p \int f_U(x) dF^{c_n, H_n}(x) = U - (p - M) \int f_U(x) dF^{c_{nM}, H_{2n}}(x),$$

$$\int f_U(x) dF^{c_{nM}, H_{2n}}(x) = \log \left(1 + \frac{c_{nM} + \sqrt{c_{nM}^2 + 4}}{2} \right) +$$

(6.22)

$$\frac{- \left(\sqrt{c_{nM}} - \frac{1}{\sqrt{c_{nM}}} \right)^2 (\log(1 - \sqrt{\tilde{c}_{nM} c_{nM}}) + \sqrt{\tilde{c}_{nM} c_{nM}}) - \sqrt{\tilde{c}_{nM}} (\sqrt{c_{nM}} - (\sqrt{c_{nM}})^3)}{1 - c_{nM}},$$

$$\sum_{k=1}^K d_k f_U(\phi_n(\alpha_k)) = \sum_{k=1}^K d_k \log(1 + \phi_n(\alpha_k)),$$

$$(6.23) \quad \frac{M}{2\pi i} \oint_{\mathcal{C}} f_U(z) \frac{\underline{m}'(z)}{\underline{m}(z)} dz = M \log(1 - \sqrt{\tilde{c}_n c_n}).$$

For consistency, we present the proofs of (6.22) and (6.23) in Section 7. According to Theorem 3.1, when $f_U(x) = \log(1+x)$, we have

$$\frac{U - (p-M) \int f_U(x) dF^{c_{nM}, H_{2n}}(x) - \mu_U}{\sqrt{\varsigma_U^2}} \xrightarrow{d} N(0, 1),$$

where

$$\begin{aligned} \mu_U &= \alpha_x I_1(f_U) + \beta_x I_2(f_U) + \sum_{k=1}^K d_k \log \left(1 + \alpha_k + \frac{\alpha_k}{\alpha_k - 1} c_{nM} \right) + M \log \left(1 - \sqrt{\tilde{c}_n c_n} \right), \\ \varsigma_U^2 &= \sum_{k=1}^K \frac{\left(\alpha_k + \frac{\alpha_k}{\alpha_k - 1} c_{nM} \right)^2}{n \left(1 + \alpha_k + \frac{\alpha_k}{\alpha_k - 1} c_{nM} \right)^2} s_k^2 + (\alpha_x + 1) J_1(f_U, f_U) + \beta_x J_2(f_U, f_U), \end{aligned} \quad (6.24)$$

$$I_1(f_U) = \log \left(1 + \frac{c_{nM} + \sqrt{c_{nM}^2 + 4}}{2} \right) - \log(2 + c_{nM}) + \sum_{k=1}^{\infty} \left(\frac{\sqrt{c_{nM}}}{2 + c_{nM}} \right)^{2k} \frac{(2k-1)!}{k!k!}, \quad (6.25)$$

$$I_2(f_U) = - \sum_{k=1}^{\infty} \left(\frac{\sqrt{c_{nM}}}{2 + c_{nM}} \right)^{2k} \frac{(2k-1)!}{(k-1)!(k+1)!}, \quad (6.26)$$

$$J_1(f_U, f_U) = J_2(f_U, f_U) = \left(\sum_{k=1}^{\infty} \left(\frac{\sqrt{c_{nM}}}{2 + c_{nM}} \right)^{2k-1} \frac{(2k-2)!}{k!(k-1)!} \right)^2,$$

$$\tilde{c}_n = \frac{4c_n}{\left(2 + c_n + \sqrt{c_n^2 + 4} \right)^2}.$$

Here for consistency we postpone the proofs of (6.24)–(6.26) to Section 7, and therefore the proof is finished.

6.4. *Proof of Corollary 4.1.* As the normalized U statistic tends to a standard normal distribution under H_0 , that is,

$$\frac{U - p \int f_U(x) dF^{c_n, H_n} - \mu_U^0}{\varsigma_U^0} \xrightarrow{d} N(0, 1),$$

where

$$p \int f_U(x) dF^{c_n, H_n} = pCT(\varrho(c_n), c_n, \tilde{c}_n),$$

$$\mu_U^0 = \alpha_x I_1^0(f_U) + \beta_x I_2^0(f_U), \varsigma_U^0 = \sqrt{(\alpha_x + 1) J_1^0(f_U, f_U) + \beta_x J_2^0(f_U, f_U)},$$

$$I_1^0(f_U) = \log \left(1 + \frac{c_n + \sqrt{c_n^2 + 4}}{2} \right) - \log(2 + c_n) + \sum_{k=1}^{\infty} \left(\frac{\sqrt{c_n}}{2 + c_n} \right)^{2k} \frac{(2k-1)!}{k!k!},$$

$$I_2^0(f_U) = - \sum_{k=1}^{\infty} \left(\frac{\sqrt{c_n}}{2 + c_n} \right)^{2k} \frac{(2k-1)!}{(k-1)!(k+1)!},$$

$$J_1^0(f_U, f_U) = J_2^0(f_U, f_U) = \left(\sum_{k=1}^{\infty} \left(\frac{\sqrt{c_n}}{2 + c_n} \right)^{2k-1} \frac{(2k-2)!}{k!(k-1)!} \right)^2.$$

Then we can obtain that $P_{H_0}(|U| > w) = P_{H_0}\left(\frac{|U| - U_0}{\varsigma_U^0} > \frac{w - U_0}{\varsigma_U^0}\right) = \xi$, where $U_0 = p \int f_U(x) dF^{c_n, H_n} + \mu_U^0$, then critical value $w = \varsigma_U^0 z_\xi + U_0$, where z_ξ is the upper quantile of standard normal distribution at level ξ . Define $U_1 = (p - M) \int f_U(x) dF^{c_{nM}, H_{2n}} + \mu_U$, then combined with Theorem 4.1, we have that the power of test U under H_1 or H_2 equals

$$\begin{aligned} P(|U| > w) &= P\left(\frac{|U| - U_1}{\varsigma_U} > \frac{w - U_1}{\varsigma_U}\right) = \Phi\left(-\frac{\varsigma_U^0 z_\xi + U_0 - U_1}{\varsigma_U}\right) \\ &= \Phi\left(\frac{U_1 - U_0 - \varsigma_U^0 z_\xi}{\varsigma_U}\right). \end{aligned}$$

Since $(p - M) \int f_U(x) dF^{c_{nM}, H_{2n}} - p \int f_U(x) dF^{c_n, H_n} = (p - M) CT(\varrho(c_{nM}), c_{nM}, \tilde{c}_{nM}) - p CT(\varrho(c_n), c_n, \tilde{c}_n)$, $\mu_U - \mu_U^0 = \alpha_x [I_1(f_U) - I_1^0(f_U)] + \beta_x [I_2(f_U) - I_2^0(f_U)] +$

$\sum_{k=1}^K \log(1 + \phi_n(\alpha_k)) + M \log(1 - \sqrt{\tilde{c}_n c_n})$ and $I_1(f_U) - I_1^0(f_U)$, $I_2(f_U) - I_2^0(f_U)$ tend to 0 as n tends to infinity, then the proof is finished.

6.5. *Proof of Theorem 4.2.* Since

$$Y = \int f_W(x) dG_n(x) - \sum_{k=1}^K d_k f_W(\phi_n(\alpha_k)) - \frac{M}{2\pi i} \oint_{\mathcal{C}} f_W(z) \frac{\underline{m}'(z)}{\underline{m}(z)} dz.$$

$$G_n(x) = p [F^{\mathbf{B}}(x) - F^{c_n, H_n}(x)].$$

When $f_W(x) = x$, we obtain

$$p \int f_W(x) dF^{c_n, H_n}(x) = (p - M) \int f_W(x) dF^{c_{nM}, H_{2n}}(x) = p - M,$$

$$(6.27) \quad \sum_{k=1}^K d_k f_W(\phi_n(\alpha_k)) = \sum_{k=1}^K d_k \phi_n(\alpha_k),$$

$$(6.28) \quad \frac{M}{2\pi i} \oint_{\mathcal{C}} f_W(z) \frac{\underline{m}'(z)}{\underline{m}(z)} dz = -Mc,$$

For consistency, we postpone the proof of (6.28) to Section 7. According to Theorem 3.1, when $f_W(x) = x$, we have

$$\frac{W - (p - M) \int f_W(x) dF^{c_{nM}, H_{2n}}(x) - \mu_W}{\sqrt{\varsigma_W^2}} \xrightarrow{d} N(0, 1),$$

where

$$\begin{aligned} \mu_W &= \sum_{k=1}^K d_k \phi_n(\alpha_k) - M c_n, \\ \varsigma_W^2 &= \sum_{k=1}^K \frac{\phi_n^2(\alpha_k)}{n} s_k^2 + \alpha_x c_{nM} + \beta_x c_{nM} + c_{nM}. \end{aligned}$$

where μ_W and ς_W^2 can be deduced from Wang and Yao (2013), (6.27), and (6.28), therefore the proof of Theorem 4.2 is finished.

6.6. *Proof of Corollary 4.2.* As the normalized W statistic tends to a standard normal distribution under H_0 , that is,

$$\frac{W - p \int f_W(x) dF^{c_n, H_n}}{\varsigma_W^0} \xrightarrow{d} N(0, 1),$$

where

$$\begin{aligned} p \int f_W(x) dF^{c_n, H_n} &= p, \\ \varsigma_W^0 &= \sqrt{(\alpha_x + 1) J_1^0(f_W, f_W) + \beta_x J_2^0(f_W, f_W)}, \\ J_1^0(f_W, f_W) &= J_2^0(f_W, f_W) = c_n. \end{aligned}$$

Then we can obtain that $P_{H_0}(|W| > w) = P_{H_0}\left(\frac{|W| - W_0}{\varsigma_W^0} > \frac{w - W_0}{\varsigma_W^0}\right) = \xi$, where $W_0 = p \int f_W(x) dF^{c_n, H_n}$, then critical value $w = \varsigma_W^0 z_\xi + W_0$, where z_ξ is the upper quantile of standard normal distribution at level ξ . Define $W_1 = (p - M) \int f_W(x) dF^{c_{nM}, H_{2n}} + \mu_W$, then combined with Theorem 4.2, we have that the power of test W under H_1 or H_2 equals

$$\begin{aligned} P(|W| > w) &= P\left(\frac{|W| - W_1}{\varsigma_W} > \frac{w - W_1}{\varsigma_W}\right) = \Phi\left(-\frac{\varsigma_W^0 z_\xi + W_0 - W_1}{\varsigma_W}\right) \\ &= \Phi\left(\frac{W_1 - W_0 - \varsigma_W^0 z_\xi}{\varsigma_W}\right). \end{aligned}$$

Since $(p - M) \int f_W(x) dF^{c_{nM}, H_{2n}} - p \int f_W(x) dF^{c_n, H_n} = -M$, $\mu_W = \sum_{k=1}^K d_k \phi_n(\alpha_k) - M c_n$, therefore $W_1 - W_0 = \sum_{k=1}^K d_k \phi_n(\alpha_k) - M c_n - M$, then the proof is finished.

6.7. *Proof of Theorem 4.3.* Since

$$Y = \int f_V(x) dG_n(x) - \sum_{k=1}^K d_k f_V(\phi_n(\alpha_k)) - \frac{M}{2\pi i} \oint_{\mathcal{C}} f_V(z) \frac{\underline{m}'(z)}{\underline{m}(z)} dz.$$

$$G_n(x) = p [F^{\mathbf{B}}(x) - F^{c_n, H_n}(x)].$$

When $f_V(x) = \frac{x}{1+x}$, we obtain

$$(6.29) \quad p \int f_V(x) dF^{c_n, H_n}(x) = (p - M) \int f_V(x) dF^{c_{nM}, H_{2n}}(x) = \frac{p - M}{1 + \varrho(c_{nM})},$$

$$\sum_{k=1}^K d_k f_V(\phi_n(\alpha_k)) = \sum_{k=1}^K d_k \frac{\phi_n(\alpha_k)}{1 + \phi_n(\alpha_k)},$$

$$(6.30) \quad \frac{M}{2\pi i} \oint_{\mathcal{C}} f_V(z) \frac{\underline{m}'(z)}{\underline{m}(z)} dz = -\frac{M(c_n - 2)}{2(1 + \varrho(c_n))(1 - \tilde{c}_n)} - \frac{M}{2}.$$

For consistency, we postpone the proofs of (6.29) and (6.30) to Section 7. According to Theorem 3.1, when $f_V(x) = \frac{x}{1+x}$, we have

$$\frac{V - (p - M) \int f_V(x) dF^{c_{nM}, H_{2n}}(x) - \mu_V}{\sqrt{\varsigma_V^2}} \xrightarrow{d} N(0, 1),$$

where

$$\mu_V = \alpha_x I_1(f_V) + \beta_x I_2(f_V) + \sum_{k=1}^K d_k \frac{\phi_n(\alpha_k)}{1 + \phi_n(\alpha_k)} - \frac{M(c_n - 2)}{(2 + c_n + \sqrt{c_n^2 + 4})(1 - \tilde{c}_n)} - \frac{M}{2},$$

$$\varsigma_V^2 = \sum_{k=1}^K \frac{\phi_n^2(\alpha_k)}{n(1 + \phi_n(\alpha_k))^4} s_k^2 + (\alpha_x + 1) J_1(f_V, f_V) + \beta_x J_2(f_V, f_V),$$

(6.31)

$$I_1(f_V) = \frac{1}{2 + c_{nM}} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c_{nM}}}{2 + c_{nM}} \right)^{2k} \frac{(2k)!}{k!k!} - \frac{1}{1 + \varrho(c_{nM})} \left(\frac{\tilde{c}_{nM}}{-(1 - \tilde{c}_{nM})^2} + \frac{1}{2(\sqrt{\tilde{c}_{nM}} - 1)^2} + \frac{1}{2(\sqrt{\tilde{c}_{nM}} + 1)^2} \right),$$

(6.32)

$$I_2(f_V) = -\frac{1}{2 + c_{nM}} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c_{nM}}}{2 + c_{nM}} \right)^{2k} \frac{(2k)!}{(k-1)!(k+1)!},$$

(6.33)

$$J_1(f_V, f_V) = J_2(f_V, f_V) = \left(\frac{1}{2 + c_{nM}} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c_{nM}}}{2 + c_{nM}} \right)^{2k+1} \frac{(2k+1)!}{k!(k+1)!} \right)^2.$$

We postpone the proofs of (6.31)–(6.33) to Section 7, then the proof is finished.

6.8. *Proof of Corollary 4.3.* As the normalized V statistic tends to a standard normal distribution under H_0 , that is,

$$\frac{V - p \int f_V(x) dF^{c_n, H_n} - \mu_V^0}{\varsigma_V^0} \xrightarrow{d} N(0, 1),$$

where

$$p \int f_V(x) dF^{c_n, H_n} = \frac{p}{1 + \varrho(c_n)},$$

$$\mu_V^0 = \alpha_x I_1^0(f_V) + \beta_x I_2^0(f_V), \varsigma_V^0 = \sqrt{(\alpha_x + 1) J_1^0(f_V, f_V) + \beta_x J_2^0(f_V, f_V)},$$

$$I_1^0(f_V) = \frac{1}{2 + c_n} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c_n}}{2 + c_n} \right)^{2k} \frac{(2k)!}{k!k!} -$$

$$\frac{1}{1 + \varrho(c_n)} \left(\frac{\tilde{c}_n}{-(1 - \tilde{c}_n)^2} + \frac{1}{2(\sqrt{\tilde{c}_n} - 1)^2} + \frac{1}{2(\sqrt{\tilde{c}_n} + 1)^2} \right),$$

$$I_2^0(f_V) = -\frac{1}{2 + c_n} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c_n}}{2 + c_n} \right)^{2k} \frac{(2k)!}{(k-1)!(k+1)!},$$

$$J_1^0(f_V, f_V) = J_2^0(f_V, f_V) = \left(\frac{1}{2 + c_n} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c_n}}{2 + c_n} \right)^{2k+1} \frac{(2k+1)!}{k!(k+1)!} \right)^2.$$

Then we can obtain that $P_{H_0}(|V| > w) = P_{H_0}\left(\frac{|V| - V_0}{\varsigma_V^0} > \frac{w - V_0}{\varsigma_V^0}\right) = \xi$, where $V_0 = p \int f_V(x) dF^{c_n, H_n} + \mu_V^0$, then critical value $w = \varsigma_V^0 z_\xi + V_0$, where z_ξ is the upper quantile of standard normal distribution at level ξ . Define $V_1 = (p - M) \int f_V(x) dF^{c_{nM}, H_{2n}} + \mu_V$, then combined with Theorem 4.3, we have that the power of test V under H_1 or H_2 equals

$$\begin{aligned} P(|V| > w) &= P\left(\frac{|V| - V_1}{\varsigma_V} > \frac{w - V_1}{\varsigma_V}\right) = \Phi\left(-\frac{\varsigma_V^0 z_\xi + V_0 - V_1}{\varsigma_V}\right) \\ &= \Phi\left(\frac{V_1 - V_0 - \varsigma_V^0 z_\xi}{\varsigma_V}\right). \end{aligned}$$

Since $(p - M) \int f_V(x) dF^{c_{nM}, H_{2n}} - p \int f_V(x) dF^{c_n, H_n} = \frac{p - M}{1 + \varrho(c_{nM})} - \frac{p}{1 + \varrho(c_n)}$, $\mu_V - \mu_V^0 = \alpha_x [I_1(f_V) - I_1^0(f_V)] + \beta_x [I_2(f_V) - I_2^0(f_V)] + \sum_{k=1}^K d_k \frac{\phi_n(\alpha_k)}{1 + \phi_n(\alpha_k)} - \frac{M(c_n - 2)}{2(1 + \varrho(c_n))(1 - \tilde{c}_n)} - \frac{M}{2}$ and $I_1(f_V) - I_1^0(f_V)$, $I_2(f_V) - I_2^0(f_V)$ tend to 0 as n tends to infinity, then the proof is finished.

6.9. *Proof of Theorem 4.4.* Under H_3 , $U = \sum_{j=1}^p \log(1 + \lambda_j) = \sum_{j=1}^p \log\left(1 + \frac{\lambda_j}{l}\right)$, where $\frac{\lambda_j}{l}$ could be treated as the sample eigenvalues with the population covariance matrix \mathbf{I}_p , and test function can be treated as $\log(1 + lx)$. Then we can use Remark 3.3 to obtain the asymptotic distribution of U under H_3 . We obtain

$$\frac{U - p \int f(x) dF^{c_n, H_n}(x) - \mu_U}{\sqrt{\varsigma_U^2}} \xrightarrow{d} N(0, 1),$$

where

$$(6.34) \quad \int f(x) dF^{c_n, H_n}(x) = \log(1 + \varepsilon(c_n)) + \frac{-\left(\sqrt{c_n} - \frac{1}{\sqrt{c_n}}\right)^2 (\log(1 - \sqrt{\tilde{c}_n c_n}) + \sqrt{\tilde{c}_n c_n}) - \sqrt{\tilde{c}_n} (\sqrt{c_n} - (\sqrt{c_n})^3)}{1 - c_n},$$

$$\mu_U = \alpha_x I_1(f_U) + \beta_x I_2(f_U), \quad \varsigma_U^2 = (\alpha_x + 1) J_1(f_U, f_U) + \beta_x J_2(f_U, f_U),$$

$$(6.35) \quad I_1(f_U) = \log(1 + \varepsilon(c_n)) - \log(1 + l + lc_n) + \sum_{k=1}^{\infty} \left(\frac{l\sqrt{c_n}}{1 + l + lc_n}\right)^{2k} \frac{(2k - 1)!}{k!k!},$$

$$(6.36) \quad I_2(f_U) = -\sum_{k=1}^{\infty} \left(\frac{l\sqrt{c_n}}{1 + l + lc_n}\right)^{2k} \frac{(2k - 1)!}{(k - 1)!(k + 1)!},$$

$$(6.37) \quad J_1(f_U, f_U) = J_2(f_U, f_U) = \left(\sum_{k=1}^{\infty} \left(\frac{l\sqrt{c_n}}{1 + l + lc_n}\right)^{2k-1} \frac{(2k - 2)!}{k!(k - 1)!}\right)^2.$$

For consistency, the proofs of formulas (6.34)–(6.37) are postponed to Section 7, then the proof is finished.

6.10. *Proof of Corollary 4.4.* As the normalized U statistic tends to a standard normal distribution under H_0 , that is,

$$\frac{U - p \int f_U(x) dF^{c_n, H_n} - \mu_U^0}{\varsigma_U^0} \xrightarrow{d} N(0, 1),$$

where the formulas of $p \int f_U(x) dF^{c_n, H_n}, \mu_U^0, \varsigma_U^0$ are given in the proof of Corollary 4.1. Then we can obtain that $P_{H_0}(|U| > w) = P_{H_0}\left(\frac{|U| - U_0}{\varsigma_U^0} > \frac{w - U_0}{\varsigma_U^0}\right) = \xi$, where $U_0 = p \int f_U(x) dF^{c_n, H_n} + \mu_U^0$, then critical value $w = \varsigma_U^0 z_\xi + U_0$, where z_ξ is the upper quantile of standard normal distribution at level ξ . Define $U_1 = p \int f(x) dF^{c_n, H_n} + \mu_U$, then combined with Theorem 4.4, we derive that the power of test U under H_3 equals

$$\begin{aligned} P_{H_3}(|U| > w) &= P_{H_3}\left(\frac{|U| - U_1}{\varsigma_U} > \frac{w - U_1}{\varsigma_U}\right) = \Phi\left(-\frac{\varsigma_U^0 z_\xi + U_0 - U_1}{\varsigma_U}\right) \\ &= \Phi\left(\frac{U_1 - U_0 - \varsigma_U^0 z_\xi}{\varsigma_U}\right). \end{aligned}$$

Since $p \int f(x) dF^{c_n, H_n} - p \int f_U(x) dF^{c_n, H_n} = pCT(\varepsilon(c_n), c_n, \check{c}_n) - pCT(\varrho(c_n), c_n, \tilde{c}_n)$, $\mu_U - \mu_U^0 = \alpha_x [I_1(f_U) - I_1^0(f_U)] + \beta_x [I_2(f_U) - I_2^0(f_U)]$, by some calculations, it equals

$$\begin{aligned} &\alpha_x \left[\log\left(\frac{1 + \varepsilon(c_n)}{1 + \varrho(c_n)}\right) + \log\left(\frac{2 + c_n}{1 + l + lc_n}\right) + \right. \\ &\quad \left. \frac{(2k-1)!}{k!k!} \left(\sum_{k=1}^{\infty} \left(\frac{l\sqrt{c_n}}{1 + l + lc_n}\right)^{2k} - \sum_{k=1}^{\infty} \left(\frac{\sqrt{c_n}}{2 + c_n}\right)^{2k} \right) \right] + \\ &\beta_x \left[\frac{(2k-1)!}{(k+1)!(k-1)!} \left(\sum_{k=1}^{\infty} \left(\frac{\sqrt{c_n}}{2 + c_n}\right)^{2k} - \sum_{k=1}^{\infty} \left(\frac{\sqrt{lc_n}}{1 + l + lc_n}\right)^{2k} \right) \right], \end{aligned}$$

then the proof is finished.

6.11. *Proof of Theorem 4.5.* Under H_3 , $W = \sum_{j=1}^p \lambda_j = l \sum_{j=1}^p \frac{\lambda_j}{l}$, where $\frac{\lambda_j}{l}$ could be treated as the sample eigenvalues with the population covariance matrix \mathbf{I}_p . From Wang and Yao (2013), we have

$$\sum_{j=1}^p \frac{\lambda_j}{l} - p \xrightarrow{d} N(0, (\alpha_x + \beta_x + 1)c).$$

Based on delta method,

$$\frac{W - lp}{l\sqrt{(\alpha_x + \beta_x + 1)c_n}} \xrightarrow{d} N(0, 1),$$

then the proof is finished.

6.12. *Proof of Corollary 4.5.* As the normalized W statistic tends to a standard normal distribution under H_0 , that is,

$$\frac{W - p}{\sqrt{(\alpha_x + 1)c_n + \beta_x c_n}} \xrightarrow{d} N(0, 1),$$

Then we can obtain that $P_{H_0}(|W| > w) = P_{H_0}\left(\frac{|W| - p}{\varsigma_W^0} > \frac{w - p}{\varsigma_W^0}\right) = \xi$, then critical value $w = \varsigma_W^0 z_\xi + p$, where z_ξ is the upper quantile of standard normal distribution at level ξ . Then combined with Theorem 4.5, we derive that the power of test W under H_3 equals

$$\begin{aligned} P_{H_3}(|W| > w) &= P_{H_3}\left(\frac{|W| - lp}{\varsigma_W} > \frac{w - lp}{\varsigma_W}\right) = \Phi\left(-\frac{\varsigma_W^0 z_\xi + p - lp}{\varsigma_W}\right) \\ &= \Phi\left(\frac{lp - p - \varsigma_W^0 z_\xi}{\varsigma_W}\right) = \Phi\left(\frac{(l-1)p}{l\sqrt{(\alpha_x + 1 + \beta_x)c_n}} - \frac{z_\xi}{l}\right), \end{aligned}$$

then the proof is finished.

6.13. *Proof of Theorem 4.6.* Under H_3 , $V = \sum_{j=1}^p \frac{\lambda_j}{1 + \lambda_j} = \sum_{j=1}^p \frac{\frac{\lambda_j}{l}}{\frac{1}{l} + \frac{\lambda_j}{l}}$, where $\frac{\lambda_j}{l}$ could be treated as the sample eigenvalues with the population covariance matrix \mathbf{I}_p , and test function can be treated as $\frac{x}{\frac{1}{l} + x}$. Then we can use Remark 3.3 to obtain the asymptotic distribution of V under H_3 . Then we have under H_3 ,

$$\frac{V - p \int f(x) dF^{c_n, H_n}(x) - \mu_V}{\sqrt{s_V^2}} \xrightarrow{d} N(0, 1),$$

where

$$(6.38) \quad \int f(x) dF^{c_n, H_n}(x) = \frac{l}{1 + \varepsilon(c_n)},$$

$$\mu_V = \alpha_x I_1(f_V) + \beta_x I_2(f_V), \quad s_V^2 = (\alpha_x + 1) J_1(f_V, f_V) + \beta_x J_2(f_V, f_V),$$

$$(6.39) \quad I_1(f_V) = \frac{1}{1 + l + lc_n} \sum_{k=0}^{\infty} \left(\frac{l\sqrt{c_n}}{1 + l + lc_n} \right)^{2k} \frac{(2k)!}{k!k!} -$$

$$\frac{1}{1 + \varepsilon(c_n)} \left(\frac{\check{c}}{-(1 - \check{c})^2} + \frac{1}{2(\sqrt{\check{c}} - 1)^2} + \frac{1}{2(\sqrt{\check{c}} + 1)^2} \right),$$

$$(6.40) \quad I_2(f_V) = -\frac{1}{1 + l + lc_n} \sum_{k=0}^{\infty} \left(\frac{l\sqrt{c_n}}{1 + l + lc_n} \right)^{2k} \frac{(2k)!}{(k-1)!(k+1)!},$$

$$(6.41) \quad J_1(f_V, f_V) = J_2(f_V, f_V) = \left(\frac{1}{1 + l + lc_n} \sum_{k=0}^{\infty} \left(\frac{l\sqrt{c_n}}{1 + l + lc_n} \right)^{2k+1} \frac{(2k+1)!}{k!(k+1)!} \right)^2,$$

For consistency, the proofs of formulas (6.38)–(6.41) are postponed to Section 7, then the proof is finished.

6.14. *Proof of Corollary 4.6.* As the normalized V statistic tends to a standard normal distribution under H_0 , that is,

$$\frac{V - p \int f_V(x) dF^{c_n, H_n} - \mu_V^0}{s_V^0} \xrightarrow{d} N(0, 1),$$

where the formulas of $p \int f_V(x) dF^{c_n, H_n}$, μ_V^0 , s_V^0 are given in the proof of Corollary 4.3. Then we can obtain that $P_{H_0}(|V| > w) = P_{H_0}\left(\frac{|V| - V_0}{s_V^0} > \frac{w - V_0}{s_V^0}\right) = \xi$, where $V_0 = p \int f_V(x) dF^{c_n, H_n} + \mu_V^0$, then critical value $w = s_V^0 z_\xi + V_0$, where z_ξ is the upper quantile of standard normal distribution at level ξ . Define $V_1 = p \int f(x) dF^{c_n, H_n} + \mu_V$, then combined with Theorem 4.6, we have that the power of test V under H_3 equals

$$\begin{aligned} P_{H_3}(|V| > w) &= P_{H_3}\left(\frac{|V| - V_1}{s_V} > \frac{w - V_1}{s_V}\right) = \Phi\left(-\frac{s_V^0 z_\xi + V_0 - V_1}{s_V}\right) \\ &= \Phi\left(\frac{V_1 - V_0 - s_V^0 z_\xi}{s_V}\right). \end{aligned}$$

Since $p \int f(x) dF^{c_n, H_n} - p \int f_V(x) dF^{c_n, H_n} = \frac{pl}{1+\varepsilon(c_n)} - \frac{p}{1+\varrho(c_n)}$, $\mu_V - \mu_V^0 = \alpha_x [I_1(f_V) - I_1^0(f_V)] + \beta_x [I_2(f_V) - I_2^0(f_V)]$, by some calculations, it equals

$$\begin{aligned} & \alpha_x \left[\frac{(2k)!}{k!k!} \left(\frac{1}{1+l+lc_n} \sum_{k=0}^{\infty} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k} - \frac{1}{2+c_n} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c_n}}{2+c_n} \right)^{2k} \right) + \right. \\ & \frac{1}{1+\varrho(c_n)} \left(\frac{\tilde{c}_n}{-(1-\tilde{c}_n)^2} + \frac{1}{2(\sqrt{\tilde{c}_n}-1)^2} + \frac{1}{2(\sqrt{\tilde{c}_n}+1)^2} \right) - \\ & \left. \frac{1}{1+\varepsilon(c_n)} \left(\frac{\check{c}_n}{-(1-\check{c}_n)^2} + \frac{1}{2(\sqrt{\check{c}_n}-1)^2} + \frac{1}{2(\sqrt{\check{c}_n}+1)^2} \right) \right] + \beta_x \frac{2k!}{(k-1)!(k+1)!} \\ & \left[\frac{1}{2+c_n} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c_n}}{2+c_n} \right)^{2k} - \frac{1}{1+l+lc_n} \sum_{k=0}^{\infty} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k} \right], \end{aligned}$$

$V_1 - V_0 = p \int f(x) dF^{c_n, H_n} - p \int f_V(x) dF^{c_n, H_n} + \mu_V - \mu_V^0$, then the proof is finished.

6.15. *Proof of Theorem 4.7.* Under H_3 , $R = \lambda_1 = l\frac{\lambda_1}{l}$, where $\frac{\lambda_1}{l}$ can be treated as the largest sample eigenvalue with population covariance matrix \mathbf{I}_p . Since under H_0 , we have

$$\frac{\frac{\lambda_1}{l} - \frac{\mu_{n,p}}{n}}{\frac{\sigma_{n,p}}{n}} \xrightarrow{d} TW_1,$$

where TW_1 denotes the type-1 Tracy-Widom distribution. Then we have

$$\frac{\lambda_1 - \frac{l\mu_{n,p}}{n}}{\frac{l\sigma_{n,p}}{n}} \xrightarrow{d} TW_1,$$

where

$$\begin{aligned} \mu_{n,p} &= (\sqrt{n-1} + \sqrt{p})^2, \\ \sigma_{n,p} &= (\sqrt{n-1} + \sqrt{p}) \left(\frac{1}{\sqrt{n-1}} + \frac{1}{\sqrt{p}} \right)^{1/3}, \end{aligned}$$

therefore the proof is finished.

6.16. *Proof of Corollary 4.7.* Since $P_{H_0}(|R| > w) = P_{H_0} \left(\frac{|R| - R_0}{\varsigma_R^0} > \frac{w - R_0}{\varsigma_R^0} \right) = \xi$, where R_0 and ς_R^0 are $\frac{\mu_{n,p}}{n}$ and $\frac{\sigma_{n,p}}{n}$, respectively. Then critical value $w = \varsigma_R^0 t_\xi + R_0$, where t_ξ is the upper quantile of type-1 Tracy-Widom distribution at level ξ . Then combined with Theorem 4.7, we derive that the power of test R under H_3 equals

$$\begin{aligned} P_{H_3}(|R| > w) &= P_{H_3} \left(\frac{|R| - R_1}{\varsigma_R} > \frac{w - R_1}{\varsigma_R} \right) = 1 - F_1 \left(\frac{\varsigma_R^0 t_\xi + R_0 - R_1}{\varsigma_R} \right) \\ &= 1 - F_1 \left(\frac{\frac{\sigma_{n,p}}{n} t_\xi + (1-l) \frac{\mu_{n,p}}{n}}{l \frac{\sigma_{n,p}}{n}} \right), \end{aligned}$$

where F_1 is the cumulative distribution function of a type-1 Tracy-Widom distribution variable, then the proof is finished.

7. Some deviations and calculations. *Proof of (6.22):* Since $f_U(x) = \log(1+x)$, then

$$\int f_U(x) dF^{c_n, H_n}(x) = \int_{a(c_n)}^{b(c_n)} \log(1+x) \frac{1}{2\pi x c_n} \sqrt{(b(c_n) - x)(x - a(c_n))} dx,$$

where $a(c_n) = (1 - \sqrt{c_n})^2$, $b(c_n) = (1 + \sqrt{c_n})^2$. By using the variable change $x = 1 + c_n - 2\sqrt{c_n} \cos(\theta)$, $0 \leq \theta \leq \pi$, we have

$$\begin{aligned} \int f_U(x) dF^{c_n, H_n}(x) &= \frac{1}{2\pi c_n} \int_0^\pi \frac{\log(2 + c_n - 2\sqrt{c_n} \cos(\theta))}{1 + c_n - 2\sqrt{c_n} \cos(\theta)} 4c_n \sin^2(\theta) d\theta \\ &= \frac{1}{2\pi} \int_0^{2\pi} \frac{2 \sin^2(\theta)}{1 + c_n - 2\sqrt{c_n} \cos(\theta)} \log(2 + c_n - 2\sqrt{c_n} \cos(\theta)) d\theta. \end{aligned}$$

Do the transformation $2 + c_n - 2\sqrt{c_n} \cos(\theta) = (1 + \varrho(c_n)) \left(\frac{2 + c_n}{1 + \varrho(c_n)} - \frac{2\sqrt{c_n}}{1 + \varrho(c_n)} \cos(\theta) \right)$,

let $\frac{2 + c_n}{1 + \varrho(c_n)} = 1 + \tilde{c}_n$, $\frac{\sqrt{c_n}}{1 + \varrho(c_n)} = \sqrt{\tilde{c}_n}$, then we obtain $\varrho(c_n) = \frac{c_n + \sqrt{c_n^2 + 4}}{2}$, $\tilde{c}_n =$

$\frac{4c_n}{(2 + c_n + \sqrt{c_n^2 + 4})^2}$. Therefore $\int f_U(x) dF^{c_n, H_n}(x)$ equals

$$\begin{aligned} &\frac{1}{2\pi} \int_0^{2\pi} \frac{2 \sin^2(\theta) \log(1 + \varrho(c_n))}{1 + c_n - 2\sqrt{c_n} \cos(\theta)} d\theta + \frac{1}{2\pi} \int_0^{2\pi} \frac{2 \sin^2(\theta)}{1 + c_n - 2\sqrt{c_n} \cos(\theta)} \\ &\log \left(\frac{2 + c_n}{1 + \varrho(c_n)} - \frac{2\sqrt{c_n}}{1 + \varrho(c_n)} \cos(\theta) \right) d\theta. \end{aligned}$$

For the first integral,

$$\begin{aligned} &\frac{1}{2\pi} \int_0^{2\pi} \frac{2 \sin^2(\theta) \log(1 + \varrho(c_n))}{1 + c_n - 2\sqrt{c_n} \cos(\theta)} d\theta \\ &= \frac{\log(1 + \varrho(c_n))}{2\pi} \int_0^{2\pi} \frac{2 \sin^2(\theta)}{1 + c_n - 2\sqrt{c_n} \cos(\theta)} d\theta \\ &= -\frac{1}{2} \frac{\log(1 + \varrho(c_n))}{2\pi i} \oint_{|z|=1} \frac{(z - \frac{1}{z})^2}{(z - \sqrt{c_n})(1 - \sqrt{c_n}z)} dz \\ &= -\frac{1}{2} \frac{\log(1 + \varrho(c_n))}{2\pi i} \oint_{|z|=1} \frac{z^4 - 2z^2 + 1}{z^2(z - \sqrt{c_n})(1 - \sqrt{c_n}z)} dz \end{aligned}$$

Since $c_n < 1$, thus $\sqrt{c_n}$ and 0 are poles. The residues are $\frac{1 - c_n}{c_n}$ and $-\frac{1 + c_n}{c_n}$, respectively.

By residue theorem, we obtain the first integral is $\log \left(1 + \frac{c_n + \sqrt{c_n^2 + 4}}{2} \right)$.

For the second integral,

$$\begin{aligned} &\frac{1}{2\pi} \int_0^{2\pi} \frac{2 \sin^2(\theta)}{1 + c_n - 2\sqrt{c_n} \cos(\theta)} \log \left(\frac{2 + c_n}{1 + \varrho(c_n)} - \frac{2\sqrt{c_n}}{1 + \varrho(c_n)} \cos(\theta) \right) d\theta \\ &= \frac{1}{2\pi} \int_0^{2\pi} \frac{2 \sin^2(\theta)}{1 + c_n - 2\sqrt{c_n} \cos(\theta)} \log \left(1 + \tilde{c}_n - 2\sqrt{\tilde{c}_n} \cos(\theta) \right) d\theta \end{aligned}$$

$$= \frac{-\left(\sqrt{c_n} - \frac{1}{\sqrt{c_n}}\right)^2 (\log(1 - \sqrt{c_n c_n}) + \sqrt{c_n c_n}) - \sqrt{c_n} (\sqrt{c_n} - (\sqrt{c_n})^3)}{1 - c_n},$$

where the last integral is calculated in [Bai and Silverstein \(2004\)](#). Note that all the c_n in the formulas above should be replaced by c_{nM} since the calculation is on the bulk part of LSS. Collecting the two integrals leads to the desired formula for $p \int f_U(x) dF^{c_n, H_n}(x)$.

Proof of (6.23): First, we consider $\oint_C f_U(z) \frac{m'(z)}{\underline{m}(z)} dz$.

$$\begin{aligned} \oint_C f_U(z) \frac{m'(z)}{\underline{m}(z)} dz &= \oint_C f_U(z) d \log \underline{m}(z) = - \oint_C f'_U(z) \log \underline{m}(z) dz \\ &= \int_{a(c)}^{b(c)} f'_U(x) [\log \underline{m}(x + i\varepsilon) - \log \underline{m}(x - i\varepsilon)] dx \\ (7.42) \quad &= 2i \int_{a(c)}^{b(c)} f'_U(x) \Im \log \underline{m}(x + i\varepsilon) dx. \end{aligned}$$

Here, $a(c) = (1 - \sqrt{c})^2$ and $b(c) = (1 + \sqrt{c})^2$. Since

$$\underline{m}(z) = -\frac{1-c}{z} + cm(z),$$

under H_1 , we have

$$\underline{m}(z) = \frac{-(z+1-c) + \sqrt{(z-1-c)^2 - 4c}}{2z}.$$

As $z \rightarrow x \in [a(c), b(c)]$, we obtain

$$\underline{m}(x) = \frac{-(x+1-c) + \sqrt{4c - (x-1-c)^2}i}{2x}.$$

Therefore,

$$\begin{aligned} &\int_{a(c)}^{b(c)} f'_U(x) \Im \log \underline{m}(x + i\varepsilon) dx \\ &= \int_{a(c)}^{b(c)} f'_U(x) \tan^{-1} \left(\frac{\sqrt{4c - (x-1-c)^2}}{-(x+1-c)} \right) dx \\ &= \left[\tan^{-1} \left(\frac{\sqrt{4c - (x-1-c)^2}}{-(x+1-c)} \right) f_U(x) \Big|_{a(c)}^{b(c)} - \int_{a(c)}^{b(c)} f_U(x) d \tan^{-1} \left(\frac{\sqrt{4c - (x-1-c)^2}}{-(x+1-c)} \right) \right]. \end{aligned}$$

It is easy to verify that the first term is 0, and we now focus on the second term,

$$\begin{aligned} &\int_{a(c)}^{b(c)} f_U(x) d \tan^{-1} \left(\frac{\sqrt{4c - (x-1-c)^2}}{-(x+1-c)} \right) \\ (7.43) \quad &= \int_{a(c)}^{b(c)} \frac{\log(1+x)}{1 + \frac{4c - (x-1-c)^2}{(x+1-c)^2}} \cdot \frac{\sqrt{4c - (x-1-c)^2} + \frac{(x-1-c)(x+1-c)}{\sqrt{4c - (x-1-c)^2}}}{(x+1-c)^2} dx. \end{aligned}$$

By substituting $x = 1 + c - 2\sqrt{c}\cos(\theta)$, we obtain

$$(7.43) = \frac{1}{2} \int_0^{2\pi} (\log(2 + c - 2\sqrt{c}\cos(\theta))) \frac{c - \sqrt{c}\cos(\theta)}{1 + c - 2\sqrt{c}\cos(\theta)} d\theta$$

$$(7.44) = \frac{1}{2} \int_0^{2\pi} \left[\log(1 + \varrho(c)) + \log\left(1 + \sqrt{c} - 2\sqrt{c}\cos(\theta)\right) \right] \frac{c - \sqrt{c}\cos(\theta)}{1 + c - 2\sqrt{c}\cos(\theta)} d\theta$$

$$(7.45) = \frac{1}{2} \int_0^{2\pi} \log(1 + \varrho(c)) \frac{c - \sqrt{c}\cos(\theta)}{1 + c - 2\sqrt{c}\cos(\theta)} d\theta +$$

$$(7.46) \quad \frac{1}{2} \int_0^{2\pi} \log\left(1 + \sqrt{c} - 2\sqrt{c}\cos(\theta)\right) \frac{c - \sqrt{c}\cos(\theta)}{1 + c - 2\sqrt{c}\cos(\theta)} d\theta.$$

For the first integral, by substituting $\cos\theta = \frac{z+z^{-1}}{2}$, we turn it into a contour integral on $|z| = 1$,

$$\begin{aligned} & \frac{1}{2} \int_0^{2\pi} \log(1 + \varrho(c)) \frac{c - \sqrt{c}\cos(\theta)}{1 + c - 2\sqrt{c}\cos(\theta)} d\theta \\ &= \frac{\log(1 + \varrho(c))}{2} \oint_{|z|=1} \frac{c - \sqrt{c}\frac{z+\frac{1}{z}}{2}}{1 + c - \sqrt{c}\left(z + \frac{1}{z}\right)} \frac{1}{iz} dz \\ &= \frac{\log(1 + \varrho(c))}{2i} \oint_{|z|=1} \frac{2cz - \sqrt{c}(z^2 + 1)}{2z(z - \sqrt{c})(1 - \sqrt{c}z)} dz. \end{aligned}$$

When $c < 1$, 0 and \sqrt{c} are poles. The residues are $\frac{1}{2}$ and $-\frac{1}{2}$, respectively. By residue theorem, the first integral is 0 .

For the second integral,

$$\begin{aligned} & \frac{1}{2} \int_0^{2\pi} \frac{\log(1 + \tilde{c} - 2\sqrt{\tilde{c}}\cos(\theta))}{1 + c - 2\sqrt{c}\cos(\theta)} (c - \sqrt{c}\cos\theta) d\theta \\ &= \frac{1}{2} \oint_{|z|=1} \log|1 - \sqrt{\tilde{c}}z|^2 \cdot \frac{c - \sqrt{c}\frac{z+z^{-1}}{2}}{1 + c - 2\sqrt{c}\cdot\frac{z+z^{-1}}{2}} \frac{dz}{iz} \\ &= \frac{1}{4i} \oint_{|z|=1} \log|1 - \sqrt{\tilde{c}}z|^2 \cdot \frac{2cz - \sqrt{c}(z^2 + 1)}{(z - \sqrt{c})(-\sqrt{c}z + 1)z} dz \\ (7.47) \quad &= \frac{1}{4i} \oint_{|z|=1} \log\left(1 - \sqrt{\tilde{c}}z\right) \frac{2cz - \sqrt{c}(z^2 + 1)}{(z - \sqrt{c})(-\sqrt{c}z + 1)z} dz + \frac{1}{4i} \oint_{|z|=1} \log\left(1 - \sqrt{\tilde{c}}\frac{1}{z}\right) \end{aligned}$$

$$(7.48) \quad \frac{2cz - \sqrt{c}(z^2 + 1)}{(z - \sqrt{c})(-\sqrt{c}z + 1)z} dz.$$

For the first term in (7.48), when $c < 1$, the pole is \sqrt{c} , and the residue is $-\log(1 - \sqrt{\tilde{c}c})$. By using the residue theorem, the integral is $-\frac{\pi}{2}\log(1 - \sqrt{\tilde{c}c})$. The same argument also holds for the second term in (7.48), and the integral is also $-\frac{\pi}{2}\log(1 - \sqrt{\tilde{c}c})$ after some calculation. Therefore the second integral equals $-\pi\log(1 - \sqrt{\tilde{c}c})$. Therefore,

$$\frac{M}{2\pi i} \oint_{\mathcal{C}} f_L(z) \frac{\underline{m}'(z)}{\underline{m}(z)} dz = M \log\left(1 - \sqrt{\tilde{c}c}\right),$$

and the result is still valid if c is replaced by c_n . Therefore, formula (6.23) holds.

Proof of (6.24): From Remark 3.3,

$$\begin{aligned} I_1(f_U) &= \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \log\left(1 + |1 + \sqrt{c}z|^2\right) \left(\frac{z}{z^2 - r^{-2}} - \frac{1}{z}\right) dz \\ &= \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \log\left(1 + |1 + \sqrt{c}z|^2\right) \frac{z}{z^2 - r^{-2}} dz - \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \log\left(1 + |1 + \sqrt{c}z|^2\right) \frac{1}{z} dz. \end{aligned}$$

For the first integral of $I_1(f_U)$,

$$\begin{aligned} & \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \log\left(1 + |1 + \sqrt{c}z|^2\right) \frac{z}{z^2 - r^{-2}} dz \\ &= \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \log\left[(1 + \varrho(c)) (1 + \sqrt{c}z) (1 + \sqrt{c}\bar{z})\right] \frac{z}{z^2 - r^{-2}} dz \\ &= \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \log(1 + \varrho(c)) \frac{z}{z^2 - r^{-2}} dz + \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \log(1 + \sqrt{c}z) \frac{z}{z^2 - r^{-2}} dz + \\ & \quad \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \log\left(1 + \sqrt{c}\frac{1}{z}\right) \frac{z}{z^2 - r^{-2}} dz, \end{aligned}$$

where \tilde{c} and $\varrho(c)$ are defined in the proof of (6.22). For the first integral, the poles are $\frac{1}{r}$ and $-\frac{1}{r}$, the residues are both $\frac{1}{2} \log(1 + \varrho(c))$. Therefore, by residue theorem, the integral is $\log(1 + \varrho(c))$. Similarly, for the second integral, the residues are $\frac{1}{2} \log\left(1 + \frac{\sqrt{c}}{r}\right)$ and $\frac{1}{2} \log\left(1 - \frac{\sqrt{c}}{r}\right)$. By the residue theorem, the integral is $\frac{1}{2} \log(1 - \tilde{c})$. For the third integral,

$$\begin{aligned} & \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \log\left(1 + \sqrt{c}\frac{1}{z}\right) \frac{z}{z^2 - r^{-2}} dz \\ &= \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|\xi|=1} \log\left(1 + \sqrt{c}\xi\right) \frac{\frac{1}{\xi}}{\xi^{-2} - r^{-2}} \frac{1}{\xi^2} d\xi \\ &= \lim_{r \downarrow 1} r^2 \frac{1}{2\pi i} \oint_{|\xi|=1} \log\left(1 + \sqrt{c}\xi\right) \frac{1}{\xi(r + \xi)(r - \xi)} d\xi, \end{aligned}$$

where the first integral results from the change of variable $\xi = \frac{1}{z}$. The poles are r and $-r$, and the residues are $-\log(1 + \sqrt{c}r) \frac{1}{2r^2}$ and $-\log(1 - \sqrt{c}r) \frac{1}{2r^2}$, respectively. Then by residue theorem, the integral is $-\frac{1}{2} \log(1 - \tilde{c})$.

Collecting the three integral above leads to $\lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \log\left(1 + |1 + \sqrt{c}z|^2\right) \frac{z}{z^2 - r^{-2}} dz = \log(1 + \varrho(c))$.

For the second integral of $I_1(f_U)$,

$$\begin{aligned} & \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \log\left(1 + |1 + \sqrt{c}z|^2\right) \frac{1}{z} dz \\ &= \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{\log(2 + c) + \log\left(1 + \frac{\sqrt{c}}{2+c} \left(\frac{1}{z} + z\right)\right)}{z} dz \\ (7.49) \quad &= \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{\log(2 + c)}{z} dz + \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{\log\left(1 + \frac{\sqrt{c}}{2+c} \left(\frac{1}{z} + z\right)\right)}{z} dz. \end{aligned}$$

By using Taylor expansion,

$$\begin{aligned}
& \oint_{|z|=1} \frac{\log\left(1 + \frac{\sqrt{c}}{2+c} \left(\frac{1}{z} + z\right)\right)}{z} dz \\
&= \oint_{|z|=1} \sum_{k=1}^{\infty} (-1)^{k+1} \left(\frac{\sqrt{c}}{2+c}\right)^k \frac{1}{k} \left(\frac{1}{z} + z\right)^k \frac{1}{z} dz \\
&= \sum_{k=1}^{\infty} (-1)^{k+1} \left(\frac{\sqrt{c}}{2+c}\right)^k \frac{1}{k} \oint_{|z|=1} \left(\frac{1}{z} + z\right)^k \frac{1}{z} dz \\
&= \sum_{k=1}^{\infty} (-1)^{2k+1} \left(\frac{\sqrt{c}}{2+c}\right)^{2k} \frac{1}{2k} \oint_{|z|=1} \left(\frac{1}{z} + z\right)^{2k} \frac{1}{z} dz \\
&= \sum_{k=1}^{\infty} (-1) \left(\frac{\sqrt{c}}{2+c}\right)^{2k} \frac{1}{2k} C_{2k}^k 2\pi i \\
&= - \sum_{k=1}^{\infty} \left(\frac{\sqrt{c}}{2+c}\right)^{2k} \frac{(2k-1)!}{k!k!} 2\pi i.
\end{aligned}$$

Therefore the second integral equals $\log(2+c) - \sum_{k=1}^{\infty} \left(\frac{\sqrt{c}}{2+c}\right)^{2k} \frac{(2k-1)!}{k!k!}$. Thus $I_1(f_U) = \log(1+\varrho) - \log(2+c) + \sum_{k=1}^{\infty} \left(\frac{\sqrt{c}}{2+c}\right)^{2k} \frac{(2k-1)!}{k!k!}$. Note that all the c in the formula above should be replaced by c_{nM} since the calculation is on the bulk part of LSS. The proof is finished.

Proof of (6.25): From Remark 3.3, by Taylor expansion,

$$\begin{aligned}
I_2(f_U) &= \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \log\left(1 + |1 + \sqrt{c}z|^2\right) \frac{1}{z^3} dz \\
&= \frac{1}{2\pi i} \oint_{|z|=1} \left(\log(2+c) + \log\left(1 + \frac{\sqrt{c}}{2+c} \left(\frac{1}{z} + z\right)\right)\right) \frac{1}{z^3} dz \\
&= \frac{1}{2\pi i} \oint_{|z|=1} \log\left(1 + \frac{\sqrt{c}}{2+c} \left(\frac{1}{z} + z\right)\right) \frac{1}{z^3} dz \\
&= \frac{1}{2\pi i} \oint_{|z|=1} \sum_{k=1}^{\infty} (-1)^{2k+1} \left(\frac{\sqrt{c}}{2+c}\right)^{2k} \frac{1}{2k} \left(\frac{1}{z} + z\right)^{2k} \frac{1}{z^3} dz \\
&= \sum_{k=1}^{\infty} (-1)^{2k+1} \left(\frac{\sqrt{c}}{2+c}\right)^{2k} \frac{1}{2k} \frac{1}{2\pi i} \oint_{|z|=1} \left(\frac{1}{z} + z\right)^{2k} \frac{1}{z^3} dz \\
&= \sum_{k=1}^{\infty} (-1)^{2k+1} \left(\frac{\sqrt{c}}{2+c}\right)^{2k} \frac{1}{2k} C_{2k}^{k-1} \\
&= - \sum_{k=1}^{\infty} \left(\frac{\sqrt{c}}{2+c}\right)^{2k} \frac{(2k-1)!}{(k-1)!(k+1)!}.
\end{aligned}$$

Notice that all the c in the formula above should be replaced by c_{nM} since the calculation is on the bulk part of LSS. The proof is finished.

Proof of (6.26): From Remark 3.3,

$$\begin{aligned} J_1(f_U, f_U) &= \lim_{r \downarrow 1} -\frac{1}{4\pi^2} \oint_{|z_1|=1} \oint_{|z_2|=1} \frac{\log\left(1 + |1 + \sqrt{c}z_1|^2\right) \log\left(1 + |1 + \sqrt{c}z_2|^2\right)}{(z_1 - rz_2)^2} dz_1 dz_2 \\ &= \lim_{r \downarrow 1} -\frac{1}{4\pi^2} \oint_{|z_2|=1} \log\left(1 + |1 + \sqrt{c}z_2|^2\right) \oint_{|z_1|=1} \frac{\log\left(1 + |1 + \sqrt{c}z_1|^2\right)}{(z_1 - rz_2)^2} dz_1 dz_2. \end{aligned}$$

Since $r > 1$, thus rz_2 is not a pole.

$$\begin{aligned} & \oint_{|z_1|=1} \frac{\log\left(1 + |1 + \sqrt{c}z_1|^2\right)}{(z_1 - rz_2)^2} dz_1 \\ &= \oint_{|z_1|=1} \frac{\sum_{k=1}^{\infty} (-1)^{k+1} \left(\frac{\sqrt{c}}{2+c}\right)^k \frac{1}{k} \left(\frac{1}{z_1} + z_1\right)^k}{(z_1 - rz_2)^2} dz_1 \\ &= \oint_{|z_1|=1} \frac{\sum_{k=1}^{\infty} (-1)^{2k} \left(\frac{\sqrt{c}}{2+c}\right)^{2k-1} \frac{1}{2k-1} \left(\frac{1}{z_1} + z_1\right)^{2k-1}}{(z_1 - rz_2)^2} dz_1 \\ &= \sum_{k=1}^{\infty} \left(\frac{\sqrt{c}}{2+c}\right)^{2k-1} \frac{1}{2k-1} \oint_{|z_1|=1} \left(\frac{1}{z_1} + z_1\right)^{2k-1} \frac{1}{(z_1 - rz_2)^2} dz_1 \\ &= \sum_{k=1}^{\infty} \left(\frac{\sqrt{c}}{2+c}\right)^{2k-1} \frac{1}{2k-1} 2\pi i \frac{(2k-1)!}{k!(k-1)!} \frac{1}{r^2 z_2^2} \\ &= \sum_{k=1}^{\infty} \left(\frac{\sqrt{c}}{2+c}\right)^{2k-1} 2\pi i \frac{(2k-2)!}{k!(k-1)!} \frac{1}{r^2 z_2^2}. \end{aligned}$$

Then

$$\begin{aligned} J_1(f_U, f_U) &= \lim_{r \downarrow 1} -\frac{1}{4\pi^2} \sum_{k=1}^{\infty} \left(\frac{\sqrt{c}}{2+c}\right)^{2k-1} 2\pi i \frac{(2k-2)!}{k!(k-1)!} \oint_{|z_2|=1} \log\left(1 + |1 + \sqrt{c}z_2|^2\right) \frac{1}{r^2 z_2^2} dz_2 \\ &= \left(\sum_{k=1}^{\infty} \left(\frac{\sqrt{c}}{2+c}\right)^{2k-1} \frac{(2k-2)!}{k!(k-1)!} \right)^2, \end{aligned}$$

where the integral about z_2 is handled the same way as z_1 . Similarly, from Remark 3.3,

$$J_2(f_U, f_U) = \lim_{r \downarrow 1} -\frac{1}{4\pi^2} \oint_{|z_1|=1} \frac{\log\left(1 + |1 + \sqrt{c}z_1|^2\right)}{z_1^2} dz_1 \oint_{|z_2|=1} \frac{\log\left(1 + |1 + \sqrt{c}z_2|^2\right)}{z_2^2} dz_2.$$

By Taylor expansion, we obtain

$$\begin{aligned} & \oint_{|z_1|=1} \frac{\log\left(1 + |1 + \sqrt{c}z_1|^2\right)}{z_1^2} dz_1 \\ &= \oint_{|z_1|=1} \frac{\log(2+c)}{z_1^2} dz_1 + \oint_{|z_1|=1} \frac{\log\left(1 + \frac{\sqrt{c}}{2+c} \left(\frac{1}{z_1} + z_1\right)\right)}{z_1^2} dz_1 \end{aligned}$$

$$\begin{aligned}
&= \sum_{k=1}^{\infty} (-1)^{k+1} \left(\frac{\sqrt{c}}{2+c} \right)^k \frac{1}{k} \oint_{|z_1|=1} \left(\frac{1}{z_1} + z_1 \right)^k \frac{1}{z_1^2} dz_1 \\
&= \sum_{k=1}^{\infty} (-1)^{2k} \left(\frac{\sqrt{c}}{2+c} \right)^{2k-1} \frac{1}{2k-1} \oint_{|z_1|=1} \left(\frac{1}{z_1} + z_1 \right)^{2k-1} \frac{1}{z_1^2} dz_1 \\
&= 2\pi i \sum_{k=1}^{\infty} \left(\frac{\sqrt{c}}{2+c} \right)^{2k-1} \frac{(2k-2)!}{k!(k-1)!},
\end{aligned}$$

thus

$$J_2(f_U, f_U) = \left(\sum_{k=1}^{\infty} \left(\frac{\sqrt{c}}{2+c} \right)^{2k-1} \frac{(2k-2)!}{k!(k-1)!} \right)^2.$$

Notice that all the c in the formula above should be replaced by c_{nM} since the calculation is on the bulk part of LSS. The proof is finished.

Proof of (6.28): Similar to proof of (6.23), we have

$$\begin{aligned}
\oint_{\mathcal{C}} f_W(z) \frac{m'(z)}{m(z)} dz &= 2i \int_{a(c)}^{b(c)} f'_W(x) \Im \log m(x+i\varepsilon) dx \\
&= -2i \int_{a(c)}^{b(c)} x d \tan^{-1} \left(\frac{\sqrt{4c - (x-1-c)^2}}{-(x+1-c)} \right).
\end{aligned}$$

Since

$$\begin{aligned}
&\int_{a(c)}^{b(c)} x d \tan^{-1} \left(\frac{\sqrt{4c - (x-1-c)^2}}{-(x+1-c)} \right) \\
&= \int_{a(c)}^{b(c)} \frac{x}{1 + \frac{4c - (x-1-c)^2}{(x+1-c)^2}} \cdot \frac{\sqrt{4c - (x-1-c)^2} + \frac{(x-1-c)(x+1-c)}{\sqrt{4c - (x-1-c)^2}}}{(x+1-c)^2} dx.
\end{aligned}$$

By substituting $x = 1 + c - 2\sqrt{c} \cos(\theta)$, it equals

$$\begin{aligned}
&\frac{1}{2} \int_0^{2\pi} (1+c-2\sqrt{c} \cos(\theta)) \frac{c - \sqrt{c} \cos(\theta)}{1+c-2\sqrt{c} \cos(\theta)} d\theta \\
&= \int_0^{\pi} (c - \sqrt{c} \cos(\theta)) d\theta = \pi c.
\end{aligned}$$

Therefore $\oint_{\mathcal{C}} f_W(z) \frac{m'(z)}{m(z)} dz = -2\pi ic$, then $\frac{M}{2\pi i} \oint_{\mathcal{C}} f_W(z) \frac{m'(z)}{m(z)} dz = -Mc$, the proof is finished.

Proof of (6.29): Since $f_V(x) = \frac{x}{1+x}$, then

$$\int f_V(x) dF^{c_n, H_n}(x) = \int_{a(c_n)}^{b(c_n)} \frac{x}{1+x} \frac{1}{2\pi x c_n} \sqrt{(b(c_n) - x)(x - a(c_n))} dx.$$

By using the variable change $x = 1 + c_n - 2\sqrt{c_n} \cos(\theta)$, $0 \leq \theta \leq \pi$, we have

$$\begin{aligned}
\int f_V(x) dF^{c_n, H_n}(x) &= \frac{1}{2\pi c_n} \int_0^{\pi} \frac{1}{2+c_n-2\sqrt{c_n} \cos(\theta)} 4c_n \sin^2(\theta) d\theta \\
&= \frac{1}{4\pi c_n} \int_0^{2\pi} \frac{1}{(1+\varrho(c_n))(1+\tilde{c}_n-2\sqrt{\tilde{c}_n} \cos(\theta))} 4c_n \sin^2(\theta) d\theta
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{4\pi c_n} \frac{1}{1+\varrho(c_n)} \int_0^{2\pi} \frac{1}{1+\tilde{c}_n-2\sqrt{\tilde{c}_n}\cos(\theta)} 4c_n \sin^2(\theta) d\theta \\
&= -\frac{1}{4\pi i} \frac{1}{1+\varrho(c_n)} \oint_{|z|=1} \frac{(z-1)^2(z+1)^2}{z^2(z-\sqrt{\tilde{c}_n})(1-\sqrt{\tilde{c}_n}z)} dz.
\end{aligned}$$

When $\tilde{c}_n < 1$, the poles are 0 and $\sqrt{\tilde{c}_n}$. The residues are $-\frac{1+\tilde{c}_n}{\tilde{c}_n}$ and $\frac{1-\tilde{c}_n}{\tilde{c}_n}$, respectively. Then by Residue theorem, $-\frac{1}{4\pi i} \frac{1}{1+\varrho(c_n)} \oint_{|z|=1} \frac{(z-1)^2(z+1)^2}{z^2(z-\sqrt{\tilde{c}_n})(1-\sqrt{\tilde{c}_n}z)} dz$ equals $\frac{1}{1+\varrho(c_n)}$. Notice that all the c_n in the formulas above should be replaced by c_{nM} . The proof is finished.

Proof of (6.30): Similar to proof of (6.23), we have

$$\begin{aligned}
&\oint_{\mathcal{C}} f_V(z) \frac{m'(z)}{m(z)} dz = 2i \int_{a(c)}^{b(c)} f'_V(x) \Im \log \underline{m}(x+i\varepsilon) dx \\
&= -2i \int_{a(c)}^{b(c)} \frac{x}{1+x} d \tan^{-1} \left(\frac{\sqrt{4c-(x-1-c)^2}}{-(x+1-c)} \right).
\end{aligned}$$

Since

$$\begin{aligned}
&\int_{a(c)}^{b(c)} \frac{x}{1+x} d \tan^{-1} \left(\frac{\sqrt{4c-(x-1-c)^2}}{-(x+1-c)} \right) \\
&= \frac{1}{2} \int_0^{2\pi} \frac{c-\sqrt{c}\cos(\theta)}{2+c-2\sqrt{c}\cos(\theta)} d\theta \\
&= \int_0^\pi \frac{1}{2} d\theta + \left(\frac{c}{2}-1\right) \int_0^\pi \frac{1}{2+c-2\sqrt{c}\cos(\theta)} d\theta.
\end{aligned}$$

The first integral is $\frac{\pi}{2}$. For the second integral, it equals

$$\begin{aligned}
&\left(\frac{c}{2}-1\right) \frac{1}{1+\varrho} \frac{1}{2} \int_0^\pi \frac{1}{1+\tilde{c}-2\sqrt{\tilde{c}}\cos(\theta)} d\theta \\
&= \frac{c-2}{4i} \frac{1}{1+\varrho} \oint_{|z|=1} \frac{1}{(z-\sqrt{\tilde{c}})(1-\sqrt{\tilde{c}}z)} dz.
\end{aligned}$$

When $\tilde{c} < 1$, the pole is $\sqrt{\tilde{c}}$, and the residue is $\frac{1}{1-\tilde{c}}$. By using Residue theorem, the integral

$$\frac{c-2}{4i} \frac{1}{1+\varrho} \oint_{|z|=1} \frac{1}{(z-\sqrt{\tilde{c}})(1-\sqrt{\tilde{c}}z)} dz \text{ equals } \frac{\pi(c-2)}{2(1+\varrho)(1-\tilde{c})}, \text{ then}$$

$$\int_{a(c)}^{b(c)} \frac{x}{1+x} d \tan^{-1} \left(\frac{\sqrt{4c-(x-1-c)^2}}{-(x+1-c)} \right) \text{ equals } \frac{\pi}{2} + \frac{\pi(c-2)}{2(1+\varrho)(1-\tilde{c})}, \text{ therefore } \oint_{\mathcal{C}} f_V(z) \frac{m'(z)}{m(z)} dz$$

equals $-\frac{1}{2} - \frac{c-2}{2(1+\varrho)(1-\tilde{c})}$, then the proof is finished.

Proof of (6.31): From Remark 3.3,

$$\begin{aligned}
I_1(f_V) &= \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{|1+\sqrt{c}z|^2}{1+|1+\sqrt{c}z|^2} \left(\frac{z}{z^2-r^{-2}} - \frac{1}{z} \right) dz \\
&= \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{|1+\sqrt{c}z|^2}{1+|1+\sqrt{c}z|^2} \frac{z}{z^2-r^{-2}} dz - \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{|1+\sqrt{c}z|^2}{1+|1+\sqrt{c}z|^2} \frac{1}{z} dz.
\end{aligned}$$

For the first integral,

$$(7.50) \quad \begin{aligned} & \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{|1 + \sqrt{cz}|^2}{1 + |1 + \sqrt{cz}|^2} \frac{z}{z^2 - r^{-2}} dz \\ &= \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{z}{z^2 - r^{-2}} dz - \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{1}{1 + |1 + \sqrt{cz}|^2} \frac{z}{z^2 - r^{-2}} dz. \end{aligned}$$

For the first term of (7.50), by using Residue theorem, the integral is 1. For the second integral,

$$(7.51) \quad \begin{aligned} & \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{1}{1 + |1 + \sqrt{cz}|^2} \frac{z}{z^2 - r^{-2}} dz \\ &= \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{1}{(1 + \varrho)(1 + \sqrt{\tilde{c}z})(1 + \sqrt{\tilde{c}}\frac{1}{z})} \cdot \frac{z}{(z - \frac{1}{r})(z + \frac{1}{r})} dz \\ &= \lim_{r \downarrow 1} \frac{1}{2\pi i} \frac{1}{1 + \varrho} \oint_{|z|=1} \frac{z^2}{(1 + \sqrt{\tilde{c}z})(z + \sqrt{\tilde{c}})(z + \frac{1}{r})(z - \frac{1}{r})} dz. \end{aligned}$$

For $\oint_{|z|=1} \frac{z^2}{(1 + \sqrt{\tilde{c}z})(z + \sqrt{\tilde{c}})(z + \frac{1}{r})(z - \frac{1}{r})} dz$, it has $-\sqrt{\tilde{c}}$, $-\frac{1}{r}$, $\frac{1}{r}$ three poles, the residues are $\frac{\tilde{c}}{(1 - \tilde{c})(-\sqrt{\tilde{c}} + \frac{1}{r})(-\sqrt{\tilde{c}} - \frac{1}{r})}$, $\frac{1/r^2}{(1 - \frac{\sqrt{\tilde{c}}}{r})(\sqrt{\tilde{c}} - \frac{1}{r}) \cdot \frac{2}{r}}$, $\frac{1/r^2}{(1 + \frac{\sqrt{\tilde{c}}}{r})(\sqrt{\tilde{c}} + \frac{1}{r}) \cdot \frac{2}{r}}$. Then the summation of residues tend to $\frac{\tilde{c}}{-(\tilde{c}-1)^2} + \frac{1}{2(\sqrt{\tilde{c}}-1)^2} + \frac{1}{2(\sqrt{\tilde{c}}+1)^2}$. Therefore the integral (7.51) equals

$$\frac{1}{1 + \varrho} \left(\frac{\tilde{c}}{-(\tilde{c}-1)^2} + \frac{1}{2(\sqrt{\tilde{c}}-1)^2} + \frac{1}{2(\sqrt{\tilde{c}}+1)^2} \right). \text{ Then the equation (7.50) equals } 1 - \frac{1}{1 + \varrho} \cdot \left(\frac{\tilde{c}}{-(\tilde{c}-1)^2} + \frac{1}{2(\sqrt{\tilde{c}}-1)^2} + \frac{1}{2(\sqrt{\tilde{c}}+1)^2} \right).$$

Then we consider the second integral of $I_1(f_V)$.

$$\begin{aligned} & \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{|1 + \sqrt{cz}|^2}{1 + |1 + \sqrt{cz}|^2} \frac{1}{z} dz \\ &= \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{1}{z} dz - \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{1}{1 + |1 + \sqrt{cz}|^2} \frac{1}{z} dz. \end{aligned}$$

The first integral is 1. By Taylor expansion, the second integral equals

$$\begin{aligned} & \lim_{r \downarrow 1} \frac{1}{2\pi i} \frac{1}{2 + c} \sum_{k=0}^{\infty} (-1)^{2k} \left(\frac{\sqrt{c}}{2 + c} \right)^{2k} \oint_{|z|=1} \left(\frac{1}{z} + z \right)^{2k} \frac{1}{z} dz. \\ &= \frac{1}{2 + c} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c}}{2 + c} \right)^{2k} \frac{(2k)!}{k!k!}, \end{aligned}$$

therefore the second integral of $I_1(f_V)$ equals $1 - \frac{1}{2+c} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c}}{2+c} \right)^{2k} \frac{(2k)!}{k!k!}$. Collecting all the integrals of $I_1(f_V)$, it equals $\frac{1}{2+c} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c}}{2+c} \right)^{2k} \frac{(2k)!}{k!k!} - \frac{1}{1+\varrho} \cdot \left(\frac{\tilde{c}}{-(\tilde{c}-1)^2} + \frac{1}{2(\sqrt{\tilde{c}}-1)^2} + \frac{1}{2(\sqrt{\tilde{c}}+1)^2} \right)$.

Notice that all the c in the formulas above should be replaced by c_{nM} since the calculation is on the bulk part of LSS. The proof is finished.

Proof of (6.32): From Remark 3.3,

$$I_2(f_V) = \frac{1}{2\pi i} \oint_{|z|=1} \frac{|1 + \sqrt{cz}|^2}{1 + |1 + \sqrt{cz}|^2} \frac{1}{z^3} dz$$

$$\begin{aligned}
&= \frac{1}{2\pi i} \oint_{|z|=1} \frac{1}{z^3} dz - \frac{1}{2\pi i} \oint_{|z|=1} \frac{1}{1 + |1 + \sqrt{c}z|^2} \frac{1}{z^3} dz \\
&= -\frac{1}{2\pi i} \oint_{|z|=1} \frac{1}{1 + |1 + \sqrt{c}z|^2} \frac{1}{z^3} dz.
\end{aligned}$$

By Taylor expansion, $\frac{1}{1+|1+\sqrt{c}z|^2} = \frac{1}{2+c} \sum_{k=0}^{\infty} (-1)^k \left(\frac{\sqrt{c}}{2+c}\right)^k \left(\frac{1}{z} + z\right)^k$, then $I_2(f_V)$ equals

$$\begin{aligned}
&-\frac{1}{2\pi i} \oint_{|z|=1} \frac{1}{2+c} \sum_{k=0}^{\infty} (-1)^k \left(\frac{\sqrt{c}}{2+c}\right)^k \left(\frac{1}{z} + z\right)^k \frac{1}{z^3} dz \\
&= -\frac{1}{2\pi i} \frac{1}{2+c} \sum_{k=0}^{\infty} (-1)^k \left(\frac{\sqrt{c}}{2+c}\right)^k \oint_{|z|=1} \left(\frac{1}{z} + z\right)^k \frac{1}{z^3} dz \\
&= -\frac{1}{2\pi i} \frac{1}{2+c} \sum_{k=0}^{\infty} (-1)^{2k} \left(\frac{\sqrt{c}}{2+c}\right)^{2k} \oint_{|z|=1} \left(\frac{1}{z} + z\right)^{2k} \frac{1}{z^3} dz \\
&= -\frac{1}{2+c} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c}}{2+c}\right)^{2k} \frac{(2k)!}{(k-1)!(k+1)!}.
\end{aligned}$$

Notice that all the c in the formulas above should be replaced by c_{nM} since the calculation is on the bulk part of LSS. The proof is finished.

Proof of (6.33): From Remark 3.3,

$$\begin{aligned}
J_1(f_V, f_V) &= \lim_{r \downarrow 1} -\frac{1}{4\pi^2} \oint_{|z_1|=1} \oint_{|z_2|=1} \frac{\frac{|1+\sqrt{c}z_1|^2}{1+|1+\sqrt{c}z_1|^2} \frac{|1+\sqrt{c}z_2|^2}{1+|1+\sqrt{c}z_2|^2}}{(z_1 - rz_2)^2} dz_1 dz_2 \\
&= \lim_{r \downarrow 1} -\frac{1}{4\pi^2} \oint_{|z_2|=1} \frac{|1+\sqrt{c}z_2|^2}{1+|1+\sqrt{c}z_2|^2} \oint_{|z_1|=1} \frac{\frac{|1+\sqrt{c}z_1|^2}{1+|1+\sqrt{c}z_1|^2}}{(z_1 - rz_2)^2} dz_1 dz_2.
\end{aligned}$$

For the integral $\oint_{|z_1|=1} \frac{\frac{|1+\sqrt{c}z_1|^2}{1+|1+\sqrt{c}z_1|^2}}{(z_1 - rz_2)^2} dz_1$, it equals

$$\begin{aligned}
&\oint_{|z_1|=1} \frac{1}{(z_1 - rz_2)^2} dz_1 - \oint_{|z_1|=1} \frac{1}{1 + |1 + \sqrt{c}z_2|^2} \frac{1}{(z_1 - rz_2)^2} dz_1 \\
&= -\oint_{|z_1|=1} \frac{1}{1 + |1 + \sqrt{c}z_2|^2} \frac{1}{(z_1 - rz_2)^2} dz_1 \\
&= -\oint_{|z_1|=1} \frac{1}{2+c} \sum_{k=0}^{\infty} (-1)^k \left(\frac{\sqrt{c}}{2+c}\right)^k \left(\frac{1}{z_1} + z_1\right)^k \frac{1}{(z_1 - rz_2)^2} dz_1 \\
&= -\frac{1}{2+c} \sum_{k=0}^{\infty} (-1)^k \left(\frac{\sqrt{c}}{2+c}\right)^k \oint_{|z_1|=1} \left(\frac{1}{z_1} + z_1\right)^k \frac{1}{(z_1 - rz_2)^2} dz_1 \\
&= \frac{1}{2+c} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c}}{2+c}\right)^{2k+1} \frac{(2k+1)!}{(k+1)!k!} \frac{1}{r^2 z_2^2} 2\pi i.
\end{aligned}$$

By using the same methods as above, then $J_1(f_V, f_V)$ equals

$$\begin{aligned} & \lim_{r \downarrow 1} -\frac{1}{4\pi^2} \oint_{|z_2|=1} \frac{|1 + \sqrt{c}z_2|^2}{1 + |1 + \sqrt{c}z_2|^2} \frac{1}{2+c} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c}}{2+c}\right)^{2k+1} \frac{(2k+1)!}{(k+1)!k!} \frac{1}{r^2 z_2^2} 2\pi i dz_2 \\ &= \lim_{r \downarrow 1} -\frac{1}{4\pi^2} 2\pi i \frac{1}{2+c} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c}}{2+c}\right)^{2k+1} \frac{(2k+1)!}{(k+1)!k!} \oint_{|z_2|=1} \frac{|1 + \sqrt{c}z_2|^2}{1 + |1 + \sqrt{c}z_2|^2} \frac{1}{r^2 z_2^2} dz_2 \\ &= \left(\frac{1}{2+c} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c}}{2+c}\right)^{2k+1} \frac{(2k+1)!}{(k+1)!k!} \right)^2. \end{aligned}$$

Then we consider $J_2(f_V, f_V)$. Since

$$J_2(f_V, f_V) = -\frac{1}{4\pi^2} \oint_{|z_1|=1} \frac{|1 + \sqrt{c}z_1|^2}{1 + |1 + \sqrt{c}z_1|^2} \frac{1}{z_1^2} dz_1 \oint_{|z_2|=1} \frac{|1 + \sqrt{c}z_2|^2}{1 + |1 + \sqrt{c}z_2|^2} \frac{1}{z_2^2} dz_2.$$

For the integral $\oint_{|z_1|=1} \frac{|1 + \sqrt{c}z_1|^2}{1 + |1 + \sqrt{c}z_1|^2} \frac{1}{z_1^2} dz_1$, by Taylor expansion, it equals

$$\begin{aligned} & -\frac{1}{2+c} \sum_{k=0}^{\infty} (-1)^k \left(\frac{\sqrt{c}}{2+c}\right)^k \oint_{|z_1|=1} \left(\frac{1}{z_1} + z_1\right)^k \frac{1}{z_1^2} dz_1 \\ &= -\frac{1}{2+c} \sum_{k=0}^{\infty} (-1)^{2k+1} \left(\frac{\sqrt{c}}{2+c}\right)^{2k+1} \oint_{|z_1|=1} \left(\frac{1}{z_1} + z_1\right)^{2k+1} \frac{1}{z_1^2} dz_1 \\ &= \frac{1}{2+c} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c}}{2+c}\right)^{2k+1} \frac{(2k+1)!}{k!(k+1)!} 2\pi i. \end{aligned}$$

Therefore $J_2(f_V, f_V) = \left(\frac{1}{2+c} \sum_{k=0}^{\infty} \left(\frac{\sqrt{c}}{2+c}\right)^{2k+1} \frac{(2k+1)!}{k!(k+1)!} \right)^2$. Notice that all the c in the formulas above should be replaced by c_{nM} since the calculation is on the bulk part of LSS. The proof is finished.

Proof of (6.34): Since $f(x) = \log(1 + lx)$, then

$$\int f(x) dF^{c_n, H_n}(x) = \int_{a(c_n)}^{b(c_n)} \log(1 + lx) \frac{1}{2\pi x c_n} \sqrt{(b(c_n) - x)(x - a(c_n))} dx,$$

where $a(c_n) = (1 - \sqrt{c_n})^2$, $b(c_n) = (1 + \sqrt{c_n})^2$. By using the variable change $x = 1 + c_n - 2\sqrt{c_n} \cos(\theta)$, $0 \leq \theta \leq \pi$, we have

$$\begin{aligned} \int f(x) dF^{c_n, H_n}(x) &= \frac{1}{2\pi c_n} \int_0^\pi \frac{\log(1 + l(1 + c_n - 2\sqrt{c_n} \cos(\theta)))}{1 + c_n - 2\sqrt{c_n} \cos(\theta)} 4c_n \sin^2(\theta) d\theta \\ &= \frac{1}{2\pi} \int_0^{2\pi} \frac{2 \sin^2(\theta)}{1 + c_n - 2\sqrt{c_n} \cos(\theta)} \log(1 + l + lc_n - 2l\sqrt{c_n} \cos(\theta)) d\theta. \end{aligned}$$

Do the transformation $1 + l + lc_n - 2l\sqrt{c_n} \cos(\theta) = (1 + \varepsilon(c_n)) \left(\frac{1 + l + lc_n}{1 + \varepsilon(c_n)} - \frac{2l\sqrt{c_n} \cos(\theta)}{1 + \varepsilon(c_n)} \right)$,

let $\frac{1 + l + lc_n}{1 + \varepsilon(c_n)} = 1 + \check{c}_n$, $\frac{l\sqrt{c_n}}{1 + \varepsilon(c_n)} = \sqrt{\check{c}_n}$, then we obtain

$$\varepsilon(c_n) = \frac{l(1+c_n)-1+\sqrt{(1-l(1+c_n)^2)-4(l^2c_n-l(1+c_n))}}{2}, \check{c}_n = \frac{l^2c_n}{(1+\varepsilon(c_n))^2}.$$

Therefore $\int f(x)dF^{c_n, H_n}(x)$ equals $\frac{1}{2\pi} \int_0^{2\pi} \frac{2\sin^2(\theta) \log(1+\varepsilon(c_n))}{1+c_n-2\sqrt{c_n}\cos(\theta)} d\theta +$

$\frac{1}{2\pi} \int_0^{2\pi} \frac{2\sin^2(\theta)}{1+c_n-2\sqrt{c_n}\cos(\theta)} \log(1+\check{c}_n-2\sqrt{\check{c}_n}\cos(\theta)) d\theta$. For the first integral and second integral, they can be handled the same way as the proof of (6.22), therefore we omit the detailed calculations and they equals $\log(1+\varepsilon(c_n))$ and $-\frac{(\sqrt{c_n}-\frac{1}{\sqrt{c_n}})^2(\log(1-\sqrt{c_n}\sqrt{c_n})+\sqrt{c_n}\sqrt{c_n})-\sqrt{\check{c}_n}(\sqrt{c_n}-(\sqrt{c_n})^3)}{1-c_n}$, respectively. Collecting the two integrals leads to the desired formula for $p \int f(x)dF^{c_n, H_n}(x)$.

Proof of (6.35): From Remark 3.3,

$$\begin{aligned} I_1(f_U) &= \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \log(1+l|1+\sqrt{c_n}z|^2) \left(\frac{z}{z^2-r^{-2}} - \frac{1}{z} \right) dz \\ &= \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \log(1+l|1+\sqrt{c_n}z|^2) \frac{z}{z^2-r^{-2}} dz - \\ &\quad \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \log(1+l|1+\sqrt{c_n}z|^2) \frac{1}{z} dz. \end{aligned}$$

From the calculations in the proof of (6.24), the first integral of $I_1(f_U)$ is $\log(1+\varepsilon(c_n))$. For the second integral of $I_1(f_U)$,

$$\lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \log(1+l|1+\sqrt{c_n}z|^2) \frac{1}{z} dz \quad (7.52)$$

$$= \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{\log(1+l+lc_n)}{z} dz + \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{\log\left(1+\frac{l\sqrt{c_n}}{1+l+lc_n}\left(\frac{1}{z}+z\right)\right)}{z} dz.$$

By using Taylor expansion, the second integral equals $-\sum_{k=1}^{\infty} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k} \frac{(2k-1)!}{k!k!}$.

Therefore the second integral of $I_1(f_U)$ equals $\log(1+l+lc_n) - \sum_{k=1}^{\infty} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k} \frac{(2k-1)!}{k!k!}$.

Thus $I_1(f_U) = \log(1+\varepsilon(c_n)) - \log(1+l+lc_n) + \sum_{k=1}^{\infty} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k} \frac{(2k-1)!}{k!k!}$. The proof is finished.

Proof of (6.36): Similar to proof of (6.25), by Remark 3.3 and Taylor expansion,

$$\begin{aligned} I_2(f_U) &= \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \log(1+l|1+\sqrt{c_n}z|^2) \frac{1}{z^3} dz \\ &= \frac{1}{2\pi i} \oint_{|z|=1} \left(\log(1+l+lc_n) + \log\left(1+\frac{l\sqrt{c_n}}{1+l+lc_n}\left(\frac{1}{z}+z\right)\right) \right) \frac{1}{z^3} dz \\ &= \frac{1}{2\pi i} \oint_{|z|=1} \log\left(1+\frac{l\sqrt{c_n}}{1+l+lc_n}\left(\frac{1}{z}+z\right)\right) \frac{1}{z^3} dz \\ &= \frac{1}{2\pi i} \oint_{|z|=1} \sum_{k=1}^{\infty} (-1)^{2k+1} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k} \frac{1}{2k} \left(\frac{1}{z}+z\right)^{2k} \frac{1}{z^3} dz \end{aligned}$$

$$= - \sum_{k=1}^{\infty} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k} \frac{(2k-1)!}{(k-1)!(k+1)!}.$$

Therefore the proof is finished.

Proof of (6.37):

$$\begin{aligned} J_1(f_U, f_U) &= \lim_{r \downarrow 1} -\frac{1}{4\pi^2} \oint_{|z_1|=1} \oint_{|z_2|=1} \frac{\log(1+l|1+\sqrt{c_n}z_1|^2) \log(1+l|1+\sqrt{c_n}z_2|^2)}{(z_1-rz_2)^2} dz_1 dz_2 \\ &= \lim_{r \downarrow 1} -\frac{1}{4\pi^2} \oint_{|z_2|=1} \log(1+l|1+\sqrt{c_n}z_2|^2) \oint_{|z_1|=1} \frac{\log(1+l|1+\sqrt{c_n}z_1|^2)}{(z_1-rz_2)^2} dz_1 dz_2. \end{aligned}$$

Similar to proof of (6.26), by using Taylor expansion,

$$\begin{aligned} & \oint_{|z_1|=1} \frac{\log(1+l|1+\sqrt{c_n}z_1|^2)}{(z_1-rz_2)^2} dz_1 \\ &= \oint_{|z_1|=1} \frac{\sum_{k=1}^{\infty} (-1)^{k+1} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^k \frac{1}{k} \left(\frac{1}{z_1} + z_1 \right)^k}{(z_1-rz_2)^2} dz_1 \\ &= 2\pi i \sum_{k=1}^{\infty} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k-1} \frac{(2k-2)!}{k!(k-1)!} \frac{1}{r^2 z_2^2}. \end{aligned}$$

Then $J_1(f_U, f_U)$ equals

$$\begin{aligned} & \lim_{r \downarrow 1} -\frac{1}{4\pi^2} \sum_{k=1}^{\infty} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k-1} 2\pi i \frac{(2k-2)!}{k!(k-1)!} \oint_{|z_2|=1} \log(1+l|1+\sqrt{c_n}z_2|^2) \frac{1}{r^2 z_2^2} dz_2 \\ &= \left(\sum_{k=1}^{\infty} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k-1} \frac{(2k-2)!}{k!(k-1)!} \right)^2. \end{aligned}$$

From Remark 3.3,

$$J_2(f_U, f_U) = \lim_{r \downarrow 1} -\frac{1}{4\pi^2} \oint_{|z_1|=1} \frac{\log(1+l|1+\sqrt{c_n}z_1|^2)}{z_1^2} dz_1 \oint_{|z_2|=1} \frac{\log(1+l|1+\sqrt{c_n}z_2|^2)}{z_2^2} dz_2.$$

Similar to the proof of (6.26), by Taylor expansion, we obtain

$$\begin{aligned} & \oint_{|z_1|=1} \frac{\log(1+l|1+\sqrt{c_n}z_1|^2)}{z_1^2} dz_1 \\ &= 2\pi i \sum_{k=1}^{\infty} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k-1} \frac{(2k-2)!}{k!(k-1)!}, \end{aligned}$$

thus

$$J_2(f_U, f_U) = \left(\sum_{k=1}^{\infty} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k-1} \frac{(2k-2)!}{k!(k-1)!} \right)^2.$$

Therefore the proof is finished.

Proof of (6.38): Since $f(x) = \frac{x}{\frac{1}{l} + x}$, then

$$\int f(x) dF^{c_n, H_n}(x) = \int_{a(c_n)}^{b(c_n)} \frac{x}{\frac{1}{l} + x} \frac{1}{2\pi x c_n} \sqrt{(b(c_n) - x)(x - a(c_n))} dx,$$

where $a(c_n) = (1 - \sqrt{c_n})^2$, $b(c_n) = (1 + \sqrt{c_n})^2$. By using the variable change $x = 1 + c_n - 2\sqrt{c_n} \cos(\theta)$, $0 \leq \theta \leq \pi$, we have

$$\begin{aligned} \int f(x) dF^{c_n, H_n}(x) &= \frac{1}{2\pi c_n} \int_0^\pi \frac{1}{\frac{1}{l} + 1 + c_n - 2\sqrt{c_n} \cos(\theta)} 4c_n \sin^2(\theta) d\theta \\ &= \frac{1}{4\pi c_n} \int_0^{2\pi} \frac{l 4c_n \sin^2(\theta)}{1 + l(1 + c_n - 2\sqrt{c_n} \cos(\theta))} d\theta. \\ &= \frac{l}{4\pi c_n} \int_0^{2\pi} \frac{4c_n \sin^2(\theta)}{(1 + \varepsilon(c_n))(1 + \check{c}_n - 2\sqrt{\check{c}_n} \cos(\theta))} d\theta. \\ &= -\frac{l}{4\pi i} \frac{1}{1 + \varepsilon(c_n)} \oint_{|z|=1} \frac{(z+1)^2 (z-1)^2}{(z - \sqrt{\check{c}_n})(1 - \sqrt{\check{c}_n} z) z^2} dz. \end{aligned}$$

From the proof of (6.34), we have $\oint_{|z|=1} \frac{(z+1)^2 (z-1)^2}{(z - \sqrt{\check{c}_n})(1 - \sqrt{\check{c}_n} z) z^2} dz = -4\pi i$. Therefore

$\int f(x) dF^{c_n, H_n}(x) = \frac{l}{1 + \varepsilon(c_n)}$. The proof is finished.

Proof of (6.39): From Remark 3.3,

$$\begin{aligned} I_1(f_V) &= \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{|1 + \sqrt{c_n} z|^2}{\frac{1}{l} + |1 + \sqrt{c_n} z|^2} \left(\frac{z}{z^2 - r^{-2}} - \frac{1}{z} \right) dz \\ &= \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{|1 + \sqrt{c_n} z|^2}{\frac{1}{l} + |1 + \sqrt{c_n} z|^2} \frac{z}{z^2 - r^{-2}} dz - \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{|1 + \sqrt{c_n} z|^2}{\frac{1}{l} + |1 + \sqrt{c_n} z|^2} \frac{1}{z} dz. \end{aligned}$$

For the first integral of $I_1(f_V)$, it equals

$$\begin{aligned} &\lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{z}{z^2 - r^{-2}} dz - \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{\frac{1}{l}}{\frac{1}{l} + |1 + \sqrt{c_n} z|^2} \frac{z}{z^2 - r^{-2}} dz \\ &= 1 - \lim_{r \downarrow 1} \frac{1}{2\pi i} \frac{1}{1 + \varepsilon(c_n)} \oint_{|z|=1} \frac{z^2}{(1 + \sqrt{\check{c}_n} z)(z + \sqrt{\check{c}_n})(z + \frac{1}{r})(z - \frac{1}{r})} dz. \\ &= 1 - \frac{1}{1 + \varepsilon(c_n)} \left(\frac{\check{c}_n}{-(\check{c}_n - 1)^2} + \frac{1}{2(\sqrt{\check{c}_n} - 1)^2} + \frac{1}{2(\sqrt{\check{c}_n} + 1)^2} \right). \end{aligned}$$

For the second integral of $I_1(f_V)$, it equals

$$\lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{1}{z} dz - \lim_{r \downarrow 1} \frac{1}{2\pi i} \oint_{|z|=1} \frac{\frac{1}{l}}{\frac{1}{l} + |1 + \sqrt{c_n} z|^2} \frac{1}{z} dz.$$

The first integral is 1. By Taylor expansion, the second integral equals

$$\begin{aligned} &\lim_{r \downarrow 1} \frac{1}{2\pi i} \frac{1}{1 + l + lc_n} \sum_{k=0}^{\infty} (-1)^{2k} \left(\frac{l\sqrt{c_n}}{1 + l + lc_n} \right)^{2k} \oint_{|z|=1} \left(\frac{1}{z} + z \right)^{2k} \frac{1}{z} dz \\ &= \frac{1}{1 + l + lc_n} \sum_{k=0}^{\infty} \left(\frac{l\sqrt{c_n}}{1 + l + lc_n} \right)^{2k} \frac{(2k)!}{k!k!}, \end{aligned}$$

Therefore the second integral of $I_1(f_V)$ equals $1 - \frac{1}{1+l+lc_n} \sum_{k=0}^{\infty} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k} \frac{(2k)!}{k!k!}$. Then

$$I_1(f_V) = \frac{1}{1+l+lc_n} \sum_{k=0}^{\infty} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k} \frac{(2k)!}{k!k!} - \frac{1}{1+\varepsilon(c_n)} \left(\frac{\check{c}_n}{-(\check{c}_n-1)^2} + \frac{1}{2(\sqrt{\check{c}_n}-1)^2} + \frac{1}{2(\sqrt{\check{c}_n}+1)^2} \right).$$

The proof is finished.

Proof of (6.40): Similar to proof of (6.32), by Remark 3.3 and Taylor expansion, we have

$$\begin{aligned} I_2(f_V) &= \frac{1}{2\pi i} \oint_{|z|=1} \frac{|1+\sqrt{c_n}z|^2}{\frac{1}{l} + |1+\sqrt{c_n}z|^2} \frac{1}{z^3} dz \\ &= \frac{1}{2\pi i} \oint_{|z|=1} \frac{1}{z^3} dz - \frac{1}{2\pi i} \oint_{|z|=1} \frac{\frac{1}{l}}{\frac{1}{l} + |1+\sqrt{c_n}z|^2} \frac{1}{z^3} dz \\ &= -\frac{1}{2\pi i} \oint_{|z|=1} \frac{\frac{1}{l}}{\frac{1}{l} + |1+\sqrt{c_n}z|^2} \frac{1}{z^3} dz. \\ &= -\frac{1}{2\pi i} \frac{1}{1+l+lc_n} \sum_{k=0}^{\infty} (-1)^k \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^k \oint_{|z|=1} \left(\frac{1}{z} + z \right)^k \frac{1}{z^3} dz \\ &= -\frac{1}{1+l+lc_n} \sum_{k=0}^{\infty} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k} \frac{(2k)!}{(k-1)!(k+1)!}. \end{aligned}$$

Therefore the proof is finished.

Proof of (6.41): From Remark 3.3,

$$\begin{aligned} J_1(f_V, f_V) &= \lim_{r \downarrow 1} -\frac{1}{4\pi^2} \oint_{|z_1|=1} \oint_{|z_2|=1} \frac{\frac{|1+\sqrt{c_n}z_1|^2}{\frac{1}{l}+|1+\sqrt{c_n}z_1|^2} \frac{|1+\sqrt{c_n}z_2|^2}{\frac{1}{l}+|1+\sqrt{c_n}z_2|^2}}{(z_1 - rz_2)^2} dz_1 dz_2 \\ &= \lim_{r \downarrow 1} -\frac{1}{4\pi^2} \oint_{|z_2|=1} \frac{|1+\sqrt{c_n}z_2|^2}{\frac{1}{l} + |1+\sqrt{c_n}z_2|^2} \oint_{|z_1|=1} \frac{\frac{|1+\sqrt{c_n}z_1|^2}{\frac{1}{l}+|1+\sqrt{c_n}z_1|^2}}{(z_1 - rz_2)^2} dz_1 dz_2. \end{aligned}$$

For the integral $\oint_{|z_1|=1} \frac{\frac{|1+\sqrt{c_n}z_1|^2}{\frac{1}{l}+|1+\sqrt{c_n}z_1|^2}}{(z_1 - rz_2)^2} dz_1$, by using Taylor expansion, it equals

$$\begin{aligned} &\oint_{|z_1|=1} \frac{1}{(z_1 - rz_2)^2} dz_1 - \oint_{|z_1|=1} \frac{\frac{1}{l}}{\frac{1}{l} + |1+\sqrt{c_n}z_1|^2} \frac{1}{(z_1 - rz_2)^2} dz_1 \\ &= -\oint_{|z_1|=1} \frac{1}{1+l|1+\sqrt{c_n}z_1|^2} \frac{1}{(z_1 - rz_2)^2} dz_1 \\ &= -\frac{1}{1+l+lc_n} \sum_{k=0}^{\infty} (-1)^k \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^k \oint_{|z_1|=1} \left(\frac{1}{z_1} + z_1 \right)^k \frac{1}{(z_1 - rz_2)^2} dz_1 \\ &= \frac{1}{1+l+lc_n} \sum_{k=0}^{\infty} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k+1} \frac{(2k+1)!}{(k+1)!k!} \frac{1}{r^2 z_2^2} 2\pi i. \end{aligned}$$

By using the same methods as above, then $J_1(f_V, f_V)$ equals

$$\lim_{r \downarrow 1} -\frac{1}{4\pi^2} \oint_{|z_2|=1} \frac{|1+\sqrt{c_n}z_2|^2}{\frac{1}{l} + |1+\sqrt{c_n}z_2|^2} \frac{1}{1+l+lc_n} \sum_{k=0}^{\infty} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k+1} \frac{(2k+1)!}{(k+1)!k!} \frac{1}{r^2 z_2^2} 2\pi i dz_2$$

$$\begin{aligned}
&= \lim_{r \downarrow 1} -\frac{1}{4\pi^2} 2\pi i \frac{1}{1+l+lc_n} \sum_{k=0}^{\infty} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k+1} \frac{(2k+1)!}{(k+1)!k!} \oint_{|z_2|=1} \frac{|1+\sqrt{c_n}z_2|^2}{\frac{1}{l} + |1+\sqrt{c_n}z_2|^2} \frac{1}{r^2 z_2^2} dz_2 \\
&= \left(\frac{1}{1+l+lc_n} \sum_{k=0}^{\infty} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k+1} \frac{(2k+1)!}{(k+1)!k!} \right)^2.
\end{aligned}$$

Then we consider $J_2(f_V, f_V)$. Since

$$J_2(f_V, f_V) = -\frac{1}{4\pi^2} \oint_{|z_1|=1} \frac{\frac{|1+\sqrt{c_n}z_1|^2}{\frac{1}{l} + |1+\sqrt{c_n}z_1|^2}}{z_1^2} dz_1 \oint_{|z_2|=1} \frac{\frac{|1+\sqrt{c_n}z_2|^2}{\frac{1}{l} + |1+\sqrt{c_n}z_2|^2}}{z_2^2} dz_2.$$

For the integral $\oint_{|z_1|=1} \frac{\frac{|1+\sqrt{c_n}z_1|^2}{\frac{1}{l} + |1+\sqrt{c_n}z_1|^2}}{z_1^2} dz_1$, by Taylor expansion, it equals

$$\begin{aligned}
& -\frac{1}{1+l+lc_n} \sum_{k=0}^{\infty} (-1)^k \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^k \oint_{|z_1|=1} \left(\frac{1}{z_1} + z_1 \right)^k \frac{1}{z_1^2} dz_1 \\
&= \frac{1}{1+l+lc_n} \sum_{k=0}^{\infty} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k+1} \frac{(2k+1)!}{k!(k+1)!} 2\pi i.
\end{aligned}$$

Therefore $J_2(f_V, f_V) = \left(\frac{1}{1+l+lc_n} \sum_{k=0}^{\infty} \left(\frac{l\sqrt{c_n}}{1+l+lc_n} \right)^{2k+1} \frac{(2k+1)!}{k!(k+1)!} \right)^2$. The proof is finished.

REFERENCES

- Zhidong Bai and Jack W. Silverstein. CLT for linear spectral statistics of large-dimensional sample covariance matrices. *The Annals of Probability*, 32(1A):553 – 605, 2004.
- Zhijun Liu, Zhidong Bai, Jiang Hu and Haiyan Song. A CLT for the LSS of large dimensional sample covariance matrices with diverging spikes. *Submitted to the Annals of Statistics*, 2022.
- Qinwen Wang and Jianfeng Yao. On the sphericity test with large-dimensional observations. *Electronic Journal of Statistics*, 7:2164–2192, 2013.
- Shurong Zheng, Zhidong Bai, and Jianfeng Yao. Substitution principle for CLT of linear spectral statistics of high-dimensional sample covariance matrices with applications to hypothesis testing. *The Annals of Statistics*, 43(2):546 – 591, 2015.