

Shallow quantum circuits for efficient preparation of Slater determinants and correlated states on a quantum computer

Chong Hian Chee,^{1,*} Daniel Leykam,¹ Adrian M. Mak,² and Dimitris G. Angelakis^{1,3,4,†}

¹Centre for Quantum Technologies, National University of Singapore, 3 Science Drive 2, Singapore 117543

²Institute of High Performance Computing, Agency for Science, Technology

& Research, 1 Fusionopolis Way, #16-16 Connexis, Singapore 138632

³School of Electrical and Computer Engineering, Technical University of Crete, Chania, Greece 73100

⁴AngelQ Quantum Computing, 531A Upper Cross Street, #04-95 Hong Lim Complex, Singapore 051531

(Dated: January 26, 2023)

Preparing quantum ansatzes is a necessary prerequisite in many quantum algorithms for quantum chemistry such as the variational quantum eigensolver. Widely-used ansatzes including the Slater determinants and Unitary Coupled Cluster, employ parameterized fermionic excitation gates, with the latter resulting in deep quantum circuits that scale at least polynomially with the system size N . Here we propose an alternate paradigm for fermionic ansatz state preparation inspired by data-loading circuits methods developed for quantum machine learning. Our approach provides a shallower, yet scalable $O(d \log^2 N)$ two-qubit gate depth preparation of d -fermion Slater determinants and correlated states, a subexponential improvement in gate depth over existing approaches. This is particularly important as it can be implemented on planar architectures without qubit swapping overheads, thereby enabling the use of larger basis sets needed for high-precision quantum chemistry studies on near-term quantum devices.

Introduction.—Quantum computers promise the ability to solve hard many-body problems in quantum chemistry and condensed matter physics, including computation of ground state energies and simulation of quantum dynamics [1–3]. The relevant quantum algorithms frequently involve quantum state preparation as a key step. For example, the success probability of quantum phase estimation is determined by the overlap of a trial ansatz state with the eigenstate of interest [4–6]. Thus, efficient preparation of high quality ansatz states is crucial for many-body applications of quantum computing [7, 8].

Most existing methods for preparing fermionic ansatzes use second quantization with Jordan-Wigner mapping [9] to efficiently represent the quantum many-body fermionic wavefunction using a number of qubits that scales linearly in the system size [1, 2]. Widely-used fermionic ansatzes typically fall into two broad classes: hardware-efficient ansätze [10], and problem-inspired ansätze. Hardware-efficient ansätze use parameterized hardware-native gates to minimize the depth of the quantum circuit, but are difficult to optimize [11] and do not guarantee an accurate representation of the desired quantum state [12]. Meanwhile, the more promising problem-inspired ansätze explicitly incorporate the physics of the system of interest but require deeper circuits which exacerbate errors due to quantum noise and decoherence [13–20], limiting state-of-the-art demonstrations to around 20 qubits [21, 22]. The polynomial scaling of existing problem-inspired ansätze unfortunately leaves studies of chemically-relevant molecular systems requiring more than 10^2 – 10^3 qubits well out of reach [23, 24].

The shallowest general-purpose problem-inspired ansatz states to date are mean-field Hartree-Fock states [13–15], which are Slater determinants that can

be prepared using a mesh of fermionic single-excitation gates which has a linear $O(N)$ two-qubit gate depth in the number of qubits N as shown in Fig. 1(a). While Hartree-Fock states are efficiently simulatable using classical computers, they nevertheless serve as a useful starting point for quantum computers to prepare more interesting, but classically-intractable correlated quantum ansatzes, such as the unitary coupled cluster ansatz which incorporates quantum correlations by applying number-conserving multi-fermion excitation operators to a reference Hartree-Fock state [16–20, 25].

Fermionic excitation operators are examples of Givens rotations that are universal quantum gates which perform rotations in a two-dimensional subspace of a larger Hilbert space [26, 27]. Givens rotations have been helpful for preparing various fermionic states in quantum chemistry and condensed matter applications [13–17, 28]. Recently, such gates have also attracted interest in the context of quantum linear algebra, where they were used to construct shallow-depth “Clifford loaders” gates as a means to efficiently encode d -dimensional subspaces of \mathbb{R}^N into a N -qubit state [29–32], enabling potential end-to-end quantum speedups for several quantum machine learning and linear algebra problems including determinant sampling and topological data analysis [29]. It is thus timely to consider whether the Clifford loader approach are useful for preparing of fermionic ansatz states.

Here we shall show how a d -fermion Slater determinant can be prepared via the Clifford loader approach by using a shallow $O(d \log^2 N)$ two-qubit gate depth quantum circuit on quantum devices with planar architectures without expensive qubit swapping overheads, that yields shallower circuits above $N \sim 100$ qubits than existing approaches. Moreover, we shall also show how this ap-




Figure 1. Different approaches for preparing Slater determinants on a quantum computer. (a) Existing approaches use a linear-depth mesh of Givens rotations to apply a fermionic basis transformation on a reference state [13–15]. (b) The proposed Clifford loader approach applies a sequence \hat{C} gates d times to an all-zero state $|0\rangle^{\otimes N}$, where \hat{C} is efficiently decomposed in (d-f). θ and \vec{A} are scalar and vector parameters respectively.

proach can be extended to prepare fermionic correlated ansatz states to capture a significant fraction of correlation energy via a numerical simulation of hydrogen chains up to $N=16$ qubits. Our results establish the Clifford loader approach as a promising method that allows efficient, yet practical and scalable preparation of fermionic ansatz states for large quantum chemistry applications on near-term quantum computers.

Slater determinant.—We begin by showing how a Slater determinant with d occupied and $N-d$ unoccupied fermionic modes can be efficiently prepared on a shallow quantum circuit. An arbitrary Slater determinant is defined as [13, 33–35]

$$|\Psi_{SD}(A)\rangle := \prod_{l=1}^d \sum_{k=1}^N A_{lk} \hat{a}_k^\dagger |\text{vac}\rangle, \quad (1)$$

where A is an $N \times d$ real matrix such that the first d columns are orthogonal and normalized, $|\text{vac}\rangle$ is a vacuum state, and \hat{a}_k^\dagger is a creation operator acting on the k^{th} mode. While the definition in Eq. (1) is pedagogically convenient, it requires non-unitary operators $\sum_{k=1}^N A_{lk} \hat{a}_k^\dagger$ which cannot be directly implemented on a quantum circuit. Consequently, the most efficient method to prepare Slater determinants to date was to simply perform a fermionic basis transformation to a reference Slater determinant state for a given skew-Hermitian parameter matrix κ as

$$|\Psi_{SD}(A)\rangle = \exp \left[\sum_{ij}^N \kappa_{ij} \hat{b}_i^\dagger \hat{b}_j \right] \prod_{l=1}^d b_l^\dagger |\text{vac}\rangle, \quad (2)$$

where under the Jordan-Wigner fermion-to-qubit mapping, the fermionic basis transformation $\exp[\sum_{ij}^N \kappa_{ij} \hat{b}_i^\dagger \hat{b}_j]$ can be implemented as a linear-depth mesh of fermionic single-excitation gates, as shown in Fig. 1(a) [14, 15].

We propose to improve the two-qubit gate depth efficiency of preparing Slater determinants using an alternative, yet equivalent definition

$$|\Psi_{SD}(A)\rangle := \prod_{l=1}^d \sum_{k=1}^N A_{lk} \left(\hat{a}_k^\dagger + \hat{a}_k \right) |\text{vac}\rangle, \quad (3)$$

where we use anti-commuting operators $\hat{p}_k = \hat{a}_k^\dagger + \hat{a}_k$, with the relation $\frac{1}{2} \{ \hat{p}_i, \hat{p}_j \} = \delta_{ij} \mathbf{I}$. We refer readers to the Supplementary Material [36] for a proof of this equivalence. Using the anti-commuting operators \hat{p}_k instead of \hat{a}_k^\dagger allow us to exploit the recent result in Ref. [29] that provides a shallow $O(\log^2 N)$ two-qubit gate depth circuit decomposition of the Clifford loader

$$\hat{C}(\vec{A}) = \sum_k A_k (\hat{a}_k^\dagger + \hat{a}_k). \quad (4)$$

for some normalized column \vec{A} , under the Jordan-Wigner mapping. When the Clifford loader $\hat{C}(\vec{A}_l)$ are applied in succession d times on an all-zero state $|0\rangle^{\otimes N}$, as shown in Fig. 1(b), each with orthogonal columns \vec{A}_l , $l=1, \dots, d$ from matrix A , it generates the desired Slater determinant

$$\prod_{l=1}^d \hat{C}(\vec{A}_l) |0\rangle^{\otimes N} = |\Psi_{SD}(A)\rangle. \quad (5)$$

Hence, we can prepare any Slater determinants using a shallow $O(d \log^2 N)$ two-qubit gate depth circuit. If $d > N/2$, we may apply the Clifford loader $N-d$ times instead, followed by a bitflip to all qubits at the end.

The Clifford loader $\hat{C}(\vec{A})$ can be decomposed as

$$\hat{C}(\vec{A}) = \hat{D}(\vec{A}) (\hat{a}_1^\dagger + \hat{a}_1) \hat{D}^\dagger(\vec{A}), \quad (6)$$




Figure 2. (a) Pairwise-correlated Clifford loader approach for preparing d -fermion pairwise correlated ansatzes on a quantum computer applying a sequence of \hat{C}_{pair} gates d times to an all-zero state $|0\rangle^{\otimes N}$, where \hat{C}_{pair} is efficiently decomposed in (b-d). θ and \vec{G} are scalar and vector parameters respectively.

as shown in Fig. 1(c), where the operator $\hat{a}_1^\dagger + \hat{a}_1$ acting on the first mode is sandwiched by unary data loaders [29, 32, 37],

$$\hat{D}(\vec{A}) = \prod_{ij \in \mathfrak{I}} \hat{U}_j^i(\theta_{ij}), \quad (7)$$

which are composed of Givens rotations arranged in a binary tree-like structure \mathfrak{I} shown in Fig. 1(d), where Givens rotations used here are fermionic single-excitation gates $\hat{U}_j^i(\theta) = \exp[\theta(\hat{a}_i^\dagger \hat{a}_j - \hat{a}_j^\dagger \hat{a}_i)]$, as shown in Fig. 1(e).

The rotation angles θ_{ij} are obtained by classically preprocessing the orthonormalized columns \vec{A} as given in Ref. [29]. Remarkably, Ref. [29] has also shown that the fermionic single-excitation \hat{U}_j^i can be decomposed in a logarithmic $O(\log N)$ two-qubit gate depth, by using a binary tree-like subcircuit \hat{B} that is composed of CNOT gates as shown in Fig. 1(f), which efficiently encodes the parity of qubits between $j+1$ and $i-2$ inclusive onto qubit $i-1$ within \hat{U}_j^i . This is a huge improvement over the existing linear two-qubit gate depth scaling gate decompositions of \hat{U}_j^i [38, 39].

Correlated Ansatz.—Next, we extended the Clifford loader approach by introducing a method to incorporate pairwise correlations into the fermionic ansatz state preparation. Under the Jordan-Wigner mapping, the anti-commuting operator $\hat{p}_k = \hat{Z}_{\otimes k-1} \hat{X}_k$, where $\hat{Z}_{\otimes k} = \hat{Z}_1 \otimes \dots \otimes \hat{Z}_k$, used to prepare the Slater determinant previously is modified to apply pairwise correlation by applying an extra Pauli \hat{X} and removing all odd labelled Pauli \hat{Z} , giving $\hat{p}'_k = \hat{Z}_{\otimes 2, 2k-2, \text{even}} \hat{X}_{2k-1} \hat{X}_{2k}$, where $\hat{Z}_{\otimes 2, 2k-2, \text{even}} = \hat{Z}_2 \otimes \hat{Z}_4 \otimes \dots \otimes \hat{Z}_{2k-2}$. Since \hat{p}'_k remains anti-commuting [36], we can reapply the Clifford loader approach as shown in Fig. 2(a) to prepare a

d -occupied pairwise-correlated state $|\Psi_{\text{pair}}(G)\rangle$ with N modes, where G is an $N/2 \times d/2$ real matrix such that the first $d/2$ columns are orthogonal and normalized. Here, for a given normalized column \vec{G} , we can define an analogous pairwise-correlated Clifford loader

$$\hat{C}_{\text{pair}}(\vec{G}) = \hat{D}_{\text{pair}}^\dagger(\vec{G}) \hat{X}_1 \hat{X}_2 \hat{D}_{\text{pair}}(\vec{G}), \quad (8)$$

as shown in Fig. 2(b), where two Pauli operators $\hat{X}_1 \hat{X}_2$ acting on the first two qubits are sandwiched by an analogous pair of pairwise-correlated unary data loader

$$\hat{D}_{\text{pair}}(\vec{G}) = \prod_{ij \in \mathfrak{I}} \hat{U}_{j-1,j}^{i-1,i}(\theta_{ij}), \quad (9)$$

which are composed of a particular type of Givens rotations gate, arranged in a binary tree-like structure \mathfrak{I} shown in Fig. 2(c). We termed the type of Givens rotations gates used here as pairwise-correlated fermionic beam splitter gate $\hat{U}_{j-1,j}^{i-1,i}(\theta_{ij})$ as shown in Fig. 2(d) [36], where the rotation angles θ_{ij} are obtained similarly as before. The pairwise-correlated fermionic beam splitter gate $\hat{U}_{j-1,j}^{i-1,i}$ is a combination of a modified fermionic double-excitation operator from Ref. [39] and a binary tree-like subcircuit \hat{B}'_+ , that is similar to \hat{B} , which efficiently encodes the parity of even-labelled qubits between $j+2$ and $i-4$ inclusive onto qubit $i-2$ within $\hat{U}_{j-1,j}^{i-1,i}$. As a result, we can efficiently prepare pairwise-correlated state $|\Psi_{\text{pair}}(G)\rangle$ with the same shallow $O(d \log^2 N)$ depth circuit.

Example.—We numerically evaluated the fraction of the electronic correlation energy $E_{\text{pair}}/E_{\text{corr}}$ captured by the optimized pairwise-correlated state $|\Psi_{\text{pair}}(G^*)\rangle$ for three linear hydrogen molecular chains: (H_2 , H_4 and H_6) with ($d=2, 4, 6$) electrons at a fixed interatomic distance of 1.4 bohr using a quantum statevector calculation. $E_{\text{pair}} = E_{\text{HF}} - \langle \hat{H}_e \rangle$ is the correlation energy beyond the mean-field energy E_{HF} of molecule captured by $|\Psi_{\text{pair}}(G^*)\rangle$, while $E_{\text{corr}} = E_{\text{HF}} - E_{\text{FCI}}$ is the exact value of correlation energy, where E_{FCI} is known as the full configuration interaction (FCI) energy. G^* is an optimized parameter matrix, obtained using a classical quasi-Newton L-BFGS-B optimizer that minimizes the expectation of the electronic Hamiltonian $\langle \hat{H}_e \rangle$. We considered different mixtures of atomic basis sets (STO-3G, 6-31G, 6-311G, cc-pvdz, aug-cc-pvdz) for each hydrogen atom, resulting in system sizes ranging from 4 to 16 qubits. Such mixing of basis sets for each atom is a common strategy in computational quantum chemistry to reduce the computational resources required to achieve a desired precision [40]. The calculations were performed using SciPy [41], PYSCF [42] and PennyLane [43]. The results are plotted in Fig. 3(a) which show that a significant fraction of the electronic correlation energy $E_{\text{pair}}/E_{\text{corr}}$ is captured by the optimized pairwise-correlated state $|\Psi_{\text{pair}}(G^*)\rangle$.

Resource Analysis.—For simplicity, we treat all types of two-qubit gates depths as equal. The fermionic single-excitation \hat{U}_j^i has a two-qubit gate depth of $3 + 2\lceil \log |i - j - 1| \rceil$, $i - j \geq 3$. The overall two-qubit gate depth required to prepare a Slater determinant is thus

$$2d (\lceil \log^2 N \rceil + 2\lceil \log N \rceil). \quad (10)$$

Similarly, the pairwise-correlated fermionic beam splitter $\hat{U}_{j-1,j}^{i-1,i}$ in Fig. 2(d) has a two-qubit gate depth of $12 + 2\lceil \log \lceil \frac{i-j-3}{2} \rceil \rceil$, $i - j \geq 7$, where we assume that the multi-controlled-rotation-y gate has a two-qubit gate depth of 8 [39]. The overall two-qubit gate depth required to prepare a pairwise-correlated state is therefore

$$d (\lceil \log^2 N \rceil + 9\lceil \log N \rceil - 10). \quad (11)$$

We plotted the estimated two-qubit gate depth required to prepare $d=2, 8, 32$ -fermion Slater determinants and pairwise-correlated state on a quantum computer in Fig. 3(b) and 3(c) respectively, and compared against the previous fermionic basis transformation approach [13–15] and method of preparing trotterized unitary coupled cluster doubles and singles (UCCSD) ansätze [16] respectively. The trotterized-UCCSD ansätze is obtained by applying trotterized evolution operators generated by fermionic single- and double-excitation operators to a d -fermion reference state. The results show that for a $d=2$ -fermion Slater determinant, the Clifford loader approach is estimated to yield shallower circuits than the previous approach at around 100 qubits with 250 two-qubit gate depth.

In addition, we observed that the binary tree structure of two-qubit gates in the Clifford loader circuit do not form K_5 complete two-qubit interactions on any five qubits or $K_{3,3}$ complete bipartite two-qubit interactions on any six qubits. We therefore conjecture that the qubit connectivity topology required by the Clifford loader approach is planar by Kuratowski’s Theorem, allowing a realizable quantum demonstration on a 2D-planar architecture without expensive qubit swapping overheads that may negate the shallow depth scaling. As an example, we show a planar 8-qubit connectivity topology required to prepare the Slater determinants and the correlated pairwise ansätze in Figs. 4(a) and (b) respectively, which are compatible with existing quantum processor architectures [44–47]

Summary and outlook.—We have proposed a Clifford loader approach to efficiently prepare d -fermion Slater determinants and correlated state using a shallow $O(d \log^2 N)$ two-qubit gate depth quantum circuit. We have also showed how the Clifford loader approach can be extended to prepare pairwise-correlated state to capture correlation energy and demonstrate its validity via a numerical simulation of hydrogen chains. As expected by the ansatz design, it does not fully capture all the electronic correlation energy. Thus, it will be interesting




Figure 3. (a) Numerical calculation of electronic correlation energy ratio $E_{\text{pair}}/E_{\text{corr}}$ captured by an optimized pairwise-correlated state for various hydrogen chains up to H_6 at fixed interatomic distance of 1.4 bohr for various number of qubits corresponding to various basis set mixtures. (b-c) Estimated two-qubit gate depth required by the Clifford loader approach to prepare $d=2, 8, 32$ -fermion Slater determinants and pairwise-correlated states respectively, against the fermionic basis transformation approach [15] and the trotterized-UCCSD [16] respectively for various number of qubits.




Figure 4. 8-qubit example of connectivity topology of the Clifford loader approach for preparing (a) a Slater determinant and (b) pairwise-correlated state.

for future work to generalize the Clifford loader approach to incorporate other types of fermionic correlation while keeping the same shallow gate depth scaling intact.

The Clifford loader approach, to the best of our knowledge, offers an subexponential improvement in gate depth for fermionic ansatz state preparation over existing approaches in second quantization. While there exist other approaches in first quantization with a similar $O(d \log N)$ Given rotation gate depth scaling, it requires the use of fault-tolerant quantum computers [48–50] which is unlikely to be available in the near-term. Nonetheless, our results have established the Clifford loader approach as an efficient, yet practical and scalable fermionic ansatz state preparation, which will enable the study of molecules and materials requiring larger basis set sizes on near-term quantum devices.

This research is supported by the National Research Foundation, Singapore and A*STAR under its CQT

Bridging Grant and Quantum Engineering Programme NRF2021-QEP2-02-P02, A*STAR (#21709) and by EU HORIZON - Project 101080085 — QCFD.

* ch.chee@u.nus.edu

† dimitris.angelakis@gmail.com

- [1] B. Bauer, S. Bravyi, M. Motta, and G. K.-L. Chan, Quantum Algorithms for Quantum Chemistry and Quantum Materials Science, *Chem. Rev.* **120**, 12685 (2020).
- [2] M. Motta and J. E. Rice, Emerging quantum computing algorithms for quantum chemistry, *WIREs Computational Molecular Science* **12**, e1580 (2022).
- [3] A. J. Daley, I. Bloch, C. Kokail, S. Flannigan, N. Pearson, M. Troyer, and P. Zoller, Practical quantum advantage in quantum simulation, *Nature* **607**, 667 (2022).
- [4] D. S. Abrams and S. Lloyd, Simulation of Many-Body Fermi Systems on a Universal Quantum Computer, *Phys. Rev. Lett.* **79**, 2586 (1997).
- [5] D. S. Abrams and S. Lloyd, Quantum Algorithm Providing Exponential Speed Increase for Finding Eigenvalues and Eigenvectors, *Phys. Rev. Lett.* **83**, 5162 (1999).
- [6] A. Aspuru-Guzik, A. D. Dutoi, P. J. Love, and M. Head-Gordon, Simulated Quantum Computation of Molecular Energies, *Science* **309**, 1704 (2005).
- [7] S. Lee, J. Lee, H. Zhai, Y. Tong, A. M. Dalzell, A. Kumar, P. Helms, J. Gray, Z.-H. Cui, W. Liu, M. Kastoryano, R. Babbush, J. Preskill, D. R. Reichman, E. T. Campbell, E. F. Valeev, L. Lin, and G. K.-L. Chan, *Is there evidence for exponential quantum advantage in quantum chemistry?* (2022), [arXiv:2208.02199](https://arxiv.org/abs/2208.02199).
- [8] K. Bharti, A. Cervera-Lierta, T. H. Kyaw, T. Haug, S. Alperin-Lea, A. Anand, M. Degroote, H. Heimonen, J. S. Kottmann, T. Menke, W.-K. Mok, S. Sim, L.-C. Kwek, and A. Aspuru-Guzik, Noisy intermediate-scale quantum algorithms, *Rev. Mod. Phys.* **94**, 015004 (2022).
- [9] P. Jordan and E. Wigner, Über das Paulische Äquivalenzverbot, *Z. Physik* **47**, 631 (1928).
- [10] A. Kandala, A. Mezzacapo, K. Temme, M. Takita, M. Brink, J. M. Chow, and J. M. Gambetta, Hardware-efficient variational quantum eigensolver for small molecules and quantum magnets, *Nature* **549**, 242 (2017).
- [11] L. Bittel and M. Kliesch, Training Variational Quantum Algorithms Is NP-Hard, *Phys. Rev. Lett.* **127**, 120502 (2021).
- [12] J. Tilly, H. Chen, S. Cao, D. Picozzi, K. Setia, Y. Li, E. Grant, L. Wossnig, I. Rungger, G. H. Booth, and J. Tennyson, The Variational Quantum Eigensolver: A review of methods and best practices, *Physics Reports The Variational Quantum Eigensolver: A Review of Methods and Best Practices*, **986**, 1 (2022).
- [13] D. Wecker, M. B. Hastings, N. Wiebe, B. K. Clark, C. Nayak, and M. Troyer, Solving strongly correlated electron models on a quantum computer, *Phys. Rev. A* **92**, 062318 (2015).
- [14] I. D. Kivlichan, J. McClean, N. Wiebe, C. Gidney, A. Aspuru-Guzik, G. K.-L. Chan, and R. Babbush, Quantum Simulation of Electronic Structure with Linear Depth and Connectivity, *Phys. Rev. Lett.* **120**, 110501 (2018).
- [15] F. Arute, K. Arya, R. Babbush, D. Bacon, J. C. Bardin, R. Barends, S. Boixo, M. Broughton, B. B. Buckley, D. A. Buell, B. Burkett, N. Bushnell, Y. Chen, Z. Chen, B. Chiaro, R. Collins, W. Courtney, S. Demura, A. Dunsworth, E. Farhi, A. Fowler, B. Foxen, C. Gidney, M. Giustina, R. Graff, S. Habegger, M. P. Harrigan, A. Ho, S. Hong, T. Huang, W. J. Huggins, L. Ioffe, S. V. Isakov, E. Jeffrey, Z. Jiang, C. Jones, D. Kafri, K. Kechedzhi, J. Kelly, S. Kim, P. V. Klimov, A. Korotkov, F. Kostritsa, D. Landhuis, P. Laptev, M. Lindmark, E. Lucero, O. Martin, J. M. Martinis, J. R. McClean, M. McEwen, A. Megrant, X. Mi, M. Mohseni, W. Mroczkiewicz, J. Mutus, O. Naaman, M. Neeley, C. Neill, H. Neven, M. Y. Niu, T. E. O'Brien, E. Ostby, A. Petukhov, H. Putterman, C. Quintana, P. Roushan, N. C. Rubin, D. Sank, K. J. Satzinger, V. Smelyanskiy, D. Strain, K. J. Sung, M. Szalay, T. Y. Takeshita, A. Vainsencher, T. White, N. Wiebe, Z. J. Yao, P. Yeh, and A. Zalcman, Hartree-Fock on a superconducting qubit quantum computer, *Science* **369**, 1084 (2020).
- [16] A. Anand, P. Schleich, S. Alperin-Lea, P. W. K. Jensen, S. Sim, M. Díaz-Tinoco, J. S. Kottmann, M. Degroote, A. F. Izmaylov, and A. Aspuru-Guzik, A quantum computing view on unitary coupled cluster theory, *Chem. Soc. Rev.* **51**, 1659 (2022).
- [17] F. A. Evangelista, G. K.-L. Chan, and G. E. Scuseria, Exact parameterization of fermionic wave functions via unitary coupled cluster theory, *J. Chem. Phys.* **151**, 244112 (2019).
- [18] Q. Wang, M. Li, C. Monroe, and Y. Nam, Resource-Optimized Fermionic Local-Hamiltonian Simulation on a Quantum Computer for Quantum Chemistry, *Quantum* **5**, 509 (2021).
- [19] J. S. Kottmann and A. Aspuru-Guzik, Optimized low-depth quantum circuits for molecular electronic structure using a separable-pair approximation, *Phys. Rev. A* **105**, 032449 (2022).
- [20] H. L. Tang, V. O. Shkolnikov, G. S. Barron, H. R. Grimley, N. J. Mayhall, E. Barnes, and S. E. Economou, Qubit-ADAPT-VQE: An Adaptive Algorithm for Constructing Hardware-Efficient Ansatzes on a Quantum Processor, *PRX Quantum* **2**, 020310 (2021).
- [21] T. E. O'Brien, G. Anselmetti, F. Gkritis, V. E. Elfving, S. Polla, W. J. Huggins, O. Oumarou, K. Kechedzhi, D. Abanin, R. Acharya, I. Aleiner, R. Allen, T. I. Andersen, K. Anderson, M. Ansmann, F. Arute, K. Arya, A. Asfaw, J. Atalaya, D. Bacon, J. C. Bardin, A. Bengtsson, S. Boixo, G. Bortoli, A. Bourassa, J. Bovaird, L. Brill, M. Broughton, B. Buckley, D. A. Buell, T. Burger, B. Burkett, N. Bushnell, J. Campero, Y. Chen, Z. Chen, B. Chiaro, D. Chik, J. Cogan, R. Collins, P. Conner, W. Courtney, A. L. Crook, B. Curtin, D. M. Debroy, S. Demura, I. Drozdov, A. Dunsworth, C. Erickson, L. Faoro, E. Farhi, R. Fatemi, V. S. Ferreira, L. F. Burgos, E. Forati, A. G. Fowler, B. Foxen, W. Giang, C. Gidney, D. Gilboa, M. Giustina, R. Gosula, A. G. Dau, J. A. Gross, S. Habegger, M. C. Hamilton, M. Hansen, M. P. Harrigan, S. D. Harrington, P. Heu, J. Hilton, M. R. Hoffmann, S. Hong, T. Huang, A. Huff, L. B. Ioffe, S. V. Isakov, J. Iveland, E. Jeffrey, Z. Jiang, C. Jones, P. Juhás, D. Kafri, J. Kelly, T. Khattar, M. Khezri, M. Kieferová, S. Kim, P. V. Klimov, A. R. Klots, R. Kothari, A. N. Korotkov, F. Kostritsa, J. M. Kreike-

- baum, D. Landhuis, P. Laptev, K. Lau, L. Laws, J. Lee, K. Lee, B. J. Lester, A. T. Lill, W. Liu, W. P. Livingston, A. Locharla, E. Lucero, F. D. Malone, S. Mandra, O. Martin, S. Martin, J. R. McClean, T. McCourt, M. McEwen, A. Megrant, X. Mi, A. Mieszala, K. C. Miao, M. Mohseni, S. Montazeri, A. Morvan, R. Movassagh, W. Mruczkiewicz, O. Naaman, M. Neeley, C. Neill, A. Nersisyan, H. Neven, M. Newman, J. H. Ng, A. Nguyen, M. Nguyen, M. Y. Niu, S. Omonije, A. Opremcak, A. Petukhov, R. Potter, L. P. Pryadko, C. Quintana, C. Rocque, P. Roushan, N. Saei, D. Sank, K. Sankaragomathi, K. J. Satzinger, H. F. Schurkus, C. Schuster, M. J. Shearn, A. Shorter, N. Shutty, V. Shvarts, J. Skruzny, V. Smelyanskiy, W. C. Smith, R. Somma, G. Sterling, D. Strain, M. Szalay, D. Thor, A. Torres, G. Vidal, B. Villalonga, C. V. Heidweiller, T. White, B. W. K. Woo, C. Xing, Z. J. Yao, P. Yeh, J. Yoo, G. Young, A. Zalcman, Y. Zhang, N. Zhu, N. Zobrist, C. Gogolin, R. Babbush, and N. C. Rubin, **Purification-based quantum error mitigation of pair-correlated electron simulations** (2022), [arXiv:2210.10799](https://arxiv.org/abs/2210.10799).
- [22] R. N. Tazhigulov, S.-N. Sun, R. Haghshenas, H. Zhai, A. T. Tan, N. C. Rubin, R. Babbush, A. J. Minnich, and G. K.-L. Chan, **Simulating Models of Challenging Correlated Molecules and Materials on the Sycamore Quantum Processor**, *PRX Quantum* **3**, 040318 (2022).
- [23] V. E. Elfving, B. W. Broer, M. Webber, J. Gavartin, M. D. Halls, K. P. Lorton, and A. Bochevarov, **How will quantum computers provide an industrially relevant computational advantage in quantum chemistry?** (2020), [arXiv:2009.12472](https://arxiv.org/abs/2009.12472).
- [24] B. Nagy and F. Jensen, **Basis Sets in Quantum Chemistry**, in *Reviews in Computational Chemistry* (John Wiley & Sons, Ltd, 2017) Chap. 3, pp. 93–149.
- [25] J. Romero, R. Babbush, J. R. McClean, C. Hempel, P. Love, and A. Aspuru-Guzik, **Strategies for quantum computing molecular energies using the unitary coupled cluster ansatz** (2018), [arXiv:1701.02691](https://arxiv.org/abs/1701.02691).
- [26] J. M. Arrazola, O. Di Matteo, N. Quesada, S. Jahangiri, A. Delgado, and N. Killoran, **Universal quantum circuits for quantum chemistry**, *Quantum* **6**, 742 (2022), [arXiv:2106.13839](https://arxiv.org/abs/2106.13839).
- [27] G.-L. R. Anselmetti, D. Wierichs, C. Gogolin, and R. M. Parrish, **Local, expressive, quantum-number-preserving VQE ansätze for fermionic systems**, *New J. Phys.* **23**, 113010 (2021).
- [28] C. H. Chee, A. M. Mak, D. Leykam, P. K. Barkoutsos, and D. G. Angelakis, **Computing Electronic Correlation Energies using Linear Depth Quantum Circuits** (2022), [arXiv:2207.03949](https://arxiv.org/abs/2207.03949).
- [29] I. Kerenidis and A. Prakash, **Quantum machine learning with subspace states** (2022), [arXiv:2202.00054](https://arxiv.org/abs/2202.00054).
- [30] G. W. Stewart, **Computing the CS decomposition of a partitioned orthonormal matrix**, *Numer. Math.* **40**, 297 (1982).
- [31] E. S. Gawlik, Y. Nakatsukasa, and B. D. Sutton, **A Backward Stable Algorithm for Computing the CS Decomposition via the Polar Decomposition**, *SIAM J. Matrix Anal. Appl.* **39**, 1448 (2018).
- [32] S. Johri, S. Debnath, A. Mocherla, A. Singk, A. Prakash, J. Kim, and I. Kerenidis, **Nearest centroid classification on a trapped ion quantum computer**, *npj Quantum Inf* **7**, 1 (2021).
- [33] K. Wan, W. J. Huggins, J. Lee, and R. Babbush, **Match-gate Shadows for Fermionic Quantum Simulation** (2022), [arXiv:2207.13723](https://arxiv.org/abs/2207.13723).
- [34] G. Ortiz, J. E. Gubernatis, E. Knill, and R. Laflamme, **Quantum algorithms for fermionic simulations**, *Phys. Rev. A* **64**, 022319 (2001).
- [35] Z. Jiang, K. J. Sung, K. Kechedzhi, V. N. Smelyanskiy, and S. Boixo, **Quantum Algorithms to Simulate Many-Body Physics of Correlated Fermions**, *Phys. Rev. Appl.* **9**, 044036 (2018).
- [36] See Supplemental Material for more details.
- [37] I. Kerenidis, J. Landman, and N. Mathur, **Classical and Quantum Algorithms for Orthogonal Neural Networks** (2021), [arXiv:2106.07198](https://arxiv.org/abs/2106.07198).
- [38] J. D. Whitfield, J. Biamonte, and A. Aspuru-Guzik, **Simulation of electronic structure Hamiltonians using quantum computers**, *Molecular Physics* **109**, 735 (2011).
- [39] Y. S. Yordanov, D. R. M. Arvidsson-Shukur, and C. H. W. Barnes, **Efficient quantum circuits for quantum computational chemistry**, *Phys. Rev. A* **102**, 062612 (2020).
- [40] C. Plascencia, J. Wang, and A. K. Wilson, **Importance of the ligand basis set in ab initio thermochemical calculations of transition metal species**, *Chemical Physics Letters* **685**, 496 (2017).
- [41] P. Virtanen, R. Gommers, T. E. Oliphant, M. Haberland, T. Reddy, D. Cournapeau, E. Burovski, P. Peterson, W. Weckesser, J. Bright, S. J. van der Walt, M. Brett, J. Wilson, K. J. Millman, N. Mayorov, A. R. J. Nelson, E. Jones, R. Kern, E. Larson, C. J. Carey, Í. Polat, Y. Feng, E. W. Moore, J. VanderPlas, D. Laxalde, J. Perktold, R. Cimrman, I. Henriksen, E. A. Quintero, C. R. Harris, A. M. Archibald, A. H. Ribeiro, F. Pedregosa, and P. van Mulbregt, **SciPy 1.0: Fundamental algorithms for scientific computing in Python**, *Nat. Methods* **17**, 261 (2020).
- [42] Q. Sun, T. C. Berkelbach, N. S. Blunt, G. H. Booth, S. Guo, Z. Li, J. Liu, J. D. McClain, E. R. Sayfutyarova, S. Sharma, S. Wouters, and G. K.-L. Chan, **PySCF: The Python-based simulations of chemistry framework**, *WIREs Computational Molecular Science* **8**, e1340 (2018).
- [43] V. Bergholm, J. Izaac, M. Schuld, C. Gogolin, S. Ahmed, V. Ajith, M. S. Alam, G. Alonso-Linaje, B. Akash-Narayanan, A. Asadi, J. M. Arrazola, U. Azad, S. Banning, C. Blank, T. R. Bromley, B. A. Cordier, J. Ceroni, A. Delgado, O. Di Matteo, A. Dusko, T. Garg, D. Guala, A. Hayes, R. Hill, A. Ijaz, T. Isaacson, D. Ittah, S. Jahangiri, P. Jain, E. Jiang, A. Khandelwal, K. Kottmann, R. A. Lang, C. Lee, T. Loke, A. Lowe, K. McKiernan, J. J. Meyer, J. A. Montañez-Barrera, R. Moyard, Z. Niu, L. J. O’Riordan, S. Oud, A. Panigrahi, C.-Y. Park, D. Polatajko, N. Quesada, C. Roberts, N. Sá, I. Schoch, B. Shi, S. Shu, S. Sim, A. Singh, I. Strandberg, J. Soni, A. Száva, S. Thabet, R. A. Vargas-Hernández, T. Vincent, N. Vitucci, M. Weber, D. Wierichs, R. Wiersema, M. Willmann, V. Wong, S. Zhang, and N. Killoran, **PennyLane: Automatic differentiation of hybrid quantum-classical computations** (2022), [arXiv:1811.04968](https://arxiv.org/abs/1811.04968).
- [44] F. Arute, K. Arya, R. Babbush, D. Bacon, J. C. Bardin, R. Barends, R. Biswas, S. Boixo, F. G. S. L. Brandao, D. A. Buell, B. Burkett, Y. Chen, Z. Chen, B. Chiaro, R. Collins, W. Courtney, A. Dunsworth, E. Farhi, B. Foxen, A. Fowler, C. Gidney, M. Giustina, R. Graff, K. Guerin, S. Habegger, M. P. Harrigan,

- M. J. Hartmann, A. Ho, M. Hoffmann, T. Huang, T. S. Humble, S. V. Isakov, E. Jeffrey, Z. Jiang, D. Kafri, K. Kechedzhi, J. Kelly, P. V. Klimov, S. Knysh, A. Korotkov, F. Kostritsa, D. Landhuis, M. Lindmark, E. Lucero, D. Lyakh, S. Mandrà, J. R. McClean, M. McEwen, A. Megrant, X. Mi, K. Michielsen, M. Mohseni, J. Mutus, O. Naaman, M. Neeley, C. Neill, M. Y. Niu, E. Ostby, A. Petukhov, J. C. Platt, C. Quintana, E. G. Rieffel, P. Roushan, N. C. Rubin, D. Sank, K. J. Satzinger, V. Smelyanskiy, K. J. Sung, M. D. Trevithick, A. Vainsencher, B. Villalonga, T. White, Z. J. Yao, P. Yeh, A. Zalcman, H. Neven, and J. M. Martinis, Quantum supremacy using a programmable superconducting processor, *Nature* **574**, 505 (2019).
- [45] G. Li, Y. Ding, and Y. Xie, Towards Efficient Superconducting Quantum Processor Architecture Design, in *Proceedings of the Twenty-Fifth International Conference on Architectural Support for Programming Languages and Operating Systems* (ACM, Lausanne Switzerland, 2020) pp. 1031–1045.
- [46] K. Wright, K. M. Beck, S. Debnath, J. M. Amini, Y. Nam, N. Grzesiak, J.-S. Chen, N. C. Pisenti, M. Chmielewski, C. Collins, K. M. Hudek, J. Mizrahi, J. D. Wong-Campos, S. Allen, J. Apisdorf, P. Solomon, M. Williams, A. M. DuCore, A. Blinov, S. M. Kreike-meier, V. Chaplin, M. Keesan, C. Monroe, and J. Kim, Benchmarking an 11-qubit quantum computer, *Nat Commun* **10**, 5464 (2019).
- [47] C. Monroe, W. C. Campbell, L.-M. Duan, Z.-X. Gong, A. V. Gorshkov, P. W. Hess, R. Islam, K. Kim, N. M. Linke, G. Pagano, P. Richerme, C. Senko, and N. Y. Yao, Programmable quantum simulations of spin systems with trapped ions, *Rev. Mod. Phys.* **93**, 025001 (2021).
- [48] Y. Su, D. W. Berry, N. Wiebe, N. Rubin, and R. Babbush, Fault-Tolerant Quantum Simulations of Chemistry in First Quantization, *PRX Quantum* **2**, 040332 (2021).
- [49] A. Delgado, P. A. M. Casares, R. dos Reis, M. S. Zini, R. Campos, N. Cruz-Hernández, A.-C. Voigt, A. Lowe, S. Jahangiri, M. A. Martin-Delgado, J. E. Mueller, and J. M. Arrazola, Simulating key properties of lithium-ion batteries with a fault-tolerant quantum computer, *Phys. Rev. A* **106**, 032428 (2022).
- [50] R. Babbush, W. J. Huggins, D. W. Berry, S. F. Ung, A. Zhao, D. R. Reichman, H. Neven, A. D. Baczewski, and J. Lee, Quantum simulation of exact electron dynamics can be more efficient than classical mean-field methods (2023), [arXiv:2301.01203](https://arxiv.org/abs/2301.01203).

Supplementary Material

Proof of Equivalence between two definitions of an Arbitrary Slater Determinant

Consider the definition of an arbitrary Slater determinant $|\Psi_{SD}(A)\rangle$ with d occupied and $N-d$ unoccupied fermionic modes

$$|\Psi_{SD}(A)\rangle = \prod_{l=1}^d \sum_{k=1}^N A_{lk} \hat{a}_k^\dagger |\text{vac}\rangle, \quad (12)$$

where A is an $N \times d$ real matrix such that the first d columns are orthogonal and normalized,, $|\text{vac}\rangle$ is a vacuum state, and \hat{a}_k^\dagger is a creation operator acting on the k^{th} mode. Expanding the product and replacing the index k in the summation with indices k_1, \dots, k_d , we have

$$|\Psi_{SD}(A)\rangle = \left(\sum_{k_d=1}^N A_{dk_d} \hat{a}_{k_d}^\dagger \right) \dots \left(\sum_{k_1=1}^N A_{1k_1} \hat{a}_{k_1}^\dagger \right) |\text{vac}\rangle \quad (13)$$

$$= \sum_{k_d, \dots, k_1=1}^N A_{dk_d} \dots A_{1k_1} \hat{a}_{k_d}^\dagger \dots \hat{a}_{k_1}^\dagger |\text{vac}\rangle, \quad (14)$$

Meanwhile, the arbitrary Slater determinant has the alternate definition

$$|\Psi_{SD,\text{alt}}(A)\rangle = \prod_{l=1}^d \sum_{k=1}^N A_{lk} \left(\hat{a}_k^\dagger + \hat{a}_k \right) |\text{vac}\rangle, \quad (15)$$

Similarly expanding the product and replacing index k in the summation with indices k_1, \dots, k_d , we have

$$|\Psi_{SD,\text{alt}}(A)\rangle = \left(\sum_{k_d=1}^N A_{dk_d} \left(\hat{a}_{k_d}^\dagger + \hat{a}_{k_d} \right) \right) \dots \left(\sum_{k_1=1}^N A_{1k_1} \left(\hat{a}_{k_1}^\dagger + \hat{a}_{k_1} \right) \right) |\text{vac}\rangle \quad (16)$$

For $d = 1$, using $\hat{a}_j |\text{vac}\rangle = 0$ for any mode j , both Eqs. (14) and (16) becomes trivially equivalent. For $d > 1$, let us evaluate the product of the rightmost two summation terms in the product in Eqs. (16), we obtain

$$\begin{aligned} & \left(\sum_{k_2=1}^N A_{2k_2} \left(\hat{a}_{k_2}^\dagger + \hat{a}_{k_2} \right) \right) \left(\sum_{k_1=1}^N A_{1k_1} \left(\hat{a}_{k_1}^\dagger + \hat{a}_{k_1} \right) \right) \\ &= \sum_{k_2, k_1=1}^N A_{2k_2} A_{1k_1} \left(\hat{a}_{k_2}^\dagger + \hat{a}_{k_2} \right) \left(\hat{a}_{k_1}^\dagger + \hat{a}_{k_1} \right) \end{aligned} \quad (17)$$

$$= \sum_{k_2, k_1=1}^N A_{2k_2} A_{1k_1} \left(\hat{a}_{k_2}^\dagger \hat{a}_{k_1}^\dagger + \hat{a}_{k_2}^\dagger \hat{a}_{k_1} + \hat{a}_{k_2} \hat{a}_{k_1}^\dagger + \hat{a}_{k_2} \hat{a}_{k_1} \right), \quad (18)$$

$$(19)$$

Applying fermionic commutation relation $\{\hat{a}_\alpha, \hat{a}_\beta^\dagger\} = \delta_{\alpha\beta} \mathbf{I}$ yields

$$= \sum_{k_2, k_1=1}^N A_{2k_2} A_{1k_1} \left(\hat{a}_{k_2}^\dagger \hat{a}_{k_1}^\dagger + \delta_{k_2 k_1} \mathbf{I} + \hat{a}_{k_2} \hat{a}_{k_1} \right), \quad (20)$$

Since the columns of A are orthogonal, the inner product between any column i, j vanishes $\sum_k A_{ik} A_{jk} = 0$, we thus have

$$= \sum_{k_2, k_1=1}^N A_{2k_2} A_{1k_1} \left(\hat{a}_{k_2}^\dagger \hat{a}_{k_1}^\dagger + \hat{a}_{k_2} \hat{a}_{k_1} \right), \quad (21)$$

Substituting the above result Eq. (21) back into Eq. (16), and using $\hat{a}_j|\text{vac}\rangle = 0$ for any mode j gives

$$|\Psi_{\text{SD,alt}}(A)\rangle = \left(\sum_{k_d=1}^N A_{dk_d} (\hat{a}_{k_d}^\dagger + \hat{a}_{k_d}) \right) \dots \left(\sum_{k_2, k_1=1}^N A_{2k_2} A_{1k_1} \hat{a}_{k_2}^\dagger \hat{a}_{k_1}^\dagger \right) |\text{vac}\rangle. \quad (22)$$

Henceforth we consider the even and odd d cases separately. First, assuming d is even we can reapply the result Eq. (21) to the rest of the pairs of summation terms in Eq. (22), which yields

$$|\Psi_{\text{SD,alt}}(A)\rangle = \left(\sum_{k_d, k_{d-1}=1}^N A_{dk_d} A_{d-1k_{d-1}} \hat{a}_{k_d}^\dagger \hat{a}_{k_{d-1}}^\dagger \right) \dots \left(\sum_{k_2, k_1=1}^N A_{2k_2} A_{1k_1} \hat{a}_{k_2}^\dagger \hat{a}_{k_1}^\dagger \right) |\text{vac}\rangle \quad (23)$$

$$= \sum_{k_d, \dots, k_1=1}^N A_{dk_d} \dots A_{1k_1} \hat{a}_{k_d}^\dagger \dots \hat{a}_{k_1}^\dagger |\text{vac}\rangle. \quad (24)$$

Alternatively, if d is odd we have

$$|\Psi_{\text{SD,alt}}(A)\rangle = \left(\sum_{k_d=1}^N A_{dk_d} (\hat{a}_{k_d}^\dagger + \hat{a}_{k_d}) \right) \sum_{k_{d-1}, \dots, k_1=1}^N A_{d-1k_{d-1}} \dots A_{1k_1} \hat{a}_{k_{d-1}}^\dagger \dots \hat{a}_{k_1}^\dagger |\text{vac}\rangle \quad (25)$$

$$= \sum_{k_d, \dots, k_1=1}^N A_{dk_d} \dots A_{1k_1} (\hat{a}_{k_d}^\dagger \hat{a}_{k_{d-1}}^\dagger \dots \hat{a}_{k_1}^\dagger + \hat{a}_{k_d} \hat{a}_{k_{d-1}}^\dagger \dots \hat{a}_{k_1}^\dagger) |\text{vac}\rangle \quad (26)$$

$$= \sum_{k_d, \dots, k_1=1}^N A_{dk_d} \dots A_{1k_1} \left(\hat{a}_{k_d}^\dagger \hat{a}_{k_{d-1}}^\dagger \dots \hat{a}_{k_1}^\dagger + \delta_{k_d k_{d-1}} \hat{a}_{k_{d-2}}^\dagger \dots \hat{a}_{k_1}^\dagger - \hat{a}_{k_{d-1}}^\dagger \hat{a}_{k_d} \hat{a}_{k_{d-2}}^\dagger \dots \hat{a}_{k_1}^\dagger \right) |\text{vac}\rangle \quad (27)$$

$$= \sum_{k_d, \dots, k_1=1}^N A_{dk_d} \dots A_{1k_1} \left(\hat{a}_{k_d}^\dagger \hat{a}_{k_{d-1}}^\dagger \dots \hat{a}_{k_1}^\dagger - \hat{a}_{k_{d-1}}^\dagger \delta_{k_d k_{d-2}} \hat{a}_{k_{d-3}}^\dagger \dots \hat{a}_{k_1}^\dagger + \hat{a}_{k_{d-1}}^\dagger \hat{a}_{k_d} \hat{a}_{k_{d-2}}^\dagger \dots \hat{a}_{k_1}^\dagger \right) |\text{vac}\rangle \quad (28)$$

$$= \sum_{k_d, \dots, k_1=1}^N A_{dk_d} \dots A_{1k_1} \hat{a}_{k_d}^\dagger \dots \hat{a}_{k_1}^\dagger |\text{vac}\rangle, \quad (29)$$

where we have applied fermionic commutation relation in Eq. (26) and matrix orthogonality in Eqs. (27), (28).

Hence, by combining the even Eq. (24) and odd Eq. (29) results, we have established the equivalence between the alternative definition in Eq. (15) to the original definition in Eq. (12).

Proof of anti-commutation relations

Here we shall show that $\hat{p}_k = \hat{a}_k^\dagger + \hat{a}_k$ has the desired anti-commutation relation $\frac{1}{2}\{\hat{p}_i, \hat{p}_j\} = \delta_{ij}\mathbf{I}$.

$$\frac{1}{2}\{\hat{p}_i, \hat{p}_j\} = \frac{1}{2}\{\hat{a}_i^\dagger + \hat{a}_i, \hat{a}_j^\dagger + \hat{a}_j\} \quad (30)$$

$$= \frac{1}{2}((\hat{a}_i^\dagger + \hat{a}_i)(\hat{a}_j^\dagger + \hat{a}_j) + (\hat{a}_j^\dagger + \hat{a}_j)(\hat{a}_i^\dagger + \hat{a}_i)) \quad (31)$$

$$= \frac{1}{2}(\cancel{\hat{a}_i^\dagger \hat{a}_j^\dagger} + \hat{a}_i \hat{a}_j^\dagger + \hat{a}_i^\dagger \hat{a}_j + \cancel{\hat{a}_i \hat{a}_j} + \cancel{\hat{a}_j^\dagger \hat{a}_i^\dagger} + \hat{a}_j \hat{a}_i^\dagger + \hat{a}_j^\dagger \hat{a}_i + \cancel{\hat{a}_j \hat{a}_i}) \quad (32)$$

$$= \frac{1}{2}(\hat{a}_i \hat{a}_j^\dagger + \hat{a}_i^\dagger \hat{a}_j + \hat{a}_j \hat{a}_i^\dagger + \hat{a}_j^\dagger \hat{a}_i) \quad (33)$$

$$= \delta_{ij}I \quad (\text{shown}) \quad (34)$$

Here we shall show that $\hat{p}_k = \hat{Z}_{\otimes 2, 2k-2, \text{even}} \hat{X}_{2k-1} \hat{X}_{2k}$, where $\hat{Z}_{\otimes 2, 2k-2, \text{even}} = \hat{Z}_2 \otimes \hat{Z}_4 \otimes \dots \otimes \hat{Z}_{2k-2}$, used to incorporate electronic correlation into the Clifford ladder approach has the desired anti-commutation relation $\{\hat{p}_i, \hat{p}_j\} = 2\delta_{ij}\mathbf{I}$. We shall split the derivation into two cases $j = 1$ and $j < i$. Let us first consider $j = i$, where

$$2\hat{p}_i^2 = (\hat{Z}_{\otimes 2, 2i-2, \text{even}} \hat{X}_{2i-1} \hat{X}_{2i})(\hat{Z}_{\otimes 2, 2i-2, \text{even}} \hat{X}_{2i-1} \hat{X}_{2i}) \quad (35)$$

$$= 2\mathbf{I} \quad (\text{shown}) \quad (36)$$

Second, without loss of generality, let us consider $j < i$,

$$\hat{p}_j \hat{p}_i = \hat{Z}_{\otimes 2, 2j-2, \text{even}} \hat{X}_{2j-1} \hat{X}_{2j} \hat{Z}_{\otimes 2, 2i-2, \text{even}} \hat{X}_{2i-1} \hat{X}_{2i} \quad (37)$$

$$= \hat{X}_{2j-1} \hat{Z}_{2j} \hat{X}_{2j} \hat{Z}_{\otimes 2j+2, 2i-2, \text{even}} \hat{X}_{2i-1} \hat{X}_{2i}, \quad (38)$$

since $\{\hat{Z}_l, \hat{X}_l\} = 0$ for any index l we thus obtain

$$\{\hat{p}_i, \hat{p}_j\} = \hat{p}_i \hat{p}_j + \hat{p}_j \hat{p}_i \quad (39)$$

$$= \hat{X}_{2i} \hat{X}_{2i-1} \hat{Z}_{\otimes 2i-2, 2j+2, \text{even}} \hat{X}_{2j} \hat{Z}_{2j} \hat{X}_{2j-1} + \hat{X}_{2j-1} \hat{Z}_{2j} \hat{X}_{2j} \hat{Z}_{\otimes 2j+2, 2i-2, \text{even}} \hat{X}_{2i-1} \hat{X}_{2i} \quad (40)$$

$$= \hat{X}_{2i} \hat{X}_{2i-1} \hat{Z}_{\otimes 2i-2, 2j+2, \text{even}} \hat{X}_{2j-1} \underbrace{\{\hat{Z}_{2j}, \hat{X}_{2j}\}}_0 \quad (41)$$

$$= 0 \quad (\text{shown}) \quad (42)$$

Pairwise-Correlated Fermionic Beam Splitter Operator

The standard fermionic double-excitation operator $U_{kj}^{li}(\theta) = \exp \left[\theta \left(\hat{a}_i^\dagger \hat{a}_l^\dagger \hat{a}_k \hat{a}_j - \hat{a}_j^\dagger \hat{a}_k^\dagger \hat{a}_l \hat{a}_i \right) \right]$ in the Pauli form under the Jordan-Wigner mapping is given by

$$U_{kj}^{li}(\theta) = \exp \left[-\mathbf{i} \frac{\theta}{8} (\hat{X}_k \hat{Y}_j \hat{X}_l \hat{X}_i + \hat{Y}_k \hat{X}_j \hat{X}_l \hat{X}_i + \hat{Y}_k \hat{Y}_j \hat{Y}_l \hat{X}_i + \hat{Y}_k \hat{Y}_j \hat{X}_l \hat{Y}_i - \hat{X}_k \hat{X}_j \hat{Y}_l \hat{X}_i - \hat{X}_k \hat{X}_j \hat{X}_l \hat{Y}_i - \hat{Y}_k \hat{Y}_j \hat{X}_l \hat{Y}_i - \hat{X}_k \hat{Y}_j \hat{Y}_l \hat{Y}_i) \otimes_{r=k+1}^{j-1} \hat{Z}_r \otimes_{r'=l+1}^{i-1} \hat{Z}_{r'} \right], \quad (43)$$

The main paper, we used the pairwise-correlated fermionic beam splitter $\hat{U}_{j-1,j}^{i-1,i}$

$$\hat{U}_{j-1,j}^{i-1,i}(\theta) = \exp \left[-\mathbf{i} \frac{\theta}{8} (\hat{X}_{j-1} \hat{Y}_j \hat{X}_{i-1} \hat{X}_i + \hat{Y}_{j-1} \hat{X}_j \hat{X}_{i-1} \hat{X}_i + \hat{Y}_{j-1} \hat{Y}_j \hat{Y}_{i-1} \hat{X}_i + \hat{Y}_{j-1} \hat{Y}_j \hat{X}_{i-1} \hat{Y}_i - \hat{X}_{j-1} \hat{X}_j \hat{Y}_{i-1} \hat{X}_i - \hat{X}_{j-1} \hat{X}_j \hat{X}_{i-1} \hat{Y}_i - \hat{Y}_{j-1} \hat{Y}_j \hat{X}_{i-1} \hat{Y}_i - \hat{X}_{j-1} \hat{Y}_j \hat{Y}_{i-1} \hat{Y}_i) \otimes_{\substack{r=j+2 \\ \text{even}}}^{i-2} \hat{Z}_r \right], \quad (44)$$

which is a modified fermionic double-excitation operator, where the difference lies in the Pauli-Z operators used in the exponent. Here, the indices of the Pauli-Z operators are between the even-labelled qubits between $j+2$ to $i-2$ inclusive, whereas in the standard operator in Eq. (43), the indices of the Pauli-Z operators are between all qubits from $k+1$ to $j-1$ and from $l+1$ to $i-1$ inclusive.