

A NOTE ON FIVE DIMENSIONAL KISSING ARRANGEMENTS

FERENC SZÖLLŐSI

ABSTRACT. The kissing number $\tau(d)$ is the maximum number of pairwise non-overlapping unit spheres each touching a central unit sphere in the d -dimensional Euclidean space. In this note we report on how we discovered a new, previously unknown arrangement of 40 unit spheres in dimension 5. Our arrangement saturates the best known lower bound on $\tau(5)$, and refutes a ‘belief’ of Cohn–Jiao–Kumar–Torquato.

1. INTRODUCTION

This paper is based on a talk given by the author during the “Workshop on Orthogonal designs and related Combinatorics” held in Meiji University in Tokyo on January 21, 2023.

Given a central unit sphere in the d -dimensional Euclidean space \mathbb{R}^d , surrounded by $n \geq 1$ pairwise non-overlapping unit spheres at distance 2 is called a *kissing arrangement*. The main problem of interest is the determination of the maximum number $\tau(d)$ of such spheres. This is a challenging open problem, called the *kissing number problem* in geometry. For background and references we refer the reader to [1], [2], [3], and [6].

It is useful to reformulate this problem in the language of algebra. The kissing number problem asks for the maximum number n of unit vectors v_1, v_2, \dots, v_n in \mathbb{R}^d , such that their pairwise inner product satisfies $\langle v_i, v_j \rangle \leq \frac{1}{2}$ for every $i \neq j \in \{1, 2, \dots, n\}$. It is easy to see that the vertices of the cross-polytope, formed by the union of the standard basis and its negative, is a kissing arrangement, thus $\tau(d) \geq 2d$. The exact value is only known for $d \in \{1, 2, 3, 4, 8, 24\}$.

In this paper we report on some computational techniques leading to the discovery of a new, previously unknown kissing arrangement of 40 spheres in \mathbb{R}^5 . To this date this is the third such known arrangement in dimension 5 (see reference [5] from 1967) and testifies our extremely limited understanding of the structure of optimal high-dimensional sphere packings.

2. THE BEST KNOWN KISSING ARRANGEMENTS IN DIMENSION FIVE

Since the value of $\tau(d)$ has been determined up to $d \leq 4$, see [6], the first open case is dimension 5. Here two kissing arrangements showing $\tau(5) \geq 40$ were previously known [1], [4], [5]. We briefly describe these arrangements as follows.

First, the set of all $2d(d-1)$ unit vectors in \mathbb{R}^d , $d \geq 2$ having exactly two non-zero coordinates of equal absolute value $1/\sqrt{2}$

$$\mathcal{D}_d := \{\sigma([\pm 1/\sqrt{2}, \pm 1/\sqrt{2}, 0, \dots, 0]): \sigma \in S_d\}$$

forms an antipodal kissing arrangement: if $v \in \mathcal{D}_d$ then $-v \in \mathcal{D}_d$. Fixing $d = 5$ shows that $\tau(5) \geq 40$. The profile (i.e., the multiset of pairwise inner products) of \mathcal{D}_5 is given by:

$$\pi(\mathcal{D}_5) = \{[-1]^{40}, [-1/2]^{480}, [0]^{560}, [1/2]^{480}, [1]^{40}\}.$$

The profile is clearly an invariant up to change of basis.

Rather interestingly, another example can be obtained by rotating the 8 vectors among \mathcal{D}_5 with last coordinate $-1/\sqrt{2}$ by the orthogonal matrix

$$H = \frac{1}{2} \begin{bmatrix} -1 & 1 & 1 & 1 & 0 \\ 1 & -1 & 1 & 1 & 0 \\ 1 & 1 & -1 & 1 & 0 \\ 1 & 1 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 2 \end{bmatrix}.$$

This gives rise to the non-antipodal kissing arrangement (see [5])

$$\mathcal{L}_5 := (\mathcal{D}_5 \setminus \{v \in \mathcal{D}_5 : v_5 = -1/\sqrt{2}\}) \cup \{vH : v \in \mathcal{D}_5, v_5 = -1/\sqrt{2}\}$$

with profile

$$\pi(\mathcal{L}_5) = \{[-1]^{24}, [-3/4]^{64}, [-1/2]^{384}, [-1/4]^{64}, [0]^{544}, [1/2]^{480}, [1]^{40}\}.$$

It was believed that these two arrangements are the only possibilities [1], [2]. In the next section we are going to mention three computer-aided methods to discover kissing arrangements, and use them to construct a new, previously unknown arrangement of 40 unit spheres in dimension 5.

3. IN SEARCH OF KISSING ARRANGEMENTS

We call the distinct unit vectors $x, y \in \mathbb{R}^d$ *compatible*, if they represent non-overlapping spheres, that is if $\langle x, y \rangle \leq \frac{1}{2}$.

3.1. Method 1: The multiangular cloud. Fix a dimension d , a (finite!) set of inner products \mathcal{A} (with $\max \mathcal{A} \leq \frac{1}{2}$), and a d -dimensional basis B_1, B_2, \dots, B_d of pairwise compatible unit vectors, represented by the rows of a $d \times k$ matrix B . The Gram matrix of the basis vectors $G = BB^T$ is an invertible $d \times d$ matrix.

Now assume that a unit vector $v \in \mathbb{R}^k$ is (i) in the span of the basis vectors; and (ii) not only compatible with each of the basis vectors, but rather $\langle v, B_i \rangle \in \mathcal{A}$ for every $i \in \{1, \dots, d\}$. Then, the first condition says that we have a coordinate vector $c \in \mathbb{R}^d$, such that $v = cB$. The second condition says that we have a d -tuple $t \in \mathcal{A}^d$, such that $Bv^T = t^T$. Combining these two we obtain $v = tG^{-1}B$. Since v must be a unit vector, we have in addition $tG^{-1}t^T = 1$. We call the (finite) set of all such vectors

$$\mathcal{C}_{\mathcal{A}, B} := \{tG^{-1}B : t \in \mathcal{A}^d, tG^{-1}t^T = 1\} \subset \mathbb{R}^k$$

the *multiangular cloud surrounding the basis B with respect to the angle set \mathcal{A}* . Clearly, any subset of pairwise compatible vectors of $\mathcal{C}_{\mathcal{A}, B}$ together with the rows of B forms a kissing arrangement.

We are interested in finding the maximum (that is: the largest) subset(s). This in turn boils down to a clique search [7] in the graph Γ , where the vertex set is represented by the vectors in the cloud, and edges are precisely between compatible vectors. The size of the maximum clique in Γ is denoted by $\omega(\Gamma)$ as usual.

Lemma 1. *With the notations introduced in Section 3.1, we have $\tau(d) \geq d + \omega(\Gamma)$.*

3.2. Method 2: Extending a basis of \mathbb{R}^d to a basis of \mathbb{R}^{d+1} . Let d, t, \mathcal{A}, B , and G be the same as defined in Section 3.1. Assume further that $k = d+1$, and therefore there is a unit vector $B_{d+1} \in \mathbb{R}^{d+1}$ orthogonal to each vectors of the linearly independent set B . Once again we consider vectors of the form $v = tG^{-1}B \in \mathbb{R}^{d+1}$ in the linear span of the vectors B . However, if this v is not a unit a vector, but shorter, then we may consider instead the two vectors $v \pm \sqrt{1 - \langle v, v \rangle} B_{d+1}$. The finite set of all such vectors

$$\mathcal{C}'_{\mathcal{A},B} := \{tG^{-1}B \pm \sqrt{1 - tG^{-1}t^T} B_{d+1} : t \in \mathcal{A}^d, tG^{-1}t^T \leq 1\} \subset \mathbb{R}^{d+1},$$

is the multiangular cloud in \mathbb{R}^{d+1} surrounding the subset B with respect to the angle set \mathcal{A} . Similarly, a maximum subset of $\mathcal{C}'_{\mathcal{A},B}$ of pairwise compatible vectors can be found using a clique search in the corresponding graph Γ' .

Lemma 2. *With the notations introduced in Section 3.2, we have $\tau(d+1) \geq d + \omega(\Gamma')$.*

3.3. Method 3: Precomputed vectors. Let d be fixed, and let $\mathcal{C}'' \subset \mathbb{R}^d$ be a finite set of unit vectors. Let Γ'' be a graph on the vertex set of \mathcal{C}'' , where two vertices are connected if and only if the vectors are compatible. A maximum clique in Γ'' yields a kissing arrangement.

Lemma 3. *With the notations introduced in Section 3.3, we have $\tau(d) \geq \omega(\Gamma'')$.*

4. A NEW KISSING ARRANGEMENT IN DIMENSION FIVE

We illustrate our Method 2 as follows. Let $d = 4$, let $\mathcal{A} = \{-1, -1/2, 0, 1/2\}$, and let

$$B = \frac{1}{\sqrt{50}} \begin{bmatrix} 5 & 5 & 0 & 0 & 0 \\ 5 & 0 & 5 & 0 & 0 \\ 5 & 0 & 0 & 5 & 0 \\ 5 & 0 & 0 & 0 & 5 \end{bmatrix}, \quad G = BB^T = \frac{1}{2} \begin{bmatrix} 2 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 1 & 1 & 2 & 1 \\ 1 & 1 & 1 & 2 \end{bmatrix},$$

and $B_5 = \frac{1}{\sqrt{5}}[-1, 1, 1, 1, 1]$. The cloud $\mathcal{C}'_{\mathcal{A},B} \subset \mathbb{R}^5$ contains 78 vectors. A routine application of the cliquer C software [7] shows that the graph Γ' contains exactly four maximum cliques of size $\omega(\Gamma') = 36$. Two of these cliques yield via Lemma 2 a set of 40 vectors with the same profile as of \mathcal{D}_5 . The other two, however, yields 40 vectors with a distinct profile. Denoting one of these two by \mathcal{Q}_5 , we obtain:

$$\pi(\mathcal{Q}_5) = \{[-1]^{20}, [-4/5]^{60}, [-1/2]^{360}, [-3/10]^{120}, [0]^{500}, [1/5]^{20}, [1/2]^{480}, [1]^{40}\}.$$

This implies that the arrangement \mathcal{Q}_5 is not isometric to \mathcal{D}_5 or \mathcal{L}_5 . We record this as our main result.

Theorem 1. *In dimension 5 there exist at least three, pairwise non-isometric kissing-arrangements of 40 vectors: \mathcal{D}_5 , \mathcal{L}_5 , and $\mathcal{Q}_5 := (\mathcal{D}_5 \setminus \mathcal{X}) \cup \mathcal{Y}$, where the sets*

\mathcal{X} and \mathcal{Y} are the row vectors of the matrices

$$X = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 & 0 & 0 & 0 \\ 1 & 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & -1 & 0 \\ 1 & 0 & 0 & 0 & -1 \\ 0 & -1 & -1 & 0 & 0 \\ 0 & -1 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 & -1 \\ 0 & 0 & -1 & -1 & 0 \\ 0 & 0 & -1 & 0 & -1 \\ 0 & 0 & 0 & -1 & -1 \end{bmatrix}, \quad Y = \frac{1}{5\sqrt{2}} \begin{bmatrix} -1 & 1 & -4 & -4 & -4 \\ -1 & -4 & 1 & -4 & -4 \\ -1 & -4 & -4 & 1 & -4 \\ -1 & -4 & -4 & -4 & 1 \\ 4 & 1 & 1 & -4 & -4 \\ 4 & 1 & -4 & 1 & -4 \\ 4 & 1 & -4 & -4 & 1 \\ 4 & -4 & 1 & 1 & -4 \\ 4 & -4 & 1 & -4 & 1 \\ 4 & -4 & -4 & 1 & 1 \end{bmatrix}.$$

Proof. One readily checks that \mathcal{Q}_5 is a set of 40 pairwise compatible unit vectors in dimension 5. Further, the profile of the listed configurations are mutually different. \square

We remark that \mathcal{Q}_5 contains only 20 antipodal vectors, therefore it does not contain an isometric copy of the 24 antipodal vectors of \mathcal{D}_4 .

It would be interesting to analyze the rigidity [1] of \mathcal{Q}_5 , and explore higher dimensional kissing arrangements containing an isometric copy of it as a subset.

REFERENCES

- [1] H. COHN, Y. JIAO, A. KUMAR, S. TORQUATO: Rigidity of spherical codes, *Geometry & Topology* 15, 2235–2273 (2011).
- [2] J.H. CONWAY, N.J.A. SLOANE: What are all the best sphere packings in low dimensions?, *Discrete Comput Geom*, 13, 383–403 (1995).
- [3] M. GANZHINOV: Highly symmetric lines, *preprint*, [arXiv:2207.08266] (2022).
- [4] A. KORKINE, G. ZOLOTAREFF: Sur les formes quadratiques, *Math. Ann.* 6, 366–389 (1873).
- [5] J. LEECH: Five dimensional non-lattice sphere packings, *Canad. Math. Bull.* 10, 387–393 (1967).
- [6] O. MUSIN: The kissing number in four dimensions, *Annals of Mathematics*, 168, 1–32 (2008).
- [7] P. ÖSTERGÅRD: A fast algorithm for the maximum clique problem, *Discrete Applied Mathematics*, 120, 197–207, (2002).

APPENDIX: THE VECTORS FORMING \mathcal{Q}_5

We list the vectors of \mathcal{Q}_5 in machine-readable form. The vectors below should be rescaled by a factor of $\frac{1}{5\sqrt{2}}$.

$\mathcal{Q}_5 = \{ \{5, 5, 0, 0, 0\}, \{5, 0, 5, 0, 0\}, \{5, 0, 0, 5, 0\}, \{5, 0, 0, 0, 5\}, \{0, 0, 0, -5, 5\},$
 $\{0, 0, 0, 5, -5\}, \{0, 0, 0, 5, 5\}, \{0, 0, -5, 0, 5\}, \{0, 0, -5, 5, 0\}, \{0, 0, 5, 0, -5\},$
 $\{0, 0, 5, 0, 5\}, \{0, 0, 5, -5, 0\}, \{0, 0, 5, 5, 0\}, \{0, -5, 0, 0, 5\}, \{0, -5, 0, 5, 0\},$
 $\{0, -5, 5, 0, 0\}, \{0, 5, 0, 0, -5\}, \{0, 5, 0, 0, 5\}, \{0, 5, 0, -5, 0\}, \{0, 5, 0, 5, 0\},$
 $\{0, 5, -5, 0, 0\}, \{0, 5, 5, 0, 0\}, \{-5, 0, 0, 0, -5\}, \{-5, 0, 0, 0, 5\}, \{-5, 0, 0, -5, 0\},$
 $\{-5, 0, 0, 5, 0\}, \{-5, 0, -5, 0, 0\}, \{-5, 0, 5, 0, 0\}, \{-5, -5, 0, 0, 0\}, \{-5, 5, 0, 0, 0\},$
 $\{-1, 1, -4, -4, -4\}, \{-1, -4, 1, -4, -4\}, \{-1, -4, -4, 1, -4\}, \{-1, -4, -4, -4, 1\},$
 $\{4, 1, 1, -4, -4\}, \{4, 1, -4, 1, -4\}, \{4, 1, -4, -4, 1\}, \{4, -4, 1, 1, -4\},$
 $\{4, -4, 1, -4, 1\}, \{4, -4, -4, 1, 1\} \};$

DEPARTMENT OF MATHEMATICAL SCIENCES, INTERDISCIPLINARY FACULTY OF SCIENCE AND ENGINEERING, SHIMANE UNIVERSITY, MATSUE, SHIMANE, 690-8504, JAPAN
Email address: szollosi@riko.shimane-u.ac.jp