

INCREASING GAMBLER'S RUIN DURATION AND BROWNIAN MOTION EXIT TIMES

STEVEN EVANS, EROL A. PEKÖZ, AND RHONDA RIGHTER

University of California, Berkeley Department of Statistics, Boston University Questrom School of Business, and University of California, Berkeley Department of Industrial Engineering and Operations Research

Abstract

In Gambler's Ruin when both players start with the same amount of money, we show the playing time stochastically increases when the games are made more fair. We give two different arguments for this fact that extend results from Peköz & Ross (2021). We then use this to show that the exit time from a symmetric interval for Brownian motion with drift stochastically increases as the drift moves closer to zero; this result is not easily obtainable from available explicit formulas for the density.

1 INTRODUCTION

For a simple random walk $S_n = \sum_{i=1}^n X_i$, where each X_i is $+1$ with probability p and -1 with probability $1 - p$, we are interested in $T^{|k|}$, the time until the random walk hits either k or $-k$, where k is a specified positive integer. It should be noted that $T^{|k|}$ is the duration of the Gambler's Ruin when both players start with k . It was recently shown in Peköz & Ross (2021) that $T^{|k|}$ for any k is stochastically maximized when $p = 1/2$, and previously in Zhang & Ross (2021) where the weaker result in expectation was shown. In this paper we prove the stronger result that $T^{|k|}$ stochastically increases as p gets closer to $1/2$ using two very different arguments. We then use this to show that the exit time from a symmetric interval for Brownian motion with drift stochastically increases when the drift moves closer to zero, which appears also to be a new result and not easily obtainable from available formulas for the exit time probability density function. We also show that, for all p and k , $T^{|k|}$ is independent of which player wins the game. It should be noted that the technical challenges involved have left a scarcity of results for Gambler's Ruin with more than two players, see Diaconis & Ethier (2021) and the references therein.

The Gambler's Ruin problem is one of the oldest problems in probability. As told by Song & Song (2013), computing the chances each player wins was solved by Pascal and Fermat and appeared with four other problems in the earliest book published on probability theory (Huygens (1657)). Study of the distribution of the duration of the game started with de Moivre (1711) and was taken up in modern times with Feller (1968)[Chapter 14.5], who derived the formula

$$\begin{aligned} \mathbb{P}(T^{|k|} = n) &= k^{-1} 2^{n+1} [p^{(n+k)/2} (1-p)^{(n-k)/2} + p^{(n-k)/2} (1-p)^{(n+k)/2}] \\ &\quad \times \sum_{j=1}^{k-1} \cos^{n-1} \left(\frac{\pi j}{2k} \right) \sin \left(\frac{\pi j}{2k} \right) \sin \left(\frac{\pi j}{2} \right) \end{aligned}$$

for point probabilities when $n - k$ is even, and Karni (1977), who derived

$$\begin{aligned} \mathbb{P}(T^{|k|} = n) &= \left[\binom{n-1}{\frac{1}{2}(n-k)} - \binom{n-1}{\frac{1}{2}(n-3k)} - \binom{n-1}{\frac{1}{2}(n+k)} + \binom{n-1}{\frac{1}{2}(n-5k)} + \binom{n-1}{\frac{1}{2}(n+3k)} \right] \\ &\quad \times [p^{(n+k)/2}(1-p)^{(n-k)/2} + p^{(n-k)/2}(1-p)^{(n+k)/2}] \end{aligned}$$

in the case where $n \geq 5k$ and gave adjustments for $k \leq n < 5k$.

Neither of the expressions above, however, seems amenable to showing that the tail probabilities $\mathbb{P}(T^{|k|} > n)$ increase (i.e., that $T^{|k|}$ stochastically increases) as p moves towards $1/2$. The expressions can be easily seen to be increasing as p moves towards $1/2$ for sufficiently large n (the derivative with respect to p of either expression has the same sign as $n(1-2p)(p^k + (1-p)^k) + k(p^k - (1-p)^k)$), and hence the corresponding tail probabilities increase too. The expressions for the point probabilities above can, however, decrease as p moves towards $1/2$ when n is small, and thus not much can be easily said about tail probabilities in general. This leaves stochastic inequalities difficult to obtain from these expressions. Our main result, which we state next, bypasses the use of these expressions and shows that moving p towards $1/2$ stochastically increases the game duration.

Theorem 1.1. $\mathbb{P}(T^{|k|} > n)$ is increasing in p for $0 \leq p \leq 1/2$ and decreasing in p for $1/2 \leq p \leq 1$.

We prove this using two different approaches in the next sections.

2 A DECOMPOSITION ARGUMENT

Let T_0 be the first time (after time 0) the walk revisits state 0. As was done in Peköz & Ross (2021), we can use the decomposition

$$T^{|k|} = Z + \sum_{i=1}^{N-1} Y_i$$

where $N \sim \text{Geometric}(\mathbb{P}(T_0 > T^{|k|}))$, $N - 1$ is distributed as the number of times the walk returns to 0 before it first hits $\pm k$, $Z \sim (T^{|k|} | T_0 > T^{|k|})$ is the time to go from 0 to $\pm k$ given the walk does not return to 0, and $Y_i \sim (T_0 | T_0 < T^{|k|})$, $i = 1, 2, \dots$ are i.i.d. random variables representing the times to return to 0 given the walk returns to 0 before hitting $\pm k$. All of these component random variables are independent. We will show the following lemma, from which Theorem 1.1 follows immediately.

Lemma 2.1. Y_i , Z , and N are all stochastically increasing (decreasing) in p for $0 \leq p \leq 1/2$ ($1/2 \leq p \leq 1$).

To prove this lemma, let $u_i(p)$ be the chance the walk goes up next when it is at level i given it returns to 0 before reaching either $+k$ or $-k$, for $i = -k+1, -k+2, \dots, k-1$. Note that, by symmetry, $u_i(p) = 1 - u_{-i}(1-p)$ is the chance the walk goes down next from level $-i$ given it returns to 0 before reaching $\pm k$, where now the probability of the walk going up is $1-p$. We show in Lemma 2.2 (2) that $u_i(p) = u_i(1-p)$, which then implies that the probability of moving away from 0 given the walk returns to 0 before reaching $\pm k$ depends only on the player imbalance, $|p - (1-p)| = 2|p - 1/2|$, not on the direction of imbalance. Moreover, from Lemma 2.2 (3), this probability is decreasing in the player imbalance.

Lemma 2.2. *The following hold for $0 \leq p \leq 1$:*

1. $\mathbb{P}(T_0 < T^{|k|})$ is strictly increasing (decreasing) in p when $p < 1/2$ (when $p > 1/2$).
2. $u_i(p) = u_i(1-p)$ for $-k+1 \leq i \leq k-1$.
3. When $1 \leq i \leq k-2$, $u_i(p)$ is strictly increasing (decreasing) in p when $p < 1/2$ (when $p > 1/2$).

Proof of Lemma 2.2. For Part 1, note that any path of length $2n$ that starts and ends at 0 without first visiting k or $-k$ must contain n upward steps and n downward steps, and thus its probability equals $(p(1-p))^n$, which is strictly increasing in p for $p < 1/2$ and strictly decreasing in p for $p > 1/2$. The result for $\mathbb{P}(T_0 < T^{|k|})$ follows, since it is the sum of the probabilities of many such paths. This gives Part 1.

For Parts 2 and 3, we use the notation \mathbb{P}_i to denote the measure conditioned on starting the walk at level i . From our observation about the symmetry of $u_i(p)$, pick $i > 0$ without loss of generality, and let $T^j(p)$ be the time to hit level $j \geq 0$. We have

$$\begin{aligned} u_i(p) &= \frac{p\mathbb{P}_{i+1}(T_0 < T^k)}{\mathbb{P}_i(T_0 < T^k)} = \frac{p\mathbb{P}_{i+1}(T^i < T^k)\mathbb{P}_i(T_0 < T^k)}{\mathbb{P}_i(T_0 < T^k)} \\ &= p\mathbb{P}_{i+1}(T^i < T^k), \end{aligned}$$

which we combine with

$$\mathbb{P}_{i+1}(T^i < T^k) = 1 - p + p\mathbb{P}_{i+2}(T^{i+1} < T^k)\mathbb{P}_{i+1}(T^i < T^k),$$

to get

$$u_i(p) = p(1-p) + u_{i+1}(p)u_i(p). \quad (2.1)$$

For Part 2 we use backward induction. First note the boundary case $u_{k-1}(p) = u_{k-1}(1-p) = 0$. We use the induction hypothesis $u_{i+1}(p) = u_{i+1}(1-p)$ in the second equality below and (2.1) in the first equality below to get

$$u_i(p) = \frac{p(1-p)}{1-u_{i+1}(p)} = \frac{p(1-p)}{1-u_{i+1}(1-p)} = u_i(1-p)$$

and thus Part 2 holds for all $i > 0$ and then for all i looking at the walk reflected across the origin.

For Part 3, we again use backward induction. First note that $u_{k-1}(p) = 0$ and (2.1) with $i = k-2$ gives $u_{k-2}(p) = p(1-p)$, which is increasing in p for $p < 1/2$ and decreasing for $p > 1/2$. Taking the derivative, from equation (2.1) we get

$$u'_i(p) = 1 - 2p + u'_i(p)u_{i+1}(p) + u_i(p)u'_{i+1}(p) = \frac{1 - 2p + u_i(p)u'_{i+1}(p)}{1 - u_{i+1}(p)}$$

which is positive when $p < 1/2$ and negative when $p > 1/2$ using the induction hypothesis that $u'_{i+1}(p) > 0$ for $p < 1/2$ and $u'_{i+1}(p) < 0$ for $p > 1/2$. Thus Part 3 holds. \square

Proof of Lemma 2.1. Let $Y_i(p)$ be the time for the walk to go from level i to 0 given it hits 0 before it hits $\pm k$, and for fixed p , $0 < i < k$. Let $Z_i(p)$ be the time for the walk to

go from level i to $\pm k$ given it hits $\pm k$ before it hits 0, and for fixed p , $0 < i < k$. Then, from the structure of the random walk, and with $I(p) \sim \text{Bernoulli}(p)$,

$$\begin{aligned} Y(j) &=_{st} I(p)Y_1(p) + (1 - I(p))Y_{-1}(p) \\ Z(j) &=_{st} I(p)Z_1(p) + (1 - I(p))Z_{-1}(p) \\ Y_i(p) &=_{st} Y_{-i}(1 - p) \\ Z_i(p) &=_{st} Y_{-(k-i)}(1 - p). \end{aligned}$$

The results for $Y(j)$ and Z will then follow once we show that $Y_i(p) \leq_{st} Y_i(p')$ for $0 \leq p \leq p' \leq 1/2$ and $0 < i < k$, and that $Y_1(p) =_{st} Y_{-1}(p)$ and $Z_1(p) =_{st} Z_{-1}(p)$ for all p . That $Y_1(p) =_{st} Y_{-1}(p)$ and $Z_1(p) =_{st} Z_{-1}(p)$ follows from Part 2 of Lemma 2.2 and the last two equations above. We will show that $Y_i(p) \leq_{st} Y_i(p')$ by coupling the walks starting in state i so that $Y_i(p) \leq Y_i(p')$ with probability 1. Whenever the walks are at different levels, we let them step independently; since the walks both started at the same level, they will remain an even number of steps apart and thus after this step can meet but not cross. When the walks are at the same level, by Part 3 of Lemma 2.2 the chance of a step up is at least as large for the second walk as it is for the first walk, so we can couple the steps so the second walk is always no closer to the origin than the first walk. This means we will have $Y_i \leq Y_i(p')$ and thus $Y_i \leq_{st} Y_i(p')$. The result for N follows from Part 1 of Lemma 2.2. \square

Proof of Theorem 1.1. We can similarly couple $T^{|k|}(p')$ together with $T^{|k|}(p)$ on the same probability space (making the dependency on p explicit in the notation), using our decomposition and Lemma 2.2, so that $T^{|k|}(p') \geq T^{|k|}$ almost surely and the Theorem follows. \square

3 AN INDUCTIVE ARGUMENT

We will use induction on k in our second proof of Theorem 1. Consider the following, alternative, decomposition, in which the time to play a game when the boundary is $\pm k$ (call this a size k game) is a random sum of times to play smaller games. In particular, we first play a size 1 game, at which time the random walk is at ± 1 with respective probabilities p and $1 - p$. Then we play a size $k - 1$ game, at the end of which the position of the random walk will be in the set $\{-k, -(k - 2), k - 2, k\}$, depending on whether it was at $+1$ or -1 at the end of the first game, and depending on whether it went up or down by $k - 1$ steps from there. If the position at the end of the second game is $\pm(k - 2)$, then we play a third game of size 2, and so on, until we finally reach $\pm k$ and stop. Let N be the total number of such games, and let $r(n) = \mathbb{P}\{N = n | N \geq n\}$ be its hazard rate. Let $y(i)$ be such that at the end of the i 'th game, the possible positions of the random walk are $\pm k$ and $\pm y(i) \neq \pm k$ and let $d(i)$ be the size of the i 'th game. That is, let $y(1) = d(1) = 1$, and, for $i = 2, 3, \dots$, if $y(i - 1) > 0$ let $d(i) = k - y(i - 1)$ and $y(i) = |y(i - 1) - d(i)|$, and if $y(i - 1) = 0$ let $d(i) = y(i) = 1$. Note that for fixed k , $y(i)$ and $d(i)$ are deterministic sequences. We therefore have, by construction and from the Markov property of our random walk, the following.

Lemma 3.1. For $k > 1$,

$$T^{|k|} = \sum_{i=1}^N T^{|d(i)|}$$

where the $T^{|d(i)|}$'s are independent.

We note that $y(i)$, $d(i)$, $r(n)$, (and N) all depend on k , but we suppress this dependency in the notation.

Let $\pi_k^+(p) = \mathbb{P}(T^k < T^{-k})$ be the probability that the walk $S = \{S_n\}_{n=0}^\infty$, with $S_0 = 0$, ends at $+k$ before hitting $-k$, so $\pi_1^+(p) = p$ and $\pi_k^+(p) = p^k / (p^k + (1-p)^k)$, from the classic Gambler's Ruin formula (Feller (1968)). Let $\tau(j) = \sum_{i=1}^j T^{|d(i)|} | N > j$ be the time to play j subgames, conditioned on S having not hit $\pm k$ in the first j games (so $S_{\tau(j)} = \pm y(j)$).

Lemma 3.2. *For any j , $\tau(j)$ is independent of the event $\{S_{\tau(j)} = +y(j)\}$, and*

$$\mathbb{P}(S_{\tau(j)} = +y(j)) = \frac{p^{y(j)}}{p^{y(j)} + (1-p)^{y(j)}} = \pi_{y(j)}^+(p).$$

Proof. Fix j , Condition on $N > j$ and $\tau(j) = t$, and for ease of notation, let $y = y(j)$. Let ρ^+ be an arbitrary path of S that ends up at $+y$ at time t without hitting $\pm k$ before time t . Then, on path ρ^+ , S has a total of $(t-y)/2 + y$ upward steps and $(t-y)/2$ downward steps, so the probability of path ρ^+ is $p^{(t-y)/2} (1-p)^{(t-y)/2} p^y$. Let ρ^- be the same path reflected through the origin (interchanging up and down steps), so the path ρ^- ends up at $-y$ at time t , also without hitting $\pm k$ before time t . The probability of path ρ^- is $p^{(t-y)/2} (1-p)^{(t-y)/2} (1-p)^y$. Let $A(\rho^\pm)$ be the event that S takes path ρ^+ or ρ^- before hitting $\pm k$. Then

$$\begin{aligned} \mathbb{P}(S_{\tau(j)} = +y, A(\rho^\pm) | \tau(j) = t) &= \mathbb{P}(S_{\tau(j)} = +y | A(\rho^\pm), \tau(j) = t) \mathbb{P}(A(\rho^\pm) | \tau(j) = t) \\ &= \frac{p^{(t-y)/2} (1-p)^{(t-y)/2} p^y}{p^{(t-y)/2} (1-p)^{(t-y)/2} p^y + p^{(t-y)/2} (1-p)^{(t-y)/2} (1-p)^y} \mathbb{P}(A(\rho^\pm) | \tau(j) = t) \\ &= \frac{p^y}{p^y + (1-p)^y} \mathbb{P}(A(\rho^\pm) | \tau(j) = t). \end{aligned}$$

Summing over all such pairs of reflected paths, ρ^+ and ρ^- , of length t , we have for each t ,

$$\mathbb{P}(S_{\tau(j)} = +y | \tau(j) = t) = \frac{p^y}{p^y + (1-p)^y} = \pi_y^+(p).$$

Thus, the event that $S_{\tau(j)} = +y$ is independent of $\tau(j)$, and $\mathbb{P}(S_{\tau(j)} = +y) = \pi_y^+(p)$. \square

A similar argument gives the following corollary.

Corollary 3.3. *The random variable $T^{|k|}(p)$ is independent of whether the walk first hits $+k$ or $-k$ for all p and for any k .*

The following are also corollaries of Lemma 3.2.

Corollary 3.4. *For $0 \leq p \leq 1/2$, $\pi_k^+(p)$ is increasing in p , from $\pi_k^+(0) = 0$ to $\pi_k^+(1/2) = 1/2$.*

Corollary 3.5. *In the decomposition of Lemma 3.1, for fixed $k > 1$, N is independent of the $T^{|d(i)|}$'s, and, for $n = 1, 2, \dots$,*

$$r(n) = \begin{cases} \pi_{y(n-1)}^+(p) \pi_{d(n)}^+(p) + (1 - \pi_{y(n-1)}^+(p))(1 - \pi_{d(n)}^+(p)) & \text{if } y(n-1) > 0 \\ 0 & \text{if } y(n-1) = 0 \end{cases}.$$

Proof. That N is independent of the $T^{|d(i)|}$'s follows from Corollary 3.3, because, for each subgame, the probability that it is the last game is independent of the time to play the game. That $r(n) = 0$ if $y(n-1) = 0$ is immediate because then $y(n) = d(n) = 1$. From Lemma 3.2, for $y(n-1) > 0$ and given $N \geq n$, the probability that the walk is at $y(n-1)$

(respectively, $-y(n-1)$) at the end of the $n-1$ 'st game, i.e., at time $\tau(n-1)$, is $\pi_{y(n-1)}^+(p)$ (respectively, $1 - \pi_{y(n-1)}^+(p)$). The probability that the walk hits $\pm k$ at the end of the n 'th game given it has not yet hit $\pm k$, is the probability of going up by $d(n)$ from $y(n-1)$ or going down by $d(n)$ from $-y(n-1)$, i.e.,

$$r(n) = \mathbb{P}\{N = n | N \geq n\} = \pi_{y(n-1)}^+(p)\pi_{d(n)}^+(p) + (1 - \pi_{y(n-1)}^+(p))(1 - \pi_{d(n)}^+(p)).$$

□

Proof of Theorem 1.1. We use induction on k , where the result follows trivially for $T^{|1|} \equiv 1$. Fix k and assume the result holds for all $j = 1, \dots, k-1$, and therefore for each subgame i because $d(i) < k$. That is, assume $T^{|d(i)|}$ is stochastically increasing in p (for $0 \leq p \leq 1/2$). Then we will have the result for k , i.e., $T^{|k|}$ is stochastically increasing in p , from Lemma 3.1 and the induction hypothesis, as long as N is stochastically increasing in p . We will show the stronger result, that N gets larger in the hazard rate sense as p increases, i.e., its hazard rate $r(n)$ is decreasing (nonstrictly) in p for all n . If $y(n-1) = 0$ then $r(n) = 0$ is trivially decreasing in p . Define, for fixed n such that $y(n-1) \neq 0$,

$$g_n(p) = 1 - r(n) = \pi_{y(n-1)}^+(p)(1 - \pi_{d(n)}^+(p)) + (1 - \pi_{y(n-1)}^+(p))\pi_{d(n)}^+(p).$$

Then, taking the derivative with respect to p and using Corollary 3.4 for $0 \leq p \leq 1/2$,

$$\begin{aligned} g'_n(p) &= \pi_{y(n-1)}^{+'}(p) + \pi_{d(n)}^{+'}(p) - 2\pi_{y(n-1)}^{+'}(p)\pi_{d(n)}^+(p) - 2\pi_{y(n-1)}^+(p)\pi_{d(n)}^{+'}(p) \\ &\geq \pi_{y(n-1)}^{+'}(p) + \pi_{d(n)}^{+'}(p) - \pi_{y(n-1)}^{+'}(p) - \pi_{d(n)}^{+'}(p) = 0. \end{aligned}$$

□

Remark 3.6. We note that the decomposition can be simplified when k is even by choosing $y(1) = k/2$ instead of $y(1) = 1$:

$$T^{|k|}(p) = \sum_{i=1}^N \left(T_i^{|k/2|} + \tilde{T}_i^{|k/2|} \right)$$

where $T_i^{|k/2|} =_{st} \tilde{T}_i^{|k/2|} =_{st} T^{|k/2|}$, and N , $T_i^{|k/2|}$, and $\tilde{T}_i^{|k/2|}$ for all i are independent, and

$$N \sim \text{Geometric}((\pi_{k/2}^+(p))^2 + (1 - \pi_{k/2}^+(p))^2).$$

4 A COROLLARY FOR BROWNIAN MOTION WITH DRIFT

Letting $B(t)$ denote standard Brownian motion at time t , for $k > 0$ consider the time

$$T_\mu = \inf\{t \geq 0 : \mu t + B(t) \notin [-k, k]\}$$

that Brownian motion with drift μ first exits the interval $[-k, k]$. Borodin, A. & Salminen, P. (2002)[p. 309 equation (3.0.2), and p. 641] gives the probability density function for T_μ as

$$\mathbb{P}(T_\mu \in dt) = (e^{-\mu k} + e^{\mu k})e^{-\mu^2 t/2} \sum_{i=-\infty}^{\infty} \frac{k + 4ik}{\sqrt{2\pi}t^{3/2}} e^{-(k+4ik)^2/(2t)} dt. \quad (4.1)$$

Using weak convergence of the paths of asymmetric simple random walk to Brownian motion with drift, we obtain the following immediate corollary from Theorem 1.1.

Corollary 4.1. *With the definitions above, if $0 \leq \mu \leq \mu'$ then $T_\mu \geq_{st} T_{\mu'}$.*

This result shows that the exit time from a symmetric interval for Brownian motion with drift is stochastically increased as the drift is moved towards zero. In particular, it is stochastically maximized when there is no drift. As far as we can tell, this result is new and does not seem easily obtainable from the explicit form (4.1) of the density.

5 ACKNOWLEDGEMENTS

We greatly appreciate the feedback from the referees, which improved the presentation.

REFERENCES

- Andrei N. Borodin & Paavo Salminen. *Handbook of Brownian Motion - Facts and Formulae*, Second Edition, Springer 2002.
- P. Diaconis & S. N. Ethier. Gambler's Ruin and the ICM, *preprint on <https://arxiv.org/abs/2011.07610>*
- Feller, W. (1968). *An introduction to probability theory and its applications*. Vol. I. New York: John Wiley & Sons Inc.
- Huygens, C. (1657). *De Ratiociniis in Ludo Aleae*, printed in *Exercitationum Mathematicarum* by F. van Schooten, Elsevirii, Leiden. Reprinted in *Oeuvres*, 14, (1920).
- Karni, E. (1977). The Probability Distribution of the Duration of the Game in the Classical Ruin Problem. *Journal of Applied Probability*, 14(2), 416-420.
- Moivre, A. de (1711). De Mensura Sortis, seu, de Probabilitate Eventuum in Ludis a Casu Fortuito Pendentibus, *Philosophical Transactions*, 27, 213–264.
- Peköz, E. & Ross, S., Fair Gambler's Ruin stochastically maximizes playing time. Accepted to appear, *Advances in Applied Probability* (2022).
- Ross, Sheldon M. (2019). *Introduction to Probability Models*, 12th Edition. Amsterdam: Academic Press. Print.
- Song, Seongjoo, & Song, Jongwoo. (2013). A Note on the History of the Gambler's Ruin Problem. *Communications for Statistical Applications and Methods*, 20(2), 157–168.
- Zhang, Z., and Ross, S. (2021). Finding the Best Dueler, *Preprint*.