

# Coresets for Clustering with General Assignment Constraints

Lingxiao Huang\*  
Nanjing University

Shaofeng H.-C. Jiang†  
Peking University

Jian Li‡  
Tsinghua University

Xuan Wu§  
Huawei TCS Lab

## Abstract

Designing small-sized *coresets*, which approximately preserve the costs of the solutions for large datasets, has been an important research direction for the past decade. We consider coreset construction for a variety of general constrained clustering problems. We significantly extend and generalize the results of a very recent paper [BCAJ<sup>+</sup>22], by demonstrating that the idea of hierarchical uniform sampling [Che09, BCAJ<sup>+</sup>22] can be applied to efficiently construct coresets for a very general class of constrained clustering problems with general assignment constraints, including capacity constraints on cluster centers, and assignment structure constraints for data points (modeled by a convex body  $\mathcal{B}$ ).

Our main theorem shows that a small-sized  $\varepsilon$ -coreset exists as long as a complexity measure  $\text{Lip}(\mathcal{B})$  of the structure constraint, and the *covering exponent*  $\Lambda_\varepsilon(\mathcal{X})$  for metric space  $(\mathcal{X}, d)$  are bounded. The complexity measure  $\text{Lip}(\mathcal{B})$  for convex body  $\mathcal{B}$  is the Lipschitz constant of a certain transportation problem constrained in  $\mathcal{B}$ , called *optimal assignment transportation problem*. We prove nontrivial upper bounds of  $\text{Lip}(\mathcal{B})$  for various polytopes, including the general matroid basis polytopes, and laminar matroid polytopes (with better bound). Even without any structure constraint (in which case  $\text{Lip}(\mathcal{B}) = 1$ ), our analysis allows us to obtain several new coreset results for capacitated clustering using existing bounds of  $\Lambda_\varepsilon(\mathcal{X})$  for several metric spaces, such as doubling metrics and shortest-path metrics of graphs that exclude a fixed minor, greatly generalizing several previous works. As an application of our general theorem, we construct the first coreset for the fault-tolerant clustering problem (with or without capacity upper/lower bound) for the above metric spaces, in which the fault-tolerance requirement is captured by a uniform matroid basis polytope.

---

\*Email: [huanglingxiao1990@126.com](mailto:huanglingxiao1990@126.com)

†Email: [shaofeng.jiang@pku.edu.cn](mailto:shaofeng.jiang@pku.edu.cn)

‡Email: [lapordge@gmail.com](mailto:lapordge@gmail.com)

§Email: [wu3412790@gmail.com](mailto:wu3412790@gmail.com)

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
1.1	Our Contributions . . . . .	2
1.2	Technical Overview . . . . .	5
1.3	Related Work . . . . .	8
<b>2</b>	<b>Preliminaries</b>	<b>9</b>
<b>3</b>	<b>Coresets for Clustering with General Assignment Constraints</b>	<b>10</b>
3.1	The Coreset Construction Algorithm . . . . .	11
3.2	Proof of Theorem 3.1: Analysis of Algorithm 1 . . . . .	12
3.3	Proof of Lemma 3.11: Error Analysis for Rings . . . . .	19
3.4	Proof of Lemma 3.12: Error Analysis for Groups . . . . .	26
<b>4</b>	<b>Bounding the Lipschitz Constant <math>\text{Lip}(\mathcal{B})</math></b>	<b>30</b>
4.1	Lipschitz Constant for (Laminar) Matroid Basis Polytopes . . . . .	30
4.1.1	Lipschitz Constant for Matroid Basis Polytopes . . . . .	33
4.1.2	Lipschitz Constant for Laminar Matroid Basis Polytopes . . . . .	37
4.2	Lipschitz Constant $\text{Lip}(\mathcal{B})$ can be Unbounded . . . . .	38
<b>5</b>	<b>Bounding the Covering Exponent <math>\Lambda_\varepsilon(\mathcal{X})</math></b>	<b>39</b>
<b>A</b>	<b>Proof of Lemma 5.3: <math>\Lambda_\varepsilon(\mathcal{X})</math> Bound via Shattering Dimension</b>	<b>47</b>
<b>B</b>	<b>Proof of Lemma 5.6: <math>\Lambda_\varepsilon(\mathcal{X})</math> Bound via Doubling Dimension</b>	<b>47</b>

# 1 Introduction

We study coresets for clustering with general assignment constraints. Clustering is a fundamental data analysis task that receives significant attention in various areas. In the (center-based) clustering problem, given as input a metric space  $(\mathcal{X}, d)$  and a weighted data set  $P \subseteq \mathcal{X}$  with weight function  $w_P : P \rightarrow \mathbb{R}_{\geq 0}$ , the clustering cost is defined for a set of clustering centers  $C$  of  $k$  points from  $\mathcal{X}$ , and an assignment function  $\sigma : P \times C \rightarrow \mathbb{R}_+$  that assigns each data point  $p$  to the centers (such that  $\|\sigma(p, \cdot)\|_1 = w_P(p)$  which ensures the total assignment of  $p$  is exactly the weight of the point  $w_P(p)$ <sup>1</sup>), as an  $\ell_z$  aggregation of the distances from the data points to the centers  $C$  weighted by  $\sigma$ , i.e.,

$$\text{cost}_z^\sigma(P, C) := \sum_{x \in P} \sum_{c \in C} \sigma(x, c) \cdot (d(x, c))^z. \quad (1)$$

The goal of the clustering problem is to find a center set  $C$  and assignment  $\sigma$  that minimizes the clustering cost. If there is no additional constraint on  $\sigma$ , then it is optimal to assign the data points to the (unique) nearest center, which is the case of vanilla  $(k, z)$ -CLUSTERING. However, the general formulation above allows a point to be assigned fractionally to multiple centers (which is sometimes called soft clustering) if we impose some additional constraint on  $\sigma$ .

**Assignment Constraints** Many types of constrained clustering can be captured by imposing constraints on the assignment function  $\sigma$ , that appear as either *capacity constraints* on cluster centers or *structure constraints* on  $\sigma(p, \cdot)$  for points  $p$  (or a combination of both). Specifically, in a collective view from the centers, a *capacity* constraint enforces certain lower and/or upper bound on the total amount assigned to every center  $c \in C$ , i.e.,  $l_c \leq \|\sigma(\cdot, c)\|_1 \leq u_c$ . Such constraint is widely considered, and notable examples include capacitated clustering [CGTS02, LSS04, CHK12, Li16] and fair clustering [CKLV17]. A *structure* constraint, on the other hand, focuses on a single point  $p$ , and puts restrictions on the subset of centers that  $p$  may be assigned to. More concretely, a *structure* constraint requires that the vector  $\sigma(p, \cdot)$  can only take values from a constrained set in addition to the aforementioned constraint  $\|\sigma(p, \cdot)\|_1 = w_P(p)$ . Various structure constraints appear abundantly in different applications. One important example is the fault tolerance constraint, which requires each point to be assigned to at least  $l \geq 1$  centers. Fault tolerance constraint has been studied extensively in the clustering/facility location literature [KPS00, SS08, HHL<sup>+</sup>16]. Structure constraints can capture more advanced fault-tolerance requirements: for example, the centers are partitioned into several groups (centers in each group located in the same geographical region), and we require that each point is connected to centers in at least  $h$  different regions [BDHR05]. For another example, in the method of model-based clustering [ZG03], each center comes from one of the  $k$  predefined classes, and prior knowledge puts restrictions on the classes that each point may be assigned to, which can be captured by a similar structure constraint: suppose  $k$  center classes are divided into some categories, and we require that a point be assigned to one center from each category, which can be modeled by a partition matroid constraint.

**Coresets** We focus on coresets for constrained clustering with general assignment constraints. Roughly speaking, an  $\varepsilon$ -coreset  $S$  is a tiny proxy of the data set  $P$ , such that the cost evaluated on both  $S$  and  $P$  are within  $(1 \pm \varepsilon)$  factor for all potential centers  $C$  and assignments. The coreset is a powerful technique that can be used to compress large datasets, speed up existing algorithms, and design efficient approximation schemes [CL19, BFS21, BJKW21a]. The concept of coreset has

---

<sup>1</sup>Here, for some function of the form  $\sigma : X \times Y \rightarrow \mathbb{R}_{\geq 0}$ , we write  $\sigma(x, \cdot)$  as the vector  $u \in \mathbb{R}^Y$  such that  $\forall y \in Y, u_y = \sigma(x, y)$ . The notation  $\sigma(\cdot, y)$  is defined similarly.

also found many applications in modern sublinear models, such as streaming algorithms [HM04], distributed algorithms [BEL13] and dynamic algorithms [HK20], by combining it with the so-called merge-and-reduce framework [HM04].

The study of coresets size bounds for clustering has been very fruitful, especially the case of vanilla  $(k, z)$ -CLUSTERING (i.e., without constraints). A series of works focus on the Euclidean spaces [HM04, HK07, FL11, FSS20, SW18, HV20, CSS21, CLSS22] and near-optimal size bounds have been obtained in more recent works for  $k$ -MEDIAN and  $k$ -MEANS [HV20, CSS21, CLSS22]. Going beyond Euclidean spaces, another series of works provide coresets of small size in other types of metrics, such as doubling metrics [HJLW18, CSS21] and graph shortest-path metrics [BBH<sup>+</sup>20, BJKW21a]. More interestingly, throughout the line of research, various fundamental techniques have been proposed, including the importance sampling framework [FL11, FSS20] which generally applies to problems other than clustering, and a more recently developed hierarchical sampling framework [Che09, CSS21, BCAJ<sup>+</sup>22] that makes use of uniform sampling in a novel way.

**Challenge: Coresets for Constrained Clustering** Unfortunately, coresets for constrained clustering has been less understood. In particular, only the capacity constraints were considered, and the research focus was on coresets for fair clustering [CKLV17] and capacitated clustering [CGTS02]. While significant progress has been made in a series of works [SSS19, HJV19, CL19, BFS21], they achieved coresets of size either depending on  $n$  or exponential in  $d$  (which is the Euclidean dimension). As noted in many of these works, the fundamental difficulty is that standard frameworks, such as importance sampling, do not seem to work well for capacity constraints. Very recently, a breakthrough has been made in [BCAJ<sup>+</sup>22] where the first coresets for both fair and capacitated  $k$ -MEDIAN in Euclidean  $\mathbb{R}^d$  with size  $\text{poly}(k\varepsilon^{-1})$  were obtained, via an improved hierarchical uniform sampling framework. However, even with this new framework, the solution is still somewhat ad-hoc, and it is unclear if the result can be adapted to more general assignment constraints such as the above-mentioned structure constraints, and/or other metrics such as doubling metrics and graph shortest-path metrics. Indeed, a perhaps more fundamental gap is that, a systematic characterization of what types of assignment constraints allow small coresets, is missing in the literature.

## 1.1 Our Contributions

Our main contribution is a characterization of families of assignment constraints that admit small coresets. Specifically, we propose a fairly general model of assignment constraints and we define a new parameter as its complexity measure. Our main result (Theorem 1.4) is a generic coreset construction, whose size bound only depends on the proposed parameter, together with another “dimension” parameter that measures the complexity of the metric space (and  $\text{poly}(k\varepsilon^{-1})$ ). We establish new bounds for the complexity of several general constraint families, and we obtain various new coreset results for constrained clustering by combining these new bounds with our main theorem. Next, we discuss our contributions in more detail.

**Modeling Assignment Constraints** We consider two types of general assignment constraints in clustering literature: *capacity constraints* on cluster centers and *structure constraints* on  $\sigma(p, \cdot)$  for each points  $p$  (or both).

The capacity constraints are modeled similarly as in previous works [SSS19, HJV19, CL19, BFS21, BCAJ<sup>+</sup>22], where we simply consider a vector  $h \in \mathbb{R}^k$  ( $k$  is the number of centers), and require that the mass assigned to each center  $c$  equals  $h_c$ .

**Definition 1.1 (Capacity constraints).** We say an assignment function  $\sigma : P \times C \rightarrow \mathbb{R}_{\geq 0}$  is consistent with  $h \in \mathbb{R}^k$ , denoted as  $\sigma \sim h$ , if  $\|\sigma(\cdot, c)\|_1 = h_c$  for every center  $c \in [k]$ , which means the total assignment to a center  $c$  is exactly  $h_c$ .<sup>2</sup> Equivalently, we can write  $\sum_p \sigma(p, \cdot) = h$ .

A structure constraint constrains the range of the assignment vector  $\sigma(p, \cdot)$  for each points  $p$ . We model a structure constraint by a convex body  $\mathcal{B} \subseteq \Delta_k$ , where  $\Delta_k := \left\{x \in \mathbb{R}_{\geq 0}^k : \sum_{i \in [k]} x_i = 1\right\}$  denotes the simplex in  $\mathbb{R}^k$ .

**Definition 1.2 (Structure constraints).** Given convex body  $\mathcal{B} \subseteq \Delta_k$ , we say an assignment function  $\sigma : P \times C \rightarrow \mathbb{R}_{\geq 0}$  is consistent with  $\mathcal{B}$ , denoted as  $\sigma \sim \mathcal{B}$ , if

$$\frac{\sigma(p, \cdot)}{\|\sigma(p, \cdot)\|_1} = \frac{\sigma(p, \cdot)}{w_P(p)} \in \mathcal{B} \quad \text{for every } p \in P,$$

where the normalizing factor appears because vector  $\sigma(p, \cdot)$  has to satisfy  $\|\sigma(p, \cdot)\|_1 = w_P(p)$ .

We list a few examples of structure constraints as follows.

1.  $\mathcal{B} = \Delta_k$ . This is no additional constraint besides  $\|\sigma(p, \cdot)\|_1 = w_P(p)$ . So the points are free to be assigned to the nearest center.
2.  $\mathcal{B} = \left\{x \in \Delta_k : x_i \leq \frac{1}{l}, \forall i \in [k]\right\}$  for some integer  $l \geq 1$ . In this case, the cheapest way to assign  $p$  is to connect it to the  $l$  nearest centers. This captures the fault-tolerant clustering where we require a point to be assigned to at least  $l$  points. Such fault-tolerant constraints have been studied extensively in a variety of clustering problems [KPS00, SS08, HHL<sup>+</sup>16].
3.  $\mathcal{B}$  is a (scaled) matroid basis polytope. This is a significant generalization of the above constraint which corresponds to a uniform matroid polytope. As alluded before [BDHR05, ZG03], more advanced fault-tolerance requirements can be captured by partition matroid polytope. As another application in data placement problem [BRS08], consider a network where each center stores one type of data object, and a point (client) demands  $c$  types of data object. Such assignment constraint can be naturally captured by a *laminar matroid* basis polytope.

In this paper, we consider the case where all points  $p$  are subject to the same structure constraint  $\mathcal{B}$ . When an assignment function  $\sigma$  satisfies both capacity constraint  $h$  and structure constraint  $\mathcal{B}$ , we denote it as  $\sigma \sim (\mathcal{B}, h)$  and call the pair of constraints  $(\mathcal{B}, h)$  a *general assignment constraint*.

**Coresets for Clustering with General Assignment Constraints** Under the general assignment constraint  $(\mathcal{B}, h)$ , we define the cost of center set  $C$  for point set  $P$  as

$$\text{cost}_z(P, C, \mathcal{B}, h) := \min_{\sigma \sim (\mathcal{B}, h)} \text{cost}_z^\sigma(P, C). \quad (2)$$

That is for the center  $C$ , the cost is computed via the min-cost assignment  $\sigma$  that satisfies both  $\mathcal{B}$  and  $h$ . Our notion of coreset is defined for this new formulation of the cost function, as follows.

---

<sup>2</sup>We only require the capacities to be exactly  $h_c$  instead of placing a lower and/or upper bound, since we would preserve the cost for all  $h$  *simultaneously* in our coreset, so the lower/upper bounds are interpreted as a family of equality.

**Definition 1.3** ( $\varepsilon$ -Coreset for  $(k, z)$ -Clustering with general assignment constraints). For a dataset  $P \subseteq \mathcal{X}$  and a structure constraint  $\mathcal{B}$ , an  $\varepsilon$ -coreset for  $(k, z)$ -CLUSTERING with general assignment constraints is a (weighted) set  $S \subseteq X$  that satisfies

$$\forall C \in \mathcal{X}^k, h \in w_P(P) \cdot \mathcal{B}, \quad \text{cost}_z(S, C, \mathcal{B}, h) \in (1 \pm \varepsilon) \cdot \text{cost}_z(P, C, \mathcal{B}, h), \quad (3)$$

where  $w_P(P) := \sum_{p \in P} w_P(p)$ , and  $\text{cost}(P, C, \mathcal{B}, h)$  is defined as in (2).

The requirement of  $h \in w_P(P) \cdot \mathcal{B}$  is to ensure  $h$  is *feasible*, i.e.,  $\exists \sigma \sim h$  such that  $\|\sigma(p, \cdot)\|_1 = w_P(p)$  for  $p \in P$ .<sup>3</sup> This notion of coreset preserves the cost for all centers  $C$  and capacity constraints  $h$  *simultaneously*, and for a structure constraint  $\mathcal{B}$  given in advance. As also noted in previous works [SSS19, HJV19, BFS21, BCAJ<sup>+</sup>22], the guarantee that the cost is preserved for all  $h$  simultaneously implies that such a coreset simultaneously captures all types of upper/lower bound capacity constraints of the form  $l_c \leq \|\sigma(\cdot, c)\|_1 \leq u_c$ .

**Main Theorem** In our main theorem (Theorem 1.4), we show that a small coreset exists, as long as the structure constraints have bounded complexity  $\text{Lip}(\mathcal{B}) = O_k(1)$  (see Definition 3.6 and Definition 3.7), which only depends on the convex body  $\mathcal{B}$ , and the *covering exponent*  $\Lambda_\varepsilon(\mathcal{X})$  (see Definition 3.8) can be bounded independent of  $n$ . Hence, the main theorem systematically reduces the problem of constructing coresets of size  $\text{poly}(k\varepsilon^{-1})$ , to the mathematical problems of bounding the parameters  $\text{Lip}(\mathcal{B})$  and  $\Lambda_\varepsilon(\mathcal{X})$ . The parameter  $\text{Lip}(\mathcal{B})$  is new, and is defined as the Lipschitz constant of a certain transportation procedure inside the convex body  $\mathcal{B}$ . Due to its conceptual and technical importance, we provide a high-level overview in Section 1.2. On the other hand, the covering exponent  $\Lambda_\varepsilon(\mathcal{X})$  for metric space  $(\mathcal{X}, d)$  is closely related to several combinatorial dimension notions such as the VC-dimension and the (fat) shattering dimension of the set system formed by all metric balls which have been extensively studied in previous works, e.g. [LS10, FL11, FSS20, HJLW18, BJKW21a].

**Theorem 1.4 (Informal; see Theorem 3.1).** *There is an algorithm that given  $0 < \varepsilon < 1$ , metric space  $(\mathcal{X}, d)$ ,  $n$ -point data set  $P \subseteq \mathcal{X}$  and the structure constraint modeled by convex body  $\mathcal{B} \subseteq \Delta_k$ , computes in  $\tilde{O}(nk)$  time an  $\varepsilon$ -coreset of size  $\tilde{O}_z(\text{Lip}(\mathcal{B})^2 \cdot \Lambda_\varepsilon(\mathcal{X}) \cdot k^3 \varepsilon^{-3z})$  for  $(k, z)$ -CLUSTERING with general assignment constraints (which preserves the clustering cost subject to the structure constraint  $\mathcal{B}$  and all capacity constraints  $h \in w_P(P) \cdot \mathcal{B}$  as in (3)) with high probability.*

Interestingly, unlike many previous works that use ad-hoc methods to deal with constraints in specific metric spaces (such as Euclidean  $\mathbb{R}^d$ ) [SSS19, HJV19, CL19, BFS21, BCAJ<sup>+</sup>22], the required bounds for  $\text{Lip}(\mathcal{B})$  and  $\Lambda_\varepsilon(\mathcal{X})$  are completely decoupled and may be dealt with independently. Hence, even without any constraint on  $\mathcal{B}$ , i.e.,  $\mathcal{B} = \Delta_k$  where we have  $\text{Lip}(\mathcal{B}) = 1$ , and only considering the capacity constraints, we can already obtain many new coreset results by simply using known bounds of  $\Lambda_\varepsilon(\mathcal{X})$  (see Remark 5.8), which greatly generalizes all previous works. Specifically, combining with the existing/immediate  $\Lambda_\varepsilon(\mathcal{X})$  bounds for various metric spaces [CSS21, BT15], we obtain coresets of size  $\text{poly}(k\varepsilon^{-1})$  for fair  $(k, z)$ -CLUSTERING and capacitated  $(k, z)$ -CLUSTERING (and potentially many more), in e.g., doubling metrics, shortest-path metrics of graphs that exclude a fixed minor. Previously, such a coreset size independent of  $n$  is only known for Euclidean spaces [HJV19, BCAJ<sup>+</sup>22].

---

<sup>3</sup>To see this, note that we require  $\sigma(p, \cdot) \in w_P(p) \cdot \mathcal{B}$ , hence  $h = \sum_{p \in P} \sigma(p, \cdot) = w_P(P) \cdot \mathcal{B}$ .

**New  $\text{Lip}(\mathcal{B})$  Bounds: Matroid Basis Polytope** In light of Theorem 1.4, one can see that proving upper bound of  $\text{Lip}(\mathcal{B})$  is crucial for bounding the coreset size. In fact, one can easily construct polytope  $\mathcal{B}$ , which is simply defined by some linear constraints, such that  $\text{Lip}(\mathcal{B})$  is unbounded (See Section 4). In this paper, we focus on an important class of polytopes, called matroid basis polytopes (see e.g., [Sch03]). A matroid basis polytope is the convex hull of all 0/1 indicator vectors of the basis of a matroid. Matroids generalize many combinatorial structures and are popular ways to model the structure of an assignment/solution in various contexts, such as online computation [BIK07, BIKK18], matching [Law01, LSV13], diversity maximization [AMT13] and variants of clustering problems [CLLW16, KKN+11, KKN+15, Swa16]. For a general matroid basis polytope  $\mathcal{B}$ , we prove  $\text{Lip}(\mathcal{B}) \leq k - 1$  (see Theorem 4.7). We also provide an improved  $\text{Lip}(\mathcal{B}) \leq \ell + 1$  bound for the special case of laminar matroids of depth  $\ell$  (see Theorem 4.11). This readily implies coresets of size  $\text{poly}(k\varepsilon^{-1})$  for any general assignment constraints with a matroid basis polytope  $\mathcal{B}$ , under various types of metric spaces.

**Simultaneous Coresets for Multiple  $\mathcal{B}$ 's** Since our coreset can handle all capacity constraints simultaneously, one may be interested in whether our coresets can also handle multiple structure constraints  $\mathcal{B}$ 's simultaneously as well. We show that this is indeed possible for several families of  $\mathcal{B}$ . A similar idea, called simultaneous or one-shot coresets, has also been considered in the literature [BLL18, BJKW19]. This type of coresets is particularly useful when some key hyperparameters, in our case  $\mathcal{B}$ , are not given/known but need to be figured out by experiments, since a single coreset can be reused to support all the experiments. Fortunately, our coreset in Theorem 1.4 has a powerful feature that it does not use any specific structure of the given  $\mathcal{B}$  except an upper bound of  $\text{Lip}(\mathcal{B})$  in the algorithm. Hence, for a family  $\mathcal{F}$  of  $m \geq 1$  different  $\mathcal{B}$ 's, one can apply Theorem 1.4 with a union bound, so that with an additional  $\text{poly}(\log(m))$  factor in the coreset size, the returned coreset is a coreset for all  $\mathcal{B}$ 's in  $\mathcal{F}$  simultaneously. Below, we discuss the size bounds of the simultaneous coresets for two useful families  $\mathcal{F}$  and their potential applications.

1. Let  $\mathcal{F}$  be the collection of all (scaled) uniform matroid basis polytopes ( $\text{Lip}(\mathcal{B}) \leq 2$ ). Since the ground set is  $[k]$ , we have  $|\mathcal{F}| = k$  (each corresponding to a cardinality constraint). Then we can achieve a simultaneous coreset  $S$  for every  $\mathcal{B} \in \mathcal{F}$  by increasing the coreset size in Theorem 1.4 by a multiplicative  $\log k$  factor, say  $\tilde{O}_z(\Lambda_\varepsilon(\mathcal{X}) \cdot k^3 \varepsilon^{-3z})$ . Consequently,  $S$  is a simultaneous coreset for all fault-tolerant clusterings.
2. Let  $\mathcal{F}$  be the collection of all (scaled) partition matroid basis polytopes ( $\text{Lip}(\mathcal{B}) \leq 3$ ). Since the ground set is  $[k]$ , there are at most  $k^k$  partition ways. For each partition, there are at most  $k^k$  different selections of rank functions (see Definition 2.5). Thus, we have  $|\mathcal{F}| \leq k^{2k}$  and we can achieve a simultaneous coreset for every  $\mathcal{B} \in \mathcal{F}$  by increasing the coreset size in Theorem 1.4 by a multiplicative  $k \log(2k)$  factor, say  $\tilde{O}_z(\Lambda_\varepsilon(\mathcal{X}) \cdot k^4 \varepsilon^{-3z})$ .

## 1.2 Technical Overview

**Complexity Measure of Structure Constraint:  $\text{Lip}(\mathcal{B})$**  We start with a more detailed discussion on the new notion of  $\text{Lip}(\mathcal{B})$  due to its conceptual and technical importance. The definition of  $\text{Lip}(\mathcal{B})$  (Definitions 3.6 and 3.7) may be interpreted in several ways. We start with an explanation from a technical perspective of coreset construction. A natural way of building a coreset which we also use, is to draw independent samples  $S$  from data points  $P$  (and re-weight). To analyze  $S$ , let us fix some capacity constraint  $h$  and some center  $C$ . Let  $\sigma^*$  be an assignment of  $h$ , i.e.,  $\text{cost}_z(P, C, \mathcal{B}, h) = \text{cost}_z^{\sigma^*}(P, C)$ . Then, one can convert  $\sigma^*$  to  $\sigma : S \times C \rightarrow \mathbb{R}_{\geq 0}$  for the sample

$S \subseteq P$ , by setting  $\sigma(p, \cdot) := w_S(p) \cdot \sigma^*(p, \cdot)$  and  $w_S(p) = \frac{w_P(P)}{|S|}$  for  $p \in S$  (where we can guarantee that  $w_S(S) = w_P(P)$ ). Even though this assignment  $\sigma$  may slightly violate the capacity constraint  $h$ , the violation, denoted as  $\|h - h'\|_1$  where  $h'$  is the capacity induced by  $\sigma$ , is typically very small (by concentration inequalities), and it may be charged to  $\text{cost}_z(P, C, \mathcal{B}, h)$ . However, we still need to transport  $\sigma$  to  $\sigma'$  so that it is consistent with  $h'$ . More precisely, we need to find  $\sigma' : S \times C \rightarrow \mathbb{R}_{\geq 0}$  for the sample  $S$  that satisfies  $\sigma' \sim (\mathcal{B}, h)$ , such that the total transportation  $\|\sigma - \sigma'\|_1$ <sup>4</sup> (which relates to the cost change), is minimized. We call this minimum transportation plan the *optimal assignment transportation* (OAT). Since we eventually wish to bound the OAT cost against the mentioned  $\|h - h'\|_1$ , our  $\text{Lip}(\mathcal{B})$  is defined as the universal upper bound of the OAT cost relative to  $\|h - h'\|_1$  over all  $h, h', \sigma$ , which can also be viewed as the Lipschitz constant of OAT.

Another perspective of interpreting the notion of OAT is via the well-known optimal transportation. Specifically, the minimum  $L_1$  transportation cost for turning  $h$  to  $h'$  without any constraint is exactly  $\|h - h'\|_1$ . Hence, compared with optimal transportation, our notion adds additional requirements that the transportation plan must be inside of  $\mathcal{B}$ , and from a given starting assignment  $\sigma$ . We care about the worst-case relative ratio  $\text{Lip}(\mathcal{B})$ , which measures how many times more expensive the constrained optimal transportation cost OAT than the optimal transportation cost  $\|h - h'\|_1$ . Even though notions of constrained optimal transportation were also considered in the literature, see e.g., [TT13, KM13, KM15, KMS15] and a survey [PC<sup>+</sup>19, Chapter 10.3] for more, we are not aware of any previous work that studies exactly the same problem. While we are able to provide nontrivial upper bound of  $\text{Lip}(\mathcal{B})$  for some specific polytope  $\mathcal{B}$  in this work, bounding  $\text{Lip}(\mathcal{B})$  for other convex set  $\mathcal{B}$ , as well as the efficient computation of it (which we do not need) may be of independent interest for future research.

**Technical Overview: Main Theorem** The proof of our main theorem makes use of a recent hierarchical uniform sampling (HUS) framework developed in [BCAJ<sup>+</sup>22] (Algorithm 1), which is inspired by an earlier idea of [Che09]. In particular, our algorithm is almost the same as in [BCAJ<sup>+</sup>22], except that the total number of samples is different. Our main contribution is an analysis of the sampling complexity of HUS for  $(k, z)$ -CLUSTERING with general assignment constraints.

In the algorithm of HUS, one starts with partitioning the input into a collection of *rings* and *groups* (Theorem 3.3), and the analysis is done for rings and groups separately. For each ring  $R$ , we take a uniform sample  $S$  and show that the estimation error  $|\text{cost}_z(R, C, \mathcal{B}, h) - \text{cost}_z(S, C, \mathcal{B}, h)|$  induced by  $S$  is small for every center set  $C$  and every capacity constraint  $h$  (Lemma 3.11). A natural idea is to discretize the hyper-parameter space  $\mathcal{X}^k \times (|R| \cdot \mathcal{B})$  into a small number of representative pairs, and show that the error is preserved for every representative pair  $(C, h)$  with very high probability (using e.g., concentration bounds). Finally, the coresets property follows by a union bound over all representative pairs. Similar ideas have been applied in [CL19, BCAJ<sup>+</sup>22] for the capacitated  $k$ -MEDIAN, but they only worked for Euclidean spaces.

Indeed, it requires several new ideas in our analysis to realize the improvements and generalizations over previous results, . We highlight our new ideas in the following, and they also constitute the main aspects of improvements over the mentioned previous works [CL19, BCAJ<sup>+</sup>22]. We first discuss a key step/observation in handling the structure constraint  $\mathcal{B}$  which is used to obtain the concentration bound for a fixed representative pair  $(C, h)$  (Lemma 3.15). This step is also useful for obtaining coresets for  $(k, z)$ -CLUSTERING for general  $z \geq 1$ . Finally, we discuss our new idea of discretizing the hyper-parameter space, which helps to decouple the covering exponent and  $\text{Lip}(\mathcal{B})$ , and also leads to several new coreset results for various metric spaces.

---

<sup>4</sup>Here, we interpret  $\sigma, \sigma'$  as vectors on  $\mathbb{R}^{S \times C}$ .

1. **(Handling Structure Constraint  $\mathcal{B}$ )** To control the coresets error  $|\text{cost}_z(R, C, \mathcal{B}, h) - \text{cost}_z(S, C, \mathcal{B}, h)|$ , the idea is to show that  $\text{cost}_z(S, C, \mathcal{B}, h)$  concentrates on its expectation  $\mathbb{E}_S[\text{cost}_z(S, C, \mathcal{B}, h)]$  (Lemma 3.16) and to show the expectation is very close to  $\text{cost}_z(R, C, \mathcal{B}, h)$  (Lemma 3.17). The proof of the former one about the expectation follows easily from concentration inequalities, but that of the latter one is much more difficult and constitutes a major part of our analysis. Let  $\sigma^*$  be an optimal assignment of  $h$  on  $R$ , i.e.,  $\text{cost}_z(R, C, \mathcal{B}, h) = \text{cost}_z^{\sigma^*}(R, C)$ . We convert  $\sigma^*$  to  $\sigma : S \times C \rightarrow_{\geq 0}$ , by setting  $\sigma(p, \cdot) := w_S(p) \cdot \sigma^*(p, \cdot)$  for  $p \in S$  (as mentioned in the previous paragraph “Complexity Measure of Structure Constraint”), and can show that  $\text{cost}_z^\sigma(S, C) \approx \text{cost}_z^{\sigma^*}(R, C)$  ((17) in the proof of Claim 3.20). We are done if  $\sigma \sim h$ , but unfortunately this does not hold in general. Hence, we turn to show the existence of an assignment  $\sigma' \sim (\mathcal{B}, h)$  on  $S$  such that  $|\text{cost}_z^\sigma(S, C) - \text{cost}_z^{\sigma'}(S, C)|$  is small enough. This existence of such  $\sigma'$  is shown in Claim 3.20, and here we sketch the main technical ideas. We reduce the problem of bounding  $|\text{cost}_z^\sigma(S, C) - \text{cost}_z^{\sigma'}(S, C)|$  to bounding the mass movement  $\|\sigma - \sigma'\|_1$  from  $\sigma$  to  $\sigma'$ , based on a novel idea that we can safely ignore the total assignment cost of  $R$  and  $S$  to remote centers (denoted by  $C_{\text{far}}$ ), and the difference between the assignment cost induced by  $\sigma$  and  $\sigma'$  to  $C \setminus C_{\text{far}}$  is proportional to  $\|\sigma - \sigma'\|_1$ , due to the generalized triangle inequality (Lemma 2.1). Now it suffices to require that  $\|\sigma - \sigma'\|_1 \leq \tilde{O}_z(\varepsilon^{z+1}|R|)$  for bounding  $|\text{cost}_z^\sigma(S, C) - \text{cost}_z^{\sigma'}(S, C)|$ . By the definition of  $\text{Lip}(\mathcal{B})$ , we only need to make sure that  $\|h - h'\|_1$  ( $h'$  is induced by  $\sigma$ ) is as small up to  $\tilde{O}_z(\varepsilon^{z+1}|R|)/\text{Lip}(\mathcal{B})$  (Claim 3.22), which is again guaranteed by McDiarmid’s Inequality (Theorem 2.2). The additional factor  $1/\text{Lip}(\mathcal{B})$  results in the term  $\text{Lip}(\mathcal{B})^2$  in our coresets size.
2. **(Dealing with General  $z \geq 1$ )** The analysis for  $k$ -MEDIAN in [CL19] relies on the fact that for any two points  $p, q \in R$  and a center  $c$ ,  $|d(p, c) - d(q, c)| \leq d(p, q)$  by the triangle inequality. However, this property does not extend to  $z > 1$ , and even replacing the usage of triangle inequality with the generalized triangle inequality (Lemma 2.1) directly does not suffice, specifically, for remote centers to ring  $R$ .<sup>5</sup> To overcome this difficulty, we again use the idea that we can safely ignore the total assignment cost of  $R$  and  $S$  to remote centers (denoted by  $C_{\text{far}}$ ), and the assignment cost of  $S$  to the remaining centers  $C \setminus C_{\text{far}}$  is highly concentrated to that of  $R$  (Lemmas 3.16 and 3.17), where we can use the generalized triangle inequality. This idea enables us to handle all  $z \geq 1$ .
3. **(Decoupling Metric Structure from Constraint Complexity)** At a high level, we discretize the center sets  $C \in \mathcal{X}^k$  and capacity constraints  $h \in |R| \cdot \mathcal{B}$  separately, and show their Cartesian products form a discretization of the entire hyper-parameter space  $\mathcal{X}^k \times (|R| \cdot \mathcal{B})$ . This high-level idea was also considered in previous works [BCAJ<sup>+</sup>22], but compared with the previous implementation of this idea, our discretization of the center space  $\mathcal{X}^k$  directly relates to the *covering exponent*  $\Lambda_\varepsilon(\mathcal{X})$  of the metric space, which is a complexity measure of general metric spaces, instead of a more geometric discretization based on the notion of metric balls which are more specific to Euclidean spaces. Another perhaps more significant difference, is that the discretization of the capacity constraint space must be subject to the structure constraint  $\mathcal{B}$ . While we use the overall plan as in [BCAJ<sup>+</sup>22, Lemma 4.4, Definition of  $\mathcal{F}$ ], where we aim to ensure that for every  $h$ , there exists some representative  $h'$  such that  $\|h - h'\|_1$  is small enough, we must deal with the structure constraint  $\mathcal{B}$  using a way similar to the first bullet (about handling structure constraint), and this additionally requires a smaller

---

<sup>5</sup>It was claimed in [CL19] that the analysis can be extended to  $z = 2$  ( $k$ -means). But the argument in their paper is incomplete and it is unclear how to fix it in a straightforward manner.

$\|h - h'\|_1 \leq O_z(\varepsilon^{z+1}|R|/\text{Lip}(\mathcal{B}))$ , which results in  $\text{poly}(\log \text{Lip}(\mathcal{B}))$  factors in the final coreset size. We implement all these ideas in Lemma 3.10, and this turns out to be a crucial step for decoupling  $\text{Lip}(\mathcal{B})$  and  $\Lambda_\varepsilon(\mathcal{X})$  in our coreset.

For each group  $G$ , we still construct a two-point coreset  $D$  (Definition 3.4) as in [BCAJ<sup>+</sup>22], and prove that the estimation error  $|\text{cost}_z(G, C, \mathcal{B}, h) - \text{cost}_z(D, C, \mathcal{B}, h)|$  is *deterministically* small for every tuple  $(C, \mathcal{B}, h)$  (Lemmas 3.25 and 3.26). The main difficulty of introducing structure constraints  $\mathcal{B}$  is to prove that  $\text{cost}_z(G, C, \mathcal{B}, h) \leq (1 + \varepsilon) \cdot \text{cost}_z(D, C, \mathcal{B}, h)$  (Item 2 of Lemma 3.26), since it is unknown what the optimal assignment  $\sigma^* \sim (\mathcal{B}, h)$  on  $D$  looks like. We handle this issue by a surprising observation for a two-point coreset (Remark 3.28):  $\text{cost}_z^\sigma(D, C) \approx \text{cost}_z(D, C, \mathcal{B}, h)$  holds for any assignment  $\sigma$  of  $h$ .

**Bounds of  $\text{Lip}(\mathcal{B})$**  For  $\mathcal{B} = \Delta_k$  which is the unconstrained case, we have  $\text{Lip}(\mathcal{B}) = 1$ . However, the geometric structure of  $\mathcal{B}$  can indeed result in a significantly larger, even unbound,  $\text{Lip}(\mathcal{B})$ . In particular, we show that the value  $\text{Lip}(\mathcal{B})$  is unbounded, even for a very simple  $\mathcal{B}$  defined by two knapsack constraints (see Theorem 4.14). On the other hand, we do manage to show  $\text{Lip}(\mathcal{B}) \leq k - 1$  for an very important family of  $\mathcal{B}$ , namely, matroid basis polytopes (Theorem 4.7). To analyze  $\text{Lip}(\mathcal{B})$  for the matroid case, we first show in Lemma 4.3 that it suffices to restrict our attention to the case where the initial assignment  $\sigma$  corresponds to vertices of the polytope (or basis) and so does terminal assignment  $\sigma'$  (which we need to find). Our argument is combinatorial and heavily relies on constructing augmenting paths (Definition 4.9). Roughly speaking, an augmenting path is defined by a sequence of basis  $(I_1, \dots, I_m)$  and a sequence of elements  $(a_0, a_1, \dots, a_m)$  where each  $I_i$  is an initial basis and the following exchange property holds:  $I_i \cup \{a_{i-1}\} \setminus \{a_i\} \in \mathcal{M}$  for all  $i$ . The exchange property allows us to perform a sequence of basis exchanges to transport the mass from  $a_0$  to  $a_m$ . Hence, the key is to find such an augmenting path in which  $a_0$  is an element with  $h_{a_0} < h'_{a_0}$  and  $a_m$  is one with  $h_{a_m} > h'_{a_m}$ , for some  $h, h' \in \mathcal{B}$ . Then performing the exchange operation reduces the value of  $\|h - h'\|_1$  by  $2\tau$  with total transportation cost  $2m\tau$  for some small  $\tau > 0$ . Thus,  $\text{Lip}(\mathcal{B})$  is bounded by the path length  $m$ . We show that  $m \leq k - 1$  when  $\mathcal{B}$  is a matroid basis polytope (Lemma 4.10). To show the existence of such an augmenting path, we leverage several combinatorial properties of a matroid, as well as the submodularity of the rank function, and the non-crossing property of tight subsets. We further improve the bound to  $m \leq \ell + 1$  for laminar matroid basis polytopes of depth at most  $\ell \geq 1$  (Lemma 4.12), which results in bounds  $\text{Lip}(\mathcal{B}) \leq 2$  for uniform matroids and  $\text{Lip}(\mathcal{B}) \leq 3$  for partition matroids.

### 1.3 Related Work

Approximation algorithms have been extensively studied for constrained clustering problems. We focus on the  $k$ -median case for several notable problems in the following discussion. For fair  $k$ -median, [BCFN19] provided a bi-criteria  $O(1)$ -approximation in general metric spaces, but the solution may violate the capacity/fair constraint by an additive error. [BIO<sup>+</sup>19] gave  $O(\log n)$ -approximation without violating the constraints in Euclidean spaces. For capacitated  $k$ -median,  $O(1)$ -approximation were known in general metrics, but they either need to violate the capacity constraint [DL16, BRU16] or the number of centers  $k$  [Li17, Li16] by a  $(1 + \varepsilon)$  factor. Both problems admit polynomial-time algorithms that have better approximation and/or no violation of constraints when  $k$  is not considered a part of the input [ABM<sup>+</sup>19, CL19, FZH<sup>+</sup>20, BFS21]. For fault-tolerant  $k$ -median, a constant approximation was given in [SS08], and  $O(1)$ -approximation also exists even when the number of centers that each data point needs to connect can be different [HHL<sup>+</sup>16]. Finally, we mention a variant of clustering called matroid  $k$ -median which also admit

$O(1)$ -approximation in general metrics [KKN<sup>+</sup>11]. In this problem, a matroid is defined on the vertices of the graph, and only the center set that form independent sets may be chosen. While this sounds different from the constraints that we consider, our coresets actually captures this trivially since our coresets preserves the cost for all centers (not only those that form independent sets).

Apart from constrained clustering, coresets were also considered for clustering with generalized objectives (but without constraints). Examples include clustering with outliers [FS12, HJLW18, HJLW22], projective clustering [FL11, FSS20, TWZ<sup>+</sup>22], clustering with missing values [BJKW21b], ordered-weighted clustering [BJKW19] and clustering with panel data [HSV21].

## 2 Preliminaries

Throughout, we are given an underlying metric space  $(\mathcal{X}, d)$ , integer  $k \geq 1$ , constant  $z \geq 1$ , and precision parameter  $\varepsilon \in (0, 1)$ . For integer  $n \geq 1$ , let  $[n] := \{1, \dots, n\}$ . For a function  $\sigma : X \times Y \rightarrow \mathbb{R}$ , for  $i \in X$  we write  $\sigma(i, \cdot)$  as the vector  $u_i \in \mathbb{R}^Y$  such that  $u_i(j) := \sigma(i, j)$ , and define  $\sigma(\cdot, j)$  similarly. Furthermore, sometimes we interpret  $\sigma$  as a vector in  $\mathbb{R}^{X \times Y}$ , so  $\|\sigma\|_1 = \sum_{i \in X, j \in Y} |\sigma(i, j)|$ . We use  $\Delta_k$  to denote the simplex  $\{x \in \mathbb{R}_{\geq 0}^k : \|x\|_1 = 1\}$ . Given a point  $a \in \mathcal{X}$  and a radius  $r > 0$ , we define  $\text{Ball}(a, r) := \{x \in \mathcal{X}, d(x, a) \leq r\}$  to be the ball or radius  $r$  centered at  $a$ .

We use the following well-known generalized triangle inequalities for  $(k, z)$ -CLUSTERING; see e.g., [MSSW19, BCAJ<sup>+</sup>22] for more variants.

**Lemma 2.1 (Generalized triangle inequality [BCAJ<sup>+</sup>22, Lemma 2.1]).** *Let  $a, b, c \in \mathcal{X}$  and  $z \geq 1$ . For every  $t \in (0, 1]$ , the following inequalities hold:*

$$d^z(a, b) \leq (1+t)^{z-1} d^z(a, c) + \left(1 + \frac{1}{t}\right)^{z-1} d^z(b, c),$$

and

$$|d^z(a, c) - d^z(b, c)| \leq t \cdot d^z(a, c) + \left(\frac{3z}{t}\right)^{z-1} \cdot d^z(a, b).$$

We also use the following concentration inequality for analysis.

**Theorem 2.2 (McDiarmid's Inequality [vH14, Theorem 3.11]).** *Let  $E$  be a ground set and  $n \geq 1$  be an integer. Let  $g : E^n \rightarrow \mathbb{R}$  be a function satisfying that for any sequence  $(x_1, x_2, \dots, x_n) \in E^n$ , there exists a universal constant  $\delta_i > 0$  for each  $i \in [n]$  such that*

$$\delta_i \geq \sup_{y \in E} g(x_1, \dots, x_{i-1}, y, x_{i+1}, \dots, x_n) - \inf_{y \in E} g(x_1, \dots, x_{i-1}, y, x_{i+1}, \dots, x_n).$$

Then for independent random variables  $X_1, \dots, X_n$ , we have for every  $t > 0$ ,

$$\Pr[g(X_1, \dots, X_n) - \mathbb{E}_{S_R}[g(X_1, \dots, X_n)] \geq t] \leq e^{-\frac{2t^2}{\sum_{i \in [n]} \delta_i^2}}.$$

We consider a general structure constraint captured by the so-called matroid basis polytope. For more information about matroid and matroid basis polytope, see the classic reference [Sch03]. We review the definitions and some useful properties as follows.

**Definition 2.3 (Matroid).** *Given a ground set  $E$ , a family  $\mathcal{M}$  of subsets of  $E$  is a matroid if*

- $\emptyset \in \mathcal{M}$ ;
- If  $I \in \mathcal{M}$  and  $I' \subset I$ , then  $I' \in \mathcal{M}$ ;

- If  $I, I' \in \mathcal{M}$  and  $|I| < |I'|$ , then there must exist an element  $a \in I' \setminus I$  such that  $I \cup \{a\} \in \mathcal{M}$ .

Each  $I \in \mathcal{M}$  is called an independent set. The maximum size of an independent set is called the rank of  $\mathcal{M}$ , denoted by  $\text{rank}(\mathcal{M})$ . Each set  $I \in \mathcal{M}$  of size equal to the rank is called a basis of  $\mathcal{M}$ .

Matroid is a very general combinatorial structure that generalizes many set systems including uniform matroid, partition matroid, laminar matroid, regular matroid, graphic matroid, transversal matroid and so on. Now we define the matroid basis polytope.

**Definition 2.4 (Matroid basis polytope).** Let  $E$  be a ground set and let  $\mathcal{M}$  be a matroid on  $E$ . For each basis  $I \in \mathcal{M}$ , let  $e_I := \sum_{i \in I} e_i$  denote the indicator vector of  $I$ , where  $e_i \in \mathbb{R}^{|E|}$  is the standard  $i$ -th unit vector. The matroid basis polytope  $P_{\mathcal{M}}$  is the convex hull of the set

$$P_{\mathcal{M}} := \{e_I : I \text{ is a basis of } \mathcal{M}\}.$$

The following definition provides another description of  $P_{\mathcal{M}}$  by rank functions.

**Definition 2.5 (Rank function and  $P_{\mathcal{M}}$ ).** Let  $E$  be a ground set of size  $n \geq 1$  and let  $\mathcal{M}$  be a matroid on  $E$ . The rank function  $\text{rank} : 2^E \rightarrow \mathbb{Z}_{\geq 0}$  of  $\mathcal{M}$  is defined as follows: for every  $A \subseteq E$ ,  $\text{rank}(A) = \max_{I \in \mathcal{M}} |I \cap A|$ . Moreover, the matroid basis polytope can be equivalently defined by the following linear program (see e.g., [Sch03]):

$$P_{\mathcal{M}} = \left\{ x \in \mathbb{R}_{\geq 0}^n : \sum_{i \in A} x_i \leq \text{rank}(A), \forall A \in \mathcal{A}; \|x\|_1 = \text{rank}(\mathcal{M}) \right\}.$$

Let  $h \in P_{\mathcal{M}}$  be a point inside the matroid basis polytope. We say a subset  $A \subseteq E$  is tight on  $h$  if  $\sum_{i \in A} h_i = \text{rank}(A)$ .

We also consider a specific type of matroid, called laminar matroid.

**Definition 2.6 (Laminar matroid).** Given a ground set  $E$ , a family of  $\mathcal{A}$  of subsets of  $E$  is called laminar if for every two subsets  $A, B \in \mathcal{A}$  with  $A \cap B \neq \emptyset$ , either  $A \subseteq B$  or  $B \subseteq A$ . Assume  $E \in \mathcal{A}$  and we define the depth of a laminar  $\mathcal{A}$  to be the largest integer  $\ell \geq 1$  such that there exists a sequence of subsets  $A_1, \dots, A_\ell \in \mathcal{A}$  with  $A_1 \subsetneq A_2 \subsetneq \dots \subsetneq A_\ell = E$ .

We say a family  $\mathcal{M}$  of subsets of  $E$  is a laminar matroid if there exists a laminar  $\mathcal{A}$  and a capacity function  $u : \mathcal{A} \rightarrow \mathbb{Z}_{\geq 0}$  such that  $\mathcal{M} = \{I \subseteq E : |I \cap A| \leq u(A), \forall A \in \mathcal{A}\}$ .

By Definition 2.5, we know that  $\text{rank}(A) \leq u(A)$  holds for every  $A \in \mathcal{A}$ . Specifically, we can see that a *uniform matroid* is a laminar matroid of depth 1 (i.e., there is a single cardinality constraint over the entire set  $E$ ) and a *partition matroid* is a laminar matroid of depth 2 (i.e., there is a partition of  $E$  and each partition has a cardinality constraint).

### 3 Coresets for Clustering with General Assignment Constraints

Throughout this paper, we assume there exists an oracle that answers  $d(p, q)$  in  $O(1)$  time for any  $p, q \in \mathcal{X}$ . Our main theorem is as follows.

**Theorem 3.1 (Coresets for  $(k, z)$ -Clustering with general assignment constraints).** Let  $(\mathcal{X}, d)$  be a metric space,  $k \geq 1$  be an integer, and  $z \geq 1$  be constant. Let  $\varepsilon, \delta \in (0, 1)$  and  $\mathcal{B}$  be a structure constraint. There exists a randomized algorithm that given a dataset  $P \subseteq \mathcal{X}$  of size  $n \geq 1$

and an  $O(1)$ -approximate solution  $C^* \in \mathcal{X}^k$  of  $P$  for  $(k, z)$ -CLUSTERING<sup>6</sup>, constructs an  $\varepsilon$ -coreset for  $(k, z)$ -CLUSTERING with general assignment constraints of size

$$2^{O(z \log z)} \cdot \tilde{O}(\text{Lip}(\mathcal{B})^2 \cdot \Lambda_\varepsilon(\mathcal{X}) \cdot k^3 \varepsilon^{-3z}) \cdot \log \delta^{-1},$$

in  $O(nk)$  time with probability at least  $1 - \delta$ , where  $\text{Lip}(\mathcal{B})$  is the Lipschitz constant of  $\mathcal{B}$  defined in Definition 3.7,  $\Lambda_\varepsilon(\mathcal{X})$  is the covering exponent of  $\mathcal{X}$  defined in Definition 3.8, and  $\tilde{O}$  hides a  $\text{poly}(\log(\text{Lip}(\mathcal{B}) \cdot \Lambda_\varepsilon(\mathcal{X}) \cdot k \varepsilon^{-1}))$  term.

For ease of analysis, we assume the given dataset  $P$  is unweighted. This assumption can be removed by a standard process of scaling the point weights to large integers,<sup>7</sup> and treating each weight as a multiplicity of a point; details can be found in [CSS21, Corollary 2.3] and [CLSS22, Section 6.1]. By this theorem, the coreset size is decided by the Lipschitz constant  $\text{Lip}(\mathcal{B})$  and the covering exponent  $\Lambda_\varepsilon(\mathcal{X})$ . If both  $\Lambda_\varepsilon(\mathcal{X})$  and  $\text{Lip}(\mathcal{B})$  are independent of  $n$ , e.g., upper bounded by  $\text{poly}(k, \varepsilon^{-1})$ , our coreset size is at most  $\text{poly}(k, \varepsilon^{-1})$ . We discuss the Lipschitz constant  $\text{Lip}(\mathcal{B})$  in Section 4, and show how to bound  $\Lambda_\varepsilon(\mathcal{X})$  in Section 5.

### 3.1 The Coreset Construction Algorithm

We introduce the algorithm used in Theorem 3.1. The algorithm is almost the same as the coreset algorithm proposed very recently by [BCAJ<sup>+</sup>22].

**Definition 3.2 (Rings).** For a point  $a \in \mathcal{X}$ , and a positive real number  $r > 0$ , define  $\text{ring}(a, r, 2r) := \text{Ball}(a, 2r) \setminus \text{Ball}(a, r)$ .

We need the following decomposition that is defined with respect to a dataset  $Q \subseteq \mathcal{X}$  and a given center  $c \in \mathcal{X}$ .

**Theorem 3.3 (Decomposition into rings and groups, [BCAJ<sup>+</sup>22]).** Let  $Q \subseteq \mathcal{X}$  be a weighted dataset and  $c \in \mathcal{X}$  be a center point. There exists an  $O(|Q|)$  time algorithm  $\text{Decom}(Q, c)$  that outputs  $(\mathcal{R}^*, \mathcal{G}^*)$  as a partition of  $Q$ , where  $\mathcal{R}^*$  and  $\mathcal{G}^*$  are two disjoint collections of sets such that  $Q = (\cup_{R \in \mathcal{R}^*} R) \cup (\cup_{G \in \mathcal{G}^*} G)$ . Moreover,  $\mathcal{R}^*$  is a collection of disjoint rings satisfying

1.  $\forall R \in \mathcal{R}^*$ ,  $R$  is a ring of form  $R = R_i(Q, c)$  for some integer  $i \in \mathbb{Z} \cup \{-\infty\}$ , where  $R_i(Q, c) := Q \cap \text{ring}(c, 2^{i-1}, 2^i)$  for  $i \in \mathbb{Z}$  and  $R_{-\infty}(Q, c) := Q \cap \{c\}$ ;
2.  $|\mathcal{R}^*| \leq 2^{O(z \log z)} \cdot \tilde{O}(k \varepsilon^{-z})$ ;

and  $\mathcal{G}^*$  is a collection of disjoint groups satisfying

1.  $\forall G \in \mathcal{G}^*$ ,  $G$  is the union of a few consecutive rings of  $(Q, c)$  and all these rings are disjoint. Formally,  $\forall G \in \mathcal{G}^*$ , there exists  $l_G, r_G \in \mathbb{Z}^*$ ,  $l_G \leq r_G$  such that  $G = \cup_{i=l_G}^{r_G} R_i(Q, c)$  and the intervals  $\{[l_G, r_G], G \in \mathcal{G}^*\}$  are disjoint;
2.  $\forall G \in \mathcal{G}^*$ ,  $\text{cost}_z(G, c) \leq (\frac{\varepsilon}{6z})^z \cdot \frac{\text{cost}_z(Q, c)}{k \cdot \log(24z\varepsilon^{-1})}$ ;
3.  $|\mathcal{G}^*| \leq 2^{O(z \log z)} \cdot \tilde{O}(k \varepsilon^{-z})$ .

<sup>6</sup>A  $2^{O(z)}$ -approximation may be obtained in  $\tilde{O}(nk)$  time for a general metric space [MP04].

<sup>7</sup>We suppose all weights are rational numbers such that we can round them to integers. If not, we can always replace it with a sufficiently close rational number such that the slight difference does not affect the clustering cost.

Roughly speaking, we decompose  $Q$  into rings w.r.t.  $c$ . The collection  $\mathcal{R}^*$  contains those rings with “heavy” costs, say  $\text{cost}_z(R, c) > \frac{\varepsilon}{6z} \cdot \frac{\text{cost}_z(Q, c)}{k \cdot \log(24z\varepsilon^{-1})}$  for  $R \in \mathcal{R}^*$ .<sup>8</sup> They also gather the remaining “light” rings and form the collection of groups  $\mathcal{G}^*$  (see [BCAJ<sup>+</sup>22, Lemma 3.4]), and ensure that the cardinality  $|\mathcal{G}^*|$  is upper bounded by  $2^{O(z \log z)} \cdot \tilde{O}(k\varepsilon^{-z})$ .

For each group  $G$ , we provide the following data structure.

**Definition 3.4 (Two-point coreset, Line 5 of Algorithm 1 in [BCAJ<sup>+</sup>22]).** *For a weighted dataset/group  $G \subset \mathbb{R}^d$  and a center point  $c \in \mathbb{R}^d$ , let  $p_{\text{far}}^G$  and  $p_{\text{close}}^G$  denote the furthest and closest point to  $c$  in  $G$ . For every  $p \in G$ , compute the unique  $\lambda_p \in [0, 1]$  such that  $d^z(p, c) = \lambda_p \cdot d^z(p_{\text{close}}^G, c) + (1 - \lambda_p) \cdot d^z(p_{\text{far}}^G, c)$ . Let  $D = \{p_{\text{far}}^G, p_{\text{close}}^G\}$ ,  $w_D(p_{\text{close}}^G) = \sum_{p \in G} \lambda_p \cdot w_G(p)$ , and  $w_D(p_{\text{far}}^G) = \sum_{p \in G} (1 - \lambda_p) \cdot w_G(p)$ .  $D$  is called the two-point coreset of  $G$  with respect to  $c$ .*

By definition, we know that  $w(D) = |G|$  and  $\text{cost}_z(D, c) = \text{cost}_z(G, c)$ , which are useful for upper bounding the error induced by such two-point coresets.

---

**Algorithm 1** HUS( $P, k, z, \Gamma$ )

---

**Input:** An unweighted dataset  $P \subseteq \mathcal{X}$  of size  $n \geq 1$ , an integer  $k \geq 1$ , constant  $z \geq 1$ , a sampling size  $\Gamma \geq 1$ , and an  $O(1)$ -approximate solution  $C^* = \{c_1^*, \dots, c_k^*\} \subseteq \mathcal{X}$  of  $P$  for  $(k, z)$ -CLUSTERING

- 1: Decompose  $P$  into  $k$  clusters  $P_1, \dots, P_k$  such that each  $P_i$  contains all points in  $P$  whose closest center in  $C^*$  is  $c_i^*$  (breaking ties arbitrarily).
  - 2: For each  $i \in [k]$ , apply the decomposition of Theorem 3.3 to  $(P_i, c_i^*)$  and obtain  $(\mathcal{R}_i, \mathcal{G}_i) \leftarrow \text{Decom}(P_i, c_i^*)$ , where  $\mathcal{R}_i$  is a collection of disjoint rings and  $\mathcal{G}_i$  is a collection of disjoint groups.
  - 3: For each  $i \in [k]$  and each ring  $R \in \mathcal{R}_i$ , take a uniform sample  $S_R$  of size  $\Gamma$  from  $R$ , and set  $w_{S_R}(x) = \frac{|R|}{\Gamma}$  for each point  $x \in S_R$ .
  - 4: For each  $i \in [k]$ , for each group  $G \in \mathcal{G}_i$  and center  $c_i^*$ , construct a two-point coreset  $D_G$  of  $G$  by Definition 3.4.
  - 5: Return  $S \leftarrow (\bigcup_R S_R) \cup (\bigcup_G D_G)$ .
- 

**Hierarchical Uniform Sampling Coreset Framework** We are now ready to introduce the hierarchical uniform sampling framework HUS( $P, k, z, \Gamma$ ) (Algorithm 1). Note that we apply Theorem 3.3 for each  $P_i$  ( $i \in [k]$ ) in Line 2 of Algorithm 1, and obtain collections  $\mathcal{R}_1, \dots, \mathcal{R}_k$  and  $\mathcal{G}_1, \dots, \mathcal{G}_k$ . Let  $\mathcal{R} = \bigcup_{i \in [k]} \mathcal{R}_i$  be the collection of all rings in different clusters and  $\mathcal{G} = \bigcup_{i \in [k]} \mathcal{G}_i$  be the collection of all groups. By Theorem 3.3, we have the following observation.

**Observation 3.5 (Bound for  $|\mathcal{R}|$  and  $|\mathcal{G}|$ ).**  $|\mathcal{R}|, |\mathcal{G}| \leq 2^{O(z \log z)} \cdot \tilde{O}(k^2 \varepsilon^{-z})$ .

[BCAJ<sup>+</sup>22] showed that by letting  $\Gamma = \tilde{O}(k\varepsilon^{-5})$  we can obtain an  $\varepsilon$ -coreset of size  $\tilde{O}(k^3 \varepsilon^{-6})$  for capacitated  $k$ -MEDIAN in Euclidean spaces  $\mathbb{R}^d$ . We show that the same algorithm HUS( $P, k, z, \Gamma$ ) is surprisingly more powerful – the output  $(S, w)$  is a coreset for  $(k, z)$ -CLUSTERING with general assignment constraints in general metric spaces, using a carefully chosen sample number  $\Gamma$ .

## 3.2 Proof of Theorem 3.1: Analysis of Algorithm 1

**Lipschitz constant  $\text{Lip}(\mathcal{B})$ .** Before we proceed to the proof, we provide the formal definitions of  $\text{Lip}(\mathcal{B})$  and  $\Lambda_\varepsilon(\mathcal{X})$  that appear in the statement of Theorem 3.1. We first define optimal assignment

---

<sup>8</sup>In [BCAJ<sup>+</sup>22], they call these rings heavy rings or marked rings.

transportation (OAT). Since this notion may be of independent interest, we present it in a slightly abstract way and we explain how it connects to our problem after the definition. Then, the key notion  $\text{Lip}(\mathcal{B})$  is defined (in Definition 3.7) as the Lipschitz constant of this OAT procedure.

**Definition 3.6 (Optimal assignment transportation).** *Given a convex body  $\mathcal{B} \subseteq \Delta_k$ , two  $k$ -dimensional vectors  $h, h' \in \mathcal{B}$ , and a function  $\sigma : [n] \times [k] \rightarrow \mathbb{R}_{\geq 0}$  such that  $\|\sigma\|_1 = 1$  and  $\sigma \sim (\mathcal{B}, h)$ , the optimal assignment transportation is defined as the minimum total mass transportation  $\|\sigma - \sigma'\|_1$  from  $\sigma$  to  $\sigma'$ , over functions  $\sigma' : [n] \times [k] \rightarrow \mathbb{R}_{\geq 0}$  such that  $\sigma' \sim (\mathcal{B}, h')$  (see Definition 1.1, 1.2) and  $\forall p \in [n], \|\sigma'(p, \cdot)\|_1 = \|\sigma(p, \cdot)\|_1$ . Namely,*

$$\text{OAT}(\mathcal{B}, h, h', \sigma) := \min_{\substack{\sigma' \sim (\mathcal{B}, h') \\ \forall p \in [n], \|\sigma'(p, \cdot)\|_1 = \|\sigma(p, \cdot)\|_1}} \|\sigma - \sigma'\|_1.$$

To see how our problem is related to Definition 3.6, we view  $\sigma$  as an assignment function, and  $[n]$  and  $[k]$  are interpreted as a data set  $P$  of  $n$  points and a center set  $C$ , respectively. Without loss of generality, we can assume  $w_P(P) = 1$  by normalization, and the requirement  $\|\sigma\|_1 = w_P(P) = 1$  in Definition 3.6 is satisfied. In fact, we choose to use  $[n]$  and  $[k]$  since the representation/identity of a point in the metric does not affect transportation; in other words, OAT is oblivious to the metric space. The requirement of  $h, h' \in \mathcal{B}$  is to ensure the feasibility of OAT. Intuitively, this OAT aims to find a transportation plan, that transports the minimum mass to turn an initial capacity vector  $h$  into a target capacity vector  $h'$ . However, due to the presence of the structure constraint  $\mathcal{B}$ , not all transportation plans are allowed. In particular, we are in addition required to start from a given assignment  $\sigma \in \mathcal{B}$  that is consistent with  $h$ , and the way we reach  $h'$  must be via another assignment  $\sigma' \in \mathcal{B}$  (which we optimize).

**Definition 3.7 (Lipschitz constant of OAT on  $\mathcal{B}$ ).** *Let  $\mathcal{B} \subseteq \Delta_k$  be a convex body. We define the Lipschitz constant of OAT on  $\mathcal{B}$  as*

$$\text{Lip}(\mathcal{B}) := \max_{\substack{h, h' \in \mathcal{B}, \\ \sigma \sim (\mathcal{B}, h): \|\sigma\|_1 = 1}} \frac{\text{OAT}(\mathcal{B}, h, h', \sigma)}{\|h - h'\|_1}.$$

Note that  $\text{Lip}(\mathcal{B}) \geq 1$  since  $\sum_{p \in [n]} (\sigma(p, \cdot) - \sigma'(p, \cdot)) = h - h'$ . It is not hard to see that the Lipschitz constant is scale-invariant, i.e.,  $\text{Lip}(\mathcal{B}) = \text{Lip}(c \cdot \mathcal{B})$  for any  $\mathcal{B}$  and  $c > 0$ .

**Covering Exponent  $\Lambda_\varepsilon(\mathcal{X})$ .** We introduce the notion of covering for all center points  $c \in \mathcal{X}$  in Definition 3.8. Furthermore, we also define a quantity called *covering number* to measure the size/complexity of the covering, and we use a more convenient *covering exponent* to capture the worst-case size of the covering, which can serve as a parameter of the complexity for the metric space.

**Definition 3.8 (Covering, covering number and covering exponent).** *Let  $P \subset \mathcal{X}$  be an unweighted set of  $n \geq 1$  points and  $a \in \mathcal{X}$  be a point. Let  $r_{\max} = \max_{p \in P} d(p, a) \geq 0$  such that  $P \subseteq \text{Ball}(a, r)$ . We say a collection  $\mathcal{C} \subseteq \text{Ball}(a, 24zr_{\max}\varepsilon^{-1})$  of points is an  $\alpha$ -covering of  $P$  if for every point  $c \in \text{Ball}(a, 24zr_{\max}\varepsilon^{-1})$ , there exists some  $c' \in \mathcal{C}$  such that*

$$\max_{p \in P} |d(p, c) - d(p, c')| \leq \frac{\alpha r_{\max}}{12z}.$$

Define  $N_{\mathcal{X}}(P, \alpha)$  to be the minimum cardinality  $|\mathcal{C}|$  of any  $\alpha$ -covering  $\mathcal{C}$  of  $P$ . Define the  $\alpha$ -covering number of  $\mathcal{X}$  to be

$$N_{\mathcal{X}}(n, \alpha) := \max_{P \subseteq \mathcal{X}: |P|=n} N_{\mathcal{X}}(P, \alpha),$$

i.e., the maximum  $N_{\mathcal{X}}(P, \alpha)$  over all possible unweighted sets  $P$  of size  $n$ .

Moreover, we define the  $\alpha$ -covering exponent of the metric space  $(\mathcal{X}, d)$ , denoted by  $\Lambda_{\alpha}(\mathcal{X})$ , to be the least integer  $\gamma \geq 1$  such that  $N_{\mathcal{X}}(n, \alpha) \leq O(n^{\gamma})$  holds for any  $n \geq 2$ .

Roughly speaking, an  $\alpha$ -covering  $\mathcal{C}$  is an discretization of the continuous space of all possible centers within  $\text{Ball}(a, 24zr\varepsilon^{-1})$  of a large radius w.r.t. the  $\ell_{\infty}$ -distance differences from points  $p \in P$  to  $c \in \text{Ball}(a, 24zr\varepsilon^{-1})$ . For those center  $c \notin \text{Ball}(a, 24zr\varepsilon^{-1})$ , we can verify that the distances between every point  $p \in P$  and  $c$  are very close, i.e.,  $d^z(p, c) \in (1 \pm \varepsilon) \cdot d^z(q, c)$  for any two points  $p, q \in P$ . This observation enables us to safely “ignore” the complexity of these remote centers in coresets construction, and hence we only need to consider centers within  $\text{Ball}(a, 24zr\varepsilon^{-1})$ .

Since  $|\mathcal{C}| \leq |\mathcal{X}|$ , we have that the covering exponent  $\Lambda_{\alpha}(\mathcal{X}) \leq \log |\mathcal{X}|$ . This notion is closely related to other combinatorial dimension notions such as the VC-dimension and the (fat) shattering dimension of the set system formed by all metric balls which have been extensively considered in previous works, e.g. [LS10, FL11, FSS20, HJLW18, BJKW21a]. We will show the relations between the covering exponent in our setting and two well-studied dimension notions, the shattering dimension and the doubling dimension, in Section 5.

For preparation, we also need to define another notion of covering (Definition 3.9), which aims to cover both the metric space  $\mathcal{X}$  and the hyper-parameter space of the feasible capacity constraints induced by  $\mathcal{B}$ . A similar idea appeared in [BCAJ<sup>+</sup>22, Lemma 4.4] but it concerns capacity constraints only in Euclidean spaces.

**Definition 3.9 (Coverings and covering numbers with structure constraints).** Let  $\mathcal{B} \subseteq \Delta_k$  be a structure constraint. Let  $P \subset \mathcal{X}$  be an unweighted set of  $n \geq 1$  points and  $a \in \mathcal{X}$  be a point. Let  $r_{\max} = \max_{p \in P} d(p, a) \geq 0$  such that  $P \subseteq \text{Ball}(a, r_{\max})$ . Define  $C_{\text{far}} := \{c \in C : d(c, a) > 24zr_{\max}\varepsilon^{-1}\}$  and for a given capacity constraint  $h \in \mathbb{R}_{\geq 0}^k$ , define

$$\phi(C, h) := \sum_{c \in C_{\text{far}}} h_c \cdot d^z(c, a).^9$$

We say a collection  $\mathcal{F} \subset \mathcal{X}^k \times n\mathcal{B}$  is an  $\alpha$ -covering w.r.t.  $(P, \mathcal{B})$  if for every  $(C, h) \in \mathcal{X}^k \times n\mathcal{B}$ , there exists  $(C', h') \in \mathcal{F}$  such that for every weighted set  $Q \subseteq P$  with  $w_Q(Q) = n$ ,

$$\text{cost}_z(Q, C, \mathcal{B}, h) \in (1 \pm (\alpha + \varepsilon)) \cdot (\text{cost}_z(Q, C', \mathcal{B}, h') + \phi(C, h)) \pm \alpha n r_{\max}^z$$

where notably  $\phi(C, h)$  is independent of the choice of  $Q$ . Define  $N_{\mathcal{X}}(P, \alpha, \mathcal{B})$  to be the minimum cardinality  $|\mathcal{F}|$  of any  $\alpha$ -covering  $\mathcal{F}$  w.r.t.  $(P, \mathcal{B})$ . Define the  $\alpha$ -covering number of  $(\mathcal{X}, \mathcal{B})$  to be

$$N_{\mathcal{X}}(n, \alpha, \mathcal{B}) := \max_{P \subseteq \mathcal{X}: |P|=n} N_{\mathcal{X}}(P, \alpha, \mathcal{B}),$$

i.e., the maximum  $N_{\mathcal{X}}(P, \alpha, \mathcal{B})$  over all possible unweighted sets  $P$  of size  $n$ .

As the covering number  $N_{\mathcal{X}}(n, \alpha, \mathcal{B})$  becomes larger, the pair  $(\mathcal{X}^k, n\mathcal{B})$  is likely to induce more types of  $\text{cost}_z(Q, C, \mathcal{B}, h)$ 's. Although the definition of  $N_{\mathcal{X}}(n, \alpha, \mathcal{B})$  is based on the clustering cost, which looks quite different from Definition 3.8, we have the following lemma that relates the two notions of covering numbers.

---

<sup>9</sup> $\phi(C, h)$  plays the same role as  $\Delta(C)$  defined in [BCAJ<sup>+</sup>22, Lemma 4.4], that captures the clustering cost of points to remote centers in  $C_{\text{far}}$ .

**Lemma 3.10 (Relating two types of covering numbers).** *Let  $\mathcal{B}$  be a structure constraint. For every  $n \geq 2$  and  $\alpha > 0$ , we have*

$$\begin{aligned} \log N_{\mathcal{X}}(n, \alpha, \mathcal{B}) &\leq O(k \cdot \log N_{\mathcal{X}}(n, \alpha)) + zk \cdot \log(\text{Lip}(\mathcal{B}) \cdot z\varepsilon^{-1}\alpha^{-1}) \\ &\leq O(\Lambda_{\alpha}(\mathcal{X}) \cdot k \log n + zk \cdot \log(\text{Lip}(\mathcal{B}) \cdot z\varepsilon^{-1}\alpha^{-1})). \end{aligned}$$

*Proof.* Fix a  $P \subseteq \text{Ball}(a, r_{\max})$  for some  $a \in \mathcal{X}$  and  $r_{\max} = \max_{p \in P} d(p, a)$  of size  $n \geq 1$ . For simplicity, we use  $r$  to represent  $r_{\max}$  in the following. Let  $\mathcal{C}$  denote the  $\alpha$ -covering of  $P$  satisfying Lemma 5.3 and for technical convenience we also add  $a$  into  $\mathcal{C}$ . We show how to construct an  $\alpha$ -covering  $\mathcal{F} \subseteq \mathcal{X}^k \times n\mathcal{B}$  w.r.t.  $(P, \mathcal{B})$  with the desired size and covering property. Let  $\mathcal{N}$  denote an  $l_1$  distance  $\frac{\alpha \cdot \varepsilon^z}{4 \cdot (24z)^z \cdot \text{Lip}(\mathcal{B})}$ -net of  $\mathcal{B}$ , namely, for every  $h \in \mathcal{B}$ , there exists  $h' \in \mathcal{N}$  such that  $\|h - h'\|_1 \leq \frac{\alpha \cdot \varepsilon^z}{4 \cdot (24z)^z \cdot \text{Lip}(\mathcal{B})}$ . Since  $\mathcal{B} \subseteq \Delta_k$ , we know that  $|\mathcal{N}| \leq (\frac{z \cdot \text{Lip}(\mathcal{B})}{\alpha \cdot \varepsilon^z})^{O(k)}$ . We let  $\mathcal{F} := \mathcal{C}^k \times n\mathcal{N}$  to be the Cartesian product of  $\mathcal{C}$  and  $n\mathcal{N}$ . By construction, we have

$$\log |\mathcal{F}| \leq k \log |\mathcal{C}| + \log |\mathcal{N}| \leq O(k \cdot \log N_{\mathcal{X}}(n, \alpha) + zk \cdot \log(\text{Lip}(\mathcal{B}) \cdot z\alpha^{-1}\varepsilon^{-1})).$$

Then it remains to show that  $\mathcal{F}$  is an  $\alpha$ -covering w.r.t.  $(P, \mathcal{B})$ .

Now we fix a center  $C \subseteq \mathcal{X}$  with  $|C| = k$ , and fix a capacity constraint  $h$  on  $C$ . Decompose  $C$  into  $C_{\text{far}} = \{c \in C \mid d(a, c) \geq 24zr/\varepsilon\}$  and  $C_{\text{close}} = C \setminus C_{\text{far}}$ . By Lemma 5.3, there exists a  $C'_{\text{close}} \in \mathcal{C}$  such that  $|C_{\text{close}}| = |C'_{\text{close}}|$  and there exists a bijection  $\pi : C_{\text{close}} \rightarrow C'_{\text{close}}$  such that  $\forall c \in C_{\text{close}}, \forall y \in P, |d(c, y) - d(\pi(c), y)| \leq \frac{\alpha r}{12z}$ . We show that there exists  $(C', h') \in \mathcal{F}$  such that for every  $Q \subseteq P, w_Q(Q) = n$ ,

$$\text{cost}_z(Q, C, \mathcal{B}, h) \in (1 \pm (\alpha + \varepsilon)) \cdot (\text{cost}_z(Q, C', \mathcal{B}, h') + \phi(C, h)) \pm \alpha n r^z. \quad (4)$$

To find such  $(C', h')$ , we first lift  $\pi$  from  $C_{\text{close}}$  to the whole  $C$  by defining  $\forall c \in C_{\text{far}}, \pi(c) = a$ . Then we let  $C' = \pi(C)$ . As  $\mathcal{N}$  is an  $\frac{\alpha \cdot \varepsilon^z}{4 \cdot (24z)^z \cdot \text{Lip}(\mathcal{B})}$ -net of  $\mathcal{B}$ , we know that there exists  $\tilde{h} \in n\mathcal{N}$  such that  $\|\tilde{h} - h\|_1 \leq \frac{\alpha \cdot \varepsilon^z}{4 \cdot (24z)^z \cdot \text{Lip}(\mathcal{B})} \cdot n$ . We define a capacity constraint  $h'$  on  $C' = \pi(C)$  such that  $\forall c \in C, h'(\pi(c)) = \tilde{h}(c)$ .

In the following, we fix a weighted  $Q \subseteq \text{Ball}(a, r)$  with total weight  $n$  and prove (4). Assume  $\sigma$  is the corresponding optimal assignment for  $\text{cost}_z(Q, C, \mathcal{B}, h)$  and expand it as

$$\text{cost}_z(Q, C, \mathcal{B}, h) = \sum_{p \in Q} \left( \sum_{c \in C_{\text{close}}} \sigma(p, c) \cdot d(p, c)^z + \sum_{c \in C_{\text{far}}} \sigma(p, c) \cdot d(p, c)^z \right).$$

Recall that  $C_{\text{far}}$  contains the portion of  $C$  that is at least  $24zr/\varepsilon$  from  $a$  and  $P \subset \text{Ball}(a, r)$ , thus we know that for every  $c \in C_{\text{far}}$  and  $p \in P, d(p, c)^z \in (1 \pm \frac{\varepsilon}{6}) \cdot d(a, c)^z$ . Recall that  $\phi(C, h) = \sum_{c \in C_{\text{far}}} h_c \cdot d(a, c)^z$ . So we have

$$\sum_{c \in C_{\text{far}}} \sigma(p, c) \cdot d(p, c)^z \in (1 \pm \frac{\varepsilon}{6}) \cdot \phi(C, h), \quad (5)$$

and conclude that

$$\text{cost}_z(Q, C, \mathcal{B}, h) \in (1 \pm \frac{\varepsilon}{6}) \cdot \phi(C, h) + \sum_{p \in Q} \sum_{c \in C_{\text{close}}} \sigma(p, c) \cdot d(p, c)^z. \quad (6)$$

So it suffices to prove the following inequality and scale  $\alpha$ ,

$$\text{cost}_z(Q, C', \mathcal{B}, h') \in (1 \pm \alpha) \cdot \sum_{p \in Q} \sum_{c \in C_{\text{close}}} \sigma(p, c) \cdot d(p, c)^z \pm \frac{\varepsilon}{2} \cdot \phi(C, h). \quad (7)$$

We prove the two directions separately. Firstly, let  $\sigma'$  denote an optimal assignment of  $\text{cost}_z(Q, C', \mathcal{B}, h')$ , we prove that

$$\text{cost}_z(Q, C', \mathcal{B}, h') \geq (1 - \alpha) \cdot \sum_{p \in Q} \sum_{c \in C_{\text{close}}} \sigma(p, c) \cdot d(p, c)^z - \frac{\varepsilon}{2} \cdot \phi(C, h) - \alpha n r^z. \quad (8)$$

For the sake of contradiction, suppose (8) does not hold. By the definition of  $\text{Lip}(\mathcal{B})$ , we know that there exists an assignment  $\sigma'' \sim (\mathcal{B}, h)$  such that

$$\sum_{p \in Q} \sum_{c \in C} |\sigma''(p, c) - \sigma'(p, \pi(c))| \leq \frac{\alpha}{4} \cdot \left(\frac{\varepsilon}{24z}\right)^z \cdot n \quad (9)$$

It suffices to show that  $\sigma''$  is actually a better assignment than  $\sigma$  to conclude a contradiction. To see this, we expand  $\text{cost}_z^{\sigma''}(Q, C)$  as

$$\begin{aligned} \text{cost}_z^{\sigma''}(Q, C) &= \sum_{p \in P} \sum_{c \in C_{\text{close}}} \sigma''(p, c) \cdot d(p, c)^z + \sum_{p \in P} \sum_{c \in C_{\text{far}}} \sigma''(p, c) \cdot d(p, c)^z \\ &\leq \sum_{p \in Q} \sum_{c \in C_{\text{close}}} \sigma''(p, c) \cdot d(p, c)^z + \left(1 + \frac{\varepsilon}{6}\right) \cdot \phi(C, h). \end{aligned} \quad (\text{Ineq. (5)}) \quad (10)$$

Moreover, we can see that

$$\begin{aligned} &\sum_{p \in Q} \sum_{c \in C_{\text{close}}} \sigma''(p, c) \cdot d(p, c)^z \\ &= \sum_{p \in Q} \sum_{c \in C_{\text{close}}} \sigma'(p, \pi(c)) \cdot d(p, c)^z + \sum_{p \in Q} \sum_{c \in C_{\text{close}}} |\sigma'(p, \pi(c)) - \sigma''(p, c)| \cdot d(p, c)^z \\ &\leq \sum_{p \in Q} \sum_{c \in C_{\text{close}}} \sigma'(p, \pi(c)) \cdot d(p, c)^z + \|\sigma'' - \sigma'\|_1 \cdot \left(\frac{24zr}{\varepsilon}\right)^z \quad (\text{Defn. of } C_{\text{close}}) \\ &\leq \sum_{p \in Q} \sum_{c \in C_{\text{close}}} \sigma'(p, \pi(c)) \cdot d(p, c)^z + \frac{\alpha}{4} \cdot n r^z. \end{aligned} \quad (\text{Ineq. (9)}) \quad (11)$$

By Lemma 2.1, we know that

$$\begin{aligned} &\sum_{p \in Q} \sum_{c \in C_{\text{close}}} \sigma'(p, \pi(c)) \cdot d(p, c)^z \\ &= \sum_{p \in Q} \sum_{c \in C_{\text{close}}} \sigma'(p, \pi(c)) \cdot (d(p, \pi(c)) + d(p, c) - d(p, \pi(c)))^z \\ &\leq (1 + \alpha) \cdot \sum_{p \in Q} \sum_{c \in C_{\text{close}}} \sigma'(p, \pi(c)) \cdot d(p, \pi(c))^z + \left(\frac{3z}{\alpha}\right)^{z-1} \cdot h'(C_{\text{close}}) \cdot \left(\frac{\alpha r}{12z}\right)^z \quad (\text{Lemma 2.1}) \\ &\leq (1 + \alpha) \cdot \text{cost}_z(Q, C', \mathcal{B}, h') + \frac{\alpha}{4} \cdot n r^z. \end{aligned} \quad (h'(C_{\text{close}}) \leq n)$$

Summing up the above three inequalities, we conclude that

$$\begin{aligned} \text{cost}_z^{\sigma''}(Q, C) &\leq (1 + \alpha) \cdot \text{cost}_z(Q, C', \mathcal{B}, h') + \alpha n r^z + \left(1 + \frac{\varepsilon}{6}\right) \cdot \phi(C, h) \quad (\text{Ineqs. (10) to (12)}) \\ &< \sum_{p \in Q} \sum_{c \in C_{\text{close}}} \sigma(p, c) \cdot d(p, c)^z + \left(1 - \frac{\varepsilon}{3}\right) \cdot \phi(C, h) \quad (\text{by assumption}) \\ &\leq \text{cost}_z^{\sigma}(Q, C), \end{aligned} \quad (\text{Ineq. (6)})$$

which has been a contradiction to the fact that  $\sigma$  is an optimal assignment for  $\text{cost}_z(Q, C, \mathcal{B}, h)$ .

To prove the other direction of (7), it suffices to construct an assignment  $\sigma' \sim (\mathcal{B}, h')$  such that

$$\text{cost}_z^{\sigma'}(Q, C) \leq (1 + \alpha) \sum_{p \in Q} \sum_{c \in C_{\text{close}}} \sigma(p, c) \cdot d(p, c)^z + \frac{\varepsilon}{2} \cdot \phi(C, h) + \alpha n r^z.$$

To this end, we simply choose  $\sigma'$  to be an assignment consistent with  $(\mathcal{B}, h')$  and

$$\sum_{p \in Q} \sum_{c \in C} |\sigma'(p, \pi(c)) - \sigma(p, c)| \leq \frac{\alpha}{4} \cdot \left(\frac{\varepsilon}{24z}\right)^z \cdot n.$$

Again, by the definition of  $\text{Lip}(\mathcal{B})$ , such  $\sigma'$  exists.

Expand  $\text{cost}_z^{\sigma'}(Q, C)$  as

$$\begin{aligned} & \text{cost}_z^{\sigma'}(Q, C) \\ &= \sum_{p \in Q} \sum_{c \in C_{\text{close}}} \sigma'(p, \pi(c)) \cdot d(p, \pi(c))^z + \sum_{p \in Q} \sum_{c \in C_{\text{far}}} \sigma'(p, \pi(c)) \cdot d(p, \pi(c))^z \\ &= \sum_{p \in Q} \sum_{c \in C_{\text{close}}} \sigma'(p, \pi(c)) \cdot d(p, \pi(c))^z + \sum_{p \in Q} \sum_{c \in C_{\text{far}}} \sigma'(p, \pi(c)) \cdot d(p, a)^z \\ &\leq \sum_{p \in Q} \sum_{c \in C_{\text{close}}} \sigma'(p, \pi(c)) \cdot d(p, \pi(c))^z + h'(C_{\text{far}}) \cdot r^z && (Q \subseteq \text{Ball}(a, r)) \\ &\leq \sum_{p \in Q} \sum_{c \in C_{\text{close}}} \sigma'(p, \pi(c)) \cdot d(p, \pi(c))^z + h(C_{\text{far}}) \cdot r^z + \frac{\alpha}{4} \cdot n r^z && (\|h - h'\|_1 \leq \frac{\alpha n}{4}) \\ &\leq \sum_{p \in Q} \sum_{c \in C_{\text{close}}} \sigma'(p, \pi(c)) \cdot d(p, \pi(c))^z + \frac{\varepsilon}{6} \cdot \phi(C, h) + \frac{\alpha}{4} \cdot n r^z. && (\text{Defn. of } \phi(C, h)) \end{aligned}$$

It remains to upper bound  $\sum_{p \in Q} \sum_{c \in C_{\text{close}}} \sigma'(p, \pi(c)) \cdot d(p, \pi(c))^z$ . Using the identical way we upper bound  $\sum_{p \in Q} \sum_{c \in C_{\text{close}}} \sigma''(p, c) \cdot d(p, c)^z$ , we have that

$$\sum_{p \in Q} \sum_{c \in C_{\text{close}}} \sigma'(p, \pi(c)) \cdot d(p, \pi(c))^z \leq (1 + \alpha) \sum_{p \in Q} \sum_{c \in C_{\text{close}}} \sigma(p, c) \cdot d(p, c)^z + \alpha n r^z$$

which finishes the proof.  $\square$

Now we are ready to prove Theorem 3.1. Like [BCAJ+22], we prove the theorem by analyzing groups and rings separately; summarized by the following lemmas.

**Lemma 3.11 (Error analysis for rings).** *For each  $i \in [k]$  and ring  $R \in \mathcal{R}_i$ , suppose  $N_{\mathcal{X}}(R, \varepsilon, \mathcal{B}) \leq O(|R|^\gamma)$  for some integer  $\gamma \geq 1$  and suppose*

$$|S_R| \geq 2^{O(z \log z)} \cdot \text{Lip}(\mathcal{B})^2 \cdot (\gamma + k) \varepsilon^{-2z} \cdot \log(\gamma \cdot k \varepsilon^{-1} \delta^{-1}).$$

*With probability at least  $1 - \frac{\delta}{|\mathcal{R}|}$ , for every  $k$ -center set  $C \in \mathcal{X}^k$  and capacity constraint  $h \in |R| \cdot \mathcal{B}$ ,*

$$|\text{cost}_z(R, C, \mathcal{B}, h) - \text{cost}_z(S_R, C, \mathcal{B}, h)| \leq O(\varepsilon) \cdot (\text{cost}_z(R, C, \mathcal{B}, h) + \text{cost}_z(R, c_i^*)).$$

*Proof.* The proof can be found in Section 3.3.  $\square$

**Lemma 3.12 (Error analysis for groups).** For each  $i \in [k]$ , let  $G[i] = \bigcup_{G \in \mathcal{G}_i} G$  be the union of all groups  $G \in \mathcal{G}_i$  and  $D[i] = \bigcup_{G \in \mathcal{G}_i} D_G$  be the union of all two-point coresets  $D_G$  with  $G \in \mathcal{G}_i$ . For every  $k$ -center set  $C \in \mathcal{X}^k$  and capacity constraint  $h \in |G[i]| \cdot \mathcal{B}$ ,

$$|\text{cost}_z(G[i], C, \mathcal{B}, h) - \text{cost}_z(D[i], C, \mathcal{B}, h)| \leq O(\varepsilon) \cdot (\text{cost}_z(G[i], C, \mathcal{B}, h) + \text{cost}_z(P_i, c_i^*)).$$

*Proof.* The proof can be found in Section 3.4.  $\square$

Note that for groups, the induced error of two-point coresets  $D[i]$  is deterministically upper bounded, which is not surprising since there is no randomness in the construction of  $D[i]$ . This property is quite powerful since we do not need to consider the complexity of center sets in different metric spaces when analyzing the performance of two-point coresets. Now we are ready to prove Theorem 3.1.

*Proof of Theorem 3.1.* By Lemma 3.10, we can apply  $\gamma = O(\Lambda_\varepsilon(\mathcal{X}) \cdot k \cdot \log(\text{Lip}(\mathcal{B}) \cdot zk\varepsilon^{-1}))$  to Lemma 3.11, and verify that the requirement of  $|S_R|$  is

$$|S_R| \geq 2^{O(z \log z)} \cdot \tilde{O}(\text{Lip}(\mathcal{B})^2 \cdot \Lambda_\varepsilon(\mathcal{X}) \cdot k\varepsilon^{-2z}) \cdot \log \delta^{-1},$$

Thus, we can select

$$\Gamma = 2^{O(z \log z)} \cdot \tilde{O}(\text{Lip}(\mathcal{B})^2 \cdot \Lambda_\varepsilon(\mathcal{X}) \cdot k\varepsilon^{-2z}) \cdot \log \delta^{-1}$$

and apply  $\text{HUS}(P, k, z, \Gamma)$  that outputs  $(S, w)$ . Now we verify that  $S$  is the desired  $O(\varepsilon)$ -coreset.

By Observation 3.5, the size  $|S|$  is dominated by

$$\Gamma \cdot |\mathcal{R}| \leq 2^{O(z \log z)} \cdot \tilde{O}(\text{Lip}(\mathcal{B})^2 \cdot \Lambda_\varepsilon(\mathcal{X}) \cdot k^3\varepsilon^{-3z}) \cdot \log \delta^{-1},$$

which matches the coreset size in Theorem 3.1. For the correctness, we can apply the union bound on all rings  $R \in \mathcal{R}$  to Lemma 3.11 and obtain that with probability at least  $1 - \delta$ , for every  $i \in [k]$ , every ring  $R \in \mathcal{R}_i$ , every  $k$ -center set  $C \in \mathcal{X}^k$ , and every capacity constraint  $h \in |R| \cdot \mathcal{B}$ , the following inequality holds:

$$|\text{cost}_z(R, C, \mathcal{B}, h) - \text{cost}_z(S_R, C, \mathcal{B}, h)| \leq O(\varepsilon) \cdot (\text{cost}_z(R, C, \mathcal{B}, h) + \text{cost}_z(R, c_i^*)).$$

Fix a  $k$ -center set  $C \in \mathcal{X}^k$ , and a capacity constraint  $h \in n\mathcal{B}$ . Suppose a collection  $\{h^R \in |R| \cdot \mathcal{B} : R \in \mathcal{R}\} \cup \{h^{(i)} \in |G[i]| \cdot \mathcal{B} : G \in \mathcal{G}\}$  of capacity constraints satisfy that

$$\sum_{R \in \mathcal{R}} h^R + \sum_{i \in [k]} h^{(i)} = h,$$

and

$$\text{cost}_z(P, C, \mathcal{B}, h) = \sum_{R \in \mathcal{R}} \text{cost}_z(R, C, \mathcal{B}, h^R) + \sum_{i \in [k]} \text{cost}_z(G[i], C, \mathcal{B}, h^{(i)}). \quad (13)$$

We have

$$\begin{aligned}
& \text{cost}_z(S, C, \mathcal{B}, h) \\
& \leq \sum_{R \in \mathcal{R}} \text{cost}_z(S_R, C, \mathcal{B}, h^R) + \sum_{i \in [k]} \text{cost}_z(D[i], C, \mathcal{B}, h^{(i)}) \quad (\text{by optimality}) \\
& \leq (1 + O(\varepsilon)) \cdot \sum_{R \in \mathcal{R}} \text{cost}_z(R, C, \mathcal{B}, h^R) + O(\varepsilon) \cdot \sum_{i \in [k], R \in \mathcal{R}_i} \text{cost}_z(R, c_i^*) \\
& \quad + \sum_{i \in [k]} (1 + O(\varepsilon)) \cdot \text{cost}_z(G[i], C, \mathcal{B}, h^{(i)}) + O(\varepsilon) \cdot \text{cost}_z(P_i, c_i^*) \quad (\text{Lemmas 3.11 and 3.12}) \\
& \leq (1 + O(\varepsilon)) \cdot \text{cost}_z(P, C, \mathcal{B}, h) + O(\varepsilon) \cdot \text{cost}_z(P, C^*) \quad (\text{Ineq. (13)}) \\
& \leq (1 + O(\varepsilon)) \cdot \text{cost}_z(P, C, \mathcal{B}, h). \quad (\text{Defn. of } C^*)
\end{aligned}$$

Similarly, we also have that  $\text{cost}_z(P, C, \mathcal{B}, h) \leq (1 + O(\varepsilon)) \cdot \text{cost}_z(S, C, \mathcal{B}, h)$ . Thus,  $(S, w)$  is indeed an  $O(\varepsilon)$ -coreset.

For the running time, Line 1 costs  $O(nk)$  time. Line 2 costs  $\sum_{i \in [k]} O(|P_i|) = O(n)$  time by Theorem 3.3. Line 3 costs  $O(n)$  time. Line 4 costs  $\sum_{G \in \mathcal{G}_i} O(|G|) = O(n)$  time by Definition 3.4. Overall, the total time is  $O(nk)$ .  $\square$

### 3.3 Proof of Lemma 3.11: Error Analysis for Rings

For preparation, we have the following observation that shows that  $\text{cost}_z(P, C, \mathcal{B}, h)$  has some Lipschitz property. The proof is the same as that in [HJLW22, Lemma 3.6].

**Observation 3.13 (Lipschitz property of  $\text{cost}_z(P, C, \mathcal{B}, h)$  on  $P$ ).** *Let  $P, Q \subseteq \mathcal{X}$  be two weighted sets with  $w_P(P) = w_Q(Q)$  and  $c \in \mathcal{X}$  be a center point. For any  $k$ -center set  $C \in \mathcal{X}^k$ , any assignment constraint  $(\mathcal{B}, h)$  and any  $\varepsilon \in (0, 1]$ , we have*

$$|\text{cost}_z(P, C, \mathcal{B}, h) - \text{cost}_z(Q, C, \mathcal{B}, h)| \leq \varepsilon \cdot \text{cost}_z(P, C, \mathcal{B}, h) + \left(\frac{6z}{\varepsilon}\right)^{z-1} \cdot (\text{cost}_z(P, c) + \text{cost}_z(Q, c)).$$

This observation is useful for providing an upper bound for  $|\text{cost}_z(P, C, \mathcal{B}, h) - \text{cost}_z(Q, C, \mathcal{B}, h)|$ . For instance, if  $z = 1$  and  $\text{cost}_1(P, c) + \text{cost}_1(Q, c) \ll \text{cost}_1(P, C, \mathcal{B}, h)$ , we may have

$$|\text{cost}_1(P, C, \mathcal{B}, h) - \text{cost}_1(Q, C, \mathcal{B}, h)| \leq O(\varepsilon) \cdot \text{cost}_1(P, C, \mathcal{B}, h).$$

Now we are ready to prove Lemma 3.11. We first prove the following weak version.

**Lemma 3.14 (A weak version of Lemma 3.11).** *For each  $i \in [k]$  and ring  $R \in \mathcal{R}_i$  of size  $n_R \geq 1$ , suppose*

$$|S_R| \geq 2^{O(z \log z)} \cdot \text{Lip}(\mathcal{B})^2 \cdot \varepsilon^{-2z} \cdot \log(N_{\mathcal{X}}(R, \varepsilon, \mathcal{B}) \cdot 2^k k \varepsilon^{-1} \delta^{-1}).$$

*With probability at least  $1 - \frac{\delta}{|\mathcal{R}|}$ , for every  $k$ -center set  $C \in \mathcal{X}^k$  and capacity constraint  $h \in n_R \cdot \mathcal{B}$ ,*

$$|\text{cost}_z(R, C, \mathcal{B}, h) - \text{cost}_z(S_R, C, \mathcal{B}, h)| \leq O(\varepsilon) \cdot (\text{cost}_z(R, C, \mathcal{B}, h) + \text{cost}_z(R, c_i^*)).$$

Suppose  $N_{\mathcal{X}}(R, \varepsilon, \mathcal{B}) \leq O(n_R^\gamma)$  for some integer  $\gamma \geq 1$ . Combining with Lemma 3.10, we know that the required sample number in Lemma 3.14 is at most

$$2^{O(z \log z)} \cdot \text{Lip}(\mathcal{B})^2 \cdot (\gamma + k) \varepsilon^{-2z} \cdot \log(n_R \cdot k \varepsilon^{-1} \delta^{-1}).$$

Compared to Theorem 3.1, there is an additional term  $\log n_R$  in the coreset size, which can be as large as  $O(\log n)$ . We will show how to remove this term later.

The proof idea of Lemma 3.14 is to first show the concentration property of  $S_R$  w.r.t. a fixed  $k$ -center set  $C \in \mathcal{X}^k$  and a fixed capacity constraint  $h \in n_R \cdot \mathcal{B}$ , and then discretize the parameter space of  $(C, n_R \cdot \mathcal{B})$  and use the union bound to handle all pairs  $(C, h)$ .

We first have the following lemma for a specific pair  $(C, h)$ , which is a generalization of [BCAJ<sup>+</sup>22, Lemma 4.3] based on [CL19, BFS21] by considering general  $z \geq 1$  and introducing structure constraints.

**Lemma 3.15 (Error analysis for rings w.r.t. a pair  $(C, h)$ ).** *Fix  $i \in [k]$  and ring  $R \in \mathcal{R}_i$  with  $R \subseteq \text{ring}(c_i^*, r, 2r)$  for some  $r > 0$  of size  $n_R \geq 1$ . Let  $C \in \mathcal{X}^k$  and  $h \in n_R \cdot \mathcal{B}$ . With probability at least  $1 - \frac{\delta}{N_{\mathcal{X}}(R, \varepsilon, \mathcal{B}) \cdot |\mathcal{R}|}$ ,*

$$|\text{cost}_z(R, C, \mathcal{B}, h) - \text{cost}_z(S_R, C, \mathcal{B}, h)| \leq O(\varepsilon) \cdot (\text{cost}_z(R, C, \mathcal{B}, h) + n_R r^z).$$

*Proof.* Note that  $R \subseteq \text{Ball}(c_i^*, 2r)$  and recall that  $C_{\text{far}} := \{c \in C : d(c, c_i^*) > \frac{48zr}{\varepsilon}\}$  is the collection of remote centers w.r.t.  $\text{Ball}(c_i^*, 2r)$ . Let  $C_{\text{close}} = C \setminus C_{\text{far}}$ . We define a function  $g$  that takes a weighted set  $Q \subseteq R$  with  $w_Q(Q) = n_R$  as input, and outputs

$$\begin{aligned} g(Q) &:= \min_{\sigma: Q \times C \rightarrow \mathbb{R}_{\geq 0}: \sigma \sim (\mathcal{B}, h)} g^\sigma(Q) \\ &= \min_{\sigma: Q \times C \rightarrow \mathbb{R}_{\geq 0}: \sigma \sim (\mathcal{B}, h)} \sum_{p \in Q} \left( \sum_{c \in C_{\text{close}}} \sigma(p, c) \cdot d^z(p, c) + \sum_{c \in C_{\text{far}}} \sigma(p, c) \cdot d^z(c, c_i^*) \right). \end{aligned}$$

Compared to  $\text{cost}_z^\sigma(Q, C)$ , we replace  $d^z(p, c)$  by  $d^z(c, c_i^*)$  for centers  $c \in C_{\text{far}}$  in the definition of  $g(Q)$ . Since for every  $c \in C_{\text{far}}$ , we have  $d^z(p, c) \in (1 \pm \frac{\varepsilon}{6}) \cdot d^z(c, c_i^*)$  by construction. We conclude that  $g(Q) \in (1 \pm \frac{\varepsilon}{6}) \cdot \text{cost}_z(Q, C, \mathcal{B}, h)$ . Thus, it suffices to prove that with probability at least  $1 - \frac{\delta}{N_{\mathcal{X}}(R, \varepsilon, \mathcal{B}) \cdot |\mathcal{R}|}$ ,

$$|g(R) - g(S_R)| \leq O(\varepsilon n_R r^z). \quad (14)$$

Actually, the construction and the following analysis of  $g$  is the key for extending the  $k$ -MEDIAN results in [CL19] to general  $z \geq 1$ . Our idea is to prove the following two lemmas, where the first one shows the concentration property of  $g(S_R)$  and the second one shows the closeness of  $\mathbb{E}_{S_R} [g(S_R)]$  and  $g(R)$ . The main difficulty is handling the additional structure constraint  $\mathcal{B}$ .

**Lemma 3.16 (Concentration of  $g(S_R)$ ).** *With probability at least  $1 - \frac{\delta}{N_{\mathcal{X}}(R, \varepsilon, \mathcal{B}) \cdot |\mathcal{R}|}$ ,*

$$|g(S_R) - \mathbb{E}_{S_R} [g(S_R)]| \leq \varepsilon n_R r^z.$$

**Lemma 3.17 (Closeness of  $\mathbb{E}_{S_R} [g(S_R)]$  and  $g(R)$ ).** *The following holds:*

$$g(R) \leq \mathbb{E}_{S_R} [g(S_R)] \leq g(R) + \varepsilon n_R r^z.$$

Inequality (14) is a direct corollary of the above lemmas since with probability at least  $1 - \frac{\delta}{N_{\mathcal{X}}(R, \varepsilon, \mathcal{B}) \cdot |\mathcal{R}|}$ , we have that

$$|g(R) - g(S_R)| \leq |g(R) - \mathbb{E}_{S_R} [g(S_R)]| + |\mathbb{E}_{S_R} [g(S_R)] - g(S_R)| \leq 2\varepsilon n_R r^z.$$

Hence, it remains to prove these two lemmas.

*Proof of Lemma 3.16.* We first show the following Lipschitz property of  $g$ .

**Claim 3.18 (Lipschitz property of  $g$ ).** *For every two realizations  $S_R$  and  $S'_R$  of size  $\Gamma$  that differ by one sample, we have*

$$|g(S_R) - g(S'_R)| \leq \frac{n_R}{\Gamma} \cdot (62z)^z \varepsilon^{-z+1} r^z.$$

*Proof.* Suppose  $S_R = S \cup \{q\}$  and  $S'_R = S \cup \{q'\}$  where  $S \subseteq R$ ,  $|S| = \Gamma - 1$  and  $q \neq q' \in R$ . Let  $\sigma : S_R \times C \rightarrow \mathbb{R}_{\geq 0}$  with  $\sigma \sim (\mathcal{B}, h)$  be an optimal assignment function such that  $g^\sigma(S_R) = g(S_R)$ . We define  $\sigma' : S'_R \times C \rightarrow \mathbb{R}_{\geq 0}$  with  $\sigma' \sim (\mathcal{B}, h)$  to be:

$$\sigma'(p, \cdot) = \sigma(p, \cdot), \forall p \in S, \text{ and } \sigma'(q', \cdot) = \sigma(q, \cdot).$$

We have

$$\begin{aligned} g(S'_R) &\leq g^{\sigma'}(S'_R) && \text{(by optimality)} \\ &= g^\sigma(S_R) + \sum_{c \in C_{\text{close}}} \sigma(q, c) \cdot (d^z(q', c) - d^z(q, c)) && \text{(Defns. of } \sigma' \text{ and } g) \\ &\leq g^\sigma(S_R) + \sum_{c \in C_{\text{close}}} \sigma(q, c) \cdot \left( \varepsilon \cdot d^z(q, c) + \left(\frac{3z}{\varepsilon}\right)^{z-1} d^z(q, q') \right) && \text{(Lemma 2.1)} \\ &\leq g^\sigma(S_R) + \sum_{c \in C_{\text{close}}} \sigma(q, c) \cdot \left( \varepsilon \cdot (d(c_i^*, c) + d(q, c_i^*))^z + \left(\frac{3z}{\varepsilon}\right)^{z-1} (4r)^z \right) && \text{(triangle ineq.)} \\ &\leq g^\sigma(S_R) + \sum_{c \in C_{\text{close}}} \sigma(q, c) \cdot \left( \varepsilon \cdot \left(\frac{48zr}{\varepsilon} + 2r\right)^z + \left(\frac{3z}{\varepsilon}\right)^{z-1} (4r)^z \right) && \text{(Defn. of } C_{\text{close}}) \\ &\leq g^\sigma(S_R) + \sum_{c \in C_{\text{close}}} \sigma(q, c) \cdot (62z)^z \varepsilon^{-z+1} r^z \\ &\leq g^\sigma(S_R) + \frac{n_R}{\Gamma} \cdot (62z)^z \varepsilon^{-z+1} r^z. && (w_{S_R}(q) = \frac{n_R}{\Gamma}) \end{aligned}$$

By symmetry, we complete the proof.  $\square$

By construction,  $S_R$  consists of  $\Gamma$  i.i.d. uniform samples. Hence, we can apply Theorem 2.2 (McDiarmid's Inequality) and obtain that for every  $t > 0$ ,

$$\Pr [|g(S_R) - \mathbb{E}_{S_R} [g(S_R)]| \geq t] \leq e^{-\frac{2t^2}{\Gamma \cdot \left(\frac{n_R}{\Gamma} \cdot (62z)^z \varepsilon^{-z+1} r^z\right)^2}}.$$

Lemma 3.16 can be verified by letting  $t = \varepsilon n_R r^z$  and the fact that  $\Gamma = \Omega(2^{O(z \log z)} \cdot \varepsilon^{-2z} \cdot \log(N_{\mathcal{X}}(R, \varepsilon, \mathcal{B})) \cdot n_R \cdot \delta^{-1})$ .  $\square$

*Proof of Lemma 3.17.* We first show the easy direction, say  $g(R) \leq \mathbb{E}_{S_R} [g(S_R)]$ , which is guaranteed by the following convexity of  $g$ . For each possible realization of  $S_R$ , let  $\mu_{S_R}$  denote the realized probability of  $S_R$  and  $\sigma_{S_R} : S_R \times C \rightarrow \mathbb{R}_{\geq 0}$  with  $\sigma_{S_R} \sim (\mathcal{B}, h)$  be an optimal assignment function such that  $g^{\sigma_{S_R}}(S_R) = g(S_R)$ . For the convenience of the argument, we extend the domain of  $\sigma_{S_R}$  to be  $\sigma_{S_R} : R \times C \rightarrow \mathbb{R}_{\geq 0}$  where  $\sigma_{S_R}(p, \cdot) = 0$  if  $p \in R \setminus S_R$ . By definition, we know that

$$\mathbb{E}_{S_R} [g(S_R)] = \mathbb{E}_{S_R} [g^{\sigma_{S_R}}(S_R)] = \sum_{S_R} \mu_{S_R} \cdot g(S_R). \quad (15)$$

Now consider an assignment function  $\sigma : R \times C \rightarrow \mathbb{R}_{\geq 0}$  obtained by adding up  $\sigma_{S_R}$ , i.e.,  $\sigma = \mathbb{E}_{S_R}[\sigma_{S_R}] = \sum_{S_R} \mu_{S_R} \cdot \sigma_{S_R}$ . We first show that  $\sigma \sim (\mathcal{B}, h)$  is a feasible assignment function on  $R$ . On one hand, since  $\sigma_{S_R} \sim \mathcal{B}$ , we know that for every  $p \in S_R$ ,  $\sigma_{S_R}(p, \cdot) \in w_{S_R}(p) \cdot \mathcal{B}$ . Thus, we have for every  $p \in R$ ,

$$\sigma(p, \cdot) = \sum_{S_R} \mu_{S_R} \cdot \sigma_{S_R}(p, \cdot) \in \sum_{S_R} \mu_{S_R} \cdot w_{S_R}(p) \cdot \mathcal{B} = \mathcal{B},$$

since  $\mathbb{E}_{S_R}[w_{S_R}(p)] = 1$ . On the other hand, since  $\sigma \sim h$ , it is obvious that  $\sigma \sim h$  since  $\sum_{S_R} \mu_{S_R} = 1$ . Thus, we obtain the following inequality

$$\begin{aligned} g(R) &\leq g^\sigma(R) && \text{(by optimality)} \\ &= \sum_{S_R} \mu_{S_R} \cdot g(S_R) && \text{(by linearity)} \\ &= \mathbb{E}_{S_R}[g(S_R)]. && \text{(Eq. (15))} \end{aligned}$$

Next, we show the difficult direction that  $\mathbb{E}_{S_R}[g(S_R)] \leq (1 + \varepsilon) \cdot g(R) + \varepsilon n_R r^z$ . We first have the following claim showing that the maximum difference between  $g(S_R)$  and  $g(R)$  is bounded.

**Claim 3.19 (Uniform upper bound of  $g(S_R)$ ).** *The following holds*

$$g(S_R) \leq g(R) + (62z)^z \varepsilon^{-z+1} n_R r^z.$$

*Proof.* The proof is almost identical to that of Claim 3.18. The only difference is that  $R$  and  $S_R$  may differ by  $n$  points instead of one point with weight  $\frac{n_R}{\Gamma}$ , which results in a total difference on weights

$$\sum_{p \in R} |w_R(p) - w_{S_R}(p)| \leq 2n_R,$$

where we let  $w_{S_R}(p) = 0$  for  $p \notin S_R$ . □

Let  $\sigma^* : R \times C \rightarrow \mathbb{R}_{\geq 0}$  with  $\sigma^* \sim (\mathcal{B}, h)$  be an optimal assignment function such that  $g^{\sigma^*}(R) = g(R)$ . We first construct  $\pi : S_R \times C \rightarrow \mathbb{R}_{\geq 0}$  as follows: for every  $p \in S_R$ ,

$$\pi(p, \cdot) = \frac{n_R}{\Gamma} \cdot \sigma^*(p, \cdot).$$

Note that  $\pi \sim \mathcal{B}$  always holds, but  $\pi \sim h$  may not hold. We then construct another assignment function  $\pi' : S_R \times C \rightarrow \mathbb{R}_{\geq 0}$  with  $\|\pi'(p, \cdot)\|_1 = \|\pi(p, \cdot)\|_1$  for every  $p \in S_R$  and  $\pi' \sim (\mathcal{B}, h)$  such that the total mass movement  $\sum_{p \in S_R} \|\pi(p, \cdot) - \pi'(p, \cdot)\|_1$  from  $\pi$  to  $\pi'$  is minimized. We have the following claim.

**Claim 3.20 (Upper bound of  $g(S_R)$  W.h.p.).** *With probability at least  $1 - \frac{\varepsilon^z}{2 \cdot (62z)^z}$ , we have*

$$g(S_R) \leq g^{\pi'}(S_R) \leq g(R) + 0.5\varepsilon n_R r^z.$$

*Proof.*  $g(S_R) \leq g^{\pi'}(S_R)$  holds by optimality, and it remains to prove  $g^{\pi'}(S_R) \leq g^{\sigma^*}(R) + 0.5\varepsilon n_R r^z$ .

Since  $g^{\sigma^*}(R) = g(R)$  by definition, note that

$$\begin{aligned}
& g^{\pi'}(S_R) - g^{\sigma^*}(R) \\
&= \sum_{c \in C_{\text{far}}} \left( \sum_{p \in S_R} \pi'(p, c) \cdot d^z(c, c_i^*) - \sum_{p \in R} \sigma^*(p, c) \cdot d^z(c, c_i^*) \right) \\
&+ \sum_{c \in C_{\text{close}}} \left( \sum_{p \in S_R} \pi'(p, c) \cdot d^z(p, c) - \sum_{p \in R} \sigma^*(p, c) \cdot d^z(p, c) \right) \quad (\text{Defn. of } g) \\
&= \sum_{c \in C_{\text{close}}} \left( \sum_{p \in S_R} \pi'(p, c) \cdot d^z(p, c) - \sum_{p \in R} \sigma^*(p, c) \cdot d^z(p, c) \right). \quad (\pi', \sigma^* \sim h)
\end{aligned}$$

Hence, it is equivalent to proving that with probability at least  $1 - \frac{\varepsilon^z}{2 \cdot (62z)^z}$ ,

$$\sum_{c \in C_{\text{close}}} \left( \sum_{p \in S_R} \pi'(p, c) \cdot d^z(p, c) - \sum_{p \in R} \sigma^*(p, c) \cdot d^z(p, c) \right) \leq 0.5 \varepsilon n_R r^z. \quad (16)$$

The idea is to use  $\pi$  as an intermediary and we define another helper function  $\phi$  that takes  $S_R$  as input, and outputs

$$\phi(S_R) := \sum_{c \in C_{\text{close}}} \sum_{p \in S_R} \pi(p, c) \cdot d^z(p, c) + \sum_{c \in C_{\text{close}}} (h_c - \sum_{p \in S_R} \pi(p, c)) \cdot d^z(c, c_i^*).$$

To prove Inequality (16), it suffices to prove that each of the following two inequalities holds with probability at least  $1 - \frac{\varepsilon^z}{4 \cdot (62z)^z}$ .

$$\phi(S_R) - \sum_{c \in C_{\text{close}}} \sum_{p \in R} \sigma^*(p, c) \cdot d^z(p, c) \leq 0.25 \varepsilon n_R r^z, \quad (17)$$

and

$$\sum_{c \in C_{\text{close}}} \sum_{p \in S_R} \pi'(p, c) d^z(p, c) - \phi(S_R) \leq 0.25 \varepsilon n_R r^z. \quad (18)$$

**Proof of Inequality (17)** By linearity and the definition of  $\pi$ , we know that  $\mathbb{E}_{S_R}[\phi(S_R)] = \sum_{c \in C_{\text{close}}} \sum_{p \in R} \sigma^*(p, c) \cdot d^z(p, c)$ . Similar to  $g$ , we show the following Lipschitz of  $\phi$ .

**Claim 3.21 (Lipschitz property of  $\phi$ ).** *For every two realizations  $S_R$  and  $S'_R$  of size  $\Gamma$  that differ by one sample, we have*

$$|\phi(S_R) - \phi(S'_R)| \leq \frac{2n_R}{\Gamma} \cdot (62z)^z \varepsilon^{-z+1} r^z.$$

*Proof.* Suppose the only different points are  $p \in S_R$  and  $q \in S'_R$ . By definition, we know that

$$\begin{aligned}
& |\phi(S_R) - \phi(S'_R)| \\
&= \frac{n_R}{\Gamma} \cdot \left| \sum_{c \in C_{\text{close}}} \sigma^*(p, c) \cdot (d^z(p, c) - d^z(c, c_i^*)) - \sigma^*(q, c) \cdot (d^z(q, c) - d^z(c, c_i^*)) \right| \quad (\text{Defns. of } \phi \text{ and } \pi') \\
&\leq \frac{2n_R}{\Gamma} \cdot (62z)^z \varepsilon^{-z+1} r^z, \quad (\text{Proof of Claim 3.18})
\end{aligned}$$

which completes the proof.  $\square$

Now similar to Lemma 3.16, we can apply Theorem 2.2 (McDiarmid's Inequality) and obtain that for every  $t > 0$ ,

$$\Pr [|\phi(S_R) - \mathbb{E}_{S_R} [\phi(S_R)]| \geq t] = \Pr [|\phi(S_R)| \geq t] \leq e^{-\frac{2t^2}{\Gamma \cdot (\frac{2n_R}{\Gamma} \cdot (62z)^z \varepsilon^{-z+1} r^z)^2}}.$$

Inequality (17) can be verified by letting  $t = 0.25\varepsilon n_R r^z$  and the fact that  $\Gamma = \Omega(2^{O(z \log z)} \cdot \varepsilon^{-2z} \cdot \log(N_{\mathcal{X}}(R, \varepsilon, \mathcal{B}) \cdot n_R \cdot \varepsilon^{-1}))$ .

**Proof of Inequality (18)** For analysis, we construct a function  $\psi$  that takes  $S_R$  as input, and outputs

$$\psi(S_R) := \sum_{c \in C} \left| \sum_{p \in S_R} \pi(p, c) - h_c \right|,$$

i.e., the total capacity difference between  $\pi$  and  $\sigma^*$ . We have the following claim showing that  $\psi(S_R)$  is likely to be small.

**Claim 3.22** ( $\psi(S_R)$  is tiny W.h.p.). *With probability at least  $1 - \frac{\varepsilon^z}{4 \cdot (62z)^z}$ , we have*

$$\psi(S_R) \leq \frac{\varepsilon^z n_R}{4 \cdot \text{Lip}(\mathcal{B}) \cdot (62z)^z}.$$

*Proof.* For every string  $s \in \{+1, -1\}^k$ , we define a function  $\psi_s$  that takes  $S_R$  as input, and outputs

$$\psi_s(S_R) := \sum_{c \in C} s_c \cdot \left( \sum_{p \in S_R} \pi(p, c) - h_c \right).$$

Note that  $\psi(S_R) = \max_{s \in \{+1, -1\}^k} \psi_s(S_R)$ . It is equivalent to proving that with probability at least  $1 - \frac{\varepsilon^z}{4 \cdot (62z)^z}$ , for all  $s \in \{+1, -1\}^k$ ,

$$\psi_s(S_R) \leq \frac{\varepsilon^z n_R}{4 \cdot \text{Lip}(\mathcal{B}) \cdot (62z)^z}. \quad (19)$$

By the union bound, it suffices to prove that for any  $s \in \{+1, -1\}^k$ , Inequality (19) holds with probability at least  $1 - \frac{\varepsilon^z}{2^{k+2} \cdot (62z)^z}$ .

Fix a string  $s \in \{+1, -1\}^k$  and note that  $\mathbb{E}_{S_R} [\psi_s(S_R)] = 0$ . Also noting that for any two realizations  $S_R$  and  $S'_R$  of size  $\Gamma$  that differ by one sample, we have

$$|\psi_s(S_R) - \psi_s(S'_R)| \leq \frac{2n_R}{\Gamma},$$

i.e.,  $\psi_s(S_R)$  is  $\frac{2n_R}{\Gamma}$ -Lipschitz. We again apply Theorem 2.2 (McDiarmid's Inequality) and obtain that for every  $t > 0$ ,

$$\Pr [|\psi_s(S_R) - \mathbb{E}_{S_R} [\psi_s(S_R)]| \geq t] = \Pr [|\psi_s(S_R)| \geq t] \leq e^{-\frac{2t^2}{\Gamma \cdot (2n_R/\Gamma)^2}}.$$

Inequality (19) can be verified by letting  $t = \frac{\varepsilon^z n_R}{4 \cdot \text{Lip}(\mathcal{B}) \cdot (62z)^z}$  and the fact that  $\Gamma = \Omega(2^{O(z \log z)} \cdot \text{Lip}(\mathcal{B})^2 \cdot k \varepsilon^{-2z} \cdot \log(\varepsilon^{-1}))$ , which ensures the success probability to be at least  $1 - \frac{\varepsilon^z}{2^{k+2} \cdot (62z)^z}$ .  $\square$

Let  $h' \in n_R \cdot \mathcal{B}$  be the capacity constraint with  $h'_c = \sum_{p \in S_R} \pi(p, c)$ . We have that  $\pi \sim (\mathcal{B}, h')$  and  $\psi(S_R) = \|h' - h\|_1$ . Let  $\tau = \sum_{p \in S_R} \|\pi(p, \cdot) - \pi'(p, \cdot)\|_1$ . Recall that  $\pi(p, \cdot), \pi'(p, \cdot) \in \frac{n_R}{\Gamma} \cdot \mathcal{B}$  for every  $p \in S_R$ . We have that with probability at least  $1 - \frac{\varepsilon^z}{4 \cdot (62z)^z}$ ,

$$\begin{aligned} \tau &\leq \text{Lip}\left(\frac{n_R}{\Gamma} \cdot \mathcal{B}\right) \cdot \|h' - h\|_1 && \text{(Defns. of Lip}(\mathcal{B}) \text{ and } \pi') \\ &= \text{Lip}(\mathcal{B}) \cdot \psi(S_R) && \text{(scale-invariant of Lip)} \\ &\leq \frac{\varepsilon^z n_R}{4 \cdot (62z)^z}, && \text{(Claim 3.22)} \end{aligned} \tag{20}$$

which implies that

$$\begin{aligned} &\sum_{c \in C_{\text{close}}} \sum_{p \in S_R} \pi'(p, c) \cdot d^z(p, c) - \phi(S_R) \\ &= \sum_{c \in C_{\text{close}}} \sum_{p \in S_R} (\pi'(p, c) - \pi(p, c)) \cdot d^z(p, c) - \sum_{c \in C_{\text{close}}} (h_c - \sum_{c \in C_{\text{close}}} \pi(p, c)) \cdot d^z(c, c_i^*) && \text{(Defn. of } \phi) \\ &= \sum_{c \in C_{\text{close}}} \sum_{p \in S_R} (\pi'(p, c) - \pi(p, c)) \cdot (d^z(p, c) - d^z(c, c_i^*)) && (\pi' \sim h) \\ &\leq \left| \sum_{c \in C_{\text{close}}} \sum_{p \in S_R} (\pi'(p, c) - \pi(p, c)) \right| \cdot \max_{p \in R, c \in C_{\text{close}}} (d^z(p, c) - d^z(c, c_i^*)) \\ &\leq \tau \cdot \max_{p \in R, c \in C_{\text{close}}} (d^z(p, c) - d^z(c, c_i^*)) && \text{(Defn. of } \tau) \\ &\leq \tau \cdot (62z)^z \varepsilon^{-z+1} r^z && \text{(Defn. of } C_{\text{close}}) \\ &\leq 0.25 \varepsilon n_R r^z, \end{aligned}$$

i.e., Inequality (18) holds. Thus, we complete the proof of Claim 3.20.  $\square$

Now we are ready to prove  $\mathbb{E}_{S_R} [g(S_R)] \leq g(R) + \varepsilon n_R r^z$ . By Claims 3.19 and 3.20,

$$\begin{aligned} &\mathbb{E}_{S_R} [g(S_R)] \\ &\leq \frac{\varepsilon^z}{2 \cdot (62z)^z} \cdot (g(R) + (62z)^z \varepsilon^{-z+1} n_R r^z) + \left(1 - \frac{\varepsilon^z}{2 \cdot (62z)^z}\right) (g(R) + 0.5 \varepsilon n_R r^z) \\ &\leq g(R) + \varepsilon n_R r^z. \end{aligned}$$

Thus, we complete the proof of Lemma 3.17.  $\square$

Overall, we complete the proof of Lemma 3.15.  $\square$

Combining with Definition 3.9, we are ready to prove Lemma 3.14.

*Proof of Lemma 3.14.* Suppose  $R \subseteq \text{ring}(c^*, r, 2r) \subseteq \text{Ball}(c^*, 2r)$  for some  $r > 0$ . We recall that  $\phi(C, h) = \sum_{c \in C_{\text{far}}} h_c \cdot d^z(c, c_i^*)$ , where  $C_{\text{far}} := \{c \in C : d(c, c_i^*) > \frac{48zr}{\varepsilon}\}$ . Let  $\mathcal{F} \subset \mathcal{X}^k \times n_R \cdot \mathcal{B}$  be an  $\varepsilon$ -covering of  $(R, \mathcal{B})$  of size at most  $N_{\mathcal{X}}(R, \varepsilon, \mathcal{B})$ . By the union bound, with probability at least  $1 - \frac{1}{\delta |\mathcal{R}|}$ , for every  $(C, h) \in \mathcal{F}$ , the following holds:

$$|\text{cost}_z(R, C, \mathcal{B}, h) - \text{cost}_z(S_R, C, \mathcal{B}, h)| \leq \varepsilon \cdot (\text{cost}_z(R, C, \mathcal{B}, h) + \text{cost}_z(R, c_i^*)). \tag{21}$$

For every  $(C, h) \in \mathcal{X}^k \times n_R \cdot \mathcal{B}$ , let  $(C', h') \in \mathcal{F}$  be a pair such that

$$\text{cost}_z(R, C, \mathcal{B}, h) \in (1 \pm \varepsilon) \cdot (\text{cost}_z(R, C', \mathcal{B}, h') + \phi(C, h)) \pm \varepsilon \cdot n_R r^z,$$

and

$$\begin{aligned} \text{cost}_z(S_R, C, \mathcal{B}, h) &\in (1 \pm \varepsilon) \cdot (\text{cost}_z(S_R, C', \mathcal{B}, h') + \phi(C, h)) \pm \varepsilon n_R r^z \\ &\in (1 \pm O(\varepsilon)) \cdot (\text{cost}_z(R, C', \mathcal{B}, h') + \phi(C, h)) \pm O(\varepsilon) \cdot n_R r^z, \end{aligned} \quad (\text{Ineq. (21)})$$

which implies that

$$\begin{aligned} \text{cost}_z(S_R, C, \mathcal{B}, h) &\in (1 \pm O(\varepsilon)) \cdot \text{cost}_z(R, C, \mathcal{B}, h) \pm O(\varepsilon) \cdot n_R r^z \\ &\in (1 \pm O(\varepsilon)) \cdot \text{cost}_z(R, C, \mathcal{B}, h) \pm O(\varepsilon) \cdot \text{cost}_z(R, c_i^*). \end{aligned} \quad (\text{Defn. of } R)$$

Hence, we have

$$|\text{cost}_z(R, C, \mathcal{B}, h) - \text{cost}_z(S_R, C, \mathcal{B}, h)| \leq O(\varepsilon) \cdot (\text{cost}_z(R, C, \mathcal{B}, h) + \text{cost}_z(R, c_i^*)).$$

□

Finally, we show how to prove Lemma 3.11 by shaving off the term  $\log n_R$  in Lemma 3.14. The main idea is to use the iterative size reduction approach [BJKW21a] that has been widely employed in recent works [CSS21, CLSS22, BCAJ<sup>+</sup>22] to obtain coresets of size independent of  $n_R$ . The technique is somewhat standard and we omit the details.

*Proof of Lemma 3.11.* We can interpret  $S_R$  as an  $L = O(\log^* |R|)$ -steps uniform sampling. Specifically, set  $T = 2^{O(z \log z)} \cdot \text{Lip}(\mathcal{B})^2 \cdot \varepsilon^{-2z} \cdot \log(k\varepsilon^{-1}\delta^{-1})$ , and for  $i = 1, \dots, L$ , let  $S_R^{(i)}$  be a uniform sample of size  $\tilde{O}((\gamma+k)T \cdot (\log^{(i)}(|R|))^3)$  from  $S_R^{(i-1)}$  where  $\log^{(i)}$  denotes the  $i$ -th iterated logarithm and  $S_R^{(0)} = R$ . We remark that  $S_R$  has the same distribution as  $S_R^{(L)}$ .

Identical to the proof of Theorem 3.1 in [BJKW21a], we can obtain that with probability at least  $1 - \frac{\delta}{|\mathcal{R}| \cdot |S_R^{(i-1)}|}$ , for every  $(C, h)$ ,

$$\left| \text{cost}_z(S_R^{(i)}, C, \mathcal{B}, h) - \text{cost}_z(S_R^{(i-1)}, C, \mathcal{B}, h) \right| \leq O\left( \frac{\varepsilon \cdot (\text{cost}_z(R, C, \mathcal{B}, h) + \text{cost}_z(R, c_i^*))}{(\log^{(i)} |R|)^{\frac{1}{2z+2}}} \right). \quad (22)$$

Summing (22) over  $i = 1, 2, \dots, L$  and applying the union bound, we finish the proof. □

### 3.4 Proof of Lemma 3.12: Error Analysis for Groups

Throughout this section, we fix  $i \in [k]$  and a  $k$ -center set  $C \in \mathcal{X}^k$ . We need the following group decomposition as in [BCAJ<sup>+</sup>22].

**Definition 3.23 (Colored and uncolored groups [BCAJ<sup>+</sup>22]).** Fix  $i \in [k]$  and a  $k$ -center set  $C \in \mathcal{X}^k$ . The collection of groups  $\mathcal{G}_i$  can be decomposed into colored groups and uncolored groups w.r.t.  $C$  such that

1. There are at most  $O(k \log(z\varepsilon^{-1}))$  colored groups;
2. For every uncolored group  $G \in \mathcal{G}_i$ , center set  $C$  can be decomposed into two parts  $C = C_{\text{close}}^G \cup C_{\text{far}}^G$  such that
  - For any  $c \in C_{\text{close}}^G$  and  $p \in G$ ,  $d(c, c_i^*) < \frac{\varepsilon}{9z} \cdot d(p, c_i^*)$ ;
  - For any  $c \in C_{\text{far}}^G$  and  $p \in G$ ,  $d(c, c_i^*) > \frac{24z}{\varepsilon} \cdot d(p, c_i^*)$ .

Intuitively, for every uncolored group  $G \in \mathcal{G}_i$ , every center  $c \in C$  is either very “close” or very “far” from  $G$ , which implies the following lemma that is useful for upper bounding the error of  $D_G$ . For simplicity, we omit the superscript  $G$  in the remaining proof for both  $p$  and  $C$  when group  $G$  is fixed.

**Lemma 3.24 (Properties of uncolored groups [BCAJ<sup>+</sup>22, Lemma 3.7]).** *For every uncolored group  $G \in \mathcal{G}_i$ , the following inequalities hold:*

1. For every  $c \in C_{\text{far}}$ ,  $d(p_{\text{far}}, c_i^*) < \frac{\varepsilon}{24z} \cdot d(c, c_i^*)$ ;
2. For every  $p \in G$  and  $c \in C_{\text{far}}$ ,  $d^z(p, c) \in (1 \pm \frac{\varepsilon}{6}) \cdot d^z(c, c_i^*)$ ;
3. For every  $p \in G$  and  $c \in C_{\text{close}}$ ,  $d^z(p, c) \in (1 \pm \frac{\varepsilon}{6}) \cdot d^z(p, c_i^*)$ .

For every colored group  $G \in \mathcal{G}_i$ , we use the following lemma to upper bound the induced error, which is a corollary of Theorem 3.3 and Observation 3.13.

**Lemma 3.25 (Error analysis of colored groups).** *Let  $G \in \mathcal{G}_i$  be a colored group and  $h \in |G| \cdot \mathcal{B}$  be a feasible capacity constraint of  $G$ . The following holds:*

$$|\text{cost}_z(G, C, \mathcal{B}, h) - \text{cost}_z(D_G, C, \mathcal{B}, h)| \leq \varepsilon \cdot \text{cost}_z(G, C, \mathcal{B}, h) + \frac{\varepsilon}{k \log(z\varepsilon^{-1})} \cdot \text{cost}_z(P_i, c_i^*).$$

*Proof.* By Theorem 3.3, we know that  $\text{cost}_z(G, c_i^*) \leq (\frac{\varepsilon}{6z})^z \cdot \frac{\text{cost}_z(P_i, c_i^*)}{k \cdot \log(24z/\varepsilon)}$  since  $G$  is a group. Then by Observation 3.13, we have

$$\begin{aligned} & |\text{cost}_z(G, C, \mathcal{B}, h) - \text{cost}_z(D_G, C, \mathcal{B}, h)| \\ & \leq \varepsilon \cdot \text{cost}_z(G, C, \mathcal{B}, h) + \left(\frac{6z}{\varepsilon}\right)^{z-1} \cdot (\text{cost}(G, c_i^*) + \text{cost}(D_G, c_i^*)) && \text{(Ob. 3.13)} \\ & = \varepsilon \cdot \text{cost}_z(G, C, \mathcal{B}, h) + 2 \cdot \left(\frac{6z}{\varepsilon}\right)^{z-1} \cdot \text{cost}(G, c_i^*) && \text{(Defn. 3.4)} \\ & \leq \varepsilon \cdot \text{cost}_z(G, C, \mathcal{B}, h) + \frac{\varepsilon}{k \log(z\varepsilon^{-1})} \cdot \text{cost}_z(P_i, c_i^*), && \text{(Thm. 3.3)} \end{aligned}$$

which completes the proof. □

Since there are at most  $O(k \log(z\varepsilon^{-1}))$  colored groups, the above lemma provides an upper bound of the error induced by all colored groups  $G \in \mathcal{G}_i$ . It remains to upper bound the error of uncolored groups; summarized by the following lemma, which is a generalization of [BCAJ<sup>+</sup>22, Lemma 3.6] by introducing structure constraints.

**Lemma 3.26 (Error analysis of uncolored groups).** *Let  $G \in \mathcal{G}_i$  denote an uncolored group and  $h \in |G| \cdot \mathcal{B}$  be a feasible capacity constraint of  $G$ . The following holds.*

1. For every  $\sigma : G \times C \rightarrow \mathbb{R}_{\geq 0}$  with  $\sigma \sim (\mathcal{B}, h)$ , there exists  $\pi : D_G \times C \rightarrow \mathbb{R}_{\geq 0}$  with  $\pi \sim (\mathcal{B}, h)$  such that  $\text{cost}_z^\pi(D_G, C) \leq (1 + \varepsilon) \cdot \text{cost}_z^\sigma(G, C)$ ;
2. For every  $\pi : D_G \times C \rightarrow \mathbb{R}_{\geq 0}$  with  $\pi \sim (\mathcal{B}, h)$ , there exists  $\sigma : G \times C \rightarrow \mathbb{R}_{\geq 0}$  with  $\sigma \sim (\mathcal{B}, h)$  such that  $\text{cost}_z^\sigma(G, C) \leq (1 + 3\varepsilon) \cdot \text{cost}_z^\pi(D_G, C)$ .

Moreover, we have

$$\text{cost}_z(D_G, C, \mathcal{B}, h) \in (1 \pm 3\varepsilon) \cdot \text{cost}_z(G, C, \mathcal{B}, h).$$

*Proof.* We discuss two items separately.

**Item 1** Recall that for every  $p \in G$ , there exists a unique  $\lambda_p \in [0, 1]$  such that  $d^z(p, c) = \lambda_p \cdot d^z(p_{\text{close}}, c) + (1 - \lambda_p) \cdot d^z(p_{\text{far}}, c)$ . Then for some  $\sigma : G \times C \rightarrow \mathbb{R}_{\geq 0}$  with  $\sigma \sim (\mathcal{B}, h)$ , we construct  $\pi : D_G \times C \rightarrow \mathbb{R}_{\geq 0}$  as follows: for every  $c \in C$ ,

$$\pi(p_{\text{close}}, c) = \sum_{p \in G} \lambda_p \cdot \sigma(p, c), \text{ and } \pi(p_{\text{far}}, c) = \sum_{p \in G} (1 - \lambda_p) \cdot \sigma(p, c).$$

Note that for every  $\pi(D_G, c) = \sigma(G, c) = h_c$ . Also, since  $\sigma \sim \mathcal{B}$ ,  $\sigma(p, \cdot) \in \mathcal{B}$  holds for every  $p \in G$ . This implies that

$$\pi(p_{\text{close}}, \cdot) \in \sum_{p \in G} \lambda_p \cdot \mathcal{B} = w_{D_G}(p_{\text{close}}) \cdot \mathcal{B}, \text{ and } \pi(p_{\text{far}}, \cdot) \in \sum_{p \in G} (1 - \lambda_p) \cdot \mathcal{B} = w_{D_G}(p_{\text{far}}) \cdot \mathcal{B}.$$

Hence, we have  $\pi \sim (\mathcal{B}, h)$ . Moreover, we have the following claim.

**Claim 3.27 (Restatement of the proof of Item (1) for [BCAJ<sup>+</sup>22, Lemma 3.6]).** *The following holds:*

$$\text{cost}_z^\pi(D_G, C) \leq (1 + \varepsilon) \cdot \text{cost}_z^\sigma(G, C), \text{ and } \text{cost}_z^\sigma(G, C) \leq (1 + \varepsilon) \cdot \text{cost}_z^\pi(D_G, C).^{10}$$

Item 1 is a direct corollary of the above claim.

**Item 2** Consider  $\pi : D_G \times C \rightarrow \mathbb{R}_{\geq 0}$  with  $\pi \sim (\mathcal{B}, h)$ . We construct  $\sigma : G \times C \rightarrow \mathbb{R}_{\geq 0}$  as follows: for every  $p \in G$ , let

$$\sigma(p, \cdot) = \frac{1}{|G|} \cdot h.$$

Since  $h \in |G| \cdot \mathcal{B}$ , we have that  $\sigma(p, \cdot) \in \mathcal{B}$ . Also note that  $\sum_{p \in G} \sigma(p, \cdot) = h$ . Hence, we have  $\sigma \sim (\mathcal{B}, h)$ . It remains to prove that  $\text{cost}_z^\sigma(G, C) \leq (1 + \varepsilon) \cdot \text{cost}_z^\pi(D_G, C)$ . For analysis, we construct another assignment function  $\pi' : D_G \times C \rightarrow \mathbb{R}_{\geq 0}$  as follows:

$$\pi'(p_{\text{close}}, \cdot) = \frac{w_{D_G}(p_{\text{close}})}{|G|} \cdot h, \text{ and } \pi'(p_{\text{far}}, \cdot) = \frac{w_{D_G}(p_{\text{far}})}{|G|} \cdot h.$$

Similarly, we have that  $\pi' \sim (\mathcal{B}, h)$  and can check that  $\pi'(p_{\text{close}}, \cdot) = \sum_{p \in G} \lambda_p \cdot \sigma(p, \cdot)$  and  $\pi'(p_{\text{far}}, \cdot) = \sum_{p \in G} (1 - \lambda_p) \cdot \sigma(p, \cdot)$ . Then by Claim 3.27, we conclude that

$$\text{cost}_z^\sigma(G, C) \leq (1 + \varepsilon) \cdot \text{cost}_z^{\pi'}(D_G, C).$$

It suffices to prove that  $\text{cost}_z^{\pi'}(D_G, C) \leq (1 + \varepsilon) \cdot \text{cost}_z^\pi(D_G, C)$ . Let  $\tau = \sum_{c \in C_{\text{close}}} \pi(p_{\text{far}}, c) - \pi'(p_{\text{far}}, c)$ . Since  $\pi, \pi' \sim h$ , we know that  $\tau = \sum_{c \in C_{\text{close}}} \pi'(p_{\text{close}}, c) - \pi(p_{\text{close}}, c)$  holds. Also, since  $\|\pi(p_{\text{far}}, \cdot)\|_1 = \|\pi'(p_{\text{far}}, \cdot)\|_1$ , we have

$$|\tau| = \left| \sum_{c \in C_{\text{far}}} \pi'(p_{\text{far}}, c) - \pi(p_{\text{far}}, c) \right| \leq \max \left\{ \sum_{c \in C_{\text{far}}} \pi(p_{\text{far}}, c), \sum_{c \in C_{\text{far}}} \pi'(p_{\text{far}}, c) \right\}. \quad (23)$$

<sup>10</sup>Note that Item (1) of [BCAJ<sup>+</sup>22, Lemma 3.6] only presents the first inequality  $\text{cost}_z^\pi(D_G, C) \leq (1 + \varepsilon) \cdot \text{cost}_z^\sigma(G, C)$ . However, the second inequality  $\text{cost}_z^\sigma(G, C) \leq (1 + \varepsilon) \cdot \text{cost}_z^\pi(D_G, C)$  can be directly obtained from their proof.

Now we have

$$\begin{aligned}
& \text{cost}_z^{\pi'}(D_G, C) \\
&= \sum_{c \in C_{\text{close}}} \pi'(p_{\text{close}}, c) \cdot d^z(p_{\text{close}}, c) + \pi'(p_{\text{far}}, c) \cdot d^z(p_{\text{far}}, c) \\
&\quad + \sum_{c \in C_{\text{far}}} \pi'(p_{\text{close}}, c) \cdot d^z(p_{\text{close}}, c) + \pi'(p_{\text{far}}, c) \cdot d^z(p_{\text{far}}, c) \\
&\leq \left(1 + \frac{\varepsilon}{6}\right) \cdot \sum_{c \in C_{\text{close}}} \pi'(p_{\text{close}}, c) \cdot d^z(p_{\text{close}}, c_i^*) + \pi'(p_{\text{far}}, c) \cdot d^z(p_{\text{far}}, c_i^*) \\
&\quad + \left(1 + \frac{\varepsilon}{6}\right) \cdot \sum_{c \in C_{\text{far}}} \pi'(p_{\text{close}}, c) \cdot d^z(c, c_i^*) + \pi'(p_{\text{far}}, c) \cdot d^z(c, c_i^*) \\
&= \left(1 + \frac{\varepsilon}{6}\right) \cdot \left( \left( \sum_{c \in C_{\text{close}}} \pi(p_{\text{close}}, c) + \tau \right) \cdot d^z(p_{\text{close}}, c_i^*) + \left( \sum_{c \in C_{\text{close}}} \pi(p_{\text{far}}, c) - \tau \right) \cdot d^z(p_{\text{far}}, c_i^*) \right) \\
&\quad + \left(1 + \frac{\varepsilon}{6}\right) \cdot \sum_{c \in C_{\text{close}}} \pi(p_{\text{close}}, c) \cdot d^z(p_{\text{close}}, c_i^*) + \pi(p_{\text{far}}, c) \cdot d^z(p_{\text{far}}, c_i^*) \\
&\leq \left(1 + \frac{\varepsilon}{2}\right) \cdot \sum_{c \in C_{\text{close}}} \pi(p_{\text{close}}, c) \cdot d^z(p_{\text{close}}, c) + \pi(p_{\text{far}}, c) \cdot d^z(p_{\text{far}}, c) \\
&\quad + \left(1 + \frac{\varepsilon}{2}\right) \sum_{c \in C_{\text{far}}} \pi(p_{\text{close}}, c) \cdot d^z(p_{\text{close}}, c) + \pi(p_{\text{far}}, c) \cdot d^z(p_{\text{far}}, c) + \left(1 + \frac{\varepsilon}{6}\right) \cdot |\tau| \cdot d^z(p_{\text{far}}, c_i^*) \\
&\leq \left(1 + \frac{\varepsilon}{2}\right) \cdot \text{cost}_z^\pi(D_G, C) + \left(1 + \frac{\varepsilon}{6}\right) \cdot |\tau| \cdot \left(\frac{\varepsilon}{12z}\right)^z \cdot \min\{d^z(p_{\text{close}}, C_{\text{far}}), d^z(p_{\text{far}}, C_{\text{far}})\} \\
&\leq \left(1 + \frac{\varepsilon}{2}\right) \cdot \text{cost}_z^\pi(D_G, C) \\
&\quad + \frac{\varepsilon}{6} \cdot \max\left\{ \sum_{c \in C_{\text{far}}} \pi(p_{\text{far}}, c), \sum_{c \in C_{\text{far}}} \pi'(p_{\text{far}}, c) \right\} \cdot \min\{d^z(p_{\text{close}}, C_{\text{far}}), d^z(p_{\text{far}}, C_{\text{far}})\} \\
&\leq \left(1 + \frac{\varepsilon}{2}\right) \cdot \text{cost}_z^\pi(D_G, C) + \frac{\varepsilon}{6} \cdot \max\left\{ \text{cost}_z^\pi(D_G, C), \text{cost}_z^{\pi'}(D_G, C) \right\},
\end{aligned}$$

where the first equality follows from the definition of  $\pi'$ , the first inequality holds since Lemma 3.24, the second equality follows from the Definition of  $\tau$  and the fact that  $\pi, \pi' \sim h$ , the second and the third inequalities hold due to Lemma 3.24, and the fourth inequality follows from the Inequality (23). This implies that  $\text{cost}_z^{\pi'}(D_G, C) \leq (1 + \varepsilon) \cdot \text{cost}_z^\pi(D_G, C)$  and completes the proof.

Finally,  $\text{cost}_z(D_G, C, \mathcal{B}, h) \in (1 \pm 3\varepsilon) \cdot \text{cost}_z(G, C, \mathcal{B}, h)$  is a direct corollary of the two items. The proof is identical to that of Item (3) for [BCAJ<sup>+</sup>22, Lemma 3.6], and we omit the details here. Overall, we complete the proof of Lemma 3.26.  $\square$

Lemma 3.12 is a direct corollary of Lemmas 3.25 and 3.26. The proof is the same as that of [BCAJ<sup>+</sup>22, Theorem 3.3] (see the paragraph ‘‘Concluding the error analysis’’ in [BCAJ<sup>+</sup>22, Section 3.2] for details).

**Remark 3.28 (Stronger conclusion for two-point coresot).** *Due to the arbitrary selection of  $\pi$  in the proof of Item 2 of Lemma 3.26, we can obtain an even stronger conclusion, say for any  $\pi, \pi' \sim h$ ,*

$$\text{cost}_z^\pi(D_G, C) \in (1 \pm \varepsilon) \cdot \text{cost}_z^{\pi'}(D_G, C).$$

*This property indicates why the two-point coresot is so powerful – any assignment function  $\pi \sim h$  that preserves the center/capacity constraint leads to a close clustering cost  $\text{cost}_z^\pi(D_G, C) \approx$*

$\text{cost}_z(D_G, C, \mathcal{B}, h)$  for every structure constraint  $\mathcal{B}$ . Moreover, for any two structure constraints  $\mathcal{B}, \mathcal{B}'$ , we have

$$\text{cost}_z(D_G, C, \mathcal{B}, h) \in (1 \pm \varepsilon) \cdot \text{cost}_z(D_G, C, \mathcal{B}', h).$$

as long as  $h \in |G| \cdot (\mathcal{B} \cap \mathcal{B}')$ . Hence, two-point coresets may be useful for handling other constraints on the assignment function beyond structure constraints.

## 4 Bounding the Lipschitz Constant $\text{Lip}(\mathcal{B})$

In this section, we give upper and lower bounds for the Lipschitz constant  $\text{Lip}(\mathcal{B})$ , for various notable cases of  $\mathcal{B}$ . In particular, we provide an upper bound  $\text{Lip}(\mathcal{B})$  for an important class of structure constraints  $\mathcal{B}$ , called matroid basis polytopes (Theorems 4.7 and 4.11). We also show that  $\text{Lip}(\mathcal{B})$  may be unbounded with knapsack constraints (Theorem 4.14).

### 4.1 Lipschitz Constant for (Laminar) Matroid Basis Polytopes

For ease of analysis, we first propose another Lipschitz constant on  $\mathcal{B}$  for general polytopes (Definition 4.2). For preparation, we need the following notion of coupling of distributions (see e.g., [MU17]).

**Definition 4.1 (Coupling of distributions).** *Given two distributions  $(\mu, \mu') \in \Delta_m$  ( $m \geq 1$ ),  $\kappa : [m] \times [m] \rightarrow \mathbb{R}_{\geq 0}$  is called a coupling of  $(\mu, \mu')$ , denoted by  $\kappa \vdash (\mu, \mu')$ , if*

- For any  $i \in [m]$ ,  $\sum_{j \in [m]} \kappa(i, j) = \mu_i$ ;
- For any  $j \in [m]$ ,  $\sum_{i \in [m]} \kappa(i, j) = \mu'_j$ .

Now we define the Lipschitz constant for a restricted assignment transportation problem, in which the initial and terminal assignments are restricted to the vertices of polytope  $\mathcal{B}$ .

**Definition 4.2 (Lipschitz constant for restricted OAT).** *Let  $\mathcal{B} \subseteq c \cdot \Delta_k$  for some  $c > 0$  be a polytope. Let  $V$  denote the collection of vertices of  $\mathcal{B}$ . Let  $h, h' \in \mathcal{B}$  be two points inside  $\mathcal{B}$ . Let  $\mu \in \Delta_{|V|}$  denote a distribution on  $V$  satisfying that*

$$\sum_{v \in V} \mu_v \cdot v = h.$$

We define

$$\widetilde{\text{OAT}}(\mathcal{B}, h, h', \mu) := \min_{\substack{\mu' \in \Delta_{|V|} \\ \kappa \vdash (\mu, \mu')}} \sum_{v, v' \in V} \kappa(v, v') \cdot \|v - v'\|_1$$

to be the optimal assignment transportation cost constrained to vertices of  $\mathcal{B}$ . Define the Lipschitz constant for the restricted OAT on  $\mathcal{B}$  to be

$$\widetilde{\text{Lip}}(\mathcal{B}) := \max_{\substack{h, h' \in \mathcal{B} \\ \mu \in \Delta_{|V|} : \sum_{v \in V} \mu_v \cdot v = h}} \frac{\widetilde{\text{OAT}}(\mathcal{B}, h, h', \mu)}{\|h - h'\|_1}.$$

By definition, we know that  $\widetilde{\text{Lip}}(\mathcal{B}) = \widetilde{\text{Lip}}(c \cdot \mathcal{B})$  for any  $c > 0$ , i.e.,  $\widetilde{\text{Lip}}$  is scale-invariant on  $\mathcal{B}$ . We will analyze  $\widetilde{\text{Lip}}(P_{\mathcal{M}})$  for matroid basis polytopes in the later sections. Now, we show the following lemma that connects two Lipschitz constants.

**Lemma 4.3 (Relation between  $\text{Lip}(\mathcal{B})$  and  $\widetilde{\text{Lip}}(\mathcal{B})$ ).** Let  $\mathcal{B} \subseteq \Delta_k$  be a polytope. We have  $\text{Lip}(\mathcal{B}) \leq \widetilde{\text{Lip}}(\mathcal{B})$ .

*Proof.* Let  $V$  denote the collection of vertices of  $\mathcal{B}$ . We define a collection  $\mathcal{H}$  of assignment functions  $\sigma$  such that for every  $\sigma \in \mathcal{H}$ : for every  $p \in [n]$ , there exists a vertex  $v \in V$  such that  $\frac{\sigma(p, \cdot)}{\|\sigma(p, \cdot)\|_1} = v$ , i.e., the assignment vector of each  $p \in [n]$  is equal to a scale of some vertex of  $\mathcal{B}$ . We have the following claim.

**Claim 4.4 (An equivalent formulation of  $\text{Lip}(\mathcal{B})$ ).**  $\text{Lip}(\mathcal{B}) = \max_{\substack{h, h' \in \mathcal{B} \\ \sigma \in \mathcal{H}}} \frac{\text{OAT}(\mathcal{B}, h, h', \sigma)}{\|h - h'\|_1}$ .

*Proof.* Fix  $h, h' \in \mathcal{B}$  and an assignment function  $\sigma \sim (\mathcal{B}, h)$ . It suffices to show the existence of another assignment function  $\pi \in \mathcal{H}$  such that

$$\text{OAT}(\mathcal{B}, h, h', \sigma) \leq \text{OAT}(\mathcal{B}, h, h', \pi). \quad (24)$$

For every  $p \in [n]$ , since  $\frac{\sigma(p, \cdot)}{\|\sigma(p, \cdot)\|_1} \in \mathcal{B}$ , we can rewrite

$$\sigma(p, \cdot) = \|\sigma(p, \cdot)\|_1 \cdot \sum_{v \in V} \alpha_v^p \cdot v$$

for some distribution  $\alpha^p \in \Delta_{|V|}$ . Now we consider a weighted set  $V$  together with weights  $w_V(v) = \sum_{p \in [n]} \|\sigma(p, \cdot)\|_1 \cdot \alpha_v^p$ , and we construct  $\pi : V \times C \rightarrow \mathbb{R}_{\geq 0}$  as follows: for every  $v \in V$ , let  $\pi(v, \cdot) = w_V(v) \cdot v$ . Since

$$\begin{aligned} \sum_{v \in V} \pi(v, \cdot) &= \sum_{v \in V, p \in [n]} \|\sigma(p, \cdot)\|_1 \cdot \alpha_v^p \cdot v && \text{(Defn. of } \pi) \\ &= \sum_{p \in [n]} \sigma(p, \cdot) && \text{(Defn. of } \alpha^p) \\ &= h, && (\sigma \sim h) \end{aligned}$$

we know that  $\pi \sim (\mathcal{B}, h)$ , which implies that  $\pi \in \mathcal{H}$ .

Let  $\pi' : V \times C \rightarrow \mathbb{R}_{\geq 0}$  with  $\pi' \sim (\mathcal{B}, h')$  be an assignment function such that

$$\text{OAT}(\mathcal{B}, h, h', \pi) = \sum_{v \in V} \|\pi(v, \cdot) - \pi'(v, \cdot)\|_1.$$

We construct an assignment function  $\sigma' : [n] \times [k] \rightarrow \mathbb{R}_{\geq 0}$  as follows: for every  $p \in [n]$ , let

$$\sigma'(p, \cdot) = \sum_{v \in V} \frac{\|\sigma(p, \cdot)\|_1 \cdot \alpha_v^p}{w_V(v)} \cdot \pi'(v, \cdot).$$

We first verify that  $\sigma' \sim (\mathcal{B}, h')$ . Note that for every  $p \in [n]$

$$\begin{aligned} \|\sigma'(p, \cdot)\|_1 &= \sum_{v \in V} \frac{\|\sigma(p, \cdot)\|_1 \cdot \alpha_v^p}{w_V(v)} \cdot \|\pi'(v, \cdot)\|_1 && \text{(Defn. of } \sigma') \\ &= \|\sigma(p, \cdot)\|_1 \cdot \sum_{v \in V} \alpha_v^p && \text{(Defn. of } \pi') \\ &= \|\sigma(p, \cdot)\|_1, && (\alpha^p \in \Delta_{|V|}) \end{aligned}$$

which implies that  $\sigma'(p, \cdot) \in \|\sigma(p, \cdot)\|_1 \cdot \mathcal{B}$ . Also, we have

$$\begin{aligned} \sum_{p \in [n]} \sigma'(p, \cdot) &= \sum_{p \in [n], v \in V} \frac{\|\sigma(p, \cdot)\|_1 \cdot \alpha_v^p}{w_V(v)} \cdot \pi'(v, \cdot) && \text{(Defn. of } \pi') \\ &= \sum_{v \in V} \pi'(v, \cdot) && \text{(Defn. of } w_V(v)) \\ &= h', && (\pi' \sim h') \end{aligned}$$

which implies that  $\sigma' \sim h'$ . Thus,  $\sigma' \sim (\mathcal{B}, h')$  holds. Finally, we have

$$\begin{aligned} &\sum_{p \in [n]} \|\sigma(p, \cdot) - \sigma'(p, \cdot)\|_1 \\ &= \sum_{p \in [n]} \|\sigma(p, \cdot)\|_1 \cdot \left\| \sum_{v \in V} \alpha_v^p \cdot v - \frac{\alpha_v^p}{w_V(v)} \cdot \pi'(v, \cdot) \right\|_1 && \text{(Defns. of } \alpha^p \text{ and } \sigma') \\ &\leq \sum_{p \in [n]} \sum_{v \in V} \frac{\|\sigma(p, \cdot)\|_1 \cdot \alpha_v^p}{w_V(v)} \cdot \|w_V(v) \cdot v - \pi'(v, \cdot)\|_1 \\ &\leq \sum_{v \in V} \sum_{p \in [n]} \frac{\|\sigma(p, \cdot)\|_1 \cdot \alpha_v^p}{w_V(v)} \cdot \|\pi(v, \cdot) - \pi'(v, \cdot)\|_1 && \text{(Defn. of } \pi) \\ &= \sum_{v \in V} \|\pi(v, \cdot) - \pi'(v, \cdot)\|_1, && \text{(Defn. of } w_V(v)) \end{aligned}$$

which completes the proof of Claim 4.4.  $\square$

Fix  $h, h' \in \mathcal{B}$  and  $\sigma : [n] \times [k] \rightarrow \mathbb{R}_{\geq 0} \in \mathcal{H}$ . We construct a distribution  $\mu \in \Delta_{|V|}$  as follows: for every  $v \in V$ , let

$$\mu_v = \sum_{p \in [n]: \sigma(p, \cdot) = \|\sigma(p, \cdot)\|_1 \cdot v} \|\sigma(p, \cdot)\|_1.$$

By construction, we know that

$$\begin{aligned} \sum_{v \in V} \mu_v \cdot v &= \sum_{v \in V} \sum_{p \in [n]: \sigma(p, \cdot) = \|\sigma(p, \cdot)\|_1 \cdot v} \|\sigma(p, \cdot)\|_1 \cdot v && \text{(Defn. of } \mu) \\ &= \sum_{p \in [n]} \sigma(p, \cdot) \\ &= h, && (\sigma \sim h). \end{aligned}$$

Let  $\mu' \in \Delta_{|V|}$  with  $\sum_{v \in V} \mu'_v \cdot v = h'$  and  $\kappa \vdash (\mu, \mu')$  satisfy that

$$\widetilde{\text{OAT}}(\mathcal{B}, h, h', \mu) = \sum_{v, v' \in V} \kappa(v, v') \cdot \|v - v'\|_1.$$

Next, we construct another assignment function  $\sigma' \sim (\mathcal{B}, h')$  as follows: for every  $p \in [n]$  with  $\sigma(p, \cdot) = \|\sigma(p, \cdot)\|_1 \cdot v$ , let

$$\sigma'(p, \cdot) = \frac{\|\sigma(p, \cdot)\|_1}{\mu_v} \cdot \sum_{v' \in V} \kappa(v, v') \cdot v'.$$

Note that for every  $p \in [n]$ ,  $\|\sigma'(p, \cdot)\|_1 = \frac{\|\sigma(p, \cdot)\|_1}{\mu_v} \cdot \sum_{v' \in V} \kappa(v, v') = \|\sigma(p, \cdot)\|_1$ , which implies that  $\sigma'(p, \cdot) \in \|\sigma(p, \cdot)\|_1 \cdot \mathcal{B}$ . Also, we have

$$\begin{aligned}
\sum_{p \in [n]} \sigma'(p, \cdot) &= \sum_{v \in V} \sum_{p \in [n]: \sigma(p, \cdot) = \|\sigma(p, \cdot)\|_1 \cdot v} \frac{\|\sigma(p, \cdot)\|_1}{\mu_v} \cdot \sum_{v' \in V} \kappa(v, v') \cdot v' && \text{(Defn. of } \sigma') \\
&= \sum_{v, v' \in V} \kappa(v, v') \cdot v' \\
&= \sum_{v' \in V} \mu_{v'} \cdot v' && (\kappa \vdash (\mu, \mu')) \\
&= h',
\end{aligned}$$

which implies that  $\sigma' \sim (\mathcal{B}, h')$ . Moreover,

$$\begin{aligned}
&\text{OAT}(\mathcal{B}, h, h', \sigma) \\
&\leq \sum_{p \in [n]} \|\sigma(p, \cdot) - \sigma'(p, \cdot)\|_1 && \text{(by optimality)} \\
&= \sum_{v \in V} \sum_{p \in [n]: \sigma(p, \cdot) = \|\sigma(p, \cdot)\|_1 \cdot v} \|\sigma(p, \cdot) - \sigma'(p, \cdot)\|_1 \\
&= \sum_{v \in V} \sum_{p \in [n]: \sigma(p, \cdot) = \|\sigma(p, \cdot)\|_1 \cdot v} \frac{\|\sigma(p, \cdot)\|_1}{\mu_v} \cdot \|\mu_v \cdot v - \sum_{v' \in V} \kappa(v, v') \cdot v'\|_1 && \text{(Defns. of } \sigma \text{ and } \sigma') \quad (25) \\
&= \sum_{v \in V} \|\mu_v \cdot v - \sum_{v' \in V} \kappa(v, v') \cdot v'\|_1 \\
&\leq \sum_{v, v' \in V} \kappa(v, v') \cdot \|v - v'\|_1 \\
&= \widetilde{\text{OAT}}(\mathcal{B}, h, h', \mu), && \text{(Defns. of } \mu' \text{ and } \kappa)
\end{aligned}$$

which implies that

$$\begin{aligned}
\text{Lip}(\mathcal{B}) &= \max_{\substack{h, h' \in \mathcal{B} \\ \sigma \in \mathcal{H}: \|\sigma\|_1 = 1}} \frac{\text{OAT}(\mathcal{B}, h, h', \sigma)}{\|h - h'\|_1} && \text{(Claim 4.4)} \\
&\leq \max_{\substack{h, h' \in \mathcal{B} \\ \mu \in \Delta_{|V|}: \sum_{v \in V} \mu_v \cdot v = h}} \frac{\widetilde{\text{OAT}}(\mathcal{B}, h, h', \mu)}{\|h - h'\|_1} && \text{(Ineq. (25))} \\
&= \widetilde{\text{Lip}}(\mathcal{B}).
\end{aligned}$$

Thus, we complete the proof.  $\square$

#### 4.1.1 Lipschitz Constant for Matroid Basis Polytopes

For preparation, we need the following lemmas that provide well-known properties of matroids; see [Sch03] for more details. The first is an easy consequence of the submodularity of the rank function and the second easily follows from the exchange property of matroid.

**Lemma 4.5 (Properties of rank function).** *Let  $E$  be a ground set and let  $\mathcal{M}$  be a matroid on  $E$  with a rank function  $\text{rank} : 2^E \rightarrow \mathcal{Z}_{\geq 0}$ . Let  $h \in P_{\mathcal{M}}$  be a point inside the matroid basis polytope. Recall that we say a subset  $A \subseteq E$  is tight on  $h$  if  $\sum_{i \in A} h_i = \text{rank}(A)$ . If two subsets  $A, B \subseteq E$  are tight on  $h$ , then both  $A \cup B$  and  $A \cap B$  are tight on  $h$ .*

**Lemma 4.6 (Circuit).** *Let  $E$  be a ground set of size  $n \geq 1$  and let  $\mathcal{M}$  be a matroid on  $E$ . Let  $I \in \mathcal{M}$  be a basis and  $a \in E \setminus I$  be an element. Let circuit  $C(I, a)$  be the smallest tight set on  $e_I$  that contains  $a$ . We have that  $C(I, a) \subseteq I$  and for every element  $b \in C(I, a)$ ,  $I \cup \{a\} \setminus \{b\} \in \mathcal{M}$ .*

We show that  $\text{Lip}(P_{\mathcal{M}}) \leq |E| - 1$  for matroid basis polytopes by the following theorem. This theorem is useful since  $|E| = k$  for every structure constraint  $\mathcal{B}$ , and hence,  $\text{Lip}(\mathcal{B}) \leq k - 1$  when  $\mathcal{B}$  is a scaled matroid basis polytope.

**Theorem 4.7 (Lipschitz constant for matroid basis polytopes).** *Let  $E$  be a ground set and let  $\mathcal{M}$  be a matroid on  $E$ . We have  $\text{Lip}(P_{\mathcal{M}}) \leq \widetilde{\text{Lip}}(P_{\mathcal{M}}) \leq |E| - 1$ .*

*Proof.* Let  $V = \{e_I : I \text{ is a basis of } \mathcal{M}\}$  be the vertex set of  $P_{\mathcal{M}}$ . Fix  $h, h' \in P_{\mathcal{M}}$  and a distribution  $\mu \in \Delta_{|V|}$  with  $\sum_{v \in V} \mu_v \cdot v = h$ . We aim to show that

$$\widetilde{\text{OAT}}(\mathcal{B}, h, h', \mu) = \min_{\substack{\mu' \in \Delta_{|V|} \\ \sum_{v \in V} \mu'_v \cdot v = h' \\ \kappa^{\dagger}(\mu, \mu')}} \sum_{v, v' \in V} \kappa(v, v') \cdot \|v - v'\|_1 \leq (|E| - 1) \cdot \|h - h'\|_1.$$

We first have the following claim.

**Claim 4.8 (Path decomposition from  $h$  to  $h'$ ).** *There exists a sequence  $h^{(0)} = h, h^{(1)}, \dots, h^{(m)} = h' \in P_{\mathcal{M}}$  satisfying*

1. For every  $i \in [m]$ ,  $\|h^{(i-1)} - h^{(i)}\|_0 = 2$ ;
2.  $\sum_{i \in [m]} \|h^{(i-1)} - h^{(i)}\|_1 = \|h - h'\|_1$ .

*Proof.* We first show how to construct  $h^{(1)}$  such that  $\|h^{(0)} - h^{(1)}\|_0 = 2$  and  $\|h^{(0)} - h^{(1)}\|_1 + \|h^{(1)} - h'\|_1 = \|h - h'\|_1$ .

Let  $H^+ = \{i \in E : h_i > h'_i\}$  and  $H^- = \{i \in E : h_i < h'_i\}$ . Consider for every element  $j \in H^-$ , the smallest tight set  $S_j$  on  $h$  that contains  $j$  (inclusion-wise). By Lemma 4.5, we know that such  $S_j$  is unique. If  $S_j \cap H^+ \neq \emptyset$  for some  $j^* \in H^-$ , we can choose an arbitrary element  $i^* \in S_{j^*} \cap H^+$ . Let  $\tau = \min_{A \subseteq E: \sum_{i \in A} h_i < \text{rank}(A)} \{\text{rank}(A) - \sum_{i \in A} h_i\}$ . We construct  $h^{(1)} = h + \tau \cdot (e_{j^*} - e_{i^*})$ . It is easy to see that  $h^{(1)} \in P_{\mathcal{M}}$ , since for every  $S \subseteq E$ ,

1. If  $S$  is not tight on  $h$ , we have  $\sum_{i \in S} h_i^{(1)} \leq \sum_{i \in S} h_i^{(0)} + \tau \leq \text{rank}(S)$  by the definition of  $\tau$ .
2. If  $S$  does not contain  $j^*$ , then  $\sum_{i \in S} h_i^{(1)} \leq \sum_{i \in S} h_i^{(0)} \leq \text{rank}(S)$  by the construction of  $h^{(1)}$ .
3. If  $S$  is tight on  $h$  and  $j^* \in S$ , then  $S$  must contain  $S_{j^*}$  by Lemma 4.5, which makes  $\sum_{i \in S} h_i^{(1)} = \sum_{i \in S} h_i^{(0)} \leq \text{rank}(S)$ .

Now, suppose for every  $j \in H^-$  we have  $S_j \cap H^+ = \emptyset$ . Consider the union  $U$  of all such  $S_j$ s. We can easily see that  $H^- \subseteq U$  and  $U \cap H^+ = \emptyset$ . By Lemma 4.5,  $U$  is also tight on  $h$ , say  $\sum_{i \in U} h_i = \text{rank}(U)$ . Since  $h' \in P_{\mathcal{M}}$ ,  $\sum_{i \in U} h'_i \leq \text{rank}(U)$ . However, by the definition of  $H^-$ , we have

$$\sum_{i \in U} h_i < \sum_{i \in U} h'_i \leq \text{rank}(U),$$

which is a contradiction. Hence, this case is impossible and we can always find  $j \in H^-$  with  $S_j \cap H^+ \neq \emptyset$  and construct  $h^{(1)} = h + \tau \cdot (e_{j^*} - e_{i^*})$ .

We then can repeat the above procedure and construct a sequence  $h^{(1)}, h^{(2)}, \dots$ . By the previous argument, at each iteration  $t$  we increase the number of tight sets on  $h^{(t)}$  by at least one from  $h^{(t-1)}$ . Since there is a finite number of tight sets (at most  $2^{|E|}$ ), the above process terminates in a finite number of times and arrives  $h'$ . Hence, we complete the proof of Claim 4.8.  $\square$

Suppose  $\widetilde{\text{OAT}}(\mathcal{B}, h, h', \mu) \leq (|E| - 1) \cdot \|h - h'\|_1$  holds for the case of  $\|h - h'\|_0 = 2$ . Let  $\mu^{(0)} = \mu$ . For every  $i \in [m]$ , we consecutively construct a distribution  $\mu^{(i)} \in \Delta_{|V|}$  with  $\sum_{v \in V} \mu_v^{(i)} \cdot v = h^{(i)}$  and a coupling  $\kappa^{(i)} \vdash (\mu^{(i-1)}, \mu^{(i)})$  such that

$$\widetilde{\text{OAT}}(\mathcal{B}, h^{(i-1)}, h^{(i)}, \mu^{(i-1)}) = \sum_{v, v' \in V} \kappa^{(i)}(v, v') \cdot \|v - v'\|_1 \leq (|E| - 1) \cdot \|h^{(i-1)} - h^{(i)}\|_1. \quad (26)$$

Then we have

$$\begin{aligned} \widetilde{\text{OAT}}(P_{\mathcal{M}}, h, h', \mu) &= \min_{\substack{\mu' \in \Delta_{|V|} : \sum_{v \in V} \mu'_v \cdot v = h' \\ \kappa \vdash (\mu, \mu')}} \sum_{v, v' \in V} \kappa(v, v') \cdot \|v - v'\|_1 \\ &\leq \sum_{i \in [m]} \sum_{v, v' \in V} \kappa^{(i)}(v, v') \cdot \|v - v'\|_1 && \text{(Defn. of } \kappa^{(i)}) \\ &\leq (|E| - 1) \cdot \sum_{i \in [m]} \|h^{(i-1)} - h^{(i)}\|_1 && \text{(Ineq. (26))} \\ &= (|E| - 1) \cdot \|h - h'\|_1. && \text{(Claim 4.8)} \end{aligned}$$

Thus, we only prove for the case that  $h$  and  $h'$  differ in exactly two entries  $\|h - h'\|_0 = 2$ . We define the following augmenting path.

**Definition 4.9 (Augmenting path from  $h$  to  $h'$ ).** *Let  $h$  and  $h'$  be two vectors in  $\mathcal{B}$  that differ in exactly two entries,  $\|h - h'\|_0 = 2$ . W.l.o.g., we assume the corresponding two elements are  $s, t \in E$  with  $h_s < h'_s$  and  $h_t > h'_t$  respectively. Let  $\mu \in \Delta_{|V|}$  be the distribution associated with  $h$  (i.e.,  $\sum_{v \in V} \mu_v \cdot v = h$ ). We say a sequence of indicator vector of basis  $(v_1 = e_{I_1}, \dots, v_m = e_{I_m})$  together with a sequence of indices of elements  $(a_0 = s, a_1, \dots, a_{m-1}, a_m = t)$  ( $a_i \in E$  for  $0 \leq i \leq m$ ) form a weak augmenting path of length  $m \geq 1$  from  $h$  to  $h'$  if*

1. For every  $i \in [m]$ ,  $\mu_{v_i} > 0$  (i.e., every  $v_i$  appears in the support of  $\mu$ );
2. For every  $i \in [m]$ ,  $I_i \cup \{a_{i-1}\} \setminus \{a_i\} \in \mathcal{M}$  (weak exchange property).

Here, we do not require that  $I_i$ s and  $a_i$ s be distinct.

Moreover, we say  $(v_1 = e_{I_1}, \dots, v_m = e_{I_m})$  and  $(a_0 = s, a_1, \dots, a_{m-1}, a_m = t)$  form a strong augmenting path if it is a weak augmenting path with the following additional properties:

1. All elements  $a_i$  are distinct;
2. For every basis  $I \in \mathcal{M}$ , letting  $A_I = \{i \in [m] : I_i = I\}$ , we have

$$\widehat{I} = I \cup \{a_{i-1} : i \in A_I\} \setminus \{a_i : i \in A_I\} \in \mathcal{M}. \quad \text{(strong exchange property)}$$

Intuitively, given a weak augmenting path, we can perform a sequence of exchange of basis according to the weak exchange property. As a result, we get a sequence of new basis so that the mass in  $a_0 = s$  is transported to  $a_m = t$ . However, there is a subtle technical problem that one basis may be used several times and we need to guarantee it is still a basis after all exchange operations, which motivates the notion of strong exchange property. The following crucial lemma shows that a strong augmenting path from  $h$  to  $h'$  exists and its length can be bounded.

**Lemma 4.10 (Existence of a strong augmenting path of length  $\leq |E| - 1$ ).** *Given  $h$  and  $h'$  in  $\mathcal{B}$  that differ in exactly two entries, for any  $\mu \in \Delta_{|V|}$  with  $\sum_{v \in V} \mu_v \cdot v = h$ , there exists a strong augmenting path of length at most  $|E| - 1$  from  $h$  to  $h'$ .*

*Proof.* For preparation, we construct a (multi-edge) directed graph  $G_{\text{ex}}$ , called the *exchange graph*.  $G_{\text{ex}}$  has vertex set  $E$  and the following set of edges: for every basis  $I \in \mathcal{M}$  with  $\mu_{e_I} > 0$ , every element  $a \in E \setminus I$  and every element  $b \in C(I, a)$ , add a directed edge  $(a, b)$  with a certificate  $C(I, a)$  to the graph  $G_{\text{ex}}$ .

**Existence of A Weak Augmenting Path** We first prove the existence of a weak augmenting path from  $h$  to  $h'$ , which is equivalent to proving that there exists a path on  $G_{\text{ex}}$  from vertex  $s$  to vertex  $t$ . By contradiction assume that there does not exist such a path. Let  $E^-$  be the collection of elements  $i \in E$  such that there exists a path from  $s$  to  $i$ , and let  $E^+ = E \setminus E^-$ . We have that  $t \in E^+$ . Since  $\sum_{i \in E^-} h_i < \sum_{i \in E^-} h'_i \leq \text{rank}(E^-)$ , we know that  $E^-$  is not a tight set on  $h$ . Consequently, there must exist a basis  $I \in \mathcal{M}$  with  $\mu_{e_I} > 0$  such that  $|I \cap E^-| < \text{rank}(E^-)$ . Let  $a \in E^- \setminus I$  be an element such that  $(I \cap E^-) \cup \{a\} \in \mathcal{M}$ . If  $C(I, a) \subseteq E^-$ , we have  $C(I, a) \cup \{a\} \notin \mathcal{M}$ . However,  $C(I, a) \cup \{a\} \subseteq (I \cap E^-) \cup \{a\} \in \mathcal{M}$ , which is a contradiction. Hence, we have that  $C(I, a) \cap E^+ \neq \emptyset$  and there exists an edge from  $a \in E^-$  to some element in  $E^+$ , which contradicts the definition of  $E^-$ .

**Existence of A Strong Augmenting Path of Length  $\leq |E| - 1$**  Among all weak augmenting path from  $h$  to  $h'$ , we select a shortest one, say  $(v_1 = e_{I_1}, \dots, v_m = e_{I_m})$  and  $(a_0 = s, a_1, \dots, a_{m-1}, a_m = t)$ . We claim that  $a_0 \neq a_1 \neq \dots \neq a_m$ . Assume that  $a_i = a_j$  for some  $0 \leq i < j \leq m$ , we can see that  $(e_{I_1}, \dots, e_{I_i}, e_{I_{j+1}}, \dots, e_{I_m})$  and  $(a_0 = s, a_1, \dots, a_i, a_{j+1}, \dots, a_{m-1}, a_m = t)$  form a shorter weak augmenting path, which is a contradiction. Since there are at most  $|E|$  different elements, we have that  $m \leq |E| - 1$ .

Now we prove that  $(v_1 = e_{I_1}, \dots, v_m = e_{I_m})$  and  $(a_0 = s, a_1, \dots, a_{m-1}, a_m = t)$  already form a strong augmenting path. Note that for every  $i \in [m]$ ,  $(a_{i-1}, a_i)$  is a direct edge on  $G_{\text{ex}}$  with a certificate  $C(I_i, a_i)$ . It is easy to verify this when all  $v_1, \dots, v_m$  are distinct (i.e., each edge  $(a_{i-1}, a_i)$  is defined by a distinct basis).

Now, we consider the more difficult case where  $v_i$ s are not all distinct. Observe  $v_i \neq v_{i+1}$  (i.e., two adjacent edges correspond to two different bases), since  $a_i \in I_i$  and  $a_i \notin I_{i+1}$  which implies  $I_i \neq I_{i+1}$ . Fix a basis  $I \in \mathcal{M}$  with  $A_I = \{i_l \in [m] : t \in [T], I_{i_l} = I\}$  and  $|A_I| \geq 2$ . W.l.o.g., assume  $0 \leq i_1 < i_2 < \dots < i_T \leq m$  and we have  $i_{l+1} - i_l \geq 2$  for  $l \in [T-1]$ . We first note that for any  $i < j \in A_I$ ,  $a_j \notin C(I, a_{i-1})$ . Suppose, for contradiction that  $a_j \in C(I, a_{i-1})$ . Based on the construction of graph  $G_{\text{ex}}$ ,  $(a_{i-1}, a_j)$  is also an edge with certificate  $C(I, a_j)$ . Consequently,  $(e_{I_1}, \dots, e_{I_i}, e_{I_{j+1}}, \dots, e_{I_m})$  and  $(a_0 = s, a_1, \dots, a_{i-1}, a_j, a_{j+1}, \dots, a_{m-1}, a_m = t)$  form a shorter weak augmenting path, which is a contradiction.

Hence, we have that  $a_j \notin C(I, a_{i-1})$  holds for any  $i < j \in A_I$ . We first swap  $a_{i_{T-1}}$  and  $a_{i_T}$  and obtain a basis  $I^{(T)} = I \cup \{a_{i_{T-1}}\} \setminus \{a_{i_T}\}$ . The circuit  $C(I, a_{i_{T-1}-1})$  is completely contained in  $I^{(T)}$  since  $a_{i_T} \notin C(I, a_{i_{T-1}-1})$ , and hence we can continuously perform the swap between  $a_{i_{T-1}-1}$  and  $a_{i_{T-1}}$ . By reduction, we have that  $I^{(l)} = I \cup \{a_{i_{l-1}}, \dots, a_{i_{T-1}}\} \setminus \{a_{i_l}, \dots, a_{i_T}\}$  for every  $l \in [T]$  is still a basis. Hence,  $\hat{I} = I^{(1)}$  is a basis, which completes the proof.  $\square$

We consider the following procedure:

1. Find a strong augmenting path of length  $m \leq |E| - 1$  from  $h$  to  $h'$ , say  $(v_1 = e_{I_1}, \dots, v_m = e_{I_m})$  and  $(a_0 = s, a_1, \dots, a_{m-1}, a_m = t)$ .
2. Let  $\tau = \min \{h'_1 - h_1, \min_{i \in [m]} \mu_{v_i}\}$ . For every basis  $I \in \mathcal{M}$ , let  $A_I := \{i \in [m] : I_i = I\}$  and let  $\hat{I} = I \cup \{a_i : i \in A_I\} \setminus \{a_{i-1} : i \in A_I\}$ .

3. Construct  $\mu'' \in \Delta_{|V|}$  as the resulting distribution of the following procedure: for every  $I \in \mathcal{M}$  with  $A_I \neq \emptyset$ , reduce  $\mu_{e_I}$  by  $\tau$  and increase  $\mu_{e_{\bar{I}}}$  by  $\tau$ .
4. Let  $h'' = \sum_{v \in V} \mu''_v \cdot v$ . If  $h'' = h'$ , we are done. Otherwise, let  $h \leftarrow h''$  and  $\mu \leftarrow \mu''$ , and iteratively run the above steps for tuple  $(h, h', \mu)$ .

After running an iteration, we can easily verify the following

1.  $\|h'' - h'\|_0 \leq 2$ ,  $h''_1 - h_1 = \tau$  and  $h_2 - h''_2 = \tau$ .
2.  $\|h'' - h'\|_1 = \|h - h'\|_1 - 2\tau$ .
3. The total assignment transportation from  $\mu$  to  $\mu''$ , say  $\min_{\kappa \vdash (\mu, \mu'')} \sum_{v, v' \in V} \kappa(v, v') \cdot \|v - v'\|_1$ , is at most  $\sum_{I \in \mathcal{M}: |I| = \text{rank}(\mathcal{M})} \tau \cdot \|e_I - e_{\bar{I}}\|_1 = 2m\tau \leq 2(|E| - 1)\tau$ .

It means that we can reduce the value  $\|h - h'\|_1$  by  $2\tau$ , by introducing at most  $2(|E| - 1)\tau$  assignment transportation for  $\widetilde{\text{OAT}}(\mathcal{B}, h, h', \mu)$ . Hence, the required assignment transportation from  $h$  to  $h'$  is at most  $(|E| - 1) \cdot \|h - h'\|_1$ , which implies that

$$\widetilde{\text{OAT}}(P_{\mathcal{M}}, h, h', \mu) \leq (|E| - 1) \cdot \|h - h'\|_1.$$

Due to the arbitrary selection of  $h, h'$  and  $\mu$ , we complete the proof of Theorem 4.11.  $\square$

#### 4.1.2 Lipschitz Constant for Laminar Matroid Basis Polytopes

Our main theorem for laminar matroid basis polytopes is as follows.

**Theorem 4.11 (Lipschitz constant for laminar matroid basis polytopes).** *Let  $E$  be a ground set and let  $\mathcal{M}$  be a laminar matroid on  $E$  of depth  $\ell \geq 1$ . We have  $\text{Lip}(P_{\mathcal{M}}) \leq \widetilde{\text{Lip}}(P_{\mathcal{M}}) \leq \ell + 1$ .*

*Proof.* Let  $V = \{e_I : I \text{ is a basis of } \mathcal{M}\}$  be the vertex set of  $P_{\mathcal{M}}$ . Let  $\mathcal{A}$  be the corresponding laminar and  $u$  be the corresponding capacity function of  $\mathcal{M}$ . W.l.o.g., we assume  $u(A) = \text{rank}(A)$  for every  $A \in \mathcal{M}$ . Let  $h, h' \in P_{\mathcal{M}}$  and  $\mu \in \Delta_{|V|}$  be a distribution on  $V$  satisfying that

$$\sum_{v \in V} \mu_v \cdot v = h.$$

By the proof of Theorem 4.7, we only need to prove for the case that  $\|h - h'\|_0 = 2$ . W.l.o.g., we assume  $h_1 < h'_1$  and  $h_2 > h'_2$ . Again, by the same argument as in Theorem 4.7, it suffices to prove the following claim.

**Lemma 4.12 (Existence of a strong augmenting path of length  $\leq \ell + 1$  for laminar matroids).** *There exists a strong augmenting path of length at most  $\ell + 1$  from  $h$  to  $h'$ .*

*Proof.* We again construct the exchange graph  $G_{\text{ex}}$  as in the proof of Lemma 4.10, and iteratively construct a strong augmenting path from  $h$  to  $h'$ . Also recall that we assume  $h_s < h'_s$  and  $h_t > h'_t$  for some  $s, t \in E$ . Let  $A^* \in \mathcal{A}$  be the minimal set that contains  $\{s, t\}$ . Firstly, we find a vector  $v_1 = e_{I_1}$  with  $\mu_{v_1} > 0$  and  $a_0 = s \notin I_1$ . Let  $A_1 \in \mathcal{A}$  be the minimal set with  $a_0 \in A_1$  and  $|I_1 \cap A_1| = u(A_1)$ . Note that such  $A_1$  must exist, since  $I_1 \in P_{\mathcal{M}}$  and we have  $|I_1 \cap E| = \text{rank}(\mathcal{M}) = u(E)$ . We discuss the following cases.

**Case 1:**  $A_1 \supseteq A^*$  Since  $h_t > h'_t \geq 0$ , there must exist a vertex  $v_2 = e_{I_2}$  with  $\mu_{v_2} > 0$  and  $a_2 = t \in I_2$ . Note that  $|(I_2 \setminus \{a_2\}) \cap A_1| < |I_1 \cap A_1| = u(A_1)$ . There must exist an element  $a_1 \in I_1 \cap A_1$  such that  $(I_2 \cap A_1) \cup \{a_1\} \setminus \{a_2\} \in \mathcal{M}$ . Due to the laminar structure, we know that  $I_2 \cup \{a_1\} \setminus \{a_2\} \in \mathcal{M}$  also holds. Also, since  $(I_1 \cap A) \setminus \cup \{a_0\} \setminus \{a_1\} \in \mathcal{M}$ , we conclude that  $I_1 \cup \{a_0\} \setminus \{a_1\} \in \mathcal{M}$ . Consequently,  $(v_1, v_2)$  and  $(a_0, a_1, a_2)$  form an augmenting path of length 2 from  $h$  to  $h'$ .

**Case 2:**  $A_1 \subsetneq A^*$  We know that  $\sum_{i \in A_1} h'_i - h_i = h'_s - h_s > 0$ . Thus, we must have

$$\begin{aligned} \sum_{e_I \in V} \mu_{e_I} \cdot |I \cap A_1| &= \sum_{i \in A_1} h_i && \text{(Defn. of } \mu) \\ &< \sum_{i \in A_1} h'_i \\ &\leq u(A_1), && (h' \in P_{\mathcal{M}}) \end{aligned}$$

which implies the existence of a vertex  $v_2 = e_{I_2} \in V$  with  $\mu_{v_2} > 0$  and  $|I_2 \cap A_1| < u(A_1)$ . Since  $u(A) = |I_1 \cap A_1| > |I_2 \cap A_1|$ , there must exist an element  $a_1 \in I_1 \cap A_1$  such that  $(I_2 \cap A_1) \cup \{a_1\} \in \mathcal{M}$ . Also note that since  $I_1 \cup \{a_0\}$  only violates the capacity constraint on sets  $A \supseteq A_1$  ( $A \in \mathcal{A}$ ) by the fact that  $\mathcal{M}$  is a laminar matroid, we have that  $I_1 \cup \{a_0\} \setminus \{a_1\} \in \mathcal{M}$ .

Note that  $I_2 \cup \{a_1\}$  can only violate the capacity constraint on sets  $A \supseteq A_1$  ( $A \in \mathcal{A}$ ). We can again find  $A_2 \in \mathcal{A}$  be the minimal set with  $a_1 \in A_2$  and  $|I_2 \cap A_2| = u(A_2)$ , and recursively apply the above argument to tuple  $(I_2, a_1, A_2)$  until arriving  $A_{m-1} \supseteq A^*$ . Since the operation on  $I_1$ , say  $\tilde{I}_1 = I_1 \cup \{a_0\} \setminus \{a_1\} \in \mathcal{M}$ , maintains the number  $|\tilde{I}_1 \cap A_2| = |I_1 \cap A_2|$ , we have that if  $A_2 \subsetneq A^*$ , the following inequality holds

$$\sum_{e_I \in V} \mu_{e_I} \cdot |I \cap A_1| + \mu_{e_{\tilde{I}_1}} (|\tilde{I}_1 \cap A_2| - |I_1 \cap A_2|) < u(A_2).$$

Hence, the above argument can be applied to tuple  $(I_2, a_1, A_2)$ .

Overall, we can get an augmenting path  $(v_1 = e_{I_1}, \dots, v_m = e_{I_m})$  and  $(a_0 = s, a_1, \dots, a_{m-1}, a_m = t)$ , together with a certificate set sequence  $A_1 \subsetneq A_2 \subsetneq \dots \subsetneq A_{m-1}$  where  $A_{m-1} \supseteq A^*$ . Since the depth of  $\mathcal{M}$  is  $\ell$ , we have  $m - 1 \leq \ell$ . Thus, we complete the proof of Claim 4.12.  $\square$

Overall, we complete the proof of Theorem 4.11.  $\square$

## 4.2 Lipschitz Constant $\text{Lip}(\mathcal{B})$ can be Unbounded

We show that  $\text{Lip}(\mathcal{B})$  can be extremely large by considering the following structure constraints  $\mathcal{B}$ , called knapsack polytopes.

**Definition 4.13 (Knapsack polytope).** Let  $A \in \mathbb{R}_{\geq 0}^{m \times k}$  be a non-negative matrix. We say a structure constraint  $\mathcal{B} \in \Delta_k$  is  $A$ -knapsack polytope if

$$\mathcal{B} = \{x \in \Delta_k : Ax \leq 1\},$$

where  $Ax \leq 1$  is called knapsack constraints.

Our result is as follows.

**Theorem 4.14 (Lipschitz constant may be unbounded for knapsack polytope).** *Let  $m = 2$  and  $k = 3$ . For any  $U > 0$ , there exists a matrix  $A \in \mathbb{R}_{\geq 0}^{m \times k}$  such that the Lipschitz constant  $\text{Lip}(\mathcal{B})$  of the  $A$ -knapsack polytope  $\mathcal{B}$  satisfies  $\text{Lip}(\mathcal{B}) \geq U$ .*

*Proof.* We construct  $A$  by  $A_1 = (\frac{10U+2}{5U+3}, \frac{4}{5U+3}, 0)$  and  $A_2 = (\frac{10U+2}{5U+3}, 0, \frac{4}{5U+3})$ . Let  $h = (\frac{1}{2}, \frac{1}{4}, \frac{1}{4})$  and  $h' = (\frac{1}{2} + \frac{1}{10U}, \frac{1}{4} - \frac{1}{20U}, \frac{1}{4} - \frac{1}{20U})$ . We can check that  $h, h' \in \mathcal{B}$  and specifically,  $h'$  is a vertex of  $\mathcal{B}$ . Let  $\sigma : [2] \times [3] \rightarrow \mathbb{R}_{\geq 0}$  be defined as follows:

$$\sigma(1, \cdot) = (\frac{1}{4}, \frac{1}{4}, 0), \text{ and } \sigma(2, \cdot) = (\frac{1}{4}, 0, \frac{1}{4}).$$

We know that  $\sigma \sim (\mathcal{B}, h)$ . Since  $h'$  is a vertex of  $\mathcal{B}$ , there is only one  $\sigma' : [2] \times [3] \rightarrow \mathbb{R}_{\geq 0}$  with  $\sigma' \sim (\mathcal{B}, h')$  and  $\|\sigma'(p, \cdot)\|_1 = \|\sigma(p, \cdot)\|_1$  for  $p \in [2]$ , say

$$\sigma'(1, \cdot) = \frac{1}{2} \cdot h', \text{ and } \sigma'(2, \cdot) = \frac{1}{2} \cdot h'$$

for some  $\alpha \in [0, 1]$ . Thus, we have

$$\sum_{p \in [2]} \|\sigma(p, \cdot) - \sigma'(p, \cdot)\|_1 = \frac{1}{2} + \frac{3}{20U}.$$

However,  $\|h - h'\|_1 = \frac{1}{5U}$ , which implies that

$$\text{Lip}(\mathcal{B}) \geq \text{OAT}(\mathcal{B}, h, h', \sigma) \cdot \|h - h'\|_1 = (\frac{1}{2} + \frac{3}{20U}) / (\frac{1}{5U}) > U.$$

This completes the proof. □

## 5 Bounding the Covering Exponent $\Lambda_\varepsilon(\mathcal{X})$

As we show in Theorem 3.1, the covering exponent  $\Lambda_\varepsilon(\mathcal{X})$  is a key parameter for the complexity of the metric space. In this section, we give several upper bounds of  $\Lambda_\varepsilon(\mathcal{X})$ , against several other well-known notions of “dimension” measure of the metric space  $(\mathcal{X}, d)$ , which enables us to obtain coresets for constrained clustering in various metric spaces (see Remark 5.8 for a summary of the concrete metric families that we can handle). We first introduce the shattering dimension (Definition 5.2), and show its relation to the covering exponent in Lemma 5.3. This relation helps to translate the upper bounds for the shattering dimension (which is rich in the literature) to small-size coresets (Corollary 5.4). Then we consider the doubling dimension (Definition 5.5), and show the covering exponent is bounded by the doubling dimension (Lemma 5.6).

**Shattering Dimension** As was also considered in recent papers [BBH<sup>+</sup>20, BJKW21a, BCAJ<sup>+</sup>22], we employ the following notion of shattering dimension of (the ball range space of) metric space (see also e.g. [Hp11]).

**Definition 5.1 (Ball range space).** *A ball  $\text{Ball}(x, r) = \{y \in \mathcal{X} \mid d(x, y) \leq r\}$  is determined by the center  $c \in \mathcal{X}$  and the radius  $r > 0$ . Let  $\text{Balls}(\mathcal{X}) = \{\text{Ball}(x, r) \mid x \in \mathcal{X}, r > 0\}$  denote the collection of all balls in  $\mathcal{X}$ .  $(\mathcal{X}, \text{Balls}(\mathcal{X}))$  is called the ball range space of  $\mathcal{X}$ . For  $P \subset \mathcal{X}$ , let*

$$P \cap \text{Balls}(\mathcal{X}) := \{P \cap \text{Ball}(x, r) \mid \text{Ball}(x, r) \in \text{Balls}(\mathcal{X})\}.$$

We are ready to define the shattering dimension.

**Definition 5.2 (Shattering dimension of ball range space).** *The shattering dimension of  $(\mathcal{X}, \text{Balls}(\mathcal{X}))$ , denoted by  $\text{dim}(\mathcal{X})$  is the minimum positive integer  $t$  such that for every  $P \subset \mathcal{X}$  with  $|P| \geq 2$ ,*

$$|P \cap \text{Balls}(\mathcal{X})| \leq |P|^t$$

A closely related notion to shattering dimension is the well-known Vapnik-Chervonenkis dimension [VC71]. Technically, they are equivalent up to only a logarithmic factor (see e.g., [Hp11]). The applications of shattering dimension in designing small-sized coresets for unconstrained  $(k, z)$ -CLUSTERING stem from the work [FL11], which are followed by [HJLW18, BBH<sup>+</sup>20, BJKW21a]. However, all these works require to bound the shattering dimension of a more complicated weighted range space. Only recently, this complication of weight in the range space is removed in the uniform sampling framework by [BCAJ<sup>+</sup>22], which only requires to consider the unweighted ball range space (Definition 5.1).

We have the following lemma that upper bounds the covering number via the shattering dimension, whose proof can be found in Section A.

**Lemma 5.3 (Bounding  $\Lambda_\alpha(\mathcal{X})$  via the shattering dimension).** *For every  $\alpha > 0$ , we have  $\Lambda_\alpha(\mathcal{X}) \leq O(\text{dim}(\mathcal{X}) \cdot z^2 \alpha^{-1} \varepsilon^{-1})$ .*

Applying  $\Lambda_\varepsilon(\mathcal{X}) = O_z(\text{dim}(\mathcal{X}) \cdot \varepsilon^{-2})$  to Theorem 3.1, we have the following corollary that bounds the coreset size via  $\text{dim}(\mathcal{X})$ .

**Corollary 5.4 (Coreset size via the shattering dimension).** *Let  $\mathcal{X}(\mathcal{X}, d)$  be a metric space,  $k \geq 1$  be an integer, and  $z \geq 1$ . Let  $\varepsilon, \delta \in (0, 1)$  and  $\mathcal{B}$  be a structure constraint. There exists a randomized algorithm that given a dataset  $P \subseteq \mathcal{X}$  of size  $n \geq 1$  and an  $O(1)$ -approximate solution  $C^* \in \mathcal{X}^k$  of  $P$  for  $(k, z)$ -CLUSTERING, in  $O(nk)$  time constructs an  $\varepsilon$ -coreset for  $(k, z)$ -CLUSTERING with general assignment constraints of size*

$$2^{O(z \log z)} \cdot \tilde{O}(\text{Lip}(\mathcal{B})^2 \cdot \text{dim}(\mathcal{X}) \cdot k^3 \varepsilon^{-3z-2}) \cdot \log \delta^{-1},$$

*with probability at least  $1 - \delta$ , where  $\tilde{O}$  hides a  $\text{poly}(\text{Lip}(\mathcal{B}) \cdot \text{dim}(\mathcal{X}) \cdot k \varepsilon^{-1})$  term.*

**Doubling Dimension** Doubling dimension is an important generalization of Euclidean and more generally  $\ell_p$  spaces with the motivation to capture the intrinsic complexity of a metric space [Ass83, GKL03]. Metric spaces with bounded doubling dimensions are known as doubling metrics. For unconstrained clustering, small-sized coresets were found in doubling metrics [HJLW18, CSS21].

**Definition 5.5 (Doubling dimension [Ass83, GKL03]).** *The doubling dimension of a metric space  $(\mathcal{X}, d)$  is the least integer  $t$ , such that every ball can be covered by at most  $2^t$  balls of half the radius.*

**Lemma 5.6 (Bounding  $\Lambda_\alpha(\mathcal{X})$  via the doubling dimension).** *For every  $\alpha > 0$ , we have  $\Lambda_\alpha(\mathcal{X}) \leq O(\text{ddim}(\mathcal{X}) \cdot \log(z \alpha^{-1} \varepsilon^{-1}))$ .*

The proof can be found in Section B. Similarly, applying  $\Lambda_\varepsilon(\mathcal{X}) = \tilde{O}(\text{ddim}(\mathcal{X}))$  to Theorem 3.1, we have the following corollary that bounds the coreset size via  $\text{ddim}(\mathcal{X})$ .

**Corollary 5.7 (Coreset size via the doubling dimension).** *Let  $\mathcal{X}(\mathcal{X}, d)$  be a metric space,  $k \geq 1$  be an integer, and  $z \geq 1$ . Let  $\varepsilon, \delta \in (0, 1)$  and  $\mathcal{B}$  be a structure constraint. There exists a randomized algorithm that given a dataset  $P \subseteq \mathcal{X}$  of size  $n \geq 1$  and an  $O(1)$ -approximate solution*

$C^* \in \mathcal{X}^k$  of  $P$  for  $(k, z)$ -CLUSTERING, in  $O(nk)$  time constructs an  $\varepsilon$ -coreset for  $(k, z)$ -CLUSTERING with general assignment constraints of size

$$2^{O(z \log z)} \cdot \tilde{O}(\text{Lip}(\mathcal{B})^2 \cdot \text{ddim}(\mathcal{X}) \cdot k^3 \varepsilon^{-3z}) \cdot \log \delta^{-1},$$

with probability at least  $1 - \delta$ , where  $\tilde{O}$  hides a poly  $\log(\text{Lip}(\mathcal{B}) \cdot \text{ddim}(\mathcal{X}) \cdot k \varepsilon^{-1})$  term.

**Remark 5.8 (Covering exponents in different metrics).** We list below several examples of metric spaces that have bounded covering exponent, which is obtained by using Corollary 5.4 or Corollary 5.7.

- Let  $(\mathcal{X}, d)$  be an Euclidean metric  $\mathbb{R}^d$ . We have  $\text{ddim}(\mathcal{X}) \leq d+1$  and hence,  $\Lambda_\varepsilon(\mathcal{X}) \leq \tilde{O}(d)$  by Lemma 5.6. Once we have the bound  $\Lambda_\varepsilon(\mathcal{X}) \leq \tilde{O}(d)$ , we can further assume  $d = \tilde{O}(\varepsilon^{-2} \log k)$  by applying a standard iterative size reduction technique introduced in [BJKW21a], which has been applied in other coreset works [CLSS22, BCAJ<sup>+</sup>22]. This idea yields a coreset of size  $2^{O(z \log z)} \cdot \tilde{O}(\text{Lip}(\mathcal{B})^2 \cdot k^3 \varepsilon^{-3z-2}) \cdot \log \delta^{-1}$ , which removes the dependence of  $d$ .
- Let  $(\mathcal{X}, d)$  be a doubling metric with bounded doubling dimension  $\text{ddim}(\mathcal{X})$ . We directly have  $\Lambda_\varepsilon(\mathcal{X}) \leq \tilde{O}(\text{ddim}(\mathcal{X}))$  by Lemma 5.6.
- Let  $(\mathcal{X}, d)$  be a general discrete metric. Note that  $\text{ddim}(\mathcal{X}) \leq \log |\mathcal{X}|$ , and hence, we have  $\Lambda_\varepsilon(\mathcal{X}) \leq \tilde{O}(\log |\mathcal{X}|)$  by Lemma 5.6.
- Let  $(\mathcal{X}, d)$  be a shortest-path metric of a graph with bounded treewidth  $t$ . By [BT15, BBH<sup>+</sup>20], we know that  $\text{dim}(\mathcal{M}) \leq O(t)$ , which implies that  $\Lambda_\varepsilon(\mathcal{X}) \leq \tilde{O}(t \varepsilon^{-2})$  by Lemma 5.3.
- Let  $(\mathcal{X}, d)$  be a shortest-path metric of a graph that excludes a fixed minor  $H$ . By [BT15], we know that  $\text{dim}(\mathcal{M}) \leq O(|H|)$ , which implies that  $\Lambda_\varepsilon(\mathcal{X}) \leq \tilde{O}(|H| \cdot \varepsilon^{-2})$  by Lemma 5.3.

## References

- [ABM<sup>+</sup>19] Marek Adamczyk, Jaroslaw Byrka, Jan Marcinkowski, Syed Mohammad Meesum, and Michal Wlodarczyk. Constant-factor FPT approximation for capacitated k-median. In *ESA*, volume 144 of *LIPICs*, pages 1:1–1:14. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2019.
- [AMT13] Zeinab Abbassi, Vahab S. Mirrokni, and Mayur Thakur. Diversity maximization under matroid constraints. In *KDD*, pages 32–40. ACM, 2013.
- [Ass83] Patrice Assouad. Plongements lipschitziens dans  $\mathbb{R}^n$ . *Bulletin de la Société Mathématique de France*, 111:429–448, 1983.
- [BBH<sup>+</sup>20] Daniel N. Baker, Vladimir Braverman, Lingxiao Huang, Shaofeng H.-C. Jiang, Robert Krauthgamer, and Xuan Wu. Coresets for clustering in graphs of bounded treewidth. In *ICML*, volume 119 of *Proceedings of Machine Learning Research*, pages 569–579. PMLR, 2020.
- [BCAJ<sup>+</sup>22] Vladimir Braverman, Vincent Cohen-Addad, Shaofeng Jiang, Robert Krauthgamer, Chris Schwiegelshohn, Mads Bech Tofttrup, and Xuan Wu. The power of uniform sampling for coresets. In *62nd IEEE Annual Symposium on Foundations of Computer Science, FOCS 2022*. IEEE Computer Society, 2022.

- [BCFN19] Suman Kalyan Bera, Deeparnab Chakrabarty, Nicolas Flores, and Maryam Negahbani. Fair algorithms for clustering. In *NeurIPS*, pages 4955–4966, 2019.
- [BDHR05] Jonathan L Bredin, Erik D Demaine, MohammadTaghi Hajiaghayi, and Daniela Rus. Deploying sensor networks with guaranteed capacity and fault tolerance. In *Proceedings of the 6th ACM international symposium on Mobile ad hoc networking and computing*, pages 309–319, 2005.
- [BEL13] Maria-Florina Balcan, Steven Ehrlich, and Yingyu Liang. Distributed  $k$ -means and  $k$ -median clustering on general communication topologies. In *NIPS*, pages 1995–2003, 2013.
- [BFS21] Sayan Bandyapadhyay, Fedor V. Fomin, and Kirill Simonov. On coresets for fair clustering in metric and Euclidean spaces and their applications. In Nikhil Bansal, Emanuela Merelli, and James Worrell, editors, *48th International Colloquium on Automata, Languages, and Programming, ICALP 2021, July 12-16, 2021, Glasgow, Scotland (Virtual Conference)*, volume 198 of *LIPICs*, pages 23:1–23:15. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021.
- [BIK07] Moshe Babaioff, Nicole Immorlica, and Robert Kleinberg. Matroids, secretary problems, and online mechanisms. In *SODA*, pages 434–443. SIAM, 2007.
- [BIKK18] Moshe Babaioff, Nicole Immorlica, David Kempe, and Robert Kleinberg. Matroid secretary problems. *J. ACM*, 65(6):35:1–35:26, 2018.
- [BIO<sup>+</sup>19] Arturs Backurs, Piotr Indyk, Krzysztof Onak, Baruch Schieber, Ali Vakilian, and Tal Wagner. Scalable fair clustering. In *ICML*, volume 97 of *Proceedings of Machine Learning Research*, pages 405–413. PMLR, 2019.
- [BJKW19] Vladimir Braverman, Shaofeng H.-C. Jiang, Robert Krauthgamer, and Xuan Wu. Coresets for ordered weighted clustering. In *ICML*, volume 97 of *Proceedings of Machine Learning Research*, pages 744–753. PMLR, 2019.
- [BJKW21a] Vladimir Braverman, Shaofeng H.-C. Jiang, Robert Krauthgamer, and Xuan Wu. Coresets for clustering in excluded-minor graphs and beyond. In *SODA*, pages 2679–2696. SIAM, 2021.
- [BJKW21b] Vladimir Braverman, Shaofeng H.-C. Jiang, Robert Krauthgamer, and Xuan Wu. Coresets for clustering with missing values. In M. Ranzato, A. Beygelzimer, Y. Dauphin, P.S. Liang, and J. Wortman Vaughan, editors, *Advances in Neural Information Processing Systems*, volume 34, pages 17360–17372. Curran Associates, Inc., 2021. URL: <https://proceedings.neurips.cc/paper/2021/file/90fd4f88f588ae64038134f1eeaa023f-Paper.pdf>
- [BLL18] Olivier Bachem, Mario Lucic, and Silvio Lattanzi. One-shot coresets: The case of  $k$ -clustering. In *AISTATS*, volume 84 of *Proceedings of Machine Learning Research*, pages 784–792. PMLR, 2018.
- [BRS08] Ivan Baev, Rajmohan Rajaraman, and Chaitanya Swamy. Approximation algorithms for data placement problems. *SIAM Journal on Computing*, 38(4):1411–1429, 2008.

- [BRU16] Jaroslaw Byrka, Bartosz Rybicki, and Sumedha Uniyal. An approximation algorithm for uniform capacitated  $k$ -median problem with  $1 + \varepsilon$  capacity violation. In *IPCO*, volume 9682 of *Lecture Notes in Computer Science*, pages 262–274. Springer, 2016.
- [BT15] Nicolas Bousquet and Stéphan Thomassé.  $V_c$ -dimension and erdős-pósa property. *Discrete Mathematics*, 338(12):2302–2317, 2015.
- [CGTS02] Moses Charikar, Sudipto Guha, Éva Tardos, and David B. Shmoys. A constant-factor approximation algorithm for the  $k$ -median problem. *J. Comput. Syst. Sci.*, 65(1):129–149, 2002.
- [Che09] Ke Chen. On coresets for  $k$ -median and  $k$ -means clustering in metric and Euclidean spaces and their applications. *SIAM Journal on Computing*, 39(3):923–947, 2009.
- [CHK12] Marek Cygan, MohammadTaghi Hajiaghayi, and Samir Khuller.  $L_p$  rounding for  $k$ -centers with non-uniform hard capacities. In *2012 IEEE 53rd Annual Symposium on Foundations of Computer Science*, pages 273–282. IEEE, 2012.
- [CKLV17] Flavio Chierichetti, Ravi Kumar, Silvio Lattanzi, and Sergei Vassilvitskii. Fair clustering through fairlets. In *NIPS*, pages 5029–5037, 2017.
- [CL19] Vincent Cohen-Addad and Jason Li. On the fixed-parameter tractability of capacitated clustering. In Christel Baier, Ioannis Chatzigiannakis, Paola Flocchini, and Stefano Leonardi, editors, *46th International Colloquium on Automata, Languages, and Programming, ICALP 2019, July 9-12, 2019, Patras, Greece*, volume 132 of *LIPICs*, pages 41:1–41:14. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2019.
- [CLLW16] Danny Z Chen, Jian Li, Hongyu Liang, and Haitao Wang. Matroid and knapsack center problems. *Algorithmica*, 75(1):27–52, 2016.
- [CLSS22] Vincent Cohen-Addad, Kasper Green Larsen, David Saulpic, and Chris Schwiegelshohn. Towards optimal lower bounds for  $k$ -median and  $k$ -means coresets. In *STOC*, pages 1038–1051. ACM, 2022.
- [CSS21] Vincent Cohen-Addad, David Saulpic, and Chris Schwiegelshohn. A new coreset framework for clustering. In *STOC*, pages 169–182. ACM, 2021.
- [DL16] H. Gökalp Demirci and Shi Li. Constant approximation for capacitated  $k$ -median with  $(1+\varepsilon)$ -capacity violation. In *ICALP*, volume 55 of *LIPICs*, pages 73:1–73:14. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2016.
- [FL11] Dan Feldman and Michael Langberg. A unified framework for approximating and clustering data. In *STOC*, pages 569–578. ACM, 2011. <https://arxiv.org/abs/1106.1379>.
- [FS12] Dan Feldman and Leonard J. Schulman. Data reduction for weighted and outlier-resistant clustering. In *SODA*, pages 1343–1354. SIAM, 2012.
- [FSS20] Dan Feldman, Melanie Schmidt, and Christian Sohler. Turning big data into tiny data: Constant-size coresets for  $k$ -means, pca, and projective clustering. *SIAM J. Comput.*, 49(3):601–657, 2020.

- [FZH<sup>+</sup>20] Qilong Feng, Zhen Zhang, Ziyun Huang, Jinhui Xu, and Jianxin Wang. A unified framework of FPT approximation algorithms for clustering problems. In *ISAAC*, volume 181 of *LIPICs*, pages 5:1–5:17. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2020.
- [GKL03] Anupam Gupta, Robert Krauthgamer, and James R. Lee. Bounded geometries, fractals, and low-distortion embeddings. In *FOCS*, pages 534–543. IEEE Computer Society, 2003.
- [HHL<sup>+</sup>16] Mohammadtaghi Hajiaghayi, Wei Hu, Jian Li, Shi Li, and Barna Saha. A constant factor approximation algorithm for fault-tolerant  $k$ -median. *ACM Transactions on Algorithms (TALG)*, 12(3):1–19, 2016.
- [HJLW18] Lingxiao Huang, Shaofeng H.-C. Jiang, Jian Li, and Xuan Wu.  $\varepsilon$ -coresets for clustering (with outliers) in doubling metrics. In *FOCS*, pages 814–825. IEEE Computer Society, 2018.
- [HJLW22] Lingxiao Huang, Shaofeng H.-C. Jiang, Jianing Lou, and Xuan Wu. Near-optimal coresets for robust clustering. *CoRR*, abs/2210.10394, 2022.
- [HJV19] Lingxiao Huang, Shaofeng H.-C. Jiang, and Nisheeth K. Vishnoi. Coresets for clustering with fairness constraints. In *NeurIPS*, pages 7587–7598, 2019.
- [HK07] Sarel Har-Peled and Akash Kushal. Smaller coresets for  $k$ -median and  $k$ -means clustering. *Discret. Comput. Geom.*, 37(1):3–19, 2007.
- [HK20] Monika Henzinger and Sagar Kale. Fully-dynamic coresets. In *ESA*, volume 173 of *LIPICs*, pages 57:1–57:21. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2020.
- [HM04] Sarel Har-Peled and Soham Mazumdar. On coresets for  $k$ -means and  $k$ -median clustering. In *36th Annual ACM Symposium on Theory of Computing*, pages 291–300, 2004.
- [Hp11] Sarel Har-peled. *Geometric Approximation Algorithms*. American Mathematical Society, USA, 2011.
- [HSV21] Lingxiao Huang, K. Sudhir, and Nisheeth K. Vishnoi. Coresets for time series clustering. In Marc’Aurelio Ranzato, Alina Beygelzimer, Yann N. Dauphin, Percy Liang, and Jennifer Wortman Vaughan, editors, *Advances in Neural Information Processing Systems 34: Annual Conference on Neural Information Processing Systems 2021, NeurIPS 2021, December 6-14, 2021, virtual*, pages 22849–22862, 2021.
- [HV20] Lingxiao Huang and Nisheeth K. Vishnoi. Coresets for clustering in Euclidean spaces: Importance sampling is nearly optimal. In *STOC*, pages 1416–1429. ACM, 2020.
- [KKN<sup>+</sup>11] Ravishankar Krishnaswamy, Amit Kumar, Viswanath Nagarajan, Yogish Sabharwal, and Barna Saha. The matroid median problem. In *SODA*, pages 1117–1130. SIAM, 2011.
- [KKN<sup>+</sup>15] Ravishankar Krishnaswamy, Amit Kumar, Viswanath Nagarajan, Yogish Sabharwal, and Barna Saha. Facility location with matroid or knapsack constraints. *Math. Oper. Res.*, 40(2):446–459, 2015.

- [KM13] Jonathan Korman and Robert J McCann. Insights into capacity-constrained optimal transport. *Proceedings of the National Academy of Sciences*, 110(25):10064–10067, 2013.
- [KM15] Jonathan Korman and Robert McCann. Optimal transportation with capacity constraints. *Transactions of the American Mathematical Society*, 367(3):1501–1521, 2015.
- [KMS15] Jonathan Korman, Robert J McCann, and Christian Seis. Dual potentials for capacity constrained optimal transport. *Calculus of Variations and Partial Differential Equations*, 54(1):573–584, 2015.
- [KPS00] Samir Khuller, Robert Pless, and Yoram J Sussmann. Fault tolerant  $k$ -center problems. *Theoretical Computer Science*, 242(1-2):237–245, 2000.
- [Law01] Eugene L Lawler. *Combinatorial optimization: networks and matroids*. Courier Corporation, 2001.
- [Li16] Shi Li. Approximating capacitated  $k$ -median with  $(1 + \varepsilon)k$  open facilities. In *SODA*, pages 786–796. SIAM, 2016.
- [Li17] Shi Li. On uniform capacitated  $k$ -median beyond the natural LP relaxation. *ACM Trans. Algorithms*, 13(2):22:1–22:18, 2017.
- [LS10] Michael Langberg and Leonard J. Schulman. Universal  $\varepsilon$ -approximators for integrals. In *SODA*, pages 598–607. SIAM, 2010.
- [LSS04] Retsef Levi, David B Shmoys, and Chaitanya Swamy. Lp-based approximation algorithms for capacitated facility location. In *International Conference on Integer Programming and Combinatorial Optimization*, pages 206–218. Springer, 2004.
- [LSV13] Jon Lee, Maxim Sviridenko, and Jan Vondrák. Matroid matching: The power of local search. *SIAM J. Comput.*, 42(1):357–379, 2013.
- [MP04] Ramgopal R. Mettu and C. Greg Plaxton. Optimal time bounds for approximate clustering. *Mach. Learn.*, 56(1-3):35–60, 2004.
- [MSSW19] Alexander Munteanu, Chris Schwiegelshohn, Christian Sohler, and David P. Woodruff. On coresets for logistic regression. In Klaus David, Kurt Geihs, Martin Lange, and Gerd Stumme, editors, *49. Jahrestagung der Gesellschaft für Informatik, 50 Jahre Gesellschaft für Informatik - Informatik für Gesellschaft, INFORMATIK 2019, Kassel, Germany, September 23-26, 2019*, volume P-294 of *LNI*, pages 267–268. GI, 2019.
- [MU17] Michael Mitzenmacher and Eli Upfal. *Probability and computing: Randomization and probabilistic techniques in algorithms and data analysis*. Cambridge university press, 2017.
- [PC<sup>+</sup>19] Gabriel Peyré, Marco Cuturi, et al. Computational optimal transport: With applications to data science. *Foundations and Trends® in Machine Learning*, 11(5-6):355–607, 2019.
- [Sch03] Lex Schrijver. *Combinatorial optimization - polyhedra and efficiency. Algorithms and Combinatorics*, 2003.

- [SS08] Chaitanya Swamy and David B. Shmoys. Fault-tolerant facility location. *ACM Trans. Algorithms*, 4(4):51:1–51:27, 2008.
- [SSS19] Melanie Schmidt, Chris Schwiegelshohn, and Christian Sohler. Fair coresets and streaming algorithms for fair  $k$ -means. In *WAOA*, volume 11926 of *Lecture Notes in Computer Science*, pages 232–251. Springer, 2019.
- [SW18] Christian Sohler and David P. Woodruff. Strong coresets for  $k$ -median and subspace approximation: Goodbye dimension. In *FOCS*, pages 802–813. IEEE Computer Society, 2018.
- [Swa16] Chaitanya Swamy. Improved approximation algorithms for matroid and knapsack median problems and applications. *ACM Trans. Algorithms*, 12(4):49:1–49:22, 2016.
- [TT13] Xiaolu Tan and Nizar Touzi. Optimal transportation under controlled stochastic dynamics. *The annals of probability*, 41(5):3201–3240, 2013.
- [TWZ<sup>+</sup>22] Murad Tukan, Xuan Wu, Samson Zhou, Vladimir Braverman, and Dan Feldman. New coresets for projective clustering and applications. In *AISTATS*, volume 151 of *Proceedings of Machine Learning Research*, pages 5391–5415. PMLR, 2022.
- [VC71] VN Vapnik and A Ya Chervonenkis. On the uniform convergence of relative frequencies of events to their probabilities. *Theory of Probability and its Applications*, 16(2):264, 1971.
- [vH14] Ramon van Handel. *Probability in high dimension*. 2014.
- [ZG03] Shi Zhong and Joydeep Ghosh. A unified framework for model-based clustering. *J. Mach. Learn. Res.*, 4:1001–1037, 2003.

## A Proof of Lemma 5.3: $\Lambda_\varepsilon(\mathcal{X})$ Bound via Shattering Dimension

**Lemma A.1** (Restatement of Lemma 5.3). *For every  $\alpha > 0$ , we have  $\Lambda_\alpha(\mathcal{X}) \leq O(z^2\alpha^{-1}\varepsilon^{-1} \cdot \dim(\mathcal{M}))$ .*

*Proof.* We fix an unweighted dataset  $P \subseteq \text{Ball}(a, r_{\max})$  of  $n \geq 2$  points. For ease of statement, we use  $r$  to represent  $r_{\max}$  in the following. We upper bound  $N_{\mathcal{X}}(P, \alpha)$  by constructing an  $\alpha$ -covering  $\mathcal{C}$ . For every  $x \in \text{Ball}(a, 24zr/\varepsilon)$ , let  $f_x(y) = d(x, y)$ ,  $y \in P$  denote its distance function. We round  $f_x(y)$  to obtain an approximation  $\tilde{f}_x$  such that  $\forall y \in P, \tilde{f}_x(y) = \lfloor \frac{24z \cdot d(x, y)}{\alpha r} \rfloor \cdot \frac{\alpha r}{24z}$ . Let  $\mathcal{F}_\alpha = \{\tilde{f}_x \mid x \in \text{Ball}(a, 24zr/\varepsilon)\}$  and for each  $\tilde{f} \in \mathcal{F}_\alpha$ , we add exactly one  $x_0$  such that  $\tilde{f}_{x_0} = \tilde{f}$  into  $\mathcal{C}$  (notice that there can be multiple  $x$ 's that have the same approximation  $\tilde{f}_x$ , we include only one of them).

We claim that  $\mathcal{C}$  is already an  $\alpha$ -covering of  $P$ . To see this, note that for every  $x \in \text{Ball}(a, 24zr/\varepsilon)$ , by construction,  $\forall y \in P, |\tilde{f}_x(y) - f_x(y)| \leq \frac{\alpha r}{24z}$  and also by construction, there exists  $c \in \mathcal{C}$  such that  $\tilde{f}_c = \tilde{f}_x$ . So we conclude that for every

$$\max_{p \in P} |d(x, y) - d(c, y)| \leq \frac{\alpha r}{24z} + \frac{\alpha r}{24z} = \frac{\alpha r}{12z}.$$

It remains to upper bound the size of  $\mathcal{C}$ . To achieve this goal, we first construct a family of distance functions  $G_\alpha$ . Let  $T = O(\frac{z^2}{\alpha \varepsilon})$ . We start with enumerating all possible collections of subsets  $\{H_i \mid i = 0, 1, \dots, T\}$  satisfying the followings

1.  $H_i \in P \cap \text{Balls}(\mathcal{X})$  (Definition 5.1),
2.  $H_T \subseteq H_{T-1} \subseteq \dots \subseteq H_0 \subseteq P$ .

By the definition of the shattering dimension, we know that  $|P \cap \text{Balls}(\mathcal{X})| \leq |P|^{\dim(\mathcal{X})}$ . So there are at most  $(|P|^{\dim(\mathcal{X})})^{O(z^2\alpha^{-1}\varepsilon^{-1})} = |P|^{O(\dim(\mathcal{X}) \cdot z^2\alpha^{-1}\varepsilon^{-1})}$  collections of such  $\{H_i \mid i = 0, \dots, T\}$ .

Fix a collection  $\mathcal{H} = \{H_i \mid i = 0, 1, \dots, T\}$ , we construct a corresponding distance function  $g_{\mathcal{H}}$  as the following. For every  $p \in P$ , let  $i_p$  denote the maximum integer  $i \in \{0, \dots, T\}$  such that  $p \in H_i$ . Let  $g_{\mathcal{H}}(p) = i_p \cdot \frac{\alpha r}{24z}$ . Let  $G_\alpha$  denote the subset of all possible  $g_{\mathcal{H}}$ 's constructed as above.

Now we claim that  $|\mathcal{C}| \leq |G_\alpha|$  since  $|G_\alpha| \leq n^{O(z^2\alpha^{-1}\varepsilon^{-1} \cdot \dim(\mathcal{M}))}$ , which completes the proof. It suffices to show that  $\mathcal{F}_\alpha \subseteq G_\alpha$ . To prove  $\mathcal{F}_\alpha \subseteq G_\alpha$ , we fix  $\tilde{f}_x \in \mathcal{F}_\alpha$  and show that there must exist a realization  $\mathcal{H} = \{H_i \mid i = 0, 1, \dots, T\}$  such that  $g_{\mathcal{H}} = \tilde{f}_x$ . To see this, we simply let  $H_i = \{y \in P \mid \tilde{f}_x(y) \leq i \cdot \frac{\alpha r}{24z}\}$ . It is obvious that  $H_i \in P \cap \text{Balls}(\mathcal{X})$  and  $H_T \subseteq \dots \subseteq H_0$ , thus  $\mathcal{H}$  is a valid realization of the enumeration. Moreover, by construction, we know that  $g_{\mathcal{H}} = \tilde{f}_x$ .  $\square$

## B Proof of Lemma 5.6: $\Lambda_\varepsilon(\mathcal{X})$ Bound via Doubling Dimension

**Lemma B.1** (Restatement of Lemma 5.6). *For every  $\alpha > 0$ , we have  $\Lambda_\alpha(\mathcal{X}) \leq O(\text{ddim}(\mathcal{X}) \cdot \log(z\alpha^{-1}\varepsilon^{-1}))$ .*

*Proof.* Fix an unweighted dataset  $P \subseteq \text{Ball}(a, r_{\max})$  of  $n \geq 2$  points. For ease of statement, we use  $r$  to represent  $r_{\max}$  in the following. By the definition of doubling dimension,  $\text{Ball}(a, 24zr/\varepsilon)$  has an  $\frac{\alpha r}{12z}$ -net of size  $(\frac{z}{\alpha \varepsilon})^{O(\text{ddim}(\mathcal{X}))}$ . By the triangle inequality, we can see that this net is an  $\alpha$ -covering of  $\mathcal{X}$ . So we have

$$N_{\mathcal{X}}(P, \alpha) \leq \left(\frac{z}{\alpha \varepsilon}\right)^{O(\text{ddim}(\mathcal{X}))} \leq n^{O(\text{ddim}(\mathcal{X}) \cdot \log(z\alpha^{-1}\varepsilon^{-1}))}.$$

Due to the arbitrary selection of  $P$  and  $n \geq 2$ , we have  $\Lambda_\alpha(\mathcal{X}) \leq O(\text{ddim}(\mathcal{X}) \cdot \log(z\alpha^{-1}\varepsilon^{-1}))$ , which completes the proof.  $\square$