

A triangle process on graphs with given degree sequence

Colin Cooper

Department of Informatics
King's College
London WC2B 4BG, U.K.
colin.cooper@kcl.ac.uk

Martin Dyer*

School of Computing
University of Leeds
Leeds LS2 9JT, U.K.
m.e.dyer@leeds.ac.uk

Catherine Greenhill[†]

School of Mathematics and Statistics
UNSW Sydney
NSW 2052, Australia
c.greenhill@unsw.edu.au

January 20 2023

Abstract

The triangle switch Markov chain is designed to generate random graphs with given degree sequence, but having more triangles than would appear under the uniform distribution. Transition probabilities of the chain depends on a parameter, called the activity, which is used to assign higher stationary probability to graphs with more triangles. In previous work we proved ergodicity of the triangle switch chain for regular graphs. Here we prove ergodicity for all sequences with minimum degree at least 3, and show rapid mixing of the chain when the activity and the maximum degree are not too large. As far as we are aware, this is the first rigorous analysis of a Markov chain algorithm for generating graphs from a known non-uniform distribution.

1 Introduction

Randomly generating graphs has been the subject of considerable research interest. See, for example, [1, 2, 3, 7, 11, 14, 17, 19, 36, 39, 40, 52]. A comprehensive survey of work to date is given in [25].

*Research supported by EPSRC grant EP/S016562/1, “Sampling in hereditary classes”.

[†]Research supported an Australian Research Council grant DP190100977.

Generation using Markov chains has been an important approach in this area, particularly Markov chains based on switches, for example [3, 11, 14, 17, 36, 38, 39, 40, 52]. Switches delete a pair of edges from the graph and insert a pair using the same four vertices. They have the important property that they preserve the degree sequence of the graph. Thus they are useful for generating regular graphs, or other graphs with a given degree sequence. Markov chains also give a dynamic reconfigurability property, which is useful in some applications, for example [11, 17, 40].

For any such Markov chain, three questions arise. Firstly, can the chain generate any graph in the chosen class? In other words, is the Markov chain ergodic? Secondly, if so what is its equilibrium distribution? Thirdly, what is the rate of convergence to equilibrium? That is, what is the mixing time of the chain?

We investigate the *triangle switch Markov chain*, which we abbreviate as “ Δ -switch chain”. The Δ -switch chain was introduced in [12] to generate regular random graphs with more triangles than would appear under the uniform distribution. See section 1.2 below for the relevant definitions. The chain was proved to be ergodic for 3-regular graphs, and a variant was shown to produce $\Omega(n)$ triangles in $O(n)$ steps of the chain. The proof of ergodicity involved finding a sequence of transitions of the chain (Δ -switches) which transform any 3-regular graph into disjoint copies of K_4 together with a disjoint 6-vertex graph, and then showing that any such graph can be transformed into another using Δ -switches. This proof is not well suited to standard methods for proving rapid mixing of the chain. A further obstacle is that some of the variants studied in [12] do not seem to be time-reversible, a requirement for most approaches to proving rapid mixing.

The proof of ergodicity in [12] was extended in [13] to d -regular graphs, for any d . The proof involved finding a sequence of Δ -switches which transform any d -regular graph into disjoint copies of K_{d+1} together with a disjoint graph, called a “fragment”, with at most $2d$ vertices; then showing that any such graph can be transformed into another using Δ -switches. This required a different approach to proving ergodicity of the fragments, by showing that any step of the flip chain of [40] can be simulated by at most three Δ -switches.

In this paper we extend and generalise the result of [12] by showing that any step of the switch chain [11] can be simulated by at most five Δ -switches. Moreover this holds for all graphs with a given degree sequence such that the minimum degree is at least 3. We prove this in section 2.

Moreover, the simulation approach allows us to prove rapid mixing for some degree sequences of the variant of the chain described in section 1.2. This chain has an equilibrium distribution in which the probability of a graph is exponentially weighted by its number of triangles. The analysis is given in section 4. Such distributions on graphs seem relevant for researchers modelling real-world networks, such as social networks, which contain many triangles [23, 35].

We state our results more formally in section 1.3. As far as we are aware, this is the first rigorous analysis of a Markov chain algorithm for generating graphs from a known non-uniform distribution.

1.1 Notation and definitions

Let $[n]$ denote the set $\{1, 2, \dots, n\}$ and let $G = (V, E)$ be an n -vertex (simple) graph with vertex set $V = [n]$. The degree sequence $\mathbf{d} = (d_1, d_2, \dots, d_n)$ of G is given by $d_i = \deg(i)$ for all $i \in [n]$. An arbitrary sequence \mathbf{d} of n nonnegative integers is called *graphical* if there is any graph which has degree sequence \mathbf{d} . Given the sequence, this property can be tested efficiently and constructively using, for example, the Havel-Hakimi algorithm [28, 31].

We assume without loss of generality that $d_1 \geq d_2 \geq \dots \geq d_n$, so d_1 is the maximum degree and d_n the minimum degree. We note that d_1, d_n are usually denoted by Δ, δ , but we will not do so here. If $d_1 = d_n = d$, the d -regular case, d is common degree of all vertices. Define

$$M = M(\mathbf{d}) = \sum_{i=1}^n d_i = 2|E| = n\bar{d}, \quad M_2 = M_2(\mathbf{d}) = \sum_{i=1}^n d_i(d_i - 1),$$

where \bar{d} is the *average* degree. We will denote the number of triangles in G by

$$t(G) = \left| \left\{ \{i, j, k\} \subseteq V : ij, jk, ki \in E \right\} \right|.$$

If G has degree sequence \mathbf{d} then the maximum value for $t(G)$ is $1/3 \sum_{i=1}^n \binom{d_i}{2} = M_2/6$. (The sum is over all ways of choosing two edges of the triangle meeting a given vertex, the divisor $1/3$ because this counts each triangle three times.) In the regular case, this maximum is achieved by a d -regular graph consisting of the disjoint union of $n/(d+1)$ cliques of size $(d+1)$.

We will make the assumption that $M_2 \geq M$, since this is implicit in some of the proofs of asymptotic distributions below. Now let $\mathcal{G}_{n,\mathbf{d}}$ be the set of all labelled graphs with degree sequence \mathbf{d} . The *switch chain* is a Markov chain which walks on $\mathcal{G}_{n,\mathbf{d}}$ by choosing two non-incident edges uniformly at random, and randomising these two edges without changing the degree sequence. Here we consider a restricted switch chain, in which a switch is only allowed if it changes the set of triangles in the graph. This is achieved by Δ -switches. A Δ^+ -switch is designed to increase the number of triangles, and a Δ^- -switch to decrease this number. See Fig. 1, where $a_2a_1va_3a_4$ is a path in G such that a_1a_3 and a_2a_4 are non-edges, which is transformed to a triangle va_1a_3 and a vertex-disjoint edge a_2a_4 and vice versa.




Fig. 1: : The Δ -switches

We show in Lemma 3.2 below that the $\mathbb{E}[t(G)]$ for a uniformly chosen random graph $G \in \mathcal{G}_{n,\mathbf{d}}$ is $\mu(\mathbf{d}) = M_2^3/M^3$ asymptotically if $d_1 = o(\sqrt{M})$. Our chain will have a non-uniform stationary distribution, where graph G has probability proportional to $\lambda^{t(G)}$, for some parameter $\lambda = \lambda(n) \geq 1$, which we call the *activity*. When $\lambda > 1$ the Δ -switch chain will generate graphs with more triangles than expected in a uniform random graph with the same degree sequence. In fact, we will show that $\mathbb{E}[t(G)] \sim \lambda\mu$ in this distribution, if

λ and d_1 are small enough.

We denote the transitions in the triangle process as Δ -switches. We may use Δ^+ -switch to denote a switch which is known to create a triangle, and Δ^- -switch for a switch which is known to break a triangle. However, a given Δ -switch can simultaneously create one or more triangles and break one or more triangles, in which case it can be viewed as either a Δ^+ -switch or a Δ^- -switch. We reserve the term “switch”, without qualification, for the transitions of the switch chain, as studied in [11], for example.

1.2 Markov chain

Recall that $t(G)$ denote the number of triangles in the graph $G \in \mathcal{G}_{n,d}$. Suppose the current state of the chain is $G \in \mathcal{G}_{n,d}$. The transition procedure of the Δ -switch chain, with parameter $\lambda \geq 1$, is shown in Fig. 2.

```

At the current state  $G \in \mathcal{G}_{n,d}$  do
  choose a pair of non-adjacent edges  $F = \{a_1a_2, a_3a_4\}$  uniformly at random
  choose a perfect matching  $F'$  of  $\{a_1, a_2, a_3, a_4\}$  uniformly at random
  let  $H := ([n], E')$  where  $E' := (E(G) \setminus F) \cup F'$ 
  if  $F' \cap (E(G) \setminus F) = \emptyset$  and
       $(N(a) \cap N(a')) \setminus \{a_1, a_2, a_3, a_4\} \neq \emptyset$  for some  $aa' \in F \cup F'$  then
    the next state is  $H$  with probability  $\min\{1, \lambda^{t(H)-t(G)}\}$ 
  otherwise
    stay at  $G$ 
  end if
end

```

Fig. 2: The transition procedure of the Δ -switch chain with parameter λ on $\mathcal{G}_{n,d}$

Comparing the transition procedure in Fig. 2 with Fig. 1, taking $aa' = a_1a_3$ satisfies the condition $(N(a) \cap N(a')) \setminus \{a_1, a_2, a_3, a_4\} \neq \emptyset$.

At every step we have $P_\lambda(G, G) \geq 1/3$ and hence the Δ -switch chain is aperiodic. We prove in section 2 below that the chain is irreducible, and hence ergodic, provided the minimum degree $d_n \geq 3$.

Define the probability distribution π by $\pi(G) = \lambda^{t(G)} / Z_\lambda(\mathbf{d})$ for all $G \in \mathcal{G}_{n,d}$, where

$$Z_\lambda(\mathbf{d}) = \sum_{H \in \mathcal{G}_{n,d}} \lambda^{t(H)}$$

is the normalising factor. Let $a(\mathbf{d})$ be the number of unordered pairs of non-incident distinct edges in any element of $\mathcal{G}_{n,d}$, given by

$$a(\mathbf{d}) = \binom{M/2}{2} - \frac{M_2}{2}. \quad (1)$$

If $P_\lambda(G, H) > 0$ and $G \neq H$ then, assuming without loss of generality that $t(H) \leq t(G)$, we have

$$\pi(G) P_\lambda(G, H) = \frac{\lambda^{t(G)}}{Z_\lambda(\mathbf{d})} \frac{1}{3a(\mathbf{d})} \lambda^{t(H)-t(G)} = \frac{\lambda^{t(H)}}{Z_\lambda(\mathbf{d})} \frac{1}{3a(\mathbf{d})} = \pi(H) P_\lambda(H, G).$$

Hence the chain satisfies the detailed balance equations with respect to π , so π is the unique stationary distribution of the chain, which is of Metropolis-Hastings type [30].

For technical reasons, we will use a modified Δ -switch chain, obtained by replacing $t(H)$ by $\min\{t(H), \nu\}$ and replacing $t(G)$ by $\min\{t(G), \nu\}$ in the transition procedure Figure 2, where $\nu(n) = \log n / \log \log n$. By the same argument as above, the modified Δ -switch chain has stationary distribution which assigns to graph G a probability proportional to $\lambda^{\min\{t(G), \nu\}}$. The modified Δ -switch chain is polynomial-time indistinguishable from the Δ -switch chain when λ and μ are small enough. See section 4.2.

1.3 Our results

In section 3 we will show that $t(G)$ is approximately $\text{Pois}(\lambda\mu)$ in the equilibrium distribution π of the Δ -switch chain if λ and μ are small enough. In section 4 we will show that the chain is rapidly mixing for smaller values of λ and μ .

However, we must first show that the Δ -switch chain is irreducible on $\mathcal{G}_{n,d}$. This was proved for regular graphs in [13], by showing that (a) every d -regular graph could be transformed by a sequence of Δ -switches into the disjoint union of $\lfloor n/(d+1) \rfloor$ copies of K_{d+1} and one other small component, called a ‘‘fragment’’, and (b) using Δ -switches, it is possible to specify the vertices in each copy of K_{d+1} and to transform fragments into other fragments. This approach is not suitable for proving rapid mixing of the Δ -switch chain by existing methods. Here we will use the ‘‘two-stage direct canonical path construction’’ method from [14] to prove rapid mixing, and this informs our approach to proving irreducibility.

It is known that all graphs with a given degree sequence are connected by switches [38, 51]. In other words, the switch chain is irreducible on $\mathcal{G}_{n,d}$ for all n and d .

Hence in order to prove that the Δ -switch chain is irreducible on $\mathcal{G}_{n,d}$, it suffices to show that every switch (G, H) can be simulated using a sequence of Δ -switches. To do this we will specify a sequence

$$\sigma_{GH} : G = X_0, X_1, \dots, X_\kappa = H$$

such that (X_i, X_{i+1}) is a Δ -switch for $i = 0, \dots, \kappa - 1$. The sequence σ_{GH} is called a *simulation path* for (G, H) of length κ . Rather than simply prove existence of such a path, we will specify one uniquely for each possible switch (G, H) , and show that all our simulation paths have length at most 5. We will do this for degree sequences with minimum degree at least 3, proving the following.

Theorem 1.1. *Let \mathbf{d} be a graphical degree sequence with $d_1 \geq d_2 \cdots \geq d_n \geq 3$ and let $\lambda \geq 1$ be a constant. Then the Δ -switch chain with parameter λ is irreducible on $\mathcal{G}_{n,d}$.*

Note that this is not true in general if $d_n < 3$, see [13]. The above result can be viewed as a *reconfiguration* result on $\mathcal{G}_{n,d}$, generalising that given in [13]. Reconfiguration problems have been a topic of much recent interest in graph theory. See Nishimura [44] for an introduction.

Our second result gives a condition under which we can deduce rapid mixing of the modified Δ -switch chain with parameter λ from rapid mixing of the switch chain.

Theorem 1.2. *Let \mathcal{D} be a family of graphical degree sequences such that for all $\mathbf{d} \in \mathcal{D}$ of length n we have $d_1 \geq d_2 \geq \cdots \geq d_n \geq 3$. Let $\lambda = \lambda(n) \geq 1$ be a function of n . Suppose*

that there exists an $\alpha \in (0, 1)$ such that $\lambda\mu(\mathbf{d}) \leq \log^\alpha n$ for every $\mathbf{d} \in \mathcal{D}$ of length n and all large enough n . If the switch chain is rapidly mixing on $\mathcal{G}_{n,\mathbf{d}}$ for all $\mathbf{d} \in \mathcal{D}$ then the same is true for the modified Δ -switch chain with parameter λ .

The switch chain is known to be rapidly mixing for large classes of graphical sequences, in particular regular sequences [11, 24] and all graphical sequences from P-stable families [16, 20]. By Theorem 1.2, the Δ -switch chain is also rapidly mixing for these graphical sequences, whenever λ and \mathbf{d} satisfy the conditions of the theorem.

To conclude this section we give some background on the use of canonical paths for bounding the mixing time of Markov chains, and the two-stage direct canonical path construction method which we will use to prove Theorem 1.2. Then in section 2 we prove Theorem 1.1, and in section 4 we prove Theorem 1.2.

1.4 Background on canonical paths and mixing time

For further details, see for example Jerrum [33] or Sinclair [48].

Let \mathcal{M} be an ergodic Markov chain with finite state space Ω , transition matrix P and stationary distribution π . Let $\mathcal{G} = (\Omega, E(\mathcal{M}))$ be the graph underlying \mathcal{M} , so the edge set of \mathcal{G} corresponds to (non-loop) transitions of \mathcal{M} .

Given two probability distributions σ, ρ on Ω , the *total variation distance* between them, denoted $d_{\text{TV}}(\sigma, \rho)$, is defined by

$$d_{\text{TV}}(\sigma, \rho) = \frac{1}{2} \sum_{x \in \Omega} |\sigma(x) - \rho(x)| = \max_{A \subseteq \Omega} (\sigma(A) - \rho(A)). \quad (2)$$

The *mixing time* $\tau(\varepsilon)$ of \mathcal{M} is given by

$$\tau(\varepsilon) = \max_{x \in \Omega} \min \{t \geq 0 \mid d_{\text{TV}}(P_x^t, \pi) \leq \varepsilon\},$$

where P_x^t is the distribution of the random state X_t of the Markov chain after t steps with initial state x .

Let the eigenvalues of P be

$$1 = \mu_0 > \mu_1 \geq \mu_2 \cdots \geq \mu_{N-1} > -1$$

and define $\mu_* = \max\{\mu_1, |\mu_{N-1}|\}$. Then, see for example [48, Proposition 1], the mixing time of \mathcal{M} satisfies

$$\tau(\varepsilon) \leq (1 - \mu_*)^{-1} (\ln(1/\pi^*) + \ln(\varepsilon^{-1})) \quad (3)$$

where $\pi^* = \min_{x \in \Omega} \pi(x)$ is the smallest stationary probability. In applications we often focus on the second-largest eigenvalue μ_1 , if necessary by making the Markov chain lazy (replacing the transition matrix P by $(I + P)/2$ to ensure that all eigenvalues are nonnegative).

Jerrum and Sinclair [34] introduced the *canonical path* method for bounding μ_1 . For each pair $x, y \in \Omega$, we choose a path γ_{xy} from x to y in the graph \mathcal{G} underlying the Markov chain. A *canonical* choice of paths seeks to produce a small value of the critical parameter $\bar{\rho}(\Gamma)$, the *congestion* of the set of paths $\Gamma = \{\gamma_{xy} \mid x, y \in \Omega\}$, defined by

$$\bar{\rho}(\Gamma) = \max_{e \in E(\mathcal{M})} Q(e)^{-1} \sum_{\substack{x, y \in \Omega \\ e \in \gamma_{xy}}} \pi(x)\pi(y) |\gamma_{xy}|$$

where $Q(e) = \pi(u) P(u, v)$ when $e = uv \in E(\mathcal{M})$. Then

$$(1 - \mu_1)^{-1} \leq \bar{\rho}(\Gamma), \quad (4)$$

so a low congestion can lead to a good bound on the mixing time, using (3). See [48, Theorem 5] or [33].

In [14], the *two-stage direct canonical path construction method* was described. Here we have two ergodic Markov chains \mathcal{M} and \mathcal{M}' , with state spaces Ω and Ω' , respectively. We assume that we have a set of canonical paths for \mathcal{M}' , and that we can simulate each transition (Y, Z) of \mathcal{M}' by a sequence σ_{YZ} of transitions of \mathcal{M} :

$$\sigma_{YZ} : Y = X_0, X_1, \dots, X_\kappa = Z$$

such that (X_i, X_{i+1}) is a transition of \mathcal{M} . We call σ_{YZ} a $(\mathcal{M}, \mathcal{M}')$ -simulation path and let $\Sigma = \{\sigma_{YZ} \mid (Y, Z) \in E(\mathcal{M}')\}$ be the set of all $(\mathcal{M}, \mathcal{M}')$ -simulation paths.

For each transition $e \in E(\mathcal{M})$, let

$$\Sigma(e) = \{\sigma \in \Sigma \mid e \in \sigma\}$$

be the set of all simulation paths which use the transition e . We define the parameters

$$\ell(\Sigma) = \max_{\sigma \in \Sigma} |\sigma|, \quad B(\Sigma) = \max_{e \in E(\mathcal{M})} |\Sigma(e)| \quad (5)$$

representing the maximum length of any simulation path and the maximum number of simulation paths through a given transition of \mathcal{M} , respectively. Finally, the *simulation gap* between the two chains is defined by

$$D(\mathcal{M}, \mathcal{M}') = \max_{\substack{uv \in E(\mathcal{M}) \\ zw \in E(\mathcal{M}')}} \frac{\pi'(z)P'(z, w)}{\pi(u)P(u, v)} \quad (6)$$

and the *stationary ratio* is defined by

$$R(\mathcal{M}, \mathcal{M}') = \max_{x, y \in \Omega} \frac{\pi(x)\pi(y)}{\pi'(x)\pi'(y)} = \left(\max_{x \in \Omega} \frac{\pi(x)}{\pi'(x)} \right)^2. \quad (7)$$

Note that π and π' are both positive on their domains as $\mathcal{M}, \mathcal{M}'$ are ergodic and reversible.

The following is a slight simplification of [14, Theorem 2.1], tailored to the case that $\Omega = \Omega'$. (If $\Omega = \Omega'$ then the surjection h mentioned in [14, Theorem 2.1] can be taken to be the identity.) We will also take the opportunity to correct a minor error in [14, Theorem 2.1], where the factor $R(\mathcal{M}, \mathcal{M}')$ was omitted.

Theorem 1.3. *Let \mathcal{M} and \mathcal{M}' be two ergodic Markov chains on a set Ω . Let Γ' be a set of canonical paths in \mathcal{M}' and let Σ be a set of $(\mathcal{M}, \mathcal{M}')$ -simulation paths. Then there exists a set of canonical paths in \mathcal{M} whose congestion satisfies*

$$\bar{\rho}(\Gamma) \leq D(\mathcal{M}, \mathcal{M}') R(\mathcal{M}, \mathcal{M}') \ell(\Sigma) B(\Sigma) \bar{\rho}(\Gamma').$$

Proof. In the proof of [14, Theorem 2.1] it is shown that

$$\bar{\rho}(\Gamma) \leq D(\mathcal{M}, \mathcal{M}') \ell(\Sigma) \max_{uv \in E(\mathcal{M})} \sum_{\substack{zw \in E(\mathcal{M}') \\ \sigma_{zw} \in \Sigma(uv)}} \frac{1}{\pi'(z)P'(z, w)} \sum_{\substack{x, y \in \Omega' \\ zw \in \gamma'_{xy}}} \pi(x)\pi(y) |\gamma'_{xy}|$$

and that

$$\max_{uv \in E(\mathcal{M})} \sum_{\substack{zw \in E(\mathcal{M}') \\ \sigma_{zw} \in \Sigma(uv)}} \frac{1}{\pi'(z)P'(z,w)} \sum_{\substack{x,y \in \Omega' \\ zw \in \gamma'_{xy}}} \pi'(x)\pi'(y) |\gamma'_{xy}| \leq B(\Sigma) \bar{\rho}(\Gamma').$$

The result follows as $\pi(x)\pi(y) \leq R(\mathcal{M}, \mathcal{M}') \pi'(x)\pi'(y)$. \square

We remark that in the analysis of the flip chain given in [14], the above theorem is applied twice. In one application the missing factor R is exactly 1, and in the other application it can be bounded above by a constant close to 1 (say 2). So the stated bound on the mixing time of the flip chain in [14] is too small by a factor of at most 2, which is of no consequence. However, in the highly non-uniform distribution on triangles below, the ratio $R(\mathcal{M}, \mathcal{M}')$ becomes rather more significant.

2 Irreducibility

To prove Theorem 1.1, we will show that any switch can be simulated by a sequence of at most five \triangle -switches. We will present the constructions diagrammatically, with the following conventions: solid lines represent known edges of G , dotted lines represent known non-edges, and dashed lines represent non-edges involved in the current \triangle -switch.

Suppose that a \triangle -switch (G, H) replaces the edges a_1a_2 , a_3a_4 with the edges a_1a_3 , a_2a_4 . We classify switches into three types, depending on whether one or both of the *diagonals* a_1a_4 , a_2a_3 is present or absent, as shown in Fig. 3. Note that the diagonals have the form a_ja_{5-j} for $j = 1, 2$.




Fig. 3: The three types of switch, depending on how many diagonals are present

Let $D = \{a_1, a_2, a_3, a_4\}$ be the vertices of the switch, and $A_i = N(a_i) \setminus D$ ($i \in [4]$) be the neighbours of a_i outside of D . Note that $|A_i| = \deg(a_i) - \deg_D(a_i)$, so $|A_i| \geq d_n - 1$ or $|A_i| \geq d_n - 2$. Thus, since $d_n \geq 3$, we have $A_i \neq \emptyset$ for all $i \in [4]$ in any switch.

Lemma 2.1. *Suppose that (G, H) is a switch which removes the edges a_1a_2 , a_3a_4 and replaces them with the edges a_1a_3 , a_2a_4 . Further suppose that at least one of the following holds:*


- (i) *The switch is Type (b) or Type (c). That is, at least one diagonal is present.*
- (ii) *The switch is Type (a), and some pair of distinct elements of $D = \{a_1, a_2, a_3, a_4\}$ have a common neighbour in $[n] \setminus D$.*

- (iii) There is a triangle on vertices a_j, u, w for some vertices u, w which satisfy $u \notin A_{5-j}$ and $\{u, w\} \cap D = \emptyset$, where $j \in [4]$.


Then there is a Δ -switch simulation path of length at most 4 which simulates the switch (G, H) .

Proof. For future reference, we will number the cases which arise when defining the simulation paths. This will be needed later in the proof of rapid mixing. We treat these cases in order, so we only move to (II) if (I) does not hold, and so on.

- (I) If the switch (G, H) is itself a Δ -switch then the simulation path is G, H which has length 1. This occurs if and only if at least one of $A_1 \cap A_2$, $A_1 \cap A_3$, $A_3 \cap A_4$, $A_2 \cap A_4$ is nonempty.
- (II) Now suppose that the switch has Type (a) (both diagonals absent) and $(A_1 \cap A_4) \cup (A_2 \cap A_3)$ is non-empty. After relabelling if necessary, assume that $A_1 \cap A_4 \neq \emptyset$. Then there is a simulation path of length 2, shown below.



- (III) Next, suppose that the switch has Type (b) (both diagonals present) and $(A_1 \cap A_4) \cup (A_2 \cap A_3)$ is non-empty. After relabelling if necessary, $A_1 \cap A_4 \neq \emptyset$. Again, there is a simulation path of length 2, as shown below.




Note that if none of (I), (II) or (III) hold then $A_i \cap A_j = \emptyset$ for all distinct $i, j \in [4]$. Next we suppose that, after relabelling if necessary,


- (IV) – at least one of the diagonal edges $a_j a_{5-j}$ is present, and the symmetric difference $A_i \oplus A_{5-i}$ is nonempty, where $\{i, j\} = \{1, 2\}$, or
- there is a triangle on vertices a_j, u, w for two distinct vertices $u, w \in [n] \setminus D$. (Here we do not care how many diagonal edges are present.)

In the first subcase, choose (i, u) to be lexicographically-least. Then after relabelling if necessary, we can assume that u is the lexicographically-least element of $A_1 \setminus A_4$. In the second subcase, choose (j, u, w) to be the lexicographically-least triple satisfying the requirements. Then after relabelling if necessary, we can assume that $j = 1$. In both cases, we have a simulation path of length 2, consisting of first replacing edges $a_1 u, a_3 a_4$ by $a_1 a_3, u a_4$ and then replacing edges $u a_4, a_1 a_3$ by edges $a_1 u, a_2 a_4$. Under the given conditions, each of these operations is a Δ -switch, as required.


We illustrate the simulation path below for the first subcase when exactly one diagonal a_2a_3 is present (noting that it this still gives a sequence of two \triangle -switches if a_1a_4 is also present), and in the second subcase when neither diagonal is present (noting that this still gives a sequence of two \triangle -switches if one or both diagonals are present).



(V) Next, suppose that the switch has Type (c), so exactly one diagonal a_ja_{5-j} is present, and that $A_i = A_{5-i}$, and that $G[A_i]$ has at least one edge, where $\{i, j\} = \{1, 2\}$. By relabelling if necessary, we may assume that diagonal a_1a_4 is present, and for ease of notation let $A = A_2 = A_3$. Let (u, v) be the lexicographically-least pair of vertices in A such that uv is an edge of G . Then a_2, u, v and a_3, u, v both determine triangles, while $u \notin A_1$ and $v \notin A_4$ as we are not in (I). Then there is a simulation path of length 4 as shown below.



(VI) Finally, suppose that the switch has Type (c), so exactly one diagonal a_ja_{5-j} is present, and that $A_i = A_{5-i}$, and that $G[A_i]$ has no edges, where $\{i, j\} = \{1, 2\}$. Again, by relabelling if necessary, we assume that diagonal a_1a_4 is present, and write $A = A_2 = A_3$. Note that $|A| \geq 2$ as $d_D(a_2) = d_D(a_3) = 1$. Let (u, v) be the lexicographically-least pair of distinct elements in A , and let w be the smallest-labelled vertex in $N(v) \setminus D$, which exists as the minimum degree is at least 3. Then $w \notin A = A_3$ as $v \in A$ and $vw \in E$. We can construct the following simulation path of length 4.



To complete the proof, observe that (I), (III) and (IV) together cover all Type (b) switches, while Type (c) switches are covered by combining (I), (IV), (V) and (VI). This proves the lemma under assumption (i). The proof for assumption (ii) is covered by (I) and (II) together, while (IV) completes the proof for assumption (iii). \square

For the remainder of the proof we assume that the assumptions of Lemma 1 do not hold. Hence the switch has Type (a), that is, no diagonals are present. Furthermore, no two vertices in D have a common neighbour outside D , and there is no triangle of the form $a_j u w$ with $\{u, w\} \cap D = \emptyset$, where $j \in [4]$.

In this situation it is possible that the switch edges $a_1 a_2$, $a_3 a_4$ lie in different components of G , so that H has one fewer components than G . Only a Δ^- -switch can reduce the number of components, and so our strategy is to “plant” a triangle of the form a_j, u, w with $u, w \notin D$ distinct. By assumption, $u a_{5-j}$ is not an edge as $A_j \cap A_{5-j} = \emptyset$. Hence, after planting the triangle, we can simulate the switch using the simulation path described in part (IV) of Lemma 2.1, then “unplant” the planted triangle. There is one exceptional situation where we will not plant a triangle, but instead will simulate the switch (G, H) with a simulation path of length three. (This is case (IXa) of Lemma 2.3 below.)


We will make use of the following lemma to plant a triangle on some element of D . This lemma will be applied to graphs in $\mathcal{G}_{n,d}$, or to graphs in $\mathcal{G}_{n,d}$ with one edge removed.

Lemma 2.2. *Let G be a graph on the vertex set $[n]$. Suppose $v \in [n]$ has a set of three distinct neighbours $R = \{r_1, r_2, r_3\}$, with $\deg(r_i) \geq 2$ for $i = 1, 2, 3$. Assume that there is no triangle in G which contains v and an element of R .*


- (i) *Then there is a Δ^+ -switch (G, \tilde{G}) such that \tilde{G} contains a triangle T with $v \in T$ and $R \cap T \neq \emptyset$.*
- (ii) *Furthermore, we can insist that $r_1 \in T$ unless for $j = 2, 3$ there is a 5-cycle which contains the path $r_1 v r_j$, but no 4-cycle in G contains this path.*

Proof. Let ℓ be the length of the shortest cycle in G which contains a path of the form $r_i v r_j$, where r_i, r_j are distinct elements of R . If no such cycle exists, set $\ell = \infty$. By assumption, $\ell \geq 4$. First suppose that $\ell = 4$. After relabelling the vertices in R if


necessary, assume that the cycle contains the path r_1vr_2 together with some vertex w . (For definiteness, let w be the least-labelled common neighbour of r_1 and r_2 .) Note that vw and r_2r_3 are not edges of G , by assumption. Then we can plant a triangle on v, r_1, w as follows:



Next suppose that $\ell = 5$. After relabelling elements of R if necessary, we can assume that the 5-cycle is $vr_1w_1w_2r_2v$, where $w_1, w_2 \notin \{v\} \cup R$ are distinct. For definiteness, choose (w_1, w_2) to be lexicographically least among all possibilities. Note that vw_1 is a non-edge in G by assumption, and w_2r_3 is a non-edge in G as $\ell \neq 4$. We can “plant” a triangle as follows:



Now suppose that $\ell \geq 6$. After relabelling vertices in R if necessary, suppose that we have a path $w_1r_1vr_2w_2$ in G , where $w_1, w_2 \notin \{v\} \cup R$ are distinct. (For definiteness, choose (w_1, w_2) to be lexicographically least among all options.) Note that r_1r_2 is a non-edge in G by assumption, and w_1w_2 is a non-edge in G as $\ell \neq 5$. Then we can plant a triangle as follows:



In all cases, we have (uniquely) identified a Δ^+ -switch which produces a graph \tilde{G} with the same degree sequence as G which contains a triangle satisfying the stated conditions. This proves (i).

To prove the second statement, adapt the above proof by letting ℓ be the length of the shortest cycle which contains the path r_1vr_j for some $j \in \{2, 3\}$. If $\ell \neq 5$ then the Δ -switch described above creates a triangle T with $r_1 \in T$, as required. If $\ell = 5$ then the Δ -switch described above works if and only if w_2r_3 is not an edge, assuming (after swapping the labels of r_2 and r_3 if necessary) that the 5-cycle $vr_1w_1w_2r_2v$ is present. This completes the proof of (ii). \square

To describe how to remove a planted triangle, we need some terminology. Let (G, \tilde{G}) be a Δ -switch, where \tilde{G} is obtained from G by deleting the edges e_1, e_2 and inserting the edges e_3, e_4 . If \tilde{H} is any graph which contains the edges e_3, e_4 and does not contain e_1, e_2 , then the *inverse* of the Δ -switch (G, \tilde{G}) , applied to \tilde{H} , is the Δ -switch (\tilde{H}, H) where H is obtained from \tilde{H} by deleting the edges e_3, e_4 and inserting the edges e_1, e_2 . This matches the usual notion of an inverse operation when $\tilde{H} = \tilde{G}$, since then $H = G$, but here we need to apply inverse Δ -switches when $\tilde{H} \neq \tilde{G}$.

For us, (G, \tilde{G}) will be a Δ^+ -switch which plants a triangle. Then we will perform two or three Δ -switches from \tilde{G} to reach a graph \tilde{H} which contains e_3, e_4 but not e_1, e_2 . Finally we will remove the planted triangle by applying the inverse of (G, \tilde{G}) to \tilde{H} . There is one exception, which is case (IXa) below, where we do not plant a triangle but instead simulate the required switch with a simulation path of length three.

Lemma 2.3. *Suppose that (G, H) is a switch which removes the edges a_1a_2, a_3a_4 and replaces them with the edges a_1a_3, a_2a_4 . Further suppose that the following conditions all hold:*

- *The switch has Type (a) (no diagonals present),*
- *no pair of vertices in $D = \{a_1, a_2, a_3, a_4\}$ have a common neighbour, and*
- *there is no triangle on vertices a_j, u, w with $\{u, w\} \cap D = \emptyset$, where $j \in [4]$.*


Then there is a Δ -switch simulation path of length at most five which simulates the switch (G, H) .

Proof. We will number the cases which arise when defining the simulation paths. This will be needed later in the proof of rapid mixing. Since Lemma 2.1 has cases (I)–(VI), here our numbering starts from (VII). We treat these cases in order, so we only move to (VIII) if (VII) does not hold, and so on.

(VII) First suppose that after at least one of a_1, a_2, a_3, a_4 has degree at least 4. Choose the least-labelled element of D which has degree at least 4: after relabelling the elements of D if necessary, we can assume that a_1 has degree at least 4. We apply Lemma 2.2(i) with $v = a_1$, and R a set of three neighbours of a_1 other than a_2 . This identifies a Δ^+ -switch (G, \tilde{G}) where \tilde{G} contains a triangle of the form a_1, u, w . From \tilde{G} we can perform the two Δ -switches described in Lemma 2.1, part (IV), to delete the edges a_1a_2, a_3a_4 and replace them with a_1a_3, a_1a_4 using two Δ -switches. From the resulting graph \tilde{H} we apply the inverse of the Δ^+ -switch (G, \tilde{G}) . This “unplants” the planted triangle and takes us to H . This gives a simulation path of length 4.


(VIII) Next, assume that $\deg(a_i) = 3$ for all $i \in [4]$ and that there is a path of length at most 2 in $G \setminus D$ between some element of A_j and some element of A_{2+j} , for some $j \in \{1, 2\}$. Take a shortest such path, and by relabelling if necessary assume that $j = 1$. Let $A_1 = \{u_1, u_2\}$ and $A_3 = \{v_1, v_2\}$ and by relabelling these vertices if necessary, assume that the path has endvertices u_1, v_1 . Note that $u_2 \notin A_4$ and $v_1 \notin A_2$, since no two elements of D have a common neighbour (by assumption).

(VIIIa): If the path has length 1 then the edge u_1v_1 is present and we can plant a triangle as follows.



We have a triangle a_1, u_1, v_2 which allows us to perform the two Δ -switches as described in Lemma 2.1, part (IV), to simulate the required switch. From the resulting graph, we apply the inverse of the triangle-planting Δ -switch, to “unplant” the triangle and produce H . This gives a simulation path of length 4.

(VIIIb): If the path has length 2 then (after possibly relabelling as described above), u_1 and v_1 have a common neighbour w , where we choose w to be the least-labelled common neighbour of u_1, v_1 . Note that $a_3w \notin E$ because $\deg(a_3) = 3$, and $u_2v_1 \notin E$ since otherwise it would be an edge between the neighbourhoods of a_1, a_3 . Then we can plant a triangle as follows:




Now we have a triangle a_1, u_1, w which we can use to perform the two Δ^+ -switches as specified in Lemma 2.1, case (IV). From the resulting graph, we apply the inverse of the triangle-planting Δ -switch, to “unplant” this triangle and produce H . This gives a simulation path of length 4.


(IX) Here we assume that $\deg(a_i) = 3$ for all $i \in [4]$ and that there is no path of length at most 2 between an element of A_j and an element of A_{2+j} , for $j = 1, 2$. Let $A_1 = \{u_1, u_2\}$.

(IXa) First suppose that there is an edge between two neighbours of u_j , and by relabelling if necessary we can assume that $j = 1$. Then $w_1w_2 \in E$ for some $w_1, w_2 \in N(u_1)$ which are distinct from a_1 . (For definiteness, let (w_1, w_2) be the lexicographically-least such pair.) In this subcase, we do not plant a triangle: instead we use the following simulation path of length 3, where v is the least-labelled element of A_3 . Note that $u_1a_3 \notin E$ since $A_1 \cap A_3 = \emptyset$, and $vw_2 \notin E$ since u_1 and v are at distance at least 3 in G .

(IXb) Now suppose that there is no edge between any two neighbours of u_j for $j \in \{1, 2\}$. Let w_1, w_2 be the two least-labelled neighbours of u_1 distinct from a_1 . We want to apply Lemma 2.2(ii) to the graph $G^{(-)} := G - \{a_1a_2\}$ with the edge a_1a_2 deleted, setting $v = u_1$ and $R = \{a_1, w_1, w_2\}$ with $r_1 = a_1$. Suppose that Lemma 2.2(ii) succeeds in identifying a Δ^+ -switch $(G^{(-)}, \tilde{G}^{(-)})$ such that $\tilde{G}^{(-)}$ contains a triangle on the vertices a_1, u_1, z for some vertex $z \notin D$. Let \tilde{G} denote the graph on $\mathcal{G}_{n,d}$ obtained by inserting the edge a_1a_2 in $\tilde{G}^{(-)}$. Then (G, \tilde{G}) is a Δ -switch which can be applied to G . From \tilde{G} , we can perform the two Δ -switches described in Lemma 2.1 case (IV), to exchange the switch edges. Then from the resulting graph, apply the inverse of the Δ -switch (G, \tilde{G}) to “unplant” the triangle and produce the graph H . This gives a simulation path of length 4.



(IXc) Finally we assume that there is no edge between any two neighbours of u_j , for $j \in \{1, 2\}$, but that Lemma 2.2(ii) did not succeed when applied to the graph $G^{(-)} := G - \{a_1 a_2\}$ with $v = u_1$, $R = \{a_1, w_1, w_2\}$ and $r_1 = a_1$. (As above, we assume that w_1, w_2 are the two least-labelled neighbours of u_1 distinct from a_1 .) In particular, this implies that $G^{(-)}$ has a 5-cycle C which contains the path $a_1 u_1 w_1$, but $G^{(-)}$ does not have any 4-cycle which contains the path $a_1 u_1 w_j$ for $j \in \{1, 2\}$. The same properties are true for G . Since a_1 has degree 2 in $G^{(-)}$, the 5-cycle C must be $a_1 u_1 w_1 z u_2$ for some vertex z . Note that $u_1 z$ cannot be an edge of G , or we are in case (IXa), and $w_2 u_2$ cannot be an edge of G as this would give rise to a 4-cycle $a_1 u_1 w_2 u_2$ in $G^{(-)}$ containing the path $a_1 u_1 w_2$. Hence we can perform the following Δ -switch in G to plant a triangle on u_1, w_1, z .



Then we have the situation in (IXa), where u_2 and its neighbours are temporarily unimportant. The three Δ -switches given in case (IXa) perform the required switch, and from the resulting graph we apply the inverse of the Δ -switch which planted $u_1 w_1 w_2$, reaching the graph H . This gives a simulation path of length five. \square

The proof of Theorem 1.1 combines Lemmas 2.1 and 2.3 and the fact that the switch chain is irreducible on $\mathcal{G}_{n,d}$ for all graphical degree sequences \mathbf{d} (see [38, 51]).

3 The asymptotic distribution of $t(G)$

We denote the Poisson distribution with mean μ by $\text{Pois}(\mu)$ and, if the random variable X has this distribution, we write in standard notation $X \sim_d \text{Pois}(\mu)$ or $\mathcal{L}(X) = \text{Pois}(\mu)$. We wish to determine the approximate distribution of $t(G)$ for large n , so we first give an appropriate definition of distributional approximation in this context, which has wider applicability. We note that the traditional approach to distributional approximation, particularly in Statistics, has been through limit distributions, usually with little or no error estimation.

3.1 Distributional approximation

Let \mathcal{X}_n be a sequence of discrete state spaces, and let X_n, X'_n be random variables on \mathcal{X}_n such that $\mathcal{L}(X_n) = p_n, \mathcal{L}(X'_n) = p'_n$. We will say that X_n *approximates* X'_n *asymptotically* if $p'_n(A) = p_n(A) + o(1)$ for all $A \subseteq \Sigma$ as $n \rightarrow \infty$, and we will denote this by $X'_n \approx_d X_n$. If X_n has a known distribution, $\text{Pois}(\mu)$ say, we also write $X'_n \approx_d \text{Pois}(\mu)$. This notation for distributional approximation is taken from [46], but was used only informally there.

i.e.

$$\max_{A \subseteq \mathcal{X}} |\Pr(X'_n \in A) - \Pr(X_n \in A)| = o(1).$$

This involves a known metric on distributions, the total variation distance, $d_{\text{TV}}(p'_n, p_n)$, as defined in (2). So $X'_n \approx_d X_n$ (and vice versa) if the total variation distance

$$d_{\text{TV}}(p'_n, p_n) = \frac{1}{2} \sum_{x \in \mathcal{X}_n} |p'_n(x) - p_n(x)| = o(1) \text{ as } n \rightarrow \infty.$$

This relation is symmetric, reflexive and transitive, since d_{TV} is a metric. However, transitivity holds only over a finite number of applications, since we must preserve the $o(1)$ term. So this is only a local equivalence relation on sequences (\mathcal{X}_n, p_n) .

This definition is not new. Janson [32] used the term ‘‘asymptotic equivalence’’ for this relationship between sequences of random variables, using the notation $(X_n) \cong (X'_n)$. However, the focus in [32] is on the case where both X'_n and X_n are unknown distributions, and the equivalence indicates a stronger relationship than contiguity. There the relationship between X_n and X'_n is symmetrical. Here we assume X_n has a known distribution which approximates the unknown distribution of X'_n , so the relationship is not symmetrical. Also, as we have noted above, this is not a true equivalence relation. Therefore, we prefer the above terminology and notation.

Note that requiring only $p'_n(x) = p_n(x) + o(1)$ for all $x \in \mathcal{X}_n$ is not a useful notion of approximation. For example, suppose $\mathcal{X}_n = [2n]$, $p_n(x) = 1/n$ ($x \in [n]$), $p'_n(x) = 1/n$, ($x \in [2n] \setminus [n]$). Then X_n approximates X'_n in this weaker sense, since $p'_n(x) = p_n(x) \pm 1/n$. However, $d_{\text{TV}}(p_n, p'_n) = 1$ for all n , so this does not seem a reasonable approximation.

Though we will only use it in the discrete case, the approach to distributional approximation outlined above can be extended to arbitrary probability measures, as is done in [32]. It was initially developed for Normal approximations [50], but later for other distributions (e.g. [10, 45, 46]). The methods used have been elementary (e.g. [47]), coupling arguments (e.g. [45]), Stein’s method [50, 6] and semigroup methods (e.g. [37, 49]). The aim has been to quantify errors in limit distributions in Statistics, or to quantify rates of convergence

to these limit distributions, rather than to quantify asymptotic approximations.

We will use only elementary methods, but with an alternative criterion, which we now show is equivalent to the definition above.

Lemma 3.1. $X_n \approx_d X'_n$ if and only if there is a set $S_n \subset \mathcal{X}_n$ such that, as $n \rightarrow \infty$, $p'_n(x) = (1 + o(1))p_n(x)$ for all $x \notin S_n$ and $\max\{p'_n(S_n), p_n(S_n)\} = o(1)$.

Proof. If S_n exists then there exist sequences $0 < \delta_n, \epsilon_n = o(1)$ so that $|p'_n(x) - p_n(x)| < \delta_n p_n(x)$ for $x \notin S_n$ and $\max\{p'_n(S_n), p_n(S_n)\} < \epsilon_n$. Hence

$$d_{\text{TV}}(p_n, p'_n) < \epsilon_n + \delta_n \sum_{x \notin S_n} p_n(x) < \epsilon_n + \delta_n = o(1),$$

so $X_n \approx_d X'_n$.

Conversely, if $X_n \approx_d X'_n$ then there is a sequence $0 < \eta_n = o(1)$ such that $d_{\text{TV}}(p_n, p'_n) < \eta_n$. Let $\epsilon_n = \sqrt{\eta_n} = o(1)$ and define

$$S_n = \left\{ x \in \mathcal{X}_n : \frac{|p'_n(x) - p_n(x)|}{\max\{p_n(x), p'_n(x)\}} \geq \epsilon_n \right\}.$$

Then

$$\epsilon_n^2 > d_{\text{TV}}(p_n, p'_n) \geq \epsilon_n \sum_{x \in S_n} \max\{p_n(x), p'_n(x)\} \geq \epsilon_n \max\{p_n(S_n), p'_n(S_n)\},$$

so $\max\{p_n(S_n), p'_n(S_n)\} < \epsilon_n = o(1)$.

If $x \notin S_n$, then we have $|p'_n(x) - p_n(x)| < \epsilon_n \max\{p_n(x), p'_n(x)\}$. By taking n sufficiently large, we can assume that $\epsilon_n < 1/2$, and hence $p'_n(x) \leq 2p_n(x)$ for all $x \notin S_n$. For all such n and all $x \notin S_n$, we conclude that $p'_n(x)/p_n(x) \in (1 - 2\epsilon_n, 1 + 2\epsilon_n)$, and hence $p'_n(x) \in (1 \pm \delta_n)p_n$ where $\delta_n = 2\epsilon_n = o(1)$ as $n \rightarrow \infty$. This completes the proof. \square

This formulation also gives explicit bounds on errors. In one direction, we can take $\eta_n = \epsilon_n + \delta_n$, while in the other we can take $\epsilon_n = \sqrt{\eta_n}$ and $\delta_n = 2\sqrt{\eta_n}$.

Note that $X'_n \approx_d X_n$ does not imply that X'_n, X_n have exactly the same properties. Suppose $Y_n = h_n(X_n), Y'_n = h_n(X'_n)$ for a function $h_n : \mathcal{X}_n \rightarrow \mathbb{C}$ (the complex numbers). Let $\|h_n\| = \max_{x \in \mathcal{X}_n} |h_n(x)|$. Then we can only claim $\mathbb{E}(Y'_n) \sim \mathbb{E}(Y_n)$ if h_n is uniformly bounded in n using this norm. This follows from the (easily proved) inequality

$$|\mathbb{E}[Y'_n] - \mathbb{E}[Y_n]| \leq \|h_n\| d_{\text{TV}}(p_n, p'_n),$$

with equality possible if $h_n(x) = \text{sign}(p'_n(x) - p_n(x))$, where $\text{sign}(x) \in \{-1, 0, 1\}$ (in the obvious way), so $\|h_n\| = 1$ (unless $p'_n = p_n$ and $d_{\text{TV}}(p'_n, p_n) = 0$). Indeed, this equality is another characterisation of d_{TV} . Note that the uniform boundedness requirement includes X_n itself, so care is needed.

As an example, suppose $\mathcal{X}_n = [n]$, $p_n = C_n/i^2$ ($i \in [n]$), where $C_n \sim \pi^2/6$. (This example is related to the Cauchy distribution.) However, $\mathbb{E}[X_n] = \Omega(\log n) \rightarrow \infty$ as $n \rightarrow \infty$. The explanation is that n is not a uniformly bounded function of n . This is a known pathology in the limit theory of distributions. In this example, X'_n converges pointwise to a fixed distribution, but $\lim_{n \rightarrow \infty} \mathbb{E}[X'_n]$ does not even exist.

In the limit distribution approach to approximation, we must restrict attention to uniformly bounded functions. Here, if h_n is not uniformly bounded, we have the additional

option of estimating the contribution to the expectation from S_n , as we do below for Poisson approximations.

However, if $X'_n \approx_d X_n$ for real-valued random variables, then X_n and X'_n do have the same characteristic function $\mathbb{E}(e^{itx})$ in the limit as $n \rightarrow \infty$, since $|e^{itx}| = 1$ for all real t and x . This could be used, for example, to prove asymptotic Normality of the Poisson approximations below in the limit distribution sense.

However, note that we cannot use this idea of approximation to compare discrete distributions with continuous distributions. Discrete points have measure zero in a continuous distribution, so we always have $d_{\text{TV}}(X_n, X'_n) = 1$. Thus, any definition of approximation in this context must use a weaker metric than d_{TV} . For real-valued random variables, the usual choice is the *Kolmogorov metric* $\max_{x \in \mathbb{R}} |F_n(x) - F'_n(x)|$, where F_n, F'_n are the respective distribution functions of X_n, X'_n . Thus, the asymptotic Normal results for triangles proved in [18] are valid for larger values of μ than the asymptotic Poisson results proved here, but they are much weaker approximations. In fact, the indirect method of proof used in [18] does not seem to allow for any quantification of errors.

In our application below we compare the distribution of $t(G)$, the number of triangles in G chosen uniformly at random from $\mathcal{G}_{n,d}$, to a Poisson distribution with the same mean μ .

The maximum number of triangles in $G \in \mathcal{G}_{n,d}$ is $M_2/6$, so we take $\mathcal{X}_n = \{0, 1, \dots, M_2/6\}$. Then X'_n will be the the number of triangles $t(G)$ when G is chosen uniformly at random from $\mathcal{G}_{n,d}$, $\mu = \mathbb{E}_G[t(G)]$ and $X_n \sim_d \text{Pois}(\mu)$. We will take $S_n = \{G : t(G) > t_0\}$, for a suitably large t_0 , to be determined. Thus S_n is the tail of the distribution of $t(G)$. Then we will show that $X_n \approx_d \text{Pois}(\mu)$, when μ is a small enough unbounded function of n . We divide the proof by analysing first “small” values of $t < t_0$ in section 3.2 and then “large” values of $t > t_0$ in section 3.2. Thus, in section 3.2 we consider $G \notin S_n$, and in section 3.3 we consider $G \in S_n$.

3.2 Small t

For $t = 0, \dots, M_2/6$, let $\mathcal{N}_t = \{G \in \mathcal{G}_{n,d} : t(G) = t\}$ be the set of graphs with exactly t triangles, and let $N_t = |\mathcal{N}_t|$. We will use \mathcal{G}_t to denote an element of \mathcal{N}_t chosen uniformly at random.

Our proof will be based on the *switchings* technique of McKay [41] and others and the conditional probability estimates of Gao and Ohapkin [21]. For reference, we paraphrase [21, Corollary 2] as follows. Let $G \in \mathcal{G}_{n,d}$ and $H^+ \in \mathcal{G}_{n,d'}$, where $d' < d$, and let $M' = \sum_{i=1}^n d'_i = 2|E(H^+)|$. Then, if G is chosen uniformly from $\mathcal{G}_{n,d}$,

$$\Pr(uv \in G | H^+ \subset G) = \left(1 + O\left(\frac{d_1^2}{M - M'}\right)\right) \frac{(d_u - d'_u)(d_v - d'_v)}{M - M' + (d_u - d'_u)(d_v - d'_v)}. \quad (8)$$

In our application H^+ will be a subgraph H_t of G containing exactly t triangles with $t = o(M)$, which will imply that the error term in (8) is $O(d_1^3/M)$.

Remark 1. In [21] there is also a subgraph H^- , edge-disjoint from H^+ , which is conditioned on being absent from G . Since we make no use of this, we omit it. Also, we make the $1 + o(1)$ term explicit in (8), since we iterate these estimates and hence exponentiate this term. We start with $H^+ = \emptyset$, $d' = \mathbf{0}$, so H^+ remains small relative to G .

Remark 2. Note that (8) is essentially the unconditioned estimate on the graph which results from deleting the edges of H_t from G , as might be expected using the configuration

model for graphs with given degree sequences.

We apply these estimates with the switching method, as follows. Consider the set S_t^+ of Δ^+ -switches from \mathcal{N}_t which create exactly one triangle and destroy none. Also define the set of the inverses of these Δ^+ -switches, namely the set S_{t+1}^- of Δ^- -switches from \mathcal{N}_{t+1} which delete exactly one triangle and create none. Let $s_t^+ = |S_t^+|$ and $s_t^- = |S_t^-|$. Then we have

$$N_t \mathbb{E}[s_t^+ | \mathcal{G}_t] = N_{t+1} \mathbb{E}[s_{t+1}^- | \mathcal{G}_{t+1}], \quad (9)$$

from which we can estimate N_{t+1}/N_t . We note that McKay and co-authors have considerably generalised this switchings idea, for example in [29], but here we only require the simple version given above.

Recall that $\mu(\mathbf{d}) = M_2^3/6M^3$. We first prove a claim made in section 1.1 above.

Lemma 3.2. *If $d_1^2 = o(M)$ and $M_2 \geq M$ then $\mathbb{E}[t(G)] \sim \mu$.*

Proof. Using (8), and calculations similar to cases (a) and (b) of Lemma 3.3 below, we have, for triangles i, j, k ,

$$\mathbb{E}[t(G)] = \left(1 + O\left(\frac{d_1^2}{M}\right)\right)^3 \sum_{\substack{i,j,k=1 \\ \text{distinct}}}^n \frac{d_i(d_i-1)d_j(d_j-1)d_k(d_k-1)}{6M^3} \sim \frac{M_2^3}{6M^3} = \mu, \quad (10)$$


where the divisor 6 counts permutations of i, j, k representing the same triangle. The exponent 3 on the error term in (10) results from 3 uses of (8) and replacing the 3 occurrences of $M + O(d_1^2)$ with M . A relative error of $O(d_1^2/M_2)$ arises from the terms where i, j, k are not distinct: these terms arise in M_2^3 but are not present in the original sum. Therefore we need to assume $M_2 = \Omega(M)$, but we make the stronger assumption $M_2 \geq M$. Note that \square

We have $\mu(\mathbf{d}) = M_2^3/6M^3 \geq 1/6$ under our assumption $M_2 \geq M$, so $\mu = \Omega(1)$ as $n \rightarrow \infty$.

Note that the numerator in the summation in (10) is independent of the order in which the edges of the triangle are exposed, and depends only on its degree structure. This observation is true for all small subgraphs, and we use it below without further comment.

Lemma 3.3. *If $td_1^3 = o(M)$ and $M_2 \geq M$ then $\mathbb{E}[s_t^+ | \mathcal{G}_t] = (1 + O(td_1^3/M)) 6M\mu$.*

Proof. We first give an estimate of $\mathbb{E}[s_t^+ | \mathcal{G}_t]$, and then show that it is asymptotically tight. A Δ^+ -switch corresponds to a path of length 5 in G , which we will denote by P_5 , though we make no assumptions about other possible edges between its vertices. The Δ^- -switch corresponding to this P_5 starts with a triangle on its 3 middle edges, and a disjoint edge between its end vertices. We will label these as illustrated below.



Then our initial estimate is the number of P_5 's, where we will temporarily ignore other restrictions:

$$\begin{aligned} \mathbb{E}[s_t^+ \mid \mathcal{G}_t] &\leq (1 + O(d_1^2/M))^4 \sum_{\substack{i,j,k,\ell,m=1 \\ \text{distinct}}}^n \frac{d_i(d_i-1)d_j(d_j-1)d_k(d_k-1)d_\ell d_m}{M^4} \\ &\leq (1 + O(d_1^2/M)) \frac{M_2^3 M^2}{M^4} = 6M\mu, \end{aligned} \quad (11)$$

where the expression in the summation is obtained using (8) with $H_t = \emptyset$.

The estimate (11) is an upper bound because we have ignored conditions that should be imposed and which reduce the expectation. We will now refine this by considering each possible overestimate in turn. This will give us a lower bound on the expectation which matches the upper bound asymptotically for small enough t and d_1 .

- (a) First, we should have $i \neq j \neq k \neq l \neq m$. There are $\binom{5}{2} = 10$ possible first-order equalities in (11), and we claim that each of these gives rise to a $1 + O(d_1^2/M)$ error factor. Consider, for example if $i = j$. If the other vertices are distinct, then (11) becomes

$$\sum_{i,k,l,m=1}^n \frac{d_i^2(d_i-1)^2 d_k(d_k-1)d_l d_m}{M^4} \leq \frac{d_1^2 M_2^2 M^2}{M^4} = \frac{d_1^2 M \mu}{M_2} \leq d_1^2 \mu.$$

Comparing this with (11) gives the claim. So overall the effect is no worse than a $(1 + O(d_1^2/M))^{10} = 1 + O(d_1^2/M)$ factor. Note that higher-order equalities are automatically included, in fact over-counted, in these calculations.

- (b) We have used M in the denominator in (11) rather than the $M - M'$ from (8). In fact, H_t is a set of t triangles, so $M' \leq 6t$, so the denominator in (8) is at most $M + d_1^2$, and at least $M - 6t$. So using M produces a $(1 + O(t + d_1^2)/M)^4 = 1 + O(t + d_1^2)/M$ error factor in (11).
- (c) We have not considered the effect of H_t on the calculation of the quantity $M_2' = \sum_{i=1}^n (d_i - d_i')(d_i - d_i' - 1)$ which corresponds to M_2 on G . We have

$$M_2 \geq M_2' = \sum_{i=1}^n (d_i - d_i')(d_i - d_i' - 1) \geq \sum_{i=1}^n d_i(d_i - 1) - 2d_1 \sum_{i=1}^n d_i' \geq M_2 - 12td_1,$$

So using M_2 produces a $(1 + O(td_1)/M_2)^3 = 1 + O(td_1)/M$ error factor in (11).

- (d) We wish to ensure that almost all P_5 's do not contain an edge of H_t . We claim that these give rise to a $1 + O(td_1^2/M)$ error factor. Consider, for example if $ij \in E(H_t)$. Then (11) becomes

$$\begin{aligned} 2 \sum_{ij \in E(H_t)} \sum_{k,l,m=1}^n \frac{(d_i-1)(d_j-1)d_k(d_k-1)d_l d_m}{M^3} &\leq \frac{2td_1^2 M_2 M^2}{M^3} = \frac{2td_1^2 M_2^3 M^2}{M^3 M_2^2} \\ &= \frac{12td_1^2 \mu M^2}{M_2^2} = O(td_1^2 \mu). \end{aligned}$$

Again, comparing this with (11) gives the claim. Since there are only four edges, the overall the effect is no worse than a factor $(1 + O(td_1^2/M))^4 = 1 + O(td_1^2/M)$.

Since it avoids edges of H_t , the Δ^+ -switch move has probability at most $O(td_1^2 \mu)/M$ of breaking any triangle.

- (e) We need to ensure that the edges jk and ℓm are absent in almost all P_5 's, so that

they give a valid Δ^+ -switches. We claim that these give rise to a $1 + O(d_1^5/M)$ error factor. Consider jk . The expected number of P_5 's in which it is present is at most

$$\sum_{i,j,k,l,m=1}^n \frac{d_i(d_i-1)d_j(d_j-1)(d_j-2)d_k(d_k-1)(d_k-2)d_l d_m}{M^5} \leq \frac{d_1^2 M_2^3 M^2}{M^5} = O(\mu d_1^2),$$

and comparison with (11) gives the claim. The calculation for ℓm is similar.

- (f) We wish to ensure that the edges jk and ℓm are not edges of H_t . Otherwise, the P_5 will not give a valid Δ^+ -switch. If $jk \in E(H_t)$, then (11) becomes

$$\begin{aligned} 2 \sum_{jk \in E(H_t)} \sum_{k,l,m=1}^n \frac{d_i(d_i-1)(d_j-2)(d_k-2)d_l d_m}{M^4} &\leq \frac{2td_1^2 M_2 M^2}{M^4} = \frac{2td_1^2 M_2}{M^2} \\ &= \frac{12td_1^2 \mu}{M} = O(td_1^2 \mu / M). \end{aligned}$$

and, if $\ell m \in E(H_t)$, (11) becomes

$$\begin{aligned} 2 \sum_{lm \in E(H_t)} \sum_{k,l,m=1}^n \frac{d_i(d_i-1)d_j(d_j-2)d_k(d_k-1)(d_\ell-1)(d_m-1)}{M^4} &\leq \frac{2td_1^2 M_2^3}{M^4} \\ &= \frac{12td_1^2 \mu}{M} = O(td_1^2 \mu / M). \end{aligned}$$

Note that these events will not even occur asymptotically if $td_1^2 \mu = o(M)$, since their expectation is then $o(1)$.

- (g) We must ensure that almost all P_5 's create only one triangle. The expected number in which the new edge jk will create a second triangle (hjk) is at most

$$\begin{aligned} \sum_{h,i,j,k,l,m=1}^n \frac{d_h(d_h-1)d_i(d_i-1)d_j(d_j-1)(d_j-2)d_k(d_k-1)(d_k-2)d_l d_m}{M^6} \\ \leq \frac{d_1^2 M_2^4 M^2}{M^6} = \frac{6d_1^2 \mu M_2}{M} = O(\mu d_1^3). \end{aligned}$$

The expected number of P_5 's in which the new edge ℓm will create any triangle (hlm) is at most

$$\begin{aligned} \sum_{h,i,j,k,l,m=1}^n \frac{d_i(d_i-1)d_j(d_j-1)d_k(d_k-1)d_\ell(d_\ell-1)d_h(d_h-1)d_m(d_m-1)}{M^6} \\ = \frac{M_2^6}{M^6} = \mu^2 = O(\mu d_1^3). \end{aligned}$$

Combining these two cases gives an error factor $1 + O(d_1^3/M)$.

The effect of combining all these cases gives an overall error factor which we may bound by $1 + O(d_1^2(d_1 + t)/M) = 1 + O(td_1^3/M)$, the critical cases arising from (d) and (g). Note also that (d) implies that almost all valid Δ^+ -switches give rise to a triangle in \mathcal{G}_{t+1} which is edge-disjoint from H_t . \square

We now consider the Δ^- -switches reversing these Δ^+ -switches.

Lemma 3.4. *If $td_1^2 = o(M)$, then $\mathbb{E}[s_{t+1}^- | \mathcal{G}_t] = (1 + O(td_1^3/M))6(t+1)M$.*

Proof. We must choose a triangle (ijk) and a labelling of it as i, j, k , which can be done in 6 ways. Then, from the proof of Lemma 3.3 we may assume that the triangle (ijk) is

edge-disjoint from any other since, from case (d) in the proof of Lemma 3.3, this will be true for all but a proportion $O(td_1^2/M)$. So deleting jk will delete exactly one triangle. Note that we could show that (ijk) will be vertex-disjoint from H_t , but edge-disjointness is sufficient here.

Deleting lm will delete no triangles if $lm \notin E(H_t)$. So there are at most M , and at least $M - 2|E(H_t)|$ choices for lm . So there are $(1 - O(t/M))M$ choices. If $\ell j \in E(H_t)$ or $mk \in E(H_t)$, the Δ^- -switch will be invalid, but again case (d) in the proof of Lemma 3.4 implies that only a proportion $O(td_1^2/M)$ of Δ^- -switches will \mathcal{G}_{t+1} can be invalid for this reason. Otherwise, the expected number of triangles (ℓjh) created by inserting ℓj will be at most

$$\sum_{h=1}^n \frac{d_h(d_h - 1)(d_j - 2)(d_\ell - 1)}{M^2} \leq \frac{d_1^2 M_2}{M^2} \leq \frac{d_1^3}{M}.$$

So the error factor in $\mathbb{E}[s_{t+1}^- | \mathcal{G}_t]$ is $1 + O(d_1^3/M)$. The same calculation for mk gives the same error, so the total effect will remain $1 + O(d_1^3/M)$.

Combining the error estimates we have an overall $1 + O(td_1^3/M)$ factor, completing the proof. \square

We can now show that $t(G)$ is asymptotically Poisson for a large range of d_1 .

Theorem 3.1. *Let $t_0 = \min\{\sqrt{M/d_1^3}, \omega(n)d_1^3\}$ for any function $\omega(n) \rightarrow \infty$ as $n \rightarrow \infty$. Then, if $t = o(t_0)$, $\Pr(t(G) = t) \sim e^{-\mu} \mu^t / t!$.*

Proof. It follows from Lemmas 3.3 and 3.4 and (9) that

$$\frac{N_{t+1}}{N_t} = (1 + O(td_1^3/M)) \frac{\mu}{t+1},$$

and, iterating this through $t, t-1, \dots, 1$,

$$\frac{\Pr(t(G) = t)}{\Pr(t(G) = 0)} = \frac{N_t}{N_0} = (1 + O(td_1^3/M))^t \frac{\mu^t}{t!} = (1 + O(t^2 d_1^3/M)) \frac{\mu^t}{t!} \sim \frac{\mu^t}{t!}, \quad (12)$$

if $t^2 d_1^3 = o(M)$, which is $t = o(\sqrt{M/d_1^3})$, so is valid for $t \leq t_0$. Summing (12)

$$\frac{\Pr(t \leq t_0)}{\Pr(t = 0)} \sim \sum_{t=0}^{t_0} \frac{\mu^t}{t!} = e^\mu - \sum_{t=t_0+1}^{\infty} \frac{\mu^t}{t!}. \quad (13)$$

Now $d_1^3 = o(t_0)$ implies $\mu = o(t_0)$, so the term on the right is

$$\sum_{t=t_0+1}^{\infty} \frac{\mu^t}{t!} < \sum_{t=t_0+1}^{\infty} \left(\frac{o(t_0)}{t_0} \right)^t \sim o(1)^{t_0} = o(1).$$

In section 3.3 below, we show that $\Pr(t \leq t_0) = 1 - o(1)$. So, since $\mu = \Omega(1)$, (13) gives

$$\frac{\Pr(t \leq t_0)}{\Pr(t = 0)} = \frac{1 - o(1)}{\Pr(t = 0)} \sim e^\mu - o(1) = (1 - o(1))e^\mu. \quad (14)$$

From (14), $\Pr(t = 0) \sim e^{-\mu}$, and so (12) gives $\Pr(t(G) = t) \sim e^{-\mu} \mu^t / t!$, and so $t(G)$ is approximately $\text{Poiss}(\mu)$. \square

The conditions of Theorem 3.1 are satisfied if $d_1 = o(M^{1/9})$ and $d_1^3 = o(t_0)$. This will be satisfied if $d_1 = o(n^{1/9})$ and $t = o(n^{1/3})$.

Remark 3. Gao [18] has shown a Normal limit distribution for $t(G)$ for d -regular graphs as $n \rightarrow \infty$, in the form $(t - \mu)/\sqrt{\mu} \xrightarrow{d} N(0, 1)$, provided $d \rightarrow \infty$ and $d = O(\sqrt{n})$. This is a limit for $\text{Pois}(\mu)$ as $\mu \rightarrow \infty$, but whether Poisson approximation holds for such large d is an open question.

3.3 Large t

Here we will use much weaker bounds, since our goal is only to show that the tail of the distribution of $t(G)$ has very small probability in the equilibrium distribution, provided λ and d_1 is small enough. Then, in combination with Theorem 3.1, we will have shown that $t(G) \approx_d \text{Pois}(\mu)$, in the notation of section 3.1. In fact, we prove something stronger, that the contribution of the tail to $\mathbb{E}[\lambda^{t(G)}]$ is negligible for small enough λ and d_1 . We will use this in section 3.4 to show that $t(G) \approx_d \text{Pois}(\lambda\mu)$ for sufficiently small λ and d_1 . Specifically, when $\lambda > 1$ we will assume that

$$d_1 \log \lambda = o(\log n) \quad \text{and} \quad d_1^3 = o(M).$$

A lower bound on the number $s^-(G)$ of valid Δ^- -switches from $G \in \mathcal{G}_i$ which create no triangle is given by taking any triangle and any edge at distance at least 3 from the triangle. There are at least $M - 6(d_1 - 1)^3$ choices for this directed edge. If $d_1 = o(M^{1/3})$, this is $(1 - o(1))M$. So $s^-(G) \geq (6 - o(1))tM$ since we must choose a triangle edge and a direction, and $s^-(\mathcal{N}_t) = \sum_{G \in \mathcal{N}_t} s^-(G) \geq (6 - o(1))tMN_t$. No triangles are created by any of these Δ^- -switches, but may be destroyed by removal of the switch edges. So we destroy at least 1 and at most $2(d_1 - 1)$ triangles. These bounds are tight, since every edge might lie in a $(d_1 + 1)$ -clique, as is the case for graphs comprising $n/(d_1 + 1)$ cliques of size $(d_1 + 1)$.

So all the Δ^- -switches from $G \in \mathcal{N}_t$ lead to a $G' \in \mathcal{N}'_t = \bigcup_{1 \leq i \leq 2(d_1 - 1)} \mathcal{N}_{t-i}$. Suppose a proportion γ_i leads to \mathcal{N}_{t-i} , so $\sum_{i=1}^{2(d_1 - 1)} \gamma_i = 1$, and note that hence $\sum_{i=1}^{2(d_1 - 1)} \gamma_i N_{t-i} \leq \max\{N_{t-i} : 1 \leq i \leq 2(d_1 - 1)\} = N_{t'}$, for some $t - 2(d_1 - 1) \leq t' \leq t - 1$. Observe that we will have $t' = t - 2(d_1 - 1)$ if G is d_1 -regular and $t = nd_1(d_1 - 1)/6$, so the lower bound can be tight. Moreover, van der Hoorn et al. [27] show that almost all d_1 -regular graphs have almost all their triangles in large cliques for $t(G) = \Omega(n)$, so the analysis we give here may be difficult to improve significantly.

An upper bound on the number $s^+(G')$ of valid Δ^+ -switches from any $G' \in \mathcal{N}'_t$ is the total number of possible switches, valid or not. There are at most $\sum_{i=1}^n 2 \binom{d_i}{2} (d_1 - 1)^2 = M_2(d_1 - 1)^2$ possibilities, so there are at most $M_2(d_1 - 1)^2 N_{t'}$ valid Δ^+ -switches from \mathcal{N}'_t to \mathcal{N}_t . Consequently, we must have $M_2(d_1 - 1)^2 N_{t'} \geq (6 - o(1))tMN_t$, and hence

$$\frac{N_t}{N_{t'}} \leq (1 + o(1)) \frac{M_2(d_1 - 1)^2}{6tM} \leq \frac{\hat{\mu}}{t},$$

where $\hat{\mu} = (d_1 - 1)^3/6 \geq M_2^3/M^3 = \mu$, since $M_2 \leq M(d_1 - 1)$. Note that equality holds if G is d_1 -regular.

We can iterate this to give

$$\frac{N_t}{N_0} \leq \frac{\hat{\mu}^\ell}{tt' \dots t^{(\ell-1)}},$$

where $t^{(j-1)} - t^{(j)} \leq 2d_1 - 2$ for $0 < j < \ell$, $t^{(0)} = t$, $t^{(\ell)} = 0$. The weakest bound results

when $t^{(j-1)} - t^{(j)} = 2d_1 - 2$ and $\ell = t/(2d_1 - 2)$, giving

$$\frac{N_t}{N_0} \leq \frac{\hat{\mu}^{t/(2d_1-2)}}{(2d_1-2)^{t/(2d_1-2)}(t/(2d_1-2))!} = \frac{(\hat{\mu}/(2d_1-2))^{t/(2d_1-2)}}{(t/(2d_1-2))!} < \left(\frac{e\hat{\mu}}{t}\right)^{t/(2d_1-2)},$$

using the bound $(t/(2d_1-2))! > (e^{-1}t/(2d_1-2))^{t/(2d_1-2)}$ from Stirling's approximation. It follows that, for $G \in \mathcal{G}_{n,d}$,

$$\Pr(t(G) = t) < \left(\frac{e\hat{\mu}}{t}\right)^{t/(2d_1-2)}.$$

We will assume that $d_1 = o(n^{1/9})$ and $t_0 = \Omega(n^{1/3})$ from section 3.2. However, we must also assume $\lambda^{2d_1-2}\hat{\mu} = o(t_0)$, which we use to imply $\lambda^{2d_1-2}\hat{\mu} \leq t_0/e^2$. This assumption is true if $d_1 \log \lambda = o(\log n)$, which is a stronger constraint on d_1 unless $\lambda \sim 1$. Then

$$\begin{aligned} \Pr(t \geq t_0) &< \sum_{t=t_0}^{M_2/6} \lambda^t \left(\frac{e\hat{\mu}}{t}\right)^{t/(2d_1-2)} < \sum_{t=t_0}^{\infty} \left(\frac{e\lambda^{2d_1-2}\hat{\mu}}{t}\right)^{t/(2d_1-2)} \\ &< \sum_{t=t_0}^{\infty} e^{-t/(2d_1-2)} = e^{-t_0/(2d_1-2)} / (1 - e^{-1/(2d_1-2)}) \\ &< 2d_1 e^{-t_0/2d_1} = O(e^{-\Omega(n^{1/4})}), \end{aligned}$$

using the inequality $1 - x < e^{-x}$ for $x > 0$.

If $\lambda = 1$, the uniform distribution π' , the constraint $d_1 \log \lambda = o(\log n)$ disappears and the tail bound is small provided $\hat{\mu} = d_1^3/6 = o(t_0)$, i.e. $d_1 = o(\sqrt[3]{t_0})$. So we can take $d_1 = o(n^{1/9})$ and $t = o(n^{1/3})$, as claimed in section 3.2. However, if $\lambda > 1$ is a constant, we must have $d_1 = o(\log n)$.

3.4 Equilibrium distribution of the chain

In Sections 3.2, 3.3, we have shown that $t(G) \approx_d \text{Pois}(\mu)$ under the uniform distribution π' on \mathcal{G} , provided $d_1 = o(n^{1/9})$. Since we know $t(G)$ is approximately Poisson, we can derive its distribution under the equilibrium distribution π of the chain on \mathcal{G} , provided that $d_1 \log \lambda = o(\log n)$, so that the tail of the distribution has probability $o(1)$, as shown in section 3.3. For the uniform distribution π' , using the discussion in section 1.2, we have

$$\mathbb{E}_{\pi'}[\lambda^t] \sim \sum_{k \geq 0} \lambda^k e^{-\mu} \frac{\mu^k}{k!} \sim e^{(\lambda-1)\mu} \sum_{k \geq 0} \frac{(\mu\lambda)^k}{k!} e^{-\mu\lambda} = e^{(\lambda-1)\mu}.$$

as claimed above. Hence, for the Metropolis distribution π , if $t(G) = o(n^\epsilon)$,

$$\Pr_{\pi}(t(G) = s) = \frac{\lambda^s \Pr_{\pi'}(t(G) = s)}{\mathbb{E}_{\pi'}[\lambda^t]} \sim \frac{\lambda^s e^{-\mu} \mu^s / s!}{e^{(\lambda-1)\mu}} = \frac{e^{-\lambda\mu} (\lambda\mu)^s}{s!}.$$

Hence we have shown that $t(G) \approx_d \text{Pois}(\lambda\mu)$ under π . In particular the mean is $\lambda\mu$, so the expected number of triangles increases by a factor λ . So, in order to increase the number of triangles as far as possible, we should take λ as large as possible, under the constraint $d_1 \log \lambda = o(\log n)$.

3.5 Assumptions on the degree sequence

In our asymptotic results we made the assumption that $M_2 \geq M$. We will now prove that this is equivalent to having average degree $\bar{d} \geq 2$. For our Markov chain analysis the condition $\bar{d} \geq 2$ is automatically satisfied, since we assume $d_n \geq 3$.

Lemma 3.5. $M_2 \geq M$ if and only if $\bar{d} \geq 2$.

Proof. We have

$$0 \leq \sum_{i=1}^n (d_i - \bar{d})^2 = \sum_{i=1}^n d_i^2 - n\bar{d}^2 = M_2 + M - M\bar{d} = M_2 - M(\bar{d} - 1),$$

so $M_2 \geq M$ if $\bar{d} \geq 2$. For the converse, suppose that $M_2 < M$. Let n_j be the number of vertices with degree j , for all $j \geq 1$. Then

$$M_2 - M = -n_1 + \sum_{j \geq 3} j(j-2)n_j < 0,$$

and

$$M = n_1 + 2n_2 + \sum_{j \geq 3} jn_j \leq n_1 + 2n_2 + \sum_{j \geq 3} j(j-2)n_j < 2n_1 + 2n_2 \leq 2n.$$

So $\bar{d} < 2$, completing the proof. \square

For our Markov chain $\bar{d} \geq 2$ is automatically satisfied, since we assume $d_n \geq 3$. However, our asymptotic distributions are valid for $\bar{d} \geq 2$. The excluded graphs are rather uninteresting. They have a preponderance of components which are paths, as we see from the following.

Lemma 3.6. $M_2 \geq M$ for any connected graph G which is not a path.

Proof. If G is not a path or a cycle, then $d_1 \geq 3$. A cycle has $\bar{d} = 2$, so $M_2 \geq M$ from Lemma 3.5. Otherwise, we can construct G from a 3-star, a graph with degree sequence $(3, 1, 1, 1)$, by successively adding edges and vertices. For a 3-star, $M_2 = M = 6$. Now, if we add an edge between vertices of degrees d_i and d_j , $M \leftarrow M + 2$, and

$M_2 \leftarrow M_2 + (d_i + 1)d_i + (d_j + 1)d_j - d_i(d_i - 1) - d_j(d_j - 1) = M_2 + 2(d_i + d_j) \geq M_2 + 2$, with equality only if we add an edge from a vertex of degree $d_i = 1$ to a new vertex of degree 0. Otherwise the inequality is strict. The claim now follows by induction. \square

If G is a path P_ℓ with ℓ vertices, then $M = 2(\ell - 1)$ and $M_2 = 2(\ell - 2)$, so $M - M_2 = 2$. Thus, if $M > M_2$, Lemma 3.6 implies that most components must be paths.

4 Rapid mixing

In this section we prove Theorem 1.2 using the two-stage direct canonical path construction method, restated above as Theorem 1.3.

4.1 The simulation path

Recall the definitions from section 1.4. Let \mathcal{M} denote the Δ -switch chain and \mathcal{M}' denote the switch chain. Let P, π be the transition matrix and stationary distribution of the Δ -switch chain \mathcal{M} , and similarly define P', π' with respect to the switch chain \mathcal{M}' . Recall that π' is uniform on $\mathcal{G}_{n, \mathbf{d}}$ while $P'(G, H) = 1/(3a(\mathbf{d}))$ if $G \neq H$ and $P'(G, H) > 0$, where $a(\mathbf{d})$ is defined in (1). We assume that the switch chain \mathcal{M}' is rapidly mixing on $\mathcal{G}_{n, \mathbf{d}}$, and let Γ' be a set of canonical paths for \mathcal{M}' with polynomial congestion. Using results of Sinclair [48], Guruswami [26, Theorem 4.9] proved that under mild conditions on \mathcal{M}' , a polynomial bound on the mixing time of \mathcal{M}' implies the existence of an appropriate set Γ' of canonical paths with $\rho(\Gamma') \leq \text{poly}(n)$.

In section 2 we constructed a set

$$\Sigma = \{\sigma_{GH} \mid (G, H) \in E(\mathcal{M}')\}$$

of $(\mathcal{M}, \mathcal{M}')$ -simulation paths. Recall the parameters defined in (5). The maximum simulation path length is $\ell(\Sigma) = 5$ from Lemmas 2.1 and 2.3. Next we prove an upper bound on the parameter $B(\Sigma)$, which measures the maximum number of simulation paths containing any given Δ -switch.

Lemma 4.1. *Let \mathbf{d} be a graphical degree sequence with $d_1 \geq d_2 \geq \dots \geq d_n \geq 3$ and let Σ be the set of $(\mathcal{M}, \mathcal{M}')$ -simulation paths defined in section 2. Then*

$$B(\Sigma) \leq 20d_1^2(2M + d_1^2)$$

where $M = M(\mathbf{d}) = \sum_{j \in [n]} d_j$.

Proof. Let $e = (X, Y)$ be a transition of the Δ -switch chain. We want an upper bound on the number of simulation paths σ_{GH} which contain e , where (G, H) is a transition of the switch chain. To do this, we will work through the cases (I)–(IX) from Lemmas 2.1 and 2.3. We will not try to optimise the bounds we determine. We are only interested here in correctness and, to a lesser extent, simplicity.

Let $D = \{a_1, a_2, a_3, a_4\}$ denote the vertices involved in the switch (G, H) that we need to identify.

- (I) In Case (I) we have $(G, H) = (X, Y)$ which happens in exactly 1 way.
- (II) In Case (II), the simulation path has length 2 and only involves the vertices in D . So the simulation path is G, G_1, H and in G_1 , the edges $a_1a_4a_2a_3$ are present and the switch edges $a_1a_2, a_1a_3, a_2a_4, a_3a_4$ are all absent. Therefore, if $(X, Y) = (G, G_1)$ then we know $G = X$ and H is obtained from G_1 by replacing edges a_1a_4, a_2a_3 with a_1a_3, a_2a_4 (the only perfect matching of D not seen in either G or G_1). This can be done in exactly one way. By a similar argument, if $(X, Y) = (G_1, H)$ then there is exactly one way to reconstruct G . So overall, there are 2 possibilities for (G, H) if (X, Y) is involved in a simulation path in Case (II).
- (III) There are exactly 2 possibilities arising from Case (III), using a very similar argument to that used for Case (II).
- (IV) The simulation path defined in Case (IV) has length 2, so it is G, G_1, H for some G_1 . Each Δ -switch in the path involves 3 vertices in D and one vertex not in D . (Note

that in the second subcase, the vertex w of the triangle on a_1, u, w is not involved in either of the Δ -switches in the simulation path.) There are 4 ways to label the vertices involved in a given Δ -switch, and in each case the remaining vertex of D is a neighbour of one of the four vertices involved in the Δ -switch. Hence there are at most $4d_1$ ways to label all vertices in D or involved in the Δ -switch, and using these vertices we can uniquely construct H from (G, G_1) , or uniquely construct G from (G_1, H) . Hence there are at most $8d_1$ possibilities for (G, H) if (X, Y) is involved in a simulation path in Case (IV).

- (V) The simulation path in Case (V) has length 4, so it is G, G_1, G_2, G_3, H for some graphs G_1, G_2, G_3 . If $(X, Y) = (G, G_1)$ then one of the edges switched out is the diagonal a_1a_4 , the other edge switched out is uv where $u, v \notin D$. Then a_2 is a common neighbour of a_1, u, v in G , and a_3 is a common neighbour of a_4, u, v in G . So there are 2 choices for which of the removed edges is a_1a_4 and at most d_1^2 ways to choose the unordered pair $\{a_2, a_3\}$. From $G = X$ and this information we can recover H . (Note that by symmetry we do not need to multiply by a factor of 2 to decide which vertex of the edge a_1a_4 is a_1 . The same argument works for the final Δ -switch in the simulation path.)

Next, if $(X, Y) = (G_1, G_2)$ then the Δ -switch involves the vertices a_1, a_2, a_3, a_4 , removing the edges a_1a_2, a_3a_4 which are present in G and inserting the diagonals a_1a_4, a_2a_3 . Hence we know that $H[D]$ is given by the only perfect matching of D not seen in either X or Y . To recover G from X we must know the vertices u, v which are both common neighbours of a_2 and a_3 , so there are at most d_1^2 choices for u, v . The same argument holds if $(X, Y) = (G_2, G_3)$. Overall, there are at most $6d_1^2$ ways that (X, Y) could be involved in a simulation path in Case (V).

- (VI) The simulation path in Case (VI) has length 4. Each Δ -switch in this simulation path involves four of the vertices $a_1, a_2, a_3, a_4, u, v, w$. There are 4 ways to label the vertices involved in each Δ -switch in this path, and then the remaining 3 vertices can be chosen in at most d_1^3 ways, since the subgraph induced by $\{a_1, a_2, a_3, a_4, u, v, w\}$ is connected at each step. Once the identities of these vertices are known, from any Δ -switch on the simulation path, we can construct (G, H) uniquely. It follows that there are at most $16d_1^3$ ways that (X, Y) could be involved in a simulation path in Case (VI).

- (VII) The simulation path in Case (VII) has length 4, consisting of a triangle-planting Δ -switch, then the simulation path from Case (IV) in the resulting graph, then the “unplanting” Δ -switch. Lemma 2.2 is used to choose the triangle-planting Δ -switch, with $v = a_1$ and $R \subseteq N(a_1) \setminus D$. If (X, Y) is the triangle-planting Δ -switch (G, \tilde{G}) then $X = G$. If (in the application of Lemma 2.2) $\ell = 4$ then the triangle-planting Δ -switch involves one vertex from D and three vertices outside D . There are 4 ways to label the vertices involved in the Δ -switch, which identifies a_1 (say). Then there are at most d_1 ways to choose a_2 from the neighbours of a_1 , and at most M ways to choose the vertices (a_3, a_4) . Using these vertices we can construct H . So there are at most $4d_1M$ ways that (X, Y) can be the triangle-planting Δ -switch when $\ell = 4$, and the same bound holds when $\ell = 5$.

However, if $\ell \geq 6$ then the triangle-planting Δ -switch does not involve a_1 , but involves two neighbours u_1, u_2 of a_1 (the neighbours form a triangle with a_1 after

the Δ -switch). Here there are 2 choices for the edge u_1u_2 which is inserted by the Δ -switch, then at most d_1^2 choices for (a_1, a_2) and at most M choices for (a_3, a_4) . Once these vertices are known, we can construct H from G . There are at most $2d_1^2M$ ways that (X, Y) can be the triangle-planting switch when $\ell \geq 6$. Hence, overall there are at most $8d_1M + 2d_1^2M$ ways that (X, Y) could be the triangle-planting Δ -switch in Case (VII). The same bound holds for the “unplanting” Δ -switch at the end of the simulation path, and we must also add the $8d_1$ choices that arise if (X, Y) is used as the second or third Δ -switch in the simulation path. (We multiply $8d_1$ by three as there are three possibilities for ℓ , as described above.) As a grand total, there are at most

$$4d_1^2M + 16d_1M + 24d_1$$

possibilities for (G, H) if (X, Y) is part of a simulation path in Case (VII).

- (VIII) In subcase (VIIIa), the triangle-planting Δ -switch involves two vertices from D , say a_1, a_3 , and two vertices outside D . There are 4 ways to label the vertices involved in the Δ -switch and then at most d_1^2 ways to choose (a_2, a_4) . So if (X, Y) is the triangle-planting Δ -switch in subcase (VIIIa) then there are at most $4d_1^2$ options for (G, H) , as $G = X$ and H can be constructed from G using these known vertices. The same bound holds for the unplanting Δ -switch in subcase (VIIIa), and again we have at most $8d_1$ options from the second or third Δ -switch in the simulation path, giving at most $8d_1^2 + 8d_1$ choices for (G, H) from subcase (VIIIa).

In subcase (VIIIb), the triangle-planting Δ -switch involves exactly one vertex from D , say a_1 , and three vertices outside D . One of the vertices involved in the triangle-planting Δ -switch is a neighbour of a_3 , so there are at most $4d_1^3$ ways to identify all vertices in the triangle-planting Δ -switch and all vertices in D . The same bound holds for the unplanting Δ -switch, and again we must include the bound $8d_1$ to cover the second and third Δ -switches in the simulation path. Hence there are at most $8d_1^3 + 8d_1$ choices for (G, H) if (X, Y) is part of a simulation path in subcase (VIIIb).

Overall, there are at most

$$8d_1^3 + 8d_1^2 + 16d_1$$

choices for (G, H) if (X, Y) is part of a simulation path in case (VIII).

- (IX) In subcase (IXa), the simulation path has length 3, so it is G, G_1, G_2, H for some graphs G_1, G_2 . (Here we do not plant a triangle.) The first Δ -switch involves one vertex of D , say a_3 , and a neighbour of a_1 . So there are at most $4d_1^3$ ways to identify all vertices in the Δ -switch as well as (a_1, a_2, a_4) . Here $G = X$ and we can construct H using the vertices already identified. The second Δ -switch (G_1, G_2) involves the four vertices of D , with u_1 a common neighbour of a_1 and a_3 (say) and where $u_1w_1w_2v$ is a path in G_1 . Hence there are at most $2d_1^4$ ways to identify each of the vertices of $D \cup \{u_1, w_1, w_2\}$. Using this we can construct G from G_1 and then construct H . Finally, the third switch involves exactly one vertex from D , say a_3 , but now it is possible that a_2 and a_4 belong to a different component than a_3 in both G_2 and H . Since $H = Y$, there are at most $4d_1M$ ways to identify the vertices of D and then construct G from H . Overall there are at most $4d_1M + 2d_1^4 + 4d_1^3$ choices for (G, H) if (X, Y) is part of a simulation path in subcase (IXa).

In subcase (IXb), we call Lemma 2.2 to identify a triangle-planting switch (G, \tilde{G}) . Then from \tilde{G} we perform two Δ -switches as described in Case (IV), then we “unplant” the triangle by performing the inverse Δ -switch in the resulting graph. There are several possibilities which we must analyse:

- If Lemma 2.2 uses a 4-cycle which includes the path $a_1u_1w_j$ for $j \in \{1, 2\}$, then the triangle-planting Δ -switch does not involve any vertex of D , but involves a neighbour of a_1 , say. Hence there are at most $4d_1^2M$ ways to identify the vertices of D . Since $G = X$, this allows us to construct H .
- If $a_1u_1w_j$ is not part of a 4-cycle or 5-cycle ($j = 1, 2$) then the triangle-planting Δ -switch involves a_1 , and hence there are at most $4d_1M$ ways to identify the vertices of D and hence construct H from $G = X$.
- If $a_1u_1w_j$ is in a 5-cycle but is not in any 4-cycle, then the triangle-planting Δ -switch does not involve any element of D but it involves neighbours of a_1 , say. So there are at most $4d_1^2M$ ways to identify the vertices of the Δ -switch and of D , which allows us to construct H uniquely from $G = X$.

In each case, the same bounds hold for the final “unplanting” Δ -switch, and there are at most $8d_1$ choices for (G, H) which arise if (X, Y) is the second or third Δ -switch in the simulation path. (We multiply $8d_1$ by 3 to account for the three possible situations for ℓ .) In total, this gives at most

$$2(8d_1^2M + 4d_1M) + 24d_1 = 16d_1^2M + 8d_1M + 24d_1$$

choices for (G, H) when (X, Y) is part of a simulation path in subcase (IXb).

In subcase (IXc), we require a Δ -switch to plant a triangle, say (G, \tilde{G}) . The Δ -switch involves no vertex from D but involves a neighbour of one vertex from D , say a_1 . Here $G = X$ and there are at most $4d_1^2M$ ways to identify the elements of the Δ -switch vertices and D , and hence construct H from G . The same bound holds for the final “unplanting” Δ -switch. The other three Δ -switches in the simulation path match those in subcase (IXa), and we proved above that there are at most $4d_1M + 2d_1^4 + 4d_1^3$ choices for (G, H) if (X, Y) is one of these three Δ -switches. This gives at most

$$4d_1M + 2d_1^4 + 4d_1^3 + 2(4d_1^2M) = 8d_1^2M + 4d_1M + 2d_1^4 + 4d_1^3$$

choices for (G, H) when (X, Y) is part of a simulation path in subcase (IXc).

Hence overall, there are at most

$$24d_1^2M + 16d_1M + 4d_1^4 + 8d_1^3 + 24d_1$$

choices for (G, H) if (X, Y) is part of a simulation path in Case (IX).

Adding up contributions from every case, we see that a Δ -switch (X, Y) can belong to the simulation path of at most

$$28d_1^2M + 32d_1M + 4d_1^4 + 32d_1^3 + 14d_1^2 + 72d_1 + 5$$

pairs (G, G') . We can simplify this using the assumption that $d_1 \geq 3$, and the fact that $M \geq 2d_1$ (as can be seen by considering a star K_{1, d_1}). This gives an upper bound of

$$\left(28 + \frac{32}{3}\right) d_1^2M + \left(4 + \frac{32}{3} + \frac{14}{9} + \frac{72}{27} + \frac{5}{81}\right) d_1^4 < 20d_1^2(2M + d_1^2).$$

This completes the proof. \square

4.2 Modifying the chain

We can now combine the results bounds to prove Theorem 1.2.

We first show that, provided $\lambda > 1$ and λ, μ are small enough, then there is a $\nu(n)$ such that $\Pr_\pi(t > \nu) = o(n^{-r})$ for any constant r , but $\lambda^\nu \leq n^s$ for some constant s . Then we can restrict the chain to the region $t \leq \nu$, since we will almost never see a G with $t(G) > \nu$ in polynomial time.

From section 3 we know that $\Pr(t > \nu) \sim \Pr(X > \nu)$, where $X \approx_d \text{Pois}(\lambda\mu)$, provided $d_1 \log \lambda = o(\log n)$ and $\nu = o(d_1^3)$.

Suppose $\lambda\mu \leq \log^\alpha n$, and $\nu = \log n / \log \log n$ for some constant $0 \leq \alpha < 1$. Then $d_1 \log \lambda = o(\log n)$ and, from above,

$$\begin{aligned} \Pr_\pi(t > \nu) &< (\lambda\mu/\nu)^\nu < (\log^\alpha n \log \log n / \log n)^{\log n / \log \log n} \\ &= (\log \log n)^{-(1-\alpha) \log n} = n^{-(1-\alpha) \log \log \log n} = o(n^{-r}), \end{aligned}$$

for any constant $r > 0$. On the other hand,

$$\lambda^\nu \leq (\log^\alpha n)^{\log n / \log \log n} = e^{\alpha \log n} = n^\alpha,$$

for large enough n .

The simplest way to use this is to modify the chain so that the equilibrium distribution is proportional to $\lambda^{\min\{t, \nu\}}$. This is easy to do using detailed balance. The total variation distance of this distribution from the original is $o(n^{-r})$ for all constant r , so it is polynomial-time indistinguishable from the true Metropolised distribution π .

Then we have

$$\frac{Z_\lambda(\mathbf{d})}{|\mathcal{G}_{n,\mathbf{d}}|} = \mathbb{E}_u[\lambda^t] < \lambda^\nu \leq n^\alpha, \quad \text{and} \quad R(\mathcal{M}, \mathcal{M}') \leq \lambda^{2\nu} \leq n^{2\alpha},$$

for the above ranges of λ and μ and where $R(\mathcal{M}, \mathcal{M}')$ is given by (7). We now apply this to the rapid mixing proof.

Proof. Let $\tau_\lambda(\varepsilon)$ denote the mixing time of the Δ -switch chain with parameter λ . Suppose that the switch chain is rapidly mixing on $\mathcal{G}_{n,\mathbf{d}}$ for all $\mathbf{d} \in \mathcal{D}$, and let $\mathbf{d} \in \mathcal{D}$ have length n . Hence we may assume (see for example Guruswami [26, Theorem 4.9]) that there exists a set of canonical paths Γ' for the switch chain with congestion $\bar{\rho}(\Gamma')$ bounded above by a polynomial in n , say

$$\bar{\rho}(\Gamma') \leq q(n) \tag{15}$$

for all $\mathbf{d} \in \mathcal{D}$ with length n .

We will bound the mixing time using (3). It follows from a result of Diaconis and Saloff-Coste [15, p. 702] that the smallest eigenvalue of the Δ -switch chain satisfies

$$(1 + \mu_{N-1})^{-1} \leq \max_{H \in \mathcal{G}_{n,\mathbf{d}}} \frac{1}{2 P_\lambda(H, H)} \leq 3/2, \tag{16}$$

where P_λ denotes the transition matrix of the Δ -switch chain with parameter λ . We will bound μ_1 by applying the direct two-step canonical path construction method (as restated in Theorem 1.3) using the set of simulation paths Σ defined in section 2.

It follows from Lemma 2.1 and Lemma 2.3 that

$$\ell(\Sigma) = 5. \tag{17}$$

An upper bound for $B(\Sigma)$ was proved in Lemma 4.1. The simulation gap $D(\mathcal{M}, \mathcal{M}')$, defined in (6), satisfies

$$\begin{aligned}
D(\mathcal{M}, \mathcal{M}') &= \max_{\substack{uv \in E(\mathcal{M}) \\ zw \in E(\mathcal{M}')}} \frac{\pi'(z)P'(z, w)}{\pi(u)P_\lambda(u, v)} \\
&= \max_{\substack{uv \in E(\mathcal{M}) \\ zw \in E(\mathcal{M}')}} \frac{|\mathcal{G}_{n, \mathbf{d}}|^{-1} (3a(\mathbf{d}))^{-1}}{\lambda^{t(u)} Z_\lambda(\mathbf{d})^{-1} \min\{1, \lambda^{t(v)-t(u)}\} (3a(\mathbf{d}))^{-1}} \\
&= \max_{uv \in E(\mathcal{M})} \frac{|\mathcal{G}_{n, \mathbf{d}}|^{-1} Z_\lambda(\mathbf{d})}{\min\{\lambda^{t(u)}, \lambda^{t(v)}\}} \leq \frac{Z_\lambda(\mathbf{d})}{|\mathcal{G}_{n, \mathbf{d}}|} \\
&\leq n^\alpha,
\end{aligned} \tag{18}$$

from above. Note that the inequality in (18) above will be an equality if

$$\min_{uv} \{\lambda^{\min\{t(u), \nu\}}, \lambda^{\min\{t(v), \nu\}}\} = 1.$$

It is easy to see that this will be the case if $\mathcal{G}_{n, \mathbf{d}}$ contains at least two graphs, such that one of them is triangle-free.

Hence we have

$$D(\mathcal{M}, \mathcal{M}') R(\mathcal{M}, \mathcal{M}') \leq n^\alpha n^{2\alpha} = n^{3\alpha}.$$

Using Theorem 1.3, Lemma 4.1 and (15)–(18), we conclude that there is a set Γ of canonical paths for the Δ -switch chain with parameter λ such that the congestion of Γ satisfies

$$\bar{\rho}(\Gamma) \leq D(\mathcal{M}, \mathcal{M}') R(\mathcal{M}, \mathcal{M}') |\ell(\Sigma) B(\Sigma) \bar{\rho}(\Gamma') \leq 100n^3 d_1^2 (2M + d_1^2) q(n),$$

using the assumption of the theorem for the final inequality. It follows from (4) that $\mu \leq 1 - 1/\bar{\rho}(\Gamma)$ and we can upper bound the mixing time using (3).

It can be proved, for example using the configuration model [8], that

$$\log |\mathcal{G}_{n, \mathbf{d}}| \leq M \ln M$$

where $M = M(\mathbf{d}) = \sum_{j \in [n]} d_j$, as above. Furthermore, if π^* denotes the smallest stationary probability of the modified Δ -switch chain, then

$$1/\pi^* = \hat{Z}_\lambda(\mathbf{d}) = \sum_{G \in \Omega_{n, \mathbf{d}}} \lambda^{\min(t(G), \nu)} \leq Z_\lambda(\mathbf{d}),$$

where $\hat{Z}_\lambda(\mathbf{d})$ is the normalisation factor for the stationary distribution of the modified Δ -switch chain. Hence (15) implies that

$$\log(1/\pi^*) \leq M \log M + \alpha \log n.$$

Putting all this together, using (3), (4) and (16), gives

$$\tau_\lambda(\varepsilon) \leq 100n^{3\alpha} d_1^2 (2M + d_1^2) q(n) (M \ln M + \alpha \log n + \ln(\varepsilon^{-1})).$$

Thus, the modified Δ -switch chain with parameter λ is rapidly mixing for all $\mathbf{d} \in \mathcal{D}$, under the assumption $\lambda\mu \leq \log^\alpha n$ for some $0 < \alpha < 1$. \square

These results seem weak, since the assumption $\lambda\mu \leq \log^\alpha n$ is quite restrictive, but improvement requires a different approach to analysing the Markov chain.

4.3 Slow mixing

It seems disappointing that we can only prove rapid mixing for rather small values of μ and λ , but this is largely a consequence of the Metropolisisation. Consider a d -regular graph G which consists of k disjoint copies of K_{d+1} . (Note that these graphs are central to the irreducibility proof in [13].) Then G contains no paths of length 4, so no Δ^+ -switch is possible. Any Δ^- -switch must take one edge from each of two disjoint copies of K_{d+1} , and hence will destroy $2(d-1)$ triangles. So any change to G will occur only with probability $\lambda^{-2(d-1)}$. Suppose that $\lambda > 1$ is any constant and $d = \gamma \log n$, where $\gamma \rightarrow \infty$ as $n \rightarrow \infty$. Then, with high probability, no change will occur within $(1/\lambda)^d = n^{\gamma \log \lambda}$ steps of the chain, which is superpolynomial.

There are exponentially few d -regular graphs of this form, as a fraction of the set of all d -regular graphs on $n = kd$ vertices, but each such graph gives a potential bottleneck for mixing. The small proportion suggests a solution by “ignoring” these graphs in some way. Unfortunately, the difficulty is not restricted to these graphs. Suppose we have only a single K_{d+1} component. In order to break the K_{d+1} , we require the two vertices of a non-edge in the remainder of the graph to have many common neighbours, at least $d/4$, say. Then a Δ^- -switch could create at most $d/2$ triangles, which is not enough to offset the $d-1$ triangles that have been destroyed. If there are $o(n)$ triangles in the current graph, then we can bound the probability of such a non-edge appearing in the graph by $O(n^{d/4+2}(d/n)^{d/2}) = o(n^{-d/5})$, which is exponentially small if $d \rightarrow \infty$ with n . Thus all Δ^- -switch’s involving the K_{d+1} will have exponentially small probability of acceptance, and hence mixing will take exponential time.

Of course, the proportion of d -regular graphs with n vertices and at least one K_{d+1} component is still small. But, if $t(G) = \Omega(n)$, it is shown in [27] that G contains many dense clique-like structures, called pseudocliques, which would require many steps to disassemble. Therefore, it seems unlikely that the Δ -switch chain could generate regular graphs with $\Omega(n)$ triangles in polynomial time, except possibly for small constant d . Nevertheless, the chain may well be rapidly mixing for a wider range of μ and λ than we are able to prove here.

References

- [1] D. Allendorf, U. Meyer, M. Penschuck, H. Tran and N. Wormald (2022), Engineering uniform sampling of graphs with a prescribed power-law degree sequence, in Proc. Symposium on Algorithm Engineering and Experiments (ALENEX 2022), pp. 27–40.
- [2] Z. Allen-Zhu, A. Bhaskara, S. Lattanzi, V. Mirrokni and L. Orecchia (2018), Expanders via local edge flips, in: *Proc. 27th Annual ACM-SIAM Symposium on Discrete Algorithms* (SODA 2016), pp. 259–269.
- [3] G. Amanatidis and P. Kleer (2020), Rapid mixing of the switch Markov chain for strongly stable degree sequences, *Random Struct. Algorithms* **57**, 637–657.
- [4] A. Arman, P. Gao and N. Wormald (2019), Fast uniform generation of random graphs with given degree sequences, in *60th Annual IEEE Symposium on Foundations of Computer Science (FOCS 2019)*, pp. 1371–1379.

- [5] T. Bannink, R. van der Hofstad and C. Stegehuis (2018), Switch chain mixing times and triangle counts in simple random graphs with given degrees, *Journal of Complex Networks* **7**, 210–225.
- [6] A. Barbour, L. Holst and S. Janson, *Poisson Approximation*, Oxford University Press, 1992.
- [7] M. Bayati, J.H. Kim and A. Saberi (2010), A sequential algorithm for generating random graphs, *Algorithmica* **58**, 860–910.
- [8] B. Bollobás (1980), A probabilistic proof of an asymptotic formula for the number of labelled regular graphs, *European Journal of Combinatorics* **1**, 311–316.
- [9] S. Chatterjee, P. Diaconis and A. Sly (2011), Random graphs with a given degree sequence, *Annals of Applied Probability* **21**, 1400–1435.
- [10] L. Chen (1975), Poisson approximation for dependent trials, *Annals of Probability* **3**, 534–545.
- [11] C. Cooper, M. Dyer and C. Greenhill (2007), Sampling regular graphs and a peer-to-peer network, *Combinatorics, Probability and Computing* **16**, 557–593.
- [12] C. Cooper, M. Dyer and C. Greenhill (2019), Triangle-creation processes on cubic graphs, [arXiv:1905.04490](https://arxiv.org/abs/1905.04490).
- [13] C. Cooper, M. Dyer and C. Greenhill (2021), A triangle process on regular graphs, in *Proc. 32nd International Workshop on Combinatorial Algorithms (IWOCA 2021)*, Springer Lecture Notes in Computer Science, vol. 12757, pp. 310–323.
- [14] C. Cooper, M. Dyer, C. Greenhill and A. Handley (2019), The flip Markov chain for connected regular graphs, *Discrete Applied Mathematics* **254**, 56–79.
- [15] P. Diaconis and L. Saloff-Coste (1993), Comparison theorems for reversible Markov chains, *Annals of Applied Probability* **3**, 696–730.
- [16] P.L. Erdős, C. Greenhill, T.R. Mezei, I. Miklos, D. Soltész, L. Soukup (2022), The mixing time of the switch Markov chains: a unified approach. *European Journal of Combinatorics*, **99**, 103421.
- [17] T. Feder, A. Guetz, M. Mihail and A. Saberi (2006), “A local switch Markov chain on given degree graphs with application in connectivity of peer-to-peer networks”, in: *Proc. 47th Annual IEEE Symposium on Foundations of Computer Science (FOCS 2006)*, pp. 69–76.
- [18] P. Gao (2021), Triangles and subgraph probabilities in random regular graphs, Preprint: [arXiv:2012.01492](https://arxiv.org/abs/2012.01492).
- [19] P. Gao and N. Wormald (2017), Uniform generation of random regular graphs, *SIAM Journal of Computing* **46**, 1395–1427.
- [20] P. Gao and C. Greenhill (2021), Mixing time of the switch Markov chain and stable degree sequences, *Discrete Applied Mathematics* **291**, 143–162.

- [21] P. Gao and Y. Osherson (2022), Subgraph probability of random graphs with specified degrees and applications to chromatic number and connectivity, *Random Structures & Algorithms*, to appear. Also [arXiv:2007.02216](https://arxiv.org/abs/2007.02216).
- [22] Z. Gao and N. Wormald (2008), Distribution of subgraphs of random regular graphs, *Random Structures & Algorithms* **32**, 38–48.
- [23] S.M. Goodreau, J.A. Kitts and M. Morris (2009), Birds of a feather, or friend of a friend? Using exponential random graph models to investigate adolescent social networks, *Demography* **46**, 103–125.
- [24] C. Greenhill and M. Sfragara (2018), The switch Markov chain for sampling irregular graphs and digraphs, *Theoretical Computer Science* **719**, 1–20.
- [25] C. Greenhill (2021), Generating graphs randomly, in *Surveys in Combinatorics 2021*, London Mathematical Society Lecture Note Series **470**, Cambridge University Press, pp. 133–186.
- [26] V. Guruswami (2000), Rapidly mixing Markov chains: A comparison of techniques (a survey). Unpublished manuscript. [arXiv:1603.01512](https://arxiv.org/abs/1603.01512)
- [27] P. van der Hoorn, G. Lippner and E. Mossel (2022), Regular graphs with many triangles are structured, *Electronic Journal of Combinatorics* **29**, Art. P1.7.
- [28] S. L. Hakimi (1962), On realizability of a set of integers as degrees of the vertices of a linear graph. I, *SIAM Journal on Applied Mathematics* **10**, 496–506.
- [29] M. Hasheminezhad and B. McKay (2010), Combinatorial estimates by the switching method, *Contemporary Mathematics* **531**, 209–221.
- [30] W.K. Hastings (1970), Monte Carlo sampling methods using Markov chains and their applications, *Biometrika* **57**, 97–109.
- [31] V. Havel (1995), A remark on the existence of finite graphs (in Czech), *Časopis Pro Pěstování Matematiky* **80**, 477–480.
- [32] S. Janson (2010), Asymptotic equivalence and contiguity of some random graphs, *Random Structures & Algorithms* **36**, 26–45.
- [33] M. Jerrum, *Counting, Sampling and Integrating: Algorithms and Complexity*, Birkhäuser, Basel, 2003.
- [34] M. Jerrum and A. Sinclair (1989), Approximating the permanent, *SIAM J. Comput.* **18(6)**, 1149–1178.
- [35] E.M. Jin, M. Girvan and M.E.J. Newman (2001), Structure of growing social networks, *Physical Review E* **64**, 046132.
- [36] R. Kannan, P. Tetali and S. Vempala (1999), “Simple Markov chain algorithms for generating random bipartite graphs and tournaments”, *Random Structures & Algorithms* **14**, 293–308.
- [37] L. Le Cam (1960), An approximation theorem for the Poisson binomial distribution, *Pacific J. Mathematics* **10**, 1181–1197.

- [38] C. Lowcay, S. Marsland and C. McCartin (2013), Constrained switching in graphs: a constructive proof, in *2013 International Conference on Signal-Image Technology and Internet-Based Systems*, pp. 599–604.
- [39] I. Miklos, P. Erdős and L. Soukup (2013), “Towards random uniform sampling of bipartite graphs with given degree sequence”, *Electronic Journal of Combinatorics* **20**, #P16.
- [40] P. Mahlmann and C. Schindelbauer (2005), Peer-to-peer networks based on random transformations of connected regular undirected graphs, in *Proc. 17th Annual ACM Symposium on Parallelism in Algorithms and Architectures (SPAA 2005)*, pp. 155–164.
- [41] B. McKay (1985), Asymptotics for symmetric 0-1 matrices with prescribed row sums, *Ars Combin.* **19**, 15–25.
- [42] B. McKay (2010), Subgraphs of random graphs with specified degrees, *Proc. International Congress of Mathematicians (ICM 2010)*, pp. 2489–2501.
- [43] B. McKay, N. Wormald and B. Wysocka (2004), Short cycles in random regular graphs, *Electronic Journal of Combinatorics* **11**, #R66.
- [44] N. Nishimura (2018), Introduction to reconfiguration, *Algorithms* **11**, 52.
- [45] E. Pekoz, A. Rollin and N. Ross (2013), Total variation error bounds for geometric approximation, *Bernoulli* **19**, 610—632.
- [46] G. Reinert (2011), A short introduction to Stein’s method, Department of Statistics, University of Oxford.
- [47] R. Serfling (1978), Some elementary results on Poisson approximation in a sequence of Bernoulli trials, *SIAM Review* **20**, 567–579.
- [48] A. Sinclair (1992), Improved bounds for mixing rates of Markov chains and multi-commodity flows, *Combinatorics, Probability and Computing* **1**, 351–370.
- [49] J.M. Steele, Le Cam’s inequality and Poisson approximations, *American Mathematical Monthly* **101**, 48–54.
- [50] C. Stein, (1986), *Approximate computation of expectations*, Lecture Notes-Monograph Series vol. 7, Institute of Mathematical Statistics.
- [51] R. Taylor (1981), Constrained switchings in graphs, in *Combinatorial Mathematics VIII* (K.L. McAvaney, ed.), Springer Lecture Notes in Mathematics, vol. 884, pp. 314–336.
- [52] K. Tikhomirov and P. Youssef (2022), Sharp Poincaré and log-Sobolev inequalities for the switch chain on regular bipartite graphs, *Probability and Related Fields* **185**, to appear. Also [arXiv:2007.02729](https://arxiv.org/abs/2007.02729).
- [53] N. Wormald (2018), Asymptotic enumeration of graphs with given degree sequence, *Proc. International Congress of Mathematicians (ICM 2018)*, pp. 3245–3264