

Separating the edges of a graph by a linear number of paths

Marthe Bonamy¹ Fábio Botler² François Dross¹
Tássio Naia^{1,3} Jozef Skokan⁴

February 20, 2023

¹ Univ. Bordeaux, CNRS, Bordeaux INP, LaBRI, UMR 5800, F-33400 Talence, France,
E-mail: {marthe.bonamy|francois.dross}@u-bordeaux.fr

² Programa de Engenharia de Sistemas e Computação
Instituto Alberto Luiz Coimbra de Pós-Graduação e Pesquisa em Engenharia
Universidade Federal do Rio de Janeiro, Brasil
E-mail: fbotler@cos.ufrj.br

³ Departamento de Ciência da Computação
Instituto de Matemática e Estatística
Universidade de São Paulo, Brasil
E-mail: tnaia@member.fsf.org

⁴ Department of Mathematics
London School of Economics and Political Science (LSE)
Houghton Street, London, WC2A 2AE, United Kingdom
E-mail: j.skokan@lse.ac.uk

Abstract

Recently, Letzter proved that any graph of order n contains a collection \mathcal{P} of $O(n \log^* n)$ paths with the following property: for all distinct edges e and f there exists a path in \mathcal{P} which contains e but not f . We improve this upper bound to $19n$, thus answering a question of Katona and confirming a conjecture independently posed by Balogh, Csaba, Martin, and Pluhár and by Falgas-Ravry, Kittipassorn, Korándi, Letzter, and Narayanan. Our proof is elementary and self-contained.

A path *separates* an edge e from an edge f if it contains the former and not the latter. How many paths do we need to separate any edge from any other?

This question was first asked by Katona in 2013 (see [6]), in line with the general study of separating systems initiated by Rényi in 1961 [10]. Note that the roles of e and f are not symmetric. There are in fact two variants of the problem, which we clarify now. A collection \mathcal{P} of paths in a graph G is a *strongly-separating path system* of G if, for any two distinct edges e and f of G , there exist paths P_e and P_f in \mathcal{P} such that $e \in P_e$, $e \notin P_f$

and $f \in P_f$, $f \notin P_e$. The collection \mathcal{P} is a *weakly-separating path system* of G if for any two distinct edges e and f , there is a path $P \in \mathcal{P}$ that contains one and not the other. The *size* of a (weakly or strongly) separating path system is the number of paths in the collection. We recommend the excellent introduction in [7] for a more complete description of the history of the problem. We are interested in the following conjecture.

Conjecture 1 ([1, 6]). *Every graph on n vertices admits a strongly-separating path system of size $O(n)$.*

This conjecture was formulated independently by Falgas-Ravry, Kittipassorn, Korándi, Letzter, and Narayanan [6] for weakly-separating path systems, and later strengthened by Balogh, Csaba, Martin, and Pluhár [1] to strongly-separating path systems. Very recently, Letzter [7] made major progress toward Conjecture 1 by proving that every graph on n vertices admits a strongly-separating system with $O(n \cdot \log^* n)$ paths. A key component in [7] is the use of sublinear expanders, motivated by the recent work of Bucić and Montgomery [3]. In this note we obtain a linear bound, thus confirming Conjecture 1.

Theorem 2. *Every graph on n vertices admits a strongly-separating path system of size $19n$.*

Our proof uses Pósa rotation–extension tools [9] as presented by Brandt, Broersma, Diestel, and Kriesell [2], and is inspired by Lemma 2.2 in Conlon, Fox, and Sudakov [4]. In essence, we reduce the general problem to *traceable graphs*, i.e., graphs with a path that spans all vertices, known as Hamiltonian path.

Before delving into technical arguments, we point out that the upper bound of $19n$ is likely far from optimal. Balogh et al. [1, Theorem 10] showed that the complete bipartite graph $K_{\varepsilon n, (1-\varepsilon)n}$ cannot be strongly separated by fewer than $2(1 - 2\varepsilon)n$ paths. However, we are not aware of any graph with a worse lower bound. Therefore, the following might be true.

Problem 3. *Does every graph on n vertices admit a strongly-separating path system of size $2n$?*

Determining the exact constant is challenging even for cliques [11]. In our initial attempts to solve Conjecture 1, we stumbled upon this question:

Problem 4. *Does every properly edge-colored graph G admit $O(n)$ rainbow paths that cover every edge of G ?*

In this context, a rainbow path is a path that does not contain two edges of the same color. To the best of our knowledge, this question has not been stated anywhere else. We pose this question here, since we believe it to be natural and interesting in its own right. Problem 4, if answered affirmatively, would yield an alternative proof of Conjecture 1. We remark that we can verify a version of Problem 4 where “paths” is replaced by “trails”.




Figure 1: a path (highlighted) obtained by an elementary exchange fixing v .

Proof of Theorem 2

From now on, every separating path system is a strongly-separating path system. We use the following standard notation. Given a graph G , and a set $S \subseteq V(G)$, we denote by $N_G(S)$ the set of vertices *not* in S adjacent in G to some vertex in S . We omit subscripts when clear from the context.

Pósa rotation-extension. Given a graph G and vertices u, v in G , let $P = u \cdots v$ be a path from u to v . If $x \in V(P)$ is a neighbor of u in G and x^- is the vertex preceding x in P , then $P' = P - xx^- + ux$ is a path in G for which $V(P') = V(P)$. We say that P' has been obtained from P by an *elementary exchange* fixing v (see Figure 1). A path obtained from P by a (possibly empty) sequence of elementary exchanges fixing v is said to be a path *derived* from P . The set of endvertices of paths derived from P distinct from v is denoted by $S_v(P)$. Since all paths derived from P have the same vertex set as P , we have $S_v(P) \subseteq V(P)$. When P is a longest path ending at v , we obtain the following.

Lemma 5 ([2]). *Let $P = u \cdots v$ be a longest path of a graph G and let $S = S_v(P)$. Then $|N_G(S)| \leq 2|S|$.*

Proof rephrased from [2, Lemma 2.6]. Since P has maximum degree 2, it suffices to show that $N_G(S) \subseteq N_P(S)$. Let $x \in S$ and $y \in N_G(x) \setminus S$ be given. As $x \in S$ there is a longest path $Q = x \cdots v$ derived from P . Note that, by the maximality of Q , we have $y \in V(Q)$. Let z denote the predecessor of y in Q , and note that $Q + xy - yz$ is obtained from Q by an elementary exchange. Now, consider the first edge e incident to y that is removed by an elementary exchange in the “derivation sequence” from P to $Q + xy - yz$, and note that $e \in E(P)$. Note that when an elementary exchange removes an edge ab of a path derived from P , then a or b belongs to S . Therefore one of the vertices of e is in S . Since $y \notin S$ and $e \in E(P)$, we obtain $y \in N_P(S)$, as desired. \square

Our argument requires a lemma stating that every n -vertex graph admits an edge-cover by $O(n)$ paths. While elementary arguments yield a cover by at most $6n$ paths (as we discuss at the end of the paper), we optimize the multiplicative constant in Theorem 2 using a result of Dean and Kouider [5] that strengthens a well-known result of Lovász [8].

Theorem 6 ([5]). *Every graph G contains $2|V(G)|/3$ edge-disjoint paths covering $E(G)$.*

Let H and G be two (not necessarily disjoint) graphs. We say that a set of paths \mathcal{P} *separates* H from G if for each pair $(e, f) \in E(H) \times E(G)$ with $e \neq f$ there is a path in \mathcal{P} that contains e and does not contain f . Now, we are able to prove our result.




Figure 2: Left: a set S in a traceable graph and its neighborhood $N(S)$; a Hamiltonian path is highlighted, dashed red edges are the edges in P_S . Right: paths that separate subgraphs of G , where $A \xrightarrow{\alpha} B$ indicates that α separates A from B (for instance, \mathcal{Q}' separates G' from H').

Proof of Theorem 2. We proceed by induction on n . Let G be a graph with n vertices. If G is empty, the result trivially holds. If not, we consider P and S as in Lemma 5. Let H be the subgraph of G induced by the edges incident to vertices of S , and note that $n' = |V(H)| \leq 3|S|$. Let $G' = G \setminus S$. Note that, since G is not empty, S is not empty. By the induction hypothesis, G' admits a separating path system \mathcal{Q}' of size $19(n - |S|)$. Since \mathcal{Q}' covers G' , \mathcal{Q}' separates G' from H . In what follows, we construct a set of paths \mathcal{P} that separates H from G . Moreover, we obtain \mathcal{P} so that $|\mathcal{P}| \leq 19|S|$, and hence $\mathcal{Q}' \cup \mathcal{P}$ is a separating path system of G with cardinality at most $19(n - |S|) + 19|S| = 19n$ as desired.

First, let $P_S = E(P) \cap E(H)$ be the set of edges of P having at least one vertex in S (see Figure 2), and let \mathcal{P}_S be the set of paths each consisting of a single edge in P_S . Now, let \mathcal{D} be the path decomposition of $H' = H \setminus P_S$ given by Theorem 6. Note that $\mathcal{P}_S \cup \mathcal{D}$ has size at most $2|S| + 2n'/3 \leq 4|S|$ and separates (i) H from G' ; (ii) H' from P ; and (iii) P_S from G . It remains to create a set of at most $5n' \leq 15|S|$ paths that separates H' from itself (see Figure 2).

Let us write $P = u_1 \cdots u_m$. For $i \in [n'] = \{1, \dots, n'\}$, we define $\sigma(i)$ to be the index of the i^{th} vertex $u_{\sigma(i)}$ of H in P . For every such i , we let $v_i = u_{\sigma(i)}$. In what follows, each edge $v_i v_j$ is written so that $i < j$. We define $5n'$ sets of edges as follows: For $k \in [2n']$ let $M_k = \{v_i v_j \in E(H') : i + j = k\}$, and for $k \in [3n']$ let $N_k = \{v_i v_j \in E(H') : i + 2j = k\}$. In fact this is a bit less than $5n'$, especially if M_k and N_k are only defined for meaningful values. However, as far as we can tell this only leads to marginal improvements. Note that $\mathcal{M} = \{M_1, \dots, M_{2n'}\}$ (resp. $\mathcal{N} = \{N_1, \dots, N_{3n'}\}$) partitions $E(H')$.

We observe that for any k , the edges in M_k are pairwise comparable (similarly for N_k), in the sense that given $v_i v_j, v_{i'} v_{j'} \in M_k$ with $i > i'$ we have $j' > j$, and in fact $j' > j > i > i'$. Moreover, the set M_k is a subset of $\{v_1 v_{k-1}, \dots, v_{\frac{k}{2}-1} v_{\frac{k}{2}+1}\}$, with the last indices tweaked to fit the parity of k . In the same manner $N_k \subseteq \{v_1 v_{\frac{k-1}{2}}, \dots, v_{\frac{k}{3}-2} v_{\frac{k}{3}+1}\}$, again with tweaks




Figure 3: Set of comparable edges M_k (dashed) and a highlighted path P_k joining them.

depending on the value of $k \bmod 6$. Now, we find, for each $k \in [2n']$, a path P_k using all edges of M_k and some edges in P (see Figure 3). Let $M_k = \{x_1y_1, \dots, x_sy_s\}$ where the x_i 's are chosen in increasing order on P (hence the y_i 's in decreasing order). By induction on i from s down to 1, there is a path R_i in $P_i \cup \{x_iy_i, \dots, x_sy_s\}$ that contains x_iy_i, \dots, x_sy_s and ends in either x_i or y_i , where P_i denotes the subpath of P between x_i and y_i . Therefore, $P_k = R_1$ is the desired path. Analogously, for $k \in [3n']$, we obtain a path Q_k that contains N_k and some edges of P .

We claim that the collection of P_k 's and Q_k 's together separates each pair of edges in H' . Indeed, let v_iv_j and $v_{i'}v_{j'}$ be two edges of H' . Each such edge belongs to exactly one M_k and exactly one N_k . If they belong to different M_k 's, they belong to different P_k 's and are separated. Similarly if they belong to different N_k 's. Therefore, we may assume that $i + j = i' + j'$ and $i + 2j = i' + 2j'$. This immediately yields $j = j'$ and $i = i'$, as desired. Therefore, any two distinct edges in H' are separated by this collection, which involves $5n'$ paths. This concludes the proof. \square

Discussion and further work

As mentioned in the introduction, the best possible constant factor in Theorem 2 might well be 2. Here we obtain 19, which can still be reduced a little with some care¹. While partial progress would only be of limited interest, we highlight here straightforward candidates for improvement. First, any improvement of the ratio $\frac{n'}{|S|}$ (currently 3) would immediately significantly improve the factor. Second, with the arguments in this paper we can prove that the edges of any traceable graph with n vertices can be separated by $(6 + 2/3)n$ paths. In fact, any improvement of that could be plugged in the proof and used to improve the bound. In contrast, we observe that Theorem 6 is tight for disjoint triangles.

We also note that by using the M_k 's together with Lemma 5, we can prove that the edges of any graph can be covered by $6n$ paths without using Theorem 6. This allows us to verify Conjecture 1 in a self-contained manner.

As a final remark, we observe that robust sublinear expanders, which play a key role in [7], have also been applied in [3] to investigate the well-known Erdős–Gallai conjecture, that says that the edges of any n vertex graph can be partitioned into $O(n)$ edges and cycles. It is natural to wonder if our approach might be useful in tackling this conjecture as well.

¹For example, only applying Theorem 6 once for the whole graph, for all edges not in any P , would give an improvement of 1.3333. We decided against implementing this and other small improvements, as they do not seem worth the loss of clarity and simplicity.

Acknowledgments

The first author expresses gratitude to Bhargav Narayanan for sharing the question with her at the Oberwolfach Workshop id:2201, and to the organizers and participants of that workshop for creating a stimulating environment. We also thank Matija Bucić, Victor Falgas-Ravry and Lyuben Lichev for insightful comments which enriched this manuscript.

This research has been partially supported by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil – CAPES – Finance Code 001. M. Bonamy is supported by ANR project DISTANCIA (Metric Graph Theory, ANR-17-CE40-0015). F. Botler is supported by CNPq (423395/2018-1), FAPERJ (211.305/2019 and 201.334/2022) and CAPES-PRINT (88887.695773/2022-00). T. Naia is supported by CNPq (201114/2014-3) and FAPESP (2019/04375-5, 2019/13364-7, 2020/16570-4). FAPERJ and FAPESP are, respectively, Research Foundations of Rio de Janeiro and São Paulo. CNPq is the National Council for Scientific and Technological Development of Brazil.

References

- [1] J. Balogh, B. Csaba, R. R. Martin, and A. Pluhár, *On the path separation number of graphs*, Discrete Appl. Math. **213** (2016), 26–33.
- [2] S. Brandt, H. Broersma, R. Diestel, and M. Kriesell, *Global connectivity and expansion: long cycles and factors in f -connected graphs*, Combinatorica **26** (2006), no. 1, 17–36.
- [3] M. Bucić and R. Montgomery, *Towards the Erdős-Gallai cycle decomposition conjecture*, 2022, preprint, arXiv:2211.07689.
- [4] D. Conlon, J. Fox, and B. Sudakov, *Cycle packing*, Random Structures Algorithms **45** (2014), no. 4, 608–626.
- [5] N. Dean and M. Kouider, *Gallai’s conjecture for disconnected graphs*, Discrete Math. **213** (2000), no. 1-3, 43–54.
- [6] V. Falgas-Ravry, T. Kittipassorn, D. Korándi, S. Letzter, and B. P. Narayanan, *Separating path systems*, J. Comb. **5** (2014), no. 3, 335–354.
- [7] S. Letzter, *Separating paths systems of almost linear size*, 2022, preprint, arXiv:2211.07732.
- [8] L. Lovász, *On covering of graphs*, Theory of Graphs (Proc. Colloq., Tihany, 1966), Academic Press, New York, 1968, pp. 231–236.
- [9] L. Pósa, *Hamiltonian circuits in random graphs*, Discrete Math. **14** (1976), no. 4, 359–364.
- [10] A. Rényi, *On random generating elements of a finite boolean algebra*, Acta Sci. Math. Szeged **22** (1961), no. 75-81, 4.
- [11] B. Wickes, *Separating path systems for the complete graph*, 2022, preprint, arXiv:2209.04302.