

On left legal semigroups¹

Attila Nagy²

Department of Algebra
Budapest University of Technology and Economics
Műgyetem rkp. 3, Budapest, 1111, Hungary

Abstract

In this paper we study semigroups satisfying the identity $aba = ab$.

1 Introduction

Many important combinatorial structures such as real and complex hyperplane arrangements, interval greedoids, matroids and oriented matroids have the structure of a left regular band (see, for example, [1, 8, 10, 21]). A band is a semigroup in which every element is an idempotent element, and a left regular band is a band which satisfies the identity $aba = ab$. Left regular bands are examined by many authors. See, for example, papers [2–4, 8, 21, 22, 24, 26] and the book [16]. In [19, 20], the authors examined semigroups which are not necessarily bands, but satisfy the identity $aba = ab$. Using the terminology of [20], such semigroups are called left legal semigroups. The class of all left legal semigroups is a variety. Therefore, it is natural to find free semigroups and subdirectly irreducible semigroups in this variety. In Section 3, we construct free left legal semigroups F_X for arbitrary nonempty sets X . The subdirectly irreducible left legal semigroups are examined in Section 4. We show that a semigroup containing at least two elements is a subdirectly irreducible left legal semigroup satisfying the identity $ab = a^2b$ if and only if it is either a subdirectly irreducible left regular band or a two-element zero semigroup. We also show that a semigroup containing at least two elements is a subdirectly irreducible left legal semigroup satisfying the identity $ab = a^2$ if and only if it is either a two-element left zero semigroup or a two-element zero semigroup. The concept of the retract ideal extension of semigroups is used effectively in many cases in the study of the structure of semigroups. In Section 4, we give a necessary and sufficient condition for a left legal semigroup to be a retract ideal extension of a left regular band by a zero semigroup. One of the basic concepts of semigroup theory is the semilattice decomposition. In Section 5, this concept is in the centre of investigations. We show that a left legal semigroup is semilattice indecomposable if and only if it satisfies the identity $a^2b = a^2$. The left (resp., right, weakly) separative semigroups are examined in many papers. At the end of Section 5, we prove theorems on left (resp., right, weakly) separative left legal semigroups. We show that a left legal semigroup is right separative if and only if it is weakly separative. It is also proved that every left legal left separative

¹Mathematics Subject Classification: 20M10, 20M12

Keywords: Semigroup, free semigroup, variety of semigroups

²email: nagyat@math.bme.hu

semigroup is commutative. In Section 6, the lattice of all left legal semigroup varieties is examined.

2 Preliminaries

By a *semigroup* we mean a multiplicative semigroup, that is, a nonempty set together with an associative multiplication. A nonempty subset I of a semigroup S is said to be an *ideal* of S if $sa, as \in I$ for every $a \in I$ and $s \in S$. If I is an ideal of a semigroup S , then the relation ϱ_I on S defined by $(a, b) \in \varrho_I$ if and only if $a = b$ or $a, b \in I$ is a congruence on S , which is called the *Rees congruence on S determined by I* . The equivalence classes of $S \bmod \varrho_I$ are I itself and every one-element set $\{a\}$ with $a \in S \setminus I$. The factor semigroup S/ϱ_I is called the *Rees factor semigroup of S modulo I* . We shall write S/I instead of S/ϱ_I . We may describe S/I as the result of collapsing I into a single (zero) element, while the elements of S outside of I retain their identity. A semigroup S containing a zero element 0 is called a *zero semigroup* if $ab = 0$ for every $a, b \in S$. It is obvious that if S is a semigroup, then S^2 is an ideal of S , and the Rees factor semigroup S/S^2 is a zero semigroup. Let B and Q be disjoint semigroups, Q having a zero element. A semigroup S is called an *ideal extension* of B by Q if S contains B as an ideal, and if the Rees factor semigroup S/B is isomorphic with Q . An ideal I of a semigroup S is called a *retract ideal* if there is a homomorphism of S onto I which leaves the elements of I fixed. Such a homomorphism is called a *retract homomorphism* of S onto I . In this case we say that S is a *retract ideal extension* of I .

We say that a semigroup S is a *subdirect product* of semigroups S_i ($i \in I$) if S is isomorphic to a subsemigroup T of the direct product of semigroups S_i ($i \in I$) such that the restriction of the projection homomorphisms to T are surjective. A semigroup S is said to be *subdirectly irreducible* if whenever S is written as a subdirect product of a family of semigroups S_i ($i \in I$), then, for at least one $j \in I$, the projection homomorphism π_j maps S onto S_j isomorphically. Subdirect decompositions of a semigroup S are closely connected with congruences on S (see, for example, [16, I.3.6]). If α_i ($i \in I$) are congruences on a semigroup S and $\bigcap_{i \in I} \alpha_i = \iota_S$, the equality relation on S , then S is a subdirect product of the factor semigroups S/α_i . Conversely, if a semigroup is a subdirect product of semigroups S_i ($i \in I$) and α_i is the congruence on S induced by the projection homomorphism π_i ($i \in I$), then $\bigcap_{i \in I} \alpha_i = \iota_S$.

If \mathcal{C} is a class of semigroups, then a congruence ϱ on a semigroup S is called a *\mathcal{C} -congruence* if the factor semigroup S/ϱ belongs to \mathcal{C} . An element e of a semigroup is called an *idempotent element* if $e^2 = e$. A semigroup in which every element is an idempotent element is called a *band*. A commutative band is called a *semilattice*. It is clear that the universal relation on an arbitrary semigroup is a semilattice congruence. A semigroup S is said to be *semilattice indecomposable* if the universal relation is the only semilattice congruence on S . By Theorem of [23], every semigroup S has a least semilattice congruence η_S whose classes are semilattice indecomposable semigroups. In other words,

every semigroup is a semilattice of semilattice indecomposable semigroups. A band satisfying the identity $aba = ab$ is called a *left regular band*. A semigroup satisfying the identity $ab = a$ is called a *left zero semigroup*. By the dual of [16, II.3.12. Proposition], a band is left regular if and only if it is a semilattice of left zero semigroups. It is clear that every left zero semigroup is left legal and semilattice indecomposable.

A semigroup S is called a left legal semigroup if it satisfies the identity $aba = ab$. Every left regular band is a left legal semigroup. The converse is also true for semigroups S containing a right identity element, because $a^2 = aea = ae = a$ is satisfied for every element a of S and a right identity element e of S .

A semigroup S is called an \mathcal{L} -commutative semigroup [13] if, for every elements $a, b \in S$, there is an element $x \in S^1$ such that $ab = xba$. A semigroup S is called a *right weakly commutative semigroup* if, for every $a, b \in S$, there exists $x \in S$ and a positive integer n such that $(ab)^n = xa$. It is clear that every \mathcal{L} -commutative semigroup is right weakly commutative.

A semigroup S is said to be a *right separative semigroup* if $ab = b^2$ and $ba = a^2$ imply $a = b$ for every $a, b \in S$. A *left separative semigroup* is defined analogously. A semigroup S is called a *weakly separative semigroup* if $a^2 = ab = b^2$ implies $a = b$ for every $a, b \in S$.

For an arbitrary semigroup S ,

$$S^1 = \begin{cases} S, & \text{if } S \text{ has an identity element;} \\ S \cup 1, & \text{otherwise,} \end{cases}$$

where $S \cup 1$ is the semigroup which is obtained by the adjunction of an identity element 1 to S .

For notions and notations not defined but used in this paper, we refer the reader to [5].

3 Free left legal semigroups

Let \mathcal{V} be a non-trivial variety of algebras, and X be a nonempty set. An algebra F_X is said to be a free algebra in \mathcal{V} with a free generating system (or basis) X if F_X is in \mathcal{V} , X generates F_X , and every mapping of X into any algebra A from \mathcal{V} can be extended to a homomorphism of F_X into A . In this section we construct free left legal semigroups F_X for arbitrary nonempty sets X . First we prove two lemmas which will be used throughout the paper.

Lemma 1. *If S is a left legal semigroup then every element of S^2 is an idempotent element. Moreover, $a^k = a^2$ for every $a \in S$ and every integer $k \geq 2$.*

Proof. Let S be a left legal semigroup. For every elements a and b of S , we have $(ab)^2 = a(bab) = aba = ab$. Thus every element of S^2 is an idempotent element. For every $a \in S$, $a^3 = a^2$ by definition, and hence $a^k = a^2$ for every integer $k \geq 2$. \square

Lemma 2. *For arbitrary elements a and b of a left legal semigroup, $ab = ab^2$.*

Proof. Using Lemma 1, we get $ab = (ab)^2 = (aba)b = ab^2$. \square

Let X be a nonempty set (whose elements are also said to be letters). Let F_X denote the set of all finite nonempty sequences (or words) of letters of X in which each letter $x \in X$ can occur at most once or twice, in the latter case in the form xx at the beginning of the word. For example, if x and y are elements of X with $x \neq y$, then $xx, xxy \in F_X$ but $xyx, yxx \notin F_X$. If ω_1 and ω_2 are words of F_X , then let $\exists_{\omega_1}(\omega_2)$ denote the word which can be obtained from ω_2 in the following way: for every $x \in X$, if x occurs in both of ω_1 and ω_2 , then we delete x from ω_2 . For example, $\exists_{ux}(xxy) = y$. We note that $\exists_{\omega_1}(\omega_2)$ may be the empty word. For an arbitrary $\omega \in F_X$, let $\omega^* = \omega$, if the letters of ω are pairwise distinct, and let ω^* be the word which can be obtained from ω by deleting its second letter if the first two letters of ω are equal. For example, if $x, y, z \in X$ are pairwise distinct, then $(xyz)^* = xyz = (xxyz)^*$. We note that if $\exists_{\omega_1}(\omega_2^*)$ is not empty, then its letters are pairwise distinct. Thus $(\omega^*)^* = \omega^*$ for every $\omega \in F_X$.

We define an operation \circ on F_X . For arbitrary words $\omega_1 = x_1 \dots x_n$ and $\omega_2 = y_1 \dots y_m$ of F_X , let

$$\omega_1 \circ \omega_2 = \begin{cases} xx\exists_{\omega_1}(\omega_2^*), & \text{if } \omega_1 = x \text{ and } y_1 = x \text{ for some } x \in X, \\ \omega_1\exists_{\omega_1}(\omega_2^*), & \text{otherwise.} \end{cases} \quad (1)$$

Remark 1. It is clear that, for arbitrary nonempty set X and arbitrary words ω_1 and ω_2 of F_X , the length of $\omega_1 \circ \omega_2$ is greater than 1.

Remark 2. The word $xx\exists_{\omega_1}(\omega_2^*)$ in the first row of (1) can also be written in the form $\omega_1 x \exists_{\omega_1}(\omega_2^*)$. Thus, for every $\omega_1, \omega_2 \in F_X$, the word ω_1 is a prefix of the word $\omega_1 \circ \omega_2$.

Example 1. If $X = \{x, y\}$, then $F_X = \{x, xx, y, yy, xy, xxy, yx, yyx\}$. The operation \circ on F_X is given by Table 1.

\circ	x	xx	y	yy	xy	xxy	yx	yyx
x	xx	xx	xy	xy	xxy	xxy	xy	xy
xx	xx	xx	xxy	xxy	xxy	xxy	xxy	xxy
y	yx	yx	yy	yy	yx	yx	yyx	yyx
yy	yyx	yyx	yy	yy	yyx	yyx	yyx	yyx
xy	xy	xy	xy	xy	xy	xy	xy	xy
xxy	xxy	xxy	xxy	xxy	xxy	xxy	xxy	xxy
yx	yx	yx	yx	yx	yx	yx	yx	yx
yyx	yyx	yyx	yyx	yyx	yyx	yyx	yyx	yyx

Table 1:

Theorem 3. For an arbitrary nonempty set X , the operation \circ is associative on F_X .

Proof. (By induction on the length of the middle word) Let $\omega_1 = x_1 \dots x_n$ and $\omega_2 = y_1 \dots y_m$ be arbitrary words of F_X . Let $x \in X$ be an arbitrary letter. First consider the case $x \notin \omega_1$. Then

$$(\omega_1 \circ x) \circ \omega_2 = \omega_1 x \exists_{\omega_1 x} (\omega_2^*) = \omega_1 \circ (x \circ \omega_2).$$

Next consider the case when $x \in \omega_1$.

If $\omega_1 = x$, then

$$(\omega_1 \circ x) \circ \omega_2 = xx \exists_{\omega_1} (\omega_2^*) = \omega_1 \circ (x \circ \omega_2).$$

If $\omega_1 \neq x$, then

$$(\omega_1 \circ x) \circ \omega_2 = \omega_1 \circ \omega_2 = \omega_1 \circ (x \circ \omega_2).$$

Let n be a positive integer greater than 1. Assume that

$$(\omega_1 \circ \omega_2) \circ \omega_3 = \omega_1 \circ (\omega_2 \circ \omega_3)$$

is satisfied for every words $\omega_1, \omega_2, \omega_3 \in F_X$ where the length of ω_2 is less than n . Let ω_2 be a word of F_X whose length is n . Denote the last letter of ω_2 by x , that is, $\omega_2 = \omega x = \omega \circ x$ ($\omega \in F_X, x \in X$). Then, for arbitrary $\omega_1, \omega_3 \in F_X$,

$$\begin{aligned} (\omega_1 \circ \omega_2) \circ \omega_3 &= (\omega_1 \circ (\omega \circ x)) \circ \omega_3 = ((\omega_1 \circ \omega) \circ x) \circ \omega_3 = (\omega_1 \circ \omega)(x \circ \omega_3) = \\ &= \omega_1 \circ (\omega \circ (x \circ \omega_3)) = \omega_1 \circ ((\omega \circ x) \circ \omega_3) = \omega_1 \circ (\omega_2 \circ \omega_3). \end{aligned}$$

Thus the operation \circ is associative, and hence F_X is a semigroup with respect to the operation \circ . \square

Theorem 4. For an arbitrary nonempty set X , the semigroup F_X is left legal.

Proof. Let ω_1 and ω_2 be arbitrary elements of F_X . By Remark 1, the length of $\omega_1 \circ \omega_2$ is greater than 1. By Remark 2, ω_1 is a prefix of $\omega_1 \circ \omega_2$. Thus

$$\omega_1 \circ \omega_2 \circ \omega_1 = \omega_1 \circ \omega_2 \exists_{(\omega_1 \circ \omega_2)} (\omega_1^*) = \omega_1 \circ \omega_2.$$

\square

Theorem 5. If X is an arbitrary nonempty set, then F_X is a free left legal semigroup with a free generating system X .

Proof. Let X be an arbitrary nonempty set. By Theorem 4, the semigroup F_X is left legal. Let S be an arbitrary left legal semigroup and f be an arbitrary mapping of X into S . Let Φ be the mapping of the semigroup F_X into the semigroup S defined by the usual way: for arbitrary $\omega = x_1 \dots x_n \in F_X$, let

$$\Phi(x_1 \dots x_n) = f(x_1) \cdots f(x_n).$$

We show that Φ is a homomorphism. Let $\omega_1 = x_1 \dots x_n$ and $\omega_2 = y_1 \dots y_m$ be arbitrary elements of F_X . First consider the case when $n = 1$ and $\omega_1 = x = y_1$ for some $x \in X$. If $y_2 \neq x$, then

$$\Phi(\omega_1 \circ \omega_2) = \Phi(xxy_2 \dots y_m) = f(x)f(x)f(y_2) \dots f(y_m) = \Phi(\omega_1)\Phi(\omega_2).$$

If $y_2 = x$, then (using also Lemma 1)

$$\begin{aligned} \Phi(\omega_1 \circ \omega_2) &= \Phi(xxy_3 \dots y_m) = f(x)f(x)f(y_3) \dots f(y_m) = \\ &= f(x)f(x)f(x)f(y_3) \dots f(y_m) = \Phi(\omega_1)\Phi(\omega_2). \end{aligned}$$

Next consider the opposite case: $\omega_1 = x$ for some $x \in X$ and $y_1 \neq x$ or $n > 1$. In this case $y_1 \in \omega_1$ implies $n > 1$, and hence

$$f(x_1) \dots f(x_n) = f(x_1) \dots f(x_n)f(y_1), \quad (2)$$

because S satisfies the identities $ab = aba$ and $ab = ab^2$. By the definition of the operation \circ ,

$$\omega_1 \circ \omega_2 = \omega_1 \exists_{\omega_1}(\omega_2^*). \quad (3)$$

We have two subcases. First assume that $\exists_{\omega_1}(\omega_2^*)$ is the empty word. Then $y_j \in \omega_1$ for every $j \in \{1, \dots, m\}$. Thus (2) holds, and hence

$$\begin{aligned} f(x_1) \dots f(x_n) &= f(x_1) \dots f(x_n)f(y_1) = \\ &= f(x_1) \dots f(x_n)f(y_1) \dots f(y_m), \end{aligned}$$

because S satisfies the identity $ab = aba$, and every factor $f(y_j)$ ($j = 1, \dots, m$) equals some factor $f(x_t)$ ($t = 1, \dots, n$). Next, assume that $\exists_{\omega_1}(\omega_2^*)$ is not the empty word. Then there are pairwise distinct letters z_1, \dots, z_k such that every z_i equals some y_j , and

$$\exists_{\omega_1}(\omega_2^*) = z_1 z_2 \dots z_k. \quad (4)$$

Using (3) and (4), we have

$$\Phi(\omega_1 \circ \omega_2) = f(x_1) \dots f(x_n)f(z_1) \dots f(z_k). \quad (5)$$

If $y_1 = z_1$, then we can complete the product $f(x_1) \dots f(x_n)f(y_1)f(z_2) \dots f(z_k)$ with the missing factors $f(y_j)$ ($y_j \in \omega_1$) to get the equation

$$f(x_1) \dots f(x_n)f(y_1)f(z_2) \dots f(z_k) = f(x_1) \dots f(x_n)f(y_1)f(y_2) \dots f(y_m). \quad (6)$$

Equations (5) and (6) together imply that $\Phi(\omega_1 \circ \omega_2) = \Phi(\omega_1)\Phi(\omega_2)$.

If $y_1 \neq z_1$, then $y_1 \in \omega_1$, and hence (2) holds. Thus

$$\begin{aligned} f(x_1) \dots f(x_n)f(z_1) \dots f(z_k) &= \\ &= f(x_1) \dots f(x_n)f(y_1)f(z_1) \dots f(z_k) = f(x_1) \dots f(x_n)f(y_1)f(y_2) \dots f(y_m), \end{aligned}$$

because we can complete the product $f(x_1) \dots f(x_n)f(y_1)f(z_1) \dots f(z_k)$ with the missing factors. Hence $\Phi(\omega_1 \circ \omega_2) = \Phi(\omega_1)\Phi(\omega_2)$.

Thus Φ is a homomorphism. It is obvious that $\Phi(x) = f(x)$ for every $x \in X$. \square

4 Retract ideal extensions of left regular bands by zero semigroups

By Lemma 1, if S is a left legal semigroup, then every element of S^2 is an idempotent element. Thus every left legal semigroup is an ideal extension of a left regular band by a zero semigroup. In this section we focus on the case when this extension is retract.

Lemma 6. *If S is a semigroup such that S^2 is a band and φ is a retract homomorphism of S onto S^2 , then $\varphi(a) = a^2$ for every $a \in S$.*

Proof. Let S be a semigroup such that S^2 is a band. Let φ be a retract homomorphism of S onto S^2 . Then, for arbitrary $a \in S$,

$$\varphi(a) = (\varphi(a))^2 = \varphi(a^2) = a^2.$$

□

Theorem 7. *The following four conditions on a semigroup S are equivalent.*

- (i) S is a left legal semigroup satisfying the identity $ab = a^2b$.
- (ii) S is a retract ideal extension of a left regular band by a zero semigroup.
- (iii) S is a subdirect product of a left regular band and a zero semigroup.
- (iv) S is a left legal semigroup such that the factor semigroup S/θ_S^{left} is a band.

Proof. It is obvious that (i) and (iv) are equivalent.

(i) \Rightarrow (ii): Assume that S is a left legal semigroup satisfying the identity $ab = a^2b$. By Lemma 1, S is an ideal extension of the left regular band S^2 by the zero semigroup S/S^2 . Let φ be the mapping of S into S^2 defined by the following way: for an arbitrary $a \in S$, $\varphi(a) = a^2$. As S^2 is a band, φ maps S onto S^2 and leaves the elements of S^2 fixed. Let $a, b \in S$ be arbitrary elements. Since S is a left legal semigroup and satisfies the identity $ab = a^2b$, we have

$$\begin{aligned} \varphi(a)\varphi(b) &= a^2b^2 = a(ab)b = a(aba)b = a(ab)^2 = \\ &= a(ab) = a^2b = ab = (ab)^2 = \varphi(ab). \end{aligned}$$

Then φ is a homomorphism. Consequently S is a retract ideal extension of the left regular band S^2 by the zero semigroup S/S^2 . Thus (ii) is satisfied.

(ii) \Rightarrow (iii): Assume that S is a retract ideal extension of a left regular band B by a zero semigroup. Then B is an ideal of S such that the Rees factor semigroup S/B is a zero semigroup. Since B is a band, $B \subseteq S^2$. Since the Rees factor semigroup S/B is a zero semigroup, $S^2 \subseteq B$. Thus $B = S^2$. Let φ be a retract homomorphism of S onto S^2 . It is easy to see that $\ker \varphi \cap \varrho_{S^2} = \iota_S$, where $\ker \varphi$ denotes the kernel of φ , ϱ_{S^2} denotes the Rees congruence on S determined by the ideal S^2 . Obviously, the factor semigroup $S/\ker \varphi$ is isomorphic to the

left regular band S^2 and the Rees factor semigroup S/S^2 is a zero semigroup. Thus S is a subdirect product of a left regular band and a zero semigroup.

(iii) \Rightarrow (i): It is obvious, because every subsemigroup of the direct product of a left regular band and a zero semigroup is a left legal semigroup satisfying the identity $ab = a^2b$. \square

Example 2. On the set $S = \{x, e, f, g, h\}$ consider the operation defined by Table 2.

	x	e	f	g	h
x	e	e	e	g	h
e	e	e	e	g	h
f	f	f	f	g	h
g	g	g	g	g	g
h	h	h	h	h	h

Table 2:

It is a matter of checking to see that this operation is associative. $S^2 = \{e, f, g, h\}$ is a semilattice of the left zero semigroups $S_1 = \{e, f\}$ and $S_2 = \{g, h\}$, and hence S^2 is a left regular band. The mapping $\varphi : s \mapsto s^2$ is a retract homomorphism of S onto S^2 . The Rees factor semigroup $S/S^2 = \{x, 0\}$ is a two-element zero semigroup. By Theorem 7, S is a left legal semigroup satisfying the identity $ab = a^2b$. S is a subdirect product of the left regular band S^2 and the two-element zero semigroup S/S^2 .

Theorem 8. *A semigroup containing at least two elements is a subdirectly irreducible left legal semigroup satisfying the identity $ab = a^2b$ if and only if it is either a subdirectly irreducible left regular band or a two-element zero semigroup.*

Proof. A left regular band is a left legal semigroup and satisfies the identity $ab = a^2b$. It is easy to see that if Z is a two-element zero semigroup, then Z is a subdirectly irreducible left legal semigroup satisfying the identity $ab = a^2b$. To prove the converse assertion, assume that S is a subdirectly irreducible left legal semigroup which has at least two elements and satisfies the identity $ab = a^2b$. Then, by (iii) of Theorem 7, S is either a subdirectly irreducible left regular band or a subdirectly irreducible zero semigroup Z . Since a zero semigroup is subdirectly irreducible if and only if it has at most two elements, we have $|Z| = 2$. \square

For subdirectly irreducible left regular bands, we refer the reader to [26, Theorem 2.16].

Theorem 9. *The following three conditions on a semigroup S are equivalent.*

- (i) S is a left legal semigroup satisfying the identity $ab = a^2$.
- (ii) S is a retract ideal extension of a left zero semigroup by a zero semigroup.

(iii) S is a subdirect product of a left zero semigroup and a zero semigroup.

Proof. (i) \Rightarrow (ii): Let S be a left legal semigroup satisfying the identity $ab = a^2$. Since S is left legal, S^2 is a band by Lemma 1. For arbitrary elements $e, f \in S^2$, we have

$$ef = e^2 = e.$$

Hence S^2 is a left zero semigroup. Let φ be the mapping of S into S^2 defined by the following way: for an arbitrary $a \in S$, $\varphi(a) = a^2$. As S^2 is a band, φ maps S onto S^2 which leaves the elements of S^2 fixed. For arbitrary $a, b \in S$,

$$\varphi(a)\varphi(b) = a^2b^2 = (ab)(ba) = ab^2a = ab^2 = (ab)b = (aba)b = (ab)^2 = \varphi(ab).$$

Thus φ is a homomorphism. Consequently S is a retract ideal extension of the left zero semigroup S^2 by the zero semigroup S/S^2 , and hence (ii) is satisfied.

(ii) \Rightarrow (iii): Assume that S is a retract ideal extension of a left zero semigroup B by a zero semigroup. Then B is an ideal of S , and the Rees factor semigroup S/B is a zero semigroup. Since B is a band, $B \subseteq S^2$. Since S/B is a zero semigroup, $S^2 \subseteq B$. Thus $B = S^2$. Let φ be a retract homomorphism of S onto S^2 . Then $\ker \varphi \cap \varrho_{S^2} = \iota_S$. Thus S is a subdirect product of the left regular band S^2 and the zero semigroup S/S^2 .

(iii) \Rightarrow (i): It is obvious, because every subsemigroup of the direct product of a left zero semigroup and a zero semigroup is a left legal semigroup satisfying the identity $ab = a^2$. \square

Example 3. On the set $S = \{x, y, b, c\}$ consider the operation defined by Table 3.

	x	y	b	c
x	b	b	b	b
y	c	c	c	c
b	b	b	b	b
c	c	c	c	c

Table 3:

It is a matter of checking to see that S is a left legal semigroup satisfying the identity $ab = a^2$. S is an ideal extension of the left zero semigroup $S^2 = \{b, c\}$ by the zero semigroup $S/S^2 = \{x, y, 0\}$. The mapping $\varphi: s \mapsto s^2$ is a retract homomorphism of S onto S^2 .

Theorem 10. A semigroup containing at least two elements is a subdirectly irreducible left legal semigroup satisfying the identity $ab = a^2$ if and only if it is either a two-element left zero semigroup or a two-element zero semigroup.

Proof. It is obvious that every two-element semigroup is subdirectly irreducible. Moreover, every left zero semigroup and every zero semigroup is left legal and satisfies the identity $ab = a^2$. To prove the converse assertion, let S be a

subdirectly irreducible left legal semigroup which has at least two elements and satisfies the identity $ab = a^2$. Then, by (iii) of Theorem 9, S is either a left zero semigroup or a zero semigroup. Since a left zero semigroup and, similarly, a zero semigroup is subdirectly irreducible if and only if it contains at most two elements, S is either a two-element left zero semigroup or a two-element zero semigroup. \square

5 The least semilattice congruence on left legal semigroups

In the study of the structure of semigroups, the semilattice decomposition plays a basic role. A semigroup S is said to be an *archimedean semigroup* if, for arbitrary $a, b \in S$, there are positive integers m and n such that $a^m \in SbS$ and $b^n \in SaS$. Every archimedean semigroup is semilattice indecomposable. A semigroup S is called a *Putcha semigroup* if, for every $a, b \in S$, the assumption $a \in S^1bS^2$ implies $a^m \in S^1b^2S^1$ for some positive integer m . It is proved in [18] that a semigroup is a semilattice of archimedean semigroups if and only if it is a Putcha semigroup.

Theorem 11. *Every left legal semigroup is a Putcha semigroup.*

Proof. Let S be a left legal semigroup. It is clear that S is \mathcal{L} -commutative, and hence right weakly commutative. By [14, Theorem 4.1] and [14, Lemma 2.2], every right weakly commutative semigroup is a Putcha semigroup, which proves our assertion. \square

Remark 3. *Theorem 11 and [14, Theorem 2.1] together imply that if S is a left legal semigroup, then,*

$$\eta_S = \{(a, b) \in S \times S : a^m \in SbS, b^n \in SaS \text{ for some positive integers } m, n\},$$

where η_S denotes the least semilattice congruence on S . This implies that a left legal semigroup is semilattice indecomposable if and only if it is archimedean.

For an arbitrary semigroup S , let τ_S denote the binary relations on S defined by the following way: $(a, b) \in \tau_S$ for elements $a, b \in S$, if and only if there is a positive integer n such that $a^n b = a^{n+1}$ and $b^n a = b^{n+1}$. By [14, Lemma 4.1], τ_S and its dual σ_S are equivalence relations on an arbitrary semigroup S . Equivalence relations τ_S and σ_S are studied by many authors in special classes of semigroups. In the book [5], Theorem 4.14 asserts that if S is a commutative semigroup, then $\tau_S = \sigma_S$ is a least weakly separative congruence on S . Chrislock shown [6, Theorem 5.4] that if S is a medial semigroup, then τ_S is the least left separative congruence on S , and σ_S is the least right separative congruence on S . Mukherjee proved [11, Theorem 6] that σ_S is a weakly separative congruence on a quasicommutative semigroup S . It is also conjectured that σ_S is the least weakly separative congruence. Pondelicek proved [17, Theorem 5] that if S is a duo semigroup, then $\tau_S \cap \sigma_S$ is the least weakly separative congruence on S .

Since every quasicommutative semigroup is a duo semigroup, this result implies that $\tau_S \cap \sigma_S$ is the least weakly separative congruence on a quasicommutative semigroup, giving a negative answer for Mukherjee's conjecture. Generalizing and completing the above results, in [12, Theorem 1] it is proved that $\tau_S \cap \sigma_S$ is the least weakly separative congruence on a weakly commutative semigroup S . Using this result, in [15] the weakly separative weakly commutative semigroups are characterized.

Since every left legal semigroup is right weakly commutative, [14, Theorem 4.7] implies that τ_S is a weakly separative congruence on a left legal semigroup. The next theorem gives an additional result about τ_S if S is a left legal semigroup.

Theorem 12. *If S is a left legal semigroup, then τ_S is the least semilattice congruence on S .*

Proof. We use Remark 3. First we show that $\eta_S \subseteq \tau_S$. Assume $(a, b) \in \eta_S$ for elements $a, b \in S$. Then, by Remark 3, $a^n = xby$ and $b^n = uav$ for a positive integer n and for some elements $x, y, u, v \in S$. We can suppose $n \geq 2$. Then $a^n b = (xby)b = x(byb) = xby = a^n = a^{n+1}$, because $a^k = a^2$ for every integer $k \geq 2$. Similarly, $b^n a = b^{n+1}$. Thus $(a, b) \in \tau_S$. Hence $\eta_S \subseteq \tau_S$. To show $\tau_S \subseteq \eta_S$, assume $(a, b) \in \tau_S$ for elements $a, b \in S$. Then $a^n b = a^{n+1}$ and $b^n a = b^{n+1}$ for some positive integer $n \geq 2$. Thus $a^{n+2} = a^n b a \in SbS$ and $b^{n+2} = b^n a b \in SaS$. Hence $(a, b) \in \eta_S$. Thus $\tau_S \subseteq \eta_S$. Consequently $\eta_S = \tau_S$. \square

Remark 4. *By Lemma 1, if a is an arbitrary element of a left legal semigroup, then $a^k = a^2$ is satisfied for every integer $k \geq 2$. Thus $(a, b) \in \tau_S$ is satisfied for elements a and b of a left legal semigroup if and only if $a^2 b = a^2$ and $b^2 a = b^2$.*

Theorem 13. *The following two conditions on a left legal semigroup S are equivalent.*

- (i) S is semilattice indecomposable.
- (ii) S satisfies the identity $a^2 b = a^2$.

Proof. A semigroup S is semilattice indecomposable if and only if η_S is the universal relation on S . By Theorem 12, $\eta_S = \tau_S$ on a left legal semigroup S . Thus a left legal semigroup S is semilattice indecomposable if and only if τ_S is the universal relation on S . By Remark 4, if S is a left legal semigroup, then $(a, b) \in \tau_S$ if and only if $a^2 b = a^2$ and $b^2 a = b^2$. Consequently a left legal semigroup is semilattice indecomposable if and only if it satisfies the identity $a^2 b = a^2$. \square

Corollary 14. *If a semigroup is a retract ideal extension of a left zero semigroup by a zero semigroup, then it is a semilattice indecomposable left legal semigroup.*

Proof. Let S be a semigroup which is a retract ideal extension of a left zero semigroup by a zero semigroup. By Theorem 9, S is a left legal semigroup satisfying the identity $ab = a^2$. Since $a^3 = a^2$ for every $a \in S$, the semigroup S satisfies the identity $a^2b = a^2$. Then S is semilattice indecomposable by Theorem 13. \square

By Theorem 13 or Corollary 14, the semigroup S defined in Example 3 is a semilattice indecomposable left legal semigroup.

At the end of the section we prove two theorems on left (resp., right, weakly) separative left legal semigroups.

Theorem 15. *The following two conditions on a left legal semigroup S are equivalent.*

(i) S is right separative.

(ii) S is weakly separative.

Proof. (i) \Rightarrow (ii): Let S be a right separative left legal semigroup. Assume $a^2 = ab = b^2$ for elements $a, b \in S$. From $ab = b^2$ we get

$$bab = b^3 = b^2,$$

and hence

$$ba = bab = b^2 = a^2.$$

Then

$$ab = b^2 \quad \text{and} \quad ba = a^2.$$

Since S is right separative, we get $a = b$. Consequently S is weakly separative.

(ii) \Rightarrow (i): Let S be a weakly separative left legal semigroup. Assume $ab = b^2$ and $ba = a^2$ for elements $a, b \in S$. Using the first equation, we get

$$ba = bab = b^3 = b^2,$$

and hence

$$a^2 = ba = b^2.$$

Since S is weakly separative, we get $a = b$. Consequently S is right separative. \square

Remark 5. *It is clear that every left zero semigroup containing at least two elements is left legal and weakly separative, but not left separative. Thus the weakly separativity of a left legal semigroup S does not imply the left separativity of S , in general.*

Theorem 16. *Every left separative left legal semigroup is commutative.*

Proof. Let S be a left separative left legal semigroup. Then, for every $a, b \in S$,

$$(ab)(ba) = (aba)(bab) = (ab)^3 = (ab)^2.$$

Similarly,

$$(ba)(ab) = (ba)^2.$$

Since S is left separative, we get $ab = ba$, and hence S is commutative. \square

6 On the lattice of all left legal semigroup varieties

The class of all varieties of semigroups forms a lattice under the following operations: for varieties \mathcal{X} and \mathcal{Y} , their join $\mathcal{X} \vee \mathcal{Y}$ is the variety generated by the set-theoretical union of \mathcal{X} and \mathcal{Y} (as classes of semigroups), and their meet $\mathcal{X} \wedge \mathcal{Y}$ coincides with the set-theoretical intersection of \mathcal{X} and \mathcal{Y} . A variety \mathcal{V} of semigroups is said to be a left legal semigroup variety, if every semigroup belonging to \mathcal{V} is left legal. The set of all left legal semigroup varieties form a sublattice of the lattice of all semigroup varieties. This sublattice denoted by $\mathcal{L}(\mathbf{LLS})$. Let \mathcal{A} (resp., \mathcal{B}) denote the class of all semigroups that are retract extensions of a left regular band (resp., left zero semigroup) by a zero semigroup. By Theorem 7 and Theorem 9, \mathcal{A} and \mathcal{B} are subvarieties of the variety \mathcal{LLS} . Using Lemma 2, $ab = a^2$ implies $ab = ab^2 = a^2b$ for arbitrary elements a and b of a left legal semigroup. Then $\mathcal{B} \subseteq \mathcal{A}$. We also use the following notations.

\mathcal{LRB} : the variety of all left regular bans ($a^2 = a, aba = ab$);

\mathcal{LNB} : the variety of all left normal bands ($a^2 = a, axy = ayx$);

\mathcal{SL} : the variety of all semilattices ($a^2 = a, ab = ba$);

\mathcal{LZ} : the variety of all left zero semigroups ($ab = a$);

\mathcal{ZM} : the variety of all zero semigroups ($ab = 0$);

\mathcal{T} : the trivial variety ($a = b$).

It is clear that $\mathcal{ZM} \subseteq \mathcal{B}$, $\mathcal{LZ} \subseteq \mathcal{B} \cap \mathcal{LNB}$, and $\mathcal{SL} \subseteq \mathcal{LNB} \subseteq \mathcal{LRB} \subseteq \mathcal{A}$. Since $\mathcal{A} \subseteq \mathcal{LLS}$, all of the above varieties are in the lattice $\mathcal{L}(\mathbf{LLS})$.

An element a of a lattice $(L; \vee, \wedge)$ with a zero 0 is said to be an atom of L if $0 < a$ and $0 \leq b < a$ implies $0 = b$ for arbitrary $b \in L$.

An element x of a lattice $(L; \vee, \wedge)$ is called a neutral element of L if, for all $y, z \in L$, the elements x, y and z generate a distributive sublattice of L .

Theorem 17. *The varieties \mathcal{SL} , \mathcal{LZ} , \mathcal{ZM} are atoms, the varieties \mathcal{T} , \mathcal{SL} , \mathcal{ZM} , $\mathcal{SL} \vee \mathcal{ZM}$, and \mathcal{LLS} are neutral elements of the lattice $\mathcal{L}(\mathbf{LLS})$.*

Proof. By [9, Theorem 2.6], \mathcal{SL} , \mathcal{LZ} and \mathcal{ZM} are atoms of the lattice of all semigroup varieties (see also the Theorem of Section IV of [7]). Thus they are atoms of the lattice $\mathcal{L}(\mathbf{LLS})$. It is clear that the varieties \mathcal{T} and \mathcal{LLS} are neutral elements of the lattice $\mathcal{L}(\mathbf{LLS})$. By [25, Proposition 4.1], varieties \mathcal{SL} , \mathcal{ZM} , and $\mathcal{SL} \vee \mathcal{ZM}$ are neutral elements of the lattice of all semigroup varieties. Thus they are neutral elements of the lattice $\mathcal{L}(\mathbf{LLS})$. \square

The relative position of the above-mentioned varieties in the lattice $\mathcal{L}(\mathbf{LLS})$ is illustrated in Figure 1. $\mathcal{X} \rightarrow \mathcal{Y}$ denotes that the variety \mathcal{X} is a subvariety of the variety \mathcal{Y} .




Figure 1:

References

- [1] Aguiar, M and Mahajan, S.: Topics in hyperplane arrangements, Mathematical Surveys and Monographs **226**, American Mathematical Society, Providence, RI, 2017.
- [2] Auinger, K., Gomes, G.M.S., Gould, V. and Steinberg, B.: An Application of a Theorem of Ash to Finite Covers. *Studia Logica* **78**, 45–57 (2004). <https://doi.org/10.1007/s11225-005-5057-6>
- [3] Billhardt, B., Marques-Smith, P. and Martins, P.M.: Extensions of left regular bands by R-unipotent semigroups. *Periodica Mathematica Hungarica* **83**, 260–271 (2021). <https://doi.org/10.1007/s10998-021-00384-z>
- [4] Branco, M.J.J., Gomes, G.M.S., Gould, V.: Extensions and covers for semigroups whose idempotents form a left regular band. *Semigroup Forum* **81-1**, 51–70 (2010). <https://doi.org/10.1007/s00233-010-9239-9>
- [5] Clifford, A.H. and Preston, G.B.: The Algebraic Theory of Semigroups I. Amer. Math. Soc. Providence R.I. (1961). <https://doi.org/10.1090/surv/007.1>
- [6] Chrislock, J.L.: On medial semigroups. *Journal of Algebra* **12**, 1–9 (1969). [https://doi.org/10.1016/0021-8693\(69\)90013-1](https://doi.org/10.1016/0021-8693(69)90013-1)
- [7] Evans, T.: The lattice of semigroup varieties. *Semigroup Forum* **2-1**, 1–43 (1971). [https://doi.org/10.1016/S1385-7258\(55\)50089-8](https://doi.org/10.1016/S1385-7258(55)50089-8)
- [8] Friedman, M. and Garber, D.: On Left regular bands and real Conic-Line arrangements. *Semigroup Forum* **98**, 83–139 (2019). <https://doi.org/10.1007/s00233-018-9966-x>

- [9] Kalicki, J. and Scott, D., Equationally completeness in abstract algebras. Proc. Koninkl. Nederl. Akad. Wetensch. Ser. A, **58**, 650–659 (1955).
- [10] Margolis, S.W., Saliola, F. and Steinberg, B.: Cell complexes, poset topology and the representation theory of algebras arising in algebraic combinatorics and discrete geometry. *Memoirs of the American Mathematical Society* **274**, pp 135 (2021).
- [11] Mukherjee, N.P.: Quasicommutative semigroups I. *Czechoslovak Mathematical Journal* **22-3**, 449–453 (1972).
<https://doi.org/10.21136/CMJ.1972.101114>
- [12] Nagy, A.: The least separative congruence on a weakly commutative semigroup. *Czechoslovak Mathematical Journal* **32**, 630–632 (1982).
<https://doi.org/10.21136/CMJ.1982.101840>
- [13] Nagy, A.: \mathcal{RC} -commutative Δ -semigroups. *Semigroup Forum* **44**, 332–340 (1992). <https://doi.org/10.1007/BF02574352>
- [14] Nagy, A.: *Special Classes of Semigroups*. Kluwer Academic Publishers, Dordrecht/Boston/London (2001).
<https://doi.org/10.1007/978-1-4757-3316-7>
- [15] Nagy, A.: Weakly separative weakly commutative semigroups. *Semigroup Forum* **89**, 475–478 (2014).
<https://doi.org/10.1007/s00233-014-9581-4>
- [16] Petrich, M.: *Lectures in Semigroups*. Akademie-Verlag-Berlin (1977).
<https://doi.org/10.1515/9783112471326>
- [17] Pondelicek, B.: On weakly commutative semigroups. *Czechoslovak Mathematical Journal* **25-1**, 20–23 (1975).
<https://doi.org/10.21136/CMJ.1975.101289>
- [18] Putcha, M.S.: Semilattice decomposition of semigroups. *Semigroup Forum* **6**, 12–34 (1973). <https://doi.org/10.1007/BF02389104>
- [19] Salam, A., Ashraf, W. and Khan, N.M.: Congruences, Green’s relations, translational hull and legal semigroups. *The Aligarh Bulletin of Mathematics* **36-1,2**, 95–107 (2017).
- [20] Salam, A., Ashraf, W. and Khan, N.M.: On sandwich sets in legal semigroups. *Southeast Asian Bulletin of Mathematics* **46-1**, 45–55 (2022).
- [21] Saliola, F.V.: The quiver of the semigroup algebra of a left regular band. *International Journal of Algebra and Computation* **17-8** 1593–1610 (2007).
<https://doi.org/10.1142/S0218196707004219>
- [22] Shevlyakov, A.S.: Finite Submonoids of Free Left Regular Bands with Unit. *Algebra and Logic* **59**, 456–470 (2021).
<https://doi.org/10.1007/s10469-021-09619-x>

- [23] Tamura, T.: Note on the greatest semilattice decomposition of semigroups. *Semigroup Forum* **4**, 255–261 (1972).
<https://doi.org/10.1007/BF02570795>
- [24] Vakhrameev, M.A.: Edit distance in free left-regular bands. *Journal of Physics: Conf. Series* **1210** (2019).
<https://doi.org/10.1088/1742-6596/1210/1/012153012153>
- [25] Volkov, M. V.: Modular elements of the lattice of semigroup varieties. *Contrib. General Algebra* **16**, 275–288 (2005).
- [26] Zheng-Pan Wang, Jing Leng, Hou-Yi Yu, On subdirectly irreducible regular bands. *Turkish Journal of Mathematics* **41-5**, 1337–1343 (2017).
<https://doi.org/10.3906/mat-1609-12>