

ON TOTALLY LINDELÖF SPACES

GABRIEL FERNANDES, GUILHERME PINTO, AND VINÍCIUS ROCHA[†]

ABSTRACT. We prove that every Alster space is totally Lindelöf and this gives a new characterization of regular Alster spaces. We construct a totally Lindelöf space that is not Alster and we prove that there exists a P -space that is not Frolik. This answers three questions asked by Noble in [Nob19].

1. INTRODUCTION

Totally Lindelöf is a property of topological spaces that was defined by Vaughan in order to unify the proof that countable products of P -spaces and countable products of σ -compact spaces are Lindelöf (see [Vau78] and Definition 1.12). Alster spaces were defined by Alster to characterize productively Lindelöf spaces (see [Als88] and Definition 1.16).

Since 2009 there has been a vigorous effort to understand Alster spaces and related spaces and their relationship to the class of productively Lindelöf spaces, see [Nob19], [TT11], [Tal11b], [Tal11a], [Tal13], [ST09], [BT12], [MZ14] and [AZ18]. Our main result is the following new characterization of Alster spaces:

Theorem 1.1. *If X is a regular topological space, then X is an Alster space if and only if X is totally Lindelöf*

That regular totally Lindelöf spaces are Alster spaces is due to Noble in [Nob19], we prove the other direction. Regarding totally Lindelöf spaces, Noble asked in [Nob19, Section 5] the following questions:

- (a) Is there an example of an Alster space which is not totally Lindelöf?
- (b) Is there an example of a totally Lindelöf (not regular) space which is not Alster?
- (c) What is the relationship of Alster/totally Lindelöf spaces and Frolik spaces?

We will study totally Lindelöf spaces and filter bases and answer the three questions above. The paper is organized as follows:

Below, in subsection 1.1, we establish our notation and main definitions, in Section 2 we prove properties of filter bases and Theorem 1.1, which answers question (a) negatively.

In Section 3 we answer question (b) positively:

Theorem 1.2. *There is a non regular space X that is a Hausdorff, Menger, totally Lindelöf space and is not Alster.*

In Section 4 we study the ω_1 discrete set property:

Definition 1.3. If X is a topological space and there is $\{x_\alpha \mid \alpha < \omega_1\} \subseteq X$ such that for all $\beta < \omega_1$ we have that $\{x_\alpha \mid \alpha < \beta\}$ is a discrete set. Then we say that X has the ω_1 discrete set property.

2020 *Mathematics Subject Classification.* 54D20, 54D10, 54D45.

Key words and phrases. filter bases, totally Lindelöf, Alster spaces, Frolik spaces.

[†] The author is funded by CAPES-88887.605742/2021-00.

Our main results in Section 4 are Theorem 1.4 and Corollary 1.5, they together with Theorem 1.1 answer question (c):

Theorem 1.4. *Let $\{X_n \mid n < \omega\}$ be a sequence of σ -compact spaces. Then the product space $\prod_{n < \omega} X_n$ does not have the ω_1 discrete set property.*

Corollary 1.5. *There is a P -space that is not Frolik, namely $L(\omega_1)$ is not Frolik.*

1.1. Notation and preliminaries . We denote by $L(\omega_1)$ the topological space $(\omega_1 + 1, \tau)$ such that $[\omega_1]^{<\omega} \subseteq \tau$ and $B_{\omega_1} = \{] \alpha, \omega_1 + 1[\mid \alpha < \omega_1\}$ is a local base for ω_1 where $] \alpha, \omega_1 + 1[= \{\beta \in \omega_1 \mid \alpha < \beta \leq \omega_1 + 1\}$.

Definition 1.6. If X is a set and $\mathcal{F} \subseteq \mathcal{P}(X)$.

- (1) We say that \mathcal{F} is a filter base on X if for all $F_0, F_1 \in \mathcal{F}$ we have that $F_0 \cap F_1 \in \mathcal{F}$ and $\emptyset \notin \mathcal{F}$.
- (2) If \mathcal{H} and \mathcal{F} are filter bases on X , we say that \mathcal{H} extends \mathcal{F} if $\mathcal{F} \subseteq \mathcal{H}$.

The definitions of totally Lindelöf presented in [Nob19] and [Vau78] look different at first sight. In [Nob19] the definition of totally Lindelöf uses the concept of δ -stable and in [Vau78] it uses the concept of stable under countable intersections. We will introduce these both notions and show in Proposition 1.15 that they are interchangeable in the definition of totally Lindelöf.

Definition 1.7. If X is a set and \mathcal{F} is a filter base on X .

- (a) We say that \mathcal{F} is stable under countable intersections if for all $S \in [\mathcal{F}]^{\leq \omega}$ there is $H \in \mathcal{F}$ such that $H \subseteq \bigcap S$.
- (b) We say that \mathcal{F} is ω -closed if for all $S \in [\mathcal{F}]^{\leq \omega}$ we have that $\bigcap S \in \mathcal{F}$.
- (c) We say that \mathcal{F} is δ -stable if for all $S \in [\mathcal{F}]^{\leq \omega}$ we have that $\bigcap S \neq \emptyset$.

Definition 1.8. If $\mathcal{F} \subseteq \mathcal{P}(X)$ and (X, τ) is a topological space we define $ad^{(X, \tau)}(\mathcal{F}) = \bigcap \{F \mid F \in \mathcal{F}\}$

Remark 1.9. When there is no risky of ambiguity we shall write $ad(\mathcal{F})$ instead of $ad^{(X, \tau)}(\mathcal{F})$. Occasionally we may also write $ad^\tau(\mathcal{F})$ or $ad^X(\mathcal{F})$ instead of $ad^{(X, \tau)}(\mathcal{F})$.

Definition 1.10. If \mathcal{F} is a filter base on a topological space X , we say that \mathcal{F} is total if and only if for every filter base $\mathcal{H} \supseteq \mathcal{F}$ we have that $ad(\mathcal{H}) \neq \emptyset$.

Remark 1.11. Notice that the definition of stable under countable intersections filter base on a set X does not depend on which topology we consider on X , nevertheless whether a filter base \mathcal{F} on X is total, it will depend on the topology that we are considering, since being total depends on whether $ad^X(\mathcal{G}) \neq \emptyset$ for all \mathcal{G} that extends \mathcal{F} .

Definition 1.12. (totally Lindelöf) We say that a topological space X is totally Lindelöf if and only if for every stable under countable intersections filter base \mathcal{F} on X there is $\mathcal{H} \supseteq \mathcal{F}$ a total stable under countable intersections filter base on X .

Definition 1.13. If X is a topological space we denote by G_δ^X the collection of all countable intersections of open subsets of X . Given $S \subseteq X$ we define the following filter:

$$G_\delta(S) = \{G \subseteq X \mid S \subseteq G \wedge G \in G_\delta^X\}$$

As mentioned above totally Lindelöf is a property that was isolated in order to unify the proof that countable products of P -spaces and countable products of σ -compact spaces remain Lindelöf. The reader interested in gaining more intuition on this definition should see [Vau78]. Nonetheless the proof of the following proposition that will be used later may help the reader gaining some intuition on the definition of totally Lindelöf.

Proposition 1.14. $L(\omega_1)$ is totally Lindelöf

Proof. Let \mathcal{F} be a stable under countable intersections filter base on $L(\omega_1)$. We first consider the uninteresting case when $\bigcap \mathcal{F} \neq \emptyset$.

Case 1: $\bigcap \mathcal{F} \neq \emptyset$.

In this case, we fix $p \in \bigcap \mathcal{F}$ and consider the stable under countable intersections filter base $\mathcal{H} = \mathcal{F} \cup \{\{p\}\}$.

It follows that if $\mathcal{G} \supseteq \mathcal{H}$ and \mathcal{G} is a filter base, then $\{p\} \subseteq G$ for all $G \in \mathcal{G}$, hence $ad(\mathcal{G}) \supseteq \bigcap \mathcal{G} \supseteq \{p\} \neq \emptyset$. Thus \mathcal{H} is total.

Case 2: $\bigcap \mathcal{F} = \emptyset$

We will prove that in this case \mathcal{F} is total. Given a countable set $G \subseteq \omega_1 + 1$, fixing, for each $\alpha < \omega_1$, $F_\alpha \in \mathcal{F}$ such that $\alpha \notin F_\alpha$ and using the definition of stable under countable intersections, we can fix $F \in \mathcal{F}$ such that $F \subseteq \bigcap_{\alpha \in G} F_\alpha$. Therefore if $\mathcal{G} \supseteq \mathcal{F}$ is a filter base, it follows that for all $G \in \mathcal{G}$ we have that $sup(G \cap \omega_1) = \omega_1$, therefore for all $G \in \mathcal{G}$ we have that $\omega_1 \in \overline{G}$. Thus $ad(\mathcal{G}) \supseteq \{\omega_1\} \neq \emptyset$. This proves that \mathcal{F} is total. \square

We observe that $L(\omega_1)$ is P -Lindelöf so one could avoid the computations in Proposition 1.14 applying Vaughan's result that every P -Lindelöf space is totally Lindelöf.

Our next two proposition addresses slight variations in the definition of totally Lindelöf:

Proposition 1.15. *Suppose X is a topological space. Then the following are equivalent:*

- (a) X is totally Lindelöf.
- (b) Every \mathcal{F} which is an ω -closed filter base on X has a extension \mathcal{H} that is a total ω -closed filter base on X .
- (c) Every \mathcal{F} which is a δ -stable filter base on X has an extension \mathcal{H} that is a total δ -stable filter base on X .

Proof. We start with (a) \Rightarrow (b). Suppose that \mathcal{F} is an ω -closed filter base on X . Then \mathcal{F} is a filter base on X that is stable under countable intersections and by (a) there is \mathcal{H} , a filter base on X such that $\mathcal{H} \supseteq \mathcal{F}$, it is stable under countable intersections and total. Let

$$\mathcal{G} = \{\bigcap S \mid S \in [\mathcal{H}]^{\leq \omega}\}$$

From the fact that \mathcal{H} is a filter base that is stable under countable intersections it follows that \mathcal{G} is an ω -closed filter base. From the fact that \mathcal{H} is total, it follows that \mathcal{G} is total.

(b) \Rightarrow (c) Suppose \mathcal{F} is a δ -stable filter base on X . Let

$$\mathcal{F}' = \{\bigcap S \mid S \in [\mathcal{F}]^{\leq \omega}\}$$

From the fact that \mathcal{F} is δ -stable, it follows that \mathcal{F}' is an ω -closed filter base. Let $\mathcal{H} \supseteq \mathcal{F}'$ be a total ω -closed filter base on X . Then \mathcal{H} is a total δ -stable filter base that extends \mathcal{F} .

(c) \Rightarrow (a) Suppose that \mathcal{F} is filter base that is stable under countable intersections. Then \mathcal{F} is a δ -stable filter base. By (c) there is $\mathcal{H} \supseteq \mathcal{F}$ which is a total δ -stable filter base. Then

$$\mathcal{G} = \{\bigcap S \mid S \in [\mathcal{H}]^{\leq \omega}\}$$

is a total stable under countable intersections filter base that extends \mathcal{F} . \square

For self-containment sake we introduce the definition of the properties of Alster and Frolik.

Definition 1.16. (Alster) If X is a topological space we say that \mathcal{G} is an Alster covering of X if $\mathcal{G} \subseteq G_\delta^X$ and for every $K \subseteq X$ compact subset of X there is $G \in \mathcal{G}$ such that $K \subseteq G$.

We say that X is an Alster space if for every \mathcal{G} that is an Alster covering of X there is $\{A_n \mid n \in \omega\} \subseteq \mathcal{G}$ such that $X = \bigcup_{n \in \omega} A_n$.

Definition 1.17. (Frolik) We say that X is a Frolik space if¹ there is a sequence $\langle S_n \mid n \in \omega \rangle$ such that for every $n \in \omega$ the space S_n is σ -compact and there is φ and F such that $\varphi : X \rightarrow F \subseteq \prod_{n \in \omega} S_n$ is a homeomorphism and F is a closed subset of the product space.

2. A CHARACTERIZATION OF REGULAR ALSTER SPACES

In this section we prove Theorem 1.1. We will start proving Lemma 2.2 which was originally stated in [Vau78] without a proof.

Definition 2.1. Let \mathcal{F} be a filter base on X and $W \subseteq X$. We say that \mathcal{F} converges to W if and only if for all open set U which contains W there exists $F \in \mathcal{F}$ such that $F \subseteq U$.

Lemma 2.2. ([Vau78, Lemma 5.1]) *If \mathcal{F} is a filter base that converges to a compact set, then \mathcal{F} is total.*

Proof. Let \mathcal{F} be a filter base that converges to a compact set K and \mathcal{H} be a filter base that extends \mathcal{F} . Towards a contradiction, suppose that $ad(\mathcal{H}) = \emptyset$. For each $p \in K$, let V_p be an open set such that for some $H_p \in \mathcal{H}$ we have that $H_p \cap V_p = \emptyset$. Since K is compact there is a finite cover $V_{p_0} \cup \dots \cup V_{p_n} \supseteq K$. Hence $H := H_{p_0} \cap \dots \cap H_{p_n} \in \mathcal{H}$ and

$$H \cap (V_{p_0} \cup \dots \cup V_{p_n}) = \emptyset$$

On the other hand, since \mathcal{F} converges to K there is $T \in \mathcal{H}$ such that $T \subseteq (V_{p_0} \cup \dots \cup V_{p_n})$, which implies that $H \cap T = \emptyset$, contradicting the fact that \mathcal{H} is a filter base. Hence $ad(\mathcal{H})$ is non-empty. \square

Proposition 2.3. *Suppose \mathcal{F} is a stable under countable intersections filter base on a topological space X and $K \subseteq X$ is a compact set. If*

$$\mathcal{F} \vee G_\delta(K) := \{F_0 \cap F_1 \mid F_0, F_1 \in \mathcal{F} \cup G_\delta(K)\}$$

is a filter base, then it is a total stable under countable intersections filter base.

Proof. If $\mathcal{F} \vee G_\delta(K)$ is a filter base, then it is a stable under countable intersections filter base. Given U an open set of X that contains K , we have that $U \in G_\delta(K)$. Therefore $\mathcal{F} \vee G_\delta(K)$ converges to K . Hence by Lemma 2.2 it follows that $\mathcal{F} \vee G_\delta(K)$ is total. \square

Theorem 2.4. *If X is an Alster space, then X is totally Lindelöf.*

Proof. Let \mathcal{F} be a stable under countable intersections filter base on X . By Proposition 2.3 in order to find a total stable under countable intersections filter base of \mathcal{F} it is enough to find $K \subseteq X$ compact such that $\mathcal{F} \vee G_\delta(K)$ is a filter base.

Suppose that for every $K \subseteq X$ compact we have that $\mathcal{F} \vee G_\delta(K)$ is not a filter base.

¹See [Fro63, Theorem 12] for a proof that Definition 1.17 is equivalent to the original definition.

Hence for each $K \subseteq X$ compact let $G_K \in G_\delta(K)$ and $F_K \in \mathcal{F}$ be such that $G_K \cap F_K = \emptyset$. We set

$$\mathcal{A} = \{G_K \mid K \subseteq X \wedge K \text{ is compact}\}$$

which is an Alster cover of X .

Since X is an Alster space, there is $\langle K_n \mid n \in \omega \rangle$, a countable sequence of compact sets of X , such that $\{G_{K_n} \mid n \in \omega\} \subseteq \mathcal{A}$ is a countable subcover of X . We have that $G_{K_n} \cap F_{K_n} = \emptyset$ for every $n \in \omega$, thus

$$\left(\bigcap_{n \in \omega} F_{K_n}\right) \cap \left(\bigcup_{n \in \omega} G_{K_n}\right) = \emptyset$$

But $X = \bigcup_{n \in \omega} G_{K_n}$. Since \mathcal{F} is stable under countable intersection there is $F \in \mathcal{F}$ such that $F \subseteq \bigcap_{n \in \omega} F_{K_n}$, which is a contradiction.

Thus there exists some $K \subseteq X$ compact such that $\mathcal{H} = \mathcal{F} \vee G_\delta(K)$ is a filter base and from Proposition 2.3 it is a total stable under countable intersections filter base. Therefore X is totally Lindelöf. \square

We end this section with results on filter bases, we will not need these results in the next sections.

In Proposition 2.6, below, we prove a sort of characterization of total stable under countable intersections filter bases that holds when we are restricted to regular totally Lindelöf spaces and which will complement Proposition 2.3.

Our proof of Proposition 2.6 relies on the following lemma due to Vaughan:

Lemma 2.5. [Vau78, Lemma 5.1] *Suppose that X is regular space. If \mathcal{F} is a total filter base on X , then there is a non empty compact set $K \subseteq X$ such that \mathcal{F} converges to K .*

Proposition 2.6. *Suppose X is a regular totally Lindelöf space, \mathcal{F} is a stable under countable intersections filter base on X . Then there is K a compact set such that*

$$\mathcal{F} \vee G_\delta(K) := \{F_0 \cap F_1 \mid F_0, F_1 \in \mathcal{F} \cup G_\delta(K)\}$$

is a total stable under countable intersections filter base which extends \mathcal{F} .

Proof. Let $\mathcal{H} \supseteq \mathcal{F}$ be a total stable under countable intersections filter base that extends \mathcal{F} . By Lemma 2.5 there is $K \subseteq X$ compact such that \mathcal{H} converges to K . We will verify that $\mathcal{F} \vee G_\delta(K)$ is a filter base and the proposition will follow from Proposition 2.3.

Let G be a G_δ set such that $G \supseteq K$ and let $F \in \mathcal{F}$.

Write $G = \bigcap_{n \in \omega} V_n$ where each V_n is an open set. Since \mathcal{H} converges to K , it follows that for every $n \in \omega$ there is $H_n \in \mathcal{H}$ such that $H_n \subseteq V_n$. Using that \mathcal{H} is stable under countable intersections there is $H \subseteq \bigcap_{n \in \omega} H_n \subseteq \bigcap_{n \in \omega} V_n = G$ such that $H \in \mathcal{H}$. From the fact that \mathcal{H} extends \mathcal{F} we have that $F \cap H \neq \emptyset$, thus $F \cap G \neq \emptyset$. This verifies that $\mathcal{F} \vee G_\delta(K)$ is a filter base. \square

Theorem 2.7. *Suppose that X is a Lindelöf space and \mathcal{F} is a stable under countable intersections filter base on X . Then \mathcal{F} converges to $ad(\mathcal{F})$*

Proof. Suppose that there is some open set U such that $ad(\mathcal{F}) \subseteq U$ and there is no $F \in \mathcal{F}$ such that $F \subseteq U$. Let $\mathcal{H} = \{F \setminus U \mid F \in \mathcal{F}\}$, and $\{F_n \mid n \in \omega\} \subseteq \mathcal{F}$. From the hypothesis that \mathcal{F} is stable under countable intersections we can fix $F \subseteq \bigcap_{n \in \omega} F_n$. Notice that

$$F \setminus U \subseteq \left(\bigcap_{n \in \omega} F_n\right) \setminus U = \bigcap_{n \in \omega} (F_n \setminus U)$$

This verifies that \mathcal{H} is stable under countable intersections. Since X is Lindelöf it follows that $ad(\mathcal{H}) \neq \emptyset$ and $ad(\mathcal{H}) \subseteq (X \setminus U)$. On the other hand $ad(\mathcal{H}) \subseteq ad(\mathcal{F}) \subseteq U$, a contradiction. \square

3. A TOTALLY LINDELÖF SPACE THAT IS NOT AN ALSTER SPACE

In this section we prove that the refinement of the usual topology of the real line which is obtained by taking complements of countable sets is a totally Lindelöf space that is not Alster.

Theorem 3.1. *There is a Hausdorff, totally Lindelöf space that is not Alster.*

Proof. Let (\mathbb{R}, τ) be the reals with the usual topology. We define

$$\rho := \{U \setminus E \mid U \in \tau_0 \wedge E \in [\mathbb{R}]^{\leq \omega}\}$$

We will verify that $X := (\mathbb{R}, \rho)$ is a totally Lindelöf space which is not an Alster space.

Let \mathcal{F} be a stable under countable intersections filter base on \mathbb{R} such that $\bigcap F = \emptyset$. Since \mathbb{R} is totally Lindelöf, there is $\mathcal{G} \supseteq \mathcal{F}$ a stable under countable intersections filter base that is total in (\mathbb{R}, τ) . We will prove that \mathcal{G} is also total in X . Let \mathcal{H} be a filter base \mathcal{H} that extends \mathcal{G} .

For each $H \in \mathcal{H}$, we define

$$V_H = \bigcup \{V \in \tau \mid |V \cap H| \leq \omega\}$$

Since the usual topology on the reals is hereditary Lindelöf it follows that V_H is Lindelöf. Therefore we can find $\mathcal{A} \subseteq \{V \in \tau_0 \mid |V \cap H| \leq \omega\}$ countable such that $\bigcup \mathcal{A} = V_H$. Hence $V_H \cap H$ is countable.

We set

$$\widehat{\mathcal{H}} = \{H \setminus V_H \mid H \in \mathcal{H}\} \cup \mathcal{H} \supseteq \mathcal{G}$$

Let us verify that $\widehat{\mathcal{H}}$ is a filter base. Given $H \in \mathcal{H}$ and $E \in [\mathbb{R}]^{\leq \omega}$, using that $\bigcap F = \emptyset$ we can fix for each $z \in E$ a set $F_z \in \mathcal{F}$ such that $z \notin F_z$. Since \mathcal{F} is stable under countable intersections there is $F' \in \mathcal{H}$ such that $F' \subseteq H \cap \bigcap_{z \in E} F_z$, therefore F' is disjoint from E . Setting $E = H \cap V_H$ it follows that for every $H \in \mathcal{H}$ there is $F' \in \mathcal{F}$ such that $F' \subseteq (H \setminus V_H) \subseteq H$. Thus since \mathcal{H} is a filter base it follows that $\widehat{\mathcal{H}}$ is a filter base.

We will prove that for every $H \in \mathcal{H}$ we have that $\overline{H \setminus V_H}^\rho = \overline{H \setminus V_H}^\tau$. Thus from the fact that \mathcal{G} is total in (\mathbb{R}, τ) it will follow that

$$\bigcap \{\overline{T}^\rho \mid T \in \widehat{\mathcal{H}}\} = \bigcap \{\overline{T}^\tau \mid T \in \widehat{\mathcal{H}}\} \neq \emptyset$$

Since \mathcal{H} is an arbitrary base filter extending \mathcal{G} , it will follow that \mathcal{G} is total in $X = (\mathbb{R}, \rho)$.

The inclusion \subseteq follows from the fact that ρ is a refinement of τ . So we are left with verifying the other inclusion. Let $x \in \overline{H \setminus V_H}^\tau$ and let V be an open set of (\mathbb{R}, τ) such that $x \in V$. If $|V \cap H| \leq \omega$, then $V \subseteq V_H$, contradicting the choice of $x \in \overline{H \setminus V_H}^\tau$. Thus for all H that is an open neighborhood of x we have $|V \cap H| \geq \omega_1$ and therefore for every $E \in [\mathbb{R}]^{\leq \omega}$ we have $|V \cap (H \setminus E)| \geq \omega_1 > 0$. In particular if $E = V_H \cap H$, we have that $|V \cap (H \setminus V_H)| \geq \omega_1$. Hence $x \in \overline{H \setminus V_H}^\rho$, which verifies that $\overline{H \setminus V_H}^\rho = \overline{H \setminus V_H}^\tau$. \square

The space (X, τ) constructed above, despite it being a non-regular space, is relatively well behaved. It is a Hausdorff space, as we have verified in Theorem 3.1, and we shall prove that it is a Menger space in Proposition 3.2 below.

Proposition 3.2. *The space (\mathbb{R}, τ) is a Menger space.*

Proof. Let $\langle A_n \mid n < \omega \rangle$ a sequence of open coverings of X . So for each $n < \omega$ and each $A \in A_n$ fix $V(A)$ an open on the usual topology and $E(A)$ a countable subset such that $A = V(A) \setminus E(A)$.

Since the usual topology of \mathbb{R} is a Menger space, fix a sequence $\langle \mathcal{B}_{k(n)} \mid n < \omega \rangle$ with $\mathcal{B}_n \subseteq A_n$ finite such that $\{V(A) \mid n < \omega \wedge A \in \mathcal{B}_{k(n)}\}$ is a covering of \mathbb{R} .

$$\mathbb{R} \setminus \bigcup \{V(A) \mid n < \omega \wedge A \in \mathcal{B}_n\} \subseteq \bigcup \{E(A) \mid n < \omega \wedge A \in \mathcal{B}_n\}$$

is countable.

Let $\pi : \omega \rightarrow \bigcup \{E(A) \mid n < \omega \wedge A \in \mathcal{B}_{k(n)}\}$ be a bijection. Since for every $m \in \omega$ we have that \mathcal{A}_m is a covering of X , we can fix for each $m \in \omega$ a set $V(A_m) \setminus E(A_m)$ in \mathcal{A}_m such that $\pi(m) \in V(A_m) \setminus E(A_m)$. Thus


$$\mathcal{C}_m = \{V(A) \setminus E(A) \mid A \in \mathcal{B}_m\} \cup \{V(A_m) \setminus E(A_m)\}$$

We have that each \mathcal{C}_m is finite, $\mathcal{C}_m \subseteq \mathcal{A}_m$ and

$$\bigcup_{m \in \omega} \bigcup \mathcal{C}_m = X$$

Hence (X, τ) is a Menger space. \square

Thus by Theorem 2.4 and Theorem 3.1 we have that the totally Lindelöf property is in between the Alster property and productively Lindelöf.



4. THE DISCRETE SET PROPERTY AND FROLIK SPACES

We will introduce the ω_1 discrete set property and prove that no countable product of σ -compact spaces has the ω_1 discrete set property.

Proposition 4.1. *Let X be a σ -compact space. Then there is no sequence $\{x_\alpha \mid \alpha < \omega_1\} \subseteq X$ such that for all $\beta < \omega_1$ we have that $\{x_\alpha \mid \alpha < \beta\}$ is a discrete set.*

Proof. Let $X = \bigcup_{n \in \omega} K_n$ where for each $n \in \omega$ we have that K_n is compact. Towards a contradiction, suppose that there is a $\{x_\alpha \mid \alpha < \omega_1\} \subseteq X$ is such that for all $\beta < \omega_1$ we have that $\{x_\alpha \mid \alpha < \beta\}$ is a discrete set. There is $n^* \in \omega$ such that $\{x_\alpha \mid \alpha < \omega_1\} \cap K_{n^*}$ is uncountable. Then for some $\beta < \omega_1$ large enough we have $\{x_\alpha \mid \alpha < \beta\} \cap K_{n^*}$ is a countable discrete set contradicting the fact that K_{n^*} is compact. \square

The proposition above is what motivates our next definition:

Definition 4.2. If X is a topological space and $\{x_\alpha \mid \alpha < \omega_1\} \subseteq X$ such that for all $\beta < \omega_1$ we have that $\{x_\alpha \mid \alpha < \beta\}$ is a discrete set. Then we say that X has the ω_1 discrete set property.

Theorem 4.3. *Let $\{X_n \mid n < \omega\}$ be a sequence of σ -compact spaces. Then there is no sequence with the ω_1 discrete property in the product space $\prod_{n < \omega} X_n$.*

Proof. Firstly, let us fix the following notation for each $m < \omega$, $\pi_m : \prod_{n < \omega} X_n \rightarrow X_m$ is the usual projection.

Let $\{x_\alpha \mid \alpha < \omega_1\} \subseteq \prod_{n < \omega} X_n$ such that $x_\alpha = x_\beta$ iff $\alpha = \beta$. For $\alpha < \omega_1$ and $n < \omega$, define $x_\alpha(n) = \pi_n(x_\alpha)$. And for each $n < \omega$, fix a sequence $\{K_n^i \mid i < \omega\}$ where for each $i < \omega$ K_n^i is a compact subspace of X_n , $K_n^i \subseteq K_n^{i+1}$ and $X_n = \bigcup_{i < \omega} K_n^i$.

We will verify that for some $\beta < \omega$ the set $\{x_\alpha \mid \alpha < \beta\}$ has an accumulation point.

Given $\alpha < \omega_1$, we have that $x_\alpha \in \prod_{n < \omega} X_n$ and there is some $i < \omega$ such that $x_\alpha(0) \in K_0^i$. Therefore $x_\alpha \in \pi_0^{-1}[K_0^i]$. Thus

$$\{x_\alpha \mid \alpha < \beta\} \subseteq \bigcup_{i < \omega} \pi_0^{-1}[K_0^i]$$

and there is a least natural number i_0 such

$$|\{x_\alpha \mid \alpha < \omega_1\} \cap \pi_0^{-1}[K_0^{i_0}]| = \aleph_1$$

Inductively, if there are $i_0, \dots, i_n < \omega$ such that

$$|\{x_\alpha \mid \alpha < \omega_1\} \cap \pi_0^{-1}[K_0^{i_0}] \cap \dots \cap \pi_n^{-1}[K_n^{i_n}]| = \aleph_1$$

it follows that there is a least natural i_{n+1} such that

$$(|(\{x_\alpha \mid \alpha < \omega_1\} \cap \pi_0^{-1}[K_0^{i_0}] \cap \dots \cap \pi_n^{-1}[K_n^{i_n}]) \cap \pi_{n+1}^{-1}[K_{n+1}^{i_{n+1}}]| = \aleph_1$$

For each $n < \omega$ define $F_n = \{x_\alpha \mid \alpha < \omega_1\} \cap \bigcap_{m \leq n} \pi_m^{-1}[K_m^{i_m}]$. Since each F_n is uncountable, fix a strictly increasing sequence $\{\alpha_n \mid n < \omega\} \subseteq \omega_1$ such that for each $n < \omega$, $x_{\alpha_n} \in F_n$.

Since $K_0^{i_0}$ is compact, if $\{x_{\alpha_n}(0) \mid n \in \omega\}$ is infinite then we can fix $p_0 \in K_0^{i_0}$ an accumulation point of $\{x_{\alpha_n}(0) \mid n \in \omega\}$. If $\{x_{\alpha_n}(0) \mid n \in \omega\}$ is not infinite, we can fix $h \in \omega$ such that $\{n \in \omega \mid x_{\alpha_n}(0) = x_{\alpha_h}(0)\}$ is infinite, in this case we let $p_0 = x_{\alpha_h}(0)$. In both cases, we have that for every neighborhood V of p_0 the set $\{n < \omega \mid x_{\alpha_n}(0) \in V\}$ is infinite and $\{n < \omega \mid x_{\alpha_n}(0) \in V\} \subseteq \{n < \omega \mid x_{\alpha_n} \in \pi_0^{-1}[V]\}$.

Claim 4.3.1. *There exists a sequence $\langle p_n \mid n \in \omega \rangle \in \prod_{n \in \omega} X_n$ such that for every $n \in \omega$, if V_m is a neighborhood of p_m in X_m for each $m \leq n$, then the set $\{k < \omega \mid x_{\alpha_k} \in \bigcap_{m \leq n} \pi_m^{-1}[V_m]\}$ is infinite.*

Proof. We will construct $\langle p_n \mid n < \omega \rangle$ by induction. We have discussed the case $n = 0$ above, so let us assume that $n + 1 < \omega$ and that we have constructed $\langle p_m \mid m \leq n \rangle \in \prod_{m \leq n} K_m^{i_m}$ such that if we fix for each $m \leq n$ a neighborhood V_m of p_m in X_m , then the set $\{k < \omega \mid x_{\alpha_k} \in \bigcap_{m \leq n} \pi_m^{-1}[V_m]\}$ is infinite.

Suppose the induction step does not hold, namely, that for all $p \in K_n^{i_n}$ and $m < n$ there is $V_m(p)$ neighborhood of p_m and $V(p)$ of p such that $\{k < \omega \mid x_{\alpha_k} \in \bigcap_{m < n} \pi_m^{-1}[V_m(p)] \cap \pi_n^{-1}[V(p)]\}$ is finite. Since $K_n^{i_n}$ is compact, fix $l \in \omega$ and $\{q_0, \dots, q_l\} \subseteq K_n^{i_n}$ such that $K_n^{i_n} \subseteq \bigcup_{j \leq l} V(q_j)$. For each $m < n$, let $V_m = \bigcap \{V_m(p) \mid p \in I\}$.

Note that $\prod_{k < \omega} X_k \subseteq \bigcup_{j \leq l} \pi_{n+1}^{-1}[V(q_j)]$, and hence for each $k < \omega$, if $x_{\alpha_k} \in \bigcap_{m < n+1} \pi_m^{-1}[V_m]$ there is some $j \leq l$ such that

$$x_{\alpha_k} \in \bigcap_{m < n+1} \pi_m^{-1}[V_m] \cap \pi_{n+1}^{-1}[V(q_j)]$$

Thus $\{k < \omega \mid x_{\alpha_k} \in \bigcap_{m < n} \pi_m^{-1}[V_m]\}$ is a subset of

$$(1) \quad \bigcup_{j \leq l} \{k < \omega \mid x_{\alpha_k} \in \bigcap_{m < n+1} \pi_m^{-1}[V_m] \cap \pi_{n+1}^{-1}[V(q_j)]\}$$

which is absurd, since for each $j \leq l$ we have that

$$\{k < \omega \mid x_{\alpha_k} \in \bigcap_{m < n+1} \pi_m^{-1}[V_m] \cap \pi_{n+1}^{-1}[V(q_j)]\}$$

is a subset of

$$\{k < \omega \mid x_{\alpha_k} \in \bigcap_{m < n+1} \pi_m^{-1}[V_m(q_j)] \cap \pi_{n+1}^{-1}[V(q_j)]\}$$

which is finite. Therefore the union in (1) is finite, but we assumed that $\{k < \omega \mid x_{\alpha_k} \in \bigcap_{m < n} \pi_m^{-1}[V_m]\}$ is infinite. This contradiction proves the claim. \square

Now fix $\langle p_n \mid n \in \omega \rangle \in \prod_{n < \omega} X_n$ given by the Claim above.

Let $\beta = \sup_{n < \omega} \alpha_n$, we have that $\{x_\alpha \mid \alpha < \beta\}$ is not a discrete set. Therefore X does not have the ω_1 discrete set property. \square

Definition 4.4. (P-space) We say that a topological space X is a P -space if every countable intersection of open sets of X is an open set of X .

Corollary 4.5. *There is a P -space that is not Frolík, namely $L(\omega_1)$ is not Frolík*

Proof. If there was a homeomorphism to a closed subspace Z of the product space

$$\rho : L(\omega_1) \rightarrow Z \subseteq \prod_{n < \omega} X_n$$

Where each X_n is σ -compact, then $\{\rho(\alpha) \mid \alpha < \omega_1\}$ would witness that $\prod_{n \in \omega} X_n$ has the ω_1 discrete set property, contradicting 4.3. \square

We recall the reader that in Proposition 1.14 we proved that $L(\omega_1)$ is totally Lindelöf and it is straightforward to verify that $L(\omega_1)$ is regular. Hence by Theorem 1.1 we have that $L(\omega_1)$ is Alster. All together $L(\omega_1)$ is totally Lindelöf, Alster and it is not a Frolík space which answers question (c).

5. ACKNOWLEDGMENTS

We would like to thank Lúcia R. Junqueira for illuminating discussions on non-regular topological spaces. We are grateful to Rodrigo Carvalho and Artur Tomita for several discussions on totally Lindelöf spaces.

REFERENCES

- [Als88] K Alster. On the class of all spaces of weight not greater than ω_1 whose cartesian product with every lindelöf space is lindelöf. *Fundamenta Mathematicae*, 129:133–140, 1988.
- [AZ18] Leandro Aurichi and Lyubomyr Zdomskyy. Internal characterizations of productively lindelöf spaces. *Proceedings of the American Mathematical Society*, 146(8):3615–3626, 2018.
- [BT12] Peter Burton and Franklin D Tall. Productive lindelöfness and a class of spaces considered by z. frolik. *Topology and its Applications*, 159(13):3097–3102, 2012.
- [Fro63] Zdeněk Frolík. On the descriptive theory of sets. *Czechoslovak Math. J.*, 13(88):335–359, 1963.
- [MZ14] Andrea Medini and Lyubomyr Zdomskyy. Productively lindelöfness of spaces of countable tightness. *arXiv preprint arXiv:1405.6107*, 2014.
- [Nob19] N Noble. Some thoughts on countable lindelöf products. *Topology and its Applications*, 259:287–310, 2019.
- [ST09] Marion Scheepers and Franklin D Tall. Lindelof indestructibility, topological games and selection principles. *arXiv preprint arXiv:0902.1944*, 2009.
- [Tal11a] Franklin D Tall. Lindelöf spaces which are menger, hurewicz, alster, productive, or d. *Topology and its Applications*, 158(18):2556–2563, 2011.
- [Tal11b] Franklin D Tall. Set-theoretic problems concerning lindelof spaces. *arXiv preprint arXiv:1104.2796*, 2011.

- [Tal13] Franklin D Tall. Productively lindelöf spaces may all be d. *Canadian Mathematical Bulletin*, 56(1):203–212, 2013.
- [TT11] Franklin D Tall and Boaz Tsaban. On productively lindelöf spaces. *Topology and its Applications*, 158(11):1239–1248, 2011.
- [Vau78] JE Vaughan. Products of topological spaces. *General Topology and its Applications*, 8(3):207–217, 1978.

INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO, RUA DO MATÃO,
1010, SÃO PAULO, SP, BRAZIL

Email address: `fernandes@ime.usp.br`, `guipullus@ime.usp.br`, `vorocha@ime.usp.br`