

A generalized Kervaire Problem in stable homotopy groups of spheres

Petr M. Akhmetiev

March 21, 2023

Abstract

We will give an elementary self-contained description of the Mahowald element in the stable homotopy group of spheres Π_{2^l} , $l \geq 3$. Using this construction, we prove that a generalized Kervaire Problem, formulated by the author in [A] is solved positively.

Introduction

The Kervaire Problem in stable homotopy groups of spheres is a problem to calculate all dimensions, for which there exist an element in the 2-component of the stable homotopy group of sphere Π_n with the Arf-invariant 1. By the statement of the Hill-Hopkins-Ravenel Theorem [H-H-R] this is possible only for $n = 2^l - 2$, $l \leq 6$ and, probably, for $l = 7$.

According to an approach [E], [A-E] the Kervaire Problem is reformulated as following. Let us assume that the integer n is even and consider the homomorphism $Imm^{fr}(n, 1) \rightarrow Imm^{sf}(n - 1, 1)$, which corresponds to an immersion $f : M^n \looparrowright \mathbb{R}^{n+1}$ in its regular cobordism class the regular cobordism class $g : N^{n-1} \looparrowright \mathbb{R}^n$ of its self-intersection manifold. Then, let us consider the cobordism group of skew-framed immersions $Imm^{sf}(\frac{n}{2}, \frac{n}{2})$ and the homomorphism

$$J : Imm^{sf}(n - 1, 1) \rightarrow Imm^{sf}(\frac{n}{2}, \frac{n}{2}), \quad (1)$$

which corresponds to an immersion g a skew-framed immersion $(g - 1, \kappa_1, \Psi)$, which is a restriction of g on the submanifold $N_1^{\frac{n}{2}} \subset N^{n-1}$, this submanifold is defined as the transversal preimage of the universal submanifold

$$\mathbb{RP}^{\infty - \frac{n}{2} - 1} \subset \mathbb{RP}^{\infty} \quad (2)$$

by the mapping $\kappa : N^{n-1} \rightarrow \mathbb{R}P^\infty$, which represents the orienting class of the manifold N^{n-1} , assuming that κ is transversal along the submanifold (2).

Let us consider the homomorphism

$$sip : Imm^{sf}\left(\frac{n}{2}, \frac{n}{2}\right) \rightarrow \mathbb{Z}/2, \quad (3)$$

which corresponds to a skew-framed immersion a parity of its self-intersection points. Let us consider the composition (1), (3):

$$\Theta = sip \circ J : Imm^{sf}(n-1, 1) \rightarrow \mathbb{Z}/2. \quad (4)$$

Theorem 1. *Kervaire Problem is solved positively for the dimension n , iff the homomorphism (4) is non-trivial.*

In [A] a generalized Kervaire Problem is formulated:

Problem 2. *To calculate all dimensions n , for which the homomorphism (3) is non-trivial. If for a dimension n the homomorphism is non-trivial, one say that the problem has a positive solution in the dimension n .*

A generalized Kervaire Problem admits important applications.

At the first, by [A-C-R] a positive solution of the problem in a dimension n implies that there exists a stably-framed immersion (this means that the normal bundle of the immersion is equipped with the stable framing) of a closed n -dimensional manifold with the Browder-Eccles invariant 1 (the Browder-Eccles invariant is the twisted Arf-invariant, associated with the stable-framed immersion). This result is motivated by calculations of the Eccles-Wood filtration of stable homotopy groups of spheres.

At the second, if we additionally assume that an immersed manifold $\zeta_l^{sf} \in Imm^{sf}\left(\frac{n}{2}, \frac{n}{2}\right)$, $\frac{n}{2} = 2^l - 1$, which determines a positive solution of the generalized Kervaire Problem has an integer orienting cohomology class, the problem was investigated in [B-J-M] and is called the strong Kervaire Problem in dimension $\frac{n-2}{2}$. From a positive solution of the strong Kervaire Problem the Brown-Peterson Theorem (a reference is given in [B-J-M]) implies, that the Kervaire Problem in the dimension $\frac{n-2}{2}$ has a positive solution.

The result of the paper is the following:

Theorem 3. *A generalized Kervaire Problem has a positive solution for all integers $n = 2^l - 2$, $l \geq 2$.*

As a conclusion, a calculation of the minimal codimension of skew-framed immersion of n -dimensional manifold with an odd number of self-intersection points, as well as a calculation of the minimal cohomology length of the

orienting class for a skew-framed immersed $\frac{n}{2}$ -manifold in the codimension $\frac{n}{2}$ are interesting.

The proof of Theorem 3 is based on a result by Mahowald [M],[E-Z]. Because our approach is different with respect to the classical approach, even at a level of statements of the results, we will consider a statement of the Mahowald Theorem (Section 2) and give a self-contained proof of this theorem. The main result is Theorem 3, which is proved geometrically by a short construction.

1 The element η_i^{sf} , $i = 3$, with the Hopf invariant 1

In the Section the first term $i = 3$ of the family of immersions $\phi_i : N^{2^i-1} \looparrowright \mathbb{R}^{2^i}$, which represents a non-trivial element is the cobordism group $Imm^{sf}(2^i - 1, 1)$, $i \geq 3$ is constructed. For this element the Hopf invariant (the Hopf invariant $h(f)$ for an r -immersion $f : N^r \looparrowright \mathbb{R}^{r+k}$ $k < r$, is the number modulo 2 of self-intersection points of a generic alteration of the following immersion: $N^r \rightarrow \mathbb{R}^{r+k} \subset \mathbb{R}^{2r}$) satisfies the equations: $h(\phi) = 1$.

We follows an approach toward stable homotopy groups of spheres, which is based on the immersion theorem by Smale, see [H]. Let us recall a definition of the cobordism groups $Imm^{sf}(n - k, k)$, $Imm^{st-sf}(n - k, k)$.

Let $\varphi : M^{n-k} \looparrowright \mathbb{R}^n$ be an immersion, $\kappa : M^{n-k} \rightarrow \mathbb{RP}^\infty$ be a characteristic class, which is called the characteristic class of a skew-framing, $\gamma \rightarrow \mathbb{RP}^\infty$ be the canonical line bundle over the real projective space \mathbb{RP}^∞ , $\Xi : \nu_\varphi \equiv \kappa^*(k\gamma)$ be an isomorphism of the normal bundle of the immersion φ with the inverse image of k -dimensional bundle $k\gamma$ over \mathbb{RP}^∞ , the Whitney sum of k copies of the line bundle γ . The isomorphism Ξ is called a skew-framing of the immersion φ in the codimension k .

Let us define the following Abelian group $Imm^{sf}(n - k, k)$ as regular cobordism classes of immersion (up to regular cobordism), the group structure is determined by disjoint union of the following triples: (φ, κ, Ξ) . The regular cobordism relation of triples is a standard relation, see [A-F] for details.

Let us say that an element $x \in Imm^{sf}(n - k, k)$ is represented by an immersion with an integer characteristic class of its skew-framing, if there exists in the regular cobordism class of the element a triple (φ, κ, Ξ) for which the image of the mapping κ is inside $\mathbb{RP}^1 \subset \mathbb{RP}^\infty$.

Let $\varphi : M^{n-k} \looparrowright \mathbb{R}^n$ be an immersion, $\kappa : M^{n-k} \rightarrow \mathbb{RP}^\infty$ be the characteristic class of its skew-framing, $\Xi^{st} : \nu_\varphi \oplus N\kappa^*(\gamma) \equiv \kappa^*((k + N)\gamma)$ be a

stable isomorphism of the normal bundle of the immersion φ with the inverse image of a k -dimensional bundle $k\gamma$ over $\mathbb{R}P^\infty$, N be a great positive integer. Such an isomorphism Ξ^{st} is called a stable skew-framing of the immersion φ in the codimension k .

Let us define the cobordism group $Imm^{st-sf}(n-k, k)$ as a set (the group structure is defined by disjoint union of triples described above) of regular cobordism classes of triples $(\varphi, \kappa, \Xi^{st})$. The equivalent relations of triples is determined by regular cobordism classes of triples.

The Mahowald element $\eta_i^{fr} \in \Pi_{2^i}$ is defined using the element η_i^{sf} , constructed below, by the Khan-Priddy homomorphism :

$$\lambda : Imm^{st}(2^i - 1, 1) \rightarrow \Pi_{2^i},$$

[K].

In this section we construct the Mahowald element η_l^{sf} , $l = 3$, which determines the base of the induction over the parameter $l \geq 3$. The following lemma explain a geometrical meaning of the following in [M]: the element hh_3 is spherical in ASS.

Lemma 4. *There exists an element $x \in Imm^{sf}(7, 1)$ with the Hopf invariant $h(x) = 1$.*

Proof of Lemma 4

Obviously, the image of the generator Π_7 by the homomorphism $forg : \Pi_7 \equiv Imm^{fr}(7, 1) \rightarrow Imm^{sf}(7, 1)$ into the group of skew-framed immersion satisfies the statement of the lemma.

To our goal the base of the induction has to be proved alternatively. Let us construct another immersion (which satisfies special properties) $\phi : N^7 \looparrowright \mathbb{R}^8$, which represents an element $x = [\phi]$, $h(x) = 1$.

Define the immersion ϕ by the composition of a 2-sheeted covering $p : N^{2^i-1} \rightarrow \hat{N}^{2^i-1}$, $i = 3$ and an immersion $\hat{\phi} : \hat{N}^{2^i-1} \looparrowright \mathbb{R}^{2^i}$. At the beginning, let us prove in this case that the Hopf invariant $h(\phi)$ is calculated by the formula:

$$\langle p^{2^i-1}; [\hat{N}^{2^i-1}] \rangle = h(\phi), \quad i = 3, \quad (5)$$

where $p \in H^1(\hat{N}^{2^i-1}; \mathbb{Z}/2)$ is a characteristic cohomology class, which represents the covering (we denote a covering and its cohomology characteristic class the same).

To get a proof, let us consider an immersion $\hat{\phi} = I \circ \hat{\phi} : \hat{N}^{2^i-1} \looparrowright \mathbb{R}^{2^i} \subset \mathbb{R}^{2^{i+1}-2}$, which to be assumed a self-transversal. Let us consider an immersion

$\phi' = \hat{\phi} \circ p : N^{2^i-1} \looparrowright \mathbb{R}^{2^{i+1}-2}$. Denote by $\phi : N^{2^i-1} \looparrowright \mathbb{R}^{2^{i+1}-2}$ a generic immersion, which is closed to ϕ' . One has to prove that $h(\phi) \pmod{2}$ (the number of self-intersection points of ϕ) is calculated by the formula: (5).

Obviously, self-intersection points of the immersion $\hat{\phi}$ give no contribution to the value $h(\phi)$, because a self-intersection point of $\hat{\phi}$ determines an even number of self-intersection points on the double covering. Therefore only self-intersection points of a deformation $\phi \mapsto \phi'$ inside a regular neighbourhood of the immersion $\hat{\phi}$ gives a contribution. It is clear (the reasoning is analogous to [Sz]), that the number (modulo 2) of such points is calculated by the formula:

$$\langle (w_1(\hat{N}) + p)p^{2^i-2}; [\hat{N}^{2^i-1}] \rangle = h(\phi).$$

Because

$$\langle w_1(\hat{N})p^{2^i-2}; [\hat{N}^{2^i-1}] \rangle = 0,$$

using the fact that one-dimensional manifold, which is Poincarè dual to the cocycle $p^{2^i-2} \in H^1(\hat{N}; \mathbb{Z}/2)$, the expression for the Hopf invariant is given by the left hand side of the formula (5).

Let us define a closed manifold $N^7 = (S^3 \times S^3) \tilde{\times} S^1$ as a semi-direct product $S^3 \times S^3$ with the circle S^1 . This manifold N^7 is fibred over S^1 with the fibre $S^3 \times S^3$. Fibres $S^3 \times S^3$ are identified (by a monodromy of a generic loop on the base S^1 of the fibre) by the involution $S^3 \times S^3 \rightarrow S^3 \times S^3$, which permutes the factors. The oriented class of the manifold N^7 , which is denoted by $\kappa_N \in H^1(N^7; \mathbb{Z}/2)$, is induced from the generator $H^1(S^1)$ by the projection $N^7 \rightarrow S^1$.

Let us consider the following manifold: $\tilde{N}^7 = (\mathbb{RP}^3 \times \mathbb{RP}^3) \tilde{\times} S^1$, this manifold is fibred over S^1 with the fibre $\mathbb{RP}^3 \times \mathbb{RP}^3$, this manifold is a result of the quotient of the manifold N^7 by a pair of free involutions. The monodromy of a fibre $\mathbb{RP}^3 \times \mathbb{RP}^3$ along the base S^1 is defined by the involution $\mathbb{RP}^3 \times \mathbb{RP}^3 \rightarrow \mathbb{RP}^3 \times \mathbb{RP}^3$, which permutes the factors. Let us define an immersion $\tilde{\varphi}_0 : \tilde{N}^7 \looparrowright \mathbb{R}^8$, the normal bundle of this immersion is a linear bundle, which coincides with the orienting bundle $\nu_{\tilde{\varphi}_0} \cong \tilde{\kappa}_N$.

Let us define an auxiliary immersion $\tilde{\varphi} : \tilde{N}^7 \looparrowright \mathbb{R}^9$. Consider an immersion $f : \mathbb{RP}^3 \looparrowright D^4$ of the codimension 1 and consider the Cartesian product $f_1 \times f_2 : \mathbb{RP}^3 \times \mathbb{RP}^3 \looparrowright D_1^4 \times D_2^4$. Let us consider the manifold $(D_1^4 \times D_2^4) \rtimes_T S^1$, which is defined as the semi-direct product $D_1^4 \times D_2^4$ with the circle, along the circle the 4-disks are permuted. Let us define an immersion $\tilde{\varphi}' : \tilde{N}^7 \looparrowright (D_1^4 \times D_2^4) \rtimes_T S^1$, as an immersion, which is induced from the immersion $f_1 \times f_2$ by a semi-direct product. The normal bundle of this immersion is 2-dimensional, this bundle is the Whitney sum $\tilde{\kappa}_N \oplus \varepsilon$. Let us define an immersion $\tilde{\varphi}$ as the composition $I \circ \tilde{\varphi}'$, where $I : (D_1^4 \times D_2^4) \rtimes_T S^1 \subset \mathbb{R}^{10}$ is an embedding of a (thin) cylinder.

By the Hirsch Theorem [H], starting by the immersion $\tilde{\varphi}$, one may define a compressed immersion $\tilde{\varphi}_0 : \tilde{N}^7 \looparrowright \mathbb{R}^8$, uniquely, up to a concordance. Let us now define the required manifold \hat{N}^7 , which is equipped by a characteristic class $p \in H^1(\hat{N}^7; \mathbb{Z}/2)$, which satisfies the condition:

$$\langle p^{2^i-1}; [\hat{N}^{2^i-1}] \rangle = 1 \pmod{2}, \quad i = 3, \quad (6)$$

and, moreover, an immersion $\hat{\varphi} : \hat{N}^7 \looparrowright \mathbb{R}^8$.

For this purpose let us consider the standard 4-sheeted covering $p : N^7 = (S^3 \times S^3) \tilde{\times} S^1 \rightarrow \hat{N}^7$, which is induced by the pair of coverings of factors: $S^3 \rightarrow \mathbb{RP}^3$.

Let us define the following manifold \hat{N}^7 , as a manifold of pairs of two-point configurations $[(x_1, x_2), (y_1, y_2)]$ in N^7 , which satisfies the condition: $x_1 = Tx_2, y_1 = Ty_2$, where T is the transposition of points in a fibre of a corresponding covering factor: $S^3 \rightarrow \mathbb{RP}^3$. The two pairs of 2-point configurations are equivalent, if one configuration is the result of the transposition of points in each 2-point configurations. As a result, the manifold \hat{N}^7 as a base of 2-sheeted covering with the covering space N^7 is well-defined. Let us denote by $p \in H^1(\hat{N}^7; \mathbb{Z}/2)$ the characteristic class of this covering.

Lemma 5. *For the characteristic number, constructed by the only cohomology class p in $H^1(\hat{N}^7)$, the equation (6) is satisfied.*

Proof of Lemma 5

Let us consider the standard 7-sphere S^7 as the join of the two standard 3-spheres S^3 : $S^7 = S^3 * S^3$. The antipodal involution T_{S^7} on S^7 is induced from the antipodal involutions T_{S^3} on the factors of the join. Consider 6-torus $P^6 \subset S^3 * S^3$, which is defined as the middle submanifold in the join. Obviously, $P^6 \cong S^3 \times S^3$ and the restriction of the involution T_{S^7} on this torus P^6 coincides with the diagonal of the two involutions T_{S^3} on the factors.

Let us construct a mapping $\hat{N}^7 \rightarrow \mathbb{RP}^7$ of the degree 1 (mod 2), and which corresponds with the cohomology class p by the stabilization $\mathbb{RP}^7 \subset \mathbb{RP}^\infty$ of the image. Consider the projection $\hat{N}^7 \rightarrow S^1$ on the generator and denote by $\hat{K}^6 \subset \hat{N}^7$ the submanifold, which is defined as the fibre of the projection. Clearly, $\hat{K}^6 = S^3 \times S^3 / T_{diag}$, where T_{diag} is the diagonal involution $T_{diag} = T_{S^3} \times T_{S^3}$. Let us consider the cutting of \hat{N}^7 along \hat{K}^6 and define a mapping $\hat{F} : \hat{N}^7 \setminus \hat{K}^6 \rightarrow \mathbb{RP}^7$, which maps the two components of the boundary into two disjoint copies $\mathbb{RP}^3 \cup \mathbb{RP}^3 \subset \mathbb{RP}^7$ as following. Describe \hat{F} using the equivariant covering mapping $F : N^7 \setminus K^6 \rightarrow S^3 * S^3$, which is mapped the two components of the boundary onto the disjoint spheres, the generators of the join.

By the construction, the restriction of the equivariant mapping F on the first (on the second, correspondingly) component of the boundary coincides with the composition of the equivariant projection $p_i : S^3 \times S^3 / T_{diag}$ on the first $i = 1$ (on the second, correspondingly) $i = 2$ coordinate with the equivariant inclusion $I_i : S^3 \subset S^3 * S^3$ of the first (the second, correspondingly) coordinate.

Let us prove that the mapping p_1 is equivariant homotopic to p_2 after the equivariant stabilization of the images $S^3 \subset S^6$ in a common manifold. To prove this, let us define the total obstruction of an equivariant homotopy of the two equivariant mappings $f_i : S^3 \times S^3 \rightarrow S^6$, $i = 1, 2$. This obstruction is defined using the standard definition, this total obstruction is denoted by $o(f_1, f_2)$ and belongs to \mathbb{Z} .

To prove this fact we recall that the involution in the preimage $S^3 \times S^3$ is defined as the diagonal antipodal involution T_{diag} on the factors and this involution preserves the orientation. In the image S^6 the involution is the standard antipodal involution and inverses the orientation. An obstruction $o(f_1, f_2)$ looks like an analogous obstruction in the problem for the covering mappings $\bar{f}_1, \bar{f}_2 : T^2 \rightarrow \mathbb{RP}^2$, where T^2 is the standard torus. This obstruction is identified with the half degree of the covering mappings in the prescribed coordinate system in the target.

In the considered situation the involution T_{diag} keeps the orientation, and the standard involution T_{S^6} reverses the orientation, the integer degree $deg(f_i)$ of the two equivariant mappings are related with the obstruction by the formula:

$$o(f_1, f_2) = \frac{1}{2}(deg(\bar{f}_1) - deg(\bar{f}_2)).$$

Because $deg(\bar{p}_1) = deg(\bar{p}_2) = 0$, the mappings p_1, p_2 are homotopic inside \mathbb{RP}^6 .

Let us consider the submanifold $\mathbb{RP}^6 \subset \mathbb{RP}^7$, which contains the images of the equivariant mappings $I_i \circ p_i$, $i = 1, 2$. The mappings f_1, f_2 are equivariant homotopic, therefore an extension of the mapping F on the manifold N^7 is well defined, and extra regular preimages are not added. The calculation proves Lemma 5, as well as Lemma 4. \square

2 The Mahowald elements η_i^{sf} , $i > 3$

In this section let us consider the next step of the induction $i = 4$, and construct an immersion $\phi : N^{15} \looparrowright \mathbb{RP}^{16}$, which represents the element $[\phi] = \eta_4^{sf}$ in the group $Imm^{sf}(15, 1)$, using the element $\eta_3^{sf} \in Imm^{sf}(7, 1)$, constructed above.

Let us repeat a reasoning and define $\tilde{M}^{15} = (\hat{N}^7 \times \hat{N}^7) \tilde{\times} S^1$. As the component of this semi-direct product let us take the manifold \hat{N}^7 , which is constructed in the section 1. Then let us define the tower of 2-sheeted coverings:

$$M^{15} \xrightarrow{p} \hat{M}^{15} \xrightarrow{\hat{p}} \tilde{M}^{15}, \quad (7)$$

the bottom covering \hat{p} is called a primary covering, the top covering p is called a secondary covering. To indicate the base of the covering we will use the denotation $p_{\hat{M}}$ e.t.c. The following formula, which is an analogue of the formula (5) in the case $i = 4$ is satisfied:

$$\langle p^{2^i-1}; [\hat{M}^{2^i-1}] \rangle = h(\phi), \quad i = 3. \quad (8)$$

In the proof one can put $\hat{N}^7 = \mathbb{R}P^7$, then to pass to the required case and prove that the equivariant composition $p \times p : N^7 \times N^7 \rightarrow S^7 \times S^7$, where the equivariant structure in the image and the preimage are defined by the free diagonal involutions. The mapping is equivariant homotopic inside S^{14} to an equivariant mapping, which is defined by the composition of the transposition of the factors.

Then, starting by the manifold \hat{M}^{15} the required manifold \hat{N}^{15} , equipped by the covering $p : N^{15} \rightarrow \hat{N}^{15}$, and by the immersion $\varphi = \hat{\varphi} \circ p : N^{15} \looparrowright \mathbb{R}^{15}$ is constructed. This immersion represents the described element $\eta_4^{sf} \in Imm^{sf}(15, 1)$ with Hopf invariant 1.

Let us consider the Cartesian product $\hat{\varphi}_7 \times \hat{\varphi}_7 : \hat{N}^7 \times \hat{N}^7 \looparrowright \mathbb{R}^8 \times \mathbb{R}^8$ of the two copies of the immersions. The normal bundle $\nu(\hat{\varphi}_7 \times \hat{\varphi}_7)$ of this immersion is 2-dimensional and is represented as the Whitney sum of the two copies of the line bundle $\nu(\hat{\varphi})$.

Let us consider the Cartesian product $D^8 \times D^8$ of two open 8-disks and the open manifold $\tilde{K}^{17} = (D^8 \times D^8) \tilde{\times} S^1$, where the semi-direct product is defined by the monodromy $m : D^8 \times D^8 \rightarrow D^8 \times D^8$ along the generator S^1 by the formula: $m(x, y) = (y, x)$. Evidently, \tilde{K}^{17} is diffeomorphic to $D^8 \times D^8 \times S^1$ and there exists an embedding $i_{\tilde{K}} : \tilde{K}^{17} \subset \mathbb{R}^{17}$ (the construction below does not depend on a choice of the embedding). Let us consider the manifold $\tilde{M}^{15} = (\hat{N}^7 \times \hat{N}^7) \tilde{\times} S^1$ (which is analogous to the manifold $(\mathbb{R}P^3 \times \mathbb{R}P^3) \tilde{\times} S^1$ constructed in the section 9). Let us consider an immersion $j : (\hat{N}^7 \times \hat{N}^7) \tilde{\times} S^1 \looparrowright \tilde{K}^{17}$ and the composition $\tilde{\psi} = j \circ i_{\tilde{K}} : (\hat{N}^7 \times \hat{N}^7) \tilde{\times} S^1 \looparrowright \mathbb{R}^{17}$. The normal bundle $\nu_{\tilde{\psi}}$ of the immersion $\tilde{\psi}$ admits a reduction of the structured group $O(2)$ to the discrete subgroup \mathbf{D} .

The dihedral group \mathbf{D} contains 8 elements and has the corepresentation $\{a, b | a^2 = b^4 = 1; [a, b] = b^2\}$. The element $b^2 \in \mathbf{D}$ is also will be denoted

denoted $-1 \in \mathbf{D}$. The group D is represented in $O(2)$ as following: the element b is represented by the rotation of the plane through the angle $\frac{\pi}{2}$, the element a is represented by the symmetry with respect to the bisector of the first-third coordinate quarter. Let us denote $A = a \circ b$, this element is represented by the symmetry with respect to the horizontal coordinate axis. Let us denote $\dot{A} = b^2 A = -A$, $\dot{a} = b^2 a = -a$.

Over the manifold \tilde{M}^{15} the following tower of 2-sheeted coverings (7) is well-defined, this tower corresponds to the following system of subgroups:

$$\mathbf{I}_a \subset \mathbf{I}_{a,\dot{a}} \subset \mathbf{D}. \quad (9)$$

The structured group of the normal bundle $\nu_{\tilde{\psi}}$ of the immersion $\tilde{M}^{15} \looparrowright \mathbb{R}^{17}$ is also isomorphic to the group \mathbf{D} . To distinguish the two isomorphic sequence of subgroups: the sequence of coverings over the manifold \tilde{M}^{15} and the sequence of reductions of the structured group of the normal bundle, the subgroup of coverings tower is denoted by (9), a subgroup of the structured group is denoted by the corresponding collection of elements. In particular, the sequence of subgroups (9) of the covering corresponds to the following sequence of monodromy:

$$\{a\} \subset \{a,\dot{a}\} \subset \{a,b\} \simeq \mathbf{D}. \quad (10)$$

The element A of the residue class of the subgroup $\mathbf{I}_{a,\dot{a}} \subset \mathbf{D}$ acts by an involution on the covering manifold \hat{M}^{15} of the covering \hat{p} , this action is denoted by T_A . The element $-1 = b^2$ in the residue class of the subgroup $\mathbf{I}_a \subset \mathbf{I}_{a,\dot{a}}$ acts on M^{15} by a covering translation $M^{15} \rightarrow \hat{M}^{15}$, this action is denoted by T_{b^2} . The restriction of T_A on the submanifold

$$M^{15} \supset (N^7 \times N^7)/\{b^2\} \quad (11)$$

is defined by the formula: $(n_1, n_2) \mapsto (T_A(n_1), n_2)$.

Let us consider the subgroups $\{a\} \subset \{a,b\}$, $\mathbf{I}_a \subset \mathbf{D}$, which are not normal subgroups. The normaliser of the subgroup $\{a\}$ is the subgroup $\{a,\dot{a}\} \subset \{a,b\}$. Analogously, the normaliser of \mathbf{I}_a is the subgroup $\mathbf{I}_{a,\dot{a}} \subset \mathbf{D}$, recall that \mathbf{D} is the structured group of the bundle $\nu_{\tilde{\psi}}$.

The subgroup $\mathbf{I}_a \subset \mathbf{D}$ is a subgroup of the covering $\hat{p} \circ p : M^{15} \rightarrow \tilde{M}^{15}$, see (7), which is not a covering with a normal subgroup. The subgroup $\mathbf{I}_{A,\dot{A}} \subset \mathbf{D}$ is visualized as a translation group of the covering $M^{15} \rightarrow \tilde{M}^{15}$, over the submanifold $\hat{N}^7 \times \hat{N}^7 \subset \tilde{M}^{15}$; this group determines transformations of sheets of the covering.

Let us define the manifold \hat{N}^{15} . Let us consider the vector bundle $\nu_{\tilde{\psi}}$ over \tilde{M}^{15} and denote

$$\nu_{\tilde{\psi}}^{tw} = \nu_{\tilde{\psi}} \otimes \lambda, \quad (12)$$

where $\lambda = \lambda(\hat{p})$ is a linear bundle over \tilde{M}^{15} , which is associated with the covering $\hat{p} : \hat{M}^{15} \rightarrow \tilde{M}^{15}$. Denote by $\tilde{L}^{13} \subset \tilde{M}^{15}$ the Euler class of the bundle $\nu_{\hat{\psi}}^{tw}$ (recall, the Euler class of a vector bundle over a base manifold is defined as a submanifold of zero sections of the vector bundle, in particular the normal bundle of the Euler class submanifold is isomorphic to the bundle itself, where $\hat{K}^{14} \subset \hat{M}^{15}$ is the boundary of the disk-bundle associated with the vector bundle (12), which is the total space of the corresponding S^1 -bundle over \tilde{L}^{13} . For the formula (12) of the Euler class of the twisted section see [K1].

The total space D_{λ}^{16} of the disk bundle, which is associated with the bundle λ , is a 16-dimensional manifold with the boundary $\hat{M}^{15} = \partial(D_{\lambda}^{16})$. In the manifold with boundary D_{λ}^{16} let us define a subspace $\hat{N}^{15} \subset D_{\lambda}^{16}$, which is the required closed 15-dimensional manifold.

Take the submanifold \tilde{L}^{13} and take its preimage by the projection $D_{\lambda}^{16} \rightarrow \tilde{M}^{15}$ on the base of the bundle. Let us cut-out fibres over the regular neighbourhood $U(\tilde{L}^{13})$ in D_{λ}^{16} . The boundary of the obtained manifold consists of a lateral surface of the handle and the bottom of the handle, the bottom is the double covering over $\hat{N}_{pred}^{15} \subset D_{\lambda}^{16} \rightarrow \tilde{M}^{15}$. Denote this closed manifold by \hat{N}^{15} (this manifold is the required manifold), the decomposition of this manifold as the lateral surface and the bottom of the handle denote by

$$\hat{N}^{15} = \hat{N}_{pred}^{15} \cup \hat{K}^{15}. \quad (13)$$

Let us consider the projection $q : \hat{N}_{pred}^{15} \rightarrow \tilde{M}^{15} \setminus U(\tilde{L}^{13})$, as the restriction of the projection $D_{\lambda}^{16} \rightarrow \tilde{M}^{15}$ on the submanifold. By this projection the image $r(\hat{K}^{15})$ coincides with $\partial(U(\tilde{L}^{13}))$, the image $r(\hat{N}^{15})$ coincides with the interior $\tilde{M}^{15} \setminus U(\tilde{L}^{13})$.

Let us consider the lateral surface of the handle \hat{K}^{15} . The middle submanifold of the lateral surface \hat{K}^{15} will be denoted by $\hat{K}_m^{14} \subset \hat{K}^{15}$. The natural projection $\hat{K}^{15} \rightarrow \hat{K}_m^{14}$ is a D^1 -bundle; the restriction of this projection on the boundary $\partial(\hat{K}^{15})$ of the lateral handle is the double covering over \hat{K}_m^{14} . Let us denote $\partial(\hat{K}^{15})$ by K_m^{14} . This covering coincides with the pull-back $\lambda \mapsto \lambda^!$, $\lambda^! \simeq \varepsilon$, by the bottom covering in (7) of the spherization $S(\nu_{\hat{L}} \otimes \lambda)$ of the normal bundle of the submanifold $\tilde{L}^{13} \subset \tilde{M}^{15}$, which is a copy the restriction of the normal bundle $\nu_{\hat{M}}$ over its own Euler class. By construction, there is a cohomology class:

$$p : K_m^{14} \rightarrow \mathbb{RP}^{\infty}, \quad (14)$$

which is the restriction of the top class in (7).

We may identify \hat{K}_m^{14} with $S(\nu_{\tilde{L}}) \otimes \lambda$, i.e. with the spherization of the normal Euler class of the bundle (12). Obviously,

$$S(\nu_{\tilde{L}} \otimes \lambda) = S(\nu_{\tilde{L}}) \otimes \lambda. \quad (15)$$

The left hand side of the formula (15) looks more naturally.

On the manifold \hat{K}_m^{14} the antipodal free involution is well-defined:

$$J_{\hat{K}_m} : \hat{K}_m^{14} \rightarrow \hat{K}_m^{14},$$

the quotient $\hat{K}_m^{14}/J_{\hat{K}_m}$ is naturally identified with the projectivization $P(\nu_{\tilde{L}} \otimes \lambda)$ corresponded with the bundle (15). Let us denote this manifold by \tilde{K}_m^{14} .

Let us consider the following diagram of the 2-sheeted coverings:

$$\begin{array}{ccc} S(\nu_{\tilde{L}} \otimes \lambda) = \hat{K}_m^{14} & \xrightarrow{\hat{\pi}} & \tilde{K}_m^{14} = P(\nu_{\tilde{L}} \otimes \lambda) \\ \hat{q} \uparrow & & \uparrow \tilde{q} \\ S(\nu_{\tilde{L}} \otimes \lambda^!) = K_m^{14} & \xrightarrow{\hat{\pi}} & K_m^{14}/J_{K_m} = P(\nu_{\tilde{L}} \otimes \lambda^!). \end{array} \quad (16)$$

Lemma 6. *There exists a cohomology class $\alpha : K_m^{14} \rightarrow \mathbb{RP}^\infty$, which is pull-back of p , defined by (14), by the projection $K_m^{14} \rightarrow \hat{L}^{13} \subset \hat{M}^{15}$.*

Corollary 7. *There exist a cohomology class $\hat{\beta} : \hat{K}_m^{14} \rightarrow \mathbb{RP}^\infty$, and the pull-back by the covering \hat{q} of this class coincides with the class $\alpha : K_m^{14} \rightarrow \mathbb{RP}^\infty$.*

Proof of Lemma 6

Take the following diagram, which contains the diagram (16) as the Euler class:

$$\begin{array}{ccc} S(\nu_{\tilde{M}} \otimes \lambda) = \hat{Q}_m^{16} & \xrightarrow{\hat{\pi}} & \tilde{Q}_m^{16} = P(\nu_{\tilde{M}} \otimes \lambda) \\ \hat{q} \uparrow & & \uparrow \tilde{q} \\ S(\nu_{\tilde{M}} \otimes \lambda^!) = Q_m^{16} & \xrightarrow{\hat{\pi}} & Q_m^{16}/J_{Q_m} = P(\nu_{\tilde{M}} \otimes \lambda^!). \end{array} \quad (17)$$

Let us consider the class $\alpha_{\hat{R}}$ on \hat{R}_m^{16} , which is defined by the standard characters \mathbf{I}_A (by the projection $\mathbf{I}_{A \times A} \rightarrow \mathbf{I}_A$). Let us prove the relation

$$T_{\hat{R}_m} \circ \alpha_{\hat{R}} = \alpha_{\hat{R}}. \quad (18)$$

In the diagram (17) spaces and coverings are defined analogously with (16), instead of the manifold \tilde{L}^{13} , the common base of the covering, we get the manifold \tilde{M}^{15} equipped by the bundles described by the formula (12).

Let us consider the dyagram of spaces, which are double coverings over the corresponding spaces of the diagram (17):

$$\begin{array}{ccc}
\hat{R}_m^{16} & \xrightarrow{\hat{\pi}} & \tilde{R}_m^{16} \\
\hat{q} \uparrow & & \uparrow \tilde{q} \\
R_m^{16} & \xrightarrow{\hat{\pi}} & R_m^{16}/J_{R_m},
\end{array} \tag{19}$$

which is defined as following.

Recall that $\tilde{M}^{15} = \hat{N}^7 \times \hat{N}^7 \rtimes_{\hat{T}} S^1$. The submanifold $\hat{N}^7 \times \hat{N}^7 \subset \tilde{M}^{15}$ is the ordered 2-points configuration space of \hat{N}^7 , the involution \tilde{T} permutes points in pairs. The covering $\hat{M}^{15} \rightarrow \tilde{M}^{15}$ is described in (7), using the diagram (9), we get $\hat{M}^{15} = (N^7 \times N^7)/T_{diag} \rtimes_{\hat{T}} S^1$, where, T_{diag} is the diagonal of the involution $N^7 \rightarrow \hat{N}^7$ on the Cartesian product, this involution corresponds to the element $b^2 \in \mathbf{D}$; the involution \hat{T} permutes points in pairs. The following covering $\hat{N}^7 \times \hat{N}^7 \times S^1 \rightarrow \hat{M}^{15}$, which is the covering with the subgroup $\mathbf{I}_{A \times A} \subset \mathbf{D}$, induces the coverings of spaces of (19) over corresponding spaces of (17).

Let us describe the cohomology classes $\alpha_R : R_m^{16} \rightarrow \mathbb{R}P^\infty$. Then let us prove that the following formula is satisfied:

$$\alpha_R = T_R \circ \alpha_R, \tag{20}$$

where T_R is the involution of the covering $R_m^{16} \rightarrow Q_m^{16}$.

Define the class α_R as the pull-back of the class $[T_{diag}]^* \in H^1((N^7 \times N^7)/T_{diag})$ by the projection $R_m^{16} \rightarrow (N^7 \times N^7)/T_{diag} \times S^1 \rightarrow (N^7 \times N^7)/T_{diag}$. The fact (20) is a corollary that α_R coincides to the standard \mathbf{I}_a -character (is invariant with respect to \mathbf{I}_a -transformation). The cohomology class $\alpha_Q \in H^1(Q_m^{16})$ is well-defined.

Lemma 6 is proved. \square

Proof of Lemma 7

Take the left-bottom manifold K_m^{14} in (16) and consider the section $\varepsilon = \lambda^!$ as the pull-back of λ of the bundle $\nu_{\tilde{L}} \otimes \lambda$, restricted outside the Euler class $\tilde{L}^{13} \subset \tilde{M}^{15}$ of the cross-section. Below we will speak of the section $\lambda^!$ as a section of the bundle $\nu_{\tilde{L}}$ instead of a section of the bundle $(\nu_{\tilde{L}} \otimes \lambda)^!$, which is equivalent.

An open neighbourhood of $K_m^{14} \subset Q_m^{16}$ is referred as the regular domain in Q_m^{16} . A regular domain of $K_m^{14} \subset Q_m^{16}$ and its preimage in R_m^{16} are denoted by $Q_{m;reg}^{16}$, $R_{m;reg}^{16}$ correspondingly.

Let us denote by $H^+(X) \subset H_1(X)$ for an non-orientable closed manifold X (we take below $X = K_m^{14}$, for example) the index 2 subgroup, which is the

kernel of the orientation homomorphism: $Ker(w) : H_1(X) \rightarrow \mathbb{Z}/2$. Analogously, define the non-trivial resedue class of the orientation homomorphism by $H^-(X)$. Obviously, $H^1(X) = H^+(X) \cup H^-(X)$. For an arbitrary homology class $[l]_* \in H^+(K_m^{(14)})$ let us consider a connected loop and the double covering l_R of the loop in $R_{m;reg}^{16}$ (the lift-bottom space in (19)). The loop l_R is visualized as two disjoint closed loops (Case I, the standard a -character (recall that the a -character coincides to the standard character \mathbf{I}_a) on l is trivial) and the covering l_R over l is disconnected, or, as a single connected loop (Case II, the \mathbf{I}_a -character on l is non-trivial). In Case II, we may see l_R as a segment on $R_{reg;m}^{16}$ with involutive end-points by T_{R_m} . Moreover, in the case the class l_* is represented by a segment l_R , such that end-points of the segment are T_{R_m} -involutive, or, the projection of l_R on the circle fibre are \mathbf{I}_a -involutive. We assume that end-points of l_R are not on the section.

Denote by $\sharp(l) \in \{0, 1\}$ the number of intersection points of l_R with the positive base vector of cross-section λ^1 . Take the involution $T_{\hat{q}}$ in the covering by the left vertical arrow of (16) and look the lift of $T_{\hat{q}}$ as an involution $T_{\hat{q};R_{m;reg}}$ on $R_{m;reg}^{16}$ inside the regular domain. Let us prove that numbers $\sharp(l)$, $\sharp(T_{\hat{q};R_{m;reg}}(l))$ coincide in Case I and different in Case II, assuming the the standard $A + \bar{A}$ -character on l is trivial. (In the Case II the numbers $\sharp(l_R)$, $\sharp(T_{\hat{q};R_{m;reg}}(l_R))$ are different and $[l]_* \in H^-(Q_{m;reg})$.)

Let us consider the Case I, recall $l \in H^+(R_{m;reg})$. The bundles $\kappa_A, \kappa_{\bar{A}}$ over l_R are trivial (Subcase 1), or, non-trivial (Subcase 2) simultaneously. Take a coordinate system at the initial point of the path. The section ε over each component of l_R is projected on the spherical fibre, which is the projection on the circle in the subcase 1. The number $\sharp(l_R)$ for the first component depends on a number of rotation of the section, but the number $\sharp(T_{\hat{q};R_{m;reg}}(l_R))$ is also depended, because the sections are antipodal.

In the subcase 1 of the trivial bundles the section λ^1 is represented by a constant section projected on the circle fiber to the point e_1 . The involution $T_{\hat{q};R_{m;reg}}$ transforms the section into its antipodal point $-e_1$ on the circle. In the subcase 2, we will see a closed loop as a twisted cylinder, where "twisted" means a rotation of its generator through the angle $\pm\pi$ with respect to the coordinates in $S(\kappa \oplus \kappa_{\bar{a}})$. In this case the section is translated by $T_{\hat{q};R_{m;reg}}$ into the antipodal twisted cylinder.

Assume a component l_R is represented by a curve at the point e_1 . Take the involution $T_{\hat{q};R_{m;reg}}$ and take the images of l_R and of λ^1 by the involution. The curve $T_{\hat{q};R_{m;reg}}(l_R)$ is a constant at the point e_1 , the section $T_{\hat{q};R_{m;reg}}(\lambda^1)$ is a constant at the point $-e_1$, (recall, the bundle $\nu_{\hat{M}} \otimes \lambda^1$ is twisted by the involution $T_{\hat{q}}$). For the curves in the example we get $\sharp(l_R) = 0$; $\sharp(T_{\hat{q};R_{m;reg}}(l_R)) = 0$.

As the conclusion, in the Case *I*, subcase 1, we get:

$$\sharp(l_R) = \sharp(T_{\hat{q}; R_{m; reg}}(l_R)).$$

The case *I*, subcase 2 is analogous.

Let us consider the Case *II*. We may assume that the both standard A , \hat{A} -characters on l_R are trivial (this case is analogous to the subcase 1 in Case *I*; the case when the both characters are non-trivial is analogous to the subcase 2, Case *I*). The curve l_R is represented by a constant segment at the point $e_1 + e_2$, the section $\lambda^!$ is represented by a constant at the point $e_1 + e_2$. The curve $T_{\hat{q}; R_{m; reg}}(l_R)$ is a constant at the point $e_1 + e_2$, the section $T_{\hat{q}; R_{m; reg}}(\lambda^!)$ is a constant at the point $-e_1 - e_2$. We get: $\sharp(l_R) = 1$; $\sharp(T_{\hat{q}; R_{m; reg}}(l_R)) = 0$ As the conclusion, in the Case *II* we get:

$$\sharp(l_R) = \sharp(T_{\hat{q}; R_{m; reg}}(l_R)) + 1.$$

The calculation above means that $\sharp : H^+(Q_{m; reg}^{16}) \rightarrow \mathbb{Z}/2$ is a homomorphism, which is an \mathbf{I}_a -character on $\mathbf{I}_{A, \hat{A}}$. This exactly means that on an arbitrary class in H^\oplus (see the denotation below) the homomorphism \sharp changes its value with respect to the involution $T_{\hat{q}} : H_1(Q_{m; reg}^{16}) \rightarrow H_1(Q_{m; reg}^{16})$, iff the class is not in the kernel of the \mathbf{I}_a -character.

Denote by

$$H^\oplus(K_m^{14}) = [H^+(K_m^{14}) \cap Ker(\mathbf{I}_a)] \cup [H^-(K_m^{14}) \cap \overline{Ker(\mathbf{I}_a)}].$$

(Similar denotations for various manifolds is used below.) Obviously, the standard character \mathbf{I}_A on $H^\oplus(K_m^{14})$, (\mathbf{I}_A is $\mathbf{I}_{A, \hat{A}} \rightarrow \mathbf{I}_A$) admits the same property. This implies that the homomorphism $\sharp + \mathbf{I}_A : H^\oplus(R_{m; reg}^{16}) \rightarrow \mathbb{Z}/2$ is invariant with respect $T_{\hat{q}} : H^\oplus(R_{m; reg}^{16}) \rightarrow H^\oplus(R_{m; reg}^{16})$, this homomorphism is a cohomology class in $H^\oplus(\hat{R}_{m; reg}^{16}) = H^\oplus(\hat{K}_m^{14})$.

As the result, there exists a homomorphism $\hat{\beta} : H_1(\hat{K}_m^{14}) \rightarrow \mathbb{Z}/2$, which is defined as the canonical extension of $\sharp + \mathbf{I}_A$. An extension is defined by two possible ways, let us take $\hat{\beta}$ with a regular condition, using α , described below.

Take the pull-back $\alpha = \hat{q}^*(\hat{\beta})$ of $\hat{\beta}$ by the covering \hat{q} . The restriction of the cocycle \sharp on $H^\oplus(K_m^{14})$ is extended to the cocycle on $H^\oplus(\hat{M}^{15} \setminus U(\hat{L}^{13}))$, because the section $\lambda^!$ on this manifold $\hat{M}^{15} \setminus U(\hat{L}^{13})$ is well-defined. To calculate the pull-back $\sharp^!$ of \sharp by the covering \hat{q} take a generic intersection of the section on $\hat{M}^{15} \setminus U(\hat{L}^{13})$ with itself. Obviously, the extension $\sharp^!$ coincides with $w_1(\hat{M}^{15} \setminus U(\hat{L}^{13}))$, which is trivial on $H^+(\hat{M}^{15} \setminus U(\hat{L}^{13}))$ and is non-trivial on $H^-(\hat{M}^{15} \setminus U(\hat{L}^{13}))$.

An extension of $\sharp^!$ determines the corresponding extension of \sharp . Because $\alpha = \sharp^! + \mathbf{I}_a^! = p$ on $H^1(\hat{M}^{15} \setminus U(L^{\hat{1}3}))$, the cocycle $\hat{\beta}$ is well-defined and satisfies the condition $\hat{q}^*(\hat{\beta}) = \alpha$. Lemma 7 is proved. \square

To get a construction of η_4^{sf} one may start with a simplification, that the Euler class \tilde{L}^{13} of $\nu_{\hat{M}}$ is empty. With this assumption we may see that $\hat{M}^{15} = \hat{N}^{15}$ is a boundary of the total space of the disk bundle associated with λ (looks like a generalized Möbius band), immersed into \mathbb{R}^{16} , and therefore the immersion $g : \hat{M}^{15} \looparrowright \mathbb{R}^{17}$ admits a section by vector inside the band). This gives the required element $\eta_4^{sf} \in Imm^{sf}(15, 1)$ by the composition $M^{15} \rightarrow \hat{M}^{15} \looparrowright \mathbb{R}^{17}$, analogously with Section 1.

Let us consider a general case. The manifold \hat{N}^{15} (13), which is equipped with the characteristic class $p_{\hat{N}} \in H^1(\hat{N}^{15})$ by Lemma 7 is well-defined. Let us prove that $p_{\hat{N}}^{15}$ is the fundamental class of this closed manifold. By the construction, \hat{N}^{15} (see the formula (13)) is the result of a surgery of \hat{M}^{15} by a generalized handle with a 14-dimensional bottom. The bottom of the lateral surface of the handle is equipped with the class $p_{\hat{K}}$, which is the pull-back of a class α by the projection of the lateral surface onto the middle submanifold \hat{K}_m^{14} , as it is proved in Lemma 7. Therefore the support of the cohomology class $p_{\hat{K}}$ on the lateral handle \hat{K}^{15} is induced by a mapping into 14-skeleton $\mathbb{R}P^{14} \subset \mathbb{R}P^\infty$. As a result, the degree of the characteristic mapping into 15-skeleton for the class of the mapping $p_{\hat{M}} \mapsto p_{\hat{N}}$ by the surgery $\hat{M}^{15} \mapsto \hat{N}^{15}$ remains fixed, and the characteristic number of $p_{\hat{N}}$ and of $p_{\hat{M}}$ coincide. The required equation

$$\langle p_{\hat{N}}^{15}; [\hat{N}^{15}] \rangle = 1 \quad (21)$$

is deduced from (8).

An immersion $\phi_{\hat{N}} : \hat{N}^{15} \looparrowright \mathbb{R}^{17}$ is well-defined, using $p_{\hat{N}}$ with the equation (21) and is determined the required element in $Imm^{sf}(15, 1)$, which will be denoted by η_4^{sf} . Let us repeat reasoning from the section 1 and prove that the element η_4^{sf} has the non-trivial Hopf invariant.

The construction of the elements $\eta_l^{sf} \in Imm^{sf}(2^l - 1, 1)$, $l \geq 5$ is completely analogous to the case $l = 4$.

The Mahowald element $\eta_4^{fr} \in Imm^{fr}(16, 1)$ is well-defined. It is not difficult to prove, using the calculation of the secondary cohomology operation by the Khan-Priddy transfer (see [E]), that the element η_4^{fr} is detected by the corresponding secondary cohomology operation, described in [M].

3 The Hopf invariant one element $\zeta_l^{sf} \in Imm^{sf}(2^l - 1, 2^l - 1)$

Let us prove Theorem 3. Consider an immersion $\hat{\phi} : \hat{N}^{2^l-1} \looparrowright \mathbb{R}^{2^l}$, where $p \in H^1(\hat{N}^{2^l-1})$, $\langle p^{2^l-1}; [\hat{N}] \rangle = 1$, $l \geq 3$, such an immersion is constructed in the section 2. Using this data we construct a self-transversal immersion $\varphi : L^{2^l-1} \looparrowright \mathbb{R}^{2^{l+1}-2}$ with an odd number of self-intersection points and a stable skew-framing $\Xi : \nu_\varphi \oplus n\kappa \simeq (2^l - 1 + n)\kappa$, where κ is the line orienting bundle over L^{2^l-1} . Then we calculate the obstruction to reduce the stable skew-framing Ξ to a skew-framing and we shall prove that there exists a modification $\varphi \mapsto \varphi_1$ of stably skew-framed immersions by a regular surgery, such that the stably skew-framed immersion φ_1 is a skew-framed immersion. The surgery preserves the set of self-intersection points of φ . This proves the theorem.

Let us consider the characteristic mappings $p : \hat{N}^{2^l-1} \rightarrow \mathbb{R}P^\infty$ $\hat{\kappa} : \hat{N}^{2^l-1} \rightarrow \mathbb{R}P^\infty$, where $\hat{\kappa}$ is the orienting cohomology class of the manifold \hat{N}^{2^l-1} , p is a classifying mapping of the given cohomology class. Let us denote by

$$\hat{\mu} : \hat{N}^{2^l-1} \rightarrow \mathbb{R}P^\infty \quad (22)$$

a classifying mapping with the characteristic class $\hat{\mu} = \hat{\kappa} + p$. The mapping $\hat{\mu}$ is the classifying mapping of the line bundle $\hat{\kappa} \otimes p$ over \hat{N}^{2^l-1} . Consider the submanifold $\hat{K}^{2^l-3} \subset \hat{N}^{2^l-1}$, which is defined as the regular preimage by $\hat{\mu}$ of the standard codimension 2 submanifold: $\mathbb{R}P^{\infty-2} \subset \mathbb{R}P^\infty$. The manifold \hat{K}^{2^l-3} is assumed to be connected. Denote by $U(\hat{K}^{2^l-3})$ a thin regular neighbourhood of this manifold in \hat{N}^{2^l-1} .

A generic immersion by a small generic (which is special near \hat{K}^{2^l-3}) alteration of the compressed mapping $\phi_1 = I \circ \hat{\phi} \circ p : N^{2^l-1} \looparrowright \mathbb{R}^{2^{l+1}-2}$, where $p : N^{2^l-1} \rightarrow \hat{N}^{2^l-1}$ is the covering with the prescribed cohomology class, $\hat{\phi} : \hat{N}^{2^l-1} \looparrowright \mathbb{R}^{2^l}$ is given, $I : \mathbb{R}^{2^l} \subset \mathbb{R}^{2^{l+1}-2}$ is the standard embedding. Define a regular surgery $\phi_1 \mapsto \phi_2$ using a generalized handle, which is defined as the disk-line bundle over the manifold $U(\hat{K}^{2^l-3})$, associated with the covering $p_K : K^{2^l-3} \rightarrow \hat{K}^{2^l-3}$. The handle is taken along with the generator along a line bundle with the class p_K . As a result, we shall prove, that the immersion ϕ_2 become stably skew-framed in the ambient space $\mathbb{R}^{2^{l+1}-2+C_{2^l-1}-(2^{l+1}-2-(2^l-1)+1)} = \mathbb{R}^{(2^l-1)+C_{2^l-1}+1}$, where C_{2^l-1} is the Hurwitz number (the minimal power of 2, such that the bundle $C_{2^l-1}\gamma \rightarrow \mathbb{R}P^{2^l-1}$ is isomorphic to the trivial bundle, where γ is the canonical line bundle over the standard skeleton $\mathbb{R}P^{2^l-1}$). Let us describe the surgery $N^{2^l-1} \mapsto N_2^{2^l-1}$.

Consider the immersions $\hat{\phi}_1 = I \circ \hat{\phi} : \hat{N}^{2^l-1} \looparrowright \mathbb{R}^{2^{l+1}-2}$, $\hat{\phi}_1^{st} = I_1 \circ I \circ \hat{\phi} : \hat{N}^{2^l-1} \looparrowright \mathbb{R}^{2^{l+1}-2} \subset \mathbb{R}^{2^l+C_{2^l-1}}$. Over the immersed manifold $\hat{\phi}_1$ let us consider

the trivial bundle $C_{2^l-1}\varepsilon$, which is the restriction of the normal bundle of the embedding $\mathbb{R}^{2^l} \subset \mathbb{R}^{2^l+C_{2^l-1}}$ and let us consider the direct first factor of the normal bundle $C_{2^l-1}p \subset \nu_{\phi_1^{st}}$ by the isomorphism $C_{2^l-1}\varepsilon \simeq C_{2^l-1}p$, restricted over the submanifold $\hat{K}^{2^l-3} \subset \hat{N}^{2^l-1}$. Without loss of a generality, let us assume, that the first factor line bundle p of the normal bundle of $\hat{\phi}_1^{st}$ is inside the hyperspace $\mathbb{R}^{2^{l+1}-2} \subset \mathbb{R}^{2^l+C_{2^l-1}}$.

Let us consider a deformation of the immersion ϕ_1 into a self-transversal immersion (without a change of the denotation) (as it is proved above with the odd number of self-intersection points, each self-intersection is outside $\phi_1(U(K^{2^l-3}))$), which over \hat{K}^{2^l-3} coincides with a small deformation along the fibre of the line bundle p . The result of the surgery by the handle is the transformation $\phi_1(N^{2^l-1}) \mapsto \phi_2(N_2^{2^l-1})$. We glue the disk bundle, associated with p , over the boundary of the disk normal bundle of the codimension 2 submanifold $i_{\hat{K}} : \hat{K}^{2^l-3} \subset \hat{N}^{2^l-1}$. The stabilization of the immersion ϕ_2 in the space $\mathbb{R}^{2^l+C_{2^l-1}}$ let us denote by ϕ_2^{st} .

The handlebody consists the boundary, which consists of the lateral handle, denoted by Q^{2^l-1} (the lateral part of the boundary of the body) and the bottom. The lateral handle contains the boundary, which is a manifold of the dimension $2^l - 2$, this manifold is connected because \hat{K}^{2^l-3} is connected and the restriction $p|_{\hat{K}^{2^l-3}}$ is non-trivial. The boundary $\partial(Q^{2^l-1})$ of the lateral handle is identified with the boundary of the regular neighbourhood of the submanifold i_K , the double covering over $i_{\hat{K}}$.

The lateral handle has the normal bundle is equipped with the line bundles p, κ , because the lateral handle is projected onto $U(\hat{K}^{2^l-3})$, the pull-back of $\hat{\kappa}$, the line bundle over the base $\partial(U(\hat{K}^{2^l-3}))$, is denoted by κ . Moreover, the terms p and $\hat{\kappa}$ over the lateral handle are isomorphic: this fact is satisfied, because $\partial(U(\hat{K}^{2^l-3}))$ admits a foliation across the meridians and the both classes are defined as rotation classes over this foliation through the angle $\pi \pmod{2\pi}$. The normal bundle of the lateral handle in $\mathbb{R}^{2^l+C_{2^l-1}}$ is the following:

$$(C_{2^l-1} - 1)p \oplus \varepsilon \oplus \kappa, \quad (23)$$

where the term ε is the tangent vector to the handlebody, which is perpendicular to the lateral handle. The factor ε in the formula (23) can be replaced into π if we assume that the boundary conditions over end-points of segments, the generator of the handle, is changed into the opposite. The formula (23) is modified as the following:

$$(C_{2^l-1} - 1)p \oplus p \oplus \hat{\kappa}. \quad (24)$$

The formula (24) looks more natural with respect to the formula (23), because the boundary condition on the bottom of the lateral handle for (24) is satisfied. The formula (24) describes the normal bundle of the immersion ϕ_2^{st} .

Let us modify the formula (24). Recall that the bundle κ is the pull-back of the line bundle $\hat{\mu}$ over $\hat{N}^{2^l-1} \setminus U(\hat{K}^{2^l-3})$, described by (22). The target bundle $\hat{\mu}$ admits an integer characteristic classes, in particular, over $\hat{N}^{2^l-1} \setminus U(\hat{K}^{2^l-3})$ the following equation

$$2\hat{\mu} = 2\varepsilon \quad (25)$$

is satisfied.

Let us do the following change:

$$C_{2^l-1}p \oplus \kappa_1 \mapsto (C_{2^l-1} + 1)\kappa_1, \quad (26)$$

because $C_{2^l-1}p$, $C_{2^l-1}\kappa_1$ are isomorphic as the trivial bundles. The skew-framing of the immersion ϕ_2^{st} is given by the formula: $\Psi_2 : \nu_{\phi_1^{st}} \equiv (C_{2^l-1} + 1)\kappa_1$. Let us prove that the skew-framing Ψ_2 is extended into a stable skew-framing of the immersion $\phi_2 : N^{2^l-1} \hookrightarrow \mathbb{R}^{2^{l+1}-2}$ inside $\mathbb{R}^{2^l+C_{2^l-1}}$.

Using the formula (25) and the modification (26), we may rewrite the formula (24) into the following:

$$\Psi_1 : \nu_{\phi_1} \oplus (C_{2^l-1} + 1 - (2^l - 1))\varepsilon = (C_{2^l-1} + 1)\kappa_1. \quad (27)$$

Because the bundle κ_1 is the pull-back of a bundle over $\hat{N}^{2^l-1} \setminus U_{\hat{K}}$, analogously to (25), one gets: $(C_{2^l-1} - 2^l + 2)\varepsilon \simeq (C_{2^l-1} - 2^l + 2)\kappa_1$.

After we apply this isomorphism to the right-hand side of the equation (27), we get the following formula for the stable skew-framing Ψ_1^{st} of the of the codimension $2^l - 1$ immersion ϕ_1 :

$$\Psi_1^{st} : \nu_{\phi_1} \oplus (C_{2^l-1} + 1 - (2^l - 1))\kappa_1 = (C_{2^l-1} + 1)\kappa_1. \quad (28)$$

Let us calculate the total obstruction for destabilization of the stable skew-framing Ψ_1^{st} into a skew-framing Ψ_1 . The total obstruction is an integer, denoted by $o(\Psi_1)$, this obstruction is, in particular, investigated in [A-C-R]. Let us recall the definition of the total obstruction for a connected $N_1^{2^l-1}$.

Let us consider an isotropy of the family of skew-sections Ψ_1^{st} to a special $C_{2^l-1} + 1 - (2^l - 1)$ -family of skew-sections. For a special family all sections, except, probably, the only skew-section in the left hand side of the equation (28) coincides with the corresponding skew-sections in the right-hand side

of the equation. Then the integer obstruction $o(\Psi_1)$ to a deformation of the last skew-section in the right-hand side of the equation into the corresponding skew-section in the left-hand side of the equation is well-defined.

Let us prove that

$$o(\Psi_1) = 0 \pmod{2}. \quad (29)$$

Take an arbitrary skew-section in the left-hand side of the formula and let us prove that the algebraic number of transversal intersection points of this skew-section with the subbundle of the codimension $2^l - 1$, generated by skew-sections in the right-hand side of the formula is an even integer. This is following from the the facts: the skew-section (in fact, this skew-section is a skew-section line bundle κ) is a pull-back from $\hat{N}^{2^l-1} \setminus \hat{K}^{2^l-3}$; the codimension 2^{l-1} family of cross-sections is a pull-back bundle from the same base manifold. This proves the formula (29). Denote the value of this obstruction by $o(\Psi_1) = 2k$, $k \in \mathbb{Z}$. Consider the standard immersion $f : S^{2^l-1} \looparrowright \mathbb{R}^{2^{l+1}-2}$ with the only self-intersection point. For the immersion f a stable framing is well-defined, this stable skew-framing can be considered as the corresponding stable skew-framing. For this skew-framing Ξ^{st} we get: $o(\Xi^{st}) = \pm 1$, where a sign is related of a coorientation of the standard immersion f . Take an even $-2k$ algebraic copies of the skew-framed immersion f , join this copies to the immersion (ϕ_1, Ψ_1) , we get a stably skew-framed immersion (ϕ_2, Ψ_2) with $o(\Psi_2) = 0$, which is connected in its stably skew-framing cobordism class. As the result, we get a skew-framed immersion $\phi_3 : N_3^{2^l-1} \looparrowright \mathbb{R}^{2^{l+1}-2}$ with an odd number of self-intersection points. Theorem 3 is proved.

4 Discussion

In the section 2 an element $\eta_l^{sf} \in Imm^{sf}(2^l - 1, 1)$, $l \geq 3$ is constructed. This element is associated by the Khan-Priddy transfer with the element $\eta_l \in \pi_{2^{l+1}+2}(S^{2^l+2}) \simeq \Pi_{2^l-2}$, which is not desuspended 5 times from the stable range into un unstable range.

The same time, for an arbitrary $d \geq 0$ there exists $l_0 = l_0(d)$ such that for an arbitrary $l \geq l_0$ an arbitrary element in the stable homotopy group of spheres Π_{2^l-2} admits a d -fold desuspension, as it is proved in [A1]. In [L-R] it is proved, that if the group Π_{126} there exists an element with the Kervaire invariant one then in the residue class of this element there exists an element, which is not desuspended 11 times into an unstable range.

The author is grateful S.A.Melikhov for discussions of the Section 2.

References

- [A] Petr M. Akhmet'ev, *Kervaire Problems in Stable Homotopy Theory*, <http://de.arxiv.org/abs/1608.01206v1>
- [A-C-R] Akhmetiev Peter M., Cencelj Matija, Repovs Dusan 'The Arf-Kervaire invariant of framed manifolds as an obstruction to embeddability', *Atti della Accademia Perloritana dei Pericolanti. Classe di Scienze Fisiche, Matematiche e Naturali* (2022) Vol 100, N2.
- [A1] P.M.Akhmet'ev, *Geometric approach to stable homotopy groups of spheres II. The Kervaire invariant*, <http://arxiv.org/abs/1011.5717v3>
- [A-E] P.M.Akhmet'ev and P.J.Eccles, *The relationship between framed bordism and skew-framed bordism*, *Bull. London Math. Soc.*, vol 39 (2007) 473-481.
- [A-F] P.M.Akhmet'ev, O.D.Frolkina, "On properties of skew-framed immersions cobordism groups", *Fundam. Prikl. Mat.*, 21:5 (2016), 19-46 <http://arxiv.org/abs/1712.00959v1>.
- [B-J-M] M.G. Barratt, J.D.S. Jones and M.E. Mahowald, *The Kervaire invariant one problem*, *Contemporary Mathematics* Vol 19, (1983) 9-22.
- [B] W.Browder, *The Kervaire invariant of framed manifolds and its generalization*, *Ann. of Math.*, (2) 90 (1969) 157-186.
- [E] P.J.Eccles, *Codimension One Immersions and the Kervaire Invariant One Problem*, *Math. Proc. Cambridge Phil. Soc.* vol 90 (1981) 483- 493.
- [E-Z] Peter J. Eccles, Hadi Zare, *The Hurewicz image of the η_i family, a polynomial subalgebra of $H_*(\Omega_0^{2^{i+1}-8+k}S^{2^i-2})$* arXiv:0909.3791v1 [math.AT] 21 Sep 2009.
- [H-H-R] M. A. Hill, M. J. Hopkins and D. C. Ravenel, *On the non-existence of elements of Kervaire invariant one*, *Ann. of Math.* (2) 184 (2016), 1-262.
- [H] M.W.Hirsch, *Immersion of manifolds*, *Trans.Amer.Math.Soc.*, 93, (1960) 242-276.
- [K] U.Koschorke, *Multiple points of immersions and the Kahn-Priddy theorem*, *Math.Z.*, 169 (1979) 223-236.

- [K1] U.Koschorke, *Nonstable and stable monomorphisms of vector bundles*.
Topology and Applications 75 (1987) 261-286
- [L-R] K.Y.Lam, D.Randall, *Block bundle obstruction to Kervaire Invariant one*, Contamporary Mathematics, Vol. 407 (2006) 163-169.
- [M] M.Mahowald, *A new infinite family in ${}_2\pi_*^S$* , Topology vol. 16 (1977) 249-256.
- [M-T] Mosher R.S. Tangora M.C., *Cohomology operations and their applications in homotopy theory*, N.Y. Harper and Row, Publishers (1968).
- [S] V.P. Snaith, *Stable homotopy around the Arf-Kervaire invariant*, Birkhauser Progress on Math. Series vol. 273 (April 2009).
- [Sz] A.Szücs, *Cobordism of immersions and singular maps, loop spaces and multiple points*, Geometric and Algebraic Topology, Banach Center Publ., 18 PWN, Warsaw (1986) 239-253.