

GENERALIZED CHOQUARD EQUATION WITH POTENTIAL VANISHING AT INFINITY

LUCAS DA SILVA* AND MARCO SOUTO

ABSTRACT. In this paper we investigate the existence of solution for the following generalized Choquard equation

$$-\Delta_{\Phi}u + V(x)\phi(|u|)u = \left(\int_{\mathbb{R}^N} \frac{K(y)F(u(y))}{|x-y|^{\lambda}} dy \right) K(x)f(u(x)), \quad x \in \mathbb{R}^N$$

where $N \geq 3$, $\lambda > 0$ is a positive parameter, $V, K \in C(\mathbb{R}^N, [0, \infty))$ are nonnegative functions that may vanish at infinity, the function $f \in C(\mathbb{R}, \mathbb{R})$ is quascritical and $F(t) = \int_0^t f(s)ds$. This work incorporates the reflexive and non-reflexive cases taking into account from Orlicz-Sobolev framework. The non-reflexive case occurs when the N -function $\tilde{\Phi}$ does not verify the Δ_2 -condition. In order to prove our main results we employ variational methods and regularity results.

1. INTRODUCTION

This paper concerns the existence of weak solution for the following generalized Choquard equation

$$\begin{cases} -\Delta_{\Phi}u + V(x)\phi(|u|)u = \left(\int_{\mathbb{R}^N} \frac{K(y)F(u(y))}{|x-y|^{\lambda}} dy \right) K(x)f(u(x)), & x \in \mathbb{R}^N \\ u \in D^{1,\Phi}(\mathbb{R}^N) \end{cases} \quad (P)$$

where $N \geq 3$, $\lambda > 0$ is a positive parameter, $V, K \in C(\mathbb{R}^N, [0, \infty))$ are nonnegative functions that may vanish at infinity, the function $f \in C(\mathbb{R}, \mathbb{R})$ is quascritical and $F(t) = \int_0^t f(s)ds$. It is important to recall that

$$\Delta_{\Phi}u = \operatorname{div}(\phi(|\nabla u|)\nabla u),$$

where $\Phi : \mathbb{R} \rightarrow \mathbb{R}$ is a N -function of the form

$$(1.1) \quad \Phi(t) = \int_0^{|t|} s\phi(s)ds,$$

and $\phi : (0, \infty) \rightarrow (0, \infty)$ is a C^1 function verifying some assumptions. This type of problem driven by a N -function Φ appears in a lot of physical applications, such as Nonlinear Elasticity, Plasticity, Generalized Newtonian Fluid, Non-Newtonian Fluid and Plasma Physics. The reader can find more details about this subject in [10], [15], [30] and their references.

Recently, motivated by the applications above, the quasilinear problems driven by a N -function Φ has been frequently studied. We would like to cite [13,14,37,34, 39,25,26,12,20,41,45] and their references. In all of these works, the so-called Δ_2 -condition was assumed on Φ and on $\tilde{\Phi}$ (Complementary function of Φ), which ensures that the Orlicz-Sobolev space $W^{1,\Phi}(\Omega)$ and $D^{1,\Phi}(\Omega)$ are reflexive Banach spaces. This assertion is used several times in order to get a

2020 *Mathematics Subject Classification*. Primary: 35J15, 35J62, 35J60.

Key words and phrases. Orlicz-Sobolev spaces; Variational methods; Choquard equation; nonreflexive spaces.

M. Souto was partially supported by CNPq/Brazil 309.692/2020-2.

*Corresponding author.

nontrivial solution for elliptic problems taking into account the weak topology and the classical variational methods to C^1 functionals.

In recent years, problems without the Δ_2 -condition of the function $\tilde{\Phi}$ are being studied. This type of problem brings us many difficulties when we intend to apply variational methods. For example, in the paper [21], the energy functional associated with the problem might not be C^1 , so classical minimax type results cannot be used here, however, Silva and Souto, showed that the energy functional verifies the hypotheses of the Ghoussoub-Preiss theorem. In [6], Alves and Carvalho make use of the minimax theory developed by Szulkin [3]. Some other works can be mentioned here, we refer the interested readers to [9].

Based on the papers above, we assume that $\phi : (0, \infty) \rightarrow (0, \infty)$ is C^1 and satisfies the following hypotheses:

$$(\phi_1) \quad t \mapsto t\phi(t) \text{ is increasing for } t > 0.$$

$$(\phi_2) \quad \lim_{t \rightarrow 0^+} t\phi(t) = 0 \quad \text{and} \quad \lim_{t \rightarrow +\infty} t\phi(t) = +\infty.$$

$$(\phi_3) \quad 1 \leq \ell = \inf_{t>0} \frac{\phi(t)t^2}{\Phi(t)} \leq \sup_{t>0} \frac{\phi(t)t^2}{\Phi(t)} = m < N, \quad \ell \leq m < \ell^* \quad \text{and} \quad m \neq 1.$$

$$(\phi_4) \quad t \mapsto \frac{\phi(t)}{t^{m-2}} \text{ is nonincreasing for } t > 0.$$

(ϕ_5) There is a constant $C > 0$ satisfying

$$\Phi(t) \geq C|t|^m, \quad \text{for } t \in \mathbb{R}.$$

In this work the parameter $\lambda \in (0, N) \cap (0, 2N - \frac{2N}{m})$ and $s = \frac{2N}{2N-\lambda}$. Having that noted, we will also assume that Φ in addition satisfies the following hypothesis:

(ϕ_6) The function $\Phi_*(|t|^{1/s})$ is convex in \mathbb{R} .

As it is known, the Choquard equation (P) appears in many physical problems. The case $N = 3$, $\Phi(t) = t^2/2$, $\lambda = 1$, $V = K = 1$ and $f(t) = t$ goes back to the description of the quantum theory of a polaron at rest by Pekar [42] in 1954 and the modeling of an electron trapped in its own hole (in the work of Choquard in 1976), in a certain approximation to Hartree–Fock theory of one-component plasma [11]. This equation was also proposed by Penrose in [38] as a model of self-gravitating matter and is known in that context as the Schrödinger–Newton equation. In the mathematical context, the existence of non-trivial solutions for this case was proved by Lieb, Lions and Menzala [11, 35, 16] using variational methods and also by ordinary differential equation techniques [33, 19, 36]. Later, the result was extended to the case $N \geq 3$, $0 < \lambda < N$, $V = K = 1$ and $f(t) = |t|^{p-2}t$ with $(2N - \lambda)/N < p < (2N - \lambda)/(N - 2)$, see Ma and Zhao [23] and Moroz and Schaftingen [44].

Recently, in [7], Alves, Rădulescu and Tavares studied the equation (P) with $V = K = 1$ using different assumptions on the N -function Φ . In this work, the authors aimed to show that the variational methods could be applied to establish the existence of solutions assuming that the N -function Φ satisfies the conditions (ϕ_1) – (ϕ_3) with $\ell > 1$. One of the main difficulties was to prove that the energy functional associated with equation (P) is differentiable. However, good conditions involving the function f made it possible to show the differentiability of the energy functional and consequently allowed to guarantee the existence of a solution through the mountain pass theorem. It is also worth mentioning that in the same work, Alves, Rădulescu and Tavares extended the result to the case where $K = 1$ and V is one of the following

potentials: periodic function, asymptotic periodic function, coercive or Bartsch-Wang-like potential.

When it comes to the non-autonomous cases of the equation (P) , an important class of problems to it is the zero mass case which occurs if the potential V vanishes at infinity. In [4], Alves, Figueiredo and Yang studied a particular case of the equation (P) with $\Phi(t) = t^2/2$ and $K = 1$, where the potential $V \in C(\mathbb{R}^N, \mathbb{R})$ is nonnegative and vanishes at infinity whose nonlinearity $f \in C(\mathbb{R}, \mathbb{R})$ has a supercritical growth and verifies some appropriate assumptions. Still in this case, the authors assumed that $\lambda \in (\frac{N-2}{2}, N)$ and showed the existence of a solution using the penalization method introduced by del Pino and Felmer in [24].

Recently, Chen and Yuan in [40], investigated the equation (P) in the case where $\Phi(t) = t^2/2$, where the function $f \in C(\mathbb{R}, \mathbb{R})$ is quascritical and the potentials $V, K \in C(\mathbb{R}^N, \mathbb{R})$ satisfy hypotheses similar to (K_0) , (K_1) , (K_2) or (K_3) described below. Based on the non-Nehari manifold methods developed by Tang in [46] and a diagonal approach, Chen and Yuan proved the existence of ground state solution under some appropriate noncritical assumptions on f . For more investigations involving the equation (P) , we refer the interested readers to [43].

To the best of our knowledge, there seem to be no results on ground state solutions for the equation (P) with V vanishing at infinity and Φ verifying or not the Δ_2 -condition. In view of this, driven by the importance of Choquard's equations and the problems in non-reflexive spaces mentioned in the works above, in particular [21], [40], the first objective of this article is to guarantee the existence of a ground state solution for the equation (P) and, after that, to prove regularity results. To state the first results, we introduce the following hypotheses about the potentials V and K :

(K_0) $V > 0$, $K \in L^\infty(\mathbb{R}^N)$ and K is positive almost everywhere.

(I) If $\{A_n\} \subset \mathbb{R}^N$ is a sequence of Borelian sets such that $\sup_n |A_n| < +\infty$, then

$$\lim_{r \rightarrow +\infty} \int_{A_n \cap B_r^c(0)} K(x)^s dx = 0, \quad \text{uniformly in } n \in \mathbb{N}. \quad (K_1)$$

(III) One of the below conditions occurs:

$$\frac{K}{V} \in L^\infty(\mathbb{R}^N) \quad (K_2)$$

or there are $b_1, b_2 \in (m, \ell^*)$ and a N -function $B(t) = \int_0^{|t|} b(\tau)\tau d\tau$ verifying the following properties:

$$(B_1) \quad t \mapsto tb(t) \text{ is increasing for } t > 0.$$

$$(B_2) \quad \lim_{t \rightarrow 0^+} tb(t) = 0 \quad \text{and} \quad \lim_{t \rightarrow +\infty} tb(t) = +\infty.$$

$$(B_3) \quad b_1 \leq \frac{b(t)t^2}{B(t)} \leq b_2, \quad \text{for all } t > 0$$

$$(B_4) \quad \text{The function } B(|t|^{1/s}) \text{ is convex in } \mathbb{R}$$

and

$$\frac{K(x)^s}{H(x)} \rightarrow 0 \quad \text{as } |x| \rightarrow +\infty \quad (K_3)$$

where $H(x) = \min_{\tau > 0} \left\{ V(x) \frac{\Phi(\tau)}{B(\tau)} + \frac{\Phi_*(\tau)}{B(\tau)} \right\}$.

Hereafter, we say that $(V, K) \in \mathcal{K}_1$ if (I), (II) and (K_2) hold. When (I), (II) and (K_3) hold, then we say that $(V, K) \in \mathcal{K}_2$. These hypotheses were inspired by the paper [5] which deals with a local case involving the Laplacian and which were generalized in [21]. In a first moment, we study the equation (P) assuming that $(V, K) \in \mathcal{K}_1$. To this end, we must assume some conditions on f .

We will consider $A : \mathbb{R} \rightarrow [0, +\infty)$ being a N -function given by $A(w) = \int_0^{|w|} ta(t)dt$ where $a : (0, +\infty) \rightarrow (0, +\infty)$ is a function satisfying the following conditions:

$$(A_1) \quad t \mapsto ta(t) \text{ is increasing for } t > 0.$$

$$(A_2) \quad \lim_{t \rightarrow 0^+} ta(t) = 0 \quad \text{and} \quad \lim_{t \rightarrow +\infty} ta(t) = +\infty.$$

(A₃) There exist $a_1, a_2 \in [m, \ell^*]$ such that

$$(1.2) \quad a_1 \leq \frac{a(t)t^2}{A(t)} \leq a_2, \quad \forall t > 0.$$

(A₄) The function $A(|t|^{1/s})$ is convex in \mathbb{R} .

Let f be a continuous function satisfying the following conditions:

$$(f_1) \quad \lim_{t \rightarrow 0} \frac{f(t)}{\left(\frac{1}{s}a(|t|)|t|^{2-s}\right)^{1/s}} = 0 \quad \text{and} \quad \lim_{t \rightarrow +\infty} \frac{f(t)}{\left(\frac{1}{s}\phi_*(|t|)|t|^{2-s}\right)^{1/s}} = 0.$$

$$(f_2) \quad tf(t) - \frac{m}{2}F(t) \text{ is an increasing function in } (0, +\infty).$$

$$(f_3) \quad f(t) \geq 0 \text{ for } t \geq 0 \text{ and } f(t) = 0 \text{ for } t \leq 0.$$

$$(f_4) \quad \lim_{|t| \rightarrow \infty} \frac{F(t)}{|t|^{\frac{m}{2}}} = +\infty.$$

Here, unlike Chen and Yuan in [40], the condition $f(t)$ being nondecreasing on \mathbb{R} is not required. Under these conditions, our first main result can be stated as follows.

Theorem 1.1. *Assume that Φ satisfies $(\phi_1) - (\phi_4)$ and $\lambda \in (0, N) \cap (0, 2N - \frac{2N}{m})$. Suppose that $(V, K) \in \mathcal{K}_1$, $(A_1) - (A_4)$ and $(f_1), (f_2), (f_3), (f_4)$ holds. Then, problem (P) possesses a nonnegative ground state solution. If $\lambda < 2m$ and (ϕ_5) hold, then the nonnegative solutions are locally bounded.*

To study the regularity of the solutions provided by Theorem 1.1, we add the following assumptions:

(ϕ_6) There are $0 < \delta < 1$, $C_1, C_2 > 0$ and $1 < \beta \leq \ell^*$ such that

$$C_1 t^{\beta-1} \leq t\phi(t) \leq C_2 t^{\beta-1}, \quad \text{for } t \in [0, \delta].$$

(ϕ_7) There are constants $\delta_0 > 0$ and $\delta_1 > 0$ such that

$$\delta_0 \leq \frac{(\phi(t)t)'}{\phi(t)} \leq \delta_1 \quad \text{for } t > 0.$$

We are in position to state the following regularity result:

Theorem 1.2. *Suppose that Φ satisfies $(\phi_5) - (\phi_6)$. Under the assumptions of Theorem 1.1, if $K \in L^1(\mathbb{R}^N)$ and $\lambda < 2m$, then the problem (P) possesses a $C_{loc}^{1,\alpha}(\mathbb{R}^N)$ positive ground state solution. Furthermore, when $\inf_{x \in \mathbb{R}^N} V(x) > 0$ the solutions $C_{loc}^{1,\alpha}(\mathbb{R}^N)$ go to zero at infinity.*

Still regarding equation (P), to study the case where $(V, K) \in \mathcal{K}_2$ we must assume that f satisfies the conditions (f_2) , (f_3) , (f_4) and

$$(f_5) \quad \limsup_{t \rightarrow 0} \frac{f(t)}{\left(\frac{1}{s}b(|t|)|t|^{2-s}\right)^{1/s}} < \infty \quad \text{and} \quad \lim_{t \rightarrow +\infty} \frac{f(t)}{\left(\frac{1}{s}\phi_*(|t|)|t|^{2-s}\right)^{1/s}} = 0.$$

Under these conditions, the next result of the existence of a nonnegative solution has the following statement:

Theorem 1.3. *Assume that Φ satisfies $(\phi_1) - (\phi_4)$ and $\lambda \in (0, N) \cap (0, 2N - \frac{2N}{m})$. Suppose that $(V, K) \in \mathcal{K}_1$, $(B_1) - (B_4)$ and (f_2) , (f_3) , (f_4) , (f_5) hold. Then, problem (P) possesses a nonnegative ground state solution. If $\lambda < 2m$ and (ϕ_5) hold, then the nonnegative solutions are locally bounded.*

As in Theorem 1.2, the following result guarantees the regularity and positiveness of nonnegative solutions provided by Theorem 1.3.

Theorem 1.4. *Suppose that Φ satisfies $(\phi_5) - (\phi_6)$. Under the assumptions of Theorem 1.3, if $K \in L^1(\mathbb{R}^N)$ and $\lambda < 2m$, then the problem (P) possesses a $C_{loc}^{1,\alpha}(\mathbb{R}^N)$ positive ground state solution. Furthermore, when $\inf_{x \in \mathbb{R}^N} V(x) > 0$ the solutions $C_{loc}^{1,\alpha}(\mathbb{R}^N)$ go to zero at infinity.*

Next, we are interested to study the following Choquard equation with zero mass case

$$-\Delta_{\Phi} u = \left(\int_{\mathbb{R}^N} \frac{K(y)F(u(y))}{|x-y|^{\lambda}} dy \right) K(x)f(u(x)), \quad u \in D^{1,\Phi}(\mathbb{R}^N) \quad (P')$$

To study this equation, we assume that f satisfies (f_2) , (f_3) , (f_4) , and the following condition:

$$(f_6) \quad \lim_{t \rightarrow 0} \frac{f(t)}{\left(\frac{1}{s}\phi_*(|t|)|t|^{2-s}\right)^{1/s}} = 0 \quad \text{and} \quad \lim_{t \rightarrow +\infty} \frac{f(t)}{\left(\frac{1}{s}\phi_*(|t|)|t|^{2-s}\right)^{1/s}} = 0.$$

And then, the results we declare are:

Theorem 1.5. *Suppose that $(\phi_1) - (\phi_4)$ and $(K_0) - (K_1)$ hold. Assume that f satisfies (f_2) , (f_3) , (f_4) and (f_6) . Then the problem (P') possesses a nonnegative ground state solution. If $\lambda < 2m$ and (ϕ_5) hold, then the nonnegative solutions are locally bounded.*

Theorem 1.6. *Suppose that Φ satisfies $(\phi_5) - (\phi_6)$. Under the premises of Theorem 1.5, if $K \in L^1(\mathbb{R}^N)$ and $\lambda < 2m$, then the problem (P') possesses a $C_{loc}^{1,\alpha}(\mathbb{R}^N)$ positive ground state solution. Furthermore, when $\inf_{x \in \mathbb{R}^N} V(x) > 0$ the solutions $C_{loc}^{1,\alpha}(\mathbb{R}^N)$ go to zero at infinity.*

2. BASICS ON ORLICZ-SOBOLEV SPACES

In this section we recall some properties of Orlicz and Orlicz-Sobolev spaces, which can be found in [1,12,27,29]. First of all, we recall that a continuous function $\Phi : \mathbb{R} \rightarrow [0, +\infty)$ is a N -function if:

- (i) Φ is convex;
- (ii) $\Phi(t) = 0 \Leftrightarrow t = 0$;
- (iii) Φ is even;
- (iv) $\lim_{t \rightarrow 0} \frac{\Phi(t)}{t} = 0$ and $\lim_{t \rightarrow +\infty} \frac{\Phi(t)}{t} = +\infty$.

We say that a N -function Φ verifies the Δ_2 -condition, and we denote by $\Phi \in (\Delta_2)$, if there are constants $K > 0$, $t_0 > 0$ such that

$$\Phi(2t) \leq K\Phi(t), \quad \forall t \geq t_0.$$

In the case of $|\Omega| = +\infty$, we will consider that $\Phi \in (\Delta_2)$ if $t_0 = 0$. For instance, it can be shown that $\Phi(t) = |t|^p/p$ for $p > 1$ satisfies the Δ_2 -condition, while $\Phi(t) = (e^{t^2} - 1)/2$.

If Ω is an open set of \mathbb{R}^N , where N can be a natural number such that $N \geq 1$, and Φ a N -function then define the Orlicz space associated with Φ as

$$L^\Phi(\Omega) = \left\{ u \in L^1_{\text{loc}}(\Omega) : \int_{\Omega} \Phi\left(\frac{|u|}{\lambda}\right) dx < +\infty \text{ for some } \lambda > 0 \right\}.$$

The space $L^\Phi(\Omega)$ is a Banach space endowed with the Luxemburg norm given by

$$\|u\|_{L^\Phi(\Omega)} = \inf \left\{ \lambda > 0 : \int_{\Omega} \Phi\left(\frac{|u|}{\lambda}\right) dx \leq 1 \right\}.$$

In the case that Φ verifies Δ_2 -condition we have

$$L^\Phi(\Omega) = \left\{ u \in L^1_{\text{loc}}(\Omega) : \int_{\Omega} \Phi(|u|) dx < +\infty \right\}.$$

The complementary function $\tilde{\Phi}$ associated with Φ is given by the Legendre transformation, that is,

$$\tilde{\Phi}(s) = \max_{t \geq 0} \{st - \Phi(t)\}, \quad \forall t \geq 0.$$

The functions Φ and $\tilde{\Phi}$ are complementary to each other and satisfy the inequality below

$$\tilde{\Phi}(\Phi'(t)) \leq \Phi(2t), \quad \forall t > 0.$$

Moreover, we also have a Young type inequality given by

$$st \leq \Phi(t) + \tilde{\Phi}(s), \quad \forall s, t \geq 0.$$

Using the above inequality, it is possible to establish the following Holder type inequality:

$$\left| \int_{\Omega} uv dx \right| \leq 2 \|u\|_{L^\Phi(\Omega)} \|v\|_{L^{\tilde{\Phi}}(\Omega)}, \quad \text{for all } u \in L^\Phi(\Omega) \text{ and } v \in L^{\tilde{\Phi}}(\Omega).$$

If $|\Omega| < \infty$, the space $E^\Phi(\Omega)$ denotes the closing of $L^\infty(\Omega)$ in $L^\Phi(\Omega)$ with respect to the norm $\|\cdot\|_\Phi$. When $|\Omega| = \infty$, the space $E^\Phi(\Omega)$ denotes the closure of $C_0^\infty(\Omega)$ in $L^\Phi(\Omega)$ with respect to norm $\|\cdot\|_\Phi$. In any of these cases, $L^\Phi(\Omega)$ is the dual space of $E^{\tilde{\Phi}}(\Omega)$, while $L^{\tilde{\Phi}}(\Omega)$ is the dual space of $E^\Phi(\Omega)$. Moreover, $E^\Phi(\Omega)$ and $E^{\tilde{\Phi}}(\Omega)$ are separable and all continuous functional $M : E^\Phi(\Omega) \rightarrow \mathbb{R}$ are of the form

$$M(v) = \int_{\Omega} v(x)g(x)dx, \quad \text{for some function } g \in L^{\tilde{\Phi}}(\Omega).$$

Another important function related to function Φ it is the Sobolev conjugate function Φ_* of Φ defined by

$$\Phi_*^{-1}(t) = \int_0^t \frac{\Phi^{-1}(s)}{s^{(N+1)/N}} ds \text{ for } t > 0 \quad \text{when} \quad \int_1^{+\infty} \frac{\Phi^{-1}(s)}{s^{(N+1)/N}} ds = +\infty.$$

Lemma 2.1. *Consider Φ a N -function of the form (1.1) and satisfying (ϕ_1) and (ϕ_2) . Set*

$$\xi_0(t) = \min\{t^\ell, t^m\} \quad \text{and} \quad \xi_1(t) = \max\{t^\ell, t^m\}, \quad \forall t \geq 0.$$

Then Φ satisfies

$$\xi_0(t)\Phi(\rho) \leq \Phi(\rho t) \leq \xi_1(t)\Phi(\rho), \quad \forall \rho, t \geq 0$$

and

$$\xi_0(\|u\|_\Phi) \leq \int_{\Omega} \Phi(u) dx \leq \xi_1(\|u\|_\Phi), \quad \forall u \in L^\Phi(\Omega).$$

Lemma 2.2. *If Φ is an N -function of the form (1.1) satisfying (ϕ_1) , (ϕ_2) and (ϕ_3) , then*

$$(2.1) \quad \xi_2(t)\Phi_*(\rho) \leq \Phi_*(\rho t) \leq \xi_3(t)\Phi_*(\rho), \quad \forall \rho, t > 0$$

and

$$\xi_2(\|u\|_{\Phi_*}) \leq \int_{\Omega} \Phi_*(u) dx \leq \xi_3(\|u\|_{\Phi_*}), \quad \forall u \in L^{\Phi_*}(\Omega),$$

where

$$\xi_2(t) = \min\{t^{\ell^*}, t^{m^*}\} \quad \text{and} \quad \xi_3(t) = \max\{t^{\ell^*}, t^{m^*}\}, \quad t \geq 0.$$

Lemma 2.3. (2.1) is equivalent to

$$\ell^* \leq \frac{\Phi'_*(t)t^2}{\Phi_*(t)} \leq m^*, \quad \forall t > 0$$

Lemma 2.4. *If Φ is an N -function of the form (1.1) satisfying (ϕ_1) , (ϕ_2) and (ϕ_3) , then*

$$\tilde{\Phi}(\rho t) \leq t^{\frac{m}{m-1}} \tilde{\Phi}(\rho), \quad \text{for all } \rho > 0 \quad \text{and} \quad 0 \leq t < 1.$$

Lemma 2.5. *If Φ is N -function and $(\int_{\Omega} \Phi(|u_n|) dx)$ is a bounded sequence, then (u_n) is a bounded sequence in $L^{\Phi}(\Omega)$. When $\Phi \in (\Delta_2)$, the equivalence is valid.*

Lemma 2.6. *If Φ is an N -function of the form (1.1) satisfying (ϕ_1) , (ϕ_2) and (ϕ_3) with $\ell = 1$, then $\tilde{\Phi} \notin (\Delta_2)$.*

Lemma 2.7. *Let $\Omega \subset \mathbb{R}^N$ be a domain, if $\Phi \in (\Delta_2)$ and (u_n) is a sequence in $L^{\Phi}(\Omega)$ with $u_n \rightarrow u$ in $L^{\Phi}(\Omega)$, there is $H \in L^{\Phi}(\Omega)$ and a subsequence (u_{n_j}) such that*

i) $|u_{n_j}(x)| \leq H(x)$ a.e. in Ω .

ii) $u_{n_j}(x) \rightarrow u(x)$ a.e. in Ω and all $j \in \mathbb{N}$.

For a N -function Φ , the corresponding Orlicz-Sobolev space is defined as the Banach space

$$W^{1,\Phi}(\Omega) = \left\{ u \in L^{\Phi}(\Omega) : \frac{\partial u}{\partial x_i} \in L^{\Phi}(\Omega), i = 1, \dots, N \right\},$$

with the norm

$$(2.2) \quad \|u\|_{1,\Phi} = \|\nabla u\|_{\Phi} + \|u\|_{\Phi}.$$

If $\Phi \in (\Delta_2)$, the space $D^{1,\Phi}(\mathbb{R}^N)$ is defined to be the complement of the space $C_0^{\infty}(\mathbb{R}^N)$ with respect to the standard

$$(2.3) \quad \|u\|_{D^{1,\Phi}(\mathbb{R}^N)} = \|u\|_{\Phi_*} + \|\nabla u\|_{\Phi}.$$

Since the Orlicz-Sobolev inequality

$$(2.4) \quad \|u\|_{\Phi_*} \leq S_N \|\nabla u\|_{\Phi},$$

holds for $u \in D^{1,\Phi}(\mathbb{R}^N)$ with a constant $S_N > 0$, the norm (2.3) is equivalent to the norm

$$(2.5) \quad \|u\|_{D^{1,\Phi}(\mathbb{R}^N)} = \|\nabla u\|_{\Phi},$$

on $D^{1,\Phi}(\mathbb{R}^N)$. In this paper, we will use (2.5) as the norm of $D^{1,\Phi}(\mathbb{R}^N)$. Clearly

$$D^{1,\Phi}(\mathbb{R}^N) \xrightarrow[\text{cont}]{} L^{\Phi_*}(\mathbb{R}^N).$$

The space $L^{\Phi}(\mathbb{R}^N)$ is separable and reflexive when the N -functions Φ and $\tilde{\Phi}$ satisfy the Δ_2 -condition. Knowing that $D^{1,\Phi}(\mathbb{R}^N)$ can be seen as a closed subspace of the space $L^{\Phi_*}(\mathbb{R}^N) \times (L^{\Phi}(\mathbb{R}^N))^N$, then $D^{1,\Phi}(\mathbb{R}^N)$ is reflexive when the N -functions Φ , $\tilde{\Phi}$, Φ_* and $\tilde{\Phi}_*$ satisfy the Δ_2 -condition.

The following lemma is an immediate consequence of the Banach-Alaoglu-Bourbaki theorem [18].

Lemma 2.8. *Assume that Φ is an N -function of the form (1.1) satisfying (ϕ_1) , (ϕ_2) and (ϕ_3) . If $(u_n) \subset D^{1,\Phi}(\mathbb{R}^N)$ is a bounded sequence, then there exists a subsequence of (u_n) , which we will still denote by (u_n) , and $u \in D^{1,\Phi}(\mathbb{R}^N)$ such that*

$$(2.6) \quad u_n \xrightarrow{*} u \quad \text{in } L^{\Phi_*}(\mathbb{R}^N) \quad \text{and} \quad \frac{\partial u_n}{\partial x_i} \xrightarrow{*} \frac{\partial u}{\partial x_i} \quad \text{in } L^{\Phi}(\mathbb{R}^N)$$

or equivalently,

$$\int_{\mathbb{R}^N} u_n v dx \longrightarrow \int_{\mathbb{R}^N} u v dx, \quad \forall v \in E^{\tilde{\Phi}_*}(\mathbb{R}^N)$$

and

$$\int_{\mathbb{R}^N} \frac{\partial u_n}{\partial x_i} w dx \longrightarrow \int_{\mathbb{R}^N} \frac{\partial u}{\partial x_i} w dx, \quad \forall w \in E^{\tilde{\Phi}}(\mathbb{R}^N).$$

From now on, we denote the limit (2.6) by $u_n \xrightarrow{*} u$ in $D^{1,\Phi}(\mathbb{R}^N)$. As an immediate consequence of the last lemma, we have the following corollary.

Corollary 2.9. *If $(u_n) \subset D^{1,\Phi}(\mathbb{R}^N)$ is a bounded sequence with $u_n \longrightarrow u$ in $L_{loc}^{\Phi}(\mathbb{R}^N)$, then $u \in D^{1,\Phi}(\mathbb{R}^N)$.*

The last lemma is crucial when the space $D^{1,\Phi}(\mathbb{R}^N)$ can be nonreflexive. And, this happens, for example, when $\Phi_{\alpha}(t) = |t| \ln(|t|^{\alpha} + 1)$, for $0 < \alpha < \frac{N}{N-1} - 1$, because $\tilde{\Phi}_{\alpha}$ does not verify the Δ_2 -condition. Here we emphasize that the condition (ϕ_3) guarantees that Φ and $\tilde{\Phi}$ verifies the Δ_2 -condition when $\ell > 1$, for more details see Fukagai and Narukawa [12].

3. PRELIMINARY RESULTS

This section focuses on preparing some preliminaries for proving Theorems 1.1, 1.2, 1.3 and 1.4. Since the potential V may vanish at infinity, we cannot study equation (P) on the Sobolev space $D^{1,\Phi}(\mathbb{R}^N)$ by variational methods. As in [21], we work in the space $E = \{u \in D^{1,\Phi}(\mathbb{R}^N) : \int_{\mathbb{R}^N} V(x)\Phi(|u|)dx < +\infty\}$ with norm

$$\|u\|_E = \|u\|_{D^{1,\Phi}(\mathbb{R}^N)} + \|u\|_{L_V^{\Phi}(\mathbb{R}^N)},$$

where $\|u\|_{L_V^{\Phi}(\mathbb{R}^N)}$ is the norm of Banach space $L_V^{\Phi}(\mathbb{R}^N)$.

Lemma 3.1. *$E = \overline{C_0^{\infty}(\mathbb{R}^N)}^{\|\cdot\|_E}$ is a Banach space and E is compactly embedded in $L_{loc}^{\Phi}(\mathbb{R}^N)$*

Lemma 3.2. *Suppose $(u_n) \subset E$ is a bounded sequence in E , then there is $u \in E$ such that $u_n \xrightarrow{*} u$ in $D^{1,\Phi}(\mathbb{R}^N)$ and*

$$\int_{\mathbb{R}^N} \Phi(|\nabla u|) dx \leq \liminf_{n \rightarrow \infty} \int_{\mathbb{R}^N} \Phi(|\nabla u_n|) dx.$$

Proof: See Lemma 3.4 and Lemma 4.7 in [21].

Remark 1. *The inequality (1.2) implies the following inequalities*

$$\xi_{0,A}(t)A(\rho) \leq A(\rho t) \leq \xi_{1,A}(t)A(\rho), \quad \forall \rho, t \geq 0$$

when

$$\xi_{0,A}(t) = \min\{t^{a_1}, t^{a_2}\} \quad \text{and} \quad \xi_{1,A}(t) = \max\{t^{a_1}, t^{a_2}\}, \quad \forall t \geq 0.$$

Besides by Lemma 2.1 and Lemma 2.2, we have

$$\lim_{t \rightarrow 0} \frac{A(t)}{\Phi(t)} = 0 \quad \text{and} \quad \lim_{|t| \rightarrow \infty} \frac{A(t)}{\Phi_*(t)} = 0$$

Remark 2. The inequality (1.2) implies the following inequalities

$$\xi_{0,B}(t)B(\rho) \leq B(\rho t) \leq \xi_{1,B}(t)B(\rho), \quad \forall \rho, t \geq 0$$

when

$$\xi_{0,B}(t) = \min\{t^{b_1}, t^{b_2}\} \quad \text{and} \quad \xi_{1,B}(t) = \max\{t^{b_1}, t^{b_2}\}, \quad \forall t \geq 0.$$

Besides by Lemma 2.1 and Lemma 2.2, we have

$$\lim_{t \rightarrow 0} \frac{B(t)}{\Phi(t)} = 0 \quad \text{and} \quad \lim_{|t| \rightarrow \infty} \frac{B(t)}{\Phi_*(t)} = 0$$

Proposition 3.3. If $(V, K) \in \mathcal{K}_1$, then the space E is continuous embedded in $L_{K^s}^A(\mathbb{R}^N)$. On the other hand, if $(V, K) \in \mathcal{K}_2$, then the space E is continuous embedded in $L_{K^s}^B(\mathbb{R}^N)$.

Proof: The case $(V, K) \in \mathcal{K}_1$ is obvious.

Now, let us assume that $(V, K) \in \mathcal{K}_2$. As E is continuously embedded in $L^{\Phi^*}(\mathbb{R}^N)$, there exists $C_1 > 0$ such that

$$(3.1) \quad \|u\|_{\Phi_*} \leq C_1 \|u\|_E, \quad \forall u \in E.$$

By the condition (K_3) , there is $r > 0$ satisfying

$$(3.2) \quad K(x)B(t) \leq V(x)\Phi(t) + \Phi_*(t), \quad \forall t > 0 \quad \text{and} \quad |x| \geq r.$$

On the other hand, by the Remark 1, there is a constant $C_2 > 0$ such that

$$B(t) \leq C_2\Phi(t) + C_2\Phi_*(t), \quad \forall t > 0.$$

Hence, for each $x \in B_r(0)$,

$$(3.3) \quad K(x)B(t) \leq C_2 \left\| \frac{K}{V} \right\|_{L^\infty(B_r(0))} V(x)\Phi(t) + C_2 \|K\|_\infty \Phi_*(t), \quad \forall t > 0.$$

Combining (3.2) and (3.3),

$$(3.4) \quad K(x)B(t) \leq C_3 V(x)\Phi(t) + C_3 \Phi_*(t), \quad \forall t > 0 \quad \text{and} \quad x \in \mathbb{R}^N$$

with $C_3 = \max\{1, C_2 \|K\|_\infty, C_2 \left\| \frac{K}{V} \right\|_{L^\infty(B_r(0))}\}$. By the inequalities (3.1) and (3.1), we get

$$\int_{\mathbb{R}^N} K(x)B \left(\frac{|u|}{C_3 \|u\|_E + C_1 \|u\|_E} \right) dx \leq C_3 \int_{\mathbb{R}^N} V(x)\Phi \left(\frac{|u|}{\|u\|_{V,\Phi}} \right) dx + C_3 \int_{\mathbb{R}^N} \Phi_* \left(\frac{|u|}{\|u\|_{\Phi_*}} \right) dx \leq C_4$$

where C_4 is a positive constant that does not depend on u . So we can conclude that $E \subset L_{K^s}^B(\mathbb{R}^N)$. Furthermore, there is a constant $C_5 > 0$ that does not depend on u , so $\|u\|_{K,B} \leq C_5 \|u\|_E$. Concluding that E is continuous embedded in $L_{K^s}^B(\mathbb{R}^N)$. \square

Noting that Choquard equation is no longer defined pointwise, to study equations (P) by variational methods, we need the following well-known Hardy-Littlewood-Sobolev inequality afterwards, whose proof can be found in [22].

Proposition 3.4. [Hardy-Littlewood-Sobolev inequality] Set $t, r > 1$ and $\lambda \in (0, N)$ with $1/t + \lambda/N + 1/r = 2$, $G_1 \in L^r(\mathbb{R}^N)$ and $G_2 \in L^t(\mathbb{R}^N)$. Then there exists a sharp constant $C(t, N, \lambda, r)$, independent of G_1, G_2 such that

$$(3.5) \quad \left| \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{G_1(x)G_2(y)}{|x-y|^\lambda} dx dy \right| \leq C(t, N, \lambda, r) \|G_1\|_{L^t(\mathbb{R}^N)} \|G_2\|_{L^t(\mathbb{R}^N)}.$$

From (f_1) and $f \in C(\mathbb{R}, \mathbb{R})$, for any $\varepsilon > 0$ there exists $\delta_0 > 0$, $\delta_1 > 0$ and $C_\varepsilon > 0$ such that

$$(3.6) \quad |f(t)|^s \leq \frac{\varepsilon}{s}(a(t)t^{2-s} + \phi_*(t)t^{2-s}) + C_\varepsilon \phi_*(t)t^{2-s} \chi_{[\delta_0, \delta_1]}(t), \quad \forall t > 0.$$

Lemma 3.5. *Suppose that $(V, K) \in \mathcal{K}_1$ and (f_1) holds. For each $u \in E$, there is a constant $C_1 > 0$ that does not depend on u , such that*

$$\left| \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))F(u(y))}{|x-y|^\lambda} dx dy \right| \leq C_1 \left[\left(\int_{\mathbb{R}^N} K(x)^s A(|u|) dx \right)^{\frac{2}{s}} + \left(\int_{\mathbb{R}^N} \Phi_*(|u|) dx \right)^{\frac{2}{s}} \right].$$

Furthermore, for each $u \in E$, there is a constant $C_2 > 0$, which does not depend on u , such that

$$(3.7) \quad \left| \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))f(u(y))v(y)}{|x-y|^\lambda} dx dy \right| \leq C \|v\|_E, \quad \forall v \in E.$$

Proof: By (3.6), there is a constant $C > 0$ such that

$$(3.8) \quad |f(t)|^s \leq C(a(t)t^{2-s} + \phi_*(t)t^{2-s}), \quad \forall t \in \mathbb{R}.$$

For each $t \geq 0$, we have

$$|F(t)| \leq \int_0^t |f(\tau)| d\tau \leq \left[\int_0^t |f(\tau)|^s d\tau \right]^{\frac{1}{s}} \left[\int_0^t d\tau \right]^{\frac{s-1}{s}} \leq t^{\frac{s-1}{s}} \left[\int_0^t |f(\tau)|^s d\tau \right]^{\frac{1}{s}}.$$

Thus,

$$\begin{aligned} |F(t)|^s &\leq t^{s-1} C \left(\int_0^t (a(\tau)\tau^{2-s} + \phi_*(\tau)\tau^{2-s}) d\tau \right) \\ &\leq C t^s (a(t)t^{2-s} + \phi_*(t)t^{2-s}) \quad (\tau h(\tau) \text{ and } \tau w(\tau) \text{ are increasing in } (0, \infty)) \\ &\leq C (A(|t|) + \Phi_*(|t|)), \quad \forall t \geq 0 \end{aligned}$$

Similarly,

$$|F(t)|^s \leq C (A(|t|) + \Phi_*(|t|)), \quad \forall t \leq 0$$

Therefore,

$$|F(t)|^s \leq C (A(|t|) + \Phi_*(|t|)), \quad \forall t \in \mathbb{R}$$

that is,

$$(3.9) \quad \int_{\mathbb{R}^N} K(x)^s |F(u)|^s dx \leq C \int_{\mathbb{R}^N} K(x)^s A(|u|) dx + C \int_{\mathbb{R}^N} \Phi_*(|u|) dx < \infty, \quad \forall u \in E.$$

By the Hardy-Littlewood-Sobolev inequality, it follows that

$$(3.10) \quad \begin{aligned} \left| \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))F(u(y))}{|x-y|^\lambda} dx dy \right| &\leq \left| \int_{\mathbb{R}^N} K(x)^s |F(u)|^s dx \right|^{\frac{2}{s}} \\ &\leq C \left(\int_{\mathbb{R}^N} K(x)^s A(|u|) dx + \int_{\mathbb{R}^N} \Phi_*(|u|) dx \right)^{\frac{2}{s}} \\ &\leq C \left[\left(\int_{\mathbb{R}^N} K(x)^s A(|u|) dx \right)^{\frac{2}{s}} + \left(\int_{\mathbb{R}^N} \Phi_*(|u|) dx \right)^{\frac{2}{s}} \right], \end{aligned}$$

for every $u \in E$ and $C > 0$ is a positive constant that does not depend on u .

Now, consider $u, v \in E$, from (3.8) we have

$$\tau_{uv} = \int_{\mathbb{R}^N} K(x)^s |f(u)|^s |v|^s dx \leq C \int_{\mathbb{R}^N} K(x)^s a(|u|) |u|^{2-s} |v|^s dx + C \int_{\mathbb{R}^N} \phi_*(|u|) |u|^{2-s} |v|^s dx.$$

Define the functions $H : \mathbb{R} \rightarrow [0, \infty)$ and $P : \mathbb{R} \rightarrow [0, \infty)$ given by $H(t) = A(|t|^{1/s})$ and $P(t) = \Phi_*(|t|^{1/s})$. Through the assumptions imposed under A and Φ_* it is possible to show that H and P are N -functions, in addition, the functions $h : (0, \infty) \rightarrow (0, \infty)$ defined by $h(t)t = \frac{1}{s} a(t^{1/s}) t^{(2/s)-1}$ and $p(t)t = \frac{1}{s} \phi_*(t^{1/s}) t^{(2/s)-1}$ are increasing and satisfy

$$(3.11) \quad H(w) = \int_0^{|w|} th(t)dt, \quad \text{and} \quad P(w) = \int_0^{|w|} tp(t)dt.$$

Since E is continuous embedded in $L_{K^s}^A(\mathbb{R}^N)$ and $L^{\Phi_*}(\mathbb{R}^N)$, we have

$$\begin{aligned} \int_{\mathbb{R}^N} K(x)^s \tilde{H}(a(|u|) |u|^{2-s}) dx &= \int_{\mathbb{R}^N} K(x)^s \tilde{H}(|u|^s h(|u|^s)) dx \\ &\leq C_1 \int_{\mathbb{R}^N} K(x)^s H(|u|^s) dx = C_1 \int_{\mathbb{R}^N} K(x)^s A(|u|) dx < +\infty, \end{aligned}$$

$$\int_{\mathbb{R}^N} \tilde{P}(\phi_*(|u|) |u|^{2-s}) dx = \int_{\mathbb{R}^N} \tilde{P}(|u|^s p(|u|^s)) dx \leq C_1 \int_{\mathbb{R}^N} P(|u|^s) dx = C_1 \int_{\mathbb{R}^N} \Phi_*(|u|) dx < +\infty,$$

$$\int_{\mathbb{R}^N} K(x)^s H\left(\frac{|v|^s}{\|v\|_{L_{K^s}^A(\mathbb{R}^N)}^s}\right) dx = \int_{\mathbb{R}^N} K(x)^s A\left(\frac{|v|}{\|v\|_{L_{K^s}^A(\mathbb{R}^N)}}\right) dx = 1,$$

$$\int_{\mathbb{R}^N} P\left(\frac{|v|^s}{\|v\|_{L^{\Phi_*}(\mathbb{R}^N)}^s}\right) dx = \int_{\mathbb{R}^N} \Phi_*\left(\frac{|v|}{\|v\|_{L^{\Phi_*}(\mathbb{R}^N)}}\right) dx = 1.$$

With this, we conclude that $a(|u|) |u|^{2-s} \in L_{K^s}^{\tilde{H}}(\mathbb{R}^N)$, $\phi_*(|u|) |u|^{2-s} \in L^{\tilde{P}}(\mathbb{R}^N)$, $|v|^s \in L_{K^s}^H(\mathbb{R}^N)$ and $|v|^s \in L^P(\mathbb{R}^N)$. Furthermore,

$$\| |v|^s \|_{L_{K^s}^H(\mathbb{R}^N)} = \|v\|_{L_{K^s}^A(\mathbb{R}^N)}^s \leq C_2 \|v\|_E^s$$

and

$$\| |v|^s \|_{L^P(\mathbb{R}^N)} = \|v\|_{L^{\Phi_*}(\mathbb{R}^N)}^s \leq C_2 \|v\|_E^s$$

where $C_i > 0$, $i = 1, 2$, are positives constants that does not depend on v . By Proposition 3.3, it follows that

$$(3.12) \quad \begin{aligned} \tau_{uv} &\leq C_3 \|a(|u|) |u|^{2-s}\|_{L_{K^s}^{\tilde{H}}(\mathbb{R}^N)} \| |v|^s \|_{L_{K^s}^H(\mathbb{R}^N)} + C_3 \|\phi_*(|u|) |u|^{2-s}\|_{L^{\tilde{P}}(\mathbb{R}^N)} \| |v|^s \|_{L^P(\mathbb{R}^N)} \\ &\leq C_4 (\|a(|u|) |u|^{2-s}\|_{L_{K^s}^{\tilde{H}}(\mathbb{R}^N)} \|v\|_E^s + \|\phi_*(|u|) |u|^{2-s}\|_{L^{\tilde{P}}(\mathbb{R}^N)} \|v\|_E^s), \end{aligned}$$

where $C_i > 0$, $i = 3, 4$, are positives constants that does not depend on u and v . From (3.12), (3.9) together with the Proposition 3.4, it follows that

$$\begin{aligned} \left| \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))f(u(y))v(y)}{|x-y|^\lambda} dx dy \right| &\leq \left| \int_{\mathbb{R}^N} W(x) |F(u)|^s dx \right|^{\frac{1}{s}} \left| \int_{\mathbb{R}^N} W(x) |f(u)|^s |v|^s dx \right|^{\frac{1}{s}} \\ &\leq C_5 \left(\int_{\mathbb{R}^N} K(x)^s A(u) dx + \int_{\mathbb{R}^N} \Phi_*(u) dx \right)^{\frac{1}{s}} (\|a(|u|) |u|^{2-s}\|_{L_{K^s}^{\tilde{H}}(\mathbb{R}^N)} + \|\phi_*(|u|) |u|^{2-s}\|_{L^{\tilde{P}}(\mathbb{R}^N)})^{\frac{1}{s}} \|v\|_E. \end{aligned}$$

□

Lemma 3.6. *Assume that $(V, K) \in \mathcal{K}_1$ and (f_1) holds. Let (u_n) be a bounded sequence in E , and consider $u \in E$ such that $u_n \xrightarrow{*} u$ in E . We will show the following limits*

$$(3.13) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} K(x)^s |F(u_n) - F(u)|^s dx = 0,$$

$$(3.14) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} K(x)^s |f(u_n)u_n - f(u)u|^s dx = 0$$

and

$$(3.15) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} K(x)^s |f(u_n)\varphi - f(u)\varphi|^s dx = 0.$$

Proof. By (4.10), for any $\varepsilon > 0$ there exists $\delta_0 > 0$, $\delta_1 > 0$ and $C_\varepsilon > 0$ such that

$$(3.16) \quad |F(t)|^s \leq \varepsilon \left(\frac{a_2}{s} A(t) + \frac{m^*}{s} \Phi_*(t) \right) + \frac{C_\varepsilon m^*}{s} \Phi_*(t) \chi_{[\delta_0, \delta_1]}(t)$$

From (3.16), Proposition 3.3 and the Sobolev inequality, it follows that the sequence $(K(\cdot)F(u_n))$ is bounded in $L^s(\mathbb{R}^N)$. It is clear that $K(x)F(u_n(x)) \rightarrow K(x)F(u(x))$ a.e. in \mathbb{R}^N in the sense of subsequence. Then, by the Brézis-Lieb lemma [28, Lemma 1.32] we obtain

$$(3.17) \quad \int_{\mathbb{R}^N} K(x)^s |F(u_n) - F(u)|^s dx = \int_{\mathbb{R}^N} K(x)^s |F(u_n)|^s dx - \int_{\mathbb{R}^N} K(x)^s |F(u)|^s dx + o_n(1).$$

In view of this fact, to verify (3.13), we only need to prove that the right side of (3.17) belongs to $o_n(1)$. Note that $F_n = \{x \in \mathbb{R}^N : |v_n(x)| \geq \delta_0\}$ is such that

$$\Phi_*(\delta_0)|F_n| \leq \int_{F_n} \Phi_*(|v_n(x)|) dx \leq \int_{\mathbb{R}^N} \Phi_*(|v_n(x)|) dx \leq C_1,$$

for some constant $C_1 > 0$ that does not depend on n . Thus, $\sup_{n \in \mathbb{N}} |F_n| < +\infty$. From (K_1) , we have

$$\lim_{r \rightarrow +\infty} \int_{F_n \cap B_r^c(0)} K(x)^s dx = 0, \quad \text{uniformly in } n \in \mathbb{N},$$

thus, there is $r_0 > 0$, so that

$$\int_{F_n \cap B_{r_0}^c(0)} K(x)^s dx < \frac{\varepsilon}{\Phi_*(\delta_1)C_\varepsilon}, \quad \forall n \in \mathbb{N}.$$

Moreover, as (v_n) is bounded in E , there is a constant $M_1 > 0$ satisfying

$$\int_{\mathbb{R}^N} K(x)^s A(|v_n|) dx \leq M_1 \quad \text{and} \quad \int_{\mathbb{R}^N} \Phi_*(|v_n|) dx \leq M_1, \quad \forall n \in \mathbb{N}.$$

By (4.11), it follows that

$$\begin{aligned} \int_{B_{r_0}^c(0)} K(x)^s |F(u_n)|^s dx &\leq \varepsilon C_1 \left(\int_{B_{r_0}^c(0)} K(x)^s A(|u_n|) dx + \int_{B_{r_0}^c(0)} \Phi_*(|u_n|) dx \right) \\ &\quad + C_\varepsilon \Phi_*(\delta_1) \int_{F_n \cap B_{r_0}^c(0)} K(x)^s dx \\ &\leq \varepsilon (C_1 M_1 + 1), \end{aligned}$$

for all $n \in \mathbb{N}$. Therefore

$$(3.18) \quad \limsup_{n \rightarrow +\infty} \int_{B_{r_0}^c(0)} K(x)^s |F(u_n)|^s dx \leq \varepsilon (C_1 M_1 + 1).$$

On the other hand, using the compactness lemma of Strauss [17, Theorem A.I, p. 338], it follows that

$$(3.19) \quad \lim_{n \rightarrow +\infty} \int_{B_{r_0}(0)} K(x)^s |F(u_n)|^s dx = \int_{B_{r_0}(0)} K(x)^s |F(u)|^s dx.$$

In light of this, we can conclude that

$$\lim_{n \rightarrow +\infty} \int_{\mathbb{R}^N} K(x)^s |F(u_n)|^s dx = \int_{\mathbb{R}^N} K(x)^s |F(u)|^s dx.$$

Through this limit together with (3.17), we will get (3.13). Similarly, the limit (3.14) is shown. Related the limit (3.15), it follows directly from the condition (f_1) together with a version of the compactness lemma of Strauss for non-autonomous problem. \square

The following result is an immediate consequence of Hardy-Littlewood-Sobolev and Lemma 3.6.

Lemma 3.7. *Assume that $(V, K) \in \mathcal{K}_1$ and (f_1) holds. Let (u_n) be a sequence bounded in E and $u \in E$ such that $u_n \xrightarrow{*} u$ in $D^{1,\Phi}(\mathbb{R}^N)$. Then*

$$(3.20) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))F(u_n(y))}{|x-y|^\lambda} dx dy = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))F(u(y))}{|x-y|^\lambda} dx dy,$$

$$(3.21) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))f(u_n(y))u_n(y)}{|x-y|^\lambda} dx dy = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))f(u(y))u(y)}{|x-y|^\lambda} dx dy$$

and

$$(3.22) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))f(u_n(y))\varphi(y)}{|x-y|^\lambda} dx dy = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))f(u(y))\varphi(y)}{|x-y|^\lambda} dx dy$$

for all $\varphi \in C_0^\infty(\mathbb{R}^N)$.

Now, suppose that the function f satisfies the condition (f_5) . Similar to (4.10), given $\varepsilon > 0$ there exists $\delta_0 > 0$, $\delta_1 > 0$ and $C_\varepsilon > 0$ such that

$$(3.23) \quad |f(t)|^s \leq \frac{\varepsilon}{s}(b(t)t^{2-s} + \phi_*(t)t^{2-s}) + C_\varepsilon \phi_*(t)t^{2-s} \chi_{[\delta_0, \delta_1]}(t), \quad \forall t > 0.$$

Repeating the same arguments used in the proof of Lemma 3.5, we can state the following result.

Lemma 3.8. *Suppose that $(V, K) \in \mathcal{K}_2$ and (f_5) holds. For each $u \in E$, there is a constant $C_1 > 0$ that does not depend on u , such that*

$$\left| \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))F(u(y))}{|x-y|^\lambda} dx dy \right| \leq C_1 \left[\left(\int_{\mathbb{R}^N} K(x)^s B(|u|) dx \right)^{\frac{2}{s}} + \left(\int_{\mathbb{R}^N} \Phi_*(|u|) dx \right)^{\frac{2}{s}} \right].$$

Furthermore, for each $u \in E$ there is a constant $C_2 > 0$, which does not depend on u , such that

$$(3.24) \quad \left| \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))f(u(y))v(y)}{|x-y|^\lambda} dx dy \right| \leq C \|v\|_E, \quad \forall v \in E.$$

Lemma 3.9. *Suppose that $(V, K) \in \mathcal{K}_2$ and (f_5) holds. Let (u_n) be a bounded sequence in E , and consider $u \in E$ such that $u_n \xrightarrow{*} u$ in E . We will show the following limits*

$$(3.25) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} K(x)^s |F(u_n) - F(u)|^s dx = 0,$$

$$(3.26) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} K(x)^s |f(u_n)u_n - f(u)u|^s dx = 0$$

and

$$(3.27) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} K(x)^s |f(u_n)\varphi - f(u)\varphi|^s dx = 0.$$

Proof. Due to similarity, it suffices to verify (3.25). By (3.23), for any $\varepsilon > 0$ there exists $\delta_0 > 0$, $\delta_1 > 0$ and $C_\varepsilon > 0$ such that

$$(3.28) \quad |F(t)|^s \leq \varepsilon \left(\frac{b_2}{s} B(t) + \frac{m^*}{s} \Phi_*(t) \right) + \frac{C_\varepsilon m^*}{s} \Phi_*(t) \chi_{[\delta_0, \delta_1]}(t)$$

By the condition (K_3) , there is $r_0 > 0$ sufficiently large satisfying

$$K(x)B(t) \leq \varepsilon (V(x)\Phi(t) + \Phi_*(t)), \quad \forall t > 0 \text{ and } |x| \geq r_0.$$

From the above inequalities, we have

$$(3.29) \quad K(x)F(t) \leq \varepsilon C_1 V(x)\Phi(t) + \varepsilon C_2 \Phi_*(t) + C_\varepsilon K(x)\Phi_*(\delta_1)\chi_{[\delta_0, \delta_1]}(t), \quad \forall t > 0 \text{ and } |x| \geq r_0.$$

From (3.28), Lemma 3.3 and the Sobolev inequality, it follows that the sequence $(K(\cdot)F(u_n))$ is bounded in $L^s(\mathbb{R}^N)$. It is clear that $K(x)F(u_n(x)) \rightarrow K(x)F(u(x))$ a.e. in \mathbb{R}^N in the sense of subsequence. Then, by the Brézis-Lieb lemma [28, Lemma 1.32] we obtain

$$(3.30) \quad \int_{\mathbb{R}^N} K(x)^s |F(u_n) - F(u)|^s dx = \int_{\mathbb{R}^N} K(x)^s |F(u_n)|^s dx - \int_{\mathbb{R}^N} K(x)^s |F(u)|^s dx + o_n(1)$$

In view of this fact, to verify (3.25), we only need to prove that the right side of (3.30) belongs to $o_n(1)$. Repeating the same arguments used in the proof of Lemma 3.6, it follows that

$$(3.31) \quad \limsup_{n \rightarrow +\infty} \int_{B_{r_0}^c(0)} K(x)^s |F(u_n)|^s dx \leq \varepsilon (C_1 M_1 + 1).$$

On the other hand, using (f_5) and the compactness lemma of Strauss [17, Theorem A.I, p. 338], it is guaranteed that

$$(3.32) \quad \lim_{n \rightarrow +\infty} \int_{B_{r_0}(0)} K(x)^s |F(u_n)|^s dx = \int_{B_{r_0}(0)} K(x)^s |F(u)|^s dx.$$

In light of this, we can conclude that

$$\lim_{n \rightarrow +\infty} \int_{\mathbb{R}^N} K(x)^s |F(u_n)|^s dx = \int_{\mathbb{R}^N} K(x)^s |F(u)|^s dx.$$

Through this limit together with (3.30), we will get (3.25). Similarly, we show the limit (3.26). Related the limit (3.27), it follows directly from the condition (f_5) together with a version of the compactness lemma of Strauss for non-autonomous problem. \square

Corollary 3.10. *Assume that $(V, K) \in \mathcal{K}_2$ and (f_5) holds. Let (u_n) be a sequence bounded in E and $u \in E$ such that $u_n \xrightarrow{*} u$ in $D^{1, \Phi}(\mathbb{R}^N)$. Then*

$$(3.33) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))F(u_n(y))}{|x-y|^\lambda} dx dy = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))F(u(y))}{|x-y|^\lambda} dx dy,$$

$$(3.34) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))f(u_n(y))u_n(y)}{|x-y|^\lambda} dx dy = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))f(u(y))u(y)}{|x-y|^\lambda} dx dy$$

and

$$(3.35) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))f(u_n(y))\varphi(y)}{|x-y|^\lambda} dx dy = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))f(u(y))\varphi(y)}{|x-y|^\lambda} dx dy$$

for all $\varphi \in C_0^\infty(\mathbb{R}^N)$.

By the inequalities (3.6) and (3.23), together with all the results presented above, it is verified that the function

$$\Psi(u) = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))F(u_n(y))}{|x-y|^\lambda} dx dy, \quad u \in E$$

is well defined, is continuously differentiable and the Gateaux derivative $\Psi' : E \rightarrow E^*$ is given by

$$\Psi'(u)v = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))f(u_n(y))v(y)}{|x-y|^\lambda} dx dy, \quad \forall u, v \in E.$$

This fact is proved similarly to Lemma 3.2, found in [7].

The necessary properties involving the functional $Q : E \rightarrow \mathbb{R}$ given by

$$Q(u) = \int_{\mathbb{R}^N} \Phi(|\nabla u|) dx + \int_{\mathbb{R}^N} V(x)\Phi(|u|) dx.$$

can be found in [21]. In summary, the energy function associated with (P) given by

$$J(u) = \int_{\mathbb{R}^N} \Phi(|\nabla u|) dx + \int_{\mathbb{R}^N} V(x)\Phi(|u|) dx - \frac{1}{2} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))F(u_n(y))}{|x-y|^\lambda} dx dy, \quad u \in E$$

is a continuous and Gateaux-differentiable functional such that $J' : E \rightarrow E^*$ given by

$$J'(u)v = \int_{\mathbb{R}^N} \phi(|\nabla u|)\nabla u \nabla v dx + \int_{\mathbb{R}^N} V(x)\phi(|u|)uv dx - \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))f(u_n(y))v(y)}{|x-y|^\lambda} dx dy$$

is continuous from the norm topology of E to the weak*-topology of E^* .

The convexity of the functional Q together with the Gateaux-differentiability of the functional J allows us to present a definition of a critical point for J . In this sense, we will say that $u \in E$ is a critical point for the functional J if

$$(3.36) \quad Q(v) - Q(u) \geq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))f(u_n(y))(v(y) - u(y))}{|x-y|^\lambda} dx dy, \quad \forall v \in E.$$

Our next lemma establishes that a critical point u in the sense (3.36) is a weak solution for (P).

Proposition 3.11. *If $u \in E$ is a critical point of J in E , then u is a weak solution to (P).*

Proof: See Lemma 4.1 in [6].

Lemma 3.12. *Suppose that $(V, K) \in \mathcal{K}_1$ and (f_1) hold. Then there are $\rho, \eta > 0$ such that $J(u) \geq \eta$ for all $u \in E \cap \partial B_\rho(0)$.*

Proof. By Lemma 3.5, there exists a positive constant $C > 0$ satisfying

$$\left| \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))F(u(y))}{|x-y|^\lambda} dx dy \right| \leq C \left[\left(\int_{\mathbb{R}^N} K(x)^s A(|u|) dx \right)^{\frac{2}{s}} + \left(\int_{\mathbb{R}^N} \Phi_*(|u|) dx \right)^{\frac{2}{s}} \right]$$

for all $u \in E$. Hence, by defining the functional J together with Proposition 3.3, we get

$$\begin{aligned}
J(u) &\geq \int_{\mathbb{R}^N} \Phi(|\nabla u|) dx + \int_{\mathbb{R}^N} V(x)\Phi(|u|) dx - C \left(\int_{\mathbb{R}^N} K(x)^s A(|u|) dx \right)^{\frac{2}{s}} - C \left(\int_{\mathbb{R}^N} \Phi_*(|u|) dx \right)^{\frac{2}{s}} \\
&\geq \xi_0(\|\nabla u\|_\Phi) + \xi_0(\|u\|_{V,\Phi}) - C \left(\int_{\mathbb{R}^N} K(x)^s A(|u|) dx \right)^{\frac{2}{s}} - C \left(\int_{\mathbb{R}^N} \Phi_*(|u|) dx \right)^{\frac{2}{s}} \\
&\geq \xi_0(\|\nabla u\|_\Phi) + \xi_0(\|u\|_{V,\Phi}) - C(\xi_5(\|u\|_{L_{K^s}^A(\mathbb{R}^N)}))^{\frac{2}{s}} - C(\xi_3(\|u\|_{L^{\Phi_*}(\mathbb{R}^N)}))^{\frac{2}{s}} \\
&\geq \|\nabla u\|_\Phi^m + \|u\|_{V,\Phi}^m - C(\|u\|_{L_{K^s}^A(\mathbb{R}^N)})^{\frac{2a_1}{s}} - C(\|u\|_{L^{\Phi_*}(\mathbb{R}^N)})^{\frac{2\ell^*}{s}},
\end{aligned}$$

for $u \in E$ with $\|u\|_E \leq 1$ where $\xi_0(t) = \min_{t>0}\{t^\ell, t^m\}$, $\xi_3(t) = \max_{t>0}\{t^{\ell^*}, t^{m^*}\}$ and $\xi_5(t) = \max_{t>0}\{t^{a_1}, t^{a_2}\}$. By using the classical inequality

$$(x + y)^\sigma \leq 2^{\sigma-1}(x^\sigma + y^\sigma), \quad x, y > 0, \text{ e } \sigma > 1,$$

and the continuous embedded of the space E in the spaces $L_{K^s}^A(\mathbb{R}^N)$ and $L^{\Phi_*}(\mathbb{R}^N)$, we get for $u \in E$ with $\|u\|_E \leq 1$ that

$$\begin{aligned}
J(u) &\geq C(\|\nabla u\|_\Phi + \|u\|_{V,\Phi})^m - C(\|u\|_{L_{K^s}^A(\mathbb{R}^N)})^{\frac{2a_1}{s}} - C(\|u\|_{L^{\Phi_*}(\mathbb{R}^N)})^{\frac{2\ell^*}{s}} \\
&= C\|u\|_E^m - C\left(\|u\|_E^{\frac{2a_1}{s}} + \|u\|_E^{\frac{2\ell^*}{s}}\right),
\end{aligned}$$

for some constant $C > 0$. As $\frac{2}{s} > 1$, then $m < \frac{2a_1}{s}$ and $m < \frac{2\ell^*}{s}$. Hence, setting $\rho = \|u\|$ small enough,

$$J(u) \geq C\|u\|_E^m - C\left(\|u\|_E^{\frac{2a_1}{s}} + \|u\|_E^{\frac{2\ell^*}{s}}\right) := \eta > 0.$$

Which completes the proof. \square

Using the Lemma 3.8, we can repeat the same arguments used in the proof of Lemma 3.12 to prove the following lemma:

Lemma 3.13. *Suppose that $(V, K) \in \mathcal{K}_2$ and (f_5) hold. Then there are $\rho, \eta > 0$ such that $J(u) \geq \eta$ for all $u \in E \cap \partial B_\rho(0)$.*

By a standard argument, the following Lemma follows from the condition (f_4) .

Lemma 3.14. *There is $e \in E$ with $\|u\|_E > 1$ and $J(e) < 0$.*

The previous lemmas establish the mountain pass geometry for the functional J in both cases. In what follows, let us denote by $c > 0$ the mountain pass level associated with J , that is,

$$c = \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} J(\gamma(t))$$

where

$$\Gamma = \{\gamma \in C([0,1], X) : \gamma(0) = 0 \text{ and } \gamma(1) = e\}.$$

Associated with c , we have a Cerami sequence $(u_n) \subset E$, that is,

$$(3.37) \quad J(u_n) \rightarrow c \quad \text{and} \quad (1 + \|u_n\|)\|J'(u_n)\|_* \rightarrow 0.$$

The above sequence is obtained from the Ghoussoub-Preiss theorem, see [8, Theorem 5.46].

Now, we are able to prove that the Cerami sequence given in (3.37) is bounded in E .

Lemma 3.15. *Let (u_n) the Cerami sequence given in (3.37). There is a constant $M > 0$ such that $J(tu_n) \leq M$ for every $t \in [0, 1]$ and $n \in \mathbb{N}$.*

Proof: Let $t_n \in [0, 1]$ be such that $J(t_n u_n) = \max_{t \in [0, 1]} J(tu_n)$. If $t_n = 0$ and $t_n = 1$, we are done.

Thereby, we can assume $t_n \in (0, 1)$, and so $J'(t_n u_n)u_n = 0$. From this,

$$\begin{aligned} mJ(t_n u_n) &= mJ(t_n u_n) - J'(t_n u_n)(t_n u_n) \\ &= \int_{\mathbb{R}^N} (m\Phi(|\nabla(t_n u_n)|) - \phi(|\nabla(t_n u_n)|)|\nabla(t_n u_n)|^2) dx \\ &\quad + \int_{\mathbb{R}^N} V(x)(m\Phi(|t_n u_n|) - \phi(|t_n u_n|)|t_n u_n|^2) dx \\ &\quad + \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)[F(t_n u_n(x))f(t_n u_n(y))t_n u_n(y) - \frac{m}{2}F(t_n u_n(x))F(t_n u_n(y))]}{|x-y|^\lambda} dx dy. \end{aligned}$$

The conditions (f_4) and (f_5) guarantee that the function $f(t)t - \frac{m}{2}F(t)$ is increasing for $t > 0$ and $F(t) \geq 0$ for $t \in \mathbb{R}$, respectively. The condition (ϕ_4) ensures that the function $t \mapsto m\Phi(t) - \phi(t)t^2$ is increasing for $t > 0$. Thus,

$$mJ(t_n u_n) \leq mJ(u_n) - J'(u_n)u_n = mJ(u_n) - o_n(1).$$

Since $(J(u_n))$ is bounded, there is $M > 0$ such that

$$J(tu_n) \leq M, \quad \forall t \in [0, 1] \quad \text{and} \quad n \in \mathbb{N}.$$

□

Proposition 3.16. *The Cerami sequence (u_n) given in (3.37) is bounded.*

Proof: Suppose by contradiction that $\|u_n\|_E \rightarrow \infty$, then we have the following cases:

i) $\|\nabla u_n\|_\Phi \rightarrow +\infty$ and $(\|u_n\|_{V,\Phi})$ is bounded

ii) $\|u_n\|_{V,\Phi} \rightarrow \infty$ and $(\|\nabla u_n\|_\Phi)$ is bounded

iii) $\|\nabla u_n\|_\Phi \rightarrow +\infty$ and $\|u_n\|_{V,\Phi} \rightarrow +\infty$.

In the case iii), consider

$$w_n = \frac{u_n}{\|u_n\|_E}, \quad \forall n \in \mathbb{N}.$$

Since $\|w_n\|_E = 1$, by Lemma 3.2, there exists $w \in E$ such that $w_n \xrightarrow{*} w$ in $D^{1,\Phi}(\mathbb{R}^N)$. There are two possible cases: $w = 0$ or $w \neq 0$.

Case: $w = 0$

Note that for every constant $\sigma > 1$ there is $n_0 \in \mathbb{N}$ such that $\frac{\sigma}{\|u_n\|_E} \in [0, 1]$, for $n \geq n_0$.

Given this, we get

$$\begin{aligned} J(t_n u_n) &\geq J\left(\frac{\sigma}{\|\nabla u_n\|_\Phi} u_n\right) \\ &= J(\sigma w_n) \\ &= \int_{\mathbb{R}^N} \Phi(\sigma|\nabla w_n|) dx + \int_{\mathbb{R}^N} V(x)\Phi(\sigma|w_n|) dx - \frac{1}{2} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(\sigma w_n(x))F(\sigma w_n(y))}{|x-y|^\lambda} dx dy \\ &\geq \sigma Q(w_n) - \frac{1}{2} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(\sigma w_n(x))F(\sigma w_n(y))}{|x-y|^\lambda} dx dy \end{aligned}$$

By definition of the sequence (w_n) , we have $\|\nabla w_n\|_\Phi \leq 1$ and $\|w_n\|_{V,\Phi} \leq 1$, for all $n \in \mathbb{N}$. Then,

$$\int_{\mathbb{R}^N} \Phi(|\nabla w_n|) dx \geq \|\nabla w_n\|_\Phi^m \quad \text{and} \quad \int_{\mathbb{R}^N} V(x)\Phi(|w_n|) dx \geq \|w_n\|_{V,\Phi}^m.$$

So there is $C > 0$ such that

$$Q(w_n) \geq \|\nabla w_n\|_{\Phi}^m + \|w_n\|_{V, \Phi}^m \geq C(\|\nabla w_n\|_{\Phi} + \|w_n\|_{V, \Phi})^m, \quad \forall n \in \mathbb{N}.$$

Thus

$$\begin{aligned} J(t_n u_n) &\geq \sigma C (\|w_n\|_E)^m - \frac{1}{2} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(\sigma w_n(x))F(\sigma w_n(y))}{|x-y|^\lambda} dx dy \\ &= \sigma C - \frac{1}{2} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(\sigma w_n(x))F(\sigma w_n(y))}{|x-y|^\lambda} dx dy \end{aligned}$$

If $w = 0$, segue de (3.20) or (3.33) that

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(\sigma w_n(x))F(\sigma w_n(y))}{|x-y|^\lambda} dx dy = 0,$$

therefore,

$$\liminf_{n \rightarrow \infty} J(t_n u_n) \geq C\sigma, \quad \forall \sigma \geq 1.$$

which constitutes a contradiction with Lemma 3.15, once that $(J(t_n u_n))$ is bounded from above.

Case: $w \neq 0$

Recalling that

$$|u_n| = |w_n| \|u_n\|_E \quad \text{and} \quad \frac{u_n(x)}{\|u_n\|_E} = w_n(x) \longrightarrow w(x), \quad a.e. \text{ in } \mathbb{R}^N$$

we will get that

$$|w_n(x)| \longrightarrow |w(x)|, \quad a.e. \text{ in } \mathbb{R}^N.$$

Furthermore, from the fact that $\|u_n\|_E \rightarrow +\infty$, we can conclude that

$$|u_n(x)| = |w_n(x)| \|u_n\|_E \rightarrow +\infty, \quad \text{as } n \rightarrow \infty \text{ for } x \in \{y \in \mathbb{R}^N : w(y) \neq 0\}.$$

By (3.37),

$$(3.38) \quad 0 = \limsup_{n \rightarrow \infty} \frac{c}{\|u_n\|_E^m} = \limsup_{n \rightarrow \infty} \frac{J(u_n)}{\|u_n\|_E^m}.$$

As $\|u_n\|_{\Phi} \geq 1$ and $\|u_n\|_{V, \Phi} \geq 1$ for every $n \geq n_0$,

$$(3.39) \quad \int_{\mathbb{R}^N} \Phi(|\nabla u_n|) dx \leq \|\nabla u_n\|_{\Phi}^m \quad \text{and} \quad \int_{\mathbb{R}^N} V(x) \Phi(|u_n|) dx \leq \|u_n\|_{V, \Phi}^m, \quad \forall n \geq n_0.$$

Thus, it follows from (f₄), (3.38), (3.39) and Fatou's Lemma that

$$\begin{aligned} 0 &= \limsup_{n \rightarrow \infty} \frac{J(u_n)}{\|u_n\|_E^m} \\ &\leq \limsup_{n \rightarrow \infty} \left[\frac{1}{\|u_n\|_E^m} \int_{\mathbb{R}^N} \Phi(|\nabla u_n|) dx + \frac{1}{\|u_n\|_E^m} \int_{\mathbb{R}^N} V(x) \Phi(|u_n|) dx \right] \\ &\quad - \liminf_{n \rightarrow \infty} \left[\frac{1}{2} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))F(u_n(y))}{|x-y|^\lambda \|u_n\|_E^m} dx dy \right] \\ &\leq 2 - \frac{1}{2} \liminf_{n \rightarrow \infty} \left[\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))}{|x-y|^\lambda} \frac{F(u_n(y))}{|u_n(x)|^{\frac{m}{2}}} |w_n(x)| \frac{F(u_n(y))}{|u_n(y)|^{\frac{m}{2}}} |w_n(y)| dx dy \right] \\ &= -\infty \end{aligned}$$

which is a contradiction. This shows that (u_n) is bounded in E .

The cases *i*) and *ii*) are analogous to the case *iii*).

□

4. GROUND STATE SOLUTIONS FOR (P)

Since that the Cerami sequence (u_n) given in (3.37) is bounded in E , by Lemma 2.8, we can assume that for some subsequence, there is $u \in E$ such that

$$(4.1) \quad u_n \xrightarrow{*} u \text{ in } D^{1,\Phi}(\mathbb{R}^N) \quad \text{and} \quad u_n(x) \rightarrow u(x) \text{ a.e. } \mathbb{R}^N.$$

Fix $v \in C_0^\infty(\mathbb{R}^N)$. By boundedness of Cerami sequence (u_n) , we have $J'(u_n)(v - u_n) = o_n(1)$, hence, since Φ is a convex function, it is possible to show that

$$(4.2) \quad Q(v) - Q(u_n) \geq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))f(u_n(y))(v(y) - u_n(y))}{|x - y|^\lambda} dx dy + o_n(1).$$

For the Lemma 3.2, we have to

$$\liminf_{n \rightarrow \infty} \int_{\mathbb{R}^N} \Phi(|\nabla u_n|) dx \geq \int_{\mathbb{R}^N} \Phi(|\nabla u|) dx.$$

Now, due to the Fatou's Lemma, we have

$$\liminf_{n \rightarrow \infty} \int_{\mathbb{R}^N} V(x)\Phi(|u_n|) dx \geq \int_{\mathbb{R}^N} V(x)\Phi(|u|) dx.$$

Therefore,

$$(4.3) \quad \liminf_{n \rightarrow \infty} Q(u_n) \geq Q(u).$$

From (4.2) and (4.3) together with the limits (3.20) and (3.22) (or the limit (3.34) and (3.35)), we get

$$Q(v) - Q(u) \geq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))f(u(y))(v(y) - u(y))}{|x - y|^\lambda} dx dy.$$

As $E = \overline{C_0^\infty(\mathbb{R}^N)}^{\|\cdot\|_E}$ and $\Phi \in (\Delta_2)$, we conclude that

$$(4.4) \quad Q(v) - Q(u) \geq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))f(u(y))(v(y) - u(y))}{|x - y|^\lambda} dx dy, \quad \forall v \in E.$$

In other words, u is a critical point of the J functional. And from Proposition 3.11, we can conclude that u is a weak solution for (P). Now, we substitute $v = u^+ := \max\{0, u(x)\}$ in (4.4) we get

$$- \int_{\mathbb{R}^N} \Phi(|\nabla u^-|) dx - \int_{\mathbb{R}^N} V(x)\Phi(u^-) dx \geq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))f(u(y))(u^-(y))}{|x - y|^\lambda} dx dy = 0,$$

which leads to

$$\int_{\mathbb{R}^N} \Phi(|\nabla u^-|) dx = 0 \quad \text{and} \quad \int_{\mathbb{R}^N} V(x)\Phi(u^-) dx = 0$$

whence it is readily inferred that $u^- = 0$, therefore, u is a weak nonnegative solution.

Note that u is nontrivial. In the sense, consider a sequence $(\varphi_k) \subset C_0^\infty(\mathbb{R}^N)$ such that $\varphi_k \rightarrow u$ in $D^{1,\Phi}(\mathbb{R}^N)$. Since (u_n) bounded, we get $J'(u_n)(\varphi_k - u_n) = o_n(1)\|\varphi_k\| - o_n(1)$. As Φ is convex, we can show that

$$(4.5) \quad Q(\varphi_k) - Q(u_n) - o_n(1) \geq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))f(u_n(y))(\varphi_k(y) - u_n(y))}{|x - y|^\lambda} dx dy.$$

Since $(\|\varphi_k\|)_{k \in \mathbb{N}}$ is a bounded sequence, it follows from (4.5) and from limits (3.21) and (3.22) (or the limit (3.34) and (3.35)) it follows that

$$Q(\varphi_k) - \limsup_{n \rightarrow \infty} Q(u_n) \geq \limsup_{n \rightarrow \infty} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))f(u_n(y))(\varphi_k(y) - u_n(y))}{|x - y|^\lambda} dx dy,$$

for every $k \in \mathbb{N}$. Now, note that being $\Phi \in (\Delta_2)$ and $\varphi_k \rightarrow u$ in E , we conclude from the inequality above that

$$(4.6) \quad Q(u) \geq \limsup_{n \rightarrow \infty} Q(u_n).$$

From (4.3) and (4.6),

$$(4.7) \quad Q(u) = \lim_{n \rightarrow \infty} Q(u_n).$$

By (3.20) (or (3.33)), we have

$$(4.8) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))F(u_n(y))}{|x - y|^\lambda} dx dy = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))F(u(y))}{|x - y|^\lambda} dx dy,$$

Therefore

$$\begin{aligned} 0 < c &= \lim_{n \rightarrow \infty} J(u_n) \\ &= \int_{\mathbb{R}^N} \Phi(|\nabla u|) dx + \int_{\mathbb{R}^N} V(x)\Phi(|u|) dx - \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))F(u(y))}{|x - y|^\lambda} dx dy \\ &= J(u), \end{aligned}$$

that is, $u \neq 0$.

4.1. Proof of Theorem 1.1 and 1.2. Assuming the assumptions of Theorem 1.1, the above argument guarantees the existence of a nonnegative solution for problem (P), thus showing the first part of Theorem 1.1. Now, to show to study the boundedness of nonnegative solutions of the problem (P) we will make heavy use of hypotheses (ϕ_5) and $\lambda < 2m$.

Now, we begin by presenting a technical result, which is an adaptation of a result that can be found in [2].

Lemma 4.1. *Let $u \in E$ be a nonnegative solution of (P), $x_0 \in \mathbb{R}^N$ and $R_0 > 0$. Then*

$$\int_{\mathcal{A}_{k,t}} |\nabla u|^m dx \leq C \left(\int_{\mathcal{A}_{k,s}} \left| \frac{u-k}{s-t} \right|^{m^*} dx + (k^{m^*} + 1)|\mathcal{A}_{k,s}| \right) + C \left(\int_{\mathcal{A}_{k,s}} \left| \frac{u-k}{s-t} \right|^{m^*} dx + (k^{m^*} + 1)|\mathcal{A}_{k,s}| \right)^{\frac{1}{s}}$$

where $0 < t < s < R_0$, $k > 1$, $\mathcal{A}_{k,\rho} = \{x \in B_\rho(x_0) : u(x) > k\}$ and $C > 0$ is a constant that does not depend on k .

Proof: Let $u \in E$ be a weak solution nonnegative of (P) and $x_0 \in \mathbb{R}^N$. Moreover, fix $0 < t < s < R_0$ and $\zeta \in C_0^\infty(\mathbb{R}^N)$ verifying

$$0 \leq \zeta \leq 1, \quad \text{supp}(\zeta) \subset B_s(x_0), \quad \zeta \equiv 1 \text{ on } B_t(x_0) \quad \text{and} \quad |\nabla \zeta| \leq \frac{2}{s-t}.$$

For $k > 1$, set $\varphi = \zeta^m(u - k)^+$ and

$$J = \int_{\mathcal{A}_{k,s}} \Phi(|\nabla u|) \zeta^m dx,$$

Using φ as a test function and $\ell\Phi(t) \leq \phi(t)t^2$, we find

$$\begin{aligned} \ell J &\leq m \int_{\mathcal{A}_{k,s}} \zeta^{m-1}(u-k)^+ \phi(|\nabla u|)|\nabla u||\nabla \zeta| dx - \int_{\mathcal{A}_{k,s}} V(x)\phi(u)u\zeta^m(u-k)^+ dx \\ &\quad + \int_{\mathcal{A}_{k,s}} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))f(u(y))\zeta^m(y)(u(y)-k)^+}{|x-y|^\lambda} dx dy \\ &\leq \int_{\mathcal{A}_{k,s}} \zeta^{m-1}(u-k)^+ \phi(|\nabla u|)|\nabla u||\nabla \zeta| dx \\ &\quad + C_1(s, \lambda, N) \left| \int_{\mathbb{R}^N} K(x)^s |F(u)|^s dx \right|^{\frac{1}{s}} \left| \int_{\mathbb{R}^N} W(x)|f(u)|^s |\zeta^m(u-k)^+|^s dx \right|^{\frac{1}{s}} \end{aligned}$$

As in (3.6), given $\eta > 0$, there exists $C_\varepsilon > 0$ such that

$$K(x)^s f(t)^s \leq \frac{\varepsilon}{s} K(x)^s a(t)t^{2-s} + C_\varepsilon \phi_*(t)t^{2-s}, \quad \forall t \geq 0 \text{ and } x \in \mathbb{R}^N.$$

Thus,

(4.9)

$$\begin{aligned} \ell J &\leq m \int_{\mathcal{A}_{k,s}} \zeta^{m-1}(u-k)^+ \phi(|\nabla u|)|\nabla u||\nabla \zeta| dx \\ &\quad + C_2 \left[\int_{\mathcal{A}_{k,s}} K^s(x)a(|u|)u^{2-s}(\zeta^m(u-k)^+)^s dx + \int_{\mathcal{A}_{k,s}} \phi_*(|u|)u^{2-s}(\zeta^m(u-k)^+)^s dx \right]^{\frac{1}{s}} \end{aligned}$$

where $C_2 = C_1(s, \lambda, N) \left(\int_{\mathbb{R}^N} K(x)^s |F(u)|^s dx \right)^{\frac{1}{s}}$. For each $\tau \in (0, 1)$, the Young's inequalities gives

$$(4.10) \quad \phi(|\nabla u|)|\nabla u||\nabla \zeta|\zeta^{m-1}(u-k)^+ \leq \tilde{\Phi}(\phi(|\nabla u|)|\nabla u|\zeta^{m-1}\tau) + C_3\Phi\left(\left|\frac{u-k}{s-t}\right|\right).$$

It follows from Lemma 2.4,

$$(4.11) \quad \tilde{\Phi}(\phi(|\nabla u|)|\nabla u|\zeta^{m-1}\tau) \leq C_4(\tau\zeta^{m-1})^{\frac{m}{m-1}}\Phi(|\nabla u|).$$

From (4.9), (4.10) and (4.11),

$$\begin{aligned} \ell J &\leq mC_4\tau^{\frac{m}{m-1}} \int_{\mathcal{A}_{k,s}} \Phi(|\nabla u|)\zeta^m + mC_3 \int_{\mathcal{A}_{k,s}} \Phi\left(\left|\frac{u-k}{s-t}\right|\right) dx \\ &\quad + C_2 \left[\int_{\mathcal{A}_{k,s}} K^s(x)a(|u|)u^{2-s}(\zeta^m(u-k)^+)^s dx + \int_{\mathcal{A}_{k,s}} \phi_*(|u|)u^{2-s}(\zeta^m(u-k)^+)^s dx \right]^{\frac{1}{s}} \end{aligned}$$

Choosing $\tau \in (0, 1)$ such that $0 < mC_4\tau^{\frac{m}{m-1}} < \ell$, we derive

(4.12)

$$J \leq C_5 \int_{\mathcal{A}_{k,s}} \Phi\left(\left|\frac{u-k}{s-t}\right|\right) dx + C_2 \left[\int_{\mathcal{A}_{k,s}} K^s(x)a(|u|)u^{2-s}(\zeta^m(u-k)^+)^s dx + \int_{\mathcal{A}_{k,s}} \phi_*(|u|)u^{2-s}(\zeta^m(u-k)^+)^s dx \right]^{\frac{1}{s}}$$

By Young's inequalities,

$$(4.13) \quad \phi_*(u)u^{2-s}(\zeta^m(u-k)^+)^s \leq C_6\Phi_*\left(\left|\frac{u-k}{s-t}\right|\right) + C_6\Phi_*(k).$$

and

$$(4.14) \quad a(u)u^{2-s}(\zeta^m(u-k)^+)^s \leq C_6 A\left(\left|\frac{u-k}{s-t}\right|\right) + C_6 A(k).$$

Therefore, a combination of (4.12) and (4.13), yields

(4.15)

$$J \leq C_7 \int_{\mathcal{A}_{k,s}} \Phi\left(\left|\frac{u-k}{s-t}\right|\right) dx + C_7 \left[\int_{\mathcal{A}_{k,s}} A\left(\left|\frac{u-k}{s-t}\right|\right) dx + \int_{\mathcal{A}_{k,s}} A(k) dx + \int_{\mathcal{A}_{k,s}} \Phi_*\left(\left|\frac{u-k}{s-t}\right|\right) dx + \int_{\mathcal{A}_{k,s}} \Phi_*(k) dx \right]^{\frac{1}{s}}.$$

Now, using that $\ell \leq m < a_2 < m^*$ and applying the Lemmas 2.1, 2.2 and the Remark 1 for functions Φ , A and Φ_* , respectively, we get

$$\Phi\left(\left|\frac{u-k}{s-t}\right|\right) \leq \Phi(1)\left(\left|\frac{u-k}{s-t}\right|^{m^*} + 1\right),$$

$$A\left(\left|\frac{u-k}{s-t}\right|\right) \leq A(1)\left(\left|\frac{u-k}{s-t}\right|^{m^*} + 1\right) \text{ and } A(k) \leq (k^{m^*} + 1).$$

and

$$\Phi_*\left(\left|\frac{u-k}{s-t}\right|\right) \leq \Phi_*(1)\left(\left|\frac{u-k}{s-t}\right|^{m^*} + 1\right) \text{ and } \Phi_*(k) \leq (k^{m^*} + 1).$$

From (4.15) and the inequality above,

$$J \leq C_8 \left(\int_{\mathcal{A}_{k,s}} \left|\frac{u-k}{s-t}\right|^{m^*} dx + (k^{m^*} + 1)|\mathcal{A}_{k,s}| \right) + C_8 \left(\int_{\mathcal{A}_{k,s}} \left|\frac{u-k}{s-t}\right|^{m^*} dx + (k^{m^*} + 1)|\mathcal{A}_{k,s}| \right)^{\frac{1}{s}}$$

The result follows from condition (ϕ_5). □

Lemma 4.2. *Let $u \in E$ be a nonnegative solution of (P). Then, $u \in L_{loc}^\infty(\mathbb{R}^N)$.*

Proof: To begin with, consider Λ a compact subset on \mathbb{R}^N . Fix $R_1 \in (0, 1)$, $x_0 \in \Lambda$ and define the sequences

$$\sigma_n = \frac{R_1}{2} + \frac{R_1}{2^{n+1}} \quad \text{and} \quad \bar{\sigma}_n = \frac{\sigma_n + \sigma_{n+1}}{2} \quad \text{for } n = 0, 1, 2, \dots.$$

Note that

$$\sigma_n \downarrow \frac{R_1}{2} \quad \text{and} \quad \sigma_{n+1} < \bar{\sigma}_n < \sigma_n < R_1.$$

Since E is continuously embedded in $W_{loc}^{1,m}(\mathbb{R}^N)$, it follows from Lebesgue dominated convergence theorem,

$$(4.16) \quad \lim_{M \rightarrow \infty} \int_{B_{R_1}(x_0)} ((u-M)^+)^{m^*} dx = 0,$$

hence, there is $M^* \geq 1$ which depends on x_0 and R_1 , such that

$$(4.17) \quad \int_{B_{R_1}(x_0)} ((u-M)^+)^{m^*} dx \leq 1, \quad \text{for } M \geq M^*.$$

Now, consider $K > 4M^*$ and for every $n \in \mathbb{N}$ define

$$K_n = \frac{K}{2} \left(1 - \frac{1}{2^{n+1}}\right) \quad \text{and} \quad J_n = \int_{\mathcal{A}_{K_n, \sigma_n}} ((u-K_n)^+)^{m^*} dx, \quad \text{for } n = 0, 1, 2, \dots.$$

and

$$\xi_n = \xi \left(\frac{2^{n+1}}{R_1} \left(|x - x_0| - \frac{R_1}{2} \right) \right), \quad x \in \mathbb{R}^N \text{ and } n = 0, 1, 2, \dots$$

where $\xi \in C^1(\mathbb{R})$ satisfies

$$0 \leq \xi \leq 1, \quad \xi(t) = 1 \text{ for } t \leq \frac{1}{2} \quad \text{and} \quad \xi(t) = 0 \text{ for } t \geq \frac{3}{4}.$$

From definition of ξ_n ,

$$\xi_n = 1 \text{ in } B_{\sigma_{n+1}}(x_0) \quad \text{and} \quad \xi_n = 0 \text{ outside } B_{\bar{\sigma}_n}(x_0),$$

consequently

$$(4.18) \quad J_{n+1} \leq \int_{B_{R_1}(x_0)} ((u - K_{n+1})^+ \xi_n)^{m^*} dx.$$

Note that

$$(4.19) \quad |\nabla((u - K_{n+1})^+ \xi_n)|^m \leq 2^m (|\nabla u|^m \xi_n^m + \frac{2^{m(n+1)}}{R_1^m} ((u - K_{n+1})^+)^m \chi_{B_{R_1}(x_0)})$$

and

$$(4.20) \quad \text{supp}(\nabla((u - K_{n+1})^+ \xi_n)) \subset \mathcal{A}_{K_{n+1}, \bar{\sigma}_n}.$$

The inequality (4.19) together with the hypothesis (ϕ_5) guarantee that $|\nabla((u - K_{n+1})^+ \xi_n)| \in L^m(\mathbb{R}^N)$. Therefore, we can conclude that $(u - K_{n+1})^+ \xi_n \in W^{1,m}(\mathbb{R}^N)$. In view of this, there is a constant $C_1 = C_1(N, m) > 0$ satisfying

$$(4.21) \quad J_{n+1} \leq C_1 \left(\int_{\mathcal{A}_{K_{n+1}, \bar{\sigma}_n}} |\nabla((u - K_{n+1})^+ \xi_n)|^m dx \right)^{\frac{m^*}{m}}.$$

From (4.19) and (4.21),

$$(4.22) \quad J_{n+1}^{\frac{m}{m^*}} \leq C_2 \left(\int_{\mathcal{A}_{K_{n+1}, \bar{\sigma}_n}} |\nabla u|^m dx + 2^{mn} \int_{\mathcal{A}_{K_{n+1}, \bar{\sigma}_n}} ((u - K_{n+1})^+)^m dx \right)$$

for some constant $C_2 = C(N, m, R_1) > 0$. Applying the lemma 4.1 to the previous inequality, we get

$$(4.23) \quad J_{n+1}^{\frac{m}{m^*}} \leq C_3 \left(\int_{\mathcal{A}_{K_{n+1}, \sigma_n}} \left| \frac{u - K_{n+1}}{\sigma_n - \bar{\sigma}_n} \right|^{m^*} dx + (K_{n+1}^{m^*} + 1) |\mathcal{A}_{K_{n+1}, \sigma_n}| + 2^{mn} \int_{\mathcal{A}_{K_{n+1}, \bar{\sigma}_n}} ((u - K_{n+1})^+)^m dx \right) + C_3 \left(\int_{\mathcal{A}_{K_{n+1}, \sigma_n}} \left| \frac{u - K_{n+1}}{\sigma_n - \bar{\sigma}_n} \right|^{m^*} dx + (K_{n+1}^{m^*} + 1) |\mathcal{A}_{K_{n+1}, \sigma_n}| \right)^{\frac{1}{s}}$$

where $C_3 > 0$ is a constant that depends only on N, m and R_1 . Being $|\sigma_n - \bar{\sigma}_n| = \frac{R_1}{2^{n+3}}$, we conclude that

$$(4.24) \quad J_{n+1}^{\frac{m}{m^*}} \leq C_4(N, m, R_1) \left(2^{mn} \int_{\mathcal{A}_{K_{n+1}, \sigma_n}} ((u - K_{n+1})^+)^{m^*} dx + (K_{n+1}^{m^*} + 1) |\mathcal{A}_{K_{n+1}, \sigma_n}| + 2^{mn} \int_{\mathcal{A}_{K_{n+1}, \bar{\sigma}_n}} ((u - K_{n+1})^+)^m dx \right) + C_4(N, m, R_1) \left(2^{mn} \int_{\mathcal{A}_{K_{n+1}, \sigma_n}} ((u - K_{n+1})^+)^{m^*} dx + (K_{n+1}^{m^*} + 1) |\mathcal{A}_{K_{n+1}, \sigma_n}| \right)^{\frac{1}{s}}$$

Combined the inequality above with $t^m \leq t^{m^*} + 1$, for $t \geq 0$ and using that $\bar{\sigma}_n < \sigma_n$, we get that

$$(4.25) \quad \begin{aligned} J_{n+1}^{\frac{m}{m^*}} &\leq C_4(N, m, R_1) \left(2^{mn} \int_{\mathcal{A}_{K_{n+1}, \sigma_{n_k}}} ((u - K_{n+1})^+)^{m^*} dx + (K^{m^*} + 2^{mn} + 1) |\mathcal{A}_{K_{n+1}, \sigma_n}| \right) \\ &\quad + C_4(N, m, R_1) \left(2^{mn} \int_{\mathcal{A}_{K_{n+1}, \sigma_n}} ((u - K_{n+1})^+)^{m^*} dx + (K^{m^*} + 2^{mn} + 1) |\mathcal{A}_{K_{n+1}, \sigma_n}| \right)^{\frac{1}{s}} \end{aligned}$$

On the other hand, since $K_{n+1} - K_n = \frac{K}{2^{n+3}}$,

$$(4.26) \quad \begin{aligned} \left(\frac{K}{2^{n+3}} \right)^{m^*} |\mathcal{A}_{K_{n+1}, \sigma_n}| &= (K_{n+1} - K_n)^{m^*} |\mathcal{A}_{K_{n+1}, \sigma_n}| \\ &\leq \int_{\mathcal{A}_{K_{n+1}, \sigma_n}} (K_{n+1} - K_n)^{m^*} dx \\ &\leq \int_{\mathcal{A}_{K_{n+1}, \sigma_n}} \Phi_*((u - K_n)^+) \chi_{\mathcal{A}_{K_{n+1}, \sigma_n}}(x) dx \leq J_n, \end{aligned}$$

which yields

$$(4.27) \quad |\mathcal{A}_{K_{n+1}, \sigma_n}| \leq \frac{1}{\left(\frac{K}{2^{n+3}} \right)^{m^*}} J_n.$$

Thus,

$$\begin{aligned} \int_{\mathcal{A}_{K_{n+1}, \sigma_n}} ((u - K_{n+1})^+)^{m^*} dx &\leq \int_{\mathcal{A}_{K_{n+1}, \sigma_n}} ((u - K_n)^+)^{m^*} dx + \int_{\mathcal{A}_{K_{n+1}, \sigma_n}} (K_{n+1} - K_n)^{m^*} dx \\ &\leq \int_{\mathcal{A}_{K_n, \sigma_n}} ((u - K_n)^+)^{m^*} dx + |K_{n+1} - K_n|^{m^*} |\mathcal{A}_{K_{n+1}, \sigma_n}| \\ &\leq 2J_n \end{aligned}$$

and consequently

$$(4.28) \quad \begin{aligned} J_{n+1}^{\frac{m}{m^*}} &\leq C_5(N, m, R_1) \left(2^{mn+1} + 2^{n(m+m^*)} + (2^{mn} + 1) 2^{n(m+m^*)} \right) J_n \\ &\quad + C_5(N, m, R_1) \left(2^{mn+1} + 2^{n(m+m^*)} + (2^{mn} + 1) 2^{n(m+m^*)} \right)^{\frac{1}{s}} J_n^{\frac{1}{s}} \end{aligned}$$

Due to the fact that $K > 4M^*$, we conclude from the inequality 4.17 that

$$J_n \leq \int_{B_{R_1}(x_0)} ((u - M^*)^+)^{m^*} dx \leq 1, \quad \text{for } n = 0, 1, 2, \dots,$$

hence,

$$(4.29) \quad J_{n+1} \leq CD^n J_n^{1+\omega},$$

where $C = 2C_5(N, m, R_1)$, $D = 2^{2(m+m^*)\frac{m}{m^*}}$ and $\omega = \frac{m^*}{sm} - 1$.

We claim that

$$J_0 \leq C^{-\frac{1}{\omega}} D^{-\frac{1}{\omega^2}}, \quad \text{for } K \geq K^*.$$

Indeed, note that,

$$(4.30) \quad J_0 = \int_{\mathcal{A}_{K_0, \sigma_0}} ((u - K_0)^+)^{m^*} dx \leq \int_{B_{R_1}(x_0)} ((u - K_0)^+)^{m^*} dx$$

Since E is continuously embedded in $W_{loc}^{1,m}(\mathbb{R}^N)$, it follows from Lebesgue dominated convergence theorem,

$$\lim_{K \rightarrow \infty} \int_{B_{R_1}(x_0)} ((u - K)^+)^{m^*} dx = 0.$$

Therefore, there exists $K^* \geq 5M^*$ that depends on x_0 , such that

$$(4.31) \quad \int_{B_{R_1}(x_0)} ((u - K)^+)^{m^*} dx \leq C^{-\frac{1}{\omega}} D^{-\frac{1}{\omega^2}}, \quad \text{for } K \geq K^*.$$

From (4.30) and (4.31),

$$(4.32) \quad J_0 \leq C^{-\frac{1}{\omega}} D^{-\frac{1}{\omega^2}}, \quad \text{for } K \geq K^*.$$

Fix $K = K^*$, by Lemma [32, Lemma 4.7], we deduce that

$$J_n \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

On the other hand,

$$\lim_{n \rightarrow \infty} J_n = \lim_{n \rightarrow \infty} \int_{\mathcal{A}_{K_n, \sigma_n}} ((u - K_n)^+)^{m^*} dx = \int_{\mathcal{A}_{\frac{K^*}{2}, \frac{R_1}{2}}} ((u - \frac{K^*}{2})^+)^{m^*} dx$$

Hence,

$$\int_{\mathcal{A}_{\frac{K^*}{2}, \frac{R_1}{2}}} ((u - \frac{K^*}{2})^+)^{m^*} dx = 0,$$

leading to

$$u(x) \leq \frac{K^*}{2} \quad \text{a.e. in } B_{\frac{R_1}{2}}(x_0).$$

Since x_0 is arbitrary and Λ is a compact subset, the last inequality ensures that

$$(4.33) \quad u(x) \leq \frac{M}{2} \quad \text{a.e. in } \Lambda$$

for some constant $M > 0$. By the arbitrariness of Λ , we conclude that $u \in L_{loc}^\infty(\mathbb{R}^N)$. \square

This Lemmas guarantees that the Theorem 1.1 is valid. The proof of Theorem 1.2 will be divided into the following Lemmas:

Lemma 4.3. $u \in C_{loc}^{1,\alpha}(\mathbb{R}^N)$.

Proof. From the definition of F , there exists $C_1 > 0$ such that

$$|F(t)|^s \leq C_1 \Phi_*(|t|), \quad \forall t \geq 1$$

Hence, by Hölder inequality

$$\begin{aligned}
\int_{[u \geq 1]} \frac{K(y)F(u(y))}{|x-y|^\lambda} dy &\leq \left(\int_{[u \geq 1]} \frac{K(y)^{\frac{s-1}{s-1}}}{|x-y|^{\frac{\lambda(s-1)}{s}}} dy \right)^{\frac{s-1}{s}} \left(\int_{[u \geq 1]} F(u(y))^s dy \right)^{\frac{1}{s}} \\
&\leq C_1 \left(\int_{\mathbb{R}^N} \frac{K(y)^{\frac{s-1}{s-1}}}{|x-y|^{\frac{\lambda(s-1)}{s}}} dy \right)^{\frac{s-1}{s}} \left(\int_{\mathbb{R}^N} \Phi_*(u) dy \right)^{\frac{1}{s}} \\
&\leq C_1 \left(\int_{\mathbb{R}^N} \Phi_*(u) dy \right)^{\frac{1}{s}} \left(\int_{|x-y| < 1} \frac{K(y)^{\frac{s-1}{s-1}}}{|x-y|^{\frac{\lambda(s-1)}{s}}} dy + \int_{|x-y| \geq 1} \frac{K(y)^{\frac{s-1}{s-1}}}{|x-y|^{\frac{\lambda(s-1)}{s}}} dy \right)^{\frac{s-1}{s}} \\
&\leq C_1 \left(\int_{\mathbb{R}^N} \Phi_*(u) dy \right)^{\frac{1}{s}} \left(\|K\|_\infty^s \int_{|x-y| < 1} \frac{1}{|x-y|^{\frac{\lambda(s-1)}{s}}} dy + \int_{|x-y| \geq 1} K(y)^{\frac{s-1}{s-1}} dy \right)^{\frac{s-1}{s}} \\
&\leq C_1 \left(\int_{\mathbb{R}^N} \Phi_*(u) dy \right)^{\frac{1}{s}} \left(\|K\|_\infty^s \int_0^1 r^{N-1-\frac{\lambda(s-1)}{s}} dy + \|K\|_{L^{\frac{s-1}{s}}(\mathbb{R}^N)}^{\frac{s-1}{s}} \right)^{\frac{s-1}{s}}
\end{aligned}$$

On the other hand,

$$\begin{aligned}
\int_{[u \leq 1]} \frac{K(y)F(u(y))}{|x-y|^\lambda} dy &\leq \|F\|_{L^\infty([0,1])} \int_{\mathbb{R}^N} \frac{K(y)}{|x-y|^\lambda} dy \\
&\leq \|F\|_{L^\infty([0,1])} \left(\int_{|x-y| < 1} \frac{K(y)}{|x-y|^\lambda} dy + \int_{|x-y| \geq 1} \frac{K(y)}{|x-y|^\lambda} dy \right) \\
&\leq \|F\|_{L^\infty([0,1])} \left(\|K\|_\infty \int_0^1 r^{N-1-\lambda} dy + \|K\|_{L^1(\mathbb{R}^N)} \right),
\end{aligned}$$

that is,

$$\int_{\mathbb{R}^N} \frac{K(y)F(u(y))}{|x-y|^\lambda} dy \leq C_2,$$

where

$$\begin{aligned}
C_2 = \left\{ C_1 \left(\int_{\mathbb{R}^N} \Phi_*(u) dy \right)^{\frac{1}{s}} \left(\|K\|_\infty^s \int_0^1 r^{N-1-\frac{\lambda(s-1)}{s}} dy + \|K\|_{L^{\frac{s-1}{s}}(\mathbb{R}^N)}^{\frac{s-1}{s}} \right)^{\frac{s-1}{s}}, \right. \\
\left. \|F\|_{L^\infty([0,1])} \left(\|K\|_\infty \int_0^1 r^{N-1-\lambda} dy + \|K\|_{L^1(\mathbb{R}^N)} \right) \right\},
\end{aligned}$$

showing that

$$(4.34) \quad \int_{\mathbb{R}^N} \frac{K(y)F(u(y))}{|x-y|^\lambda} dy \in L^\infty(\mathbb{R}^N).$$

Consider the scalar measurable function $Z : \Omega \times \mathbb{R} \times \mathbb{R}^N \rightarrow \mathbb{R}$ given by

$$Z(x, t, p) = V(x)\varphi(|t|)t - \left(\int_{\mathbb{R}^N} \frac{K(y)F(u(y))}{|x-y|^\lambda} dy \right) K(x)\tilde{f}(t)$$

with

$$\varphi(t) = \begin{cases} \phi(t) & , \quad \text{for } 0 < t \leq M/2 \\ \phi(M/2) & , \quad \text{for } t \geq M/2 \end{cases},$$

and

$$\varphi(t) = \begin{cases} f(t) & , \quad \text{for } 0 < t \leq M/2 \\ f(M/2) & , \quad \text{for } t \geq M/2 \end{cases} ,$$

where $M > 0$ is the constant satisfying (4.33). By (4.34), there exists a constant C_3 such that

$$|B(x, t, p)| \leq C_3$$

This fact together with the hypothesis (ϕ_5) allows us to apply the theorem of regularity due to Lieberman [32, Theorem 1.7]. Thus showing the result \square

Corollary 4.4. *If $\inf_{x \in \mathbb{R}^N} V(x) > 0$, then $\lim_{|x| \rightarrow \infty} u(x) = 0$.*

Proof: Repeating the same arguments used in Lemma 4.2 with $K = \frac{\delta}{2}$ and $K_0 = \frac{\delta}{4}$ for $\delta > 0$, we find

$$J_0 = \int_{\mathcal{A}_{K_0, \sigma_0}} ((u - K_0)^+)^{m^*} dx \leq \int_{B_{\sigma_0}(x_0)} \left((u - \frac{\delta}{4})^+ \right)^{m^*} dx$$

As $\inf_{x \in \mathbb{R}^N} V(x) > 0$, then by the condition (ϕ_5) we have that E is continuously embedded in $W^{1, m^*}(\mathbb{R}^N)$. By Lebesgue dominated convergence theorem,

$$\lim_{|x_0| \rightarrow \infty} \int_{B_{\sigma_0}(x_0)} \left((u - \frac{\delta}{4})^+ \right)^{m^*} dx = 0$$

we also have

$$\lim_{|x_0| \rightarrow \infty} J_0 = 0.$$

Then, there exists $R^* > 0$, such that

$$J_0 \leq C^{-\frac{1}{\omega}} D^{-\frac{1}{\omega^2}} \quad \text{for } |x_0| \geq R^*.$$

where the constants C , D and ω are given in (4.29). Applying the Lemma [32, Lemma 4.7],

$$\lim_{n \rightarrow \infty} J_n = 0 \quad \text{if } |x_0| \geq R^*.$$

that is,

$$\int_{\mathcal{A}_{\frac{\delta}{4}, \frac{R_1}{2}}} \left((u - \frac{\delta}{4})^+ \right)^{m^*} dx = 0 \quad \text{for } |x_0| \geq R^*.$$

The continuity of u yields

$$u(x) \leq \frac{\delta}{4}, \quad \forall x \in \mathbb{R}^N \setminus B_{R^*}(0),$$

finishing the proof of the lemma. \square

Corollary 4.5. *Let $u \in E$ be a nonnegative solution of (P). Then, u is positive solution.*

Proof: If $\Omega \subset \mathbb{R}^N$ is a bounded domain, the Lemma 4.2 implies that $u \in C^1(\overline{\Omega})$. Using this fact, in the sequel, we fix $M_1 > \max \{ \|\nabla u\|_{L^\infty(\overline{\Omega})}, 1 \}$ and

$$\varphi(t) = \begin{cases} \phi(t) & , \quad \text{for } 0 < t \leq M_1 \\ \frac{\phi(M_1)}{M_1^{\beta-2}} t^{\beta-2} & , \quad \text{for } t \geq M_1 \end{cases} ,$$

where β is given in the hypothesis (ϕ_7) . Still by condition (ϕ_7) , there are $\alpha_1, \alpha_2 > 0$ satisfying

$$(4.35) \quad \varphi(|y|)|y|^2 = \phi(|y|)|y|^2 \geq \alpha_1|y|^\beta \quad \text{and} \quad |\varphi(|y|)y| \leq \alpha_2|y|^{\beta-1}, \quad \forall y \in \mathbb{R}^N.$$

Now, consider the vector measurable functions $G : \Omega \times \mathbb{R} \times \mathbb{R}^N \rightarrow \mathbb{R}^N$ given by $G(x, t, p) = \frac{1}{\alpha_1}\varphi(|p|)p$. From (4.35),

$$(4.36) \quad |G(x, t, p)| \leq \frac{\alpha_2}{\alpha_1}|p|^{\beta-1} \quad \text{and} \quad pG(x, t, p) \geq |p|^{\beta-1}, \quad \text{for all } (x, t, p) \in \Omega \times \mathbb{R} \times \mathbb{R}^N.$$

We next will consider the scalar measurable function $Z : \Omega \times \mathbb{R} \times \mathbb{R}^N \rightarrow \mathbb{R}$ given by

$$Z(x, t, p) = \frac{1}{\alpha_1}(V(x)\phi(|t|)t - \left(\int_{\mathbb{R}^N} \frac{K(y)F(u(y))}{|x-y|^\lambda} dy \right) K(x)f(t)).$$

Remember that from the inequality (3.6), there will be a constant $C_1 > 0$ satisfying

$$(4.37) \quad K(x)|f(t)| \leq C_1K(x)a(|t|)|t| + C_1\phi_*(|t|)|t|, \quad \forall t \in \mathbb{R}^N \quad \text{and} \quad x \in \mathbb{R}^N.$$

Fix $M \in (0, \infty)$. Through the condition (ϕ_7) and by a simple computation yields there exists $C_2 = C_2(M) > 0$ verifying

$$|Z(x, t, p)| \leq C_2|t|^{\beta-1}, \quad \text{for every } (x, t, p) \in \Omega \times (-M, M) \times \mathbb{R}^N.$$

By the arbitrariness of M , we can conclude that functions G and B fulfill the structure required by Trudinger [31]. Also, as u is a weak solution of (P) , we infer that u is a quasilinear problem solution

$$-\operatorname{div} G(x, u, \nabla u(x)) + Z(x, u, \nabla u(x)) = 0 \quad \text{in } \Omega.$$

By [31, Theorem 1.1], we deduce that $u > 0$ in Ω . By the arbitrariness of Ω , we conclude that $u > 0$ in \mathbb{R}^N . □

4.2. Proof of Theorem 1.3 and 1.4. Assuming the assumptions of Theorem 1.3, the arguments at the beginning of this section 4, guarantees the existence of a nonnegative solution for problem (P) . A result analogous to Lemma 4.1 can also be shown here. Therefore, considering $u \in E$ a nonnegative solution to (P) the Lemma 4.2 guarantees that $u \in L_{loc}^\infty(\mathbb{R}^N)$. With that, the Theorem 1.3 is proved. Through small changes, the proof of Theorem 1.4 will be similar to the proof presented in Theorem 1.2. □

5. ZERO MASS CASE

In this section, we will study the problem (P') and give the proof of Theorem 1.5. Here, the coefficient K only satisfies the conditions (K_0) , (K_1) and f satisfies (f_2) , (f_3) , (f_4) , (f_6) . By (f_6) , there is $C > 0$ such that

$$(5.1) \quad K(x)|f(t)| \leq C\phi_*(|t|)|t|, \quad x \in \mathbb{R}^N \quad \text{and} \quad t \in \mathbb{R}.$$

The following results are proved using the same arguments as presented in proof of Lemmas 3.5 and 3.6.

Lemma 5.1. *Suppose that (f_6) holds. For each $u \in D^{1,\Phi}(\mathbb{R}^N)$, there is a constant $C_1 > 0$ that does not depend on u , such that*

$$\left| \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))F(u(y))}{|x-y|^\lambda} dx dy \right| \leq C_1 \left(\int_{\mathbb{R}^N} \Phi_*(|u|) dx \right)^{\frac{2}{s}}.$$

Also, for each $u \in E$, there is a constant $C_2 > 0$, which does not depend on u , such that

$$(5.2) \quad \left| \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))f(u(y))v(y)}{|x-y|^\lambda} dx dy \right| \leq C \|v\|_{D^{1,\Phi}(\mathbb{R}^N)}, \quad \forall v \in D^{1,\Phi}(\mathbb{R}^N).$$

Lemma 5.2. *Suppose that (f₆) holds. Let (u_n) be a bounded sequence in $D^{1,\Phi}(\mathbb{R}^N)$, and consider $u \in D^{1,\Phi}(\mathbb{R}^N)$ such that $u_n \xrightarrow{*} u$ in $D^{1,\Phi}(\mathbb{R}^N)$. We will show the following limits*

$$(5.3) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} K(x)^s |F(u_n) - F(u)|^s dx = 0,$$

$$(5.4) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} K(x)^s |f(u_n)u_n - f(u)u|^s dx = 0$$

and

$$(5.5) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} K(x)^s |f(u_n)\varphi - f(u)\varphi|^s dx = 0$$

for $\varphi \in C_0^\infty(\mathbb{R}^N)$.

And as a consequence of the previous Lemma we have the following result.

Corollary 5.3. *Assume that (f₆) holds. Let (u_n) be a sequence bounded in $D^{1,\Phi}(\mathbb{R}^N)$ and $u \in D^{1,\Phi}(\mathbb{R}^N)$ such that $u_n \xrightarrow{*} u$ in $D^{1,\Phi}(\mathbb{R}^N)$. Then*

$$(5.6) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))F(u_n(y))}{|x-y|^\lambda} dx dy = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))F(u(y))}{|x-y|^\lambda} dx dy,$$

$$(5.7) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))f(u_n(y))u_n(y)}{|x-y|^\lambda} dx dy = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))f(u(y))u(y)}{|x-y|^\lambda} dx dy$$

and

$$(5.8) \quad \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))f(u_n(y))\varphi(y)}{|x-y|^\lambda} dx dy = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))f(u(y))\varphi(y)}{|x-y|^\lambda} dx dy$$

for all $\varphi \in C_0^\infty(\mathbb{R}^N)$.

By the inequalities (5.1) together with all the results presented above, it is verified that the function

$$\Psi(u) = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))F(u_n(y))}{|x-y|^\lambda} dx dy, \quad u \in D^{1,\Phi}(\mathbb{R}^N)$$

is well defined, is continuously differentiable and the Gateaux derivative $\Psi' : D^{1,\Phi}(\mathbb{R}^N) \rightarrow D^{1,\Phi}(\mathbb{R}^N)^*$ is given by

$$\Psi'(u)v = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))f(u_n(y))v(y)}{|x-y|^\lambda} dx dy, \quad \forall u, v \in D^{1,\Phi}(\mathbb{R}^N).$$

As in the previous sections, the energy functional $J : D^{1,\Phi}(\mathbb{R}^N) \rightarrow \mathbb{R}$ associated with the problem (P), which is given by

$$J(u) = \int_{\mathbb{R}^N} \Phi(|\nabla u|) dx - \frac{1}{2} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))F(u_n(y))}{|x-y|^\lambda} dx dy,$$

is continuous and Gateaux-differentiable with derivative $J' : D^{1,\Phi}(\mathbb{R}^N) \rightarrow (D^{1,\Phi}(\mathbb{R}^N))^*$ defined by

$$J'(u)v = \int_{\mathbb{R}^N} \phi(|\nabla u|) \nabla u \nabla v dx - \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))f(u_n(y))v(y)}{|x-y|^\lambda} dx dy$$

continuous from the norm topology of $D^{1,\Phi}(\mathbb{R}^N)$ to the weak*-topology of $(D^{1,\Phi}(\mathbb{R}^N))^*$.

We will say that $u \in D^{1,\Phi}(\mathbb{R}^N)$ is a critical point for the functional $J : D^{1,\Phi}(\mathbb{R}^N) \rightarrow \mathbb{R}$ if

$$(5.9) \quad \int_{\mathbb{R}^N} \Phi(\nabla v) dx - \int_{\mathbb{R}^N} \Phi(\nabla u) dx \geq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))f(u_n(y))(v(y) - u(y))}{|x - y|^\lambda} dx dy,$$

for all $v \in D^{1,\Phi}(\mathbb{R}^N)$.

Similar to the proof of [6, Lemma 4.1], we give the following result.

Proposition 5.4. *If $u \in D^{1,\Phi}(\mathbb{R}^N)$ is a critical point of J in $D^{1,\Phi}(\mathbb{R}^N)$, then u is a weak solution to (P).*

Also due to the inequality (5.1), it follows that J satisfies the geometry of the mountain pass. Therefore, by [8, Theorem 5.46], there is a sequence $(u_n) \subset D^{1,\Phi}(\mathbb{R}^N)$ such that

$$(5.10) \quad J(u_n) \rightarrow c \quad \text{and} \quad (1 + \|u_n\|)\|J'(u_n)\| \rightarrow 0.$$

where c is the mountain pass level given by

$$c = \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} J(\gamma(t))$$

with

$$\Gamma = \{\gamma \in C([0, 1], X) : \gamma(0) = 0 \text{ and } \gamma(1) = e\}.$$

Now, similar to the proof of Lemma 3.15, we give the following Lema.

Lemma 5.5. *Let (u_n) the Cerami sequence given in (5.10). There is a constant $M > 0$ such that $J(tu_n) \leq M$ for every $t \in [0, 1]$ and $n \in \mathbb{N}$.*

Knowing that the Corollary 5.3 holds, we can repeat the same arguments used in the proof of Proposition 2.1 to show that the Cerami sequence (u_n) given in (5.10) is bounded in $D^{1,\Phi}(\mathbb{R}^N)$.

Proposition 5.6. *The Cerami sequence (u_n) given in (5.10) is bounded in $D^{1,\Phi}(\mathbb{R}^N)$.*

Since that the Cerami sequence (u_n) given in (5.10) is bounded in $D^{1,\Phi}(\mathbb{R}^N)$, by Lemma 2.8, we can assume that for some subsequence, there is $u \in D^{1,\Phi}(\mathbb{R}^N)$ such that

$$(5.11) \quad u_n \xrightarrow{*} u \text{ in } D^{1,\Phi}(\mathbb{R}^N), \quad \text{and} \quad u_n(x) \rightarrow u(x) \text{ a.e. } \mathbb{R}^N.$$

Given these limits, we can obtain, as in Lemma 3.2, the inequality

$$(5.12) \quad \int_{\mathbb{R}^N} \Phi(|\nabla u|) dx \leq \liminf_{n \rightarrow \infty} \int_{\mathbb{R}^N} \Phi(|\nabla u_n|) dx.$$

Fix $v \in C_0^\infty(\mathbb{R}^N)$. By boundedness of Cerami sequence (u_n) , we have $J'(u_n)(v - u_n) = o_n(1)$, hence, since Φ is a convex function, it is possible to show that

$$(5.13) \quad \int_{\mathbb{R}^N} \Phi(\nabla v) dx - \int_{\mathbb{R}^N} \Phi(\nabla u_n) dx \geq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))f(u_n(y))(v(y) - u_n(y))}{|x - y|^\lambda} dx dy + o_n(1).$$

For the Lemma 3.2, we have to

$$\liminf_{n \rightarrow \infty} \int_{\mathbb{R}^N} \Phi(|\nabla u_n|) dx \geq \int_{\mathbb{R}^N} \Phi(|\nabla u|) dx.$$

Now, due to the Fatou's Lemma, we have

$$\liminf_{n \rightarrow \infty} \int_{\mathbb{R}^N} V(x)\Phi(|u_n|) dx \geq \int_{\mathbb{R}^N} V(x)\Phi(|u|) dx.$$

Therefore,

$$(5.14) \quad \liminf_{n \rightarrow \infty} \int_{\mathbb{R}^N} \Phi(u_n) \geq \int_{\mathbb{R}^N} \Phi(u).$$

From (5.13) and (5.14) together with the limits (5.7) and (5.8), we get

$$\int_{\mathbb{R}^N} \Phi(|\nabla v|) dx - \int_{\mathbb{R}^N} \Phi(|\nabla u|) dx \geq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))f(u(y))(v(y) - u(y))}{|x - y|^\lambda} dx dy.$$

As $E = \overline{C_0^\infty(\mathbb{R}^N)}^{\|\cdot\|_E}$ and $\Phi \in (\Delta_2)$, we conclude that

$$(5.15) \quad \int_{\mathbb{R}^N} \Phi(|\nabla v|) - \int_{\mathbb{R}^N} \Phi(|\nabla u|) \geq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))f(u(y))(v(y) - u(y))}{|x - y|^\lambda} dx dy, \quad \forall v \in E.$$

In other words, u is a critical point of the J functional. And from Proposition 5.4, we can conclude that u is a weak solution for (P). Now, we substitute $v = u^+ := \max\{0, u(x)\}$ in (5.15) we get

$$- \int_{\mathbb{R}^N} \Phi(|\nabla u^-|) dx \geq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))f(u(y))(u^-(y))}{|x - y|^\lambda} dx dy = 0,$$

which leads to

$$\int_{\mathbb{R}^N} \Phi(|\nabla u^-|) dx = 0$$

whence it is readily inferred that $u^- = 0$, therefore, u is a weak nonnegative solution.

Note that u is nontrivial. In that sense, let us consider a sequence $(\varphi_k) \subset C_0^\infty(\mathbb{R}^N)$ such that $\varphi_k \rightarrow u$ in $D^{1,\Phi}(\mathbb{R}^N)$. Since (u_n) bounded, we get $J'(u_n)(\varphi_k - u_n) = o_n(1)\|\varphi_k\| - o_n(1)$. As Φ is convex, we can show that

$$(5.16) \quad \int_{\mathbb{R}^N} \Phi(|\nabla \varphi_k|) dx - \int_{\mathbb{R}^N} \Phi(|\nabla u_n|) dx - o_n(1) \geq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))f(u_n(y))(\varphi_k(y) - u_n(y))}{|x - y|^\lambda} dx dy.$$

Since $(\|\varphi_k\|)_{k \in \mathbb{N}}$ is a bounded sequence, it follows from (5.16) and from limits (5.7) and (5.8),

$$Q(\varphi_k) - \limsup_{n \rightarrow \infty} Q(u_n) \geq \limsup_{n \rightarrow \infty} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))f(u_n(y))(\varphi_k(y) - u_n(y))}{|x - y|^\lambda} dx dy,$$

for every $k \in \mathbb{N}$. Now, note that being $\Phi \in (\Delta_2)$ and $\varphi_k \rightarrow u$ in $D^{1,\Phi}(\mathbb{R}^N)$, we conclude from the inequality above that

$$(5.17) \quad Q(u) \geq \limsup_{n \rightarrow \infty} Q(u_n).$$

From (5.14) and (5.17),

$$Q(u) = \lim_{n \rightarrow \infty} Q(u_n).$$

By (5.6), we have

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u_n(x))F(u_n(y))}{|x - y|^\lambda} dx dy = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))F(u(y))}{|x - y|^\lambda} dx dy,$$

Therefore,

$$\begin{aligned} 0 < c &= \lim_{n \rightarrow \infty} J(u_n) \\ &= \int_{\mathbb{R}^N} \Phi(|\nabla u|) dx + \int_{\mathbb{R}^N} V(x) \Phi(|u|) dx - \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{K(x)K(y)F(u(x))F(u(y))}{|x-y|^\lambda} dx dy \\ &= J(u), \end{aligned}$$

that is, $u \neq 0$.

Through the assumptions required in Theorem 3.4, it is possible to show an inequality similar to that provided by Lemma 4.1. With that, repeating the same arguments used in the proof of Lemma 4.2, we have to $u \in L_{loc}^\infty(\mathbb{R}^N)$. Therefore, the Theorem 3.4 is finished.

The proof of Theorem 1.6 closely follows the ideas presented in the proof of Theorem 1.2.

6. ACKNOWLEDGMENTS

The authors are grateful to the Paraíba State Research Foundation (FAPESQ), Brazil, and the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Brazil, whose funds partially supported this paper.

REFERENCES

- [1] A. Adams and J. F. Fournier, *Sobolev Spaces*, Academic Press (2003). 5
- [2] A.R.D. Silva and C.O. Alves *Multiplicity and concentration of positive solutions for a class of quasilinear problems through Orlicz-Sobolev space*, J. Math. Phys. 57, 143-162 (2016) 20
- [3] A. Szulkin, *Minimax principle for lower semicontinuous functions and applications to nonlinear boundary value problems*, Ann. Inst. H. Poincaré 3 (1986), 77-109. 2
- [4] C.O. Alves, G.M. Figueiredo, M. Yang, *Existence of solutions for a nonlinear Choquard equation with potential vanishing at infinity*, Adv. Nonlinear Anal. 5 (2016) 1–15. 3
- [5] C.O. Alves and M.A.S. Souto, *Existence of solutions for a class of nonlinear Schrödinger equations with potential vanishing at infinity*, J. Differential Equations 254 (2013) 1977–1991 4
- [6] C. O. Alves and M. L. M. Carvalho, *A Lions Type Result For a Large Class of Orlicz-Sobolev Space and Applications*, Moscow Mathematical Journal, 2021. 2, 15, 30
- [7] C.O. Alves, V. Rădulescu, L.S. Tavares, *Generalized Choquard equations driven by nonhomogeneous operators*, Mediterr J Math. 2019;16.1, 20, 24pp. 2, 15
- [8] D. Montreano, D. Montreano, N.S. Papageorgiou, *Topological and Variational Methods with Applications to Nonlinear Boundary Value Problems*, Springer, New York (2014) 16, 30
- [9] E.D. da Silva, M. L. M. Carvalho, K. Silva and J. V.A. Gonçalves, *Quasilinear elliptic problems on non-reflexive Orlicz-Sobolev spaces*, Topol. Methods Nonlinear Anal. 54 (2019), 587–612 2
- [10] E. DiBenedetto, *$C^{1,\gamma}$ local regularity of weak solutions of degenerate elliptic equations*, Nonlinear Anal. 7, no. 8, (1985) 827-850. 1
- [11] E.H. Lieb, *Existence and uniqueness of the minimizing solution of Choquard's nonlinear equation*, Stud. Appl. Math. 57 (1977) 93–105. 2
- [12] Fukagai, M. Ito and K. Narukawa, *Positive solutions of quasilinear elliptic equations with critical Orlicz-Sobolev nonlinearity on \mathbb{R}^N* , Funkscial. Ekvac. 49 (2006), 235-267. 1, 5, 8
- [13] G. Bonanno, G.M. Bisci and V. Radulescu, *Quasilinear elliptic non-homogeneous Dirichlet problems through Orlicz-Sobolev spaces*, Nonl. Anal. 75 (2012), 4441-4456. 1
- [14] G. Bonanno, G. M. Bisci and V. Radulescu, *Arbitrarily small weak solutions for a nonlinear eigenvalue problem in Orlicz-Sobolev spaces*, Monatshefte für Mathematik 165 (2012), 305-318. 1
- [15] G.M. Figueiredo, *Existence and multiplicity of solutions for a class of p&q elliptic problems with critical exponent*, Math. Nachr. 286, no. 11-12, (2013) 1129-1141. 1
- [16] G.P. Menzala, *On regular solutions of a nonlinear equation of Choquard's type*, Proc. Roy. Soc. Edinburgh Sect. A 86 (1980) 291–301. 2
- [17] H. Berestycki, P.L. Lions, *Nonlinear scalar field equations. I. Existence of a ground state*, Arch. Ration. Mech. Anal. 82 (1983) 313–346. 13, 14
- [18] H. Brezis, *Functional Analysis, Sobolev Spaces and Partial Differential Equations*. Springer, New York (2011) 8

- [19] I.M. Moroz, R. Penrose, P. Tod, *Spherically-symmetric solutions of the Schrödinger–Newton equations*, Classical Quantum Gravity 15 (1998) 2733–2742. [2](#)
- [20] J.P. Gossez, *Nonlinear Elliptic boundary value problems for equations with rapidly(or slowly) increasing coefficients*, Trans. Amer. Math. Soc. 190 (1974), 753–758. [1](#)
- [21] L. da Silva and M. Souto, *Existence of positive solution for a class of quasilinear Schrödinger equations with potential vanishing at infinity on nonreflexive Orlicz-Sobolev spaces*. (In preparation.) [2](#), [3](#), [4](#), [8](#), [15](#)
- [22] Lieb, E., Loss, M., *Analysis, Graduate Studies in Mathematics*. Amer. Math. Soc, Providence (2001) [9](#)
- [23] L. Ma, L. Zhao, *Classification of positive solitary solutions of the nonlinear Choquard equation*, Arch. Ration. Mech. Anal. 195 (2010) 455–467. [2](#)
- [24] M. del Pino, P. Felmer, *Local mountain passes for semilinear elliptic problems in unbounded domains*, Calc. Var. Partial Differential Equations 4 (1996) 121–137. [3](#)
- [25] M. Fuchs and G. Li, *Variational inequalities for energy functionals with nonstandard growth conditions*, Abstr. Appl. Anal. 3 (1998), 405-412. [1](#)
- [26] M. Fuchs and V. Osmolovski, *Variational integrals on Orlicz Sobolev spaces*. Z. Anal. Anwendungen 17, 393-415 (1998) [6](#) [1](#)
- [27] M.N. Rao and Z.D. Ren, *Theory of Orlicz Spaces*, Marcel Dekker, New York (1985). [5](#)
- [28] M. Willem, *Minimax Theorems*, Progress in Nonlinear Differential Equations and Their Applications, vol. 24, Birkhäuser, Boston, MA, 1996. [12](#), [14](#)
- [29] M. Tienari *A degree theory for a class of mappings of monotone type in Orlicz-Sobolev spaces*, Annales Academiae Scientiarum Fennicae. Series A. I, Mathematica. Dissertationes ; 97 [5](#)
- [30] N. Fukagai and K. Narukawa, *On the existence of multiple positive solutions of quasilinear elliptic eigenvalue problems*, Ann. Mat. Pura Appl. 186, no. 3, (2007) 539-564. [1](#)
- [31] N.S. Trudinger, *On Harnack type inequalities and their applicatoin to quasilinear elliptic equations*, Communication on Pure and Applied Mathematics, Vol. XX, 721-747, (1967). [28](#)
- [32] O.A. Ladyzhenskaya and N.N. Ural'tseva, *Linear and quasilinear elliptic equations*, Acad. Press (1968). [25](#), [27](#)
- [33] P. Choquard, J. Stubbe, M. Vuffray, *Stationary solutions of the Schrödinger–Newton model – an ODE approach*, Differ-ential Integral Equations 21 (2008) 665–679. [2](#)
- [34] PH. Clément, M. Garcia-Huidobro, R. Manásevich and K. Schmitt, *Mountain pass type solutions for quasilinear elliptic equations*, Calc. Var. 11 (2000), 33-62. [1](#)
- [35] P.L. Lions, *P.L. Lions, The Choquard equation and related questions*, Nonlinear Anal. 4 (1980) 1063–1072 [2](#)
- [36] P. Tod, I.M. Moroz,, *An analytical approach to the Schrödinger–Newton equations*, Nonlinearity 12 (1999) 201–216. [2](#)
- [37] R. Cerný, *Generalized Moser-Trudinger inequality for unbounded domains and its application* , Nonlinear Differ. Equ. Appl. DOI 10.1007/s00030-011-0143-0. [1](#)
- [38] R. Penrose, *On gravity’s role in quantum state reduction*, Gen. Relativity Gravitation 28 (1996) 581–600. [2](#)
- [39] T. Donaldson, *Nonlinear elliptic boundary value problems in Orlicz-Sobolev spaces*, J. Diff. Equations 10 (1971), 507-528. [1](#)
- [40] S. Chen, S. Yuan, *Ground state solutions for a class of Choquard equations with potential vanishing at infinity*, J. Math. Anal. Appl. 463 (2018) 880–894. [3](#), [4](#)
- [41] V.K. Le and K. Schmitt, *Quasilinear elliptic equations and inequalities with rapidly growing coefficients*, J. London Math. Soc. 62 (2000) 852-872 [1](#)
- [42] S. Pekar, *Untersuchung Über die Elektronentheorie der Kristalle*, Berlin Akademie, 1954. [2](#)
- [43] V. Moroz, J. Van Schaftingen,, *A guide to the Choquard equation*, J. Fixed Point Theory Appl. 19 (2017) 773–813. [3](#)
- [44] V. Moroz, J. Van Schaftingen,, *Groundstates of nonlinear Choquard equations: existence, qualitative properties and decay asymptotics*, J. Funct. Anal. 265 (2013) 153–184. [2](#)
- [45] V. Mustonen and M. Tienari, *An eigenvalue problem for generalized Laplacian in Orlicz-Sobolev spaces*, Proc. R. Soc. Edinburgh, 129A (1999), 153-163. [1](#)
- [46] X. Tang,, *Non-Nehari manifold method for asymptotically periodic Schrödinger equation*, Sci. China Math. 58 (2015) 715–728. [3](#)

(Lucas da Silva)

UNIDADE ACADÊMICA DE MATEMÁTICA
UNIVERSIDADE FEDERAL DE CAMPINA GRANDE,
58429-970, CAMPINA GRANDE - PB - BRAZIL
Email address: ls3@academico.ufpb.br

(Marco Souto)

UNIDADE ACADÊMICA DE MATEMÁTICA
UNIVERSIDADE FEDERAL DE CAMPINA GRANDE,
58429-970, CAMPINA GRANDE - PB - BRAZIL
Email address: marco@dme.ufcg.edu.br