

Efficient Flow-based Approximation Algorithms for Submodular Hypergraph Partitioning via a Generalized Cut-Matching Game

Konstantinos Ameranis
kameranis@uchicago.edu
University of Chicago

Antares Chen
antaresc@uchicago.edu
University of Chicago

Lorenzo Orecchia
orecchia@uchicago.edu
University of Chicago

Erasmus Tani
etani@uchicago.edu
University of Chicago

January 24, 2023

Abstract

In the past 20 years, increasing complexity in real world data has lead to the study of higher-order data models based on partitioning hypergraphs. However, hypergraph partitioning admits multiple formulations as hyperedges can be cut in multiple ways. Building upon a class of hypergraph partitioning problems introduced by Li & Milenkovic [36], we study the problem of minimizing ratio-cut objectives over hypergraphs given by a new class of cut functions, *monotone submodular cut functions* (MSCFs), which captures hypergraph expansion and conductance as special cases.

We first define the ratio-cut improvement problem, a family of local relaxations of the minimum ratio-cut problem. This problem is a natural extension of the Andersen & Lang cut improvement problem [5] to the hypergraph setting. We demonstrate the existence of efficient algorithms for approximately solving this problem. These algorithms run in almost-linear time for the case of hypergraph expansion, and when the hypergraph rank is at most $O(1)$.

Next, we provide an efficient $O(\log n)$ -approximation algorithm for finding the minimum ratio-cut of G . We generalize the cut-matching game framework of Khandekar *et al.* [31] to allow for the cut player to play unbalanced cuts, and matching player to route approximate single-commodity flows. Using this framework, we bootstrap our algorithms for the ratio-cut improvement problem to obtain approximation algorithms for minimum ratio-cut problem for all MSCFs. This also yields the first almost-linear time $O(\log n)$ -approximation algorithms for the case of hypergraph expansion, and constant hypergraph rank.

Finally, we extend a result of Louis & Makarychev [41] to a broader set of objective functions by giving a polynomial time $O(\sqrt{\log n})$ -approximation algorithm for the minimum ratio-cut problem based on rounding ℓ_2^2 -metric embeddings.

1 Introduction

Consider an undirected graph $G = (V, E, \mathbf{w} \in \mathbb{Z}_{>0}^E, \boldsymbol{\mu} \in \mathbb{Z}_{\geq 0}^V)$ with positive edge weights \mathbf{w} and non-negative vertex weights $\boldsymbol{\mu}$. For an edge $\{i, j\} \in E$ and a subset $S \subseteq V$, the edge cut-function $\delta_{ij} : 2^{\{i, j\}} \rightarrow \{0, 1\}$ takes value 0 if i and j are either both in S or both in \bar{S} and value 1 otherwise, i.e. $\delta_{ij}(S \cap \{i, j\}) = |\mathbf{1}_i^S - \mathbf{1}_j^S|$. For $S \subseteq V$, the graph cut function $\delta_G : 2^V \rightarrow \mathbb{Z}$ is then defined as $\delta_G(S) \stackrel{\text{def}}{=} \sum_{e \in E} w_e \delta_e(S \cap e)$ and the ratio-cut objective function $\Psi_G : 2^V \rightarrow \mathbb{Q}$ is

$$\Psi_G(S) \stackrel{\text{def}}{=} \frac{\delta_G(S)}{\min\{\boldsymbol{\mu}(S), \boldsymbol{\mu}(\bar{S})\}}. \quad (1)$$

The ratio-cut problem requires finding the cut $S \subseteq V$ minimizing $\Psi_G(S)$. We denote its optimum value $\Psi_G^* \stackrel{\text{def}}{=} \min_{S \subseteq V} \Psi_G(S)$. The minimum ratio-cut problem is NP-hard as it captures the classical NP-hard problems of minimizing graph expansion ($\boldsymbol{\mu} = \mathbf{1}$) and graph conductance ($\boldsymbol{\mu}$ equal to the weighted degree measure) as special cases [43].

Most algorithmic work in graph partitioning has thus focused on producing efficient and accurate approximation algorithms. These typically fall into two categories. The first category leverages the discrete Cheeger’s inequality [4, 52] relating the conductance of a graph to the spectrum of its Laplacian matrix. In particular, if the graph conductance is ϕ , and the second eigenvalue of the normalized Laplacian is λ_2 , then $\frac{\lambda_2}{2} \leq \phi(G) \leq O(\sqrt{\lambda_2})$. Spectral-based graph partitioning algorithms are known to be efficiently implementable, running in nearly-linear time [53], and are heavily used across a number of practical domains. However, spectral methods also suffer from poor approximation guarantees, particularly in the case where ϕ is small as using the discrete Cheeger inequality to approximate ϕ produces an $O(\sqrt{\phi})$ -approximation.

The second category of algorithms rely on solving metric relaxations of the ratio-cut problem via convex programming and rounding the fractional solutions back to cuts via metric embeddings. Leighton and Rao [34] obtain a polynomial time $O(\log n)$ -approximation by relaxing to general metrics and solving a linear program. The current best known approximation of $O(\sqrt{\log n})$ is due to Arora, Rao and Vazirani [7] using a positive semidefinite relaxation to metrics of negative-type, i.e., ℓ_2^2 -metrics.

Despite enjoying excellent approximation guarantees and applying to all ratio-cut problems, rather than just conductance, metric embedding methods did not a priori find broad practical use because of the high computational complexity of solving the metric relaxation problems, which originally required solving maximum multicommodity flow problems with $\Omega(n^2)$ demands over the instance graph. This situation was significantly ameliorated by Khandekar *et al.* [32], who introduced the cut-matching game framework to reduce the approximate solution of metric relaxations of graph partitioning to a polylogarithmic number of single-commodity maximum flow problems over the instance graph. While Khandekar *et al.* only achieved a $O(\log^2 n)$ approximation ratio, a subsequent series of papers based on the same framework yielded a $O(\log n)$ -approximation in polylogarithmic single-commodity maxflow computations [49] and a $O\left(\sqrt{\frac{\log n}{\epsilon}}\right)$ -approximation¹ in $O(n^\epsilon)$ single-commodity maxflow computations [51]. All these algorithms now run in time equal or close to almost-linear, thanks to recent advances on the computation of exact maximum flows [18].

The cut-matching game has itself become a fundamental primitive in the design of graph algorithms [22, 20, 16, 17, 38] and, in particular, for deterministic algorithms that are crucial in the dynamic setting [21]. However, a number of unresolved issues with the current formulation of the cut-matching game have hindered its wider deployment and application as an algorithmic design framework. We highlight major ones here.

- The standard cut-matching approach only applies to expansion ($\boldsymbol{\mu} = \mathbf{1}$). Saranurak and Wang [50] extend it to conductance by a direct reduction that turns every edge into a vertex. However, no cut-matching game is known for general vertex measures $\boldsymbol{\mu}$.
- Currently, the reduction from the minimum ratio-cut problem to the cut-matching game requires exact single-commodity flow computations over the instance graph. While almost-linear-time algorithms for this task are now available [19], it remains an interesting question, and a promising avenue towards

¹This relies on a more complex framework than the original cut-matching game.

simpler and faster algorithms, to develop a version of the cut-matching game for which approximate flow solvers suffice without compromising the approximation guarantee by additional logarithmic factor, as is the case in [22].

- The cut-matching game only applies to edge-based graph partitioning problems. To tackle vertex-based objectives [22] and mixed-vertex-edge objectives [46], it is necessary to develop algorithms for the hypergraph partitioning problem [36]. While hypergraph partitioning, both from the perspective of spectral techniques [15] and metric relaxations [41], has been well studied, we currently have no fast algorithms for these tasks, as existing results rely on generic semidefinite solvers.

In this paper, we resolve the above issues. We propose a simple generalization of the cut-matching game that supports approximate flow computation, while still achieving a $O(\log n)$ -approximation (Section 5). In the process, we develop a novel perspective on the cut-matching game based on local convex relaxations of the non-convex ratio-cut objective (Section 4). We also start a methodical investigation of fast flow-based algorithms for general submodular hypergraph ratio-cut problems and identify a natural class of hypergraph models that is amenable to this approach (Section 3). As the rest of the paper builds language and theory for constructing algorithms on general submodular hypergraphs, we dedicate the next subsection towards introducing these objects. We then discuss our contributions in detail in Section 1.2.

1.1 Submodular Cut Functions over Hypergraphs

A hypergraph is a tuple $G = (V, E)$, where $E \subseteq 2^V$ is a set of hyperedges. The rank of a hyperedge $h \in E$ is its cardinality $|h|$. The sparsity of a hypergraph G is defined as $\text{sp}(G) = \sum_{h \in E} |h|$. Note that the input size required to specify a hypergraph is $O(|V| + \text{sp}(G))$. If every $h \in E$ satisfies $|h| = k$, the hypergraph G is called k -uniform. In this way, an undirected graph is simply a 2-uniform hypergraph. A key step to formalize a hypergraph cut problem is the definition of a suitable cut function $\delta_h : 2^h \rightarrow [0, 1]$ of a hyperedge $h \in E$. Li and Milenkovic [35, 36] propose the following general definition.

Definition 1 (Cut Function). A function $\delta_h : 2^h \rightarrow [0, 1]$ is a *cut function* if it is symmetric ($\forall S \subseteq h, \delta_h(S) = \delta_h(\bar{S})$), it satisfies $\delta_h(\emptyset) = \delta_h(h) = 0$, and it achieves value 1 ($\exists T \subset h, \delta_h(T) = 1$).

While the canonical edge-cut function $\delta_{ij}(S \cap \{i, j\}) = |\mathbf{1}_i^S - \mathbf{1}_j^S|$ is the unique cut function respecting for rank-2 hyperedges, higher-rank hyperedges support other choices of cut functions. A natural additional requirement is that the cut functions be *submodular*. This sufficiently restricts the intractability of the problem, while preserving some of the crucial analytical and features of cut problems over graphs, such as Cheeger’s inequality [37], connectivity-based decompositions and flow-cut duality, which we strongly exploit in this work.

Indeed, all examples of cut functions in previous work on hypergraph partitioning, both theoretical and applied as far as we are aware, fall in the class of submodular cut functions:

- **Clique-expansion cut function** [1, 57]: $\delta_h(S) = |S| \cdot |h \setminus S| = \delta_{K_h}(S)$, where K_h is the (2-uniform) complete graph over hyperedge h . This version reduces to the standard graph version by replacing each hyperedge h with the complete graph K_h . The same reduction applies when δ_h is the cut function of any 2-uniform graph over h .
- **Star-expansion cut function**[1, 57]: $\delta_h(S) = \min\{|S|, |h \setminus S|\}$. This version reduces to the standard graph version by adding an auxiliary vertex for each hyperedge h , connected to each vertex $i \in h$ by a unit weight edge.
- **Standard hypergraph cut function** [14, 39]: $\delta_h^{\text{cut}}(S) \stackrel{\text{def}}{=} \min\{1, |S|, |h \setminus S|\}$. Analogously to the graph case, this function takes value 1 if the hyperedge is cut and 0 otherwise. This is taken to be the standard notion of hypergraph partitioning in the existing literature. It cannot be expressed as the cut function of any undirected graph, even if partial minimization over auxiliary nodes is allowed.

In the following, we let a *weighted submodular hypergraph* G be a weighted hypergraph $(V, E, \mathbf{w} \in \mathbb{Z}_{>0}^E, \boldsymbol{\mu} \in \mathbb{Z}_{\geq 0}^V)$ with positive edge weights \mathbf{w} and non-negative vertex weights $\boldsymbol{\mu}$, together with a choice of submodular cut functions $\{\delta_h\}_{h \in E}$. In the special case when all cut functions are standard, we refer to G as a *weighted standard hypergraph*. The ratio-cut objective on G is defined analogously to Equation (1), i.e. for all $S \subseteq V$:

$$\Psi_G(S) \stackrel{\text{def}}{=} \frac{\delta_G(S)}{\min\{\mu(S), \mu(\bar{S})\}} = \frac{\sum_{h \in E} w_h \delta_h(S \cap h)}{\min\{\mu(S), \mu(\bar{S})\}}.$$

Analogous to the graph case, we use the terms hypergraph expansion ($\boldsymbol{\mu} = \mathbf{1}$) and hypergraph conductance ($\forall i \in V, \mu_i = \sum_{h \in E: i \in h} w_h$) to refer to specific choice of vertex measures.

Previous Work The standard hypergraph partitioning problem was first studied in the contexts of parallel numerical algorithms [13] and scientific computing [23]. Louis and Makarychev [41] provided the first randomized polynomial-time $O(\sqrt{\log n})$ -approximation algorithm, matching the best known approximation for the graph version of the problem [7] by using the same relaxation to negative-type metrics. In more recent work, Louis [39] and Chan *et al.* [14] developed a spectral approach to the conductance version of the problem, achieving a Cheeger-like approximation guarantee: their algorithms return a partition with ratio-cut at most $O(\sqrt{\Psi_G^* \log \max_{h \in E} |h|})$, which is shown to be tight under the small-set expansion conjecture. Their approach is based on defining a variational quantity analogue to the spectral gap of a graph, whose optimizer, just like the Fiedler vector of the graph, can be rounded to a cut of ratio-cut at most $O(\sqrt{\Psi_G^*})$. Unfortunately, in the hypergraph case, such vector is itself NP-hard to compute [42], requiring a SDP relaxation which must provably lose a factor of $\log \max_{h \in E} |h|$ under the Small-Set-Expansion Conjecture [39].

Much less is known about the approximability of hypergraph ratio-cut objectives for general submodular cut functions. For submodular hypergraph conductance, Li and Milenkovic [36] show that the same SDP approach yields cuts with objective $O(\sqrt{\Psi_G^* \cdot \max_{h \in E} |h|})$. They also conjecture that improving on such linear dependence on hypergraph rank is NP-hard. It is important to note that algorithms for the above settings require solving SDP relaxations over $|V| \times |V|$ symmetric matrices, with at least $|E|$ constraints. No faster algorithms are currently known beyond generic SDP solvers [30], which run in $\Omega(|V|^2|E|)$ time. Even for the simplest form of standard hypergraph partitioning, the resulting SDP relaxation is a mixed packing-and-covering program for which no almost-linear time algorithm is currently known [29].

1.2 Our Contributions

Novel Hypergraph Partitioning Objectives Our first contribution is to define *monotone submodular cut functions*, a subclass of submodular cut functions given by the infimal symmetrizations of non-decreasing submodular functions:

Definition 2 (Monotone Submodular Cut Function). A cut function $\delta_h : 2^h \rightarrow \mathbb{R}_{\geq 0}$ is MSCF if, for all $S \subseteq h$, it can be expressed as

$$\delta_h(S) = \min \{F_h(S), F_h(h \setminus S)\},$$

where $F_h : 2^h \rightarrow \mathbb{R}$ is a non-decreasing submodular function such that $F_h(\emptyset) = 0$.

It is clear that both the star-expansion cut function ($F_h(S) = |S|$) and the standard hypergraph cut function ($F_h(S) = \min\{1, |S|\}$) are monotone submodular cut functions. However, this class extends beyond these two examples. We highlight two additional examples that are relevant in applications:

- F_h is a cardinality-based non-decreasing submodular function [56], i.e., a non-decreasing concave function of the cardinality. This is useful when we wish to modify the star-expansion cut function by diminishing the penalty on balanced partitions of h , e.g., by taking $\delta_h(S) = \min\{|S|^p, |h \setminus S|^p\}$ for $p \in (0, 1)$.
- F_h is the entropy of a subset of random variables associated with vertices of h . When the hypergraph G is the factor graph of an undirected graphical model, this yields a variant of the minimum information partition [27, 44].

We refer to weighted submodular hypergraphs with MSCF cut functions as MSCF-hypergraphs. Section 3 establishes the fundamental properties of MSCFs. The following primal-dual properties are critical to our subsequent results:

1. Primal property [Theorem 5 in Section 3]: the Lovász extension of a MSCF can be approximated within a constant by an *absolute norm*² over $\mathbb{R}^h \perp \mathbf{1}$. The main use of this property will be in rounding ℓ_2^2 -metric relaxations for ratio-cut problems over MSCF-hypergraphs in Section 6.
2. Dual property [Theorems 6 and 7 in Section 3.1]: dual vectors in the base polytopes of MSCFs can be extrapolated to construct *flow embeddings of graphs into hypergraphs*. The main use of this property will be in Section 5, where we establish the proving the viability of the cut-matching approach for constructing approximation algorithms for ratio-cut problems over MSCF-hypergraphs.

These properties allow us to leverage metric and flow arguments to obtain $\text{polylog}(|V|)$ -approximation algorithms for ratio-cut problems over MSCF-hypergraphs. To the best of our knowledge, this paper is the first to draw attention to this class of symmetrized submodular functions. The most common notion of symmetrization in submodular optimization is the standard additive symmetrization ($F_{\text{sym}}(S) \stackrel{\text{def}}{=} F(S) + F(\bar{S}) - F(h)$ for every $S \subseteq h$) [9]. However, the fundamental primal-dual properties above fail for such symmetrization.

Local Relaxations and Cut Improvement Our next contribution is a new perspective on the cut-matching game. We interpret the cut-matching game as a generic approach for minimizing a (non-convex) ratio of submodular functions [10] via a sequence of local convex relaxations. We introduce the *ratio-cut improvement* objective, a local convex objective parametrized by a *seed* \mathbf{s} satisfying $\sum_{i \in V} \mu_i \mathbf{s}_i = 0$ and $\|\mathbf{s}\|_\infty \leq 1$:

$$\Psi_{G,\mathbf{s}}(S) \stackrel{\text{def}}{=} \frac{\delta_G(S)}{\left| \sum_{i \in S} \mu_i \mathbf{s}_i \right|},$$

This problem can be thought of as attempting to find a cut (S, \bar{S}) that simultaneously has both a small boundary, and high correlation with the input seed \mathbf{s} simultaneously. Indeed, Appendix B.1 demonstrates that the problem of minimizing $\Psi_{G,\mathbf{s}}$ is precisely the generalization of the cut-improvement problem of Andersen and Lang [5] to weighted submodular hypergraphs and fractional seeds.

We give algorithms that minimize this objective under different assumptions (see Theorem 8 and Table 8). If G has constant rank, or δ_G is the sum of standard hypergraph cut functions, the algorithm outputs an ϵ -approximate primal-dual solution to the ratio-cut improvement problem in time that is almost-linear with respect to the input size of G (Section 4.3). Furthermore, these results do not require the cut functions to be monotone, and apply to all weighted submodular hypergraphs.

Efficient algorithms for the hypergraph ratio-cut improvement problem are of particular interest as they allow us to locally probe the cut space of the instance hypergraph G and find sparse cuts near the input seed \mathbf{s} . Concurrently, these algorithms, via a dual solution, provide local lower bounds on the cut function δ_G . Crucially, by the dual property of MSCF cut functions, such lower bounds yield flow embeddings of a (2-uniform) demand graph D into the hypergraph G , so that $\delta_D(S) \leq \delta_G(S)$ for all $S \subseteq V$. Consequently, if one wishes to boost the local ratio-cut improvement procedure to produce a global approximation for the global ratio-cut problem, then one need only find an adaptive sequence of T cut-improvement problems such that the cut function of the sum $\sum_{t=1}^T D_t$ of local demand graphs yields a global lower bound on δ_G . This is precisely the problem solved by the cut-matching framework of Khandekar *et al.* [32]

Generalized Cut-Matching Game The classical cut-matching game of Khandekar *et al.* [31] is a two-player game between a cut player \mathcal{C} and a matching player \mathcal{M} . At every iteration, \mathcal{C} outputs a balanced bipartition, and \mathcal{M} outputs a matching across the bipartition. The goal of the cut player is to construct an expander, in as few rounds as possible, using only matchings returned by \mathcal{M} . This cut-matching game was originally intended to guarantee lower bounds to graph expansion. Consequently, the intended matching

²A norm $\|\cdot\|$ over \mathbb{R}^n is absolute if, for all $\mathbf{x} \in \mathbb{R}^n$ we have $\|\mathbf{x}\| = \|\|\mathbf{x}\|\|$. See Definition 3 and related discussion in Section 2.

player \mathcal{M} can be cast as an exact solution to the ratio-cut improvement problem for vertex weighting $\boldsymbol{\mu} = \mathbf{1}$, and seed vector $\mathbf{s} = \mathbf{1}_S - \mathbf{1}_{\bar{S}}$ for S the cut played by \mathcal{C} .

A main contribution of our work is to define a generalized cut-matching game where the matching strategy can be replaced with an approximate solver for the ratio-cut improvement problem. Even in the restricted case where $\boldsymbol{\mu} = \mathbf{1}$, replacing an exact solver with an approximate solver presents insurmountable issues: an ϵ -approximate solver can be modeled as a matching player that is allowed to drop an ϵ -fraction of matching edges (see Definition 10 in Section 5 for more details). Subsequently, the matching player could select a set T with $|T| = \Theta(\epsilon n)$ and always drop matching edges adjacent to T . Even if the cut player can quickly identify such set T , it would have no action that can add edges to it, as it is restricted to play bisections.

For this reason, for approximate matching players, we define in Section 5 a generalized cut-matching game that allows the cut player to play unbalanced cuts. Widening the set of cut actions also solves another immediate difficulty in generalizing the cut-matching game from expansion ($\boldsymbol{\mu} = \mathbf{1}$) to generic ratio-cut objectives: the fact that no $\boldsymbol{\mu}$ -bisection of V may exist when considering $\boldsymbol{\mu}$ -conductance. We demonstrate how to use our generalized cut-matching game to approximate arbitrary ratio-cut objectives.

Theorem 1 (Informal). *Let $V = [n]$. There exists an algorithm `approx-rc` which, on input of a MSCF-hypergraph $G = (V, E_G, \mathbf{w}^G, \boldsymbol{\mu})$, and ratio cut-improvement error parameter $\epsilon > 0$, outputs a $O\left(\frac{\log n}{1-\epsilon}\right)$ -approximation to the minimum ratio-cut. Furthermore `approx-rc` requires at most $O\left(\frac{\log^2 n}{1-\epsilon}\right)$ calls to an ϵ -approximate solver³ for the ratio-cut improvement problem.*

This theorem is given in Section 5 where the exact form of the statement is captured via Theorem 9, then proven in Corollary 1.

To the best of our knowledge, this is the first formal generalization of the cut-matching game to ratio-cut objectives beyond expansion. Previous work [50] had generalized the cut-matching game to conductance via a sophisticated, recursive combinatorial reduction. Our approach shows how to generalize the original analytical argument of Orecchia *et al.* [49].

Restricting to the case of the standard hypergraph cut function, and constant rank MSCFs, we also note that this theorem yields an $O(\log n)$ approximation algorithm whose running time is dominated by a polylogarithmic number of s - t approximate maximum flow computations. Leveraging [12] produces the first almost-linear time algorithm for the minimum ratio-cut problem. If we wish to only route exact s - t maximum flows, then recent advances due to [19] still recovers an almost-linear time algorithm. In general, the generalized cut-matching game provides a fast approximation algorithm for minimum ratio-cut provided a fast approximate ratio-cut improvement algorithm exists.

Our description of the generalized cut-matching game abstracts what is required from the cut player to yield an approximation algorithm. In section 5, we define a so-called $(f(n), g(n, \epsilon))$ -good cut strategy, and prove

Theorem 2. *Given a $(f(n), g(n, \epsilon))$ -good cut strategy, there exists an algorithm which, on input of a MSCF-hypergraph G , and $\epsilon > 0$, outputs an $O\left(\frac{g(n, \epsilon)}{f(n)}\right)$ -approximate minimum ratio-cut to G .*

restated in Theorem 9. Theorem 1, in fact, follows from then demonstrating the following result.

Theorem 3. *There exists a randomized cut strategy \mathcal{C} where, for any $\epsilon \geq 0$, \mathcal{C} is $\left(\Omega(\log n), O\left(\frac{\log^2 n}{1-\epsilon}\right)\right)$ -good with probability $1 - o_n(1)$. Furthermore, \mathcal{C} outputs a cut in time that is nearly-linear in n .*

which is restated in Theorem 10. The proof of this Theorem extracts a robust separation property that allows for the matching strategy to play approximate primal-dual solutions to the ratio-cut improvement problem.

³See Section 4 for an exact definition.

Metric Embedding Relaxations for MSCF cut functions In Section 6, we investigate the polynomial-time approximability of ratio-cut problems over submodular hypergraphs with MSCF cut functions. We leverage the primal property of MSCFs to prove the following theorem.

Theorem 4. *There exists a randomized polynomial-time $O(\sqrt{\log n})$ -approximation algorithm for solving the minimum ratio-cut problem on weighted submodular hypergraphs equipped with monotone submodular cut functions.*

This result is restated in Theorem 13 and proved in Section 6. It extends the $O(\sqrt{\log n})$ -approximation of Louis and Makarychev [41] for standard hypergraph sparsest cut to all ratio-cut problems for the more general class of monotone submodular cut functions.

Our algorithm for Theorem 4 solves an ℓ_2^2 -metric relaxation of the ratio-cut problem for weighted submodular hypergraphs, given by the program **RC-Metric**. This relaxation crucially exploits the geometric connection between a monotone submodular cut function δ_h and the norm induced by the associated non-decreasing submodular function F_h . We note that our techniques can be applied in a straightforward manner to the sparsest cut problem with general demands to yield rank-independent $O(\sqrt{\log k})$ -approximation ratios, where k is the number of demands. This improves on previous work by Louis [40], which achieves a $O\left(\sqrt{\log(k) \cdot \log(\max_{h \in E} |h|)}\right)$ -approximation also via an ℓ_2^2 -metric relaxation, by removing the dependence on the hypergraph rank.

1.3 Additional Remarks and Open Problems

Introducing MSCF helps provide a finer-grained picture of submodular cut functions by highlighting a meaningful subclass where it is possible to obtain approximations in time, and quality, independent of hypergraph rank. It would be interesting to understand whether such approximations exist for cut functions whose Lovász extensions are not closely approximated by an absolute norm. The results in this paper are not immediately generalizable as the flow-decomposition statement strongly depends on this property.

The rounding analysis for our metric relaxation also relies on the monotonicity of absolute norms, but it is uncertain if this is necessary for producing roundable ℓ_2^2 -metric relaxations for ratio-cut problems over hypergraphs in general. Is there a natural, larger, class of submodular hypergraph cut functions for which rounding negative-type metric relaxations produce good approximation algorithms?

It would be interesting to see if techniques developed in this paper can help in formalizing, and extending recent flow-based algorithms for local graph partitioning [55, 28] to the setting of submodular hypergraphs.

Finally, implementations of the cut-matching game [47, 33], and decomposable submodular minimization [24] are readily available. Consequently, the algorithmic framework presented here may be deployable in practice. How effective is this framework when applied to real world test loads?

In general, we hope that this work assists in pursuing a more methodical investigation, both theoretical and practical, of hypergraph partitioning and the rich landscape of submodular cut functions.

2 Preliminaries

Linear Algebra Throughout this paper, we adopt the convention of representing scalars x in unbolded lowercase, vectors \mathbf{x} in boldface lowercase, and matrices \mathbf{X} as boldface uppercase.

Given vectors $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$, we use $\mathbf{x} \circ \mathbf{y}$ to denote the Hadamard (entry-wise) product between \mathbf{x} and \mathbf{y} . We also use $|\mathbf{x}|$ to denote the vector whose entries are the absolute value of the entries in \mathbf{x} . We let $\langle \mathbf{x}, \mathbf{y} \rangle$ denote the standard Euclidean inner product, $\|\cdot\|$ the Euclidean norm, and $\|\cdot\|_p$ the ℓ_p -norm. We define an inner product between $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ weighted by a non-negative vector $\boldsymbol{\mu} \in \mathbb{R}_{\geq 0}^n$ as follows:

$$\langle \mathbf{x}, \mathbf{y} \rangle_{\boldsymbol{\mu}} \stackrel{\text{def}}{=} \sum_{i=1}^n \mu_i \cdot x_i \cdot y_i.$$

The quantity $\|\mathbf{x}\|_{\boldsymbol{\mu}} \stackrel{\text{def}}{=} \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle_{\boldsymbol{\mu}}}$ denotes the norm induced by the inner product $\langle \cdot, \cdot \rangle_{\boldsymbol{\mu}}$, while for any $p > 0$, we use $\|\mathbf{x}\|_{\boldsymbol{\mu}, p}$ to denote the ℓ_p -norm weighted by $\boldsymbol{\mu}$

$$\|\mathbf{x}\|_{\boldsymbol{\mu}, p} \stackrel{\text{def}}{=} \left(\sum_{i=1}^n \mu_i \cdot |x_i|^p \right)^{1/p}.$$

Note consequently that $\|\mathbf{x}\|_{\boldsymbol{\mu}} = \|\mathbf{x}\|_{\boldsymbol{\mu}, 2}$. Given a vector $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$, we write $\mathbf{x} \perp \mathbf{y}$ to indicate the relation: $\langle \mathbf{x}, \mathbf{y} \rangle = 0$. We also define the following notation for the orthogonal complement (in \mathbb{R}^V) of the span of the all-one vector: $\mathbf{1}^\perp \stackrel{\text{def}}{=} \text{span}\{\mathbf{1}\}^\perp$.

If $\mathbf{X}, \mathbf{Y} \in \mathbb{R}^{n \times n}$, then $\langle \mathbf{X}, \mathbf{Y} \rangle = \sum_{i,j=1}^n X_{ij} Y_{ij}$ denotes the Frobenius inner product. We denote by $\mathbf{X} \succeq 0$ the statement that \mathbf{X} is positive semidefinite, and by $\mathbf{X} \preceq \mathbf{Y}$ the relation $0 \preceq \mathbf{Y} - \mathbf{X}$. We use $\text{diag}(\mathbf{x}) \in \mathbb{R}^{n \times n}$ to denote the real matrix with the coordinates of \mathbf{x} placed along its diagonal. We reserve specifically the matrix $\mathbf{M} \stackrel{\text{def}}{=} \text{diag}(\boldsymbol{\mu})$ for the non-negative vector $\boldsymbol{\mu} \in \mathbb{R}_{\geq 0}^n$. We assume that $\min_{i \in V} \left\{ \frac{\mu_i}{\boldsymbol{\mu}(V)} \right\} \geq \frac{1}{\text{poly}(n)}$.

Given a finite set V , a vector $\mathbf{x} \in \mathbb{R}^V$, and a subset $S \subseteq V$, we let $\mathbf{x}(S) = \sum_{v \in S} x_v$. We also denote by $\mathbf{x}^S \in \mathbb{R}^V$, the restriction of \mathbf{x} to coordinates in S :

$$\mathbf{x}^S \stackrel{\text{def}}{=} \sum_{i \in S} x_i \mathbf{e}_i$$

where \mathbf{e}_i is the i -th standard unit vector. In this way, the vector $\mathbf{1}^S$ denotes the $\{0, 1\}$ -indicator of set S .

Absolute and Monotonic Norms We often refer to the following properties for norms over \mathbb{R}^V .

Definition 3. A norm $\|\cdot\|$ over \mathbb{R}^V is *absolute* if $\|\mathbf{x}\| = \|\mathbf{x}\|$ for all $\mathbf{x} \in \mathbb{R}^V$.

Definition 4. A norm $\|\cdot\|$ over \mathbb{R}^V is *monotonic* if, for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^V$, $|\mathbf{x}| \leq |\mathbf{y}|$ implies $\|\mathbf{x}\| \leq \|\mathbf{y}\|$.

We also use the following basic facts about absolute and monotonic norms.

Fact 1. *If a norm $\|\cdot\|$ over \mathbb{R}^V is absolute, then it is monotonic.*

Fact 2. *The dual norm $\|\cdot\|_*$ of an absolute norm $\|\cdot\|$ over \mathbb{R}^V is also absolute.*

Fact 3. *The following stronger form of Hölder's Inequality holds for absolute norms:*

$$\langle |\mathbf{x}|, |\mathbf{y}| \rangle \leq \|\mathbf{x}\|_* \cdot \|\mathbf{y}\|$$

Proofs of these facts can be found in [11]. We remark that absoluteness and monotonicity are not true for general norms, as can be seen by considering the norm induced by an ellipsoid whose axis are not coordinate-parallel.

Submodular Functions A set function $F : 2^V \rightarrow \mathbb{R}$ is submodular if for all $A, B \subseteq V$, we have:

$$F(A \cup B) \leq F(A) + F(B) - F(A \cap B).$$

From the above, one can see that the set of submodular functions is closed under taking conic combinations. Following standard conventions, we always assume that $F(\emptyset) = 0$ unless specified otherwise.

Any submodular set function F can be associated with a convex body $\mathcal{B}(F)$, known as its base polytope, which is defined as follows:

$$\mathcal{B}(F) \stackrel{\text{def}}{=} \{ \mathbf{x} \in \mathbb{R}^V : \langle \mathbf{x}, \mathbf{1} \rangle = F(V) \wedge \langle \mathbf{x}, \mathbf{1}_S \rangle \leq F(S) \quad \forall S \subseteq V \}.$$

Crucially, a submodular set function F can be extended to its convex closure, known as its Lovász extension and denoted by $\bar{F} : \mathbb{R}^V \rightarrow \mathbb{R}$. There are a number of equivalent characterizations of the Lovász extension (Propositions 3.1 and 3.2 in [9]).

Fact 4. For any submodular function $F : 2^V \rightarrow \mathbb{R}$ such that $F(\emptyset) = 0$, $\bar{F} : \mathbb{R}^V \rightarrow \mathbb{R}$ satisfies the following.

1. \bar{F} is the conjugate dual function of the indicator of $\mathcal{B}(F)$, i.e.,

$$\bar{F}(\mathbf{x}) = \max_{\mathbf{y} \in \mathcal{B}(F)} \langle \mathbf{x}, \mathbf{y} \rangle.$$

2. Given a vector $\mathbf{x} \in [0, 1]^V$, let $S_t(\mathbf{x}) = \{i \in V : x_i \geq t\}$. Then

$$\bar{F}(\mathbf{x}) = \mathbb{E}_t [F(S_t(\mathbf{x}))],$$

where t is sampled uniformly from $[0, 1]$.

3. Given $\mathbf{x} \in \mathbb{R}^V$, order the coordinates of \mathbf{x} by $x_1 \geq \dots \geq x_n$ and define $S_i = [i]$. Then

$$\bar{F}(\mathbf{x}) = \sum_{i=1}^n (x_i - x_{i+1}) \cdot F(S_i) = \sum_{i=1}^n x_i \cdot (F(S_i) - F(S_{i-1})).$$

From these facts, one can also see that the Lovász extension of a conic combination of submodular functions is the same conic combination of the Lovász extensions of those functions. That is, given $F = \sum_{i \in I} a_i F_i$ for some index set I , where each F_i is submodular and $a_i \in \mathbb{R}_{>0}$, we have:

$$\bar{F} = \sum_{i \in I} a_i \bar{F}_i.$$

As we will be dealing with cut functions that are symmetric, we make use of the following fact.

Fact 5. For any submodular symmetric function $F : 2^V \rightarrow \mathbb{R}$ such that $F(\emptyset) = 0$, the Lovász extension \bar{F} is an even function, i.e., $\forall \mathbf{x} \in \mathbb{R}^V$, $\bar{F}(\mathbf{x}) = \bar{F}(-\mathbf{x})$.

Monotone Submodular Functions and Associated Norms A submodular function is (monotone) non-decreasing if $F(S) \geq F(T)$ for all $T \subseteq S$. From Fact 4, one can see that the Lovász extension \bar{F} of such a function is also non-decreasing. We often work with a submodular non-decreasing function by considering the following associated norm. Any non-decreasing submodular function $F : 2^V \rightarrow \mathbb{R}$, with $F(\emptyset) = 0$ and $F(\{i\}) > 0$ for all singleton sets i , gives rise to the norm $\|\cdot\|_F$ on \mathbb{R}^V (see Propositions 3.5 and 5.2 in [9]):

$$\|\mathbf{x}\|_F \stackrel{\text{def}}{=} \bar{F}(|\mathbf{x}|)$$

The unit ball of the dual norm to $\|\cdot\|_F$ defines a polyhedron known as the symmetric submodular polyhedron of F and denoted $\mathcal{P}_{\text{sym}}(F)$, so that we can write:

$$\|\mathbf{x}\|_F \stackrel{\text{def}}{=} \max_{\mathbf{y} \in \mathcal{P}_{\text{sym}}(F)} \langle \mathbf{y}, \mathbf{x} \rangle \quad \text{and} \quad \forall \mathbf{y} \in \mathbb{R}^V, \|\mathbf{y}\|_{*,F} = \min\{\lambda \geq 0 : \lambda \mathbf{y} \in \mathcal{P}_{\text{sym}}(F)\}.$$

Like the base polytope, the symmetric submodular polyhedron admits an alternative characterization (Proposition 3.5 in [9]):

$$\mathcal{P}_{\text{sym}}(F) = \left\{ \mathbf{x} \in \mathbb{R}^V : \sum_{i \in S} |x_i| \leq F(S) \quad \forall S \subseteq V \right\}.$$

The following fact is crucial in establishing our main theorems. It is an immediate consequence of the definition of $\|\cdot\|_F$ and of Facts 1 and 2.

Fact 6. The norms $\|\cdot\|_F$ and $\|\cdot\|_{*,F}$ are both absolute and monotonic.




Figure 1: A hypergraph (left) and its factor graph (right). In the factor graph, variable vertices are shown in solid black, while factor vertices are shown as empty circles.

Graphs We represent a weighted graph with a measure placed on its vertices as the tuple $G = (V, E_G, \mathbf{w}_G, \boldsymbol{\mu})$. Here, $\mathbf{w}_G \in \mathbb{R}_{>0}^{E_G}$ is a vector of edge weights, and $\boldsymbol{\mu} \in \mathbb{Z}_{>0}^V$ is a vector of vertex weights. We may omit the subscript G whenever the graph in reference is clear from context.

Given a graph G , we use $\mathbf{L}(G)$ to denote its combinatorial Laplacian:

$$\mathbf{L}(G) \stackrel{\text{def}}{=} \mathbf{D}(G) - \mathbf{A}(G),$$

where the $\mathbf{D}(G)$ is the diagonal matrix of the weighted degrees of the vertices, and $\mathbf{A}(G)$ is the graph adjacency matrix. We denote $\mathcal{L}_\mu(G)$ as its Laplacian matrix normalized by the vertex measure $\boldsymbol{\mu}$:

$$\mathcal{L}_\mu(G) \stackrel{\text{def}}{=} \mathbf{M}^{-1/2} \mathbf{L}(G) \mathbf{M}^{-1/2},$$

where $\mathbf{M} = \text{diag}(\boldsymbol{\mu})$. Observe that, if G is $\boldsymbol{\mu}$ -regular, meaning $\sum_{j:ij \in E_G} w_{ij} = \boldsymbol{\mu}_i$ for every $i \in V$, then $\mathbf{0} \preceq \mathcal{L}_\mu(G) \preceq 2\mathbf{I}_n$. We call a cut $S \subseteq V$ a $\boldsymbol{\mu}$ -bisection if $\mu(S) = \mu(\bar{S})$.

Hypergraph Flows Given a weighted submodular hypergraph $G = (V, E, \mathbf{w} \in \mathbb{Z}_{>0}^E, \boldsymbol{\mu} \in \mathbb{Z}_{>0}^V)$, a *hyperedge flow* over a hyperedge $h \in E$ is a vector $\mathbf{y}_h \in \mathbb{R}^h \perp \mathbf{1}_h$. A *hypergraph flow* $\mathbf{Y} \in \bigoplus \mathbb{R}^h \perp \mathbf{1}_h$ is a direct sum of hyperedge flows $\mathbf{Y} = (\mathbf{y}_h)_{h \in E}$. The demand vector $\text{dem}(\mathbf{Y}) \in \mathbb{R}^V \perp \mathbf{1}$ of a hypergraph flow $\mathbf{Y} = (\mathbf{y}_h)_{h \in E}$ is defined as:

$$\text{dem}_i(\mathbf{Y}) = \sum_{h \in E: h \ni i} (\mathbf{y}_h)_i$$

We use the vector space structure inherited from $\mathbf{Y} \in \bigoplus \mathbb{R}^h \perp \mathbf{1}_h$ to define linear combinations of hypergraph flows $c\mathbf{Y}_1 + \mathbf{Y}_2$ for any $c \in \mathbb{R}$. The demand vector also behaves linearly as $\text{dem}(c\mathbf{Y}_1 + \mathbf{Y}_2) = c \cdot \text{dem}(\mathbf{Y}_1) + \text{dem}(\mathbf{Y}_2)$.

Finally, notice that hypergraph flows over hyperedges of rank 2 are equivalent to graph flows as, for an edge $\{i, j\} \in E$, $(\mathbf{y}_{ij})_i$ represents the flow from i to j , while $(\mathbf{y}_{ij})_j = -(\mathbf{y}_{ij})_i$ is the flow from j to i .

Factor Graphs A hypergraph $G = (V, E_G, \mathbf{w}, \boldsymbol{\mu})$ is naturally associated with its *factor graph*, an unweighted graph $\hat{G} = (V \cup E_G, \hat{E}_G)$ that is bipartite between the *variable* vertices representing vertices $i \in V$, and *factor* vertices representing hyperedges $h \in E_G$. For $i \in V$ and $h \in E_G$, the edge $\{i, h\}$ belongs to \hat{E}_G if and only if $i \in h$.

$$\hat{E}_G = \{\{i, h\} \in V \times E_G : i \in h\}$$

A hypergraph flow \mathbf{Y} over G can now be equivalently interpreted as a graph flow over the factor graph \hat{G} , where the flow from vertex i to factor vertex h over the edge $\{i, h\}$ is given by $(\mathbf{Y}^h)_i$.

3 Monotone Submodular Cut Functions

We start by verifying that any MSCF is indeed symmetric and submodular. A proof of this fact can be found in Appendix A:

Fact 7. *Let V be an arbitrary finite set and $F : 2^V \rightarrow \mathbb{R}$ be a monotone non-decreasing submodular function. If $\delta : 2^V \rightarrow \mathbb{R}$ is the function defined by:*

$$\delta(S) \stackrel{\text{def}}{=} \min\{F(S), F(V \setminus S)\},$$

then δ is symmetric (in the sense of Definition 1) and submodular.

We next establish the primal property of MSCF. This will allow us to approximate the Lovász extension $\bar{\delta}(\mathbf{x})$ by the F -norm of the projection of \mathbf{x} onto $\mathbb{R}^h \perp \mathbf{1}$. The following theorem proves a stronger property stating that the base polytope $\mathcal{B}(\delta_h)$ is closely approximated by the symmetric polytope $\mathcal{P}_{\text{sym}}(F_h)$ of the non-decreasing function F_h inducing δ_h . Its proof can also be found in Appendix A.

Theorem 5. *Suppose $\delta_h(S) \stackrel{\text{def}}{=} \min\{F_h(S), F_h(\bar{S})\}$ where $F_h : 2^h \rightarrow \mathbb{R}$ is a non-decreasing submodular function such that $F(\emptyset) = 0$ and $F(\{i\}) > 0$ for all $i \in V$. We have:*

$$(\mathcal{P}_{\text{sym}}(F_h) \cap \mathbf{1}^\perp) \subseteq \mathcal{B}(\delta_h) \subseteq 2 \cdot (\mathcal{P}_{\text{sym}}(F_h) \cap \mathbf{1}^\perp).$$

In particular, for all $\mathbf{x} \in \mathbb{R}^h$

$$\min_u \|\mathbf{x} - u\mathbf{1}\|_{F_h} \leq \bar{\delta}_h(\mathbf{x}) \leq 2 \min_u \|\mathbf{x} - u\mathbf{1}\|_{F_h}.$$

It is instructive to consider how the previous results are instantiated for the standard hypergraph cut function $\delta_h^{\text{cut}}(S) \stackrel{\text{def}}{=} \min\{F^{\text{cut}}(S), F^{\text{cut}}(\bar{S})\}$ with $F^{\text{cut}}(S) \stackrel{\text{def}}{=} \min\{1, |S|\}$. In this case, the norm $\|\cdot\|_F$ is the ℓ_∞ -norm and the symmetric submodular polyhedron $\mathcal{P}_{\text{sym}}(F^{\text{cut}})$ is the unit ℓ_1 -ball. Hence the relaxation in Theorem 5 is equivalent to:

$$\min_u \|\mathbf{x} - u\mathbf{1}\|_\infty \leq \bar{\delta}_h(\mathbf{x}) \leq 2 \min_u \|\mathbf{x} - u\mathbf{1}\|_\infty.$$

Note that, in this case, the right-hand side is actually tight, as can be seen by letting $\mathbf{x} = \mathbf{1}^S$ and $u = 1/2$ for a non-trivial $S \subset h$.

By comparison, the denominator of the ratio-cut function is given by the cut function $\delta_\mu(S) \stackrel{\text{def}}{=} \min\{\mu(S), \mu(\bar{S})\}$. In this case, the $\mathcal{P}_{\text{sym}}(\mu)$ is given by the unit ℓ_∞ -ball, and the relaxation is given by the ℓ_1 -norm.

$$\min_u \|\mathbf{x} - u\mathbf{1}\|_{\mu,1} \leq \bar{\delta}_V(\mathbf{x}) \leq 2 \min_u \|\mathbf{x} - u\mathbf{1}\|_{\mu,1}$$

In this case, plugging in $\mathbf{x} = \mathbf{1}^S$ shows that the left hand side is tight. Hence, the coefficients in Theorem 5 cannot be improved.

3.1 Flow Embedding of Graphs into Weighted Submodular Hypergraphs

Let us now introduce a mechanism for flow embedding graphs into weighted submodular hypergraphs with MSCF cut function. We will show how these embeddings yield lower bounds on the hypergraph cut functions in terms of the graph cuts of the embedded graph. Analogously to graphs, we also show that any hypergraph flow can be converted into a flow embedding via flow-path decompositions. It is in this step that we crucially make use of the absoluteness of the norms $\|\cdot\|_{F_h}$ associated with a MSCF cut function δ_h .

Hypergraph Flow Embeddings We begin with a notion of congestion for hypergraph flows over MSCF-hypergraphs.

Definition 5. Given a weighted submodular hypergraph $G = (V, E_G, \mathbf{w}^G, \boldsymbol{\mu})$ equipped with a collection of MSCFs $\{\delta_h\}_{h \in E_G}$ and a hypergraph flow $\mathbf{Y} = (\mathbf{y}_h \in \mathbb{R}^h \perp \mathbf{1}_h)_{h \in E}$, the congestion of \mathbf{Y} over a hyperedge $h \in E$ is defined as

$$\text{cong}_h(\mathbf{Y}) \stackrel{\text{def}}{=} \frac{\|\mathbf{y}_h\|_{*, F_h}}{w_h}.$$

The congestion of hypergraph flow \mathbf{Y} over the hypergraph G is:

$$\text{cong}_G \stackrel{\text{def}}{=} \min_{h \in E} \text{cong}_h(\mathbf{Y}).$$

The following definition then captures what is meant by a weighted graph being flow-embeddable into a given weighted submodular hypergraph.

Definition 6 (Hypergraph Flow Embedding). Let $G = (V, E_G, \mathbf{w}^G, \boldsymbol{\mu})$ be a weighted hypergraph equipped with a collection of MSCFs $\{\delta_h\}_{h \in E_G}$. We say a weighted graph $H = (V, E^H, \mathbf{w}^H)$ embeds into G as a flow with congestion $\rho \geq 0$, denoted $H \preceq_\rho G$, if there exist hypergraph flows $\{\mathbf{Y}^e\}_{e \in E^H}$ such that:

1. For all $e = \{i, j\} \in E^H$, the flow \mathbf{Y}^e routes w_e^H units of flow between vertex i and vertex j , i.e.

$$|\text{dem}(\mathbf{Y}^e)| = w_e^H \cdot (\mathbf{1}_i + \mathbf{1}_j).$$

2. For each $h \in E_G$, we have:

$$\left\| \sum_{e \in E^H} |\mathbf{y}_h^e| \right\|_{*, F_h} \leq \rho \cdot w_h^G.$$

There is a slight subtlety in this definition. One may be inclined, by analogy with graphs, to replace the last condition in the definition with the more natural bound on the total congestion $\sum_{e \in E^H} \text{cong}_h(\mathbf{Y}^e) \leq \rho$. However, for hypergraphs of rank larger than 2, this latter quantity can be significantly larger than the former quantity in the definition, preventing the proof of Theorem 7.

We will use this notion of embeddability to certify lower bounds to the ratio-cut objective for general monotone submodular cut functions, and not just standard cut functions. We will crucially make use of this theorem in our development of the cut-matching game for solving ratio-cut problems over weighted submodular hypergraphs.

Theorem 6. Let $G = (V, E_G, \mathbf{w}^G, \boldsymbol{\mu})$ be a weighted hypergraph equipped with a collection of monotone submodular cut functions $\{\delta_h\}_{h \in E_G}$, and $H = (V, E^H, \mathbf{w}^H)$ be a weighted graph. If $H \preceq_\rho G$, then

$$\sum_{\{i, j\} \in E^H} w_{ij}^H \cdot |\mathbf{x}_i - \mathbf{x}_j| \leq \rho \cdot \sum_{h \in E_G} w_h^G \cdot \min_{u \in \mathbb{R}} \|\mathbf{x}_h - u\mathbf{1}\|_{F_h}$$

for all $\mathbf{x} \in \mathbb{R}^V$. In particular, setting $\mathbf{x} = \mathbf{1}^S$ yields a lower bound on the hypergraph cut in terms of the graph cut function of H for any $S \subseteq V$:

$$\delta_H(S) \leq \rho \cdot \delta_G(S).$$

Proof. Consider the hypergraph flows $\{\mathbf{Y}^e\}_{e \in E^H}$ associated with the embedding $H \preceq_\rho G$ and let $\mathbf{u} \in \mathbb{R}_G^E$ be an arbitrary vector over the hyperedges. As each \mathbf{Y}^e routes demand w_e^H between the endpoints of $e = \{u, v\} \in E^H$, we have

$$w_e^H \cdot |\mathbf{x}_u - \mathbf{x}_v| = |\langle \text{dem}(\mathbf{Y}^e), \mathbf{x} \rangle| = \left| \sum_{h \in E_G} \langle \mathbf{y}_h^e, \mathbf{x}_h \rangle \right| = \left| \sum_{h \in E_G} \langle \mathbf{y}_h^e, \mathbf{x}_h - u_h \mathbf{1}_h \rangle \right| \leq \sum_{h \in E_G} \langle |\mathbf{y}_h^e|, |\mathbf{x}_h - u_h \mathbf{1}_h| \rangle,$$

where the last equality relied on the fact that $\mathbf{y}_h^e \in \mathbb{R}^h \perp \mathbf{1}_h$. Summing over all edges in E_h yields:

$$\begin{aligned}
\sum_{e=\{u,v\} \in E_H} w_{uv}^H \cdot |\mathbf{x}_u - \mathbf{x}_v| &\leq \sum_{e \in E_H} \sum_{h \in E_G} \langle |\mathbf{y}_h^e|, |\mathbf{x}_h - u_h \mathbf{1}_h| \rangle \\
&= \sum_{h \in E_G} \left\langle \sum_{e \in E_H} |\mathbf{y}_h^e|, |\mathbf{x}_h - u_h \mathbf{1}_h| \right\rangle \\
&\leq \sum_{h \in E_G} \left\| \sum_{e \in E_H} |\mathbf{y}_h^e| \right\|_{*, F_h} \cdot \|\mathbf{x}_h - u_h \mathbf{1}_h\|_{F_h} \\
&\leq \rho \cdot \sum_{h \in E_G} w_h^G \cdot \|\mathbf{x}_h - u \mathbf{1}\|_{F_h}.
\end{aligned}$$

The first inequality is an application of Fact 3. The second inequality follows from Definition 5. Finally, the first statement in the theorem follows by setting the arbitrary \mathbf{u} to be the required minimizer. The specialization to cuts follows by plugging in $\mathbf{x} = \mathbf{1}^S$ and applying Theorem 5. \square

We now state the main algorithmic result of this section, which allows us to construct hypergraph flow embedding from any single hypergraph flow. Just like in the case of graphs, this result is based on constructing a flow-path decomposition of the original flow, with the additional complication that we need to appeal to the absoluteness of the $\|\cdot\|_{F_h}$ -norms to derive the required congestion bounds. The proof appears in Appendix A.

Theorem 7. *Let $G = (V, E_G, \mathbf{w}^G, \boldsymbol{\mu})$ be a weighted hypergraph equipped with a collection of MSCFs $\{\delta_h\}_{h \in E_G}$. Given a hypergraph flow \mathbf{Y} over G , there is an algorithm that, in time $\tilde{O}(\text{sp}(G))$, computes a weighted graph $H = (V, E_h, \mathbf{w}^H)$ such that $H \preceq_{\text{cong}_G(\mathbf{Y})} G$. Furthermore, the following hold:*

1. *The graph H is bipartite over the partition $(\{i \in V : \text{dem}_i(\mathbf{Y}) \geq 0\}, \{i \in V : \text{dem}_i(\mathbf{Y}) < 0\})$.*
2. *The degree of vertex i in H equals $\text{dem}_i(\mathbf{Y})$.*

4 Local Relaxations of Minimum Ratio-cut Over Hypergraphs

Next, we introduce a family of locally-biased convex relaxations and study how it approximates the minimum ratio-cut problem. Prior to this, we state an equivalent formulation of minimum ratio-cut as a continuous optimization problem. Given a weighted submodular hypergraph $G = (V, E, \mathbf{w}, \boldsymbol{\mu})$, and cut functions $\{\delta_h\}_{h \in E}$, we define (RC-NonCvx) as the optimization problem $\bar{\Psi}_G^*$ over \mathbb{R}^V :

$$\begin{aligned}
\bar{\Psi}_G(\mathbf{x}) &\stackrel{\text{def}}{=} \frac{\bar{\delta}_G(\mathbf{x})}{\bar{\delta}_V(\mathbf{x})} = \frac{\bar{\delta}_G(\mathbf{x})}{\min_u \|\mathbf{x} - u \mathbf{1}\|_{\boldsymbol{\mu}, 1}} \\
\bar{\Psi}_G^* &\stackrel{\text{def}}{=} \min_{\mathbf{x} \in \mathbb{R}^V} \bar{\Psi}_G(\mathbf{x}) \tag{RC-NonCvx}
\end{aligned}$$

The fact that $\bar{\Psi}_G^*$ is equivalent to solving the minimum ratio-cut problem (expressed in the following lemma) is a direct consequence of Fact 4. It is proved in full in Appendix A.

Lemma 1. *Given a weighted hypergraph $G^* = (V, E, \boldsymbol{\mu}, \mathbf{w})$, we have $\Psi_G^* = \bar{\Psi}_G^*$. Furthermore, there exists an algorithm that, given any $\mathbf{x} \in \mathbb{R}^V$, recovers a cut $S \subseteq V$ satisfying*

$$\Psi_G(S) \leq \bar{\Psi}_G(\mathbf{x})$$

in time $O(|V| \log |V| + \text{sp}(G))$.

4.1 The ratio-cut improvement problem

The impetus for studying a family of local convex relaxations for the minimum ratio-cut problem lies in the fact that $\bar{\Psi}_G^*$ is non-convex: the denominator is not concave. Consequently, it is unclear that this problem can be solved to near-optimality in polynomial time.

Our family of relaxations is parameterized by a *seed vector* $\mathbf{s} \in \mathbb{R}^V$. We first define this relaxation, *the ratio-cut improvement problem*, as a combinatorial problem over cuts.

Definition 7 (Ratio-Cut Improvement Problem). Given a weighted submodular hypergraph $G = (V, E, \mathbf{w}, \boldsymbol{\mu})$, cut functions $\{\delta_h\}_{h \in E}$, and a *seed vector* $\mathbf{s} \in \mathbb{R}^V$ with $\langle \mathbf{s}, \mathbf{1} \rangle_{\boldsymbol{\mu}} = 0$ and $\|\mathbf{s}\|_{\infty} \leq 1$, the ratio-cut improvement objective $\Psi_{G,\mathbf{s}}$ is defined for all $S \subseteq V$ as:

$$\Psi_{G,\mathbf{s}}(S) \stackrel{\text{def}}{=} \frac{\delta_G(S)}{|\langle \mathbf{s}, \mathbf{1}^S \rangle_{\boldsymbol{\mu}}|} = \frac{\sum_{h \in E} w_h \cdot \delta_h(S \cap h)}{|\langle \mathbf{s}, \mathbf{1}^S \rangle_{\boldsymbol{\mu}}|}$$

$$\Psi_{G,\mathbf{s}}^* \stackrel{\text{def}}{=} \min_{S \subseteq V} \Psi_{G,\mathbf{s}}(S)$$

The ratio-cut improvement problem asks to find $S \subseteq V$ achieving $\Psi_{G,\mathbf{s}}^* = \Psi_{G,\mathbf{s}}(S)$.

The combinatorial formulation is equivalent to a continuous, convex optimization problem where the non-concave denominator of $\bar{\Psi}_G^*$ is replaced with the correlation $|\langle \mathbf{s}, \mathbf{x} \rangle_{\boldsymbol{\mu}}|$. We define $\bar{\Psi}_{G,\mathbf{s}}^*$ to be this objective:

$$\bar{\Psi}_{G,\mathbf{s}}(\mathbf{x}) \stackrel{\text{def}}{=} \frac{\bar{\delta}_G(\mathbf{x})}{|\langle \mathbf{s}, \mathbf{x} \rangle_{\boldsymbol{\mu}}|}, \quad \text{and} \quad \bar{\Psi}_{G,\mathbf{s}}^* \stackrel{\text{def}}{=} \min_{\mathbf{x} \in \mathbb{R}^V} \bar{\Psi}_{G,\mathbf{s}}(\mathbf{x}).$$

The fact that the ratio-cut improvement problem $\Psi_{G,\mathbf{s}}^*$ is a relaxation of the minimum ratio-cut problem over hypergraphs Ψ_G^* follows immediately from observing $\bar{\Psi}_G^* \leq \bar{\Psi}_{G,\mathbf{s}}^*$ when the seed satisfies conditions derived from duality of conjugate norms.

Lemma 2. For all $\mathbf{x} \in \mathbb{R}^V$, and seeds $\mathbf{s} \in \mathbb{R}^V$ satisfying $\|\mathbf{s}\|_{\infty} \leq 1$ and $\langle \mathbf{s}, \mathbf{1} \rangle_{\boldsymbol{\mu}} = 0$, we have:

$$\bar{\Psi}_G(\mathbf{x}) \leq \bar{\Psi}_{G,\mathbf{s}}(\mathbf{x}).$$

In particular, setting $\mathbf{x} = \mathbf{1}^S$ yields $\Psi_{G,\mathbf{s}}^*$ is a relaxation of Ψ_G^* for any $S \subseteq V$:

$$\Psi_G(S) \leq \Psi_{G,\mathbf{s}}(S).$$

Proof. By the duality of conjugate norms, we have

$$\min_{u \in \mathbb{R}} \|\mathbf{x} - u\mathbf{1}\|_{1,\boldsymbol{\mu}} = \min_{u \in \mathbb{R}} \max_{\|\mathbf{y}\|_{\infty} \leq 1} \langle \mathbf{y}, \mathbf{x} - u\mathbf{1} \rangle_{\boldsymbol{\mu}} = \max_{\substack{\|\mathbf{y}\|_{\infty} \leq 1 \\ \langle \mathbf{y}, \mathbf{1} \rangle_{\boldsymbol{\mu}} = 0}} \langle \mathbf{y}, \mathbf{x} \rangle_{\boldsymbol{\mu}} \geq |\langle \mathbf{s}, \mathbf{x} \rangle_{\boldsymbol{\mu}}|$$

for all $\mathbf{x} \in \mathbb{R}^V$. Consequently,

$$\bar{\Psi}_G(\mathbf{x}) = \frac{\sum_{h \in E} w_h \bar{\delta}_h(\mathbf{x})}{\min_{u \in \mathbb{R}} \|\mathbf{x} - u\mathbf{1}\|_{1,\boldsymbol{\mu}}} \leq \frac{\sum_{h \in E} w_h \bar{\delta}_h(\mathbf{x})}{|\langle \mathbf{s}, \mathbf{x} \rangle_{\boldsymbol{\mu}}|} = \bar{\Psi}_{G,\mathbf{s}}(\mathbf{x})$$

as required. \square

Typically, we will be interested in the case where the seed vector \mathbf{s} is given by

$$\mathbf{1}^{A,\bar{A}} \stackrel{\text{def}}{=} \mathbf{1}^A - \frac{\mu(A)}{\mu(\bar{A})} \cdot \mathbf{1}^{\bar{A}}. \quad (3)$$

for a cut $A \subseteq V$. This choice of seed vector certainly satisfies $\langle \mathbf{s}, \mathbf{1} \rangle_{\boldsymbol{\mu}} = 0$ and $\|\mathbf{s}\|_{\infty} \leq 1$. Using this seed vector for the ratio-cut improvement problem also immediately recovers the cut improvement problem of Andersen and Lang [5] as discussed in Appendix B.1.

Of critical, algorithmic, consequence is the fact that computing the value of $\bar{\Psi}_{G,\mathbf{s}}^*$ can be written as a convex program. We refer to this program as **(CI-Primal)**:

$$\begin{aligned} \min_{\mathbf{x} \in \mathbb{R}^V} \quad & \sum_{h \in E} w_h \cdot \bar{\delta}_h(\mathbf{x}) \\ \text{s.t.} \quad & \langle \mathbf{s}, \mathbf{x} \rangle_\mu = 1 \end{aligned} \tag{CI-Primal}$$

The following lemma is a simple consequence of the submodularity and symmetry of δ_G .

Lemma 3. *The optimum value of (CI-Primal) equals $\Psi_{G,\mathbf{s}}^*$.*

With this convex program, one can leverage convex duality to construct lower bounds for $\bar{\Psi}_{G,\mathbf{s}}^*$, and consequently $\Psi_{G,\mathbf{s}}^*$. We consider a dual formulation of **(CI-Primal)** as a problem over hypergraph flows:

$$\begin{aligned} \max_{\substack{\alpha \in \mathbb{R}, \\ \{\mathbf{Y} = \{\mathbf{y}_h\}_{h \in E} \in \bigoplus_{h \in E} \mathbb{R}^h\}}} \quad & \alpha \\ \text{s.t.} \quad & \text{dem}(\mathbf{Y}) = \alpha \cdot (\boldsymbol{\mu} \circ \mathbf{s}) \\ & \mathbf{y}_h \in w_h \cdot \mathcal{B}(\delta_h) \quad \forall h \in E \end{aligned} \tag{CI-Dual}$$

In this setting, strong duality between **(CI-Primal)** and **(CI-Dual)** holds by virtue of the **(CI-Primal)** having a finite number of polyhedral constraints. The full proof of the following lemma can be found in Appendix B.2.

Lemma 4. *The programs (CI-Primal) and (CI-Dual) are dual to each other. Strong duality holds.*

4.2 From Primal-Dual Solutions to Hypergraph Flow Embeddings

One of our main contributions will be to apply a generalized cut-matching game to combine local lower bounds into global lower bounds for the non-convex problem $\bar{\Psi}_G^*$. Our local certificate is computed using an approximate solutions to **(CI-Primal)** and **(CI-Dual)**. Let us define an ϵ -approximate primal-dual solution as a cut $A \subseteq V$ paired with a hypergraph flow $\mathbf{Y} \in \bigoplus_{h \in E} w_h \cdot \mathcal{B}(\delta_h)$ that approximately satisfies the demand constraint in **(CI-Dual)** up to multiplicative ℓ_1 -error.

Definition 8 (ϵ -approximate primal-dual solution). Let $G = (V, E_G, \mathbf{w}^G, \boldsymbol{\mu})$ be a MSCF-hypergraph equipped with cut functions $\{\delta_h\}_{h \in E}$, and $\mathbf{s} \subseteq V$ a seed vector satisfying $\mathbf{s} \in \mathbb{R}^V$ with $\langle \mathbf{s}, \mathbf{1} \rangle_\mu = 0$ and $\|\mathbf{s}\|_\infty \leq 1$. An ϵ -approximate primal-dual solution for (G, \mathbf{s}) is a pair

$$\left(A \subseteq V, \mathbf{Y} \in \bigoplus_{h \in H} (\mathbb{R}^h \perp \mathbf{1}_h) \right)$$

such that the following hold simultaneously.

1. $\mathbf{y}_h \in w_h \cdot \mathcal{B}(\delta_h)$ for all $h \in E_G$,
2. $\|\text{dem}(\mathbf{Y}) - \Psi_{G,\mathbf{s}}^* \cdot (\boldsymbol{\mu} \circ \mathbf{s})\|_1 \leq \frac{\epsilon}{8} \cdot \|\Psi_{G,\mathbf{s}}^* \cdot (\boldsymbol{\mu} \circ \mathbf{s})\|_1$,
3. $\text{dem}(\mathbf{Y}) \leq \Psi_{G,\mathbf{s}}^* \cdot (\boldsymbol{\mu} \circ \mathbf{s})$ in an entry-wise sense,
4. $\Psi_{G,\mathbf{s}}(A) \leq 2 \cdot \Psi_{G,\mathbf{s}}^*$,
5. $\text{dem}(\mathbf{Y})_i \cdot \mathbf{s}_i \geq 0$ for every $i \in V$.

The local certificate then takes the form of a demand graph given by running a flow-path decomposition on the hypergraph flow in an ϵ -approximate primal-dual solutions yields a hypergraph flow embedding. The fact that the demand graph certifies a lower bound to the minimum ratio-cut objective Ψ_G^* is a direct consequence of the algorithmic result in Section 3, Theorem 7.

Lemma 5. *Given an ϵ -approximate primal-dual solution (A, \mathbf{Y}) to the ratio-cut improvement problem on (G, \mathbf{s}) , let H be the graph obtained by applying the flow-path decomposition algorithm of Theorem 7 to the hypergraph embedding \mathbf{Y} . Then*

$$H \preceq_2 G$$

Furthermore, graph H is bipartite over the partition $(\{i \in V : \mathbf{s}_i(\mathbf{Y}) \geq 0\}, \{i \in V : \mathbf{s}_i(\mathbf{Y}) < 0\})$, and the degree of vertex i in H equals $\text{dem}_i(\mathbf{Y})$.

Proof. The statements on the bipartiteness and degree distribution of H follow directly from by Theorem 7. By the same theorem, we also have $H_t \preceq_{\text{cong}_G(\mathbf{Y}^t)} G$ for each t , so it only remains to bound $\text{cong}_G(\mathbf{Y}^t)$.

Since \mathbf{Y}^t is part of an ϵ -approximate primal-dual definition output, Definition 8 and Theorem 5 imply that, for all $h \in E_G$,

$$\mathbf{y}_h^t \in w_h \cdot \mathcal{B}(\delta_h) \subseteq 2w_h \cdot (\mathcal{P}_{\text{sym}}(F_h) \cap \mathbf{1}^\perp).$$

Because $\mathcal{P}_{\text{sym}}(F_h)$ is the unit-norm polytope for the dual norm $\|\cdot\|_{*,F_h}$, this yields, for all $h \in E_G$:

$$\text{cong}_h(\mathbf{Y}^t) = \frac{\|\mathbf{y}_h^t\|_{*,F_h}}{w_h} \leq 2$$

as required. □

4.3 Approximation algorithms for the ratio-cut improvement problem

Let us now discuss algorithms for approximating (CI-Primal) and (CI-Dual). For convenience, we introduce the following oracle to abstract algorithmic details in the rest of the paper.

Definition 9. An approximate primal-dual oracle for the ratio-cut improvement problem is an algorithm \mathcal{A}_{CI} which outputs an ϵ -approximate primal-dual solution when given a MSCF-hypergraph G equipped with $\{\delta_h\}_{h \in E}$, a seed $\mathbf{s} \in \mathbb{R}^V$ satisfying $\langle \mathbf{s}, \mathbf{1} \rangle_\mu = 0$ and $\|\mathbf{s}\|_\infty \leq 1$, and an $\epsilon > 0$.

We can now state our main theorem on efficient approximation algorithms for the ratio-cut improvement problem on weighted submodular hypergraphs. The running time of approximating ratio-cut improvement depends on what assumptions are made regarding the hypergraph G , and the cut functions $\{\delta_h\}_{h \in E}$. The running time also leverages different known results on computing approximate solutions to maximum flow and decomposable submodular minimization. Its proof is given in Appendix B.3.

Theorem 8. *Given a MSCF-hypergraph $G = (V, E, \mathbf{w}, \boldsymbol{\mu})$ with $\{\delta_h\}_{h \in E}$, let $T_\epsilon(G, \{\delta_h\}_{h \in E})$ be the time required by an approximate primal-dual oracle \mathcal{A}_{CI} to output an ϵ -approximate primal-dual solution for seed \mathbf{s} . For each set of assumptions in the table below, there exists \mathcal{A}_{CI} with the stated bounds on $T_\epsilon(G, \{\delta_h\}_{h \in E})$.*

Assumptions on G and $\{\delta_h\}_{h \in E}$	Upperbound on $T_\epsilon(G, \{\delta_h\}_{h \in E})$	Relevant Results
$\forall h \in E, \delta_h = \delta_h^{\text{cut}}$	$\tilde{O}\left(\left(V + \text{sp}(G)^{1+o(1)}\right) \cdot \log\left(\max_{h \in E} w_h\right)\right)$	Appendix B.3 and [12]
$\forall h \in E, h = O(1)$	$\tilde{O}\left(\sum_{h \in E} T_h + T_{\text{flow}}(n, n + E)\right)$	Axiotis <i>et al.</i> [8]
General G and $\{\delta_h\}_{h \in E}$	$\tilde{O}\left(V \cdot \sum_{h \in E} \Theta_h \cdot \log\left(V \cdot \max_{h \in E} w_h\right)\right)$	Ene <i>et al.</i> [24]

Here, T_h is the running time of a linear optimization oracle for δ_h , Θ_h is the running time for a quadratic optimization oracle for δ_h , and $T_{\text{flow}}(n, m)$ is the running time required to solve a maximum flow problem on an undirected graph with n vertices and m edges.

5 An $O(\log n)$ -Approximation via Generalized Cut-Matching Games

We now describe a generalized version of the cut-matching game, and discuss how it can be used to design an approximation algorithm for the minimum ratio-cut problem.

A key aspect of our cut-matching game is the possibility for the cut player to play unbalanced cuts. This is really a consequence of wanting the matching player to be implementable by an approximate solver for the ratio-cut improvement problem. Let us formally define a valid action from such a matching player as follows.

Definition 10 (ϵ -approximate matching action). Given $\epsilon > 0$, an ϵ -approximate matching action with respect to a cut $S \subseteq V$ is a weighted bipartite graph $D = (S \cup \bar{S}, E_D, \mathbf{w}^D, \boldsymbol{\mu})$ such that the following hold:

1. $\|\mathbf{d} - \boldsymbol{\mu} \circ |\mathbf{s}|\|_1 \leq \epsilon \cdot \|\boldsymbol{\mu} \circ |\mathbf{s}|\|_1$,
2. $d_i \leq (\boldsymbol{\mu} \circ |\mathbf{s}|)_i$ for each $i \in V$,

where $\mathbf{d} \in \mathbb{R}^V$ is the vector of vertex degrees in D , i.e. $\mathbf{d}_i = \deg_D(i)$ and $\mathbf{s} = \mathbf{1}^{S, \bar{S}}$ as defined in equation 3.

Notice that Lemma 5 demonstrates such a graph D is immediately obtained via the demand graph of a hypergraph flow given by an approximate primal-dual solution to the ratio-cut improvement problem on the seed $\mathbf{1}^{S, \bar{S}}$. Since D certifies a local lower bound to the ratio-cut objective, it is natural to use the cut-matching games framework compose different certificates D . The hope is that this will produce a certificate that globally lower bounds the ratio-cut objective with a quantity depending only on the number of vertices.

Now, applying the standard cut-matching game is insufficient as an approximate solution to the cut-improvement problem may produce D that is not a perfect matching. In fact, D may even fail to contain a matching between a constant fraction of vertex pairs, and thus recursive application of the standard cut-matching game may not suffice for fast algorithms either. We amend the standard cut-matching games framework to allow for the cut player to play potentially unbalanced cuts.

Definition 11 (Generalized cut-matching game). Let $n > 0$, $\epsilon > 0$, and $\boldsymbol{\mu} \in \mathbb{R}_{\geq 0}^V$ be a weighting over a set of vertices V such that $|V| = n$. An $(n, \epsilon, \boldsymbol{\mu})$ -generalized cut-matching game is a two-player game between a cut strategy \mathcal{C} , and a matching strategy \mathcal{M} that proceeds as follows. In the t -th round of the game:

1. \mathcal{C} plays a cut $S_t \subset V$ such that $S_t \neq \emptyset$, and $\boldsymbol{\mu}(S_t) \leq \frac{1}{2} \cdot \boldsymbol{\mu}(V)$, and
2. \mathcal{M} responds with an ϵ -approximate matching action D_t .

Note that this definition also allows for the specification of an arbitrary vertex weighting $\boldsymbol{\mu}$. Consequently, algorithms built on this notion of the cut-matching game can approximate not only expansion, but arbitrary ratio-cut objectives, such as $\boldsymbol{\mu}$ -conductance, without the need for recursion.

Even though the cut player is given more power in the generalized cut-matching game, it must also produce S with stronger structural guarantees in order for the cut-matching game to produce an approximation algorithm for the minimum ratio-cut problem. The next definition formalizes what is required of the cut strategy so that one may leverage this cut-matching game to approximate minimum ratio-cut.

Definition 12 ($(f(n), g(n, \epsilon))$ -good cut strategy). Let $f(n)$ and $g(n, \epsilon)$ be non-decreasing functions, and suppose \mathcal{C} plays the sequence of cuts $S_1, \dots, S_{g(n, \epsilon)} \subseteq V$. We say that \mathcal{C} is $(f(n), g(n, \epsilon))$ -good if, for any sequence weighted graphs $D_1, \dots, D_{g(n, \epsilon)}$ such that each D_t is an ϵ -approximate matching action with respect to S_t , the graph $H = \sum_{t=1}^{g(n, \epsilon)} D_t$ (taken in an edge-wise manner) satisfies

$$\Psi_H \geq f(n).$$

Using this requirement, we can now state the first main result concerning cut-matching games.

Theorem 9. *Let $f(n)$ and $g(n, \epsilon)$ be non-decreasing functions. Given an $(f(n), g(n, \epsilon))$ -good cut strategy, there exists an algorithm **approx-rc** which, on input of a MSCF-hypergraph $G = (V, E_G, \mathbf{w}, \boldsymbol{\mu})$, and $\epsilon > 0$, outputs a cut $A \subseteq V$ that is an $O\left(\frac{g(n, \epsilon)}{f(n)}\right)$ -approximate minimum ratio-cut:*

$$\Psi_G^* \leq \Psi_G(A) \leq O\left(\frac{g(n, \epsilon)}{f(n)}\right) \cdot \Psi_G^*.$$

We prove this theorem in Section 5.1. We then show a good cut strategy exists. Doing so illuminates what additional structure is required from the cut player in the generalized cut-matching game, and yields

Theorem 10. *There exists a randomized cut strategy \mathcal{C} where, for any $\epsilon > 0$, \mathcal{C} is $(\Omega(\log n), O(\frac{\log^2 n}{1-\epsilon}))$ -good with probability $1 - o_n(1)$. Furthermore, \mathcal{C} outputs a cut in time that is nearly-linear in n .*

Its proof is given in Section 5.2. Finally, taking the two aforementioned results in conjunction demonstrates our algorithmic guarantee for the minimum ratio-cut problem.

Corollary 1. *Given an approximate primal-dual oracle \mathcal{A}_{CI} for the ratio-cut improvement problem, there exists an algorithm \mathcal{A} which, on input of a MSCF-hypergraph $G = (V, E_G, \mathbf{w}, \boldsymbol{\mu})$, and $\epsilon > 0$, outputs a $O(\frac{\log n}{1-\epsilon})$ -approximate minimum ratio-cut. Furthermore, \mathcal{A} requires at most $O(\frac{\log^2 n}{1-\epsilon})$ calls to \mathcal{A}_{CI} .*

Proof. Theorem 10 yields a cut strategy that is $(f(n), g(n, \epsilon))$ -good cut strategy where $O\left(\frac{g(n, \epsilon)}{f(n)}\right) = O\left(\frac{\log n}{1-\epsilon}\right)$. Applying Theorem 9 then gives the desired approximation ratio. For the running time, note \mathcal{A} requires only $g(n, \epsilon) = O(\frac{\log^2 n}{1-\epsilon})$ iterations. \square

In light of the results of Theorem 8, the above corollary gives an algorithm to solve the minimum ratio-cut problem over any weighted hypergraph with cut function $\delta_h = \delta_h^{\text{cut}}$ for every $h \in E$ in time:

$$\tilde{O}\left(\left(|V| + \text{sp}(G)^{1+o(1)}\right) \cdot \log \max_{h \in E} w_h\right),$$

and similarly, the running times appearing in the table in Theorem 8 yield equivalent (up to constant factors) running times for approximation algorithm for the minimum ratio-cut problem in their respective settings.

5.1 Proof of Theorem 9: the cut-matching game reduction

We now prove that, given a $(f(n), g(n, \epsilon))$ -good cut strategy, there exists an algorithm that outputs an $O\left(\frac{g(n, \epsilon)}{f(n)}\right)$ -approximation for minimum ratio-cut. The reduction is given by the algorithm **approx-rc** which is summarized in Algorithm 1 below.

Proof of Theorem 9. Consider the cut A_{t^*} returned by **approx-rc**, and let $\alpha_t = \Psi_{G, \text{si}}(A_t)$ be the ratio-cut improvement objective of A_t returned by \mathcal{A}_{CI} for each $t = 1, \dots, g(n)$. The lower bound $\Psi_G(A_{t^*}) \geq \Psi_G^*$ follows by Ψ_G^* being a minimum. For the upper bound, we obtain from Lemma 5, the following guarantee:

$$D_t \preceq_{\alpha_t} \frac{2}{\alpha_t} G.$$

If we let $H = \sum_{t=1}^{g(n)} D_t$, then H must satisfy $H \preceq_{\rho} G$ for $\rho = \sum_{t=1}^{g(n)} \frac{2}{\alpha_t}$. Theorem 6 then implies:

$$\Psi_H^* \leq \rho \cdot \Psi_G^*.$$

However, \mathcal{C} is $(f(n, \epsilon), g(n))$ -good and hence

$$f(n, \epsilon) \leq \Psi_H^* \leq \rho \cdot \Psi_G^* \iff \frac{1}{\rho} \leq \frac{1}{f(n, \epsilon)} \cdot \Psi_G^*$$

Algorithm 1 approx-rc

Parameters: $G = (V, E_G, \mathbf{w}^G, \boldsymbol{\mu})$, $\epsilon > 0$, \mathcal{A}_{CI} an approximate primal-dual cut-improvement oracle, and \mathcal{C} an $(f(n, \epsilon), g(n))$ -good cut strategy.

Initialize: $D_0 = (V, \emptyset, \mathbf{0})$ the empty graph.

For: $t = 1, \dots, g(n)$, do the following:

1. Run $\mathcal{C}(G, D_0, \dots, D_{t-1})$ to receive cut $S_t \subseteq V$.
2. Set the seed vector $\mathbf{s}_t = \boldsymbol{\mu}(S_t) \cdot \mathbf{1}^{S_t} - \frac{\boldsymbol{\mu}(S_t)}{\boldsymbol{\mu}(\overline{S_t})} \cdot \mathbf{1}^{\overline{S_t}}$.
3. Compute $\mathcal{A}_{\text{CI}}(G, \mathbf{s}_t, \epsilon)$ from Theorem 8 to receive an ϵ -approximate primal-dual solution

$$\left(A_t \subseteq V, \mathbf{Y}^t \in \bigoplus_{h \in E_G} (\mathbb{R}^h \perp \mathbf{1}_h) \right).$$

4. Apply the flow-path decomposition algorithm of Theorem 7 on \mathbf{Y}^t to obtain a graph H_t .
5. Denote $\alpha_t = \Psi_{G, \mathbf{s}_t}(A_t)$, and compute the graph $D_t = \frac{1}{\alpha_t} \cdot H_t$

Output: A_{t^*} for $t^* = \operatorname{argmin}_{t=1, \dots, T} \Psi_{G, \mathbf{s}_t}(A_t)$.

Focusing on ρ , we can upper bound an average by its largest element

$$\rho = g(n) \cdot \frac{1}{g(n)} \cdot \sum_{t=1}^{g(n)} \frac{2}{\alpha_t} \leq g(n) \cdot \max_{t=1, \dots, g(n)} \left\{ \frac{2}{\alpha_t} \right\} = \frac{2 \cdot g(n)}{\alpha_{t^*}} = \frac{2 \cdot g(n)}{\Psi_{G, \mathbf{s}_{t^*}}(A_{t^*})},$$

and thus conclude

$$\Psi_{G, \mathbf{s}_{t^*}}(A_{t^*}) \leq 2 \cdot \frac{g(n)}{f(n, \epsilon)} \cdot \Psi_G^*.$$

Finally, Lemma 2 implies that the ratio-cut improvement problem is a relaxation of the minimum ratio-cut problem for any seed $\mathbf{s} \in \mathbb{R}^V$ satisfying $\langle \mathbf{s}, \mathbf{1} \rangle \boldsymbol{\mu} = 0$, and $\|\mathbf{s}\|_\infty \leq 1$. Hence,

$$\Psi_G(A_{t^*}) \leq \Psi_{G, \mathbf{s}_{t^*}}(A_{t^*}) \leq O\left(\frac{g(n)}{f(n, \epsilon)}\right) \cdot \Psi_G^*$$

as required. □

5.2 Proof of Theorem 10: an $\left(\Omega(\log n), O\left(\frac{\log^2 n}{1-\epsilon}\right)\right)$ -good cut strategy

We now turn our attention towards constructing a good cut strategy. In the standard cut-matching game, the cut player seeks to construct a certificate graph that has large expansion in as few rounds as possible. To do this, cut strategies typically maintain a vector embedding of the vertices in the current certificate graph. The cut strategy then plays a cut rounded from the embedding using spectral methods. Our cut strategy adopts an analogous approach, and we further discuss how the embedding is updated in Section 5.2.2.

The novelty in our construction lies in extracting a robust separation property which allows for updates to the certificate to be given by the demand graph of a flow produced by approximately solving a ratio-cut improvement problem. We discuss this property in detail in Section 5.2.3. The final construction of the cut strategy is given in Section 5.2.4. Before proceeding, we introduce some notation and facts that will be relevant when analyzing a vector embedding.

5.2.1 Facts about vector embeddings

We denote a *vector embedding* of a set of vertices $V = [n]$ into dimension $d > 0$ as the set $\{\mathbf{v}_i\}_{i \in V} \subseteq \mathbb{R}^d$. When provided a vertex weighting $\boldsymbol{\mu} \in \mathbb{R}_{\geq 0}^V$, one can consider the *expected embedding vector* $\mathbf{v}_{\text{avg}} \stackrel{\text{def}}{=} \sum_{i \in V} \frac{\mu_i}{\boldsymbol{\mu}(V)} \cdot \mathbf{v}_i$, and the *variance of the embedding* under when i is randomly sampled proportionally to $\boldsymbol{\mu}$:

$$\text{Var}_{\boldsymbol{\mu}}(\mathbf{v}_i) \stackrel{\text{def}}{=} \sum_{i \in V} \frac{\mu_i}{\boldsymbol{\mu}(V)} \cdot \|\mathbf{v}_i - \mathbf{v}_{\text{avg}}\|_2^2.$$

When $\mathbf{v}_{\text{avg}} = \mathbf{0}$, the variance is just the expectation of ℓ_2^2 -lengths $\|\mathbf{v}_i\|_2^2$. In such a case, we use the following fact to lower bound the average ℓ_2^2 -lengths when i is restricted to be contained in some set. We remark that [48] writes this in terms of trace inner products with the Gram matrix associated with $\{\mathbf{v}_i\}_{i \in V}$.

Lemma 6 (Fact 5.19 from [48]). *Let $\{\mathbf{v}_i\}_{i \in V}$ be a vector embedding such that $\sum_{i \in V} \mu_i \mathbf{v}_i = \mathbf{0}$. We have:*

$$\sum_{i \in S} \frac{\mu_i}{\boldsymbol{\mu}(V)} \cdot \|\mathbf{v}_i\|_2^2 \geq \frac{\boldsymbol{\mu}(S)}{\boldsymbol{\mu}(V)} \left(\text{Var}_{\boldsymbol{\mu}}(\mathbf{v}_i) - \frac{1}{\boldsymbol{\mu}(V)} \cdot \sum_{i,j \in \binom{S}{2}} \frac{\mu_i \mu_j}{\boldsymbol{\mu}(V)} \cdot \|\mathbf{v}_i - \mathbf{v}_j\|_2^2 \right).$$

5.2.2 Updating the vector embedding via mirror descent

Khandekar *et al.* [32] describe a vector embedding update based on computing the matrix exponential of the current certificate graph’s normalized Laplacian.

It is convenient to cast this update using mirror descent. Suppose at time t , the cut player has received the sequence of matching actions D_0, \dots, D_t . In this setting, the update takes in a sequence of “feedback” matrices corresponding to $\mathcal{L}_{\boldsymbol{\mu}}(D_0), \dots, \mathcal{L}_{\boldsymbol{\mu}}(D_t)$, then computes a matrix \mathbf{X}_t given by the mirror descent step over the set of density matrices $\Delta_{n \times n} \stackrel{\text{def}}{=} \{\mathbf{X} \succeq \mathbf{0} : \text{tr } \mathbf{X} = 1\}$ using the Von Neumann entropy, $-\langle \mathbf{X}, \log \mathbf{X} \rangle$, as a regularizer. We summarize this update in `update-embedding` below.

Algorithm 2 `update-embedding`

Parameters: Graph Laplacians $\mathcal{L}_{\boldsymbol{\mu}}(D_0), \dots, \mathcal{L}_{\boldsymbol{\mu}}(D_t)$, and step size $\eta > 0$.

Do: The following:

1. Compute the matrix multiplicative update:

$$\mathbf{X}_t = \frac{\exp\left(-\eta \sum_{k=0}^t \mathcal{L}_{\boldsymbol{\mu}}(D_k)\right)}{\text{tr} \exp\left(-\eta \sum_{k=0}^t \mathcal{L}_{\boldsymbol{\mu}}(D_k)\right)}$$

Output: the matrix \mathbf{X}_t

It is important to note that our cut strategy ultimately does not use `update-embedding` as an explicit computational step; if one hopes for a cut strategy that runs in nearly-linear time, then the matrix exponential cannot be computed in full dimension. Instead, we use fast matrix exponentiation algorithms with Johnson–Lindenstrauss to compute the embedding given by low-dimensional approximations of $\mathbf{X}_t \mathbf{u}$ for $\mathbf{u} \in \mathbb{R}^V$ a unit vector. We describe this in more detail when we summarize our cut strategy in section 5.2.4

Contextualizing this update using mirror descent is convenient for analytic purposes: one can leverage regret bounds to lower bound the spectral gap, and subsequently the expansion, of the certificate graph. The following is a straightforward application of the regret bound given by Theorem 3.1 from [3]. Further exposition on mirror descent, and a full proof of the following lemma are provided in Appendix C.1.

Lemma 7. Let $T > 0$, and D_1, \dots, D_T be any sequence of ϵ -approximate matching actions. For each $t \in [T]$, suppose `update-embedding` outputs $\mathbf{X}_t \in \Delta_{n \times n}$ when given the feedback matrices $\mathcal{L}_\mu(D_0), \dots, \mathcal{L}_\mu(D_t)$. Finally, consider the graph $H = \frac{1}{T} \sum_{t=1}^T D_t$. Then for any $\eta > 0$,

$$\lambda_2(\mathcal{L}_\mu(H_T)) \geq \frac{1-2\eta}{T} \sum_{t=1}^T \langle \mathcal{L}_\mu(D_t), \mathbf{X}_t \rangle - \frac{\log n}{T\eta}.$$

Finally, we require a fact regarding the vector embedding whose Gram matrix is the output of `update-embedding`. The following lemma determines where the average embedding vector, and what the variance of the embedding is. Its proof is also in Appendix C.1.

Lemma 8. Suppose `update-embedding` outputs \mathbf{X}_t when given the feedback matrices $\mathcal{L}_\mu(D_0), \dots, \mathcal{L}_\mu(D_t)$. If $\{\mathbf{v}_i\}_{i \in V}$ is the vector embedding whose Gram matrix is \mathbf{X}_t , then the following hold:

1. $\sum_{i \in V} \|\mathbf{v}_i\|^2 = 1$,
2. $\sum_{i \in V} \sqrt{\mu_i} \cdot \mathbf{v}_i = \mathbf{0}$.

5.2.3 Rounding a robustly separated set

Recall that the cut player seeks to make the expansion of $\sum_{t=1}^T D_t$ large, in as few rounds as possible. The regret bound given by Lemma 7 implies that it suffices for the cut player to compute an embedding such that the term $\langle \mathcal{L}_\mu(D_t), \mathbf{X}_t \rangle$ is large for each t . It is illuminating to rewrite $\langle \mathcal{L}_\mu(D_t), \mathbf{X}_t \rangle$ using the vector embedding whose Gram matrix is \mathbf{X}_t . Supposing that $\mathbf{X}_t = \mathbf{V}_t \mathbf{V}_t^\top$ where the i -th row of \mathbf{V}_t is given by \mathbf{v}_i for each $i \in V$, we have

$$\langle \mathcal{L}_\mu(D_t), \mathbf{X}_t \rangle = \langle \mathbf{L}(D_t), \mathbf{M}^{-1/2} \mathbf{V}_t \mathbf{V}_t^\top \mathbf{M}^{-1/2} \rangle = \sum_{\{i,j\} \in E_{D_t}} w_{ij}^{D_t} \cdot \left\| \frac{\mathbf{v}_i}{\sqrt{\mu_i}} - \frac{\mathbf{v}_j}{\sqrt{\mu_j}} \right\|_2^2. \quad (6)$$

The term $\langle \mathcal{L}_\mu(D_t), \mathbf{X}_t \rangle$ must be large regardless of what ϵ -approximate matching action is returned by the matching strategy. Recall that our matching strategy is to output the demand graph corresponding to a hypergraph flow computed from an ϵ -approximate solution to the ratio-cut improvement problem. Our cut strategy must then output a cut such that, no matter what approximate flow the matching player routes, D will be forced to route a large amount of flow between vertices whose embedding vectors are far apart.

If the cut player is not careful in choosing $S \subseteq V$, then D being the demand graph of an *approximate* solution may, in the worst-case, possess no flow routed between vertices whose embedding vectors are far in ℓ_2^2 -distance. Our fix is to round a set S such that a constant fraction of the total flow routed across S will route into another subset of vertices $T \subseteq \bar{S}$ whose embedding vectors are well separated, *independent* of what $(1-\epsilon)$ -fraction of vertices in S possess flow routed from it. We capture this property using a (σ, c) -robustly separated set.

Definition 13 ((σ, c) -robustly separated set). Let $\sigma, c > 0$. We say $S \subseteq V$ is a (σ, c) -robustly separated set with respect to a vector embedding $\{\mathbf{v}_i\}_{i \in V} \subseteq \mathbb{R}^d$ if there exists $T \subseteq \bar{S}$ satisfying the following conditions.

1. $\mu(T) \geq c \cdot \mu(V)$.
2. $\|\mathbf{v}_i - \mathbf{v}_j\|_2^2 \geq \frac{\sigma}{\mu(S)}$ for all $i \in S$ and $j \in T$.

The rest of this section is devoted to demonstrating how to round a robustly separated set from a vector embedding $\{\mathbf{v}_i\}_{i \in V} \subseteq \mathbb{R}^d$. Before proceeding, we remark on how the embedding is normalized. In light of the calculation in (6), we can focus our attention on the vector embeddings of the form $\hat{\mathbf{v}}_i = \frac{\mathbf{v}_i}{\sqrt{\mu_i}}$ where \mathbf{v}_i comes from `update-embedding`. Lemma 8 then implies that $\hat{\mathbf{v}}_{\text{avg}} = \mathbf{0}$, and $\text{Var}_\mu(\hat{\mathbf{v}}_i) = \frac{1}{\mu(V)}$. Consequently, the statements in this section assume that $\mathbf{v}_{\text{avg}} = \mathbf{0}$, and $\text{Var}_\mu(\mathbf{v}_i) = \frac{1}{\mu(V)}$.

Our algorithm for rounding a robustly separated set is split between two cases: (1) when the vector embedding is *balanced*, and (2) when it is *unbalanced*. Given a $\{\mathbf{v}_i\}_{i \in V}$, we define the sweep cut $R_t \subseteq V$

$$R_t \stackrel{\text{def}}{=} \left\{ i \in V : \|\mathbf{v}_i\|_2^2 \leq t \cdot \mathbb{E}_{i \sim \mu} [\|\mathbf{v}_i\|_2^2] \right\}.$$

We then say that an embedding is (t, b) -balanced if the variance due to vertices in R_t captures a b -fraction of the total embedding variance.

Definition 14 ((t, b) -Balanced Embedding). Given $t > 0$, and $b > 0$, a vector embedding $\{\mathbf{v}_i\}_{i \in V} \subseteq \mathbb{R}^d$ is (t, b) -balanced if cut R_t satisfies

$$\frac{1}{\mu(V)} \cdot \sum_{i, j \in \binom{R_t}{2}} \frac{\mu_i \mu_j}{\mu(V)} \cdot \|\mathbf{v}_i - \mathbf{v}_j\|_2^2 \geq b \cdot \text{Var}_{\mu}(\mathbf{v}_i).$$

We remark that, given t and b , one can check if the embedding is (t, b) -balanced in $O(nd)$ time by computing distances to the average vector in R_t , and V .

Let us now describe our rounding algorithm **round-cut** summarized below. The constants in what follows are chosen for convenience and could, otherwise, be further optimized.

Algorithm 3 round-cut.

Parameters: $d > 0$, a vector embedding $\{\mathbf{v}_i\}_{i \in V} \subseteq \mathbb{R}^d$ such that $\mathbf{v}_{\text{avg}} = \mathbf{0}$, and $\text{Var}_{\mu}(\mathbf{v}_i) = \frac{1}{\mu(V)}$.

Do: the following

1. If $\{\mathbf{v}_i\}_{i \in V} \subseteq \mathbb{R}^d$ is $(3, \frac{1}{10})$ -balanced, do the following.

- Sample \mathbf{g} uniformly at random from \mathcal{S}^{d-1} .
- Compute the projections $r_i = \sqrt{d} \cdot \langle \mathbf{g}, \mathbf{v}_i \rangle$ for each $i \in V$.
- Sort r_i . Assume without loss of generality that $r_1 \geq \dots \geq r_n$ and define sweep cuts

$$S_t = \{i \in V : r_i \geq r_t\}$$

- Output (S_t, \overline{S}_t) such that $\mu(S_t) \geq \frac{1}{3} \cdot \mu(V)$ and $(r_t - r_{t+1}) \geq \frac{r}{\mu(V)}$ for the constant $r > 0$ guaranteed by Lemma 19.

2. If $\{\mathbf{v}_i\}_{i \in V} \subseteq \mathbb{R}^d$ is not $(3, \frac{1}{10})$ -balanced, do the following.

- Compute $\ell_i = \|\mathbf{v}_i\|_2^2$.
- Sort ℓ_i . Assume without loss of generality that $\ell_1 \geq \dots \geq \ell_n$ and define sweep cuts

$$S_t = \{i \in V : \ell_i \geq \ell_t\}$$

- Output (S_t, \overline{S}_t) as guaranteed by Lemma 10

Output: The cut $S_t \subseteq V$.

In the balanced case, R_3 captures a constant fraction of the vector embedding's variance. R_3 , a set of short vectors, also captures a large fraction of the total volume $\mu(V)$ via Markov's inequality. One can

subsequently expect that random projection coupled with a search for a balanced sweep cut will work as a bulk of the vectors are well separated from one another. The balanced case analysis is thus very similar to the analysis of SDP-based approximation algorithms for balanced separator. In fact, it is simpler as one does not need to bound the conductance of such a cut. We leave the proof of the following lemma to Appendix C.2.

Lemma 9. *Given $V = [n]$, let $\{\mathbf{v}_i\}_{i \in V} \subseteq \mathbb{R}^d$ be a vector embedding satisfying $\mathbf{v}_{\text{avg}} = \mathbf{0}$ and $\text{Var}_{\mu}(\mathbf{v}_i) = \frac{1}{\mu(V)}$. If $\{\mathbf{v}_i\}_{i \in V}$ is $(3, \frac{1}{10})$ -balanced embedding, then algorithm **round-cut** outputs S that is $(\Omega(\frac{1}{\log n}), \frac{1}{3})$ -robustly separated set with probability $1 - o_n(\frac{1}{n})$ over the uniformly random choice of $\mathbf{g} \in \mathcal{S}^{d-1}$.*

In the unbalanced case, R_3 fails to capture a constant fraction of the variance. R_3 is a set of long vectors with small volume and so one might hope that R_3 and $\overline{R_3}$ are already robustly separated. However, this is not always the case as there may be many vertex pairs near the defining threshold of R_3 whose embedding vectors are close to each other. Routing an approximate flow may thus route only between such pairs causing $\langle \mathcal{L}_{\mu}(D_t), \mathbf{X}_t \rangle$ to be small.

The key insight is that picking a smaller threshold, say $R_{3/2}$, and performing a region growing argument demonstrates the existence of another sweep cut $\overline{R_t}$ contained within $\overline{R_3}$ that is robustly separated from $R_{3/2}$.

Lemma 10. *Given $V = [n]$, let $\{\mathbf{v}_i\}_{i \in V} \subseteq \mathbb{R}^d$ be a vector embedding satisfying $\mathbf{v}_{\text{avg}} = \mathbf{0}$ and $\text{Var}_{\mu}(\mathbf{v}_i) = \frac{1}{\mu(V)}$. If $\{\mathbf{v}_i\}_{i \in V}$ is not $(3, \frac{1}{10})$ -balanced, then **round-cut** deterministically outputs a $(\Omega(\frac{1}{\log n}), \frac{1}{3})$ -robustly separated set S .*

Proof. Fix $T = R_{3/2}$ where $R_t = \{i \in V : \|\mathbf{v}_i\|_2^2 \leq \frac{t}{\mu(V)}\}$. We first demonstrate that there exists an $S \subseteq \overline{R_3}$ that is $(\Omega(\frac{1}{\log n}), \frac{1}{3})$ -robustly separated with respect to T . The first requirement for any $S \subseteq \overline{R_3}$ to be a robustly separated, $\mu(T) \geq \frac{1}{3} \cdot \mu(V)$, is satisfied since Markov's inequality dictates:

$$\mu(T) = \mu(R_{3/2}) \geq \left(1 - \frac{1}{3/2}\right) \cdot \mu(V) = \frac{1}{3} \cdot \mu(V).$$

All that remains to show is that there exists some S where every $i \in S, j \in T$ satisfies $\|\mathbf{v}_i - \mathbf{v}_j\|_2^2 \geq \frac{\sigma}{\mu(S)}$ for $\sigma \geq \Omega(\frac{1}{\log n})$. Call such an S valid.

We begin by making two observations. First, the average length of vectors in $\overline{R_3}$ is at least as large as a constant. By Lemma 6, we have that:

$$\sum_{i \in \overline{R_3}} \frac{\mu_i}{\mu(V)} \cdot \|\mathbf{v}_i\|_2^2 \geq \frac{\mu(R_3)}{\mu(V)} \left(\text{Var}_{\mu}(\mathbf{v}_i) - \frac{1}{\mu(V)} \cdot \sum_{i,j \in \binom{R_3}{2}} \frac{\mu_i \mu_j}{\mu(V)} \cdot \|\mathbf{v}_i - \mathbf{v}_j\|_2^2 \right).$$

By Markov's inequality, we have that $\frac{\mu(R_3)}{\mu(V)} \geq \frac{2}{3}$. Furthermore, since the vector embedding $\{\mathbf{v}_i\}_{i \in V} \subseteq \mathbb{R}^d$ is not $(3, \frac{1}{10})$ -balanced, we have:

$$\frac{1}{\mu(V)} \cdot \sum_{i,j \in \binom{R_3}{2}} \frac{\mu_i \mu_j}{\mu(V)} \cdot \|\mathbf{v}_i - \mathbf{v}_j\|_2^2 < \frac{1}{10} \cdot \text{Var}_{\mu}(\mathbf{v}_i).$$

Combining the three inequalities above, and using that $\text{Var}_{\mu}(\mathbf{v}_i) = \frac{1}{\mu(V)}$, we derive:

$$\sum_{i \in \overline{R_3}} \frac{\mu_i}{\mu(V)} \cdot \|\mathbf{v}_i\|_2^2 \geq \frac{2}{3} \cdot \left(1 - \frac{1}{10}\right) \cdot \frac{1}{\mu(V)} \iff \sum_{i \in \overline{R_3}} \mu_i \cdot \|\mathbf{v}_i\|_2^2 \geq \frac{3}{5}. \quad (7)$$

Second, consider any $t \geq 3$, and note that not only $\overline{R_t} \subseteq \overline{R_3}$, but also that we can lower bound $\|\mathbf{v}_i - \mathbf{v}_j\|_2^2$ for any $i \in \overline{R_t}$ and $j \in T$ as follows. Since $i \in \overline{R_t}$, we have $\|\mathbf{v}_i\|_2^2 \geq \frac{t}{\mu(V)}$. Similarly as $j \in T$, we have $\|\mathbf{v}_j\|_2^2 \leq \frac{3/2}{\mu(V)}$. Hence, the triangle inequality implies:

$$\|\mathbf{v}_i - \mathbf{v}_j\|_2 \geq \|\mathbf{v}_i\|_2 - \|\mathbf{v}_j\|_2 \geq \sqrt{\frac{t}{\mu(V)}} - \sqrt{\frac{3/2}{\mu(V)}} \iff \|\mathbf{v}_i - \mathbf{v}_j\|_2^2 \geq \frac{(\sqrt{t} - \sqrt{3/2})^2}{\mu(V)} \quad (8)$$

Now, fix σ , and for the sake of contradiction, suppose no valid $S \subseteq \overline{R_3}$ exists. That is, for every $S \subseteq \overline{R_3}$, there exists $i \in S, j \in T$ such that $\|\mathbf{v}_i - \mathbf{v}_j\|_2^2 < \frac{\sigma}{\mu(S)}$. Since $\overline{R_t} \subseteq \overline{R_3}$ for every $t \geq 3$, this implies

$$\|\mathbf{v}_i - \mathbf{v}_j\|_2^2 < \frac{\sigma}{\mu(\overline{R_t})}.$$

for some $i \in \overline{R_t}$ and $j \in T$. However, inequality (8) holds for every $t \geq 3$, $i \in \overline{R_t}$, and $j \in T$. Consequently,

$$\frac{(\sqrt{t} - \sqrt{3/2})^2}{\mu(V)} \leq \|\mathbf{v}_i - \mathbf{v}_j\|_2^2 < \frac{\sigma}{\mu(\overline{R_t})} \iff \mu(\overline{R_t}) < \frac{\sigma}{(\sqrt{t} - \sqrt{3/2})^2} \cdot \mu(V) \leq \frac{30\sigma}{\sqrt{t}} \cdot \mu(V) \quad (9)$$

where we use the fact that $(\sqrt{t} - \sqrt{3/2})^2 \geq \frac{t}{30}$ for all $t \geq 3$. Using this, we can strike the following upper bound on $\sum_{i \in \overline{R_3}} \mu_i \cdot \|\mathbf{v}_i\|$. If we let $M \stackrel{\text{def}}{=} \max_{i \in V} \{\mu(V) \cdot \|\mathbf{v}_i\|_2^2\}$, then we can write

$$\begin{aligned} \sum_{i \in \overline{R_3}} \mu_i \cdot \|\mathbf{v}_i\| &= \frac{1}{\mu(V)} \cdot \int_3^M \mu(\overline{R_t}) dt + \frac{3}{\mu(V)} \cdot \mu(\overline{R_3}) \\ &< 30\sigma \cdot \left(\int_3^M \frac{1}{t} dt + \frac{1}{\sqrt{3}} \right) \\ &= 30\sigma \cdot \left(\ln\left(\frac{M}{3}\right) + \frac{1}{\sqrt{3}} \right) \end{aligned}$$

Because $\sum_{i \in V} \mu_i \cdot \|\mathbf{v}_i\|_2^2 = 1$, we have $M \leq \max_{i \in V} \left\{ \frac{\mu(V)}{\mu_i} \right\} \leq \text{poly}(n)$. Consequently,

$$\sum_{i \in \overline{R_3}} \mu_i \cdot \|\mathbf{v}_i\| < \sigma \cdot O(\log n).$$

However, inequality (7) implied that $\sum_{i \in \overline{R_3}} \mu_i \cdot \|\mathbf{v}_i\| \geq \Omega(1)$ which reaches a contradiction unless $\sigma \geq \Omega\left(\frac{1}{\log n}\right)$.

Consequently, there exists some $S \subseteq \overline{R_3}$ that is $\left(\Omega\left(\frac{1}{\log n}\right), \frac{1}{3}\right)$ -robustly separated from T . Furthermore, $S = \overline{R_t}$ for some $t \geq 3$, and since **round-cut** step (2) searches through all of at most n such cuts, we conclude that **round-cut** outputs the required robustly separated set in the unbalanced case. \square

This section concludes with a lemma capturing both the fact that **round-cut** produces a $\left(\Omega\left(\frac{1}{\log n}\right), \frac{1}{3}\right)$ -robustly separated set in all cases, and the running time of **round-cut**.

Lemma 11. *Given $V = [n]$, let $\{\mathbf{v}_i\}_{i \in V} \subseteq \mathbb{R}^d$ be a vector embedding satisfying $\mathbf{v}_{\text{avg}} = \mathbf{0}$ and $\text{Var}_{\mu}(\mathbf{v}_i) = \frac{1}{\mu(V)}$. Then **round-cut** outputs a $\left(\Omega\left(\frac{1}{\log n}\right), \frac{1}{3}\right)$ -robustly separated set S with $1 - o_n\left(\frac{1}{n}\right)$ probability over the uniformly random choice of $\mathbf{g} \in \mathcal{S}^{d-1}$. Furthermore, **round-cut** runs in time $\tilde{O}(nd)$.*

Proof. If $\{\mathbf{v}_i\}_{i \in V}$ is $(3, \frac{1}{10})$ -balanced, then Lemma 9 implies **round-cut** outputs a $\left(\Omega\left(\frac{1}{\log n}\right), \frac{1}{3}\right)$ -robustly separated set with probability $1 - o_n\left(\frac{1}{n}\right)$. Otherwise, Lemma 10 provides the same guarantee deterministically.

To bound the running time, consider the following. Checking if the embedding is balanced takes $O(nd)$ time. Computing ℓ_2^2 -distances and vector projections takes $O(d)$ time per vector, of which there are n . Sorting takes $\tilde{O}(n)$ time. There are at most $O(n)$ sweep cuts to check in both the balanced and unbalanced case, and finding the appropriate cut in both cases takes $O(1)$ time per cut. \square

5.2.4 Completing the cut strategy

As alluded to in Section 5.2.2, the matrix exponential required to compute the mirror descent update cannot be computed in full if one wants a nearly-linear time algorithm. Here, we can take advantage of the fact that each feedback matrix provided to mirror descent is given by the Laplacian of a sparse graph D_t . So long as at most $\text{polylog}(n)$ iterations of mirror descent are required, the sum of all D_t , throughout the lifetime of the algorithm, will remain sparse. We denote algorithm EXPV as some generic algorithm that computes the matrix vector product $\mathbf{X}_t \mathbf{u}$ for \mathbf{u} some unit vector. One can find an explicit description of this algorithm in Theorem 3.2 of [48] which takes advantage of the Laplacian system solver of Spielman and Teng [54].

Next, we use Johnson–Lindenstrauss to project the embedding into $O\left(\frac{\log n}{\delta^2}\right)$ dimensions for $\delta = O(1)$,

Lemma 12 (Johnson–Lindenstrauss). *Given an embedding $\{\mathbf{v}_i\}_{i \in V} \subseteq \mathbb{R}^d$ for $V = [n]$, and $\delta > 0$, let $k > \Omega\left(\frac{\log n}{\delta^2}\right)$ and $\mathbf{U} \in \mathbb{R}^{k \times d}$ be a matrix where the i -th row is a vector sampled independently and uniformly from the $(n-1)$ -dimensional sphere of radius $\sqrt{\frac{n}{k}}$. Let $\tilde{\mathbf{v}}_i = \mathbf{U} \mathbf{v}_i$ for each $i \in V$. With probability $1 - o_n\left(\frac{1}{n}\right)$, the following holds for all $i, j \in V$:*

$$(1 - \delta) \cdot \|\mathbf{v}_i - \mathbf{v}_j\|^2 \leq \|\tilde{\mathbf{v}}_i - \tilde{\mathbf{v}}_j\|^2 \leq (1 + \delta) \cdot \|\mathbf{v}_i - \mathbf{v}_j\|^2.$$

allowing us to compute ℓ_2^2 -distances and projections in logarithmic time. Since the error is multiplicative in terms of pairwise ℓ_2^2 -distances, picking δ to be constant only causes a constant factor loss in the number of cut strategy plays, and the final approximation ratio.

Once we compute the low-dimensional vector embedding, we use **round-cut**, to produce a robustly separated set and output that as the cut strategy’s action. This procedure is summarized in algorithm 4 \mathcal{C}_{MD} .

Algorithm 4 \mathcal{C}_{MD} .

Parameters: ϵ -approximate matching actions D_0, \dots, D_t , step size $\eta > 0$, $1 > \delta \geq \frac{1}{10}$.

Do: The following.

1. Pick $d = O\left(\frac{\log n}{\delta^2}\right)$ random unit vectors $\mathbf{u}_1, \dots, \mathbf{u}_d \in \mathbb{R}^V$.
2. Denote \mathbf{X}_t as the matrix update specified in **update-embedding** $(\mathcal{L}_\mu(D_0), \dots, \mathcal{L}_\mu(D_t), \eta)$, and use EXPV to compute embedding $\{\tilde{\mathbf{v}}_i\}_{i \in V}$ given by

$$(\tilde{\mathbf{v}}_i)_j = \sqrt{\frac{n}{d}} \cdot \langle \mathbf{X}_t \mathbf{u}_j, \mathbf{e}_i \rangle \quad \forall i \in V, j \in [d]$$

3. Scale the embedding $\hat{\mathbf{v}}_i^{(t)} = \frac{\tilde{\mathbf{v}}_i^{(t)}}{\sqrt{\mu_i}}$ for each $i \in V$.
4. Execute **round-cut** $(d, \{\hat{\mathbf{v}}_i^{(t)}\}_{i \in V})$ to produce the cut $S_t \subseteq V$.

Output: The cut S_t .

Let us proceed with the analysis. First, we demonstrate that returning a robustly separated set forces $\langle \mathcal{L}_\mu(D), \mathbf{X} \rangle$ to be large for any ϵ -approximate matching action D .

Lemma 13. *Let $S \subseteq V$ be a (σ, c) -robustly separated set with respect to a vector embedding $\{\mathbf{v}_i\}_{i \in V} \subseteq \mathbb{R}^d$. For any $\epsilon > 0$, if $D = (S \cup \bar{S}, E_D, \mathbf{w}^D)$ is an ϵ -approximate matching action with respect to S , then*

$$\sum_{i \in S, j \in \bar{S}} w_{ij}^D \cdot \|\mathbf{v}_i - \mathbf{v}_j\|_2^2 \geq \sigma \cdot \left(\frac{1}{3} - 4\epsilon\right).$$

Proof. Since $S \subseteq V$ is (σ, c) -robustly separated, there exists $T \subseteq \bar{S}$ such that $\boldsymbol{\mu}(T) \geq c \cdot \boldsymbol{\mu}(V)$ and $\|\mathbf{v}_i - \mathbf{v}_j\|_2^2 \geq \frac{\sigma}{\boldsymbol{\mu}(S)}$ for all $i \in S$ and $j \in T$. Thus we can bound

$$\sum_{i \in S, j \in \bar{S}} w_{ij}^D \cdot \|\mathbf{v}_i - \mathbf{v}_j\|_2^2 \geq \sum_{i \in S, j \in T} w_{ij}^D \cdot \|\mathbf{v}_i - \mathbf{v}_j\|_2^2 \geq \frac{\sigma}{\boldsymbol{\mu}(S)} \cdot \sum_{i \in S, j \in T} w_{ij}^D. \quad (10)$$

Observing that D is bipartite between (S, \bar{S}) , we have

$$\sum_{i \in S, j \in T} w_{ij}^D = \sum_{i \in S} \deg_i(D) - \sum_{i \in \bar{S} \setminus T} \deg_i(D). \quad (11)$$

Now we can lower bound the first term in the RHS of (11) like so. Notice that

$$\sum_{i \in S} (\mu_i \cdot |s_i| - \deg_i(D)) \leq \sum_{i \in S} |\mu_i \cdot |s_i| - \deg_i(D)| \leq \|\boldsymbol{\mu} \circ |s| - \mathbf{d}\|_1 \leq \epsilon \cdot \|\mathbf{s}\|_{1, \boldsymbol{\mu}},$$

where the last inequality follows as D is an ϵ -approximate matching action. Consequently,

$$\sum_{i \in S} \deg_i(D) \geq \sum_{i \in S} \mu_i \cdot |s_i| - \epsilon \cdot \|\mathbf{s}\|_{1, \boldsymbol{\mu}} = \boldsymbol{\mu}(S) - 2\epsilon \cdot \boldsymbol{\mu}(S). \quad (12)$$

Meanwhile, the second term in the RHS of (11) can be upper bounded via the following.

$$\sum_{i \in \bar{S} \setminus T} (\deg_i(D) - \mu_i \cdot |s_i|) \leq \sum_{i \in \bar{S} \setminus T} |\deg_i(D) - \mu_i \cdot |s_i|| \leq \|\mathbf{d} - (\boldsymbol{\mu} \circ |s|)\|_1 \leq \epsilon \cdot \|\mathbf{s}\|_{1, \boldsymbol{\mu}}$$

where the last inequality again comes from D being an ϵ -approximate matching action. Re-arranging gives

$$\sum_{i \in \bar{S} \setminus T} \deg_i(D) \leq \sum_{i \in \bar{S} \setminus T} \mu_i \cdot |s_i| + \epsilon \cdot \|\mathbf{s}\|_{1, \boldsymbol{\mu}} = \frac{\boldsymbol{\mu}(S)}{\boldsymbol{\mu}(\bar{S})} \cdot \boldsymbol{\mu}(\bar{S} \setminus T) + 2\epsilon \cdot \boldsymbol{\mu}(S) \leq \frac{2}{3} \cdot \boldsymbol{\mu}(S) + 2\epsilon \cdot \boldsymbol{\mu}(S). \quad (13)$$

The last inequality comes from the fact that $\boldsymbol{\mu}(T) \geq \frac{1}{3} \cdot \boldsymbol{\mu}(V) \geq \frac{1}{3} \boldsymbol{\mu}(\bar{S})$, and the fact that $T \subseteq \bar{S}$ and hence $\boldsymbol{\mu}(\bar{S} \setminus T) = \boldsymbol{\mu}(\bar{S}) - \boldsymbol{\mu}(T)$. Substituting inequalities (12) and (13) into (11) then implies

$$\sum_{i \in S, j \in T} w_{ij}^D \geq \left(\frac{1}{3} - 4\epsilon\right) \cdot \boldsymbol{\mu}(S).$$

Inequality (10) then reduces to:

$$\sum_{i \in S, j \in \bar{S}} w_{ij}^D \cdot \|\mathbf{v}_i - \mathbf{v}_j\|_2^2 \geq \sigma \cdot \left(\frac{1}{3} - 4\epsilon\right)$$

as required. \square

Using Lemma 13 to analyze the regret bound in Lemma 7 yields a proof a main theorem, Theorem 10.

Proof of Theorem 10. Let $T = g(n)$, and suppose \mathcal{C}_{MD} plays the cuts S_1, \dots, S_T . Consider any sequence of weighted graphs D_1, \dots, D_T such that D_t is an ϵ -approximate matching action with respect to S_t for each $t = 1, \dots, T$. Define \mathcal{E}_t to be the event where the following occurs in iteration t :

1. the random projection in step (2) of \mathcal{C}_{MD} produces an embedding satisfying the Johnson–Lindenstrauss error bounds, and
2. calling **round-cut** in step (4) of \mathcal{C}_{MD} produces a $\left(\Omega\left(\frac{1}{\log n}\right), \frac{1}{3}\right)$ -robustly separated set.

First, we prove that if \mathcal{E}_t occur jointly over all $t = 1, \dots, T$, then \mathcal{C}_{MD} is an $\left(\Omega(\log n), \left(\frac{\log^2 n}{1-\epsilon}\right)\right)$ -good cut strategy. For the graph $H = \sum_{t=1}^T D_t$, Lemma 7 implies the lower bound on the spectral gap of $\frac{1}{T}H$

$$\lambda_2\left(\mathcal{L}_{\boldsymbol{\mu}}\left(\frac{1}{T} \cdot H\right)\right) \geq \frac{1-2\eta}{T} \sum_{t=1}^T \langle \mathcal{L}_{\boldsymbol{\mu}}(D_t), \mathbf{X}_t \rangle - \frac{\log n}{T\eta}.$$

Let us now consider the t -th cut played by \mathcal{C}_{MD} . Suppose \mathbf{X}_t , the matrix update specified by step (2) in \mathcal{C}_{MD} , is also the Gram matrix of an embedding $\{\mathbf{v}_i^{(t)}\}_{i \in V}$. Denote \mathbf{V}_t to be the matrix whose i -th row is $\mathbf{v}_i^{(t)}$. We can then write $\mathbf{X}_t = \mathbf{V}_t \mathbf{V}_t^\top$ and hence

$$\langle \mathcal{L}_{\boldsymbol{\mu}}(D_t), \mathbf{X}_t \rangle = \langle \mathbf{L}(D_t), \mathbf{M}^{-1/2} \mathbf{V}_t \mathbf{V}_t^\top \mathbf{M}^{-1/2} \rangle = \sum_{\{i,j\} \in E_{D_t}} w_{ij}^{D_t} \cdot \left\| \frac{\mathbf{v}_i}{\sqrt{\mu_i}} - \frac{\mathbf{v}_j}{\sqrt{\mu_j}} \right\|_2^2.$$

Using Lemma 12, we have

$$\begin{aligned} \sum_{\{i,j\} \in E_{D_t}} w_{ij}^{D_t} \cdot \left\| \frac{\mathbf{v}_i}{\sqrt{\mu_i}} - \frac{\mathbf{v}_j}{\sqrt{\mu_j}} \right\|_2^2 &\geq \frac{1}{1+\delta} \cdot \sum_{\{i,j\} \in E_{D_t}} w_{ij}^{D_t} \cdot \left\| \frac{\hat{\mathbf{v}}_i}{\sqrt{\mu_i}} - \frac{\hat{\mathbf{v}}_j}{\sqrt{\mu_j}} \right\|_2^2 \\ &= \frac{1}{1+\delta} \cdot \sum_{\{i,j\} \in E_{D_t}} w_{ij}^{D_t} \cdot \|\hat{\mathbf{v}}_i - \hat{\mathbf{v}}_j\|_2^2 \geq \frac{1}{2} \cdot \sum_{\{i,j\} \in E_{D_t}} w_{ij}^{D_t} \cdot \|\hat{\mathbf{v}}_i - \hat{\mathbf{v}}_j\|_2^2. \end{aligned}$$

with the last inequality following as $\delta < 1$. Focusing on the embedding $\{\hat{\mathbf{v}}_i\}_{i \in V}$, note that it satisfies

$$\sum_{i \in V} \frac{\mu_i}{\boldsymbol{\mu}(V)} \cdot \|\hat{\mathbf{v}}_i\|_2^2 = \frac{1}{\boldsymbol{\mu}(V)} \sum_{i \in V} \|\hat{\mathbf{v}}_i\|_2^2 = \frac{1}{\boldsymbol{\mu}(V)}$$

where the last equality follows by Lemma 8 item (1). Furthermore,

$$\sum_{i \in V} \mu_i \cdot \hat{\mathbf{v}}_i = \sum_{i \in V} \sqrt{\mu_i} \cdot \mathbf{v}_i = \mathbf{0}$$

where the last equality follows by Lemma 8 item (2). Consequently, the embedding that is passed into **round-cut** in iteration t is centered appropriately. Lemma 11 then implies that **round-cut** and \mathcal{C}_{MD} output $S_t \subseteq V$ that is $\left(\Omega\left(\frac{1}{\log n}\right), \frac{1}{3}\right)$ -robustly separated with respect to $\{\hat{\mathbf{v}}_i\}_{i \in V}$. Lemma 13 then implies

$$\frac{1}{2} \cdot \sum_{\{i,j\} \in E_{D_t}} w_{ij}^{D_t} \cdot \|\hat{\mathbf{v}}_i - \hat{\mathbf{v}}_j\|_2^2 = \frac{1}{2} \cdot \sum_{i \in S, j \in \bar{S}} w_{ij}^{D_t} \cdot \|\hat{\mathbf{v}}_i - \hat{\mathbf{v}}_j\|_2^2 \geq \Omega\left(\frac{1}{\log n}\right) \cdot \left(\frac{1-12\epsilon}{6}\right) = \Omega\left(\frac{1-\epsilon}{\log n}\right).$$

Returning to our regret bound, this implies that there exists an absolute constant $C > 0$ such that

$$\lambda_2\left(\mathcal{L}_{\boldsymbol{\mu}}\left(\frac{1}{T} \cdot H\right)\right) \geq \frac{1-2\eta}{T} \sum_{t=1}^T C \cdot \frac{1-\epsilon}{\log n} - \frac{\log n}{T\eta} = C \cdot (1-2\eta) \cdot \frac{1-\epsilon}{\log n} - \frac{\log n}{T\eta}.$$

Choosing $\eta = \sqrt{\frac{\log^2 n}{2C \cdot T \cdot (1-\epsilon)}}$ and $T = \frac{32}{C} \cdot \frac{\log^2 n}{1-\epsilon}$, we derive:

$$\lambda_2\left(\mathcal{L}_{\boldsymbol{\mu}}\left(\frac{1}{T} \cdot H\right)\right) \geq \frac{C}{2} \cdot \frac{1-\epsilon}{\log n}.$$

This implies the following bound on the spectral gap of H :

$$\lambda_2(\mathcal{L}_{\boldsymbol{\mu}}(H)) \geq T \cdot \frac{C}{2} \cdot \frac{1-\epsilon}{\log n} = 16 \cdot \log n = \Omega(\log n).$$

Finally, Cheeger's inequality implies $\Psi_H \geq \frac{\lambda_2(\mathcal{L}_\mu(H))}{2} \geq \Omega(\log n)$ hence \mathcal{C}_{MD} is a $(\Omega(\log n), O(\frac{\log^2 n}{1-\epsilon}))$ -good cut strategy as required.

Now, to show that \mathcal{C}_{MD} is a $(\Omega(\log n), O(\frac{\log^2 n}{1-\epsilon}))$ -good cut strategy with probability $1 - o_n(1)$, note that \mathcal{C}_{MD} requires at most $O(\frac{\log^2 n}{1-\epsilon})$ iterations. By Lemma 12, the random projection step fails to satisfy the requisite error bound with probability smaller than $o_n(\frac{1}{n})$, while by Lemma 9 states that **round-cut** fails to output a $(\Omega(\frac{1}{\log n}), \frac{1}{3})$ -robustly separated set in the balanced case with probability smaller than $o_n(\frac{1}{n})$. Performing a union bound over all iterations of \mathcal{C}_{MD} implies \mathcal{C}_{MD} is a good cut strategy with $1 - o_n(1)$ probability. \square

6 An $O(\sqrt{\log n})$ -Approximation via Metric Embeddings

Finally, we prove Theorem 4: there exists a polynomial time randomized $O(\sqrt{\log n})$ -approximation algorithm for the minimum ratio-cut problem over hypergraphs.

We begin by constructing a convex relaxation of the minimum ratio-cut problem that embeds the vertex set V into a metric space of negative type. Recall that a metric space (\mathcal{X}, d) is said to be of negative type if there exists a function $f : \mathcal{X} \rightarrow \mathbb{R}^n$ such that:

$$d(x, y) = \|f(x) - f(y)\|_2^2.$$

Given a weighted hypergraph $G = (V, E, \mathbf{w}, \boldsymbol{\mu})$ equipped with MSCFs $\{\delta_h\}_{h \in E}$, we define two types of variables:

1. *Embedding variables* – the program assigns a vector variable $\mathbf{v}_i \in \mathbb{R}^n$ to each vertex $i \in V$. Each \mathbf{v}_i represents the position of i embedded into the metric space $(\mathbb{R}^n, \|\cdot\|_2^2)$.
2. *Length variables* – the program also assigns a vector variable $\boldsymbol{\ell}^h \in \mathbb{R}^h$ to each hyperedge $h \in E$. The coordinates ℓ_i^h can be interpreted as giving a length to the edge $\{i, h\}$ in the factor graph representation \hat{G} of G .

Given a weighted submodular hypergraph $G = (V, E, \mathbf{w}, \boldsymbol{\mu})$, we consider the following convex program:

$$\begin{aligned} \min_{\mathbf{v}_i, \boldsymbol{\ell}^h} \quad & \sum_{h \in E} w_h \cdot \|\boldsymbol{\ell}^h\|_{F_h} \\ \text{s.t.} \quad & \sum_{\{i, j\} \subseteq V} \frac{\mu_i \mu_j}{\mu(V)} \cdot \|\mathbf{v}_i - \mathbf{v}_j\|_2^2 \geq 1 \\ & \|\mathbf{v}_i - \mathbf{v}_j\|_2^2 \leq \sum_{(u, h, w) \in p} \ell_u^h + \ell_w^h \quad \forall \{i, j\} \in \binom{V}{2}, \forall p \in \mathcal{P}_{ij}(G) \\ & \|\mathbf{v}_i - \mathbf{v}_j\|_2^2 \leq \|\mathbf{v}_i - \mathbf{v}_k\|_2^2 + \|\mathbf{v}_k - \mathbf{v}_j\|_2^2 \quad \forall i, j, k \in V \\ & \mathbf{v}_i \in \mathbb{R}^n, \boldsymbol{\ell}^h \in \mathbb{R}^h \quad \forall i \in V, \forall h \in E \end{aligned} \tag{RC-Metric}$$

Here, \mathcal{P}_{ij} is the set of all paths in the factor graph \hat{G} . We note that even though the number of paths is exponential, the program has an equivalent characterization with a polynomial number of constraints. This is obtained by replacing the set of constraints

$$\|\mathbf{v}_i - \mathbf{v}_j\|_2^2 \leq \sum_{(u, h, w) \in p} \ell_u^h + \ell_w^h \quad \forall \{i, j\} \in \binom{V}{2}, \forall p \in \mathcal{P}_{ij}(G)$$

with

$$\|\mathbf{v}_i - \mathbf{v}_j\|_2^2 \leq \ell_i^h + \ell_j^h \quad \forall h \in E, \forall \{i, j\} \in \binom{h}{2},$$

which are equivalent whenever the triangle inequality constraints are enforced.

The vector constraints can alternatively be represented as positive semidefinite constraints using the Gram matrix of the vector embedding $\{\mathbf{v}_i\}_{i \in V}$. The program (**RC-Metric**) is, up to a constant factor, a relaxation of the minimum ratio cut problem, as expressed in the following theorem. This is proved in Appendix D.

Theorem 11. *The (**RC-Metric**) vector program is, up to a constant, a relaxation of the minimum ratio-cut program for the hypergraph $G = (V, E, \mathbf{w}, \boldsymbol{\mu})$ with MSCFS $\{\delta_h\}_{h \in E}$.*

6.1 Rounding the Metric Embedding Relaxation

The crucial property we rely on for rounding the program is that any feasible set of vectors $\{\mathbf{v}_i\}_{i \in V}$ produces a metric of negative type. Our argument then uses the following theorem from [7], which guarantees the existence of an efficient algorithm to embed a metric of negative type onto \mathbb{R} . This version of the theorem is given by Feige, Hajiaghayi and Lee [25].

Theorem 12 (Theorem 3.3 from [25]). *If (\mathcal{X}, d) is a n -point metric space of negative type $d: \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$, then for every product weight $\omega_0: \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}_+$ there exists an efficiently computable 1-Lipschitz⁴ map $\phi: \mathcal{X} \rightarrow \mathbb{R}$ satisfying:*

$$\frac{\sum_{\mathbf{x}, \mathbf{y} \in \mathcal{X}} \omega_0(\mathbf{x}, \mathbf{y}) \cdot d(\mathbf{x}, \mathbf{y})}{\sum_{\mathbf{x}, \mathbf{y} \in \mathcal{X}} \omega_0(\mathbf{x}, \mathbf{y}) \cdot |\phi(\mathbf{x}) - \phi(\mathbf{y})|} = O(\sqrt{\log n})$$

Before analyzing our rounding algorithm, we require a technical lemma that is proved Appendix D.

Lemma 14. *Let $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ be such that $\mathbf{y} \geq 0$ and $|x_i - x_j| \leq y_i + y_j$ for all $i, j \subseteq [n]$. Then, there exists some $\nu \in \mathbb{R}$ such that, for every $i \in [n]$:*

$$|x_i - \nu| \leq y_i.$$

Let us now show the main theorem of this section: rounding the convex program (**RC-Metric**) produces a $O(\sqrt{\log n})$ -approximation for finding the minimum ratio-cut of a hypergraph.

Theorem 13. *There is a polynomial-time algorithm that, given an optimal solution $(\{\ell^h\}_{h \in E}, \{\mathbf{v}_i\}_{i \in V})$ to (**RC-Metric**) with objective value $\text{opt} \stackrel{\text{def}}{=} \sum_{h \in E} w_h \cdot \|\ell^h\|_{F_h}$, returns a cut $S \subseteq V$ such that*

$$\Psi_G(S) \leq \text{opt} \cdot O(\sqrt{\log n}).$$

Proof. The vectors $\{\mathbf{v}_i\}$ form a metric space of negative type. Hence, we can efficiently find an embedding $\phi: \{\mathbf{v}_i\}_{i \in V} \rightarrow \mathbb{R}$ with the guarantees of Theorem 12, where ω_0 is taken to be:

$$\omega_0(i, j) = \frac{\mu_i \mu_j}{\mu(V)}.$$

Let $\mathbf{x} \in \mathbb{R}^n$ be the vector given by $x_i \stackrel{\text{def}}{=} \phi(\mathbf{v}_i)$. Since $(\{\ell^h\}_{h \in E}, \{\mathbf{v}_i\}_{i \in V})$ is feasible, 1-lipschitzness of ϕ guarantees:

$$|\phi(\mathbf{v}_i) - \phi(\mathbf{v}_j)| = |x_i - x_j| \leq \ell_i^h + \ell_j^h.$$

Hence, by Lemma 14, for every hyperedge $h \in E$, there exists some choice of $\nu_h \in \mathbb{R}$, such that:

$$(\mathbf{x} - \nu_h \mathbf{1})_i \leq \ell_i^h,$$

for every $i \in V$. By the monotonicity of the $\|\cdot\|_{F_h}$ norm (Fact 6), we have:

$$\sum_{h \in E} w_h \cdot \min_{\nu \in \mathbb{R}} \|\mathbf{x} - \nu \mathbf{1}\|_{F_h} \leq \sum_{h \in E} w_h \cdot \|\mathbf{x} - \nu_h \mathbf{1}\|_{F_h} \leq \sum_{h \in E} w_h \cdot \|\ell^h\|_{F_h} = \text{opt}. \quad (15)$$

⁴Recall that, given a metric space (X, d) , a function $f: X \rightarrow \mathbb{R}$ is said to be K -Lipschitz for some constant K if $|f(x) - f(y)| \leq K \cdot d(x, y)$ for every $x, y \in X$.

We now proceed to obtain a lower bound on the denominator of the objective. Begin by choosing

$$\gamma^* \stackrel{\text{def}}{=} \operatorname{argmin}_{\gamma \in \mathbb{R}} \|\mathbf{x} - \gamma \mathbf{1}\|_{1, \mu}$$

and let $\hat{\mathbf{x}} := \mathbf{x} - \gamma^* \mathbf{1}$. We can write the denominator as the expectation of $|\hat{x}_i|$ where $i \in V$ are sampled randomly with probability proportional to μ like so.

$$\min_{\gamma \in \mathbb{R}} \|\mathbf{x} - \gamma \mathbf{1}\|_{1, \mu} = \|\hat{\mathbf{x}}\|_{1, \mu} = \mu(V) \mathbb{E}_{i \sim \mu} [|\hat{x}_i|]$$

Notice $\mathbb{E}_{i \sim \mu} [|\hat{x}_i|]$ can then be bounded below by the following inequality:

$$\mu(V) \mathbb{E}_{i \sim \mu} [|\hat{x}_i|] = \frac{\mu(V)}{2} \cdot \left(\mathbb{E}_{i \sim \mu} [|\hat{x}_i|] + \mathbb{E}_{j \sim \mu} [|\hat{x}_j|] \right) = \frac{\mu(V)}{2} \cdot \mathbb{E}_{i, j \sim \mu \times \mu} [|\hat{x}_i| + |\hat{x}_j|] \geq \frac{\mu(V)}{2} \cdot \mathbb{E}_{i, j \sim \mu \times \mu} [|\hat{x}_i - \hat{x}_j|].$$

This expectation evaluates to:

$$\frac{\mu(V)}{2} \cdot \mathbb{E}_{i, j \sim \mu \times \mu} [|\hat{x}_i - \hat{x}_j|] = \frac{1}{2} \sum_{i, j \in V} \frac{\mu_i \mu_j}{\mu(V)} \cdot |x_i - x_j| = \frac{1}{2} \sum_{i, j \in V} \frac{\mu_i \mu_j}{\mu(V)} \cdot |x_i - x_j|.$$

Consequently, the above calculation gives the following bound.

$$\min_{\gamma \in \mathbb{R}} \|\mathbf{x} - \gamma \mathbf{1}\|_{1, \mu} \geq \frac{1}{2} \sum_{i, j \in V} \frac{\mu_i \mu_j}{\mu(V)} \cdot |x_i - x_j| \geq O\left(\frac{1}{\sqrt{\log n}}\right) \sum_{i, j \in V} \frac{\mu_i \mu_j}{\mu(V)} \|\mathbf{v}_i - \mathbf{v}_j\|_2^2 \geq O\left(\frac{1}{\sqrt{\log n}}\right) \quad (16)$$

Finally, Theorem 5, together with Equations (15) and (16), yields:

$$\frac{\sum_{h \in E} w_h \bar{\delta}_h(\mathbf{x})}{\min_{\gamma \in \mathbb{R}} \|\mathbf{x} - \gamma \mathbf{1}\|_{1, \mu}} \leq 2 \cdot \frac{\sum_{h \in E} w_h \min_{\nu \in \mathbb{R}} \|\mathbf{x} - \nu \mathbf{1}\|_{F_h}}{\min_{\gamma \in \mathbb{R}} \|\mathbf{x} - \gamma \mathbf{1}\|_{1, \mu}} \leq \text{opt} \cdot O(\sqrt{\log n}).$$

The vector \mathbf{x} can then be efficiently rounded to a subset $S \subseteq V$ with:

$$\Psi_G(S) \leq \frac{\sum_{h \in E} w_h \bar{\delta}_h(\mathbf{x})}{\min_{\gamma \in \mathbb{R}} \|\mathbf{x} - \gamma \mathbf{1}\|_{1, \mu}},$$

as per Lemma 1. □

7 Acknowledgements

AC is supported by NSF DGE 1746045. AC would like to thank Luca Trevisan for his support while visiting Bocconi University; part of this work was completed while AC was a visiting student at Bocconi. LO is supported by NSF CAREER 1943510. The authors would like to thank Jeffrey Negrea for assisting on topics related to online learning.

References

- [1] Sameer Agarwal, Kristin Branson, and Serge Belongie. Higher order learning with graphs. In *Proceedings of the 23rd International Conference on Machine Learning, ICML '06*, pages 17–24, New York, NY, USA, June 2006. Association for Computing Machinery.
- [2] Ahuja, Ravindra K and Magnanti, Thomas L and Orlin, James B. *Network Flows: Theory, Algorithms, and Applications*. Prentice Hall, 1993.
- [3] Zeyuan Allen-Zhu, Zhenyu Liao, and Lorenzo Orecchia. Spectral sparsification and regret minimization beyond matrix multiplicative updates. In *Proceedings of the forty-seventh annual ACM symposium on Theory of computing*, pages 237–245, 2015.
- [4] Noga Alon and Vitali D Milman. Eigenvalues, expanders and superconcentrators. In *25th Annual Symposium on Foundations of Computer Science, 1984.*, pages 320–322. IEEE, 1984.
- [5] Reid Andersen and Kevin J. Lang. An algorithm for improving graph partitions. In *Proceedings of the Nineteenth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA '08*, page 651–660, USA, 2008. Society for Industrial and Applied Mathematics.
- [6] Sanjeev Arora, Elad Hazan, and Satyen Kale. The multiplicative weights update method: a meta-algorithm and applications. *Theory of computing*, 8(1):121–164, 2012.
- [7] Sanjeev Arora, Satish Rao, and Umesh Vazirani. Expander flows, geometric embeddings and graph partitioning. *Journal of the ACM*, 56(2):5:1–5:37, April 2009.
- [8] Kyriakos Axiotis, Adam Karczmarz, Anish Mukherjee, Piotr Sankowski, and Adrian Vladu. Decomposable Submodular Function Minimization via Maximum Flow. In *Proceedings of the 38th International Conference on Machine Learning*, pages 446–456. PMLR, July 2021.
- [9] Francis Bach. Learning with Submodular Functions: A Convex Optimization Perspective, October 2013.
- [10] Wenruo Bai, Rishabh Iyer, Kai Wei, and Jeff Bilmes. Algorithms for Optimizing the Ratio of Submodular Functions. In *Proceedings of The 33rd International Conference on Machine Learning*, pages 2751–2759. PMLR, June 2016.
- [11] F. L. Bauer, J. Stoer, and C. Witzgall. Absolute and monotonic norms. *Numerische Mathematik*, 3(1):257–264, December 1961.
- [12] Aaron Bernstein, Maximilian Probst Gutenberg, and Thatchaphol Saranurak. Deterministic decremental sssp and approximate min-cost flow in almost-linear time. In *2021 IEEE 62nd Annual Symposium on Foundations of Computer Science (FOCS)*, pages 1000–1008. IEEE, 2022.
- [13] U.V. Catalyurek and C. Aykanat. Hypergraph-partitioning-based decomposition for parallel sparse-matrix vector multiplication. *IEEE Transactions on Parallel and Distributed Systems*, 10(7):673–693, July 1999.
- [14] T.-H. Hubert Chan, Anand Louis, Zhihao Gavin Tang, and Chenzi Zhang. Spectral Properties of Hypergraph Laplacian and Approximation Algorithms. *arXiv:1605.01483 [cs]*, May 2016.
- [15] T-H Hubert Chan, Anand Louis, Zhihao Gavin Tang, and Chenzi Zhang. Spectral properties of hypergraph laplacian and approximation algorithms. *Journal of the ACM (JACM)*, 65(3):1–48, 2018.
- [16] Chandra Chekuri and Alina Ene. Poly-logarithmic approximation for maximum node disjoint paths with constant congestion. In *Proceedings of the twenty-fourth annual ACM-SIAM symposium on Discrete algorithms*, pages 326–341. SIAM, 2013.

- [17] Chandra Chekuri, Alina Ene, and Marcin Pilipczuk. Constant congestion routing of symmetric demands in planar directed graphs. *SIAM Journal on Discrete Mathematics*, 32(3):2134–2160, 2018.
- [18] Li Chen, Rasmus Kyng, Yang P. Liu, Richard Peng, Maximilian Probst Gutenberg, and Sushant Sachdeva. Maximum Flow and Minimum-Cost Flow in Almost-Linear Time, April 2022.
- [19] Li Chen, Rasmus Kyng, Yang P. Liu, Richard Peng, Maximilian Probst Gutenberg, and Sushant Sachdeva. Maximum Flow and Minimum-Cost Flow in Almost-Linear Time, April 2022.
- [20] Julia Chuzhoy. Routing in undirected graphs with constant congestion. In *Proceedings of the forty-fourth annual ACM symposium on Theory of computing*, pages 855–874, 2012.
- [21] Julia Chuzhoy, Yu Gao, Jason Li, Danupon Nanongkai, Richard Peng, and Thatchaphol Saranurak. A deterministic algorithm for balanced cut with applications to dynamic connectivity, flows, and beyond. In *2020 IEEE 61st Annual Symposium on Foundations of Computer Science (FOCS)*, pages 1158–1167. IEEE, 2020.
- [22] Julia Chuzhoy and Sanjeev Khanna. A new algorithm for decremental single-source shortest paths with applications to vertex-capacitated flow and cut problems. In *Proceedings of the 51st Annual ACM SIGACT Symposium on Theory of Computing*, pages 389–400, 2019.
- [23] K.D. Devine, E.G. Boman, R.T. Heaphy, R.H. Bisseling, and U.V. Catalyurek. Parallel hypergraph partitioning for scientific computing. In *Proceedings 20th IEEE International Parallel & Distributed Processing Symposium*, pages 10 pp.–, April 2006.
- [24] Alina Ene, Huy L. Nguyen, and László A. Végh. Decomposable Submodular Function Minimization: Discrete and Continuous, March 2017.
- [25] Uriel Feige, MohammadTaghi Hajiaghayi, and James R Lee. Improved approximation algorithms for minimum-weight vertex separators. In *Proceedings of the thirty-seventh annual ACM symposium on Theory of computing*, pages 563–572, 2005.
- [26] Elad Hazan et al. Introduction to online convex optimization. *Foundations and Trends[®] in Optimization*, 2(3-4):157–325, 2016.
- [27] Shohei Hidaka and Masafumi Oizumi. Fast and exact search for the partition with minimal information loss. *PLOS ONE*, 13(9):e0201126, September 2018.
- [28] Rania Ibrahim and David F. Gleich. Local hypergraph clustering using capacity releasing diffusion. *PLOS ONE*, 15(12):e0243485, December 2020.
- [29] Arun Jambulapati, Yin Tat Lee, Jerry Li, Swati Padmanabhan, and Kevin Tian. Positive Semidefinite Programming: Mixed, Parallel, and Width-Independent, July 2021.
- [30] Haotian Jiang, Tarun Kathuria, Yin Tat Lee, Swati Padmanabhan, and Zhao Song. A Faster Interior Point Method for Semidefinite Programming. In *2020 IEEE 61st Annual Symposium on Foundations of Computer Science (FOCS)*, pages 910–918, November 2020.
- [31] Rohit Khandekar, Subhash Khot, Lorenzo Orecchia, and Nisheeth K Vishnoi. On a cut-matching game for the sparsest cut problem. *Univ. California, Berkeley, CA, USA, Tech. Rep. UCB/EECS-2007-177*, 6(7):12, 2007.
- [32] Rohit Khandekar, Satish Rao, and Umesh Vazirani. Graph partitioning using single commodity flows. *Journal of the ACM*, 56(4):19:1–19:15, July 2009.
- [33] Kevin J. Lang, Michael W. Mahoney, and Lorenzo Orecchia. Empirical Evaluation of Graph Partitioning Using Spectral Embeddings and Flow. In Jan Vahrenhold, editor, *Experimental Algorithms*, Lecture Notes in Computer Science, pages 197–208, Berlin, Heidelberg, 2009. Springer.

- [34] Tom Leighton and Satish Rao. Multicommodity max-flow min-cut theorems and their use in designing approximation algorithms. *Journal of the ACM*, 46(6):787–832, November 1999.
- [35] Pan Li and Olgica Milenkovic. Inhomogeneous Hypergraph Clustering with Applications. In *Advances in Neural Information Processing Systems*, volume 30. Curran Associates, Inc., 2017.
- [36] Pan Li and Olgica Milenkovic. Submodular Hypergraphs: P-Laplacians, Cheeger Inequalities and Spectral Clustering. In *Proceedings of the 35th International Conference on Machine Learning*, pages 3014–3023. PMLR, July 2018.
- [37] Pan Li and Olgica Milenkovic. Submodular hypergraphs: p-laplacians, cheeger inequalities and spectral clustering. In *International Conference on Machine Learning*, pages 3014–3023. PMLR, 2018.
- [38] Anand Louis. Cut-matching games on directed graphs. *arXiv preprint arXiv:1010.1047*, 2010.
- [39] Anand Louis. Hypergraph Markov Operators, Eigenvalues and Approximation Algorithms. In *Proceedings of the Forty-Seventh Annual ACM Symposium on Theory of Computing*, STOC '15, pages 713–722, New York, NY, USA, June 2015. Association for Computing Machinery.
- [40] Anand Louis. Hypergraph markov operators, eigenvalues and approximation algorithms. In *Proceedings of the forty-seventh annual ACM symposium on Theory of computing*, pages 713–722, 2015.
- [41] Anand Louis and Yury Makarychev. Approximation Algorithms for Hypergraph Small-Set Expansion and Small-Set Vertex Expansion. *Theory of Computing*, 12(17):1–25, October 2016.
- [42] Anand Louis, Prasad Raghavendra, and Santosh Vempala. The Complexity of Approximating Vertex Expansion. *arXiv:1304.3139 [cs]*, November 2013.
- [43] David W. Matula and Farhad Shahrokhi. Sparsest cuts and bottlenecks in graphs. *Discrete Applied Mathematics*, 27(1):113–123, May 1990.
- [44] Mukund Narasimhan, Nebojsa Jojic, and Jeff A Bilmes. Q-Clustering. In *Advances in Neural Information Processing Systems*, volume 18. MIT Press, 2005.
- [45] Lorenzo Orecchia. *Fast Approximation Algorithms for Graph Partitioning using Spectral and Semidefinite-Programming Techniques*. PhD thesis, EECS Department, University of California, Berkeley, May 2011.
- [46] Lorenzo Orecchia, Konstantinos Ameranis, Charalampos Tsourakakis, and Kunal Talwar. Practical almost-linear-time approximation algorithms for hybrid and overlapping graph clustering. In *International Conference on Machine Learning*, pages 17071–17093. PMLR, 2022.
- [47] Lorenzo Orecchia, Konstantinos Ameranis, Charalampos Tsourakakis, and Kunal Talwar. Practical Almost-Linear-Time Approximation Algorithms for Hybrid and Overlapping Graph Clustering. In *Proceedings of the 39th International Conference on Machine Learning*, pages 17071–17093. PMLR, June 2022.
- [48] Lorenzo Orecchia, Sushant Sachdeva, and Nisheeth K Vishnoi. Approximating the exponential, the lanczos method and an $o(m)$ -time spectral algorithm for balanced separator. In *Proceedings of the forty-fourth annual ACM symposium on Theory of computing*, pages 1141–1160, 2012.
- [49] Lorenzo Orecchia, Leonard J. Schulman, Umesh V. Vazirani, and Nisheeth K. Vishnoi. On partitioning graphs via single commodity flows. In *Proceedings of the Fortieth Annual ACM Symposium on Theory of Computing*, STOC '08, pages 461–470, New York, NY, USA, May 2008. Association for Computing Machinery.
- [50] Thatchaphol Saranurak and Di Wang. Expander Decomposition and Pruning: Faster, Stronger, and Simpler, December 2021.

- [51] Jonah Sherman. Breaking the Multicommodity Flow Barrier for $\sqrt{\log(n)}$ -Approximations to Sparsest Cut, August 2009.
- [52] Alistair Sinclair and Mark Jerrum. Approximate counting, uniform generation and rapidly mixing Markov chains. *Information and Computation*, 82(1):93–133, July 1989.
- [53] Daniel A. Spielman and Shang-Hua Teng. Nearly-linear time algorithms for graph partitioning, graph sparsification, and solving linear systems. In *Proceedings of the Thirty-Sixth Annual ACM Symposium on Theory of Computing*, STOC '04, pages 81–90, New York, NY, USA, June 2004. Association for Computing Machinery.
- [54] Daniel A Spielman and Shang-Hua Teng. Nearly-linear time algorithms for graph partitioning, graph sparsification, and solving linear systems. In *Proceedings of the thirty-sixth annual ACM symposium on Theory of computing*, pages 81–90, 2004.
- [55] Nate Veldt, Austin R. Benson, and Jon Kleinberg. Minimizing Localized Ratio Cut Objectives in Hypergraphs. In *Proceedings of the 26th ACM SIGKDD International Conference on Knowledge Discovery & Data Mining*, KDD '20, pages 1708–1718, New York, NY, USA, August 2020. Association for Computing Machinery.
- [56] Nate Veldt, Austin R. Benson, and Jon Kleinberg. Approximate Decomposable Submodular Function Minimization for Cardinality-Based Components, October 2021.
- [57] J.Y. Zien, M.D.F. Schlag, and P.K. Chan. Multilevel spectral hypergraph partitioning with arbitrary vertex sizes. *IEEE Transactions on Computer-Aided Design of Integrated Circuits and Systems*, 18(9):1389–1399, September 1999.

A An Optimization View on Submodular Hypergraph Partitioning

In this section, we prove some technical lemmata used in the body of the paper. We begin with a proof of Fact 7.

Proof of Fact 7. Symmetry of δ follows from definition. To show that δ is submodular, let $S, T \subseteq V$. We have one of two cases:

1. $F(S) \leq F(\bar{S})$ and $F(T) \leq F(\bar{T})$, or $F(\bar{S}) \leq F(S)$ and $F(\bar{T}) \leq F(T)$,
2. $F(S) \leq F(\bar{S})$ and $F(\bar{T}) \leq F(T)$, or $F(\bar{S}) \leq F(S)$ and $F(T) \leq F(\bar{T})$.

For Case 1, we may assume without loss of generality that $F(S) \leq F(\bar{S})$ and $F(T) \leq F(\bar{T})$. We then have:

$$\begin{aligned} \delta(S \cup T) + \delta(S \cap T) &= \min\{F(S \cup T), F(\bar{S} \cap \bar{T})\} + \min\{F(S \cap T), F(\bar{S} \cup \bar{T})\} \\ &\leq F(S \cup T) + F(S \cap T) \\ &\leq F(S) + F(T) = \delta(S) + \delta(T). \end{aligned}$$

Similarly, for Case 2, we may assume $F(S) \leq F(\bar{S})$ and $F(\bar{T}) \leq F(T)$. We then have:

$$\begin{aligned} \delta(S \cup T) + \delta(S \cap T) &= \min\{F(S \cup T), F(\bar{S} \cap \bar{T})\} + \min\{F(S \cap T), F(\bar{S} \cup \bar{T})\} \\ &\leq F(S \cup T) + F(\bar{S} \cup \bar{T}) \\ &\leq F(S) + F(\bar{T}) = \delta(S) + \delta(T). \end{aligned}$$

Showing that for every $S, T \subseteq V$:

$$\delta(S \cup T) + \delta(S \cap T) \leq \delta(S) + \delta(T),$$

and hence δ is submodular as needed. \square

Let us next show that the ratio-cut improvement problem is equivalent to (RC-NonCvx) introduced in Section 4. We first require that the Lovász extension is unaffected by a shift of the input by $\mathbf{1}$.

Lemma 15. *Suppose $\delta_h : 2^h \rightarrow \mathbb{R}$ is a monotone submodular cut function. For any $u \in \mathbb{R}$,*

$$\bar{\delta}_h(\mathbf{x} + u \cdot \mathbf{1}) = \bar{\delta}_h(\mathbf{x}).$$

Proof. Since δ_h is a monotone submodular cut function, there exists a non-decreasing submodular function $F_h : 2^h \rightarrow \mathbb{R}$ such that $F_h(\emptyset) = 0$ and $\delta_h(S) = \min\{F_h(S), F_h(\bar{S})\}$. Observe then that if $\mathbf{y} \in \mathcal{B}(\delta_h)$ is in the base polytope, we have

$$\sum_{i \in h} y_i \leq \delta_h(h) = \min\{F_h(h), F_h(\emptyset)\} = 0.$$

This implies that for any $u \in \mathbb{R}$, the Lovász extension of δ_h satisfies:

$$\bar{\delta}_h(\mathbf{x} + u \cdot \mathbf{1}) = \max_{\mathbf{y} \in \mathcal{B}(\delta_h)} \langle \mathbf{y}, \mathbf{x} + u \cdot \mathbf{1} \rangle = \max_{\mathbf{y} \in \mathcal{B}(\delta_h)} \langle \mathbf{y}, \mathbf{x} \rangle = \bar{\delta}_h(\mathbf{x}),$$

since $\mathbf{y} \in \mathcal{B}(\delta_h)$ implies $\langle \mathbf{y}, \mathbf{1} \rangle = 0$. \square

Proof of Lemma 1. Observe that the denominator of the ratio-cut objective is a monotone submodular cut function applied over the entire vertex set

$$\delta_{\boldsymbol{\mu}}(S) = \min\{\boldsymbol{\mu}(S), \boldsymbol{\mu}(\bar{S})\},$$

Consequently, for every $S \subseteq V$:

$$\Psi_G(S) = \frac{\sum_{h \in E} w_h \delta_h(S \cap h)}{\min\{\boldsymbol{\mu}(S), \bar{\boldsymbol{\mu}}(S)\}} = \frac{\delta_G(S)}{\delta_{\boldsymbol{\mu}}(S)} = \frac{\bar{\delta}_G(\mathbf{1}^S)}{\bar{\delta}_{\boldsymbol{\mu}}(\mathbf{1}^S)},$$

which gives:

$$\Psi_G^* \geq \min_{\mathbf{x} \in \mathbb{R}^n} \frac{\bar{\delta}_G(\mathbf{x})}{\bar{\delta}_{\boldsymbol{\mu}}(\mathbf{x})}.$$

We now prove the opposite inequality by showing the second part of the lemma. By Lemma 15, the value of the ratio in the right-hand side of (RC-NonCvx) is invariant under shifting all the coordinates by a fixed constant. Furthermore, both the numerator and the denominator of the right-hand side of (RC-NonCvx) are homogeneous and hence their ratio is invariant under scaling \mathbf{x} by a constant, i.e., for any $u \in \mathbb{R}$:

$$\frac{\bar{\delta}_G(u \cdot \mathbf{x})}{\bar{\delta}_{\boldsymbol{\mu}}(u \cdot \mathbf{x})} = \frac{u \cdot \bar{\delta}_G(\mathbf{x})}{u \cdot \bar{\delta}_{\boldsymbol{\mu}}(\mathbf{x})} = \frac{\bar{\delta}_G(\mathbf{x})}{\bar{\delta}_{\boldsymbol{\mu}}(\mathbf{x})}.$$

Hence, for any \mathbf{x} , by letting $\hat{\mathbf{x}}$ be defined as:

$$\hat{x}_i = \frac{x_i - \min_j x_j}{\max_{j,k} (x_j - x_k)},$$

we have $\max_i \hat{x}_i = 1$, $\min_i \hat{x}_i = 0$, and:

$$\frac{\bar{\delta}_G(\mathbf{x})}{\bar{\delta}_{\boldsymbol{\mu}}(\mathbf{x})} = \frac{\bar{\delta}_G(\hat{\mathbf{x}})}{\bar{\delta}_{\boldsymbol{\mu}}(\hat{\mathbf{x}})}.$$

We then have, as per Fact 4:

$$\frac{\bar{\delta}_G(\mathbf{x})}{\bar{\delta}_{\boldsymbol{\mu}}(\mathbf{x})} = \frac{\bar{\delta}_G(\hat{\mathbf{x}})}{\bar{\delta}_{\boldsymbol{\mu}}(\hat{\mathbf{x}})} = \frac{\mathbb{E}_{t \sim [0,1]}[\delta_G(S_t(\hat{\mathbf{x}}))]}{\mathbb{E}_{t \sim [0,1]}[\delta_{\boldsymbol{\mu}}(S_t(\hat{\mathbf{x}}))]}.$$

Hence, there must exist some $t \in [0, 1]$ such that:

$$\frac{\bar{\delta}_G(\mathbf{x})}{\bar{\delta}_{\boldsymbol{\mu}}(\mathbf{x})} \geq \frac{\delta_G(S_t(\hat{\mathbf{x}}))}{\delta_{\boldsymbol{\mu}}(S_t(\hat{\mathbf{x}}))}.$$

This gives:

$$\min_{\mathbf{x} \in \mathbb{R}^n} \frac{\bar{\delta}_G(\mathbf{x})}{\bar{\delta}_{\boldsymbol{\mu}}(\mathbf{x})} = \Psi_G^*,$$

as needed for the first part of the lemma. Moreover, computing and checking all of the sets $\{S_t(\mathbf{x})\}_{t \in [0,1]}$ only requires time $O(n \log n)$ for sorting the entries of \mathbf{x} and time $O(\text{sp}(G) + |V|)$ for evaluating the numerator and denominator. \square

A critical step towards deploying the cut-matching game was to restrict our attention to monotone submodular cut functions, for which Theorem 5 provides a specific geometric characterization. Towards proving this lemma, we first demonstrate that the base polytope of a monotone submodular cut function δ_h is enclosed within a scaling of the symmetric submodular polyhedron of F .

Lemma 16. *Suppose $\delta_h(S) \stackrel{\text{def}}{=} \min\{F(S), F(\bar{S})\}$ where $F : 2^h \rightarrow \mathbb{R}$ is a non-decreasing submodular function such that $F(\emptyset) = 0$ and $F(\{i\}) > 0$ for all $i \in V$. Let $\mathbf{y} \in \mathcal{B}(\delta_h)$. For all $A \subseteq h$, we have*

$$\left| \sum_{i \in A} y_i \right| \leq \delta_h(A)$$

Proof. Observe that since $\mathbf{y} \in \mathcal{B}(\delta_h)$, we immediately have

$$\sum_{i \in A} y_i \leq \delta_h(A)$$

On the other hand, since $\sum_{i \in h} y_i = \delta_h(h) = 0$, we have

$$\sum_{i \in A} y_i = - \sum_{i \in \bar{A}} y_i \geq -\delta_h(\bar{A}) = -\delta_h(A)$$

as required. \square

We are now ready to prove Theorem 5.

Proof of Theorem 5. First, suppose $\mathbf{y} \in \mathcal{P}_{\text{sym}}(F)$, and fix any $S \subseteq h$. Note that:

$$\sum_{i \in S} y_i \leq \sum_{i \in S} |y_i| \leq F(S).$$

Simultaneously, since $\langle \mathbf{y}, \mathbf{1} \rangle = 0$, we have:

$$\sum_{i \in S} y_i = - \sum_{i \in \bar{S}} y_i \leq \sum_{i \in \bar{S}} |y_i| \leq F(\bar{S}).$$

Consequently, we have $\sum_{i \in S} y_i \leq \min\{F(S), F(\bar{S})\} = \delta_h(S)$, giving $\mathcal{P}_{\text{sym}}(F) \subseteq \mathcal{B}(\delta_h)$.

Now, suppose $\mathbf{y} \in \mathcal{B}(\delta_h)$, and fix any $S \subseteq h$. Let $S^+ = \{i \in S : y_i \geq 0\}$, and $S^- = \{i \in S : y_i < 0\}$. Observe

$$\sum_{i \in S} |y_i| = \sum_{i \in S^+} y_i - \sum_{i \in S^-} y_i \leq \left| \sum_{i \in S^+} y_i \right| + \left| \sum_{i \in S^-} y_i \right| \leq \delta_h(S^+) + \delta_h(S^-),$$

where the last inequality follows by Lemma 16. However, this means:

$$\begin{aligned} \delta_h(S^+) + \delta_h(S^-) &= \min\{F(S^+), F(\bar{S}^+)\} + \min\{F(S^-), F(\bar{S}^-)\} \\ &\leq F(S^+) + F(S^-) \\ &\leq 2 \cdot F(S), \end{aligned}$$

since both $S^+, S^- \subseteq S$ and F is non-decreasing. Consequently, $\sum_{i \in S} |y_i| \leq 2 \cdot F(S)$ allowing us to conclude $\mathcal{B}(\delta_h) \subseteq \mathcal{P}_{\text{sym}}(2 \cdot F)$ as required. The lowerbound to the Lovász extension $\bar{\delta}_h(\mathbf{x}) = \max_{\mathbf{y} \in \mathcal{B}(\delta_h)} \langle \mathbf{y}, \mathbf{x} \rangle$ is given by $\mathcal{P}_{\text{sym}}(F) \subseteq \mathcal{B}(\delta_h)$. On the other hand, $\mathcal{B}(\delta_h) \subseteq \mathcal{P}_{\text{sym}}(2 \cdot F)$ implies the upperbound:

$$\max_{\mathbf{y} \in \mathcal{B}(\delta_h)} \langle \mathbf{y}, \mathbf{x} \rangle \leq \max_{\substack{\mathbf{y} \in \mathcal{P}_{\text{sym}}(2 \cdot F) \\ \langle \mathbf{y}, \mathbf{1} \rangle = 0}} \langle \mathbf{y}, \mathbf{x} \rangle = \max_{\substack{\mathbf{y} \in \mathcal{P}_{\text{sym}}(F) \\ \langle \mathbf{y}, \mathbf{1} \rangle = 0}} \langle \mathbf{y}, \mathbf{x} \rangle = 2 \cdot \max_{\substack{\mathbf{y} \in \mathcal{P}_{\text{sym}}(F) \\ \langle \mathbf{y}, \mathbf{1} \rangle = 0}} \langle \mathbf{y}, \mathbf{x} \rangle,$$

as required. \square

We now prove Theorem 7. This requires the notion of factor graph introduced in Section 2.

Proof Theorem 7. Consider the hypergraph flow as a flow over the factor graph \hat{G} . By a standard application of dynamic trees (see for example Lemma 3.8 in [51]) to the computation of flow-path decompositions [2] over \hat{G} , we can compute, in time $\tilde{O}(|\hat{E}_G|) = \tilde{O}(\text{sp}(G))$, a graph $H = (V, E_H, \mathbf{w}^H)$ such that there exist hypergraph flows $\{\mathbf{Y}^e\}_{e \in E_H}$ in G with the following properties:

- $\sum_{e \in E_H} \mathbf{Y}^e = \mathbf{Y}$,

- For any $e \in E_H$, for any factor graph edge $\{i, h\}$, the flow $(\mathbf{Y}_h^e)_i$ has the same sign as $(\mathbf{Y}_h)_i$. In particular:

$$\sum_{e \in E_H} |\mathbf{Y}^e| = |\mathbf{Y}| \quad (17)$$

- For all $e = \{i, j\} \in E^H$, we have $\text{dem}_i(\mathbf{Y}) \geq 0$ and $\text{dem}_j(\mathbf{Y}) < 0$. The flow \mathbf{Y}^e routes w_e^H units of flow between vertex i and vertex j , i.e.,

$$\text{dem}(\mathbf{Y}^e) = w_e^H (\mathbf{1}_i - \mathbf{1}_j).$$

The last point proves the required statements about the bipartiteness of H and its degrees. To complete the proof, we verify that $H \preceq_{\text{cong}_G(Y)} G$. For every hyperedge $h \in E_G$, we have by Equation 17 and the absoluteness of $\|\cdot\|_{*,F_h}$:

$$\left\| \sum_{e \in E_H} |\mathbf{y}_h^e| \right\|_{*,F_h} = \|\mathbf{y}_h\|_{*,F_h} = \|\mathbf{y}_h\|_{*,F_h} = w_h^G \text{cong}_h(\mathbf{Y}) \leq w_h^G \text{cong}_G(\mathbf{Y}).$$

□

B Local Relaxations of Minimum Ratio-cut Over Hypergraphs

B.1 Comparison to the Cut Improvement of Andersen & Lang

In this section, we briefly show that our formulation of the ratio-cut improvement problem in Definition 7 generalizes the definition of the modified quotient cut score, introduced by Andersen and Lang [5], to potentially non-integral seed vectors (rather than cut seeds) and to arbitrary hypergraphs (rather than graphs). Consider the following seed vector. Let $A \subseteq V$ be a cut, and define the vector $\mathbf{s}^{A,\bar{A}} \in \mathbb{R}^V$ to be

$$\mathbf{s}^{A,\bar{A}} \stackrel{\text{def}}{=} \mathbf{1}^A - \mu(A)/\mu(\bar{A}) \cdot \mathbf{1}^{\bar{A}}. \quad (18)$$

Assuming $\mu(A) \leq \mu(\bar{A})$, note that $\langle \mathbf{s}^{A,\bar{A}}, \mathbf{1} \rangle_\mu = 0$ and $\|\mathbf{s}^{A,\bar{A}}\|_\infty \leq 1$, so that $\mathbf{s}^{A,\bar{A}}$ is a valid seed vector. With this choice of seed vector, the ratio-cut improvement objective on a graph G becomes:

$$\Psi_{G,\mathbf{s}^{A,\bar{A}}}(S) = \frac{\sum_{e \in E} w_e \cdot \delta_e(S)}{\left| \mu(A \cap S) - \frac{\mu(A)}{\mu(\bar{A})} \cdot \mu(\bar{A} \cap S) \right|},$$

which is equal to the *modified quotient cut score* considered by Andersen and Lang.

B.2 Dual to the ratio-cut improvement problem

In this section we prove Lemma 4.

Proof of Lemma 4. Let us proceed with taking the conjugate dual. Starting with (CI-Primal):

$$\begin{aligned} \min_{\substack{\mathbf{x} \in \mathbb{R}^V \\ \langle \mathbf{s}, \mathbf{x} \rangle_\mu = 1}} \sum_{h \in E} w_h \cdot \bar{\delta}_h(\mathbf{x}) &= \min_{\mathbf{x} \in \mathbb{R}^V} \sum_{h \in E} w_h \cdot \bar{\delta}_h(\mathbf{x}) + \max_{\alpha \in \mathbb{R}} \left(1 - \langle \mathbf{s}, \mathbf{x} \rangle_\mu \right) \\ &= \min_{\mathbf{x} \in \mathbb{R}^V} \max_{\alpha \in \mathbb{R}} \left\{ \alpha + \sum_{h \in E} w_h \cdot \bar{\delta}_h(\mathbf{x}) - \langle \alpha(\boldsymbol{\mu} \circ \mathbf{s}), \mathbf{x} \rangle \right\}. \end{aligned}$$

Recall that the Lovász extension is given by

$$\bar{\delta}_h(\mathbf{x}) = \max_{\mathbf{f} \in \mathcal{B}(\delta_h)} \langle \mathbf{f}, \mathbf{x} \rangle,$$

and hence:

$$\begin{aligned}
\min_{\mathbf{x} \in \mathbb{R}^V} \max_{\alpha \in \mathbb{R}} \left\{ \alpha + \sum_{h \in E} w_h \cdot \bar{\delta}_h(\mathbf{x}) - \langle \alpha(\boldsymbol{\mu} \circ \mathbf{s}), \mathbf{x} \rangle \right\} &= \min_{\mathbf{x} \in \mathbb{R}^V} \max_{\alpha \in \mathbb{R}} \left\{ \alpha + \sum_{h \in E} \max_{\mathbf{f}_h \in w_h \cdot \mathcal{B}(\delta_h)} \langle \mathbf{f}_h, \mathbf{x} \rangle - \langle \alpha(\boldsymbol{\mu} \circ \mathbf{s}), \mathbf{x} \rangle \right\} \\
&= \min_{\mathbf{x} \in \mathbb{R}^V} \max_{\substack{\alpha \in \mathbb{R}, \mathbf{f}_h \in \mathbb{R}^V \\ \mathbf{f}_h \in w_h \cdot \mathcal{B}(\delta_h)}} \left\{ \alpha + \sum_{h \in E} \langle \mathbf{f}_h, \mathbf{x} \rangle - \langle \alpha(\boldsymbol{\mu} \circ \mathbf{s}), \mathbf{x} \rangle \right\} \\
&= \min_{\mathbf{x} \in \mathbb{R}^V} \max_{\substack{\alpha \in \mathbb{R}, \mathbf{f}_h \in \mathbb{R}^V \\ \mathbf{f}_h \in w_h \cdot \mathcal{B}(\delta_h)}} \left\{ \alpha + \left\langle \sum_{h \in E} \mathbf{f}_h - \alpha(\boldsymbol{\mu} \circ \mathbf{s}), \mathbf{x} \right\rangle \right\}.
\end{aligned}$$

Strong duality always holds when the feasible region is polyhedron. Hence, we have:

$$\min_{\mathbf{x} \in \mathbb{R}^V} \max_{\substack{\alpha \in \mathbb{R}, \mathbf{f}_h \in \mathbb{R}^V \\ \mathbf{f}_h \in w_h \cdot \mathcal{B}(\delta_h)}} \left\{ \alpha + \left\langle \sum_{h \in E} \mathbf{f}_h - \alpha(\boldsymbol{\mu} \circ \mathbf{s}), \mathbf{x} \right\rangle \right\} = \max_{\substack{\alpha \in \mathbb{R}, \mathbf{f}_h \in \mathbb{R}^V \\ \mathbf{f}_h \in w_h \cdot \mathcal{B}(\delta_h)}} \alpha + \min_{\mathbf{x} \in \mathbb{R}^V} \left\langle \sum_{h \in E} \mathbf{f}_h - \alpha(\boldsymbol{\mu} \circ \mathbf{s}), \mathbf{x} \right\rangle,$$

which is equivalent to the required linear program. \square

B.3 Approximate Ratio-Cut Improvement: Proof of Theorem 8

In this section, we address the problem of obtaining an ϵ -approximate primal-dual solution for (G, \mathbf{s}) . This makes up the core implementation of the matching player we use in the algorithm guaranteed by Theorem 9 and Corollary 1. Firstly, we will show in the next section that there are efficient algorithms that allow us to find ϵ primal-dual solutions that work for arbitrary choices of δ_h , leveraging recent results by Axiotis *et al.* [8] and Ene *et al.* [24].

Then, in the section that follows, we will show that when the objective functions δ_h are all standard hypergraph cut functions (i.e. $\delta_h = \delta_h^{\text{cut}}$ for every $h \in E$), the problem can be solved in almost-linear time. We do so by solving a sequence of mixed capacitated maximum flow problems. These are simply maximum flow problems over an undirected graph in which both edges and vertices can have capacity constraints. Our algorithm leverages the fact that one can find an approximate solution to this type of problem in almost-linear time following a recent result of Bernstein *et al.* [12].

B.3.1 Solvers for Arbitrary Monotone Submodular Cut Functions

In this section, we discuss how to obtain an ϵ -approximate primal-dual solution, given input (G, \mathbf{s}) where G is equipped with an arbitrary MSCF $\{\delta_h\}_{h \in E}$.

The first step is to reduce the problem from optimization to feasibility via binary search. Let $\alpha^* = \Psi_{G, \mathbf{s}}^*$ be the optimal value of (CI-Primal). In the feasibility version, for a fixed $\alpha \geq 0$, we consider the problem:

$$\exists \mathbf{x} : \sum_{h \in E} w_h \cdot \bar{\delta}_h(\mathbf{x}) - \alpha \cdot \langle \mathbf{s}, \mathbf{x} \rangle_{\mu} < 0 \tag{19}$$

By the properties of the Lovász extension, this is equivalent to considering:

$$\exists S \subseteq V : \sum_{h \in E} w_h \cdot \bar{\delta}_h(\mathbf{1}_S) - \alpha \cdot \langle \mathbf{s}, \mathbf{1}_S \rangle_{\mu} < 0 \tag{(\alpha\text{-CI-Primal-Feasibility})}$$

The following fact is a simple consequence of the evenness of δ_h (Fact 5).

Fact 8. *Problem $(\alpha\text{-CI-Primal-Feasibility})$ is feasible if and only if $\alpha \geq \alpha^*$.*

$$\min_{T \subseteq V} \sum_{h \in E} w_h \cdot \delta_h(T) - \alpha \cdot \langle \mathbf{s}, \mathbf{1}_S \rangle_\mu < 0 \quad (\alpha\text{-CI-Primal})$$

We note that for any given choice of α , the above problem is an instance of a decomposable submodular function minimization problem. That is, it's the minimization of a submodular function which can be written as the sum of simpler submodular functions.

In the case in which $\{\delta_h\}_{h \in E}$ are arbitrary monotone submodular cut functions, we consider two parameter settings: one in which the rank of each hyperedge is bounded by some constant and one in which it is not. For these two settings, we will solve the above two problems exactly using the algorithms of Axiotis *et al.* and Ene *et al.* respectively.

For the algorithm of Ene *et al.* we require access to the following for every $h \in H$:

1. A *value oracle*, which on input a subset $S \subseteq V$ returns $\delta_h(S)$,
2. A *quadratic minimization oracle* $\mathcal{O}_h(\mathbf{w})$, which on input \mathbf{w} returns the solution to the problem:

$$\min_{\mathbf{y} \in \mathcal{B}(\delta_h)} \|\mathbf{y} + \mathbf{w}\|_2^2.$$

The running time of the algorithm of Ene *et al.* is upperbounded by:

$$\mathcal{O} \left(|V| \cdot \sum_{h \in E} \Theta_h \log \left(|V| \cdot \max_h w_h \right) \right),$$

where Θ_h is the running time of a quadratic minimization oracle for the function δ_h . This running time bound follows from Corollary 3.6 in [24] with $\epsilon = \frac{1}{n}$.

In the case of bounded rank, we require a *linear optimization oracle* \mathcal{O}_h for each $h \in E$ which, on input any vector $\mathbf{w} \in \mathbb{R}^V$ supported on $|h|$, finds a set S minimizing $\delta_h(S) - \mathbf{w}(S)$ in time T_h . Given this, we can get an exact solution in time :

$$\tilde{\mathcal{O}} \left(\max_{h \in E} |h|^2 \left(\sum_{h \in E} |h|^2 T_h + T_{\text{flow}} \left(n, n + \sum_{h \in E} |h|^2 \right) \right) \right),$$

where $T_{\text{flow}}(n, m)$ is the time required to solve an instance of the maximum flow problem in a graph with n vertices and m edges. Note that since we are running the above algorithm in the case in which $|h|$ is at most some constant, its running time becomes:

$$\tilde{\mathcal{O}} \left(\sum_{h \in E} T_h + T_{\text{flow}}(n, n + |E|) \right).$$

This result follows from Theorem 1.1 with $\epsilon = \frac{1}{n}$.

In both cases the algorithm outputs both an optimal primal and an optimal dual solution $\text{dem}(\mathbf{Y}) = \alpha \mu \circ \mathbf{s}$ and $\mathbf{Y} \in \bigoplus_{h \in E} \mathbb{R}^h$ to the problem above.

We can then do binary search over the choice of α in the range $[1/\mu(V), \sum_{h \in E} w_h]$, to find a value of α such that the algorithms finds a cut A such that:

$$\Psi_{G, \mathbf{s}}(A) \leq 2\alpha^*,$$

and a hypergraph flow $\mathbf{Y} \in \bigoplus_{h \in E} \mathbb{R}^h$ such that:

$$\left(1 - \frac{\epsilon}{8}\right) \alpha^* \cdot (\mu \circ \mathbf{s}) \leq \text{dem}(\mathbf{Y}) \leq \alpha^* (\mu \circ \mathbf{s}),$$

so that \mathbf{Y} and A make up an ϵ -approximate primal-dual solution. Since the range of possible values for α is $[1/\mu(V), \sum_{h \in E} w_h]$, we can get the above guarantees by running the above algorithms at most:

$$\mathcal{O}(\log |V| + \log \mu(V) + \max_h \log w_h)$$

times.




Figure 2: The figure above shows the interpretation of a vector \mathbf{y}_h in the (CI-Dual- α -Feasibility) problem as a flow in a star graph. On the left side, we drew a hyperedge h of G and the values of $\mathbf{y}_h(i)$ for $i \in h$, and on the right side we draw its interpretation as a flow through a star graph which has replaced the hyperedge. Note how the amount of flow entering the newly-introduced factor vertex equals the amount of flow exiting it.

B.3.2 A Fast Algorithm for the Standard Cut Function

In this subsection, we focus on finding an ϵ -approximate primal-dual solution for the case in which the cut function chosen is the standard hypergraph cut function, i.e. $\delta_h = \delta_h^{\text{cut}}$ for all $h \in E$. We will do this by constructing a (2-uniform) flow graph G_{flow} and running on it an approximate maximum flow algorithm, and then turning the output of this algorithm into an ϵ -approximate primal-dual solution for (G, \mathbf{s}) .

Under this assumption, consider the following feasibility problem parametrized by α corresponding to (CI-Dual):

$$\begin{aligned}
 \text{Find } \mathbf{Y} = \{\mathbf{y}_h\}_{h \in \mathbb{R}^h} &\in \bigoplus_{h \in E} \mathbb{R}^h \\
 \text{s.t. } \sum_{h \in E} \mathbf{y}_h &\leq \alpha \cdot (\boldsymbol{\mu} \circ \mathbf{s}) && \text{(CI-Dual-}\alpha\text{-Feasibility)} \\
 \mathbf{y}_h &\in w_h \cdot \mathcal{B}(\delta_h) && \forall h \in E
 \end{aligned}$$

We will produce an ϵ -approximate primal-dual solution for (G, \mathbf{s}) by finding an approximately feasible point $\mathbf{Y} \in \bigoplus_{h \in E} \mathbb{R}^h$ for the above program for a suitable choice of α . In particular, \mathbf{Y} will satisfy the last constraint exactly and the first constraint approximately.

The problem asks one to find a vector \mathbf{y}_h for each hyperedge of G . The last constraint in the program enforces: $\mathbf{y}_h \in w_h \cdot \mathcal{B}(\delta_h^{\text{cut}})$. Following the definition of δ_h^{cut} and of the base polytope, it is easy to see that the vectors \mathbf{y}_h satisfying this constraint are those vectors such that:

1. $\sum_{i \in h} \mathbf{y}_h(i) = 0$,
2. and $\sum_{i \in h} |\mathbf{y}_h(i)| \leq 2 \cdot w_h$.

This allows us to interpret each $\mathbf{y}_h(i)$ as an amount of flow that is leaving vertex i through hyperedge h , and the constraints ensure that the total flow going through any hyperedge is not dissipated (item 2 above) and is at most w_h (item 3 above).

In this way, \mathbf{Y} corresponds to a flow routed in the factor graph \hat{G} of G (See Figure 2, and the Preliminaries Section discussion on hypergraph flows). Under this interpretation, the first constraint in CI-Dual- α -Feasibility asks one to route a specific demand in the factor graph, equal to $\alpha \cdot \boldsymbol{\mu}_i \cdot \mathbf{s}_i$ for every $i \in V$, while the second constraint imposes flow conservation constraints and vertex-capacity constraints on the factor vertices. We will actually not enforce that the exact demand imposed by the first constraint is met, but rather that we route a sufficiently high amount of demand to meet the requirements of an ϵ -approximate

primal-dual solution. In this setting, the above problem can be cast as a mixed edge-vertex capacitated single-commodity maximum flow problem in a graph G_{flow} purpose-built starting from the factor graph \hat{G} to enforce the above constraints.

Armed with this perspective, we now give an algorithm (Algorithm 5 below) which on input (G, \mathbf{s}) , constructs a flow graph G_{flow} which can be used to find an ϵ -approximate primal-dual solution (A, \mathbf{Y}) for (G, \mathbf{s}) by applying the algorithm of Bernstein *et al.* [12]

Algorithm 5 Flow Graph Construction

Parameters: $\alpha > 0$, $G = (V, E, \mathbf{w}, \boldsymbol{\mu})$, and $\mathbf{s} \in \mathbb{R}^V$ satisfying $\langle \mathbf{s}, \mathbf{1} \rangle_{\boldsymbol{\mu}} = 0$ and $\|\mathbf{s}\|_{\infty} \leq 1$.

Do:

1. Starting with $G_{\text{flow}} = \hat{G} = (V \cup E, \hat{E})$, construct a flow network as follows (see Figure 3):
 - Add source vertex s , and sink vertex t
(Note the distinction between the source vertex s and the seed vector \mathbf{s}).
 - For each $i \in V$ s.t. $\mathbf{s}_i \geq 0$, add an edge between s and i to \hat{E} with capacity $\alpha \boldsymbol{\mu}_i \cdot \mathbf{s}_i$.
(So that for any feasible flow $\mathbf{f} \in \mathbb{R}^{E(G_{\text{flow}})}$, $|\mathbf{f}_{si}| < \alpha \cdot \boldsymbol{\mu}_i \cdot \mathbf{s}_i$).
 - For each $j \in V$ s.t. $\mathbf{s}_j < 0$, add an edge between t and j to \hat{E} with capacity $\alpha \cdot \boldsymbol{\mu}_j \cdot \mathbf{s}_j$
(So that for any feasible flow $\mathbf{f} \in \mathbb{R}^{E(G_{\text{flow}})}$, $|\mathbf{f}_{tj}| < \alpha \cdot \boldsymbol{\mu}_j \cdot \mathbf{s}_j$).
 - For each factor vertex representing $h \in E$, enforce a vertex capacity constraint of w_h .
2. Return G_{flow} .

In order to solve the feasibility problem, Algorithm 5 builds the following flow network G_{flow} . Starting from $\hat{G} = (V \cup E, \hat{E})$ the factor graph representation of G , we add a vertex capacity constraint of value w_h on each factor vertex representing a hyperedge $h \in E$. We then seek to use a single commodity flow to route multiple demands placed on variable vertices $i \in V$ given by the vector $\alpha \cdot \boldsymbol{\mu} \circ \mathbf{s} \in \mathbb{R}^V$. Routing this demand can be done via standard flow reduction techniques which works as follows. Add a source vertex s and sink vertex t to G_{flow} . For every $i \in V$ such that $\mathbf{s}_i \geq 0$, add an undirected edge si with a capacity constraint $\alpha \cdot \boldsymbol{\mu}_i \cdot \mathbf{s}_i$. For every $j \in V$ such that $\mathbf{s}_j < 0$, add an undirected edge jt with a capacity constraint of $\alpha \cdot \boldsymbol{\mu}_j \cdot \mathbf{s}_j$. Finally, route a maximum flow from s to t , and let \mathbf{f}_{ih} to be the amount of flow sent on each edge $ih \in \hat{E}$.

We note that, since we are enforcing vertex constraints, the algorithm involves solving a mixed edge-vertex-capacitated version of the undirected maximum flow problem, and that this version of the problem is solved by [12].

On input a mixed capacitated flow network G_{flow} , we can use the Algorithm of Bernstein *et al.* to route find a flow in the network of value $(1 - \epsilon)f$, as well as an s - t mixed vertex-edge cut of capacity at most $(1 + \epsilon)f$, where f is the value of the maximum flow in G_{flow} . Note that for the case of the G_{flow} returned by Algorithm 5, the mixed vertex-edges cut returned by the algorithm must be comprised entirely of factor vertices (i.e. vertices that arose from hyperedges in G) and edges connecting V to s or t , since these are the only objects with finite capacity.

We argue now that obtaining a $(1 - \epsilon)$ -approximate solution of the kind guaranteed by Bernstein *et al.* , suffices to return an ϵ -approximate primal-dual solution on input (G, \mathbf{s}) .

Note that, when G_{flow} is constructed with a choice of α that is smaller than $\Psi_{G, \mathbf{s}}^*$, the value of the maximum flow is α and the minimum cut is just given by $\{\mathbf{s}\} \subseteq V(G^{\text{flow}})$. On the other hand, when the value of α used to construct G^{flow} is larger than $\Psi_{G, \mathbf{s}}^*$, the value of the maximum flow is at returned by Bernstein *et al.* is at least $(1 - \epsilon)\alpha$.




Figure 3: An illustration of the construction of the maximum flow instance corresponding to (CI-Dual- α -Feasibility). On the left is the original hypergraph G . The middle of the figure shows the factor graph. The maximum flow algorithm is run on the graph G_{flow} on the right. In this graph only the dashed edges (in red) and the white (non-solid) vertices have finite capacity constraints which depend on the values of $\mu_i \cdot s_i$ and the values of w_h respectively. Here, we assume that we are given an \mathbf{s} such that $s_1 > 0$ and $s_2 > 0$, but $s_3 < 0$, $s_4 < 0$ and $s_5 < 0$.

Like we did in the previous section, we will perform binary search for a value of α in the range $[1/\mu(V), \sum_{h \in E} w_h]$, to find a value of α such that the algorithm finds a cut A such that the algorithm approximate maxflow algorithm routes at least $(1 - \epsilon)\eta\alpha\|\mathbf{s}\|_{1,\mu}$ units of flow in G_{flow} , and returns it together with a cut T of a capacity:

$$\text{cap}(T) = (1 + \epsilon)\eta\alpha\frac{\|\mathbf{s}\|_{1,\mu}}{2}$$

for some choice of η and some accuracy parameter ϵ . We then have the following:

Lemma 17. *In the setting described above, we have:*

$$\langle \mathbf{s}, \mathbf{1}_T \rangle_{\mu} \geq \frac{\|\mathbf{s}\|_{1,\mu}}{2} (1 - (1 + \epsilon)\eta).$$

Proof. We have:

$$\alpha((\mu \circ \mathbf{s})(S \cap \bar{T}) - (\mu \circ \mathbf{s})(\bar{S} \cap T)) \leq \frac{(1 + \epsilon)\eta\alpha\|\mathbf{s}\|_{1,\mu}}{2},$$

which gives:

$$((\mu \circ \mathbf{s})(S \cap \bar{T}) - (\mu \circ \mathbf{s})(\bar{S} \cap T)) \leq \frac{(1 + \epsilon)\eta\|\mathbf{s}\|_{1,\mu}}{2},$$

which in turn implies:

$$\frac{\|\mathbf{s}\|_{1,\mu}}{2} - (\mu \circ \mathbf{s})(S \cap T) - (\mu \circ \mathbf{s})(\bar{S} \cap T) \leq \frac{(1 + \epsilon)\eta\|\mathbf{s}\|_{1,\mu}}{2}.$$

From the above, we obtain:

$$\|\mathbf{s}\|_{1,\mu} - \langle \mathbf{s}, \mathbf{1}_T \rangle_{\mu} \leq (1 + \epsilon)\eta\|\mathbf{s}\|_{1,\mu},$$

yielding the wanted result. \square

Lemma 17 implies that:

$$\Psi_G(T \cap V) \leq \frac{\sum_{h \in E} w_h \delta_h^{\text{cut}}(T \cap V)}{|\langle \mathbf{s}, \mathbf{1}_T \rangle_{\mu}|} \leq \frac{\text{cap}(T)}{|\langle \mathbf{s}, \mathbf{1}_T \rangle_{\mu}|} \leq \frac{(1 + \epsilon)\eta\alpha\|\mathbf{s}\|_{1,\mu}}{|\langle \mathbf{s}, \mathbf{1}_T \rangle_{\mu}|} \leq \frac{(1 + \epsilon)\eta\alpha\|\mathbf{s}\|_{1,\mu}}{\|\mathbf{s}\|_{1,\mu} (1 - (1 + \epsilon)\eta)} \leq 2\alpha.$$

for small enough values of η and ϵ . This in particular guarantees that flow and cut pair $(\mathbf{Y}, T \cap V)$ built by taking the output of the algorithm of Bernstein *et al.* on G^{flow} , and restricting it to G is a valid ϵ -approximate primal-dual response.

C An $O(\log n)$ -Approximation via Generalized Cut-Matching Games

In this section, we include additional details from our construction of an $O(\log n)$ -approximation algorithm for minimum ratio-cut based on the generalized cut-matching game.

C.1 Updating the vector embedding via mirror descent

We now describe the operation of Mirror Descent in the context of online linear optimization over the set of density matrices [26]. Using known performance guarantees, we then prove Lemmata 7 and 8 as stated in Section 5.2.2.

Let $\Delta_{n \times n} = \{\mathbf{X} \succeq \mathbf{0} : \text{tr } \mathbf{X} = 1\}$ be the set of $n \times n$ density matrices. Online linear optimization considers the following game played over T rounds. In each round t , a player plays a single action $\mathbf{X}_t \in \Delta_{n \times n}$, receives a feedback matrix $\mathbf{F}_t \in \Delta_{n \times n}$, and incurs a loss of $\langle \mathbf{X}_t, \mathbf{F}_t \rangle$. The regret minimization problem asks for a sequence of actions which minimize the regret: the difference between total loss incurred by the player, and the loss suffered by the best fixed action $\mathbf{U} \in \Delta_{n \times n}$ in hindsight:

$$\min_{\mathbf{X}_1, \dots, \mathbf{X}_T \in \Delta_{n \times n}} \max_{\mathbf{U} \in \Delta_{n \times n}} \left\{ \sum_{t=1}^T \langle \mathbf{X}_t, \mathbf{F}_t \rangle - \sum_{t=1}^T \langle \mathbf{U}, \mathbf{F}_t \rangle \right\}.$$

A standard approach towards minimizing regret is to use Mirror Descent. This corresponds setting

$$\mathbf{X}_{t+1} = \operatorname{argmin}_{\mathbf{X} \in \Delta_{n \times n}} \left\{ D_\psi(\mathbf{X}, \mathbf{X}_t) + \eta \cdot \langle \mathbf{X}, \mathbf{F}_t \rangle \right\}.$$

where $\psi : \Delta_{n \times n} \rightarrow \mathbb{R}$ is a regularization function, that is to be specified, and $D_\psi(\cdot, \cdot)$ is the Bregman Divergence associated with ψ .

$$D_\psi(\mathbf{Z}, \mathbf{X}) \stackrel{\text{def}}{=} \psi(\mathbf{Z}) - (\psi(\mathbf{X}) + \langle \nabla \psi(\mathbf{X}), \mathbf{Z} - \mathbf{X} \rangle).$$

When ψ is taken to be the Von Neumann entropy, i.e. $\psi(\mathbf{X}) = -\langle \mathbf{X}, \log \mathbf{X} \rangle$, \mathbf{X}_{t+1} becomes the following.

$$\mathbf{X}_{t+1} = \frac{\exp(-\eta \sum_{k=1}^t \mathbf{F}_k)}{\text{tr} \exp(-\eta \sum_{k=1}^t \mathbf{F}_k)},$$

This choice of update is also known as the Matrix Multiplicative Weight Update [6], and enjoys the following regret guarantee.

Theorem 14 (Theorem 3.1 from [3]). *Suppose $\mathbf{X}_1, \dots, \mathbf{X}_T \in \Delta_{n \times n}$ are matrices output by update-embedding using the Von Neumann entropy regularizer, and $\mathbf{F}_1, \dots, \mathbf{F}_T$ is a sequence of losses such that:*

$$\mathbf{F}_t \succeq -\frac{1}{\eta} \cdot \mathbf{I}_n$$

for all $t \in [T]$. Then, for any $\mathbf{U} \in \Delta_{n \times n}$, we have:

$$\frac{1}{T} \sum_{t=1}^T \langle \mathbf{F}_t, \mathbf{U} \rangle \geq \frac{1}{T} \sum_{t=1}^T \langle \mathbf{F}_t, \mathbf{X}_t \rangle - \frac{\eta}{T} \sum_{t=1}^T \langle |\mathbf{F}_t|, \mathbf{X}_t \rangle \cdot \|\mathbf{F}_t\|_{\text{spec}} - \frac{\log n}{\eta T}$$

where if $\mathbf{F}_t = \sum_{i=1}^n \lambda_i \cdot \mathbf{v}_i \mathbf{v}_i^\top$, then $|\mathbf{F}_t| = \sum_{i=1}^n |\lambda_i| \cdot \mathbf{v}_i \mathbf{v}_i^\top$.

Using this regret guarantee, we can now prove Lemma 7.

Proof of Lemma 7. Using the regret bound in Theorem 14 with feedback matrices $\mathbf{F}_t = \mathcal{L}_\mu(D_t)$, we have

$$\frac{1}{T} \sum_{t=1}^T \langle \mathcal{L}_\mu(D_t), \mathbf{U} \rangle \geq \frac{1}{T} \sum_{t=1}^T \langle \mathcal{L}_\mu(D_t), \mathbf{X}_t \rangle - \frac{\eta}{T} \sum_{t=1}^T \langle |\mathcal{L}_\mu(D_t)|, \mathbf{X}_t \rangle \cdot \|\mathcal{L}_\mu(D_t)\|_{\text{spec}} - \frac{\log n}{\eta T}$$

Since D_t is an ϵ -approximate matching action, we have that D_t is at most $\mu \circ |\mathbf{s}|$ -regular. Since $\|\mathbf{s}\|_\infty \leq 1$, we have that D_t is at most μ -regular and hence $\mathbf{0} \preceq \mathcal{L}_\mu(D_t) \preceq 2 \cdot \mathbf{I}$. Our regret bound reduces to:

$$\frac{1}{T} \sum_{t=1}^T \langle \mathcal{L}_\mu(D_t), \mathbf{U} \rangle \geq \frac{1-2\eta}{T} \sum_{t=1}^T \langle \mathcal{L}_\mu(D_t), \mathbf{X}_t \rangle - \frac{\log n}{\eta T}.$$

holding for every $\mathbf{U} \in \Delta_{n \times n}$, and $\eta > 0$. Now take $H_T = \frac{1}{T} \sum_{t=1}^T D_t$. We have

$$\langle \mathcal{L}_\mu(H_T), \mathbf{U} \rangle = \left\langle \frac{1}{T} \sum_{t=0}^T \mathcal{L}_\mu(D_t), \mathbf{U} \right\rangle = \frac{1}{T} \sum_{t=0}^T \langle \mathcal{L}_\mu(D_t), \mathbf{U} \rangle \geq \frac{1-2\eta}{T} \sum_{t=1}^T \langle \mathcal{L}_\mu(D_t), \mathbf{X}_t \rangle - \frac{\log n}{T\eta}.$$

Choosing $\mathbf{U} = \mathbf{u}\mathbf{u}^\top$ where \mathbf{u} is the normalized eigenvector of $\mathcal{L}_\mu(H_t)$ corresponding to $\lambda_2(\mathcal{L}_\mu(H_t))$ gives the desired bound. \square

Let us now prove the properties required by Lemma 8 regarding the vector embedding $\{\mathbf{v}_i^{(t)}\}_{i \in V}$ associated with the matrix \mathbf{X}_t generated by `update-embedding`.

Proof of Lemma 8. Let $\mathbf{X}_t = \mathbf{V}_t \mathbf{V}_t^\top$ where the i -th row of $\mathbf{V} \in \mathbb{R}^{n \times n}$ is given by \mathbf{v}_i . To see item (1), observe that $\mathbf{X}_t \in \Delta_{n \times n}$ implies $\text{tr } \mathbf{X}_t = 1$ and hence

$$\sum_{i \in V} \|\mathbf{v}_i\|_2^2 = \text{tr } \mathbf{V}_t \mathbf{V}_t^\top = \text{tr } \mathbf{X}_t = 1$$

To see item (2), observe that

$$\left\| \sum_{i \in V} \sqrt{\mu_i} \cdot \mathbf{v}_i \right\|_2^2 = \left\langle \sum_{i \in V} \sqrt{\mu_i} \cdot \mathbf{v}_i, \sum_{i \in V} \sqrt{\mu_i} \cdot \mathbf{v}_i \right\rangle = \sum_{i, j \in V} \sqrt{\mu_i \mu_j} \cdot \langle \mathbf{v}_i, \mathbf{v}_j \rangle = \langle \mathbf{X}_t, \bar{\boldsymbol{\mu}} \bar{\boldsymbol{\mu}}^\top \rangle$$

where $\bar{\boldsymbol{\mu}}_i = \sqrt{\mu_i}$. Now, since $\bar{\boldsymbol{\mu}} \in \text{kern } \mathcal{L}_\mu(D_t)$ for every t , we have that $\bar{\boldsymbol{\mu}} \in \text{kern } \mathbf{X}_t$ as well. Consequently,

$$\langle \mathbf{X}_t, \bar{\boldsymbol{\mu}} \bar{\boldsymbol{\mu}}^\top \rangle = 0$$

and since $\left\| \sum_{i \in V} \sqrt{\mu_i} \cdot \mathbf{v}_i \right\|_2^2 = 0$, we can conclude $\sum_{i \in V} \sqrt{\mu_i} \cdot \mathbf{v}_i = \mathbf{0}$ as required. \square

C.2 Rounding a robustly separated set

We now complete the argument that `round-cut` outputs a robustly separated set when the given vector embedding is balanced. We begin by recalling a fact about uniformly sampled vectors on the sphere.

Lemma 18. *Given $\mathbf{v} \in \mathbb{R}^d$, and \mathbf{g} a uniformly sampled random vector in \mathcal{S}^{d-1} , the following hold.*

1. $\mathbb{E}_{\mathbf{g}} \langle \mathbf{g}, \mathbf{v} \rangle^2 = \frac{\|\mathbf{v}\|_2^2}{d}$
2. For any $\delta > \frac{d}{16}$, we have $\Pr_{\mathbf{g}} \left(\langle \mathbf{g}, \mathbf{v} \rangle^2 \geq \delta \cdot \frac{\|\mathbf{v}\|_2^2}{d} \right) \leq e^{-\delta/4}$.

We also require a lemma proved in [45] which states that, with constant probability, one can extract two balanced partitions where projection lengths are well separated when projecting along a random direction.

Lemma 19 (Lemma 5.5.6 item (3) of [45]). *Given a set V such that $|V| = n$, and a $(3, \frac{1}{10})$ -balanced embedding $\{\mathbf{v}_i\}_{i \in V} \subseteq \mathbb{R}^d$, let $r_i = \sqrt{d} \cdot \langle \mathbf{g}, \mathbf{v}_i \rangle$ for a $\mathbf{g} \in \mathcal{S}^{d-1}$ and each $i \in V$. Assume, without loss of generality, that $r_1 \geq \dots \geq r_n$. Then, there exists an absolute constant $r > 0$ such that, with $O(1)$ probability over a uniformly random choice of $\mathbf{g} \in \mathcal{S}^{d-1}$, there exists $1 \leq a < b \leq n$ satisfying the following:*

1. $\mu(\{1, \dots, a\}) \geq \frac{1}{3} \cdot \mu(V)$,
2. $\mu(\{b, \dots, n\}) \geq \frac{1}{3} \cdot \mu(V)$,
3. $(r_a - r_b)^2 \geq r \cdot \text{Var}_{\mu}(\mathbf{v}_i)$.⁵

Using the above two lemmata, we can now prove that **round-cut** produces a robustly separated set in the balanced case.

Proof of Lemma 9. Define \mathcal{E}_1 to be the event where there exists $1 \leq a < b \leq n$ such that the following hold simultaneously: (1) $\mu(\{1, \dots, a\}) \geq \frac{1}{3} \cdot \mu(V)$, (2) $\mu(\{b, \dots, n\}) \geq \frac{1}{3} \cdot \mu(V)$, and (3) $(r_a - r_b)^2 \geq \frac{r}{\mu(V)}$. Next, for a constant $c_1 > 0$ to be chosen subsequently, and any $i, j \in V$, define the event $\mathcal{E}_2(i, j)$ as

$$\mathcal{E}_2(i, j) \stackrel{\text{def}}{=} \left\{ (r_i - r_j)^2 < (4c_1 \cdot \log n) \cdot \|\mathbf{v}_i - \mathbf{v}_j\|_2^2 \right\} \quad \text{and} \quad \mathcal{E}_2 \stackrel{\text{def}}{=} \bigcap_{i, j \in V} \mathcal{E}_2(i, j).$$

To begin, we observe that if \mathcal{E}_1 and \mathcal{E}_2 occur jointly, then choosing the sweep cut $S = \{1, \dots, a\}$ gives an $(\Omega(\frac{1}{\log n}), \frac{1}{3})$ -robustly separated set with $T = \{b, \dots, n\}$, and, consequently, executing step (1) of **round-cut** outputs a robustly separated set in the balanced case.

The first requirement in demonstrating S is a robustly separated is immediately satisfied by item (2) defining event \mathcal{E}_1 since $\mu(T) \geq \frac{1}{3} \cdot \mu(V)$. For the second requirement, event \mathcal{E}_2 implies that

$$\|\mathbf{v}_i - \mathbf{v}_j\|_2^2 > \frac{(r_i - r_j)^2}{4c_1 \cdot \log n}$$

holds for all $i \in S$, and $j \in T$. Item (3) defining event \mathcal{E}_1 then determines:

$$\frac{(r_i - r_j)^2}{4c_1 \cdot \log n} \geq r \cdot \text{Var}_{\mu}(\mathbf{v}_i) \cdot \frac{1}{4c_1 \cdot \log n} = \frac{r}{\mu(V)} \cdot \frac{1}{4c_1 \cdot \log n}.$$

Finally, item (1) in event \mathcal{E}_1 implies that $\frac{1}{\mu(V)} \geq \frac{1}{3\mu(S)}$ and hence:

$$\|\mathbf{v}_i - \mathbf{v}_j\|_2^2 \geq \frac{r}{\mu(V)} \cdot \frac{1}{4c_1 \cdot \log n} \geq \frac{1}{\mu(S)} \cdot \frac{r}{12c_1 \cdot \log n} = \frac{1}{\mu(S)} \cdot \Omega\left(\frac{1}{\log n}\right)$$

Let us next bound the probability that \mathcal{E}_1 and \mathcal{E}_2 occur jointly. By Lemma 19, there exists a constant $p > 0$ such that $\Pr_{\mathbf{g}}(\mathcal{E}_1) = p$ and hence $\Pr_{\mathbf{g}}(\overline{\mathcal{E}_1}) = 1 - p$. By Lemma 18, we have

$$\begin{aligned} \Pr_{\mathbf{g}}(\overline{\mathcal{E}_2(i, j)}) &= \Pr_{\mathbf{g}}\left((r_i - r_j)^2 \geq (4c_1 \cdot \log n) \cdot \|\mathbf{v}_i - \mathbf{v}_j\|_2^2\right) \\ &= \Pr_{\mathbf{g}}\left(\langle \mathbf{g}, \mathbf{v}_i - \mathbf{v}_j \rangle^2 \geq (4c_1 \cdot \log n) \cdot \frac{\|\mathbf{v}_i - \mathbf{v}_j\|_2^2}{d}\right) \leq \frac{1}{n^{c_1}} \end{aligned}$$

Performing a union bound over $\overline{\mathcal{E}_1}$ and $\overline{\mathcal{E}_2(i, j)}$ for each $i, j \in V$ derives

$$\Pr_{\mathbf{g}}(\mathcal{E}_1 \cap \mathcal{E}_2) \geq p - \binom{n}{2} \frac{1}{n^{c_1}} \geq p - \frac{1}{n^{c_1-2}} = \Omega(1)$$

for c_1 large enough. We conclude that with constant probability, a single execution of step (1) in **round-cut** outputs a $(\Omega(\frac{1}{\log n}), \frac{1}{3})$ -robustly separated set. Repeating step (1) $O(\log n)$ times and performing another union bound then allows us to conclude **round-cut** produces a robustly separated set with probability $1 - o_n(\frac{1}{n})$. \square

⁵We note that in [45], the variance of the vector embedding is normalized to be 1, and hidden in the $\Omega_b(1)$ term.

D An $O(\sqrt{\log n})$ -Approximation via Metric Embeddings

In this section, we prove the technical results used in to obtain the $O(\sqrt{\log n})$ -approximation algorithm in Theorem 4. To begin, we complete the proof that the vector program (RC-Metric) is, up to a factor of 2, a relaxation of the original problem.

Proof of Theorem 11. Given any cut (S, \bar{S}) with $\mu(S) \leq \mu(\bar{S})$, for each $i \in V$, set the vector embedding of the vertices to

$$\mathbf{v}_i = \begin{cases} \left(\sqrt{\frac{\mu(V)}{\mu(S)\mu(\bar{S})}} & 0 & \dots & 0 \right)^\top & \text{if } i \in S \\ \mathbf{0} & & & & \text{otherwise.} \end{cases}$$

Assign the length vectors as follows:

$$\ell^h = \frac{\mu(V)}{\mu(S)\mu(\bar{S})} \cdot \begin{cases} \mathbf{1}^{S \cap h} & \text{if } F_h(S \cap h) \leq F_h(\bar{S} \cap h), \\ \mathbf{1}^{\bar{S} \cap h} & \text{otherwise.} \end{cases}$$

Then, $(\{\ell^h\}_{h \in E}, \{\mathbf{v}_i\}_{i \in V})$ is a feasible solution to the vector program as, for $i \in S \cap h$ and $j \in \bar{S} \cap h$:

$$\|\mathbf{v}_i - \mathbf{v}_j\|^2 = \frac{\mu(V)}{\mu(S)\mu(\bar{S})} = \ell_i^h + \ell_j^h.$$

Furthermore, this solution has objective value:

$$\sum_{h \in E} w_h \|\ell^h\|_F = \frac{\mu(V)}{\mu(S)\mu(\bar{S})} \cdot \sum_{h \in E} w_h \min\{\|\mathbf{1}^{S \cap h}\|_F, \|\mathbf{1}^{\bar{S} \cap h}\|_F\} \leq 2 \cdot \frac{\sum_{h \in E} w_h \delta_h(S \cap h)}{\min\{\mu(S), \mu(\bar{S})\}} = 2\Psi_G(S).$$

□

We now prove the technical Lemma 14 required by our rounding argument.

Proof of Lemma 14. Let $Y = \text{diag}(\mathbf{y})$, then the above statement is equivalent to proving that there exists some $\nu \in \mathbb{R}$ such that:

$$\min_{\nu \in \mathbb{R}} \|Y^{-1}(\mathbf{x} - \nu \mathbf{1})\|_\infty \leq 1.$$

We have:

$$\begin{aligned} \min_{\nu \in \mathbb{R}} \|Y^{-1}(\mathbf{x} - \nu \mathbf{1})\|_\infty &= \min_{\nu \in \mathbb{R}} \max_{\|\mathbf{z}\|_1 \leq 1} \langle \mathbf{z}, Y^{-1}(\mathbf{x} - \nu \mathbf{1}) \rangle = \min_{\nu \in \mathbb{R}} \max_{\|Y\mathbf{z}\|_1 \leq 1} \langle \mathbf{z}, \mathbf{x} - \nu \mathbf{1} \rangle = \max_{\|Y\mathbf{z}\|_1 \leq 1} \min_{\nu \in \mathbb{R}} \langle \mathbf{z}, \mathbf{x} - \nu \mathbf{1} \rangle \\ &= \max_{\substack{\|Y\mathbf{z}\|_1 \leq 1 \\ \mathbf{z} \perp \mathbf{1}}} \langle \mathbf{z}, \mathbf{x} \rangle. \end{aligned}$$

For any \mathbf{z} , let $Z^+ \stackrel{\text{def}}{=} \{i : z_i > 0\}$ and $Z^- \stackrel{\text{def}}{=} \{i : z_i < 0\}$. The above orthogonality constraint, implies:

$$-\sum_{i \in Z^-} z_i = \sum_{i \in Z^+} z_i,$$

so that the positive entries of \mathbf{z} can be charged to the negative entries of \mathbf{z} , i.e. there exists some choice of non-negative numbers $\{w_{i,j}\}_{(i,j) \in Z^+ \times Z^-}$ such that:

$$\forall i \in Z^- : \sum_{j \in Z^+} w_{i,j} = |z_i| \quad \text{and} \quad \forall j \in Z^+ : \sum_{i \in Z^-} w_{i,j} = |z_j|.$$

This gives:

$$\max_{\substack{\|Y\mathbf{z}\|_1 \leq 1 \\ \mathbf{z} \perp \mathbf{1}}} \langle \mathbf{z}, \mathbf{x} \rangle = \max_{\substack{\|Y\mathbf{z}\|_1 \leq 1 \\ \mathbf{z} \perp \mathbf{1}}} \left\{ \sum_{i \in Z^+} |z_i| x_i - \sum_{j \in Z^-} |z_j| x_j \right\} = \max_{\substack{\|Y\mathbf{z}\|_1 \leq 1 \\ \mathbf{z} \perp \mathbf{1}}} \left\{ \sum_{i \in Z^+} \sum_{j \in Z^-} w_{i,j} x_i - \sum_{j \in Z^-} \sum_{i \in Z^+} w_{i,j} x_j \right\}$$

$$\begin{aligned}
&= \max_{\substack{\|Y_{\mathbf{z}}\|_1 \leq 1 \\ \mathbf{z} \perp \mathbf{1}}} \left\{ \sum_{(i,j) \in Z^+ \times Z^-} w_{i,j} (x_i - x_j) \right\} \leq \max_{\substack{\|Y_{\mathbf{z}}\|_1 \leq 1 \\ \mathbf{z} \perp \mathbf{1}}} \left\{ \sum_{(i,j) \in Z^+ \times Z^-} w_{i,j} |x_i - x_j| \right\} \\
&\leq \max_{\substack{\|Y_{\mathbf{z}}\|_1 \leq 1 \\ \mathbf{z} \perp \mathbf{1}}} \left\{ \sum_{(i,j) \in Z^+ \times Z^-} w_{i,j} (y_i + y_j) \right\} = \max_{\substack{\|Y_{\mathbf{z}}\|_1 \leq 1 \\ \mathbf{z} \perp \mathbf{1}}} \left\{ \sum_{i \in Z^+} |z_i| y_i + \sum_{j \in Z^-} |z_j| y_j \right\} \\
&\leq \max_{\substack{\|Y_{\mathbf{z}}\|_1 \leq 1 \\ \mathbf{z} \perp \mathbf{1}}} \|Y_{\mathbf{z}}\|_1 \leq 1,
\end{aligned}$$

which yields the desired result. □