

A central limit theorem for a sequence of conditionally centered and α -mixing random fields

Abdollah Jalilian
Razi University, Iran
stat4aj@gmail.com

Ganggang Xu
University of Miami, United States of America
gangxu@bus.miami.edu

Rasmus Waagepetersen
Aalborg University, Denmark
rw@math.aau.dk

January 24, 2023

Abstract

A central limit theorem is established for a sum of random variables belonging to a sequence of random fields. The fields are assumed to have zero mean conditional on the past history and to satisfy certain α -mixing conditions in space or/and time. The limiting normal distribution is obtained for increasing spatial domain, increasing length of the sequence or a combination of these. The applicability of the theorem is demonstrated by examples regarding estimating functions for a space-time point process and a space-time Markov process.

Keywords: central limit theorem; conditional centering; conditional α -mixing; estimating function; random field; spatio-temporal; space-time
MSC2020 subject classifications: 60F05; 60G55; 60G60; 62M30

1 Introduction

While the literature on central limit theorems for spatial stochastic processes is extensive [Bolthausen \(1982\)](#); [Jensen and Künsch \(1994\)](#); [Karácsony \(2006\)](#); [Biscio and Waagepetersen \(2019\)](#), to mention a few, the literature on central limit theorems for spatio-temporal processes seems much less developed. The objective of this paper is to derive a central limit theorem for a sequence of random fields that satisfy a certain conditional centering condition. This condition is motivated by applications of estimating functions in space-time statistics (see [Section 3](#)) where the estimating functions based on a random field at one point in time are unbiased (have zero mean expectation at the true parameter value) conditional on the past.

In the case of space-time processes, asymptotics can be based on increasing spatial domain, increasing time horizon, or both, and our result covers all these possibilities. Our proof is based on the approach developed in the spatial setting by [Bolthausen \(1982\)](#), see also [Biscio et al. \(2018\)](#). This requires to bound a large variety of covariances. In the case of spatial random fields, this is typically done by imposing appropriate weak spatial dependence conditions like m -dependence, e.g. [Leonenko \(1975\)](#); [Heinrich \(2013\)](#) or α -mixing [Bolthausen \(1982\)](#). [Leonenko \(1975\)](#) considers a central limit theorem for a sequence of statistics each obtained as a sum of variables for a random field in a sequence of increasing domain m -dependent random fields. [Jensen and Künsch \(1994\)](#) considers a spatial random field of conditionally centered random variables that have expectation zero given all other variables. By exploiting the conditional centering, they are able to dispense with spatial mixing assumptions.

Different from [Jensen and Künsch \(1994\)](#), we assume conditional centering given the past history in time, which is useful for eliminating covariances involving variables at distinct time points. This technique is related to the key property of independent increments for martingales. Nevertheless, we need further additional assumptions of weak spatial or temporal dependence and impose conditions of conditional α -mixing in space and/or time. The reason for considering α -mixing conditional on the past, instead of unconditional α -mixing, is that the conditional property may be easier to verify in some applications, see [Section 3](#). We note that our notion of conditional α -mixing is very different from the notion of conditional m -dependence in [Biscio and Svane \(2022\)](#) for spatial point processes conditional on an underlying random field.

To give a concise statement of the various conditions needed, we introduce the novel concept of a space-time filtration, that is an increasing sequence of σ -algebras that are indexed both by time points and subsets of space.

2 The space-time central limit theorem

Let \mathbb{N} and \mathbb{Z} denote respectively the set of natural and integer numbers. Consider for each $k, n \in \mathbb{N}$ a random field $E_{k,n} = (E_{k,n}(\mathbf{1}))_{\mathbf{1} \in \mathcal{D}_n}$, on $\mathcal{D}_n \subseteq \mathbb{Z}^d$, $d \geq 1$, where $E_{k,n}(\mathbf{1}) = \left(E_{k,n}^{(1)}(\mathbf{1}), \dots, E_{k,n}^{(q)}(\mathbf{1}) \right)^\top$, $q \geq 1$, and $(\mathcal{D}_n)_{n \in \mathbb{N}}$ is a sequence of finite observation lattices while the index k can be viewed as a time index. Depending on the context (see examples in [Section 3](#)), it may or may not be reasonable to assume properties like stationarity or mixing in time for the sequence $E_{k,n}$, $k \in \mathbb{N}$. If the sequence $E_{k,n}$, $k \in \mathbb{N}$, can not be viewed as a part of a stationary sequence, it is convenient to consider it conditional on an initial condition represented by a σ -algebra \mathcal{H}_0 , see [Section 2.1](#). Defining for $K, n \in \mathbb{N}$,

$$T_{K,n} = \sum_{k=1}^K \sum_{\mathbf{1} \in \mathcal{D}_n} E_{k,n}(\mathbf{1}), \quad (1)$$

we formulate in [Section 2.2](#) a central limit theorem for $\Sigma_{K,n}^{-1/2} T_{K,n}$ with $\Sigma_{K,n} = \text{Var}_{\mathcal{H}_0}[T_{K,n}]$, that is conditional on the initial condition \mathcal{H}_0 .

2.1 Notation, space-time filtration and α -mixing

We define

$$d(x, y) = \max\{|x_i - y_i| : 1 \leq i \leq d\}, \quad x, y \in \mathbb{R}^d,$$

and, reusing notation,

$$d(A, B) = \inf\{d(x, y) : x \in A, y \in B\}, \quad A, B \subset \mathbb{R}^d.$$

For a subset $A \subseteq \mathbb{R}^d$ we denote by $|A|$ the cardinality or Lebesgue measure of A . The meaning of $|\cdot|$ and $d(\cdot, \cdot)$ will be clear from the context.

Conditioning on the past history plays a crucial role for our result. We represent the history by what we call a space-time filtration whose definition not only involves time points but also spatial regions. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be the underlying probability space on which all random objects in this paper are defined. Letting $\mathbb{Z}_+ = \{0, 1, 2, \dots\}$, we coin a set $\mathcal{H} = \{\mathcal{H}_{l,k,A} \mid l, k \in \mathbb{Z}_+, A \subseteq \mathbb{R}^d\}$ of σ -algebras $\mathcal{H}_{l,k,A} \subseteq \mathcal{F}$ a space time filtration if $\mathcal{H}_{l,k,A} \subseteq \mathcal{H}_{l',k',A}$ for $l' \leq l \leq k \leq k'$ and $\mathcal{H}_{l,k,A} \subseteq \mathcal{H}_{l,k,B}$ for $A \subseteq B$. For brevity we let $\mathcal{H}_k = \mathcal{H}_{0,k,\mathbb{R}^d}$. One can think of the space-time filtration as being generated by random vectors $E_{k,n}(\mathbf{l})$ for subsets of time and space indices k and \mathbf{l} as well as possible space-time exogeneous variables characterizing a space-time environment for the random fields $E_{k,n}$, $k \in \mathbb{N}$.

For any σ -algebras $\mathcal{G}_0, \mathcal{G}_1, \mathcal{G}_2 \subseteq \mathcal{F}$, the conditional α -mixing coefficient of \mathcal{G}_1 and \mathcal{G}_2 given \mathcal{G}_0 is defined in [Prakasa Rao \(2009\)](#) as

$$\alpha_{\mathcal{G}_0}(\mathcal{G}_1, \mathcal{G}_2) = \sup_{\substack{F_1 \in \mathcal{G}_1 \\ F_2 \in \mathcal{G}_2}} |\mathbb{P}_{\mathcal{G}_0}(F_1 \cap F_2) - \mathbb{P}_{\mathcal{G}_0}(F_1)\mathbb{P}_{\mathcal{G}_0}(F_2)|,$$

where $\mathbb{P}_{\mathcal{G}_0}(F) = \mathbb{E}_{\mathcal{G}_0}[I_F]$, $\mathbb{E}_{\mathcal{G}_0}$ denotes the conditional expectation with respect to \mathcal{G}_0 , $I_F(\omega) = I[\omega \in F]$ and $I[\cdot]$ is the indicator function. Similarly, $\text{Var}_{\mathcal{G}_0}$ and $\text{Cov}_{\mathcal{G}_0}$ denote variance and covariance conditional on \mathcal{G}_0 . Note that $0 \leq \alpha_{\mathcal{G}_0}(\mathcal{G}_1, \mathcal{G}_2) \leq 1/4$ is a \mathcal{G}_0 -measurable random variable.

For $m_1, m_2 \geq 0$, $c_1, c_2 \geq 0$, $\mathcal{D} \subseteq \mathbb{R}^d$, $K \in \mathbb{N}$ and $r \in \mathbb{Z}_+$, we define

$$\alpha_{c_1, c_2, r}^{(K, \mathcal{D})}(m_1, m_2) = \sup_{\substack{l, k \in \mathbb{Z}_+ \\ 0 \leq l, k \leq K \\ k-l \geq m_1}} \sup_{\substack{A, B \subseteq \mathcal{D} \\ |A| \leq c_1, |B| \leq c_2 \\ d(A, B) \geq m_2}} \sup_{i \in \{0, l-1\}} \alpha_{\mathcal{H}_i}(\mathcal{H}_{0,l,A}, \mathcal{H}_{(k-r)^+, k, B}), \quad (2)$$

where $(x)^+ = \max\{x, 0\}$.

Remark 1. *The definition (2) involves in a standard way sup over space and time lags as well as over sizes of spatial regions. However, it also involves a sup over conditional α -mixing coefficients where conditioning is either on the initial condition \mathcal{H}_0 or the ‘immediate’ past \mathcal{H}_{l-1} for a time point l . In real applications, dependence may propagate over space as time goes by. In such cases it usually is not more restrictive to consider sup over $\alpha_{\mathcal{H}_0}$ and $\alpha_{\mathcal{H}_{l-1}}$ instead of just considering $\alpha_{\mathcal{H}_0}$.*

2.2 Statement of result

We assume the existence of a space-time filtration \mathcal{H} so that the following conditional centering condition holds:

(A1) for any $k, n \in \mathbb{N}$, the random vectors $E_{k,n}(\mathbf{1})$, $\mathbf{1} \in \mathcal{D}_n$, are conditionally centered; i.e.,

$$\mathbb{E}_{\mathcal{H}_{k-1}}[E_{k,n}(\mathbf{1})] = 0.$$

We further need certain ‘localization’ properties, either in space or time, or both:

(A2) Consider $n, k \in \mathbb{N}$, $\mathbf{1} \in \mathbb{Z}^d$, $\mathcal{I} \subset \mathbb{Z}^d$, $\mathcal{T} \subseteq \{k, k+1, \dots, K\}$. Let $f : \mathbb{R}^{q|\mathcal{I}||\mathcal{T}|} \rightarrow \mathbb{R}$ be any measurable function and define $Y_n^{(\mathcal{T}, \mathcal{I})} = f((E_{k,n}(\mathbf{1}))_{(k, \mathbf{l}) \in \mathcal{T} \times \mathcal{I}})$.

(A2-1) there exists a distance $R > 0$ so that $E_{k,n}(\mathbf{1})$ and $\mathbb{E}_{\mathcal{H}_{k-1}}[Y_n^{(\mathcal{T}, \mathcal{I})}]$ are measurable with respect to respectively $\mathcal{H}_{0,k,C(\mathbf{1})}$ and $\mathcal{H}_{0,k-1, \cup_{\mathbf{l} \in \mathcal{I}} C(\mathbf{l})}$, where for any $\mathbf{l} \in \mathbb{Z}^d$, $C(\mathbf{l})$ is a cube of sidelength R centered at \mathbf{l} .

(A2-2) there exists a lag $r \in \mathbb{N}$ so that $E_{k,n}(\mathbf{1})$ and $\mathbb{E}_{\mathcal{H}_{k-1}}[Y_n^{(\mathcal{T}, \mathcal{I})}]$ are measurable with respect to respectively $\mathcal{H}_{0,k,\mathcal{D}_n}$ and $\mathcal{H}_{(k-r)^+, k-1, \mathcal{D}_n}$.

(A2-3) there exists an $R > 0$ and $r \in \mathbb{N}$ so that $E_{k,n}(\mathbf{1})$ and $\mathbb{E}_{\mathcal{H}_{k-1}}[Y_n^{(\mathcal{T}, \mathcal{I})}]$ are measurable with respect to respectively $\mathcal{H}_{0,k,C(\mathbf{1})}$ and $\mathcal{H}_{(k-r)^+, k-1, \cup_{\mathbf{l} \in \mathcal{I}} C(\mathbf{l})}$ where $C(\mathbf{l})$ is defined as in (A2-1).

Condition (A1) implies that $\mathbb{E}_{\mathcal{H}_{k'}}[E_{k,n}(\mathbf{1})] = 0$ for any $k' < k$. Combined with condition (A2) it further holds for any $0 \leq k' < k < l$, and $\mathbf{l}, \mathbf{j} \in \mathbb{Z}^d$,

$$\begin{aligned} \text{Cov}_{\mathcal{H}_{k'}}[E_{k,n}(\mathbf{1}), E_{l,n}(\mathbf{j})] &= \mathbb{E}_{\mathcal{H}_{k'}}[E_{k,n}(\mathbf{1})E_{l,n}^T(\mathbf{j})] \\ &= \mathbb{E}_{\mathcal{H}_{k'}}[\mathbb{E}_{\mathcal{H}_{l-1}}[E_{k,n}(\mathbf{1})E_{l,n}^T(\mathbf{j})]] = 0. \end{aligned}$$

Moreover, it holds that $\mathbb{E}_{\mathcal{H}_0}[T_{K,n}] = 0$ and

$$\Sigma_{K,n} = \sum_{k=1}^K \text{Var}_{\mathcal{H}_0} \left[\sum_{\mathbf{l} \in \mathcal{D}_n} E_{k,n}(\mathbf{l}) \right] = \sum_{k=1}^K \sum_{\mathbf{l}, \mathbf{j} \in \mathcal{D}_n} \mathbb{E}_{\mathcal{H}_0} [\text{Cov}_{\mathcal{H}_{k-1}}[E_{k,n}(\mathbf{l}), E_{k,n}(\mathbf{j})]].$$

Next, we introduce the following conditional α -mixing conditions for the filtration \mathcal{H} :

(A3) there exists a real number $p > 2$ such that one of the following holds almost surely.

(A3-1) for any $K \in \mathbb{N}$, there exists a function $\bar{\alpha}_K : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that

$$\left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{2R^2, \infty, K}^{(K, \mathbb{R}^d)}(0, m) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}} \leq \bar{\alpha}_K(m)$$

and $\bar{\alpha}_K(m) = O(m^\eta)$, where $\eta + d < 0$.

(A3-2) for any $n \in \mathbb{N}$, there exists a function $\tilde{\alpha}_n : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that

$$\left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{|\mathcal{D}_n|, |\mathcal{D}_n|, r}^{(\infty, \mathcal{D}_n)}(m, 0) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}} \leq \tilde{\alpha}_n(m)$$

and $\tilde{\alpha}_n(m) = O(m^\psi)$, where $\psi < -1$.

(A3-3) there exists a function $\check{\alpha} : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$ such that

$$\begin{aligned} & \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{\infty, \infty, r}^{(\infty, \mathbb{R}^d)}(m_1, m_2) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}} \leq \check{\alpha}(m_1, m_2) \\ & \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{2R^2, 2R^2, K}^{(K, \mathbb{R}^d)}(0, m_2) \right)^{p-2} \right] \right)^{\frac{1}{p}} \mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{|\mathcal{D}_n|, |\mathcal{D}_n|, r}^{(\infty, \mathcal{D}_n)}(m_1, 0) \right)^{\frac{p-2}{p}} \right] \leq \check{\alpha}(m_1, m_2) \end{aligned}$$

and $\check{\alpha}(m_1, m_2) = O(m_1^\psi m_2^\eta)$, where $\eta + d < 0$ and $\psi < -1$.

Finally, to obtain a central limit theorem for $\Sigma_{K,n}^{-1/2} T_{K,n}$, we need the following sets of conditions:

(A4) there exist an $\epsilon \geq 3(p-2)$ and a constant $0 < M_\epsilon < \infty$ such that for

$$d_{K,n} = \sup_{1 \leq k' \leq k \leq K} \sup_{\mathbf{l} \in \mathcal{D}_n} \sup_{i=1, \dots, q} \mathbb{E}_{\mathcal{H}_0} \left[\left| E_{k,n}^{(i)}(\mathbf{l}) \right|^{6+\epsilon} \right],$$

one of the following holds almost surely

$$(A4-1) \sup_{n \in \mathbb{N}} d_{K,n} \leq M_\epsilon^{6+\epsilon}.$$

$$(A4-2) \sup_{K \in \mathbb{N}} d_{K,n} \leq M_\epsilon^{6+\epsilon}.$$

$$(A4-3) \sup_{K \in \mathbb{N}} \sup_{n \in \mathbb{N}} d_{K,n} \leq M_\epsilon^{6+\epsilon}.$$

and, with $\lambda_{\min}(M)$ being the smallest eigen value of a symmetric matrix M ,

(A5) one of the following holds almost surely.

$$(A5-1) 0 < \liminf_{n \rightarrow \infty} \lambda_{\min}(\Sigma_{K,n}) / |\mathcal{D}_n|.$$

$$(A5-2) 0 < \liminf_{K \rightarrow \infty} \lambda_{\min}(\Sigma_{K,n}) / K.$$

$$(A5-3) 0 < \liminf_{K, n \rightarrow \infty} \lambda_{\min}(\Sigma_{K,n}) / (K |\mathcal{D}_n|).$$

We then obtain the following theorem.

Theorem 1. *Let $(T_{K,n})_{K,n \in \mathbb{N}}$ be a sequence of q -dimensional statistics of the form (1) and assume that condition (A1) holds. Assume further that either*

(i) *condition (A2-1), (A3-1), (A4-1), and (A5-1) hold, $n \rightarrow \infty$ and $\lim_{n \rightarrow \infty} |\mathcal{D}_n| = \infty$,*

(ii) *condition (A2-2), (A3-2), (A4-2), and (A5-2) hold, and $K \rightarrow \infty$, or*

(iii) *condition (A2-3), (A3-3), (A4-3), and (A5-3) hold, $n \rightarrow \infty$, $K \rightarrow \infty$, and $\lim_{n \rightarrow \infty} |\mathcal{D}_n| = \infty$.*

Then for any $t \in \mathbb{R}^q$,

$$\mathbb{E}_{\mathcal{H}_0} \left[\exp(it^\top \Sigma_{K,n}^{-\frac{1}{2}} T_{K,n}) \right] \rightarrow \exp(\|t\|^2/2)$$

almost surely.

A proof of Theorem 1 is given in section 4. We discuss the various assumptions in relation to the examples of applications in Section 3.

Remark 2. The conclusion of [Theorem 1](#) is exactly the same for all three cases (i)-(iii). Therefore, from a practical point of view, case (iii) is superfluous since the assumptions for this case imply the assumptions for (i) and (ii). Nevertheless, for sake of completeness, we include (iii) in the statement of the result.

Remark 3. There exists some measurable function X_0 so that $\mathcal{H}_0 = \sigma(X_0)$. Let $\phi_{K,n}(x_0) = \mathbb{E}[\exp(it^\top \Sigma_{K,n}^{-\frac{1}{2}} T_{K,n}) | X_0 = x_0]$ where expectation is with respect to a regular conditional distribution of the $E_{k,n}$, $1 \leq k \leq K$, $n \in \mathbb{N}$, given $X_0 = x_0$. Then $\mathbb{E}_{\mathcal{H}_0}[\exp(it^\top \Sigma_{K,n}^{-\frac{1}{2}} T_{K,n})] = \phi_{K,n}(X_0)$ almost surely. In our proof we thus show that $\phi_{K,n}(x_0)$ converges to $\exp(\|t\|^2/2)$ for almost all values x_0 of X_0 .

Remark 4. The index n for $E_{k,n}(\mathbf{1})$ may arise in applications due to boundary effects for indices $\mathbf{1}$ close to the boundary of \mathcal{D}_n . In that case taking sup over n in conditions (A5-1)-(A5-3) is usually not restrictive.

Remark 5. The result in case (ii) ($K \rightarrow \infty$, n fixed) could alternatively under the same conditions be established using existing martingale central limit theorems, e.g. [Brown \(1971\)](#); [Hall et al. \(2014\)](#). To apply these theorems it is required to control a sum of conditional variances. This, however, involves all the results from the supplementary material and lengthy derivations analogous to those used to handle the term A_1 in our proof. We therefore find it more natural to use [Bolthausen's](#) approach as a unified framework for all cases (i)-(iii) in our theorem.

Remark 6. In the proof of [Theorem 1](#), we need to bound a large number, of the order $K^4 |\mathcal{D}_n|^4$, of certain covariances, see [Section 4.1](#) and the supplementary material. The conditional centering property (A1) is crucial in this regard since it immediately reduces the number of the non-zero covariances to be of the order $K^3 |\mathcal{D}_n|^4$. However, it does not affect the order $|\mathcal{D}_n|^4$. This is why spatial mixing assumptions are required in the K fixed case (i).

3 Examples of applications

In the following sections we provide illustrative examples of the use of [Theorem 1](#), where we try to strike a balance between accessibility and practical relevance of the examples. Following [Remark 2](#) we focus on cases (i) and (ii) in the following [Sections 3.1](#) and [3.2](#). We also sketch how case (iii) can cover a modification of the example in [Section 3.1](#).

3.1 A discrete time spatial birth-death point process

As a specific example of application of [Theorem 1](#) in case (i), consider a sequence of point processes X_0, \dots, X_K and spatial covariates Z_0, \dots, Z_K on \mathbb{R}^d , where the latter are considered non-random. For each n , we observe $X_k \cap W_n$ and $Z_k(v)$, $v \in W_n$, $k = 0, \dots, K$, where $W_1 \subset W_2 \subset \dots \subseteq \mathbb{R}^2$ is a sequence of increasing observation windows. We consider a model for the X_k corresponding to a discrete-time spatial birth-death process that mimics the seasonal behaviour of a plant community.

Given X_{k-1} and Z_{k-1} , $X_k = S_k \cup B_k \cup I_k$ where $B_k = \cup_{u \in X_{k-1}} Y_u^k$ is a union of offspring clusters Y_u^k given by independent Poisson processes. For each

$u \in X_{k-1}$, Y_u^k has an intensity function with support given by the disc $b(u, \omega)$ with center u and radius ω . The intensity function depends on X_{k-1} only through u . The point process S_k represents plants from X_{k-1} that survived to time k and is given by $S_k = \{u \in X_{k-1} | R_u = 0\}$, where the R_u are Bernoulli variables conditionally independent given X_{k-1} and Z_{k-1} and with ‘survival’ probability $\mathbb{P}(R_u = 0 | X_{k-1}, Z_{k-1})$ depending on X_{k-1} and Z_{k-1} only through $Z_{k-1}(u)$. The ‘immigrant’ point processes I_k , $k \geq 1$, are independent Poisson processes of intensity $\rho \geq 0$. For any $A \subseteq \mathbb{R}^d$ and $k \in \mathbb{Z}_+$, we define $\mathcal{H}_{l,k,A} = \sigma((X_{k'} \cap A), l \leq k' \leq k)$.

For each k we let $T_k^{W_n}$ denote the concatenation of likelihood score functions for the conditional distributions of $B_k \cup I_k$ and S_k given Z_{k-1} and X_{k-1} . Then $T_k^{W,n}$ can be additively decomposed as

$$T_k^{W_n} = \sum_{\mathbf{l} \in \mathcal{D}_n} E_{k,n}(\mathbf{l}),$$

where $\mathcal{D}_n = \{\mathbf{l} \in \mathbb{Z}^d : c(\mathbf{l}) \cap W_n \neq \emptyset\}$, $c(\mathbf{l})$ is the unit cube centered at \mathbf{l} and the likelihood score component $E_{k,n}(\mathbf{l})$ depends on $Z_{k-1}, X_{k-1}, B_k, S_k$ only through $Z_{k-1}(v)$, $v \in c(\mathbf{l})$ and through the intersections of $X_{k-1}, B_k \cup I_k$, and S_k with $c(\mathbf{l})$.

By the first Bartlett identity (unbiasedness of score functions), $E_{k,n}(\mathbf{l})$ is conditionally centered. Since $B_k \cup I_k = X_k \setminus X_{k-1}$ and $S_k = X_k \cap X_{k-1}$, $E_{k,n}(\mathbf{l})$ is $\mathcal{H}_{0,k,c(\mathbf{l})}$ measurable. Defining $Y_n^{(\mathcal{T}, \mathcal{I})}$ as in (A2), $\mathbb{E}_{\mathcal{H}_{k-1}}[Y_n^{(\mathcal{T}, \mathcal{I})}]$ is a function only of X_{k-1} restricted to $(\cup_{\mathbf{i} \in \mathcal{I}c(\mathbf{i})) \oplus (K-k)\omega$ and is hence $\mathcal{H}_{0,k-1,(\cup_{\mathbf{i} \in \mathcal{I}c(\mathbf{i})) \oplus (K-k)\omega}$ measurable, which means that (A2-1) holds (here $A \oplus s = \{x \in \mathbb{R}^d | d(\{x\}, A) \leq s\}$ for $A \subset \mathbb{R}^d$ and $s > 0$). Condition (A3-1) holds since $(X_l \cap A)$, $l = 1, \dots, K$, and $(X_k \cap B)$, $k = 1, \dots, K$, are independent given \mathcal{H}_0 whenever $d(A, B) > 2K\omega$. It is easy to check that (A4-1) holds and assuming further that (A5-1) holds, case (i) of our Theorem 1 is applicable to

$$T_{K,n} = \sum_{k=1}^K T_k^{W_n} = \sum_{k=1}^K \sum_{\mathbf{l} \in \mathcal{D}_n} E_{k,n}(\mathbf{l}),$$

for any fixed K . The validity of (A5-1) depends on X_0 that e.g. must not be confined to a bounded region.

Suppose further that the immigrant intensity ρ is strictly positive, ensuring that the ‘forest’ never becomes extinct. Also for u in X_0 or $\cup_{k \geq 1} I_k$ let C_u denote the cluster consisting of u and all its descendants, c.f. the birth-death mechanism above. If the probability of survival is small enough compared to the birth intensities, C_u will be finite almost surely and hence stochastically bounded in space and time. Thereby techniques similar to those in Cheyssson and Lang (2021) can be used to establish the required α -mixing for case (iii) of Theorem 1.

3.2 Conditional autoregressive model

In this section, we apply Theorem 1 case (ii) to a spatio-temporal autoregressive model. The model can also be viewed as a multivariate time series with graphical interactions at each time instance (see Dahlhaus (2000)). Let $\mathcal{D}_n \subset \mathbb{R}^d$ be a finite lattice and $G_n = (\mathcal{D}_n, \mathcal{E}_n)$ be an undirected graph with vertices in \mathcal{D}_n and edges \mathcal{E}_n where $\mathbf{l} \neq \mathbf{j}$ in \mathcal{D}_n are connected if $\{\mathbf{l}, \mathbf{j}\} \in \mathcal{E}_n$ Besag (1974).

For example, elements of \mathcal{D}_n may represent n sub-regions (provinces, counties or municipalities) of a specific geographical region and edges in \mathcal{E}_n determine inter-connections between these sub-regions due to e.g. adjacency or transportation routes.

Let $X_{k,n} = (X_{k,n}(\mathbf{l}))_{\mathbf{l} \in \mathcal{D}_n}$, $k \in \mathbb{Z}_+$, be a sequence of random fields with induced filtration

$$\mathcal{H}_{l,k,A} = \sigma(\{X_{k',n}(\mathbf{l}) : \mathbf{l} \in A, l \leq k' \leq k\}),$$

for $l, k \in \mathbb{Z}_+$, $l \leq k$, and $A \subseteq \mathcal{D}_n$. For any $k \in \mathbb{Z}_+$ and $\mathbf{l} \in \mathcal{D}_n$, we assume that given the past history $\mathcal{H}_{k-1} = \mathcal{H}_{0,k-1,\mathcal{D}_n}$ and neighborhood information $\mathcal{H}_{k,k,\mathcal{D}_n \setminus \{\mathbf{l}\}}$, $X_{k,n}(\mathbf{l})$ follows the conditional distribution

$$X_{k,n}(\mathbf{l}) | \sigma(\mathcal{H}_{k-1} \cup \mathcal{H}_{k,k,\mathcal{D}_n \setminus \{\mathbf{l}\}}) \sim \mathcal{N}(\beta \xi_{k-1,n}^{\text{temp}}(\mathbf{l}) + \gamma \xi_{k,n}^{\text{spat}}(\mathbf{l}), 1/a_n(\mathbf{l})), \quad (3)$$

where

$$\xi_{k-1,n}^{\text{temp}}(\mathbf{l}) = \sum_{j=1}^r \sum_{\mathbf{j} \in \mathcal{D}_n} b_j(\mathbf{l}, \mathbf{j}) X_{k-j,n}(\mathbf{j}),$$

with $X_{k-j,n}(\mathbf{l}) = 0$ for $k < j$ and

$$\xi_{k,n}^{\text{spat}}(\mathbf{l}) = \sum_{\mathbf{j} \in \mathcal{D}_n \setminus \{\mathbf{l}\}} b_0(\mathbf{l}, \mathbf{j}) (X_{k,n}(\mathbf{j}) - \beta \xi_{k-1,n}^{\text{temp}}(\mathbf{j})).$$

Let $\xi_{k-1,n}^{\text{temp}} = (\xi_{k-1,n}^{\text{temp}}(\mathbf{l}))_{\mathbf{l} \in \mathcal{D}_n}$ and $\xi_{k,n}^{\text{spat}} = (\xi_{k,n}^{\text{spat}}(\mathbf{l}))_{\mathbf{l} \in \mathcal{D}_n}$, $A_n = \text{diag}(a_n(\mathbf{l}) : \mathbf{l} \in \mathcal{D}_n)$ and $B_{j,n} = [b_j(\mathbf{l}, \mathbf{j})]_{\mathbf{l}, \mathbf{j} \in \mathcal{D}_n}$, $j = 0, \dots, r$, with diagonal entries $b_0(\mathbf{l}, \mathbf{l}) = 0$ for $B_{0,n}$. Then,

$$\xi_{k-1,n}^{\text{temp}} = \sum_{j=1}^r B_{j,n} X_{k-j,n} \quad \text{and} \quad \xi_{k,n}^{\text{spat}} = B_{0,n} (X_{k,n} - \beta \xi_{k-1,n}^{\text{temp}}).$$

Given \mathcal{H}_{k-1} and by Brook's factorization Lemma [Besag \(1974\)](#), $X_{k,n}$ is a Gaussian Markov random field with mean vector $\mu_{k-1,n} = \beta \xi_{k-1,n}^{\text{temp}}$ and precision matrix

$$Q_n = A_n (I_{|\mathcal{D}_n|} - \gamma B_{0,n}),$$

where $I_{|\mathcal{D}_n|}$ is the $|\mathcal{D}_n| \times |\mathcal{D}_n|$ identity matrix. We assume Q_n is positive definite and symmetric where the latter follows from symmetry of $A_n B_{0,n}$. Given \mathcal{H}_{k-1} ,

$$X_{k,n} \sim \beta \xi_{k-1,n}^{\text{temp}} + Q_n^{-1/2} Z_{k,n}, \quad k \in \mathbb{N}, \quad (4)$$

where the $Z_{k,n}$, $k \geq 1$, are independent standard normal vectors.

Define for $m \in \mathbb{N}$ and $i < l \leq k$ the maximal conditional correlation coefficient

$$\rho_{\mathcal{H}_i}(\mathcal{H}_{0,l,\mathcal{D}_n}, \mathcal{H}_{(k-r)^+,k,\mathcal{D}_n}) = \sup |\text{Corr}_{\mathcal{H}_i}[Z, Y]|,$$

where the supremum is over all random variables Z and Y that are respectively $\mathcal{H}_{0,l,\mathcal{D}_n}$ and $\mathcal{H}_{(k-r)^+,k,\mathcal{D}_n}$ measurable, and where $\mathbb{E}_{\mathcal{H}_i}[Z^2] < \infty$ and $\mathbb{E}_{\mathcal{H}_i}[Y^2] < \infty$. It is shown in Section 5.1 in the supplementary material that for a $\psi < -1$ and any $\mathbf{l}, \mathbf{j} \in \mathcal{D}_n$, $\text{Cov}_{\mathcal{H}_i}[X_{l,n}(\mathbf{l}), X_{k,n}(\mathbf{j})] = O(m^\psi)$ as $m \rightarrow \infty$ so that

$$\sup_{\substack{l,k \in \mathbb{Z}_+ \\ k-l \geq m}} \sup_{A, B \subseteq \mathcal{D}_n} \sup_{i \in \{0, l-1\}} \rho_{\mathcal{H}_i}(\mathcal{H}_{0,l,A}, \mathcal{H}_{(k-r)^+,k,B}) \leq \text{constant } m^\psi.$$

Since (see inequality (1.12) in Bradley (2005))

$$\alpha_{\mathcal{H}_i}(\mathcal{H}_{0,l,\mathcal{D}_n}, \mathcal{H}_{(k-r)^+,k,\mathcal{D}_n}) \leq \frac{1}{4} \rho_{\mathcal{H}_i}(\mathcal{H}_{0,l,\mathcal{D}_n}, \mathcal{H}_{(k-r)^+,k,\mathcal{D}_n}),$$

we have

$$\alpha_{|\mathcal{D}_n|,|\mathcal{D}_n|,r}^{\infty,\mathcal{D}_n}(m, 0) \leq \frac{1}{4} \sup_{\substack{l,k \in \mathbb{Z}_+ \\ k-l \geq m}} \sup_{A,B \subseteq \mathcal{D}_n} \sup_{i \in \{0,l-1\}} \rho_{\mathcal{H}_i}(\mathcal{H}_{0,l,A}, \mathcal{H}_{(k-r)^+,k,B}) \leq \text{constant } m^\psi,$$

which means that

$$\left(\mathbb{E}_{\mathcal{H}_0} \left[\alpha_{|\mathcal{D}_n|,|\mathcal{D}_n|,r}^{\infty,\mathcal{D}_n}(m, 0)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}}$$

is dominated by an $O(m^\psi)$ term and hence condition (A3-2) holds.

We assume that $b_0(\mathbf{l}, \mathbf{j}), \dots, b_p(\mathbf{l}, \mathbf{j})$ and $a_n(\mathbf{l})$ are known and $\beta, \gamma \in \mathbb{R}$ are unknown parameters. We further assume that $X_{k,n}$ is only observed for $k = 1, \dots, K$, and based on these partial observations, a score function for estimation of β and γ is given by (see Section 5.2 of the supplementary material for details)

$$T_{K,n} = \sum_{k=1}^K \sum_{\mathbf{l} \in \mathcal{D}_n} E_{k,n}(\mathbf{l}),$$

where

$$E_{k,n}(\mathbf{l}) = \varepsilon_{k,n}(\mathbf{l}) \left(\xi_{k-1,n}^{\text{temp}}(\mathbf{l}), \xi_{k,n}^{\text{spat}}(\mathbf{l}) \right)^\top$$

and $\varepsilon_{k,n} = X_{k,n} - \beta \xi_{k-1,n}^{\text{temp}} \sim Q_n^{-1/2} Z_{k,n}$.

By definition, $\xi_{k-1,n}^{\text{temp}}(\mathbf{l})$ and $\xi_{k,n}^{\text{spat}}(\mathbf{l})$ are respectively measurable with respect to $\mathcal{H}_{(k-r)^+,k-1,\mathcal{D}_n}$ and $\mathcal{H}_{k,k,\mathcal{D}_n \setminus \{\mathbf{l}\}}$. The conditional centering (A1) holds because

$$\mathbb{E}_{\mathcal{H}_{k-1}}[E_{k,n}(\mathbf{l})] = \mathbb{E}_{\mathcal{H}_{k-1}} \left[\mathbb{E}_{\sigma(\mathcal{H}_{k-1} \cup \mathcal{H}_{k,k,\mathcal{D}_n \setminus \{\mathbf{l}\}})}[\varepsilon_{k,n}(\mathbf{l})] \left(\xi_{k-1,n}^{\text{temp}}(\mathbf{l}), \xi_{k,n}^{\text{spat}}(\mathbf{l}) \right)^\top \right] = 0.$$

Hence, $T_{K,n}$ provides an unbiased estimating function. In addition, $E_{k,n}(\mathbf{l})$ is $\mathcal{H}_{(k-r)^+,k,\mathcal{D}_n}$ measurable and for any $\mathcal{T} \subset \{k, k = 1, \dots, K\}$, $\mathcal{I} \subset \mathcal{D}_n$ and measurable function $f : \mathbb{R}^{2^{|\mathcal{T}||\mathcal{I}|}} \rightarrow \mathbb{R}$, $\mathbb{E}_{\mathcal{H}_{k-1}}[f((E_{l,n}(\mathbf{l}))_{l \in \mathcal{T}, \mathbf{l} \in \mathcal{I}})]$ is $\mathcal{H}_{(k-r)^+,k-1,\mathcal{D}_n}$ measurable. This means that the localization condition (A2-2) holds.

It is shown in Section 5.3 of the supplementary material that

$$\Sigma_{K,n} = \text{Var}_{\mathcal{H}_0}[T_{K,n}] = \sum_{k=1}^K \mathbb{E}_{\mathcal{H}_0} \text{Var}_{H_{k-1}} \sum_{\mathbf{l} \in \mathcal{D}_n} E_{k,n}(\mathbf{l}) = \text{diag}(\lambda_{K,n}^{(1)}, \lambda_{K,n}^{(2)}),$$

where $\lambda_{K,n}^{(1)} \geq K \text{trace}(B_{1,n}^2)$ and $\lambda_{K,n}^{(2)} = 2K \text{trace}(B_{0,n}^2)$. We assume that the spatial coefficient matrix $B_{0,n}$ and the first order temporal coefficient matrix $B_{1,n}$ are such that

$$\begin{aligned} \text{trace}(B_{0,n}^2) &= \sum_{\mathbf{l}, \mathbf{j} \in \mathcal{D}_n} b_0(\mathbf{l}, \mathbf{j}) b_0(\mathbf{j}, \mathbf{l}) > 0, \\ \text{trace}(B_{1,n}^2) &= \sum_{\mathbf{l}, \mathbf{j} \in \mathcal{D}_n} b_1(\mathbf{l}, \mathbf{j}) b_1(\mathbf{j}, \mathbf{l}) > 0. \end{aligned}$$

Then, $\liminf_{K \rightarrow \infty} \lambda_{K,n}^{(1)}/K > 0$ and $\liminf_{K \rightarrow \infty} \lambda_{K,n}^{(2)}/K > 0$ and hence condition (A5-2) holds.

Following Section 5.4 in the supplementary material, for $1 \leq k' \leq k \leq K$ and $\nu > 0$,

$$\begin{aligned} \mathbb{E}_{\mathcal{H}_{k'-1}} \left[\sup_{\mathbf{l} \in \mathcal{D}_n} \mathbb{E}_{\mathcal{H}_{k-1}} \left[\left| \varepsilon_{k,n}(\mathbf{l}) \xi_{k-1,n}^{\text{temp}}(\mathbf{l}) \right|^\nu \right] \right] &\leq \text{constant} \mathbb{E}_{\mathcal{H}_{k'-1}} \left[\left\| \xi_{k-1,n}^{\text{temp}} \right\|^\nu \right], \\ \mathbb{E}_{\mathcal{H}_{k'-1}} \left[\sup_{\mathbf{l} \in \mathcal{D}_n} \mathbb{E}_{\mathcal{H}_{k-1}} \left[\left| \varepsilon_{k,n}(\mathbf{l}) \xi_{k,n}^{\text{spat}}(\mathbf{l}) \right|^\nu \right] \right] &\leq \text{constant}. \end{aligned}$$

Using the representation (5.1) in the supplementary material and the Minkowski inequality,

$$\begin{aligned} \mathbb{E}_{\mathcal{H}_{k'-1}} \left[\left\| \xi_{k-1,n}^{\text{temp}} \right\|^\nu \right] &= \mathbb{E}_{\mathcal{H}_{k'-1}} \left[\left\| C_{k,n} X_{0,n} + \sum_{j=1}^{k-1} C_{j,n} Q_n^{-1/2} Z_{k-j,n} \right\|^\nu \right] \\ &\leq \mathbb{E}_{\mathcal{H}_{k'-1}} \left[\left\| C_{k,n} X_{0,n} \right\|^\nu \right] + \mathbb{E}_{\mathcal{H}_{k'-1}} \left[\left\| \sum_{j=1}^{k-1} C_{j,n} Q_n^{-1/2} Z_{k-j,n} \right\|^\nu \right] \\ &\leq \|C_{k,n}\|^\nu \|X_{0,n}\|^\nu + \sum_{j=1}^{k-1} \|C_{j,n}\|^\nu \mathbb{E}_{\mathcal{H}_{k'-1}} \left[\left\| Q_n^{-1/2} Z_{k-j,n} \right\|^\nu \right] \\ &\leq \text{constant}_1 k^{\psi\nu} + \text{constant}_2 \sum_{j=1}^{\infty} j^{\psi\nu} < \infty \end{aligned}$$

because

$$\mathbb{E}_{\mathcal{H}_{k'-1}} \left[\left\| Q_n^{-1/2} Z_{k-j,n} \right\|^\nu \right] = \mathbb{E}_{\mathcal{H}_{k'-1}} \left[\left(Z_{k-j,n}^\top Q_n^{-1} Z_{k-j,n} \right)^{\frac{\nu}{2}} \right]$$

is a bounded constant that does not depend on j and k and $C_{k,n} = O(k^\psi)$, as $k \rightarrow \infty$. Thus, condition (A4-2) holds for $\nu = 6 + \epsilon$ and $\epsilon \geq 3(p-2)$. Therefore, the case (ii) of Theorem 1 is applicable for the statistic $T_{K,n}$.

4 Proof of the central limit theorem

In the proof of Theorem 1 in Section 4.1, we need to bound covariances of the form

$$\text{Cov}_{\mathcal{H}_0} [E_{k,n}(\mathbf{l}) E_{l,n}(\mathbf{j}), E_{k',n}(\mathbf{l}') E_{l',n}(\mathbf{j}')] \quad (5)$$

for $1 \leq k, k' \leq K$ and $k \leq l \leq K, k' \leq l' \leq K$ in the one-dimensional case $q = 1$.

The covariance (5) can be rewritten as

$$\begin{aligned} &\mathbb{E}_{\mathcal{H}_0} [\text{Cov}_{\mathcal{H}_{k^*-1}} [E_{k,n}(\mathbf{l}) E_{l,n}(\mathbf{j}), E_{k',n}(\mathbf{l}') E_{l',n}(\mathbf{j}')]] \\ &+ \text{Cov}_{\mathcal{H}_0} [\mathbb{E}_{\mathcal{H}_{k^*-1}} [E_{k,n}(\mathbf{l}) E_{l,n}(\mathbf{j})], \mathbb{E}_{\mathcal{H}_{k^*-1}} [E_{k',n}(\mathbf{l}') E_{l',n}(\mathbf{j}')]], \end{aligned}$$

where $k^* = \max\{k, l, k', l'\}$. If precisely one index coincides with k^* , then the covariance (5) is zero. Thus, the covariance (5) can only be non-zero in the cases a) $k^* = k = l > l' \geq k'$ or $k^* = k' = l' > l \geq k$, b) $k^* = l = l' > k, k'$,

Table 1: Bounds on the covariance (5) in different cases.

a)	case (i)	$8M_\epsilon^4 \bar{\alpha}_K (\text{d}(\{\mathbf{1}, \mathbf{j}\}, \{\mathbf{1}', \mathbf{j}'\}) - 2R)$
	case (ii)	$8M_\epsilon^4 \tilde{\alpha}_n (k - 1 - l')$
	case (iii)	$8M_\epsilon^4 \check{\alpha} (k - 1 - l', \text{d}(\{\mathbf{1}, \mathbf{j}\}, \{\mathbf{1}', \mathbf{j}'\}) - 2R)$
b)	case (i)	$8M_\epsilon^4 \bar{\alpha}_K (\text{d}(\{\mathbf{1}, \mathbf{j}\}, \{\mathbf{1}', \mathbf{j}'\}) - 2R)$
	case (ii)	$M_\epsilon^4 (8\tilde{\alpha}_n (l - 1 - \max\{k, k'\}))^{I[k \neq k']}$
	case (iii)	$8M_\epsilon^4 \check{\alpha} (I[k \neq k'] (l - 1 - \max\{k, k'\}), \text{d}(\{\mathbf{1}, \mathbf{j}\}, \{\mathbf{1}', \mathbf{j}'\}) - 2R)$
c)	case (i)	$8M_\epsilon^4 \bar{\alpha}_K (\text{d}(\{\mathbf{1}, \mathbf{j}\}, \{\mathbf{1}', \mathbf{j}'\}) - 2R)$
	case (ii)	$8M_\epsilon^4 \tilde{\alpha}_n (k - 1 - k')$
	case (iii)	$64M_\epsilon^4 \check{\alpha} (k - 1 - k', \text{d}(\{\mathbf{1}, \mathbf{j}\}, \{\mathbf{1}', \mathbf{j}'\}) - 2R)$
d)	case (i)	$8M_\epsilon^4 \bar{\alpha}_K (\text{d}(\{\mathbf{1}, \mathbf{j}\}, \{\mathbf{1}', \mathbf{j}'\}) - 2R)$
	case (ii)	$2M_\epsilon^4$
	case (iii)	$8M_\epsilon^4 \check{\alpha} (0, \text{d}(\{\mathbf{1}, \mathbf{j}\}, \{\mathbf{1}', \mathbf{j}'\}) - 2R)$

c) $k^* = k = l = l' > k'$ or $k^* = l' = k' = l > k$, and d) $k^* = k = l = k' = l'$. The consideration of each of these cases is rather tedious and the derivations are given in the supplementary material, where we obtain the bounds for (5) summarized in Table 1.

4.1 Proof of Theorem 1

Conditions (A4-1), (A4-2), or (A4-3) ensure that $\limsup_{n \rightarrow \infty} \lambda_{\max}(\Sigma_{K,n})/|D_n| < \infty$, $\limsup_{K \rightarrow \infty} \lambda_{\max}(\Sigma_{K,n})/K < \infty$, or $\limsup_{K,n \rightarrow \infty} \lambda_{\max}(\Sigma_{K,n})/(K|D_n|) < \infty$, where $\lambda_{\max}(M)$ denotes the largest eigen value of a symmetric matrix M . By the extension of the Cramér-Wold device (see Lemma 2.1 in Biscio et al. (2018)), the conditional (given \mathcal{H}_0) characteristic function of $\Sigma_{K,n}^{-\frac{1}{2}} T_{K,n}$ converges to the characteristic function of $\mathcal{N}(0, I_q)$ if and only if for any $a \in \mathbb{R}^q$, the conditional characteristic function of $(a^T \Sigma_{K,n} a)^{-1/2} a^T T_{K,n}$ converges to that of $\mathcal{N}(0, 1)$. Note that

$$a^T T_{K,n} = \sum_{k=1}^K \sum_{\mathbf{1} \in \mathcal{D}_n} \tilde{E}_{k,n}(\mathbf{1}),$$

where $\tilde{E}_{k,n}(\mathbf{1}) = a^T E_{k,n}(\mathbf{1})$ are components of a sequence of univariate random fields that satisfy the conditions of Theorem 1 when the conditions hold for the multivariate random fields $E_{k,n}(\mathbf{1})$. Thus to prove Theorem 1, it suffices to verify Theorem 1 in the univariate case $q = 1$.

Proof. In the univariate case, $\Sigma_{K,n} = \sigma_{K,n}^2 = \text{Var}_{\mathcal{H}_0}[T_{K,n}]$ is a scalar and by conditional centering (A1),

$$\sigma_{K,n}^2 = \sum_{k,k'=1}^K \sum_{\mathbf{1}, \mathbf{1}' \in \mathcal{D}_n} \mathbb{E}_{\mathcal{H}_0}[E_{k,n}(\mathbf{1}) E_{k',n}(\mathbf{1}')] = \sum_{k=1}^K \sum_{\mathbf{1}, \mathbf{1}' \in \mathcal{D}_n} \mathbb{E}_{\mathcal{H}_0}[E_{k,n}(\mathbf{1}) E_{k,n}(\mathbf{1}')].$$

In case (iii), we define truncation distances l_K and m_n such that as $n \rightarrow \infty$,

$$m_n \rightarrow \infty \quad \text{and} \quad \frac{m_n^{2d}}{\sqrt{|\mathcal{D}_n|}} \rightarrow 0, \quad (6)$$

and as $K \rightarrow \infty$,

$$l_K \rightarrow \infty \quad \text{and} \quad \frac{l_K}{\sqrt{K}} \rightarrow 0. \quad (7)$$

In case (i) we modify the definition by setting $l_K = \infty$ and only assume (6), and in case (ii) we set $m_n = \infty$ and only assume (7).

We further define

$$V_{K,n}^2 = \sum_{k=1}^K \sum_{(\mathbf{l}, \mathbf{j}) \in \mathcal{J}_{K,n}} \mathbb{E}_{\mathcal{H}_0} [E_{k,n}(\mathbf{l}) E_{k,n}(\mathbf{j})],$$

where $\mathcal{J}_{K,n} = \{(\mathbf{l}, \mathbf{j}) : \mathbf{l}, \mathbf{j} \in \mathcal{D}_n, d(\mathbf{l}, \mathbf{j}) \leq m_n\}$.

We define $\bar{S}_{K,n} = T_{K,n}/V_{K,n}$ and

$$\bar{S}_{K,n}(k, \mathbf{l}) = \frac{1}{V_{K,n}} \sum_{(l, \mathbf{j}) \in \mathcal{I}_{K,n}(k, \mathbf{l})} E_{l,n}(\mathbf{j}),$$

where

$$\mathcal{I}_{K,n}(k, \mathbf{l}) = \{(l, \mathbf{j}) : k \leq l \leq K, l - k \leq l_K, (\mathbf{l}, \mathbf{j}) \in \mathcal{J}_{K,n}\}.$$

Then,

$$\mathbb{E}_{\mathcal{H}_0} \left[\left(\frac{V_{K,n}}{\sigma_{K,n}} \bar{S}_{K,n} \right)^2 \right] = \mathbb{E}_{\mathcal{H}_0} \left[\left(\frac{T_{K,n}}{\sigma_{K,n}} \right)^2 \right] = \frac{1}{\sigma_{K,n}^2} \mathbb{V}\text{ar}_{\mathcal{H}_0} [T_{K,n}] = 1$$

and in Section 4.3 of the supplementary material it is shown that, uniformly in K ,

$$\left| 1 - \frac{V_{K,n}^2}{\sigma_{K,n}^2} \right| \xrightarrow[n \rightarrow \infty]{a.s.} 0. \quad (8)$$

Moreover, using $|e^{ih} - 1| \leq |h|$, $h \in \mathbb{R}$,

$$\begin{aligned} & \left| \mathbb{E}_{\mathcal{H}_0} \left[\exp \left(it \frac{T_{K,n}}{\sigma_{K,n}} \right) \right] - \mathbb{E}_{\mathcal{H}_0} \left[\exp (it \bar{S}_{K,n}) \right] \right| \\ &= \left| \mathbb{E}_{\mathcal{H}_0} \left[\exp (it \bar{S}_{K,n}) \left(\exp \left\{ it \bar{S}_{K,n} \left(\frac{V_{K,n}}{\sigma_{K,n}} - 1 \right) \right\} - 1 \right) \right] \right| \\ &\leq \mathbb{E}_{\mathcal{H}_0} \left[\left| \exp \left\{ it \bar{S}_{K,n} \left(\frac{V_{K,n}}{\sigma_{K,n}} - 1 \right) \right\} - 1 \right| \right] \leq \mathbb{E}_{\mathcal{H}_0} \left[\left| t \bar{S}_{K,n} \left(\frac{V_{K,n}}{\sigma_{K,n}} - 1 \right) \right| \right] \\ &\leq |t| \left| \frac{V_{K,n}}{\sigma_{K,n}} - 1 \right| \left(\mathbb{E}_{\mathcal{H}_0} [\bar{S}_{K,n}^2] \right)^{\frac{1}{2}} = |t| \left| 1 - \frac{V_{K,n}}{\sigma_{K,n}} \right| \frac{\sigma_{K,n}}{V_{K,n}}. \end{aligned}$$

It therefore follows from (8) that $\mathbb{E}_{\mathcal{H}_0} \left[\exp \left(it \frac{T_{K,n}}{\sigma_{K,n}} \right) \right] \rightarrow \exp(t^2/2)$ for any $t \in \mathbb{R}$ if and only if $\mathbb{E}_{\mathcal{H}_0} \left[\exp (it \bar{S}_{K,n}) \right] \rightarrow \exp(t^2/2)$. However, by Lemma 2 in Bolthausen (1982) (see also the discussion in Biscio et al. (2018)), the latter convergence holds if for any $t \in \mathbb{R}$,

$$\mathbb{E}_{\mathcal{H}_0} \left[(it - \bar{S}_{K,n}) e^{it \bar{S}_{K,n}} \right] \rightarrow 0. \quad (9)$$

Following [Bolthausen \(1982\)](#),

$$(it - \bar{S}_{K,n}) e^{it\bar{S}_{K,n}} = A_1 - A_2 - A_3,$$

where

$$A_1 = ite^{it\bar{S}_{K,n}} \left(1 - \frac{1}{V_{K,n}} \sum_{k=1}^K \sum_{\mathbf{l} \in \mathcal{D}_n} E_{k,n}(\mathbf{l}) \bar{S}_{K,n}(k, \mathbf{l}) \right), \quad (10)$$

$$A_2 = \frac{e^{it\bar{S}_{K,n}}}{V_{K,n}} \sum_{k=1}^K \sum_{\mathbf{l} \in \mathcal{D}_n} E_{k,n}(\mathbf{l}) (1 - it\bar{S}_{K,n}(k, \mathbf{l}) - \exp\{-it\bar{S}_{K,n}(k, \mathbf{l})\}), \quad (11)$$

$$A_3 = \frac{1}{V_{K,n}} \sum_{k=1}^K \sum_{\mathbf{l} \in \mathcal{D}_n} E_{k,n}(\mathbf{l}) \exp\{it(\bar{S}_{K,n} - \bar{S}_{K,n}(k, \mathbf{l}))\}. \quad (12)$$

Hence, (9) follows from the convergences to zero of $\mathbb{E}_{\mathcal{H}_0}[A_1]$, $\mathbb{E}_{\mathcal{H}_0}[A_2]$, and $\mathbb{E}_{\mathcal{H}_0}[A_3]$.

4.1.1 Convergence of $\mathbb{E}_{\mathcal{H}_0}[A_1]$

First note that

$$\begin{aligned} \mathbb{E}_{\mathcal{H}_0} \left[\frac{1}{V_{K,n}} \sum_{k=1}^K \sum_{\mathbf{l} \in \mathcal{D}_n} E_{k,n}(\mathbf{l}) \bar{S}_{K,n}(k, \mathbf{l}) \right] &= \frac{1}{V_{K,n}^2} \sum_{k=1}^K \sum_{\mathbf{l} \in \mathcal{D}_n} \sum_{(l, \mathbf{j}) \in \mathcal{I}_{K,n}(k, \mathbf{l})} \mathbb{E}_{\mathcal{H}_0} [E_{k,n}(\mathbf{l}) E_{l,n}(\mathbf{j})] \\ &= \frac{1}{V_{K,n}^2} \sum_{(\mathbf{l}, \mathbf{j}) \in \mathcal{J}_{K,n}} \mathbb{E}_{\mathcal{H}_0} [E_{k,n}(\mathbf{l}) E_{k,n}(\mathbf{j})] = 1, \end{aligned}$$

where the second equality is due to conditional centering. Thus,

$$\begin{aligned} \mathbb{E}_{\mathcal{H}_0} [|A_1|^2] &\leq \frac{t^2}{V_{K,n}^4} \text{Var}_{\mathcal{H}_0} \left[\sum_{k=1}^K \sum_{\mathbf{l} \in \mathcal{D}_n} \sum_{(l, \mathbf{j}) \in \mathcal{I}_{K,n}(k, \mathbf{l})} E_{k,n}(\mathbf{l}) E_{l,n}(\mathbf{j}) \right] \\ &= \frac{t^2}{V_{K,n}^4} \sum_{k, k'=1}^K \sum_{\mathbf{l}, \mathbf{l}' \in \mathcal{D}_n} \sum_{\substack{(l, \mathbf{j}) \in \mathcal{I}_{K,n}(k, \mathbf{l}) \\ (l', \mathbf{j}') \in \mathcal{I}_{K,n}(k', \mathbf{l}')}} \text{Cov}_{\mathcal{H}_0} [E_{k,n}(\mathbf{l}) E_{l,n}(\mathbf{j}), E_{k',n}(\mathbf{l}') E_{l',n}(\mathbf{j}')] \\ &= \frac{t^2}{V_{K,n}^4} (2a + b + 2c + d), \end{aligned} \quad (13)$$

where

$$\begin{aligned}
a &= \sum_{k=2}^K \sum_{l'=1}^{k-1} \sum_{\substack{k'=1 \\ l'-k' \leq l_K}}^{l'} \sum_{(\mathbf{l}, \mathbf{j}), (\mathbf{l}', \mathbf{j}') \in \mathcal{J}_{K,n}} \text{Cov}_{\mathcal{H}_0} [E_{k,n}(\mathbf{l})E_{k,n}(\mathbf{j}), E_{k',n}(\mathbf{l}')E_{l',n}(\mathbf{j}')] \\
b &= \sum_{l=2}^K \sum_{\substack{k, k'=1 \\ l-k, l-k' \leq l_K}}^{l-1} \sum_{(\mathbf{l}, \mathbf{j}), (\mathbf{l}', \mathbf{j}') \in \mathcal{J}_{K,n}} \text{Cov}_{\mathcal{H}_0} [E_{k,n}(\mathbf{l})E_{l,n}(\mathbf{j}), E_{k',n}(\mathbf{l}')E_{l,n}(\mathbf{j}')] \\
c &= \sum_{k=2}^K \sum_{\substack{k'=1 \\ k-k' \leq l_K}}^{k-1} \sum_{(\mathbf{l}, \mathbf{j}), (\mathbf{l}', \mathbf{j}') \in \mathcal{J}_{K,n}} \text{Cov}_{\mathcal{H}_0} [E_{k,n}(\mathbf{l})E_{k,n}(\mathbf{j}), E_{k',n}(\mathbf{l}')E_{k,n}(\mathbf{j}')] \\
d &= \sum_{k=1}^K \sum_{(\mathbf{l}, \mathbf{j}), (\mathbf{l}', \mathbf{j}') \in \mathcal{J}_{K,n}} \text{Cov}_{\mathcal{H}_0} [E_{k,n}(\mathbf{l})E_{k,n}(\mathbf{j}), E_{k,n}(\mathbf{l}')E_{k,n}(\mathbf{j}')]
\end{aligned}$$

are corresponding to cases a)-d) in Section 4 in the supplementary material. We control a-d in each of the cases (i)-(iii) using the bounds in Table 1.

Case (i)

In case (i)

$$\begin{aligned}
|a| &\leq 8K^3 M_\epsilon^4 \sum_{(\mathbf{l}, \mathbf{j}), (\mathbf{l}', \mathbf{j}') \in \mathcal{J}_{K,n}} \bar{\alpha}_K (d(\{\mathbf{l}, \mathbf{j}\}, \{\mathbf{l}', \mathbf{j}'\}) - 2R) \\
|b| &\leq 8K^3 M_\epsilon^4 \sum_{(\mathbf{l}, \mathbf{j}), (\mathbf{l}', \mathbf{j}') \in \mathcal{J}_{K,n}} \bar{\alpha}_K (d(\{\mathbf{l}, \mathbf{j}\}, \{\mathbf{l}', \mathbf{j}'\}) - 2R) \\
|c| &\leq 8K^2 M_\epsilon^4 \sum_{(\mathbf{l}, \mathbf{j}), (\mathbf{l}', \mathbf{j}') \in \mathcal{J}_{K,n}} \bar{\alpha}_K (d(\{\mathbf{l}, \mathbf{j}\}, \{\mathbf{l}', \mathbf{j}'\}) - 2R) \\
|d| &\leq 8K M_\epsilon^4 \sum_{(\mathbf{l}, \mathbf{j}), (\mathbf{l}', \mathbf{j}') \in \mathcal{J}_{K,n}} \bar{\alpha}_K (d(\{\mathbf{l}, \mathbf{j}\}, \{\mathbf{l}', \mathbf{j}'\}) - 2R).
\end{aligned}$$

If $(\mathbf{l}, \mathbf{j}), (\mathbf{l}', \mathbf{j}') \in \mathcal{J}_{K,n}$, then by the triangular inequality,

$$d(\mathbf{l}, \mathbf{l}') \leq \min \{d(\mathbf{l}, \mathbf{j}') + m_n, d(\mathbf{l}', \mathbf{j}) + m_n, d(\mathbf{j}, \mathbf{j}') + 2m_n\},$$

which means that $d(\mathbf{l}, \mathbf{l}') - 2m_n \leq d(\{\mathbf{l}, \mathbf{j}\}, \{\mathbf{l}', \mathbf{j}'\})$. We split the sum over $(\mathbf{l}, \mathbf{j}), (\mathbf{l}', \mathbf{j}') \in \mathcal{J}_{K,n}$ in the above inequalities into two terms, those with $d(\mathbf{l}, \mathbf{l}') \geq 3m_n$ and the rest. Since for any $\mathbf{l} \in \mathbb{Z}^d$, the cardinality of $\{\mathbf{j} \in \mathbb{Z}^d : d(\mathbf{l}, \mathbf{j}) \leq m_n\}$ is at most $(2\lceil m_n \rceil + 1)^d$ and the cardinality of $\{\mathbf{l}' \in \mathbb{Z}^d : d(\mathbf{l}, \mathbf{l}') = l\}$ is less than or equal to $3^d l^{d-1}$ for $l \in \mathbb{N}$ (Lemma F4 in [Biscio and Waagepetersen \(2019\)](#)),

we have

$$\begin{aligned}
& \sum_{\substack{(\mathbf{l}, \mathbf{j}), (\mathbf{l}', \mathbf{j}') \in \mathcal{J}_{K,n} \\ d(\mathbf{l}, \mathbf{l}') \geq 3m_n}} \bar{\alpha}_K (d(\{\mathbf{l}, \mathbf{j}\}, \{\mathbf{l}', \mathbf{j}'\}) - 2R) \\
& \leq (2\lceil m_n \rceil + 1)^{2d} \sum_{\substack{\mathbf{l}, \mathbf{l}' \in \mathcal{D}_n \\ d(\mathbf{l}, \mathbf{l}') \geq 3m_n}} \bar{\alpha}_K (d(\mathbf{l}, \mathbf{l}') - 2m_n - 2R) \\
& \leq (2\lceil m_n \rceil + 1)^{2d} \sum_{\mathbf{l} \in \mathcal{D}_n} \sum_{\substack{\mathbf{l}' \in \mathbb{Z}^d \\ d(\mathbf{l}, \mathbf{l}') \geq 3m_n}} \bar{\alpha}_K (d(\mathbf{l}, \mathbf{l}') - 2m_n - 2R) \\
& \leq (2\lceil m_n \rceil + 1)^{2d} \sum_{\mathbf{l} \in \mathcal{D}_n} \sum_{l=\lceil 3m_n \rceil}^{\infty} \sum_{\substack{\mathbf{l}' \in \mathbb{Z}^d \\ d(\mathbf{l}, \mathbf{l}')=l}} \bar{\alpha}_K (l - 2m_n - 2R) \\
& \leq 3^d |\mathcal{D}_n| (2\lceil m_n \rceil + 1)^{2d} \sum_{l=\lceil 3m_n \rceil}^{\infty} l^{d-1} \bar{\alpha}_K (l - 2m_n - 2R).
\end{aligned}$$

On the other hand, if $(\mathbf{l}, \mathbf{j}), (\mathbf{l}', \mathbf{j}') \in \mathcal{J}_{K,n}$ and $d(\mathbf{l}, \mathbf{l}') < 3m_n$, then $d(\mathbf{l}, \mathbf{j}') < 4m_n$, $d(\mathbf{l}', \mathbf{j}) < 4m_n$, $d(\mathbf{j}, \mathbf{j}') < 5m_n$ and $d(\{\mathbf{l}, \mathbf{j}\}, \{\mathbf{l}', \mathbf{j}'\}) < 3m_n$. Hence,

$$\begin{aligned}
& \sum_{\substack{(\mathbf{l}, \mathbf{j}), (\mathbf{l}', \mathbf{j}') \in \mathcal{J}_{K,n} \\ d(\mathbf{l}, \mathbf{l}') < 3m_n}} \bar{\alpha}_K (d(\{\mathbf{l}, \mathbf{j}\}, \{\mathbf{l}', \mathbf{j}'\}) - 2R) \\
& \leq \sum_{\substack{\mathbf{l}, \mathbf{l}' \in \mathcal{D}_n \\ d(\mathbf{l}, \mathbf{l}') < 3m_n}} \sum_{\substack{\mathbf{j} \in \mathcal{D}_n \\ d(\mathbf{l}, \mathbf{j}) \leq m_n}} \sum_{\substack{\mathbf{j}' \in \mathcal{D}_n \\ d(\mathbf{l}', \mathbf{j}') \leq m_n}} \sum_{l=0}^{\lceil 3m_n \rceil} I[d(\{\mathbf{l}, \mathbf{j}\}, \{\mathbf{l}', \mathbf{j}'\}) = l] \bar{\alpha}_K (l - 2R)
\end{aligned}$$

We can further split the sum according to whether $d(\{\mathbf{l}, \mathbf{j}\}, \{\mathbf{l}', \mathbf{j}'\})$ is equal to $d(\mathbf{l}, \mathbf{l}')$, $d(\mathbf{l}, \mathbf{j}')$, $d(\mathbf{j}, \mathbf{l}')$, or $d(\mathbf{j}, \mathbf{j}')$. Considering the first possibility,

$$\begin{aligned}
& \sum_{\substack{\mathbf{l}, \mathbf{l}' \in \mathcal{D}_n \\ d(\mathbf{l}, \mathbf{l}') < 3m_n}} \sum_{\substack{\mathbf{j} \in \mathcal{D}_n \\ d(\mathbf{l}, \mathbf{j}) \leq m_n}} \sum_{\substack{\mathbf{j}' \in \mathcal{D}_n \\ d(\mathbf{l}', \mathbf{j}') \leq m_n}} \sum_{l=0}^{\lceil 3m_n \rceil} I[d(\mathbf{l}, \mathbf{l}') = l] \bar{\alpha}_K (l - 2R) \\
& \leq 3^d |\mathcal{D}_n| (2\lceil m_n \rceil + 1)^{2d} \sum_{l=0}^{\lceil 3m_n \rceil} l^{d-1} \bar{\alpha}_K (l - 2R).
\end{aligned}$$

In the other cases we just replace $(2\lceil m_n \rceil + 1)^{2d}$ by $(6\lceil m_n \rceil + 1)^d (2\lceil m_n \rceil + 1)^d$. We can thus conclude

$$\begin{aligned}
& \sum_{\substack{(\mathbf{l}, \mathbf{j}), (\mathbf{l}', \mathbf{j}') \in \mathcal{J}_{K,n} \\ d(\mathbf{l}, \mathbf{l}') < 3m_n}} \bar{\alpha}_K (d(\{\mathbf{l}, \mathbf{j}\}, \{\mathbf{l}', \mathbf{j}'\}) - 2R) \\
& \leq 4(3^d) |\mathcal{D}_n| (6\lceil m_n \rceil + 1)^{2d} \sum_{l=0}^{\lceil 3m_n \rceil} l^{d-1} \bar{\alpha}_K (l - 2R).
\end{aligned}$$

By condition [\(A3-1\)](#) $\bar{\alpha}_K(l) = O(l^\eta)$, $\eta + d < 0$, and hence

$$\sum_{l=\lceil 3m_n \rceil}^{\infty} l^{d-1} \bar{\alpha}_K (l - 2m_n - 2R) \leq \text{constant} \sum_{l=\lceil 3m_n \rceil}^{\infty} l^{\eta+d-1} \left(1 - \frac{2m_n}{l} - \frac{2R}{l}\right)^\eta \xrightarrow{n \rightarrow \infty} 0$$

and

$$\sum_{l=0}^{\lfloor 3m_n \rfloor} l^{d-1} \bar{\alpha}_K(l-2R) \leq \sum_{l=0}^{\infty} l^d \bar{\alpha}_K(l-2R) \leq \text{constant} \sum_{l=0}^{\infty} l^{d+\eta} < \infty.$$

Thus, $a = O(K^3 |\mathcal{D}_n| m_n^{2d})$, $b = O(K^3 |\mathcal{D}_n| m_n^{2d})$, $c = O(K^2 |\mathcal{D}_n| m_n^{2d})$ and $d = O(K |\mathcal{D}_n| m_n^{2d})$ which ensures that $V_{K,n}^4 \mathbb{E}_{\mathcal{H}_0} [|A_1|^2] = O(K^3 |\mathcal{D}_n| m_n^{2d})$ as $n \rightarrow \infty$.

Case (ii)

In case (ii)

$$\begin{aligned} |a| &\leq 8M_\epsilon^4 |\mathcal{D}_n|^4 l_K \sum_{l'=1}^{K-1} \sum_{z=0}^{K-1-l'} \bar{\alpha}_n(z) \leq 8M_\epsilon^4 |\mathcal{D}_n|^4 l_K K \sum_{z=0}^{\infty} \bar{\alpha}_n(z) \\ |c| &\leq 8M_\epsilon^4 |\mathcal{D}_n|^4 \sum_{k'=1}^{K-1} \sum_{k=k'+1}^K \bar{\alpha}_n(k-1-k') \leq 8M_\epsilon^4 |\mathcal{D}_n|^4 K \sum_{z=0}^{\infty} \bar{\alpha}_n(z) \\ |d| &\leq 2M_\epsilon^4 |\mathcal{D}_n|^4 K \end{aligned}$$

and

$$\begin{aligned} |b| &\leq M_\epsilon^4 |\mathcal{D}_n|^4 \sum_{l=2}^K \sum_{\substack{k,k'=1 \\ l-k, l-k' \leq l_K}}^{l-1} (8\bar{\alpha}_n(l-1-\max\{k, k'\}))^{I[k \neq k']} \\ &\leq M_\epsilon^4 |\mathcal{D}_n|^4 \left(16 \sum_{l=2}^K \sum_{\substack{1 \leq k < k' \leq l-1 \\ l-k, l-k' \leq l_K}} \bar{\alpha}_n(l-1-k') + Kl_K \right) \\ &\leq M_\epsilon^4 |\mathcal{D}_n|^4 Kl_K \left(16 \sum_{z=0}^{\infty} \bar{\alpha}_n(z) + 1 \right). \end{aligned}$$

By condition (A3-2), $\bar{\alpha}_n(z) = O(z^\psi)$, $\psi < -1$, and $\sum_{z=0}^{\infty} \bar{\alpha}_n(z) \leq \text{constant} \sum_{z=0}^{\infty} z^\psi < \infty$. Hence a, b, c, d are all $O(Kl_K |\mathcal{D}_n|^4)$.

Case (iii)

In case (iii)

$$\begin{aligned} |a| &\leq 8M_\epsilon^4 \sum_{k=2}^K \sum_{l'=1}^{k-1} \sum_{\substack{1 \leq k' \leq l' \\ l'-k' \leq l_K}} \sum_{(\mathbf{l}, \mathbf{j}), (\mathbf{l}', \mathbf{j}') \in \mathcal{J}_{K,n}} \check{\alpha}(k-1-l', d(\{\mathbf{l}, \mathbf{j}\}, \{\mathbf{l}', \mathbf{j}'\}) - 2R) \\ |b| &\leq 8M_\epsilon^4 \sum_{l=2}^K \sum_{\substack{1 \leq k, k' \leq l-1 \\ l-k, l-k' \leq l_K}} \sum_{(\mathbf{l}, \mathbf{j}), (\mathbf{l}', \mathbf{j}') \in \mathcal{J}_{K,n}} \check{\alpha}(I[k \neq k'](l-1-\max\{k, k'\}, d(\{\mathbf{l}, \mathbf{j}\}, \{\mathbf{l}', \mathbf{j}'\}) - 2R) \\ |c| &\leq 64M_\epsilon^4 \sum_{k=2}^K \sum_{\substack{1 \leq k' \leq k-1 \\ k-k' \leq l_K}} \sum_{(\mathbf{l}, \mathbf{j}), (\mathbf{l}', \mathbf{j}') \in \mathcal{J}_{K,n}} \check{\alpha}(k-1-k', d(\{\mathbf{l}, \mathbf{j}\}, \{\mathbf{l}', \mathbf{j}'\}) - 2R) \\ |d| &\leq 8M_\epsilon^4 K \sum_{(\mathbf{l}, \mathbf{j}), (\mathbf{l}', \mathbf{j}') \in \mathcal{J}_{K,n}} \check{\alpha}(0, d(\{\mathbf{l}, \mathbf{j}\}, \{\mathbf{l}', \mathbf{j}'\}) - 2R). \end{aligned}$$

By condition [\(A3-3\)](#), using similar arguments as for cases [\(i\)](#) and [\(ii\)](#), it follows that $V_{K,n}^4 \mathbb{E}_{\mathcal{H}_0}[|A_1|^2]$ is $O(l_K K m_n^{2d} |\mathcal{D}_n|)$.

In summary

$$V_{K,n}^4 \mathbb{E}_{\mathcal{H}_0}[|A_1|^2] = \begin{cases} O(K^3 m_n^{2d} |\mathcal{D}_n|) & \text{in case [\(i\)](#)} \\ O(l_K K |\mathcal{D}_n|^4) & \text{in case [\(ii\)](#)} \\ O(l_K K m_n^{2d} |\mathcal{D}_n|) & \text{if case [\(iii\)](#)} \end{cases} .$$

By assumption [\(A5-1\)](#), [\(A5-2\)](#) or [\(A5-3\)](#), $|\mathcal{D}_n|/\sigma_{K,n}^2$, $K/\sigma_{K,n}^2$ or $|\mathcal{D}_n|K/\sigma_{K,n}^2$ are $O(1)$ as K , n or both tends to infinity. It follows from [\(8\)](#) that

$$\mathbb{E}_{\mathcal{H}_0}[|A_1|^2] = \begin{cases} O(K^3 m_n^{2d} |\mathcal{D}_n|^{-1}) & \text{in case [\(i\)](#)} \\ O(l_K K^{-1} |\mathcal{D}_n|^4) & \text{in case [\(ii\)](#)} \\ O(l_K K^{-1} m_n^{2d} |\mathcal{D}_n|^{-1}) & \text{in case [\(iii\)](#)} \end{cases}$$

Thus by [\(6\)](#) and [\(7\)](#), $\mathbb{E}_{\mathcal{H}_0}[A_1]$ converges to zero in all of cases [\(i\)](#)-[\(iii\)](#).

4.1.2 Convergence of $\mathbb{E}_{\mathcal{H}_0}[A_2]$

According to Taylor's formula, there exists a constant c so that

$$|1 - it\bar{S}_{K,n}(k, \mathbf{1}) - \exp\{-it\bar{S}_{K,n}(k, \mathbf{1})\}| \leq ct^2 \bar{S}_{K,n}^2(k, \mathbf{1}).$$

We thus get

$$\begin{aligned} \mathbb{E}_{\mathcal{H}_0}[|A_2|] &\leq \frac{ct^2}{V_{K,n}} \sum_{k=1}^K \sum_{\mathbf{1} \in \mathcal{D}_n} \mathbb{E}_{\mathcal{H}_0}[|E_{k,n}(\mathbf{1})| \bar{S}_{K,n}^2(k, \mathbf{1})] \\ &= \frac{ct^2}{V_{K,n}^3} \sum_{k=1}^K \sum_{\mathbf{1} \in \mathcal{D}_n} \sum_{(l, \mathbf{j}), (l', \mathbf{j}') \in \mathcal{I}_{K,n}(k, \mathbf{1})} \mathbb{E}_{\mathcal{H}_0}[|E_{k,n}(\mathbf{1})| E_{l,n}(\mathbf{j}) E_{l',n}(\mathbf{j}')]. \end{aligned}$$

By conditional centering and since $l, l' \geq k$, $\mathbb{E}_{\mathcal{H}_0}[|E_{k,n}(\mathbf{1})| E_{l,n}(\mathbf{j}) E_{l',n}(\mathbf{j}')]$ is zero unless $l = l'$. By Hölder's inequality

$$|\mathbb{E}_{\mathcal{H}_0}[|E_{k,n}(\mathbf{1})| E_{l,n}(\mathbf{j}) E_{l,n}(\mathbf{j}')]| \leq (\mathbb{E}_{\mathcal{H}_0}[|E_{k,n}(\mathbf{1})|^3] \mathbb{E}_{\mathcal{H}_0}[|E_{l,n}(\mathbf{j})|^3] \mathbb{E}_{\mathcal{H}_0}[|E_{l,n}(\mathbf{j}')|^3])^{\frac{1}{3}},$$

and hence in case [\(i\)](#) from [\(A4-1\)](#), we obtain

$$\begin{aligned} \mathbb{E}_{\mathcal{H}_0}[|A_2|] &\leq \frac{ct^2}{V_{K,n}^3} \sum_{k=1}^K \sum_{\mathbf{1} \in \mathcal{D}_n} \sum_{l=k}^K \sum_{\substack{\mathbf{j}, \mathbf{j}' \in \mathcal{D}_n \\ d(\mathbf{1}, \mathbf{j}) \leq m_n, d(\mathbf{1}, \mathbf{j}') \leq m_n}} \mathbb{E}_{\mathcal{H}_0}[|E_{k,n}(\mathbf{1})| E_{l,n}(\mathbf{j}) E_{l,n}(\mathbf{j}')] \\ &\leq \frac{ct^2}{V_{K,n}^3} \sum_{k=1}^K \sum_{\mathbf{1} \in \mathcal{D}_n} \sum_{l=k}^K \sum_{\substack{\mathbf{j}, \mathbf{j}' \in \mathcal{D}_n \\ d(\mathbf{1}, \mathbf{j}) \leq m_n, d(\mathbf{1}, \mathbf{j}') \leq m_n}} M_\epsilon^3 \\ &\leq \frac{ct^2}{V_{K,n}^3} K^2 |\mathcal{D}_n| (2\lceil m_n \rceil + 1)^{2d} M_\epsilon^3 \end{aligned}$$

since the cardinality of $\{\mathbf{j} \in \mathbb{Z}^d : d(\mathbf{1}, \mathbf{j}) \leq m_n\}$ is at most $(2\lceil m_n \rceil + 1)^d$. In the case (ii), (A4-2) implies that

$$\mathbb{E}_{\mathcal{H}_0}[|A_2|] \leq \frac{ct^2}{V_{K,n}^3} \sum_{k=1}^K \sum_{\mathbf{1}, \mathbf{j}, \mathbf{j}' \in \mathcal{D}_n} \sum_{\substack{k \leq l \leq K \\ l-k \leq l_K}} M_\epsilon^3 \leq \frac{ct^2}{V_{K,n}^3} K |\mathcal{D}_n|^3 l_K M_\epsilon^3.$$

In case (iii), it follows from (A4-3) that

$$\mathbb{E}_{\mathcal{H}_0}[|A_2|] \leq \frac{ct^2}{V_{K,n}^3} K |\mathcal{D}_n| l_K (2\lceil m_n \rceil + 1)^{2d} M_\epsilon^3.$$

Since $|\mathcal{D}_n|^{3/2}/V_{K,n}^3$, $K^{3/2}/V_{K,n}^3$ and $|\mathcal{D}_n|^{3/2}K^{3/2}/V_{K,n}^3$ are $O(1)$ in respectively cases (i)-(iii) and using (6) and (7), $\mathbb{E}_{\mathcal{H}_0}[|A_2|]$ converges to zero in each of the cases (i)-(iii).

4.1.3 Convergence of $\mathbb{E}_{\mathcal{H}_0}[A_3]$

Note that

$$|\mathbb{E}_{\mathcal{H}_0}[A_3]| \leq \frac{1}{V_{K,n}} \sum_{\mathbf{1} \in \mathcal{D}_n} \sum_{k=1}^K |\text{Cov}_{\mathcal{H}_0} [E_{k,n}(\mathbf{1}), \exp \{it (\bar{S}_{K,n} - \bar{S}_{K,n}(k, \mathbf{1}))\}]|.$$

We can partition the difference $\bar{S}_{K,n} - \bar{S}_{K,n}(k, \mathbf{1})$ as $B_1 + B_2$, where

$$B_1 = \frac{1}{V_{K,n}} \sum_{l=1}^{k-1} \sum_{\mathbf{j} \in \mathcal{D}_n} E_{l,n}(\mathbf{j})$$

and

$$B_2 = \begin{cases} \frac{1}{V_{K,n}} \sum_{l=k}^K \sum_{\mathbf{j} \in \mathcal{D}_n : d(\mathbf{1}, \mathbf{j}) > m_n} E_{l,n}(\mathbf{j}), & \text{in case (i)} \\ \frac{1}{V_{K,n}} \sum_{l=k: l-k > l_K}^K \sum_{\mathbf{j} \in \mathcal{D}_n} E_{l,n}(\mathbf{j}), & \text{in case (ii)} \\ \frac{1}{V_{K,n}} \sum_{l=k: l-k > l_K}^K \sum_{\mathbf{j} \in \mathcal{D}_n : d(\mathbf{1}, \mathbf{j}) > m_n} E_{l,n}(\mathbf{j}) & \text{in case (iii)} \end{cases}.$$

Since B_1 is \mathcal{H}_{k-1} measurable and using conditional centering,

$$\begin{aligned} \text{Cov}_{\mathcal{H}_0} [E_{k,n}(\mathbf{1}), \exp \{it (\bar{S}_{K,n} - \bar{S}_{K,n}(k, \mathbf{1}))\}] &= \text{Cov}_{\mathcal{H}_0} [E_{k,n}(\mathbf{1}), e^{it(B_1+B_2)}] \\ &= \mathbb{E}_{\mathcal{H}_0} [e^{itB_1} \mathbb{E}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1}) e^{itB_2}]]. \end{aligned}$$

In case (i), B_2 is a function of $\{E_{l,n}(\mathbf{j}) : l = k, \dots, K, \mathbf{j} \in \mathcal{D}_n, d(\mathbf{1}, \mathbf{j}) > m_n\}$ and

$$\begin{aligned} \mathbb{E}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1}) e^{itB_2}] &= \mathbb{E}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1}) \mathbb{E}_{\mathcal{H}_k} [e^{itB_2}]] \\ &= \text{Cov}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1}), \mathbb{E}_{\mathcal{H}_k} [e^{itB_2}]] \end{aligned}$$

The localization assumption (A2-1) implies that $\mathbb{E}_{\mathcal{H}_k} [e^{itB_2}]$ is measurable with respect to $\mathcal{H}_{0,k,\cup_{\mathbf{j} \in \mathcal{D}_n: d(\mathbf{1},\mathbf{j}) > m_n} C(\mathbf{j})}$ and hence

$$\begin{aligned} & \left| \mathbb{E}_{\mathcal{H}_0} [e^{itB_1} \text{Cov}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1}), e^{itB_2}]] \right| \\ & \leq 8M_\epsilon \mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{\mathcal{H}_{k-1}} \left(\mathcal{H}_{0,k,C(\mathbf{1})}, \mathcal{H}_{0,k,\cup_{\mathbf{j} \in \mathcal{D}_n: d(\mathbf{1},\mathbf{j}) > m_n} C(\mathbf{j})} \right) \right)^{\frac{p-2}{p}} \right] \\ & \leq 8M_\epsilon \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{R^2, \infty, K}^{(K, \mathbb{R}^d)} (0, d(C(\mathbf{1}), \cup_{\mathbf{j} \in \mathcal{D}_n: d(\mathbf{1},\mathbf{j}) > m_n} C(\mathbf{j})) \right)^{\frac{p-2}{p-1}} \right) \right]^{\frac{p-1}{p}} \right) \\ & \leq 8M_\epsilon \bar{\alpha}_K (m_n - 2R) \leq \text{constant } m_n^\eta. \end{aligned}$$

using (2.1) from the supplementary material for the first inequality and (A3-1) for the last.

In case (ii), B_2 is a function of $\{E_{l,n}(\mathbf{j}) : l = k + \lfloor l_K \rfloor + 1, \dots, K, \mathbf{j} \in \mathcal{D}_n\}$ and

$$\begin{aligned} \mathbb{E}_{\mathcal{H}_{k-1}} [E_{k,n} e^{itB_2}] &= \mathbb{E}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1}) \mathbb{E}_{\mathcal{H}_{k+\lfloor l_K \rfloor}} [e^{itB_2}]] \\ &= \text{Cov}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1}), \mathbb{E}_{\mathcal{H}_{k+\lfloor l_K \rfloor}} [e^{itB_2}]]. \end{aligned}$$

By the localization assumption (A2-2), $\mathbb{E}_{\mathcal{H}_{k+\lfloor l_K \rfloor}} [e^{itB_2}]$ is $\mathcal{H}_{(k+l_K-r)^+, k+\lfloor l_K \rfloor, \mathcal{D}_n}$ measurable and using same type of arguments as for (i),

$$\begin{aligned} & \left| \mathbb{E}_{\mathcal{H}_0} [e^{itB_1} \text{Cov}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1}), e^{itB_2}]] \right| \\ & \leq 8M_\epsilon \mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{\mathcal{H}_{k-1}} (\mathcal{H}_{0,k,\mathcal{D}_n}, \mathcal{H}_{(k+l_K-r)^+, k+\lfloor l_K \rfloor, \mathcal{D}_n}) \right)^{\frac{p-2}{p}} \right] \\ & \leq 8M_\epsilon \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{|\mathcal{D}_n|, |\mathcal{D}_n|, r}^{\infty, \mathcal{D}_n} (l_K) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}} \\ & \leq 8M_\epsilon \bar{\alpha}_n (l_K) \leq \text{constant } l_K^\psi. \end{aligned}$$

Similarly, in case (iii),

$$\begin{aligned} \left| \mathbb{E}_{\mathcal{H}_0} [e^{itB_1} \text{Cov}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1}), e^{itB_2}]] \right| &\leq 8M_\epsilon \check{\alpha}(l_K, m_n - 2R) \\ &\leq \text{constant } l_K^\psi m_n^\eta. \end{aligned}$$

Thus $\mathbb{E}_{\mathcal{H}_0} [A_3] \rightarrow 0$ in all cases (i)-(iii). □

Supplementary material

Supplementary material for ‘A central limit theorem for a sequence of conditionally centered and α -mixing random fields’ provides supporting results and computations for the proof of Theorem 1 and for Section 3.2.

References

Besag, J. (1974). Spatial interaction and the statistical analysis of lattice systems. *Journal of the Royal Statistical Society: Series B (Methodological)*, 36(2):192–225.

- Biscio, C. A. N., Poinas, A., and Waagepetersen, R. (2018). A note on gaps in proofs of central limit theorems. *Statistics and Probability Letters*, pages 7–10.
- Biscio, C. A. N. and Svane, A. M. (2022). A functional central limit theorem for the empirical Ripley’s K-function. *Electronic Journal of Statistics*, 16(1):3060 – 3098.
- Biscio, C. A. N. and Waagepetersen, R. (2019). A general central limit theorem and a subsampling variance estimator for α -mixing point processes. *Scandinavian Journal of Statistics*, 46(4):1168–1190.
- Bolthausen, E. (1982). On the central limit theorem for stationary mixing random fields. *The Annals of Probability*, 10(4):1047–1050.
- Bradley, R. C. (2005). Basic properties of strong mixing conditions. a survey and some open questions. *Probability surveys*, 2:107–144.
- Brown, B. M. (1971). Martingale central limit theorems. *The Annals of Mathematical Statistics*, pages 59–66.
- Cheysson, F. and Lang, G. (2021). Strong-mixing rates for Hawkes processes and application to Whittle estimation from count data. 29 pages, 10 figures.
- Dahlhaus, R. (2000). Graphical interaction models for multivariate time series. *Metrika*, 51(2):157–172.
- Hall, P., Heyde, C., Birnbaum, Z., and Lukacs, E. (2014). *Martingale Limit Theory and Its Application*. Communication and Behavior. Elsevier Science.
- Heinrich, L. (2013). Asymptotic methods in statistics of random point processes. In Spodarev, E., editor, *Stochastic Geometry, Spatial Statistics and Random Fields: Asymptotic Methods*, pages 115–150. Springer Berlin Heidelberg, Berlin, Heidelberg.
- Jensen, J. L. and Künsch, H. R. (1994). On asymptotic normality of pseudo likelihood estimates for pairwise interaction processes. *Annals of the Institute of Statistical Mathematics*, 46:475–486.
- Karácsony, Z. (2006). A central limit theorem for mixing random fields. *Miskolc Mathematical Notes. A Publication of the University of Miskolc*, 7(2):147–160.
- Leonenko, N. N. (1975). Central limit theorem for m -dependent random fields in schemes related to series schemes. *Ukrainian Mathematical Journal*, 27:556–559.
- Prakasa Rao, B. (2009). Conditional independence, conditional mixing and conditional association. *Annals of the Institute of Statistical Mathematics*, 61:441–460.

Supplementary material for ‘A central limit theorem for a sequence of conditionally centered and α -mixing random fields’.

Abdollah Jalilian
Razi University, Iran
stat4aj@gmail.com

Ganggang Xu
University of Miami, United States of America
gangxu@bus.miami.edu

Rasmus Waagepetersen
Aalborg University, Denmark
rw@math.aau.dk

January 24, 2023

Abstract

This document contains supplementary material for the paper ‘A central limit theorem for a sequence of conditionally centered and α -mixing random fields’.

Keywords: central limit theorem; conditional centering; conditional α -mixing; estimating function; random field; spatio-temporal; space-time
MSC2020 subject classifications: 60F05; 60G55; 60G60; 62M30

The following sections contain results on moments, mixing coefficients, and covariances of relevance for the paper [Jalilian et al. \(2023\)](#). Cross-references starting with ‘M-’ refer to that paper to which we also refer for notational definitions. In Section 1 and Section 4 we consider the one-dimensional case $q = 1$.

1 Bounds on moments

If assumption M-(A4) holds, applications of Lyapunov’s and Hölder’s inequalities imply that for any $1 \leq k' \leq k \leq l \leq K$,

$$\mathbb{E}_{\mathcal{H}_{k'-1}} [|E_{k,n}(\mathbf{1})|^p] \leq \left(\mathbb{E}_{\mathcal{H}_{k'-1}} |E_{k,n}(\mathbf{1})|^{6+\epsilon} \right)^{\frac{p}{6+\epsilon}} \leq (d_{K,n})^{\frac{p}{6+\epsilon}} \leq M_\epsilon^p \quad (1)$$

if $1 \leq p \leq 6 + \epsilon$,

$$\begin{aligned} \mathbb{E}_{\mathcal{H}_{k'-1}} [|E_{k,n}(\mathbf{1})E_{l,n}(\mathbf{j})|^p] &\leq \left(\mathbb{E}_{\mathcal{H}_{k'-1}} [|E_{k,n}(\mathbf{1})|^{2p}] \mathbb{E}_{\mathcal{H}_{k'-1}} [|E_{l,n}(\mathbf{j})|^{2p}] \right)^{\frac{1}{2}} \\ &\leq (M_\epsilon^{4p})^{\frac{1}{2}} \leq M_\epsilon^{2p} \end{aligned} \quad (2)$$

if $1 \leq p \leq 3 + \epsilon/2$, and

$$\begin{aligned}
& \mathbb{E}_{\mathcal{H}_{k'-1}} [|E_{k,n}(\mathbf{1})E_{l,n}(\mathbf{j})E_{l',n}(\mathbf{j}')|^p] \\
& \leq \left(\mathbb{E}_{\mathcal{H}_{k'-1}} [|E_{k,n}(\mathbf{1})|^{3p}] \right)^{\frac{1}{3}} \left(\mathbb{E}_{\mathcal{H}_{k'-1}} [|E_{l,n}(\mathbf{j})E_{l',n}(\mathbf{j}')|^{\frac{3p}{2}}] \right)^{\frac{2}{3}} \\
& \leq \left(\mathbb{E}_{\mathcal{H}_{k'-1}} [|E_{k,n}(\mathbf{1})|^{3p}] \mathbb{E}_{\mathcal{H}_{k'-1}} [|E_{l,n}(\mathbf{j})|^{3p}] \mathbb{E}_{\mathcal{H}_{k'-1}} [|E_{l',n}(\mathbf{j}')|^{3p}] \right)^{\frac{1}{3}} \\
& \leq (M_\epsilon^{9p})^{\frac{1}{3}} \leq M_\epsilon^{3p} \tag{3}
\end{aligned}$$

if $1 \leq p \leq 2 + \epsilon/3$. Further, by Jensen's inequality,

$$\mathbb{E}_{\mathcal{H}_0} \left[\left| \mathbb{E}_{\mathcal{H}_{k'-1}} [E_{k,n}(\mathbf{1})E_{l,n}(\mathbf{j})]^p \right| \right] \leq \mathbb{E}_{\mathcal{H}_0} \left[\mathbb{E}_{\mathcal{H}_{k'-1}} [|E_{k,n}(\mathbf{1})E_{l,n}(\mathbf{j})|^p] \right] \leq M_\epsilon^{2p}. \tag{4}$$

2 Covariance inequalities

Let Y and Z be random variables with $\mathbb{E}_{\mathcal{H}_k} [|Y|^p] < \infty$ and $\mathbb{E}_{\mathcal{H}_k} [|Z|^p] < \infty$ for some $p > 2$ and $k \in \mathbb{Z}_+$. Then (see [Yuan and Lei \(2013\)](#), Theorem 3.4),

$$|\text{Cov}_{\mathcal{H}_k} [Y, Z]| \leq 8 (\mathbb{E}_{\mathcal{H}_k} [|Y|^p] \mathbb{E}_{\mathcal{H}_k} [|Z|^p])^{\frac{1}{p}} (\alpha_{\mathcal{H}_k} (\sigma(Y), \sigma(Z)))^{\frac{p-2}{p}}, \tag{5}$$

where $\sigma(Y) \subseteq \mathcal{F}$ denotes the σ -algebra generated by Y . In particular, using the Lyapunov's inequality,

$$\begin{aligned}
|\text{Cov}_{\mathcal{H}_0} [Y, Z]| & \leq 8 (\mathbb{E}_{\mathcal{H}_0} [|Y|^p] \mathbb{E}_{\mathcal{H}_0} [|Z|^p])^{\frac{1}{p}} (\alpha_{\mathcal{H}_0} (\sigma(Y), \sigma(Z)))^{\frac{p-2}{p}} \\
& = 8 (\mathbb{E}_{\mathcal{H}_0} [|Y|^p] \mathbb{E}_{\mathcal{H}_0} [|Z|^p])^{\frac{1}{p}} \left(\mathbb{E}_{\mathcal{H}_0} \left[(\alpha_{\mathcal{H}_0} (\sigma(Y), \sigma(Z)))^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}}. \tag{6}
\end{aligned}$$

(where the last redundant $\mathbb{E}_{\mathcal{H}_0}$ is just added for later notational convenience). In addition, by an application of Hölder's inequality,

$$\begin{aligned}
|\mathbb{E}_{\mathcal{H}_0} [\text{Cov}_{\mathcal{H}_{k-1}} [Y, Z]]| & \leq \mathbb{E}_{\mathcal{H}_0} [|\text{Cov}_{\mathcal{H}_{k-1}} [Y, Z]|] \\
& \leq 8 \mathbb{E}_{\mathcal{H}_0} \left\{ (\mathbb{E}_{\mathcal{H}_{k-1}} [|Y|^p] \mathbb{E}_{\mathcal{H}_{k-1}} [|Z|^p])^{\frac{1}{p}} (\alpha_{\mathcal{H}_{k-1}} (\sigma(Y), \sigma(Z)))^{\frac{p-2}{p}} \right\} \\
& \leq 8 \left(\mathbb{E}_{\mathcal{H}_0} \left\{ \left[(\mathbb{E}_{\mathcal{H}_{k-1}} [|Y|^p] \mathbb{E}_{\mathcal{H}_{k-1}} [|Z|^p])^{\frac{1}{p}} \right]^p \right\} \right)^{\frac{1}{p}} \left(\mathbb{E}_{\mathcal{H}_0} \left\{ \left[(\alpha_{\mathcal{H}_{k-1}} (\sigma(Y), \sigma(Z)))^{\frac{p-2}{p}} \right]^{\frac{1}{1-\frac{1}{p}}} \right\} \right)^{1-\frac{1}{p}} \\
& = 8 \left(\mathbb{E}_{\mathcal{H}_0} \left\{ \mathbb{E}_{\mathcal{H}_{k-1}} [|Y|^p] \mathbb{E}_{\mathcal{H}_{k-1}} [|Z|^p] \right\} \right)^{\frac{1}{p}} \left(\mathbb{E}_{\mathcal{H}_0} \left[(\alpha_{\mathcal{H}_{k-1}} (\sigma(Y), \sigma(Z)))^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}}. \tag{7}
\end{aligned}$$

3 Monotonicity of the α -mixing coefficients

If $m_2 < m'_2$, then

$$\{A, B \subset \mathcal{D} : d(A, B) \geq m'_2\} \subseteq \{A, B \subset \mathcal{D} : d(A, B) \geq m_2\}$$

and hence

$$\alpha_{c_1, c_2, r}^{(K, \mathcal{D})}(m_1, m_2') \leq \alpha_{c_1, c_2, r}^{(K, \mathcal{D})}(m_1, m_2).$$

Also, if $m_1 < m_1'$, then

$$\{l, k \in \mathbb{Z}_+ : 0 \leq l, k \leq K, l + m_1' \leq k\} \subseteq \{l, k \in \mathbb{Z}_+ : 0 \leq l, k \leq K, l + m_1 \leq k\}$$

and hence

$$\alpha_{c_1, c_2, r}^{(K, \mathcal{D})}(m_1', m_2) \leq \alpha_{c_1, c_2, r}^{(K, \mathcal{D})}(m_1, m_2).$$

Using the same arguments,

$$\begin{aligned} \text{for } c_1 < c_1' : \quad & \alpha_{c_1, c_2, r}^{(K, \mathcal{D})}(m_1, m_2) \leq \alpha_{c_1', c_2, r}^{(K, \mathcal{D})}(m_1, m_2), \\ \text{for } c_2 < c_2' : \quad & \alpha_{c_1, c_2, r}^{(K, \mathcal{D})}(m_1, m_2) \leq \alpha_{c_1, c_2', r}^{(K, \mathcal{D})}(m_1, m_2), \\ \text{for } r < r' : \quad & \alpha_{c_1, c_2, r}^{(K, \mathcal{D})}(m_1, m_2) \leq \alpha_{c_1, c_2, r'}^{(K, \mathcal{D})}(m_1, m_2), \\ \text{for } K < K' : \quad & \alpha_{c_1, c_2, r}^{(K, \mathcal{D})}(m_1, m_2) \leq \alpha_{c_1, c_2, r}^{(K', \mathcal{D})}(m_1, m_2), \\ \text{for } \mathcal{D} \subset \mathcal{D}' : \quad & \alpha_{c_1, c_2, r}^{(K, \mathcal{D})}(m_1, m_2) \leq \alpha_{c_1, c_2, r}^{(K, \mathcal{D}')}(m_1, m_2). \end{aligned}$$

4 Bounds on covariances

In this section we provide bounds for covariances of the form

$$\text{Cov}_{\mathcal{H}_0} [E_{k,n}(\mathbf{1})E_{l,n}(\mathbf{j}), E_{k',n}(\mathbf{1}')E_{l',n}(\mathbf{j}')] \quad (8)$$

for $1 \leq k, k' \leq K$ and $k \leq l \leq K, k' \leq l' \leq K$ in the one-dimensional case $q = 1$.

The covariance (4.1) can be rewritten as

$$\begin{aligned} & \mathbb{E}_{\mathcal{H}_0} [\text{Cov}_{\mathcal{H}_{k^*-1}} [E_{k,n}(\mathbf{1})E_{l,n}(\mathbf{j}), E_{k',n}(\mathbf{1}')E_{l',n}(\mathbf{j}')]] \\ & + \text{Cov}_{\mathcal{H}_0} [\mathbb{E}_{\mathcal{H}_{k^*-1}} [E_{k,n}(\mathbf{1})E_{l,n}(\mathbf{j})], \mathbb{E}_{\mathcal{H}_{k^*-1}} [E_{k',n}(\mathbf{1}')E_{l',n}(\mathbf{j}')]], \end{aligned}$$

where $k^* = \max\{k, l, k', l'\}$. If precisely one index coincides with k^* , then the covariance (8) is zero. Thus, the covariance (8) can only be non-zero in the cases a) $k^* = k = l > l' \geq k'$ or $k^* = k' = l' > l \geq k$, b) $k^* = l = l' > k \geq k'$, c) $k^* = k = l = l' > k'$ or $k^* = l' = k' = l > k$, and d) $k^* = k = l = k' = l'$.

We consider these cases in the following subsections. For each of the cases a) and c) we just consider one of the sub cases since the other sub cases are handled analogously. Moreover, in each subsection we consider three cases depending on whether we use condition M-(A2-1), M-(A2-2), or M-(A2-3). The cases and associated bounds are summarized in Table 1 in the main text.

4.1 Case a)

Consider the case $k^* = k = l > l' \geq k'$. Then

$$\text{Cov}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j}), E_{k',n}(\mathbf{1}')E_{l',n}(\mathbf{j}')] = 0$$

and hence the covariance (8) reduces to

$$\text{Cov}_{\mathcal{H}_0} [\mathbb{E}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j})], E_{k',n}(\mathbf{1}')E_{l',n}(\mathbf{j}')] .$$

(i): If condition (A2-1) holds, $\mathbb{E}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j})]$ and $E_{k',n}(\mathbf{l}')E_{l',n}(\mathbf{j}')$ are respectively measurable with respect to $\mathcal{H}_{0,k-1,C(\mathbf{1})\cup C(\mathbf{j})}$ and $\sigma(\mathcal{H}_{0,k',C(\mathbf{l}')}\cup\mathcal{H}_{0,l',C(\mathbf{j}')}) \subseteq \mathcal{H}_{0,l',C(\mathbf{l}')\cup C(\mathbf{j}')$. Thus, since

$$\begin{aligned} & \alpha_{\mathcal{H}_0} (\mathcal{H}_{0,k-1,C(\mathbf{1})\cup C(\mathbf{j})}, \mathcal{H}_{0,l',C(\mathbf{l}')\cup C(\mathbf{j}')}) \\ & \leq \alpha_{2R^2,2R^2,K}^{(K,\mathbb{R}^d)} (k-1-l', d(C(\mathbf{1})\cup C(\mathbf{j}), C(\mathbf{l}')\cup C(\mathbf{j}')))) \\ & \leq \alpha_{2R^2,2R^2,K}^{(K,\mathbb{R}^d)} (k-1-l', d(\{\mathbf{1},\mathbf{j}\}, \{\mathbf{l}',\mathbf{j}'\}) - 2R), \end{aligned}$$

from inequalities (2) and (6) and condition M-(A3-1) we obtain

$$\begin{aligned} & \text{Cov}_{\mathcal{H}_0} [\mathbb{E}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j})], E_{k',n}(\mathbf{l}')E_{l',n}(\mathbf{j}')] \\ & \leq 8M_\epsilon^4 \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{\mathcal{H}_0} (\mathcal{H}_{0,k-1,C(\mathbf{1})\cup C(\mathbf{j})}, \mathcal{H}_{0,l',C(\mathbf{l}')\cup C(\mathbf{j}')}) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}} \\ & \leq 8M_\epsilon^4 \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{2R^2,2R^2,K}^{(K,\mathbb{R}^d)} (k-1-l', d(\{\mathbf{1},\mathbf{j}\}, \{\mathbf{l}',\mathbf{j}'\}) - 2R) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}} \\ & \leq 8M_\epsilon^4 \bar{\alpha}_K (d(\{\mathbf{1},\mathbf{j}\}, \{\mathbf{l}',\mathbf{j}'\}) - 2R). \end{aligned}$$

(ii): If condition M-(A2-2) holds, $\mathbb{E}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j})]$ and $E_{k',n}(\mathbf{l}')E_{l',n}(\mathbf{j}')$ are respectively measurable with respect to $\mathcal{H}_{(k-r)^+,k-1,\mathcal{D}_n}$ and $\mathcal{H}_{0,l',\mathcal{D}_n}$. Thus, since

$$\alpha_{\mathcal{H}_0} (\mathcal{H}_{(k-r)^+,k-1,\mathcal{D}_n}, \mathcal{H}_{0,l',\mathcal{D}_n}) \leq \alpha_{|\mathcal{D}_n|,|\mathcal{D}_n|,r}^{(\infty,\mathcal{D}_n)} (k-1-l', 0),$$

from condition M-(A3-2) we obtain

$$\begin{aligned} & \text{Cov}_{\mathcal{H}_0} [\mathbb{E}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j})], E_{k',n}(\mathbf{l}')E_{l',n}(\mathbf{j}')] \\ & \leq 8M_\epsilon^4 \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{|\mathcal{D}_n|,|\mathcal{D}_n|,r}^{(\infty,\mathcal{D}_n)} (k-1-l', 0) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}} \\ & \leq 8M_\epsilon^4 \bar{\alpha}_n (k-1-l'). \end{aligned}$$

(iii): If condition M-(A2-3) holds, $\mathbb{E}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j})]$ and $E_{k',n}(\mathbf{l}')E_{l',n}(\mathbf{j}')$ are respectively measurable with respect to $\mathcal{H}_{(k-r)^+,k-1,C(\mathbf{1})\cup C(\mathbf{j})}$ and $\mathcal{H}_{0,l',C(\mathbf{l}')\cup C(\mathbf{j}')}$. Thus, following the same arguments as above and from condition M-(A3-3) we obtain

$$\begin{aligned} & \text{Cov}_{\mathcal{H}_0} [\mathbb{E}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j})], E_{k',n}(\mathbf{l}')E_{l',n}(\mathbf{j}')] \\ & \leq 8M_\epsilon^4 \bar{\alpha} (k-l-l', d(\{\mathbf{1},\mathbf{j}\}, \{\mathbf{l}',\mathbf{j}'\}) - 2R). \end{aligned}$$

4.2 Case b)

Consider $k^* = l = l' > k_* = \max\{k, k'\}$. Then

$$\mathbb{E}_{\mathcal{H}_{l-1}} [E_{k,n}(\mathbf{1})E_{l,n}(\mathbf{j})] = \mathbb{E}_{\mathcal{H}_{l-1}} [E_{k',n}(\mathbf{l}')E_{l,n}(\mathbf{j}')] = 0$$

and hence the covariance (8) becomes

$$\begin{aligned} & \mathbb{E}_{\mathcal{H}_0} [\text{Cov}_{\mathcal{H}_{l-1}} [E_{k,n}(\mathbf{1})E_{l,n}(\mathbf{j}), E_{k',n}(\mathbf{1}')E_{l,n}(\mathbf{j}')]] \\ &= \mathbb{E}_{\mathcal{H}_0} [E_{k,n}(\mathbf{1})E_{k',n}(\mathbf{1}')\mathbb{E}_{\mathcal{H}_{l-1}} [E_{l,n}(\mathbf{j})E_{l,n}(\mathbf{j}')]]. \end{aligned}$$

By Hölder's inequality,

$$\begin{aligned} & \left| \mathbb{E}_{\mathcal{H}_0} [E_{k,n}(\mathbf{1})E_{k',n}(\mathbf{1}')\mathbb{E}_{\mathcal{H}_{l-1}} [E_{l,n}(\mathbf{j})E_{l,n}(\mathbf{j}')]] \right| \\ & \leq (\mathbb{E}_{\mathcal{H}_0} [|E_{k,n}(\mathbf{1})E_{k',n}(\mathbf{1}')|^p])^{\frac{1}{p}} \left(\mathbb{E}_{\mathcal{H}_0} [|\text{Cov}_{\mathcal{H}_{l-1}} [E_{l,n}(\mathbf{j}), E_{l,n}(\mathbf{j}')] |^{\frac{p}{p-1}}] \right)^{\frac{p-1}{p}}. \end{aligned}$$

(i): If condition M-(A2-1) holds, $E_l(\mathbf{j})$ and $E_l(\mathbf{j}')$ are respectively $\mathcal{H}_{0,l,C(\mathbf{j})}$ and $\overline{\mathcal{H}}_{0,l,C(\mathbf{j}')}$ measurable and inequalities (1) and (5) implies that

$$|\text{Cov}_{\mathcal{H}_{l-1}} [E_{l,n}(\mathbf{j}), E_{l,n}(\mathbf{j}')] | \leq 8M_\epsilon^2 (\alpha_{\mathcal{H}_{l-1}}(\mathcal{H}_{0,l,C(\mathbf{j})}, \mathcal{H}_{0,l,C(\mathbf{j}')})) \frac{p-2}{p}.$$

It follows from the inequality (2) that

$$(\mathbb{E}_{\mathcal{H}_0} [|E_{k,n}(\mathbf{1})E_{k',n}(\mathbf{1}')|^p])^{\frac{1}{p}} \leq M_\epsilon^2.$$

Thus, since

$$\begin{aligned} \alpha_{\mathcal{H}_{l-1}} (\mathcal{H}_{0,l,C(\mathbf{j})}, \mathcal{H}_{0,l,C(\mathbf{j}')})) & \leq \alpha_{R^2, R^2, K}^{(K, \mathbb{R}^d)} (0, d(C(\mathbf{j}), C(\mathbf{j}'))) \\ & \leq \alpha_{R^2, R^2, K}^{(K, \mathbb{R}^d)} (0, d(\mathbf{j}, \mathbf{j}') - 2R) \\ & \leq \alpha_{R^2, R^2, K}^{(K, \mathbb{R}^d)} (0, d(\{\mathbf{1}, \mathbf{j}\}, \{\mathbf{1}', \mathbf{j}'\}) - 2R), \end{aligned}$$

from condition M-(A3-1) we obtain

$$\begin{aligned} & \left| \mathbb{E}_{\mathcal{H}_0} [E_{k,n}(\mathbf{1})E_{k',n}(\mathbf{1}')\mathbb{E}_{\mathcal{H}_{l-1}} [E_{l,n}(\mathbf{j})E_{l,n}(\mathbf{j}')]] \right| \\ & \leq 8M_\epsilon^4 \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{R^2, R^2, K}^{(K, \mathbb{R}^d)} (0, d(\{\mathbf{1}, \mathbf{j}\}, \{\mathbf{1}', \mathbf{j}'\}) - 2R) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}} \\ & \leq 8M_\epsilon^4 \tilde{\alpha}_K (d(\{\mathbf{1}, \mathbf{j}\}, \{\mathbf{1}', \mathbf{j}'\}) - 2R). \end{aligned}$$

(ii): On the other hand,

$$\begin{aligned} & \mathbb{E}_{\mathcal{H}_0} [E_{k,n}(\mathbf{1})E_{k',n}(\mathbf{1}')\mathbb{E}_{\mathcal{H}_{l-1}} [E_{l,n}(\mathbf{j})E_{l,n}(\mathbf{j}')]] \\ &= I[k \neq k'] \text{Cov}_{\mathcal{H}_0} [E_{k,n}(\mathbf{1})E_{k',n}(\mathbf{1}'), \mathbb{E}_{\mathcal{H}_{l-1}} [E_{l,n}(\mathbf{j})E_{l,n}(\mathbf{j}')]] \\ &+ I[k = k'] \mathbb{E}_{\mathcal{H}_0} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{1}')\mathbb{E}_{\mathcal{H}_{l-1}} [E_{l,n}(\mathbf{j})E_{l,n}(\mathbf{j}')]] \end{aligned}$$

If condition M-(A2-2) holds, $\mathbb{E}_{\mathcal{H}_{l-1}} [E_{l,n}(\mathbf{j})E_{l,n}(\mathbf{j}')]]$ and $E_{k,n}(\mathbf{1})E_{k',n}(\mathbf{1}')$ are respectively $\mathcal{H}_{(l-r)^+, l-1, \mathcal{D}_n}$ and $\mathcal{H}_{0, \max\{k, k'\}, \mathcal{D}_n}$ measurable and inequalities (2) and (6) and condition M-(A3-2) imply that

$$\begin{aligned} & \left| \text{Cov}_{\mathcal{H}_0} [E_{k,n}(\mathbf{1})E_{k',n}(\mathbf{1}'), \mathbb{E}_{\mathcal{H}_{l-1}} [E_{l,n}(\mathbf{j})E_{l,n}(\mathbf{j}')]] \right| \\ & \leq 8M_\epsilon^4 \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{\mathcal{H}_0}(\mathcal{H}_{(l-r)^+, l-1, \mathcal{D}_n}, \mathcal{H}_{0, \max\{k, k'\}, \mathcal{D}_n}) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}} \\ & \leq 8M_\epsilon^4 \tilde{\alpha}_n (l-1 - \max\{k, k'\}). \end{aligned}$$

Moreover, from the inequality (2) we obtain

$$\begin{aligned} & \mathbb{E}_{\mathcal{H}_0} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{1}')\mathbb{E}_{\mathcal{H}_{l-1}} [E_{l,n}(\mathbf{j})E_{l,n}(\mathbf{j}')]] \\ & \leq \left(\mathbb{E}_{\mathcal{H}_0} [|E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{1}')|^2] \mathbb{E}_{\mathcal{H}_0} \left[\left| \mathbb{E}_{\mathcal{H}_{l-1}} [E_{l,n}(\mathbf{j})E_{l,n}(\mathbf{j}')] \right|^2 \right] \right)^{\frac{1}{2}} \leq M_\epsilon^4. \end{aligned}$$

Thus,

$$\begin{aligned} & \left| \mathbb{E}_{\mathcal{H}_0} [E_{k,n}(\mathbf{1})E_{k',n}(\mathbf{1}')\mathbb{E}_{\mathcal{H}_{l-1}} [E_{l,n}(\mathbf{j})E_{l,n}(\mathbf{j}')]] \right| \\ & \leq M_\epsilon^4 (8\tilde{\alpha}_n(l-1-\max\{k,k'\}))^{I[k \neq k']}. \end{aligned}$$

(iii): If condition M-(A2-3) and M-(A3-3) hold and $k \neq k'$, $\mathbb{E}_{\mathcal{H}_{l-1}} [E_{l,n}(\mathbf{j})E_{l,n}(\mathbf{j}')]$ and $E_{k,n}(\mathbf{1})E_{k',n}(\mathbf{1}')$ are respectively $\mathcal{H}_{(l-r)^+, l-1, C(\mathbf{j}) \cup C(\mathbf{j}')}$ and $\mathcal{H}_{0, \max\{k, k'\}, C(\mathbf{1}) \cup C(\mathbf{1}')}$ measurable and

$$\begin{aligned} & \left| \text{Cov}_{\mathcal{H}_0} [E_{k,n}(\mathbf{1})E_{k',n}(\mathbf{1}'), \mathbb{E}_{\mathcal{H}_{l-1}} [E_{l,n}(\mathbf{j})E_{l,n}(\mathbf{j}')]] \right| \\ & \leq 8M_\epsilon^4 \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{\mathcal{H}_0}(\mathcal{H}_{(l-r)^+, l-1, C(\mathbf{j}) \cup C(\mathbf{j}')}), \mathcal{H}_{0, \max\{k, k'\}, C(\mathbf{1}) \cup C(\mathbf{1}')} \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}} \\ & \leq 8M_\epsilon^4 \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{2R^2, 2R^2, r}^{(\infty, \mathcal{D}_n)}(l-1-\max\{k, k'\}, d(\{\mathbf{1}, \mathbf{j}\}, \{\mathbf{1}', \mathbf{j}'\}) - 2R) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}} \\ & \leq 8M_\epsilon^4 \tilde{\alpha}(l-1-\max\{k, k'\}, d(\{\mathbf{1}, \mathbf{j}\}, \{\mathbf{1}', \mathbf{j}'\}) - 2R). \end{aligned}$$

However, if condition M-(A2-3) and M-(A3-3) hold and $k = k'$, we have

$$\begin{aligned} & \left| \mathbb{E}_{\mathcal{H}_0} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{1}')\mathbb{E}_{\mathcal{H}_{l-1}} [E_{l,n}(\mathbf{j})E_{l,n}(\mathbf{j}')]] \right| \\ & \leq M_\epsilon^2 \left(\mathbb{E}_{\mathcal{H}_0} \left[\left| \text{Cov}_{\mathcal{H}_{l-1}} [E_{l,n}(\mathbf{j}), E_{l,n}(\mathbf{j}')] \right|^{\frac{p}{p-1}} \right] \right)^{\frac{p-1}{p}} \\ & \leq 8M_\epsilon^4 \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{R^2, R^2, K}^{(K, \mathcal{D}_n)}(0, d(\{\mathbf{1}, \mathbf{j}\}, \{\mathbf{1}', \mathbf{j}'\}) - 2R) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}} \\ & \leq 8M_\epsilon^4 \tilde{\alpha}(0, d(\{\mathbf{1}, \mathbf{j}\}, \{\mathbf{1}', \mathbf{j}'\}) - 2R). \end{aligned}$$

Thus,

$$\begin{aligned} & \left| \mathbb{E}_{\mathcal{H}_0} [E_{k,n}(\mathbf{1})E_{k',n}(\mathbf{1}')\mathbb{E}_{\mathcal{H}_{l-1}} [E_{l,n}(\mathbf{j})E_{l,n}(\mathbf{j}')]] \right| \\ & \leq 8M_\epsilon^4 \tilde{\alpha}(I[k \neq k'])(l-1-\max\{k, k'\}, d(\{\mathbf{1}, \mathbf{j}\}, \{\mathbf{1}', \mathbf{j}'\}) - 2R). \end{aligned}$$

4.2.1 Case c)

Consider $k^* = k = l = l' > k_* = k'$. Then $\mathbb{E}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1})E_{k',n}(\mathbf{1}')] = 0$ and hence the covariance (8) is given by

$$\begin{aligned} & \mathbb{E}_{\mathcal{H}_0} [\text{Cov}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j}), E_{k',n}(\mathbf{1}')E_{k,n}(\mathbf{j}')]] \\ & = \mathbb{E}_{\mathcal{H}_0} [E_{k',n}(\mathbf{1}')\mathbb{E}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j})E_{k,n}(\mathbf{j}')]]. \end{aligned}$$

(i): Since

$$\begin{aligned} & \left| \mathbb{E}_{\mathcal{H}_0} [E_{k',n}(\mathbf{1}')\mathbb{E}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j})E_{k,n}(\mathbf{j}')]] \right| \\ & \leq \left(\mathbb{E}_{\mathcal{H}_0} [|E_{k',n}(\mathbf{1}')|^p] \right)^{\frac{1}{p}} \left(\mathbb{E}_{\mathcal{H}_0} \left[\left| \text{Cov}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j}), E_{k,n}(\mathbf{j}')] \right|^{\frac{p}{p-1}} \right] \right)^{\frac{p-1}{p}}, \end{aligned}$$

if conditions M-(A2-1) and M-(A3-1) hold, by inequalities (1), (3) and (5)

$$\begin{aligned}
& \left| \mathbb{E}_{\mathcal{H}_0} \left[E_{k',n}(\mathbf{l}') \text{Cov}_{\mathcal{H}_{k-1}} \left[E_{k,n}(\mathbf{l}) E_{k,n}(\mathbf{j}), E_{k,n}(\mathbf{j}') \right] \right] \right| \\
& \leq 8M_\epsilon^4 \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{\mathcal{H}_{k-1}} \left(\mathcal{H}_{0,k,C(\mathbf{l}) \cup C(\mathbf{j})}, \mathcal{H}_{0,k,C(\mathbf{j}')} \right) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}} \\
& \leq 8M_\epsilon^4 \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{2R^2, R^2, K}^{(K, \mathbb{R}^d)} \left(0, d(C(\mathbf{l}) \cup C(\mathbf{j}), C(\mathbf{j}')) \right) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}} \\
& \leq 8M_\epsilon^4 \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{2R^2, R^2, K}^{(K, \mathbb{R}^d)} \left(0, d(\{\mathbf{l}, \mathbf{j}\}, \{\mathbf{l}', \mathbf{j}'\} - 2R) \right) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}} \\
& \leq 8M_\epsilon^4 \bar{\alpha}_K(d(\{\mathbf{l}, \mathbf{j}\}, \{\mathbf{l}', \mathbf{j}'\} - 2R)).
\end{aligned}$$

(ii): On the other hand,

$$\begin{aligned}
& \left| \mathbb{E}_{\mathcal{H}_0} \left[E_{k',n}(\mathbf{l}') \mathbb{E}_{\mathcal{H}_{k-1}} \left[E_{k,n}(\mathbf{l}) E_{k,n}(\mathbf{j}) E_{k,n}(\mathbf{j}') \right] \right] \right| \\
& = \left| \text{Cov}_{\mathcal{H}_0} \left[E_{k',n}(\mathbf{l}'), \mathbb{E}_{\mathcal{H}_{k-1}} \left[E_{k,n}(\mathbf{l}) E_{k,n}(\mathbf{j}) E_{k,n}(\mathbf{j}') \right] \right] \right|.
\end{aligned}$$

If conditions M-(A2-2) and M-(A3-2) hold,

$$\begin{aligned}
& \left| \text{Cov}_{\mathcal{H}_0} \left[E_{k',n}(\mathbf{l}'), \mathbb{E}_{\mathcal{H}_{k-1}} \left[E_{k,n}(\mathbf{l}) E_{k,n}(\mathbf{j}) E_{k,n}(\mathbf{j}') \right] \right] \right| \\
& \leq 8M_\epsilon^4 \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{\mathcal{H}_0} \left(\mathcal{H}_{0,k', \mathcal{D}_n}, \mathcal{H}_{(k-r)^+, k-1, \mathcal{D}_n} \right) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}} \\
& \leq 8M_\epsilon^4 \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{|\mathcal{D}_n|, |\mathcal{D}_n|, r}^{(\infty, \mathcal{D}_n)} (k-1-k', 0) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}} \\
& \leq 8M_\epsilon^4 \tilde{\alpha}_n(k-1-k').
\end{aligned}$$

(iii): Finally, if condition M-(A2-3) holds

$$\begin{aligned}
& \left| \mathbb{E}_{\mathcal{H}_0} \left[E_{k',n}(\mathbf{l}') \mathbb{E}_{\mathcal{H}_{k-1}} \left[E_{k,n}(\mathbf{l}) E_{k,n}(\mathbf{j}) E_{k,n}(\mathbf{j}') \right] \right] \right| \\
& \leq 8 \left(\mathbb{E}_{\mathcal{H}_0} \left[|E_{k',n}(\mathbf{l}')|^p \right] \mathbb{E}_{\mathcal{H}_0} \left[\left| \text{Cov}_{\mathcal{H}_{k-1}} \left[E_{k,n}(\mathbf{l}) E_{k,n}(\mathbf{j}), E_{k,n}(\mathbf{j}') \right] \right|^p \right] \right)^{\frac{1}{p}} \\
& \quad \left(\alpha_{\mathcal{H}_0} \left(\mathcal{H}_{0,k', C(\mathbf{l}')}, \mathcal{H}_{(k-r)^+, k-1, C(\mathbf{l}) \cup C(\mathbf{j}) \cup C(\mathbf{j}')} \right) \right)^{\frac{p-2}{p}} \\
& \leq 8M_\epsilon \left(\mathbb{E}_{\mathcal{H}_0} \left[\left| \text{Cov}_{\mathcal{H}_{k-1}} \left[E_{k,n}(\mathbf{l}) E_{k,n}(\mathbf{j}), E_{k,n}(\mathbf{j}') \right] \right|^p \right] \right)^{\frac{1}{p}} \\
& \quad \mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{|\mathcal{D}_n|, |\mathcal{D}_n|, r}^{(\infty, \mathcal{D}_n)} (k-1-k', 0) \right)^{\frac{p-2}{p}} \right]
\end{aligned}$$

and

$$\begin{aligned}
& \mathbb{E}_{\mathcal{H}_0} \left[\left| \text{Cov}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j}), E_{k,n}(\mathbf{j}')] \right|^p \right] \\
& \leq 8M_\epsilon^4 \left[\left(\mathbb{E}_{\mathcal{H}_{k-1}} [|E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j})|^p] \mathbb{E}_{\mathcal{H}_{k-1}} [|E_{k,n}(\mathbf{j}')|^p] \right. \right. \\
& \quad \left. \left. (\alpha_{\mathcal{H}_{k-1}}(\mathcal{H}_{0,k,C(\mathbf{1}) \cup C(\mathbf{j})}, \mathcal{H}_{0,k,C(\mathbf{j}')}))^{p-2} \right) \right] \\
& \leq 8M_\epsilon^{3p} \mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{2R^2, R^2, K}^{(K, \mathbb{R}^d)}(0, d(C(\mathbf{1}) \cup C(\mathbf{j}), C(\mathbf{j}'))) \right)^{p-2} \right] \\
& \leq 8M_\epsilon^{3p} \mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{2R^2, R^2, K}^{(K, \mathbb{R}^d)}(0, d(\{\mathbf{1}, \mathbf{j}\}, \{\mathbf{1}', \mathbf{j}'\}) - 2R) \right)^{p-2} \right],
\end{aligned}$$

which combined with condition M-(A3-3) implies that

$$\begin{aligned}
& \left| \mathbb{E}_{\mathcal{H}_0} [E_{k',n}(\mathbf{1}') \mathbb{E}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j})E_{k,n}(\mathbf{j}')]] \right| \\
& \leq 64M_\epsilon^4 \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{2R^2, R^2, K}^{(K, \mathbb{R}^d)}(0, d(\{\mathbf{1}, \mathbf{j}\}, \{\mathbf{1}', \mathbf{j}'\}) - 2R) \right)^{p-2} \right] \right)^{\frac{1}{p}} \\
& \quad \mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{|\mathcal{D}_n|, |\mathcal{D}_n|, r}^{(\infty, \mathcal{D}_n)}(k-1-k', 0) \right)^{\frac{p-2}{p}} \right] \\
& \leq 64M_\epsilon^4 \check{\alpha}(k-1-k', d(\{\mathbf{1}, \mathbf{j}\}, \{\mathbf{1}', \mathbf{j}'\}) - 2R).
\end{aligned}$$

4.2.2 Case d)

When $k^* = k_* = k = l = k' = l'$, the covariance (8) becomes

$$\begin{aligned}
& \mathbb{E}_{\mathcal{H}_0} [\text{Cov}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j}), E_{k,n}(\mathbf{1}')E_{k,n}(\mathbf{j}')]] \\
& + \text{Cov}_{\mathcal{H}_0} [\mathbb{E}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j})], \mathbb{E}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1}')E_{k,n}(\mathbf{j}')]].
\end{aligned}$$

(i) and (iii): By condition M-(A2-1), $E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j})$ and $E_{k,n}(\mathbf{1}')E_{k,n}(\mathbf{j}')$ are respectively measurable with respect to $\mathcal{H}_{0,k,C(\mathbf{1}) \cup C(\mathbf{j})}$ and $\mathcal{H}_{0,k,C(\mathbf{1}') \cup C(\mathbf{j}')}$. Thus, if condition M-(A2-1) or M-(A2-3) holds, from inequalities (2), (7), (4), and (6) it follows that

$$\begin{aligned}
& \left| \mathbb{E}_{\mathcal{H}_0} [\text{Cov}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j}), E_{k,n}(\mathbf{1}')E_{k,n}(\mathbf{j}')]] \right| \\
& \leq 8M_\epsilon^4 \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{\mathcal{H}_{k-1}}(\mathcal{H}_{0,k,C(\mathbf{1}) \cup C(\mathbf{j})}, \mathcal{H}_{0,k,C(\mathbf{1}') \cup C(\mathbf{j}')}) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}}
\end{aligned}$$

and

$$\begin{aligned}
& \left| \text{Cov}_{\mathcal{H}_0} [\mathbb{E}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j})], \mathbb{E}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1}')E_{k,n}(\mathbf{j}')]] \right| \\
& \leq 8M_\epsilon^4 \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{\mathcal{H}_0}(\mathcal{H}_{0,k-1,C(\mathbf{1}) \cup C(\mathbf{j})}, \mathcal{H}_{0,k-1,C(\mathbf{1}') \cup C(\mathbf{j}')}) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}}.
\end{aligned}$$

Thus,

$$\begin{aligned}
& |\text{Cov}_{\mathcal{H}_0} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j}), E_{k,n}(\mathbf{l}')E_{k,n}(\mathbf{j}')] | \\
& \leq 16M_\epsilon^4 \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{2R^2, 2R^2, K}^{(K, \mathbb{R}^d)}(0, d(C(\mathbf{1}) \cup C(\mathbf{j}), C(\mathbf{l}') \cup C(\mathbf{j}')) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}} \\
& \leq 16M_\epsilon^4 \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{2R^2, 2R^2, K}^{(K, \mathbb{R}^d)}(0, d(\{\mathbf{1}, \mathbf{j}\}, \{\mathbf{l}', \mathbf{j}'\} - 2R) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}},
\end{aligned}$$

and if condition M-(A3-1) or M-(A3-1) also holds, then

$$\begin{aligned}
& |\text{Cov}_{\mathcal{H}_0} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j}), E_{k,n}(\mathbf{l}')E_{k,n}(\mathbf{j}')] | \\
& \leq 16M_\epsilon^4 \bar{\alpha}(d(\{\mathbf{1}, \mathbf{j}\}, \{\mathbf{l}', \mathbf{j}'\}) - 2R)
\end{aligned}$$

and

$$\begin{aligned}
& |\text{Cov}_{\mathcal{H}_0} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j}), E_{k,n}(\mathbf{l}')E_{k,n}(\mathbf{j}')] | \\
& \leq 16M_\epsilon^4 \check{\alpha}(0, d(\{\mathbf{1}, \mathbf{j}\}, \{\mathbf{l}', \mathbf{j}'\}) - 2R),
\end{aligned}$$

respectively.

(ii): In case (ii) we simply appeal to Hölder's inequality and bound $|\text{Cov}_{\mathcal{H}_0} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j}), E_{k,n}(\mathbf{l}')E_{k,n}(\mathbf{j}')] |$ by $2M_\epsilon^4$.

4.3 Convergence of variance

In the case (ii), $m_{K,n} = \infty$ and hence $\mathcal{J}_{K,n} = \mathcal{D}_n \times \mathcal{D}_n$ and $\sigma_{K,n}^2 = V_{K,n}^2$. But in cases (i) and (iii),

$$\sigma_{K,n}^2 = V_{K,n}^2 + \sum_{k=1}^K \sum_{\substack{\mathbf{1}, \mathbf{j} \in \mathcal{D}_n \\ d(\mathbf{1}, \mathbf{j}) > m_{K,n}}} \mathbb{E}_{\mathcal{H}_0} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j})]$$

and for any $k, n \in \mathbb{N}$, $\mathbf{1}, \mathbf{l}', \mathbf{j}, \mathbf{j}' \in \mathbb{Z}^d$ and $2 < p \leq 2 + \epsilon/3$, from inequalities (1) and (5), we obtain

$$\begin{aligned}
& |\mathbb{E}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1})E_{k,n}(\mathbf{j})] | = |\text{Cov}_{\mathcal{H}_{k-1}} [E_{k,n}(\mathbf{1}), E_{k,n}(\mathbf{j})] | \\
& \leq 8 \left(\mathbb{E}_{\mathcal{H}_{k-1}} [|E_{k,n}(\mathbf{1})|^p] \mathbb{E}_{\mathcal{H}_{k-1}} [|E_{k,n}(\mathbf{j})|^p] \right)^{\frac{1}{p}} \\
& \quad \left(\alpha_{\mathcal{H}_{k-1}}(\mathcal{H}_{0,k,C(\mathbf{1})}, \mathcal{H}_{0,k,C(\mathbf{j})}) \right)^{\frac{p-2}{p}} \\
& \leq 8M_\epsilon^2 \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{\mathcal{H}_{k-1}}(\mathcal{H}_{0,k,C(\mathbf{1})}, \mathcal{H}_{0,k,C(\mathbf{j})}) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}}.
\end{aligned}$$

Thus, in case (i), using M-(A3-1),

$$\begin{aligned}
|\sigma_{K,n}^2 - V_{K,n}^2| &\leq \sum_{k=1}^K \sum_{\substack{\mathbf{1}, \mathbf{j} \in \mathcal{D}_n \\ d(\mathbf{1}, \mathbf{j}) > m_{K,n}}} \mathbb{E}_{\mathcal{H}_0} |\text{Cov}_{\mathcal{H}_{k-1}}[E_{k,n}(\mathbf{1}), E_{k,n}(\mathbf{j})]| \\
&\leq 8M_\epsilon^2 \sum_{k=1}^K \sum_{\substack{\mathbf{1}, \mathbf{j} \in \mathcal{D}_n \\ d(\mathbf{1}, \mathbf{j}) > m_{K,n}}} \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{\mathcal{H}_{k-1}}(\mathcal{H}_{0,k,C(\mathbf{1})}, \mathcal{H}_{0,k,C(\mathbf{j})}) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}} \\
&\leq 8KM_\epsilon^2 \sum_{\mathbf{l} \in \mathcal{D}_n} \sum_{\substack{\mathbf{j} \in \mathbb{Z}^d \\ d(\mathbf{1}, \mathbf{j}) > m_{K,n}}} \left(\mathbb{E}_{\mathcal{H}_0} \left[\left(\alpha_{R^2, R^2, K}^{(K, \mathbb{R}^d)}(0, d(\mathbf{1}, \mathbf{j}) - 2R) \right)^{\frac{p-2}{p-1}} \right] \right)^{\frac{p-1}{p}} \\
&\leq 8KM_\epsilon^2 \sum_{\mathbf{l} \in \mathcal{D}_n} \sum_{l=\lfloor m_{K,n} \rfloor}^{\infty} \sum_{\substack{\mathbf{j} \in \mathbb{Z}^d \\ d(\mathbf{1}, \mathbf{j})=l}} \bar{\alpha}_K(l - 2R) \\
&\leq 8(3^d)K|\mathcal{D}_n|M_\epsilon^2 \sum_{l=\lfloor m_{K,n} \rfloor}^{\infty} l^{d-1} \bar{\alpha}_K(l - 2R)
\end{aligned}$$

because the cardinality of $\{\mathbf{l}' \in \mathbb{Z}^d : d(\mathbf{1}, \mathbf{l}') = l\}$ is less than or equal to $3^d l^{d-1}$ for $l \in \mathbb{N}$ (see [Biscio and Waagepetersen \(2019\)](#), Lemma F4). Thus

$$\left| 1 - \frac{V_{K,n}^2}{\sigma_{K,n}^2} \right| \leq 8(3^d)K \frac{|\mathcal{D}_n|}{\sigma_{K,n}^2} M_\epsilon^2 \sum_{l=\lfloor m_{K,n} \rfloor}^{\infty} l^{d-1} \bar{\alpha}_K(l - 2R)$$

As $n \rightarrow \infty$, assumption M-(A5-1) implies that $|\mathcal{D}_n|/\sigma_{K,n}^2$ is bounded. Condition M-(A3-1) implies that

$$\sum_{l=\lfloor m_{K,n} \rfloor}^{\infty} l^{d-1} \bar{\alpha}_K(l - 2R) \leq \text{constant} \sum_{l=\lfloor m_{K,n} \rfloor}^{\infty} l^{\eta+d-1}$$

and since $\eta+d < 0$, the tail series $\sum_{l=\lfloor m_{K,n} \rfloor}^{\infty} l^{\eta+d-1}$ converges to zero as $n \rightarrow \infty$. Therefore

$$\left| 1 - \frac{V_{K,n}^2}{\sigma_{K,n}^2} \right| \xrightarrow[n \rightarrow \infty]{a.s.} 0. \quad (9)$$

In case (iii) we obtain the same conclusion using assumptions M-(A3-3), M-(A4-3), and M-(A5-3).

5 Spatio-temporal autoregressive model

In this section we provide details regarding Section M-3.2.

5.1 Representation as Vector Autoregression

By rewriting M-(3.2) as a Vector Autoregression (VAR) equation of order one (see [Hamilton \(1994\)](#), p. 259) and recursively applying the resulting VAR(1)

representation, it follows that for any $0 \leq l < k$,

$$X_{k,n} = C_{k-l,n}X_{l,n} + \sum_{j=0}^{k-l-1} C_{j,n}Q_n^{-1/2}Z_{k-j,n} \quad (10)$$

where $C_{j,n} = H_n^\top F_n^j H_n$ and (see [Yamamoto \(1981\)](#), Lemma 1)

$$F_n = \begin{bmatrix} \beta B_{1,n} & \beta B_{2,n} & \cdots & \beta B_{r-1,n} & \beta B_{r,n} \\ I_{|\mathcal{D}_n|} & 0 & \cdots & 0 & 0 \\ 0 & I_{|\mathcal{D}_n|} & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & I_{|\mathcal{D}_n|} & 0 \end{bmatrix}_{r|\mathcal{D}_n| \times r|\mathcal{D}_n|}, \quad H_n = \begin{bmatrix} I_{|\mathcal{D}_n|} \\ 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}_{r|\mathcal{D}_n| \times |\mathcal{D}_n|}.$$

Thus, for any $0 \leq l < k$, the conditional distribution of $X_{k,n}$ given \mathcal{H}_l is multivariate normal with $\mathbb{E}_{\mathcal{H}_l}[X_{k,n}] = C_{k-l,n}X_{l,n}$ and

$$\text{Var}_{\mathcal{H}_l}[X_{k,n}] = \sum_{j=0}^{k-l-1} C_{j,n}Q_n^{-1}C_{j,n}^\top.$$

Moreover, for any $1 \leq l \leq k$ and $m \in \mathbb{Z}_+$,

$$\begin{aligned} \text{Cov}_{\mathcal{H}_{l-1}}[X_{k,n}, X_{k+m,n}] &= \mathbb{E}_{\mathcal{H}_{l-1}}[\text{Cov}_{\mathcal{H}_k}[X_{k,n}, X_{k+m,n}]] \\ &\quad + \text{Cov}_{\mathcal{H}_{l-1}}[\mathbb{E}_{\mathcal{H}_k}[X_{k,n}], \mathbb{E}_{\mathcal{H}_k}[X_{k+m,n}]] \\ &= \text{Cov}_{\mathcal{H}_{l-1}}[X_{k,n}, C_{m,n}X_{k,n}] \\ &= \text{Cov}_{\mathcal{H}_{l-1}}[X_{k,n}, X_{k,n}]C_{m,n}^\top = \left(\sum_{j=1}^{k-l} C_{j,n}Q_n^{-1}C_{j,n}^\top \right) C_{m,n}^\top. \end{aligned}$$

If the eigenvalues of F_n have modulus less than one, or equivalently the roots of the equation

$$\det \left(I_{|\mathcal{D}_n|} - \sum_{j=1}^r z^j \beta B_{j,n} \right) = 0$$

are outside the complex unit circle, then $F_n^m \rightarrow 0$ as $m \rightarrow \infty$ (see [Hamilton \(1994\)](#), p. 259). We assume that as $m \rightarrow \infty$, $\|F_n^m\| = O(m^\psi)$ for some $\psi < -1$, where for a matrix $A = [a_{i,j}]_{n_1 \times n_2}$,

$$\|A\| = \left(\sum_{i=1}^{n_1} \sum_{j=1}^{n_2} a_{i,j}^2 \right)^{\frac{1}{2}}$$

is the Frobenius norm. Then, $\|C_{m,n}\| = O(m^\psi)$ because

$$\|C_{m,n}\| = \|H_n^\top F_n^m H_n\| \leq \|H_n^\top\| \|F_n^m\| \|H_n\| \leq |\mathcal{D}_n|^2 \|F_n^m\|.$$

Consequently for any $1 \leq l \leq k$ and $\mathbf{l}, \mathbf{j} \in \mathcal{D}_n$,

$$|\text{Cov}_{\mathcal{H}_{l-1}}[X_{k,n}(\mathbf{l}), X_{k+m,n}(\mathbf{j})]| = O(m^\psi), \quad (11)$$

as $m \rightarrow \infty$, since

$$\|\text{Cov}_{\mathcal{H}_{l-1}}[X_{k,n}, X_{k+m,n}]\| \leq \left\| \sum_{j=1}^{k-l} C_{j,n} Q_n^{-1} C_{j,n}^\top \right\| \|C_{m,n}\|$$

and

$$\left\| \sum_{j=1}^{k-l} C_{j,n} Q_n^{-1} C_{j,n}^\top \right\| \leq \sum_{j=1}^{k-l} \|C_{j,n}\|^2 \|Q_n^{-1}\| \leq \text{constant} \sum_{j=1}^{\infty} j^{2\psi} < \infty.$$

For $i, l, k \in \mathbb{Z}_+$, $0 \leq i < l \leq k$, and $A, B \subseteq \mathcal{D}_n$, let

$$\rho_{\mathcal{H}_i}(\mathcal{H}_{0,l,A}, \mathcal{H}_{(k-r)^+,k,B}) = \sup |\text{Corr}_{\mathcal{H}_i}[Z, Y]|$$

denote the maximal conditional correlation coefficient between $\mathcal{H}_{0,l,A}$ and $\mathcal{H}_{(k-r)^+,k,B}$ given \mathcal{H}_i , where the supremum is over all random variables Z and Y that are respectively $\mathcal{H}_{0,l,A}$ and $\mathcal{H}_{(k-r)^+,k,B}$ measurable, and where $\mathbb{E}_{\mathcal{H}_i}[Z^2] < \infty$ and $\mathbb{E}_{\mathcal{H}_i}[Y^2] < \infty$. It follows from (11) that as $m \rightarrow \infty$,

$$\sup_{\substack{l, k' \in \mathbb{Z}_+ \\ (k-r)^+ \leq k' \leq k \\ k-l \geq m}} \sup_{\substack{A, B \subseteq \mathcal{D}_n \\ l \in A, \mathbf{j} \in B}} \sup_{i \in \{0, \dots, l-1\}} |\text{Corr}_{\mathcal{H}_i}[X_{l,n}(\mathbf{1}), X_{k',n}(\mathbf{j})]| \leq \text{constant } m^\psi,$$

which means that

$$\sup_{\substack{l, k \in \mathbb{Z}_+ \\ k-l \geq m}} \sup_{A, B \subseteq \mathcal{D}_n} \sup_{i \in \{0, \dots, l-1\}} \rho_{\mathcal{H}_i}(\mathcal{H}_{0,l,A}, \mathcal{H}_{(k-r)^+,k,B}) \leq \text{constant } m^\psi.$$

5.2 A pseudo-likelihood score function

Given \mathcal{H}_0 , the joint distribution of $X_{1,n}, \dots, X_{K,n}$ is given by

$$[X_{1,n}, \dots, X_{K,n}]_{\mathcal{H}_0} = \prod_{k=1}^K [X_{k,n}]_{\mathcal{H}_{k-1}},$$

where $[\cdot]_{\mathcal{H}_{k-1}}$ denotes conditional distribution given \mathcal{H}_{k-1} . However, given \mathcal{H}_0 , a pseudo-likelihood function for $[X_{k,n}]_{\mathcal{H}_{k-1}}$ is given by

$$\prod_{\mathbf{1} \in \mathcal{D}_n} [X_{k,n}(\mathbf{1})]_{\sigma(\mathcal{H}_{k-1} \cup \mathcal{H}_{k,k,\mathcal{D}_n \setminus \{\mathbf{1}\}})}.$$

Thus, a log pseudo-likelihood for the observations $X_{1,n}, \dots, X_{K,n}$ is given by

$$\ell(\beta, \gamma) = -\frac{1}{2} \sum_{k=1}^K \sum_{\mathbf{1} \in \mathcal{D}_n} a_n(\mathbf{1}) \left(X_{k,n}(\mathbf{1}) - \beta \xi_{k-1,n}^{\text{temp}}(\mathbf{1}) - \gamma \xi_{k,n}^{\text{spat}}(\mathbf{1}) \right)^2$$

It follows from M-(3.2) that given \mathcal{H}_{k-1} ,

$$X_{k,n} - \beta \xi_{k-1,n}^{\text{temp}} = Q_n^{-1/2} Z_{k,n}$$

and

$$\xi_{k,n}^{\text{spat}} = B_{0,n} \left(X_{k,n} - \beta \xi_{k-1,n}^{\text{temp}} \right) = B_{0,n} Q_n^{-1/2} Z_{k,n},$$

which means that

$$\begin{aligned} A_n \left(X_{k,n} - \beta \xi_{k-1,n}^{\text{temp}} - \gamma \xi_{k,n}^{\text{spat}} \right) &= A_n \left(I_{|\mathcal{D}_n|} - \gamma B_{0,n} \right) Q_n^{-1/2} Z_{k,n} \\ &= Q_n Q_n^{-1/2} Z_{k,n} = Q_n^{1/2} Z_{k,n} \end{aligned}$$

and hence the log pseudo-likelihood can be written as

$$\ell(\beta, \gamma) = -\frac{1}{2} \sum_{k=1}^K Z_{k,n}^\top Q_n^{1/2} A_n^{-1} Q_n^{1/2} Z_{k,n}.$$

The corresponding score function of $\ell(\beta, \gamma)$ is given by

$$T_{K,n} = \sum_{k=1}^K \sum_{\mathbf{l} \in \mathcal{D}_n} E_{k,n}(\mathbf{l}),$$

where

$$E_{k,n}(\mathbf{l}) = \varepsilon_{k,n}(\mathbf{l}) \left(\xi_{k-1,n}^{\text{temp}}(\mathbf{l}), \xi_{k,n}^{\text{spat}}(\mathbf{l}) \right)^\top$$

and $\varepsilon_{k,n} = Q_n^{1/2} Z_{k,n}$.

5.3 Covariance matrix of the score function

Turning to $\mathbb{E}_{\mathcal{H}_0} \text{Var}_{H_{k-1}} \sum_{\mathbf{l} \in \mathcal{D}_n} E_{k,n}(\mathbf{l})$, we have

$$\sum_{\mathbf{l} \in \mathcal{D}_n} E_{k,n}(\mathbf{l}) = \left(Z_{k,n}^\top Q_n^{1/2} \xi_{k-1,n}^{\text{temp}}, Z_{k,n}^\top K_n Z_{k,n} \right)^\top,$$

where $K_n = [K_n(\mathbf{i}, \mathbf{j})]_{\mathbf{i}, \mathbf{j}} = Q_n^{1/2} B_{0,n} Q_n^{-1/2}$. Obviously, $\mathbb{E}_{\mathcal{H}_{k-1}} \left[Z_{k,n}^\top Q_n^{1/2} \xi_{k-1,n}^{\text{temp}} \right] = 0$ and

$$\text{Var}_{\mathcal{H}_{k-1}} \left[Z_{k,n}^\top Q_n^{1/2} \xi_{k-1,n}^{\text{temp}} \right] = \left(\xi_{k-1,n}^{\text{temp}} \right)^\top Q_n \xi_{k-1,n}^{\text{temp}}.$$

Since $Z_{k,n}^\top K_n Z_{k,n}$ is a quadratic form in the standard Gaussian random vector $Z_{k,n}$, we have (see [Rencher and Schaalje \(2008\)](#), Theorems 5.2a, 5.2c and 5.2d)

$$\begin{aligned} \mathbb{E}_{\mathcal{H}_{k-1}} \left[Z_{k,n}^\top K_n Z_{k,n} \right] &= \text{trace}(K_n) = \text{trace}(Q_n^{1/2} B_{0,n} Q_n^{-1/2}) \\ &= \text{trace}(B_{0,n}) = \sum_{\mathbf{l} \in \mathcal{D}_n} b_0(\mathbf{l}, \mathbf{l}) = 0, \end{aligned}$$

$$\text{Var}_{\mathcal{H}_{k-1}} Z_{k,n}^\top K_n Z_{k,n} = 2 \text{trace}(K_n^2) = 2 \text{trace}(Q_n^{1/2} B_{0,n} B_{0,n} Q_n^{-1/2}) = 2 \text{trace}(B_{0,n}^2)$$

and

$$\text{Cov}_{\mathcal{H}_{k-1}} \left[Z_{k,n}^\top Q_n^{1/2} \xi_{k-1,n}^{\text{temp}}, Z_{k,n}^\top K_n Z_{k,n} \right] = 2 \left(\xi_{k-1,n}^{\text{temp}} \right)^\top Q_n^{1/2} K_n \mathbb{E}_{\mathcal{H}_{k-1}} [Z_{k,n}] = 0.$$

Thus $T_{K,n}$ provides an unbiased estimating function and

$$\Sigma_{K,n} = \text{Var}_{\mathcal{H}_0} [T_{K,n}] = \sum_{k=1}^K \mathbb{E}_{\mathcal{H}_0} \text{Var}_{H_{k-1}} \sum_{\mathbf{l} \in \mathcal{D}_n} E_{k,n}(\mathbf{l}) = \text{diag}(\lambda_{K,n}^{(1)}, \lambda_{K,n}^{(2)}),$$

where

$$\lambda_{K,n}^{(1)} = \sum_{k=1}^K \mathbb{E}_{\mathcal{H}_0} \left[(\xi_{k-1,n}^{\text{temp}})^\top Q_n \xi_{k-1,n}^{\text{temp}} \right]$$

and $\lambda_{K,n}^{(2)} = 2K \text{trace}(B_{0,n}^2)$.

On the other hand, given \mathcal{H}_0 , from (10),

$$\beta \xi_{k-1,n}^{\text{temp}} = X_{k,n} - Q_n^{-1/2} Z_{k,n} = C_{k,n} X_{0,n} + \sum_{j=1}^{k-1} C_{j,n} Q_n^{-1/2} Z_{k-j,n}$$

and hence $\mathbb{E}_{\mathcal{H}_0} [\beta \xi_{k-1,n}^{\text{temp}}] = C_{k,n} X_{0,n}$ and $\text{Var}_{\mathcal{H}_0} [\beta \xi_{k-1,n}^{\text{temp}}] = \sum_{j=1}^{k-1} C_{j,n} Q_n^{-1} C_{j,n}^\top$.

Thus

$$\begin{aligned} & \beta^2 \mathbb{E}_{\mathcal{H}_0} \left[(\xi_{k-1,n}^{\text{temp}})^\top Q_n \xi_{k-1,n}^{\text{temp}} \right] \\ &= \text{trace} \left(Q_n \text{Var}_{\mathcal{H}_0} [\beta \xi_{k-1,n}^{\text{temp}}] \right) + \left(\mathbb{E}_{\mathcal{H}_0} [\beta \xi_{k-1,n}^{\text{temp}}] \right)^\top Q_n \mathbb{E}_{\mathcal{H}_0} [\beta \xi_{k-1,n}^{\text{temp}}] \\ &= \sum_{j=1}^{k-1} \text{trace} (Q_n C_{j,n} Q_n^{-1} C_{j,n}^\top) + X_{0,n}^\top C_{k,n}^\top Q_n C_{k,n} X_{0,n} \end{aligned}$$

and because the precision matrix Q_n is positive definite,

$$\begin{aligned} \beta^2 \mathbb{E}_{\mathcal{H}_0} \left[(\xi_{k-1,n}^{\text{temp}})^\top Q_n \xi_{k-1,n}^{\text{temp}} \right] &\geq \sum_{j=1}^{k-1} \text{trace} (Q_n C_{j,n} Q_n^{-1} C_{j,n}^\top) \\ &\geq \sum_{j=1}^{k-1} \text{trace} (C_{j,n} C_{j,n}^\top) \geq \text{trace} (C_{1,n} C_{1,n}^\top). \end{aligned}$$

But $C_{1,n} = H_n^\top F_n H_n = \beta B_{1,n}$, which means that $\text{trace}(C_{1,n} C_{1,n}^\top) = \beta^2 \text{trace}(B_{1,n}^2)$ and $\lambda_{K,n}^{(1)} \geq K \text{trace}(B_{1,n}^2)$.

5.4 Moment bounds

For any $k \in \mathbb{N}$ and $\nu > 1$,

$$\begin{aligned} \sup_{\mathbf{1} \in \mathcal{D}_n} \mathbb{E}_{\mathcal{H}_{k-1}} \left[\left| \varepsilon_{k,n}(\mathbf{1}) \xi_{k-1,n}^{\text{temp}}(\mathbf{1}) \right|^\nu \right] &\leq \left\| \xi_{k-1,n}^{\text{temp}} \right\|^\nu \mathbb{E}_{\mathcal{H}_{k-1}} \left[\left\| \varepsilon_{k,n} \right\|^\nu \right], \\ \sup_{\mathbf{1} \in \mathcal{D}_n} \mathbb{E}_{\mathcal{H}_{k-1}} \left[\left| \varepsilon_{k,n}(\mathbf{1}) \xi_{k,n}^{\text{spat}}(\mathbf{1}) \right|^\nu \right] &\leq \left(\mathbb{E}_{\mathcal{H}_{k-1}} \left[\left\| \varepsilon_{k,n} \right\|^{2\nu} \right] \mathbb{E}_{\mathcal{H}_{k-1}} \left[\left\| \xi_{k,n}^{\text{spat}} \right\|^{2\nu} \right] \right)^{\frac{1}{2}} \end{aligned}$$

and

$$\begin{aligned} \mathbb{E}_{\mathcal{H}_{k-1}} \left[\left\| \varepsilon_{k,n} \right\|^\nu \right] &= \mathbb{E}_{\mathcal{H}_{k-1}} \left[\left\| Q_n^{1/2} Z_{k,n} \right\|^\nu \right] = \mathbb{E}_{\mathcal{H}_{k-1}} \left[\left(Z_{k,n}^\top Q_n Z_{k,n} \right)^{\frac{\nu}{2}} \right], \\ \mathbb{E}_{\mathcal{H}_{k-1}} \left[\left\| \varepsilon_{k,n} \right\|^{2\nu} \right] &= \mathbb{E}_{\mathcal{H}_{k-1}} \left[\left\| Q_n^{1/2} Z_{k,n} \right\|^{2\nu} \right] = \mathbb{E}_{\mathcal{H}_{k-1}} \left[\left(Z_{k,n}^\top Q_n Z_{k,n} \right)^\nu \right], \\ \mathbb{E}_{\mathcal{H}_{k-1}} \left[\left\| \xi_{k,n}^{\text{spat}} \right\|^{2\nu} \right] &= \mathbb{E}_{\mathcal{H}_{k-1}} \left[\left(Z_{k,n}^\top Q_n^{-1/2} B_{0,n}^\top B_{0,n} Q_n^{-1/2} Z_{k,n} \right)^\nu \right] \end{aligned}$$

are moments of quadratic forms in a standard normal random vector that are uniformly bounded in k .

References

- Biscio, C. A. N. and Waagepetersen, R. (2019). A general central limit theorem and a subsampling variance estimator for α -mixing point processes. *Scandinavian Journal of Statistics*, 46(4):1168–1190.
- Hamilton, J. D. (1994). *Time series analysis*. Princeton University Press.
- Jalilian, A., Xu, G., and Waagepetersen, R. (2023). A central limit theorem for a sequence of conditionally centered and α -mixing random fields.
- Rencher, A. C. and Schaalje, G. B. (2008). *Linear models in statistics*. John Wiley & Sons.
- Yamamoto, T. (1981). Predictions of multivariate autoregressive-moving average models. *Biometrika*, 68(2):485–492.
- Yuan, D. and Lei, L. (2013). Some conditional results for conditionally strong mixing sequences of random variables. *Science China Mathematics*, 56(4):845–859.