

BOUNDEDNESS OF THE DYADIC MAXIMAL FUNCTION ON GRADED LIE GROUPS

DUVÁN CARDONA, JULIO DELGADO, AND MICHAEL RUZHANSKY

ABSTRACT. Let $1 < p \leq \infty$ and let $n \geq 2$. It was proved independently by C. Calderón, R. Coifman and G. Weiss that the dyadic maximal function

$$\mathcal{M}_D^{d\sigma} f(x) = \sup_{j \in \mathbb{Z}} \left| \int_{\mathbb{S}^{n-1}} f(x - 2^j y) d\sigma(y) \right|$$

is a bounded operator on $L^p(\mathbb{R}^n)$ where $d\sigma(y)$ is the surface measure on \mathbb{S}^{n-1} . In this paper we prove an analogue of this result on arbitrary graded Lie groups. More precisely, to any finite Borel measure $d\sigma$ with compact support on a graded Lie group G , we associate the corresponding dyadic maximal function $\mathcal{M}_D^{d\sigma}$ using the homogeneous structure of the group. Then, we prove a criterion in terms of the order (at zero and at infinity) of the group Fourier transform $\widehat{d\sigma}$ of $d\sigma$ with respect to a fixed Rockland operator \mathcal{R} on G that assures the boundedness of $\mathcal{M}_D^{d\sigma}$ on $L^p(G)$ for all $1 < p \leq \infty$.

CONTENTS

1. Introduction	2
1.1. Outline	2
1.2. The main result	3
2. Fourier analysis on graded groups	4
2.1. Homogeneous and graded Lie groups	4
2.2. Fourier analysis on nilpotent Lie groups	5
2.3. Homogeneous linear operators and Rockland operators	7
3. Boundedness of the dyadic maximal function	9
3.1. The key lemma	9
3.2. The probabilistic argument	23
3.3. Proof of the main theorem	24
References	24

2010 *Mathematics Subject Classification.* 35S30, 42B20; Secondary 42B37, 42B35.

Key words and phrases. Dyadic maximal function, nilpotent Lie groups, graded Lie groups, Calderón theorem, Coifman-Weiss theory.

The authors are supported by the FWO Odysseus 1 grant G.0H94.18N: Analysis and Partial Differential Equations and by the Methusalem programme of the Ghent University Special Research Fund (BOF) (Grant number 01M01021). J. Delgado is also supported by Vice. Inv. Universidad del Valle Grant CI 71329, MathAmSud and Minciencias-Colombia under the project MATHAMSUD21-MATH-03. Michael Ruzhansky is also supported by EPSRC grant EP/R003025/2.

1. INTRODUCTION

1.1. Outline. For more than fifty years (dating back to the work of Folland and Stein [20] in the 1970s), there has been an extensive program to generalise the techniques from the real-variable Euclidean harmonic analysis to the more general setting of nilpotent Lie groups. This is particularly motivated by applications to degenerate partial differential operators (see e.g. Rothschild and Stein [33]), where the usual methods on the Euclidean space are not completely suitable. Among the fundamental operators of the Euclidean harmonic analysis, are the full maximal function and its dyadic counterpart. Contributing to the aforementioned program, the aim of this work is to study the L^p -boundedness of the dyadic maximal function on a nilpotent Lie group G . Since our criteria involve that the group admits left-invariant hypoelliptic partial differential operators we assume that G is a graded Lie group in view of the Helffer and Nourrigat solution of the Rockland conjecture, see [24].

To the best of our knowledge, the study of the L^p -boundedness of the spherical averages started in a satisfactory way with Stein (inspired by the 1976's work of Nagel, Rivière and Wainger [30]). Indeed, consider the full-maximal function on \mathbb{R}^n

$$\mathcal{M}_F^{d\sigma} f(x) = \sup_{r>0} \left| \int_{\mathbb{S}^{n-1}} f(x - ry) d\sigma(y) \right|, \quad (1.1)$$

where σ is the surface measure on the sphere \mathbb{S}^{n-1} . A remarkable result due to Stein in [42], proved the boundedness of $\mathcal{M}_F^{d\sigma}$ from $L^p(\mathbb{R}^n)$ to itself, if and only if $p > \frac{n}{n-1}$, for all $n \geq 3$. Then, the lower dimensional case $n = 2$ was proved by Bourgain in [3]. Additionally, C. Calderón in [7], proved that the dyadic maximal function

$$\mathcal{M}_D^{d\sigma} f(x) = \sup_{j \in \mathbb{Z}} \left| \int_{\mathbb{S}^{n-1}} f(x - 2^j y) d\sigma(y) \right|, \quad (1.2)$$

can be extended to a bounded operator on $L^p(\mathbb{R}^n)$, for all $1 < p \leq \infty$, for any $n \geq 2$. It was observed by S. Wainger that the L^p -boundedness of the dyadic maximal function was also proved independently by Coifman and Weiss in [10]. Other proofs for the L^p -boundedness of (1.1) when $n \geq 3$ can be found e.g. in Carbery [8], Cowling and Maceuri [11], Rubio de Francia [34], Duoandikoetxea and Rubio de Francia [15], and for Bourgain's result for $n = 2$ a new proof was given by Mockenhaupt, Seeger, and Sogge [28].

The study of the full maximal function and its dyadic version has been mainly concentrated in the context of the Heisenberg group \mathbb{H}_n and on two-steps nilpotent Lie groups. More precisely, by considering the unit sphere $\mathbb{S}_{\mathbb{H}_n, K}$ with respect to the Koranyi norm $|(z, t)| = (|z|^4 + 16t^2)^{\frac{1}{4}}$ there is a unique Radón measure $d\sigma$ that makes valid the polar decomposition formula. Then, Cowling in [12] proved the L^p -boundedness of the corresponding full-maximal function $\mathcal{M}_F^{d\sigma}$. Extensions of Cowling's result have been obtained e.g. by Schmidt in [41] to hypersurfaces with non-vanishing rotational curvature on nilpotent Lie groups and for two steps nilpotent Lie groups by Fischer [18]. The L^p -boundedness for the Lacunary maximal function in this setting has been proved recently by Ganguly and Thangavelu [21] on \mathbb{H}_n for all $n \geq 2$. The approach in [21] combines the ideas of Cowling [12] with the sparse techniques developed (in the Euclidean setting) by Lacey in [27]. As for the spherical maximal function on the Heisenberg group for the surface measure on the complex

sphere $\mathbb{S}_{\mathbb{H}_n, r} := \{(z, 0) : |z| = r\}$, we refer the reader to Nevo and Thangavelu [32]. The result in [32] was improved independently by Narayanan and Thangavelu [31] and by Müller and Seeger [29]. Moreover, in [1], Bagchi, Hait, Roncal and Thangavelu have proved an analogue of Calderon's theorem for the associated lacunary spherical maximal function on \mathbb{H}_n with $n \geq 2$. For $n = 1$ the L^p -boundedness of the corresponding full maximal function (restricted to a class of radial functions) has been proved by Beltrán, Guo, Hickman and Seeger in [2].

1.2. The main result. Let G be a homogeneous Lie group. Let us consider its corresponding family of dilations (see Definition 2.4)

$$D_r : G \rightarrow G, \quad x \mapsto D_r(x) \equiv r \cdot x, \quad x \in G.$$

Consider the dyadic maximal function

$$\mathcal{M}_D^{d\sigma} f(x) = \sup_{j \in \mathbb{Z}} \left| \int_G f((2^j \cdot y)^{-1}x) d\sigma(y) \right|, \quad (1.3)$$

associated to an arbitrary finite Borel measure $d\sigma$ with compact support on G . We require the existence of left-invariant hypoelliptic partial differential operators on the group and then the group has to be graded (see [19, Page 172] and Definition 2.6). Then, a positive Rockland operator \mathcal{R} is a group Fourier multiplier by the operator-valued function $\pi(\mathcal{R})$, (that is the infinitesimal representation of the operator, defined at any irreducible and unitary representation π of the unitary dual \widehat{G} of G as in (2.3)). In order to analyse the L^p -boundedness of the Dyadic maximal function (1.3) we will assume that for some $a > 0$, the measure $d\sigma$ in (1.3) satisfies the group Fourier transform condition

$$\max_{\pm} \sup_{\pi \in \widehat{G}} \|\pi(\mathcal{R})^{\pm \frac{a}{\nu}} \widehat{d\sigma}(\pi)\|_{\text{op}} < \infty, \quad (1.4)$$

where ν is the homogeneous degree of the operator \mathcal{R} , (in the case of the positive Laplacian $-\Delta$, $\nu = 2$). Then, according to the discussion above, (1.4) says that $\widehat{d\sigma}(\pi)$ has order $+a$ at infinity and $-a$ at zero (with respect to the spectrum of any $\pi(\mathcal{R})$).

The following criterion is the main theorem of this work.

Theorem 1.1. *Let $d\sigma$ be a finite Borel measure of compact support on a graded Lie group G . Let \mathcal{R} be a positive Rockland operator on G of homogeneous degree $\nu > 0$. Assume that for some $a > 0$ the Fourier transform of $d\sigma$ satisfies the growth estimate*

$$\max_{\pm} \sup_{\pi \in \widehat{G}} \|\pi(\mathcal{R})^{\pm \frac{a}{\nu}} \widehat{d\sigma}(\pi)\|_{\text{op}} < \infty. \quad (1.5)$$

Then, the dyadic maximal function $\mathcal{M}_D^{d\sigma} : L^p(G) \rightarrow L^p(G)$ can be extended to a bounded operator for all $1 < p \leq \infty$.

Remark 1.2. For the ℓ^p -boundness of the discrete dyadic maximal function we refer the reader to Bourgain, Mirek, Stein, and Wróbel [4, 5].

Remark 1.3. Our proof of Theorem 1.1 is inspired by the approach developed by Duoandikoetxea and Rubio De Francia in [15]. In the setting of non-commutative nilpotent Lie groups many difficulties arise. One is that the Fourier transform of distributions is operator valued. Also, the Fourier transform condition (1.5) is motivated by conditions of the same nature arising for homogeneous structures (nonisotropic

structures) on the Euclidean space, namely, the condition of non-vanishing curvature at infinite order for Euclidean hypersurfaces $\Sigma \subset \mathbb{R}^n$. Such a condition implies a decay estimate for the Fourier transform $\widehat{d\Sigma}$ of the corresponding surface measure $d\Sigma$, see e.g. Seeger, Tao and Wright [45].

Remark 1.4. The criterion given in Theorem 1.1 is new even on the Heisenberg group. Then also on Heisenberg type groups, stratified groups, etc. Examples of Rockland operators on stratified groups are Hörmander sub-Laplacians and their integer powers, see [19].

Remark 1.5. Let $d\sigma$ be a finite measure of compact support on $G = \mathbb{R}^n$. In the case where $\mathcal{R} = -\Delta_x$ is the positive Laplacian, the inequality in (1.5) becomes equivalent to the Fourier transform condition,

$$\forall \xi \neq 0, |\widehat{d\sigma}(\xi)| \lesssim \min\{|\xi|^a, |\xi|^{-a}\}, \quad a > 0, \quad (1.6)$$

to guarantee the L^p -boundedness of the dyadic maximal operator $\mathcal{M}_D^{d\sigma}$ on $L^p(\mathbb{R}^n)$, for all $1 < p \leq \infty$, see e.g. Theorem 6.3.4 in Grafakos [22, Page 455] or Duoandikoetxea and Rubio De Francia [15]. Let $d\mu$ be the surface measure on the sphere \mathbb{S}^{n-1} . Coifman and Weiss [10, Page 246] deduced the L^p -boundedness of $\mathcal{M}_D^{d\mu}$, $1 \leq p < \infty$, using the fact that

$$\forall \xi \neq 0, |\widehat{d\mu}(\xi)| \lesssim |\xi|^{-(n-1)/2}, \quad n \geq 2. \quad (1.7)$$

Indeed, if $\phi \in C_0^\infty(\mathbb{R}^n)$ is such that $\widehat{\phi}(0) = 1$, for the measure $d\sigma = d\mu - \widehat{\mu}(0)\phi$, Coifman and Weiss proved that the L^p -boundedness of $\mathcal{M}_D^{d\sigma}$, implies the L^p -boundedness of $\mathcal{M}_D^{d\mu}$. This latter argument due to Coifman and Weiss is an alternative proof to the one given in the classical manuscript [7] due to C. Calderón.

Remark 1.6. Let $d\sigma$ be a finite Borel measure of compact support on a graded Lie group G . Let \mathcal{R} be a positive Rockland operator on G of homogeneous degree $\nu > 0$. By the Riesz-representation theorem, we have that $d\sigma = Kdx$ for some compactly supported function K on G . Note that if $\mathcal{R}^{\pm \frac{a}{\nu}} K \in L^1(G)$, (this means that K belongs to the Sobolev space $\dot{L}_{\pm a}^1(G)$, that is $\|K\|_{\dot{L}_{\pm a}^1(G)} := \|\mathcal{R}^{\pm \frac{a}{\nu}} K\|_{L^1(G)} < \infty$) then

$$\sup_{\pi \in \widehat{G}} \|\pi(\mathcal{R})^{\pm \frac{a}{\nu}} \widehat{d\sigma}(\pi)\|_{\text{op}} = \sup_{\pi \in \widehat{G}} \left\| \int_G \mathcal{R}^{\pm \frac{a}{\nu}} K(g) \pi(g)^* dg \right\|_{\text{op}} \leq \|K\|_{\dot{L}_{\pm a}^1(G)} < \infty,$$

showing that the class of compactly supported finite Borel measures $d\sigma = Kdx$ with $K \in \dot{L}_a^1(G) \cap \dot{L}_{-a}^1(G)$, for some $a > 0$, provides examples of measures satisfying (1.5).

2. FOURIER ANALYSIS ON GRADED GROUPS

For the aspects of the Fourier analysis on nilpotent Lie groups we follow Folland and Stein [20] and the notation is taken from [19]. For the aspects about the theory of Rockland operators on graded Lie groups we follow [19].

2.1. Homogeneous and graded Lie groups. Let G be a homogeneous Lie group, that is a connected and simply connected Lie group whose Lie algebra \mathfrak{g} is endowed with a family of dilations $D_{r,\mathfrak{g}}$. We define it as follows.

Definition 2.1. A family of dilations $\text{Dil}(\mathfrak{g}) := \{D_{r,\mathfrak{g}} : r > 0\}$ on the Lie algebra \mathfrak{g} is a family of automorphisms on \mathfrak{g} satisfying the following two compatibility conditions:

1. For every $r > 0$, $D_{r,\mathfrak{g}}$ is a map of the form $D_{r,\mathfrak{g}} = \text{Exp}(\ln(r)A)$, for some diagonalisable linear operator $A \equiv \text{diag}[\nu_1, \dots, \nu_n] : \mathfrak{g} \rightarrow \mathfrak{g}$.
2. $\forall X, Y \in \mathfrak{g}$, and $r > 0$, $[D_{r,\mathfrak{g}}X, D_{r,\mathfrak{g}}Y] = D_{r,\mathfrak{g}}[X, Y]$.

Remark 2.2. We call the eigenvalues of A , $\nu_1, \nu_2, \dots, \nu_n$, the dilations weights or weights of G .

In our analysis the notion of the homogeneous dimension of the group is crucial. We introduce it as follows.

Definition 2.3. The homogeneous dimension of a homogeneous Lie group G whose dilations are defined via $D_{r,\mathfrak{g}} = \text{Exp}(\ln(r)A)$, is given by $Q = \text{Tr}(A) = \nu_1 + \dots + \nu_n$, where $\nu_i, i = 1, 2, \dots, n$, are the eigenvalues of A .

Definition 2.4 (Dilations on the group). The family of dilations $\text{Dil}(\mathfrak{g})$ of the Lie algebra \mathfrak{g} induces a family of mappings on G defined via,

$$\text{Dil}(G) := \{D_r := \exp_G \circ D_{r,\mathfrak{g}} \circ \exp_G^{-1} : r > 0\},$$

where $\exp_G : \mathfrak{g} \rightarrow G$ is the usual exponential mapping associated to the Lie group G . We refer to the elements of the family $\text{Dil}(G)$ as dilations on the group.

Remark 2.5. If we use the notation $r \cdot x = D_r(x)$, $x \in G$, $r > 0$, then the effect of the dilations of the group on the Haar measure dx on G is determined by the identity

$$\int_G (f \circ D_r)(x) dx = r^{-Q} \int_G f(x) dx.$$

Definition 2.6. A connected, simply connected Lie group G is graded if its Lie algebra \mathfrak{g} may be decomposed as the direct sum of subspaces $\mathfrak{g} = \mathfrak{g}_1 \oplus \mathfrak{g}_2 \oplus \dots \oplus \mathfrak{g}_s$ such that the following bracket conditions are satisfied: $[\mathfrak{g}_i, \mathfrak{g}_j] \subset \mathfrak{g}_{i+j}$, where $\mathfrak{g}_{i+j} = \{0\}$ if $i + j \geq s + 1$, for some s .

Examples of graded Lie groups are the Heisenberg group \mathbb{H}^n and more generally any stratified group where the Lie algebra \mathfrak{g} is generated by the first stratum \mathfrak{g}_1 . Here, n is the topological dimension of G , $n = n_1 + \dots + n_s$, where $n_k = \dim \mathfrak{g}_k$. For more examples, see [19].

Remark 2.7 (Not every nilpotent Lie group is homogeneous). A Lie algebra admitting a family of dilations is nilpotent, and hence so is its associated connected, simply connected Lie group. The converse does not hold, i.e., not every nilpotent Lie group is homogeneous although they exhaust a large class, see [19] for details. Indeed, the main class of Lie groups under our consideration is that of graded Lie groups.

2.2. Fourier analysis on nilpotent Lie groups. Let G be a simply connected nilpotent Lie group. Then the adjoint representation $\text{ad} : \mathfrak{g} \rightarrow \text{End}(\mathfrak{g})$ is nilpotent. Next, we define unitary and irreducible representations.

Definition 2.8 (Continuous, unitary and irreducible representations of G). We say that π is a continuous, unitary and irreducible representation of G , if the following properties are satisfied,

1. $\pi \in \text{Hom}(G, \text{U}(H_\pi))$, for some separable Hilbert space H_π , i.e. $\pi(xy) = \pi(x)\pi(y)$ and for the adjoint of $\pi(x)$, $\pi(x)^* = \pi(x^{-1})$, for every $x, y \in G$. This property says that the representation is compatible with the group operation.

2. The map $(x, v) \mapsto \pi(x)v$, from $G \times H_\pi$ into H_π is continuous. This says that the representation is a strongly continuous mapping.
3. For every $x \in G$, and $W_\pi \subset H_\pi$, if $\pi(x)W_\pi \subset W_\pi$, then $W_\pi = H_\pi$ or $W_\pi = \{0\}$. This means that the representation π is irreducible if its only invariant subspaces are $W = \{0\}$ and $W = H_\pi$, the trivial ones.

Definition 2.9 (Equivalent representations). Two unitary representations

$$\pi \in \text{Hom}(G, \text{U}(H_\pi)) \text{ and } \eta \in \text{Hom}(G, \text{U}(H_\eta))$$

are equivalent if there exists a bounded linear mapping $Z : H_\pi \rightarrow H_\eta$ such that for any $x \in G$, $Z\pi(x) = \eta(x)Z$. The mapping Z is called an intertwining operator between π and η . The set of all the intertwining operators between π and η is denoted by $\text{Hom}(\pi, \eta)$.

Definition 2.10 (The unitary dual). The relation \sim on the set of unitary and irreducible representations $\text{Rep}(G)$ defined by: $\pi \sim \eta$ if and only if π and η are equivalent representations, is an equivalence relation. The quotient

$$\widehat{G} := \text{Rep}(G)/\sim$$

is called the unitary dual of G .

The unitary dual encodes all the Fourier analysis on the group. So, we are going to define the Fourier transform.

Definition 2.11 (Group Fourier Transform). The Fourier transform of $f \in L^1(G)$, at $\pi \in \widehat{G}$, is defined by

$$\widehat{f}(\pi) = \int_G f(x)\pi(x)^* dx : H_\pi \rightarrow H_\pi.$$

Remark 2.12. The Schwartz space $\mathcal{S}(G)$ is defined by the smooth functions $f : G \rightarrow \mathbb{C}$, such that via the exponential mapping $f \circ \exp_G : \mathfrak{g} \cong \mathbb{R}^n \rightarrow \mathbb{C}$ can be identified with functions on the Schwartz class $\mathcal{S}(\mathbb{R}^n)$. Then, the Schwartz space on the dual \widehat{G} is defined by the image under the Fourier transform of the Schwartz space $\mathcal{S}(G)$, that is $\mathcal{F}_G : \mathcal{S}(G) \rightarrow \mathcal{S}(\widehat{G}) := \mathcal{F}_G(\mathcal{S}(G))$.

Remark 2.13 (Fourier Inversion Formula and Plancherel Theorem). If we identify one representation π with its equivalence class, $[\pi] = \{\pi' : \pi \sim \pi'\}$, for every $\pi \in \widehat{G}$, the Kirillov trace character Θ_π defined by

$$[\Theta_\pi, f] := \text{Tr}(\widehat{f}(\pi)),$$

is a tempered distribution on $\mathcal{S}(G)$. The Kirillov character allows one to write the Fourier inversion formula

$$\forall f \in L^1(G) \cap L^2(G), \quad f(x) = \int_{\widehat{G}} \text{Tr}[\pi(x)\widehat{f}(\pi)] d\pi, \quad x \in G.$$

The L^2 -space on the dual is defined by the completion of $\mathcal{S}(G)$ with respect to the norm

$$\|\sigma\|_{L^2(\widehat{G})} := \left(\int_G \|\sigma(\pi)\|_{\text{HS}}^2 d\pi \right)^{\frac{1}{2}}, \quad \sigma(\pi) = \widehat{f}(\pi) \in \mathcal{S}(\widehat{G}), \quad (2.1)$$

where $\|\cdot\|_{\text{HS}}$ denotes the Hilbert-Schmidt norm of operators on every representation space. The corresponding inner product on $L^2(\widehat{G})$ is given by

$$(\sigma, \tau)_{L^2(\widehat{G})} := \int_G \text{Tr}[\sigma(\pi)\tau(\pi)^*]d\pi, \quad \sigma, \tau \in L^2(\widehat{G}), \quad (2.2)$$

where the notation $\tau(\pi)^*$ indicates the adjoint operator. Then, the Plancherel theorem says that $\|f\|_{L^2(G)} = \|\widehat{f}\|_{L^2(\widehat{G})}$ for all $f \in L^2(G)$.

2.3. Homogeneous linear operators and Rockland operators. Homogeneous operators interact with the dilations of the group. We introduce them in the following definition.

Definition 2.14 (Homogeneous operators). A continuous linear operator $T : C^\infty(G) \rightarrow \mathcal{D}'(G)$ is homogeneous of degree $\nu_T \in \mathbb{C}$ if for every $r > 0$ the equality

$$T(f \circ D_r) = r^{\nu_T}(Tf) \circ D_r$$

holds for every $f \in \mathcal{D}(G)$.

Now, we introduce the main class of differential operators in the context of nilpotent Lie groups. The existence of these operators classifies the family of graded Lie groups. We call them Rockland operators.

Definition 2.15 (Rockland operators). If for every representation $\pi \in \widehat{G}$, $\pi : G \rightarrow U(H_\pi)$, we denote by H_π^∞ the set of smooth vectors (also called Gårding vectors), that is, the space of vectors $v \in H_\pi$ such that the function $x \mapsto \pi(x)v$, $x \in \widehat{G}$, is smooth, a Rockland operator is a left-invariant partial differential operator

$$\mathcal{R} = \sum_{|\alpha| \leq m} a_\alpha X^\alpha : C^\infty(G) \rightarrow C^\infty(G)$$

which is homogeneous of positive degree $\nu = \nu_{\mathcal{R}}$ and such that, for every unitary irreducible non-trivial representation $\pi \in \widehat{G}$, its symbol $\pi(\mathcal{R})$ defined via the Fourier inversion formula by

$$\mathcal{R}f(x) = \int_{\widehat{G}} \text{Tr}[\pi(x)\pi(\mathcal{R})\widehat{f}(\pi)]d\pi, \quad x \in G, \quad (2.3)$$

is injective on H_π^∞ ; $\sigma_{\mathcal{R}}(\pi) = \pi(\mathcal{R})$ coincides with the infinitesimal representation of \mathcal{R} as an element of the universal enveloping algebra $\mathfrak{U}(\mathfrak{g})$.

Example 2.16. Let G be a graded Lie group of topological dimension n . We denote by $\{D_r\}_{r>0}$ the natural family of dilations of its Lie algebra $\mathfrak{g} := \text{Lie}(G)$, and by ν_1, \dots, ν_n its weights. We fix a basis $Y = \{X_1, \dots, X_n\}$ of \mathfrak{g} satisfying $D_r X_j = r^{\nu_j} X_j$, for $1 \leq j \leq n$, and all $r > 0$. If ν_o is any common multiple of ν_1, \dots, ν_n , the operator

$$\mathcal{R} = \sum_{j=1}^n (-1)^{\frac{\nu_o}{\nu_j}} c_j X_j^{\frac{2\nu_o}{\nu_j}}, \quad c_j > 0,$$

is a positive Rockland operator of homogeneous degree $2\nu_o$ on G (see Lemma 4.1.8 of [19]).

Remark 2.17. It can be shown that a Lie group G is graded if and only if there exists a differential Rockland operator on G . Also, in view of the Schwartz kernel theorem, the operator \mathcal{R} admits a right-convolution kernel $k_{\mathcal{R}}$, that is for any $f \in C_0^\infty(G)$,

$$\mathcal{R}f(x) = \int_G f(y)k_{\mathcal{R}}(y^{-1}x)dy, \quad x \in G.$$

In terms of the convolution of functions

$$f * g(x) := \int_G f(y)g(y^{-1}x)dy, \quad f, g \in L^1(G), \quad (2.4)$$

and in view of the action of the Fourier transform on convolutions $\widehat{f * g} = \widehat{g}\widehat{f}$, the Fourier inversion formula shows that $\forall \pi \in \widehat{G}$, $\pi(\mathcal{R}) = \widehat{k_{\mathcal{R}}}(\pi)$.

Next, we record for our further analysis some aspects of the functional calculus for Rockland operators.

Remark 2.18 (Functional calculus for Rockland operators). If the Rockland operator is formally self-adjoint, then \mathcal{R} and $\pi(\mathcal{R})$ admit self-adjoint extensions on $L^2(G)$ and H_π , respectively. Now if we preserve the same notation for their self-adjoint extensions and we denote by E and E_π their spectral measures, we will denote by

$$\psi(\mathcal{R}) = \int_{-\infty}^{\infty} \psi(\lambda)dE(\lambda), \quad \text{and} \quad \pi(\psi(\mathcal{R})) \equiv \psi(\pi(\mathcal{R})) = \int_{-\infty}^{\infty} \psi(\lambda)dE_\pi(\lambda), \quad (2.5)$$

the functions defined by the functional calculus. In general, we will reserve the notation $\{dE_A(\lambda)\}_{0 \leq \lambda < \infty}$ for the spectral measure associated with a positive and self-adjoint operator A on a Hilbert space H .

We now recall a lemma on dilations on the unitary dual \widehat{G} , which will be useful in our analysis of spectral multipliers. For the proof, see Lemma 4.3 of [19].

Lemma 2.19. *For every $\pi \in \widehat{G}$ let us define*

$$D_r(\pi)(x) \equiv (r \cdot \pi)(x) := \pi(r \cdot x) \equiv \pi(D_r(x)), \quad (2.6)$$

for every $r > 0$ and all $x \in G$. Then, if $f \in L^\infty(\mathbb{R})$ then $f(\pi^{(r)}(\mathcal{R})) = f(r^\nu \pi(\mathcal{R}))$.

Remark 2.20. Note that if $f_r := r^{-Q}f(r^{-1}\cdot)$, then

$$\widehat{f_r}(\pi) = \int_G r^{-Q}f(r^{-1} \cdot x)\pi(x)^*dx = \int_G f(y)\pi(r \cdot y)^*dy = \widehat{f}(r \cdot \pi), \quad (2.7)$$

for any $\pi \in \widehat{G}$ and all $r > 0$, with $(r \cdot \pi)(y) = \pi(r \cdot y)$, $y \in G$, as in (2.6).

The following lemma present the action of the dilations of the group G into the kernels of bounded functions of a Rockland operator \mathcal{R} , see [19, Page 179].

Lemma 2.21. *Let $f \in L^\infty(\mathbb{R}_0^+)$ be a bounded Borel function and let $r > 0$. Then, we have*

$$\forall x \in G, \quad f(r^\nu \mathcal{R})\delta(x) = r^{-Q}[f(\mathcal{R})\delta](r^{-1} \cdot x), \quad (2.8)$$

where Q is the homogeneous dimension of G .

3. BOUNDEDNESS OF THE DYADIC MAXIMAL FUNCTION

In this section we establish the L^p -boundedness of the dyadic maximal function associated to a compactly supported Borel measure on a graded Lie group satisfying some additional Fourier transform conditions according to the hypothesis in Theorem 1.1. First, we start with our main lemma in the next subsection.

3.1. The key lemma. The following Lemma 3.1 is our main tool for the proof of Theorem 1.1. Indeed, it will be used to use the L^p -boundedness of the square function operator in Lemma 3.2 from which we will deduce the boundedness of the dyadic maximal function (1.3).

Lemma 3.1. *Let $K \in L^1(G)$ be a distribution with compact support such that for some $a > 0$ the group Fourier transform of K satisfies the growth estimate*

$$\max_{\pm} \sup_{\pi \in \widehat{G}} \|\pi(\mathcal{R})^{\pm \frac{a}{\nu}} \widehat{K}(\pi)\|_{\text{op}} < \infty. \quad (3.1)$$

For any $j \in \mathbb{Z}$, let us consider the kernel $K_j(x) = 2^{-jQ} K(2^{-j} \cdot x)$, $x \in G$, and define T as follows

$$Tf(x) := \sum_{j=-\infty}^{\infty} f * K_j(x), \quad f \in C_0^\infty(G). \quad (3.2)$$

Then, $T : L^p(G) \rightarrow L^p(G)$ admits a bounded extension for all $1 < p < \infty$.

Proof. We start the proof by considering a suitable partition of unity for the spectrum of the Rockland operator \mathcal{R} . For this, let us take a function $\Phi \in \mathcal{S}(\mathbb{R})$ such that (see Lemma 3.13 in [26])

$$\forall \lambda \in (0, \infty), \quad \sum_{j=-\infty}^{\infty} \Phi(2^{j\nu} \lambda) = 1. \quad (3.3)$$

Moreover, we can assume that Φ generates a dyadic partition in the sense that $\text{supp}(\Phi) \subset [1/2^\nu, 2^\nu]$. As a consequence we have that

$$\sum_{j=-\infty}^{\infty} \Phi(2^{j\nu} \mathcal{R}) = I = \text{identity operator on } L^2(G), \quad (3.4)$$

and the convergence in $\mathcal{S}'(G)$ to the Dirac distribution

$$\sum_{j=-\infty}^{\infty} \Phi(2^{j\nu} \mathcal{R}) \delta = \delta. \quad (3.5)$$

In view of the property in (2.5) for the functional calculus of \mathcal{R} , taking the group Fourier transform of (3.5) in both sides we obtain that

$$\forall \pi \in \widehat{G}, \quad \sum_{j=-\infty}^{\infty} \mathcal{F}_G[\Phi(\mathcal{R})\delta](2^j \cdot \pi) = \sum_{j=-\infty}^{\infty} \Phi[(2^j \cdot \pi)(\mathcal{R})] = I_{H_\pi}, \quad (3.6)$$

where we have used that, for any j , $(2^j \cdot \pi)(\mathcal{R}) = 2^{j\nu} \pi(\mathcal{R})$. To simplify the notation, let us define

$$\forall x \in G, \forall j \in \mathbb{Z}, \quad \Phi_j(x) := 2^{-jQ} (\Phi(\mathcal{R})\delta)(2^{-j} \cdot), \quad \Phi(x) := (\Phi(\mathcal{R})\delta)(x). \quad (3.7)$$

Then, we have

$$\forall \pi \in \widehat{G}, \forall j \in \mathbb{Z}, \widehat{\Phi}_j(\pi) = \mathcal{F}_G[\Phi(\mathcal{R})\delta](2^j \cdot \pi) = \Phi[(2^j \cdot \pi)(\mathcal{R})] = \Phi[2^{j\nu}\pi(\mathcal{R})]. \quad (3.8)$$

In view of (3.6), we conclude that

$$\forall \pi \in \widehat{G}, \sum_{j=-\infty}^{\infty} \widehat{\Phi}_j(\pi) = I_{H_\pi}, \quad (3.9)$$

and then, in the sense of distributions we have the identity

$$\sum_{j=-\infty}^{\infty} \Phi_j = \delta. \quad (3.10)$$

Note also, that if $\{dE_{\pi(\mathcal{R})}\}_{\lambda>0}$ is the spectral measure of the operator $\pi(\mathcal{R})$, then

$$\forall j \in \mathbb{Z}, \forall \pi \in \widehat{G}, \widehat{\Phi}_j(\pi) = \Phi[(2^j \cdot \pi)(\mathcal{R})] = \Phi[2^{j\nu}\pi(\mathcal{R})] = \int_0^\infty \Phi(2^{j\nu}\lambda) dE_{\pi(\mathcal{R})}(\lambda). \quad (3.11)$$

These properties of the partition of the unity Φ_j , $j \in \mathbb{Z}$, will be used in our further analysis. Indeed, we start using (3.10) to decompose any K_j as follows:

$$K_j = K_j * \delta = \sum_{k=-\infty}^{\infty} K_j * \Phi_{j+k}. \quad (3.12)$$

Define the linear operator \tilde{T}_k for any $k \in \mathbb{Z}$ as follows,

$$\forall f \in C_0^\infty(G), \tilde{T}_k f := \sum_{j=-\infty}^{\infty} f * K_j * \Phi_{j+k}. \quad (3.13)$$

Note that

$$\sum_{k=-\infty}^{\infty} \tilde{T}_k f = \sum_{k=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} f * K_j * \Phi_{j+k} = \sum_{k=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} f * K_k * \Phi_{k+j} = \sum_{k=-\infty}^{\infty} f * K_k =: Tf.$$

In view of the last identity we will split our proof in the following steps.

- Step 1. To estimate the norm of the operator $\tilde{T}_k : L^2(G) \rightarrow L^2(G)$. Moreover, we will prove that

$$\exists C > 0, \forall f \in C_0^\infty(G), \|\tilde{T}_k f\|_{L^2(G)} \leq C 2^{-a|k|} \|f\|_{L^2(G)}. \quad (3.14)$$

- Step 2. To estimate the norm of the operator $\tilde{T}_k : L^1(G) \rightarrow L^{1,\infty}(G)$. Moreover, we will prove that

$$\exists C > 0, \forall \lambda > 0, \forall f \in C_0^\infty(G), |\{x \in G : |\tilde{T}_k f(x)| > \lambda\}| \leq \frac{C(1+|k|)}{\lambda} \|f\|_{L^1(G)}. \quad (3.15)$$

- Step 3. To use Marcinkiewicz interpolation theorem to prove that for any $1 < p < 2$,

$$\exists C > 0, \forall f \in C_0^\infty(G), \|\tilde{T}_k f\|_{L^p(G)} \leq C_p 2^{-a|k|\theta} (1+|k|)^{1-\theta} \|f\|_{L^p(G)}, \quad (3.16)$$

where $1/p = \theta/2 + (1-\theta)$.

- Final Step. The proof of Theorem 3.1 follows if we sum over $k \in \mathbb{Z}$ both sides of (3.16), in the case when $1 < p < 2$, and then by the duality argument we complete then L^p -boundedness of T for $2 < p < \infty$.

Once proved Steps 1 and 2 all the other steps above are clear. It remains, therefore, to prove Steps 1 and 2.

3.1.1. *Step 1.* Let us prove that

$$\exists C > 0, \forall f \in C_0^\infty(G), \|\tilde{T}_k f\|_{L^2(G)} \leq C 2^{-a|k|} \|f\|_{L^2(G)}. \quad (3.17)$$

From (3.12) we have the identity

$$\forall j \in \mathbb{Z}, \forall \pi \in \widehat{G}, \widehat{K}_j(\pi) = \sum_{m=-\infty}^{\infty} \widehat{\Phi}_{m+j}(\pi) \widehat{K}_j(\pi), \quad (3.18)$$

in the strong topology on H_π . Using this fact and Plancherel theorem we deduce that

$$\|\tilde{T}_k f\|_{L^2(G)}^2 = (\tilde{T}_k f, \tilde{T}_k f)_{L^2(G)} = (\widehat{\tilde{T}_k f}, \widehat{\tilde{T}_k f})_{L^2(\widehat{G})}, \quad (3.19)$$

and by writing the right-hand side of this identity in integral form we have that

$$\begin{aligned} \|\tilde{T}_k f\|_{L^2(G)}^2 &= \sum_{m=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} (\widehat{\Phi}_{m+k}(\pi) \widehat{K}_m(\pi) \widehat{f}(\pi), \widehat{\Phi}_{j+k}(\pi) \widehat{K}_j(\pi) \widehat{f}(\pi))_{L^2(\widehat{G})} \\ &\leq \sum_{m=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \int_{\widehat{G}} |\mathrm{Tr}[\widehat{\Phi}_{m+k}(\pi) \widehat{K}_m(\pi) \widehat{f}(\pi) (\widehat{\Phi}_{j+k}(\pi) \widehat{K}_j(\pi) \widehat{f}(\pi))^*]| d\pi \\ &= \sum_{j=-\infty}^{\infty} \sum_{m=-\infty}^{\infty} \int_{\widehat{G}} |\mathrm{Tr}[\widehat{\Phi}_{m+k}(\pi) \widehat{K}_m(\pi) \widehat{f}(\pi) \widehat{f}(\pi)^* \widehat{K}_j(\pi)^* \widehat{\Phi}_{j+k}(\pi)^*]| d\pi. \end{aligned}$$

Because of (3.8), $\widehat{\Phi}_{j+k}(\pi)$ is self-adjoint in every representation space, and the operators $\widehat{\Phi}_{m+k}(\pi)$, $\widehat{\Phi}_{j+k}(\pi)$, commute with each other and with other Borel functions of $\pi(\mathcal{R})$, and we can write

$$\begin{aligned} \|\tilde{T}_k f\|_{L^2(G)}^2 &\leq \sum_{j=-\infty}^{\infty} \sum_{m=-\infty}^{\infty} \int_{\widehat{G}} |\mathrm{Tr}[\widehat{\Phi}_{m+k}(\pi) \widehat{K}_m(\pi) \widehat{f}(\pi) \widehat{f}(\pi)^* \widehat{K}_j(\pi)^* \widehat{\Phi}_{j+k}(\pi)]| d\pi \\ &= \sum_{j=-\infty}^{\infty} \sum_{m=-\infty}^{\infty} \int_{\widehat{G}} |\mathrm{Tr}[\widehat{K}_j(\pi)^* \widehat{\Phi}_{j+k}(\pi) \widehat{\Phi}_{m+k}(\pi) \widehat{K}_m(\pi) \widehat{f}(\pi) \widehat{f}(\pi)^*]| d\pi \\ &= \sum_{j=-\infty}^{\infty} \sum_{m=-\infty}^{\infty} \int_{\widehat{G}} |\mathrm{Tr}[\widehat{K}(2^j \cdot \pi)^* [(2^j \cdot \pi)(\mathcal{R})^{\pm \frac{a}{\nu}}] [(2^j \cdot \pi)(\mathcal{R})^{\mp \frac{a}{\nu}}] \\ &\quad \widehat{\Phi}_{j+k}(\pi) \widehat{\Phi}_{m+k}(\pi) \widehat{K}_m(\pi) \widehat{f}(\pi) \widehat{f}(\pi)^*]| d\pi \\ &\leq \sum_{j=-\infty}^{\infty} \sum_{m=-\infty}^{\infty} \int_{\widehat{G}} \|\widehat{K}(2^j \cdot \pi)^* [(2^j \cdot \pi)(\mathcal{R})^{\pm \frac{a}{\nu}}]\|_{\mathrm{op}} |\mathrm{Tr}[(2^j \cdot \pi)(\mathcal{R})^{\mp \frac{a}{\nu}}] \\ &\quad \widehat{\Phi}_{j+k}(\pi) \widehat{\Phi}_{m+k}(\pi) \widehat{K}_m(\pi) \widehat{f}(\pi) \widehat{f}(\pi)^*]| d\pi \end{aligned}$$

$$\begin{aligned}
&= \sum_{j=-\infty}^{\infty} \sum_{m=-\infty}^{\infty} \int_{\widehat{G}} f \left\| \left[(2^j \cdot \pi)(\mathcal{R})^{\pm \frac{a}{\nu}} \widehat{K}(2^j \cdot \pi) \right]^* \right\|_{\text{op}} |\text{Tr} \left[(2^j \cdot \pi)(\mathcal{R})^{\mp \frac{a}{\nu}} \right. \\
&\quad \left. \widehat{\Phi}_{j+k}(\pi) \widehat{\Phi}_{m+k}(\pi) \widehat{K}_m(\pi) \widehat{f}(\pi) \widehat{f}(\pi)^* \right]| d\pi \\
&= \sum_{j=-\infty}^{\infty} \sum_{m=-\infty}^{\infty} \int_{\widehat{G}} f \left\| \left[(2^j \cdot \pi)(\mathcal{R})^{\pm \frac{a}{\nu}} \widehat{K}(2^j \cdot \pi) \right] \right\|_{\text{op}} |\text{Tr} \left[(2^j \cdot \pi)(\mathcal{R})^{\mp \frac{a}{\nu}} \right. \\
&\quad \left. \widehat{\Phi}_{j+k}(\pi) \widehat{\Phi}_{m+k}(\pi) \widehat{K}_m(\pi) \widehat{f}(\pi) \widehat{f}(\pi)^* \right]| d\pi \\
&\lesssim \sum_{j=-\infty}^{\infty} \sum_{m=-\infty}^{\infty} \int_{\widehat{G}} |\text{Tr} \left[(2^j \cdot \pi)(\mathcal{R})^{\mp \frac{a}{\nu}} \right. \\
&\quad \left. \widehat{\Phi}_{j+k}(\pi) \widehat{\Phi}_{m+k}(\pi) \widehat{K}_m(\pi) \widehat{f}(\pi) \widehat{f}(\pi)^* \right]| d\pi \\
&= \sum_{j=-\infty}^{\infty} \sum_{m=-\infty}^{\infty} \int_{\widehat{G}} |\text{Tr} \left[\widehat{\Phi}_{j+k}(\pi) \widehat{\Phi}_{m+k}(\pi) \left[(2^j \cdot \pi)(\mathcal{R})^{\mp \frac{a}{\nu}} \widehat{K}_m(\pi) \widehat{f}(\pi) \widehat{f}(\pi)^* \right] \right]| d\pi \\
&= \sum_{j=-\infty}^{\infty} \sum_{m=-\infty}^{\infty} \int_{\widehat{G}} |\text{Tr} \left[\widehat{f}(\pi) \widehat{f}(\pi)^* \widehat{\Phi}_{j+k}(\pi) \widehat{\Phi}_{m+k}(\pi) \left[(2^j \cdot \pi)(\mathcal{R})^{\mp \frac{a}{\nu}} \widehat{K}_m(\pi) \right] \right]| d\pi,
\end{aligned}$$

where we have used the kernel condition

$$\sup_{j \in \mathbb{Z}, \pi \in \widehat{G}} \left\| \left[(2^j \cdot \pi)(\mathcal{R})^{\mp \frac{a}{\nu}} \widehat{K}(2^j \cdot \pi) \right] \right\|_{\text{op}} \leq \sup_{\pi \in \widehat{G}} \left\| \pi(\mathcal{R})^{\mp \frac{a}{\nu}} \widehat{K}(\pi) \right\|_{\text{op}} < \infty. \quad (3.20)$$

In view of (3.11), the properties of the functional calculus allow us to write

$$\widehat{\Phi}_{j+k}(\pi) \widehat{\Phi}_{m+k}(\pi) = \int_0^{\infty} \Phi(2^{(j+k)\nu} \lambda) \Phi(2^{(m+k)\nu} \lambda) dE_{\pi(\mathcal{R})}(\lambda). \quad (3.21)$$

Since the support of Φ lies in the interval $[1/2^\nu, 2^\nu]$ we have that

$$\forall \ell \neq \ell', \text{ such that } |\ell - \ell'| \geq 2, \quad \Phi_\ell(\lambda) \Phi_{\ell'}(\lambda) = 0.$$

In consequence the representation in (3.21) shows that $\widehat{\Phi}_{j+k}(\pi) \widehat{\Phi}_{m+k}(\pi) \equiv 0_{H_\pi}$ if $|m - j| \geq 2$. So, we have that

$$\|\tilde{T}_k f\|_{L^2(G)}^2 \lesssim \sum_{j=-\infty}^{\infty} \sum_{m=j-1}^{j+1} \int_{\widehat{G}} |\text{Tr} \left[\widehat{f}(\pi) \widehat{f}(\pi)^* \widehat{\Phi}_{j+k}(\pi) \widehat{\Phi}_{m+k}(\pi) \left[(2^j \cdot \pi)(\mathcal{R})^{\mp \frac{a}{\nu}} \widehat{K}_m(\pi) \right] \right]| d\pi. \quad (3.22)$$

In view of (3.20) we have that

$$\begin{aligned}
\mathcal{A}(\pi) &:= |\text{Tr} \left[\widehat{f}(\pi) \widehat{f}(\pi)^* \widehat{\Phi}_{j+k}(\pi) \widehat{\Phi}_{m+k}(\pi) \left[(2^j \cdot \pi)(\mathcal{R})^{\mp \frac{a}{\nu}} \widehat{K}_m(\pi) \right] \right]| \\
&= |\text{Tr} \left[\widehat{f}(\pi) \widehat{f}(\pi)^* \widehat{\Phi}_{j+k}(\pi) \widehat{\Phi}_{m+k} \left[(2^j \cdot \pi)(\mathcal{R}) \right]^{\mp \frac{a}{\nu}}(\pi) \widehat{K}(2^m \cdot \pi) \right]| \\
&= |\text{Tr} \left[\widehat{f}(\pi) \widehat{f}(\pi)^* \widehat{\Phi}_{j+k}(\pi) \widehat{\Phi}_{m+k}(\pi) \left[2^{j\nu} \times \pi(\mathcal{R}) \right]^{\mp \frac{a}{\nu}} \widehat{K}(2^m \cdot \pi) \right]| \\
&= |\text{Tr} \left[\widehat{f}(\pi) \widehat{f}(\pi)^* \widehat{\Phi}_{j+k}(\pi) \widehat{\Phi}_{m+k}(\pi) \left[2^{(j-m)\nu} \times 2^{m\nu} \times \pi(\mathcal{R}) \right]^{\mp \frac{a}{\nu}} \widehat{K}(2^m \cdot \pi) \right]| \\
&= 2^{\mp(j-m)a} |\text{Tr} \left[\widehat{f}(\pi) \widehat{f}(\pi)^* \widehat{\Phi}_{j+k}(\pi) \widehat{\Phi}_{m+k}(\pi) \left[2^{m\nu} \times \pi(\mathcal{R}) \right]^{\mp \frac{a}{\nu}} \widehat{K}(2^m \cdot \pi) \right]|.
\end{aligned}$$

Since $j - m = 0, \pm 1$, $2^{\mp 2(j-m)a} \simeq 1$, we have

$$\begin{aligned} \mathcal{A}(\pi) &:= |\mathrm{Tr}[[\widehat{f}(\pi)\widehat{f}(\pi)^*]\widehat{\Phi}_{j+k}(\pi)\widehat{\Phi}_{m+k}(\pi)[(2^j \cdot \pi)(\mathcal{R})]^{\mp \frac{a}{\nu}}\widehat{K}_m(\pi)]| \\ &= 2^{\mp 2(j-m)a} |\mathrm{Tr}[[\widehat{f}(\pi)\widehat{f}(\pi)^*]\widehat{\Phi}_{j+k}(\pi)\widehat{\Phi}_{m+k}(\pi)[2^{m\nu} \times \pi(\mathcal{R})]^{\mp \frac{a}{\nu}}\widehat{K}(2^m \cdot \pi)]| \\ &\asymp |\mathrm{Tr}[[\widehat{f}(\pi)\widehat{f}(\pi)^*]\widehat{\Phi}_{j+k}(\pi)\widehat{\Phi}_{m+k}(\pi)[2^{m\nu} \times \pi(\mathcal{R})]^{\mp \frac{a}{\nu}}\widehat{K}(2^m \cdot \pi)]|. \end{aligned}$$

Until now we have proved that the term

$$\mathcal{A}(\pi) := |\mathrm{Tr}[[\widehat{f}(\pi)\widehat{f}(\pi)^*]\widehat{\Phi}_{j+k}(\pi)\widehat{\Phi}_{m+k}(\pi)[(2^j \cdot \pi)(\mathcal{R})]^{\mp \frac{a}{\nu}}\widehat{K}_m(\pi)]|$$

can be estimated as follows

$$\mathcal{A}(\pi) \asymp |\mathrm{Tr}[[\widehat{f}(\pi)\widehat{f}(\pi)^*]\widehat{\Phi}_{j+k}(\pi)\widehat{\Phi}_{m+k}(\pi)[2^{m\nu} \times \pi(\mathcal{R})]^{\mp \frac{a}{\nu}}\widehat{K}(2^m \cdot \pi)]|. \quad (3.23)$$

To estimate the right-hand side of (3.23) let us use (3.20) again as follows

$$\begin{aligned} \mathcal{A}(\pi) &\asymp |\mathrm{Tr}[[\widehat{f}(\pi)\widehat{f}(\pi)^*]\widehat{\Phi}_{j+k}(\pi)\widehat{\Phi}_{m+k}(\pi)[(2^m \cdot \pi)(\mathcal{R})]^{\mp \frac{2a}{\nu}}[(2^m \cdot \pi)(\mathcal{R})]^{\pm \frac{a}{\nu}}\widehat{K}(2^m \cdot \pi)]| \\ &\leq \mathrm{Tr}[[\widehat{f}(\pi)\widehat{f}(\pi)^*]\widehat{\Phi}_{j+k}(\pi)\widehat{\Phi}_{m+k}(\pi)[(2^m \cdot \pi)(\mathcal{R})]^{\mp \frac{2a}{\nu}}] \|[(2^m \cdot \pi)(\mathcal{R})]^{\pm \frac{a}{\nu}}\widehat{K}(2^m \cdot \pi)\|_{\mathrm{op}} \\ &\lesssim \mathrm{Tr}[[\widehat{f}(\pi)\widehat{f}(\pi)^*]\widehat{\Phi}_{j+k}(\pi)\widehat{\Phi}_{m+k}(\pi)[(2^m \cdot \pi)(\mathcal{R})]^{\mp \frac{2a}{\nu}}]. \end{aligned}$$

Now, we have the better estimate (because it only depends on the functional calculus of the symbol $\pi(\mathcal{R})$ of the Rockland operator \mathcal{R} and of \widehat{f})

$$\mathcal{A}(\pi) \lesssim \mathrm{Tr}[[\widehat{f}(\pi)\widehat{f}(\pi)^*]\widehat{\Phi}_{j+k}(\pi)\widehat{\Phi}_{m+k}(\pi)[(2^m \cdot \pi)(\mathcal{R})]^{\mp \frac{2a}{\nu}}]. \quad (3.24)$$

Note that when composing $\widehat{\Phi}_{j+k}(\pi)\widehat{\Phi}_{m+k}(\pi)$ with the operator $[(2^m \cdot \pi)(\mathcal{R})]^{\pm \frac{2a}{\nu}}$ we remove the zero from the spectrum of the new operator

$$\widehat{\Phi}_{j+k}(\pi)\widehat{\Phi}_{m+k}(\pi)[(2^m \cdot \pi)(\mathcal{R})]^{\pm \frac{2a}{\nu}} \quad (3.25)$$

because of the spectral identity in (3.8) and the properties of the support of Φ . From (3.24) we also have the estimate

$$\mathcal{A}(\pi) \lesssim \mathrm{Tr}[[\widehat{f}(\pi)\widehat{f}(\pi)^*]\widehat{\Phi}_{j+k}(\pi)\widehat{\Phi}_{m+k}(\pi)[(2^m \cdot \pi)(\mathcal{R})]^{-\frac{2a}{\nu}}]. \quad (3.26)$$

Now, using the functional calculus for Rockland operators we can estimate (3.26). Indeed, we have

$$\begin{aligned} &\mathrm{Tr}[[\widehat{f}(\pi)\widehat{f}(\pi)^*]\widehat{\Phi}_{j+k}(\pi)\widehat{\Phi}_{m+k}(\pi)[(2^m \cdot \pi)(\mathcal{R})]^{-\frac{2a}{\nu}}] \\ &= \mathrm{Tr}[[\widehat{f}(\pi)\widehat{f}(\pi)^*]\widehat{\Phi}_{j+k}(\pi)\widehat{\Phi}_{m+k}(\pi)[(2^{m\nu} \pi)(\mathcal{R})]^{-\frac{2a}{\nu}}] \\ &= \mathrm{Tr}[[\widehat{f}(\pi)\widehat{f}(\pi)^*]\widehat{\Phi}_{j+k}(\pi)\widehat{\Phi}_{m+k}(\pi)2^{-2ma}[\pi(\mathcal{R})]^{-\frac{2a}{\nu}}] \\ &= 2^{-2ma} \mathrm{Tr}[[\widehat{f}(\pi)\widehat{f}(\pi)^*]\widehat{\Phi}_{j+k}(\pi)\widehat{\Phi}_{m+k}(\pi)[\pi(\mathcal{R})]^{-\frac{2a}{\nu}}] \\ &\asymp 2^{-2ja} \mathrm{Tr}[[\widehat{f}(\pi)\widehat{f}(\pi)^*]\widehat{\Phi}_{j+k}(\pi)\widehat{\Phi}_{m+k}(\pi)[\pi(\mathcal{R})]^{-\frac{2a}{\nu}}]. \end{aligned}$$

Summarising we have proved that

$$\mathcal{A}(\pi) \lesssim 2^{-2ja} \mathrm{Tr}[[\widehat{f}(\pi)\widehat{f}(\pi)^*]\widehat{\Phi}_{j+k}(\pi)\widehat{\Phi}_{m+k}(\pi)[\pi(\mathcal{R})]^{-\frac{2a}{\nu}}]. \quad (3.27)$$

Note that in the same way that (3.24) implies (3.27) by changing $-a$ by $+a$ in the argument of this implication we also have that (3.26) implies

$$\mathcal{A}(\pi) \lesssim 2^{2ja} \mathrm{Tr}[[\widehat{f}(\pi)\widehat{f}(\pi)^*]\widehat{\Phi}_{j+k}(\pi)\widehat{\Phi}_{m+k}(\pi)[\pi(\mathcal{R})]^{\frac{2a}{\nu}}]. \quad (3.28)$$

Then, from (3.27) and (3.28) we have the following similar bounds

$$\mathcal{A}(\pi) \lesssim 2^{-2ja} \max_{\pm} \text{Tr} \left[\left| \left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] \widehat{\Phi}_{j+k}(\pi) \widehat{\Phi}_{m+k}(\pi) [\pi(\mathcal{R})]^{\pm \frac{2a}{\nu}} \right| \right], \quad (3.29)$$

and

$$\mathcal{A}(\pi) \lesssim 2^{2ja} \max_{\pm} \text{Tr} \left[\left| \left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] \widehat{\Phi}_{j+k}(\pi) \widehat{\Phi}_{m+k}(\pi) [\pi(\mathcal{R})]^{\pm \frac{2a}{\nu}} \right| \right]. \quad (3.30)$$

Note that (3.29) and (3.30) imply the estimate

$$\mathcal{A}(\pi) \lesssim \min\{2^{2ja}, 2^{-2ja}\} \max_{\pm} \text{Tr} \left[\left| \left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] \widehat{\Phi}_{j+k}(\pi) \widehat{\Phi}_{m+k}(\pi) [\pi(\mathcal{R})]^{\pm \frac{2a}{\nu}} \right| \right].$$

Since $\min\{2^{2ja}, 2^{-2ja}\} = 2^{-2|j|a}$, we have deduced the estimate

$$\mathcal{A}(\pi) \lesssim 2^{-2|j|a} \max_{\pm} \text{Tr} \left[\left| \left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] \widehat{\Phi}_{j+k}(\pi) \widehat{\Phi}_{m+k}(\pi) [\pi(\mathcal{R})]^{\pm \frac{2a}{\nu}} \right| \right]. \quad (3.31)$$

Now in order to estimate (3.31) we can use the functional calculus for Rockland operators. Indeed, since $m \asymp j$ note that

$$\begin{aligned} & \text{Tr} \left[\left| \left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] \widehat{\Phi}_{j+k}(\pi) \widehat{\Phi}_{m+k}(\pi) \pi(\mathcal{R})^{\frac{\pm 2a}{\nu}} \right| \right] \\ &= \text{Tr} \left[\left| \left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] \left(\int_0^{\infty} \Phi(2^{(j+k)\nu} \lambda) \Phi(2^{(m+k)\nu} \lambda) dE_{\pi(\mathcal{R})}(\lambda) \right) \pi(\mathcal{R})^{\frac{\pm 2a}{\nu}} \right| \right] \\ &= \text{Tr} \left[\left| \left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] \left(\int_0^{\infty} \Phi(2^{(j+k)\nu} \lambda) \Phi(2^{(m+k)\nu} \lambda) \lambda^{\frac{\pm 2a}{\nu}} dE_{\pi(\mathcal{R})}(\lambda) \right) \right| \right] \\ &\asymp \text{Tr} \left[\left| \left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] \left(\int_{\lambda \sim 2^{-(j+k)\nu}} \Phi(2^{(j+k)\nu} \lambda) \Phi(2^{(m+k)\nu} \lambda) \lambda^{\frac{\pm 2a}{\nu}} dE_{\pi(\mathcal{R})}(\lambda) \right) \right| \right], \end{aligned}$$

where we have used the notation $\lambda \sim 2^{-(j+k)\nu}$ to indicate that $\lambda \in [2^{-(j+k+1)\nu}, 2^{-(j+k-1)\nu}]$. Using the properties of the spectral projections $E_{\pi(\mathcal{R})}$ we have that

$$\begin{aligned} & \text{Tr} \left[\left| \left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] \left(\int_{\lambda \sim 2^{-(j+k)\nu}} \Phi(2^{(j+k)\nu} \lambda) \Phi(2^{(m+k)\nu} \lambda) dE_{\pi(\mathcal{R})}(\lambda) \right) \right| \right] \\ &= \text{Tr} \left[\left| \left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] E_{\pi(\mathcal{R})}[2^{-(j+k+1)\nu}, 2^{-(j+k-1)\nu}] \left(\int_{\lambda \sim 2^{-(j+k)\nu}} \Phi(2^{(j+k)\nu} \lambda) \Phi(2^{(m+k)\nu} \lambda) dE_{\pi(\mathcal{R})}(\lambda) \right) \right| \right] \\ &\leq \text{Tr} \left[\left| \left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] E_{\pi(\mathcal{R})}[2^{-(j+k+1)\nu}, 2^{-(j+k-1)\nu}] \right| \right] \\ &\quad \times \left\| \left(\int_{\lambda \sim 2^{-(j+k)\nu}} \Phi(2^{(j+k)\nu} \lambda) \Phi(2^{(m+k)\nu} \lambda) dE_{\pi(\mathcal{R})}(\lambda) \right) \right\|_{\text{op}}. \end{aligned}$$

Let us estimate the last operator norm:

$$\begin{aligned} & \left\| \left(\int_{\lambda \sim 2^{-(j+k)\nu}} \Phi(2^{(j+k)\nu} \lambda) \Phi(2^{(m+k)\nu} \lambda) \lambda^{\frac{\pm 2a}{\nu}} dE_{\pi(\mathcal{R})}(\lambda) \right) \right\|_{\text{op}} \\ &\asymp 2^{\mp 2(j+k)a} \left\| \left(\int_{\lambda \sim 2^{-(j+k)\nu}} \Phi(2^{(j+k)\nu} \lambda) \Phi(2^{(m+k)\nu} \lambda) dE_{\pi(\mathcal{R})}(\lambda) \right) \right\|_{\text{op}} \\ &\leq 2^{\mp 2(j+k)a} \left\| \left(\int_{\lambda \geq 0} \Phi(2^{(j+k)\nu} \lambda) \Phi(2^{(m+k)\nu} \lambda) dE_{\pi(\mathcal{R})}(\lambda) \right) \right\|_{\text{op}} \\ &= 2^{\mp 2(j+k)a} \left\| \Phi(2^{(j+k)\nu} \pi(\mathcal{R})) \Phi(2^{(m+k)\nu} \pi(\mathcal{R})) \right\|_{\text{op}} \end{aligned}$$

$$\begin{aligned} &\leq 2^{\mp 2(j+k)a} \sup_{\lambda \geq 0} (\Phi(2^{(j+k)\nu} \lambda)) \Phi(2^{(m+k)\nu} \lambda) \\ &\leq 2^{\mp 2(j+k)a} \|\Phi\|_{L^\infty}^2. \end{aligned}$$

So, we have proved the inequality

$$\begin{aligned} &\mathrm{Tr} \left[\left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] \left(\int_{\lambda \sim 2^{-(j+k)\nu}} \Phi(2^{(j+k)\nu} \lambda) \Phi(2^{(m+k)\nu} \lambda) dE_{\pi(\mathcal{R})}(\lambda) \right) \right] \\ &\leq 2^{\mp 2(j+k)a} \|\Phi\|_{L^\infty}^2 \mathrm{Tr} \left[\left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] E_{\pi(\mathcal{R})} [2^{-(j+k+1)\nu}, 2^{-(j+k-1)\nu}] \right], \end{aligned}$$

from where we deduce the estimate

$$\begin{aligned} &\max_{\pm} \mathrm{Tr} \left[\left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] \left(\int_{\lambda \sim 2^{-(j+k)\nu}} \Phi(2^{(j+k)\nu} \lambda) \Phi(2^{(m+k)\nu} \lambda) dE_{\pi(\mathcal{R})}(\lambda) \right) \right] \\ &\lesssim_{\Phi} 2^{-2|j+k|a} \mathrm{Tr} \left[\left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] E_{\pi(\mathcal{R})} [2^{-(j+k+1)\nu}, 2^{-(j+k-1)\nu}] \right]. \end{aligned}$$

The previous analysis allows us to estimate (3.31) as follows

$$\begin{aligned} \mathcal{A}(\pi) &\lesssim 2^{-2|j|a} \max_{\pm} \mathrm{Tr} \left[\left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] \left(\int_{\lambda \sim 2^{-(j+k)\nu}} \Phi(2^{(j+k)\nu} \lambda) \Phi(2^{(m+k)\nu} \lambda) dE_{\pi(\mathcal{R})}(\lambda) \right) \right] \\ &\lesssim_{\Phi, K, a} 2^{-2|j|a} \times 2^{-2|j+k|a} \mathrm{Tr} \left[\left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] E_{\pi(\mathcal{R})} [2^{-(j+k+1)\nu}, 2^{-(j+k-1)\nu}] \right]. \end{aligned}$$

Using the triangle inequality $|j+k| \geq |k| - |j|$ we have the reverse inequality

$$-|j+k| \leq |j| - |k|,$$

and then

$$\begin{aligned} \mathcal{A}(\pi) &\lesssim_{\Phi, K, a} 2^{-2|j|a} \times 2^{-2|k|a+2|j|a} \mathrm{Tr} \left[\left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] E_{\pi(\mathcal{R})} [2^{-(j+k+1)\nu}, 2^{-(j+k-1)\nu}] \right] \\ &= 2^{-2|k|a} \mathrm{Tr} \left[\left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] E_{\pi(\mathcal{R})} [2^{-(j+k+1)\nu}, 2^{-(j+k-1)\nu}] \right]. \end{aligned}$$

In consequence, coming back to (3.22) we have that

$$\begin{aligned} \|\tilde{T}_k f\|_{L^2(G)}^2 &\leq \sum_{j=-\infty}^{\infty} \sum_{m=j-1}^{j+1} \int_{\widehat{G}} \|\widehat{\Phi}_{j+k}(\pi) \widehat{\Phi}_{m+k}(\pi) [(2^j \cdot \pi)(\mathcal{R})^{\pm \frac{a}{\nu}}] \widehat{K}_m(\pi)\|_{\mathrm{op}} \|\widehat{f}(\pi)\|_{\mathrm{HS}}^2 d\pi \\ &\lesssim \sum_{j=-\infty}^{\infty} \sum_{m=j-1}^{j+1} 2^{-2|k|a} \int_{\widehat{G}} \mathrm{Tr} \left[\left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] E_{\pi(\mathcal{R})} [2^{-(j+k+1)\nu}, 2^{-(j+k-1)\nu}] \right] d\pi \\ &\lesssim \sum_{j=-\infty}^{\infty} 2^{-2|k|a} \int_{\widehat{G}} \mathrm{Tr} \left[\left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] E_{\pi(\mathcal{R})} [2^{-(j+k+1)\nu}, 2^{-(j+k-1)\nu}] \right] d\pi. \end{aligned}$$

Using the fact that the mapping $t_j := j + \cdot : \mathbb{Z} \rightarrow \mathbb{Z}$, $k \mapsto j+k$, is a bijection on \mathbb{Z} , we have the estimates

$$\begin{aligned} \|\tilde{T}_k f\|_{L^2(G)}^2 &\lesssim \sum_{j=-\infty}^{\infty} 2^{-2|k|a} \int_{\widehat{G}} \mathrm{Tr} \left[\left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] E_{\pi(\mathcal{R})} [2^{-(j+1)\nu}, 2^{-(j-1)\nu}] \right] d\pi \\ &\asymp 2^{-2|k|a} \int_{\widehat{G}} \mathrm{Tr} \left[\left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] E_{\pi(\mathcal{R})}(-\infty, \infty) \right] d\pi \\ &= 2^{-2|k|a} \int_{\widehat{G}} \mathrm{Tr} \left[\left[\widehat{f}(\pi) \widehat{f}(\pi)^* \right] I_{H_\pi} \right] d\pi \end{aligned}$$

$$= 2^{-2|k|a} \|\widehat{f}\|_{L^2(\widehat{G})}^2.$$

Then, using the last inequality and the Plancherel theorem we deduce that the L^2 -norm of $\widetilde{T}_k f$ satisfies the estimate

$$\|\widetilde{T}_k f\|_{L^2(G)}^2 \lesssim_{\Phi, K, a} 2^{-2|k|a} \|f\|_{L^2(G)}^2,$$

which implies

$$\|\widetilde{T}_k f\|_{L^2(G)} \lesssim_{\Phi, K, a} 2^{-|k|a} \|f\|_{L^2(G)}. \quad (3.32)$$

The proof of (3.14) is complete and we have concluded Step 1 of the proof.

3.1.2. *Step 2.* For the proof of the *weak (1,1) estimate* we will use the fundamental theorem for singular integrals due to Coifman and Weiss [9]. We write

$$\forall f \in C_0^\infty(G), \widetilde{T}_k f := f * (\widetilde{T}_k \delta), (\widetilde{T}_k \delta) := \sum_{j=-\infty}^{\infty} K_j * \Phi_{j+k}.$$

We shall prove the estimate

$$[\widetilde{T}_k \delta]_H := \sup_{|y| \leq 1} \int_{|x| > 2|y|} |\widetilde{T}_k \delta(y^{-1}x) - \widetilde{T}_k \delta(x)| dx \lesssim_K (1 + |k|), \quad (3.33)$$

and then the constant $(1 + |k|)$ of the right-hand side of (3.33) gives the estimate in (3.15). Indeed, note that the estimate in (3.33) says that the kernel $\widetilde{T}_k \delta$ of \widetilde{T}_k satisfies the Hörmander condition $[\widetilde{T}_k \delta]_H \lesssim_K (1 + |k|)$. Then, from the fundamental theorem for singular integrals due to Coifman and Weiss [9], one has that the $(L^1, L^{1,\infty})$ -operator norm of T satisfies the estimate

$$\|\widetilde{T}_k\|_{L^1 \rightarrow L^{1,\infty}} \lesssim \|\widetilde{T}_k\|_{L^2 \rightarrow L^2} + [\widetilde{T}_k \delta]_H \lesssim 2^{-|k|a} + (1 + |k|) \lesssim (1 + |k|), \quad (3.34)$$

which proves (3.15).

Note that for any $y \in G$ with $|y| \leq 1$, we have

$$\begin{aligned} \int_{|x| > 2|y|} |\widetilde{T}_k \delta(y^{-1}x) - \widetilde{T}_k \delta(x)| dx &= \int_{|x| > 2|y|} \left| \sum_{j=-\infty}^{\infty} (K_j * \Phi_{j+k})(y^{-1}x) - (K_j * \Phi_{j+k})(x) \right| dx \\ &\leq \sum_{j=-\infty}^{\infty} \int_{|x| > 2|y|} |(K_j * \Phi_{j+k})(y^{-1}x) - (K_j * \Phi_{j+k})(x)| dx \\ &= \sum_{j=-\infty}^{\infty} I_{j,k}, \end{aligned}$$

where

$$I_{j,k} = \int_{|x| > 2|y|} |(K_j * \Phi_{j+k})(y^{-1}x) - (K_j * \Phi_{j+k})(x)| dx.$$

Note that, by the Hausdorff-Young inequality, we have the following immediate estimate

$$\begin{aligned} I_{j,k} &\leq \int_G |(K_j * \Phi_{j+k})(y^{-1}x) - (K_j * \Phi_{j+k})(x)| dx \leq 2 \int_G |(K_j * \Phi_{j+k})(z)| dz \\ &\leq \|K_j\|_{L^1(G)} \|\Phi_{j+k}\|_{L^1(G)} = \|K\|_{L^1(G)} \|\Phi(\mathcal{R})\delta\|_{L^1(G)}. \end{aligned}$$

Indeed, since

$$K_j := 2^{-jQ}K(2^{-j}\cdot) \text{ and } \Phi_{j+k} := 2^{-(j+k)Q}(\Phi(\mathcal{R})\delta)(2^{-(j+k)}\cdot),$$

the changes of variables $z = 2^{-j}\cdot x$ and $z' = 2^{-(j+k)}\cdot x'$ imply the equality

$$\|K_j\|_{L^1(G)} = \int_G |K_j(x)|dx = 2^{-jQ} \int_G |K(2^{-j}\cdot x)|dx = \int_G |K(z)|dz = \|K\|_{L^1(G)},$$

as well as that

$$\begin{aligned} \|\Phi_{j+k}\|_{L^1(G)} &= 2^{-(j+k)Q} \int_G |(\Phi(\mathcal{R})\delta)(2^{-(j+k)}\cdot x')|dx' = \int_G |(\Phi(\mathcal{R})\delta)(z')|dz' \\ &= \|(\Phi(\mathcal{R})\delta)\|_{L^1(G)}. \end{aligned}$$

Then, we have proved that

$$\forall 0 < |y| \leq 1, \forall (j, k), \quad I_{j,k} \lesssim_{\Phi} \|K\|_{L^1(G)} \quad (3.35)$$

Also, we will provide other estimates for $I_{j,k}$ as follows. First, we use the dilation property

$$\forall r > 0, \forall x, y \in G, \quad r \cdot (xy) = (r \cdot x)(r \cdot y), \text{ and } r \cdot x^{-1} = (r \cdot x)^{-1}. \quad (3.36)$$

Then

$$\begin{aligned} I_{j,k} &= \int_{|x|>2|y|} |(K_j * \Phi_{j+k})(y^{-1}x) - (K_j * \Phi_{j+k})(x)| dx \\ &= \int_{|x|>2|y|} \left| \int_G K_j(y^{-1}xz^{-1})\Phi_{j+k}(z)dz - \int_G K_j(xz^{-1})\Phi_{j+k}(z)dz \right| dx \\ &= \int_{|x|>2|y|} 2^{-jQ} \left| \int_G K(2^{-j}\cdot(y^{-1}xz^{-1}))2^{-(j+k)Q}(\Phi(\mathcal{R})\delta)(2^{-j-k}z)dz \right. \\ &\quad \left. - \int_G K(2^{-j}\cdot(xz^{-1}))2^{-(j+k)Q}(\Phi(\mathcal{R})\delta)(2^{-j-k}\cdot z)dz \right| dx \\ &= \int_{|x|>2|y|} 2^{-jQ} \left| \int_G K((2^{-j}\cdot y^{-1})(2^{-j}\cdot x)(2^{-j}\cdot z^{-1}))2^{-(j+k)Q}(\Phi(\mathcal{R})\delta)(2^{-j-k}z)dz \right. \\ &\quad \left. - \int_G K((2^{-j}\cdot x)(2^{-j}\cdot z^{-1}))2^{-(j+k)Q}(\Phi(\mathcal{R})\delta)(2^{-j-k}\cdot z)dz \right| dx \\ &= \int_{|x|>2|y|} 2^{-jQ} \left| \int_G K((2^{-j}\cdot y^{-1})(2^{-j}\cdot x)(2^{-j}\cdot z)^{-1})2^{-(j+k)Q}(\Phi(\mathcal{R})\delta)(2^{-j-k}z)dz \right. \\ &\quad \left. - \int_G K((2^{-j}\cdot x)(2^{-j}\cdot z)^{-1})2^{-(j+k)Q}(\Phi(\mathcal{R})\delta)(2^{-j-k}\cdot z)dz \right| dx. \end{aligned}$$

The change of variables $w := 2^{-j}\cdot z$ gives the new volume element $dw = 2^{-jQ}dz$, and we have that

$$\begin{aligned} I_{j,k} &= \int_{|x|>2|y|} 2^{-jQ} \left| \int_G K((2^{-j}\cdot y^{-1})(2^{-j}\cdot x)(2^{-j}\cdot z)^{-1})2^{-(j+k)Q}(\Phi(\mathcal{R})\delta)(2^{-j-k}z)dz \right. \\ &\quad \left. - \int_G K((2^{-j}\cdot x)(2^{-j}\cdot z)^{-1})2^{-(j+k)Q}(\Phi(\mathcal{R})\delta)(2^{-j-k}\cdot z)dz \right| dx \\ &= \int_{|x|>2|y|} \left| \int_G K((2^{-j}\cdot y)^{-1}(2^{-j}\cdot x)w^{-1})2^{-kQ}2^{-jQ}(\Phi(\mathcal{R})\delta)(2^{-k}w)dw \right. \\ &\quad \left. - \int_G K((2^{-j}\cdot x)w^{-1})2^{-kQ}2^{-jQ}(\Phi(\mathcal{R})\delta)(2^{-k}\cdot w)dw \right| dx \end{aligned}$$

$$\begin{aligned}
&= \int_{|x|>2|y|} 2^{-jQ} \left| \int_G K((2^{-j} \cdot y)^{-1}(2^{-j} \cdot x)w^{-1})2^{-kQ}(\Phi(\mathcal{R})\delta)(2^{-k}w)dw \right. \\
&\quad \left. - \int_G K((2^{-j} \cdot x)w^{-1})2^{-kQ}(\Phi(\mathcal{R})\delta)(2^{-k} \cdot w)dw \right| dx.
\end{aligned}$$

On the other hand, the change of variables $x' = 2^{-j} \cdot x$ gives the new volume element $dx' = 2^{-jQ}dx$, the zone $\{x \in G : |x| > 2|y|\}$ is mapped to the set

$$\{x' \in G : |x'| > 2^{-j+1}|y|\}$$

allowing us to write the following identities

$$\begin{aligned}
&I_{j,k} \\
&= \int_{|x'|>2^{1-j}|y|} 2^{-jQ} \left| \int_G K((2^{-j} \cdot y)^{-1}x'w^{-1})2^{-kQ}(\Phi(\mathcal{R})\delta)(2^{-k}w)dw \right. \\
&\quad \left. - \int_G K(x'w^{-1})2^{-kQ}(\Phi(\mathcal{R})\delta)(2^{-k} \cdot w)dw \right| 2^{jQ} dx' \\
&= \int_{|x|>2^{1-j}|y|} \left| \int_G K((2^{-j} \cdot y)^{-1}xw^{-1})\Phi_k(w)dw - \int_G K(xw^{-1})\Phi_k(w)dw \right| dx \\
&= \int_{|x|>2^{1-j}|y|} |K * \Phi_k((2^{-j} \cdot y)^{-1}x) - K * \Phi_k(x)| dx \\
&= \int_{|x|>2^{1-j}|y|} \left| \int_G K(z)[\Phi_k(z^{-1}(2^{-j} \cdot y)^{-1}x) - \Phi_k(z^{-1}x)] dz \right| dx \\
&= \int_{|x|>2^{1-j}|y|} 2^{-kQ} \left| \int_G K(z)[\Phi((2^{-k} \cdot z)^{-1}(2^{-j-k} \cdot y)^{-1}(2^{-k} \cdot x)) - \Phi((2^{-k} \cdot z)^{-1}(2^{-k} \cdot x))] dz \right| dx \\
&\leq \int_{|x|>2^{1-j}|y|} 2^{-kQ} \int_G |K(z)| \left| [\Phi((2^{-k} \cdot z)^{-1}(2^{-j-k} \cdot y)^{-1}(2^{-k} \cdot x)) - \Phi((2^{-k} \cdot z)^{-1}(2^{-k} \cdot x))] \right| dz dx,
\end{aligned}$$

where we have used the dilation property in (3.36) with $r = 2^{-k}$.

The change of variables $x' = 2^{-k} \cdot x$ and the new volume element $dx' = 2^{-kQ}dx$ implies that

$$\begin{aligned}
&I_{j,k} \\
&\leq \int_{|x'|>2^{1-j-k}|y|} \int_G |K(z)| \left| [\Phi((2^{-k} \cdot z)^{-1}(2^{-j-k} \cdot y)^{-1}(x')) - \Phi((2^{-k} \cdot z)^{-1}(x'))] \right| dz dx'.
\end{aligned}$$

Now, let us make the change of variables $z'' = 2^{-k} \cdot z$. We have that the new volume element $2^{kQ}dz'' = dz$. Then,

$$\begin{aligned}
&I_{j,k} \\
&\leq \int_{|x'|>2^{1-j-k}|y|} \int_G 2^{kQ} |K(2^k \cdot z'')| \left| [\Phi((z'')^{-1}(2^{-j-k} \cdot y)^{-1}(x')) - \Phi((z'')^{-1}(x'))] \right| dz'' dx'.
\end{aligned}$$

Now, the mean value theorem (see [19, Page 119]) implies that

$$\begin{aligned}
&\left| [\Phi((z'')^{-1}(2^{-j-k} \cdot y)^{-1}(x')) - \Phi((z'')^{-1}(x'))] \right| \\
&\lesssim \sum_{\ell=1}^n |(2^{-j-k} \cdot y)^{-1}|^{\nu_\ell} \sup_{|z'| \lesssim |(2^{-j-k} \cdot y)^{-1}|} |(X_{z',\ell}(\Phi((z'')^{-1}z'(x'))))|,
\end{aligned}$$

where we have written that $|z'| \lesssim |(2^{-j-k} \cdot y)^{-1}|$ to indicate that the inequality

$$|z'| \leq c|(2^{-j-k} \cdot y)^{-1}| = c2^{-j-k}|y| \quad (3.37)$$

is valid for some universal constant $c > 1$ as in the mean value theorem of [19, Page 119]. Using that $\int_G 2^{kQ} |K(2^k \cdot z'')| dz'' = \|K\|_{L^1(G)}$, we can estimate $I_{j,k}$ as follows

$$\begin{aligned}
& I_{j,k} \\
& \leq 2^{kQ} \int_G |K(2^{kQ} \cdot z'')| dz'' \\
& \times \sup_{z \in G} \int_{|x'| > 2^{1-j-k}|y|} \left(\sum_{\ell=1}^n 2^{-(j+k)\nu_\ell} |y|^{\nu_\ell} \sup_{|z'| \lesssim 2^{-j-k}|y|} |X_{z,\ell}(\Phi(z^{-1}z'x'))| \right) dx' \\
& = \|K\|_{L^1(G)} \\
& \times \sum_{\ell=1}^n 2^{-(j+k)\nu_\ell} |y|^{\nu_\ell} \sup_{z \in G} \int_{|x'| > 2^{1-j-k}|y|} \sup_{|z'| \lesssim 2^{-j-k}|y|} |X_{z,\ell}(\Phi(z^{-1}z'x'))| dx'.
\end{aligned}$$

Using the Sobolev embedding theorem on G (see Theorem 4.4.25 of [19, page 241]), we can estimate for $M_0 > Q/2$,

$$\begin{aligned}
& \sup_{z \in G} \int_{|x'| > 2^{1-j-k}|y|} \sup_{|z'| \lesssim 2^{-j-k}|y|} |X_{z,\ell}(\Phi(z^{-1}z'x'))| dx' \\
& \lesssim \sup_{z \in G} \int \sup_{z' \in G} |X_{z,\ell}(\Phi(z^{-1}z'x'))| dx' \lesssim \sup_{z \in G} \sum_{|\beta| \leq M_0} \int_G \|X_{z'}^\beta X_{z,\ell}(\Phi(z^{-1}z'x'))\|_{L^2(G, dz')} dx' \\
& \lesssim \sup_{z \in G} \sum_{|\beta| \leq M_0} \int_G \|X_{z'}^\beta X_{z,\ell}(\Phi(z^{-1}z'x'))\|_{L^2(G, dz')} dx' \\
& \lesssim \sup_{z \in G} \sum_{|\beta| \leq M_0} \left(\int_G (1 + |x'|)^{2M_0} \|X_{z'}^\beta X_{z,\ell}(\Phi(z^{-1}z'x'))\|_{L^2(G, dz')}^2 dx' \right)^{\frac{1}{2}} \left(\int_G (1 + |x'|)^{-2M_0} dx' \right)^{\frac{1}{2}} \\
& \lesssim \sup_{z \in G} \sum_{|\beta| \leq M_0} \left(\int_G \int_G (1 + |x'|)^{2M_0} |X_{z'}^\beta X_{z,\ell}(\Phi(z^{-1}z'x'))|^2 dx' dz' \right)^{\frac{1}{2}} < \infty,
\end{aligned}$$

where the convergence of the last integral is justified by Hulanicki theorem, see [19, page 251]. All the analysis above shows that for all $j, k \in \mathbb{Z}$,

$$\forall y \in G : 0 < |y| \leq 1, \quad I_{j,k} \lesssim \sum_{\ell=1}^n 2^{-(j+k)\nu_\ell} |y|^{\nu_\ell} \|K\|_{L^1(G)}. \quad (3.38)$$

In terms of $R > 0$ defined by

$$R = \inf\{R' > 0 : \text{supp}(K) \subset B(e, R')\}, \quad (3.39)$$

and of $0 < |y| \leq 1$, let us make a more precise estimate of $I_{j,k}$ in the case where $2^{-j}|y| \geq 2R$. This will be used in further analysis. To do this, let us come back to the estimate

$$\begin{aligned}
I_{j,k} & \lesssim \sum_{\ell=1}^n 2^{-(j+k)\nu_\ell} |y|^{\nu_\ell} \int_{|x'| > 2^{1-j-k}|y|} \int_G |K(z)| \sup_{|z'| \lesssim 2^{-j-k}|y|} |X_{z,\ell}(\Phi((2^{-k} \cdot z)^{-1}z'x'))| dx' dz \\
& = \sum_{\ell=1}^n 2^{-(j+k)\nu_\ell} |y|^{\nu_\ell} \mathcal{J}_{j,k,\ell},
\end{aligned}$$

where

$$\mathcal{I}_{j,k,\ell} = \int_{|x'| > 2^{1-j-k}|y|} \int_G |K(z)| \sup_{|z'| \lesssim 2^{-j-k}|y|} |X_{z,\ell}(\Phi((2^{-k} \cdot z)^{-1} z' x'))| dx' dz.$$

To estimate this double integral let us split it as follows

$$\begin{aligned} \mathcal{I}_{j,k,\ell} &= \int_{|x'| > 2^{1-j-k}|y|} \int_G |K(z)| \sup_{|z'| \lesssim 2^{-j-k}|y|} |X_{z,\ell}(\Phi((2^{-k} \cdot z)^{-1} z' x'))| dx' dz \\ &= \int_{|x'| > 2^{1-j-k}|y|} \int_{\{z: |x'| \geq 2^{-k+2}|z|\}} |K(z)| \sup_{|z'| \lesssim 2^{-j-k}|y|} |X_{z,\ell}(\Phi((2^{-k} \cdot z)^{-1} z' x'))| dx' dz \\ &\quad + \int_{|x'| > 2^{1-j-k}|y|} \int_{\{z: |x'| < 2^{-k+2}|z|\}} |K(z)| \sup_{|z'| \lesssim 2^{-j-k}|y|} |X_{z,\ell}(\Phi((2^{-k} \cdot z)^{-1} z' x'))| dx' dz \\ &= \mathcal{I}_{j,k,\ell}^I + \mathcal{I}_{j,k,\ell}^{II}. \end{aligned}$$

Observe that for the integral

$$\mathcal{I}_{j,k,\ell}^I = \int_{|x'| > 2^{1-j-k}|y|} \int_{\{z: |x'| \geq 2^{-k+2}|z|\}} |K(z)| \sup_{|z'| \lesssim 2^{-j-k}|y|} |X_{z,\ell}(\Phi((2^{-k} \cdot z)^{-1} z' x'))| dx' dz,$$

the integral with respect to dz is computed on the zone $\{z : |x'| \geq 2^{-k+2}|z|\}$, where one has $-|x'|/4 \leq -2^{-k}|z|$. On the other hand for the integral with respect to dx' , on the region $\{x' : |x'| > 2^{1-j-k}|y|\}$, one has that $|x'|/2 > 2^{-j-k}|y|$. In consequence, for some constant $0 < C < c$, where c is the universal constant in (3.37), and then independent of j and k , one has that

$$\begin{aligned} |(2^{-k} \cdot z)^{-1} z' x'| &\geq |x'| - |z'| - 2^{-k}|z| \geq C(|x'| - 2^{-j-k}|y| - \frac{1}{4}|x'|) \\ &= C(|x'|/2 - 2^{-j-k}|y| + |x'|/2 - \frac{1}{4}|x'|) \\ &> C(|x'|/2 - \frac{1}{4}|x'|) \\ &= C \frac{|x'|}{4}. \end{aligned}$$

Then, estimating again $|X_{z,\ell}(\Phi((2^{-k} \cdot z)^{-1} z' x'))| \leq C_L(1 + |((2^{-k} \cdot z)^{-1} z' x')|)^{-L}$, with L to be determined later, we have

$$\begin{aligned} &\int_{|x'| > 2^{1-j-k}|y|} \int_{\{z: |x'| \geq 2^{-k+2}|z|\}} |K(z)| \sup_{|z'| \lesssim 2^{-j-k}|y|} |X_{z,\ell}(\Phi((2^{-k} \cdot z)^{-1} z' x'))| dx' dz \\ &\leq \int_{|x'| > 2^{1-j-k}|y|} \int_{\{z: |x'| \geq 2^{-k+2}|z|\}} |K(z)| \sup_{|z'| \lesssim 2^{-j-k}|y|} C_L(1 + |((2^{-k} \cdot z)^{-1} z' x')|)^{-L} dx' dz \\ &\leq \int_{|x'| > 2^{1-j-k}|y|} \int_G |K(z)| dz C_L(1 + \frac{1}{4}|x'|)^{-L} dx'. \end{aligned}$$

Then, with $L = Q + 1 + 2\nu_\ell$, we have

$$\begin{aligned} &\int_{|x'| > 2^{1-j-k}|y|} \int_G |K(z)| dz C_L(1 + \frac{1}{4}|x'|)^{-L} dx' \\ &= \|K\|_{L^1(G)} \int_{|x'| > 2^{1-j-k}|y|} C_L(1 + \frac{1}{4}|x'|)^{-Q-1} (1 + |x'|)^{-2\nu_\ell} dx' \end{aligned}$$

$$\begin{aligned} &\lesssim \|K\|_{L^1(G)} (2^{1-j-k}|y|)^{-2\nu_\ell} \int_{|x'| > 2^{1-j-k}|y|} C_L (1 + \frac{1}{4}|x'|)^{-Q-1} dx' \\ &\leq \|K\|_{L^1(G)} (2^{1-j-k}|y|)^{-2\nu_\ell} \int_G C_L (1 + \frac{1}{4}|x'|)^{-Q-1} dx'. \end{aligned}$$

In consequence we have that

$$\mathcal{J}_{j,k,\ell}^I \lesssim \|K\|_{L^1(G)} (2^{1-j-k}|y|)^{-2\nu_\ell}.$$

Now, note that when $2^{-j}|y| \geq 2R$, and for $|x'| < 2^{-k+2}|z|$, (and then $|x'| < 2^{-k+2}R$, since z belongs to the support of K) we have that

$$\forall 0 < |y| \leq 1, \quad 2^{-j}|y| \geq 2R, \quad \{(x', z) : |x'| > 2^{1-j-k}|y|\} \cap \{(x', z) : |x'| < 2^{-k+2}|z|\} = \emptyset. \quad (3.40)$$

This implies that when $0 < |y| \leq 1$, $2^{-j}|y| \geq 2R$,

$$\mathcal{J}_{j,k,\ell}^{II} = \int_{|x'| > 2^{1-j-k}|y|} \int_{\{z: |x| < 2^{-k+2}|z|\}} |K(z)| \sup_{|z'| \lesssim 2^{-j-k}|y|} |X_{z,\ell}(\Phi((2^{-k} \cdot z)^{-1} z' x'))| dx' dz = 0. \quad (3.41)$$

In summarising we have proved the following estimates

- $\forall 0 < |y| \leq 1, \forall j, k \in \mathbb{Z}$

$$I_{j,k} \lesssim \sum_{\ell=1}^n 2^{-(j+k)\nu_\ell} |y|^{\nu_\ell} \|K\|_{L^1(G)}. \quad (3.42)$$

Moreover, in view of (3.35) we have that $\forall 0 < |y| \leq 1, \forall j, k \in \mathbb{Z}$

$$I_{j,k} \lesssim \sum_{\ell=1}^n \min\{1, 2^{-(j+k)\nu_\ell} |y|^{\nu_\ell}\} \|K\|_{L^1(G)}. \quad (3.43)$$

- $\forall 0 < |y| \leq 1, 2^{-j}|y| \geq 2R,$

$$I_{j,k} \lesssim \sum_{\ell=1}^n 2^{-(j+k)\nu_\ell} |y|^{\nu_\ell} (2^{1-j-k}|y|)^{-2\nu_\ell} \|K\|_{L^1(G)} \lesssim \sum_{\ell=1}^n (2^{-(j+k)\nu_\ell} |y|^{\nu_\ell})^{-1} \|K\|_{L^1(G)}. \quad (3.44)$$

Now, let us use these inequalities to estimate the Hörmander condition

$$\sup_{|y| \leq 1} \int_{|x| > 2|y|} |\tilde{T}_k \delta(y^{-1}x) - \tilde{T}_k \delta(x)| dx \leq \sum_{j \in \mathbb{Z}} I_{j,k}. \quad (3.45)$$

Indeed, we consider the cases where $2^{-k} > (2R)^{-1}$ and where $2^{-k} \leq (2R)^{-1}$.

- Case (i): $2^{-k} > (2R)^{-1}$. Observe that when $\frac{1}{2^{-k}|y|} < 2^{-j} \leq 2R/|y|$, we have that $1 \leq 2^{-j-k}|y|$. In this situation, the fact that $\min\{1, 2^{-j-k}|y|\} = 1$, and the estimate in (3.43) imply that

$$\begin{aligned} \sum_j I_{j,k} &\lesssim \sum_{\ell=1}^n \|K\|_{L^1(G)} \left(\sum_{2^{-j} \leq \frac{1}{2^{-k}|y|}} 2^{-(j+k)\nu_\ell} |y|^{\nu_\ell} + \sum_{\frac{1}{2^{-k}|y|} < 2^{-j} \leq \frac{2R}{|y|}} 1 + \sum_{2^{-j} \geq \frac{2R}{|y|}} (2^{-(j+k)\nu_\ell} |y|^{\nu_\ell})^{-1} \right) \\ &\lesssim \sum_{\ell=1}^n \|K\|_{L^1(G)} (\log(R) + |k|) \lesssim \|K\|_{L^1(G)} (\log(R) + |k|) \lesssim_{n,K} \|K\|_{L^1(G)} (1 + |k|). \end{aligned}$$

Indeed, observe that the sums $\sum_{2^{-j} \leq \frac{1}{2^{-k}|y|}} 2^{-(j+k)\nu_\ell} |y|^{\nu_\ell}$ and $\sum_{2^{-j} \geq \frac{2R}{|y|}} (2^{-(j+k)\nu_\ell} |y|^{\nu_\ell})^{-1}$ can be handled as geometric sums in order to get

$$\sum_{2^{-j} \leq \frac{1}{2^{-k}|y|}} 2^{-(j+k)\nu_\ell} |y|^{\nu_\ell} = 2^{-k\nu_\ell} |y|^{\nu_\ell} \sum_{2^{-j} \leq \frac{1}{2^{-k}|y|}} 2^{-j\nu_\ell} \asymp 2^{-k\nu_\ell} |y|^{\nu_\ell} (2^{-k\nu_\ell} |y|^{\nu_\ell})^{-1} = 1,$$

and

$$\begin{aligned} \sum_{2^{-j} \geq \frac{2R}{|y|}} (2^{-(j+k)\nu_\ell} |y|^{\nu_\ell})^{-1} &= (2^{-k\nu_\ell} |y|^{\nu_\ell})^{-1} \sum_{2^j \leq \frac{|y|}{2R}} 2^{j\nu_\ell} \leq (2^{-k\nu_\ell} |y|^{\nu_\ell})^{-1} \sum_{2^j \leq \frac{|y|}{2R}} 2^{j\nu_\ell} \\ &\lesssim (2^{-k\nu_\ell} |y|^{\nu_\ell})^{-1} 2^{-k\nu_\ell} |y|^{\nu_\ell} = 1. \end{aligned}$$

As for the sum $\sum_{\frac{1}{2^{-k}|y|} < 2^{-j} \leq \frac{2R}{|y|}} 1$, let $j_0 \in \mathbb{Z}$ and $\ell_0 \in \mathbb{N}$, be such that

$$2^{-j_0} \leq \frac{1}{2^{-k}|y|} \leq 2^{-j_0+1}, \quad 2^{-j_0+\ell_0} \leq \frac{2R}{|y|} \leq 2^{-j_0+\ell_0+1}.$$

Then

$$\sum_{\frac{1}{2^{-k}|y|} < 2^{-j} \leq \frac{2R}{|y|}} 1 \lesssim |\{j \in \{0, 1, \dots, \ell_0\} : 2^{-j_0+j} \in [2^{-j_0}, 2^{-j_0+\ell_0}]\}| = \ell_0 + 1 \lesssim \ell_0.$$

Now, to estimate ℓ_0 in terms of R and of $|k|$, note that $2^{-j_0} \asymp \frac{1}{2^{-k}|y|}$, and $2^{-j_0+\ell_0} \asymp \frac{2R}{|y|}$. We have that

$$2^{-j_0-k} \asymp \frac{1}{|y|} \asymp \frac{2^{-j_0+\ell_0}}{2R}.$$

From this estimate we deduce that $2^{\ell_0} \asymp 2^{-k} \times 2R \lesssim 2^{|k|+\log_2(R)}$, from where we have that $\ell_0 \lesssim \log(R) + |k| \lesssim_R (1 + |k|)$.

- Case (ii): $2^{-k} \leq (2R)^{-1}$. Again we divide the sum, but this time in two terms as follows

$$\begin{aligned} \sum_j I_{j,k} &\lesssim \sum_{\ell=1}^n \|K\|_{L^1(G)} \left(\sum_{2^{-j} \leq \frac{1}{2^{-k}|y|}} 2^{-(j+k)\nu_\ell} |y|^{\nu_\ell} + \sum_{2^{-j} > \frac{1}{2^{-k}|y|}} (2^{-(j+k)\nu_\ell} |y|^{\nu_\ell})^{-1} \right) \\ &\lesssim \sum_{\ell=1}^n \|K\|_{L^1(G)} = n \|K\|_{L^1(G)}. \end{aligned}$$

All the estimates above prove that

$$\sup_{|y| \leq 1} \int_{|x| > 2|y|} |\tilde{T}_k \delta(y^{-1}x) - \tilde{T}_k \delta(x)| dx \lesssim_{n,K} \|K\|_{L^1(G)} (1 + |k|), \quad (3.46)$$

as desired. So, for any $k \in \mathbb{Z}$ we have proved that $\tilde{T}_k : L^2(G) \rightarrow L^2(G)$ and $\tilde{T}_k : L^1(G) \rightarrow L^{1,\infty}(G)$ are bounded operators with the operator norms satisfying the bounds

$$\|\tilde{T}_k\|_{\mathcal{B}(L^2(G))} \lesssim 2^{-a|k|}, \quad \|\tilde{T}_k\|_{\mathcal{B}(L^1(G), L^{1,\infty}(G))} \lesssim (1 + |k|). \quad (3.47)$$

By using the Marcinkiewicz interpolation theorem, we have for all $1 < p < 2$, the bound

$$\|\tilde{T}_k\|_{\mathcal{B}(L^p(G))} \lesssim 2^{-a|k|\theta_p} (1 + |k|)^{1-|\theta_p|}, \quad (3.48)$$

where $1/p = \theta/2 + (1 - \theta_p)$. Consequently

$$\|T\|_{\mathcal{B}(L^p(G))} \lesssim \sum_{k \in \mathbb{Z}} \|\tilde{T}_k\|_{\mathcal{B}(L^p(G))} \lesssim \sum_{k \in \mathbb{Z}} 2^{-a|k|\theta_p} (1 + |k|)^{1-|\theta_p|} < \infty, \quad (3.49)$$

for all $1 < p < 2$. The boundedness of T on $L^p(G)$ for all $2 < p < \infty$ now follows from the duality argument. \square

3.2. The probabilistic argument. Now we present a lemma about the G -function associated to the family K_j of kernels in Lemma 3.1.

Lemma 3.2. *Let K be a distribution as in Lemma 3.1. Then the square function*

$$(\mathcal{G}(K))f(x) := \left(\sum_{j \in \mathbb{Z}} |f * K_j(x)|^2 \right)^{\frac{1}{2}} \quad (3.50)$$

is a bounded operator from $L^p(G)$ to $L^p(G)$ for all $1 < p < \infty$.

Proof. Let us give the classical probabilistic argument involving the Rademacher functions. So, given a sequence $\varepsilon = \{\varepsilon_j\}_{j \in \mathbb{Z}}$, where $\varepsilon_j = \pm 1$, consider the operator

$$T_\varepsilon f(x) = \sum_{j \in \mathbb{Z}} \varepsilon_j f * K_j. \quad (3.51)$$

In view of Lemma 3.1, T_ε is bounded on $L^p(G)$ for all $1 < p < \infty$, and

$$\|T_\varepsilon\|_{\mathcal{B}(L^p)} \leq C_p, \quad (3.52)$$

where C_p is independent of the choice of the sequence ε . Let us consider the orthonormal system on $L^2[0, 1]$ determined by the Rademacher functions r_j , $j \in \mathbb{Z}$. These are functions defined via $r_j(t) = r_0(2^j t)$, where r_0 is defined via

$$r_0(s) = -1, \quad 0 \leq s < 1/2, \quad r_0(s) = 1, \quad 1/2 \leq s \leq 1. \quad (3.53)$$

The Rademacher functions r_j , $j \in \mathbb{Z}$, satisfy the Khintchine inequality (see e.g. Grafakos [22, Appendix C.1]):

- If $F = \sum_j a_j r_j \in L^2[0, 1]$, then there exist positive constants $A_p, B_p > 0$ such that

$$A_p \|F\|_{L^p[0,1]} \leq \left(\sum_{j \in \mathbb{Z}} |a_j|^2 \right)^{\frac{1}{2}} \leq B_p \|F\|_{L^2[0,1]}. \quad (3.54)$$

Let $x \in G$. If we apply this property with $F(t) = (\mathcal{G}(K))f = \sum_{j \in \mathbb{Z}} f * K_j(x) r_j(t)$, then

$$\left(\sum_{j \in \mathbb{Z}} |f * K_j(x)|^2 \right)^{\frac{p}{2}} \leq B_p^p \int_{[0,1]} \left| \sum_{j \in \mathbb{Z}} f * K_j(x) r_j(t) \right|^p dt.$$

Now, integrating both sides of this inequality with respect to the Haar measure dx , then for any $t \in [0, 1]$, we have an operator like $T_\varepsilon = T_{\{r_j(t)\}}$ and the desired inequality

$$\|(\mathcal{G}(K))f\|_{L^p(G)} \leq C_p \|f\|_{L^p(G)}, \quad (3.55)$$

now follows from (3.52). The proof of Lemma 3.2 is complete. \square

3.3. Proof of the main theorem. Let $1 < p \leq \infty$. In this subsection, we prove the $L^p(G)$ -boundedness of the dyadic maximal function (1.3) associated to an finite Borel measure $d\sigma$ with compact support on G . We assume that for some $a > 0$ the group Fourier transform of $d\sigma$ satisfies the growth estimate in (1.5).

Proof of Theorem 1.1. By the Riesz representation theorem we have that $d\sigma = Kdx$, where K is a distribution with compact support on G that satisfies the group Fourier transform inequalities: $\sup_{\pi \in \widehat{G}} \|\pi(\mathcal{R})^{\pm \frac{a}{v}} \widehat{K}(\pi)\|_{\text{op}} < \infty$. Note that

$$\begin{aligned} \mathcal{M}_D^{d\sigma} f(x) &= \sup_{j \in \mathbb{Z}} \left| \int_G f((2^j \cdot y)^{-1}x) d\sigma(y) \right| = \sup_{j \in \mathbb{Z}} \left| \int_G f((2^j \cdot y)^{-1}x) K(y) dy \right| \\ &= \sup_{j \in \mathbb{Z}} \left| \int_G f(y^{-1}x) K_j(y) dy \right|, \quad \forall y \in G, \quad K_j(y) := 2^{-jQ} K(2^{-j}y). \end{aligned}$$

In view of Lemma 3.2, we have that

$$\mathcal{M}_D^{d\sigma} f(x) = \sup_{j \in \mathbb{Z}} |f * K_j(x)| \leq (\mathcal{G}(K))f(x) := \left(\sum_{j \in \mathbb{Z}} |f * K_j(x)|^2 \right)^{\frac{1}{2}}, \quad (3.56)$$

and in consequence we have proved the boundedness of $\mathcal{M}_D^{d\sigma}$ on $L^p(G)$ for all $1 < p < \infty$. The boundedness of $\mathcal{M}_D^{d\sigma}$ on $L^\infty(G)$ is clearly satisfied and in that case the $L^\infty(G)$ -operator norm of $\mathcal{M}_D^{d\sigma}$ is bounded by the variation of the finite measure $d\sigma$. The proof of Theorem 1.1 is complete. \square

REFERENCES

1. Bagchi, S., Hait, S., Roncal, L. Thangavelu, S. On the maximal function associated to the spherical means on the Heisenberg group, *New York J. Math.* 27, 631–675, (2021).
2. Beltran, D. Guo, S, Hickman, J. Seeger, A. The circular maximal operator on Heisenberg radial functions, to appear in *Ann. Sc. Norm. Super. Pisa Cl. Sci.*
3. Bourgain, J. Averages in the plane over convex curves and maximal operators, *J. Anal. Math.* 47, 69–85, (1986).
4. Bourgain, J. Mirek, M. Stein, E. Wróbel, B. On discrete Hardy-Littlewood maximal functions over the balls in \mathbb{Z}^d : dimension-free estimates. *Geometric aspects of functional analysis. Vol. I*, 127–169, *Lecture Notes in Math.*, 2256, Springer, Cham, 2020.
5. Bourgain, J. Mirek, M. Stein, E. Wróbel, B. On the Hardy-Littlewood maximal functions in high dimensions: continuous and discrete perspective. *Geometric aspects of harmonic analysis*, 107–148, *Springer INdAM Ser.*, 45, Springer, Cham, 2021.
6. Calderón, A. P., Zygmund, A. On the existence of certain singular integrals. *Acta Math.*, 88, 85–139, (1952).
7. Calderón, C. P. Lacunary spherical means, *Ill. J. Math.* 23, 476–484, (1979).
8. Carbery, A. Radial Fourier multipliers and associated maximal functions, *Recent progress in Fourier analysis*, *North-Holland Mathematical Studies*, 111 (North-Holland, Amsterdam, 1985) 49–56.
9. Coifman, R., Weiss, G. *Analyse harmonique non commutative sur certains espaces homogènes.* *Lect. Notes Math.* 242. (1971)
10. Coifman, R., Weiss, G. Review of the book *Littlewood-Paley and multiplier theory.* *Bull. Am. Math. Soc.* 84, 242–250, (1978).
11. Cowling, M. Maceuri, G. Inequalities for some maximal functions I. *Trans. Amer. Math. Soc.* 287, 431–455, (1985).
12. Cowling, M. On Littlewood-Paley-Stein theory, *Rend. Semin. Mat. Fis. Milano XLIX*, 79–87, (1979).

13. Duoandikoetxea, J. *Fourier Analysis*. 29, American Mathematical Society, Providence (2000).
14. Fefferman, C. Inequalities for strongly singular integral operators, *Acta Math.* 24, 9–36, (1970).
15. Duoandikoetxea, J. Rubio de Francia, J. L. Maximal and singular integral operators via Fourier transform estimates, *Invent. Math.* 84, 541–561, (1986).
16. Fefferman, C., Stein, E. H^p spaces of several variables. *Acta Math.*, 129, 137–193, (1972).
17. Fefferman, C. L^p -bounds for pseudo-differential operators, *Israel J. Math.* 14, 413–417, (1973).
18. Fischer, V. The spherical maximal function on the free two-step nilpotent Lie group, *Math. Scand.* 99, 99–118, (2006).
19. Fischer V., Ruzhansky M., *Quantization on nilpotent Lie groups*, Progress in Mathematics, Vol. 314, Birkhauser, 2016. xiii+557pp.
20. Folland, G. Stein, E. *Hardy Spaces on Homogeneous Groups*, Princeton University Press, Princeton, N.J., 1982.
21. Ganguly, P. Thangavelu, S. On the lacunary spherical maximal function on the Heisenberg group. *J. Funct. Anal.* 280, no. 3, Paper No. 108832, 32 pp. (2021).
22. Grafakos, L. *Classical Fourier Analysis*.vol. 249, 3rd edn. Springer, New York (2014).
23. Hardy, G. H. A theorem concerning Taylor’s series, *Quart. J. Pure Appl. Math.*, 44, 147–160, (1913).
24. Helffer, B. Nourrigat, J. Characterisation des operateurs hypoelliptiques homogenes invariants a gauche sur un groupe de Lie nilpotent gradue, *Comm. Partial Differential Equations*, 4(8), 899–958, (1979).
25. Hörmander, L. Estimates for translation invariant operators in L^p spaces, *Acta Math.*, 104, 93–139, (1960).
26. Hong Q., Hu G., Ruzhansky M., Fourier multipliers for Hardy spaces on graded Lie groups. To appear in *Proc. Royal Soc. Edinburgh*. arXiv:2101.07499.
27. Lacey, M. T. Sparse bounds for spherical maximal functions, *J. Anal. Math.* 139, (2019), 613–635.
28. Mockenhaupt, G. Seeger, A., Sogge, C. Wave front sets, local smoothing and Bourgain’s circular maximal theorem, *Ann. of Math.* 136, (1992), 207–218.
29. Müller, D., Seeger, A. Singular spherical maximal operators on a class of two-step nilpotent Lie groups, *Isr. J. Math.* 141, 315–340, (2004).
30. Nagel, A. Rivière, N. M., Wainger, S. On Hilbert transforms along curves, II, *Amer. J. Math.* 98, 395–403, (1976).
31. Narayanan, E. K., Thangavelu, S. An optimal theorem for the spherical maximal operator on the Heisenberg group, *Isr. J. Math.* 144, 211–219, (2004).
32. Nevo, A. Thangavelu, S. Pointwise ergodic theorems for radial averages on the Heisenberg group, *Adv. Math.* 127, 307–339, (1997).
33. Rothschild, L. P., Stein, E. M. Hypoelliptic differential operators and nilpotent groups. *Acta Math.*, 137(3–4), 247–320, (1976).
34. Rubio De Francia, J. L. Maximal functions and Fourier transforms, *Duke Math. J.* 53, 395–404, (1986).
35. Seeger, A. Some inequalities for singular convolution operators in L^p -spaces, *Trans. Amer. Math. Soc.*, 308(1), 259–272, (1988).
36. Seeger, A. Remarks on singular convolution operators. *Studia Math.* 97(2), 91–114, (1990).
37. Seeger, A. Endpoint estimates for multiplier transformations on compact manifolds. *Indiana Univ. Math. J.* 40(2), 471–533, (1991).
38. Seeger, A. Sogge, C. D. On the boundedness of functions of (pseudo-) differential operators on compact manifolds. *Duke Math. J.* 59(3), 709–736, (1989).
39. Seeger, A., Sogge, C. D., Stein, E. M. Regularity properties of Fourier integral operators. *Ann. of Math.*, 134(2), 231–251, (1991).
40. Seddik, A. Some results related to the Corach-Porta-Recht inequality. *Proc. Amer. Math. Soc.* 129, no. 10, pp. 3009–3015, (2001).
41. Schmidt, O. *Maximaloperatoren zu Hyperflächen in Gruppen vom homogenen Typ*, Diplomarbeit an der Christian-Albrechts-Universität zu Kiel, Mai 1998.

42. Stein, E. M. Maximal functions. I. Spherical means, Proc. Natl. Acad. Sci. USA 73, 2174–2175, (1976).
43. Stein, E. M., Singular integrals and differentiability properties of functions. Princeton 1970.
44. Taylor, M. Pseudodifferential Operators, Princeton Univ. Press, Princeton, N.J., 1981.
45. Seeger, A. Tao, T. Wright, J. Singular maximal functions and Radon transforms near L^1 , Amer. J. Math. 126, 607–647, (2004).
46. Wainger, S. Special trigonometric series in k -dimensions, Mem. Amer. Math. Soc., 59, (1965).

DUVÁN CARDONA:

DEPARTMENT OF MATHEMATICS: ANALYSIS, LOGIC AND DISCRETE MATHEMATICS

GHENT UNIVERSITY, BELGIUM

E-mail address duvanc306@gmail.com, duvan.cardonasanchez@ugent.be

JULIO DELGADO:

DEPARTAMENTO DE MATEMATICAS

UNIVERSIDAD DEL VALLE

CALI-COLOMBIA

E-mail address delgado.julio@correounivalle.edu.co

MICHAEL RUZHANSKY:

DEPARTMENT OF MATHEMATICS: ANALYSIS, LOGIC AND DISCRETE MATHEMATICS

GHENT UNIVERSITY, BELGIUM

AND

SCHOOL OF MATHEMATICAL SCIENCES

QUEEN MARY UNIVERSITY OF LONDON

UNITED KINGDOM

E-mail address michael.ruzhansky@ugent.be, m.ruzhansky@qmul.ac.uk