

On the determination of optimal tuning parameters for a space-variant LASSO problem using geometric and convex analysis techniques

Gianluca Giacchi, Bastien Milani, Benedetta Franceschiello

Abstract

Compressed Sensing (CS) comprises a wide range of theoretical and applied techniques to recover signals given a partial knowledge of their coefficients. It finds its applications in several fields, such as mathematics, physics, engineering, and many medical sciences, to name a few, cf. [17, 7, 13, 25, 33, 38, 36, 56, 57]. Driven by our interest in the mathematics behind Magnetic Resonance Imaging (MRI) and Compressed Sensing (CS), we use convex analysis techniques to determine analytically the optimal tuning parameters of the space-variant LASSO with voxel-wise weighting, under assumptions on the fidelity term, either on the sign of its gradient or orthogonality-like conditions on its matrix. Finally, we conclude conjecturing what the explicit form of optimal parameters should be in the most general setting (hypotheses-free) of the space-variant LASSO.

1 Introduction

The main topic of this work is the analytic determination of the optimal parameters of a space-variant version of LASSO, a widely studied inverse problem in the field of Compressed-Sensing (CS). Namely, we investigate the dependence of these parameters on local estimates of the unknown variable, exploiting results of convex analysis and optimization theory. The motivation behind our study is the applicability of such results in the field of Magnetic Resonance Imaging (MRI).

Basis Pursuit is a well-known convex minimization problem that was first introduced by F. Santosa and W. W. Symes in 1986, cf. [50] and it was applied to signal processing by S. S. Chen and D. Donoho in 1994, cf. [15]. In 1996, R. Tibshirani re-introduced it as linear regression method, under the name of LASSO, cf. [52]. Convex analysis approaches to LASSO problem are not new, cf. [54, 55, 3].

In its simplest definition, LASSO can be formulated as

$$\arg \min_{x \in \mathbb{R}^n} \|Ax - b\|_2^2 + \lambda \|x\|_1, \quad (1)$$

where $A \in \mathbb{R}^{m \times n}$ and $b \in \mathbb{R}^m$ is a measurement vector. (1) can be interpreted as a regularization problem, where the aim is to minimize simultaneously the fidelity term $\|Ax - b\|_2^2$ - that measures noise - and the regularization term $\|x\|_1$ - that enforces sparsity. We stress that (1) can be formulated in the more general case in which x is sparse with respect to some sparsifying transform $\Phi : \mathbb{R}^n \rightarrow \mathbb{R}^N$, replacing the ℓ^1 -norm with $\|\Phi x\|_1$ (we will explain better what a sparsifying transform is in the following).

Under a convex-analysis point of view, λ is a Lagrange multiplier and the analytic dependence $\lambda = \lambda(\tau)$ is usually hard to express, since the function $\|x\|_1$ is non-differentiable and the constraint in (3) is an inequality constraint. Several estimating procedures have been tried so far: CNN and other learning methods were employed in [30, 41], while a more statistical point of view was adopted in [9]. Nonetheless, not only the explicit dependence (4) would shed light on the actual relationship between λ and noise or sparsity estimates, but it could also be useful in concrete applications, such as imaging moving organs through MRI, cf. [18, 24], where a more general definition of LASSO is defined up to exploit different sparsifying transforms (e.g. wavelets and Total Variation), i.e. linear mappings $\Phi : \mathbb{R}^n \rightarrow \mathbb{R}^N$ such that Φx is sparse for all $x \in \mathbb{R}^n$ of interest. In MRI, for instance, vectors of interest are MR images, which tend to be approximatively sparse with respect to discrete Fourier transform (DFT), discrete cosine transform (DCT), discrete wavelets (DWT), cf. [34]. This means that a solution of the generalized LASSO problem

$$\min_{x \in \mathbb{R}^n} \|Ax - b\|_2^2 + \lambda \|\Phi x\|_1, \quad (2)$$

(where $A = UFC$ is a matrix that usually embodies properties of the scanner (coil sensitivity), undersampling schemes and DFT, and b is the underdetermined, possibly noisy measurement) will have sparse regularization term, i.e. sparse Φx . We stress that this problem always admits a solution for any choice of A , b , $\lambda \geq 0$ and Φ , and for the sake of completeness we report the proof in the appendix. In order to exploit more a-priori knowledge on the structure of MRI data, (2) can be generalized further up to considering target functions that are sum of more than one regularizing term. We refer to [27, 22, 46] for examples of these and other generalizations.

However, not always sparsity is the correct assumption in MRI. For instance, dynamic MRI images (e.g. a sequence of images of a moving organ, cf. [18, 24]) are highly compressible, rather than sparse, cf. [35]. This means that most of their coefficients with respect to some sparsifying transform are small or negligible, but they are not sparse.

The parameter $\lambda > 0$ in (1) acts as a tuning parameter that balances the contributions of these two terms: small values of λ lower the contribution of the regularization, strengthening the effect of the fidelity term; vice-versa, large values of λ make $\|Ax - b\|_2^2$ negligible and force $\|x\|_1$ to be small in order for the overall sum to be small. Consequently, solutions corresponding to $\lambda \ll 1$ will be noisy, being close to the set $A^{-1}b$,

while solutions corresponding to $\lambda \gg 1$ will be more sparse, as they are close to 0.

From this perspective, estimates of tuning parameters for inverse problems can be performed pursuing three different approaches, cf. [1]:

- **a-posteriori:** when a-priori knowledge on the amplitude of noise $e \in \mathbb{R}^m$ is accessible, say $\|e\|_2 \leq \varepsilon$. For instance, using Morozov's discrepancy principle, λ can be chosen so that a solution x_λ of (1) satisfies $\|Ax_\lambda - b\|_2 \leq \varepsilon$, cf. [29, 8].
- **A-priori:** these methods require knowledge of noise level, as before, but also a-priori information on the regularity of the solution. For this reason, a-priori approaches are usually bad suited for applications.
- **Heuristic:** they are methods such as the L-curve, cf. [10], where the optimal parameter is chosen empirically by tracing a trade-off curve (the L-curve), or the generalized cross-validation (GCV), a well-performing method that requires high-dimensional matrix calculus, cf. [31, 53, 26].

Other non-standard methods can be found in [32, 44], where the parameter is chosen so that statistical properties of noise, such as whiteness, are optimized; an implementation that avoids the computation of matrix inverses can be found in [5].

However, the very reason why λ is interpreted as a trade-off between noise and sparsity is that it depends on estimates that are usually unavailable, such as a-priori upper bounds for the ℓ^1 norm of the unknown, i.e. a-priori information on the sparsity of the solution, or upper bounds for the noise, cf. [42]. This dependence can be highlighted working on the expression of (1). The function

$$L(x, \lambda) = \|x\|_1 + \lambda(\|Ax - b\|_2^2 - \eta^2)$$

is the Lagrangian associated to the constrained minimization problem

$$\begin{cases} \min_{x \in \mathbb{R}^n} \|x\|_1, \\ \|Ax - b\|_2^2 \leq \eta^2. \end{cases} \quad (3)$$

Therefore, (1) and (3) are equivalent, up to choosing

$$\lambda = \lambda(\eta) \quad (4)$$

or, equivalently, $\eta = \eta(\lambda)$, properly. $\|x\|_1$ is said to *enforce sparsity* because a solution of (3), if unique, must be a sparse vector.

Surprisingly, it is easier to identify the relation between parameter and a-priori estimate when the more general space-variant version of LASSO is considered. Namely, we aim to compute the optimal parameters for the problem

$$\arg \min_{x \in \mathbb{R}^n} \|Ax - b\|_2^2 + \sum_{j=1}^n \lambda_j |x_j| \quad (5)$$

using convex and optimization problems analysis, techniques that have been also used in [4], for instance.

Other space-variant versions of this problem have been considered in literature, for instance, in [41] the authors present a Total Variation (TV) regularization-based space-variant LASSO for image denoising. Other references are [8, 45].

Again, problem (5) can be interpreted as the unconstrained equivalent formulation of

$$\begin{cases} \min_{x \in \mathbb{R}^n} \|Ax - b\|_2^2, \\ |x_j| \leq \tau_j \end{cases} \quad j = 1, \dots, n, \quad (6)$$

up to choose the τ_j 's and the λ_j 's ($j = 1, \dots, n$) properly. As we shall see, the relationship between these parameters gets harder and harder as A moves away from being orthogonal, due to the complicated geometry of (6). Loosely speaking, this is due to the fact that A shuffles the coordinates of x if $A^T A$ is non-diagonal. In this situation, each couple of sets $M_j := \{x \in \mathbb{R}^n : x_j = -\tau_j\}$ and $N_j := \{x \in \mathbb{R}^n : \frac{\partial}{\partial x_j} (\|Ax - b\|_2^2)\}$ are not parallel anymore.

As for the applicability of these results, the dependence on τ_1, \dots, τ_n of the optimal parameters may turn to be too restrictive, because these estimates are not in general available. However, we claim that in certain situations, such as denoising, the solution obtained using the estimated τ_j 's leads to high-quality denoising.

We also point out that, unless $\tau_j = 0$ for some $j = 1, \dots, n$, a solution of (6) does not need to be sparse, since the sparsity granted by (3) is due to the geometry of the unit ball of ℓ^1 . Clearly, according to the size of τ_j 's, problem (6), and therefore problem (5), can lead to compressible solutions.

This work investigates the explicit relationship between the parameters λ_j in (5) (as Lagrange multipliers) and the local estimates τ_j in (6). The choice of working with (6) instead of its equivalent form

$$\begin{cases} \min_{x \in \mathbb{R}^n} \|Ax - b\|_2^2, \\ x_j^2 \leq \tau_j^2 \end{cases} \quad j = 1, \dots, n,$$

which has the advantage of having smooth constraints, is that the strategy we follow does not exploit the differentiability of the constraints. Another technique can be followed that exploits this smoothness, but it turns to be ineffective, as we will specify later.

In Section 2, we establish the notation we use in this work. In Section 3 we compute the deterministic relations between λ_j 's and τ_j 's in order for problems (6) and (5) to be equivalent. Using the general theory, we prove that if $A^T A$ is diagonal, for instance if A is a subsampling matrix, the Fourier transform matrix or even the identity matrix, then the optimal parameters are explicitly given by

$$\lambda_j^\# = 2 \|a_{*,j}\|_2^2 \left(\frac{|\langle b, a_{*,j} \rangle|}{\|a_{*,j}\|_2^2} - \tau_j \right) \chi_{\left[0, \frac{|\langle b, a_{*,j} \rangle|}{\|a_{*,j}\|_2^2}\right]}(\tau_j), \quad (7)$$

where $a_{*,j}$ denotes the j -th column of A . We also provide deterministic results for those cases where we have a control on the sign of the gradient of $\|Ax - b\|_2^2$, proving an analogous of (7) in that situation. The conclusions are reported in Section 4, where a conjecture on the optimal parameters in the most general situation is presented.

We point out that our result is interesting for two main reasons: to the best of our knowledge, the analytic dependence that we investigate was never fully understood, neither computed. Formula (7) can be applied directly for denoising in some transform domain, i.e. if A is orthogonal or the identity itself. Furthermore, if A is the matrix of an undersampling pattern, $A^T A$ is diagonal and (7) can be exploited to control the tuning parameters of the space-variant LASSO problem (5) in terms of voxel-wise estimates. We presume that such estimates are relatively easy to obtain. For example, one may first reconstruct a highly undersampled image, apply filters using convolution techniques and estimate the tuning parameters λ from the filtered image, therefore obtaining a denoised image.

Finally we conjecture how this theory could be applied to compute the analytical expression of optimal parameters in the case of $A^T A$ non diagonal, solving an open problem in optimisation (5).

2 Preliminaries and notation

2.1 Lagrange duality

We consider a constrained optimization problem in the form

$$\begin{cases} \inf_{x \in \mathbb{R}^n} F_0(x), \\ Ax = y, \\ F_l(x) \leq b_l \quad l = 1, \dots, M, \end{cases} \quad (8)$$

where $A \in \mathbb{R}^{m \times n}$, $y \in \mathbb{R}^m$ and $F_0, F_1, \dots, F_M : \mathbb{R}^n \rightarrow (-\infty, +\infty]$. Throughout this work, we identify \mathbb{R}^k either with $\mathbb{R}^{1 \times k}$ or $\mathbb{R}^{k \times 1}$, as this shall not cause confusion. Also, we always assume that the infimum in (8) is attained.

A point $x \in \mathbb{R}^n$ is called **feasible** if it belongs to the constraints, that is if

$$x \in K := \left\{ \zeta \in \mathbb{R}^n : A\zeta = y \text{ and } F_l(\zeta) \leq b_l, l = 1, \dots, M \right\} \quad (9)$$

and K is called the **set of feasible points**. To avoid triviality, we always assume $K \neq \emptyset$, in the which case the problem (8) is called **feasible**. In view of the definition of K , the problem (8) can be implicitly written as

$$\min_{x \in K} F_0(x).$$

Problems such as (3) and (6) can be approached by considering their Lagrange formulation. The **Lagrange function** related to (8) is the function $L : \mathbb{R}^n \times \mathbb{R}^m \times$

$[0, +\infty)^M \rightarrow (-\infty, +\infty]$ defined as

$$L(x, \xi, \lambda) := F_0(x) + \langle \xi, Ax - y \rangle + \sum_{l=1}^M \lambda_l (F_l(x) - b_l),$$

where $\langle \cdot, \cdot \rangle$ denotes the canonical inner product of \mathbb{R}^n . Namely,

$$\langle x, y \rangle = x^T y = \sum_{j=1}^n x_j y_j.$$

Observe that for all ξ, λ and $x \in K$

$$L(x, \xi, \lambda) = F_0(x) + \underbrace{\langle \xi, Ax - y \rangle}_{=0} + \sum_{l=1}^M \underbrace{\lambda_l}_{\geq 0} \underbrace{(F_l(x) - b_l)}_{\leq 0} \leq F_0(x),$$

so that

$$\inf_{x \in \mathbb{R}^n} L(x, \xi, \lambda) \leq \inf_{x \in K} L(x, \xi, \lambda) \leq \inf_{x \in K} F_0(x). \quad (10)$$

Definition 2.1. The function $H : \mathbb{R}^m \times [0, +\infty)^M \rightarrow [-\infty, +\infty]$ defined as

$$H(\xi, \lambda) := \inf_{x \in \mathbb{R}^n} L(x, \xi, \lambda)$$

is called **Lagrange dual function**.

Inequalities (10) read as

$$H(\xi, \lambda) \leq \inf_{x \in K} F_0(x) \quad (11)$$

for all $\xi \in \mathbb{R}^m$ and all $\lambda \in [0, +\infty)^M$. For the rest of this work, if $x, y \in \mathbb{R}^n$, the notation $x \succeq y$ stands for $x_j \geq y_j$ for all $j = 1, \dots, n$. Stating (11) differently, we have the **weak duality inequality**

$$\sup_{\substack{\xi \in \mathbb{R}^m \\ \lambda \succeq 0}} H(\xi, \lambda) \leq \inf_{x \in K} F_0(x). \quad (W)$$

We point out that (W) is equivalent to

$$\sup_{\xi, \lambda} \inf_x L(x, \xi, \lambda) \leq \inf_x \sup_{\xi, \lambda} L(x, \xi, \lambda) \quad (12)$$

(see [6, Subsection 5.4.1]).

We are interested in computing the parameters (ξ, λ) such that (W) is an equality, in the which case (W) becomes

$$\sup_{\substack{\xi \in \mathbb{R}^m \\ \lambda \succeq 0}} H(\xi, \lambda) = \inf_{x \in K} F_0(x), \quad (S)$$

which is the so-called **strong duality**.

2.2 Subdifferential

Definition 2.2 (Subdifferential). Let $\Omega \subseteq \mathbb{R}^n$ be open and $g : \Omega \rightarrow \mathbb{R}$. Let $x_0 \in \Omega$. The **subdifferential** of g at x_0 is the set

$$\partial g(x_0) := \{v \in \mathbb{R}^n : g(x) \geq g(x_0) + v^T(x - x_0) \forall x \in \Omega\}.$$

We refer to any $v \in \partial g(x_0)$ as a **subgradient** of g at x_0 .

We will use the following proposition.

Proposition 2.3. *Let $\Omega \subseteq \mathbb{R}^n$ be open and $g : \Omega \rightarrow \mathbb{R}$ be convex and continuous on Ω . Let $x_0 \in \Omega$. Then, $\partial g(x_0) \neq \emptyset$.*

2.3 Optimal parameters as Lagrange multipliers

Let $F(x) := (F_1(x), \dots, F_M(x))$. In the convex framework, if the constraint $F(x) \preceq b$ does not reduce to $F(x) = b$, namely if for all $l = 1, \dots, M$ the inequality $F_l(x) < b_l$ holds for some $x \in \mathbb{R}^n$, then the strong duality holds.

Theorem 2.4 (Cf. [6], Section 5.3.2). *Assume that F_0, F_1, \dots, F_M are convex functions defined on \mathbb{R}^n . Let $x^\#$ be such that $F_0(x^\#) = \inf_{x \in \mathbb{R}^n} F_0(x)$. If*

- (i) *there exists $\tilde{x} \in \mathbb{R}^n$ such that $A\tilde{x} = y$ and $F(\tilde{x}) \preceq b$ or,*
- (ii) *in absence of inequality constraints, if $K \neq \emptyset$ (i.e. if there exists $\tilde{x} \in \mathbb{R}^n$ such that $A\tilde{x} = y$),*

then, there exists $(\xi^\#, \lambda^\#) \in \mathbb{R}^m \times [0, +\infty)^M$ such that $H(\xi^\#, \lambda^\#) = \sup_{\xi, \lambda} H(\xi, \lambda)$ and $H(\xi^\#, \lambda^\#) = F_0(x^\#)$.

The proof of Theorem 2.4 contains the fundamental construction we will use in the next sections and we report it for this reason. We refer to [6, Subsection 5.3.2]. First, we need a result from functional analysis, which is well-known as (geometrical) *Hahn-Banach theorem*.

Definition 2.5 (Separating Hyperplane). Consider two subsets $\mathcal{A}, \mathcal{B} \subseteq \mathbb{R}^n$. A hyperplane $\Gamma := \{x \in \mathbb{R}^n : \langle \xi, x \rangle = \alpha\}$ satisfying

$$\langle \xi, x \rangle \leq \alpha \leq \langle \xi, y \rangle \tag{13}$$

is a **separating hyperplane** between \mathcal{A} and \mathcal{B} .

Theorem 2.6 (Cf. [49] Theorem 3.4). *Let $\mathcal{A}, \mathcal{B} \subset \mathbb{R}^n$ be two convex and disjoint subsets. If at least one of them is open, there exists $\xi \in \mathbb{R}^n$ and $\alpha \in \mathbb{R}$ such that (13) holds for all $x \in \mathcal{A}$ and all $y \in \mathcal{B}$.*

Idea of the proof of Theorem 2.4. First, one assumes that A has full row-rank. Moreover, observing that in the other case the assertion is essentially trivial, one reduces to consider the situation in which $p^* := \inf_{x \in K} F_0(x) > -\infty$.

Consider the set

$$\mathcal{G} := \left\{ (F(x) - b, Ax - y, F_0(x)) \in \mathbb{R}^M \times \mathbb{R}^m \times \mathbb{R} : x \in \mathbb{R}^n \right\}, \quad (14)$$

where, with an abuse of notation, $Ax - y$ denotes the row vector with the same (ordered) entries of $Ax - y$, and \mathcal{A} be defined as the epigraph

$$\begin{aligned} \mathcal{A} &:= \mathcal{G} + ((\mathbb{R}_{\geq 0})^M \times \mathbb{R}^m \times \mathbb{R}_{\geq 0}) = \\ &= \left\{ (u, v, t) \in \mathbb{R}^M \times \mathbb{R}^m \times \mathbb{R} : u \succeq F(x) - b, \right. \\ &\quad \left. v = Ax - y, t \geq F_0(x) \text{ for some } x \in \mathbb{R}^n \right\}. \end{aligned} \quad (15)$$

It is easy to verify that if F_0, F_1, \dots, F_M are convex, than \mathcal{A} is convex. Then, consider the set

$$\mathcal{B} := \left\{ (0, 0, s) \in \mathbb{R}^M \times \mathbb{R}^m \times \mathbb{R} : s < p^* \right\}.$$

\mathcal{A} and \mathcal{B} are clearly disjoint, \mathcal{B} (which is an open half-line) being trivially convex and open. Therefore, the assumptions of Theorem 2.6 are satisfied and we conclude that there exists a triple of parameters $(\tilde{\lambda}, \tilde{\xi}, \mu) \neq 0$ and $\alpha \in \mathbb{R}$ such that

$$(u, v, t) \in \mathcal{A} \implies \tilde{\lambda}^T u + \langle \tilde{\xi}, v \rangle + \mu t \geq \alpha, \quad (16)$$

$$(u, v, t) \in \mathcal{B} \implies \tilde{\lambda}^T u + \langle \tilde{\xi}, v \rangle + \mu t \leq \alpha. \quad (17)$$

It is easy to see that the definition of \mathcal{A} , together with (16), imply that $\tilde{\lambda}_l \geq 0$ for all $l = 1, \dots, M$ and $\mu \geq 0$. Also, applying the definition of \mathcal{B} to (17), one finds that $\mu t \leq \alpha$ for all $t < p^*$, which implies that $\mu p^* \leq \alpha$. Therefore, for all $x \in \mathbb{R}^n$,

$$\sum_{l=1}^M \tilde{\lambda}_l (F_l(x) - b_l) + \langle \tilde{\xi}, Ax - y \rangle + \mu F_0(x) \geq \alpha \geq \mu p^*. \quad (18)$$

If $\mu > 0$, then (18) gives that $L(x, \tilde{\xi}/\mu, \tilde{\lambda}/\mu) \geq p^*$ for all $x \in \mathbb{R}^n$, which implies that $H(\tilde{\xi}/\mu, \tilde{\lambda}/\mu) \geq p^*$. Since the other inequality is implicit in weak duality inequality, we conclude that $H(\tilde{\xi}/\mu, \tilde{\lambda}/\mu) = p^*$. Finally, using the assumption on the existence of a point satisfying the strict inequality constraint, one proves by contradiction that it must be $\mu > 0$. \square

Definition 2.7 (Lagrange Optimal Parameters). We refer to a couple $(\xi^\#, \lambda^\#) \in \mathbb{R}^m \times [0, +\infty)^M$ as to as **optimal parameters** for the problem (8) if $(\xi^\#, \lambda^\#)$ attend the supremum in (S).

As a consequence of Theorem 2.4, we have the following result, which relates the minimizers of (8) and that of the dual problem $\max_{\xi, \lambda} H(\xi, \lambda)$, providing also the optimal parameters.

Corollary 2.8 (Cf. [23] Theorem B.28). *Let $F_0 : \mathbb{R}^n \rightarrow [0, +\infty)$ and $\phi : [0, +\infty) \rightarrow \mathbb{R}$ be such that ϕ is monotonically increasing and $\phi \circ F_0$ is convex. Let $\tau_j > 0$ ($j = 1, \dots, M$) and $\Psi_j : \mathbb{R}^n \rightarrow \mathbb{R}$ ($j = 1, \dots, M$) be convex functions such that $\Psi_j^{-1}([0, \tau_j]) \neq \emptyset$ for all $j = 1, \dots, M$. Let $x^\#$ is a minimizer of the problem*

$$\min_{x \in \mathbb{R}^n} F_0(x) \quad \text{subject to } \Psi(x) \preceq \tau, \quad (19)$$

being $\tau = (\tau_1, \dots, \tau_M)$. Then, there exist $\lambda_j \geq 0$ ($j = 1, \dots, M$) such that $x^\#$ is a minimizer of

$$\min_{x \in \mathbb{R}^n} \phi(F_0(x)) + \sum_{j=1}^M \lambda_j \Psi_j(x). \quad (20)$$

Proof. Since ϕ is monotonically increasing, (19) is obviously equivalent to

$$\min_{x \in \mathbb{R}^n} \phi(F_0(x)) \quad \text{subject to } \Psi_j(x) \leq \tau_j, \quad (21)$$

($j = 1, \dots, M$) whose Lagrangian is given by

$$L(x, \lambda) = \phi(F_0(x)) + \sum_{j=1}^M \lambda_j (\Psi_j(x) - \tau_j). \quad (22)$$

By the assumption, $\phi \circ F_0$ and each Ψ_j are convex and the inequalities $\Psi_j(\tilde{x}) < \tau_j$ are satisfied by some $\tilde{x} \in \mathbb{R}^n$ (observe that here we need $\tau_j > 0$), so we can apply Theorem 2.4 to get $H(\lambda^\#) = \phi(F_0(x^\#))$ for some $\lambda^\# \in [0, +\infty)^M$. By (12), for all $x \in \mathbb{R}^n$

$$L(x^\#, \lambda^\#) \leq L(x, \lambda^\#),$$

so that $x^\#$ is also a minimizer of the function $x \in \mathbb{R}^n \mapsto L(x, \lambda^\#)$. Since the constant terms $-\lambda_j \tau_j$ in (22) do not affect the minimum, we have that $x^\#$ is a minimizer of

$$\min_{x \in \mathbb{R}^n} \phi(F_0(x)) + \sum_{j=1}^M \lambda_j^\# (\Psi_j(x) - \tau).$$

□

Remark 2.9. Theorem 2.6 has a complex version that holds with $\Re\langle z, w \rangle = \Re\left(\sum_{j=1}^n \bar{z}_j w_j\right)$ (\Re denotes the real part of a complex number) instead of $\langle \cdot, \cdot \rangle$. In particular, the whole theory of this work holds in the complex framework as well, up to replacing the canonical real inner product of \mathbb{R}^n with the real inner product on \mathbb{C}^n defined above. Therefore, we don't need to study the complex case separately, as only the structure of \mathbb{C}^n as a real vector space is involved.

Remark 2.10. To sum up, Theorem 2.4 and Corollary 2.8 together tell that, up to the sign, the coefficients of any hyperplane that separates the two sets

$$\mathcal{A} = \left\{ (u, t) \in \mathbb{R}^{M+1} : u \succeq F(x) - b, t \geq F_0(x) \text{ for } x \in \mathbb{R}^n \right\}$$

and

$$\mathcal{B} = \left\{ (0, t) \in \mathbb{R}^{M+1} : t < \inf_{x \in K} F_0(x) \right\}$$

define optimal parameters for problem (8), in absence of equality constraints, i.e. if $y = 0$ and $A = 0$ in (8). This is the geometric idea we will apply in the following sections to the space-variant basis pursuit.

3 The space-variant basis pursuit

Let $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$ and $\tau_1, \dots, \tau_n \geq 0$. We denote with $a_{*,j}$ the j -th column of A and set $b = (b_1, \dots, b_m)$. We consider the constrained minimization problem

$$\begin{cases} \min_{x \in \mathbb{R}^n} \|Ax - b\|_2^2, \\ |x_j| \leq \tau_j \end{cases} \quad j = 1, \dots, n. \quad (23)$$

We also assume that $\tau_j \neq 0$ for all $j = 1, \dots, n$. In fact, if $\tau_j = 0$ for some $j = 1, \dots, n$, then the solution $x = (x_1, \dots, x_n)$ has $x_j = 0$. In this case, problem (23) reduces to

$$\begin{cases} \min_{x \in \mathbb{R}^n} \|\tilde{A}y - b\|_2^2, \\ |y_{i_j}| \leq \tau_{i_j} \end{cases} \quad j = 1, \dots, n-r, \quad (24)$$

where $r = \text{card}\{j : \tau_j = 0\} \leq m$, $J = \{1 \leq i_1 < \dots < i_{n-r} \leq n\} := \{j : \tau_j \neq 0\}$ and $\tilde{A} = (a_{*,j})_{j \in J} \in \mathbb{R}^{m \times (n-r)}$.

Let K denote the set of the feasible points, that is

$$K = \{x \in \mathbb{R}^n : |x_j| \leq \tau_j \forall j = 1, \dots, n\} \quad (25)$$

and consider the associated Lagrange function, i.e.

$$L(x, \lambda_1, \dots, \lambda_n) = \|Ax - b\|_2^2 + \sum_{j=1}^n \lambda_j (|x_j| - \tau_j). \quad (26)$$

We are interested in a vector of optimal parameters $\lambda^\# \succeq 0$, where the concept of optimality is that of Definition 2.7. Following the proofs of Theorem 2.4 and Corollary 2.8, $\lambda^\#$ can be taken as the direction of an hyperplane that separates

$$\mathcal{A} = \left\{ (u, t) \in \mathbb{R}^n \times \mathbb{R} : u_l \geq |x_l| - \tau_l \ (l = 1, \dots, n), t \geq \|Ax - b\|_2^2 \text{ for some } x \in \mathbb{R}^n \right\}$$

and

$$\mathcal{B} = \left\{ (0, t) \in \mathbb{R}^n \times \mathbb{R} : t < p^* \right\},$$

where $p^* := \inf_{x \in K} \|Ax - b\|_2^2$.

3.1 The scalar case

To clarify the general procedure, we focus on the simple case $m = n = 1$ first, in the which case (23) becomes

$$\begin{cases} \min_{x \in \mathbb{R}} (Ax - b)^2, \\ |x| \leq \tau, \end{cases} \quad (27)$$

where $A \in \mathbb{R} \setminus \{0\}$ and $b, x \in \mathbb{R}$. To find the optimal parameter, we consider the set \mathcal{G} of points $(u, t) \in \mathbb{R}^2$ that satisfy

$$\begin{cases} u = |x| - \tau, \\ t = (Ax - b)^2, \end{cases}$$

which give a curve of the half-plane $U = \{(u, t) \in \mathbb{R}^2 : u \geq \tau, t \geq 0\}$ parametrized by $x \in \mathbb{R}$. More precisely:

- if $x \geq 0$,

$$\begin{cases} x = u + \tau, \\ t = (A(u + \tau) - b)^2 = (Au + (A\tau - b))^2, \end{cases}$$

which is a branch of parabola in U with vertex in $(\frac{b}{A} - \tau, 0)$.

- If $x < 0$

$$\begin{cases} x = -u - \tau, \\ t = (-A(u + \tau) - b)^2 = (Au + (A\tau + b))^2, \end{cases}$$

which is, again, a branch of parabola in U , having its vertex in $(-\frac{b}{A} - \tau, 0)$.

Writing explicitly the sets \mathcal{A} and \mathcal{B} , a simple study of all the possible scenarios, leads to the following result.

Proposition 3.1. *Let $\tau > 0$, $A \in \mathbb{R} \setminus \{0\}$, $b \in \mathbb{R}$. The optimal parameter for (27) is given by:*

$$\lambda^\# = \begin{cases} 2A^2(|b/A| - \tau) & \text{if } 0 < \tau < |b/A|, \\ 0 & \text{if } \tau \geq |b/A| \end{cases} = 2A^2(|b/A| - \tau) \chi_{0 < \tau < |b/A|}(\tau).$$

3.2 Properties of \mathcal{A}

Consider $A \in \mathbb{R}^{m \times n}$ and $b = (b_1, \dots, b_m) \in \mathbb{R}^m$, with

$$A = \begin{pmatrix} a_{11} & \dots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{m1} & \dots & a_{mn} \end{pmatrix}.$$

We consider the problem (23) and the associated Lagrange function

$$L(x, \lambda) := \|Ax - b\|_2^2 + \sum_{j=1}^n \lambda_j (|x_j| - \tau_j). \quad (28)$$

Recall that p^* was defined as $p^* := \min_{x \in K} \|Ax - b\|_2^2$, being K the set of the points $x \in \mathbb{R}^n$ such that $|x_j| \leq \tau_j$ for all $j = 1, \dots, n$. It is not difficult to verify that

$$p^* = \inf \left\{ t \in \mathbb{R} : (u, t) \in \mathcal{G}, u_j \leq 0 \quad \forall j = 1, \dots, n \right\}. \quad (29)$$

Let \mathcal{M}_n be the set of the n -dimensional signature matrices, that are the diagonal matrices $S = (s_{ij})_{i,j=1}^n \in \mathbb{R}^{n \times n}$ such that $|s_{jj}| = 1$ for all $j = 1, \dots, n$. We stress that \mathcal{M}_n is a subgroup of the orthogonal group $O(n)$. Moreover, if $S \in \mathcal{M}_n$, then $S^2 = I_{n \times n}$, where $I_{n \times n}$ denotes the identity matrix in $\mathbb{R}^{n \times n}$, in particular S is invertible with $S^{-1} = S$. If $x \in \mathbb{R}^n$ and $S \in \mathcal{M}_n$ is such that $Sx \in \prod_{j=1}^n [0, +\infty)$, we write $S \in \text{sgn}(x)$.

Lemma 3.2. *Let $A \in \mathbb{R}^{m \times n}$. Let $b \in \mathbb{R}^m$ and $\tau_j > 0$ for $j = 1, \dots, n$. Let $S \in \mathcal{M}_n$. There exists $u \in \prod_{j=1}^n [-\tau_j, 0]$ such that $ASu + AS\tau - b = 0$ if and only if $S \in \text{sgn}(x)$ for some $x \in \mathbb{R}^n$ such that $Ax = b$ and $|x_j| \leq \tau_j$.*

Proof. Assume that there exists $u \in \prod_{j=1}^n [-\tau_j, 0]$ such that $ASu + AS\tau - b = 0$ and let $x := S(u + \tau)$. Then, $Sx = u + \tau \in \prod_{j=1}^n [0, \tau_j]$, so that $S \in \text{sgn}(x)$, $|x_j| \leq \tau_j$ for all $j = 1, \dots, n$ and

$$0 = AS(u + \tau) - b = Ax - b.$$

Vice versa, assume that $Ax = b$ for some $x \in \prod_{j=1}^n [0, \tau_j]$. Let $S \in \text{sgn}(x)$ and $u := Sx - \tau$. Then, $u \in \prod_{j=1}^n [-\tau_j, 0]$ and

$$0 = Ax - b = A(Su + \tau) - b = ASu + AS\tau - b.$$

□

Recall the definitions of the two sets \mathcal{A} and \mathcal{B} . First, \mathcal{G} is the set of the points $(u, t) \in \mathbb{R}^{n+1}$ such that

$$\begin{cases} u_j = |x_j| - \tau_j & j = 1, \dots, n, \\ t = \|Ax - b\|_2^2, \end{cases} \quad (30)$$

for some $x \in \mathbb{R}^n$. Then,

$$\mathcal{A} = \mathcal{G} + [0, +\infty)^{n+1},$$

that is, $(u, t) \in \mathcal{A}$ if and only if

$$\begin{cases} u_j \geq |x_j| - \tau_j & j = 1, \dots, n, \\ t \geq \|Ax - b\|_2^2, \end{cases} \quad (31)$$

for some $x \in \mathbb{R}^n$. Finally, $(u, t) \in \mathcal{B}$ if and only if $t < p^* = \min_{x_j \leq \tau_j} \|Ax - b\|_2^2$.

We will prove that the equations (31) defining \mathcal{G} can be written in terms of \mathcal{M}_n .

Lemma 3.3. *Let $\tau_1, \dots, \tau_n > 0$ and let \mathcal{G} be as in (31). Then,*

(i) \mathcal{G} is closed;

(ii) $(u, p^*) \in \mathcal{G}$ for some $u \in \mathbb{R}^n$ such that $-\tau_j \leq u_j \leq 0$ for all $j = 1, \dots, n$; moreover, $p^* = \min \left\{ t \in \mathbb{R} : (u, t) \in \mathcal{G}, u_j \leq 0 \ \forall j = 1, \dots, n \right\}$;

(iii) for all $(u, t) \in \mathcal{G}$ there exists $S \in \mathcal{M}_n$ such that $t = \|ASu + (AS\tau - b)\|_2^2$. Viceversa, if $t = \|ASu + (AS\tau - b)\|_2^2$ for some $u \in \mathbb{R}^n$ such that $u_j \geq -\tau_j$ and some $S \in \mathcal{M}_n$, then $(u, t) \in \mathcal{G}$.

Proof. We prove that \mathcal{G} is closed. For, let $(u^k, t^k) \in \mathcal{G}$ converge to $(u, t) \in \mathbb{R}^{n+1}$. We prove that $(u, t) \in \mathcal{G}$. Let $x^k \in \mathbb{R}^n$ be such that (31) is satisfied for (u^k, t^k) . Then, $|x_j^k| = u_j^k - \tau_j \leq u_j + 1 - \tau_j$ for j sufficiently large. In particular, the sequence $\{x^k\}_k$ is bounded and, thus, it converges up to subsequences. Without loss of generality, we may assume that $(x^k)_k$ converges to $x := \lim_{k \rightarrow +\infty} x^k$ in \mathbb{R}^n . Then, for all $j = 1, \dots, n$,

$$|x_j| = \lim_{k \rightarrow +\infty} |x_j^k| = \lim_{k \rightarrow +\infty} u_j^k + \tau_j = u_j + \tau_j$$

and, by continuity,

$$\|Ax - b\|_2^2 = \lim_{k \rightarrow +\infty} \|Ax^k - b\|_2^2 = \lim_{k \rightarrow +\infty} t^k = t.$$

This proves that $(u, t) \in \mathcal{G}$ and, thus, that \mathcal{G} is closed. (ii) follows by (i) and (29).

It remains to check (iii). If $(u, t) \in \mathcal{G}$, there exists $x \in \mathbb{R}^n$ satisfying (31). Let $S \in \mathcal{M}_n$ be such that $|x| = Sx$, where $|x| := (|x_1|, \dots, |x_n|)$. Then, using the fact that $S^{-1} = S$,

$$|x| = u + \tau \implies Sx = (u + \tau) \implies x = S(u + \tau).$$

By the last equation of (31), we have:

$$t = \|Ax - b\|_2^2 = \|ASu + (AS\tau - b)\|_2^2.$$

Viceversa, assume that $t = \|ASu + (AS\tau - b)\|_2^2$ for some $S \in \mathcal{M}_n$ and $u \in \mathbb{R}^n$ is such that $u \succeq -\tau$. Let $x := S(u + \tau)$, then $|x_j| = |u_j + \tau_j| = u_j + \tau_j$ for all $j = 1, \dots, n$ and $t = \|Ax - b\|_2^2$. This proves that $(u, t) \in \mathcal{G}$ and the proof of (iii) is concluded. \square

Lemma 3.4. *Let $u \in \prod_{j=1}^n [-\tau_j, +\infty)$,*

$$h_G(u) := \min_{S \in \mathcal{M}_n} \|ASu + AS\tau - b\|_2^2$$

and

$$g_G(u) := \min_{(u, s) \in \mathcal{G}} s.$$

Then, $h_G(u) = g_G(u)$.

Proof. By Lemma 3.3 (iii), if $(u, s) \in \mathcal{G}$, then $s = \|AS_0u + AS_0\tau - b\|_2^2$ for some $S_0 \in \mathcal{M}_n$. Hence,

$$h_G(u) = \min_{S \in \mathcal{M}_n} \|ASu + AS\tau - b\|_2^2 \leq \|AS_0u + AS_0\tau - b\|_2^2 = s$$

for all s such that $(u, s) \in \mathcal{G}$. Passing to the minimum, we get $h_G(u) \leq g_G(u)$. On the other hand, $(u, h_G(u)) \in \mathcal{G}$ by Lemma 3.3 (iii). Therefore, $g_G(u) \leq h_G(u)$ by definition of g_G . \square

Lemma 3.5. (i) $\mathcal{G} \subseteq \mathcal{A}$;

(ii) \mathcal{A} is closed.

Proof. (i) is obvious. We prove (ii).

Let $(u^k, t^k) \in \mathcal{A}$ be a sequence such that $(u^k, t^k) \xrightarrow[k \rightarrow +\infty]{} (u, t)$ in \mathbb{R}^{n+1} . We need to prove that $(u, t) \in \mathcal{A}$. For all k , let $x^k \in \mathbb{R}^n$ be such that

$$\begin{cases} u_1^k \geq |x_1^k| - \tau_1, \\ \vdots \\ u_n^k \geq |x_n^k| - \tau_n, \\ t^k \geq \|Ax^k - b\|_2^2. \end{cases}$$

The sequence $\{x^k\}_k$ is bounded, in fact for all $j = 1, \dots, n$, $|x_j^k| \leq u_j^k + \tau_j \leq u_j + 1 + \tau_j$ for k sufficiently large. Therefore, up to subsequences, we can assume $x^k \xrightarrow[k \rightarrow +\infty]{} x$ in \mathbb{R}^n . For all $j = 1, \dots, n$,

$$|x_j| = \lim_{k \rightarrow +\infty} |x_j^k| \leq \lim_{k \rightarrow +\infty} u_j^k + \tau_j = u_j + \tau_j.$$

Moreover, by continuity,

$$\|Ax - b\|_2^2 = \lim_{k \rightarrow +\infty} \|Ax^k - b\|_2^2 \leq \lim_{k \rightarrow +\infty} t^k = t.$$

\square

Lemma 3.6. (i) \mathcal{A} is the epigraph of a convex non-negative function $g : \prod_{j=1}^n [-\tau_j, +\infty) \rightarrow \mathbb{R}$ which is continuous in $\prod_{j=1}^n (-\tau_j, +\infty)$;

(ii) $\partial g(0) \neq \emptyset$;

(iii) $g(u) = 0$ if and only if $(u, t) \in \mathcal{A}$ for all $t \geq 0$.

Proof. First, observe that $\mathcal{A} \subseteq \{(u, t) : t \geq 0\}$ since $t \geq \|Ax - b\|_2^2 \geq 0$ for some $x \in \mathbb{R}^n$ whenever $(u, t) \in \mathcal{A}$.

For the sake of completeness, we check that \mathcal{A} is the epigraph of the function

$$g(u) = \min_{(u,s) \in \mathcal{A}} s \quad \left(u \in \prod_j [-\tau_j, +\infty) \right), \quad (32)$$

which is well defined by Lemma 3.5.

By the observation at the beginning of the proof, $g(u) \geq 0$.

Let

$$\text{epi}(g) := \{(u, t) : t \geq g(u)\}$$

be the epigraph of g . If $(u, t) \in \mathcal{A}$, then $t \geq \min_{(u,s) \in \mathcal{A}} s = g(u)$, this means that $(u, t) \in \text{epi}(g)$. On the other hand, if $(u, t) \in \text{epi}(g)$, then $t \geq s$ for some $(u, s) \in \mathcal{A}$. But, if $t \geq s$ (and $(u, s) \in \mathcal{A}$), then $(u, t) \in \mathcal{A}$ as well, since \mathcal{A} contains the vertical upper half-lines having their origins in (u, s) , namely $(u, s) + (\{0\} \times [0, +\infty))$.

This proves that \mathcal{A} is an epigraph. Moreover, g is convex because \mathcal{A} is convex (see [48] Proposition 2.4). The continuity of g on $\prod_j (-\tau_j, +\infty)$ follows from [47], Theorem 10.1. This proves (i).

Moreover, since $\tau_j > 0$ for all $j = 1, \dots, n$, $0 \in \mathbb{R}^n$ is an interior point of $\prod_j [-\tau_j, +\infty)$. Since g is continuous and convex in $\prod_j (-\tau_j, +\infty)$, the subdifferential of g in 0 is non-empty and (ii) follows.

To prove (iii), assume that $g(u) = 0$. Then, $\min_{(u,s) \in \mathcal{A}} s = 0$ implies $(u, 0) \in \mathcal{A}$. Since for all $t \geq 0$, $(u, 0) + (\{0\} \times [0, +\infty)) \in \mathcal{A}$, we have that $(u, t) \in \mathcal{A}$ for all $t \geq 0$. For the converse, assume that $(u, t) \in \mathcal{A}$ for all $t \geq 0$. Then, $(u, 0) \in \mathcal{A}$, so that (by the non-negativity of g) $0 \leq g(u) \leq 0$. This proves the equivalence in (iii). \square

Remark 3.7. As we observed in the general theory situation, $(0, s) \in \mathcal{A}$ if and only if $s \geq p^*$. This tells that $g(0) = p^*$ and $(0, p^*) \in \mathcal{A}$.

We want to prove formally that $g(u)$ defines the boundary $\partial\mathcal{A}$ of \mathcal{A} in a neighborhood of $u = 0$ and, then, find an explicit formula for $g(u)$. Observe that, $\mathcal{A} = \partial\mathcal{A} \cup \overset{\circ}{\mathcal{A}}$, where $\overset{\circ}{\mathcal{A}}$ denotes the topologic interior of \mathcal{A} . Since \mathcal{A} is closed and convex in \mathbb{R}^n , $\overset{\circ}{\mathcal{A}}$ coincides with the algebraic interior of \mathcal{A} , which is defined as follows:

Definition 3.8. Let X be a vector space and $\mathcal{A} \subseteq X$ be a subset. The **algebraic interior** of \mathcal{A} is defined as

$$\text{a-int}(\mathcal{A}) := \{a \in \mathcal{A} : \forall x \in X \exists \varepsilon_x > 0 \text{ s.t. } a + tx \in \mathcal{A} \forall t \in (-\varepsilon_x, \varepsilon_x)\}.$$

Remark 3.9. To clarify the notion of algebraic interior, observe that, for $X \subseteq \mathbb{R}^n$, $x \in \text{a-int}(X)$ if and only if for any fixed direction there exists a segment centered in x having that direction. In general the topologic interior is contained in the algebraic interior, but the reverse inclusion fails, e.g. if $X = \{(x, y, z) \in \mathbb{R}^3 : |z| = x^2 + y^2\} \cup \{(x, y, z) \in \mathbb{R}^3 : z = 0\}$, then $0 \in \text{a-int}(X) \setminus \overset{\circ}{X}$.

Lemma 3.10. Let \mathcal{A} be as in Lemma 3.5. Then,

$$\begin{aligned} \partial\mathcal{A} = & \{(u, t) \in \mathcal{A} : t = g(u), u_j > -\tau_j \forall j = 1, \dots, n\} \cup \\ & \cup \{(u, t) \in \mathcal{A} : u_j = -\tau_j \text{ for some } j = 1, \dots, n\} \end{aligned} \quad (33)$$

and the union is disjoint. Moreover,

$$\{(u, t) \in \mathcal{A} : u_j = -\tau_j \text{ for some } j = 1, \dots, n\} = \{(u, t) \in \partial\mathcal{A} : (u, t + \alpha) \in \partial\mathcal{A} \forall \alpha \geq 0\}.$$

Proof. Observe that the union in (33) is clearly disjoint. We first prove (33).

(\supseteq) Neither of the sets on the RHS of (33) is contained in \mathring{A} . In fact,

- the definition of $g(u)$ implies that for all $\varepsilon > 0$, $-\varepsilon < t < \varepsilon$, $(u, t) \in \mathcal{A}$ if and only if $t \geq 0$, so that $(u, g(u)) \notin \text{a-int}(\mathcal{A}) = \mathring{A}$. This proves that the graph of g in $\prod_j(-\tau_j, +\infty)$ is a subset of $\partial\mathcal{A}$.
- Analogously, assume that $u_j = -\tau_j$ for some $j = 1, \dots, n$, and for all $\varepsilon > 0$ consider the point (u_ε, t) , where $(u_\varepsilon)_l = u_l$ for all $l \neq j$ and $(u_\varepsilon)_j = -\tau_j - \varepsilon$. But g is defined on $\prod_j[-\tau_j, +\infty)$ and \mathcal{A} is its epigraph, hence all the points of \mathcal{A} must be in the form $(u, g(u) + \alpha)$ for some $u \in \prod_j[-\tau_j, +\infty)$, $t = g(u) + \alpha$ ($\alpha \geq 0$), hence $(u_\varepsilon, t) \notin \mathcal{A}$ and this proves that $(u, t) \notin \text{a-int}(\mathcal{A})$.

The fact that $\partial\mathcal{A} = \mathcal{A} \setminus \mathring{A}$ proves the first inclusion.

(\subseteq) We prove that the complementary of the RHS of (33) in \mathbb{R}^{n+1} is contained in \mathring{A} . Let (u, t) be such that $u > -\tau_j$ for all j and $t > g(u)$ (as it is easy to check, these are the conditions for (u, t) to belong to the complementary of the union of the two set at the LHS of (33)).

Let $d := t - g(u) > 0$. Since g is continuous on $\prod_j(-\tau_j, +\infty)$, there exists $\delta > 0$ such that $|g(u) - g(v)| < d/4$ for all $v \in B_\delta(u) := \{w \in \mathbb{R}^n : |w - u| < \delta\}$. In particular, for all $v \in B_\delta(u)$, $g(v) < t - \frac{3}{4}d < t$. Then, $B_\delta(u) \times (t - \frac{3}{4}d, +\infty)$ is all contained in \mathcal{A} (because \mathcal{A} is the epigraph of g) and it is an open neighborhood of (u, t) . Hence, $(u, t) \in \mathring{A} = \mathcal{A} \setminus \partial\mathcal{A}$.

Next, we check the second part of the lemma:

- (\subseteq) assume $(u, t) \in \mathcal{A}$ is such that $u_j = -\tau_j$ for some j . Then, by the first part of this Lemma, $(u, t + \alpha) \in \partial\mathcal{A}$ for all $\alpha \geq 0$, since (33) is a partition of $\partial\mathcal{A}$.
- (\supseteq) Assume that $(u, t + \alpha) \in \partial\mathcal{A}$ for all $\alpha \geq 0$. Then, $(u, t) \in \partial\mathcal{A}$. Assume by contradiction that $u_j > -\tau_j$ for all j . Then, since (33) is a partition of $\partial\mathcal{A}$, $g(u) = t + \alpha$ for all $\alpha \geq 0$, which cannot be the case.

□

Theorem 3.11. *Let \mathcal{A} be as in Lemma 3.5 and g be the function defined as in (32). For $u \in \prod_{j=1}^n[-\tau_j, +\infty)$, $u = (u_1, \dots, u_n)$, let $Q(u) := \prod_{j=1}^n[-\tau_j, u_j]$ and*

$$h(u) := \min_{S \in \mathcal{M}_n, v \in Q(u)} \|AS(v + \tau) - b\|_2^2 = \min_{v \in Q(u)} h_G(v). \quad (34)$$

Then, $h(u) = g(u)$ for all $u \in \prod_j[-\tau_j, +\infty)$.

Proof. We first prove that $g(u) \leq h(u)$. For, it is enough to prove that $(u, h(u)) \in \mathcal{A}$, so that $g(u) \leq h(u)$ would follow by the definition of g . By definition of h , there exist $S_0 \in \mathcal{M}_n$ and $v \in Q(u)$

$$h(u) = \|AS_0v + AS_0\tau - b\|_2^2.$$

By Lemma 3.3 (iii), $(v, h(u)) \in \mathcal{G}$. Since $u_j \geq v_j$ for all $j = 1, \dots, n$, it follows that $(u, h(u)) \in \mathcal{A}$ by definition of \mathcal{A} .

For the converse, since $(u, g(u)) \in \mathcal{A}$, there exists $(v', t) \in \mathcal{G}$ such that $v'_j \leq u_j$ for all $j = 1, \dots, n$ and $g(u) \geq t$. In particular, $v' \in Q(u)$. By Lemma 3.3 (iii), $t = \|AS_1v' + AS_1\tau - b\|_2^2$ for some $S_1 \in \mathcal{M}_n$. Therefore,

$$g(u) \geq \|AS_1v' + AS_1\tau - b\|_2^2 \geq \min_{S \in \mathcal{M}_n, v \in Q(u)} \|ASv + AS\tau - b\|_2^2 = h(u).$$

This concludes the proof. □

Even if $g = h$, in what follows we still distinguish h and g when we want to stress the explicit definitions of both. Namely, we write $g(u)$ when we refer to $\min_{(u,s) \in \mathcal{A}} s$ and $h(u)$ when we refer to (34).

Corollary 3.12. *Under the same notation as above, for all $S \in \mathcal{M}_n$,*

$$g(u) = \min_{-u-\tau \preceq v \preceq u+\tau} \|Av - b\|_2^2. \quad (35)$$

Proof. Using the second expression in (34),

$$g(u) = \min_{S \in \mathcal{M}_n} \min_{v \in Q(u)} \|AS(v + \tau) - b\|_2^2.$$

But,

$$f_S(v) = \|AS(v + \tau) - b\|_2^2 = f(S(v + \tau)),$$

for $f(v) = \|Av - b\|_2^2$, that gives

$$\min_{-\tau \preceq v \preceq u} f_S(v) = \min_{v \in Q(u)} f(S(v + \tau)) = \min_{v \in S(Q(u) + \tau)} \|Av - b\|_2^2,$$

so that

$$\min_{S \in \mathcal{M}_n} \min_{-\tau \preceq v \preceq u} f_S(v) = \min_{\cup_{S \in \mathcal{M}_n} S(Q(u) + \tau)} \|Av - b\|_2^2$$

and the assertion follows by observing that

$$\bigcup_{S \in \mathcal{M}_n} S(Q(u) + \tau) = \{v \in \mathbb{R}^n : -u - \tau \preceq v \preceq u + \tau\}.$$

□

3.3 A result under conditions on the gradient of $\|Ax - b\|_2^2$

In general, the geometry of \mathcal{A} is so complicated that expressing g explicitly may turn into a tough task. Nevertheless, it is obvious that if u is itself one of the minimizers of (34), then $g(u) = h_G(u) = \min_{S \in \mathcal{M}_n} \|ASu + AS\tau - b\|_2^2$. So, under further assumptions on $\nabla(\|ASu - b\|_2^2)$ granting the equality $g(u) = h_G(u)$ holds in a neighborhood of 0, we can compute explicitly the optimal parameters.

Theorem 3.13. *Let $f(v) = \|Av - b\|_2^2$ and assume that for all $k = 1, \dots, n$ the condition*

$$\sum_{j=1}^n u_j \langle a_{*,k}, a_{*,j} \rangle \leq \langle b, a_{*,k} \rangle \quad (-\tau \preceq u \preceq \tau) \quad (36)$$

holds. Then, $g(u) = f(u + \tau)$ for all $u \in Q(0)$ and $\lambda^\# = A^T(b - A\tau)$ is a set of optimal parameters for problem (23).

Proof. The set of conditions (36) is equivalent to $(Au - b)^T A \preceq b^T A$ for all $-\tau \preceq u \preceq \tau$, that is $\nabla f(u) \preceq 0$ for $-\tau \preceq u \preceq \tau$. We prove that, under this further condition, $g(u) = f(u + \tau)$ for all $u \in Q(0)$. Let $u \in Q(0)$ and $\mathbf{n} \succ 0$ be a unit vector. For all $t \in \mathbb{R}$, define

$$\begin{aligned} f_{\mathbf{n}}(t) &:= f(u + \tau + t\mathbf{n}) = \|A(u + \tau + t\mathbf{n}) - b\|_2^2 \\ &= \|\mathbf{A}\mathbf{n}\|_2^2 t^2 + 2\langle A(u + \tau) - b, \mathbf{A}\mathbf{n} \rangle t + \|A(u + \tau) - b\|_2^2, \end{aligned}$$

which is the restriction of f to the line $\{u + t\mathbf{n} : t \in \mathbb{R}\}$. If $\mathbf{n} \in \ker(A)$, then $f_{\mathbf{n}} \equiv 0$ and it has a global minimum in $t = 0$. Assume $\mathbf{n} \notin \ker(A)$. The intersection of this line with $\{-\tau \preceq v \preceq \tau\}$ is contained in $(-\infty, 0]$. If we prove that, for all $\mathbf{n} \succ 0$, $f_{\mathbf{n}}$ has a constrained minimum in $t = 0$, we get the first assertion. For, it's enough to observe that

$$f'_{\mathbf{n}}(0) = \nabla f(u + \tau) \cdot \mathbf{n} \leq 0,$$

because if $u \in Q(0)$, then $\{-u - \tau \preceq v \preceq u + \tau\} \subseteq \{-\tau \preceq v \preceq \tau\}$. This proves that $g(u) = f(u + \tau)$ for all $u \in Q(0)$. In particular,

$$-\nabla g(0) = -\nabla f(\tau) = (b - A\tau)^T A \succeq 0$$

is a set of optimal parameters for (23). \square

Remark 3.14. It is not difficult to generalize Theorem 3.13 a bit further. If the hyperparallelogram $\{-\tau \preceq u \preceq \tau\}$ is all contained in the region $\{u \in \mathbb{R}^n : S\nabla f(u) \preceq 0\}$ for some $S \in \mathcal{M}_n$, then $g(u) = f(S(u + \tau))$ for all $u \in Q(0)$ and

$$\lambda^\# = -\nabla g(0) = -S\nabla f(S(u + \tau))^T$$

defines a vector of optimal parameters for (23). The proof goes exactly as in Theorem 3.13.

3.4 Decoupling the variables

In this subsection, we focus on the situation in which $A^T A$ is a diagonal matrix. Since

$$A^T A = \begin{pmatrix} \|a_{*,1}\|_2^2 & \langle a_{*,1}, a_{*,2} \rangle & \cdots & \langle a_{*,1}, a_{*,n} \rangle \\ \langle a_{*,2}, a_{*,1} \rangle & \|a_{*,2}\|_2^2 & \cdots & \langle a_{*,2}, a_{*,n} \rangle \\ \vdots & \vdots & \ddots & \vdots \\ \langle a_{*,n}, a_{*,1} \rangle & \langle a_{*,n}, a_{*,2} \rangle & \cdots & \|a_{*,n}\|_2^2 \end{pmatrix}$$

and the rank of $A^T A$ is equal to that of A , it follows that in this case

$$A^T A = \text{diag}(\|a_{*,1}\|_2^2, \dots, \|a_{*,n}\|_2^2). \quad (37)$$

Remark 3.15. If $m \leq n$ and $A^T A$ is diagonal, $n - m$ of the norms in (37) above vanish. In this case, we assume that $a_{*,m+1} = \dots = a_{*,n} = 0$, so that A can be written in terms of its columns as

$$A = (A' | 0_{m \times (n-m)}),$$

where $A' = (a_{*,1} | \dots | a_{*,m}) \in GL(m, \mathbb{R})$. Observe that

$$\|Ax - b\|_2^2 = \|A'x' - b\|_2^2,$$

where $x' = (x_1, \dots, x_m)^T$, so that $x^\#$ is a minimizer of (23) if and only if $(x^\#)' = (x_1^\#, \dots, x_m^\#)$ is a minimizer of

$$\begin{cases} \min_{y \in \mathbb{R}^m} \|A'y - b\|_2^2, \\ |y_j| \leq \tau_j \end{cases} \quad j = 1, \dots, m, \quad (38)$$

under the further condition that the remaining coordinates of x vanish.

For this reason, for the rest of this subsection, we focus on (38), both for the cases $n \leq m$ and $m \leq n$, and provide the optimal parameters.

Remark 3.16. We point out that in this situation the optimal parameters can be computed directly from Proposition 3.1. Indeed, under the *orthogonality* assumption on A , the target function in problem (38) becomes

$$\sum_{j=1}^m (\|a_{*,j}\|_2^2 y_j^2 - 2y \langle a_{*,j}, n \rangle y_j) + \|b\|_2^2.$$

Since the variables of all the addenda are decoupled, and the addenda are non-negative,

$$\min_y \sum_{j=1}^m (\|a_{*,j}\|_2^2 y_j^2 - 2y \langle a_{*,j}, n \rangle y_j) + \|b\|_2^2 = \sum_{j=1}^m \min_{y_j} \left(\|a_{*,j}\|_2^2 y_j^2 - 2y \langle a_{*,j}, n \rangle y_j + \frac{\|b\|_2^2}{m} \right)$$

and a minimizer of (38) also minimizes

$$\begin{cases} \min_y \|a_{*,j}\|_2^2 y_j^2 - 2y \langle a_{*,j}, n \rangle y_j + \frac{\|b\|_2^2}{m}, \\ |y_j| \leq \tau_j \end{cases}$$

for all $j = 1, \dots, m$. In other words, it is enough to treat (38) as m 1-dimensional constrained minimization problems. However, our interest is testing the tools presented in the previous paragraph, computing the function g and the separating hyperplane.

To exhibit a vector of optimal parameters, we start by the set

$$\mathcal{G} := \{(u, t) \in \mathbb{R}^{m+1} : u_j = |y_j| - \tau_j \ (j = 1, \dots, m), t = \|A'y - b\|_2^2 \text{ for some } y \in \mathbb{R}^m\}.$$

By Lemma 3.3 (iii), $(u, t) \in \mathcal{G}$ if and only if $u \succeq -\tau$ and $t = \|A'S(u + \tau) - b\|_2^2$ for some $S \in \mathcal{M}_m$. Let $f_S(u) = \|A'S(u + \tau) - b\|_2^2$ and observe that

$$f_S(u) = \sum_{j=1}^m \|a_{*,j}\|_2^2 (u_j + \tau_j)^2 - 2 \sum_{j=1}^m s_{jj} \langle b, a_{*,j} \rangle (u_j + \tau_j) + \|b\|_2^2.$$

The functions f_S are the equivalent of the parabolas in the 1-dimensional case and they describe elliptic paraboloids. As it clear by Subsection 3.1, we need to understand what is $h_G(u) := \min_{S \in \mathcal{M}_m} f_S(u)$. Observe that for all $S \in \mathcal{M}_m$,

$$f_S(u) \geq \sum_{j=1}^m \|a_{*,j}\|_2^2 (u_j + \tau_j)^2 - 2 \sum_{j=1}^m |\langle b, a_{*,j} \rangle| (u_j + \tau_j) + \|b\|_2^2 = f_{S_\beta}(u), \quad (39)$$

where $S_\beta = (s_j^\beta)_{j=1}^m \in \mathcal{M}_m$ is a diagonal matrix such that $s_j^\beta \langle b, a_{*,j} \rangle \geq 0$.

Lemma 3.17. *Under the notation and the assumptions of this subsection,*

$$h_G(u) = f_{S_\beta}(u) = \sum_{j=1}^m \|a_{*,j}\|_2^2 (u_j + \tau_j)^2 - 2 \sum_{j=1}^m |\langle b, a_{*,j} \rangle| (u_j + \tau_j) + \|b\|_2^2.$$

h_G defines an elliptic paraboloid whose vertex $V = (c, 0) \in \mathbb{R}^{m+1}$ is characterized both by $c = -\tau + S_\beta(A')^{-1}b$ and

$$c_j = -\tau_j + \frac{|\langle b, a_{*,j} \rangle|}{\|a_{*,j}\|_2^2}$$

($j = 1, \dots, m$). Moreover,

$$h_G(u) = \sum_{j=1}^m \|a_{*,j}\|_2^2 (u_j - c_j)^2. \quad (40)$$

Proof. We already proved the first part of the Lemma. We only need to compute the vertex of f_{S_β} . For, observe that the minimum of f_{S_β} is $(c, 0)$, where c satisfies $f_S(c) = 0$. This equation is satisfied if and only if $c = -\tau + S_\beta(A')^{-1}b$. Moreover, the minimum of f_{S_β} is also characterized by $\nabla f_{S_\beta}(c) = 0$, that is

$$c_j + \tau_j - \frac{|\langle b, a_{*,j} \rangle|}{\|a_{*,j}\|_2^2} = 0$$

($j = 1, \dots, m$). Finally, using the first characterization of c ,

$$\begin{aligned} h_G(u) &= \|AS_\beta(u + \tau) - b\|_2^2 = \|AS_\beta(u - c) + AS_\beta(c + \tau) - b\|_2^2 = \|AS_\beta(u - c)\|_2^2 = \\ &= \sum_{j=1}^m \|a_{*,j}\|_2^2 (u_j - c_j)^2. \end{aligned}$$

This concludes the proof. \square

In order to compute the optimal parameters for the decoupled problem, we observe that $\mathcal{A} + [0, +\infty)^{m+1}$ is the epigraph of the function $g(u)$ whose first properties are proved in Lemma 3.6. Hence, this function describes the lower boundary of \mathcal{A} , that is the part of \mathcal{A} we need to compute a separating hyperplane. By (34), $g(u) = \min_{v \in Q(u)} h_G(v)$, where $Q(u) = \prod_{j=1}^m [-\tau_j, u_j]$.

Theorem 3.18. *Under the notation and the assumptions of this subsection,*

$$g(u) = h_G(Pu),$$

where $P : \prod_{j=1}^m [-\tau_j, +\infty) \rightarrow Q(c)$ is the projection defined for all $u \in \prod_{j=1}^m [-\tau_j, +\infty)$ by

$$(Pu)_j = \begin{cases} u_j & \text{if } -\tau_j \leq u_j \leq c_j, \\ c_j & \text{if } u_j > c_j \end{cases} = \min\{c_j, u_j\} \quad (41)$$

($j = 1, \dots, m$). Explicitly, under the assumptions of this subsection,

$$g(u) = \sum_{j=1}^m \|a_{*,j}\|_2^2 (u_j - c_j)^2 \chi_{[-\tau_j, c_j]}(u_j). \quad (42)$$

In particular, $g \in \mathcal{C}^1(\prod_{j=1}^m (-\tau_j, +\infty))$ with

$$\frac{\partial g}{\partial u_j}(u) = 2 \|a_{*,j}\|_2^2 (u_j - c_j) \chi_{[-\tau_j, c_j]}(u_j) \quad (43)$$

for all $u \in \prod_{j=1}^m (-\tau_j, +\infty)$.

Proof. Obviously, P is a projection of $\prod_{j=1}^m [-\tau_j, +\infty)$ onto $Q(c)$. For all $j = 1, \dots, m$,

$$\operatorname{argmin}_{-\tau_j \leq v_j \leq u_j} \|a_{*,j}\|_2^2 (v_j - c_j)^2 = \begin{cases} u_j & \text{if } -\tau_j \leq u_j \leq c_j, \\ c_j & \text{otherwise} \end{cases} = (Pu)_j.$$

Hence,

$$\begin{aligned} g(u) &= \min_{v \in Q(u)} h_G(v) = \sum_{j=1}^m \min_{-\tau_j \leq v_j \leq u_j} \|a_{*,j}\|_2^2 (v_j - c_j)^2 = \\ &= \sum_{j=1}^m \|a_{*,j}\|_2^2 ((Pu)_j - c_j)^2 = h_G(Pu). \end{aligned}$$

The explicit definition of Pu gives (42) and (43). The differentiability and formula (43) are obvious by the expression (42) of g . \square

Remark 3.19. As a consequence of Theorem 3.18,

$$p^* = g(0) = \sum_{j=1}^m \|a_{*,j}\|_2^2 \left(-\tau_j + \frac{|\langle b, a_{*,j} \rangle|}{\|a_{*,j}\|_2^2} \right)^2 \chi_{[-\tau_j, c_j]}(0).$$

Then, observe that

$$-\tau_j \leq 0 \leq -\tau_j + \frac{|\langle b, a_{*,j} \rangle|}{\|a_{*,j}\|_2^2} \iff 0 \leq \tau_j \leq \frac{|\langle b, a_{*,j} \rangle|}{\|a_{*,j}\|_2^2}, \quad (44)$$

so that

$$p^* = \sum_{j=1}^m \|a_{*,j}\|_2^2 \left(-\tau_j + \frac{|\langle b, a_{*,j} \rangle|}{\|a_{*,j}\|_2^2} \right)^2 \chi_{\left[0, \frac{|\langle b, a_{*,j} \rangle|}{\|a_{*,j}\|_2^2}\right]}(\tau_j).$$

Theorem 3.20. Under the notation of this subsection, the vector $\lambda^\# \in [0, +\infty)^m$ given by

$$\lambda_j^\# = 2 \|a_{*,j}\|_2^2 \left(\frac{|\langle b, a_{*,j} \rangle|}{\|a_{*,j}\|_2^2} - \tau_j \right) \chi_{\left[0, \frac{|\langle b, a_{*,j} \rangle|}{\|a_{*,j}\|_2^2}\right]}(\tau_j)$$

defines a vector of optimal parameters for (38).

Proof. We apply (43) to $u = 0$ and use (44). Namely,

$$t = p^* + \langle \nabla g(0), u \rangle$$

is the tangent hyperplane of g in $u = 0$, which is also the hyperplane that separates \mathcal{A} and \mathcal{B} . The direction of this hyperplane is $(\nabla g(0), -1)$, so that

$$\lambda^\# = -\nabla g(0),$$

i.e. the assertion. □

Remark 3.21. As far as the original problem (23) with $m \leq n$ is concerned, we get the optimal parameters for free by Theorem 3.20 simply observing that if $A = (a_{*1} | \dots | a_{*m} | 0 | \dots | 0) \in \mathbb{R}^{m \times n}$, $A' = (a_{*1} | \dots | a_{*m})$ and $x = (x', x'') \in \mathbb{R}^m \times \mathbb{R}^{n-m}$, then

$$\begin{aligned} \min_{x \in \mathbb{R}^n} \|Ax - b\|_2^2 + \sum_{j=1}^n \lambda_j (|x_j| - \tau_j) &= \min_{x' \in \mathbb{R}^m} \|A'x' - b\|_2^2 + \sum_{j=1}^m \lambda_j (|x'_j| - \tau_j) + \\ &+ \underbrace{\min_{x'' \in \mathbb{R}^{n-m}} \sum_{j=m+1}^n \lambda_j (|x_j| - \tau_j)}_{= -\sum_{j=m+1}^n \lambda_j \tau_j}, \end{aligned}$$

so that, if $\lambda^\# \in \mathbb{R}^m$ defines a vector of optimal parameters for (38), then $(\lambda^\# | 0) \in \mathbb{R}^m \times \mathbb{R}^{n-m}$ defines a vector of optimal parameters for (23).

3.5 Explicit solution

The conditions $|x_j| \leq \tau_j$ are equivalent to $x_j^2 \leq \tau_j^2$. Under this point of view, (23) can be restated as

$$\begin{cases} \min_x \|Ax - b\|_2^2, \\ x_j^2 \leq \tau_j^2, \end{cases} \quad (45)$$

that can be interpreted as a space-variant Tikhonov problem. Assume that $\lambda^\#$ is a vector of optimal parameters for (23) or, equivalently, for (45). We are interested in computing

$$x^\# = \arg \min_x L(x, \lambda),$$

where L is the Lagrange function associated to (45), i.e.

$$L(x, \lambda^\#) = \|Ax - b\|_2^2 + \sum_{j=1}^n \lambda_j^\# (x_j^2 - \tau_j^2).$$

Since $L \in \mathcal{C}^\infty(\mathbb{R}^n)$ and it is convex, they satisfy $\nabla L(x, \lambda^\#) = 0$, that is

$$(A^T A + \Delta_\lambda)x = A^T b,$$

where $\Delta_\lambda = \text{diag}(\lambda_1^\#, \dots, \lambda_n^\#)$. Hence, $x^\#$ satisfies

$$(A^T A + \Delta_\lambda)x^\# = A^T b, \quad (46)$$

that is, $x^\# \in (A^T A + \Delta_\lambda)^{-1} A^T b$.

Remark 3.22. Another way to compute the optimal parameters associated to (23), or equivalently to (45), could be by means of the strong duality condition, namely using

$$\lambda^\# = \arg \max_{\lambda \geq 0} \min_x L(x, \lambda).$$

However, we stress that the explicit value of $\min_x L(x, \lambda)$ is still hard to compute since the implicit relation (46) satisfied by $x^\#$ cannot be made explicit by means of Dini's theorem.

4 Considerations and conclusions

4.1 Applications

Despite the apparently heavy assumptions on A , Theorem 3.20 has itself interesting applications. For instance, it can be applied to denoising problems, where $A = I_{n \times n}$, i.e. problems in the form

$$\begin{cases} \min_{x \in \mathbb{R}^n} \|x - b\|_2^2, \\ |x_j| \leq \tau_j \end{cases} \quad j = 1, \dots, n. \quad (47)$$

By Theorem 3.20, $\lambda^\# = (\lambda_j^\#)_{j=1}^n$ is a vector of optimal parameters for (47), where

$$\lambda_j^\# = 2(|b_j| - \tau_j)^+, \quad (48)$$

where z^+ denotes the positive part of $z \in \mathbb{R}^m$, i.e. $z^+ = \max\{z, 0\}$.

We can also apply Theorem 3.20 to the discrete Fourier transform, i.e. given a noisy fully-sampled signal $b \in \mathbb{C}^n$, we want to find a vector $z \in \mathbb{C}^n$ such that $\|\Phi z - b\|_2^2$ is minimized under the constrains $|z_j| \leq \tau_j$, where $\Phi \in \mathbb{C}^{n \times n}$ denotes the (complex) DFT matrix. Since $\Phi^* \Phi = I_{n \times n}$, we can apply Theorem 3.20 to deduce that a set of optimal parameters for this problem is

$$\lambda_j^\# = 2\left(|\langle b, \phi_{*,j} \rangle| - \tau_j\right)^+,$$

($j = 1, \dots, n$), being $\phi_{*,j}$ the j -th column of Φ .

The question that naturally arises in the applications is whether the dependence of $\lambda_1, \dots, \lambda_n$ on τ_1, \dots, τ_n can be a critical issue in the applicability of the theory. Indeed, τ_1, \dots, τ_n are upper bounds for $|x_1|, \dots, |x_n|$ respectively, which are not available in the practice. However, whenever it is possible to estimate these local upper bounds, our result may lead to high-quality imaging performances. For instance, for denoising, (48) may be approximated by replacing τ_1, \dots, τ_n with the voxel values obtained by applying a Gaussian filter (or other types of filtering) to the noisy image. This opens the question of which filtering technique could lead to optimal approximations of the τ_1, \dots, τ_n depending on the field of research in which (23) can be implemented. We intend to investigate this topic in the immediate future.

4.2 Open problems

As we pointed out, as long as $A^T A$ is not a diagonal matrix, the geometry of the sets \mathcal{G} and \mathcal{A} become more involved, along with the possible casuistry. However, the general case in which $A^T A$ is not diagonal would be of great importance in applications. Indeed, the behavior of optimal parameters for the space-variant LASSO problem (8) in terms of voxel-wise estimates would provide a way to control the optimal parameters via estimates of the τ_j . This led us to formulate a conjecture that we are going to verify experimentally in our future works in an MRI framework. We conjecture that in general, the optimal parameters of problem (23) are given by

$$\lambda^\# = (-S \nabla f(S\tau))^+ \quad (49)$$

for a signature matrix $S \in \mathcal{M}_n$ that we believe to be related to the signatures of all the possible solutions of $Ax = b$. The reason why we believe that (49) could be correct is that, for the right choice of S and $u \in Q(0)$, $g(u)$ shall be either $\|AS(u + \tau) - b\|_2^2$, for some S , or $\|AS(v(u) + \tau) - b\|_2^2$, where $v(u)$ is the projection of $\{Ax = b\}$ onto $\{-\tau - u \preceq v \preceq u + \tau\}$. The conjecture on S being one of the signatures of $\{Ax = b\}$ follows the expression $g(u) = \min_{-u-\tau \preceq v \preceq u+\tau} \|Av - b\|_2^2$: intuitively, the closest of the points of $\{Ax = b\}$ to $\{-u - \tau \preceq v \preceq u + \tau\}$ is the one realizing the minimum, i.e. its

projection onto the hyperparallelogram, which is symmetric with respect to the origin. In conclusion, we expect that $\nabla f(v(0)) \in \partial g(S\tau)$ holds by construction of \mathcal{A} .

Another open problem is whether it is possible to apply the same procedure to compute the optimal parameter for (1). Clearly, the corresponding sets \mathcal{G} and \mathcal{A} lie in \mathbb{R}^2 so that $g : \mathbb{R} \rightarrow \mathbb{R}$. Despite this simplifying fact, the set \mathcal{G} is characterized by

$$\begin{cases} u = s(x)^T x - \tau, \\ t = \|Ax - b\|_2^2 \end{cases} \quad \text{for some } x \in \mathbb{R}^n,$$

where $s(x) \in \mathbb{R}^n$ is the vector such that $(s(x))_j = \text{sgn}(x_j)$ and, in this case, u and x belong to different spaces and a closed form for $t = t(u)$ is even more difficult to provide.

Finally, we stress that it would be important to generalize (23) up to consider different inner products on \mathbb{R}^n . Namely, this is the situation that occurs in MRI when the undersampling pattern is non-cartesian. Problem (8) in this case becomes

$$\min_x \|Ax - b\|_W^2 + \sum_j \lambda_j |x_j|,$$

where

$$\|x\|_W = x^T W^T W x \quad (x \in \mathbb{R}^n),$$

for a definite positive diagonal matrix W . Since this topic falls beyond the purpose of this work, we limit ourselves to mention the very mathematical reason why the weighted norm shall definitely replace the Euclidean norm over \mathbb{R}^n when sampling is not performed on a cartesian grid. Indeed, non-cartesian sampling patterns require appropriate discretizations of the Fourier transform integral. Roughly speaking,

$$\hat{f}(\xi) \approx \sum_j f(x_j) e^{-2\pi i \xi \cdot x_j} \Delta x_j = \langle f, e^{2\pi i \xi \cdot} \rangle_W,$$

where Δx_j is the Lebesgue measure of an adequate neighborhood of x_j , weighting the contribution of the sample x_j , and W is the diagonal matrix whose entries are $\sqrt{\Delta x_j}$. The inversion formula of the Fourier transform shall be modified accordingly. For instance, if the sampling follows a spiral trajectory, Δx_j shall be bigger the further x_j is from the origin, since this value serves as an average of f on a portion of sphere that is larger as x_j is far from the origin. All the above-mentioned problems will be object of our future investigations.

Acknowledgements

Financial support for this work has been provided by University of Bologna and the Institute of Systems Engineering at HES-SO Valais-Wallis. We are thankful to the discussion with Prof. Nicola Arcozzi, Dr. Luca Calatroni, Dr. Fabian Pedregosa and Mr. Pasquale Sirignano, who contributed to highly improve the quality of the manuscript. We are also very grateful to Prof. Micah M. Murray for his support.

Authors contributions

B.F. conceptualised the problem. B.F. and G.G. framed the hypothesis to be proved and tested. G.G. developed the proofs under B.F.'s supervision and drafted a first version of the manuscript, and B.F. shaped the article in its final version. The idea of adding the Appendix is due to B.M., who has perfected the proof of the existence provided by G.G. and contributed to the draft of the manuscript.

Appendix

Since we did not find one in the existing literature, we provide a formal proof of the existence of the minimizer of the generalized LASSO problem

$$\arg \min_{x \in \mathbb{R}^n} \|Ax - b\|_2^2 + \lambda \|\Phi x\|_1, \quad (50)$$

where $b \in \mathbb{R}^m$, $A \in \mathbb{R}^{m \times n}$ and $\Phi \in \mathbb{R}^{N \times n}$.

We state the result in the general framework of finite-dimensional vector spaces: we denote by X, Y, Z three finite-dimension real vector spaces. We denote by $\langle \cdot, \cdot \rangle_X$ an inner product on X and with $\|\cdot\|_X$ the induced norm. Analogous notation will be used for Y , whereas we set

$$\|z\|_p = \left(\sum_{j=1}^{\dim(Z)} |z_j|^p \right)^{1/p}, \quad z \in Z$$

for $0 < p \leq \infty$. Recall that $\|\cdot\|_p$ is a Banach quasi-norm (meaning that there exists $C_p \geq 1$ such that $\|x + y\|_p \leq C_p(\|x\|_p + \|y\|_p)$ for all $x, y \in Z$) replaces the triangular inequality for $0 < p < 1$, and it is a norm for $1 \leq p \leq \infty$.

Then, for given $\lambda \geq 0$, $b \in Y$, $A : X \rightarrow Y$ and $\Phi : X \rightarrow Z$ linear, we define for all $x \in X$

$$f(x) = \|Ax - b\|_Y^2 + \lambda \|\Phi x\|_p.$$

Theorem. *For all $0 < p \leq \infty$, $\lambda \geq 0$ there exists $x^\# \in X$ such that*

$$\inf_{x \in X} f(x) = f(x^\#).$$

In particular, the generalized LASSO problem (50) has at least one solution.

Proof. Clearly, the function $f(x) = \|b\|_Y^2 + \lambda \|\Phi x\|_p$ attains its minimum in $x^\# = 0$. Hence, we may assume

$$Im(A) = \{Ax : x \in X\} \neq \{0\}.$$

Since $Im(A)$ is a vector subspace of Y , for all $b \in Y$ there exists a unique $y^\# \in Im(A)$ such that $\inf_{y \in Y} \|y - b\|_Y^2 = \|y^\# - b\|_Y^2$. By definition, $y^\# = Ax^\#$ for some $x^\# \in X$. Hence,

$$\inf_{x \in X} \|Ax - b\|_Y^2 = \inf_{y \in Im(A)} \|y - b\|_Y^2 = \min_{y \in Y} \|y - b\|_2^2 = \|Ax^\# - b\|_Y^2$$

and the assertion follows also for the case in which $\lambda = 0$ or $Im(B) = \{\Phi x : x \in X\} = \{0\}$. We will thereby assume that $Im(A) \neq \{0\}$, $Im(B) \neq \{0\}$ and $\lambda > 0$. Let $L := \ker(A) \cap \ker(B) = \{x \in X : Ax = 0, \Phi x = 0\}$ and denote the closed ball of X of center 0 and radius $r > 0$ by $B_X(0, r) = \{x \in X : \|x\|_X \leq r\}$. The rest of the proof is divided into three graded steps.

Step 1. We prove that if $L = \{0\}$, then $\lim_{\|x\|_X \rightarrow +\infty} f(x) = +\infty$.

By convexity of $\|\cdot\|_Y$,

$$\|y_1\|_Y^2 \leq 2(\|y_1 - y_2\|_Y^2 + \|y_2\|_Y^2)$$

for all $y_1, y_2 \in Y$. Therefore,

$$\|Ax - b\|_Y^2 \geq \frac{1}{2} \|Ax\|_Y^2 - \|b\|_Y^2,$$

so that

$$\|Ax - b\|_Y^2 + \lambda \|\Phi x\|_p \geq \frac{1}{2} \|Ax\|_Y^2 + \lambda \|\Phi x\|_p - \|b\|_Y^2.$$

Let

$$\mathbb{S}_X := \{x \in X : \|x\|_X = 1\}$$

denote the unit sphere of X and set $\eta := \min_{x \in \mathbb{S}_X} \frac{1}{2} \|Ax\|_Y^2 + \lambda \|\Phi x\|_p$. If $\eta = 0$, then

$$\min_{x \in \mathbb{S}_X} \frac{1}{2} \|Ax\|_Y^2 + \lambda \|\Phi x\|_p = 0,$$

together with the assumptions on $Im(A)$, $Im(B)$ and λ , yields to the existence of $x^\# \in \mathbb{S}_X$ such that $\frac{1}{2} \|Ax^\#\|_Y^2 + \lambda \|\Phi x^\#\|_p = 0$. But $\|\cdot\|_Y$ and $\|\cdot\|_p$ are (quasi-)norms, so $x^\# = 0 \notin \mathbb{S}_X$. This is a contradiction. Hence, $\eta > 0$.

Next, for $\|x\|_X > 1$, we have

$$\begin{aligned} \frac{1}{2} \|Ax\|_Y^2 + \lambda \|\Phi x\|_p - \|b\|_Y^2 &= \frac{1}{2} \|x\|_X^2 \left\| A \frac{x}{\|x\|_X} \right\|_Y^2 + \lambda \|x\|_X \left\| \Phi \frac{x}{\|x\|_X} \right\|_p - \|b\|_Y^2 \\ &> \|x\|_X \left(\frac{1}{2} \left\| A \frac{x}{\|x\|_X} \right\|_Y^2 + \lambda \left\| \Phi \frac{x}{\|x\|_X} \right\|_p \right) - \|b\|_Y^2 \\ &\geq \eta \|x\|_X - \|b\|_Y^2. \end{aligned}$$

Therefore, for all $x \in X$ such that $\|x\|_X > 1$,

$$f(x) = \|Ax - b\|_Y^2 + \lambda \|\Phi x\|_p > \eta \|x\|_X + \|b\|_Y^2$$

and the assertion follows, since $\eta > 0$ implies that the right hand-side goes to $+\infty$ as $\|x\|_X \rightarrow +\infty$.

Step 2. We prove the assertion for $L = \{0\}$.

Let $m := \inf_{x \in X} f(x)$. By Step 1, there exist $R > 0$ such that $f(x) > m + 1$ for $\|x\|_X > R$. $B_X(0, R)$ is compact and convex, and $\inf_{x \in X} f(x) = \inf_{x \in B_X(0, R)} f(x)$ by definition of R . Let $(x_j)_j \subseteq B_X(0, R)$ be a minimizing sequence. By compactness, it admits a converging subsequence and, without loss of generality, we may assume that $\lim_{j \rightarrow +\infty} x_j = x^\# \in B_X(0, R)$. By continuity, $f(x^\#) = \lim_{j \rightarrow +\infty} f(x_j) = m$.

Step 3. We prove the assertion for $L \neq \{0\}$.

Recall that $X = L \oplus L^\perp$, where the orthogonality is defined with respect to the inner product $\langle \cdot, \cdot \rangle_X$. By definition of direct sum, for all $x \in X$ there exist unique $x_1 \in L$ and $x_2 \in L^\perp$ such that $x = x_1 + x_2$. Observe that since $x_1 \in L$,

$$f(x) = \|Ax_2 - b\|_Y^2 + \lambda \|\Phi x_2\|_p = f(x_2).$$

In particular,

$$\inf_{x \in X} f(x) = \inf_{x \in L^\perp} f(x).$$

The restrictions of A and Φ to L^\perp are linear mappings between vector spaces. We denote them with $A|_{L^\perp} : L^\perp \rightarrow Y$ and $\Phi|_{L^\perp} : L^\perp \rightarrow Z$ respectively and set $f|_{L^\perp} : L^\perp \rightarrow \mathbb{R}$ as the restriction of f to L^\perp . Obviously,

$$f|_{L^\perp}(x) = \|A|_{L^\perp}x - b\|_Y^2 + \lambda \|\Phi|_{L^\perp}x\|_p = f(x)$$

for all $x \in L^\perp$, so that

$$\inf_{x \in X} f(x) = \inf_{x \in L^\perp} f(x) = \inf_{x \in L^\perp} f|_{L^\perp}(x).$$

Obviously,

$$L^\perp := \ker(A|_{L^\perp}) \cap \ker(\Phi|_{L^\perp}) = \ker(A) \cap \ker(B) \cap L^\perp = L \cap L^\perp = \{0\}.$$

Therefore, by Step 2, it follows that there exists $x^\# \in L^\perp$ such that

$$\inf_{x \in L^\perp} f|_{L^\perp}(x) = f|_{L^\perp}(x^\#).$$

This implies that

$$\inf_{x \in X} f(x) = f|_{L^\perp}(x^\#) = f(x^\#),$$

since $x^\# \in L^\perp$. □

References

- [1] S. Arridge, P. Maass, O. Öktem, C. Schönlieb (2019). *Solving inverse problems using data-driven models*. *Acta Numerica*, 28, 1-174. doi:10.1017/S0962492919000059.

- [2] A. Berk, Y. Plan, Ö. Yilmaz. *Parameter instability regimes in sparse proximal denoising programs*. In 2019 13th International conference on Sampling Theory and Applications (SampTA) (pp. 1-5). IEEE (2019, July).
- [3] A. Berk, S. Brugiapaglia, T. Hoheisel. *LASSO reloaded: a variational analysis perspective with applications to compressed sensing*. arXiv preprint arXiv:2205.06872.
- [4] A. Berk, Y. Plan, Ö. Yilmaz. *Sensitivity of ℓ^1 minimization to parameter choice*. Information and Inference: A Journal of the IMA, 10(2), 397-453 (2021).
- [5] Q. Bertrand, Q. Klopfenstein, M. Blondel, S. Vaiter, A. Gramfort, J. Salmon. (2020, November). *Implicit differentiation of Lasso-type models for hyperparameter optimization*. International Conference on Machine Learning (pp. 810-821). PMLR.
- [6] S. Boyd, L. Vandenberghe, *Convex Optimization*, Cambridge University Press, 2004.
- [7] D. J. Brady, K. Choi, D. L. Marks, R. Horisaki, S. Lim. *Compressive holography*. Optics express, 17(15), 13040-13049, (2009).
- [8] L. Calatroni, et al, 2020, *Adaptive parameter selection for weighted-TV image reconstruction problems*, J. Phys.: Conf. Ser. 1476 012003.
- [9] L. Calatroni, A. Lanza, M. Pragliola, F. Sgallari, (2019). *A flexible space-variant anisotropic regularization for image restoration with automated parameter selection*. SIAM Journal on Imaging Sciences, 12(2), 1001-1037.
- [10] D. Calvetti, S. Morigi, L. Reichel, F. Sgallari, (2000). *Tikhonov regularization and the L-curve for large discrete ill-posed problems*, Journal of computational and applied mathematics, 123(1-2), 423-446.
- [11] E. J. Candès, *The Restricted Isometry Property and its Implications for Compressed Sensing*, C. R. Acad. Sci. Paris, Ser. I 346 (2008) 589-592.
- [12] E. J. Candès, J. Romberg, T. Tao, *Robust Uncertainty Principles: Exact Signal Reconstruction From Highly Incomplete Frequency Information*, Ieee Transactions on Information Theory, Vol. 52, No. 2, February 2006.
- [13] W. L. Chan. *Terahertz imaging with compressive sensing*. Rice University (2010).
- [14] S. Chatterjee, K. Steinhaeuser, A. Banerjee, S. Chatterjee, A. Ganguly. (2012, April). *Sparse group lasso: Consistency and climate applications*. Proceedings of the 2012 SIAM International Conference on Data Mining (pp. 47-58). Society for Industrial and Applied Mathematics.
- [15] S. Chen, D. Donoho (1994). *Basis pursuit*. Proceedings of 1994 28th Asilomar Conference on Signals, Systems and Computers. doi:10.1109/ACSSC.1994.471413.

- [16] S. S. Chen, D. L. Donoho, M. A. Saunders (2001), *Atomic decomposition by basis pursuit*. SIAM review, 43(1), 129-159.
- [17] C. V. Correa, H. Arguello, G. R. Arce. Compressive spectral imaging with colored-patterned detectors. In 2014 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP) (pp. 7789-7793). (2014, May) IEEE.
- [18] L. Di Sopra, D. Piccini, S. Coppo, M. Stuber, J. Yerly. *An automated approach to fully self-gated free-running cardiac and respiratory motion-resolved 5D whole-heart MRI*. Magnetic resonance in medicine, 82(6), 2118-2132 (2019).
- [19] C. Deledalle, S. Vaiteer, J. M. Fadili, G. Peyré. *Stein Unbiased GrAdient estimator of the Risk (SUGAR) for multiple parameter selection*, SIAM Journal on Imaging Sciences, Society for Industrial and Applied Mathematics, 2014, 7 (4), pp.2448-2487. hal-00987295v2.
- [20] D. L. Donoho, *Compressed Sensing*, Ieee Transactions on Information Theory, Vol. 52, No. 4, April 2006.
- [21] D. L. Donoho, M. Elad, V. N. Temlyakov, *Stable Recovery of Sparse Overcomplete Representations in the Presence of Noise*, Ieee Transactions on Information Theory, Vol. 52, No. 1, January 2006.
- [22] L. I. Feng, S. Coppo, D. Piccini, J. Yerly, R. P. Lim, P. G. Masci, R. Otazo. *5D whole-heart sparse MRI*. Magnetic resonance in medicine, 79(2), 826-838 (2018).
- [23] S. Foucart, H. Rauhut, *A Mathematical Introduction to Compressive Sensing*, Birkhäuser, 2013.
- [24] B. Franceschiello, L. Di Sopra, A. Minier, S. Ionta, D. Zeugin, M. P. Notter, J. A. M. Bastiaansen, J. Jorge, J. Yerly, S. Stuber, M. M. Murray. *3-Dimensional magnetic resonance imaging of the freely moving human eye*. Progress in Neurobiology, 194, 101885 (2020).
- [25] L. Gao, J. Liang, C. Li, L. V. Wang. Single-shot compressed ultrafast photography at one hundred billion frames per second. Nature, 516(7529), 74-77 (2014).
- [26] G. H. Golub, M. Heath, G. Wahba, (1979). *Generalized cross-validation as a method for choosing a good ridge parameter*. Technometrics, 21(2), 215-223.
- [27] M. Guerquin-Kern, D. Van De Ville, C. Vonesch, J. C. Baritoux, K. P. Pruessmann, M. Unser. *Wavelet-regularized reconstruction for rapid MRI*. In 2009 IEEE International Symposium on Biomedical Imaging: From Nano to Macro (pp. 193-196) (2009, June). IEEE.
- [28] P. C. Hasen, *The L-curve and its use in the numerical treatment of inverse problems*, <https://www.sintef.no/globalassets/project/evitameeting/2005/lcurve.pdf>.

- [29] C. He, C. Hu, W. Zhang, B. Shi, *A Fast Adaptive Parameter Estimation for Total Variation Image Restoration*, IEEE Transactions on Image Processing, vol. 23, no. 12, pp. 4954-4967, Dec. 2014, doi: 10.1109/TIP.2014.2360133.
- [30] M. Hintermüller, K. Papafitsoros. (2019). *Generating structured nonsmooth priors and associated primal-dual methods*. Handbook of numerical analysis (Vol. 20, pp. 437-502). Elsevier.
- [31] Y. C. Kim, S. S. Narayanan, K. S. Nayak. *Accelerated three-dimensional upper airway MRI using compressed sensing*. Magnetic Resonance in Medicine: An Official Journal of the International Society for Magnetic Resonance in Medicine, 61(6), 1434-1440 (2009).
- [32] A. Lanza, M. Pragliola, F. Sgallari, (2020). *Residual whiteness principle for parameter-free image restoration*.
- [33] X. Liu, J. U. Kang. *Compressive SD-OCT: the application of compressed sensing in spectral domain optical coherence tomography*. Optics express, 18(21), 22010-22019, (2010).
- [34] M. Lustig, D. Donoho, J. M. Pauly, *Sparse MRI: The Application of Compressed Sensing for Rapid MR Imaging*, Magn. Res. Med. 58:1182-1195 (2007).
- [35] M. Lustig, D. L. Donoho, J. M. Santos, J. M. Pauly. *Compressed sensing MRI*. IEEE signal processing magazine, 25(2), 72-82 (2008).
- [36] J. D. McEwen, Y. Wiaux. *Compressed sensing for wide-field radio interferometric imaging*. Monthly Notices of the Royal Astronomical Society, 413(2), 1318-1332 (2011).
- [37] S. Mallat, *A Wavelet Tour in Signal Processing* Third edition, Academic Press (2009).
- [38] M. M. Marim, M. Atlan, E. Angelini, J. C. Olivo-Marin. *Compressed sensing with off-axis frequency-shifting holography*. Optics letters, 35(6), 871-873, (2010).
- [39] B. Milani, C. Roy, J. Ledoux, D. C. Rotzinger, S. Si-mohamed, A. Masi, J. Yerly, T. Rutz, T. Estelle, M. Prsa, J. Schwitter, M. Stuber. *A new whole-heart 3D-CINE-reconstruction of a 3D-radial free-running acquisition with a deformation-fields corrected temporal total variation regularization*. Manuscript in preparation.
- [40] R. R. D. Nekhili, X. Descombes, L. Calatroni. *A hybrid approach combining cnns and variational modelling for blind image denoising* (2022).
- [41] R. Rekik Dit Nekhili, X. Descombes, L. Calatroni. *A hybrid approach combining cnns and variational modelling for blind image denoising*. 2022.

- [42] S. Oymak, C. Thrampoulidis, B. Hassibi. *The squared-error of generalized lasso: A precise analysis*. In 2013 51st Annual Allerton Conference on Communication, Control, and Computing (Allerton) (pp. 1002-1009). IEEE (2013, October).
- [43] Peajcariaac J. E., Tong Y. L., *Convex Functions, Partial Orderings, and Statistical Applications*, Mathematics in Science and Engineering (1992).
- [44] M. Pragliola, L. Calatroni, A. Lanza, F. Sgallari. *ADMM-based residual whiteness principle for automatic parameter selection in super-resolution problems*. 2021.
- [45] M. Pragliola, L. Calatroni, A. Lanza, F. Sgallari, *On and beyond Total Variation regularisation in imaging: the role of space variance*. 2021, arXiv preprint 2104.03650.
- [46] Y. Plan, R. Vershynin. *The Generalized Lasso With Non-Linear Observations*. IEEE Transactions on information theory, 62(3), 1528-1537 (2016).
- [47] R. T. Rockafellar, *Convex Analysis* Second Printing, Princeton University Press (1972).
- [48] R. T. Rockafellar, R. J-B Wets: *Variational Analysis* 3rd edition, Springer-Verlag Berlin Heidelberg (2009).
- [49] W. Rudin, *Functional Analysis* 2nd edition, McGraw Hill (1990).
- [50] F. Santosa, W. W. Symes (1986). *Linear inversion of band-limited reflection seismograms*. SIAM Journal on Scientific and Statistical Computing. SIAM. 7 (4): 1307–1330. doi:10.1137/0907087.
- [51] E. Soubies, L. Blanc-Féraud, G. Aubert, (2015). *A continuous exact ℓ_0 penalty (CEL0) for least squares regularized problem*. SIAM Journal on Imaging Sciences, 8(3), 1607-1639.
- [52] R. Tibshirani (1996). *Regression shrinkage and selection via the lasso*. Journal of the Royal Statistical Society: Series B (Methodological), 58(1), 267-288.
- [53] G. Varela-Mattatal, C. A. Baron, R. S. Menon, *Automatic determination of the regularization weighting for wavelet-based compressed sensing MRI reconstructions*, Magn. Reson. Med. 2021;86:1403-1419.
- [54] M. Unser. *A unifying representer theorem for inverse problems and machine learning*. Foundations of Computational Mathematics, 21(4), 941-960 (2021).
- [55] M. Unser, S. Aziznejad. *Convex optimization in sums of Banach spaces*. Applied and Computational Harmonic Analysis, 56, 1-25 (2022).
- [56] H. Yu, G. Wang. *Compressed sensing based interior tomography*. Physics in medicine & biology, 54(9), 2791 (2009).

- [57] L. Zhu, W. Zhang, D. Elnatan, B. Huang. Faster STORM using compressed sensing. *Nature methods*, 9(7), 721-723 (2012).