

Testing Many Zero Restrictions in a High Dimensional Linear Regression Setting

Jonathan B. Hill*

Dept. of Economics, University of North Carolina

This draft: January 24, 2023

Abstract

We propose a test of many zero parameter restrictions in a high dimensional linear iid regression model. The test statistic is formed by estimating key parameters one at a time based on many low dimension regression models with nuisance terms. The parsimoniously parametrized models identify whether the original parameter of interest is or is not zero. Estimating fixed low dimension sub-parameters ensures greater estimator accuracy, does not require a sparsity assumption, and using only the largest in a sequence of weighted estimators reduces test statistic complexity and therefore estimation error. We provide a parametric wild bootstrap for p-value computation, and prove the test is consistent and has non-trivial $\sqrt{n/\ln(n)}$ -local-to-null power.

Key words and phrases: high dimensional inference, max-test, linear regression

AMS classifications : 62G10, 62M99, 62F35.

JEL classifications : C12, C55.

1 Introduction

Regression settings where the number k of covariates x_t may be much larger than the sample size n ($k \gg n$) is a natural possibility in cross sections, panels and spatial settings where an enormous amount of information is available. In economics and finance, for example, reasons include the size of surveys (e.g. U.S. Census, Current Population Survey, National Longitudinal Survey of Youth), increasingly sophisticated survey techniques (e.g. text records

*Department of Economics, University of North Carolina, Chapel Hill. E-mail: jbhill@email.unc.edu; <https://jbhill.web.unc.edu>. This paper was previously circulated under the title “*Testing (Infinitely) Many Zero Restrictions*”.

and word counts, household transaction data), the unit of observation (e.g. county wide house prices are dependent on neighbor county prices), use of dummy variables for groups (e.g. age, race, region, etc.), and panel data fixed effects. See, for instance, [Fan, Lv, and Qi \(2011\)](#) and [Belloni, Chernozhukov, and Hansen \(2014\)](#) for recent surveys. In non-social sciences high dimensionality arises with, for instance, genomic data and network and communication data.¹

Inference in this case is typically done post estimation based on a regression or moment condition model with an imposed sparsity condition, or related structural assumption. Sparsity in practice is achieved by a shrinkage or regularized estimator, while recent theory allows for an increasing number of candidate and relevant covariates as the sample size increases (see, e.g., [Fan and Peng, 2004](#); [Huang, Horowitz, and Ma, 2008](#)). Valid inference, however, typically exists only for the non-zero valued parameter subset, although there is a nascent literature for gaining inference on the (number of) zero valued parameters (e.g. [Carpentier and Verzelen, 2019](#)). Conversely, [Bühlmann \(2013\)](#) develops a test of zero restrictions for a high dimensional linear model with fixed design $\{x_1, \dots, x_n\}'$ using bias-corrected ridge regression. Their p-value construction, however, hinges on a Gaussian upper bound. Low dimension projection methods offer another solution (e.g. [Zhang and Zhang, 2014](#)). As all cases above, inference is performed *post* estimation. Imposing sparsity at all, however, may be too restrictive, in particular without justification via a pre-estimation test procedure.

Consider a sample of iid observations $\{x_t, y_t\}_{t=1}^n$ with scalar y_t , and covariates or design points $x_t = [x'_{\delta,t}, x'_{\theta,t}]'$, and a linear regression model for simplicity and to fix ideas:

$$y_t = \delta'_0 x_{\delta,t} + \theta'_0 x_{\theta,t} + \epsilon_t = \beta'_0 x_t + \epsilon_t. \tag{1}$$

This paper presents a test of $H_0 : \theta_0 = 0$ vs. $H_1 : \theta_{0,i} \neq 0$ for at least one i . We assume the nuisance parameter δ_0 has a fixed dimension $k_\delta < n$, but θ_0 may have dimension $k_\theta \gg$

¹As a testament to the increasing amount of available data, in November 2022 the 27th General Conference on Weights and Measures presented new numerical prefixes handling, among other magnitudes, 10^{30} .

n , including $k_\theta \rightarrow \infty$ as $n \rightarrow \infty$ (we do not show that θ_0 may depend on n). We assume k_θ is either fixed or $k_\theta \rightarrow \infty$, but $k_\theta \rightarrow k_\theta^* \in \mathbb{N}$ can be easily allowed. Let β_0 be the unique minimizer of the squared error loss $E[(y_t - \beta'x_t)^2]$, where $E[\epsilon_t] = 0$ and (ϵ_t, x_t) are mutually independent to focus ideas, but heterogeneity including (conditional) heteroscedasticity is a simple extension. See the appendix for all assumptions. We can allow for general hypotheses $H_0 : \theta_0 = \tilde{\theta}$ vs. $H_1 : \theta_0 \neq \tilde{\theta}$, but testing $\theta_0 = 0$ naturally saves notation.

Although $k_\theta \gg n$, we do not require a sparsity condition common in the high dimensional estimation literature (see, e.g., [Hastie, Tibshirani, and Friedman, 2009](#)). There are many examples of needing to work with (1), including panels with many fixed effects, estimation with many instrumental variables, and linear projections of nonlinear response, e.g. series expansions from flexible functional forms in machine learning (e.g. [Andrews, 1991](#); [Royston and Altman, 1994](#); [Belloni, Chernozhukov, Fernandez-Val, and Hansen, 2017](#); [Cattaneo, Jansson, and Newey, 2018](#); [Gautier and Rose, 2021](#)).

There is a large literature on high dimensional linear regression with an increasing number of covariates; consult [Cattaneo, Jansson, and Newey \(2018\)](#) and [Li and Müller \(2021\)](#), amongst others. This literature concerns estimation of high dimensional models. We, however, approach inference on θ_0 *pre-estimation*, in the sense that (1) is not estimated, hence a penalized estimator, projection method, or partialing out parameters are not required. Instead, we operate on many low dimension regression models, each with one key parameter $\theta_{0,i}$ to be tested, and any other nuisance parameters δ_0 to be estimated. Parsimony improves estimation of each $\theta_{0,i}$ when k_θ is large, and allows us to sidestep sparsity considerations under H_1 . Pre-estimation itself allows one to test a sparsity assumption based on a target set of parameters.

[Ghysels, Hill, and Motegi \(2020\)](#) exploit a similar setup, but for low dimension linear models with fixed $k_\theta < n$, and a simulation-based method for p-value computation. We allow $k_\theta \gg n$ including $k_\theta \rightarrow \infty$, and therefore an increasing number of estimated parameters. The latter leads to both greater generality, and a substantially different asymptotic theory

approach detailed below. We also exploit a more compact proof of hypothesis identification and a parametric wild bootstrap for p-value computation.

Our test approach is as follows. Write $\beta_{(i)} \equiv [\delta', \theta_i]'$, and construct parsimonious models:

$$y_t = \delta_{(i)}^* x_{\delta,t} + \theta_i^* x_{\theta,i,t} + v_{(i),t} = \beta_{(i)}^* x_{(i),t} + v_{(i),t}, \quad i = 1, \dots, k_{\theta,n}, \quad (2)$$

where (pseudo-true) $\beta_{(i)}^*$ for each i satisfy $E[(y_t - \beta_{(i)}^* x_{(i),t})x_{(i),t}] = 0$. Thus the i^{th} model corresponds to a unique regressor $x_{\theta,i,t}$ from $x_{\theta,t} = [x_{\theta,i,t}]_{i=1}^{k_{\theta}}$, and regressor subset $x_{\delta,t}$. Notice $k_{\theta,n} \leq k_{\theta}$ is the number of parameterizations used in practice. In general assume monotonically $k_{\theta,n} \rightarrow k_{\theta}$ if $k_{\theta} < \infty$, else $k_{\theta,n} \rightarrow \infty$ if $k_{\theta} \rightarrow \infty$. The methods proposed here can be directly extended to more general settings, including nonlinear regression models with non-identically distributed observations, conditional moment models, random choice and quantile regression to name a few.

A key result shows that under mild conditions $\theta^* \equiv [\theta_i^*]_{i=1}^{k_{\theta}} = 0$ if and only if $\theta_0 = 0$, thus each nuisance parameter $\delta_{(i)}^*$ is identically the true δ_0 . Obviously we cannot generally identify θ_0 (or δ_0) under the alternative, but (2) can identify whether $\theta_0 = 0$ is true or not.

Now let $\hat{\beta}_{(i)} \equiv [\hat{\delta}_{(i)}', \hat{\theta}_i']'$ minimize the parsimonious least squares loss $1/n \sum_{t=1}^n (y_t - \beta_{(i)}' x_{(i),t})^2$. The test statistic is the normalized maximum weighted $\hat{\theta}_i$:

$$\mathcal{T}_n = \max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{n} \mathcal{W}_{n,i} \hat{\theta}_i \right|,$$

where $\{\mathcal{W}_{n,i}\}_{n \geq 1}$ are sequences of possibly stochastic weights, $\mathcal{W}_{n,i} > 0$ a.s. for each i , with non-random (probability) limits $\mathcal{W}_i \in (0, \infty)$. The max-test rejects H_0 at level α when an appropriate p-value approximation \hat{p}_n satisfies $\hat{p}_n < \alpha$. The weights $\mathcal{W}_{n,i}$ allow for a variety of test statistics, including $\max_{1 \leq i \leq k_{\theta,n}} |\sqrt{n} \hat{\theta}_i|$, or a max-t-statistic. Bootstrap inference for the max-statistic does not require a covariance matrix inversion like Wald and LM tests, and operates like a shrinkage estimator by using only the most relevant (weighted) estimator. The max-test

is asymptotically correctly sized, and consistent because $k_{\theta,n} \rightarrow k_\theta$ or $k_{\theta,n} \rightarrow \infty$. Further, it has non-trivial power against a sequence of $\sqrt{n/\ln(n)}$ -local alternatives. Bootstrapping in high dimension has a rich history and recent offerings. Consider [Bickel and Freedman \(1983\)](#), [Mammen \(1993\)](#), and [Chernozhukov, Chetverikov, and Kato \(2013, 2017\)](#) and their references.

Our method and setting are significantly different from [Hill and Motegi \(2020\)](#) who propose a max-correlation test of a white noise hypothesis. We test zero restrictions in a regression setting that, in a time series extension, can couch a white noise test. They use a general high dimensional first order approximation *and* Gaussian approximation theory that allows for *some* $\{k_{\theta,n}\}$, $k_{\theta,n} \rightarrow \infty$, but which sequences $\{k_{\theta,n}\}$ are not characterized. We exploit a smooth-max approximation for a non-Gaussian approximation theory in order to characterize sequences $\{k_{\theta,n}\}$ that satisfy a high dimensional first order expansion, and then extant high dimensional Gaussian approximation theory.

The theory developed here applies whether k_θ is fixed, or increases with the sample size. The latter is common in the shrinkage literature (e.g. [va de Greer, 2008](#)), and has a deep history (e.g. [Huber, 1973](#)). See [Cattaneo, Jansson, and Newey \(2018\)](#) and their references. The dimension k_δ of δ_0 is assumed fixed and $k_\delta < n$ to focus ideas, but can in principle vary with n , or be unbounded. In a more general framework, if δ_0 is sparse then a penalized estimator can be used to estimate $[\delta_{(i)}^*, \theta_i^*]$, like lasso, ridge or Dantzig. If δ_0 is a (infinite dimensional) function, e.g. $y_t = \delta_0(x_{\delta,t}) + \theta_0' x_{\theta,t} + \epsilon_t$, then $[\delta_{(i)}^*, \theta_i^*]$ can be estimated by sieves (e.g. [Chen and Pouzo, 2015](#)). In our setting inference is on the high dimensional θ_0 , contrary to [Chernozhukov et al \(2016\)](#) and [Cattaneo, Jansson, and Newey \(2018\)](#) who test a fixed low dimension parameter after partialing out a high dimensional term.

Conventional test statistics typically exhibit size distortions due both to poor estimator sample properties, and from inverting a high dimension covariance matrix estimator that may be a poor approximation of the small or asymptotic variance. The challenge of high dimensional covariance matrix estimation is well documented (e.g. [Chen, Xu, and Wu, 2013](#)). A

bootstrap method is therefore typically applied, but bootstrap tests may only have size corrected power equal to the original test, which may be low under parameter proliferation (e.g. Ghysels, Hill, and Motegi, 2020, and their references). Indeed, the bootstrapped Wald test is undersized with low power in linear iid models when k_δ is even mildly large (e.g. $k_\delta = 10$)², and potentially significantly under-sized with low power when $k_{\theta,n}$ is large, becoming acute when there are nuisance parameters. Max-tests yield better (typically sharp) size, while power improvements over the Wald test can be sizable when deviations from the null are small.

Reduced dimension regression models in the high dimensional parametric statistics and machine learning literatures are variously called *marginal regression*, *correlation learning*, and *sure screening* (e.g. Fan and Lv, 2008; Genovese, Jin, Wasserman, and Yao, 2023). McKeague and Qian (2015), for example, regress y against one covariate x_i at a time, $i = 1, \dots, k < \infty$, and compute the most relevant index $\hat{k} \equiv \arg \max_{1 \leq i \leq k} |\hat{\theta}_i|$ where $\hat{\theta}_i \equiv \widehat{cov}(y, x_i) / \widehat{var}(x_i)$, and test $\tilde{H}_0 : \theta_0 = [cov(y, x_i) / var(x_i)]_{i=1}^k = 0$ using $\hat{\theta}_{\hat{k}}$. Thus they work outside a general regression setting, consider only a fixed number k of possible covariates, do not consider a subset of regressor predictability (e.g. $x_{\theta,t}$) with other nuisance parameters (δ_0), and use linear dependence for defining predictability. Related projections to low dimension settings are considered in Bühlmann (2013) and Zhang and Zhang (2014), amongst others,

We also work pre-estimation, but within a high dimensional regression model that allows for nuisance parameters, and test the parameter subset $H_0 : \theta_0 = 0$ where θ_0 may have a diverging dimension. Our method also has a straightforward extension to nonlinear models and a broad class of extremum estimators so that predictability may not align with linear dependence between y_t and the $x'_{i,t}$ s. Consider the nonlinear model $y_t = f(x_t, \beta_0) + \epsilon_t$ with known response f , and parsimonious representations $y_t = f(x_t, [\delta_{(i)}^*; 0, \dots, \theta_{(i)}^*, \dots, 0]) + v_{(i),t}$ where $[0, \dots, \theta_{(i)}, \dots, 0]'$ is a $k_\theta \times 1$ zero vector with $\theta_{(i)}$ in the i^{th} row. Thus θ_0 need not align one-for-one with a regressor subset $x_{\cdot,t}$. Examples abound in the financial and macro-

²This may carry over to (semi)nonparametric settings with fixed low dimension δ and infinite dimensional function θ_0 . See Chen and Pouzo (2015) for simulation evidence of sieve based bootstrapped t-tests.

econometrics, neural network and machine learning literatures, including switching models, flexible functional forms, and basis expansions. If there are no nuisance parameters ($k_\delta = 0$) then our theory should allow for a high dimensional expansion of McKeague and Qian’s (2015) method, but we leave that arc for future consideration.

Forward stepwise regressions exploit parsimonious models with increasing dimension. In the estimation literature low dimension, typically linear, models are built by adding one covariate at a time (e.g. Amemiya, 1980; Stone, 1981; Royston and Altman, 1994; Castle, Qin, and Reed, 2009). Our approach is different because we do not build up a model; rather, we use a different class of parsimonious models that combines just one key covariate with the nuisance covariates. Our models are true under the null, whereas stepwise methods typically do not hinge on a presumed hypothesis. Classically Kabe (1963) tests whether a low dimension subset $\theta_0 = 0$ with stepwise regressions that partials out δ_0 , effectively using parsimonious models. This idea has been treated many times since, e.g. Cattaneo, Jansson, and Newey (2018) and Chernozhukov et al (2016). Our method does not partial out low or high dimensional components, nor add covariates one at time based on predictor power.

Adaptive minimax methods often utilize sequences of sieves for specification tests. In the functional linear regression $y_i = \int_{\mathcal{T}} x_i(t)\theta_0(t)dt + \epsilon_i$, for example, tests of $\theta_0 = 0$ may rely on projecting y_i onto linear subspaces based on x_t (e.g. Hilgert, Mas, and Verzelen, 2013). Other techniques involve functional principle components (e.g., Hall and Horowitz, 2007), or reproducing kernel Hilbert spaces (Cai and Yuan, 2012).

The goal of minimax testing against sparse alternatives is to seek optimal or sharp convergence rates of (type I and II) error probabilities (cf. Ingster, 1997). See, e.g., Baraud, Giraud, and Huat (2009) and their references for treatment of mean models without covariates. Carpentier et al (2019) study minimax rates for tests of $\theta_0 = 0$ in the linear model $y_t = \theta'_0 x_t + \epsilon_t$ against an l_2 s -sparse alternative $0 < \tau \leq \|\theta_0\|_2$, and $\|\theta_0\|_0 \leq s$ with $\|\theta_0\|_0 \equiv \sum_{i=1}^{k_\theta} I(\theta_{0,i} \neq 0)$ for some $s \in \{1, \dots, k_\theta\}$. Ingster and Sapatinas (2009) treat $y_i = \theta_0(x_i) + \epsilon_i$ with square integrable θ_0 ,

iid uniform design points x_i , and Gaussian error ϵ_i . They present an adaptive minimax test of $H_0 : \theta_0 = \tilde{\theta}$ for given $\tilde{\theta}$, against a sequence of local alternatives $H_1 : \|\theta_0 - \theta\| \geq r_n \rightarrow 0$. See [Arias-Castro, Candès, and Plan \(2011\)](#) and [Zhong, Chen, and Xu \(2013\)](#). As noted above, we do not require sparsity under H_1 , and we do not estimate the original model.

We could, broadly speaking, exploit sieves in a semiparametric environment where δ_0 is an unknown function, but we otherwise differ sharply. In terms of (1) we do not estimate θ_0 , and our parameterizations all have the same non-increasing dimension $k_\delta + 1$, but the number of such parameterizations increases with n . Conversely, sieve spaces increase in complexity with n . In (1), for example, a simple linear sieve would be $y_t = \delta^{*'} x_{\delta,t} + \sum_{j=1}^{k_{\theta,n}} \theta_{n,j} x_{\theta,t,j} + v_{n,t}$ with increasing dimension $k_\delta + k_{\theta,n}$.

The remainder of the paper is as follows. Section 2 contains test statistic asymptotics and bootstrap p-value theory. A Monte Carlo study is presented in Section 3, and parting comments are left for Section 4. Assumptions, proofs of main results, and a subset of simulation results are relegated to the appendix. The supplemental material [SM] [Hill \(2023\)](#) contains omitted proofs and all simulation results.

We assume all random variables exist on a complete measure space³. $|x| = \sum_{i,j} |x_{i,j}|$ is the l_1 -norm, $|x|_2 = (\sum_{i,j} x_{i,j}^2)^{1/2}$ is the Euclidean or l_2 norm, and $\|A\| = \max_{|\lambda|_2} \{|A\lambda|_2 / |\lambda|_2\}$ is the spectral norm for finite dimensional square matrices A (and the Euclidean norm for vectors). $\|\cdot\|_p$ denotes the L_p -norm. *a.s.* is *almost surely*. $\mathbf{0}_k$ denotes a zero vector with dimension $k \geq 1$. Write r -vectors as $x \equiv [x_i]_{i=1}^r$. $[\cdot]$ rounds to the nearest integer. $K > 0$ is non-random and finite, and may take different values in different places. $a_n \propto b_n$ implies $a_n/b_n \rightarrow K$. *awp1* = asymptotically with probability approaching one. Differing from classic usage, we say z has *sub-exponential* distribution tails when $P(|z| > \varepsilon) \leq b \exp\{c\varepsilon\}$ for some $(b, c) > 0$ and all $\varepsilon > 0$ (see, e.g., [Vershynin, 2018](#), Chap.2.7).

³Completeness ensures majorants and integrals over uncountable families of measurable functions are measurable, and probabilities where applicable are outer probability measures. Cf. [Pollard \(1984, Appendix C: permissibility criteria\)](#).

2 Max-Test and p-Value Computation

We show the use of parsimonious models (2) leads to correct identification of the hypotheses. Collect θ_i^* across all possible parsimonious models: $\theta^* \equiv [\theta_i^*]_{i=1}^{k_\theta}$. We show $\theta^* = \mathbf{0}_{k_\theta}$ if and only if $\theta_0 = \mathbf{0}_{k_\theta}$. Thus, the models $y_t = \delta_{(i)}^* x_{\delta,t} + \theta_i^* x_{\theta,i,t} + v_{(i),t}$ may be used to construct a test of $\theta_0 = \mathbf{0}_{k_\theta}$ in (1). The proof may be of independent interest and is therefore presented here.

Theorem 2.1. $\theta_0 = \mathbf{0}_{k_\theta}$ if and only if $\theta^* = \mathbf{0}_{k_\theta}$. Under $H_0 : \theta_0 = \mathbf{0}_{k_\theta}$ hence $\delta_{(i)}^* = \delta_0 \forall i$, and under $H_1 : H_1 : \theta_{0,i} \neq 0$ for at least one $i \in \{1, \dots, k_\theta\}$ there exists an i such that $\theta_i^* \neq 0$.

Proof. By supposition $E[(y_t - \beta'x_t)x_{i,t}] = 0 \forall i$ if and only if $\beta = \beta_0 = [\delta_0', \theta_0']'$. Therefore, if $\theta_0 = \mathbf{0}_{k_\theta}$ then $E[(y_t - \delta_0'x_{\delta,t})x_{i,t}] = 0 \forall i$ hence $E[(y_t - \delta_0'x_{\delta,t})x_{(i),t}] = \mathbf{0}_{k_\delta+1}$ for each $i = 1, \dots, k_\theta$. This instantly yields $\theta_i^* = 0$ for each i and $\delta^* = \delta_0$ by construction and uniqueness of $\beta_{(i)}^* = [\delta_{(i)}^*, \theta_i^*]'$.

Conversely, if $\theta^* = \mathbf{0}_{k_\theta}$ then from the parsimonious risk it follows $E[(y_t - \delta_{(i)}^*x_{\delta,t})x_{(i),t}] = \mathbf{0}_{k_\delta+1}$ for each i by identification of $\beta_{(i)}^*$. Therefore $E[(y_t - \delta_{(i)}^*x_{\delta,t})x_{j,t}] = 0 \forall j$ by the definition of $x_{(i),t}$. But this implies the $\delta_{(i)}^*$ s are equivalent $\forall i$, hence $\delta_{(i)}^* = \delta^*$ for each i . This yields $E[(y_t - \delta^*x_{\delta,t})x_{i,t}] = 0 \forall i$ and therefore $[\delta_0, \theta_0] = [\delta^*, \mathbf{0}_{k_\theta}]$ by identification of $[\delta_0, \theta_0]$. \mathcal{QED} .

Identification of the hypotheses carries over asymptotically when fewer parsimonious models are used $k_{\theta,n} < k_\theta$ as long as $k_{\theta,n} \rightarrow k_\theta$ if $k_\theta < \infty$, else $k_{\theta,n} \rightarrow \infty$. Collect $\theta^* = [\theta_i^*]_{i=1}^{k_{\theta,n}}$ which implicitly depends on n . By the same argument as above, $\theta_0 = \mathbf{0}_{k_\theta}$ if and only if $\theta^* = \mathbf{0}_{k_{\theta,n}} \forall n$. Under H_1 it therefore follows $\liminf_{n \rightarrow \infty} \max_{1 \leq i \leq k_{\theta,n}} |\theta_i^*| > 0$. In the sequel we simply assume $k_\theta \rightarrow \infty$ and $k_{\theta,n} \rightarrow \infty$ to reduce cases.

2.1 Max-Test Gaussian Approximation

Define parsimonious least squares loss gradient and Hessians,

$$\widehat{\mathcal{G}}_{(i)} \equiv -\frac{1}{n} \sum_{t=1}^n v_{(i),t} x_{(i),t}, \quad \widehat{\mathcal{H}}_{(i)} \equiv \frac{1}{n} \sum_{t=1}^n x_{(i),t} x'_{(i),t}, \quad \mathcal{H}_{(i)} \equiv E[x_{(i),t} x'_{(i),t}],$$

and define the usual least squares first order terms: $\hat{\mathbf{Z}}_{(i)} \equiv \sqrt{n}\mathcal{H}_{(i)}^{-1}\hat{\mathcal{G}}_{(i)} = -\mathcal{H}_{(i)}^{-1}n^{-1/2}\sum_{t=1}^n v_{(i),t}x_{(i),t}$.

Under H_0 of course $v_{(i),t} = \epsilon_t$ for each i by Theorem 2.1. By standard least squares arguments

$0 = \hat{\mathcal{G}}_{(i)} + \hat{\mathcal{H}}_{(i)}(\hat{\beta}_{(i)} - \beta_{(i)}^*)$, hence

$$\sqrt{n}\left(\hat{\beta}_{(i)} - \beta_{(i)}^*\right) = -\sqrt{n}\mathcal{H}_{(i)}^{-1}\hat{\mathcal{G}}_{(i)} - \left\{\hat{\mathcal{H}}_{(i)}^{-1} - \mathcal{H}_{(i)}^{-1}\right\}\sqrt{n}\hat{\mathcal{G}}_{(i)} \equiv \hat{\mathbf{Z}}_{(i)} + \hat{\mathcal{R}}_i.$$

We want a high dimensional Gaussian approximation $|\max_{1 \leq i \leq k_{\theta,n}} |\sqrt{n}\mathcal{W}_{n,i}\hat{\theta}_i| - \max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_i\mathbf{Z}_{(i)}| | \xrightarrow{p} 0$ where $\{\mathbf{Z}_{(i)}\}_{i \in \mathbb{N}}$ is Gaussian. Set all weights $\mathcal{W}_{n,i} = 1$ to ease notation for now. Then:

$$\left| \max_{1 \leq i \leq k_{\theta,n}} |\sqrt{n}\hat{\theta}_i| - \max_{1 \leq i \leq k_{\theta,n}} |\mathbf{Z}_{(i)}| \right| \leq \max_{1 \leq i \leq k_{\theta,n}} |[\mathbf{0}'_{k_\delta}, 1]\hat{\mathcal{R}}_i| + \left| \max_{1 \leq i \leq k_{\theta,n}} |[\mathbf{0}'_{k_\delta}, 1]\hat{\mathbf{Z}}_{(i)}| - \max_{1 \leq i \leq k_{\theta,n}} |\mathbf{Z}_{(i)}| \right|.$$

A Gaussian approximation theory under H_0 extends to $\max_{1 \leq i \leq k_{\theta,n}} |[\mathbf{0}'_{k_\delta}, 1]\hat{\mathbf{Z}}_{(i)}| - \max_{1 \leq i \leq k_{\theta,n}} |\mathbf{Z}_{(i)}|$ by Lemma 2.3, below. However, the remainder $\max_{1 \leq i \leq k_{\theta,n}} |[\mathbf{0}'_{k_\delta}, 1]\hat{\mathcal{R}}_i|$ is the usual first order approximation difference (in high dimension):

$$\max_{1 \leq i \leq k_{\theta,n}} |[\mathbf{0}'_{k_\delta}, 1]\hat{\mathcal{R}}_i| = \left| \max_{1 \leq i \leq k_{\theta,n}} |\sqrt{n}\mathcal{W}_{n,i}[\mathbf{0}'_{k_\delta}, 1]\hat{\theta}_i| - \max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_i[\mathbf{0}'_{k_\delta}, 1]\hat{\mathbf{Z}}_{(i)}| \right|$$

where generally $\hat{\mathbf{Z}}_{(i)} = -\mathcal{H}_{(i)}^{-1}n^{-1/2}\sum_{t=1}^n \epsilon_t x_{(i),t}$ is not Gaussian. If $\epsilon_t \sim N(0, \sigma^2)$, and $x_{i,t}$ are non-random design points, then $\hat{\mathbf{Z}}_{(i)} \sim N(0, \sigma^2\mathcal{H}_{(i)}^{-1})$ and a Gaussian theory applies. Otherwise, generally $\epsilon_t x_{(i),t}$ is non-Gaussian. Moreover, if we have stochastic weights $\{\mathcal{W}_{n,i}\}$ then we must contend with $\max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_{n,i} - \mathcal{W}_i|$ for non-Gaussian $\{\mathcal{W}_{n,i}, \mathcal{W}_i\}$.

We have the following general result proving $\max_{1 \leq i \leq k_{\theta,n}} |\hat{\mathcal{R}}_i| \xrightarrow{p} 0$. Assumption 1 is discussed in Appendix A.1.

Lemma 2.2. *Let H_0 , and Assumption 1 hold. Let $\{k_{\theta,n}\}$ be any monotonic sequence of positive integers satisfying $k_{\theta,n} = o(\sqrt{n}/(\ln(n))^4)$, and let the weight sequences $\{\mathcal{W}_{n,i}\}_{n \in \mathbb{N}}$ satisfy*

$\max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_{n,i} - \mathcal{W}_i| = o_p(1/(\ln(n))^2)$ for non-stochastic $\mathcal{W}_i \in (0, \infty)$. Then:

$$\left| \max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{n} \mathcal{W}_{n,i} \hat{\theta}_i \right| - \max_{1 \leq i \leq k_{\theta,n}} \left| \mathcal{W}_i [\mathbf{0}'_{k_\delta}, 1] \hat{\mathbf{Z}}_{(i)} \right| \right| \xrightarrow{p} 0. \quad (3)$$

Remark 2.1. The number of models must satisfy $k_{\theta,n} = o(\sqrt{n}/(\ln(n))^4)$, a rate significantly slower than the exponential rate available for *Gaussian* approximations (e.g. Chernozhukov, Chetverikov, and Wernic, 2013, amongst others), cf. Lemma 2.3 below. This arises from the fact that $\hat{\mathbf{Z}}_{(i)}$ in general is not Gaussian, hence a Gaussian approximation theory does not apply here. In the Gaussian case $\hat{\mathbf{Z}}_{(i)} \sim N(0, \sigma^2 \mathcal{H}_{(i)}^{-1})$, however, a smooth-max approximation and Slepian-type and Sudakov-Fernique comparison inequalities can be exploited to yield an exponential upper bound (e.g. Chernozhukov, Chetverikov, and Kato, 2013). We also use a smooth-max approximation to prove (3) under $k_{\theta,n} = o(\sqrt{n}/(\ln(n))^4)$, but in a limited way due to non-Gaussianity. See Lemma A.2 in the appendix, and its proof in the SM.

Remark 2.2. \mathcal{W}_i is non-random because the Gaussian approximation theory below requires $\mathcal{W}_i \mathbf{Z}_{(i)}(\cdot)$ to be Gaussian. The requirement $\max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_{n,i} - \mathcal{W}_i| = o_p(1/(\ln(n))^2)$ is the result of a non-Gaussian approximation theory leading to (3) with $\ln(k_{\theta,n}) = o(\ln(n))$, cf. Lemma A.2.

In the lemma below we derive a class of sequences $\{k_{\theta,n}\}$, and an array of scalar normal random variables $\{\mathbf{Z}_{(i)}(\lambda) : n \in \mathbb{N}\}_{i=1}^\infty$, $\mathbf{Z}_{(i)}(\lambda) \sim N(0, E[\epsilon_i^2] \lambda' \mathcal{H}_{(i)}^{-1} \lambda)$, such that for each $\lambda \in \mathbb{R}^{k_\delta+1}$, $\lambda' \lambda = 1$, the Kolmogorov distance

$$\rho_n(\lambda) \equiv \sup_{z \geq 0} \left| P \left(\max_{1 \leq i \leq k_{\theta,n}} \left| \lambda' \hat{\mathbf{Z}}_{(i)} \right| \leq z \right) - P \left(\max_{1 \leq i \leq k_{\theta,n}} \left| \mathbf{Z}_{(i)}(\lambda) \right| \leq z \right) \right| \rightarrow 0. \quad (4)$$

In terms of the linear contrasts $\lambda' \hat{\mathbf{Z}}_{(i)}$ we specifically care about $\lambda = [\mathbf{0}'_{k_\delta}, 1]'$ in view of expansion (3), hence we only require a pointwise limit theory vis-à-vis λ . The approximation does not require standardized $\lambda' \hat{\mathbf{Z}}_{(i)}$ and $\mathbf{Z}_{(i)}(\lambda)$ because we assume $\lambda' \mathcal{H}_{(i)}^{-1} \lambda$ lies in a

compact subset of $(0, \infty)$ asymptotically uniformly in i and λ . We work in the setting of Chernozhukov, Chetverikov, and Kato (2013, (E.1)) in order to prove (4).

Lemma 2.3. *Let H_0 and Assumption 1 hold. Then $\rho_n(\lambda) = O(n^{-c})$ for some $c > 0$ and any $\{k_{\theta,n}\}$ satisfying $k_{\theta,n} \leq \exp\{\mathcal{K}n^d\}/n$ with $\mathcal{K} > 0$ and $d \in (0, 1/7)$ (hence $k_{\theta,n} = O(\exp\{O(n^d)\})$).*

We now have the main result of this section. The statistic $\mathcal{T}_n = \max_{1 \leq i \leq k_{\theta,n}} |\sqrt{n}\mathcal{W}_{n,i}\hat{\theta}_i|$ can be well approximated by the max-Gaussian process $\{\max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_i \mathbf{Z}_{(i)}([\mathbf{0}'_{k_\delta}, 1])|\}_{n \in \mathbb{N}}$.

Theorem 2.4. *Let Assumption 1 hold, let $\{k_{\theta,n}\}$ be any monotonic sequence of positive integers satisfying $k_{\theta,n} = o(\sqrt{n}/(\ln(n))^4)$, and assume the weight sequences satisfy $\max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_{n,i} - \mathcal{W}_i| = o_p(1/(\ln(n))^2)$ for non-stochastic $\mathcal{W}_i \in (0, \infty)$ and $\max_{i \in \mathbb{N}} \mathcal{W}_i < \infty$.*

a. *Under H_0 , for the Gaussian random variables $\mathbf{Z}_{(i)}(\lambda) \sim N(0, E[\epsilon_t^2] \lambda' \mathcal{H}_{(i)}^{-1} \lambda)$ in (4),*

$$\left| P \left(\max_{1 \leq i \leq k_{\theta,n}} |\sqrt{n}\mathcal{W}_{n,i}\hat{\theta}_i| \leq z \right) - P \left(\max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_i \mathbf{Z}_{(i)}([\mathbf{0}'_{k_\delta}, 1])| \leq z \right) \right| \rightarrow 0 \quad \forall z \geq 0. \quad (5)$$

Furthermore, $\max_{1 \leq i \leq k_{\theta,n}} |\sqrt{n}\mathcal{W}_{n,i}\hat{\theta}_i| \xrightarrow{d} \max_{i \in \mathbb{N}} |\mathcal{W}_i \mathbf{Z}_{(i)}([\mathbf{0}'_{k_\delta}, 1])|$.

b. *Under H_1 , $\max_{1 \leq i \leq k_{\theta,n}} |\sqrt{n}\mathcal{W}_{n,i}\hat{\theta}_i| \rightarrow \infty$ for any monotonic sequence of positive integers $\{k_{\theta,n}\}$, $k_{\theta,n} \rightarrow \infty$.*

Remark 2.3. We can attempt to characterize the max-statistic \mathcal{T}_n null limit sequence $\{\max_{1 \leq i \leq k} |\mathcal{W}_i \mathbf{Z}_{(i)}([\mathbf{0}'_{k_\delta}, 1])|\}_{k \in \mathbb{N}}$ after re-scaling. In Appendix D of the SM we present an environment such that under H_0 for some positive non-random $b_n \sim \sqrt{2 \ln(k_{\theta,n})}$ and any $\{k_{\theta,n}\}$, $\sqrt{\ln(k_{\theta,n})} k_{\theta,n} = o(\sqrt{n}/(\ln(n))^4)$, we have $\sqrt{2 \ln(k_{\theta,n})/E[\epsilon_t^2]}(\mathcal{T}_n - b_n) \xrightarrow{d} \mathfrak{G}$, a Gumbel law. This is derived under fixed weights $\mathcal{W}_{n,i} = \mathcal{W}_i = 1$ and mutually uncorrelated covariates. The slightly decreased rate $k_{\theta,n} \rightarrow \infty$ follows from the first order approximation theory and a sacrifice for working in the max-domain of attraction. If stochastic weights are used then there is a significant sacrifice in $k_{\theta,n} \rightarrow \infty$ to account for $P(\mathcal{W}_{n,i} \neq \mathcal{W}_i) > 0$: we require $k_{\theta,n} = o(\exp\{(\ln(n))^{2/3}\})$, hence $k_{\theta,n} = o(n^c) \quad \forall c > 0$. See comments following Lemma D.1. The next

section delivers a wild bootstrapped p-value approximation without sacrificing $k_{\theta,n}$, allowing for nuisance parameters and arbitrary covariate dependence, and naturally affording better small sample inference.

2.2 Parametric Bootstrap

We utilize a variant of the wild (multiplier) bootstrap for inference (see [Bose, 1988](#); [Liu, 1988](#); [Gonçalves and Kilian, 2004](#)). Define the restricted estimator $\hat{\beta}^{(0)} \equiv [\hat{\delta}^{(0)'}, \mathbf{0}'_{k_\theta}]'$ under $H_0 : \theta_0 = \mathbf{0}_{k_\theta}$, where $\hat{\delta}^{(0)}$ minimizes the restricted least squares criterion $\sum_{t=1}^n (y_t - \delta' x_{\delta,t})^2$. Define residuals $\epsilon_{n,t}^{(0)} \equiv y_t - \hat{\delta}^{(0)'} x_{\delta,t}$, draw iid $\{\eta_t\}_{t=1}^n$ from $N(0, 1)$, and generate $y_{n,t}^* \equiv \hat{\delta}^{(0)'} x_{\delta,t} + \epsilon_{n,t}^{(0)} \eta_t$. Now construct $k_{\theta,n}$ regression models $y_{n,t}^* = \beta'_{(i)} x_{(i),t} + v_{n,(i),t}$, and let $\tilde{\beta}_{(i)} = [\tilde{\delta}'_{(i)}, \tilde{\theta}'_i]'$ be the least squares estimator of $\beta_{(i)}$. The bootstrapped test statistic is

$$\tilde{\mathcal{T}}_n \equiv \max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{n} \mathcal{W}_{n,i} \tilde{\theta}_i \right|. \quad (6)$$

Repeat the above steps \mathcal{M} times, each time drawing a new iid sequence $\{\eta_{j,t}\}_{t=1}^n$, $j = 1 \dots \mathcal{M}$. This results in a sequence of bootstrapped estimators $\{\tilde{\theta}_j\}_{j=1}^{\mathcal{M}}$ and test statistics $\{\tilde{\mathcal{T}}_{n,j}\}_{j=1}^{\mathcal{M}}$ that are iid conditional on the sample $\{x_t, y_t\}_{t=1}^n$. The approximate p-value is

$$\tilde{p}_{n,\mathcal{M}} \equiv \frac{1}{\mathcal{M}} \sum_{j=1}^{\mathcal{M}} I(\tilde{\mathcal{T}}_{n,j} > \mathcal{T}_n). \quad (7)$$

The above algorithm varies from [Gonçalves and Kilian \(2004\)](#) since they operate on the *unrestricted* estimator in order to generate residuals. We impose the null when we estimate β_0 : this reduces dimensionality without affecting asymptotics under either hypothesis.

Now write $\beta^{(0)} = [\delta^{(0)'}, \mathbf{0}'_{k_\theta}]'$ and $\beta_{(i)}^{(0)} = [\delta^{(0)'}, 0]'$, where $\delta^{(0)}$ minimizes $E[(y_t - \delta' x_{\delta,t})^2]$ on

compact $\mathcal{D} \subset \mathbb{R}^{k_\delta}$. Define the normalized bootstrapped gradient under the null:

$$\tilde{\mathbf{Z}}_{(i)}^{(0)} \equiv - \left(E [x_{\delta,t} x'_{\delta,t}] \right)^{-1} \frac{1}{\sqrt{n}} \sum_{t=1}^n \eta_t (y_t - \delta^{(0)' } x_{\delta,t}) x_{\delta,t}. \quad (8)$$

We prove the following asymptotic expansion in Lemma A.6: if $k_{\theta,n} = o(n)$ then

$$\left| \max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{n} \mathcal{W}_{n,i} \hat{\theta}_i \right| - \max_{1 \leq i \leq k_{\theta,n}} \left| \mathcal{W}_i [\mathbf{0}'_{k_\delta}, 1] \tilde{\mathbf{Z}}_{(i)}^{(0)} \right| \right| \xrightarrow{P} 0. \quad (9)$$

The improvement $k_{\theta,n} = o(n)$ over $k_{\theta,n} = o(\sqrt{n}/(\ln(n))^4)$ in Lemma 2.2 arises from the Gaussian multiplier η_t , and non-Gaussian approximation theory, cf. Lemma C.2 in the SM. In order to characterize a conditional high dimensional Gaussian approximation for $\tilde{\mathbf{Z}}_{(i)}^{(0)}$, define

$$\tilde{\sigma}_{(i)}^2(\lambda) \equiv \lambda' \left(\bar{\mathcal{H}}_{(i)}^{(0)} \right)^{-1} E \left[(y_t - \delta^{(0)' } x_{\delta,t})^2 x_{\delta,t} x'_{\delta,t} \right] \left(\bar{\mathcal{H}}_{(i)}^{(0)} \right)^{-1} \lambda \quad (10)$$

where $\bar{\mathcal{H}}_{(i)}^{(0)} \equiv E[x_{\delta,t} x'_{\delta,t}]$. Notice $\tilde{\sigma}_{(i)}^2(\lambda) = E[\epsilon_t^2] \lambda' \mathcal{H}_{(i)}^{-1} \lambda$ under H_0 follows from Theorem 2.1. As usual, we only care about $\lambda = [\mathbf{0}'_{k_\delta}, 1]'$. Let \Rightarrow^P denote *weak convergence in probability* (Giné and Zinn, 1990: Section 3), a useful notion for multiplier bootstrap asymptotics. We have the following conditional multiplier bootstrap central limit theorem.

Lemma 2.5. *Let Assumption 1 hold. Let $\{\tilde{\mathbf{Z}}_{(i)}(\lambda)\}_{i \in \mathbb{N}}$, $\tilde{\mathbf{Z}}_{(i)}(\lambda) \sim N(0, \tilde{\sigma}_{(i)}^2(\lambda))$, be an independent copy of the Theorem 2.4.a null distribution process $\{\mathbf{Z}_{(i)}(\lambda)\}_{i \in \mathbb{N}}$, that is independent of the asymptotic draw $\{x_t, y_t\}_{t=1}^\infty$. Then for any monotonic sequence of positive integers $\{k_{\theta,n}\}$ that satisfies $k_{\theta,n} = o(\sqrt{n}/(\ln(n))^2)$: $\sup_{z \geq 0} |P(\max_{1 \leq i \leq k_{\theta,n}} |\lambda' \tilde{\mathbf{Z}}_{(i)}^{(0)}| \leq z | \mathfrak{G}_n) - P(\max_{1 \leq i \leq k_{\theta,n}} |\tilde{\mathbf{Z}}_{(i)}(\lambda)| \leq z)| \xrightarrow{P} 0$. Furthermore $\max_{1 \leq i \leq k_{\theta,n}} |[\mathbf{0}'_{k_\delta}, 1] \tilde{\mathbf{Z}}_{(i)}^{(0)}| \Rightarrow^P \max_{i \in \mathbb{N}} |\tilde{\mathbf{Z}}_{(i)}([\mathbf{0}'_{k_\delta}, 1])|$.*

Remark 2.4. The result is based on a (conditional) Slepian-type inequality, following from Lemma 3.1 in Chernozhukov, Chetverikov, and Kato (2013) (cf. Chernozhukov, Chetverikov, and Kato, 2015, Theorem 2). We ultimately exploit $k_{\theta,n} = o(\sqrt{n}/(\ln(n))^2)$ which is just slightly better than $k_{\theta,n} = o(\sqrt{n}/(\ln(n))^4)$ in the asymptotic expansion Lemma 2.2 and Theorem 2.4 due to

the iid Gaussian draw η_t .

Lemmas 2.5 and asymptotic approximation (9) yield the following fundamental result.

Lemma 2.6. *Let Assumption 1 hold, and let $\max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_{n,i} - \mathcal{W}_i| = o_p(1/(\ln(n))^2)$ for non-stochastic $\mathcal{W}_i \in (0, \infty)$. Let $\{\tilde{\mathbf{Z}}_{(i)}(\lambda)\}_{i \in \mathbb{N}}$, $\tilde{\mathbf{Z}}_{(i)}(\lambda) \sim N(0, \tilde{\sigma}_{(i)}^2(\lambda))$, be an independent copy of the Theorem 2.4.a null distribution process $\{\mathbf{Z}_{(i)}(\lambda)\}_{i \in \mathbb{N}}$, that is independent of the asymptotic draw $\{x_t, y_t\}_{t=1}^\infty$. Then $\max_{1 \leq i \leq k_{\theta,n}} |\sqrt{n}\mathcal{W}_{n,i}\hat{\theta}_i| \Rightarrow^p \max_{i \in \mathbb{N}} |\mathcal{W}_i \tilde{\mathbf{Z}}_{(i)}([\mathbf{0}'_{k_\delta}, 1])|$ for any monotonic sequence of positive integers $\{k_{\theta,n}\}$ that satisfies $k_{\theta,n} = o(\sqrt{n}/(\ln(n))^2)$.*

Remark 2.5. The same argument presented in Lemma D.1 of the SM, cf. Remark 2.3 above, yields $\sqrt{2 \ln(k_{\theta,n})/E[\epsilon_t^2]}(\max_{1 \leq i \leq k_{\theta,n}} |\sqrt{n}\mathcal{W}_{n,i}\hat{\theta}_i| - b_n) \xrightarrow{d} \mathfrak{G}$ under additional regularity conditions. Trivially, however, a bootstrapped p-value for $\max_{1 \leq i \leq k_{\theta,n}} |\sqrt{n}\mathcal{W}_{n,i}\hat{\theta}_i|$ is identical to one built from $\sqrt{2 \ln(k_{\theta,n})/E[\epsilon_t^2]}(\max_{1 \leq i \leq k_{\theta,n}} |\sqrt{n}\mathcal{W}_{n,i}\hat{\theta}_i| - b_n)$, hence there is no advantage here of working in the max-domain of attraction.

We now have the main result of this section: the approximate p-value $\tilde{p}_{n,\mathcal{M}_n}$ promotes a correctly sized and consistent test asymptotically. Notice again $k_{\theta,n} = o(\sqrt{n}/(\ln(n))^4)$ is required because $\tilde{p}_{n,\mathcal{M}_n}$ is computed from $\hat{\theta}_i$ and $\hat{\theta}_i$, cf. Theorem 2.4.

Theorem 2.7. *Let Assumption 1 hold, let $k_{\theta,n} = o(\sqrt{n}/(\ln(n))^2)$ and assume $\max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_{n,i} - \mathcal{W}_i| = o_p(1/(\ln(n))^2)$ for non-stochastic $\mathcal{W}_i \in (0, \infty)$. Let $\{k_{\theta,n}\}_{n \geq 1}$ be any monotonic sequence of positive integers with $k_{\theta,n} = o(\sqrt{n}/(\ln(n))^4)$.*

a. Under H_0 , $P(\tilde{p}_{n,\mathcal{M}_n} < \alpha) \rightarrow \alpha$.

b. Under H_1 , $P(\tilde{p}_{n,\mathcal{M}_n} < \alpha) \rightarrow 1$.

Remark 2.6. The test is consistent against a global alternative despite the fact that \mathcal{T}_n increases in magnitude in n even under H_0 . The proof reveals the p-value approximation $\tilde{p}_{n,\mathcal{M}_n}$ is asymptotically in probability equivalent to

$$P\left(\max_{1 \leq i \leq k_{\theta,n}} \left| \mathcal{W}_i \tilde{\mathbf{Z}}_{(i)}([\mathbf{0}'_{k_\delta}, 1]) \right| > \max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{n}\mathcal{W}_{n,i}(\hat{\theta}_i - \theta_{0,i}) + \sqrt{n}\mathcal{W}_i\theta_{0,i} + o_p(1) \right|\right)$$

where $\{\tilde{\mathbf{Z}}_{(i)}([\mathbf{0}'_{k_\delta}, 1])\}$ is an independent copy of $\{\mathbf{Z}_{(i)}(\lambda)\}_{i \in \mathbb{N}}$. The max-statistic therefore increases in magnitude at rate \sqrt{n} under H_1 . Conversely standard Gaussian concentration bounds yield the null process satisfies $\max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_i \tilde{\mathbf{Z}}_{(i)}([\mathbf{0}'_{k_\delta}, 1])| = O_p(\sqrt{\ln(k_{\theta,n})})$, and by assumption $\ln(k_{\theta,n}) \propto \ln(n)$. See (A.2) in the appendix. This yields consistency $\tilde{p}_{n, \mathcal{M}_n} \xrightarrow{p} 0$.

Finally, the bootstrapped test has non-trivial power against a sequence of local alternatives $H_1^L : \theta_0 = c\sqrt{\ln(k_{\theta,n})/n}$ where $c = [c_i]_{i=1}^{k_\theta}$. The local-to-null rate is generally less than \sqrt{n} because the test statistic under the null is, *awp1*, the maximum of an increasing sequence of Gaussian random variables $\max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_i \tilde{\mathbf{Z}}_{(i)}([\mathbf{0}'_{k_\delta}, 1])|$, and $\max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_i \tilde{\mathbf{Z}}_{(i)}([\mathbf{0}'_{k_\delta}, 1])| = O_p(\sqrt{\ln(k_{\theta,n})})$ as discussed above.

Theorem 2.8. *Let Assumption 1 hold, let $\max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_{n,i} - \mathcal{W}_i| = o_p(1/(\ln(n))^2)$ for non-stochastic $\mathcal{W}_i \in (0, \infty)$ and let $k_{\theta,n} = o(\sqrt{n}/(\ln(n))^2)$. Then $\lim_{n \rightarrow \infty} P(\tilde{p}_{n, \mathcal{M}_n} < \alpha) \nearrow 1$ under H_1^L monotonically as the maximum local drift $\max_{i \in \mathbb{N}} |c_i| \nearrow \infty$.*

Remark 2.7. Asymptotic local power therefore does not depend on the degree, if any, of sparsity. By comparison, for example, [Zhong, Chen, and Xu \(2013\)](#) consider a high dimensional mean model $y_i = \theta_0 + \epsilon_i$ and test $H_0 : \theta_0 = \mathbf{0}_p$, where $y_i \in \mathbb{R}^p$. Under their alternative $p^{1-\xi}$ elements $\theta_{0,i} \neq 0$ for some $\xi \in (1/2, 1)$, where $\theta_{0,i} = \sqrt{r \ln(p)/n}$ for some $r > 0$, and by assumption $\ln(p) = o(n^{1/3})$, hence $\theta_{0,i} = o(1/n^{2/3})$. See also [Arias-Castro, Candès, and Plan \(2011\)](#) and their references. Our max-test is consistent against local deviations $\theta_0 = c(\ln(k_{\theta,n})/n)^\zeta$ for any $\zeta \in [0, 1/2)$ and $c \neq \mathbf{0}_{k_\theta}$, and has non-trivial power when $\zeta = 1/2$ and $c \neq \mathbf{0}_{k_\theta}$. Further, we allow for any positive number of elements $\theta_{0,i} \neq 0$ under H_1 or H_1^L (i.e. $1 \leq \sum_{i=0}^{k_\theta} I(\theta_{0,i} \neq 0) \leq k_\theta$). Thus, in the simple setting of [Zhong, Chen, and Xu \(2013\)](#), alternatives can be closer to the null than allowed here in a max-test setting, but sparsity is enforced. Here, any degree of (non)sparsity is permitted under H_1 , (as long as some $\theta_{0,i} \neq 0$) or H_1^L (provided $c_i \neq 0$ for some i).

3 Monte Carlo Experiments

In our simulation study we use 10,000 independently drawn samples of sizes $n \in \{100, 250, 500, 1000\}$. In each case two max-tests are performed: one uses a flat weight $\mathcal{W}_{n,i} = 1$, and the other $\mathcal{W}_{n,i}$ is equal to the inverted least squares standard error of $\hat{\theta}_{n,i}$. We call these the *max-test* and *max-t-test* respectively. Other tests are discussed below. The number of parsimonious regression models is $k_{\theta,n} \in \{10, 35, [5n^{1/2-\iota}]\}$ where $\iota = 10^{-10}$. Note $[5n^{1/2-\iota}] \in \{50, 79, 112, 158\}$ for $n \in \{100, 250, 500, 1000\}$. There are $k_{\delta} \in \{0, 10\}$ nuisance parameters. All bootstrapped p-values are based on $\mathcal{M} = 1000$ independently drawn samples from $N(0, 1)$. Significance levels are $\alpha \in \{.01, .05, .10\}$.

The following presents our benchmark design, after which we discuss robustness checks.

3.1 Design and Tests of Zero Restrictions

3.1.1 Set-Up

The DGP is $y_t = \delta'_0 x_{\delta,t} + \theta'_0 x_{\theta,t} + \epsilon_t$ where ϵ_t is iid standard normal. We consider three cases for the regressors. In the first $[x_{\delta,t}, x_{\theta,t}]$ are serially and mutually independent standard normals. In the second they are block-wise dependent normals ($x_{\delta,t}$ and $x_{\theta,t}$ are mutually independent). In the third they are within and across block dependent normals. Not surprisingly all tests perform better under the first two cases. We therefore only report results for the third case.

Covariates in case three are drawn as follows. Define $k \equiv k_{\delta} + k_{\theta,n}$. Combine $x_t \equiv [x'_{\delta,t}, x'_{\theta,t}]' \in \mathbb{R}^k$, and let (w_t, v_t) be mutually independent draws from $N(0, I_k)$. The regressors are $x_t = Aw_t + v_t$. We randomly draw each element of $A \in \mathbb{R}^k$ from a uniform distribution on $[-1, 1]$. If A does not have full column rank then we add a randomly drawn ι from $[0, 1]$ to each diagonal component (in every case the resulting A had full column rank).

We fix $\delta = \mathbf{1}_{k_{\delta}}$, a $k_{\delta} \times 1$ vector of ones. The benchmark models are as follows. Under the null $\theta_0 = [\theta_{0,i}]_{i=1}^{k_{\theta,n}} = \mathbf{0}_{k_{\theta,n}}$. The alternatives are (i) $\theta_{0,1} = .001$ with $[\theta_{0,i}]_{i=2}^{k_{\theta,n}} = \mathbf{0}_{k_{\theta,n}-1}$; (ii) $\theta_{0,i} =$

$i/k_{\theta,n}$ for $i = 1, \dots, k_{\theta,n}$; and (iii) $\theta_{0,i} = .001$ for $i = 1, \dots, k_{\theta,n}$. See Table 1 for reference. Under (i) we have a small deviation from the null. At large deviations (e.g. $\theta_{0,1} = 1/2$) all tests are comparable. Small deviations like $\theta_{0,1} \in (0, .1]$, however, reveal test differences. That said, as we will see below the max-test generally dominates by size, and by power for small deviations.

Table 1: Alternative Models

Alternative (i) :	$\theta_{0,1} = .001$ and $[\theta_{0,i}]_{i=2}^{k_{\theta,n}} = 0$
Alternative (ii) :	$\theta_{0,i} = i/k_{\theta,n}$ for $i = 1, \dots, k_{\theta,n}$
Alternative (iii) :	$\theta_{0,i} = .001$ for $i = 1, \dots, k_{\theta,n}$

In case (iii) all $\theta_{0,i}$ have the same value. A max-test might plausibly be sub-optimal to a bootstrapped Wald test since the latter combines each estimated parameter, while the max-statistic works with the single largest estimate of a sequence of identical parameters. We will see below that contrary to this logic, the max-test works very well, precisely because it incorporates so little, yet relevant, information. A Wald statistic for large $k_{\theta,n}$ adds significant sampling error due to the inverted variance matrix, and due to low degrees of freedom by estimating the original high dimensional model.

Benchmark results are discussed in Section 3.2.1. In Section 3.2.2 we investigate additional models as robustness checks against the benchmark cases. We discuss the various tests next.

3.1.2 Max-Tests

We estimate $k_{\theta,n}$ parsimonious models $y_t = \delta_{(i)}^* x_{\delta,t} + \theta_i^* x_{\theta,i,t} + v_{(i),t}$ by least squares. Denote by \mathcal{T}_n the resulting max-test or max-t-test statistic. The bootstrapped test statistic $\tilde{\mathcal{T}}_n$ and p-value $\tilde{p}_{n,\mathcal{M}}$ are computed as in (6) and (7). We reject H_0 when $\tilde{p}_{n,\mathcal{M}} < \alpha$.

3.1.3 Wald Tests

We also perform asymptotic and bootstrapped Wald tests, and asymptotic and bootstrapped normalized Wald tests. We estimate $y_t = \delta'_0 x_{\delta,t} + \theta'_0 x_{\theta,t} + \epsilon_t$ by least squares, where $x_{\theta,t}$ has dimension $k_{\theta,n}$. The asymptotic Wald test is based on the $\chi^2(k_{\theta,n})$ distribution, which is only valid (asymptotically) for fixed $k_{\theta,n} = k$ or $k_{\theta,n} \rightarrow (0, \infty)$. The test leads to large empirical size distortions, discussed above, and is therefore not reported here.

The bootstrap Wald test is based on the fixed-design parametric wild bootstrap in [Gonçalves and Kilian \(2004\)](#). The algorithm is as follows. Let \mathcal{W}_n be the Wald statistic for a test of $H_0 : \theta_0 = \mathbf{0}_{k_{\theta,n}}$. Write $y_t = \delta'_0 x_{\delta,t} + \theta'_0 x_{\theta,t} + \epsilon_t = \beta'_0 x_t + \epsilon_t$ and $\epsilon_t(\beta) \equiv y_t - \beta' x_t$, and let $\hat{\beta}_n$ be the unrestricted least squares estimator of β_0 . Generate a sample $\{\hat{y}_t^*\}_{t=1}^n$ with the null imposed: $\hat{y}_t^* = \hat{\delta}'_n x_{\delta,t} + \epsilon_t(\hat{\beta}_n)\eta_t$ where η_t is iid $N(0,1)$. Construct the model $\hat{y}_t^* = b' x_t + v_t^*$, estimate b by least squares and compute a Wald statistic \mathcal{W}_n^* for the null hypothesis. Repeat \mathcal{M} times resulting in the sequence of bootstrapped Wald statistics $\{\mathcal{W}_{n,i}^*\}_{i=1}^{\mathcal{M}}$. The p-value approximation is $p_n = 1/M \sum_{i=1}^{\mathcal{M}} I(\mathcal{W}_n > \mathcal{W}_{n,i}^*)$, and we reject H_0 when $p_n < \alpha$.

The normalized Wald statistic is $\mathcal{W}_n^s \equiv (\mathcal{W}_n - k_{\theta,n})/\sqrt{2k_{\theta,n}}$. Under the null $\mathcal{W}_n^s \xrightarrow{d} N(0,1)$ as $k_{\theta,n} \rightarrow \infty$, and as long as $k_{\theta,n}/n \rightarrow 0$ then $\mathcal{W}_n^s \xrightarrow{p} \infty$ under H_1 . This asymptotic one-sided test rejects the null when $\mathcal{W}_n^s > Z_\alpha$, where Z_α is the standard normal upper tail α -level critical value. The bootstrapped test is performed as above: we compute $\mathcal{W}_{n,i}^{s*} \equiv (\mathcal{W}_{n,i}^* - k_{\theta,n})/\sqrt{2k_{\theta,n}}$ with p-value approximation $p_n = 1/M \sum_{i=1}^{\mathcal{M}} I(\mathcal{W}_n^s > \mathcal{W}_{n,i}^{s*})$. Trivially $\mathcal{W}_n^s > \mathcal{W}_{n,i}^{s*}$ if and only if $\mathcal{W}_{n,i}^* > \mathcal{W}_n$, hence the bootstrapped normalized Wald test and bootstrapped Wald test are identical. We therefore only discuss the bootstrapped Wald test.

3.2 Simulation Results

3.2.1 Benchmark Results

We report rejection frequencies in Tables 2 and 3 under zero nuisance terms $k_\delta = 0$. See Appendix D of the SM for complete simulation results.

Empirical Size Test results under the three covariate cases are similar, although all tests perform slightly less well under correlated regressors within and across blocks (case three) which we focus on.

The asymptotic Wald test and asymptotic normalized Wald test are generally severely over-sized due to the magnitude of $k_\delta + k_{\theta,n}$. The bootstrapped Wald test is strongly under-sized when n is small, but also when $k_\delta = 10$ and/or when $k_{\theta,n}$ is large.

The max-tests typically lead to qualitatively similar results with empirical size close to nominal size (see Table 2).

Empirical Power Consider Tables 2 and 3. Under alternative (*i*) only one parameter deviates from the null, $\theta_{0,1} = .001$. The difficulty in detecting this deviation is clearly apparent at smaller sample sizes. Across sample sizes and size of k_δ , moreover, the max-tests dominate the bootstrapped Wald test. The max-tests can achieve a power gain of as much as 40% over the Wald tests. For example, at $n = 100$, $k_\delta = 10$ and $k_{\theta,n} = 50$ the Wald test has zero power above nominal size while the max-test and max-t-test have power $\{.039, .149, .266\}$ and $\{.099, .267, .427\}$ for sizes $\{1\%, 5\%, 10\%\}$. At $n = 250$, $k_\delta = 10$ and $k_{\theta,n} = 79$, max-test and max-t-test power are $\{.514, .742, .838\}$ and $\{.617, .799, .873\}$, yet Wald test power is only $\{.008, .096, .260\}$. A similar result occurs at $n = 500$.

When there are zero nuisance covariates $k_\delta = 0$, the differences are just as stark. Max-test and max-t-test power when $n = 100$ and $k_{\theta,n} = 35$, for example, are $\{.451, .762, .870\}$ and $\{.531, .795, .890\}$, while Wald test power is $\{.020, .133, .251\}$. Only when $n = 1000$ does power

across tests become comparable. Thus, the max-tests dominate in terms of detecting a singular small deviation from the null.

Under alternative *(ii)* we have $\theta_{0,i} = i/k_{\theta,n}$ for $i = 1, \dots, k_{\theta,n}$: all parameters deviate from zero at values between $(0, 1]$. In this case the max- and max-t-tests, and the Wald test, are generally similar. This is not surprising since the (bootstrapped) Wald test becomes competitive under *(i)* when $\theta_{0,1}$ has values farther from zero.

Under *(iii)*, where $\theta_{0,i} = .001$ for $i = 1, \dots, k_{\theta,n}$, all tests are naturally better able to detect a deviation from the null than under *(i)*. The max-tests, however, dominate the remaining tests at $n \leq 250$, in particular when there are nuisance parameters $k_\delta > 0$. As with alternative *(i)*, under *(iii)* all tests are comparable when $\theta_{0,i}$ is larger, e.g. $\theta_{0,i} = 1/2$ for each i (not shown).

3.2.2 Robustness Checks

As robustness checks we now allow heterogeneous covariates, and large(r) k_δ .

a. Regressor Dispersion While the covariates $x_{\theta,i,t}$ are not homogeneous in all benchmark cases, their variances are fairly similar or identical depending on case. The dispersion of $x_{\theta,i,t}$, however, has a direct impact on the dispersion of the key parsimonious model estimators $\hat{\theta}_i$.

We therefore add three covariate cases allowing for different variances across $x_{\theta,t} = [x_{\theta,i,t}] \sim N(0, \Psi)$. We bypass covariate dependence and let $[x_{\delta,t}, x_{\theta,t}]$ be within and across block iid, in order to focus on the pure effects of covariate dispersion. When present, each nuisance $x_{\delta,i,t}$ is iid $N(0, 1)$. In each case we use a diagonal variance matrix $\Psi = [\Psi_{i,j}]$. In case *(a)* $\Psi_{i,i} = 1 + 100(i - 1)/k_{\theta,n} \in [1, 100]$ hence $x_{\theta,i,t}$ has a monotonically larger variance as i increases. In this setting a max-t-test may perform better due to self-scaling. In the remaining two cases diagonal Ψ has entries *(b)* $\Psi_{1,1} = 10$ or *(c)* $\Psi_{1,1} = 100$, and all other $\Psi_{i,i} = 1$, hence there is a single potentially influential regressor.

We only inspect test performance under the null. See Tables 5-7 in [Hill \(2023, Appendix](#)

E). Covariate dispersion matters more when there are large differences. The first case with increasing variance from 1 to 100 does not lead to major size distortions generally. The second and third cases, however, where there is one influential covariate, the Wald test can suffer major size distortions, while the max-test and max-t-test are similar, or in some cases the max-t-test has slightly better empirical size as anticipated. The max-tests are mildly oversized. The slight size distortions still exist when $n = 1000$, although they are not as attenuated as at smaller sample sizes. The Wald test suffers major size distortions even at $n = 1000$ when $k_{\theta,n}$ is large (in particular when $k_\delta > 0$).

b. Large k_δ Finally, we study two cases $k_\delta \in \{20, 40\}$ under the null, hence k_δ is larger than in the baseline cases $\{0, 10\}$. The covariates are within and across group iid standard normal. In view of the large number of parameters estimated, computation time is heavy, hence we only report results for $n \in \{100, 250, 500\}$. Results are contained in Table 8 in [Hill \(2023, Appendix E\)](#). The max-tests are slightly over-sized, in particular when $k_\delta = 40$, while the bootstrapped Wald test has greater size distortions. At lower sample sizes the max-test dominates, but the max-test and max-t-test are comparable at $n = 500$.

4 Conclusion

We present a class of tests of zero restrictions for a high dimensional parameter in a regression setting with nuisance parameters. We focus on a linear model and least squares estimation, but the method and theory presented here are generalizable to a broad class of models and estimators. We show how the hypotheses can be identified by using many potentially vastly lower dimension parameterizations. We use a test statistic that is the maximum in absolute value of the weighted key estimated parameters across the many low dimension settings. The lower dimension helps improve estimation accuracy, while a max-statistic alleviates the need for a multivariate normalization used in Wald and score statistics. Thus, we avoid inverting

a potentially large dimension variance estimator that may be a poor proxy for the true sampling dispersion. Indeed, our asymptotic theory sidesteps traditional extreme value theory arguments, while we instead focus on an approximate p-value computed by parametric wild bootstrap. In simulation experiments the max-tests generally have good or sharp size, and dominate a bootstrapped (and normalized) Wald test in terms of size and power.

An interesting alternative Hausman-version of our test could focus on the low dimensional difference $\hat{\delta}_{(i)} - \hat{\delta}^{(0)}$ (which under H_0 is zero asymptotically in probability across i). Notice $\hat{\delta}^{(0)}$ is the nuisance parameter estimator from the null model. This idea may be interesting to applied social scientists. Other future work may (i) lean toward allowing for a broader class of semi-nonparametric models, including models with a high dimensional parameter θ_0 and infinite dimensional unknown function h and/or high dimensional nuisance term δ_0 ; (ii) verify key asymptotic theory arguments in a general time series setting; and (iii) expand McKeague and Qian's (2015) significant predictor method to a high dimensional setting.

A Appendix

Throughout $O_p(1)$ and $o_p(1)$ are not functions of model counter i . $\{k_{\theta,n}, k_n\}$ are monotonically increasing sequences of positive integers. Define parsimonious parameter spaces $\mathcal{B}_{(i)} \equiv \mathcal{D} \times \Theta_{(i)}$ where $\mathcal{D} \subset \mathbb{R}^{k_\delta}$ and $\Theta_{(i)} \subset \mathbb{R}$ are compact subsets, δ_0 is an interior point of \mathcal{D} , and 0 and $\theta_{0,i}$ are interior points of $\Theta_{(i)}$. Recall $\mathcal{H}_{(i)} \equiv E[x_{(i),t}x'_{(i),t}]$.

A.1 Assumptions

Assumption 1.

- a. (ϵ_t, x_t) are iid over t , and mutually independent; $E[\epsilon_t] = 0$; $E[\epsilon_t^2] \geq \underline{c}$ for some $\underline{c} \in (0, \infty)$; $\limsup_{n \rightarrow \infty} |\theta_0| < \infty$.
- b. $E[(y_t - \beta_{(i)}^{*'} x_{(i),t})x_{(i),t}] = \mathbf{0}_{k_\delta+1}$ for all i .
- c. $\min_{i \in \mathbb{N}} \inf_{\lambda' \lambda = 1} E[(\lambda' x_{(i),t})^2] > 0$ and $\max_{i \in \mathbb{N}} \|\mathcal{H}_{(i)}^{-1}\| < \infty$.
- d. $\inf_{\lambda' \lambda = 1} \min_{i \in \mathbb{N}} \{1/n \sum_{t=1}^n (\lambda' x_{(i),t})^2\} > 0$ awp1; $\max_{1 \leq i \leq k} \|(1/n \sum_{t=1}^n x_{(i),t}x'_{(i),t})^{-1}\| = O_p(k)$, $k \geq 1$.

e. Let $P(|\epsilon_t| > c) \leq \mathcal{C} \exp\{-\mathcal{K}c\}$ and $\max_{i \in \mathbb{N}} P(|x_{i,t}| > c) \leq \mathcal{C} \exp\{-\mathcal{K}c\}$ for some finite $\mathcal{C}, \mathcal{K} > 0$ that may be different in different places.

Remark A.1. (a)-(d) are standard for linear iid regression models, augmented to allow for high dimensionality. $\limsup_{n \rightarrow \infty} |\theta_0| < \infty < \infty$ (under H_1) is used solely to ensure $\beta'_0 x_t$ has sub-exponential tails in order to ensure first and second order parsimonious least squares components have sub-exponential tails, cf. Lemma C.1 in the SM. Ultimately this yields required generic bounds for the first order expansion. Of course it only matters when $k_\theta \rightarrow \infty$ as $n \rightarrow \infty$.

Remark A.2. Recall $\widehat{\mathcal{H}}_{(i)} \equiv 1/n \sum_{t=1}^n x_{(i),t} x'_{(i),t}$. The (d) bound $\max_{1 \leq i \leq k} \|\widehat{\mathcal{H}}_{(i)}^{-1}\| = O_p(k)$ is mild considering $\|\widehat{\mathcal{H}}_{(i)}^{-1}\| = O_p(1)$ for each i by the iid property, tail bound (e) and the Slutsky theorem. Indeed, Markov's inequality, the iid property and (e) yield $P(\|\widehat{\mathcal{H}}_{(i)}\| > c) \leq K/c$ which does not depend on i . Now sub-additivity implies $P(\max_{1 \leq i \leq k} \|\widehat{\mathcal{H}}_{(i)}\| > c) \leq Kk/c$ hence $\max_{1 \leq i \leq k} \|\widehat{\mathcal{H}}_{(i)}\| = O_p(k)$.

Remark A.3. (e) imposes (uniform) sub-exponential tails for the error and regressors, which we use to yield a non-Gaussian first and second order approximation theory based on a *smooth-max* technique. See Lemmas A.1 and A.3, below, and their proofs in the SM. The tail property implies $(\epsilon_t, x_{i,t})$ have moment generating functions and therefore all moments (e.g. [Vershynin, 2018](#), Proposition 2.7.1). Sub-Gaussian and sub-exponential distribution classes provide a popular realm for developing high dimensional probability theory, like concentration bounds required here, covering at least distributions with bounded support, Gaussian, Laplace, exponential, Poisson, and Gamma random variables. See, e.g., [Chernozhukov, Chetverikov, and Kato \(2013\)](#) and [Vershynin \(2018\)](#), and see Lemma A.4.b, below. This ensures first and second order terms are locally sub-Gaussian, which ultimately suffices for our non-Gaussian expansion theory (cf. [Chareka, Chareka, and Kennedy, 2006](#)).

A.2 Proofs of Main Results

The proof of Lemma 2.2 exploits the following three results: bounds on gradient and Hessian moment generating function; a general bound on a sequence of random variables with a bounded moment generating function; and a high dimensional bound on the parsimonious gradient, Hessian and least squares estimator. See [Hill \(2023, Appendix C\)](#) for a proof of these and all subsequent supporting lemmas. Recall $\widehat{\mathcal{G}}_{(i)} \equiv -1/n \sum_{t=1}^n v_{(i),t} x_{(i),t}$ and $\widehat{\mathcal{H}}_{(i)} \equiv 1/n \sum_{t=1}^n x_{(i),t} x'_{(i),t}$.

Lemma A.1. *Let Assumption 1 hold.*

a. $\max_{1 \leq i \leq k_{\theta,n}} E[\exp\{\zeta |\sqrt{n} \widehat{\mathcal{G}}_{(i)}|\}] = O(n^{\xi \ln(k_n)})$ for some $\zeta, \xi > 0$;

b. $\max_{1 \leq i \leq k_{\theta,n}} E[\exp\{\zeta \sqrt{n} |\widehat{\mathcal{H}}_{(i)} - \mathcal{H}_{(i)}|\}] = O(n^{\xi \ln(k_n)})$ for some $\zeta, \xi > 0$.

Remark A.4. The proof actually shows each max-moment is bounded by $\exp\{\zeta^2 \mathcal{K}\}$. We only require the $O(n^{\xi \ln(k_n)})$ upper bound in order to satisfy the following lemma's generic high dimensional probability bound, and the Lemma A.3 high dimensional gradient and Hessian bound. This technically allows for heterogeneity in the tails of $(\epsilon_t, x_{j,t})$ ignored here.

The proof of the next general result relies on a smooth-max argument that exploits sub-exponential tails.

Lemma A.2. Let $\{\mathcal{X}_{n,i}\}_{n \geq 1}$ be random variables on a common probability space, $i = 1, 2, \dots$. For any sequence of positive integers $\{k_n\}$ if $\max_{1 \leq i \leq k_n} E[\exp\{\vartheta |\mathcal{X}_{n,i}|\}] = O(n^{\xi \ln(k_n)})$ for some $\vartheta, \xi > 0$ then $\max_{1 \leq i \leq k_n} |\mathcal{X}_{n,i}| = O_p(\ln(k_n) \ln(n))$.

Lemma A.3. Let Assumption 1 hold, and let $\{k_{\theta,n}\}$ be any monotonically increasing sequence of integers with $k_{\theta,n} = O(\sqrt{n}/(\ln(n))^2)$. Then $\max_{1 \leq i \leq k_{\theta,n}} |\sqrt{n} \widehat{\mathcal{G}}_{(i)}| = O_p(\ln(k_{\theta,n}) \ln(n))$, $\max_{1 \leq i \leq k_{\theta,n}} |\widehat{\mathcal{H}}_{(i)}^{-1} - \mathcal{H}_{(i)}^{-1}| = O_p(k_{\theta,n} \ln(k_{\theta,n}) \ln(n)/\sqrt{n})$ and $\max_{1 \leq i \leq k_{\theta,n}} |\sqrt{n}(\widehat{\beta}_{(i)} - \beta_{(i)}^*)| = O_p((\ln(n))^2)$.

Proof of Lemma 2.2. Under H_0 recall $v_{(i),t} = \epsilon_t$. By multiple uses of the triangle inequality, and $\widehat{\mathcal{Z}}_{(i)} \equiv \sqrt{n} \mathcal{H}_{(i)}^{-1} \widehat{\mathcal{G}}_{(i)} = -\mathcal{H}_{(i)}^{-1} n^{-1/2} \sum_{t=1}^n \epsilon_t x_{(i),t}$:

$$\begin{aligned} & \left| \max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{n} \mathcal{W}_{n,i} \widehat{\theta}_i \right| - \max_{1 \leq i \leq k_{\theta,n}} \left| \mathcal{W}_i [\mathbf{O}'_{k_\delta}, 1] \widehat{\mathcal{Z}}_{(i)} \right| \right| \\ & \leq \max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{n} \mathcal{W}_{n,i} \widehat{\theta}_i - \mathcal{W}_i [\mathbf{O}'_{k_\delta}, 1] \widehat{\mathcal{Z}}_{(i)} \right| \\ & \leq \max_{i \in \mathbb{N}} |\mathcal{W}_i| \times \max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{n} \widehat{\theta}_i - [\mathbf{O}'_{k_\delta}, 1] \widehat{\mathcal{Z}}_{(i)} \right| \\ & \quad + \max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_{n,i} - \mathcal{W}_i| \times \max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{n} \widehat{\theta}_i - [\mathbf{O}'_{k_\delta}, 1] \widehat{\mathcal{Z}}_{(i)} \right| \\ & \quad + \max_{1 \leq i \leq k_{\theta,n}} \left| [\mathbf{O}'_{k_\delta}, 1] \widehat{\mathcal{Z}}_{(i)} \right| \times \max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_{n,i} - \mathcal{W}_i|. \end{aligned}$$

By assumption $\max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_{n,i} - \mathcal{W}_i| = o_p(1/(\ln(n))^2)$ and $\max_{i \in \mathbb{N}} |\mathcal{W}_i| < \infty$. Furthermore $\max_{i \in \mathbb{N}} \|\mathcal{H}_{(i)}^{-1}\| < \infty$ under Assumption 1.c. Lemmas A.1 and A.2 yield $\max_{1 \leq i \leq k_{\theta,n}} |\sqrt{n} \widehat{\mathcal{G}}_{(i)}| = O_p(\ln(k_{\theta,n}) \ln(n))$. Hence since $k_{\theta,n} = o(n)$, $\max_{1 \leq i \leq k_{\theta,n}} |[\mathbf{O}'_{k_\delta}, 1] \widehat{\mathcal{Z}}_{(i)}| \times \max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_{n,i} - \mathcal{W}_i| = O_p(\ln(k_{\theta,n}) \ln(n)/(\ln(n))^2) = o_p(1)$. Similarly, from Lemma A.3 and $k_{\theta,n} = o(\sqrt{n}/(\ln(n))^4)$:

$$\max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{n} \widehat{\theta}_i - [\mathbf{O}'_{k_\delta}, 1] \widehat{\mathcal{Z}}_{(i)} \right| \leq \max_{1 \leq i \leq k_{\theta,n}} \left| \widehat{\mathcal{H}}_{(i)}^{-1} - \mathcal{H}_{(i)}^{-1} \right| \max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{n} \widehat{\mathcal{G}}_{(i)} \right| = O_p\left(\frac{k_{\theta,n} (\ln(n))^4}{\sqrt{n}}\right) = o_p(1).$$

Therefore

$$\begin{aligned}
& \left| \max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{n} \mathcal{W}_{n,i} \hat{\theta}_i \right| - \max_{1 \leq i \leq k_{\theta,n}} \left| \mathcal{W}_i [\mathbf{0}'_{k_\delta}, 1] \hat{\mathcal{Z}}_{(i)} \right| \right| \\
& \leq K \max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{n} \hat{\theta}_i - [\mathbf{0}'_{k_\delta}, 1] \hat{\mathcal{Z}}_{(i)} \right| + O_p \left(\frac{k_{\theta,n} (\ln(n))^4}{\sqrt{n}} \right) + O_p \left(\frac{\ln(k_{\theta,n})}{\ln(n)} \right) \\
& = \max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{n} \hat{\theta}_i - [\mathbf{0}'_{k_\delta}, 1] \hat{\mathcal{Z}}_{(i)} \right| + o_p(1).
\end{aligned}$$

It remains to prove $\max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{n} \hat{\theta}_i - [\mathbf{0}'_{k_\delta}, 1] \hat{\mathcal{Z}}_{(i)} \right| \xrightarrow{p} 0$.

By positive definiteness of $\widehat{\mathcal{H}}_{(i)}$ *awp1* we may write $\sqrt{n}(\hat{\beta}_{(i)} - \beta_{(i)}^*) = -\sqrt{n} \widehat{\mathcal{H}}_{(i)}^{-1} \widehat{\mathcal{G}}_{(i)}$ *awp1*, hence the parsimonious linear contrasts $\sqrt{n} \lambda'(\hat{\beta}_{(i)} - \beta_{(i)}^*)$ satisfy *awp1*:

$$\max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{n} \lambda' \left(\hat{\beta}_{(i)} - \beta_{(i)}^* \right) - \lambda' \hat{\mathcal{Z}}_{(i)} \right| \leq \max_{1 \leq i \leq k_{\theta,n}} \left| \lambda' \left\{ \widehat{\mathcal{H}}_{(i)}^{-1} - \mathcal{H}_{(i)}^{-1} \right\} \right| \times \max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{n} \widehat{\mathcal{G}}_{(i)} \right|.$$

By Lemma A.3 we have both $\max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{n} \widehat{\mathcal{G}}_{(i)} \right| = O_p(\ln(k_{\theta,n}) \ln(n))$, and

$$\max_{1 \leq i \leq k_{\theta,n}} \left| \widehat{\mathcal{H}}_{(i)}^{-1} - \mathcal{H}_{(i)}^{-1} \right| = O_p \left(\frac{k_{\theta,n} \ln(k_{\theta,n}) \ln(n)}{\sqrt{n}} \right) = O_p \left(\frac{k_{\theta,n} (\ln(n))^2}{\sqrt{n}} \right).$$

Thus, for any $\{k_{\theta,n}\}$, $k_{\theta,n} = o(\sqrt{n}/(\ln(n))^4)$:

$$\left| \sqrt{n} \lambda' \left(\hat{\beta}_{(i)} - \beta_{(i)}^* \right) - \lambda' \hat{\mathcal{Z}}_{(i)} \right| = O_p \left(\frac{k_{\theta,n} (\ln(n))^2}{\sqrt{n}} \times (\ln(n))^2 \right) = o_p(1).$$

The proof is complete by putting $\lambda = [\mathbf{0}'_{k_\delta}, 1]$ and imposing $H_0 : \theta_0 = \mathbf{0}_{k_\theta}$. *QED*.

The next result is used in the proof of Lemma 2.3.

Lemma A.4. *Assumption 1 yields the following:*

- $\inf_{\lambda' \lambda = 1} \lambda' \mathcal{H}_{(i)}^{-1} \lambda \in (\underline{c}, \bar{c})$ for some $0 < \underline{c} \leq \bar{c} < \infty$ uniformly in i .
- Write $z_{(i),t}(\lambda) \equiv -\lambda' \mathcal{H}_{(i)}^{-1} \epsilon_t x_{(i),t}$. Then $\max_{i \in \mathbb{N}} \{E|z_{(i),t}(\lambda)|^4 + E[\exp\{|z_{(i),t}(\lambda)|\}]\} \leq \mathcal{K}$ for $\lambda' \lambda = 1$, and some finite $\mathcal{K} > 1$.

Proof of Lemma 2.3. Let $\{k_{\theta,n}\}$ be any sequence of positive integers satisfying $(\ln(k_{\theta,n}n))^7/n \leq \mathcal{K}n^{-\zeta}$ with $\zeta \in (0, 1)$ and $\mathcal{K} > 0$. Coupled with Lemma A.4, the conditions of Corollary 2.1 of Chernozhukov, Chetverikov, and Kato (2013) hold based on their (E.1) with $B_n = B$ and sufficiently large $B > 0$. Therefore $\rho_n(\lambda) = O(n^{-c})$ for some $c > 0$. This yields the desired result for any $\{k_{\theta,n}\}$ satisfying $k_{\theta,n} \leq \exp\{\mathcal{K}n^d\}/n$ for some $\mathcal{K} > 0$ a function of B , and some $d \in (0, 1/7)$. *QED*.

Proof of Theorem 2.4. For (a), convergence in distribution (5) follows from the imposed properties on $\mathcal{W}_{n,i}$ with asymptotic expansion Lemma 2.2, and convergence in Kolmogorov distance (4), cf. Lemma 2.3 with $\lambda = [\mathbf{0}'_{k_\delta}, 1]'$. Then $\max_{1 \leq i \leq k_{\theta,n}} |\sqrt{n} \mathcal{W}_{n,i} \hat{\theta}_i| \xrightarrow{d} \max_{i \in \mathbb{N}} |\mathcal{W}_i \mathbf{Z}_{(i)}([\mathbf{0}'_{k_\delta}, 1])|$ follows immediately.

(b) follows from Lemma A.3 consistency of $\hat{\theta}_i$, and $\mathcal{W}_{n,i} \xrightarrow{p} \mathcal{W}_i$ with $\mathcal{W}_i \in (0, \infty)$. \mathcal{QED} .

The bootstrap theory requires general high dimensional bounds, and a first order approximation for the bootstrapped estimator $\tilde{\theta}_i$, similar to Lemma 2.2.

Lemma A.5. *Let $\{k_n\}_{n \geq 1}$ be any sequence of positive integers with $k_n = O(n)$. Let $\{a_{i,t}\}_{t=1}^\infty$ be random variables on a common probability space, independent across t , for $i = 1, 2, \dots$. If $\max_{i \in \mathbb{N}} E[a_{i,t}^2] < \infty$ then $\max_{1 \leq i \leq k_n} |1/n \sum_{t=1}^n (\eta_t^2 a_{i,t} - E[a_{i,t}])| = O_p(k_n/\sqrt{n})$.*

Recall the bootstrapped first order term $\tilde{\mathbf{Z}}_{(i)}^{(0)}$ in (8).

Lemma A.6. *Let Assumption 1 hold, and let the weights satisfy $\max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_{n,i} - \mathcal{W}_i| = o_p(1/(\ln(n))^2)$ for non-stochastic $\mathcal{W}_i \in (0, \infty)$. Then for any monotonic sequence of positive integers $\{k_{\theta,n}\}$, $k_{\theta,n} = o(n)$:*

$$\left| \max_{1 \leq i \leq k_{\theta,n}} |\sqrt{n} \mathcal{W}_{n,i} \hat{\theta}_i| - \max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_i [\mathbf{0}'_{k_\delta}, 1] \tilde{\mathbf{Z}}_{(i)}^{(0)}| \right| = O_p\left(\frac{(\ln(n))^4}{\sqrt{n}}\right).$$

Proof of Lemma 2.5.

Step 1. Recall $\tilde{\mathbf{Z}}_{(i)}^{(0)} = -\bar{\mathcal{H}}_{(i)}^{(0)-1} n^{-1/2} \sum_{t=1}^n \eta_t G_{i,t}^{(0)}$ where $G_{i,t}^{(0)} \equiv (y_t - \delta^{(0)' } x_{\delta,t}) x_{\delta,t}$ and $\bar{\mathcal{H}}_{(i)}^{(0)} \equiv E[x_{\delta,t} x'_{\delta,t}]$, and η_t is iid $N(0, 1)$ and independent of the sample $\mathfrak{S}_n \equiv \{x_t, y_t\}_{t=1}^n$. As always we only care about $\lambda = [\mathbf{0}'_{k_\delta}, 1]'$, hence we may assume $\|\lambda\| = 1$. Write

$$\hat{\sigma}_{n,(ij)}^2(\lambda) \equiv \lambda' E \left[\tilde{\mathbf{Z}}_{(i)}^{(0)} \tilde{\mathbf{Z}}_{(j)}^{(0)} | \mathfrak{S}_n \right] \lambda = \lambda' \bar{\mathcal{H}}_{(i)}^{(0)-1} \frac{1}{n} \sum_{t=1}^n \eta_t^2 G_{i,t}^{(0)} G_{j,t}^{(0)' } \bar{\mathcal{H}}_{(i)}^{(0)-1} \lambda$$

$$\tilde{\sigma}_{(ij)}^2(\lambda) \equiv \lambda' \bar{\mathcal{H}}_{(i)}^{(0)-1} E \left[G_{i,t}^{(0)} G_{j,t}^{(0)' } \right] \bar{\mathcal{H}}_{(j)}^{(0)-1} \lambda.$$

By construction $\lambda' \tilde{\mathbf{Z}}_{(i)}^{(0)} | \mathfrak{S}_n \sim N(0, \hat{\sigma}_{n,(ii)}^2(\lambda))$. Recall $\{\tilde{\mathbf{Z}}_{(i)}(\lambda)\}_{n \in \mathbb{N}}$ are sequences with $\tilde{\mathbf{Z}}_{(i)}(\lambda) \sim N(0, \tilde{\sigma}_{(ii)}^2(\lambda))$ independent of \mathfrak{S}_n . Define $\Delta_{k_{\theta,n}}(\lambda) \equiv \max_{1 \leq i, j \leq k_{\theta,n}} |\hat{\sigma}_{n,(ij)}^2(\lambda) - \tilde{\sigma}_{(ij)}^2(\lambda)|$. Lemma 3.1 in Chernozhukov, Chetverikov, and Kato (2013) therefore yields the following conditional Slepian-type inequality (cf. Chernozhukov, Chetverikov, and Kato, 2015, Theorem 2, Proposition 1):

$$\begin{aligned} \mathcal{E}_{k_{\theta,n}} &\equiv \sup_{z \geq 0} \left| P \left(\max_{1 \leq i \leq k_{\theta,n}} |[\mathbf{0}'_{k_\delta}, 1] \tilde{\mathbf{Z}}_{(i)}^{(0)}| \leq z | \mathfrak{S}_n \right) - P \left(\max_{1 \leq i \leq k_{\theta,n}} |\tilde{\mathbf{Z}}_{(i)}([\mathbf{0}'_{k_\delta}, 1]')| \leq z \right) \right| \\ &= O_p \left(\Delta_{k_{\theta,n}}^{1/3}(\lambda) \max \{1, \ln(k_{\theta,n}/\Delta_{k_{\theta,n}}(\lambda))\}^{2/3} \right). \end{aligned}$$

The proof is therefore complete if we show $\mathcal{E}_{k_{\theta,n}} = o_p(1)$. We need to bound $\Delta_{k_{\theta,n}}(\lambda)$. By adding and subtracting terms, and the triangle inequality:

$$\Delta_{k_{\theta,n}}(\lambda) \leq \max_{i \in \mathbb{N}} \left\| \bar{\mathcal{H}}_{(i)}^{-1} \right\|^2 \times \max_{1 \leq i, j \leq k_{\theta,n}} \left| \frac{1}{n} \sum_{t=1}^n \left(\eta_t^2 G_{i,t}^{(0)} G_{j,t}^{(0)'} - E \left[G_{i,t}^{(0)} G_{j,t}^{(0)'} \right] \right) \right|.$$

Under Assumption 1 $\max_{i \in \mathbb{N}} \|\bar{\mathcal{H}}_{(i)}^{-1}\| < \infty$, and $E[\eta_t^2 G_{i,t}^{(0)} G_{j,t}^{(0)'}] = E[G_{i,t}^{(0)} G_{j,t}^{(0)'}]$ by mutual independence and $\eta_t \sim N(0, 1)$. Furthermore, by Lemma C.1 in the SM $|(y_t - \delta^{(0)'} x_{\delta,t}) x_{(i),t}|$ have uniformly sub-exponential tails, and therefore all moments uniformly over i in view of (B.2) in the SM. The Cauchy-Schwartz inequality therefore yields

$$\sup_{\lambda' \lambda = 1} E \left[\left(\lambda' G_{i,t}^{(0)} G_{j,t}^{(0)'} \lambda \right)^2 \right] \leq \sup_{\lambda' \lambda = 1} E \left[(y_t - \delta^{(0)'} x_{\delta,t})^2 (\lambda' x_{(i),t})^4 \right] < \infty.$$

Lemma A.5 thus applies to $G_{i,t}^{(0)} G_{j,t}^{(0)'}$, yielding: $\max_{1 \leq i, j \leq k_{\theta,n}} |1/n \sum_{t=1}^n (\eta_t^2 G_{i,t}^{(0)} G_{j,t}^{(0)'} - E[G_{i,t}^{(0)} G_{j,t}^{(0)'}])| = O_p(k_{\theta,n}/\sqrt{n})$. Hence $\Delta_{k_{\theta,n}}(\lambda) = O_p(k_{\theta,n}/\sqrt{n})$. Now use $k_{\theta,n} = o(\sqrt{n}/(\ln(n))^2)$ to conclude:

$$\mathcal{E}_{k_{\theta,n}} = O_p \left(\Delta_{k_{\theta,n}}^{1/3}(\lambda) \max \{1, \ln(k_{\theta,n}/\Delta_{k_{\theta,n}}(\lambda))\}^{2/3} \right) = O_p \left(\left(\frac{k_{\theta,n}}{\sqrt{n}} \right)^{1/3} \ln(n)^{2/3} \right) = o_p(1)$$

Step 2. Step 1 yields $\max_{1 \leq i \leq k_{\theta,n}} |[\mathbf{0}'_{k_{\delta}}, 1] \tilde{\mathcal{Z}}_{(i)}^{(0)}| | \mathfrak{S}_n \xrightarrow{d} \max_{i \in \mathbb{N}} | \tilde{\mathcal{Z}}_{(i)}([\mathbf{0}'_{k_{\delta}}, 1])|$ *awp1* with respect to $\{x_t, y_t\}_{t=1}^{\infty}$. Therefore $\max_{1 \leq i \leq k_{\theta,n}} |[\mathbf{0}'_{k_{\delta}}, 1] \tilde{\mathcal{Z}}_{(i)}^{(0)}| \Rightarrow^p \max_{i \in \mathbb{N}} | \tilde{\mathcal{Z}}_{(i)}([\mathbf{0}'_{k_{\delta}}, 1])|$; see. [Gine and Zinn \(1990, Section 3\)](#). \mathcal{QED} .

Proof of Lemma 2.6. Bootstrap asymptotic expansion Lemma A.6, and bootstrap Gaussian approximation Lemma 2.5, coupled with $\max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_{n,i} - \mathcal{W}_i| = o_p(1/(\ln(n))^2)$ for non-stochastic $\mathcal{W}_i \in (0, \infty)$, prove the claim. \mathcal{QED} .

Proof of Theorem 2.7.

Claim (a). We have $\max_{1 \leq i \leq k_{\theta,n}} |\sqrt{n} \mathcal{W}_{n,i} \hat{\theta}_i| \xrightarrow{d} \max_{i \in \mathbb{N}} |\mathcal{W}_i \mathbf{Z}_{(i)}([\mathbf{0}'_{k_{\delta}}, 1])|$ from Theorem 2.4.a. Lemma 2.6 yields $\max_{1 \leq i \leq k_{\theta,n}} \tilde{\mathcal{T}}_n \Rightarrow^p \max_{i \in \mathbb{N}} |\mathcal{W}_i \tilde{\mathcal{Z}}_{(i)}([\mathbf{0}'_{k_{\delta}}, 1])|$, where $\{\tilde{\mathcal{Z}}_{(i)}(\lambda)\}_{i \in \mathbb{N}}$ is an independent copy of $\{\mathbf{Z}_{(i)}(\lambda)\}_{i \in \mathbb{N}}$, $\mathbf{Z}_{(i)}(\lambda) \sim N(0, E[\epsilon_t^2] \lambda' \mathcal{H}_{(i)}^{-1} \lambda)$, independent of the asymptotic draw $\{x_t, y_t\}_{t=1}^{\infty}$. The proof now follows from arguments in [Hansen \(1996, p. 427\)](#).

Claim (b). Let H_1 hold, suppose $\theta_{0,j} \neq 0$ for some j , define $\mathfrak{S}_n \equiv \{x_t, y_t\}_{t=1}^n$, and write $\mathcal{P}_{\mathfrak{S}_n}(\mathcal{A}) \equiv \mathcal{P}(\mathcal{A} | \mathfrak{S}_n)$. In view of $\max_{1 \leq i \leq k_{\theta,n}} \tilde{\mathcal{T}}_n \Rightarrow^p \max_{i \in \mathbb{N}} |\mathcal{W}_i \tilde{\mathcal{Z}}_{(i)}([\mathbf{0}'_{k_{\delta}}, 1])|$ we have ([Gine and Zinn, 1990, eq. \(3.4\)](#))

$$\sup_{z > 0} \left| \mathcal{P}_{\mathfrak{S}_n} \left(\tilde{\mathcal{T}}_n \leq z \right) - P \left(\max_{1 \leq i \leq k_{\theta,n}} \left| \mathcal{W}_i \tilde{\mathcal{Z}}_{(i)}([\mathbf{0}'_{k_{\delta}}, 1]) \right| \leq z \right) \right| \xrightarrow{p} 0,$$

where $\{\check{\mathbf{Z}}_{(i)}(\cdot)\}_{i \in \mathbb{N}}$ is an independent copy of $\{\mathbf{Z}_{(i)}(\cdot)\}_{i \in \mathbb{N}}$. Moreover, $\tilde{p}_{n, \mathcal{M}_n} = \mathcal{P}_{\mathfrak{S}_n}(\tilde{\mathcal{T}}_{n,1} > \mathcal{T}_n) + o_p(1)$ by the Glivenko-Cantelli theorem and independence across bootstrap samples. Further, $\mathcal{T}_n \geq |\sqrt{n}\mathcal{W}_{n,j}(\hat{\theta}_j - \theta_{0,j}) + \sqrt{n}\mathcal{W}_{n,j}\theta_{0,j}| = \sqrt{n}\mathcal{W}_j|\theta_{0,j}|(|\hat{\theta}_j/\theta_{0,j}| + o_p(1)) \xrightarrow{p} \infty$ by Theorem 2.4.b and $\max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_{n,i} - \mathcal{W}_i| = o_p(1)$. Therefore

$$\tilde{p}_{n, \mathcal{M}_n} \leq P \left(\max_{1 \leq i \leq k_{\theta,n}} \left| \mathcal{W}_i \check{\mathbf{Z}}_{(i)}([\mathbf{0}'_{k_\delta}, 1]) \right| > \sqrt{n}\mathcal{W}_j|\theta_{0,j}| \left(\left| \frac{\hat{\theta}_j}{\theta_{0,j}} \right| + o_p(1) \right) \right) + o_p(1) \rightarrow 0$$

as long as $\max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_i \check{\mathbf{Z}}_{(i)}([\mathbf{0}'_{k_\delta}, 1])| = o_p(\sqrt{n})$. We show below given Gaussianicity of $\check{\mathbf{Z}}_{(i)}([\mathbf{0}'_{k_\delta}, 1])$:

$$\max_{1 \leq i \leq k_{\theta,n}} \left| \mathcal{W}_i \check{\mathbf{Z}}_{(i)}([\mathbf{0}'_{k_\delta}, 1]) \right| = O_p \left(\sqrt{\ln(k_{\theta,n})} \right) \quad (\text{A.1})$$

hence $\tilde{p}_{n, \mathcal{M}_n} \xrightarrow{p} 0$ under H_1 and therefore $P(\tilde{p}_{n, \mathcal{M}_n} < \alpha) \rightarrow 1$ as claimed.

We now prove (A.1). Put $\zeta = \sqrt{2\mathcal{W}E[\epsilon_t^2]\mathcal{H}^{-1, \theta, \theta}} > 0$ where $\mathcal{W} \equiv \max_{i \in \mathbb{N}} \{\mathcal{W}_i\} < \infty$, and $\mathcal{H}^{-1, \theta, \theta} \equiv \max_{1 \leq i \leq k_{\theta,n}} [\mathbf{0}'_{k_\delta}, 1] \mathcal{H}_{(i)}^{-1}[\mathbf{0}'_{k_\delta}, 1]' \in (0, \infty)$ under positive definiteness Assumption 1.c. Then Chernoff's Gaussian concentration bound $P(|Y| > c) \leq e^{-c^2/(2\sigma^2)}$ for $Y \sim N(0, \sigma^2)$, and Boole's inequality yield for any $\xi > 0$:

$$\begin{aligned} P \left(\max_{1 \leq i \leq k_{\theta,n}} \left| \mathcal{W}_i \check{\mathbf{Z}}_{(i)}([\mathbf{0}'_{k_\delta}, 1]) \right| > \zeta \sqrt{\ln(k_{\theta,n})} + \xi \right) & \quad (\text{A.2}) \\ & \leq \sum_{i=1}^{k_{\theta,n}} P \left(\left| \mathcal{W}_i \check{\mathbf{Z}}_{(i)}([\mathbf{0}'_{k_\delta}, 1]) \right| > \zeta \sqrt{\ln(k_{\theta,n})} + \xi \right) \\ & \leq \sum_{i=1}^{k_{\theta,n}} \exp \left\{ -\frac{\left(\zeta \sqrt{\ln(k_{\theta,n})} + \xi \right)^2}{2\mathcal{W}E[\epsilon_t^2]\mathcal{H}^{-1, \theta, \theta}} \right\} \\ & = \frac{1}{k_{\theta,n}} \sum_{i=1}^{k_{\theta,n}} \exp \left\{ -\frac{2\xi \sqrt{\ln(k_{\theta,n})}}{\zeta} \right\} \exp \left\{ -\frac{\xi^2}{\zeta^2} \right\} \leq \exp \{ -\xi^2/\zeta^2 \}. \quad \mathcal{QED}. \end{aligned}$$

Proof of Theorem 2.8. Let $k_{\theta,n} = o(\sqrt{n}/(\ln(n))^4)$. Under H_1^L by the same arguments used to prove Lemma 2.8 and Theorem 2.4,

$$\left| \max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{n} \left(\mathcal{W}_{n,i}(\hat{\theta}_i - \frac{c_i}{\sqrt{n/\ln(k_{\theta,n})}}) \right) \right| - \max_{1 \leq i \leq k_{\theta,n}} \left| \mathbf{Z}_{(i)}([\mathbf{0}'_{k_\delta}, 1]) \right| \right| \xrightarrow{p} 0, \quad (\text{A.3})$$

where $\mathbf{Z}_{(i)}([\mathbf{0}'_{k_\delta}, 1]) \sim N(0, 2E[\epsilon_t^2]\mathcal{H}^{-1, \theta, \theta})$ where $\mathcal{H}^{-1, \theta, \theta} \equiv \max_{1 \leq i \leq k_{\theta,n}} [\mathbf{0}'_{k_\delta}, 1] \mathcal{H}_{(i)}^{-1}[\mathbf{0}'_{k_\delta}, 1]'$.

Now define for arbitrary $d = [d_i]_{i=1}^{k_\theta}$ and sample $\mathfrak{S}_n = \{x_t, y_t\}_{t=1}^n$:

$$\mathcal{P}_n(d) \equiv P \left(\max_{1 \leq i \leq k_{\theta,n}} \left| \mathcal{W}_i \dot{\mathbf{Z}}_{(i)}([\mathbf{0}'_{k_\delta}, 1]) \right| / \sqrt{\ln(k_{\theta,n})} > \max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{\frac{n}{\ln(k_{\theta,n})}} \mathcal{W}_{n,i} \left(\hat{\theta}_i - c_i \sqrt{\frac{\ln(k_{\theta,n})}{n}} \right) + d_i \right| \middle| \mathfrak{S}_n \right),$$

where $\{\dot{\mathbf{Z}}_{(i)}(\cdot)\}_{i \in \mathbb{N}}$ is an independent copy of $\{\mathbf{Z}_{(i)}(\cdot)\}_{i \in \mathbb{N}}$. By the Glivenko-Cantelli Theorem with $\mathcal{M} = \mathcal{M}_n \rightarrow \infty$, the triangle inequality, and $\max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_{n,i} - \mathcal{W}_i| = o_p(1/(\ln(n))^2)$, and arguments in the proof of Theorem 2.7.b, the bootstrapped p-value approximation satisfies:

$$\begin{aligned} \tilde{p}_{n,\mathcal{M}} &= P \left(\max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{n} \mathcal{W}_{n,i} \hat{\theta}_i \right| > \max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{n} \mathcal{W}_{n,i} \left(\hat{\theta}_i - \frac{c_i}{\sqrt{n/\ln(k_{\theta,n})}} \right) + \mathcal{W}_i c_i \sqrt{\ln(k_{\theta,n})} \right| \middle| \mathfrak{S}_n \right) + o_p(1) \\ &= P \left(\max_{1 \leq i \leq k_{\theta,n}} \left| \mathcal{W}_i \dot{\mathbf{Z}}_{(i)}([\mathbf{0}'_{k_\delta}, 1]) \right| / \sqrt{\ln(k_{\theta,n})} > \max_{1 \leq i \leq k_{\theta,n}} \left| \sqrt{\frac{n}{\ln(k_{\theta,n})}} \mathcal{W}_{n,i} \left(\hat{\theta}_i - \frac{c_i}{\sqrt{n/\ln(k_{\theta,n})}} \right) + \mathcal{W}_i c_i \right| \middle| \mathfrak{S}_n \right) + o_p(1) \\ &= \mathcal{P}_n(\mathcal{W}_j c_j) + o_p(1). \end{aligned}$$

The proof of Theorem 2.7.a and arguments in Hansen (1996, p. 427) yield that $\mathcal{P}_n(\mathbf{0}_{k_{\theta,n}})$ is asymptotically uniformly distributed. Now, in view of (A.1) $\max_{1 \leq i \leq k_{\theta,n}} |\mathcal{W}_i \dot{\mathbf{Z}}_{(i)}([\mathbf{0}'_{k_\delta}, 1])| = O_p(\sqrt{\ln(k_{\theta,n})})$, and $\max_{1 \leq i \leq k_{\theta,n}} |\sqrt{n/\ln(k_{\theta,n})} \mathcal{W}_{n,i} (\hat{\theta}_i - c_i/\sqrt{n/\ln(k_{\theta,n})})| \xrightarrow{p} 0$ from (A.3). Thus $\lim_{n \rightarrow \infty} P(\tilde{p}_{n,\mathcal{M}_n} < \alpha) \nearrow 1$ monotonically as the maximum local drift $\max_{i \in \mathbb{N}} |c_i| \nearrow \infty$ since $\mathcal{W}_i \in (0, \infty) \forall i$. \mathcal{QED}

References

- AMEMIYA, T. (1980): “Selection of Regressors,” *International Economic Review*, 21, 331–354.
- ANDREWS, D. W. K. (1991): “Asymptotic Normality of Series Estimators for Nonparametric and Semiparametric Regression Models,” *Econometrica*, 59, 307–345.
- ARIAS-CASTRO, E., E. J. CANDÈS, AND Y. PLAN (2011): “Global Testing Under Sparse Alternative: ANOVA, Multiple Comparisons and the Higher Criticism,” *Annals of Statistics*, 39, 2533–2556.
- BARAUD, Y., C. GIRAUD, AND S. HUAT (2009): “Gaussian Model Selection with an Unknown Variance,” *Annals of Statistics*, 37, 630–672.
- BELLONI, A., V. CHERNOZHUKOV, I. FERNANDEZ-VAL, AND C. HANSEN (2017): “Program Evaluation and Causal Inference With High-Dimensional Data,” *Econometrica*, 85, 233–298.

- BELLONI, A., V. CHERNOZHUKOV, AND C. HANSEN (2014): “High-Dimensional Methods and Inference on Structural and Treatment Effects,” *Journal of Economic Perspectives*, 28, 29–50.
- BICKEL, P. J., AND D. A. FREEDMAN (1983): “Bootstrapping Regression Models with Many Parameters,” in *A Festschrift for Erich L. Lehmann*, ed. by P. Bickel, K. Doksum, J. Hodges, and R. BocaRaton. Chapman and Hall.
- BOSE, A. (1988): “Edgeworth Correction by Bootstrap in Autoregressions,” *Annals of Statistics*, 16, 1709–1722.
- BÜHLMANN, P. (2013): “Statistical Significance in High-Dimensional Linear Models,” *Bernoulli*, 19, 1212–1242.
- CAI, T., AND M. YUAN (2012): “Minimax and Adaptive Prediction for Functional Linear Regression,” *Journal of the American Statistical Association*, 107, 1201–1216.
- CARPENTIER, A., O. COLLIER, L. COMMINGES, A. B. TSYBAKOV, AND Y. WANG (2019): “Minimax Rate of Testing in Sparse Linear Regression,” *Automation and Remote Control*, 80, 1817–1834.
- CARPENTIER, A., AND N. VERZELEN (2019): “Adaptive Estimation of the Sparsity in the Gaussian Vector Model,” *Annals of Statistics*, 47, 93–126.
- CASTLE, J., X. QIN, AND W. R. REED (2009): “How To Pick The Best Regression Equation: A Review And Comparison Of Model Selection Algorithms,” Discussion Paper 13/2009, Dept. of Economics and Finance, University of Canterbury.
- CATTANEO, M. D., M. JANSSON, AND W. NEWEY (2018): “Inference in Linear Regression Models with Many Covariates and Heteroscedasticity,” *Journal of the American Statistical Association*, 113, 1350–1361.
- CHAREKA, P., O. CHAREKA, AND S. KENNEDY (2006): “Locally Sub-Gaussian Random Variables and the Strong Law of Large Numbers,” *Atlantic Electronic Journal of Mathematics*, 1, 75–81.
- CHEN, X., AND D. POUZO (2015): “Sieve Wald and QLR Inference on Semi-Nonparametric Conditional Moment Models,” *Econometrica*, 83, 1013–1079.
- CHEN, X., M. XU, AND W. B. WU (2013): “Covariance and Precision Matrix Estimation for High-Dimensional Time Series,” *Annals of Statistics*, 41, 2994–3021.
- CHERNOZHUKOV, V., D. CHETVERIKOV, M. DEMIRER, E. DUFLO, C. HANSEN, W. NEWEY, AND J. ROBINS (2016): “Double/Debiased Machine Learning for Treatment and Structural Parameters,” *Econometrics Journal*, 21, C1–C68.
- CHERNOZHUKOV, V., D. CHETVERIKOV, AND K. KATO (2013): “Gaussian Approximations and Multiplier Bootstrap for Maxima of Sums of High-Dimensional Random Vectors,” *Annals of Statistics*, 41, 2786–2819.
- (2015): “Comparison and Anti-Concentration Bounds for Maxima of Gaussian Random Vectors,” *Probability Theory and Related Fields*, 162, 47–70.
- (2017): “Central Limit Theorems and Bootstrap in High Dimensions,” *Annals of Probability*, 45, 2309–2352.

- FAN, J., AND J. LV (2008): “Sure Independence Screening for Ultrahigh Dimensional Feature Space,” *Journal of the Royal Statistical Society, Series B*, 70, 849–911.
- FAN, J., J. LV, AND . L. QI (2011): “Sparse High-Dimensional Models in Economics,” *Annual Economic Review*, 3, 291–317.
- FAN, J., AND H. PENG (2004): “Nonconcave Penalized Likelihood with a Diverging Number of Parameters,” *Annals of Statistics*, 32, 928–961.
- GAUTIER, E., AND C. ROSE (2021): “High-Dimensional Instrumental Variables Regression and Confidence Sets,” hal-00591732v7.
- GENOVESE, C. R., J. JIN, L. WASSERMAN, AND Z. YAO (2023): “A Comparison of the Lasso and Marginal Regression,” *Journal of Machine Learning Research*, 13, 2107–2143.
- GHYSELS, E., J. B. HILL, AND K. MOTEGI (2020): “Testing a Large Set of Zero Restrictions in Regression Models, with an Application to Mixed Frequency Granger Causality,” *Journal of Econometrics*, 218, 633–654.
- GINE, E., AND J. ZINN (1990): “Bootstrapping General Empirical Measures,” *Annals of Probability*, 18, 851–869.
- GONÇALVES, S., AND L. KILIAN (2004): “Bootstrapping Autoregressions with Conditional Heteroskedasticity of Unknown Form,” *Journal of Econometrics*, 123, 89–120.
- HALL, P., AND J. HOROWITZ (2007): “Methodology and Convergence Rates for Functional Linear Regression,” *Annals of Statistics*, 35, 70–91.
- HANSEN, B. E. (1996): “Inference When a Nuisance Parameter Is Not Identified Under the Null Hypothesis,” *Econometrica*, 64, 413–430.
- HASTIE, T., R. TIBSHIRANI, AND J. FRIEDMAN (2009): *The Elements of Statistical Learning: Data Mining, Inference, and Prediction*, Springer Series in Statistics. Springer, New York, 2nd edn.
- HILGERT, N., A. MAS, AND N. VERZELEN (2013): “Minimax Adaptive Tests for the Functional Linear Model,” *Annals of Statistics*, 41, 838–869.
- HILL, J. B. (2023): “Supplemental Material for *Testing Many Zero Restrictions in a High Dimensional Linear Regression Setting*,” Dept. of Economics, University of North Carolina.
- HILL, J. B., AND K. MOTEGI (2020): “Max-Correlation White Noise Test for Weakly Dependent Time Series,” *Econometric Theory*, 36, 907–960.
- HUANG, J., J. L. HOROWITZ, AND S. MA (2008): “Asymptotic Properties of Bridge Estimators in Sparse High-Dimensional Regression Models,” *Annals of Statistics*, 36, 587–613.
- HUBER, P. (1973): “Robust Regression: Asymptotics, Conjectures and Monte Carlo,” *Annals of Statistics*, 1, 799–821.
- INGSTER, Y. I. (1997): “Some Problems of Hypothesis Testing Leading to Infinitely Divisible Distributions,” *Mathematical Methods of Statistics*, 6, 47–49.
- INGSTER, Y. I., AND T. SAPATINAS (2009): “Minimax Goodness-of-Fit Testing in Multivariate Nonparametric Regression,” *Mathematical Methods of Statistics*, 18, 241–269.

- KABE, D. G. (1963): “Stepwise Multivariate Linear Regression,” *Journal of the American Statistical Association*, 58, 770–773.
- LI, C., AND U. K. MÜLLER (2021): “Linear Regression with Many Controls of Limited Explanatory Power,” *Quantitative Economics*, 12, 405–442.
- LIU, R. Y. (1988): “Bootstrap Procedures under some Non-I.I.D. Models,” *Annals of Statistics*, 16, 1696–1708.
- MAMMEN, E. (1993): “Bootstrap and Wild Bootstrap for High Dimensional Linear Models,” *Annals of Statistics*, 21, 255–285.
- MCKEAGUE, I., AND M. QIAN (2015): “An Adaptive Resampling Test for Detecting the Presence of Significant Predictors,” *Journal of the American Statistical Association*, 110, 1422–1433.
- POLLARD, D. (1984): *Convergence of Stochastic Processes*. Springer, New York.
- ROYSTON, P., AND D. G. ALTMAN (1994): “Regression Using Fractional Polynomials of Continuous Covariates: Parsimonious Parametric Modelling,” *Journal of the Royal Statistical Society, Series B*, 43, 429–467.
- STONE, C. J. (1981): “Admissible Selection of an Accurate and Parsimonious Normal Linear Regression Model,” *Annals of Statistics*, 9, 475–485.
- VA DE GREER, S. A. (2008): “High-Dimensional Generalized Linear Models and The Lasso,” *Annals of Statistics*, 36, 614–645.
- VERSHYNIN, R. (2018): *High-Dimensional Probability*. Cambridge University Press, Cambridge, UK.
- ZHANG, C.-H., AND S. S. ZHANG (2014): “Confidence Intervals for Low Dimensional Parameters in High Dimensional Linear Models,” *Journal of the Royal Statistical Society, Series B*, 76, 217–242.
- ZHONG, P.-S., S. X. CHEN, AND M. XU (2013): “Tests Alternative to Higher Criticism High-Dimensional Mean under Sparsity and Column-Wise Dependence,” *Annals of Statistics*, 41, 2820–2851.

Table 2: Rejection Frequencies under H_0 and $H_1(i)$

	$H_0 : \theta_0 = 0$									$H_1(i) : \theta_{0,1} = .001$ with $[\theta_{0,i}]_{i=2}^{k_{\theta,n}} = \mathbf{0}_{k_{\theta,n}-1}$								
	$n = 100$									$n = 100$								
	$\check{k}_{\theta,n} = 10$			$\check{k}_{\theta,n} = 35$			$\check{k}_{\theta,n} = 50$			$\check{k}_{\theta,n} = 10$			$\check{k}_{\theta,n} = 35$			$\check{k}_{\theta,n} = 50$		
Test / Size	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%
Max-Test	.008	.051	.101	.006	.051	.107	.008	.041	.086	.101	.307	.466	.451	.762	.870	.605	.862	.938
Max-t-Test	.010	.048	.110	.010	.055	.114	.003	.042	.089	.142	.388	.536	.531	.795	.890	.673	.876	.937
Wald	.003	.035	.076	.000	.001	.016	.000	.000	.005	.020	.133	.251	.002	.033	.131	.000	.001	.038
	$n = 250$									$n = 250$								
	$\check{k}_{\theta,n} = 10$			$\check{k}_{\theta,n} = 35$			$\check{k}_{\theta,n} = 79$			$\check{k}_{\theta,n} = 10$			$\check{k}_{\theta,n} = 35$			$\check{k}_{\theta,n} = 79$		
Test / Size	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%
Max-Test	.003	.046	.094	.010	.046	.095	.007	.052	.111	.248	.460	.600	.924	.978	.992	1.00	1.00	1.00
Max-t-Test	.005	.043	.090	.008	.052	.094	.006	.040	.082	.219	.462	.584	.914	.975	.990	1.00	1.00	1.00
Wald	.005	.041	.091	.001	.019	.059	.001	.003	.022	.068	.224	.338	.212	.522	.685	.309	.850	.963
	$n = 500$									$n = 500$								
	$\check{k}_{\theta,n} = 10$			$\check{k}_{\theta,n} = 35$			$\check{k}_{\theta,n} = 112$			$\check{k}_{\theta,n} = 10$			$\check{k}_{\theta,n} = 35$			$\check{k}_{\theta,n} = 112$		
Test / Size	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%
Max-Test	.015	.049	.099	.006	.054	.107	.007	.045	.098	.833	.963	.987	1.00	1.00	1.00	1.00	1.00	1.00
Max-t-Test	.013	.041	.093	.008	.048	.104	.006	.042	.105	.931	.985	.994	1.00	1.00	1.00	1.00	1.00	1.00
Wald	.005	.034	.082	.003	.028	.070	.000	.004	.023	.693	.876	.939	.971	.994	.98	1.00	1.00	1.00
	$n = 1000$									$n = 1000$								
	$\check{k}_{\theta,n} = 10$			$\check{k}_{\theta,n} = 35$			$\check{k}_{\theta,n} = 158$			$\check{k}_{\theta,n} = 10$			$\check{k}_{\theta,n} = 35$			$\check{k}_{\theta,n} = 158$		
Test / Size	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%
Max-Test	.006	.064	.115	.005	.044	.094	.004	.005	.101	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Max-t-Test	.012	.056	.116	.010	.045	.089	.006	.050	.102	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Wald	.015	.045	.099	.007	.034	.080	.000	.017	.056	.987	.999	1.00	1.00	1.00	1.00	1.00	1.00	1.00

All p-values are bootstrapped, based on 1,000 independently drawn samples. $k_\delta = 0$.

Table 3: Rejection Frequencies under H_1 (ii) and (iii)

	$H_1(ii) : \theta_{0,i} = i/k_{\theta,n}$ for $i = 1, \dots, k_{\theta,n}$									$H_1(iii) : \theta_{0,i} = .001$ for $i = 1, \dots, k_{\theta,n}$								
	$n = 100$									$n = 100$								
	$k_{\theta,n} = 10$			$k_{\theta,n} = 35$			$k_{\theta,n} = 50$			$k_{\theta,n} = 10$			$k_{\theta,n} = 35$			$k_{\theta,n} = 50$		
Test / Size	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%
Max-Test	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Max-t-Test	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Wald	1.00	1.00	1.00	1.00	1.00	1.00	.869	1.00	1.00	1.00	1.00	1.00	.996	1.00	1.00	.885	1.00	1.00
	$n = 250$									$n = 250$								
	$k_{\theta,n} = 10$			$k_{\theta,n} = 35$			$k_{\theta,n} = 79$			$k_{\theta,n} = 10$			$k_{\theta,n} = 35$			$k_{\theta,n} = 79$		
Test / Size	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%
Max-Test	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Max-t-Test	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Wald	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
	$n = 500$									$n = 500$								
	$k_{\theta,n} = 10$			$k_{\theta,n} = 35$			$k_{\theta,n} = 112$			$k_{\theta,n} = 10$			$k_{\theta,n} = 35$			$k_{\theta,n} = 112$		
Test / Size	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%
Max-Test	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Max-t-Test	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Wald	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
	$n = 1000$									$n = 1000$								
	$k_{\theta,n} = 10$			$k_{\theta,n} = 35$			$k_{\theta,n} = 158$			$k_{\theta,n} = 10$			$k_{\theta,n} = 35$			$k_{\theta,n} = 158$		
Test / Size	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%	1%	5%	10%
Max-Test	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Max-t-Test	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Wald	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00

All p-values are bootstrapped, based on 1,000 independently drawn samples. $k_\delta = 0$.