

THE PERIOD-INDEX PROBLEM FOR COMPLEX TORI

JAMES HOTCHKISS

ABSTRACT. We solve the period-index problem for the Brauer group of a general complex torus of dimension at least three, giving an explicit formula for the index of each Brauer class. As a consequence, the complex-analytic version of the period-index conjecture is false for infinitely many Brauer classes on a general complex torus of dimension at least three.

1. INTRODUCTION

Let X be a connected complex manifold. An *Azumaya algebra* \mathcal{A} over X is a sheaf of \mathcal{O}_X -algebras which is locally isomorphic to a matrix \mathcal{O}_X -algebra $M_n(\mathcal{O}_X)$. Since the rank of an Azumaya algebra \mathcal{A} is a square, one defines the *degree* of \mathcal{A} to be $\sqrt{\mathrm{rk}\mathcal{A}}$. Two Azumaya algebras \mathcal{A} and \mathcal{B} are *Morita equivalent* if there exist vector bundles E and F , and an isomorphism of \mathcal{O}_X -algebras

$$\mathcal{A} \otimes \mathcal{E}nd(E) \simeq \mathcal{B} \otimes \mathcal{E}nd(F).$$

The Brauer group $\mathrm{Br}(X)$ is the group of Morita-equivalence classes of Azumaya algebras with tensor product.

Given a Brauer class α , the *period* $\mathrm{per}(\alpha)$ of α is its order in $\mathrm{Br}(X)$, which is a torsion group. The *index* $\mathrm{ind}(\alpha)$ of α is the greatest common divisor of the set of degrees of Azumaya algebras of class α . It follows from a general result of Antieau and Williams that $\mathrm{per}(\alpha)$ divides $\mathrm{ind}(\alpha)$, and that they share prime factors [AW15].

The *period-index problem* is the problem of determining an integer ϵ so that $\mathrm{ind}(\alpha)$ divides $\mathrm{per}(\alpha)^\epsilon$. If X is projective, then $\mathrm{Br}(X)$ is a subgroup of $\mathrm{Br}(\mathbf{C}(X))$, with $\mathbf{C}(X)$ the function field of X , and the longstanding *period-index conjecture* about Brauer groups of function fields (cf. [CT06], [Lie08]) specializes to the following statement:

Conjecture 1.1 (Global period-index conjecture). *Let X be a smooth, connected, projective variety over \mathbf{C} . For any $\alpha \in \mathrm{Br}(X)$,*

$$\mathrm{ind}(\alpha) \mid \mathrm{per}(\alpha)^{\dim X - 1}.$$

The conjecture is trivial for $\dim X = 0$, and holds for $\dim X = 1$ by Tsen's theorem. The case $\dim X = 2$ is due to de Jong [dJ04]. According to the discriminant-avoidance theorem of de Jong and Starr [SdJ10], Conjecture 1.1 for all complex projective varieties of dimension $\leq d$ is equivalent to the period-index conjecture for complex function fields of transcendence degree $\leq d$. Notably, the analogue of Conjecture 1.1 for topological Azumaya algebras on finite CW complexes has been introduced and studied by Antieau and Williams [AW14a], [AW14b], [AW21].

Little is known about the period-index problem for Brauer groups of non-algebraic complex manifolds beyond the case of surfaces. When X is an analytic K3 surface, an analysis of the argument given in [HS03] shows that period and index coincide, in accordance with de Jong's theorem. Moreover, the same argument applies to the case when X is a 2-dimensional complex torus. When X is a Stein manifold, the period-index problem is equivalent to the topological period-index problem by the Grauert–Oka principle, and one may obtain bounds from the work of Antieau and Williams [AW21] (Remark 2.2).

If X is a complex torus of dimension g with $\alpha \in \text{Br}(X)$, then there is a naïve upper bound on $\text{ind}(\alpha)$ given by the *annihilator* $\text{Ann}(\alpha)$, which is the least degree of a finite isogeny $f : X' \rightarrow X$ such that $f^*\alpha = 0$. When $\text{NS}(X) = 0$, one may compute $\text{Ann}(\alpha)$ explicitly (Lemma 3.3), and as α ranges over $\text{Br}(X)[n]$, $\text{Ann}(\alpha)$ attains any positive integer value which divides n^g . Our main result is that for X general, $\text{Ann}(\alpha)$ is also a lower bound for $\text{ind}(\alpha)$:

Theorem 1.2. *Let X be a general complex torus of dimension $g \geq 3$, and let $\alpha \in \text{Br}(X)$. Then*

$$\text{ind}(\alpha) = \text{Ann}(\alpha)$$

More precisely, Theorem 1.2 holds for a complex torus X with $\text{NS}(X) = \text{Hdg}^4(X) = 0$. The proof of Theorem 1.2 is based on an argument of Voisin [Voi02] showing that a general complex torus of dimension at least three does not satisfy the resolution property, along with an analysis of the Hodge-theoretic properties of α -twisted sheaves in the manner of [Hot22]. As an immediate consequence of Theorem 1.2, we obtain the following result:

Corollary 1.3. *Let X be a general complex torus of dimension $g \geq 3$. For each $n > 0$, there exists a Brauer class α with $\text{per}(\alpha) = n$, $\text{ind}(\alpha) = n^g$.*

In particular, the extension of Conjecture 1.1 to even the most familiar compact Kähler manifolds is false. By contrast, we prove the period-index conjecture for abelian threefolds in forthcoming work with Perry [HP23].

One may interpret Theorem 1.2 as the statement that the rank of any locally free α -twisted sheaf is divisible by $\text{Ann}(\alpha)$. In fact, we show that the rank of an arbitrary α -twisted coherent sheaf on X is divisible by $\text{Ann}(\alpha)$ (Theorem 4.1), so that using a *coherent index* (Definition 2.4) does not correct for the failure of the expected period-index bounds.

1.1. Organization of the paper. In §2, we review background on Brauer groups and Severi–Brauer varieties. In §3, we establish some linear algebraic properties of the annihilator of a Brauer class. In §4, we prove our main result, Theorem 4.1.

1.2. Acknowledgements. I am grateful to Stefan Schreieder, whose suggestion that the period-index conjecture might fail for non-projective compact Kähler manifolds led to the present paper. In addition, I thank Alex Perry for several useful discussions.

2. PRELIMINARIES

For the definition of the Brauer group of a complex manifold and its basic properties, we refer to [Sch05]. If X is a connected complex manifold and $\alpha \in \text{Br}(X)$, we recall from §1 that the *period* $\text{per}(\alpha)$ is the order of α in $\text{Br}(X)$, and the *index* of α is given by

$$\text{ind}(\alpha) = \gcd(\deg \mathcal{A} : [\mathcal{A}] = \alpha),$$

where \mathcal{A} runs over the Azumaya algebras of class α . (We recall that the *degree* of an Azumaya algebra is the square root of its rank.)

Lemma 2.1. *Let X be a connected complex manifold, and $\alpha \in \text{Br}(X)$. Then $\text{per}(\alpha)$ divides $\text{ind}(\alpha)$, and $\text{per}(\alpha)$ and $\text{ind}(\alpha)$ share prime factors.*

Proof. See [AW15]. □

Remark 2.2. Suppose that X is a connected Stein manifold, and consider the homomorphism

$$\text{Br}(X) \rightarrow \text{H}^3(X, \mathbf{Z})^{\text{tors}}.$$

By the Grauert–Oka principle [For17, §8.2], the set of isomorphism classes of holomorphic and continuous torsors under PGL_{n+1} coincide, for each n . It follows that the map above is an isomorphism,

and that the period-index problem for X is equivalent to the topological period-index problem for the CW complex underlying X . Hence, one may obtain bounds from [AW21]. For instance,

$$\text{ind}(\alpha) \mid \text{per}(\alpha)^{\dim X - 1}$$

if $\text{per}(\alpha)$ is prime to $(\dim X - 1)!$.

It is frequently useful to work with α -twisted sheaves in the sense of Căldăraru [Că100]. We write $\text{Coh}^\alpha(X)$ for the abelian category of α -twisted sheaves (where, as is customary, we often suppress the choice of a cocycle representative for α). One has the following well-known result:

Lemma 2.3. *Let X be a complex manifold, and let $\alpha \in \text{Br}(X)$. Then*

$$F \mapsto \text{End}(F)$$

gives a bijection between the set of isomorphism classes of locally free α -twisted sheaves, and the set of isomorphism classes of Azumaya algebras on X of class α .

Proof. We refer to [Că100, Theorem 1.3.5], which holds without change in the analytic case. \square

From Lemma 2.3, the index of α coincides with the greatest common divisor of the ranks of locally free α -twisted sheaves. It is also natural to consider the ranks of arbitrary α -twisted coherent sheaves, which leads to the following variant of the index:

Definition 2.4. Let X be a complex manifold, with $\alpha \in \text{Br}(X)$. The *coherent index* $\text{ind}_{\text{Coh}}(\alpha)$ is the greatest common divisor of the ranks of α -twisted coherent sheaves on X .

We note that $\text{ind}_{\text{Coh}}(\alpha)$ divides $\text{ind}(\alpha)$. It follows from Theorem 4.1 below that $\text{ind}_{\text{Coh}}(\alpha)$ and $\text{ind}(\alpha)$ coincide when X is a general complex torus, but we do not know if they coincide for Brauer classes on arbitrary complex manifolds.

Remark 2.5. A famous question of Grothendieck [Gro95] asks when the inclusion

$$\text{Br}(X) \subseteq \text{H}^2(X, \mathcal{O}_X^\times)^{\text{tors}}$$

is an equality. While the answer is positive in the projective case [dJ03], the question remains widely open in the compact Kähler case. It is known, however, for compact Kähler surfaces (cf. [HS03] and [Sch05]), as well as complex tori (cf. [EN83], or one may deduce it from the proof of Lemma 2.6 below).

Lemma 2.6. *Let $f : X' \rightarrow X$ be a finite cover of complex manifolds. Given $\alpha \in \text{Br}(X)$ such that $f^*\alpha = 0$,*

$$\text{ind}(\alpha) \mid \deg f.$$

Proof. Let $\tilde{\alpha}$ be a cocycle representative for α , and let L be a $f^*\tilde{\alpha}$ -twisted line bundle on X' . Then f_*L is a locally free $\tilde{\alpha}$ -twisted coherent sheaf of rank $\deg f$ on X , and we conclude by Lemma 2.3. \square

In order to apply an analytic result in the proof of Theorem 4.1 below, it will be convenient to treat α -twisted sheaves on X as genuine sheaves on a Severi–Brauer variety. Recall that a Severi–Brauer variety over a connected complex manifold X is a smooth, proper PGL_{n+1} -equivariant morphism $\mathbf{P} \rightarrow X$ (where the action on X is trivial), which is locally on X of the form $\mathbf{P}^n \times U \rightarrow U$, for $U \subseteq X$. A Severi–Brauer variety $\mathbf{P} \rightarrow X$ has a Brauer class $[\mathbf{P}] \in \text{Br}(X)$, which obstructs the existence of a holomorphic vector bundle E on X such that $\mathbf{P} \simeq \mathbf{P}(E)$.

We warn the reader that we follow Giraud’s convention [Gir71, Example V.4.8] for the Brauer class of a Severi–Brauer variety. If one adopts the opposite convention, then the category of weight- k sheaves defined below is equivalent to the category of α^{-k} -twisted sheaves.

Definition 2.7. Let X be a complex manifold, and let $\mathbf{P} \rightarrow X$ be a Severi–Brauer variety. A coherent sheaf E on \mathbf{P} has *weight* k if, for any $x \in X$, the restriction of E to the fiber \mathbf{P}_x is isomorphic to $V \otimes \mathcal{O}_{\mathbf{P}_x}(k)$ for a \mathbf{C} -vector space V .

Let $\text{Coh}^k(\mathbf{P}/X)$ be the abelian category of weight- k coherent sheaves on \mathbf{P} . For simplicity, we often write $\text{Coh}^k(\mathbf{P})$, with the morphism to X implicit.

Remark 2.8 (Descent). In the setting above, let E be a weight-0 coherent sheaf on \mathbf{P} . Then the natural morphism

$$\pi^* \pi_* E \rightarrow E$$

is an isomorphism.

Lemma 2.9. Let X be a complex manifold, and let $\alpha \in \text{Br}(X)$, and let $\pi : \mathbf{P} \rightarrow X$ be a Severi–Brauer variety of class α . There exists a rank-preserving equivalence

$$\text{Coh}^\alpha(X) \simeq \text{Coh}^1(\mathbf{P}).$$

Proof. Let $\tilde{\alpha}$ be a cocycle representative for α . Since $\pi^* \alpha$ is trivial, there exists an $\pi^* \tilde{\alpha}^{-1}$ -twisted line bundle $\mathcal{O}_{\mathbf{P}}^{\text{tw}}(1)$ on \mathbf{P} . The equivalence is given by $E \mapsto \pi^* E \otimes \mathcal{O}_{\mathbf{P}}^{\text{tw}}(1)$, with inverse equivalence $F \mapsto \pi_*(F \otimes \mathcal{O}_{\mathbf{P}}^{\text{tw}}(-1))$. \square

Remark 2.10 (Topologically trivial Brauer classes). Let $\mathbf{P} \rightarrow X$ be a Severi–Brauer variety, and suppose that the Brauer class α of \mathbf{P} is *topologically trivial*, i.e., lies in the kernel of the homomorphism

$$\text{Br}(X) \subseteq \text{H}^2(X, \mathcal{O}_X^\times)^{\text{tors}} \rightarrow \text{H}^3(X, \mathbf{Z})^{\text{tors}}$$

arising from the exponential sequence. Then $\mathbf{P} \rightarrow X$ is the projectivization of a topological complex vector bundle.

In particular, there exists a *weight-1 topological line bundle* $\mathcal{O}_{\mathbf{P}}^{\text{top}}(1)$ on \mathbf{P} , whose restriction to the fiber \mathbf{P}_x over $x \in X$ is isomorphic to $\mathcal{O}_{\mathbf{P}_x}(1)$. Moreover, by the Leray–Hirsch theorem, there is a decomposition

$$\text{K}_0^{\text{top}}(\mathbf{P}) = \text{K}_0^{\text{top}}(X) \oplus \text{K}_0^{\text{top}}(X) \cdot [\mathcal{O}_{\mathbf{P}}^{\text{top}}(1)] \oplus \cdots \oplus \text{K}_0^{\text{top}}(X) \cdot [\mathcal{O}_{\mathbf{P}}^{\text{top}}(1)^{\otimes d}],$$

where d is the relative dimension of $\mathbf{P} \rightarrow X$.

Definition 2.11. Let X be a complex manifold, and let $\alpha \in \text{Br}(X)$. A *rational B-field* for α is a class $B \in \text{H}^2(X, \mathbf{Q})$ whose image under the homomorphism

$$\exp(2\pi i \cdot -) : \text{H}^2(X, \mathbf{Q}) \rightarrow \text{H}^2(X, \mathcal{O}_X^\times)^{\text{tors}}$$

is α . We note that a rational B -field for α exists if and only if α is topologically trivial.

Remark 2.12. Let X be a compact Kähler manifold, and let $\pi : \mathbf{P} \rightarrow X$ be a Severi–Brauer variety of class α . Suppose that α is topologically trivial. For any weight-1 topological line bundle $\mathcal{O}_{\mathbf{P}}^{\text{top}}(1)$ on \mathbf{P} ,

$$c_1(\mathcal{O}_{\mathbf{P}}^{\text{top}}(1)) = H + \pi^* B,$$

where $H \in \text{NS}(\mathbf{P})_{\mathbf{Q}}$ is a class whose restriction to any fiber \mathbf{P}_x is $\mathcal{O}_{\mathbf{P}_x}(1)$, and $B \in \text{H}^2(X^{\text{an}}, \mathbf{Q})$ is a rational B -field for α . We refer to [dP22, Lemma 5.9], which however uses the opposite convention for the class of a Severi–Brauer variety (hence the difference in sign), and which is stated for a smooth, proper variety X but holds in the compact Kähler case by an identical argument.

3. THE ANNIHILATOR OF A BRAUER CLASS

Definition 3.1. Let X be a complex torus.

- (1) The *annihilator* of a class $\omega \in H^2(X, \mathbf{Z}/n)$ is the least degree of a finite isogeny $f : X' \rightarrow X$ such that $f^*\omega = 0$.
- (2) The *annihilator* of a Brauer class $\alpha \in \text{Br}(X)$ is the least degree of a finite isogeny $f : X' \rightarrow X$ such that $f^*\alpha = 0$.

By Lemma 2.6, $\text{ind}(\alpha)$ divides $\text{Ann}(\alpha)$.

Remark 3.2. Suppose that X is a complex torus with $\text{NS}(X) = 0$. From the Kummer sequence, the exponential map

$$H^2(X, \mathbf{Z}/n) \rightarrow \text{Br}(X)[n]$$

is an isomorphism. If $\alpha \in \text{Br}(X)[n]$, then α lies in the image of a unique class $\omega \in H^2(X, \mathbf{Z}/n)$, and $\text{Ann}(\alpha) = \text{Ann}(\omega)$.

The following lemma computes $\text{Ann}(\omega)$ in terms of linear algebra. For an element $a \in \mathbf{Z}/n$, we write $\text{Ann}_{\mathbf{Z}}(a) \in \mathbf{Z}$ for the positive generator of the annihilator ideal of a .

Lemma 3.3. *Let X be a complex torus of dimension g , with $\omega \in H^2(X, \mathbf{Z}/n)$. Suppose that e_1, \dots, e_{2g} is a basis for $H^1(X, \mathbf{Z})$ such that*

$$\omega = \sum_{i=1}^r a_i e_i \wedge e_{i+g},$$

for $0 \neq a_i \in \mathbf{Z}/n$ and $0 \leq r \leq g$. Then

$$\text{Ann}(\omega) = \prod_{i=1}^r \text{Ann}_{\mathbf{Z}}(a_i).$$

Proof. First, consider an isogeny $f : X' \rightarrow X$ of degree $\prod \text{Ann}_{\mathbf{Z}}(a_i)$ such that

$$f^*e_i = \text{Ann}(a_i) \cdot e'_i, \quad 1 \leq i \leq r$$

where e'_1, \dots, e'_{2g} is a basis for $H^1(X', \mathbf{Z})$. Then $f^*\omega = 0$, so $\text{Ann}(\omega) \leq \prod \text{Ann}_{\mathbf{Z}}(a_i)$.

In the other direction, we may suppose that $n = p^e$ is a prime power. Let

$$\eta = \sum_{i=1}^r a'_i e_i \wedge e_{i+g} \in H^2(X, \mathbf{Z})$$

be a lift of ω , and let $f : X' \rightarrow X$ be a finite isogeny such that $f^*\omega = 0$. Then $f^*\eta$ is divisible by n . In particular,

$$\begin{aligned} f_* f^* \eta^r / r! &= \deg f \cdot \prod_{i=1}^r a'_i \\ &\equiv 0 \pmod{n^r}. \end{aligned}$$

Since n is a prime power, it follows that $\prod \text{Ann}_{\mathbf{Z}}(a_i)$ divides $\deg f$. □

If X is a complex torus, then $H^3(X, \mathbf{Z})^{\text{tors}} = 0$, so any Brauer class is topologically trivial and admits a rational B -field.

Lemma 3.4. *Let X be a complex torus with $\text{NS}(X) = 0$, and let $\alpha \in \text{Br}(X)$. For any rational B -field B for α , let N be the least positive integer such that*

$$N \cdot \exp(B) \in H^{\text{ev}}(X, \mathbf{Q})$$

lies in $H^{\text{ev}}(X, \mathbf{Z})$. Then $N = \text{Ann}(\alpha)$.

Proof. Let $n = \text{per}(\alpha)$. We write

$$B = \sum_{i=1}^r \frac{m_i}{n} e_i \wedge e_{i+g}, \quad m_i \neq 0 \in \mathbf{Z}, \quad 0 \leq r \leq g.$$

Then the image ω of $n \cdot B$ in $H^2(X, \mathbf{Z}/n)$ maps to α under the exponential. By Remark 3.2, $\text{Ann}(\alpha)$ is equal to $\text{Ann}(\omega)$. By Lemma 3.3, $\text{Ann}(\omega) = \prod \text{Ann}_{\mathbf{Z}}(m_i \bmod n)$.

With N as in the statement of the lemma, we observe that $N \cdot B^r / r!$ is integral. For each prime factor p of n , with $v_p(m) = e$, it follows that

$$N \cdot \prod m_i \equiv 0 \pmod{p^{er}},$$

so $\prod \text{Ann}_{\mathbf{Z}}(m_i \bmod p^e)$ divides N for all such p , which implies that $\prod \text{Ann}_{\mathbf{Z}}(m_i \bmod n)$ divides N . In the other direction, it is straightforward to show that

$$\prod \text{Ann}_{\mathbf{Z}}(m_i \bmod n) \cdot \exp(B)$$

is integral, so that N divides $\prod \text{Ann}_{\mathbf{Z}}(m_i \bmod n)$. \square

4. THE MAIN RESULT

In this section, we prove our main result, phrased in terms of the coherent index of Definition 2.4.

Theorem 4.1. *Let X be a complex torus such that $\text{NS}(X) = \text{Hdg}^4(X) = 0$, with $\alpha \in \text{Br}(X)$. Then*

$$\text{ind}_{\text{Coh}}(\alpha) = \text{Ann}(\alpha)$$

Theorem 4.1 implies Theorem 1.2, and also implies Corollary 1.3 by the formula given for $\text{Ann}(\alpha)$ through Remark 3.2 and Lemma 3.3. We note that if a complex torus X satisfies the condition that $\text{NS}(X) = \text{Hdg}^4(X) = 0$, and E is a locally free coherent sheaf on X , then $c_i(E) = 0$ for each $i > 0$ [Voi02]. In particular, X does not have the resolution property.

Proof. First, from the definition of $\text{ind}_{\text{Coh}}(\alpha)$ and Lemma 2.6,

$$\text{ind}_{\text{Coh}}(\alpha) \mid \text{ind}(\alpha) \mid \text{Ann}(\alpha),$$

so it remains to show that $\text{Ann}(\alpha)$ divides $\text{ind}_{\text{Coh}}(\alpha)$, i.e., that the rank of any α -twisted coherent sheaf is divisible by $\text{Ann}(\alpha)$.

Let $\mathbf{P} \rightarrow X$ be a Severi–Brauer variety of class α , and let E be a coherent sheaf of weight 1 on \mathbf{P} . By Lemma 2.9, it is enough to show the rank of E is divisible by $\text{Ann}(\alpha)$. We may suppose that E is torsion-free, and by induction on the rank, we may suppose that E contains no nonzero subsheaves of smaller rank. Then $E^{\vee\vee}$ contains no nonzero subsheaves of smaller rank as well, and is reflexive. We replace E with $E^{\vee\vee}$.

Let $f : X' \rightarrow X$ be a finite isogeny such that $f^* \alpha = 0$, and consider a pullback diagram

$$\begin{array}{ccc} \mathbf{P}' & \xrightarrow{g} & \mathbf{P} \\ \downarrow & & \downarrow \\ X' & \xrightarrow{f} & X. \end{array}$$

We note that E is μ -stable with respect to a Kähler form ω on \mathbf{P} , so by [BS09, Prop. 2.3], $g^* E$ is μ -polystable with respect to $g^* \omega$. Since $\mathbf{P}' \rightarrow X'$ is a projective bundle, there is a holomorphic relative hyperplane bundle $\mathcal{O}_{\mathbf{P}'}(1)$ so that $E' = g^* E \otimes \mathcal{O}_{\mathbf{P}'}(-1)$ has weight 0, and in particular descends to a coherent sheaf E'_0 on X' (Remark 2.8).

Our assumption on X implies that $\text{NS}(X') = \text{Hdg}^4(X') = 0$, so that $c_1(E') = c_2(E') = 0$. Since E' is reflexive and μ -polystable, the theorem of Bando and Siu [BS94, Corollary 3] implies that E' is a flat holomorphic vector bundle, and in particular, $c_i(E') = 0$ for all $i > 0$.

Next, we compute Chern characters. Let $\mathcal{O}_{\mathbf{P}}^{\text{top}}(1)$ be a 1-twisted topological line bundle on \mathbf{P} . From the Leray–Hirsch theorem (Remark 2.10), we may write

$$\text{ch}(E) = \text{ch}(\mathcal{O}_{\mathbf{P}}^{\text{top}}(1)) \cdot \text{ch}(\xi),$$

where $\xi \in K_0^{\text{top}}(X)$. From Remark 2.12, we may write $c_1(\mathcal{O}_{\mathbf{P}}^{\text{top}}(1)) = H + B$, where H is a rational Hodge class and B is a rational B -field for α . Since H is Hodge,

$$\zeta = \exp(B) \cdot \text{ch}(\xi) \in H^{\text{ev}}(X, \mathbf{Q})$$

is Hodge in each degree.

We claim that $f^*\zeta = \text{ch}(E'_0)$. To prove the claim, first observe that f^*B is a rational B -field for $f^*\alpha = 0$, hence must be integral because $\text{NS}(X') = 0$. Therefore, g^*h is an integral Hodge class, and generates $\text{NS}(\mathbf{P}')$. It follows that $c_1(\mathcal{O}_{\mathbf{P}'}(1)) = g^*h$, which implies the claim.

From the first part of the proof,

$$\text{ch}(E'_0) = N \cdot \mathbf{1} \in H^{\text{ev}}(X', \mathbf{Z}),$$

for an integer $N \geq 0$, where $\mathbf{1}$ is the multiplicative unit. It follows that $\zeta = N \cdot \mathbf{1}$.

It remains to show that $\text{Ann}(\alpha)$ divides N . But

$$\text{ch}(\xi) = N \cdot \exp(-B)$$

lies in $H^{\text{ev}}(X, \mathbf{Z})$, so we conclude by Lemma 3.4. \square

REFERENCES

- [AW14a] Benjamin Antieau and Ben Williams, *The period-index problem for twisted topological K-theory*, *Geom. Topol.* **18** (2014), no. 2, 1115–1148. MR 3190609
- [AW14b] ———, *The topological period-index problem over 6-complexes*, *J. Topol.* **7** (2014), no. 3, 617–640. MR 3252958
- [AW15] ———, *The prime divisors of the period and index of a Brauer class*, *J. Pure Appl. Algebra* **219** (2015), no. 6, 2218–2224. MR 3299728
- [AW21] ———, *The topological period-index conjecture*, *Math. Res. Lett.* **28** (2021), no. 5, 1307–1317. MR 4471709
- [BS94] Shigetoshi Bando and Yum-Tong Siu, *Stable sheaves and Einstein-Hermitian metrics*, *Geometry and analysis on complex manifolds*, World Sci. Publ., River Edge, NJ, 1994, pp. 39–50. MR 1463962
- [BS09] Indranil Biswas and S. Subramanian, *Semistability and finite maps*, *Arch. Math. (Basel)* **93** (2009), no. 5, 437–443. MR 2563589
- [Că100] Andrei Căldăraru, *Derived categories of twisted sheaves on Calabi-Yau manifolds*, Ph.D. thesis, Cornell University, May 2000.
- [CT06] Jean-Louis Colliot-Thélène, *Algèbres simples centrales sur les corps de fonctions de deux variables (d’après A. J. de Jong)*, no. 307, 2006, Séminaire Bourbaki. Vol. 2004/2005, pp. Exp. No. 949, ix, 379–413. MR 2296425
- [dJ03] Aise Johan de Jong, *A result of Gabber*, available at <https://www.math.columbia.edu/~dejong/papers/2-gabber.pdf>, 2003.
- [dJ04] A. J. de Jong, *The period-index problem for the Brauer group of an algebraic surface*, *Duke Math. J.* **123** (2004), no. 1, 71–94. MR 2060023
- [dP22] Aise Johan de Jong and Alexander Perry, *The period-index problem and Hodge theory*, arXiv e-prints (2022), <https://arxiv.org/abs/2212.12971>.
- [EN83] G. Elencwajg and M. S. Narasimhan, *Projective bundles on a complex torus*, *J. Reine Angew. Math.* **340** (1983), 1–5. MR 691957
- [For17] Franc Forstnerič, *Stein manifolds and holomorphic mappings*, second ed., *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*, vol. 56, Springer, Cham, 2017, The homotopy principle in complex analysis. MR 3700709
- [Gir71] Jean Giraud, *Cohomologie non abélienne*, *Die Grundlehren der mathematischen Wissenschaften, Band 179*, Springer-Verlag, Berlin-New York, 1971. MR 0344253
- [Gro95] Alexander Grothendieck, *Le groupe de Brauer. II. Théorie cohomologique*, Séminaire Bourbaki, Vol. 9, Soc. Math. France, Paris, 1995, pp. Exp. No. 297, 287–307. MR 1608805

- [Hot22] James Hotchkiss, *Hodge theory of twisted derived categories and the period-index problem*, arXiv e-prints (2022), <https://arxiv.org/abs/2212.10638>.
- [HP23] James Hotchkiss and Alexander Perry, *The period-index conjecture for abelian threefolds via Donaldson–Thomas theory*, in preparation, 2023.
- [HS03] Daniel Huybrechts and Stefan Schröer, *The Brauer group of analytic K3 surfaces*, Int. Math. Res. Not. (2003), no. 50, 2687–2698. MR 2017247
- [Lie08] Max Lieblich, *Twisted sheaves and the period-index problem*, Compos. Math. **144** (2008), no. 1, 1–31. MR 2388554
- [Sch05] Stefan Schröer, *Topological methods for complex-analytic Brauer groups*, Topology **44** (2005), no. 5, 875–894. MR 2153976
- [SdJ10] Jason Starr and Johan de Jong, *Almost proper GIT-stacks and discriminant avoidance*, Doc. Math. **15** (2010), 957–972. MR 2745688
- [Voi02] Claire Voisin, *A counterexample to the Hodge conjecture extended to Kähler varieties*, Int. Math. Res. Not. (2002), no. 20, 1057–1075. MR 1902630

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109

Email address: `htchkss@umich.edu`