

# Goppa codes over Edwards curves

Giuseppe Filippone  \*

*Department of Mathematics and Computer Science,  
University of Palermo, Via Archirafi 34, 90123 Palermo, Italy*

## Abstract

Given an Edwards curve, we determine a basis for the Riemann-Roch space of any divisor whose support does not contain any of the two singular points. This basis allows us to compute a generating matrix for an algebraic-geometric Goppa code over the Edwards curve.

**Keywords:** Algebraic Geometric Goppa code; Edwards curve; Riemann-Roch space

**AMS MSC (2010) codes:** 94B27; 94B05; 11T71

## 1 Introduction

The literature on elliptic curves and their applications in [cryptography](#) is well consolidated. Besides the well-known ECC (Elliptic Curve Cryptography) in which the group law, defined on these curves, is exploited to encrypt messages, and the ECDSA (Elliptic Curve Digital Signature Algorithm), another example can be found in the Lenstra algorithm for the factorization of integers. Moreover, there are as well applications to [coding theory](#) based on the [Riemann-Roch space](#)  $\mathcal{L}(D)$  associated with a rational divisor  $D$  of these curves. In particular, this space is a fundamental ingredient to construct [Goppa codes](#), first introduced in 1983 [9]. Goppa codes over the Hermitian curve, as well as over maximal curves and hyperelliptic curves, have been extensively studied in [1, 6–8, 11, 12, 15], as they have become an important topic both in coding theory and in cryptography, where they play a central role in [McEliece public-key cryptographic systems](#) [14].

To the best of our knowledge, AG Goppa codes for Edwards curves have not been considered until now. In this paper, we compute the generating matrices for AG Goppa codes over Edwards curves. These curves are already the subject of many papers in cryptography [2–5, 10, 13], in particular in their [twisted](#) version. Compared to the classic elliptic curves in Weierstrass form, they can be more efficient for cryptographic use and for the (single or multiple) digital signature.

In [section 2](#) we describe Edwards curves and their relationship with elliptic curves in Weierstrass form. In [section 3](#) we compute a basis for  $\mathcal{L}(D)$  over Edwards curves, while in [section 4](#) we construct AG Goppa codes over Edwards curves and their generating matrices. In particular, in [subsec. 4.3](#) we give a small example of a [Goppa MDS code](#) where we use the AG Goppa code defined in [subsec. 4.1](#) over Edwards curves.

## 2 Edwards curves and elliptic curves in Weierstrass form

In this section we introduce Edwards curves  $\mathcal{E}$ , that is, algebraic curves, defined over a field  $\mathbb{K}$ , which can be represented in a suitable coordinate system by the equation  $\hat{x}^2 + \hat{y}^2 = 1 + d\hat{x}^2\hat{y}^2$ , with  $d(d-1) \neq 0$ . We present these curves as a birationally equivalent version of elliptic curves in Weierstrass form.

Recall that, over a field  $\mathbb{K}$  of characteristic different from 2, a (smooth) elliptic curve (possessing at least a  $\mathbb{K}$ -rational point) can be represented in a suitable coordinate system by the Weierstrass equation  $y^2 = x^3 + a'x^2 + b'x$ , having one point at infinity  $\Omega = [Z : X : Y] = [0 : 0 : 1]$  on the  $y$  axis.

---

\*Electronic address: [giuseppe.filippone01@unipa.it](mailto:giuseppe.filippone01@unipa.it)

**Remark 2.1.** Note that, unlike those in Weierstrass form, curves in Edwards form  $\mathcal{E}$  have two points at infinity, that is,  $\Omega_1 = [\hat{Z} : \hat{X} : \hat{Y}] = [0 : 1 : 0]$  on the  $x$  axis and  $\Omega_2 = [\hat{Z} : \hat{X} : \hat{Y}] = [0 : 0 : 1]$  on the  $y$  axis, which are *ordinary singular points* for  $\mathcal{E}$  as this curve is *non-smooth*.

**Remark 2.2.** Edwards curves have four remarkable points:  $O = (0, 1)$ ,  $O' = (0, -1)$ ,  $H = (1, 0)$ ,  $H' = (-1, 0)$ . In particular,  $O - O$  is the identity element of the group law defined on them, that is,  $(P - O) + (O - O) = P - O$ , for any point  $P = (a, b) \in \mathcal{E}(\mathbb{K})$ , and these four points form the cyclic group  $C_4$ , where  $O' - O$  have order 2, while  $2(H - O) = 2(H' - O) = O' - O$ .

Edwards curves and elliptic curves in Weierstrass form are closely related. In particular, over a field  $\mathbb{K}$  of characteristic different from 2, one has that an elliptic curve  $\mathcal{W}$  defined by the equation  $y^2 = x^3 + a'x^2 + b'x$ , and an Edwards curve  $\mathcal{E}$  defined by the equation  $\hat{x}^2 + \hat{y}^2 = 1 + d\hat{x}^2\hat{y}^2$ , where  $d$  is not a square, are *birationally equivalent* (cf.[2]). Furthermore, this equivalence is given by the following two rational maps:

$$\alpha: \mathcal{E}(\mathbb{K}) \longrightarrow \mathcal{W}(\mathbb{K}) \quad (2.1a)$$

$$(\hat{x}, \hat{y}) \longmapsto (x, y) = \left( x_1 \frac{1 + \hat{y}}{1 - \hat{y}}, y_1 \frac{(1 + \hat{y})}{\hat{x}(1 - \hat{y})} \right)$$

$$\beta: \mathcal{W}(\mathbb{K}) \longrightarrow \mathcal{E}(\mathbb{K}) \quad (2.1b)$$

$$(x, y) \longmapsto (\hat{x}, \hat{y}) = \left( \frac{y_1 x}{x_1 y}, \frac{x - x_1}{x + x_1} \right),$$

where  $P = (x_1, y_1) \in \mathcal{W}(\mathbb{K})$  is such that the divisor  $2(P - O) = (0, 0) - \Omega$ .

**Remark 2.3.** The two rational maps  $\alpha$  and  $\beta$  defines a *birational equivalence* between  $\mathcal{W}$  and  $\mathcal{E}$ . Moreover, one extends the definition of  $\alpha$  and  $\beta$  by putting  $\alpha((0, 1)) = \Omega$ ,  $\beta(\Omega) = (0, 1)$ ,  $\alpha((0, -1)) = (0, 0)$  and  $\beta((0, 0)) = (0, -1)$ ; and  $\beta((t_1, 0)) = \beta((t_2, 0)) = \Omega_1$ ,  $\beta((-x_1, \pm s_1)) = \Omega_2$ , where  $(t_1, 0), (t_2, 0), (-x_1, \pm s_1) \in \mathcal{W}(\mathbb{K})$ , with  $t_1, t_2 \neq 0$ .

The value  $\Omega_1$  of the two images  $\beta((t_1, 0)), \beta((t_2, 0))$  and the value  $\Omega_2$  of the two images  $\beta((-x_1, \pm s_1))$  can be directly found by passing to homogeneous coordinates. As for the value of  $\beta(\Omega)$ , we have that, for  $\mathbb{K} = \mathbb{C}$ , if  $\wp(z)$  and  $\wp'(z)$  are the *Weierstrass elliptic functions* and  $P = \left( \wp(z) + \frac{a'}{3}, \frac{1}{2}\wp'(z) \right) \in \mathcal{W}(\mathbb{K})$ , then,  $\lim_{z \rightarrow 0} \beta(P) = \lim_{z \rightarrow 0} \left( \frac{2y_1 (3\wp(z) + a')}{3x_1 \wp'(z)}, \frac{3\wp(z) + a' - 3x_1}{3\wp(z) + a' + 3x_1} \right) = (0, 1) = O$ , as  $\frac{3\wp(z) + a'}{3\wp'(z)} = o(z)$ , that is,  $\beta$  is continuous in  $P = \Omega$ .

**Remark 2.4.** Since there are two points mapped by  $\beta$  onto  $\Omega_1$  and two points onto  $\Omega_2$ , one sees that it is not possible to coherently define  $\alpha(\Omega_1)$  and  $\alpha(\Omega_2)$ . For this reason the maps  $\alpha$  and  $\beta$  define a *birational equivalence* between the two forms. Note that  $\mathcal{W}$  is, indeed, a *smooth projective resolution* of the non-smooth curve  $\mathcal{E}$ .

**Remark 2.5.** Since the map  $\beta$  in (2.1b) transforms a line through  $P \in \mathcal{W}(\mathbb{K})$  and  $Q \in \mathcal{W}(\mathbb{K})$  onto the hyperbola through  $\beta(P), \beta(Q), O', 2\Omega_1$  and  $2\Omega_2$ , and maps vertical lines onto horizontal lines, then  $\beta$  induces a group homomorphism of the corresponding divisor classes group (cf. [2] for further details about the group law of Edwards curves).

### 3 The Riemann-Roch space $\mathcal{L}(D)$ over Edwards curves

In this section, given a divisor  $D \in \text{Div}(\mathcal{E})$ , we provide a basis of the Riemann-Roch vector space

$$\mathcal{L}(D) = \{f \in \overline{\mathbb{K}}(\mathcal{E})^* : \text{div}(f) + D \text{ is effective}\} \cup \{0\}$$

for an Edwards curve  $\mathcal{E}$ , under the assumption that the support of  $D$  does not contain the two singular points  $\Omega_1$  and  $\Omega_2$ .

We recall that a *divisor* is, in this context, an element of the free abelian group  $\text{Div}(\mathcal{E})$  on the points of  $\mathcal{E}$ , that is, a formal sum  $D = \sum_{n_P \in \mathbb{Z}} n_P P$ , with  $P \in \mathcal{E}(\mathbb{K})$ , where only finitely many integers  $n_P$  are not zero, and that a *principal divisor*  $D = \text{div}(g)$  of a function  $g$  is the sum of the zeros of  $g$  on  $\mathcal{E}$  minus the poles of  $g$  on  $\mathcal{E}$ . The integer  $\delta = \sum n_P$  is the *degree* of the

divisor  $D$  and principal divisors give a subgroup of the subgroup  $\text{Div}^0(\mathcal{E})$  of divisors having degree equal to zero, because any function  $g$  on  $\mathcal{E}$  has by Bezout theorem the same number of zeros and poles on  $\mathcal{E}$ . The group taken into account is formally the quotient group  $\frac{\text{Div}^0(\mathcal{E})}{\text{Princ}(\mathcal{E})}$ .

Also, we recall that any divisor  $D'$  on  $\mathcal{E}$  of degree  $k+1$ , such that  $\Omega_1$  and  $\Omega_2$  do not belong to the support of  $D'$ , is linearly equivalent to  $P+kO$ , for a suitable point  $P \in \mathcal{E}(\mathbb{K})$  (or  $(k+1)O$ , in the case where  $P=O$ ), that is,  $D' = P+kO + \text{div}(g)$ , for a suitable function  $g$ . Since the map

$$\begin{aligned} \chi : \mathcal{L}(D') &\longrightarrow \mathcal{L}(P+kO) \\ F &\longmapsto gF \end{aligned}$$

is an isomorphism between  $\mathcal{L}(D')$  and  $\mathcal{L}(P+kO)$ , we confine ourselves to the latter space.

**Theorem 3.1.** *Let  $\mathcal{E}$  be an Edwards curve defined, over a field  $\mathbb{K}$  of characteristic different from 2, by the equation  $x^2 + y^2 = 1 + dx^2y^2$ , where  $d$  is not a square. If  $P+kO = D \in \text{Div}(\mathcal{E})$  is a divisor of positive degree  $k+1$ , where  $P = (a, b)$ , then  $\dim(\mathcal{L}(D)) = k+1$  and*

$$\mathcal{L}(D) = \begin{cases} \langle F_0, F_1, \dots, F_k \rangle & \text{if } P \neq O \\ \langle F_0, F_2, \dots, F_{k+1} \rangle & \text{if } P = O \end{cases}$$

where  $F_0, F_1, \dots, F_{k+1}$  are rational homogeneous functions defined as follows:

$$\begin{aligned} F_0 &= \frac{Z}{Z} \\ F_1 &= \begin{cases} \frac{Z}{X} & \text{if } P = O' = (0, -1) \\ \frac{(X+Z)(Y+Z)}{XY} & \text{if } P = H = (1, 0) \\ \frac{(X-Z)(Y+Z)}{XY} & \text{if } P = H' = (-1, 0) \\ \frac{(Y+bZ) \cdot X}{(X-aZ) \cdot (Y-Z)} & \text{if } P \notin \{O', H, H'\} \end{cases} \\ F_i &= \begin{cases} \frac{Z^h}{(Y-Z)^h} & \text{if } i = 2h \\ \frac{(Y+Z)Z^h}{X(Y-Z)^h} & \text{if } i = 2h+1 \end{cases} \end{aligned}$$

for  $2 \leq i \leq k+1$ .

*Proof.* Since  $P$  is different from  $\Omega_1$  and  $\Omega_2$ , we can take the point  $\alpha(P) \in \mathcal{W}(\mathbb{K})$ , where  $\alpha : \mathcal{E}(\mathbb{K}) \longrightarrow \mathcal{W}(\mathbb{K})$  is the map defined in (2.1a). Hence, the (surjective) map

$$\alpha^{-1} = \beta : \mathcal{W}(\mathbb{K}) \longrightarrow \mathcal{E}(\mathbb{K})$$

in (2.1b), induces an (injective) homomorphism  $g \mapsto g \circ \beta$  from  $\mathcal{L}(P+kO)$  to  $\mathcal{L}(\alpha(P)+k\Omega)$ , because  $\beta(\text{div}(g \circ \beta)) = \text{div}(g)$  for any function  $g \in \mathcal{L}(P+kO)$ .

Since  $\mathcal{W}$  is smooth, by the formula of Riemann-Roch, the dimension of  $\mathcal{L}(\alpha(P)+k\Omega)$  is  $k+1$ , and we are left with exhibiting  $k+1$  linearly independent functions in  $\mathcal{L}(P+kO)$ , as manifestly  $\mathcal{L}(P+(i-1)O)$  is contained in  $\mathcal{L}(P+iO)$ , for  $i = 1, \dots, k+1$ .

For  $i=0$  the assertion follows, because  $\text{div}\left(\frac{Z}{Z}\right) = 0$  and for every  $P \in \mathcal{E}(\mathbb{K})$  we have that  $\text{div}(F_0) + P$  is effective.




Figure 3.1: Edwards real curve with  $d = -8$ . The points  $P, R, R'$  have the following coordinates  $(a, b), (a, -b), (-a, -b)$ , respectively

Recalling that  $O' = (0, -1), H = (1, 0), H' = (-1, 0)$ , and putting  $R = (a, -b)$  and  $R' = (-a, -b)$  for  $P = (a, b)$  as in [figure 3.1](#), for  $i = 1$  we have that:

$$\begin{aligned} \operatorname{div} \left( \frac{Z}{X} \right) &= 2H' + 2\Omega_2 + 2O' + 2\Omega_1 - (H + H' + 2\Omega_1 + O + O' + 2\Omega_2) \\ &= H' + O' - H - O, \\ \operatorname{div} \left( \frac{(X+Z)(Y+Z)}{XY} \right) &= (2H' + 2\Omega_2 + 2O' + 2\Omega_1) - (H + H' + 2\Omega_1 + O + O' + 2\Omega_2) = \\ &= H' + O' - H - O, \\ \operatorname{div} \left( \frac{(X-Z)(Y+Z)}{XY} \right) &= (2H + 2\Omega_2 + 2O' + 2\Omega_1) - (H + H' + 2\Omega_1 + O + O' + 2\Omega_2) = \\ &= H + O' - H' - O, \\ \operatorname{div} \left( \frac{(Y+bZ) \cdot X}{(X-aZ) \cdot (Y-Z)} \right) &= (R + R' + 2\Omega_1 + O + O' + 2\Omega_2) - (P + R + 2\Omega_2 + 2O + 2\Omega_1) = \\ &= R' + O' - P - O. \end{aligned}$$

Hence, we have that  $\operatorname{div}(F_1) + P + O$  is effective for any suitable  $P$ . Additionally, for  $i \geq 2$ , we have that:

$$\begin{aligned} \operatorname{div} \left( \frac{Z^h}{(Y-Z)^h} \right) &= (2h\Omega_1 + 2h\Omega_2) - (2hO + 2h\Omega_1) = 2h\Omega_2 - 2hO, \\ \operatorname{div} \left( \frac{(Y+Z)Z^h}{X(Y-Z)^h} \right) &= (2O' + 2\Omega_1 + 2h\Omega_1 + 2h\Omega_2) - (O + O' + 2\Omega_2 + 2hO + 2h\Omega_1) = \\ &= O' + 2\Omega_1 + (2h-2)\Omega_2 - (2h+1)O, \end{aligned}$$

hence,  $\operatorname{div}(F_i) + P + iO$  is effective in both the cases  $i = 2h$  and  $i = 2h + 1$ .

So, every function  $F_i$  is such that  $\operatorname{div}(F_i) + D$  is effective if  $D = P + kO$  or  $D = (k+1)O$ . In order to complete the proof, it is necessary to show that all these functions are linearly independent, but this follows from standard, elementary, arguments of linear algebra.

We note that in the case  $D = (k+1)O$  we simply remove  $F_1$  and we add  $F_{k+1}$ , thus, also in this case, we have  $k+1$  linearly independent functions.  $\blacksquare$

**Remark 3.1.** We note that it is not possible to extend the proof about  $\mathcal{L}(D)$  in [theorem 3.1](#) when  $P$  is equal to  $\Omega_1$  or  $\Omega_2$ , because the map  $\beta$  is not invertible on these points (see [remark 2.4](#)). Riemann-Roch spaces on curves having singular points are the subject of § IV.2 in [\[16\]](#).

### 3.1 Computational cost

Recalling that the costs of modular addition, multiplication, and inversion over  $\operatorname{GF}(q)$  are  $\mathcal{O}(\ln(q)), \mathcal{O}(\ln^2(q)), \mathcal{O}(\ln^3(q))$ , respectively, we now compute the cost of evaluating at a point  $P \in \mathcal{E}(\operatorname{GF}(q))$  each element of the basis of  $\mathcal{L}(D)$ .

We firstly note that we can compute  $F_{2h}$  from  $F_{2h-2}$ , and  $F_{2h+1}$  from  $F_{2h}$ , as

$$\begin{cases} F_{2h} = F_2 F_{2h-2} & \text{if } h \geq 2, \\ F_{2h+1} = \frac{Y+Z}{X} F_{2h} & \text{if } h \geq 1, \end{cases}$$

that is, at each step we have to perform a single multiplication times the last (or the second-last) value. Moreover, we can pre-calculate the value of the function  $\frac{Y+Z}{X}$  at  $P$  with a cost  $C\left(\frac{Y+Z}{X}\right) = \mathcal{O}(\ln(q)) + \mathcal{O}(\ln^2(q)) + \mathcal{O}(\ln^3(q)) \approx \mathcal{O}(\ln^3(q))$  to further speed up the computation.

Therefore, the maximal global cost  $C(\mathcal{L}(D))$  of evaluating the first  $k$  functions  $F_i$  is  $C(\mathcal{L}(D)) = \mathcal{O}(k \cdot \ln^2(q) + \ln^3(q))$ .

## 4 AG Goppa codes on Edwards curves

In this section, we construct the generating matrix and the parity-check matrix for a  $[n, k, d]_q$  AG Goppa code for an Edwards curve  $\mathcal{E}$  over  $\text{GF}(q)$ , compute the computational cost, and give a small example.

### 4.1 Goppa code for an Edwards curve

In the following, we adapt the definition of a Goppa code to our case.

**Definition 4.1.** Let  $D = P + (k-1)O$  be a divisor of positive degree  $\delta D = k$  of the Edwards curve  $\mathcal{E}$  over  $\text{GF}(q)$ , where  $q = p^t$  and  $p$  is an odd prime number. Let  $\mathcal{L}(D)$  be the Riemann-Roch space, let  $T = \{P_1, \dots, P_n\}$  be a set of  $n > k-1$  points such that, for  $i = 1, \dots, k$  and  $j = 1, \dots, n$ ,  $G_{ij} = F_{i-1}(P_j)$ , where  $\{F_i\}$  is a basis of  $\mathcal{L}(D)$ ,  $P_j \in \mathcal{E}$ , and  $P_j \notin \text{supp}(D)$ . Let  $(G_{ij}) = G \in \text{GF}(q)^{k \times n}$  be the  $k \times n$  matrix, we define the  $[n, k, d]_q$  AG Goppa code  $\mathcal{C}_G = \{\mathbf{c} \in \text{GF}(q)^n : \mathbf{c} = \mathbf{a} \cdot G, \mathbf{a} \in \text{GF}(q)^k\}$ .

**Remark 4.1.** We note that  $G$  is well defined because all points  $P_j \in T$  do not belong to the support of  $D$  which contains the poles of each  $F_i$ .

**Theorem 4.1.** If  $\mathcal{C}_G$  is the AG Goppa code of [definition 4.1](#), then the minimum distance  $d$  of this code is such that  $d \geq n - \delta D = n - k$ .

*Proof.* It follows from the same, classic, proof of AG Goppa codes over curves. ■

If we order the points in  $T$  so that the first  $k$  columns of the generating matrix

$$G = \begin{pmatrix} F_0(P_1) & F_0(P_2) & \cdots & F_0(P_n) \\ F_1(P_1) & F_1(P_2) & \cdots & F_1(P_n) \\ \vdots & \vdots & \ddots & \vdots \\ F_{k-1}(P_1) & F_{k-1}(P_2) & \cdots & F_{k-1}(P_n) \end{pmatrix} \in \text{GF}(q)^{k \times n}$$

of the Goppa code  $\mathcal{C}_G$  are linearly independent, e.g. by applying the Gauss-Jordan method, then  $G$  can be reduced in its [standard form](#)  $[I_k | M]$ , where  $I_k$  is the identity matrix of order  $k$  and  $M \in \text{GF}(q)^{k \times (n-k)}$ . Once  $G$  is in standard form, the parity-check matrix  $H \in \text{GF}(q)^{(n-k) \times n}$  of this Goppa code, that is, the matrix such that  $G \cdot H^T = \mathbf{0}$  and  $H \cdot \mathbf{y}^T = \mathbf{0}$  for every code word  $\mathbf{y} \in \mathcal{C}_G$ , is simply  $H = [-M^T | I_{n-k}]$ . Thus, the code  $\mathcal{C}_G$  is also defined as  $\{\mathbf{y} \in \text{GF}(q)^n : H \cdot \mathbf{y}^T = \mathbf{0}\}$ .

### 4.2 Computational cost of constructing a Goppa code

In order to compute the generating matrix  $G$  we need to evaluate each of the  $n$  points in the set  $T$  for each element of the basis of  $\mathcal{L}(D)$ , that is, we have a computational cost of  $\mathcal{O}(n \cdot C(\mathcal{L}(D)))$  because  $G$  is a matrix of size  $k \times n$ . Moreover, the cost of computing the parity-check matrix depends on the method used to solve the linear system  $G \cdot \mathbf{x} = \mathbf{0}$ . For instance, if we used the

**Gauss-Jordan method** to reduce the matrix  $G$  to its standard form, then the cost would be  $\mathcal{O}\left(\max(n, k)^3\right) = \mathcal{O}(n^3)$ .

Hence, the global computational cost of constructing a Goppa code over  $\mathcal{E}(\text{GF}(q))$  is:

$$\mathcal{O}(n \cdot C(\mathcal{L}(D))) + \mathcal{O}\left(\max(n, k)^3\right) = \mathcal{O}\left(n \cdot (k \cdot \ln^2(q) + \ln^3(q))\right) + \mathcal{O}(n^3).$$

In particular, for  $k \cdot \ln^2(q) + \ln^3(q) < n^2$ , the computational cost is  $\mathcal{O}(n^3)$ . However, if  $k \cdot \ln^2(q) + \ln^3(q) > n^2$ , for instance if we were working with very large finite fields ( $q \gg 2$ ), the overall computational cost would be  $\mathcal{O}\left(n \cdot (k \cdot \ln^2(q) + \ln^3(q))\right)$ .

### 4.3 A small example

Let  $\mathbb{K} = \text{GF}(17)$  and let  $\mathcal{E}$  be the Edwards curve defined by the equation  $x^2 + y^2 = 1 + 10x^2y^2$ . There are 24 affine points on this curve:

$$\{(0, 1), (0, 16), (1, 0), (2, 2), (2, 15), (3, 6), (3, 11), (5, 8), (5, 9), (6, 3), (6, 14), (8, 5), (8, 12), (9, 5), (9, 12), (11, 3), (11, 14), (12, 8), (12, 9), (14, 6), (14, 11), (15, 2), (15, 15), (16, 0)\},$$

and the two points at infinity.

Let  $D = (2, 15) + 4O$  be the divisor defining  $\mathcal{L}(D)$ , so the degree  $k$  is 5. Let  $T = \{P_1 = (5, 8), P_2 = (5, 9), P_3 = (6, 3), P_4 = (6, 14), P_5 = (8, 5), P_6 = (8, 12), P_7 = (9, 5)\}$  be the set of points such that  $P_j \notin \text{supp}(D)$  defining the generating matrix  $G$  of  $\mathcal{C}_G$ , thus  $n = 7$ . Applying [theorem 3.1](#), the vector space  $\mathcal{L}(D)$  has the following basis:

$$\begin{aligned} \mathcal{L}(D) &= \langle F_0, F_1, F_2, F_3, F_4 \rangle = \\ &= \left\langle 1, \frac{x(y+15)}{(x-2)(y-1)}, \frac{1}{y-1}, \frac{y+1}{x(y-1)}, \frac{1}{(y-1)^2} \right\rangle, \end{aligned}$$

whereas the generating matrix  $G = (G_{ij})$  of  $\mathcal{C}_G$  is defined by putting  $G_{ij} = F_{i-1}(P_j)$ , that is,

$$G = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 16 & 5 & 5 & 4 & 1 & 11 & 4 \\ 5 & 15 & 9 & 4 & 13 & 14 & 13 \\ 9 & 13 & 6 & 10 & 14 & 10 & 3 \\ 8 & 4 & 13 & 16 & 16 & 9 & 16 \end{pmatrix}.$$

We now compute the parity-check matrix  $H$  by solving the linear system  $G \cdot \mathbf{x} = \mathbf{0}$ , which reduces to:

$$\begin{cases} x_1 = 7x_6 + 2x_7 \\ x_2 = 3x_6 + 12x_7 \\ x_3 = x_6 + 9x_7 \\ x_4 = 13x_6 + 12x_7 \\ x_5 = 9x_6 + 15x_7 \end{cases},$$

that is,

$$H = \left( \begin{array}{cccccc|cc} 7 & 3 & 1 & 13 & 9 & 1 & 0 \\ 2 & 12 & 9 & 12 & 15 & 0 & 1 \end{array} \right). \quad (4.1)$$

Finally, the minimum distance for this code is  $d \geq n - \delta D = 7 - 5 = 2$ . Moreover, we know from the Singleton theorem that  $d \leq n - k + 1$  for a  $[n, k, d]_q$  code, that is,  $d \leq 7 - 5 + 1 = 3$  as  $k = \delta D = 5$ . Hence,  $d = 2$  or  $d = 3$ , but one can easily check that 2 columns of  $H$  are always linearly independent, so  $d = 3$ . Therefore, we have a  $[7, 5, 3]_{17}$ -Goppa [MDS](#) code.

## 5 Conclusions

Edwards curves have been recently introduced for their applications in cryptography. In this paper, we provided a basis for the Riemann-Roch space of a divisor on these curves, and we used this basis for the construction of the generating matrices of the AG Goppa codes, thus providing a possible application of Edwards curves to Coding theory, as well.

## References

- [1] Daniele Bartoli, Matteo Bonini, and Massimo Giulietti. “Constant dimension codes from Riemann-Roch spaces”. In: *Advances in Mathematics of Communications* 11.4 (2017), pp. 705–713. doi: [10.3934/amc.2017051](https://doi.org/10.3934/amc.2017051) (cit. on p. 1).
- [2] Daniel J. Bernstein and Tanja Lange. “Faster Addition and Doubling on Elliptic Curves”. In: *Advances in Cryptology – ASIACRYPT 2007*. Ed. by Kaoru Kurosawa. Berlin, Heidelberg: Springer Berlin Heidelberg, 2007, pp. 29–50. ISBN: 978-3-540-76900-2. doi: [10.1007/978-3-540-76900-2\\_3](https://doi.org/10.1007/978-3-540-76900-2_3) (cit. on pp. 1, 2).
- [3] Daniel J. Bernstein and Tanja Lange. “Inverted Edwards Coordinates”. In: *Applied Algebra, Algebraic Algorithms and Error-Correcting Codes*. Ed. by Serdar Boztaş and Hsiao-Feng (Francis) Lu. Berlin, Heidelberg: Springer Berlin Heidelberg, 2007, pp. 20–27. ISBN: 978-3-540-77224-8. doi: [10.1007/978-3-540-77224-8\\_4](https://doi.org/10.1007/978-3-540-77224-8_4) (cit. on p. 1).
- [4] Daniel J. Bernstein et al. “Optimizing Double-Base Elliptic-Curve Single-Scalar Multiplication”. In: *Progress in Cryptology – INDOCRYPT 2007*. Ed. by K. Srinathan, C. Pandu Rangan, and Moti Yung. Berlin, Heidelberg: Springer Berlin Heidelberg, 2007, pp. 167–182. ISBN: 978-3-540-77026-8. doi: [10.1007/978-3-540-77026-8\\_13](https://doi.org/10.1007/978-3-540-77026-8_13) (cit. on p. 1).
- [5] Harold Edwards. “A normal form for elliptic curves”. In: *Bulletin of The American Mathematical Society - BULL AMER MATH SOC* 44 (July 2007), pp. 393–423. doi: [10.1090/S0273-0979-07-01153-6](https://doi.org/10.1090/S0273-0979-07-01153-6) (cit. on p. 1).
- [6] Giovanni Falcone, Ágota Figula, and Carolin Hannusch. *On the generating matrix of Goppa codes over hyperelliptic curves*. (to appear on J. Ramanujan Mathe. Soc.) 2020. arXiv: [2012.08870](https://arxiv.org/abs/2012.08870) [[math.AG](https://arxiv.org/abs/2012.08870)] (cit. on p. 1).
- [7] S. Fanali and M. Giulietti. “One-Point AG Codes on the GK Maximal Curves”. In: *IEEE Transactions on Information Theory* 56.1 (2010), pp. 202–210. doi: [10.1109/TIT.2009.2034826](https://doi.org/10.1109/TIT.2009.2034826) (cit. on p. 1).
- [8] Massimo Giulietti and Gábor Korchmáros. “On automorphism groups of certain Goppa codes”. In: *Designs, Codes and Cryptography* 47.1 (June 2008), pp. 177–190. ISSN: 1573-7586. doi: [10.1007/s10623-007-9110-5](https://doi.org/10.1007/s10623-007-9110-5). URL: <https://doi.org/10.1007/s10623-007-9110-5> (cit. on p. 1).
- [9] V D Goppa. “ALGEBRAICO-GEOMETRIC CODES”. In: *Mathematics of the USSR-Izvestiya* 21.1 (Feb. 1983), pp. 75–91. doi: [10.1070/im1983v021n01abeh001641](https://doi.org/10.1070/im1983v021n01abeh001641) (cit. on p. 1).
- [10] Huseyin Hisil et al. “Faster Group Operations on Elliptic Curves”. In: *Proceedings of the Seventh Australasian Conference on Information Security - Volume 98*. AISC '09. Wellington, New Zealand: Australian Computer Society, Inc., 2009, pp. 7–20. ISBN: 9781920682798. doi: [10.5555/1862758.1862762](https://doi.org/10.5555/1862758.1862762) (cit. on p. 1).
- [11] G. Korchmáros, G. P. Nagy, and M. Timpanella. “Codes and Gap Sequences of Hermitian Curves”. In: *IEEE Transactions on Information Theory* 66.6 (2020), pp. 3547–3554. doi: [10.1109/TIT.2019.2950207](https://doi.org/10.1109/TIT.2019.2950207) (cit. on p. 1).
- [12] Gábor Korchmáros and Pietro Speziali. “Hermitian codes with automorphism group isomorphic to  $\text{PGL}(2, q)$  with  $q$  odd”. In: *Finite Fields and Their Applications* 44 (2017), pp. 1–17. ISSN: 1071-5797. doi: [10.1016/j.ffa.2016.11.003](https://doi.org/10.1016/j.ffa.2016.11.003) (cit. on p. 1).
- [13] Tanja Lange. “Edwards Curves”. In: *Encyclopedia of Cryptography and Security*. Ed. by Sushil van Tilborg Henk C. A. and Jajodia. Boston, MA: Springer US, 2011, pp. 380–382. ISBN: 978-1-4419-5906-5. doi: [10.1007/978-1-4419-5906-5\\_243](https://doi.org/10.1007/978-1-4419-5906-5_243) (cit. on p. 1).
- [14] R. J. McEliece. “A Public-Key Cryptosystem Based On Algebraic Coding Theory”. In: *Deep Space Network Progress Report* 44 (Jan. 1978), pp. 114–116. URL: <https://ui.adsabs.harvard.edu/abs/1978DSNPR..44..114M> (cit. on p. 1).
- [15] A. Sepúlveda Castellanos and G. C. Tizziotti. “Two-Point AG Codes on the GK Maximal Curves”. In: *IEEE Transactions on Information Theory* 62.2 (2016), pp. 681–686. doi: [10.1109/TIT.2015.2511787](https://doi.org/10.1109/TIT.2015.2511787) (cit. on p. 1).
- [16] Jean-Pierre Serre. “Singular Algebraic Curves”. In: *Algebraic Groups and Class Fields*. New York, NY: Springer New York, 1988, pp. 58–73. ISBN: 978-1-4612-1035-1. doi: [10.1007/978-1-4612-1035-1\\_4](https://doi.org/10.1007/978-1-4612-1035-1_4) (cit. on p. 4).