

THE EXACT SOLUTIONS OF CERTAIN LINEAR PARTIAL DIFFERENCE EQUATIONS

CHUN-KAI HWANG AND TZON-TZER LU

ABSTRACT. Difference equations have many applications and play an important role in numerical analysis, probability, statistics, combinatorics, computer science, and quantum consciousness, etc. We first prove that the partial differential equation is equivalent to partial difference equation with an example of heat equation. Additionally, we use generating functions to find the exact solutions of some simple linear partial difference equations. Then we extend it to more general partial difference equations of higher dimensions and obtain their solutions. We conclude that using multivariable power series as generating function is a very efficient method to solve partial difference equations.

1. INTRODUCTION

Many phenomena or problems occur in the natural world can be formulated by the mathematical model of partial difference equations, such as discrete potential theories, random walk problems, finite difference schemes, gambling problems, chemical structures, discrete heat diffusion equations,....etc. Roughly speaking, a partial difference equation is a functional relation involving functions with multivariate discrete variables over a discrete set.

Next, we take random work problems and heat diffusion as examples. Suppose that an ant creeps along the one dimensional straight line, x-axis. Assume the probability it doesn't move in a given unit of time is d , the probability that it moves one unit to the positive and negative directions along the x-axis in a unit of time are p and q respectively. Let $a(i, j)$ be the probability that the ant is at the point $x = i$ after j -th units of time. Then solving $a(i, j)$ is equivalent to solving the following difference equation

$$a(i, j + 1) = pa(i - 1, j) + da(i, j) + qa(i + 1, j).$$

If the ant starts from the origin at time 0, then we have initial conditions

$$a(i, 0) = \begin{cases} 1, & \text{if } i = 0, \\ 0, & \text{otherwise.} \end{cases}$$

Next, we discuss the discrete version of the heat equation

$$\frac{\partial u}{\partial t} = b \frac{\partial^2 u}{\partial x^2}, t > 0, -\infty < x < \infty.$$

Consider the distribution of heat through a long rod. Assume that the rod is long enough so that we can sample it with the discrete integer set. Let $U_{i,j}$ be the temperature at the position i after j -th units of time. At time j , if the temperature

Received by the editors January 18, 2023 .

Key words and phrases. partial difference equations, generating functions.

$U_{i-1,j}$ is higher than $U_{i,j}$, heat will flow from the position $i - 1$ to i . From the experiment, the increase is proportional to the $U_{i-1,j} - U_{i,j}$, say $r(U_{i-1,j} - U_{i,j})$, where r is a physical parameter depending on the properties of the rod. Similarly, heat will flow from the point $i + 1$ to i . Hence, the total effects can be written as the following discrete Newton law of cooling

$$U_{i,j+1} - U_{i,j} = r(U_{i-1,j} - U_{i,j}) + r(U_{i+1,j} - U_{i,j}) = r\Delta_1^2 U_{i-1,j}.$$

We can regard the equation above as a special case of the difference equation

$$U_{i,j+1} = aU_{i-1,j} + bU_{i,j} + cU_{i+1,j},$$

where $i \in Z, j \in N \cup \{0\}$. Although they have different interpretation, they correspond to the same type of equation. The exact solution of this equation will be computed in Theorem 2.1.

There are many methods developed to solve the difference equation, such as the symbolic method, Lagrange method [1], generating functions [2], symbolic calculus [4], characteristic equations, matrices iterations[5],...,etc. We will use multivariable power series as generating function to solve the exact solutions of certain partial difference equations. We demonstrate its usage for some simple cases in section 2, then extend it to general case in section 3 and multidimensional cases in section 4. It can be seen that this method is very efficient to solve the partial difference equations.

2. SOME SIMPLE CASES

Consider the following linear partial difference equations:

$$(2.1) \quad U_{i,j+1} = aU_{i-1,j} + bU_{i,j} + cU_{i+1,j}$$

$$(2.2) \quad U_{i+m,j+1} = c_1U_{i,j} + c_2U_{i+1,j} + \dots + c_nU_{i+(n-1),j}$$

$$(2.3) \quad U_{i+1,j+1} = aU_{i,j+1} + bU_{i+1,j} + cU_{i,j}$$

We will obtain the explicit solution of $U_{i,j}$ under the assumption of $U_{i,0}$ given for all $i \in Z$. That is, if we know all of the physical quantity on the x-axis, we can predict the other quantity $U_{i,j}$ at any position in the upper plane.

We first consider the equation(2.1).Lin[3] deals with more general nonhomogeneous case of this equation.

Theorem 2.1. *Let*

$$U_{i,j+1} = aU_{i-1,j} + bU_{i,j} + cU_{i+1,j}$$

with $i \in Z, j \in N \cup \{0\}$. Given $U_{i,0} = \Psi(i)$, then its solution is of the form

$$U_{i,j} = \sum_{m=0}^j \sum_{n=0}^m \binom{j}{m} \binom{m}{n} a^n b^{m-n} c^{j-n} \Psi(i + j - m - n).$$

Proof. Let

$$h(x, y) = \sum_{i=-\infty}^{\infty} \sum_{j=0}^{\infty} U_{i,j} x^i y^j.$$

Then we have

$$\sum_{i=-\infty}^{\infty} \sum_{j=0}^{\infty} U_{i,j+1} x^i y^j = a \sum_{i=-\infty}^{\infty} \sum_{j=0}^{\infty} U_{i-1,j} x^i y^j + b \sum_{i=-\infty}^{\infty} \sum_{j=0}^{\infty} U_{i,j} x^i y^j + c \sum_{i=-\infty}^{\infty} \sum_{j=0}^{\infty} U_{i+1,j} x^i y^j,$$

$$\frac{1}{y} [h(x, y) - \sum_{i=-\infty}^{\infty} \Psi(i) x^i] = axh(x, y) + bh(x, y) + \frac{c}{x} h(x, y),$$

$$h(x, y) [1 - (ax + b + \frac{c}{x})y] = \sum_{i=-\infty}^{\infty} \Psi(i) x^i,$$

$$\begin{aligned} h(x, y) &= \frac{\sum_{i=-\infty}^{\infty} \Psi(i) x^i}{1 - (ax + b + cx^{-1})y} \\ &= \sum_{i=-\infty}^{\infty} \Psi(i) x^i \sum_{j=0}^{\infty} (ax + b + cx^{-1})^j y^j \\ &= \sum_{i=-\infty}^{\infty} \Psi(i) x^i \sum_{j=0}^{\infty} \left[\sum_{\substack{0 \leq \alpha, \beta, \gamma \leq j, \\ \alpha + \beta + \gamma = j}} \binom{j}{\alpha \beta \gamma} a^\alpha b^\beta c^\gamma x^{\alpha - \gamma} \right] y^j \\ &= \sum_{j=0}^{\infty} \sum_{i=-\infty}^{\infty} \Psi(i) x^i \left[\sum_{\substack{0 \leq \alpha, \beta, \gamma \leq j, \\ \alpha + \beta + \gamma = j}} \binom{j}{\alpha \beta \gamma} a^\alpha b^\beta c^\gamma x^{\alpha - \gamma} \right] y^j. \end{aligned}$$

Let $\alpha = n, \beta = m - n, \gamma = j - m$, then we have

$$\begin{aligned} h(x, y) &= \sum_{j=0}^{\infty} \left\{ \sum_{i=-\infty}^{\infty} \Psi(i) x^i \sum_{m=0}^j \sum_{n=0}^m \binom{j}{n, m-n, j-n} a^n b^{m-n} c^{j-n} x^{n-j+m} \right\} y^j \\ &= \sum_{j=0}^{\infty} \left\{ \sum_{i=-\infty}^{\infty} \sum_{m=0}^j \sum_{n=0}^m \binom{j}{n, m-n, j-n} a^n b^{m-n} c^{j-n} \Psi(i) x^{i-j+m+n} \right\} y^j. \end{aligned}$$

Change of variable $k = i - j + m + n$ to obtain

$$h(x, y) = \sum_{j=0}^{\infty} \left\{ \sum_{i=-\infty}^{\infty} \sum_{m=0}^j \sum_{n=0}^m \binom{j}{n, m-n, j-n} a^n b^{m-n} c^{j-n} \Psi(k + j - m - n) x^k \right\} y^j.$$

Thus the solution is of the form

$$U_{i,j} = \sum_{m=0}^j \sum_{n=0}^m \binom{j}{m} \binom{m}{n} a^n b^{m-n} c^{j-n} \Psi(i + j - m - n).$$

□

Theorem 2.2. *Let*

$$U_{i+m,j+1} = c_1 U_{i,j} + c_2 U_{i+1,j} + \dots + c_n U_{i+(n-1),j}$$

with $i \in Z, j \in N \cup \{0\}$. Given $U_{i,0} = \Psi(i)$, then its solution is of the form

$$U_{i,j} = \sum_{s_r \in I} \binom{j}{s_1 s_2 \dots s_n} \prod_{r=1}^n c_r^{s_r} \Psi(i + \sum_{r=1}^n r s_r - (m+1)j),$$

where $I = \{s_1, s_2, \dots, s_n | 0 \leq s_k \leq j, \sum_{k=1}^n s_k = j\}$.

Proof. Let

$$h(x, y) = \sum_{i=-\infty}^{\infty} \sum_{j=0}^{\infty} U_{i,j} x^i y^j.$$

Then we have

$$\begin{aligned} & \sum_{i=-\infty}^{\infty} \sum_{j=0}^{\infty} U_{i+m,j+1} x^i y^j \\ = & c_1 \sum_{i=-\infty}^{\infty} \sum_{j=0}^{\infty} U_{i,j} x^i y^j + c_2 \sum_{i=-\infty}^{\infty} \sum_{j=0}^{\infty} U_{i+1,j} x^i y^j + \dots + c_n \sum_{i=-\infty}^{\infty} \sum_{j=0}^{\infty} U_{i+(n-1),j} x^i y^j, \end{aligned}$$

$$\frac{1}{x^m y} [h(x, y) - \sum_{i=-\infty}^{\infty} \Psi(i) x^i] = \sum_{t=1}^n c_t x^{1-t} h(x, y),$$

$$h(x, y) (1 - \sum_{t=1}^n c_t x^{m-t+1} y) = \sum_{i=-\infty}^{\infty} \Psi(i) x^i,$$

$$\begin{aligned} h(x, y) &= \frac{\sum_{i=-\infty}^{\infty} \Psi(i) x^i}{1 - (c_1 x^m + c_2 x^{m-1} + \dots + c_n x^{m-(n-1)}) y} \\ &= \sum_{i=-\infty}^{\infty} \Psi(i) x^i \sum_{j=0}^{\infty} (c_1 x^m + c_2 x^{m-1} + \dots + c_n x^{m-(n-1)})^j y^j \\ &= \sum_{i=-\infty}^{\infty} \Psi(i) x^i \sum_{j=0}^{\infty} \sum_{s_r \in I} \binom{j}{s_1 s_2 \dots s_n} (c_1^{s_1} x^{m s_1}) (c_2^{s_2} x^{(m-1) s_2}) \dots (c_n^{s_n} x^{[m-(n-1)] s_n}) y^j \\ &= \sum_{i=-\infty}^{\infty} \Psi(i) x^i \sum_{j=0}^{\infty} \sum_{s_r \in I} \binom{j}{s_1 s_2 \dots s_n} \prod_{r=1}^n c_r^{s_r} x^a y^j \\ &= \sum_{j=0}^{\infty} \left\{ \sum_{i=-\infty}^{\infty} \Psi(i) x^i \sum_{s_r \in I} \binom{j}{s_1 s_2 \dots s_n} \prod_{r=1}^n c_r^{s_r} x^a \right\} y^j \\ &= \sum_{j=0}^{\infty} \left\{ \sum_{i=-\infty}^{\infty} \sum_{s_r \in I} \binom{j}{s_1 s_2 \dots s_n} \prod_{r=1}^n c_r^{s_r} \Psi(i) x^{a+i} \right\} y^j, \end{aligned}$$

where $a = \sum_{r=1}^n [m - (r-1)] s_r$, $I = \{s_1, s_2, \dots, s_n | 0 \leq s_k \leq j, \sum_{k=1}^n s_k = j\}$. Let

$$k = a + i.$$

Then we have

$$\begin{aligned}
 h(x, y) &= \sum_{j=0}^{\infty} \sum_{k=-\infty}^{\infty} \sum_{s_r \in I} \binom{j}{s_1 s_2 \dots s_n} \prod_{r=1}^n c_r^{s_r} \Psi(k-a) x^k y^j \\
 &= \sum_{j=0}^{\infty} \sum_{k=-\infty}^{\infty} \sum_{s_r \in I} \binom{j}{s_1 s_2 \dots s_n} \prod_{r=1}^n c_r^{s_r} \Psi(k - \sum_{r=1}^n [m - (r-1)] s_r) x^k y^j \\
 &= \sum_{j=0}^{\infty} \sum_{k=-\infty}^{\infty} \sum_{s_r \in I} \binom{j}{s_1 s_2 \dots s_n} \prod_{r=1}^n c_r^{s_r} \Psi(k - \sum_{r=1}^n (m+1) s_r + \sum_{r=1}^n r s_r) x^k y^j \\
 &= \sum_{j=0}^{\infty} \sum_{k=-\infty}^{\infty} \sum_{s_r \in I} \binom{j}{s_1 s_2 \dots s_n} \prod_{r=1}^n c_r^{s_r} \Psi(k + \sum_{r=1}^n r s_r - (m-1)j) x^k y^j.
 \end{aligned}$$

Thus the solution is of the form

$$U_{i,j} = \sum_{s_r \in I} \binom{j}{s_1 s_2 \dots s_n} \prod_{r=1}^n c_r^{s_r} \Psi(i + \sum_{r=1}^n r s_r - (m+1)j),$$

where $I = \{s_1, s_2, \dots, s_n | 0 \leq s_k \leq j, \sum_{k=1}^n s_k = j\}$. □

Next, we turn to (2.3). Suppose $U_{i,j}$ is known on x-axis, in general (2.3) has infinitely many solutions. Consider the typical example where $a = 1$, $b = 1$, and $c = -1$ then $U_{i,j} = c_{i,j}$ is a solution. Since there are infinitely many choices of constants c_j for $j = 0, 1, \dots$, we then have infinite solutions. Using generating functions, we can find one special solution of (2.3).

Theorem 2.3. *Let*

$$U_{i+1,j+1} = aU_{i,j+1} + bU_{i+1,j} + cU_{i,j}$$

with $i \in Z$, $j \in N \cup \{0\}$ Given $U_{i,0} = \Psi(i)$, then it has following solution

$$U_{i,j} = \sum_{s=0}^{\infty} \Omega(i-s) p(s, j),$$

where

$$\begin{aligned}
 \Omega(k) &= \Psi(k) - a\Psi(k-1), \\
 p(j, t) &= \sum_{r=0}^t \binom{j}{t} \binom{t}{r} a^{j-t} b^{t-r} c^t.
 \end{aligned}$$

Proof. Let

$$\sum_{i=-\infty}^{\infty} \sum_{j=0}^{\infty} U_{i,j} x^i y^j = h(x, y).$$

Then we have

$$\begin{aligned}
 &\sum_{i=-\infty}^{\infty} \sum_{j=0}^{\infty} U_{i+1,j+1} x^i y^j \\
 &= a \sum_{i=-\infty}^{\infty} \sum_{j=0}^{\infty} U_{i,j+1} x^i y^j + b \sum_{i=-\infty}^{\infty} \sum_{j=0}^{\infty} U_{i+1,j} x^i y^j + c \sum_{i=-\infty}^{\infty} \sum_{j=0}^{\infty} U_{i,j} x^i y^j,
 \end{aligned}$$

$$\begin{aligned}
\frac{1}{xy}[h(x, y) - \sum_{i=-\infty}^{\infty} \Psi(i)x^i] &= \frac{a}{y}[h(x, y) - \sum_{i=-\infty}^{\infty} \Psi(i)x^i] + \frac{b}{x}h(x, y) + ch(x, y), \\
h(x, y)[1 - (ax + by + cxy)] &= \sum_{i=-\infty}^{\infty} \Psi(i)x^i - a \sum_{i=-\infty}^{\infty} \Psi(i)x^{i+1} \\
&= \sum_{k=-\infty}^{\infty} [\Psi(k) - a\Psi(k-1)]x^k,
\end{aligned}$$

$$\begin{aligned}
h(x, y) &= \frac{\sum_{k=-\infty}^{\infty} [\Psi(k) - a\Psi(k-1)]x^k}{1 - (ax + by + cxy)} \\
&= \sum_{k=-\infty}^{\infty} [\Psi(k) - a\Psi(k-1)]x^k \sum_{j=0}^{\infty} (ax + by + cxy)^j \\
&= \sum_{k=-\infty}^{\infty} \sum_{j=0}^{\infty} [\Psi(k) - a\Psi(k-1)]x^k (ax + (b + cx)y)^j \\
&= \sum_{k=-\infty}^{\infty} \sum_{j=0}^{\infty} [\Psi(k) - a\Psi(k-1)]x^k \sum_{t=0}^j \binom{j}{t} (ax)^{j-t} (b + cx)^t y^t \\
&= \sum_{k=-\infty}^{\infty} \sum_{j=0}^{\infty} [\Psi(k) - a\Psi(k-1)]x^k \sum_{t=0}^j \binom{j}{t} \left[\sum_{r=0}^t \binom{t}{r} b^{t-r} c^r x^t \right] a^{j-t} x^{j-t} y^t \\
&= \sum_{k=-\infty}^{\infty} \sum_{j=0}^{\infty} [\Psi(k) - a\Psi(k-1)]x^k \sum_{t=0}^j \sum_{r=0}^t \binom{j}{t} \binom{t}{r} b^{t-r} c^r x^t a^{j-t} x^{j-t} y^t \\
&= \sum_{k=-\infty}^{\infty} \sum_{j=0}^{\infty} [\Psi(k) - a\Psi(k-1)]x^k \sum_{t=0}^j \sum_{r=0}^t \binom{j}{t} \binom{t}{r} a^{j-t} b^{t-r} c^r x^j y^t \\
&= \sum_{k=-\infty}^{\infty} \sum_{j=0}^{\infty} [\Psi(k) - a\Psi(k-1)] \sum_{t=0}^j \sum_{r=0}^t \binom{j}{t} \binom{t}{r} a^{j-t} b^{t-r} c^r x^{j+k} y^t.
\end{aligned}$$

Let

$$\begin{aligned}
\Psi(k) - a\Psi(k-1) &= \Omega(k), \\
\sum_{r=0}^t \binom{j}{t} \binom{t}{r} a^{j-t} b^{t-r} c^r &= p(j, t), \\
j + k &= u.
\end{aligned}$$

Then we have

$$\begin{aligned}
 h(x, y) &= \sum_{u=-\infty}^{\infty} \sum_{j=0}^{\infty} \Omega(u-j) \sum_{t=0}^j p(j, t) x^u y^t \\
 &= \sum_{u=-\infty}^{\infty} x^u \sum_{j=0}^{\infty} \Omega(u-j) \sum_{t=0}^j p(j, t) y^t \\
 &= \sum_{u=-\infty}^{\infty} x^u \Omega(u) p(0, 0) + \\
 &\quad \sum_{u=-\infty}^{\infty} x^u \Omega(u-1) [p(1, 0) + p(1, 1)y] + \dots + \\
 &\quad \sum_{u=-\infty}^{\infty} x^u \Omega(u-k) [p(k, 0) + p(k, 1)y + \dots + p(k, k)y^k] + \dots \\
 &= \sum_{u=-\infty}^{\infty} x^u \sum_{j=0}^{\infty} \sum_{s=0}^{\infty} \Omega(u-s) p(s, j) y^j \\
 &= \sum_{i=-\infty}^{\infty} \sum_{j=0}^{\infty} \sum_{s=0}^{\infty} \Omega(i-s) p(s, j) x^i y^j
 \end{aligned}$$

Thus the solution is

$$U_{i,j} = \sum_{s=0}^{\infty} \Omega(i-s) p(s, j),$$

where

$$\begin{aligned}
 \Omega(k) &= \Psi(k) - a\Psi(k-1), \\
 p(j, t) &= \sum_{r=0}^t \binom{j}{t} \binom{t}{r} a^{j-t} b^{t-r} c^t.
 \end{aligned}$$

□

Notice that in above theorem, we have the initial condition on x-axis and then obtain the solution on the upper half plane. Suppose we are only given the initial data on an interval or ray of x-axis. On its domain of influence, the solutions of (2.1), (2.2), (2.3) are also given by the corresponding theorems.

3. THE GENERAL CASE

In this section, we discuss the general solution of the partial difference equation

$$(3.1) \quad U_{i+m, j+k} = \sum_{s=1}^k \sum_{r=1}^n c_{rs} U_{i+(r-1), j+(s-1)},$$

which is an extension of the equation(2.2) in the previous section. We try to derive its explicit solution with given initial conditions data $U_{i,t} = \Psi_t(i)$, $t = 1, 2, \dots, k-1$. The theorem below is the case for $k = 2$.

Theorem 3.1. *Let*

$$U_{i+m,j+k} = \sum_{s=1}^k \sum_{r=1}^n c_{rs} U_{i+(r-1),j+(s-1)}$$

with $i \in Z$, $j \in N \cup \{0\}$. Given $U_{i,t} = \Psi_t(i)$, for $t = 0, 1$, then its solution $U_{i,j}$ is of the form

$$\begin{aligned} & \sum_{l_{rs} \in I_1} \left(\frac{j + \sum_{s=1}^2 \sum_{r=1}^n s l_{rs}}{3} \right) \prod_{\substack{s=1,2, \\ r=1,2,\dots,n}} c_{rs}^{lrs} \\ & \Psi_0 \left(i + \sum_{s=1}^2 \sum_{r=1}^n r l_{rs} - (m+1) \frac{j + \sum_{s=1}^2 \sum_{r=1}^n s l_{rs}}{3} \right) + \\ & \sum_{l_{rs} \in I_2} \left(\frac{(j-1) + \sum_{s=1}^2 \sum_{r=1}^n s l_{rs}}{3} \right) \prod_{\substack{s=1,2, \\ r=1,2,\dots,n}} c_{rs}^{lrs} \Psi_1 \left(i + \sum_{s=1}^2 \sum_{r=1}^n r l_{rs} - (m+1) \right. \\ & \left. \frac{(j-1) + \sum_{s=1}^2 \sum_{r=1}^n s l_{rs}}{3} \right) - \\ & \sum_{e=1}^n c_{e2} \sum_{l_{rs} \in I_2} \left(\frac{(j-1) + \sum_{s=1}^2 \sum_{r=1}^n s l_{rs}}{3} \right) \prod_{\substack{s=1,2, \\ r=1,2,\dots,n}} c_{rs}^{lrs} \\ & \Psi_0 \left(i + e - 1 - m + \sum_{s=1}^2 \sum_{r=1}^n r l_{rs} - (m+1) \frac{(j-1) + \sum_{s=1}^2 \sum_{r=1}^n s l_{rs}}{3} \right), \end{aligned}$$

where

$$\begin{aligned} I_1 &= \{l_{11}, l_{12}, \dots, l_{n2} \mid 0 \leq l_{rs} \leq \frac{j + \sum_{s=1}^2 \sum_{r=1}^n s l_{rs}}{3}, \\ \sum_{s=1}^2 \sum_{r=1}^n l_{rs} &= \frac{j + \sum_{s=1}^2 \sum_{r=1}^n s l_{rs}}{3}\}, \\ I_2 &= \{l_{11}, l_{12}, \dots, l_{n2} \mid 0 \leq l_{rs} \leq \frac{(j-1) + \sum_{s=1}^2 \sum_{r=1}^n s l_{rs}}{3}, \\ \sum_{s=1}^2 \sum_{r=1}^n l_{rs} &= \frac{(j-1) + \sum_{s=1}^2 \sum_{r=1}^n s l_{rs}}{3}\}, \end{aligned}$$

for $s=1, 2, r=1, 2, \dots, n$.

Proof. Let $h(x, y) = \sum_{i=-\infty}^{\infty} \sum_{j=0}^{\infty} U_{i,j} x^i y^j$ and $k = 2$. Then we have

$$\sum_{i=-\infty}^{\infty} \sum_{j=0}^{\infty} U_{i+m,j+k} x^i y^j = \sum_{s=1}^k \sum_{r=1}^n c_{rs} U_{i+(r-1),j+(s-1)} x^i y^j,$$

$$\begin{aligned}
 & \frac{1}{x^m y^k} [h(x, y) - \sum_{r=0}^{k-1} \sum_{i=-\infty}^{\infty} \Psi_r(i) x^i y^r] \\
 = & \sum_{r=1}^n c_{r1} x^{-(r-1)} h(x, y) + \sum_{r=1}^n c_{r2} x^{-(r-1)} y^{-1} [h(x, y) - \sum_{i=-\infty}^{\infty} \Psi_0(i) x^i] + \\
 & \sum_{r=1}^n c_{r3} x^{-(r-1)} y^{-2} [h(x, y) - \sum_{t=0}^1 \sum_{i=-\infty}^{\infty} \Psi_t(i) x^i y^t] + \dots + \\
 & \sum_{r=1}^n c_{rk} x^{-(r-1)} y^{-(k-1)} [h(x, y) - \sum_{t=0}^{k-2} \sum_{i=-\infty}^{\infty} \Psi_t(i) x^i y^t],
 \end{aligned}$$

$$\begin{aligned}
 & L(x, y) \\
 = & h(x, y) [1 - (\sum_{r=1}^n c_{r1} x^{m-(r-1)} y^k + \sum_{r=1}^n c_{r2} x^{m-(r-1)} y^{k-1} + \dots + \sum_{r=1}^n c_{rk} x^{m-(r-1)} y)] \\
 = & \sum_{r=0}^{k-1} \sum_{i=-\infty}^{\infty} \Psi_r(i) x^i y^r - \sum_{s=2}^k [\sum_{r=1}^n c_{rs} x^{m-(r-1)} y^{k-s+1} \sum_{t=0}^{s-2} \sum_{i=-\infty}^{\infty} \Psi_t(i) x^i y^t] \\
 = & \sum_{i=-\infty}^{\infty} \sum_{r=0}^{k-1} \Psi_r(i) x^i y^r - \sum_{i=-\infty}^{\infty} \sum_{s=2}^k \sum_{t=0}^{s-2} \sum_{r=1}^n c_{rs} \Psi_t(i) x^{m-(r-1)+i} y^{k-s+1+t} \\
 = & \sum_{i=-\infty}^{\infty} \sum_{r=0}^{k-1} \Psi_r(i) x^i y^r - \sum_{i=-\infty}^{\infty} \sum_{h=2}^k \sum_{t=0}^{h-2} \sum_{e=1}^n c_{eh} \Psi_t(i) x^{i+m-e+1} y^{k-h+1+t},
 \end{aligned}$$

$$\begin{aligned}
 & h(x, y) \\
 = & L(x, y) [1 - \sum_{s=1}^k \sum_{r=1}^n c_{rs} x^{m-(r-1)} y^{k-s+1}]^{-1} \\
 = & L(x, y) \sum_{j=0}^{\infty} [\sum_{s=1}^k \sum_{r=1}^n c_{rs} x^{m-(r-1)} y^{k-s+1}]^j \\
 = & L(x, y) \sum_{j=0}^{\infty} \sum_{l_{rs} \in I} \binom{j}{l_{11} l_{12} \dots l_{1n} l_{21} \dots l_{nk}} \prod_{\substack{r=1,2,\dots,n, \\ s=1,2,\dots,k}} c_{rs}^{l_{rs}} x^a y^b \\
 = & \sum_{j=0}^{\infty} L(x, y) \sum_{l_{rs} \in I} \binom{j}{l_{11} l_{12} \dots l_{1n} l_{21} \dots l_{nk}} \prod_{\substack{r=1,2,\dots,n, \\ s=1,2}} c_{rs}^{l_{rs}} x^a y^b,
 \end{aligned}$$

where

$$\begin{aligned}
I &= \{l_{11}, l_{12}, \dots, l_{nk} \mid 0 \leq l_{rs} \leq j, \sum_{s=1}^k \sum_{r=1}^n l_{rs} = j\}, \\
a &= \sum_{s=1}^k \sum_{r=1}^n (m-r+1)l_{rs} \\
&= (m+1)j - \sum_{s=1}^k \sum_{r=1}^n r l_{rs}, \\
b &= \sum_{s=1}^k \sum_{r=1}^n (k-s+1)l_{rs} \\
&= (k+1) \sum_{s=1}^k \sum_{r=1}^n l_{rs} - \sum_{s=1}^k \sum_{r=1}^n s l_{rs} \\
&= (k+1)j - \sum_{s=1}^k \sum_{r=1}^n s l_{rs},
\end{aligned}$$

for $r=1, 2, \dots, n, s=1, 2, \dots, k$. When $k=2$, we have

$$\begin{aligned}
I &= \{l_{11}, l_{12}, \dots, l_{n \times 2} \mid 0 \leq l_{rs} \leq j, \sum_{s=1}^2 \sum_{r=1}^n l_{rs} = j\}, \\
a &= (m+1)j - \sum_{s=1}^2 \sum_{r=1}^n r l_{rs}, \\
b &= 3j - \sum_{s=1}^2 \sum_{r=1}^n s l_{rs},
\end{aligned}$$

and

$$\begin{aligned}
L(x, y) &= \sum_{i=-\infty}^{\infty} \sum_{r=0}^1 \Psi_r(i) x^i y^r - \sum_{i=-\infty}^{\infty} \sum_{e=1}^n c_{e2} \Psi_0(i) x^{i+m-e+1} y \\
&= \sum_{i=-\infty}^{\infty} \Psi_0(i) x^i + \sum_{i=-\infty}^{\infty} \Psi_1(i) x^i y - \sum_{i=-\infty}^{\infty} \sum_{e=1}^n c_{e2} \Psi_0(i) x^{i+m-e+1} y
\end{aligned}$$

Hence

$$\begin{aligned}
 h(x, y) &= \sum_{j=0}^{\infty} \left\{ \sum_{i=-\infty}^{\infty} \Psi_0(i) x^i + \sum_{i=-\infty}^{\infty} \Psi_1(i) x^i y - \sum_{i=-\infty}^{\infty} \sum_{e=1}^n c_{e2} \Psi_0(i) x^{i+m-e+1} y \right\} \\
 &\quad \sum_{l_{rs} \in I} \binom{j}{l_{11} l_{12} \dots l_{1n} l_{21} \dots l_{nk}} \prod_{\substack{r=1,2,\dots,n, \\ s=1,2,\dots,k}} c_{rs}^{l_{rs}} x^a y^b \\
 &= \sum_{j=0}^{\infty} \sum_{i=-\infty}^{\infty} \Psi_0(i) x^i \sum_{l_{rs} \in I} \binom{j}{l_{11} l_{12} \dots l_{1n} l_{21} \dots l_{nk}} \prod_{\substack{r=1,2,\dots,n, \\ s=1,2,\dots,k}} c_{rs}^{l_{rs}} x^a y^b + \\
 &\quad \sum_{j=0}^{\infty} \sum_{i=-\infty}^{\infty} \Psi_1(i) x^i y \sum_{l_{rs} \in I} \binom{j}{l_{11} l_{12} \dots l_{1n} l_{21} \dots l_{nk}} \prod_{\substack{r=1,2,\dots,n, \\ s=1,2,\dots,k}} c_{rs}^{l_{rs}} x^a y^b - \\
 &\quad \sum_{j=0}^{\infty} \sum_{i=-\infty}^{\infty} \left[\sum_{e=1}^n c_{e2} \Psi_0(i) x^{i+m-e+1} \right] y \sum_{l_{rs} \in I} \binom{j}{l_{11} l_{12} \dots l_{1n} l_{21} \dots l_{nk}} \prod_{\substack{r=1,2,\dots,n, \\ s=1,2,\dots,k}} c_{rs}^{l_{rs}} x^a y^b \\
 &= \sum_{j=0}^{\infty} \sum_{i=-\infty}^{\infty} \sum_{l_{rs} \in I} \binom{j}{l_{11} l_{12} \dots l_{1n} l_{21} \dots l_{nk}} \Psi_0(i) \prod_{\substack{r=1,2,\dots,n, \\ s=1,2,\dots,k}} c_{rs}^{l_{rs}} x^{a+i} y^b + \\
 &\quad \sum_{j=0}^{\infty} \sum_{i=-\infty}^{\infty} \sum_{l_{rs} \in I} \binom{j}{l_{11} l_{12} \dots l_{1n} l_{21} \dots l_{nk}} \Psi_1(i) \prod_{\substack{r=1,2,\dots,n, \\ s=1,2,\dots,k}} c_{rs}^{l_{rs}} x^{a+i} y^{b+1} - \\
 &\quad \sum_{j=0}^{\infty} \sum_{i=-\infty}^{\infty} \sum_{e=1}^n c_{e2} \sum_{l_{rs} \in I} \binom{j}{l_{11} l_{12} \dots l_{1n} l_{21} \dots l_{nk}} \Psi_0(i) \prod_{\substack{r=1,2,\dots,n, \\ s=1,2,\dots,k}} c_{rs}^{l_{rs}} x^a y^b x^{a+i+m-e+1} y^{b+1}
 \end{aligned}$$

Let

$$a + i = \alpha, b + 1 = 3j - \sum_{s=1}^2 \sum_{r=1}^n s l_{rs} + 1 = \beta, a + i + m - e + 1 = \gamma.$$

Then

$$\begin{aligned}
 j &= \frac{b + \sum_{s=1}^2 \sum_{r=1}^n s l_{rs}}{3} = \frac{(\beta - 1 + \sum_{s=1}^2 \sum_{r=1}^n s l_{rs})}{3}, \\
 i &= \alpha - a \\
 &= \alpha - (m + 1)j + \sum_{s=1}^2 \sum_{r=1}^n r l_{rs} \\
 &= \alpha + \sum_{s=1}^2 \sum_{r=1}^n r l_{rs} - (m + 1) \frac{b + \sum_{s=1}^2 \sum_{r=1}^n s l_{rs}}{3} \\
 &= \alpha + \sum_{s=1}^2 \sum_{r=1}^n r l_{rs} - (m + 1) \frac{(\beta - 1 + \sum_{s=1}^2 \sum_{r=1}^n s l_{rs})}{3} \\
 &= \gamma + e - 1 - m - a \\
 &= \gamma + e - 1 - m + \sum_{s=1}^2 \sum_{r=1}^n r l_{rs} - (m + 1) \frac{(\beta - 1 + \sum_{s=1}^2 \sum_{r=1}^n s l_{rs})}{3},
 \end{aligned}$$

and

$$\begin{aligned}
h(x, y) &= \sum_{b=0}^{\infty} \sum_{\alpha=-\infty}^{\infty} \sum_{l_{rs} \in I_1} \left(\frac{b + \sum_{s=1}^2 \sum_{r=1}^n sl_{rs}}{3} \right) \prod_{\substack{s=1,2, \\ r=1,2,\dots,n}} c_{rs}^{lrs} x^\alpha y^b \\
\Psi_0(\alpha + \sum_{s=1}^2 \sum_{r=1}^n rl_{rs} - (m+1) \frac{b + \sum_{s=1}^2 \sum_{r=1}^n sl_{rs}}{3}) &+ \\
\sum_{\beta=0}^{\infty} \sum_{\alpha=-\infty}^{\infty} \sum_{l_{rs} \in I_2} \left(\frac{(\beta-1 + \sum_{s=1}^2 \sum_{r=1}^n sl_{rs})}{3} \right) &\prod_{\substack{s=1,2, \\ r=1,2,\dots,n}} c_{rs}^{lrs} x^\alpha y^\beta \\
\Psi_1(\alpha + \sum_{s=1}^2 \sum_{r=1}^n rl_{rs} - (m+1) \frac{(\beta-1 + \sum_{s=1}^2 \sum_{r=1}^n sl_{rs})}{3}) &- \\
\sum_{\beta=0}^{\infty} \sum_{\gamma=-\infty}^{\infty} \sum_{\epsilon=1}^n c_{e2} \sum_{l_{rs} \in I_2} \left(\frac{(\beta-1 + \sum_{s=1}^2 \sum_{r=1}^n sl_{rs})}{3} \right) &\prod_{\substack{s=1,2, \\ r=1,2,\dots,n}} c_{rs}^{lrs} x^\gamma y^\beta \\
\Psi_0(\gamma + \epsilon - 1 - m + \sum_{s=1}^2 \sum_{r=1}^n rl_{rs} - (m+1) \frac{(\beta-1 + \sum_{s=1}^2 \sum_{r=1}^n sl_{rs})}{3}) & \\
= \sum_{u=0}^{\infty} \sum_{v=-\infty}^{\infty} \sum_{l_{rs} \in I_1} \left(\frac{u + \sum_{s=1}^2 \sum_{r=1}^n sl_{rs}}{3} \right) \prod_{\substack{s=1,2, \\ r=1,2,\dots,n}} c_{rs}^{lrs} & \\
\Psi_0(v + \sum_{s=1}^2 \sum_{r=1}^n rl_{rs} - (m+1) \frac{u + \sum_{s=1}^2 \sum_{r=1}^n sl_{rs}}{3}) &+ \\
\sum_{l_{rs} \in I_2} \left(\frac{(u-1 + \sum_{s=1}^2 \sum_{r=1}^n sl_{rs})}{3} \right) \prod_{\substack{s=1,2, \\ r=1,2,\dots,n}} c_{rs}^{lrs} \Psi_1(v + \sum_{s=1}^2 \sum_{r=1}^n rl_{rs} - (m+1) & \\
\frac{(u-1 + \sum_{s=1}^2 \sum_{r=1}^n sl_{rs})}{3}) &- \\
\sum_{e=1}^n c_{e2} \sum_{l_{rs} \in I_2} \left(\frac{(u-1 + \sum_{s=1}^2 \sum_{r=1}^n sl_{rs})}{3} \right) \prod_{\substack{s=1,2, \\ r=1,2,\dots,n}} c_{rs}^{lrs} & \\
\Psi_0(v + \epsilon - 1 - m + \sum_{s=1}^2 \sum_{r=1}^n rl_{rs} - (m+1) \frac{(u-1 + \sum_{s=1}^2 \sum_{r=1}^n sl_{rs})}{3}) &x^v y^u.
\end{aligned}$$

Thus for $k=2$, the solution $U_{i,j}$ has the desired form. \square

4. THE MULTIDIMENSIONAL CASES

In this section, we discuss the linear partial difference equations in higher dimensions. At the begining, we solve the following equations in three dimensions where $i, j \in Z$, and $k \in N \cup \{0\}$.

$$(4.1) \quad \begin{aligned} U_{i,j,k+1} = & c_1 U_{i-1,j-1,k} + c_2 U_{i,j-1,k} + c_3 U_{i+1,j-1,k} + c_4 U_{i-1,j,k} + \\ & c_5 U_{i,j,k} + c_6 U_{i+1,j,k} + c_7 U_{i-1,j+1,k} + c_8 U_{i,j+1,k} + c_9 U_{i+1,j+1,k} \end{aligned}$$

$$(4.2) \quad U_{i+s,j+t,k+1} = \sum_{u=1}^n \sum_{v=1}^m c_{uv} U_{i+u-1,j+v-1,k}$$

Note that equation(2.1) in section 2 is a special case for equation(4.1) and equation(2.2) is a special case for equation(4.2). We will obtain the explicit solution under the assumption of given $U_{i,j,0} = \Psi(i, j)$.

Theorem 4.1. *Let*

$$\begin{aligned} U_{i,j,k+1} = & c_1 U_{i-1,j-1,k} + c_2 U_{i,j-1,k} + c_3 U_{i+1,j-1,k} + \\ & c_4 U_{i-1,j,k} + c_5 U_{i,j,k} + c_6 U_{i+1,j,k} + \\ & c_7 U_{i-1,j+1,k} + c_8 U_{i,j+1,k} + c_9 U_{i+1,j+1,k} \end{aligned}$$

with $i, j \in Z, k \in N \cup \{0\}$. Given $U_{i,j,0} = \Psi(i, j)$, then its solution is of the form

$$U_{i,j,k} = \sum_{s_r \in I} \binom{k}{s_1, s_2, \dots, s_9} \Psi(i-a, j-b) \prod_{r=1}^9 c_r^{s_r},$$

where $I = \{s_1, s_2, \dots, s_9 | 0 \leq s_r \leq k, \sum_{r=1}^9 s_r = k\}$, $a = s_1 - s_3 + s_4 - s_6 + s_7 - s_9$, and $b = s_1 + s_2 + s_3 - s_7 - s_8 - s_9$.

Proof. Let

$$h(x, y, z) = \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \sum_{k=0}^{\infty} U_{i,j,k} x^i y^j z^k.$$

Then we have

$$\begin{aligned}
\sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \sum_{k=0}^{\infty} U_{i,j,k+1} x^i y^j z^k &= c_1 \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \sum_{k=0}^{\infty} U_{i-1,j-1,k} x^i y^j z^k + \\
&c_2 \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \sum_{k=0}^{\infty} U_{i,j-1,k} x^i y^j z^k + \\
&c_3 \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \sum_{k=0}^{\infty} U_{i+1,j-1,k} x^i y^j z^k + \\
&c_4 \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \sum_{k=0}^{\infty} U_{i-1,j,k} x^i y^j z^k + \\
&c_5 \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \sum_{k=0}^{\infty} U_{i,j,k} x^i y^j z^k + \\
&c_6 \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \sum_{k=0}^{\infty} U_{i+1,j,k} x^i y^j z^k + \\
&c_7 \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \sum_{k=0}^{\infty} U_{i-1,j+1,k} x^i y^j z^k + \\
&c_8 \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \sum_{k=0}^{\infty} U_{i,j+1,k} x^i y^j z^k + \\
&c_9 \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \sum_{k=0}^{\infty} U_{i+1,j+1,k} x^i y^j z^k,
\end{aligned}$$

$$\begin{aligned}
\frac{1}{z} [h(x, y, z) - \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \Psi(i, j) x^i y^j] &= (c_1 x y + c_2 y + c_3 x^{-1} y + c_4 x + c_5 + c_6 x^{-1} + \\
&c_7 x y^{-1} + c_8 y^{-1} + c_9 x^{-1} y^{-1}) h(x, y, z),
\end{aligned}$$

$$\begin{aligned}
\sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \Psi(i, j) x^i y^j &= [1 - (c_1 x y + c_2 y + c_3 x^{-1} y + c_4 x + c_5 + \\
&c_6 x^{-1} + c_7 x y^{-1} + c_8 y^{-1} + c_9 x^{-1} y^{-1}) z] h(x, y, z),
\end{aligned}$$

$$\begin{aligned}
 h(x, y, z) &= \frac{\sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \Psi(i, j) x^i y^j}{1 - (c_1 xy + c_2 y + c_3 x^{-1} y + c_4 x + c_5 + c_6 x^{-1} + c_7 xy^{-1} + c_8 y^{-1} + c_9 x^{-1} y^{-1}) z} \\
 &= \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \Psi(i, j) x^i y^j \sum_{k=0}^{\infty} (c_1 xy + c_2 y + c_3 x^{-1} y + c_4 x + c_5 + \\
 &\quad c_6 x^{-1} + c_7 xy^{-1} + c_8 y^{-1} + c_9 x^{-1} y^{-1})^k z^k \\
 &= \sum_{k=0}^{\infty} \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \Psi(i, j) x^i y^j \sum_{s_r \in I} \binom{k}{s_1, s_2, \dots, s_9} c_1^{s_1} (xy)^{s_1} c_2^{s_2} y^{s_2} \dots c_9^{s_9} (xy)^{-s_9} z^k \\
 &= \sum_{k=0}^{\infty} \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \Psi(i, j) x^i y^j \sum_{s_r \in I} \binom{k}{s_1, s_2, \dots, s_9} \prod_{r=1}^9 c_r^{s_r} x^a y^b z^k \\
 &= \sum_{k=0}^{\infty} \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \sum_{s_r \in I} \binom{k}{s_1, s_2, \dots, s_9} \Psi(i, j) \prod_{r=1}^9 c_r^{s_r} x^{a+i} y^{b+j} z^k,
 \end{aligned}$$

where $I = \{s_1, s_2, \dots, s_9 | 0 \leq s_r \leq k, \sum_{r=1}^9 s_r = k\}$, $a = s_1 - s_3 + s_4 - s_6 + s_7 - s_9$, and $b = s_1 + s_2 + s_3 - s_7 - s_8 - s_9$. Let

$$a + i = u, \quad b + j = v.$$

Then we have

$$h(x, y, z) = \sum_{k=0}^{\infty} \sum_{u=-\infty}^{\infty} \sum_{v=-\infty}^{\infty} \sum_{s_r \in I} \binom{k}{s_1, s_2, \dots, s_9} \Psi(u - a, v - b) \prod_{r=1}^9 c_r^{s_r} x^u y^v z^k.$$

Hence the solution is as follows

$$U_{i,j,k} = \sum_{s_r \in I} \binom{k}{s_1, s_2, \dots, s_9} \Psi(i - a, j - b) \prod_{r=1}^9 c_r^{s_r},$$

□

Theorem 4.2. *Let*

$$U_{i+s,j+t,k+1} = \sum_{u=1}^n \sum_{v=1}^m c_{uv} U_{i+u-1, j+v-1, k}$$

with $i \in Z, j \in Z, k \in N \cup \{0\}$. Given $U_{i,j,0} = \Psi(i, j)$, then its solution is of the form

$$U_{i,j} = \sum_{s_{qr} \in I} \binom{k}{s_{11}, s_{12}, \dots, s_{1m}, \dots, s_{nm}} \Psi(i - a, j - b) \prod_{\substack{u=1,2,\dots,n, \\ v=1,2,\dots,m}} c_{uv}^{s_{uv}},$$

where $I = \{s_{11}, s_{12}, \dots, s_{1m}, \dots, s_{nm} | 0 \leq s_{uv} \leq j - b, \sum_{u=1}^n \sum_{v=1}^m s_{uv} = j - b\}$, $a = (s + 1)k - \sum_{u=1}^n \sum_{v=1}^m u s_{uv}$, and $b = (t + 1)k - \sum_{u=1}^n \sum_{v=1}^m v s_{uv}$.

Proof. Let

$$h(x, y, z) = \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \sum_{k=0}^{\infty} U_{i,j} x^i y^j z^k.$$

Then we have

$$\sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \sum_{k=0}^{\infty} U_{i+s, j+t, k+1} x^i y^j z^k = \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \sum_{k=0}^{\infty} \sum_{u=1}^n \sum_{v=1}^m c_{uv} U_{i+u-1, j+v-1, k} x^i y^j z^k,$$

$$\frac{1}{x^s y^t z} [h(x, y, z) - \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \Psi(i, j) x^i y^j] = \sum_{u=1}^n \sum_{v=1}^m c_{uv} x^{-(u-1)} y^{-(v-1)} h(x, y, z),$$

$$h(x, y) (1 - \sum_{u=1}^n \sum_{v=1}^m c_{uv} x^{s-(u-1)} y^{t-(v-1)} z) = \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \Psi(i, j) x^i y^j,$$

$$\begin{aligned} h(x, y) &= \frac{\sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \Psi(i, j) x^i y^j}{1 - \sum_{u=1}^n \sum_{v=1}^m c_{uv} x^{s-(u-1)} y^{t-(v-1)} z} \\ &= \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \Psi(i, j) x^i y^j \sum_{k=0}^{\infty} \left(\sum_{u=1}^n \sum_{v=1}^m c_{uv} x^{s-(u-1)} y^{t-(v-1)} \right)^k z^k \\ &= \sum_{k=0}^{\infty} \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \Psi(i, j) x^i y^j \left(\sum_{u=1}^n c_{u1} x^{s-(u-1)} y^t + c_{u2} x^{s-(u-1)} y^{t-1} + \dots + \right. \\ &\quad \left. c_{um} x^{s-(u-1)} y^{t-(m-1)} \right)^k z^k \\ &= \sum_{k=0}^{\infty} \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \Psi(i, j) x^i y^j (c_{11} x^s y^t + c_{12} x^s y^{t-1} + \dots + c_{1m} x^s y^{t-(m-1)} + \\ &\quad c_{21} x^{s-1} y^t + c_{22} x^{s-1} y^{t-1} + \dots + c_{2m} x^{s-1} y^{t-(m-1)} + \dots + \\ &\quad c_{n1} x^{s-(n-1)} y^t + c_{n2} x^{s-(n-1)} y^{t-1} + \dots + c_{nm} x^{s-(n-1)} y^{t-(m-1)})^k z^k \\ &= \sum_{k=0}^{\infty} \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \Psi(i, j) x^i y^j c_{11}^{s_{11}} x^{s s_{11}} y^{t s_{11}} c_{12}^{s_{12}} x^{s s_{12}} y^{(t-1) s_{12}} \dots c_{1m}^{s_{1m}} x^{s s_{1m}} y^{[t-(m-1)] s_{1m}} \\ &\quad c_{21}^{s_{21}} x^{(s-1) s_{21}} y^{t s_{21}} c_{22}^{s_{22}} x^{(s-1) s_{22}} y^{(t-1) s_{22}} \dots c_{2m}^{s_{2m}} x^{(s-1) s_{2m}} y^{[t-(m-1)] s_{2m}} \dots \\ &\quad c_{n1}^{s_{n1}} x^{[s-(n-1)] s_{n1}} y^{t s_{n1}} c_{n2}^{s_{n2}} x^{[s-(n-1)] s_{n2}} y^{(t-1) s_{n2}} \dots c_{nm}^{s_{nm}} x^{[s-(n-1)] s_{nm}} y^{[t-(m-1)] s_{nm}} z^k \\ &= \sum_{k=0}^{\infty} \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \Psi(i, j) x^i y^j \sum_{s_{qr} \in I} \binom{k}{s_{11}, s_{12}, \dots, s_{1m}, \dots, s_{nm}} \prod_{\substack{u=1, 2, \dots, n, \\ v=1, 2, \dots, m}} c_{uv}^{s_{uv}} x^a y^b z^k \\ &= \sum_{k=0}^{\infty} \sum_{i=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \sum_{s_r \in I} \binom{k}{s_{11}, s_{12}, \dots, s_{1m}, \dots, s_{nm}} \Psi(i, j) \prod_{\substack{u=1, 2, \dots, n, \\ v=1, 2, \dots, m}} c_{uv}^{s_{uv}} x^{a+i} y^{b+j} z^k, \end{aligned}$$

where

$$\begin{aligned}
 I &= \{s_{11}, s_{12}, \dots, s_{1m}, \dots, s_{nm} | 0 \leq s_{uv} \leq k, \sum_{u=1}^n \sum_{v=1}^m s_{uv} = k\}, \\
 a &= s \sum_{v=1}^m s_{1v} + (s-1) \sum_{v=1}^m s_{2v} + \dots + [s - (n-1)] \sum_{v=1}^m s_{nv} \\
 &= \sum_{u=1}^n [s - (u-1)] \sum_{v=1}^m s_{uv} \\
 &= \sum_{u=1}^n \sum_{v=1}^m [s - (u-1)] s_{uv} \\
 &= \sum_{u=1}^n \sum_{v=1}^m [(s+1) - u] s_{uv} \\
 &= (s+1)j - \sum_{u=1}^n \sum_{v=1}^m u s_{uv}, \\
 b &= \sum_{u=1}^n \sum_{v=1}^m [t - (v-1)] s_{uv} \\
 &= \sum_{u=1}^n \sum_{v=1}^m [(t+1) - v] s_{uv} \\
 &= (t+1)k - \sum_{u=1}^n \sum_{v=1}^m v s_{uv}.
 \end{aligned}$$

Let

$$a + i = l, \quad b + j = p.$$

Then we have

$$h(x, y, z) = \sum_{k=0}^{\infty} \sum_{l=-\infty}^{\infty} \sum_{p=-\infty}^{\infty} \sum_{s_{qr} \in I} \binom{k}{s_{11}, s_{12}, \dots, s_{1m}, \dots, s_{nm}} \Psi(l-a, p-b) \prod_{\substack{u=1,2,\dots,n, \\ v=1,2,\dots,m}} c_{uv}^{s_{uv}} x^l y^p z^k.$$

So the solution is of the form

$$U_{i,j} = \sum_{s_{qr} \in I} \binom{k}{s_{11}, s_{12}, \dots, s_{1m}, \dots, s_{nm}} \Psi(i-a, j-b) \prod_{\substack{u=1,2,\dots,n, \\ v=1,2,\dots,m}} c_{uv}^{s_{uv}},$$

□

Now, we extend eq(4.2) to n-dimensional case as follows, and find out it's exact solution.

Lemma 4.3.

$$\left(\sum_{u_1=1}^{k_1} \sum_{u_2=1}^{k_2} \cdots \sum_{u_{n-1}=1}^{k_{n-1}} c_{u_1 u_2 \dots u_{n-1}} \prod_{i=1}^{n-1} x_i^{s_i - (u_i - 1)} \right)^j = \sum_{r_{u_1 u_2 \dots u_{n-1}} \in I} \binom{j}{r_{11\dots 1}, r_{11\dots 2}, \dots, r_{11\dots k_{n-1}}, \dots, r_{k_1 k_2 \dots k_{n-1}}} \prod_{\substack{u_1=1, 2, \dots, k_1. \\ u_2=1, 2, \dots, k_2. \\ \vdots \\ u_{n-1}=1, 2, \dots, k_{n-1}.}} c_{u_1 u_2 \dots u_{n-1}}^{r_{u_1 u_2 \dots u_{n-1}}} \prod_{i=1}^{n-1} x_i^{a_i},$$

where

$$I = \{ r_{11\dots 1}, r_{11\dots 2}, \dots, r_{11\dots k_{n-1}}, \dots, r_{k_1 k_2 \dots k_{n-1}} \mid 0 \leq r_{u_1 u_2 \dots u_{n-1}} \leq j, \sum_{u_1=1}^{k_1} \sum_{u_2=1}^{k_2} \cdots \sum_{u_{n-1}=1}^{k_{n-1}} r_{u_1 u_2 \dots u_{n-1}} = j \},$$

$$a_i = (s_i + 1)j - \sum_{u_1=1}^{k_1} \sum_{u_2=1}^{k_2} \cdots \sum_{u_{n-1}=1}^{k_{n-1}} u_i r_{u_1 u_2 \dots u_{n-1}}, \forall i = 1, 2, \dots, n-1.$$

Proof.

$$\begin{aligned}
 & \left(\sum_{u_1=1}^{k_1} \sum_{u_2=1}^{k_2} \cdots \sum_{u_{n-1}=1}^{k_{n-1}} c_{u_1 u_2 \dots u_{n-1}} \prod_{i=1}^{n-1} x_i^{s_i - (u_i - 1)} \right)^j \\
 = & \sum_{u_1=1}^{k_1} \sum_{u_2=1}^{k_2} \cdots \sum_{u_{n-2}=1}^{k_{n-2}} \left(c_{u_1 u_2 \dots u_{n-2} 1} \prod_{i=1}^{n-2} x_i^{s_i - (u_i - 1)} x_{n-1}^{s_{n-1}} + c_{u_1 u_2 \dots u_{n-2} 2} \prod_{i=1}^{n-2} x_i^{s_i - (u_i - 1)} x_{n-1}^{s_{n-1} - 1} + \right. \\
 & \left. \cdots + c_{u_1 u_2 \dots u_{n-2} k_{n-1}} \prod_{i=1}^{n-2} x_i^{s_i - (u_i - 1)} x_{n-1}^{s_{n-1} - (k_{n-1} - 1)} \right)^j \\
 = & \left[c_{11 \dots 11} \prod_{i=1}^{n-1} x_i^{s_i} + c_{11 \dots 12} \prod_{i=1}^{n-2} x_i^{s_i - 1} x_{n-1}^{s_{n-1} - 1} + \cdots + c_{11 \dots 1 k_{n-1}} \prod_{i=1}^{n-2} x_i^{s_i - 1} x_{n-1}^{s_{n-1} - (k_{n-1} - 1)} + \right. \\
 & \left. \cdots + c_{k_1 k_2 \dots k_{n-1}} \prod_{i=1}^{n-1} x_i^{s_i - (k_i - 1)} \right]^j \\
 = & \sum_{r_{u_1 u_2 \dots u_{n-1}} \in I} \binom{j}{r_{11 \dots 11}, r_{11 \dots 12}, \dots, r_{11 \dots 1 k_{n-1}}, \dots, r_{k_1 k_2 \dots k_{n-1}}} \left(c_{11 \dots 11} \prod_{i=1}^{n-1} x_i^{s_i} \right)^{r_{11 \dots 11}} \\
 & \left(c_{11 \dots 12} \prod_{i=1}^{n-2} x_i^{s_i - 1} x_{n-1}^{s_{n-1} - 1} \right)^{r_{11 \dots 12}} \left(c_{11 \dots 1 k_{n-1}} \prod_{i=1}^{n-2} x_i^{s_i - 1} x_{n-1}^{s_{n-1} - (k_{n-1} - 1)} \right)^{r_{11 \dots 1 k_{n-1}}} \\
 & \cdots \left(c_{k_1 k_2 \dots k_{n-1}} \prod_{i=1}^{n-1} x_i^{s_i - (k_i - 1)} \right)^{r_{k_1 k_2 \dots k_{n-1}}} \\
 = & \sum_{r_{u_1 u_2 \dots u_{n-1}} \in I} \binom{j}{r_{11 \dots 11}, r_{11 \dots 12}, \dots, r_{11 \dots 1 k_{n-1}}, \dots, r_{k_1 k_2 \dots k_{n-1}}} \prod_{\substack{u_1=1, 2, \dots, k_1. \\ u_2=1, 2, \dots, k_2. \\ \vdots \\ u_{n-1}=1, 2, \dots, k_{n-1}.}} \left(c_{u_1 u_2 \dots u_{n-1}} \prod_{i=1}^{n-1} x_i^{a_i} \right)^{r_{u_1 u_2 \dots u_{n-1}}}
 \end{aligned}$$

where

$$\begin{aligned}
I &= \{r_{11\dots 1}, r_{11\dots 12}, \dots, r_{11\dots 1k_{n-1}}, \dots, r_{k_1 k_2 \dots k_{n-1}}\} \\
0 &\leq r_{u_1 u_2 \dots u_{n-1}} \leq j, \sum_{u_1=1}^{k_1} \sum_{u_2=1}^{k_2} \dots \sum_{u_{n-1}=1}^{k_{n-1}} r_{u_1 u_2 \dots u_{n-1}} = j\}, \\
\forall i &= 1, 2, \dots, n-1, \\
a_i &= \sum_{u_1=1}^{k_1} \sum_{u_2=1}^{k_2} \dots \sum_{u_{n-1}=1}^{k_{n-1}} [s_i - (u_i - 1)] r_{u_1 u_2 \dots u_{n-1}} \\
&= \sum_{u_1=1}^{k_1} \sum_{u_2=1}^{k_2} \dots \sum_{u_{n-1}=1}^{k_{n-1}} [(s_i + 1) - u_i] r_{u_1 u_2 \dots u_{n-1}} \\
&= (s_i + 1)j - \sum_{u_1=1}^{k_1} \sum_{u_2=1}^{k_2} \dots \sum_{u_{n-1}=1}^{k_{n-1}} u_i r_{u_1 u_2 \dots u_{n-1}}.
\end{aligned}$$

□

Theorem 4.4. *Let*

$$U_{e_1+s_1, e_2+s_2, \dots, e_{n-1}+s_{n-1}, e_n+1} = \sum_{u_1=1}^{k_1} \sum_{u_2=1}^{k_2} \dots \sum_{u_{n-1}=1}^{k_{n-1}} c_{u_1 u_2 \dots u_{n-1}} U_{e_1+u_1-1, e_2+u_2-1, \dots, e_{n-1}+u_{n-1}-1, e_n}$$

be a linear difference equation in e_1, e_2, \dots, e_n , where $e_i \in Z$ for $i \in \{1, 2, \dots, n-1\}$, $e_n \in N \cup \{0\}$. Given $U_{e_1, e_2, \dots, e_{n-1}, 0} = \Psi(e_1, e_2, \dots, e_{n-1})$, then its solution is of the form

$$\begin{aligned}
U_{e_1, e_2, \dots, e_n} &= \sum_{r_{u_1 u_2 \dots u_{n-1}} \in I} \left(\begin{matrix} j \\ r_{11\dots 1}, r_{11\dots 2}, \dots, r_{11\dots k_{n-1}}, \dots, r_{k_1 k_2 \dots k_{n-1}} \end{matrix} \right) \\
&\quad \Psi(e_1 - a_1, e_2 - a_2, \dots, e_{n-1} - a_{n-1}) \prod_{\substack{u_1=1, 2, \dots, k_1. \\ u_2=1, 2, \dots, k_2. \\ u_{n-1}=1, 2, \dots, k_{n-1}.}} c_{u_1 u_2 \dots u_{n-1}},
\end{aligned}$$

where

$$\begin{aligned}
I &= \{r_{11\dots 1}, r_{11\dots 2}, \dots, r_{11\dots k_{n-1}}, \dots, r_{k_1 k_2 \dots k_{n-1}}\} \\
0 &\leq r_{u_1 u_2 \dots u_{n-1}} \leq j, \sum_{u_1=1}^{k_1} \sum_{u_2=1}^{k_2} \dots \sum_{u_{n-1}=1}^{k_{n-1}} r_{u_1 u_2 \dots u_{n-1}} = j\}, \\
a_i &= (s_i + 1)j - \sum_{u_1=1}^{k_1} \sum_{u_2=1}^{k_2} \dots \sum_{u_{n-1}=1}^{k_{n-1}} u_i r_{u_1 u_2 \dots u_{n-1}}, \forall i = 1, 2, \dots, n-1.
\end{aligned}$$

Proof. Let

$$\begin{aligned}
\vec{x} &= (x_1, x_2, \dots, x_n), \\
h(\vec{x}) &= \sum_{e_1=-\infty}^{\infty} \sum_{e_2=-\infty}^{\infty} \dots \sum_{e_{n-1}=-\infty}^{\infty} \sum_{e_n=0}^{\infty} U_{e_1, e_2, \dots, e_n} \prod_{i=1}^n x_i^{e_i}.
\end{aligned}$$

Then we have

$$\begin{aligned}
 & \sum_{e_1=-\infty}^{\infty} \sum_{e_2=-\infty}^{\infty} \cdots \sum_{e_{n-1}=-\infty}^{\infty} \sum_{e_n=0}^{\infty} U_{e_1+s_1, e_2+s_2, \dots, e_{n-1}+s_{n-1}, e_n+1} \prod_{i=1}^n x_i^{e_i} \\
 = & \sum_{e_1=-\infty}^{\infty} \sum_{e_2=-\infty}^{\infty} \cdots \sum_{e_{n-1}=-\infty}^{\infty} \sum_{e_n=0}^{\infty} \sum_{u_1=1}^{k_1} \sum_{u_2=1}^{k_2} \cdots \sum_{u_{n-1}=1}^{k_{n-1}} \\
 & c_{u_1 u_2 \dots u_{n-1}} U_{e_1+u_1-1, e_2+u_2-1, \dots, e_{n-1}+u_{n-1}-1, e_n} \prod_{i=1}^n x_i^{e_i}, \\
 & \frac{1}{\prod_{i=1}^{n-1} x_i^{s_i} x_n} [h(\vec{x}) - \sum_{e_1=-\infty}^{\infty} \sum_{e_2=-\infty}^{\infty} \cdots \sum_{e_{n-1}=-\infty}^{\infty} \Psi(e_1, e_2, \dots, e_{n-1}) \prod_{i=1}^{n-1} x_i^{e_i}] \\
 = & \sum_{u_1=1}^{k_1} \sum_{u_2=1}^{k_2} \cdots \sum_{u_{n-1}=1}^{k_{n-1}} c_{u_1 u_2 \dots u_{n-1}} \prod_{i=1}^{n-1} x_i^{-(u_i-1)} h(\vec{x}),
 \end{aligned}$$

$$\begin{aligned}
 & h(\vec{x}) \left(1 - \sum_{u_1=1}^{k_1} \sum_{u_2=1}^{k_2} \cdots \sum_{u_{n-1}=1}^{k_{n-1}} c_{u_1 u_2 \dots u_{n-1}} \prod_{i=1}^{n-1} x_i^{s_i - (u_i-1)} x_n \right) \\
 = & \sum_{e_1=-\infty}^{\infty} \sum_{e_2=-\infty}^{\infty} \cdots \sum_{e_{n-1}=-\infty}^{\infty} \Psi(e_1, e_2, \dots, e_{n-1}) \prod_{i=1}^{n-1} x_i^{e_i},
 \end{aligned}$$

$$\begin{aligned}
 h(\vec{x}) &= \frac{\sum_{e_1=-\infty}^{\infty} \sum_{e_2=-\infty}^{\infty} \cdots \sum_{e_{n-1}=-\infty}^{\infty} \Psi(e_1, e_2, \dots, e_{n-1}) \prod_{i=1}^{n-1} x_i^{e_i}}{\left(1 - \sum_{u_1=1}^{k_1} \sum_{u_2=1}^{k_2} \cdots \sum_{u_{n-1}=1}^{k_{n-1}} c_{u_1 u_2 \dots u_{n-1}} \prod_{i=1}^{n-1} x_i^{s_i - (u_i-1)} x_n \right)} \\
 &= \sum_{e_1=-\infty}^{\infty} \sum_{e_2=-\infty}^{\infty} \cdots \sum_{e_{n-1}=-\infty}^{\infty} \Psi(e_1, e_2, \dots, e_{n-1}) \prod_{i=1}^{n-1} x_i^{e_i} \\
 & \quad \sum_{j=0}^{\infty} \left(\sum_{u_1=1}^{k_1} \sum_{u_2=1}^{k_2} \cdots \sum_{u_{n-1}=1}^{k_{n-1}} c_{u_1 u_2 \dots u_{n-1}} \prod_{i=1}^{n-1} x_i^{s_i - (u_i-1)} \right)^j x_n^j.
 \end{aligned}$$

Applying above lemma, we have

$$\begin{aligned}
h(\vec{x}) &= \sum_{j=0}^{\infty} \sum_{e_1=-\infty}^{\infty} \sum_{e_2=-\infty}^{\infty} \cdots \sum_{e_{n-1}=-\infty}^{\infty} \Psi(e_1, e_2, \dots, e_{n-1}) \prod_{i=1}^{n-1} x_i^{e_i} \\
&\quad \sum_{r_{u_1 u_2 \dots u_{n-1}} \in I} \binom{j}{r_{11\dots 1}, r_{11\dots 2}, \dots, r_{11\dots k_{n-1}}, \dots, r_{k_1 k_2 \dots k_{n-1}}} \\
&\quad \prod_{\substack{u_1=1,2,\dots,k_1. \\ u_2=1,2,\dots,k_2. \\ \vdots \\ u_{n-1}=1,2,\dots,k_{n-1}.}} C_{u_1 u_2 \dots u_{n-1}}^{r_{u_1 u_2 \dots u_{n-1}}} \prod_{i=1}^{n-1} x_i^{a_i} x_n^j \\
&= \sum_{j=0}^{\infty} \sum_{e_1=-\infty}^{\infty} \sum_{e_2=-\infty}^{\infty} \cdots \sum_{e_{n-1}=-\infty}^{\infty} \sum_{r_{u_1 u_2 \dots u_{n-1}} \in I} \binom{j}{r_{11\dots 1}, r_{11\dots 2}, \dots, r_{11\dots k_{n-1}}, \dots, r_{k_1 k_2 \dots k_{n-1}}} \\
&\quad \Psi(e_1, e_2, \dots, e_{n-1}) \prod_{\substack{u_1=1,2,\dots,k_1. \\ u_2=1,2,\dots,k_2. \\ \vdots \\ u_{n-1}=1,2,\dots,k_{n-1}.}} C_{u_1 u_2 \dots u_{n-1}}^{r_{u_1 u_2 \dots u_{n-1}}} \prod_{i=1}^{n-1} x_i^{a_i + e_i} x_n^j.
\end{aligned}$$

Let

$$a_i + e_i = b_i, \forall i = 1, 2, \dots, n-1.$$

Then we have

$$\begin{aligned}
h(\vec{x}) &= \sum_{j=0}^{\infty} \sum_{b_1=-\infty}^{\infty} \sum_{b_2=-\infty}^{\infty} \cdots \sum_{b_{n-1}=-\infty}^{\infty} \sum_{r_{u_1 u_2 \dots u_{n-1}} \in I} \binom{j}{r_{11\dots 1}, r_{11\dots 2}, \dots, r_{11\dots k_{n-1}}, \dots, r_{k_1 k_2 \dots k_{n-1}}} \\
&\quad \Psi(b_1 - a_1, b_2 - a_2, \dots, b_{n-1} - a_{n-1}) \prod_{\substack{u_1=1,2,\dots,k_1. \\ u_2=1,2,\dots,k_2. \\ \vdots \\ u_{n-1}=1,2,\dots,k_{n-1}.}} C_{u_1 u_2 \dots u_{n-1}}^{r_{u_1 u_2 \dots u_{n-1}}} \prod_{i=1}^{n-1} x_i^{b_i} x_n^j,
\end{aligned}$$

Thus the solution is of the form

$$\begin{aligned}
U_{e_1, e_2, \dots, e_n} &= \sum_{r_{u_1 u_2 \dots u_{n-1}} \in I} \binom{j}{r_{11\dots 1}, r_{11\dots 2}, \dots, r_{11\dots k_{n-1}}, \dots, r_{k_1 k_2 \dots k_{n-1}}} \\
&\quad \Psi(e_1 - a_1, e_2 - a_2, \dots, e_{n-1} - a_{n-1}) \prod_{\substack{u_1=1,2,\dots,k_1. \\ u_2=1,2,\dots,k_2. \\ \vdots \\ u_{n-1}=1,2,\dots,k_{n-1}.}} C_{u_1 u_2 \dots u_{n-1}}^{r_{u_1 u_2 \dots u_{n-1}}},
\end{aligned}$$

□

REFERENCES

- [1] Ronald E. Mickens, Difference Equations Theory and Application, 1990.
- [2] Gerald Berman and K. D. Fryer, Introduction To Combinatorics, 1972.
- [3] Sui Sun Cheng, Stability of Partial Difference Equations : An Introduction, 1997.
- [4] Sui Sun Cheng, A symbolic Cauculus for sequence, 1997.
- [5] Jin Yu Lin, Stablility Criteria for a discrete Reaction-Diffusion Equation, 1997.

[6] Yi Feng Lu, General Solution of a Partial Difference Equation, 1998.

DEPARTMENT OF COMPUTER SCIENCE AND INFORMATION ENGINEERING, NATIONAL TAIWAN UNIVERSITY, TAIPEI, TAIWAN, 10617, R.O.C.

Email address: chunkai.hwang@gmail.com

DEPARTMENT OF APPLIED MATHEMATICS, NATIONAL SUN YAT-SEN UNIVERSITY, KAOHSIUNG, TAIWAN, 80424, R.O.C.

Email address: ttlu@math.nsysu.edu.tw