

CORNER REPLACEMENT FOR MORITA CONTEXTS

RAPHAEL BENNETT-TENNENHAUS

ABSTRACT. We consider how Morita equivalences are compatible with the notion of a corner subring. Namely, we outline a canonical way to replace a corner subring of a given ring with one which is Morita equivalent, and look at how such an equivalence ascends.

We use the language of Morita contexts, and then specify these more general results. We work in the setting of rings with local units and unital bimodules, to extend the reach of potential applications, say to categories of functors.

1. INTRODUCTION

Celebrated results of Morita from [15, §3] characterise the existence of an equivalence $R\text{-Mod} \simeq S\text{-Mod}$ between module categories in terms of the existence of a pair ${}_R N_S, {}_S L_R$ of bimodules subject to compatibility conditions. These conditions include the existence of bimodule homomorphisms of the form $\theta: N \otimes L \rightarrow R$ and $\zeta: L \otimes N \rightarrow S$ satisfying certain associativity conditions. A *Morita context*, also known as *pre-equivalence data*, refers to the tuple $(R, S; N, L; \theta, \zeta)$. We note, in particular, that such a context defines an equivalence of the form above when the maps θ, ζ are surjective; see for example Theorem 2.1. The results of Morita are presented in an illustrious book by Bass [2, Chapters II, III].

Morita contexts have since been studied in various other generalities. In particular, so-called *Morita theory* has been developed by Ánh and Márki [1] to unital modules over rings with local units, and more generally, to categories of modules over idempotent rings in work of García and Simón [9]. An idea we exploit here is that any Morita context as above gives rise to a *ring of Morita context*, a ring of matrices denoted $\begin{bmatrix} R & N \\ L & S \end{bmatrix}$ whose ring structure is canonically defined by information from the context.

The general purpose of this note is to record the statement and a proof of Proposition 1.1 below, which the author suspects was already well-known ‘folklore’.

Proposition 1.1. *Let R and S be unital associative rings and let $e \in R$ be an idempotent.*

If ${}_e R_e M_S$ and ${}_S N_{e R_e}$ are bimodules defining a Morita equivalence $e R e \sim S$, then

$$\begin{pmatrix} (1-e)R(1-e) & (1-e)Re \otimes M \\ (1-e)Re & M \end{pmatrix}, \quad \begin{pmatrix} (1-e)R(1-e) & (1-e)Re \\ N \otimes (1-e)Re & N \end{pmatrix}$$

are bimodules defining a Morita equivalence $R \sim \begin{bmatrix} (1-e)R(1-e) & (1-e)Re \otimes M \\ N \otimes (1-e)Re & S \end{bmatrix}$.

Note that the rings R and S are unital, however, in Theorem 3.14, we work in the setting of Morita contexts for (possibly non-unital) rings with local units, and unital bimodules over them. Working in this more general setting extends the reach of potential applications.

2020 *Mathematics Subject Classification.* Primary 16D90.

Key words and phrases. Morita context, Morita equivalence, generalised matrix ring.

2. BACKGROUND ON MORITA CONTEXTS

2.1. Rings with local units. [1, Definition 1] By a *ring* we will mean one which is associative, but non-unital, meaning we do not require it to have a multiplicative identity.

If R is a ring we write $\mu_R: R \otimes_R R \rightarrow R$ for the universal R - R -bimodule homomorphism given by the (balanced) multiplication map $R \times R \rightarrow R$. If M is a left module over R we likewise write $\lambda_M^R: R \otimes_R M \rightarrow M$ for the universal left R -module homomorphism given by the left action map $R \times M \rightarrow M$. Similarly define $\rho_M^S: M \otimes_S S \rightarrow M$ if M is a right S -module for some ring S , meaning that M is an R - S bimodule if and only if one has $\lambda_M^R(\mathbf{1}_R \otimes \rho_M^S) = \rho_M^S(\lambda_M^R \otimes \mathbf{1}_S)$ for a pair of such actions.

A ring R is said to *have local units* if every finite subset of R is contained in a (unital) subring of the form eRe for some $e = e^2 \in R$. For a left module M over a ring R with local units, we say M is *unital* if λ_M^R is surjective. Similarly if S has local units then we call a right S -module M unital if ρ_M^S is surjective, and we call a bimodule unital if it is both unital as a left module and as a right module.

2.2. Matrix notation. In what remains in the article we use the following notation.

Let $(X_{ij} \mid m, n)$ be a collection of additive groups X_{ij} for each $i = 1, \dots, m$ and each $j = 1, \dots, n$. Throughout we consider the following group under component-wise addition,

$$\left(\begin{array}{ccc} X_{11} & \cdots & X_{1m} \\ \vdots & \ddots & \vdots \\ X_{l1} & \cdots & X_{lm} \end{array} \right) = \left\{ \left(\begin{array}{ccc} x_{11} & \cdots & x_{1m} \\ \vdots & \ddots & \vdots \\ x_{l1} & \cdots & x_{lm} \end{array} \right) \mid x_{ij} \in X_{ij} \text{ for each } i \text{ and } j \right\},$$

which is isomorphic to the direct sum of the X_{ij} .

Consider now collections $(X_{ik} \mid l, m)$, $(Y_{kj} \mid m, n)$ and $(Z_{ij} \mid l, n)$ of additive groups. Suppose additionally there are rings A_1, \dots, A_l such that each X_{ik} is a right A_k -module and each Y_{kj} is a left A_k -module. Furthermore, suppose there is a collection of group homomorphisms $\alpha_{ikj}: X_{ik} \otimes_{A_k} Y_{kj} \rightarrow Z_{ij}$ uniquely defined by A_k -balanced maps of the form $X_{ik} \times Y_{kj} \rightarrow Z_{ij}$. Combining the functions α_{ikj} we use the additive structure of the groups involved to consider a biadditive map $|\alpha_{ikj}|$, denoted in detail by

$$\left| \begin{array}{ccc} \alpha_{1k1} & \cdots & \alpha_{1km} \\ \vdots & \ddots & \vdots \\ \alpha_{lk1} & \cdots & \alpha_{lkm} \end{array} \right| : \left(\begin{array}{ccc} X_{11} & \cdots & X_{1m} \\ \vdots & \ddots & \vdots \\ X_{l1} & \cdots & X_{lm} \end{array} \right) \times \left(\begin{array}{ccc} Y_{11} & \cdots & Y_{1n} \\ \vdots & \ddots & \vdots \\ Y_{m1} & \cdots & Y_{mn} \end{array} \right) \rightarrow \left(\begin{array}{ccc} Z_{11} & \cdots & Z_{1n} \\ \vdots & \ddots & \vdots \\ Z_{l1} & \cdots & Z_{ln} \end{array} \right)$$

which is defined by the assignment $((x_{ik}), (y_{kj})) \mapsto (z_{ij})$ where $z_{ij} = \sum_k \alpha_{ikj}(x_{ik} \otimes y_{kj})$.

In this notation we can consider some familiar situations.

- In §2.3 we take $l = m = n$ and $X_{ij} = Y_{ij} = Z_{ij}$ where each X_{ii} is a ring so that X_{ij} is an X_{ii} - X_{jj} -bimodule, such that the map $|\alpha_{ikj}|$ defines a ring multiplication.
- In §2.4 we take $l = m$, $n = 1$, $Y_{ik} = Z_{ik}$ so that $|\alpha_{ikj}|$ defines a left module action over a ring from the item above. Dually one can define right modules.
- In §2.5, §2.6, §2.7 and §2.8 we similarly use the notation above to consider module homomorphisms, balanced maps between modules, bimodules and associativity conditions, respectively. This sets up notation for the remainder of the article.

2.3. Matrix ring of a generalised Morita context. By a *generalised Morita context* we will mean a triple $(A_i; M_{ij}; \varphi_{ikj})$ where the following conditions hold.

- A_i ($i = 1, \dots, n$) are associative rings.
- M_{ij} ($i, j = 1, \dots, n$) are A_i - A_j -bimodules, where $M_{ii} = A_i$.
- $\varphi_{ikj}: M_{ik} \otimes_{A_k} M_{kj} \rightarrow M_{ij}$ ($i, j, k = 1, \dots, n$) are A_i - A_j -bimodule maps, where

$$\varphi_{kkj} = \lambda_{M_{kj}}^{A_k}, \quad \varphi_{kkk} = \mu_{A_k}, \quad \varphi_{ikk} = \rho_{M_{ik}}^{A_k}.$$

- $\varphi_{ihj}(\mathbf{1} \otimes \varphi_{hjk}) = \varphi_{ikj}(\varphi_{ihk} \otimes \mathbf{1})$ ($h, i, j, k = 1, \dots, n$) as maps $M_{ih} \otimes M_{hk} \otimes M_{kj} \rightarrow M_{ij}$.

The equations in the last item above are referred to as the *mixed associativity conditions*. Consequently, as in §2.3, with respect to $(A_i; M_{ij}; \varphi_{ikj})$ there is a *generalised matrix ring*

$$[A_i, M_{ij}] = \begin{bmatrix} A_1 & \cdots & M_{1n} \\ \vdots & \ddots & \vdots \\ M_{n1} & \cdots & A_n \end{bmatrix} \text{ with multiplication map } |\varphi_{ikj}| = \begin{bmatrix} \varphi_{1k1} & \cdots & \varphi_{1kn} \\ \vdots & \ddots & \vdots \\ \varphi_{nk1} & \cdots & \varphi_{nkn} \end{bmatrix}.$$

Note that if each ring A_i has local units and each bimodule M_{ij} is unital, then it is straightforward to see that $[A_i, M_{ij}]$ has local units. Morita contexts are usually considered as above, but for $n = 2$; see for example [9, p. 41]. The following result of Ánh and Márki [1] is part of a Morita theory developed for rings with local units.

Theorem 2.1. [1, Theorem 2.2] *Let $(R, S; N, L; \theta, \zeta)$ be a Morita context where R and S have local units and ${}_R N_S$ and ${}_S L_R$ are unital, and let $R\text{-Mod}$ and $S\text{-Mod}$ be the categories of unital left modules over R and S respectively.*

Then $N \otimes_S -: S\text{-Mod} \rightarrow R\text{-Mod}$ and $L \otimes_R -: R\text{-Mod} \rightarrow S\text{-Mod}$ are mutually quasi-inverse equivalences if and only if the bimodule homomorphisms θ and ζ are surjective.

2.4. Modules over matrix rings. Let $(A_i; M_{ij}; \varphi_{ikj})$ be a generalised Morita context in the sense of §2.3. By a *column collection* for $(A_i; M_{ij}; \varphi_{ikj})$ we mean an n -tuple $P_{\mathbf{c}} = (P_1, \dots, P_n)$ of left A_i -modules P_i ($i = 1, \dots, n$) where the following conditions hold.

- $\beta_{ik}: M_{ik} \otimes_{A_k} P_k \rightarrow P_i$ ($i, k = 1, \dots, n$) are left A_i -module homomorphisms, where $\beta_{ii} = \lambda_{P_i}^{A_i}$ for each i , which define the biadditive map $|\beta_{ik}|$, denoted in detail by

$$\begin{bmatrix} \beta_{1k} \\ \vdots \\ \beta_{nk} \end{bmatrix} : \begin{bmatrix} A_1 & \cdots & M_{1n} \\ \vdots & \ddots & \vdots \\ M_{n1} & \cdots & A_n \end{bmatrix} \times \begin{pmatrix} P_1 \\ \vdots \\ P_n \end{pmatrix} \rightarrow \begin{pmatrix} P_1 \\ \vdots \\ P_n \end{pmatrix}$$

- $\beta_{ih}(\varphi_{ikh} \otimes \mathbf{1}) = \beta_{ik}(\mathbf{1} \otimes \beta_{kh})$ ($h = 1, \dots, n$) as maps $M_{ik} \otimes M_{kh} \otimes P_h \rightarrow P_i$.

These conditions imply that the modules (P_1, \dots, P_n) together with the (left) *structure maps* β_{ik} define an object in the category of (left) *representations* of the given generalised Morita context; see [3, §2, (2.1)]. In other words, $P_{\mathbf{c}}$ together with the map $|\beta_{ik}|$ defines a left module over the generalised matrix ring $[A_i, M_{ij}]$.

By a result of Chen and Liu [3, Proposition 2.1], which generalises a result of Green [10, Theorem 1.5], if A_1, \dots, A_n are unital then the correspondence above defines a categorical equivalence. In [3, §2] these authors consider right $[A_i, M_{ij}]$ -modules, constructed dually to the column collection $P_{\mathbf{c}}$, which we refer to as a *row collection*.

2.5. Module homomorphisms over matrix rings. Let $(A_i; M_{ij}; \varphi_{ikj})$ be a generalised Morita context in the sense of §2.3. As in §2.4, let $P_{\mathbf{c}} = (P_1, \dots, P_n)$ and $P'_{\mathbf{c}} = (P'_1, \dots, P'_n)$ be column collections for $(A_i; M_{ij}; \varphi_{ikj})$, corresponding to left $[A_i; M_{ij}]$ -modules via left structure maps $\beta_{ik}: M_{ik} \otimes_{A_k} P_k \rightarrow P_i$ and $\beta'_{ik}: M_{ik} \otimes_{A_k} P'_k \rightarrow P'_i$ respectively. By a *column morphism* we mean a n -tuple $f_{\mathbf{c}} = (f_1, \dots, f_n)$ where the following conditions hold.

- $f_i: N_i \rightarrow N'_i$ ($i = 1, \dots, n$) are left A_i -module homomorphisms.
- $f_i \beta_{ij} = \beta'_{ij}(\mathbf{1} \otimes f_j)$ ($i, j = 1, \dots, n$) as maps $M_{ij} \otimes P_j \rightarrow P'_i$.

Then, as in §2.4, by following dual arguments in [3, §2] the tuple (f_1, \dots, f_n) defines an $[A_i; M_{ij}]$ -module homomorphism $f_{\mathbf{c}}: P_{\mathbf{c}} \rightarrow P'_{\mathbf{c}}$ by $f_{\mathbf{c}}(p_1, \dots, p_n) = (f_1(p_1), \dots, f_n(p_n))$.

Following [3, §2] directly gives a dual construction of a right $[A_i; M_{ij}]$ -module homomorphism between row collections, which we refer to as a *row morphism*.

2.6. Balanced homomorphisms over matrix rings. As in §2.3–2.4, for a generalised Morita context $(A_i; M_{ij}; \varphi_{ikj})$, fix a row collection $P_{\mathbf{r}} = (P_1, \dots, P_n)$ with respect to right structure maps $\beta_{ik}: P_i \otimes M_{ik} \rightarrow P_k$ and fix a column collection $Q_{\mathbf{c}} = (Q_1, \dots, Q_n)$ with respect to left structure maps $\gamma_{kj}: M_{kj} \otimes Q_j \rightarrow Q_k$, where the following conditions hold.

- Z is an additive group.
- $\alpha_k: P_k \otimes_{A_k} Q_k \rightarrow Z$ ($k = 1, \dots, n$) are additive maps defining a biadditive map

$$|\alpha_k|: \left(P_1 \quad \cdots \quad P_n \right) \times \begin{pmatrix} Q_1 \\ \vdots \\ Q_n \end{pmatrix} \rightarrow Z \quad ((p_1, \dots, p_n), (q_1, \dots, q_n)) \mapsto \sum_k \alpha_k(p_k \otimes q_k)$$

- $\alpha_j(\mathbf{1} \otimes \gamma_{ij}) = \alpha_i(\beta_{ij} \otimes \mathbf{1})$ ($i, j = 1, \dots, n$) as maps $P_i \otimes M_{ij} \otimes Q_j \rightarrow Z$.

Then $|\alpha_k|$ is $[A_i; M_{ij}]$ -balanced, so factors by an additive map $|\underline{\alpha}_k|: P_{\mathbf{r}} \otimes_{[A_i; M_{ij}]} Q_{\mathbf{c}} \rightarrow Z$.

2.7. Matrix bimodules over matrix rings. Let $(A_i; M_{ij}; \varphi_{ikj}), (A'_i; M'_{ij}; \varphi'_{ikj})$ be generalised Morita contexts. Fix additive groups T_{ij} where the following holds.

- $T_{\mathbf{c}j} = (T_{1j}, \dots, T_{nj})$ ($j = 1, \dots, n$) are column collections for $(A_i; M_{ij}; \varphi_{ikj})$ with respect to left structure maps $\gamma_{ikj}: M_{ik} \otimes T_{kj} \rightarrow T_{ij}$, summarised by

$$\begin{array}{c} \left| \begin{array}{c} \gamma_{1kj} \\ \vdots \\ \gamma_{nkj} \end{array} \right| : \begin{bmatrix} A_1 & \cdots & M_{1n} \\ \vdots & \ddots & \vdots \\ M_{n1} & \cdots & A_n \end{bmatrix} \times \begin{pmatrix} T_{1j} \\ \vdots \\ T_{nj} \end{pmatrix} \rightarrow \begin{pmatrix} T_{1j} \\ \vdots \\ T_{nj} \end{pmatrix}, \\ ((m_{ik}), (t_{kj})) \mapsto (\sum_k \gamma_{ikj}(m_{ik} \otimes t_{kj})). \end{array}$$

- $T_{\mathbf{r}i} = (T_{i1}, \dots, T_{in})$ ($i = 1, \dots, n$) are row collections for $(A'_i; M'_{ij}; \varphi'_{ikj})$ with respect to right structure maps $\beta_{ikj}: T_{ik} \otimes M'_{kj} \rightarrow T_{ij}$, summarised by

$$\left| \begin{array}{ccc} \beta_{ik1} & \cdots & \beta_{ikn} \end{array} \right| : \begin{pmatrix} T_{i1} & \cdots & T_{in} \end{pmatrix} \times \begin{bmatrix} A'_1 & \cdots & M'_{1n} \\ \vdots & \ddots & \vdots \\ M'_{n1} & \cdots & A'_n \end{bmatrix} \rightarrow \begin{pmatrix} T_{i1} & \cdots & T_{in} \end{pmatrix}, \\ ((t_{ik}), (m'_{kj})) \mapsto (\sum_k \beta_{ikj}(t_{ik} \otimes m'_{kj})).$$

- $\beta_{ikj}(\gamma_{ihk} \otimes \mathbf{1}) = \gamma_{ihj}(\mathbf{1} \otimes \beta_{hkj})$ ($h, i, j, k = 1, \dots, n$) as maps $M_{ih} \otimes T_{hk} \otimes M'_{kj} \rightarrow T_{ij}$.

These conditions imply that the collection (T_{ij}) defines an $[A_i; M_{ij}]$ - $[A'_i; M'_{ij}]$ -bimodule, which we denote $T_{\mathbf{m}}$. Since $T_{\mathbf{m}} = \bigoplus_i T_{ir} = \bigoplus_j T_{cj}$ we refer to $T_{\mathbf{m}}$ as a *matrix bimodule*.

2.8. Morita contexts for generalised matrix rings. As above fix generalised Morita contexts $(A_i; M_{ij}; \varphi_{ikj})$ and $(A'_i; M'_{ij}; \varphi'_{ikj})$. Let $N_{\mathbf{m}} = (N_{ij})$ be an $[A_i; M_{ij}]$ - $[A'_i; M'_{ij}]$ -matrix-bimodule as in §2.7 with respect to left and right structure maps

$$\beta_{ilk}: M_{il} \otimes_{A_l} N_{lk} \rightarrow N_{ik}, \quad \beta'_{lkj}: N_{lk} \otimes_{A_k} M'_{kj} \rightarrow N_{lj}.$$

Likewise let $L_{\mathbf{m}} = (L_{ij})$ be a $[A'_i; M'_{ij}]$ - $[A_i; M_{ij}]$ -matrix-bimodule with respect to

$$\gamma'_{ilk}: M'_{il} \otimes_{A_l} L_{lk} \rightarrow L_{ik}, \quad \gamma_{lkj}: L_{lk} \otimes_{A_k} M_{kj} \rightarrow L_{lj}.$$

Suppose in addition the following conditions hold.

- $\alpha_{ikj}: N_{ik} \otimes L_{kj} \rightarrow M_{ij}$ ($i, j, k = 1, \dots, n$) are additive maps such that the map

$$\begin{vmatrix} \alpha_{1k1} & \cdots & \alpha_{1kn} \\ \vdots & \ddots & \vdots \\ \alpha_{nk1} & \cdots & \alpha_{nkn} \end{vmatrix} : \begin{pmatrix} N_{11} & \cdots & N_{1n} \\ \vdots & \ddots & \vdots \\ N_{n1} & \cdots & N_{nn} \end{pmatrix} \times \begin{pmatrix} L_{11} & \cdots & L_{1n} \\ \vdots & \ddots & \vdots \\ L_{n1} & \cdots & L_{nn} \end{pmatrix} \rightarrow \begin{pmatrix} A_1 & \cdots & M_{1n} \\ \vdots & \ddots & \vdots \\ M_{n1} & \cdots & A_n \end{pmatrix}$$

is an $[A'_i; M'_{ij}]$ -balanced $[A_i; M_{ij}]$ - $[A_i; M_{ij}]$ -bimodule homomorphism.

- $\alpha'_{ikj}: L_{ik} \otimes N_{kj} \rightarrow M'_{ij}$ are additive maps such that the map

$$\begin{vmatrix} \alpha'_{1k1} & \cdots & \alpha'_{1kn} \\ \vdots & \ddots & \vdots \\ \alpha'_{lk1} & \cdots & \alpha'_{lkn} \end{vmatrix} : \begin{pmatrix} L_{11} & \cdots & L_{1n} \\ \vdots & \ddots & \vdots \\ L_{n1} & \cdots & L_{nn} \end{pmatrix} \times \begin{pmatrix} N_{11} & \cdots & N_{1n} \\ \vdots & \ddots & \vdots \\ N_{n1} & \cdots & N_{nn} \end{pmatrix} \rightarrow \begin{pmatrix} A'_1 & \cdots & M'_{1n} \\ \vdots & \ddots & \vdots \\ M'_{n1} & \cdots & A'_n \end{pmatrix}$$

is an $[A_i; M_{ij}]$ -balanced $[A'_i; M'_{ij}]$ - $[A'_i; M'_{ij}]$ -bimodule homomorphism.

- $\beta_{ikj}(\alpha_{ihk} \otimes \mathbf{1}) = \beta'_{ihj}(\mathbf{1} \otimes \alpha'_{hjk})$ ($h, i, j, k = 1, \dots, n$) as maps $N_{ih} \otimes L_{hk} \otimes N_{kj} \rightarrow N_{ij}$.
- $\gamma_{ihj}(\mathbf{1} \otimes \alpha_{hjk}) = \gamma'_{ikj}(\alpha'_{ihk} \otimes \mathbf{1})$ ($h, i, j, k = 1, \dots, n$) as maps $L_{ih} \otimes N_{hk} \otimes L_{kj} \rightarrow L_{ij}$.

By the first two items above there are induced bimodule homomorphisms

$$|\underline{\alpha}_{ikj}| : N_{\mathbf{m}} \otimes_{[A'_i; M'_{ij}]} L_{\mathbf{m}} \rightarrow [A_i; M_{ij}], \quad |\underline{\alpha}'_{ikj}| : L_{\mathbf{m}} \otimes_{[A_i; M_{ij}]} N_{\mathbf{m}} \rightarrow [A'_i; M'_{ij}].$$

By the second two items above the mixed associativity conditions hold for the pair of maps $|\underline{\alpha}_{ikj}|$ and $|\underline{\alpha}'_{ikj}|$, meaning there is a (classical) Morita context

$$([A_i; M_{ij}], [A'_i; M'_{ij}]; N_{\mathbf{m}}, L_{\mathbf{m}}; |\underline{\alpha}_{ikj}|, |\underline{\alpha}'_{ikj}|).$$

In our main result, Theorem 3.14, we establish such a Morita context. In Corollary 3.15 we provide sufficient conditions for $|\underline{\alpha}_{ikj}|$ and $|\underline{\alpha}'_{ikj}|$ to be surjective, in which case

$$\begin{aligned} L_{\mathbf{m}} \otimes_{[A_i; M_{ij}]} - &: [A_i; M_{ij}]\text{-Mod} \rightarrow [A'_i; M'_{ij}]\text{-Mod}, \\ N_{\mathbf{m}} \otimes_{[A'_i; M'_{ij}]} - &: [A'_i; M'_{ij}]\text{-Mod} \rightarrow [A_i; M_{ij}]\text{-Mod}, \end{aligned}$$

define mutually quasi-inverse functors by Theorem 2.1, yielding a Morita equivalence

$$\begin{pmatrix} A_1 & \cdots & M_{1n} \\ \vdots & \ddots & \vdots \\ M_{n1} & \cdots & A_n \end{pmatrix} \sim \begin{pmatrix} A'_1 & \cdots & M'_{1n} \\ \vdots & \ddots & \vdots \\ M'_{n1} & \cdots & A'_n \end{pmatrix}.$$

3. EXCISION AND LIGATION

In §3 we fix a particular ring-theoretic situation, and set-up the following notation.

- * A_1, \dots, A_n, R and S are rings which are associative but not necessarily unital.
- * $R = A_t$ as rings for some $t = 1, \dots, n$ which is fixed but arbitrary.
- * $(R, S; {}_R N_S, {}_S L_R; \zeta: N \otimes L \rightarrow R, \theta: L \otimes N \rightarrow S,)$ is a classical Morita context.
- * $(A_i; M_{ij}; \varphi_{ikj}: M_{ik} \otimes M_{kj} \rightarrow M_{ij})$ is a generalised Morita context.
- * $[A_i; M_{ij}]$ is the generalised matrix ring for $(A_i; M_{ij}; \varphi_{ikj})$, denoted in detail by

$$[A_i; M_{ij}] = \begin{bmatrix} A_1 & \cdots & M_{1t} & \cdots & M_{1n} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ M_{t1} & \cdots & R & \cdots & M_{tn} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ M_{n1} & \cdots & M_{nt} & \cdots & A_n \end{bmatrix}$$

In Definition 3.1 we construct a new Morita context (see Lemma 3.2) whose matrix ring is found by *excising* entries of $[A_i; M_{ij}]$ in row t or column t , and *ligating* the unchanged entries with substituted bimodules, such that R changes to S .

3.1. Excision. In Definition 3.1 we generalise the *composition* of a pair of classical Morita contexts from the thesis of of Marín [14, Proposition 4.11] (c.f. Völjako and Laan [16, §4]).

Definition 3.1. The *composition* of $(R, S; N, L; \zeta, \theta)$ and $(A_i; M_{ij}; \varphi_{ikj})$ is the tuple

$$(R, S; N, L; \zeta, \theta) \circ (A_i; M_{ij}; \varphi_{ikj}) = (A'_i; M'_{ij}; \varphi'_{ijk}),$$

which is given by the rings, bimodules and bimodule homomorphisms defined below.

- * A'_i , the rings, are given by $A'_t = S$ for $i = t$, and $A'_i = A_i$ for $i \neq t$.
- * M'_{ij} , the A'_i - A'_j -bimodules, are defined as follows

$$\begin{aligned} M'_{tt} &= S & (i = j = t), \\ M'_{it} &= M_{it} \otimes_R N & (j = t \neq i), \\ M'_{tj} &= L \otimes_R M_{tj} & (i = t \neq j), \\ M'_{ij} &= M_{ij} & (i, j \neq t). \end{aligned}$$

- * $\varphi'_{ijk}: M'_{ik} \otimes_{A'_k} M'_{kj} \rightarrow M'_{ij}$, the bimodule homomorphisms, are defined as follows

$$\begin{aligned} \mu_S: S \otimes S &\rightarrow S & (i = k = j = t), \\ \mathbf{1}_{it}^M \otimes \rho_N^S: M_{it} \otimes N \otimes S &\rightarrow M_{it} \otimes N & (k = j = t \neq i), \\ \lambda_L^S \otimes \mathbf{1}_{tj}^M: S \otimes L \otimes M_{tj} &\rightarrow L \otimes M_{tj} & (i = k = t \neq j), \\ \theta \circ (\rho_L^R \otimes \mathbf{1}_N) \circ (\mathbf{1}_L \otimes \varphi_{tkt} \otimes \mathbf{1}_N): L \otimes M_{tr} \otimes M_{rt} \otimes N &\rightarrow S & (i = j = t \neq k), \\ \varphi_{ikt} \otimes \mathbf{1}_N: M_{ik} \otimes M_{kt} \otimes N &\rightarrow M_{it} \otimes N & (j = t \neq i, k), \\ \mathbf{1}_L \otimes \varphi_{tkj}: L \otimes_R M_{tk} \otimes M_{kj} &\rightarrow L \otimes M_{tj} & (i = t \neq k, j), \\ \varphi_{itj} \circ (\rho_{M_{it}}^R \otimes \mathbf{1}_{tj}^M) \circ (\mathbf{1}_{it}^M \otimes \zeta \otimes \mathbf{1}_{tj}^M): M_{it} \otimes N \otimes L \otimes M_{tj} &\rightarrow M_{ij} & (k = t \neq i, j), \\ \varphi_{ikj}: M_{ik} \otimes M_{kj} &\rightarrow M_{ij} & (i, j, k \neq t). \end{aligned}$$

Lemma 3.2. *The composition $(A'_i; M'_{ij}; \varphi'_{ikj})$ of $(R, S; N, L; \zeta, \theta)$ with $(A_i; M_{ij}; \varphi_{ikj})$ is a generalised Morita context. Furthermore, if the rings R, S, A_i have local units and the bimodules N, L, M_{ij} are unital, then the bimodules M'_{ij} are unital.*

Proof. By construction we have that the first four conditions in §2.3 hold. To see that the composition is a generalised Morita context it suffices to check the fifth condition holds.

Hence we begin by checking that $\varphi'_{ihj} \circ (\mathbf{1}_{ih}^{M'} \otimes \varphi'_{hjk}) = \varphi'_{ikj} \circ (\varphi'_{ihk} \otimes \mathbf{1}_{kj}^{M'})$ holds for all $h, i, j, k = 1, \dots, n$. When $h = i = j = k = t$ this equation follows from the associativity of multiplication in the ring S . When $t \neq h, i, j, k$ this equation follows from the fact that $(A_i; M_{ij}; \varphi_{ikj})$ is a generalised Morita context.

We now consider the case where $h = j = t \neq i, k$. By definition we have

$$\varphi'_{itj}((\mathbf{1}_{it}^{M'} \otimes \varphi'_{tkj})(m \otimes n \otimes l' \otimes m' \otimes m'')) = m \otimes n \theta(l' \varphi_{tkt}(m' \otimes m'')) \otimes n''$$

for all $m \otimes n \in M_{it} \otimes_R N$ and $l' \otimes m' \in L \otimes_R M_{tk}$ and $m'' \otimes n'' \in M_{kt} \otimes_R N$. Since ζ is right R -linear, since φ_{tkt} is left R -linear, and by the associativity conditions for the classical Morita context $(R, S; N, L; \zeta, \theta)$, we have that $n \theta(l' \varphi_{tkt}(m' \otimes m'')) \otimes n''$ is equal to $\varphi_{tkt}(\zeta(n \otimes l') m' \otimes m'') n''$ which means that

$$\varphi'_{itj}((\mathbf{1}_{it}^{M'} \otimes \varphi'_{tkj})(m \otimes n \otimes l' \otimes m' \otimes m'')) = \varphi_{itt}(m \otimes \varphi_{tkt}(\zeta(n \otimes l') m' \otimes m'')) \otimes n''$$

which is the image of $m \otimes n \otimes l' \otimes m' \otimes m'' \otimes n''$ under $\varphi'_{tkj} \circ (\varphi'_{ttk} \otimes \mathbf{1}_{kj}^{M'})$ by definition.

Now the remaining cases are similar, and so we omit the details.

For the second assertion in the statement, consider how the right S -action on N defines the right action on each M'_{pt} and similarly consider the left action of S on each M'_{tq} . ■

By Lemma 3.2 we can and do fix the following additional notation for what remains.

- * $(A'_i; M'_{ij}; \varphi'_{ikj}; M'_{ik} \otimes M'_{kj} \rightarrow M'_{ij})$ is the generalised Morita context found by the composition of $(R, S; N, L; \zeta, \theta)$ and $(A_i; M_{ij}; \varphi_{ikj})$ at $t = 1, \dots, n$.
- * $[A'_i; M'_{ij}]$ is the generalised matrix ring for $(A'_i; M'_{ij}; \varphi'_{ikj})$, depicted in detail by

$$[A'_i; M'_{ij}] = \begin{bmatrix} A_1 & \cdots & M_{1t} \otimes_R N & \cdots & M_{1n} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ L \otimes_R M_{t1} & \cdots & S & \cdots & L \otimes_R M_{tn} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ M_{n1} & \cdots & M_{nt} \otimes_R N & \cdots & A_n \end{bmatrix}$$

Definition 3.3. By the *column-excision* (N_{ij}) of $(A_i; M_{ij}; \varphi_{ikj})$ by $(R, S; N, L; \zeta, \theta)$ at $t = 1, \dots, n$ we mean the direct sum of additive groups defined and denoted by

$$\begin{pmatrix} N_{11} & \cdots & N_{1n} \\ \vdots & \ddots & \vdots \\ N_{n1} & \cdots & N_{nn} \end{pmatrix} = \begin{pmatrix} A_1 & \cdots & M_{1t} \otimes_R N & \cdots & M_{1n} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ M_{t1} & \cdots & R \otimes_R N & \cdots & M_{tn} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ M_{n1} & \cdots & M_{nt} \otimes_R N & \cdots & A_n \end{pmatrix}$$

That is, (N_{ij}) is found by tensoring of each component in column t of $[A_i; M_{ij}]$ with N .

Lemma 3.4 explains the reason for depicting the column-excision as we have. As seen in the proof, the bimodule is canonical in that the left and right actions correspond, as in §2.3–2.8, to matrix multiplication with respect to the bimodule maps involved so far.

For each $j \neq t$, the additive group given by column j in (N_{ij}) is equal to that found by taking column j in the regular module for $[A_i; M_{ij}]$. In Lemma 3.4 we equip such a group the inherited left $[A_i; M_{ij}]$ -module structure given by multiplication in $[A_i; M_{ij}]$.

For column t in (N_{ij}) , we note that this left module is essentially constructed in items (i) and (iii) from [11, pp 1490–1491] in case $n = 2$. Since the rings and modules we consider need not be unital, we recall this construction for completeness, and repeat the check that the given action is a left module action.

Lemma 3.4. *The column-excision (N_{ij}) has the following additional (canonical) structure.*

- (i) (N_{ij}) is a left $[A_i; M_{ij}]$ -module ($[A_i; M_{ij}]$ is the generalised matrix ring).
- (ii) (N_{ij}) is a right $[A'_i; M'_{ij}]$ -module ($(A'_i; M'_{ij}; \varphi'_{ijk})$ is the composition).
- (iii) (N_{ij}) is an $[A_i; M_{ij}]$ - $[A'_i; M'_{ij}]$ -matrix-bimodule with respect to (i) and (ii).

Proof. Firstly note that (N_{ij}) can be decomposed into additive groups, either consisting of columns, or consisting of rows.

Following §2.4 we show, in parts (i) and (ii) respectively, that these columns are in fact column collections for $[A_i; M_{ij}]$ and row collections for $[A'_i; M'_{ij}]$. Showing this shall respectively verify the first two conditions from §2.7. Noting that the third condition holds by construction, in part (iii) it will be sufficient to check the fourth condition holds.

(i) For each $j \neq t$ let $N_{ij} = M_{ij}$ and consider the column $N_{cj} = (N_{1j}, \dots, N_{nj})$ equipped with the left structure maps $\beta_{ikj} = \varphi_{ikj}$. This gives N_{cj} the structure of a left $[A_i; M_{ij}]$ -module. We now deal with column t , which is more involved.

So, consider $N_{ct} = (N_{1t}, \dots, N_{nt})$ where $N_{it} = M_{it} \otimes_R N$ for each i , depicted as a column $(M_{1t} \otimes_R N, \dots, R \otimes_R N, \dots, M_{nt} \otimes_R N)$ of additive groups. Define the left structure maps $\beta_{ikt}: M_{ik} \otimes_{A_k} N_{kt} \rightarrow N_{it}$ ($i, k = 1, \dots, n$) by $\beta_{ikt} = \varphi_{ikt} \otimes \mathbf{1}_N$.

Again by the associativity conditions for $(A_i; M_{ij}; \varphi_{ikj})$, we have

$$\begin{aligned} \beta_{iht}(\varphi_{ikh} \otimes \mathbf{1}_h^X) &= (\varphi_{iht} \otimes \mathbf{1}_N)(\varphi_{ikh} \otimes \mathbf{1}_{ht}^M \otimes \mathbf{1}_N) = (\varphi_{iht}(\varphi_{ikh} \otimes \mathbf{1}_{ht}^M)) \otimes \mathbf{1}_N \\ &= (\varphi_{ikt}(\mathbf{1}_{ik}^M \otimes \varphi_{kht})) \otimes \mathbf{1}_N = (\varphi_{ikt} \otimes \mathbf{1}_N)(\mathbf{1}_{ik}^M \otimes \varphi_{kht} \otimes \mathbf{1}_N) = \beta_{ikt}(\mathbf{1}_{ih}^M \otimes \beta_{kht}), \end{aligned}$$

which verifies the fourth condition from §2.4, and thus the first condition from §2.7.

(ii) As in the proof of (i) above, we need only consider the additive groups coming from row t since the other rows are identical to the corresponding row in the generalised matrix ring $[A'_i; M'_{ij}]$. That is, we are letting $\beta'_{ikj} = \varphi'_{ikj}$ whenever $i \neq t$.

So, now consider the direct sum of the R - A_i -bimodules N_{ti} given by $N_{ti} = M_{ti}$ for $i \neq t$ and $N_{tt} = R \otimes N$. Define the right structure maps $\beta'_{thi}: N_{th} \otimes_{A'_h} M'_{hi} \rightarrow N_{ti}$ by

$$\begin{aligned} \mathbf{1}_R \otimes \rho_N^S: R \otimes N \otimes_S S &\rightarrow R \otimes N & (h = t = i), \\ \lambda_{M_{ti}}^R \circ (\mu_R \otimes \mathbf{1}_{ti}^M) \circ (\mathbf{1}_R \otimes \zeta \otimes \mathbf{1}_{ti}^M): R \otimes_R N \otimes_S L \otimes_R M_{ti} &\rightarrow M_{ti} & (h = t \neq i), \\ \varphi_{tht} \otimes \mathbf{1}_N: M_{th} \otimes_{A_h} M_{ht} \otimes_R N &\rightarrow R \otimes_R N & (h \neq t = i), \\ \varphi_{thi}: M_{th} \otimes_{A_h} M_{hi} &\rightarrow M_{ti} & (h \neq t \neq i). \end{aligned}$$

We check that the equation $\beta'_{thj}(\mathbf{1}_{th}^Y \otimes \varphi'_{hij}) = \beta'_{tij}(\beta'_{thi} \otimes \mathbf{1}_{ij}^{M'})$ holds for the case $h = j = t \neq i$. The cases where $h = t \neq i, j$ and $i = t \neq h, j$ are similar, and the remaining cases are more straightforward, and so such details are omitted.

Begin by noting that for any $r \in R$, $n, n' \in N$, $l \in L$, $m \in M_{it}$ and $m' \in M_{ti}$ we have

$$\begin{aligned} r \otimes n\theta(l\varphi_{tit}(m \otimes m') \otimes n') &= r \otimes \zeta(n \otimes l\varphi_{tit}(m \otimes m'))n' \\ &= r \otimes \zeta(n \otimes l)\varphi_{tit}(m \otimes m')n' = r \otimes \varphi_{tit}(\zeta(n \otimes l)m \otimes m')n' \\ &= r\varphi_{tit}(\zeta(n \otimes l)m \otimes m') \otimes n' = \varphi_{tit}(r\zeta(n \otimes l)m \otimes m') \otimes n' \\ &= \varphi_{tit}(\lambda_{M_{ti}}^R(\mu_R(r \otimes \zeta(n \otimes l)) \otimes m) \otimes m') \otimes n'. \end{aligned}$$

As a result we have the following equalities,

$$\begin{aligned} (\mathbf{1}_R \otimes \rho_N^S)(\mathbf{1}_{tt}^Y \otimes \varphi'_{tit}) &= (\mathbf{1}_R \otimes \rho_N^S)(\mathbf{1}_{tt}^Y \otimes (\theta(\rho_L^R \otimes \mathbf{1}_N)(\mathbf{1}_L \otimes \varphi_{tit} \otimes \mathbf{1}_N))) \\ &= (\mathbf{1}_R \otimes \rho_N^S)((\mathbf{1}_R \otimes \mathbf{1}_N) \otimes (\theta(\rho_L^R \otimes \mathbf{1}_N)(\mathbf{1}_L \otimes \varphi_{tit} \otimes \mathbf{1}_N))) \\ &= (\varphi_{tit}(\lambda_{M_{ti}}^R(\mu_R \otimes \mathbf{1}_{ti}^M)(\mathbf{1}_R \otimes \zeta \otimes \mathbf{1}_{ti}^M)) \otimes \mathbf{1}_{it}^M) \otimes \mathbf{1}_N = (\varphi_{tit} \otimes \mathbf{1}_N)(\beta'_{ti} \otimes \mathbf{1}_{it}^M \otimes \mathbf{1}_N). \end{aligned}$$

Thus we have $\beta'_{itt}(\mathbf{1}_{tt}^Y \otimes \varphi'_{tit}) = \beta'_{tit}(\beta'_{tti} \otimes \mathbf{1}_{it}^{M'})$, which verifies the dual of the fourth condition from §2.4 (for the case we cover), and thus the second condition from §2.7.

(iii) As discussed above, the third condition from §2.7 is automatic and it is sufficient from here to check the fourth condition holds. Hence we will show $\beta'_{ilk}(\mathbf{1}_{il}^M \otimes \beta'_{lkj}) = \beta'_{ikj}(\beta'_{ilk} \otimes \mathbf{1}_{kj}^{M'})$ for all $i, j, k, l = 1, \dots, n$. First consider cases based on which elements of $\{i, j, k, l\}$ are equal to t . The proof in the case $k, l = t \neq i, j$ follows from similar arguments to those used in the proofs of (i) and (ii) above, noting that we have

$$\varphi_{ihj}(\mathbf{1}_{ih}^M \otimes \varphi_{hjk}) = \varphi_{ikj}(\varphi_{ihk} \otimes \mathbf{1}_{kj}^M), \quad \varphi'_{ihj}(\mathbf{1}_{ih}^{M'} \otimes \varphi'_{hjk}) = \varphi'_{ikj}(\varphi'_{ihk} \otimes \mathbf{1}_{kj}^{M'})$$

by the assumption that $(A_i; M_{ij}; \varphi_{ikj})$ is a generalised Morita context, and by Lemma 3.2. The remaining cases either again follow from similar arguments as used in (i) and (ii), or are more straightforward, and so we omit them. \blacksquare

Definition 3.5. By the *row-excision* (L_{ij}) we mean the direct sum of additive groups defined and denoted by

$$\begin{pmatrix} L_{11} & \cdots & L_{1n} \\ \vdots & \ddots & \vdots \\ L_{n1} & \cdots & L_{nn} \end{pmatrix} = \begin{pmatrix} A_1 & \cdots & M_{1t} & \cdots & M_{1n} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ L \otimes_R M_{t1} & \cdots & L \otimes_R R & \cdots & L \otimes_R M_{tn} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ M_{n1} & \cdots & M_{nt} & \cdots & A_n \end{pmatrix}.$$

That is, (L_{ij}) is found by tensoring of each component in row t of $[A_i; M_{ij}]$ with L .

Dually to Lemma 3.4, we have the following.

Lemma 3.6. *The row-excision (L_{ij}) has the following additional (canonical) structure.*

- (i) (L_{ij}) is a left $[A'_i; M'_{ij}]$ -module.
- (ii) (L_{ij}) is a right $[A_i; M_{ij}]$ -module.
- (iii) (L_{ij}) is an $[A'_i; M'_{ij}]-[A_i; M_{ij}]$ -matrix-bimodule with respect to (i) and (ii).

Proof. For each $i = 1, \dots, n$ define $L_{i\mathbf{r}} = (L_{i1}, \dots, L_{in})$ where $L_{tj} = L \otimes M_{tj}$ and $L_{ij} = M_{ij}$ for $i \neq t$. Define the right structure maps $\gamma_{ikj}: L_{ik} \otimes_{A_k} M_{kj} \rightarrow L_{ij}$ by $\gamma_{tkj} = \mathbf{1}_L \otimes \varphi_{tkj}$ and $\gamma_{ikj} = \varphi_{ikj}$ for $i \neq t$. For each $j = 1, \dots, n$ define $L_{\mathbf{c}j} = (L_{1j}, \dots, L_{nj})$ where $L_{tj} = L \otimes M_{tj}$ and $L_{ij} = M_{ij}$ for $i \neq t$. Hence $X = Y$ which is the additive group (L_{ij}) . Define the left structure maps $\gamma'_{ikj}: M'_{ik} \otimes_{A_k} L_{kj} \rightarrow L_{ij}$ ($i, j, k = 1, \dots, n$) by $\gamma'_{ikj} = \varphi'_{ikj}$ for $j \neq t$, and otherwise as follows. Define the maps $\gamma'_{ikt}: M'_{ik} \otimes_{A'_k} L_{kt} \rightarrow L_{it}$ ($i, k = 1, \dots, n$) by

$$\begin{aligned} \lambda_L^S \otimes \mathbf{1}_R: S \otimes_S L \otimes_R R &\rightarrow L \otimes_R R & (i = t = k), \\ \mathbf{1}_L \otimes \varphi_{ikt}: L \otimes_R M_{tk} \otimes_{A_k} M_{kt} &\rightarrow L \otimes_R R & (i = t \neq k), \\ \rho_{M_{it}}^R \circ (\mathbf{1}_{it}^M \otimes \mu_R) \circ (\mathbf{1}_{it}^M \otimes \zeta \otimes \mathbf{1}_R): M_{it} \otimes_R N \otimes_S L \otimes_R R &\rightarrow M_{it} & (i \neq t = k), \\ \varphi_{ikt}: M_{ik} \otimes_{A_k} M_{kt} &\rightarrow M_{it} & (i \neq t \neq k). \end{aligned}$$

We leave the verification of (i)–(iii) to the reader, since such arguments are similar to those used in the proof of Lemma 3.4. Note that $(L_{ij}) = \bigoplus_i L_{i\mathbf{r}}$ as right $[A_i; M_{ij}]$ -modules and $(L_{ij}) = \bigoplus_j L_{\mathbf{c}j}$ as left $[A'_i; M'_{ij}]$ -modules. \blacksquare

3.2. Ligation. By Lemmas 3.4 and 3.6 the column-excision (N_{ij}) and the row-excision (L_{ij}) define matrix bimodules over the generalised matrix rings $[A_i; M_{ij}]$ and $[A'_i; M'_{ij}]$ of the generalised Morita contexts $(A_i; M_{ij}; \varphi_{ikj})$ and $(A'_i; M'_{ij}; \varphi'_{ikj})$.

From now on we relabel using the notation $N_{\mathbf{m}} = (N_{ij})$ and $L_{\mathbf{m}} = (L_{ij})$ to keep track of the fact these additive groups are matrix bimodules.

Definition 3.7. The *column-row ligation* is the biadditive map $|\alpha_{ikj}|$ of the form

$$\left| \begin{array}{ccc} \alpha_{1k1} & \cdots & \alpha_{1kn} \\ \vdots & \ddots & \vdots \\ \alpha_{nk1} & \cdots & \alpha_{nkn} \end{array} \right| : \begin{pmatrix} N_{11} & \cdots & N_{1n} \\ \vdots & \ddots & \vdots \\ N_{n1} & \cdots & N_{nn} \end{pmatrix} \times \begin{pmatrix} L_{11} & \cdots & L_{1n} \\ \vdots & \ddots & \vdots \\ L_{n1} & \cdots & L_{nn} \end{pmatrix} \rightarrow \begin{pmatrix} A_1 & \cdots & M_{1n} \\ \vdots & \ddots & \vdots \\ M_{n1} & \cdots & A_n \end{pmatrix},$$

given by the A_i - A_j -bimodule homomorphisms $\alpha_{ikj}: N_{ik} \otimes_{A'_k} L_{kj} \rightarrow M_{ij}$ defined by

$$\begin{aligned} \varphi_{ikj}: M_{ik} \otimes_{A_k} M_{kj} &\rightarrow M_{ij} & (k \neq t), \\ \varphi_{itj} \circ (\rho_{M_{it}}^R \otimes \mathbf{1}_{tj}^M) \circ (\mathbf{1}_{it}^M \otimes \zeta \otimes \mathbf{1}_{tj}^M): M_{it} \otimes_R N \otimes_S L \otimes_R M_{tj} &\rightarrow M_{ij} & (k = t). \end{aligned}$$

In what follows we show that the column-row ligation is balanced, and hence factors through the corresponding tensor product; see Lemma 3.8.

Lemma 3.8. *The column-row-ligation $|\alpha_{ikj}|: N_{\mathbf{m}} \times L_{\mathbf{m}} \rightarrow [A_i; M_{ij}]$ is $[A'_i; M'_{ij}]$ -balanced.*

Proof. Recall the right structure maps $\beta'_{ihk}: N_{ih} \otimes_{A'_h} M'_{hk} \rightarrow N_{ik}$ from Lemma 3.4(ii) and the left structure maps $\gamma'_{ijl}: M'_{ij} \otimes_{A'_j} L_{jl} \rightarrow L_{il}$ from Lemma 3.6(i). Considering the final condition from 2.6, to show the given map is balanced we require that $\alpha_{ihj}(\mathbf{1}_{ih}^N \otimes \gamma'_{hkj}) = \alpha_{ikj}(\beta'_{ihk} \otimes \mathbf{1}_{kj}^L)$ ($h, i, j, k = 1, \dots, n$) as maps $N_{ih} \otimes M'_{hk} \otimes L_{kj} \rightarrow M_{ij}$. As was done in the proofs of the aforementioned lemmas, we break our claim down into cases.

When $h \neq t \neq k$ the required equation follows from the mixed associativity conditions held by the generalised Morita context $(A_i; M_{ij}; \varphi_{ikj})$. When $i \neq t \neq j$ we can similarly apply the conditions held by the context $(A'_i; M'_{ij}; \varphi'_{ikj})$, which hold by Lemma 3.2. When $h = t = k$ or $i = t = j$ the required equation is straightforward. There are four cases left: when $h, i = t \neq j, k$; when $h, j = t \neq i, k$; when $k, i = t \neq h, j$; and when $k, j = t \neq h, i$.

We show the required equation holds in the first two cases, since the arguments for the other two cases are similar. So suppose $h, i = t \neq j, k$. The required equation then becomes

$$(\lambda_{M_{tj}}^R(\mu_R \otimes \mathbf{1})(\mathbf{1}_R \otimes \zeta \otimes \mathbf{1}))(\mathbf{1} \otimes \mathbf{1} \otimes \mathbf{1} \otimes \varphi_{tkj}) = \varphi_{tkj}((\lambda_{M_{tk}}^R(\mu_R \otimes \mathbf{1})(\mathbf{1} \otimes \zeta \otimes \mathbf{1})) \otimes \mathbf{1})$$

considered as maps of the form $R \otimes N \otimes L \otimes M_{tk} \otimes M_{kj} \rightarrow M_{tj}$. The proof from here is straightforward, since it can be shown that both sides send any pure tensor $r \otimes n \otimes l \otimes m \otimes m'$ to $r\zeta(n \otimes l)\varphi_{tkj}(m \otimes m')$ using the linearity of the involved homomorphisms.

Suppose instead $h, j = t \neq i, k$. This time the required equation becomes

$$\rho_{M_{it}}^R(\rho_{M_{it}}^R \otimes \mathbf{1})(\mathbf{1} \otimes \zeta \otimes \mathbf{1})(\mathbf{1} \otimes \mathbf{1} \otimes \mathbf{1} \otimes \varphi_{tkj}) = \varphi_{tkj}((\varphi_{itk}(\rho_{M_{it}}^R \otimes \mathbf{1})(\mathbf{1} \otimes \zeta \otimes \mathbf{1})) \otimes \mathbf{1})$$

considered as maps of the form $M_{it} \otimes N \otimes L \otimes M_{tk} \otimes M_{kt} \rightarrow M_{it}$. In this case one can show both sides send a pure tensor $m \otimes n \otimes l \otimes m' \otimes m''$ to $m\zeta(n \otimes l)\varphi_{tkj}(m' \otimes m'')$, using that $\varphi_{itk}(\varphi_{itk} \otimes \mathbf{1}) = \rho_{M_{it}}^R(\mathbf{1} \otimes \varphi_{tkj})$ by the associativity conditions for $(A_i; M_{ij}; \varphi_{ikj})$. \blacksquare

Note that the bimodule M_{ii} is unital if and only if the map $\mu_{A_i}: A \otimes A \rightarrow A$ is surjective.

Lemma 3.9. *The canonical additive map $|\underline{\alpha}_{ikj}|: N_{\mathbf{m}} \otimes_{[A_i; M'_{ij}]} L_{\mathbf{m}} \rightarrow [A_i; M_{ij}]$, defined by the column-row ligation, is an $[A_i; M_{ij}]\text{-}[A_i; M_{ij}]$ -bimodule homomorphism. Furthermore, $|\underline{\alpha}_{ikj}|$ is surjective provided ζ is surjective and each $A_i\text{-}A_j$ -bimodule M_{ij} is unital.*

Proof. Our first assertion is that $|\underline{\alpha}_{ikj}|$ is a left $[A_i; M_{ij}]$ -module homomorphism. Recall the left structure maps $\beta_{ihk}: M_{ih} \otimes_{A_h} N_{hk} \rightarrow N_{ik}$ from Lemma 3.4(i). As in §2.5, for our first assertion we will show that $\alpha_{ikj}(\beta_{ihk} \otimes \mathbf{1}_{L_{kj}}^M) = \varphi_{ihj}(\mathbf{1}_{ih}^M \otimes \alpha_{ikj})$ for all $h, i, j, k = 1, \dots, n$.

When $k \neq t$ this equality follows from the mixed associativity conditions held by $(A_i; M_{ij}; \varphi_{ikj})$. When $k = t$, the required equality becomes

$$\varphi_{itj}(\rho_{M_{it}}^R \otimes \mathbf{1}_{L_{tj}}^M)(\mathbf{1}_{it}^M \otimes \zeta \otimes \mathbf{1}_{L_{tj}}^M)(\varphi_{iht} \otimes \mathbf{1}_N \otimes \mathbf{1}_L \otimes \mathbf{1}_{L_{tj}}^M) = \varphi_{ihj}(\mathbf{1}_{ih}^M \otimes \varphi_{htj}(\rho_{M_{ht}}^M \otimes \mathbf{1}_{L_{tj}}^M)(\mathbf{1}_{ht}^M \otimes \zeta \otimes \mathbf{1}_{L_{tj}}^M))$$

which follows from the equality $\varphi_{ihj}(\mathbf{1}_{it}^M \otimes \varphi_{htj}) = \varphi_{itj}(\varphi_{iht} \otimes \mathbf{1}_{L_{tj}}^M)$, which as above is just one of the mixed associativity conditions held by $(A_i; M_{ij}; \varphi_{ikj})$.

Our second assertion is that $|\underline{\alpha}_{ikj}|$ is a right $[A_i; M_{ij}]$ -module homomorphism. So instead consider the right structure maps $\gamma_{ikj}: L_{ik} \otimes_{A_k} M_{kj} \rightarrow L_{ij}$ from Lemma 3.6(ii). For our second assertion we show that $\alpha_{ikj}(\mathbf{1}_{ih}^N \otimes \gamma_{hkj}) = \varphi_{ikj}(\alpha_{ihk} \otimes \mathbf{1}_{L_{kj}}^M)$ for all h, i, j, k .

As above, the proof that this equality holds is straightforward. In case $h = t$ the equation follows automatically as a consequence of the mixed associativity conditions for $(A_i; M_{ij}; \varphi_{ikj})$. In case $h \neq t$, it follows from the equation $\varphi_{itj}(\mathbf{1}_{it}^M \otimes \varphi_{tkj}) = \varphi_{ikj}(\varphi_{itk} \otimes \mathbf{1}_{L_{kj}}^M)$.

For the final claim in the statement, it suffices to fix i and j arbitrary and choose some k such that the map $\alpha_{ikj}: N_{ik} \otimes_{A_k} L_{kj} \rightarrow M_{ij}$ is surjective.

Since the $A_i\text{-}A_j$ bimodules M_{ij} are unital the maps $\rho_{M_{ik}}^{A_k} = \varphi_{ikk}$ and $\lambda_{M_{kj}}^{A_k} = \varphi_{kkj}$ are surjective. When $k \neq t$ we have $\alpha_{ikj} = \varphi_{ikj}$. Hence if $i \neq t$ then it suffices to choose $k = i$, and if $j \neq t$ it suffices to choose $k = j$. So we can assume $i = t = j$, and we choose $k = t$, so that α_{ikj} is the composition of φ_{ttt} , $\rho_{M_{tt}}^R \otimes \mathbf{1}$ and $\mathbf{1} \otimes \zeta \otimes \mathbf{1}$. As above, $\varphi_{ttt} = \mu_R = \rho_{M_{tt}}^R$ is surjective, and ζ is surjective by assumption. \blacksquare

Later we use Lemma 3.9, together with results from discussed in the sequel, in conjunction with Theorem 2.1. We start with a dual construction to the column-row ligation from Definition 3.7, completing the ingredients needed to state Theorem 3.14.

Definition 3.10. The *row-column ligation* is the biadditive map $|\alpha'_{ikj}|$ of the form

$$\left| \begin{array}{ccc} \alpha'_{1k1} & \cdots & \alpha'_{1kn} \\ \vdots & \ddots & \vdots \\ \alpha'_{nk1} & \cdots & \alpha'_{nkn} \end{array} \right| : \begin{pmatrix} L_{11} & \cdots & L_{1n} \\ \vdots & \ddots & \vdots \\ L_{n1} & \cdots & L_{nn} \end{pmatrix} \times \begin{pmatrix} N_{11} & \cdots & N_{1n} \\ \vdots & \ddots & \vdots \\ N_{n1} & \cdots & N_{nn} \end{pmatrix} \rightarrow \begin{pmatrix} A'_1 & \cdots & M'_{1n} \\ \vdots & \ddots & \vdots \\ M'_{n1} & \cdots & A'_n \end{pmatrix},$$

given by the A'_i - A'_j -bimodule homomorphisms $\alpha'_{ikj} : L_{ik} \otimes_{A_k} N_{kj} \rightarrow M'_{ij}$ defined by

$$\begin{aligned} \varphi_{ikj} : M_{ik} \otimes_{A_k} M_{kj} &\rightarrow M_{ij} & (i \neq t \neq j), \\ \varphi_{ikt} \otimes \mathbf{1}_N : M_{ik} \otimes_{A_k} M_{kt} \otimes_R N &\rightarrow M_{it} \otimes_R N & (i \neq t = j), \\ \mathbf{1}_L \otimes \varphi_{tkj} : L \otimes_R M_{tk} \otimes_{A_k} M_{kj} &\rightarrow L \otimes_R M_{tj} & (i = t \neq j), \\ \theta \circ (\rho_L^R \otimes \mathbf{1}_N) \circ (\mathbf{1}_L \otimes \varphi_{tkl} \otimes \mathbf{1}_N) : L \otimes_R M_{tk} \otimes_{A_k} M_{kt} \otimes_R N &\rightarrow S & (i = t = j). \end{aligned}$$

Lemma 3.11. *The row-column ligation $|\alpha'_{ikj}| : L_{\mathbf{m}} \times N_{\mathbf{m}} \rightarrow [A'_i; M'_{ij}]$ is $[A_i; M_{ij}]$ -balanced.*

Proof. We follow the strategy from the proof of Lemma 3.8. Recall the left structure maps $\beta_{hjk} : M_{hk} \otimes_{A_k} N_{kj} \rightarrow N_{hj}$ from Lemma 3.4(i), and the right structure maps $\gamma_{ihk} : L_{ih} \otimes_{A_h} M_{hk} \rightarrow L_{ik}$ from Lemma 3.6(ii). As in Lemma 3.8, but dually, it suffices check that $\alpha'_{ihj}(\mathbf{1}_{ih}^L \otimes \beta_{hjk}) = \alpha'_{ikj}(\gamma_{ihk} \otimes \mathbf{1}_{kj}^N)$ as maps $L_{ih} \otimes M_{hk} \otimes N_{kj} \rightarrow M'_{ij}$. To do so is particularly straightforward, and only uses the associativity conditions for $(A_i; M_{ij}; \varphi_{ikj})$. \blacksquare

Lemma 3.12. *The canonical additive map $|\underline{\alpha}'_{ikj}| : L_{\mathbf{m}} \otimes_{[A_i; M_{ij}]} N_{\mathbf{m}} \rightarrow [A'_i; M'_{ij}]$, defined by the row-column ligation, is an $[A'_i; M'_{ij}]$ - $[A'_i; M'_{ij}]$ -bimodule homomorphism. Furthermore, $|\underline{\alpha}'_{ikj}|$ is surjective provided θ is surjective and each A_i - A_j -bimodule M_{ij} is unital.*

Proof. Recall the left and right structure maps $\gamma'_{ijl} : M'_{ij} \otimes_{A'_j} L_{jl} \rightarrow L_{il}$ and $\beta'_{ihk} : N_{ih} \otimes_{A'_h} M'_{hk} \rightarrow N_{ik}$ from Lemmas 3.6(i) and 3.4(ii). As in the proof of Lemma 3.9, to show $|\underline{\alpha}'_{ikj}|$ is a left $[A'_i; M'_{ij}]$ -module homomorphism it suffices to show $\alpha'_{ikj}(\gamma'_{ihk} \otimes \mathbf{1}_{kj}^N) = \varphi'_{ihj}(\mathbf{1}_{ih}^{M'} \otimes \alpha'_{hkj})$.

The proof that this equation holds is similar to the arguments seen so far in the article. Namely, consider cases, and when appropriate, apply the mixed associativity conditions for $(A_i; M_{ij}; \varphi_{ikj})$ or apply Lemma 3.2. The proof that $|\underline{\alpha}'_{ikj}|$ is a right $[A'_i; M'_{ij}]$ -module homomorphism follows a similar procedure.

The proof that $|\underline{\alpha}'_{ikj}|$ is surjective given the stated provisions hold follows similar arguments to those in the proof of the second statement in Lemma 3.9, and omitted. \blacksquare

3.3. Corner replacement. We conclude §3 by showing the canonical additive maps considered in Lemmas 3.9 and 3.12 satisfy mixed associativity conditions, and hence give rise to a classical Morita context of matrix rings and matrix bimodules; see Theorem 3.14.

The title of §3.3 refers to the following ‘surgery’ to be performed on $(A_i; M_{ij}; \varphi_{ikj})$.

- ‘Cut out’ the ring R and the bimodules M_{ij} ($i = t \neq j$ or $j = t \neq i$) using the column-excision and row-excision matrix bimodules from §3.1.
- ‘Graft back’ a new context from the column-row and row-column ligations in §3.2.
- Using the items above we ‘corner replace’ R in $[A_i; M_{ij}]$ with S in $[A'_i; M'_{ij}]$.

Lemma 3.13. *The maps $|\underline{\alpha}_{ikj}|$ and $|\underline{\alpha}'_{ikj}|$ satisfy the mixed associativity conditions.*

Proof. Considering the conditions from §2.8, we need to check the equations in the final two items hold. That is, we need to check that the equations

$$\begin{aligned} (*) \quad & \beta_{ikj}(\alpha_{ihk} \otimes \mathbf{1}_{kj}^N) = \beta'_{ihj}(\mathbf{1}_{ih}^N \otimes \alpha'_{hkj}), \\ (**) \quad & \gamma_{ihj}(\mathbf{1}_{ih}^L \otimes \alpha_{hkj}) = \gamma'_{ikj}(\alpha'_{ihk} \otimes \mathbf{1}_{kj}^L), \end{aligned}$$

hold for all $h, i, j, k = 1, \dots, n$. As usual we break the proof into cases.

When $h \neq t \neq j$ the required equality (*) follows directly from the mixed associativity conditions for $(A_i; M_{ij}; \varphi_{ikj})$. Similarly when $i \neq t \neq k$ the equality (**) follows directly.

When $h \neq t = j$ we can rewrite (*) as $\sigma \otimes \mathbf{1}_N = \tau \otimes \mathbf{1}_N$ for A_i - R -bimodule homomorphisms σ and τ , and the aforementioned associativity conditions give $\sigma = \tau$. Likewise when $i = t \neq k$ there exist R - A_j bimodule homomorphisms σ' and τ' where (**) has the form $\mathbf{1}_L \otimes \sigma' = \mathbf{1}_L \otimes \tau'$ and where $\sigma' = \tau'$ can be seen following a similar argument.

We complete the proof by showing (*) and (**) hold in the case where $h = i = k = t \neq j$. All the cases that remain for checking (*) holds, and all those that remain for (**), can be seen using similar arguments: and hence we omit such details.

So, suppose $h, i, k = t \neq j$. Here we have

$$\begin{aligned} \beta_{ikj} &= \varphi_{ttj}, & \alpha_{ihk} &= \varphi_{ttt}(\rho_{M_{tt}}^R \otimes \mathbf{1}_{tt}^M)(\mathbf{1}_{tt}^M \otimes \zeta \otimes \mathbf{1}_{tt}^M), \\ \mathbf{1}_{kj}^N &= \mathbf{1}_{tj}^M, & \mathbf{1}_{ih}^N &= \mathbf{1}_{tt}^M \otimes \mathbf{1}_N, \\ \beta'_{ihj} &= \lambda_{M_{tj}}^R(\mu_R \otimes \mathbf{1}_{tj}^M)(\mathbf{1}_R \otimes \zeta \otimes \mathbf{1}_{tj}^M), & \alpha'_{hkj} &= \mathbf{1}_L \otimes \varphi_{ttj}. \end{aligned}$$

Consequently, substituting these values, the equation (*) becomes

$$\varphi_{ttj}((\varphi_{ttt}(\rho_{M_{tt}}^R \otimes \mathbf{1}_{tt}^M)(\mathbf{1}_{tt}^M \otimes \zeta \otimes \mathbf{1}_{tt}^M)) \otimes \mathbf{1}_{tj}^M) = \lambda_{M_{tj}}^R(\mu_R \otimes \mathbf{1}_{tj}^M)(\mathbf{1}_R \otimes \zeta \otimes \mathbf{1}_{tj}^M)(\mathbf{1}_{tt}^M \otimes \mathbf{1}_N \otimes \mathbf{1}_L \otimes \varphi_{ttj}).$$

That this equality holds is straightforward, since for any $r, r' \in R$, $n \in N$, $l \in L$ and $m \in M_{tj}$ both sides send the pure tensor $r \otimes n \otimes l \otimes r' \otimes m$ to $r\zeta(n \otimes l)r'm$.

We now check (**) holds, which is more involved. For the case we are in we also have

$$\begin{aligned} \gamma_{ihj} &= \mathbf{1}_L \otimes \varphi_{ttj}, & \alpha_{hkj} &= \varphi_{ttj}(\rho_{M_{tt}}^R \otimes \mathbf{1}_{tj}^M)(\mathbf{1}_{tt}^M \otimes \zeta \otimes \mathbf{1}_{tj}^M), \\ \mathbf{1}_{ih}^L &= \mathbf{1}_L \otimes \mathbf{1}_{tt}^M, & \gamma'_{ikj} &= \varphi'_{ttj} = \lambda_L^S \otimes \mathbf{1}_{tj}^M, \\ \alpha'_{ihk} &= \theta(\rho_L^R \otimes \mathbf{1}_N)(\mathbf{1}_L \otimes \varphi_{ttt} \otimes \mathbf{1}_N), & \mathbf{1}_{kj}^L &= \mathbf{1}_L \otimes \mathbf{1}_{tj}^M. \end{aligned}$$

As was done above by substituting these values into (**) we require the equation

$$\mathbf{1}_L \otimes (\varphi_{ttj}(\mathbf{1}_{tt}^M \otimes (\varphi_{ttj}(\rho_{M_{tt}}^R \otimes \mathbf{1}_{tj}^M)(\mathbf{1}_{tt}^M \otimes \zeta \otimes \mathbf{1}_{tj}^M)))) = (\lambda_L^S \theta(\rho_L^R \otimes \mathbf{1}_N)(\mathbf{1}_L \otimes \varphi_{ttt} \otimes \mathbf{1}_N) \otimes \mathbf{1}_L) \otimes \mathbf{1}_{tj}^M.$$

Fix $l, l' \in L$, $r, r' \in R$, $n \in N$ and $m \in M_{tj}$. By using the mixed associativity conditions for $(R, S; N, L; \zeta, \theta)$, the map on the right hand side sends $l \otimes r \otimes r' \otimes n \otimes l' \otimes m$ to

$$\begin{aligned} & \lambda_L^S(\theta(\rho_L^R \otimes \mathbf{1}_N)(\mathbf{1}_L \otimes \varphi_{ttt} \otimes \mathbf{1}_N)(l \otimes r \otimes r' \otimes n) \otimes l') \otimes m \\ &= \lambda_L^S(\theta(\rho_L^R \otimes \mathbf{1}_N)(l \otimes rr' \otimes n) \otimes l') \otimes m = \lambda_L^S(\theta(lrr' \otimes n) \otimes l') \otimes m \\ &= \theta(lrr' \otimes n)l' \otimes m = lrr'\zeta(n \otimes l') \otimes m = l \otimes rr'\zeta(n \otimes l')m \end{aligned}$$

which is the image of $l \otimes r \otimes r' \otimes n \otimes l' \otimes m$ under the map on the left hand side. \blacksquare

We are now ready to state and prove our main result.

Theorem 3.14. *Let $(A_i; M_{ij}; \varphi_{ikj})$ be a generalised Morita context, and for each $i = 1, \dots, n$ let $(A_i, B_i; N_i, L_i; \theta_i, \zeta_i)$ be a classical Morita context.*

There is a (classical) Morita context for the rings

$$[A_i; M_{ij}] = \begin{bmatrix} A_1 & \cdots & M_{1n} \\ \vdots & \ddots & \vdots \\ M_{n1} & \cdots & A_n \end{bmatrix}, \quad [B_i; P_{ij}] = \begin{bmatrix} B_1 & \cdots & P_{1n} \\ \vdots & \ddots & \vdots \\ P_{n1} & \cdots & B_n \end{bmatrix}.$$

where $P_{ij} = L_i \otimes_{A_i} M_{ij} \otimes_{A_j} N_j$. Namely, this context is given by: the $[A_i; M_{ij}]$ - $[B_i; P_{ij}]$ -bimodule $N_{\mathbf{m}}$ and the $[B_i; P_{ij}]$ - $[A_i; M_{ij}]$ -bimodule $L_{\mathbf{m}}$ defined by

$$N_{\mathbf{m}} = \begin{pmatrix} A_1 \otimes_{A_1} N_1 & \cdots & M_{1n} \otimes_{A_n} N_n \\ \vdots & \ddots & \vdots \\ M_{n1} \otimes_{A_1} N_1 & \cdots & A_n \otimes_{A_n} N_n \end{pmatrix}, \quad L_{\mathbf{m}} = \begin{pmatrix} L_1 \otimes_{A_1} A_1 & \cdots & L_1 \otimes_{A_1} M_{1n} \\ \vdots & \ddots & \vdots \\ L_n \otimes_{A_n} M_{n1} & \cdots & L_n \otimes_{A_n} A_n \end{pmatrix};$$

and the bimodule homomorphisms $N_{\mathbf{m}} \otimes L_{\mathbf{m}} \rightarrow [A_i; M_{ij}]$, $L_{\mathbf{m}} \otimes N_{\mathbf{m}} \rightarrow [B_i; P_{ij}]$ defined by

$$\begin{bmatrix} \varphi_{1k1}(\theta_1) & \cdots & \varphi_{1kn}(\theta_n) \\ \vdots & \ddots & \vdots \\ \varphi_{nk1}(\theta_1) & \cdots & \varphi_{nkn}(\theta_n) \end{bmatrix}, \quad \begin{bmatrix} \zeta_1(\varphi_{1k1}) & \cdots & \zeta_1(\varphi_{1kn}) \\ \vdots & \ddots & \vdots \\ \zeta_n(\varphi_{nk1}) & \cdots & \zeta_n(\varphi_{nkn}) \end{bmatrix}$$

where the maps $\varphi_{ikj}(\theta_j)$ and $\zeta_i(\varphi_{ikj})$ are respectively defined as the composites

$$\begin{aligned} M_{ik} \otimes N_k \otimes L_k \otimes M_{kj} &\xrightarrow{\mathbf{1}_{ik}^M \otimes \theta_k \otimes \mathbf{1}_{kj}^M} M_{ik} \otimes A_k \otimes M_{kj} \xrightarrow{\rho \otimes \mathbf{1} = \mathbf{1} \otimes \lambda} M_{ik} \otimes M_{kj} \xrightarrow{\varphi_{ikj}} M_{ij} \\ L_i \otimes M_{ik} \otimes M_{kj} \otimes N_j &\xrightarrow{\mathbf{1}_i^L \otimes \varphi_{ikj} \otimes \mathbf{1}_j^N} L_i \otimes M_{ij} \otimes N_j \xrightarrow[\text{(i=j)}]{\lambda \otimes \mathbf{1} = \mathbf{1} \otimes \rho} L_i \otimes N_i \xrightarrow[\text{(i=j)}]{\zeta_i} B_i \end{aligned}$$

where the dashed arrows occur in the composition of $\zeta_i(\varphi_{ikj})$ if and only if $i = j$.

Proof. Apply Lemmas 3.4, 3.6, 3.9, 3.12 and 3.13. Now iterate over $t = 1, \dots, n$. ■

Corollary 3.15. *In the notation of Theorem 3.14, assume additionally: the rings A_i, B_i have local units; the bimodules N_i, L_i, M_{ij} are unital; and the maps θ_i, ζ_i are surjective (hence giving Morita equivalences $A_i \sim B_i$ by Theorem 2.1).*

Then the classical Morita context from Theorem 3.14 defines a Morita equivalence

$$\begin{bmatrix} A_1 & \cdots & M_{1n} \\ \vdots & \ddots & \vdots \\ M_{n1} & \cdots & A_n \end{bmatrix} \sim \begin{bmatrix} B_1 & \cdots & P_{1n} \\ \vdots & \ddots & \vdots \\ P_{n1} & \cdots & B_n \end{bmatrix}, \quad (P_{ij} = L_i \otimes_{A_i} M_{ij} \otimes_{A_j} N_j).$$

Proof. Immediate by Lemmas 3.12 and 3.13 and Theorems 2.1 and 3.14. ■

Proof of Proposition 1.1. Immediate by specifying Corollary 3.15 to $n = 2$, $(1-e)R(1-e) = A_1 = B_1$, $A_2 = eRe$ and $B_2 = S$. ■

We end §3 with some remarks. Firstly, the proof of our main result can probably be found using an inductive argument, however our direct approach is still rather elementary (and perhaps illustrative). Secondly, our results likely hold even after extending the class of rings we are considering, rings with local units, to the class of all idempotent rings. Note that Morita theory for idempotent rings was developed by García and Simón [9].

4. FUTURE DIRECTIONS

We now recall three ideas. The first idea, which is well-known, is that of a ring with *enough idempotents*; see §4.1. The second idea is that of a *functor ring* following terminology of Fuller [6], using a result due to Gabriel [7]; see §4.2. The third idea is a result due to Dung and García [5] relating the first two ideas.

These ideas have been combined before; see for example work of García, Gómez Sánchez, and Martínez Hernández [8, §1]. Functor rings were also used by Harada [12, 13] where they studied *perfect* categories. Furthermore, Yamagata [18] also used this idea to give a Morita duality theory for certain functor categories.

4.1. Rings with enough idempotents. Let A be a ring with a complete set $E = \{e_i \mid i \in I\}$ of orthogonal idempotents $e_i = e_i^2 \in A$, so that $e_i e_j = 0$ for $i \neq j$ and $A = \bigoplus_{i \in I} e_i A = \bigoplus_{i \in I} A e_i$. Then A has local units. Partition $I = I_- \sqcup I_+$ to give a partition $E = E_- \sqcup E_+$ and hence a Morita context $(A_-, A_+; M_{-+}, M_{+-}; \varphi_-, \varphi_+)$ where

$$A_{\pm} = \bigoplus_{e, e' \in E_{\pm}} e' A e, \quad M_{\pm\mp} = \bigoplus_{e' \in E_{\pm}, e \in E_{\mp}} e' A e, \quad \varphi_{\pm}(a \otimes b) = ab.$$

By construction we have $A \cong \begin{bmatrix} A_+ & M_{+-} \\ M_{-+} & A_- \end{bmatrix}$ as rings. Note also each bimodule $M_{\pm\mp}$ is unital.

Suppose $(A_{\pm}, B_{\pm}; N_{\pm}, L_{\pm}; \theta_{\pm}, \zeta_{\pm})$ are Morita contexts where N_{\pm}, L_{\pm} are unital. Note $L_{\pm} e' \otimes_{A_{\pm}} e A e'' = 0$ and $e'' A e \otimes_{A_{\pm}} e' N_{\pm} = 0$ for any distinct $e, e' \in E_{\pm}$. Hence, as tensor products commute over direct sums (see for example [17, §12.4]), we have

$$L_{\pm} \otimes_{A_{\pm}} M_{\pm\mp} = \bigoplus_{e \in E_{\pm}, e' \in E_{\mp}} L_{\pm} e \otimes_{A_{\pm}} e A e', \quad M_{\pm\mp} \otimes_{A_{\mp}} N_{\mp} = \bigoplus_{e \in E_{\pm}, e' \in E_{\mp}} e A e' \otimes_{A_{\mp}} e' N_{\mp}.$$

Hence we obtain a more explicit description of the column excision and row excision from §3.1. Note that the column-row ligation and row-column-ligation from §3.2 are defined using the multiplication in A . As in Corollary 3.15, when the maps $\theta_{\pm}, \zeta_{\pm}$ are all surjective the classical Morita context from Theorem 3.14 gives a Morita equivalence

$$A \sim \left[\begin{array}{ccc} & B_+ & \bigoplus_{e_{\pm} \in E_{\pm}} L_+ e_+ \otimes e_+ A e_- \otimes e_- N_- \\ \bigoplus_{e_{\pm} \in E_{\pm}} L_- e_- \otimes e_- A e_+ \otimes e_+ N_+ & & B_- \end{array} \right].$$

4.2. Functor rings. Let \mathcal{C} be a skeletally small additive category and let \mathcal{A} be a chosen *skeleton*, meaning a set of unique representatives of isomorphism classes of objects in \mathcal{C} . The *functor ring* of \mathcal{C} (with respect to \mathcal{A}) is the direct sum of additive groups

$$\text{Func}_{\mathcal{A}}(\mathcal{C}) = \bigoplus_{X, Y \in \mathcal{A}} \text{Hom}_{\mathcal{C}}(X, Y)$$

where the multiplication map $\text{Func}_{\mathcal{A}} \times \text{Func}_{\mathcal{A}} \rightarrow \text{Func}_{\mathcal{A}}$ is defined as the universal morphism given by the diagram of additive maps $\text{Hom}_{\mathcal{C}}(Y, Z) \times \text{Hom}_{\mathcal{C}}(X, Y) \rightarrow \text{Hom}_{\mathcal{C}}(X, Z)$ given by the composition in \mathcal{C} . It is straightforward to see that $\{\mathbf{1}_X \mid X \in \mathcal{A}\}$ defines a complete set $E_{\mathcal{A}}$ of orthogonal idempotents in $\text{Func}_{\mathcal{A}}(\mathcal{C})$. By [7, p. 347, Proposition 2], if $A = \text{Func}_{\mathcal{A}}(\mathcal{C})$ then there is an equivalence between the category $A\text{-Mod}$ of unital left R -modules and the category $(\mathcal{C}, \mathbf{Ab})$ of functors $\mathcal{C} \rightarrow \mathbf{Ab}$ where \mathbf{Ab} is the category of abelian groups. The author is interested in translating the discussion in §4.1 to the language of functor rings, with a view towards applying Corollary 3.15.

4.3. Locally finitely presented additive categories. Following Crawley-Boevey [4, p. 1642] an object Z in an additive category \mathcal{C} with direct limits is *finitely presented* if $\mathcal{C}(Z, -)$ commutes with direct limits, and \mathcal{C} is *locally finitely presented* if the finitely presented objects form a skeletally small subcategory where every object in \mathcal{C} is a direct limit of finitely presented ones. By [5, Theorem 1.1] there is a one-to-one correspondence between rings with enough idempotents up to Morita equivalence and locally finitely presented additive categories up to equivalence. Hence one can use the discussion so far to reinterpret Corollary 3.15 in terms of equivalence classes of locally finitely presentable additive categories.

ACKNOWLEDGEMENTS. The author is grateful for support by the Danish National Research Foundation (DNRF156); the Independent Research Fund Denmark (1026-00050B); and the Aarhus University Research Foundation (AUFF-F-2020-7-16). The author is also grateful to Benjamin Briggs, Peter Jørgensen and Toby Stafford for helpful discussions.

REFERENCES

- [1] P. N. Ánh and L. Márki. *Morita equivalence for rings without identity*. Tsukuba J. Math., **11**(1):1–16, 1987.
- [2] H. Bass. *Algebraic K-theory*. W. A. Benjamin, Inc., New York-Amsterdam, 1968.
- [3] Q. Chen and H. Liu. *Derived equivalences between generalized matrix algebras*. Czechoslovak Math. J., **70**(145)(1):147–160, 2020.
- [4] W. Crawley-Boevey. *Locally finitely presented additive categories*. Comm. Algebra, **22**(5):1641–1674, 1994.
- [5] N. V. Dung and J. L. García. *Additive categories of locally finite representation type*. J. Algebra, **238**(1):200–238, 2001.
- [6] K. R. Fuller. *On rings whose left modules are direct sums of finitely generated modules*. Proc. Amer. Math. Soc., **54**:39–44, 1976.
- [7] P. Gabriel. *Des catégories abéliennes*. Bull. Soc. Math. France, **90**:323–448, 1962.
- [8] J. L. García, P. L. Gómez Sánchez, and J. Martínez Hernández. *Locally finitely presented categories and functor rings*. Osaka J. Math., **42**(1):173–187, 2005.
- [9] J. L. García and J. J. Simón. *Morita equivalence for idempotent rings*. J. Pure Appl. Algebra, **76**(1):39–56, 1991.
- [10] E. L. Green. *On the representation theory of rings in matrix form*. Pacific J. Math., **100**(1):123–138, 1982.
- [11] E. L. Green and C. Psaroudakis. *On Artin algebras arising from Morita contexts*. Algebr. Represent. Theory, **17**(5):1485–1525, 2014.
- [12] M. Harada. *Perfect categories. I*. Osaka Math. J., **10**:329–341, 1973.
- [13] M. Harada. *Perfect categories. II. Hereditary categories*. Osaka Math. J., **10**:343–355, 1973.
- [14] L. Marín. *Categories of modules for idempotent rings and morita equivalences*. Master’s thesis, University of Glasgow, 1997.
- [15] K. Morita. *Duality for modules and its applications to the theory of rings with minimum condition*. Sci. Rep. Tokyo Kyoiku Daigaku Sect. A, **6**:83–142, 1958.
- [16] K. Völjako and V. Laan. *Morita contexts and unitary ideals of rings*. Proc. Est. Acad. Sci., **70**(2):122–134, 2021.
- [17] R. Wisbauer. *Foundations of module and ring theory*, volume 3 of *Algebra, Logic and Applications*. Gordon and Breach Science Publishers, Philadelphia, PA, german edition, 1991.
- [18] K. Yamagata. *On Morita duality for additive group valued functors*. Comm. Algebra, **7**(4):367–392, 1979.