

On the analytic version of the Mitiagin - DeLeeuw - Mirkhil non-inequality on bi-disc

Krystian Kazaniecki^{1,2} and Michał Wojciechowski³

¹Institute of Analysis, Johannes Kepler University Linz

²Institute of Mathematics, University of Warsaw

³Institute of Mathematics, Polish Academy of Sciences

January 24, 2023

Abstract

Using the method of Rudin-Shapiro polynomials we prove the analytic version of the Mitiagin - DeLeeuw - Mirkhil non-inequality for complex partial differential operators with constant coefficients on bi-disc.

1 Introduction

The classical results of Mitiagin, Mirkhil and DeLeeuw and Ornstein ([4, 12, 13], see also [2, 8, 10]) say that for $p = 1$ or $p = \infty$, and homogeneous differential operators of the same degree with constant coefficients $P_0(D), P_1(D), \dots, P_k(D)$ the inequality

$$\|P_0(D)f\|_p \lesssim \sum_{i=1}^k \|P_i(D)f\|_p \quad (1)$$

holds for every smooth function f with bounded support iff

$$P_0 \in \text{span}\{P_1, \dots, P_k\}.$$

Similar problem for differential operators with variable coefficients was considered in [11]. Analogous characterization holds for functions on tori. In the present paper we deal with the simplest case $d = 2$, $P_0(x, y) = xy$, $P_1(x, y) = x^2$ and $P_2(x, y) = y^2$ with additional restriction that f is analytic, i.e. $\hat{f}(n) = 0$ for n outside the set $\{n \in \mathbb{Z}^2; n_1 \geq 0, n_2 \geq 0\}$. Under such restriction for $p = 1$ the inequality (1) holds true. Indeed $D_{xy}^2 f = T_m(D_{x^2}^2 f + D_{y^2}^2 f)$ where T_m is an invariant operator given by the multiplier $m(n_1, n_2) = \frac{n_1 n_2}{n_1^2 + n_2^2}$. By ([16], see also [5], [1]) it is bounded in L^1 norm for analytic functions. Surprisingly, in this note we show that the situation in L^∞ norm is different. Namely we have:

Theorem 1. *For every $C > 0$, there exists an analytic trigonometric polynomial f on the torus \mathbb{T}^2 such that*

$$\|D_{xx}^2 f\|_\infty + \|D_{yy}^2 f\|_\infty \leq 1$$

but

$$\|D_{xy}^2 f\|_\infty > C.$$

This research was partially supported by the National Science Centre, Poland, and Austrian Science Foundation FWF joint CEUS programme. National Science Centre project no. 2020/02/Y/ST1/00072 and FWF project no. I5231.

It is not difficult to see that the Poisson extensions of $D_{xx}^2 f$, $D_{yy}^2 f$ and $D_{xy}^2 f$ onto unit bi-disc are respectively $z_2 \frac{\partial}{\partial z_2} \left(z_1 \frac{\partial}{\partial z_1} \right) F$, $\left(z_1 \frac{\partial}{\partial z_1} \right)^2 F$ and $\left(z_2 \frac{\partial}{\partial z_2} \right)^2 F$, where F is the Poisson extension of f . Using the maximum principle we can then reformulate Theorem 1:

Theorem 2. *For every $c > 0$, there exists an analytic polynomial F on the unit bi-disc such that*

$$\left\| \left(z_1 \frac{\partial}{\partial z_1} \right)^2 F \right\|_{A(D^2)} \leq 1 \quad \text{and} \quad \left\| \left(z_2 \frac{\partial}{\partial z_2} \right)^2 F \right\|_{A(D^2)} \leq 1$$

but

$$\left\| z_2 \frac{\partial}{\partial z_2} \left(z_1 \frac{\partial}{\partial z_1} \right) F \right\|_{A(D^2)} \geq c.$$

Moreover, we provide the dependence of the above constant c on the degree of F (Theorem 3 and Corollary 4). A double logarithmic bound on the growth of the constant in Theorem 1 (without analyticity) could be derived from the construction of Curcă (cf.[3] sections 5.1 and 5.2);. Our construction not only gives analytic example but also provides (almost) logarithmic bound.

Theorem 3. *There exists an analytic polynomial P_n of degree n of two complex variables such that*

$$\left\| \left(z_1 \frac{\partial}{\partial z_1} \right)^2 P_n \right\|_{A(D^2)} \leq 1 \quad \text{and} \quad \left\| \left(z_2 \frac{\partial}{\partial z_2} \right)^2 P_n \right\|_{A(D^2)} \leq 1$$

but

$$\left\| z_2 \frac{\partial}{\partial z_2} \left(z_1 \frac{\partial}{\partial z_1} \right) P_n \right\|_{A(D^2)} \gtrsim \log^{1/8} n.$$

The similar quantitative result follows immediately on tori for boundary values of analytic functions.

Corollary 4. *There exists an analytic trigonometric polynomial P_n of degree n of two complex variables such that*

$$\|D_{xx}^2 f\|_\infty + \|D_{yy}^2 f\|_\infty \leq 1$$

but

$$\|D_{xy}^2 f\|_\infty \gtrsim \log^{1/8} n.$$

For some other problems related to quantitative estimates of the norm of derivatives from below see [15].

With a little effort we can replace the Euler derivatives from Theorem 2 by the ordinary derivatives (lower order derivatives appear here to compensate for the lack of compactly embedded support):

Theorem 5. *For every $c > 0$, there exists an analytic polynomial F on unit bi-disc such that*

$$\|F\|_{A(D^2)} \leq 1, \quad \left\| \frac{\partial}{\partial z_1} F \right\|_{A(D^2)} \leq 1, \quad \left\| \frac{\partial}{\partial z_2} F \right\|_{A(D^2)} \leq 1,$$

and

$$\left\| \frac{\partial^2}{\partial z_1^2} F \right\|_{A(D^2)} \leq 1, \quad \left\| \frac{\partial^2}{\partial z_2^2} F \right\|_{A(D^2)} \leq 1$$

but

$$\left\| \frac{\partial^2}{\partial z_1 \partial z_2} F \right\|_{A(D^2)} \geq c.$$

Despite we are mostly interested in the quantitative estimates, in the last section of the article we provide another purely qualitative proof.

Remark 6. Our qualitative method (Theorem 1, Section 3) extends to the general situation i.e. homogeneous differential operators of the same degree with constant coefficients $P_0(D), P_1(D), \dots, P_k(D)$ such that $P_0 \notin \text{span}\{P_1, \dots, P_k\}$. However when it comes to quantitative results (Theorem 3, Corollary 4) we only know how to extend them under some geometrical conditions as in [9, Theorem 2].

Motivated by Theorem 5 we propose the following definition.

Definition 7. Let $\Omega \subset \mathbb{C}^2$ and $z \in \partial\Omega$ be its boundary point. We say that z is a local MDM point for Ω provided there exists a ball B centered at z such that for every $c > 0$, there exists an analytic polynomial F on $\Omega \cap B$ such that

$$\|F\|_{A(\Omega \cap B)} \leq 1, \quad \left\| \frac{\partial}{\partial z_1} F \right\|_{A(\Omega \cap B)} \leq 1, \quad \left\| \frac{\partial}{\partial z_2} F \right\|_{A(\Omega \cap B)} \leq 1$$

and

$$\left\| \frac{\partial^2}{\partial z_1^2} F \right\|_{A(\Omega \cap B)} \leq 1, \quad \left\| \frac{\partial^2}{\partial z_2^2} F \right\|_{A(\Omega \cap B)} \leq 1$$

but

$$\left| \frac{\partial^2}{\partial z_1 \partial z_2} F(z) \right| \geq c.$$

Similarly for bounded domains Ω we define the global MDM point by replacing B by \mathbb{C}^2 in the above definition. Using this notion we can reformulate Theorem 5 saying that point $\mathbf{1} := (1, 1)$ is a (global) MDM point of the unit bi-disc (actually it is easy to see that any Shilov boundary point of a bi-disc is its MDM point). Note that MDM property is inherited by inclusion i.e. if $z \in \overline{\Omega} \subset \overline{\Omega'}$ is an MDM point of Ω' then z is an MDM point of Ω . There are some natural restrictions on $z \in \partial\Omega$ to be an MDM point. Indeed, suppose $z \in \partial\Omega$ is an internal point of $\hat{\Omega}$ - the polynomial convex hull of $\overline{\Omega}$. Then there exists a ball B centered at z contained in $\hat{\Omega}$. For every analytic polynomial f we have

$$\left| \frac{\partial^2}{\partial z_1 \partial z_2} f(z) \right| \leq C_B \sup_{\omega \in B} |f(\omega)| \leq C_B \sup_{\omega \in \Omega} |f(\omega)|.$$

The first inequality is a consequence of the Cauchy formula [14, 1.2.2. (v)] and second follows by definition of polynomial convex hull [6, p. 53]. We conjecture that above obstacle is the only one which prevents the MDM property. However even in the case of the unit ball of \mathbb{C}^2 we do not know whether any MDM point exists.

Acknowledgments We would like to thank Paul F.X. Müller and Fedor Nazarov for valuable comments and suggestions.

2 Proofs of quantitative results

Proof of Theorem 2. Let $(n_k) \subset \mathbb{Z}_+^2$ be a lacunary sequence which will be defined later. Let $(a_k)_{k=1}^n$ be a sequence of scalars. Define the sequence of Rudin - Shapiro polynomials:

$$p_0(z) = a_0 z^{n_0} \quad q_0 = 1$$

and for $n > 0$

$$\begin{aligned} p_n(z) &= p_{n-1}(z) + a_n z^{n_k} q_{n-1}(z^{-1}) \\ q_n(z) &= q_{n-1}(z) - \bar{a}_n z^{n_k} p_{n-1}(z^{-1}) \end{aligned}$$

(here we adopt the notation $z^{-1} = (z_1^{-1}, z_2^{-1})$ for $z = (z_1, z_2) \in \mathbb{C}^2$). By the simple induction, p_k 's and q_k 's are holomorphic polynomials. By the elementary identity valid for any $x, y, a \in \mathbb{C}$,

$$|x + ay|^2 + |y - \bar{a}x|^2 = (1 + |a|^2)(|x|^2 + |y|^2)$$

we get for any $|z_1| = |z_2| = 1$,

$$|p_n(z)|^2 + |q_n(\bar{z})|^2 = \prod_{k=0}^n (1 + |a_k|^2)$$

Indeed, this follows inductively since for $|z_1| = |z_2| = 1$,

$$\begin{aligned}
|p_k(z)|^2 + |q_k(\bar{z})|^2 &= |p_{k-1}(z) + a_n z^{n_k} q_{k-1}(z^{-1})|^2 \\
&\quad + |q_{k-1}(\bar{z}) - \bar{a}_n \bar{z}^{n_k} p_{k-1}(\bar{z}^{-1})|^2 \\
&= |p_{k-1}(z) + a_k z^{n_k} q_{k-1}(z^{-1})|^2 \\
&\quad + |z^{n_k} q_{k-1}(z^{-1}) - \bar{a}_k p_{k-1}(z)|^2 \\
&= (1 + |a_k|^2)(|p_k(z)|^2 + |q_k(\bar{z})|^2)
\end{aligned}$$

By the construction we get

$$p_n(z) = a_0 z^{n_0} + \sum_{k=1}^n a_k z^{n_k} q_{k-1}(z^{-1})$$

and

$$q_n(z) = 1 - \sum_{k=0}^n \bar{a}_k z^{n_k} p_{k-1}(z^{-1})$$

For any sequence of complex numbers $\{\omega_k\}_{k=1}^n$ there exists a subset $A \subset \{0, 1, 2, \dots, n\}$ such that

$$\left| \sum_{k \in A} \omega_k \right| \geq \frac{1}{\pi} \sum_{k=1}^n |\omega_k|$$

Note that the values $p_k(\mathbf{1})$ and $q_k(\mathbf{1})$ are independent on the choice sequence (n_k) . They only depend on the choice of the sequence $(a_k)_{k=1}^n$. For fixed sequence (a_k) we choose larger of two sums $\sum_{k=0}^n |a_k q_{k-1}(\mathbf{1})|$, $\sum_{k=1}^n |\bar{a}_k p_{k-1}(\mathbf{1})|$. Without loss of generality we assume it is the first one and we put $\omega_k = a_k q_{k-1}(\mathbf{1})$. Therefore there exists $A := A((a_k)) \subset \{0, 1, 2, \dots, n\}$ such that

$$\begin{aligned}
&\left| \sum_{k \in A} a_k q_{k-1}(\mathbf{1}) \right| + \left| \sum_{k \in A} \bar{a}_k p_{k-1}(\mathbf{1}) \right| \\
&\geq \frac{1}{2\pi} \left(\sum_{k=0}^n |a_k q_{k-1}(\mathbf{1})| + \sum_{k=1}^n |\bar{a}_k p_{k-1}(\mathbf{1})| \right) \\
&= \frac{1}{2\pi} \sum_{k=0}^n |a_k| (|q_{k-1}(\mathbf{1})| + |p_{k-1}(\mathbf{1})|) \\
&\geq \frac{1}{2\pi} \sum_{k=0}^n |a_k| (|q_{k-1}(\mathbf{1})|^2 + |p_{k-1}(\mathbf{1})|^2)^{\frac{1}{2}} \\
&\geq \frac{1}{2\pi} \sum_{k=1}^n |a_k| \prod_{j=0}^{k-1} (1 + |a_j|^2)^{1/2}
\end{aligned}$$

On the other hand, for $|z_1| = |z_2| = 1$

$$\begin{aligned}
&\left| \sum_{k \in A} a_k z^{n_k} q_{k-1}(z^{-1}) \right| + \left| \sum_{k \in A} \bar{a}_k \bar{z}^{n_k} p_{k-1}(\bar{z}^{-1}) \right| \\
&\leq \sum_{k=0}^n |a_k| \cdot |q_{k-1}(z^{-1})| + \sum_{k=0}^n |\bar{a}_k| \cdot |p_{k-1}(\bar{z}^{-1})| \\
&\leq \sum_{k=0}^n |a_k| \cdot (|q_{k-1}(z^{-1})| + |p_{k-1}(\bar{z}^{-1})|) \\
&\leq \sqrt{2} \sum_{k=1}^n |a_k| \prod_{j=0}^n (1 + |a_j|^2)^{1/2}
\end{aligned}$$

Put now $(z_1 \frac{\partial}{\partial z_1})^2 F = p_n$ and $(z_1 \frac{\partial}{\partial z_1})^2 G = q_n - 1$. Clearly

$$\begin{aligned} \max\{\|(z_1 \frac{\partial}{\partial z_1})^2 F\|_{A(D^2)}, \|(z_1 \frac{\partial}{\partial z_1})^2 G\|_{A(D^2)}\} &\leq \max\{\|p_n\|_{A(D^2)}, \|q_n\|_{A(D^2)}\} + 1 \\ &\leq \left(\sup_{z \in D^2} |p_n(z)|^2 + |q_n(z)|^2 \right)^{\frac{1}{2}} + 1 \\ &= \left(\sup_{|z_1|=|z_2|=1} |p_n(z_1, z_2)|^2 + |q_n(z_1, z_2)|^2 \right)^{\frac{1}{2}} + 1 \\ &\leq \prod_{n=1}^{\infty} (1 + a_n^2)^{\frac{1}{2}} + 1. \end{aligned} \quad (2)$$

Function p_k is a polynomial which is a sum of 2^k monomials of the form $b_M z^{m(M)}$, where $M = \{i_0, i_1, \dots, i_{2s}\} \subset \{0, 1, 2, \dots, n\}$ and $m(M) = \sum_{j=0}^{2s} (-1)^j n_{i_j}$ and $|b_M| = \prod_{j \in M} |a_{n_j}|$. Similarly q_k is a polynomial which is a sum of 1 and $2^k - 1$ monomials of the form $b_M z^{m(M)}$, where $M = \{i_0, i_1, \dots, i_{2s+1}\} \subset \{0, 1, 2, \dots, n\}$ and $m(M) = \sum_{j=0}^{2s+1} (-1)^{j+1} n_{i_j}$ and $|b_M| = \prod_{j \in M} |a_{n_j}|$. Therefore

$$z_2 \frac{\partial}{\partial z_2} (z_1 \frac{\partial}{\partial z_1}) F(z) = \sum_{2 \nmid \#M} b_M \frac{m(M)_2}{m(M)_1} z^{m(M)}$$

and

$$(z_2 \frac{\partial}{\partial z_2})^2 F(z) = \sum_{2 \nmid \#M} b_M \left(\frac{m(M)_2}{m(M)_1} \right)^2 z^{m(M)}$$

Similarly

$$z_2 \frac{\partial}{\partial z_2} (z_1 \frac{\partial}{\partial z_1}) G(z) = \sum_{2 \nmid \#M} b_M \frac{m(M)_2}{m(M)_1} z^{m(M)}$$

and

$$(z_2 \frac{\partial}{\partial z_2})^2 G(z) = \sum_{2 \nmid \#M} b_M \left(\frac{m(M)_2}{m(M)_1} \right)^2 z^{m(M)}$$

Suppose that the sequence (n_k) was chosen in such a way that for $0 < b < 1$

- (i) $\sum_{\max M \in A} \left| \frac{m(M)_2}{m(M)_1} - \kappa \right| < 1$
- (ii) $\sum_{\max M \in A} \left| \left(\frac{m(M)_2}{m(M)_1} \right)^2 - \kappa^2 \right| < 1$
- (iii) $\sum_{\max M \notin A} \left| \frac{m(M)_2}{m(M)_1} \right| < 1$
- (iv) $m(M) \in \mathbb{Z}_+^2$ for $M \subset \{0, 1, 2, \dots, n\}$

The above conditions are satisfied if the ratio $\frac{(n_k)_2}{(n_k)_1}$ is suitably chosen with respect to the fact $k \in A$ or not, and $\frac{|n_{k+1}|}{|n_k|}$ is big enough. Simple calculations show that one can choose $|n_k| \simeq 3^{\lambda n_k}$ for suitable λ cf. [9]. For this choice

$$\log \deg F \simeq \log \deg G \simeq n^2. \quad (3)$$

From now on we will assume that

$$|a_k| \leq 1$$

Therefore $|b_M| \leq 1$. Then, by the triangle inequality,

$$\begin{aligned}
& \left| z_2 \frac{\partial}{\partial z_2} \left(z_1 \frac{\partial}{\partial z_1} \right) F(\mathbf{1}) \right| + \left| z_2 \frac{\partial}{\partial z_2} \left(z_1 \frac{\partial}{\partial z_1} \right) G(\mathbf{1}) \right| \\
& \quad > \left| \kappa \sum_{k \in A} a_k q_{k-1}(\mathbf{1}) \right| + \left| \kappa \sum_{k \in A} \bar{a}_k p_{k-1}(\mathbf{1}) \right| \\
& \quad - \sum_{\max M \in A} |b_M| \cdot \left| \frac{m(M)_2}{m(M)_1} - \kappa \right| - \sum_{\max M \notin A} |b_M| \cdot \left| \frac{m(M)_2}{m(M)_1} \right| \\
& \quad \geq \frac{\kappa}{2\pi} \sum_{k=1}^n |a_k| \prod_{j=0}^{k-1} (1 + |a_j|)^{1/2} - 2
\end{aligned} \tag{4}$$

Moreover, for $|z_1| = |z_2| = 1$,

$$\begin{aligned}
& \left| \left(z_2 \frac{\partial}{\partial z_2} \right)^2 F(z) \right| + \left| \left(z_2 \frac{\partial}{\partial z_2} \right)^2 G(\bar{z}) \right| \\
& \quad < \left| \kappa^2 \sum_{k \in A} a_k z^{n_k} q_{k-1}(z) \right| + \left| \kappa^2 \sum_{k \in A} \bar{a}_k \bar{z}^{n_k} p_{k-1}(\bar{z}) \right| \\
& \quad + \sum_{\max M \in A} |b_M| \cdot \left| \frac{m(M)_2}{m(M)_1} - \kappa^2 \right| + \sum_{\max M \notin A} |b_M| \cdot \left| \frac{m(M)_2}{m(M)_1} \right| \\
& \quad \leq \sqrt{2} \kappa^2 \sum_{k=1}^n |a_k| \prod_{j=0}^{k-1} (1 + |a_j|)^{1/2} + 2
\end{aligned} \tag{5}$$

We put $a_k = k^{-\frac{1}{2}}$ and $\kappa = n^{-\frac{1}{4}}$. Substituting aforementioned values in (2) gives us

$$\begin{aligned}
\max \left\{ \left\| \left(z_1 \frac{\partial}{\partial z_1} \right)^2 F \right\|_{A(D^2)}, \left\| \left(z_1 \frac{\partial}{\partial z_1} \right)^2 G \right\|_{A(D^2)} \right\} & \leq \prod_{k=1}^n \left(1 + \frac{1}{k} \right)^{\frac{1}{2}} + 1 = \prod_{k=1}^n \left(\frac{k+1}{k} \right)^{\frac{1}{2}} + 1 \\
& = \sqrt{n+1} + 1.
\end{aligned}$$

By (5)

$$\begin{aligned}
\max \left\{ \left\| \left(z_2 \frac{\partial}{\partial z_2} \right)^2 F \right\|_{A(D^2)}, \left\| \left(z_2 \frac{\partial}{\partial z_2} \right)^2 G \right\|_{A(D^2)} \right\} & \leq \sqrt{2} n^{-\frac{1}{2}} \sum_{k=1}^n k^{-\frac{1}{2}} \prod_{j=0}^{k-1} \left(1 + \frac{1}{j} \right)^{1/2} + 2 \\
& \leq \sqrt{2n} + 2.
\end{aligned}$$

Finally by (4) for mixed derivative we get

$$\begin{aligned}
\max \left\{ \left\| z_2 \frac{\partial}{\partial z_2} \left(z_1 \frac{\partial}{\partial z_1} \right) F \right\|_{A(D^2)}, \left\| z_2 \frac{\partial}{\partial z_2} \left(z_1 \frac{\partial}{\partial z_1} \right) G \right\|_{A(D^2)} \right\} & \geq \frac{n^{\frac{1}{4}}}{2\pi} \sum_{k=1}^n k^{-\frac{1}{2}} \prod_{j=0}^{k-1} \left(1 + \frac{1}{j} \right)^{1/2} - 2 \\
& \geq \frac{n^{\frac{3}{4}}}{2\pi} - 2.
\end{aligned}$$

Hence for fixed n we get in Theorem 2

$$c = \left(\frac{1}{2\sqrt{2\pi}} + o(1) \right) n^{\frac{1}{4}}$$

□

Considering now either $F(z)$ or $G(z)$ and a sequence $\{n_k\}$ such that (i) – (iv) and (3) are satisfied we obtain Theorem 3.

Now we can deduce Theorem 5 from the proof of Theorem 2.

Proof of Theorem 5. From the construction $(n_k)_1 > 2^{k^2}$. Then

$$\begin{aligned} |F(z)| + |G(z)| &\leq \sum_{M \subset \{1, \dots, n\}} |b_M| \frac{1}{m(M)_1^2} \leq \sum_{k=1}^n \sum_{\max M=k} \frac{1}{(n_k)_1^2} \\ &\leq \sum_{k=1}^n \frac{2^k}{2^{2k^2}} < 2. \end{aligned}$$

Similarly

$$\begin{aligned} \left| z_1 \frac{\partial}{\partial z_1} F(z) \right| + \left| z_1 \frac{\partial}{\partial z_1} G(z) \right| &\leq \sum_{M \subset \{1, \dots, n\}} |b_M| \frac{1}{m(M)_1} \\ &\leq \sum_{k=1}^n \sum_{\max M=k} \frac{1}{|(n_k)_1|} \leq \sum_{k=1}^n \frac{2^k}{2^{k^2}} < 2. \end{aligned}$$

Since we may assume $\frac{m(M)_2}{m(M)_1} \leq 1$ we get

$$\begin{aligned} \left| z_2 \frac{\partial}{\partial z_2} F(z) \right| + \left| z_2 \frac{\partial}{\partial z_2} G(z) \right| &\leq \sum_{M \subset \{1, \dots, n\}} |b_M| \frac{m(M)_2}{m(M)_1^2} \\ &\leq \sum_{k=1}^n \sum_{\max M=k} \frac{1}{|(n_k)_1|} \leq \sum_{k=1}^n \frac{2^k}{2^{k^2}} < 2. \end{aligned}$$

Now since $\frac{\partial}{\partial z_2} F(z)$ is an analytic function we have following identity

$$\left\| \frac{\partial}{\partial z_2} F(z) \right\|_{\infty} = \sup_{|z_1|=|z_2|=1} \left| \frac{\partial}{\partial z_2} F(z) \right| = \sup_{|z_1|=|z_2|=1} \left| z_1 \frac{\partial}{\partial z_2} F(z) \right| < 2.$$

On the other hand for $j \in \{1, 2\}$

$$z_j \frac{\partial}{\partial z_j} \left(z_j \frac{\partial}{\partial z_j} \right) F(z) = z_j^2 \frac{\partial^2}{\partial z_j^2} F(z) + z_j \frac{\partial}{\partial z_j} F.$$

Once again because partial derivative is analytic we have

$$\begin{aligned} \left\| \frac{\partial^2}{\partial z_i^2} F(z) \right\|_{\infty} &= \sup_{|z_1|=|z_2|=1} \left| \frac{\partial^2}{\partial z_i^2} F(z) \right| = \sup_{|z_1|=|z_2|=1} \left| z_i^2 \frac{\partial^2}{\partial z_i^2} F(z) \right| \\ &\leq \left\| z_i \frac{\partial}{\partial z_i} \left(z_i \frac{\partial}{\partial z_i} \right) F \right\|_{\infty} + \left\| z_i \frac{\partial}{\partial z_i} F \right\|_{\infty} < 1 + 2 = 3. \end{aligned}$$

Finally for mixed derivatives we have

$$\begin{aligned} \left\| \frac{\partial}{\partial z_1} \left(\frac{\partial}{\partial z_2} \right) F \right\|_{\infty} &= \sup_{|z_1|=|z_2|=1} \left| \frac{\partial}{\partial z_1} \left(\frac{\partial}{\partial z_2} \right) F \right| \\ &= \sup_{|z_1|=|z_2|=1} \left| z_1 \frac{\partial}{\partial z_1} \left(z_2 \frac{\partial}{\partial z_2} \right) F \right| \\ &= \left\| z_1 \frac{\partial}{\partial z_1} \left(z_2 \frac{\partial}{\partial z_2} \right) F \right\|_{\infty}. \end{aligned}$$

By analogous arguments we obtain estimates for the function G . One of the functions F or G is the wanted polynomial. \square

3 Qualitative approach

In this section we give a qualitative proof of Theorem 1. It is simpler but in contrast to Theorem 3, does not provide any dependence of the constant on the degree of involved polynomials. It is based, as the proof by Mirkhil and DeLeeuw [4], on the Hahn-Banach theorem. The new ingredient is the use of Wiener's singularity criterion.

Qualitative proof of Theorem 1. Denote by $C_{\mathbb{Z}_+^2}^{\{\partial_1^2, \partial_2^2\}}(\mathbb{T}^2)$ the closure of analytic trigonometric polynomials of mean zero with respect to the norm

$$\|f\| = \|D_{xx}^2 f\|_\infty + \|D_{yy}^2 f\|_\infty$$

This space embeds isometrically as closed subspace into $C(\mathbb{T}^2, \mathbb{C}^2)$ by the formula

$$f \rightarrow (D_{xx}^2 f, D_{yy}^2 f)$$

Suppose that the assertion of Theorem 1 does not hold then the functional $\Lambda : C_{\mathbb{Z}_+^2}^{\{\partial_1^2, \partial_2^2\}}(\mathbb{T}^2) \rightarrow \mathbb{C}$ given by the formula

$$\Lambda(f) = D_{xy}^2 f(0, 0)$$

is continuous. By the Hahn-Banach theorem it extends to $C(\mathbb{T}^2, \mathbb{C}^2)$. Thus there exists a pair of bounded measures μ and ν such that

$$D_{xy}^2 f(0, 0) = \int_{\mathbb{T}} D_{xx}^2 f \, d\mu + \int_{\mathbb{T}} D_{xx}^2 f \, d\nu$$

Applying the above formula to characters $(t, s) \rightarrow e^{i(nt+ms)}$ we get

$$\frac{nm}{n^2 + m^2} = \frac{n^2}{n^2 + m^2} \hat{\mu}(n, m) + \frac{m^2}{n^2 + m^2} \hat{\nu}(n, m). \quad (6)$$

Let

$$K_r(n, m) = \{(k, l) \in \mathbb{Z}_+^2 : \max\{|n - k|, |m - l|\} < r\}.$$

It follows from (6) that

$$\begin{aligned} \hat{\mu}(k, l) + \hat{\nu}(k, l) &= 1 + o(1) && \text{for } (k, l) \in K_n(n^2, n^2), \\ \hat{\mu}(k, l) &= o(1) && \text{for } (k, l) \in K_n(n^2, n), \\ \hat{\nu}(k, l) &= o(1) && \text{for } (k, l) \in K_n(n, n^2). \end{aligned} \quad (7)$$

Then

$$\begin{aligned} A &= \liminf_{n \rightarrow \infty} \frac{1}{\#K_n(n^2, n^2)} \sum_{(k, l) \in K_n(n^2, n^2)} |\hat{\mu}(k, l)|^2 + |\hat{\nu}(k, l)|^2 \geq \frac{1}{4}, \\ B &= \lim_{n \rightarrow \infty} \frac{1}{\#K_n(n^2, n)} \sum_{(k, l) \in K_n(n^2, n)} |\hat{\mu}(k, l)|^2 = 0, \\ C &= \lim_{n \rightarrow \infty} \frac{1}{\#K_n(n, n^2)} \sum_{(k, l) \in K_n(n, n^2)} |\hat{\nu}(k, l)|^2 = 0. \end{aligned}$$

By Wiener's theorem on the discrete parts of measures [7, p. 45], A is a genuine limit and we get following contradiction

$$A = \sum_{\tau \in \mathbb{T}^2} |\mu(\{\tau\})|^2 + |\nu(\{\tau\})|^2 = \sum_{\tau \in \mathbb{T}^2} |\mu(\{\tau\})|^2 + \sum_{\tau \in \mathbb{T}^2} |\nu(\{\tau\})|^2 = B + C.$$

□

In the end we present yet another approach to Theorem 1, which could be used to indicate wider class of MDM points.

Proposition 8. *Let $\Omega \subset \mathbb{C}^2$ and $(z_0, w_0) \in \partial\Omega$ be such that*

1. *there are two different supporting hyperplanes L, K at (z_0, w_0) ;*
2. *$L \cap K$ is not a complex line;*
3. *$L \cap \overline{\Omega} = \{(z_0, w_0)\}$.*

Then (z_0, w_0) is an MDM point of Ω .

Proof. Since we have two supporting hyperplanes at point (z_0, w_0) there are two different affine functions $\gamma_i : \mathbb{C}^2 \rightarrow \mathbb{C}$ given by

$$\gamma_j(z, w) = x_j(z - z_0) + y_j(w - w_0), \quad j \in \{1, 2\}$$

such that

$$\operatorname{Re}(\gamma_j(z, w)) \geq 0 \quad \text{for } (z, w) \in \overline{\Omega}.$$

Note that if we have two such functions there is another one γ_3 with above properties (in fact there are infinitely many). Since $K \cap L$ is not a complex line we may assume that

$$x_j \neq 0, \quad y_j \neq 0, \quad \frac{x_i}{y_i} \neq \frac{x_j}{y_j} \quad \text{for } i \neq j, \quad i, j \in \{1, 2, 3\}. \quad (8)$$

Let us consider holomorphic functions

$$f_{j,N}(z, w) = N^{-2} e^{-N\gamma_j(z, w)}, \quad j = \{1, 2, 3\}.$$

We check that

$$\begin{aligned} \partial_z f_{j,N}(z, w) &= -N^{-1} x_j e^{-N\gamma_j(z, w)}, & \partial_z^2 f_{j,N}(z, w) &= x_j^2 e^{-N\gamma_j(z, w)}, \\ \partial_w f_{j,N}(z, w) &= -N^{-1} y_j e^{-N\gamma_j(z, w)}, & \partial_w^2 f_{j,N}(z, w) &= y_j^2 e^{-N\gamma_j(z, w)}, \end{aligned}$$

and

$$z \partial_z (w \partial_w) f_{j,N}(z, w) = x_j y_j e^{-N\gamma_j(z, w)}.$$

Therefore we get

$$\lim_{N \rightarrow \infty} \partial_z f_{j,N}^N(z, w) = \lim_{N \rightarrow \infty} \partial_w f_{j,N}^N(z, w) = \lim_{N \rightarrow \infty} f_{j,N}^N(z, w) = 0 \quad \forall (z, w) \in \overline{\Omega}$$

and

$$\lim_{N \rightarrow \infty} \partial_z^2 f_{j,N}^N(z, w) = \begin{cases} x_j^2 & (z, w) = (z_0, w_0); \\ 0 & (z, w) \neq (z_0, w_0); \end{cases} \quad \lim_{N \rightarrow \infty} \partial_w^2 f_{j,N}^N(z, w) = \begin{cases} y_j^2 & (z, w) = (z_0, w_0); \\ 0 & (z, w) \neq (z_0, w_0). \end{cases}$$

Similarly as in the case of bi-disc we obtain with the help of Hahn-Banach theorem, that if (z_0, w_0) is not an MDM point there are bounded Borel measures $\mu_1, \mu_2, \nu_1, \nu_2, \nu_3$ such that

$$\begin{aligned} \partial_z \partial_w f_{j,N}(z_0, w_0) &= \int_{\partial\Omega} \partial_z^2 f_{j,N}(z, w) \, d\mu_1 + \int_{\partial\Omega} \partial_w^2 f_{j,N}(z, w) \, d\mu_2 \\ &+ \int_{\partial\Omega} \partial_z f_{j,N}(z, w) \, d\nu_1 + \int_{\partial\Omega} \partial_w f_{j,N}(z, w) \, d\nu_2 + \int_{\partial\Omega} f_{j,N}(z, w) \, d\nu_3. \end{aligned}$$

Let $\mu_1 = \alpha \delta_{(z_0, w_0)} + \tilde{\mu}_1$ and $\mu_2 = \beta \delta_{(z_0, w_0)} + \tilde{\mu}_2$, where $\delta_{(z_0, w_0)} \perp \tilde{\mu}_1$ and $\delta_{(z_0, w_0)} \perp \tilde{\mu}_2$. By Lebesgue's dominated convergence theorem, for $j \in \{1, 2, 3\}$ we get

$$\begin{aligned} &\int_{\partial\Omega} \partial_z^2 f_{j,N}(z, w) \, d\mu_1 + \int_{\partial\Omega} \partial_w^2 f_{j,N}(z, w) \, d\mu_2 \\ &= \alpha \partial_z^2 f_{j,N}(z_0, w_0) + \int_{\partial\Omega} \partial_z^2 f_{j,N}(z, w) \, d\tilde{\mu}_1 \\ &+ \beta \partial_w^2 f_{j,N}(z_0, w_0) + \int_{\partial\Omega} \partial_w^2 f_{j,N}(z, w) \, d\tilde{\mu}_2 \xrightarrow{N \rightarrow \infty} \alpha x_j^2 + \beta y_j^2. \end{aligned}$$

and

$$\int_{\partial\Omega} \partial_z f_{j,N}(z, w) \, d\nu_1 + \int_{\partial\Omega} \partial_w f_{j,N}(z, w) \, d\nu_2 + \int_{\partial\Omega} f_{j,N}(z, w) \, d\nu_3 \xrightarrow{N \rightarrow \infty} 0$$

On the other hand

$$\lim_{N \rightarrow \infty} \partial_z \partial_w f_{i,N}(z_0, w_0) = x_j y_j.$$

Hence

$$x_j y_j = \alpha x_j^2 + \beta y_j^2 \quad \text{for } j \in \{1, 2, 3\}. \quad (9)$$

By (8) we have $p_j = x_j y_j^{-1} \neq 0$ and

$$1 - \alpha p_j - \beta p_j^{-1} = 0, \quad j \in \{1, 2, 3\} \quad (10)$$

Hence p_1, p_2, p_3 are roots of the quadratic equation $x - \alpha x^2 - \beta = 0$ which stands in contradiction with (8). Therefore (z_0, w_0) is an MDM point of Ω . \square

References

- [1] L. Carleson. Two remarks on H^1 and BMO. *Advances in Math.*, 22(3):269–277, 1976.
- [2] S. Conti, D. Faraco, and F. Maggi. A new approach to counterexamples to L^1 estimates: Korn’s inequality, geometric rigidity, and regularity for gradients of separately convex functions. *Archive for Rational Mechanics and Analysis*, 175(2):287–300, dec 2004.
- [3] E. Curcă. On the continuity of Fourier multipliers on $\dot{W}^{l,1}(\mathbb{R}^d)$ and $\dot{W}^{l,\infty}(\mathbb{R}^d)$. *Journal of Functional Analysis*, page 109573, 2022.
- [4] K. de Leeuw and H. Mirkil. A priori estimates for differential operators in L_∞ norm. *Illinois Journal of Mathematics*, 8(1), mar 1964.
- [5] R. A. Fefferman. Some topics from harmonic analysis and partial differential equations. In *Essays on Fourier analysis in honor of Elias M. Stein (Princeton, NJ, 1991)*, volume 42 of *Princeton Math. Ser.*, pages 175–210. Princeton Univ. Press, Princeton, NJ, 1995.
- [6] L. Hörmander. *An introduction to complex analysis in several variables*. D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto, Ont.-London, 1966.
- [7] Y. Katznelson. *An introduction to harmonic analysis*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, third edition, 2004.
- [8] K. Kazaniecki, D. M. Stolyarov, and M. Wojciechowski. Anisotropic ornstein noninequalities. *Analysis & PDE*, 10(2):351–366, feb 2017.
- [9] K. Kazaniecki and M. Wojciechowski. On Bernstein type quantitative estimates for Ornstein non-inequalities. <https://arxiv.org/abs/2206.13666>, 2022.
- [10] B. Kirchheim and J. Kristensen. On rank one convex functions that are homogeneous of degree one. *Archive for Rational Mechanics and Analysis*, 221(1):527–558, feb 2016.
- [11] D. V. Limanskiĭ and M. M. Malamud. On an analogue of the de Leeuw and Mirkil theorem for operators with variable coefficients. *Mat. Zametki*, 83(5):783–786, 2008.
- [12] B. S. Mityagin. On the second mixed derivative. *Dokl. Akad. Nauk SSSR*, 123:606–609, 1958.
- [13] D. Ornstein. A non-equality for differential operators in the L_1 norm. *Arch. Rational Mech. Anal.*, 11:40–49, 1962.
- [14] W. Rudin. *Function theory in the unit ball of \mathbb{C}^n* . Classics in Mathematics. Springer-Verlag, Berlin, 2008. Reprint of the 1980 edition.
- [15] S. A. Teljakovskiĭ. Bounds for the derivatives of trigonometric polynomials of several variables. *Sibirsk. Mat. Ž.*, 4:1404–1411, 1960.
- [16] M. Wojciechowski. A Marcinkiewicz type multiplier theorem for H^1 spaces on product domains. *Studia Math.*, 140(3):273–287, 2000.

Krystian Kazaniecki
Institute of Analysis, JKU Linz
Institute of Mathematics, University of Warsaw
krystian.kazaniecki@jku.at

Michał Wojciechowski
Institute of Mathematics, Polish Academy of Sciences
miwoj@impan.pl