

# Random models for singular SPDEs

I. BAILLEUL and Y. BRUNED

**Abstract.** We give a simple proof of the convergence of the BHZ renormalized model associated with the generalized (KPZ) equation. Our method pushes the spectral gap strategy for the construction of renormalized models to its natural limit in the form of Stroock formula. Then the convergence boils down to controlling Feynman diagrams via an extension of the Hairer-Quastel convergence theorem for models.

## 1 – Introduction

Let  $\zeta$  stand for a spacetime distribution of negative regularity. Random or not we talk about it as a ‘noise’. We call *singular*, an elliptic or parabolic semilinear partial differential equation (PDE) with real-valued unknown  $u$ , an equation of the form

$$\mathcal{L}u = f(u, \nabla u; \zeta) \quad (1.1)$$

where a formal application of Schauder estimates do not give sufficient regularity to  $u$  for the term  $f(u, \nabla u; \zeta)$  to make sense. Recall indeed that loosely speaking the product of a distribution of regularity  $a$  with a function of regularity  $b$  is well-defined if and only if  $a + b > 0$ . A typical example of a singular PDE is provided by the two dimensional parabolic Anderson model equation

$$(\partial_t - \Delta)u = u\xi,$$

where  $\xi$  stands for the random realization of a space white noise on the two dimensional torus. The distribution  $\xi$  has almost surely Hölder regularity  $-1 - \varepsilon$  for all  $\varepsilon > 0$ ; this gives  $u$  an a priori Hölder regularity  $1 - \varepsilon$  which is not sufficient to make sense of the product  $u\xi$ . Coupled systems of singular PDEs are characterized by the same product problem. Such equations/systems typically appear as the probabilistic large scale limit of random microscopic equations/systems where the strength of the nonlinearity and the randomness balance each other. The study of singular stochastic PDEs was launched by the groundbreaking works of M. Hairer on regularity structures [16] and Gubinelli, Imkeller & Perkowski on paracontrolled calculus [15]. While [16] and [15] use different tools to build an equation-dependent setting where to make sense of it and provide a robust solution theory for a large family of equations, both works share the same guidelines.

1. Work in a restricted space of potential solutions characterized by the fact that they ‘look like’ some linear combination of reference objects that depend only on the noise  $\zeta$ . The choice of language used to give sense to the expression ‘look like’ distinguishes regularity structures from paracontrolled calculus.
2. Point 1 brings back the initial problem of making sense of the ill-defined products involving  $u$  in (1.1) to the problem of giving sense to some ill-defined products of quantities that only involve the noise  $\zeta$ , typically some polynomial expression  $P(\zeta, \dots, \zeta)$  of  $\zeta$ . This is where working with *random* distributions  $\zeta$  saves the day. While the random variable  $\omega \mapsto P(\zeta(\omega), \dots, \zeta(\omega))$  does not make sense one can construct in a consistent way a *random variable*, suggestively denoted here  $\omega \mapsto P(\zeta, \dots, \zeta)(\omega)$ , that plays the role of  $P(\zeta(\omega), \dots, \zeta(\omega))$ . This is what *renormalization* is about and the collection  $\widehat{\zeta}(\omega)$  of all the random variables  $P(\zeta, \dots, \zeta)(\omega)$  needed in the analysis of a given equation is called the *enhanced noise*. On a technical level the above polynomial expressions of the noise come under the form of *models over a regularity structure*.
3. Replacing the formal quantities  $P(\zeta(\omega), \dots, \zeta(\omega))$  by the renormalized quantities

$$P(\zeta, \dots, \zeta)(\omega)$$

in a naive formulation of the equation turns out to provide a well-defined equation that can be solved uniquely in an appropriate function space. We talk of a ‘renormalized equation’ and its solution.

4. Of course a proper interpretation of the solution of the renormalized equation is needed. In the end one can prove that it is the limit, in a probabilistic sense, of solutions to equations of the form

$$\mathcal{L}u^\varepsilon = f(u^\varepsilon, \nabla u^\varepsilon; \zeta^\varepsilon) + g^\varepsilon(u^\varepsilon)$$

in which  $\zeta$  has been regularized into  $\zeta^\varepsilon$  and some  $\varepsilon$ -dependent, a priori diverging, counterterm  $g^\varepsilon(\cdot)$  has been added into the dynamics. The parameter  $\varepsilon > 0$  goes to 0 in the limit.

The analytic machinery introduced by Hairer in [16] was complemented in subsequent joint works with Bruned & Zambotti [8], Chandra [11] and Bruned, Chandra & Chevyrev [6] to provide a robust solution theory for a large class of (systems of) equations. The work [8] provided a systematic study of the algebraic setting needed to analyse singular stochastic PDEs from a regularity structures point of view. The work [11] proved the convergence of a general algorithm used to construct the random variables  $P(\zeta, \dots, \zeta)(\omega)$  as limits in some appropriate spaces of quantities of the form

$$P(\zeta^\varepsilon(\omega), \dots, \zeta^\varepsilon(\omega)) = \sum_Q c_Q^\varepsilon Q(\zeta^\varepsilon(\omega), \dots, \zeta^\varepsilon(\omega))$$

for some finite sums of polynomial expressions  $Q$  of  $\zeta^\varepsilon$  and  $\varepsilon$ -dependent, a priori diverging, constants  $c_Q^\varepsilon$ . We call *BHZ renormalization* this construction, after the initials of the authors of the work [8]. (Amusingly these initials are very close to the initials BPHZ used to name the renormalization algorithm of Feynman graphs introduced by Bogoliubov & Parasiuk in the 50s, and clarified later by some works of Hepp and Zimmermann. The BHZ renormalization algorithm is an elaboration of the BPHZ renormalization prescription on different types of objects.) The work [6] proved point 4 above. In the end for any given equation of a well-identified class of subcritical singular elliptic/parabolic semilinear stochastic PDEs there exists, under a small parameter condition, a unique solution to that equation in an appropriate equation-dependent function space. This solution depends continuously on the realization  $\widehat{\zeta}(\omega)$  of the random enhanced noise. To contrast things, note that  $\widehat{\zeta}(\omega)$  itself is only a measurable function of  $\zeta(\omega)$ . We refer the reader to the overviews [12, 13] of Chandra & Weber and Corwin & Shen for non-technical introductions to the domain of semilinear singular SPDEs, to the books [14, 3] of Friz & Hairer and Berglund for a mildly technical introduction to regularity structures, and to Bailleul & Hoshino's *Tourist's Guide* [2] for a thorough tour of the analytic and algebraic sides of the theory. Our previous work [1] gives a short proof of a generalization of point 4, and some aspects of the renormalization problem are described in Hairer's lecture notes [17].

The fantastic work [11] of Chandra & Hairer on the convergence of the BHZ renormalized models is fairly difficult. It holds under a very general set of conditions on the law of the translation invariant (possibly multi-dimensional) noise that involve its cumulants. Recently Linares, Otto, Tempelmayr & Tsatsoulis [20] devised a different approach to the renormalization problem within the setting devised by Otto, Sauer, Smith & Weber [21] for the study of a class of quasilinear singular stochastic PDEs. While the latter shares many features with the setting of regularity structures devised for the study of semilinear singular equations the setting introduced in [21] is different; it involves in particular no tree indexed objects even though its construction is inductive. More importantly for us they prove their convergence result for the renormalized regularized models assuming a spectral gap inequality for the law of the random noise rather than making assumptions on its cumulants. We push this idea to its natural limit and give in the present work a simple and short proof of the convergence of the BHZ renormalized model used in a regularity structures analysis of the iconic generalized (KPZ) equation.

Denote by  $M^\varepsilon = (\Pi^\varepsilon, \mathbf{g}^\varepsilon)$  the naive admissible model built from the regularized noise  $\zeta^\varepsilon$ , and denote by  $\overline{M}^\varepsilon = (\overline{\Pi}^\varepsilon, \overline{\mathbf{g}}^\varepsilon)$  the BHZ renormalized naive admissible model built from  $\zeta^\varepsilon$ . (All the objects will be properly introduced in Section 2.1.) On a technical side, the convergence of  $\overline{\Pi}^\varepsilon$  is encoded in inequalities of the form

$$\|\langle \overline{\Pi}_z^\varepsilon \tau - \overline{\Pi}_z^{\varepsilon'} \tau, \varphi_z^\lambda \rangle\|_{L^2(\Omega)} \lesssim o_{\varepsilon \vee \varepsilon'}(1) \lambda^{|\tau|}, \quad (1.2)$$

for all symbols  $\tau$  of the regularity structure with negative homogeneity  $|\tau|$ . Think here of  $\varphi_z^\lambda$  as a regular  $\lambda$ -approximation of a Dirac distribution at the point  $z$  – proper notations will be introduced

below. Linares, Otto, Tempelmayr & Tsatsoulis take profit from their spectral gap assumption to bound the  $L^2(\Omega)$  norm of the quantity of interest

$$\|\langle \bar{\Pi}_z^\varepsilon \tau - \bar{\Pi}_z^{\varepsilon'} \tau, \varphi_z^\lambda \rangle\|_{L^2(\Omega)} \leq |\mathbb{E}[\langle \bar{\Pi}_z^\varepsilon \tau - \bar{\Pi}_z^{\varepsilon'} \tau, \varphi_z^\lambda \rangle]| + \|\langle d(\bar{\Pi}_z^\varepsilon \tau - \bar{\Pi}_z^{\varepsilon'} \tau), \varphi_z^\lambda \rangle\|_{L^2(\Omega)}$$

by its expectation and the norm of its Malliavin derivative. In the setting of [20] the control of the above expectation comes at low cost from their choice of renormalization procedure. The control of the  $L^2$ -norm of the derivative is way more involved and done inductively by seeing first  $d\bar{\Pi}^\varepsilon$  as part of an extended model. (The very recent work [22] of Otto, Seong and Tempelmayr gives a reader's guide digest of [20].)

We follow in the present work the insight of [20] and push the spectral gap strategy to its natural limit in the form of *Stroock formula*. For a symbol  $\tau$  built from  $|\tau|_\zeta$  noise symbols one has

$$\|\langle \bar{\Pi}_z^\varepsilon \tau - \bar{\Pi}_z^{\varepsilon'} \tau, \varphi_z^\lambda \rangle\|_{L^2(\Omega)} \leq \sum_{k=0}^{|\tau|_\zeta - 1} \|\mathbb{E}[\langle d^k(\bar{\Pi}_z^\varepsilon \tau - \bar{\Pi}_z^{\varepsilon'} \tau), \varphi_z^\lambda \rangle]\| + \|\langle d^{|\tau|_\zeta}(\bar{\Pi}_z^\varepsilon \tau - \bar{\Pi}_z^{\varepsilon'} \tau), \varphi_z^\lambda \rangle\|. \quad (1.3)$$

The norms  $\|\cdot\|$  are Hilbert-Schmidt type norms. Note that the  $|\tau|_\zeta$ -th derivative term in the right hand side is not random anymore. As in [20] the control of the size of the terms given by the expectations comes at low cost from the renormalization prescription, here the BHZ renormalization procedure from [8]. Note that

$$d^{|\tau|_\zeta}(\bar{\Pi}_z^\varepsilon \tau - \bar{\Pi}_z^{\varepsilon'} \tau) = d^{|\tau|_\zeta}(\Pi_z^\varepsilon \tau - \Pi_z^{\varepsilon'} \tau).$$

The inductive structures of the trees  $\tau$  and the naive model  $\Pi^\varepsilon$  allow to control recursively the norm of the corresponding term in the above estimate. We treat the BPHZ renormalization by hand in our study of the generalized (KPZ) equation; this is due to the fact that we only have to deal with one subdivergence involving two noises. We treat these divergences drawing ideas from [4, 9] where a renormalization by telescopic sums has been proposed.

We describe in Section 2 the generalized (KPZ) equation, its associated regularity structure and the BHZ renormalized model. We prove its convergence in Section 3. The useful notion of mirror graph is introduced in Section 2.2 and we talk in Section 2.3 of Malliavin derivatives of trees and their expectation as this is what appears in Stroock's formula (1.3). The proof of convergence of the BHZ model to a limit model is done in Section 3.2; its main ingredient is recalled in Section 3.1.

**Notations** – We denote by  $z = (t, x)$  a generic spacetime point, with parabolic norm

$$\|z\|_{\mathfrak{s}} := \sqrt{|t| + |x|^2}.$$

We will freely use all the notions and tools of regularity structures. Denote by  $H$  the Hilbert space  $L^2([0, 1] \times \mathbb{T})$ . All random variables below will be defined on a fixed probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ .

## 2 – The generalized (KPZ) equation

Let  $\mathbb{T}$  stand for the one dimension torus and  $\zeta$  stand for a spacetime white noise on  $\mathbb{R} \times \mathbb{T}$ . It has almost surely parabolic Hölder regularity  $-3/2 - \kappa$ , for every  $\kappa > 0$ . We fix  $\kappa$  once and for all. The generalized (KPZ) equation has unknown a function  $u : [0, T] \times \mathbb{T} \rightarrow \mathbb{R}$  and reads

$$(\partial_t - \partial_x^2)u = f(u)\zeta + g(u)(\partial_x u)^2, \quad (2.1)$$

for  $C^4$  bounded real valued functions  $f, g$  with bounded derivatives. The time horizon  $T$  is not fixed and is part of the problem. As we expect from Schauder estimates that  $u$  has parabolic regularity  $1/2 - \kappa$  the products  $f(u)\zeta$  and  $(\partial_x u)^2$  are not well-defined, let alone  $g(u)(\partial_x u)^2$ .

Recall here for later use Hairer's notation

$$\|K\|_{a,m} := \sup_{|k|_{\mathfrak{s}} \leq m} \sup_z \|z\|_{\mathfrak{s}}^{|k|_{\mathfrak{s}} - a} |\partial^k K(z)|$$

for the size of a kernel with a singularity at 0. Here  $|k|_{\mathfrak{s}}$  is defined as

$$|k|_{\mathfrak{s}} = |(k_0, k_1)|_{\mathfrak{s}} := 2k_0 + k_1.$$

The heat kernel  $P$  has a finite  $\|\cdot\|_{1,m}$  size for any fixed  $m \geq 0$ . Pick a smooth non-increasing function  $\chi : [0, \infty) \rightarrow [0, 1]$  equal to 1 on  $[0, 1/2]$  and 0 on  $(1, \infty)$ . Rather than working with the heat kernel  $P$  we work with

$$K := (P - c)\chi,$$

with the constant  $c$  is chosen for  $K$  to have null mean – a useful technical property that does not destroy the behaviour of  $K$  as  $z$  approaches the spacetime point 0. The corresponding two convolutions operators coincide up to a smoothing operator and  $\|K\|_{1,m} < \infty$ .

## 2.1 The regularity structure and the BHZ renormalized model

We describe in this section the regularity structure associated with the generalized (KPZ) equation and the BHZ renormalized model. A reader familiar with the theory can skip this section.

**The regularity structure** – Let  $\mathcal{I}$  and  $\mathcal{I}_{(0,1)}$  stand for abstract ‘integration’ symbols; they are formal placeholders for the convolution operator with  $K$  and its space derivative. Let  $\Xi$  stand for the symbol playing in the regularity structure the role of  $\zeta$ . The regularity structure associated with the generalized (KPZ) equation (2.1) is generated by the following rule  $\mathcal{R}$  that describes the local structure of its (decorated) trees. Write  $[\mathcal{I}]_\ell$  for an  $\ell$ -tuple of operators  $\mathcal{I}$ . One has

$$\mathcal{R}(\Xi) = \{\emptyset\}, \quad \mathcal{R}(\mathcal{I}) = \mathcal{R}(\mathcal{I}_{(0,1)}) = \left\{ [\mathcal{I}]_\ell, ([\mathcal{I}]_\ell, \mathcal{I}_{(0,1)}), ([\mathcal{I}]_\ell, \mathcal{I}_{(0,1)}, \mathcal{I}_{(0,1)}), (\mathcal{I}_{(0,1)}, \Xi) \right\}_{\ell \in \mathbb{N}}$$

Polynomial decorations ( $k = (k_0, k_1) \in \mathbb{N}^2$ ) are added on the nodes of these trees:  $k_0$  is for time monomial and  $k_1$  is for spatial monomial. We denote by  $\mathcal{B}$  the family of all (polynomially) decorated trees satisfying the rule  $\mathcal{R}$ . The empty tree is added to  $\mathcal{B}$  and denoted by  $\mathbf{1}$ . To define the homogeneity  $|\tau|$  of an element  $\tau \in \mathcal{B}$  we set

$$|\mathbf{1}| := 0, \quad |\Xi| := -3/2 - \kappa,$$

and define inductively for  $k \in \mathbb{N}^2, \sigma, \tau \in \mathcal{B}$


$$|X^k \tau| := |\tau| + |k|_{\mathfrak{s}}, \quad |\mathcal{I}(\tau)| := |\tau| + 2, \quad |\mathcal{I}_{(0,1)}(\tau)| := |\tau| + 1, \quad |\sigma\tau| := |\sigma| + |\tau|.$$

One has  $X^k = X_0^{k_0} X_1^{k_1}$  where the  $X_i$  are identified with the canonical basis of  $\mathbb{N}^2$ . In the sequel, we will write  $X$  instead of  $X_1$ . Call ‘noises’ the elements of  $\mathcal{B}$  of negative homogeneity and denote their collection by  $\mathcal{B}^-$ . Recall that one can build inductively the set of noises using the operators


$$\begin{aligned} \mu &\mapsto \mathcal{I}_{(0,1)}(\mu), \\ (\mu, \nu) &\mapsto \mu\mathcal{I}(\nu), \\ (\mu, \nu) &\mapsto \mathcal{I}_{(0,1)}(\mu)\mathcal{I}_{(0,1)}(\nu), \end{aligned} \tag{2.2}$$

starting from the initial collection of noises  $\{\Xi, X\Xi\}$ . Write  $\circ$  for  $\Xi$  and use the blue symbol  $\bullet$  for the product symbol  $X\circ$ . With a plain line for the operator  $\mathcal{I}$  and a dotted line for the operator  $\mathcal{I}_{(0,1)}$  we can list the elements of  $\mathcal{B}^-$  according to their homogeneity.


- Elements of  $\mathcal{B}^-$  of homogeneity  $-1 - 2\kappa$




- Elements of  $\mathcal{B}^-$  of homogeneity  $-1/2 - 3\kappa$



- Elements of  $\mathcal{B}^-$  of homogeneity  $-4\kappa$



- Elements of  $\mathcal{B}^-$  of homogeneity  $-2\kappa$



For  $\tau \in T$  denote by  $|\tau|_\zeta$  the number of noises in  $\tau$ . We have  $|\tau|_\zeta \leq 4$  for the trees of  $\mathcal{B}$  with  $|\tau| < 0$ . Denote by  $((T, \Delta), (T^+, \Delta^+))$  and  $((T, \delta), (T^-, \delta^-), (T^+, \delta^+))$  the BHZ regularity and renormalization structures associated to that equation – see e.g. [17, 2].

**The BHZ model** – Let  $\zeta^\varepsilon$  stand for a regularization of  $\zeta$  by convolution with a smooth function of the form

$$\rho^\varepsilon(z) = \varepsilon^{-3} \rho(\varepsilon^{-2}t, \varepsilon^{-1}x),$$

with  $\rho \geq 0$  of unit integral and equal to 1 in 0. Denote by  $M^\varepsilon = (\Pi^\varepsilon, \mathbf{g}^\varepsilon)$  the naive admissible model on the BHZ regularity structure of the generalized (KPZ) equation associated with the smooth noise  $\zeta^\varepsilon$ . Given a character  $\ell$  on the algebra  $T^-$  generated by the trees of  $T$  of non-positive homogeneity the maps

$$\tilde{\ell} = (\ell \otimes \text{Id})\delta : T \rightarrow T, \quad \tilde{\ell}^+ = (\ell \otimes \text{Id})\delta^+$$

turn the pair

$$\overline{M}^\varepsilon := (\Pi^\varepsilon \circ \tilde{\ell}, \mathbf{g}^\varepsilon \circ \tilde{\ell}^+)$$

into a model; this is part of Theorem 6.15 in [8] – see also Theorem 2.6 in Section 5 of [2]. Bruned, Hairer & Zambotti proved in Theorem 6.15 of [8] the existence of a unique character  $\ell^\varepsilon$  on  $T^-$  such that  $\overline{M}^\varepsilon$  is admissible and satisfies for all  $\tau \in \mathcal{B}^-$  and all  $y$  the identity

$$\mathbb{E}[(\overline{M}^\varepsilon \tau)(y)] = 0. \quad (2.3)$$

The map  $\overline{M}^\varepsilon$  can be described using a preparation map  $R^\varepsilon$  defined from a splitting map  $\delta_r$ , with the index ‘ $r$ ’ for ‘root’, defined similarly as the splitting map  $\delta$  but extracting from any  $\tau \in T$  only one diverging subtree of  $\tau$  with the same root as  $\tau$  at a time, and summing over all possible such subtrees – see Definition 4.2 in [5]. One can also provide a recursive definition of this map by first defining

$$\begin{aligned} \delta'_r(\bullet) &:= \bullet \otimes \mathbf{1} + \mathbf{1} \otimes \bullet, \quad \text{for } \bullet \in \{\mathbf{1}, X_i, \Xi\}, \\ \delta'_r(\mathcal{I}_a \tau) &:= (\mathcal{I}_a \otimes \text{Id})\delta'_r \tau + \sum_{k \in \mathbb{N}^2} \frac{X^k}{k!} \otimes \mathcal{I}_{a+k}(\tau). \end{aligned}$$

where the  $\mathcal{I}_a$  are abstract integration symbol encoding the convolution operator with  $D^a K$ . We define also an abstract derivative  $\mathcal{D}^a$  satisfying the Leibniz rule and such that:

$$\mathcal{D}^a \mathcal{I}_b(\tau) = \mathcal{I}_{a+b}(\tau), \quad \mathcal{D}^a X^k = X^{k-a}.$$

This derivative corresponds to  $D^a$ . In the sequel, we will denote  $\mathcal{D}^a$  by  $\mathcal{D}$ . The infinite sum in the definition of  $\delta'_r$  makes perfect sense using a bigraduation introduced in [8, Section 2.3]. Then  $\delta_r$  is obtained by applying the projection  $\pi_-$  that keeps only decorated trees with negative homogeneity

$$\delta_r := (\pi_- \otimes \text{Id})\delta'_r.$$

Using Sweedler’s notations

$$\delta_r \tau = \tau \otimes \mathbf{1} + \sum_{(\tau)} \tau^{(1)} \otimes \tau^{(2)},$$

the  $\tau^{(1)}$  have strictly less edges than  $\tau$  due to the extraction of a subtree of negative homogeneity. Given any character  $\ell$  on the algebra  $T^-$  set

$$R = R_\ell := (\ell(\cdot) \otimes \text{Id})\delta_r \quad (2.4)$$

with  $\Pi^R$  defined by

$$\Pi^R \Xi = \Pi^{R, \times} \Xi = \xi^\varepsilon,$$

and the maps  $\Pi^R$  and  $\Pi^{R, \times}$  defined inductively by the relations

$$\Pi^R = \Pi^{R, \times} R, \quad \Pi^{R, \times}(\tau\sigma) = (\Pi^{R, \times} \tau)(\Pi^{R, \times} \sigma), \quad \Pi^{R, \times}(\mathcal{I}_a \tau) = D^a K \star (\Pi^R \tau).$$

It follows from this definition and the fact that  $R$  leaves fixed the elements of  $T$  of the form  $\mathcal{I}_a \tau$  that the map  $\Pi^R$  satisfies the admissibility condition

$$\Pi^R(\mathcal{I}_a \tau) = D^a K \star (\Pi^R \tau).$$


We recall the definition of the space  $T^+$  which is the linear span of




Its formal limit as  $\varepsilon$  goes to 0 would be a simple integral of the form

$$\int \varphi(y)^2 \mathcal{M}^0(\tau)(y, y) dy,$$

due to the  $\delta$  correlation of spacetime white noise. We use here the generic notation  $\mathcal{M}$  for ‘mirror’. Here are two other examples of tree representations of test operations together with their mirror graphs.



Here again the two diagrams on the right hand side represent a double integral of the form (2.5). More generally mirror diagrams are defined in the obvious way from any connected finite graphs built with the bricks  $\circ, \bullet, \bullet$  and the solid and dotted edges. We talk of a *tree-like graph* if it corresponds to the quantity obtained by testing a polynomial function of the regularized noise against a test function  $\varphi$ . We call it ‘tree-like’ rather than ‘tree’ as is associated analytic object does not necessarily have a tree structure. As an example the following quantities are tree-like graphs




One can define more generally tree-like graphs where each edge represents a kernel that may depend not only on the two integration variables associated with the vertices of the edge but also on the other integration variables. This is what happens when we replace a kernel by a Taylor remainder of that kernel based at some point different from the two vertices variables. The introduction here of the mirror graphs is motivated by the following useful observation from Section 10 of Hairer’s seminal work [16].

**Lemma 1** – For a tree-like graph  $\tau$  one has


$$\mathbb{E}[(\int \varphi(y) dy)^2] \leq \iint \varphi(y)\varphi(y') \mathcal{M}^\varepsilon(\tau)(y, y') dy dy'.$$

### 2.3 Malliavin derivatives of trees and their expectation


For  $0 \leq k \leq |\tau|_\zeta - 1$  denote by  $d^k \overline{\Pi}_x^\varepsilon \tau$  the  $k$ th order Malliavin derivative of  $\overline{\Pi}_x^\varepsilon \tau$ ; this is an element of  $L(H^{\otimes k}, L^2(\Omega, \mathbb{P}))$ . On the example of the ‘integrated’ tree




here is a piece of  $\langle (d\Pi^\varepsilon \tau)(h), \varphi \rangle$



As  $\zeta$  has the same law as  $-\zeta$  only quantities with an even number of noises have possibly non-null expectation. The first Malliavin derivative of a 4-linear functions of the noise being 3-linear it has null expectation. Here are pieces of  $\langle (d^2 \Pi^\varepsilon \tau)(h_1, h_2), \varphi \rangle$  for the preceding tree



We are interested in the expectation of such quantities, which produce elements of the space of Hilbert-Schmidt operators on  $H^{\otimes k}$  that are tree-like graph maps where the former noise arguments  $\circ, \bullet$  from Section 2.2 are now elements of  $H$ . We will use a common notation  $\|\cdot\|$  for the norms on these spaces. The square norm of a tree-like graph map is bounded above by its corresponding mirror diagram, where the pairing of an element of  $H$  with its mirror element produces a  $\bullet$  vertex in the mirror diagram. So while the mirror diagrams of trees involve only the convolution operator  $\bullet$  in addition to the convolution operator with  $K$  and  $\partial_x K$ , the mirror diagrams of a derivative tree also involve the convolution operator  $\bullet$ . As an example here are pieces of  $\mathbb{E}[\langle (d^2 \Pi^\varepsilon \tau)(h_1, h_2), \varphi \rangle]$ , for the same tree  $\tau$  as above



The loop in the third diagram is a constant annihilated by the kernel  $K$ , so the diagram represents the null function. We see on that example that looking at  $\mathbb{E}[\langle (d^2 \bar{\Pi}^\varepsilon \tau)(h_1, h_2), \varphi \rangle]$  for the BHZ renormalized model  $\bar{\Pi}^\varepsilon$  removes exactly the diverging contribution of the first graph in the sum giving  $\mathbb{E}[\langle (d^2 \Pi^\varepsilon \tau)(h_1, h_2), \varphi \rangle]$ ; the other terms are converging as the regularization parameter  $\varepsilon$  is sent to 0.

**Noise derivative operators** – We formalize the noise differentiation at the level of the regularity structure by introducing derivative operators on trees that replace a noise with new noise symbols  $\Xi_j$  with homogeneity equal to

$$|\Xi_j| = -\kappa.$$

In the context of the generalized (KPZ) equation we are interested in stochastic iterated integrals having at most four noises, so the index  $j$  will belong to the finite set  $\{1, 2, 3, 4\}$ . We define the derivative operators  $D_{\Xi_j}$  by

$$\begin{aligned} D_{\Xi_j} \bullet &:= 0, \quad \text{for } \bullet \in \{\mathbf{1}; X_i; \Xi; \Xi_\ell, 1 \leq \ell \leq 4\}, \\ D_{\Xi_j}(\tau \tau') &= (D_{\Xi_j} \tau) \tau' + \tau (D_{\Xi_j} \tau'), \\ D_{\Xi_j} \mathcal{I}_a(\tau) &= \mathcal{I}_a(D_{\Xi_j} \tau). \end{aligned}$$

We define  $\mathcal{B}_j^-$  as the decorated trees obtained by applying the iterated derivative operator  $D_{\Xi_j} \cdots D_{\Xi_1}$  to an element of  $\mathcal{B}^-$ . As no element of  $\mathcal{B}_j^-$  has a subtree of negative homogeneity containing a special noise  $\Xi_j$  one has

$$\delta_r D_{\Xi_j} = (\text{Id} \otimes D_{\Xi_j}) \delta_r,$$

which implies that

$$RD_{\Xi_j} = D_{\Xi_j} R.$$

We extend the definition of the pre-model and the model to  $\mathcal{B}_j^-$  by associating to each noise  $\Xi_j$  a variable  $h_j \in H$ . For the pre-model we set

$$(\Pi^{R, \times} \Xi_j)(h_1, \dots, h_4) = (\Pi^{R, \times} \Xi_j)(h_1, \dots, h_4) = h_j,$$

For the model we use a different homogeneity  $|\cdot|_+$  that assigns the same number to  $\Xi_j$  as to  $\Xi$

$$|\Xi_j|_+ := -\frac{3}{2} - \kappa,$$

and is defined as  $|\cdot|$  on the rest of the decorated trees. For  $\tau \in \mathcal{B}_j^-$ , one has

$$\begin{aligned} (\Pi_z^{R, \times} (\mathcal{I}_a \tau))(y, h_1, \dots, h_j) &= (D^a K \star \Pi_z^R \tau)(y, h_1, \dots, h_j) \\ &- \sum_{|k| \leq |\mathcal{I}_a \tau|_+} \frac{(y-z)^k}{k!} (D^{a+k} K \star \Pi_z^R \tau)(z, h_1, \dots, h_j). \end{aligned} \quad (2.6)$$



Note the difference with the formula giving  $\Pi_z^{R,\times}(\mathcal{I}_a\tau)(y)$ .

Except in the trivial case where  $\tau$  has only one noise, a *derivative tree-like graph map*

$$\mathbb{E}[\langle d^k \Pi^\varepsilon \tau, \varphi \rangle] \quad \text{or} \quad \mathbb{E}[\langle d^k \bar{\Pi}^\varepsilon \tau, \varphi \rangle]$$

with  $k \geq 1$  is the sum of several tree-like graph maps. We will always bound the norm of the sum by the sum of the norms.

**Lemma 2** – For  $\tau \in \mathcal{B}^-$ ,  $0 \leq k \leq |\tau|_\zeta$  and all  $z$  one has

$$d^k \bar{\Pi}_z^\varepsilon \tau = \bar{\Pi}_z^\varepsilon (D_{\Xi_k} \dots D_{\Xi_1} \tau).$$

**Proof** – One can proceed by induction on  $k$  and use [10, Prop 4.1] that says  $d \bar{\Pi}_z^\varepsilon \tau = \bar{\Pi}_z^\varepsilon (D_{\Xi_1} \tau)$ .  $\triangleright$

The extension of the character  $\ell$  to the regularity structure built for the generalized (KPZ) with the five noises  $\Xi, \Xi_1, \dots, \Xi_4$  retains its expression. Since we only take expectation in the analytic object with respect to the  $\Xi$  noise any counterterm involving a  $\Xi_i$  noise is actually null.

### 3 – Convergence of the BHZ model

We prove in this section the convergence of the BHZ model after recalling a use criterion to get scaling bounds on mirror graphs. For a test function  $\varphi$  write  $\varphi_z$  for  $\varphi(\cdot - z)$  and  $\varphi_z^\lambda$  for  $\lambda^{-3} \varphi(\mathfrak{s}(\lambda)(\cdot - z))$ .

#### 3.1 A workhorse

We will use as a workhorse a refinement due to Bruned & Nadeem [9] of a result of Hairer & Quastel [19] giving some scaling bounds on Feynman graphs – Theorem A.3 therein. We will use it in our setting to deal with the mirror graphs that come from estimating the terms  $\|\mathbb{E}[\langle d^k \bar{\Pi}_z^\varepsilon \tau, \varphi_z^\lambda \rangle]\|$ . We restrict its statement to our setting, in which our graphs always have two green vertices, with each of them attached to the remainder of the graph by one green edge that represents integration against a scaled test function  $\varphi^\lambda$ . Except from the green edges each edge  $e = (e_+, e_-)$  in our graphs represents a kernel of the form

$$L_e(z_{e_-}, z_{e_+}) = K_e(z_{e_+} - z_{e_-}) - \sum_{|j|_s < r_e} \frac{(z_{e_+} - z_{v_e})^j}{j!} \partial^j K_e(z_{v_e} - z_{e_-}),$$

a Taylor remainder of the function  $K_e(\cdot - z_{e_-})$  around an edge-dependent point  $z_{v_e}$ . Note the strict inequality  $< r_e$ , so there is no Taylor expansion at all if  $r_e = 0$ . We will have

$$r_e \in \{0, 1, 2\},$$

so at most a first order Taylor remainder, and our edges will be either  $K$ ,  $\partial K$  or  $\rho^\varepsilon \star \rho^\varepsilon$ . We note that we always have  $r_e = 0$  for edges for which  $K_e = \rho^\varepsilon \star \rho^\varepsilon$ . We assign the green integration edge a number  $a_e = 0$  and set

$$a_e := 1 \text{ if } K_e = K, \text{ and } a_e := -2 \text{ if } K_e = \partial K, \text{ and } a_e := -3 \text{ if } K_e = \rho^\varepsilon \star \rho^\varepsilon.$$

This is the scaling degree of divergence of the kernel  $K_e$  near 0. For a set  $V$  of vertices in our graph we define  $E_{\text{int}}(V)$  as the set of edges  $e$  in the graph with  $e_+$  and  $e_-$  in  $V$ . We define  $E^\uparrow(V)$ , resp.  $E^\downarrow(V)$  as the set of edges of the graph with  $e_- \in V$ , resp.  $e_+ \in V$ . Last we define  $E_{r>0}^\uparrow(V)$  as the set of edges  $e$  of  $E^\uparrow(V)$  such that  $r_e > 0$ . The following two conditions jointly define **Condition (C)**.

- For every subset  $V$  of vertices of our graph with at least two elements we have

$$\sum_{e \in E_{\text{int}}(V)} a_e + \sum_{e \in E_{r>0}^\uparrow(V)} \mathbf{1}_{v_e \in V} (a_e + r_e - 1) - \sum_{e \in E^\downarrow(V)} \mathbf{1}_{v_e \in V} r_e < 3(|V| - 1). \quad (3.1)$$

- For every subset  $V$  of vertices of our graph not containing the green vertices one has

$$\sum_{e \in E_{\text{int}}(V)} a_e + \sum_{e \in E^\downarrow(V)} (\mathbf{1}_{\{v_e \in V\} \cup \{r_e=0\}} (a_e + r_e - 1) - (r_e - 1)) + \sum_{e \in E^\uparrow(V)} (a_e + r_e \mathbf{1}_{v_e \notin V}) > 3|V|. \quad (3.2)$$

The number 3 in the two inequalities is the scaling dimension of the one dimensional spacetime. We denote by  $\mathcal{E}$  the set of non-green edges of our graph  $G$  and write

$$\mathcal{J}^G(\lambda) := \int \prod_{e \in \mathcal{E}} L_e(z_{e_-}, z_{e_+}) \varphi^\lambda(z_1) \varphi^\lambda(z_2) dz$$

with  $dz$  a shorthand notation for all the integration variables. We denote by  $V_0$  the set of non-green vertices of  $G$ .

**Theorem 3** – *If our graph  $G$  satisfies assumption **(C)** then there exists a constant  $c$  depending only on  $G$*

$$|\mathcal{J}^G(\lambda)| \leq c \lambda^\alpha \prod_{e \in \mathcal{E}} \|K_e\|_{a_e, 1}$$

where  $\alpha = 3|V_0| - \sum_{e \in \mathcal{E}} a_e$ .

The condition (3.1) checks the integrability of the Feynman diagram. It can be understood as proving the integrability on all subsets of variables by selecting a subset  $V$  of vertices. It is also an equivalent of the power counting given by [23] on Feynman diagrams. One can see easily that the diagrams that may fail this bound are coming from trees of negative homogeneity. The second condition (3.2) allows to get the correct scaling of  $\lambda$  in the end. This is the reason why the nodes in green which are already scaled in  $\lambda$  are excluded from the subset  $V$ .

For a spacetime point  $z$  denote by  $\mathfrak{T}_z$  the translation operator

$$h \in H \mapsto h(\cdot + z) \in H$$

on  $H$ ; this is an isometry. As one has

$$\langle d^k \bar{\Pi}_z^\varepsilon, \varphi_z^\lambda \rangle(h_1, \dots, h_k) = \langle d^k \bar{\Pi}_0^\varepsilon, \varphi_0^\lambda \rangle(\mathfrak{T}_z h_1, \dots, \mathfrak{T}_z h_k)$$

the mirror graph of  $\mathbb{E}[\langle d^k \bar{\Pi}_0^\varepsilon, \varphi_0^\lambda \rangle]$  provides a  $z$ -independent upper bound for the norm of

$$\mathbb{E}[\langle d^k \bar{\Pi}_z^\varepsilon, \varphi_z^\lambda \rangle].$$

It is then for instance elementary to check that the mirror graph associated to  $\langle d^{|\tau|_\zeta} \Pi_z^\varepsilon \tau, \varphi_z^\lambda \rangle$  satisfies condition **(C)** for every  $\tau \in \mathcal{B}^-$ . One first notices that

$$d^{|\tau|_\zeta} \bar{\Pi}_z^\varepsilon \tau = d^{|\tau|_\zeta} \Pi_z^\varepsilon \tau,$$

a consequence of the fact that  $\bar{\Pi}_z^\varepsilon \tau - \Pi_z^\varepsilon \tau$  has a number of noises strictly smaller than  $|\tau|_\zeta$ . One gets as a direct consequence of Theorem 3 the following result.

**Corollary 4** – *One has for all  $0 < \lambda \leq 1$  and every  $\tau \in \mathcal{B}^-$  the estimates*

$$\|\langle d^{|\tau|_\zeta} \Pi_z^\varepsilon \tau, \varphi_z^\lambda \rangle\| \lesssim \lambda^{|\tau|},$$

for an implicit multiplicative constant independent of  $\varepsilon > 0$  and  $z$ .

**Proof** – The condition (3.1) is satisfied due to the fact one does not see any subdivergence in a full mirror graph without contraction. For the second bound (3.2) one can proceed by induction on the construction of the diagram as in Section 4 in [9]. The result is straightforward for a product of diagrams. (By product of diagrams we mean identifying the green vertices of two graphs  $G_1$  and  $G_2$ .) The most difficult part is the integration of a diagram that is the adjunction of new green vertices that are connected to the old ones (which are no longer green vertices). The new edges have their node  $v_e$  equal to zero and the  $r_e$  is bigger to the homogeneity of the half diagram used for constructing the mirror graph. This operation correspond to the recentering operation in the model – see (2.6). One observes that the correct choice of  $r_e$  allows to get the bound (3.2). The proof here is easier than the proof in [9] as there are no  $\rho^\varepsilon \star \rho^\varepsilon$  edges in the complete mirror graph. The quantity  $\alpha = 3|V_0| - \sum_{e \in \mathcal{E}} a_e$  is indeed equal to  $|\tau|$  in that case.  $\triangleright$

### 3.2 Proof of convergence

**Proposition 5** – *There exists a positive finite constant  $C$  such that one has for all  $\tau \in \mathcal{B}^-$  and  $0 \leq k \leq |\tau|_\zeta - 1$ , and all test functions  $\varphi$  of a given regularity and support in the parabolic ball of*

radius  $1/2$ , the bound

$$\|\mathbb{E}[\langle d^k \bar{\Pi}_z^\varepsilon \tau, \varphi_z^\lambda \rangle]\| \leq C \lambda^{|\tau|}. \quad (3.3)$$

**Proof** – Pick  $\tau \in \mathcal{B}^-$ . Note first that the  $k = 3$  case is trivial as it concerns only the trees with four noises, in which case the third Malliavin derivative of  $\bar{\Pi}_z^\varepsilon \tau$  has only one noise and null expectation. We are thus left with the  $0 \leq k \leq 2$  cases. We consider the different cases in three steps. Recall that

$$(\bar{\Pi}_z^\varepsilon \tau)(y) = \sum_{\sigma \leq \tau} (\bar{\mathfrak{g}}_z^\varepsilon)^{-1}(\tau/\sigma) (\bar{\Pi}^\varepsilon \sigma)(y). \quad (3.4)$$

(a) *Case  $\sigma = \tau$  and  $k = 0$ .* In the sum (3.4), the term  $\sigma = \tau$  has null mean from the fundamental property (2.3) of the BHZ renormalization map.

(b) *Case  $1 < \sigma < \tau$ .* Both  $\tau/\sigma$  and  $\sigma$  contain at least one noise in that case. The terms that need to be controlled are given by

$$\sum_{1 < \sigma < \tau} \mathbb{E} \left[ (\bar{\mathfrak{g}}_z^\varepsilon)^{-1}(\tau/\sigma) (\bar{\Pi}^\varepsilon \sigma)(y) \right]. \quad (3.5)$$

When one considers a tree  $\tau$  with four noises it means that  $\tau/\sigma$  and  $\sigma$  can contain at most three noises. In this case the only possible subtrees of negative homogeneity with at least two noises are

$$\begin{array}{c} \circ \diagdown \circ \\ \circ \diagup \circ \end{array}, \begin{array}{c} \circ \\ | \\ \circ \end{array}, \begin{array}{c} \circ \diagdown \circ \\ | \\ \circ \end{array}, \begin{array}{c} \circ \diagup \circ \\ | \\ \circ \end{array}, \begin{array}{c} \circ \diagdown \circ \\ \circ \diagup \circ \end{array}. \quad (3.6)$$

The first two terms are of homogeneity  $(-1)^-$  whereas the last three terms are of homogeneity  $0^-$ . They appear as subtrees of some other trees  $\sigma$  with other branches attached to them for trees  $\sigma$  of the form

$$\tau_1 \Xi \mathcal{I}(\Xi \tau_2), \quad \mathcal{I}_1(\Xi \tau_1) \mathcal{I}_1(\Xi \tau_2), \quad \mathcal{I}(\Xi \tau_1) \mathcal{I}_1(\Xi \tau_2), \quad \Xi \mathcal{I}(\tau_1 \mathcal{I}_1(\Xi \tau_2)), \quad \mathcal{I}_1(\Xi) \mathcal{I}_1(\tau_1 \mathcal{I}_1(\Xi \tau_2)) \quad (3.7)$$

where  $\tau_1$  or  $\tau_2$  could be the empty tree. The tree  $\tau_1 \tau_2$  contains at most one noise. Note that the diverging subtrees of (3.6) have at most one incoming edge when seen as a subset of  $\sigma$ .

We briefly explain in a situation where  $\sigma = \mathcal{I}_1(\Xi) \mathcal{I}_1(\Xi \mathcal{I}(\Xi))$  and  $\tau/\sigma = \mathcal{I}(\Xi)$  how such a renormalization works via a mere telescopic sum. The other cases are similar and left to the reader. We start by considering  $(\bar{\mathfrak{g}}_z^\varepsilon)^{-1}(\tau/\sigma) (\bar{\Pi}^\varepsilon \sigma)(y)$  instead of  $(\bar{\mathfrak{g}}_z^\varepsilon)^{-1}(\tau/\sigma) (\bar{\Pi}_z^\varepsilon \sigma)(y)$ . We will see below how the second term builds upon the first. After applying the expectation operator on  $(\bar{\mathfrak{g}}_z^\varepsilon)^{-1}(\tau/\sigma) (\bar{\Pi}^\varepsilon \sigma)(y)$  one gets, among various diagrams, a diagram of the form

$$\begin{array}{c} G \\ | \\ \text{---} \\ \diagdown \\ \circ \\ | \\ \text{---} \\ \diagup \\ \circ \\ | \\ y \end{array} \quad (3.8)$$

where  $G$  is a convergent diagram corresponding to the pairing of  $\tau/\sigma$  with the deeper noise  $\Xi$  in  $\sigma$ . Let mention here that the other diagrams are fine as they will not contain a subgraph that fails condition 3.1; we come back on that point later in the proof. To make the following explanations more visual it will be useful to trade our notation  $\bullet$  for the spacetime convolution operator with  $\rho^\varepsilon \star \rho^\varepsilon$  for a **purple edge**. The point  $z$  is the integration variable associated to the root of the subtree with negative homogeneity. The subdivergent diagram is given by

$$\begin{array}{c} \text{---} \\ \diagdown \\ \circ \\ | \\ \text{---} \\ \diagup \\ \circ \\ | \\ y \end{array} = \int K'(y - z_1) K'(y - z_2) (\rho^\varepsilon \star \rho^\varepsilon)(z_1 - z_2) dz_1 dz_2.$$

We compensate the subdivergence in (3.8) by a **Taylor expansion** on some incoming edge. We use a telescopic sum to make appear this renormalisation

$$\begin{array}{c} G \\ | \\ \text{---} \\ \diagdown \\ \circ \\ | \\ \text{---} \\ \diagup \\ \circ \\ | \\ y \end{array} = \begin{array}{c} G \\ | \\ \text{---} \\ \diagdown \\ \circ \\ | \\ \text{---} \\ \diagup \\ \circ \\ | \\ y \end{array} + \begin{array}{c} G \\ | \\ \text{---} \\ \diagdown \\ \circ \\ | \\ \text{---} \\ \diagup \\ \circ \\ | \\ y \end{array} + \begin{array}{c} G \\ | \\ \text{---} \\ \diagdown \\ \circ \\ | \\ \text{---} \\ \diagup \\ \circ \\ | \\ y \end{array} + \begin{array}{c} G \\ | \\ \text{---} \\ \diagdown \\ \circ \\ | \\ \text{---} \\ \diagup \\ \circ \\ | \\ y \end{array} \quad (3.9)$$

The first term is well defined as the decoration  $(z, 2)$  means a kernel of the form

$$K(z_1 - z_3) - K(y - z_3) - (x_1 - x) \partial K(y - z_3)$$

that behaves like  $(x_1 - x)^2$  when  $z_1$  is close to  $y = (t, x)$ . For the first term of (3.9) one can check the integrability condition (3.1) or apply alternatively Lemma 10.14 of [16] on the spacetime convolution of singular kernels with singularity at 0 of the form  $\|\cdot\|_s^{-\alpha}$ . In the third term of (3.9) the orange line is encoding a term of the form  $(x_1 - x)$

$$\begin{array}{c} \text{---} \\ \diagdown \\ \circ \\ | \\ \text{---} \\ \diagup \\ \circ \\ | \\ y \end{array} = \int (x_1 - x) \partial K(y - z_1) \partial K(y - z_2) (\rho^\varepsilon \star \rho^\varepsilon)(z_1 - z_2) dz_1 dz_2.$$

Now the two problematic terms are the last two terms of (3.9). They are removed by the renormalization. Indeed one has

$$R(y) \sigma = \sigma + \ell(\mathcal{I}_1(\Xi)^2)(y) \mathcal{I}(\Xi) + \ell(\mathcal{I}_1(\Xi) \mathcal{I}_1(X \Xi))(y) \mathcal{I}_1(\Xi)$$

by using the definition of  $R$  given by (2.4). Then we conclude by noticing that

$$\ell(\mathcal{I}_1(\Xi)^2)(y) = - \begin{array}{c} \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ y \end{array}, \quad \ell(\mathcal{I}_1(\Xi)\mathcal{I}_1(X\Xi))(y) = - \begin{array}{c} \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ y \end{array}$$

(We kept track of the point  $y$  in the above computations to stress that the formulation is robust in the non-translation invariant case. In this case we obtain renormalization functions instead of renormalization constants. We could have removed the dependency in  $y$ .) One can observe that the telescopic sums are a local procedure that detaches incoming edges from a diverging subgraph associated in fact to a subtree of negative homogeneity. These edges are therefore attached to the root of the diverging subtree and then we obtain a diverging constant that could be detached if one has translation-invariance. Moreover one has also the same changes in the decorations provided by the order of the telescopic sum. One gets in the end a one to one correspondance with the procedure of extraction/contraction of a subtree given by  $R$ . This has been previously observed in [9].

It remains to check that the diagrams have the good scaling in  $\lambda$  by looking at the correct quantity which is  $(\bar{\mathfrak{g}}_z^\varepsilon)^{-1}(\tau/\sigma)(\bar{\Pi}_z^\varepsilon\sigma)(\varphi_z^\lambda)$ . We can perform the same telescopic sum with a priori recentering on some of the edges. One has a new telescopic sum given by

$$K(z_1 - z_3) - K(z - z_3) = K(z_1 - z_3) - K(z_0 - z_3) - (x_1 - x_0)\partial K(z_0 - z_3) \\ + K(z_1 - z_3) - K(z - z_3) + (x_1 - x_0)\partial K(z_0 - z_3)$$

where  $z_0 = (t_0, x_0)$ . Now (3.9) becomes

$$\begin{array}{c} G \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ z_0 \end{array}^{(z,1)} = \begin{array}{c} G \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ z_0 \end{array}^{(z_0,2)} + \begin{array}{c} G \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ z_0 \end{array}^{(z,1)} + \begin{array}{c} G \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ z_0 \end{array}^{(z,1)} \quad (3.10)$$

where  $z_0$  is a green variable coming from  $\varphi_z^\lambda$ . From Corollary 4, one can deduce that the right hand side satisfies (3.2) but may a priori fail (3.1) due to sub graphs coming from subtrees of negative homogeneity. From Proposition 5.7 in [9] one sees that recentering bounds are preserved for this transformation. Then, this implies by Proposition 5.4 in [9] that the bounds are also satisfied for the other terms of (3.11) if one includes the correct recentering which is coming from (3.4). Indeed in the telescopic sum we can move along the way the recentering – see beginning of Section 5 in [9]. The crucial point is to have  $r' \geq r$  which always happens in the diagrams we have to treat. The second term above is well recentered with the same length of the Taylor expansion as the right hand side. The third term does not need any recentering due to the extra derivative added.

We let apart a number of diagrams to concentrate on diagram that have the form (3.8). For other trees, one may have some recentering inside the sub divergent graph. In that case, it is removed at the end in order to make appear the renormalisation constant. The condition (3.2) is easily satisfied on the extra terms produced – see Proposition 5.10 in [9]. One moves from  $(\bar{\mathfrak{g}}_z^\varepsilon)^{-1}(\tau/\sigma)(\bar{\Pi}_z^\varepsilon\sigma)(\varphi_z^\lambda)$  to  $(\bar{\mathfrak{g}}_z^\varepsilon)^{-1}(\tau/\sigma)(\bar{\Pi}_z^\varepsilon\sigma)(\varphi_z^\lambda)$  by collecting the correct recentering in (3.5).

Such reasoning works the same when we apply one or two Malliavin derivatives as they commute with the model by replacing noises of type  $\Xi$  by  $\Xi_1$  and  $\Xi_2$ , from Lemma 2.

(c) *Case  $k = 1$ .* One looks at trees with three noises and the divergence are of same nature as in the case  $k = 0$  and  $\tau \neq \sigma$ . The only difference is that we have to work with a mirror graph as a noise  $\Xi$  has been replaced by its Malliavin derivative.

(d) *Case  $k = 2$ .* The trees in the regularity structure which have diverging subtrees in  $\mathcal{B}^-$  have the same form as in (3.7) but now  $\tau_1\tau_2$  is at most a product of two planted trees as we consider only trees with four noises. Note that these diverging diagrams have at most two incoming edges.

We now explain on the example of the tree  $\mathcal{I}_1(\Xi\tau_1)\mathcal{I}_1(\Xi\tau_2)$  how such a renormalization works via a mere telescopic sum. One deals with the other terms of the list (3.7) in exactly the same way. Indeed one gets the following diagram for  $\mathbb{E}[(\Pi^\varepsilon\mathcal{I}_1(\Xi\tau_1)\mathcal{I}_1(\Xi\tau_2))(y)]$

$$\begin{array}{c} h_1 \quad h_2 \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ y \end{array}$$

where here, without loss of generality, we have supposed that  $\tau_1 = \mathcal{I}(\Xi_1)$  and  $\tau_2 = \mathcal{I}(\Xi_2)$ . We recognise the same subdivergent diagram but this time, we have two incoming edges. We proceed with several telescopic sums as below

$$\begin{array}{c} h_1 \quad h_2 \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ y \end{array} = \begin{array}{c} h_1 \quad h_2 \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ y \end{array}^{(z_2,2)} + \begin{array}{c} h_1 \quad h_2 \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ y \end{array} + \begin{array}{c} h_1 \quad h_2 \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ y \end{array} \quad (3.11)$$

The first term is well-defined as the edge decoration  $(z_2, 2)$  means a kernel of the form

$$K(z_1 - z_3) - K(z_2 - z_3) - (x_1 - x_2)\partial K(z_2 - z_3)$$

that behaves like  $(x_1 - x_2)^2$  when  $z_1$  is close to  $z_2$ . This is sufficient for making sense of the first term in (3.11) when the  $\varepsilon$ -regularization is removed. More precisely, one can check the bounds (3.1). The **orange line** is encoding a multiplicative term of the form  $(x_2 - x_1)$

$$\text{orange line} = \int (x_2 - x_1)K'(y - z_1)K'(y - z_2)(\rho^\varepsilon \star \rho^\varepsilon)(z_1 - z_2)dz_2dz_1.$$

Note that this term is equal to zero when the mollifier  $\rho$  is symmetric. The last two terms in (3.11) are still ill-defined and one has to move the two edges to the root. One gets

$$\begin{array}{c} h_1 \quad h_2 \\ \diagdown \quad \diagup \\ (y, 2) \\ \diagup \quad \diagdown \\ y \end{array} = \begin{array}{c} h_1 \quad h_2 \\ \diagdown \quad \diagup \\ (y, 2) \\ \diagup \quad \diagdown \\ y \end{array} + h_1 \begin{array}{c} h_2 \\ \diagup \\ y \end{array} + h_1 \begin{array}{c} h_2 \\ \diagdown \\ y \end{array}$$

Then we apply the same reduction to the last term and obtain

$$\begin{array}{c} h_2 \\ \diagup \\ h_1 \text{---} y \end{array} = \begin{array}{c} h_2 \\ \diagup \\ (y, 1) \\ \diagdown \\ h_1 \text{---} y \end{array} + \begin{array}{c} h_1 \quad h_2 \\ \diagdown \quad \diagup \\ y \end{array}$$

Repeating the operation, one has in the end

$$\begin{array}{c} h_1 \quad h_2 \\ \diagdown \quad \diagup \\ y \end{array} = (\dots) + \begin{array}{c} h_1 \quad h_2 \\ \diagdown \quad \diagup \\ y \end{array} + \begin{array}{c} h_1 \quad h_2 \\ \diagdown \quad \diagup \\ y \end{array} + \begin{array}{c} h_1 \quad h_2 \\ \diagdown \quad \diagup \\ y \end{array}$$

The terms  $(\dots)$  are well-defined with the correct Taylor expansions added. The other terms correspond exactly to the counter-terms that are removed by the BPHZ renormalization. Indeed if we apply a 'local product'  $R(y)$  depending possibly on the spacetime point one has

$$R(y)(\mathcal{I}_1(\Xi\tau_1)\mathcal{I}_1(\Xi\tau_2)) = \mathcal{I}_1(\Xi\tau_1)\mathcal{I}_1(\Xi\tau_2) + \ell(\mathcal{I}_1(\Xi)\mathcal{I}_1(\Xi))(y)\tau_1\tau_2 + \ell(\mathcal{I}_1(\Xi X)\mathcal{I}_1(\Xi))(y)\mathcal{D}(\tau_1\tau_2).$$

where

$$\ell(\mathcal{I}_1(\Xi)\mathcal{I}_1(\Xi))(y) = - \begin{array}{c} h_1 \quad h_2 \\ \diagdown \quad \diagup \\ y \end{array}, \quad \ell(\mathcal{I}_1(\Xi X)\mathcal{I}_1(\Xi))(y) = - \begin{array}{c} h_1 \quad h_2 \\ \diagdown \quad \diagup \\ y \end{array}$$

Extraction elsewhere in the trees is not possible as  $\tau_1$  and  $\tau_2$  may contain only the noises  $\Xi_1$  and  $\Xi_2$ . For controlling the quantity  $\|\mathbb{E}[\langle \overline{\Pi}_z^\varepsilon \mathcal{I}_1(\Xi\tau_1)\mathcal{I}_1(\Xi\tau_2), \varphi_z^\lambda \rangle]\|$  one has to check that the recentering bounds are not modified by the telescopic sum. This is verified in Proposition 5.4 and 5.7 in [9].  $\triangleright$

The reader may wonder why we did not directly moved the edges toward the base point  $z$ . Actually the blue line corresponding to the mollifier can be understood as being as singular as a Dirac mass at a first approximation and therefore it is a sub-divergence of order 0. By adding the red line it becomes well-defined and if there is no incoming edge on the node associated with the variable  $z$  one can perform the convolution and say that  $K \star \rho^\varepsilon \star \rho^\varepsilon$  has the same behaviour as  $K$ .

Given the results of Corollary 4 and Proposition 5 the following fact follows as a consequence of Stroock formula

$$\|\langle \overline{\Pi}_z^\varepsilon \tau, \varphi_z^\lambda \rangle\|_{L^2(\Omega)} \leq \sum_{k=0}^{|\tau|_\zeta - 1} \|\mathbb{E}[\langle d^k \overline{\Pi}_z^\varepsilon \tau, \varphi_z^\lambda \rangle]\| + \|\langle d^{|\tau|_\zeta} \overline{\Pi}_z^\varepsilon \tau, \varphi_z^\lambda \rangle\|.$$

**Corollary 6** – One has for all  $0 < \lambda \leq 1$  and every  $\tau \in \mathcal{B}^-$  the estimates

$$\|\langle \overline{\Pi}_z^\varepsilon \tau, \varphi_z^\lambda \rangle\|_{L^2(\Omega)} \lesssim \lambda^{|\tau|}, \quad (3.12)$$

for an implicit multiplicative constant independent of  $\varepsilon$ .

We know from Theorem 10.7 of [16] that the convergence of the BHZ model to a limit admissible model follows from estimates of the form

$$\|\langle \overline{\Pi}_z^\varepsilon \tau - \overline{\Pi}_z^{\varepsilon'} \tau, \varphi_z^\lambda \rangle\|_{L^2(\Omega)} \lesssim o_{\varepsilon \vee \varepsilon'}(1) \lambda^{|\tau|}, \quad (3.13)$$

for  $\tau \in \mathcal{B}^-$ . To prove these bounds from the previous bounds we note that the scaling bound (3.3) from Proposition 5 holds if the kernel  $K$  behaved not as  $|z|_s^{-1}$ , as  $z$  goes to 0, but as  $|z|_s^{-1-\eta}$ , for

$\eta > 0$  small enough. Recall from Proposition 10.17 of [16] the bound

$$\|\rho^\varepsilon \star \rho^\varepsilon - \rho^{\varepsilon'} \star \rho^{\varepsilon'}\|_{3+\eta,1} \lesssim \max(\varepsilon, \varepsilon')^\eta. \quad (3.14)$$

As in Section 5.2 of Hairer & Pardoux work [18] one can decompose  $\overline{\Pi}_z^\varepsilon \tau - \overline{\Pi}_z^{\varepsilon'}$  as a telescopic sum where we replace one after the other the  $\zeta^\varepsilon$  by the  $\zeta^{\varepsilon'}$ . This amounts to replacing some convolution operators with  $\rho^\varepsilon$  by the corresponding convolution operator with  $\rho^{\varepsilon'}$ , so the corresponding term in the telescopic sum involves the difference of two terms that differ at only one place. We gain the  $o_{\varepsilon \vee \varepsilon'}(1)$  term in (3.13) from using the estimate (3.14) for this edge while repeating the proof of (3.12). We leave the details to the reader.

We note that it is straightforward to adapt the present work to prove the convergence of the BHZ model associated with a system of generalized (KPZ) equations, such as the system describing the random motion of a rubber on a manifold [7].

## References

- [1] I. Bailleul and Y. Bruned, *Locality for singular stochastic PDEs*. arXiv:2109.00399, (2021).
- [2] I. Bailleul and M. Hoshino, *A tourist guide to regularity structures and singular stochastic PDEs*. arXiv:2006.03525, (2020).
- [3] N. Berglund, *An Introduction to Singular Stochastic PDEs. Allen-Cahn Equations, Metastability, and Regularity Structures*. EMS Series of Lectures in Mathematics, (2022).
- [4] Y. Bruned. *Singular KPZ Type Equations*. 205 pages, PhD thesis, Université Pierre et Marie Curie - Paris VI, (2015). <https://tel.archives-ouvertes.fr/tel-01306427>.
- [5] Y. Bruned, *Recursive formulae for regularity structures*. Stoch. PDEs: Anal. Comp., **6**(4):525–564, (2018).
- [6] Y. Bruned and A. Chandra and I. Chevyrev and M. Hairer, *Renormalising SPDEs in regularity structures*. J. Europ. Math. Soc., **23**(3):869–947, (2021).
- [7] Y. Bruned and F. Gabriel and M. Hairer and L. Zambotti, *Geometric stochastic heat equations*. J. Am. Math. Soc., **35**(1):1–80, (2022).
- [8] Y. Bruned and M. Hairer, and L. Zambotti, *Algebraic renormalization of regularity structures*, Invent. Math., **215**(3):1039–1156, (2019).
- [9] Y. Bruned and U. Nadeem, *Convergence of space-discretised gKPZ via Regularity Structures*. arXiv:2207.09946, (2022).
- [10] Y. Bruned and U. Nadeem, *Diagram-free approach for convergence of tree-based models in regularity structures*. arXiv:2211.11428, (2022).
- [11] A. Chandra and M. Hairer, *An analytic BPHZ theorem for regularity structures*. arXiv:1612.08138, (2016).
- [12] A. Chandra and H. Weber, *Stochastic PDEs, regularity structures and interacting particle systems*. Ann. Fac. Sci. Toulouse, **26**(4):847–909, (2017).
- [13] I. Corwin and H. Shen, *Some recent progress in singular stochastic PDEs*. Bull. Amer. Math. Soc., **57**(3):409–454, (2020).
- [14] P. Friz and M. Hairer, *A Course on Rough Paths, With an introduction to regularity structures*. Universitext, Springer, (2020).
- [15] M. Gubinelli and P. Imkeller and N. Perkowski, *Paracontrolled distributions and singular PDEs*. Forum Math. Pi, **3**(e6):1–75, (2015).
- [16] M. Hairer, *A theory of regularity structures*. Invent. Math., **198**:269–504, (2014).
- [17] M. Hairer, *Renormalization of parabolic stochastic PDEs*. The 20th Takagi Lectures, arXiv:1803.03044, (2018).
- [18] M. Hairer and E. Pardoux. *A Wong-Zakai theorem for stochastic PDEs*. J. Math. Soc. Japan, **67**(4):1551–1604, (2015).
- [19] M. Hairer and J. Quastel. *A class of growth models rescaling to KPZ*. Forum Math. Pi, **6**(e3):1–112, (2018).
- [20] P. Linares and F. Otto and M. Tempelmayr and P. Tsatsoulis, *A diagram-free approach to the stochastic estimates in regularity structures*. arXiv:2112.10739, (2021).
- [21] F. Otto and J. Sauer and S. Smith and H. Weber, *A priori bounds for quasi-linear SPDEs in the full sub-critical regime*. arXiv:2103.11039, (2021).
- [22] F. Otto and K. Seong and M. Tempelmayr, *Lecture notes on tree-free regularity structures*. arXiv:2301.00778, (2023).
- [23] S. Weinberg. *High-energy behavior in quantum field-theory*. Phys. Rev., **118**(2):838–849, (1960).

• I. Bailleul – Univ. Brest, CNRS, LMBA - UMR 6205, 29200 Brest, France.

*E-mail:* ismael.bailleul@univ-brest.fr

• Y. Bruned – IECL (UMR 7502), Université de Lorraine, 54506 Vandœuvre-lès-Nancy, France.

*E-mail:* yvain.bruned@univ-lorraine.fr