

MSC 34C10

**Kamenev type conditions for oscillation of third order
linear ordinary differential equations**

G. A. Grigorian

0019 Armenia c. Yerevan, str. M. Bagramian 24/5

Institute of Mathematics of NAS of Armenia

E - mail: mathphys2@instmath.sci.am, phone: 098 62 03 05, 010 35 48 61

Abstract. The Riccati equation method is used to establish Kamenev-type conditions for the existence of oscillatory solutions to third order linear ordinary differential equations. Three oscillatory theorems are proved, which generalize the Lazer's oscillation criterion.

Key words: third order linear ordinary differential equations, oscillation, Kamenev type conditions, the Riccati equation method, comparison criterion.

1. Introduction. Let $p(t)$, $q(t)$ and $r(t)$ be real-valued continuous functions on $[t_0, +\infty)$. Consider the third order linear ordinary differential equation

$$\phi''' + p(t)\phi'' + q(t)\phi' + r(t)\phi = 0, \quad t \geq t_0.$$

Throughout this paper we will assume that $p(t)$ is continuously differentiable on $[t_0, +\infty)$.

Definition 1.1. *A solution of Eq. (1.1) is called oscillatory if it has arbitrarily large zeroes on $[t_0, +\infty)$.*

The study of the oscillatory behavior of solutions to linear third-order ordinary differential equations is an important problem in the qualitative theory of differential equations, and many works are devoted to it (see [1,2,4] and the works cited therein). Among them notice the following result due to Lazer.

Theorem 1.1([4, Theorem 1.3]). *If $p(t) \equiv 0$, $q(t) \leq 0$, $r(t) > 0$, $t \geq t_0$ and*

$$\int_{t_0}^{+\infty} \left[r(t) - \frac{2}{3\sqrt{3}}(-q(t))^{3/2} \right] dt = +\infty,$$

then Eq. (1.1) has oscillatory solutions. ■

In this paper, we use the Riccati equation method to establish Kamenev-type conditions for the existence of oscillatory solutions to the equation. (1.1). Three oscillatory theorems generalizing Theorem 1.1 are proved.

Auxiliary propositions. The following two lemmas are of fundamental importance for the proof of the main results.

Lemma 2.1.([1, Lemma 2.2]) *If $q(t) \leq 0$, $r(t) > 0$, $t \geq t_0$ and $\phi(t) \neq 0$ is a solution of Eq. (1.1) with $\phi(t) \geq 0$ or $\phi(t) \leq 0$ eventually, then there exists $T \in [t_0, +\infty)$ such that either*

$$\phi(t)\phi'(t) \leq 0, \quad t \geq T \quad (2.1)$$

or

$$\phi(t)\phi'(t) \geq 0, \quad t \geq T \quad \text{and} \quad \phi(t) > 0 \quad \text{for} \quad t \geq T.$$

Furthermore if (2.1) holds, then

$$\phi(t)\phi'(t)\phi''(t) \neq 0, \quad \text{sgn} \phi(t) = \text{sgn} \phi''(t) \neq \text{sgn} \phi'(t), \quad t \geq t_0 \quad (2.2)$$

and

$$\lim_{t \rightarrow +\infty} \phi'(t) = \lim_{t \rightarrow +\infty} \phi''(t) = 0, \quad \lim_{t \rightarrow +\infty} \phi(t) = k \neq \pm\infty. \quad (2.3)$$

■

Lemma 2.2([1, Lemma 2.3]). *Let $q(t) \leq 0$, $r(t) > 0$, $t \geq t_0$. A necessary and sufficient condition for Eq. (1.1) to have oscillatory solutions is that for any nontrivial solution (2.2) and (2.3) hold.*

■

Let $f_k(t)$, $g_k(t)$, $h_k(t)$, $k = 1, 2$, be real-valued continuous functions on $[t_0, +\infty)$. Consider the Riccati equations

$$y' + f_k(t)y^2 + g_k(t)y + h_k(t) = 0, \quad t \geq t_0, \quad (2.4_k)$$

$k = 1, 2$ and the differential inequalities

$$\eta' + f_k(t)\eta^2 + g_k(t)\eta + h_k(t) \geq 0, \quad t \geq t_0, \quad (2.5_k)$$

$k = 1, 2$.

Remark 2.1. *Every solution of Eq. (2.4₂) on $[t_0, t_1)$ is also a solution of the inequality (2.5₂) on $[t_0, t_1)$.*

Remark 2.2. *If $f_1(t) \geq 0$, $t \in [t_0, t_1)$, then every solution of the linear equation*

$$\zeta' + g_1(t)\zeta + h_1(t) = 0, \quad t \in [t_0, t_1)$$

is also a solution of the inequality (2.5₁) on $[t_0, t_1)$.

Theorem 2.1 [3, Theorem 3.1]. Let $y_2(t)$ be a solution of Eq. (2.4₂) on $[t_0, \tau_0)$ ($t_0 < \tau_0 \leq +\infty$) and let $\eta_1(t)$ and $\eta_2(t)$ be solutions of the inequalities (2.5₁) and (2.5₂) respectively on $[t_0, \tau_0)$ such that $y_2(t_0) \leq \eta_k(t_0)$ $k = 1, 2$. In addition let the following conditions be satisfied: $f_1(t) \geq 0$, $\gamma - y_2(t_0) + \int_{t_0}^t \exp \left\{ \int_{t_0}^{\tau} [f_1(s)(\eta_1(s) + \eta_2(s)) + g_1(s)] ds \right\} \left[(f_2(\tau) - f_1(\tau))^2 y_2^2(\tau) + (g_2(\tau) - g_1(\tau)) y_2(\tau) + h_2(\tau) - h_1(\tau) \right] d\tau \geq 0$, $t \in [t_0, \tau_0)$ for some $\gamma \in [y_2(t_0), \eta_1(t_0)]$. Then Eq. (2.4₁) has a solution $y_1(t)$ on $[t_0, \tau_0)$ with $y_1(t_0) \geq \gamma$ and $y_1(t) \geq y_2(t)$, $t \in [t_0, \tau_0)$. ■

3. Oscillation criteria. We set:

$$p_-(t) \equiv \min\{p(t), 0\}, \quad t \geq t_0, \quad F(t, u) \equiv p(t)u^2 + (q(t) - p'(t))u + r(t), \quad t \geq t_0, \quad u \geq 0$$

One can easily verify that for every fixed $t \geq t_0$ the minimum of the function F is

$$D(t) \equiv \begin{cases} r(t), & \text{if } p^2(t) + 3(p'(t) - q(t)) < 0, \\ F_{min}(t), & \text{if } p^2(t) + 3(p'(t) - q(t)) \geq 0, \quad t \geq t_0, \end{cases}$$

where $F_{min}(t) \equiv \min \left\{ r(t), \left[\frac{\sqrt{p^2(t) + 3(p'(t) - q(t))} - p(t)}{3} \right]^3 + p(t) \left[\frac{\sqrt{p^2(t) + 3(p'(t) - q(t))} - p(t)}{3} \right]^2 + (q(t) - p'(t)) \frac{\sqrt{p^2(t) + 3(p'(t) - q(t))} - p(t)}{3} + r(t) \right\}$, $t \geq t_0$.

In the case $p(t) \equiv 0$, $q(t) \leq 0$, $r(t) > 0$ we have

$$D(t) = r(t) - \frac{2}{3\sqrt{3}}(-q(t))^{3/2}, \quad t \geq t_0. \quad (3.1)$$

Theorem 3.1. Let the following conditions be satisfied.

(a) $r(t) > 0$, $q(t) \leq 0$, $t \geq t_0$.

(b) $\limsup_{t \rightarrow +\infty} \frac{1}{t^{\alpha+1}} \int_{t_0}^t (t-\tau)^\alpha \left[(t-\tau)D(\tau) - \frac{\alpha+1}{9} p_-^2(\tau) \right] d\tau = +\infty$, for some $\alpha > 0$.

Then Eq. (1.1) has oscillatory solutions.

Proof. By Lemma 2.1 it follows from the conditions (a) that for every nontrivial solution $\phi(t)$ of Eq. (1.1) with $\phi(t) \geq 0$ or $\phi(t) \leq 0$ eventually there exists $T \geq t_0$ such that either

$$\phi'(t)\phi(t) \leq 0, \quad t \geq T$$

or

$$\phi'(t)\phi(t) \geq 0, \quad t \geq T \text{ and } \phi(t) > 0 \text{ for } t \geq T. \quad (3.2)$$

By virtue of Lemma 2.2 to prove the theorem it is enough to show that the relations (3.2) are impossible. Suppose, at contrary, that (3.2) is valid. Then $y(t) \equiv \frac{\phi'(t)}{\phi(t)}$, $t \geq T$ is a solution to the second order Riccati equation

$$y'' + 3y[p(t) + y]y' + y^3 + p(t)y^2 + q(t)y + r(t) = 0, \quad t \geq T.$$

Therefore, for $y(t)$ the following equality holds.

$$y'(t) + \frac{3}{2}y^2(t) + p(t)y(t) + \int_T^t [y^3(\tau) + p(\tau)y^2(\tau) + (q(\tau) - p'(\tau))y(\tau) + r(\tau)]d\tau = c_T, \quad t \geq T,$$

where $c_T \equiv y'(T) + \frac{3}{2}y^2(T) + p(T)y(T)$. Since $y(t) \geq 0$, $t \geq T$ from the last equality we obtain

$$y'(t) + \frac{3}{2}y^2(t) + p_-(t)y(t) \leq - \int_T^t D(\tau)d\tau + c_T, \quad t \geq T. \quad (3.3).$$

This inequality we can rewrite in the form

$$y'(t) + \frac{3}{2}\left(y(t) + \frac{p_-(t)}{3}\right)^2 \leq - \int_T^t D(\tau)d\tau + \frac{p_-^2(t)}{9} + c_T, \quad t \geq T.$$

Therefore,

$$y(t) \leq - \int_T^t d\tau \int_T^\tau D(\xi)d\xi + \frac{1}{9} \int_T^t p_-^2(\tau)d\tau + c_T(t - T), \quad t \geq T. \quad (3.4)$$

Without loss of generality we can take that $T > 0$. For any $\alpha > 0$ and $M > 0$ consider the integral operator

$$K_{M,\alpha}\phi(t) \equiv \frac{\alpha(\alpha + 1)}{t^{\alpha+1}} \int_T^t (t - \tau)^{\alpha-1} \phi(\tau)d\tau, \quad t \geq M \quad \phi \in \mathbb{C}([M, +\infty)).$$

Obviously this operator is monotone in the sense that if $\phi_j(t) \in \mathbb{C}([M, +\infty))$, $j = 1, 2$, $\phi_1(t) \geq \phi_2(t)$, $t \geq M$, then $(K_{M,\alpha}\phi_1)(t) \geq (K_{M,\alpha}\phi_2)(t)$, $t \geq M$. Due to this, acting on both sides of (3.4) by operator $K_{T,\alpha}$ and making some simplifications we obtain

$$\begin{aligned} \frac{\alpha(\alpha+1)}{t^{\alpha+1}} \int_T^t (t-\tau)^{\alpha-1} y(\tau) d\tau &\leq -\frac{1}{t^{\alpha+1}} \int_T^t (t-\tau)^{\alpha+1} D(\tau) d\tau + \frac{(\alpha+1)}{9t^{\alpha+1}} \int_T^t (t-\tau)^\alpha p_-^2(\tau) d\tau + \\ &+ \frac{\alpha(\alpha+1)c_T}{t^{\alpha+1}} \int_T^t (t-\tau)^{\alpha-1} (\tau-T) d\tau + \frac{\alpha(\alpha+1)}{t^{\alpha+1}} y(T) \int_T^t (t-\tau)^{\alpha-1} d\tau, \quad t \geq T. \end{aligned} \quad (3.5)$$

Consider the function

$$\begin{aligned} \Delta(t) &\equiv \frac{1}{t^{\alpha+1}} \int_{t_0}^T (t-\tau)^{\alpha+1} D(\tau) d\tau - \frac{\alpha+1}{9t^{\alpha+1}} \int_{t_0}^T (t-\tau)^\alpha p_-^2(\tau) d\tau + \\ &+ \frac{\alpha(\alpha+1)c_T}{t^{\alpha+1}} \int_T^t (t-\tau)^{\alpha-1} (\tau-T) d\tau + \frac{\alpha(\alpha+1)}{t^{\alpha+1}} y(T) \int_T^t (t-\tau)^\alpha d\tau \quad t \geq T. \end{aligned}$$

We have

$$\begin{aligned} |\Delta(t)| &\leq \frac{(t-T)^{\alpha+1}}{t^{\alpha+1}} \int_{t_0}^T |D(\tau)| d\tau + \frac{(\alpha+1)(t-t_0)^\alpha}{9t^{\alpha+1}} \int_{t_0}^T p_-^2(\tau) d\tau + \\ &+ \alpha(\alpha+1)c_T \frac{(t-T)^{\alpha-1} (t-T)^2}{t^{\alpha+1} \cdot 2} + \frac{\alpha(\alpha+1)y(T)(t-T)^{\alpha+1}}{t^{\alpha+1}}, \quad t \geq T. \end{aligned} \quad (3.6)$$

Rewrite (3.5) in the form

$$\begin{aligned} \frac{\alpha(\alpha+1)}{t^{\alpha+1}} \int_{t_0}^t (t-\tau)^{\alpha-1} y(\tau) d\tau &\leq -\frac{1}{t^{\alpha+1}} \int_{t_0}^t (t-\tau)^\alpha \left[(t-\tau)D(\tau) - \frac{(\alpha+1)}{9} p_-^2(\tau) \right] d\tau + \\ &+ \Delta(t), \quad t \geq T. \end{aligned} \quad (3.7)$$

It follows from (3.5) that $\Delta(t)$ is a bounded function. Then from the condition (b) of the theorem it follows that the right part of (3.7) takes negative values in some points $t \geq T$,

whereas the hypothesis (3,2) implies that the left part of (3.2) is nonnegative for all $t \geq T$. We have obtained a contradiction, completing the proof of the theorem.

Due to (3.1) from Theorem 3.1 we obtain immediately

Corollary 3.1. *If $p(t) \equiv 0$, $q(t) \leq 0$, $r(t) > 0$, $t \geq t_0$ and for some $\alpha > 1$*

$$\limsup_{t \rightarrow +\infty} \frac{1}{t^\alpha} \int_{t_0}^t (t - \tau)^\alpha \left[r(\tau) - \frac{2}{3\sqrt{3}} (-q(\tau))^{3/2} \right] d\tau = +\infty,$$

then Eq. (1.1) has oscillatory solutions.

Theorem 3.2. *Let the condition (a) of Theorem 3.1 and the following conditions be satisfied*

(c) $\int_{t_0}^{+\infty} D(\tau) d\tau = +\infty,$

(d) *there exists $\alpha > 0$ such that*

$$\limsup_{t \rightarrow +\infty} \frac{1}{t^{\alpha+1}} \int_{t_0}^t (t - \tau)^\alpha \left[(t - \tau) D(\tau) - (\alpha + 1) p_-(\tau) \right] d\tau = +\infty.$$

Then Eq. (1.1) has oscillatory solutions.

Proof. Let the condition (a) of Theorem 3.1 holds. To prove the theorem it is enough, as in the proof of Theorem 3.1, to show that (3.2) cannot be satisfied. Assume the contrary, that (3.2) is true. Then (3.3) holds. We set

$$\lambda(t) \equiv - \int_T^t D(\tau) d\tau + c_T - y'(t) - \frac{3}{2} y^2(t) - p_-(t) y(t), \quad t \geq T.$$

It follows from (3.3) that

$$\lambda(t) \geq 0, \quad t \geq T. \tag{3.8}$$

Consider the Riccati equations

$$y' + \frac{3}{2} y^2 + p_-(t) y + \lambda(t) = - \int_T^t D(\tau) d\tau + c_T, \quad t \geq T, \tag{3.9}$$

$$u' + \frac{3}{2} u^2 + p_-(t) u = 0, \quad t \geq T. \tag{3.10}$$

It follows from the condition (c) that there exists $T_1 \geq T$ such that

$$\int_T^t D(\tau)d\tau \geq c_T, \quad t \geq T_1. \quad (3.11)$$

Obviously $y(t)$ is a solution of Eq. (3.9) on $[T_1, +\infty)$. Then applying Theorem 2.1 to the pair of equations (3.9) and (3.10) and taking into account (3.8) and (3.11) we conclude that the solution $u(t)$ of Eq. (3.10) with $u(T_1) > y(T_1)$ exists on $[T_1, +\infty)$ and

$$u(t) \geq y(t), \quad t \geq T_1. \quad (3.12)$$

It is well known that $u(t)$ can be represent in the following explicit form (as a solution of a Bernoulli equation)

$$u(t) = \frac{u(T_1) \exp\left\{-\int_{T_1}^t p_-(\tau)d\tau\right\}}{1 + \frac{3}{2}u(T_1) \int_{T_1}^t \exp\left\{-\int_{T_1}^{\tau} p_-(s)ds\right\}d\tau}, \quad t \geq T_1.$$

Then from (3.12) it follows

$$p_-(t)y(t) - \frac{p_-(t)u(T_1) \exp\left\{-\int_{T_1}^t p_-(\tau)d\tau\right\}}{1 + \frac{3}{2}u(T_1) \int_{T_1}^t \exp\left\{-\int_{T_1}^{\tau} p_-(s)ds\right\}d\tau} \geq 0, \quad t \geq T_1.$$

This together with (3.3) implies

$$y'(t) \leq -\int_T^t D(\tau)d\tau + c_T - \frac{p_-(t)u(T_1) \exp\left\{-\int_{T_1}^t p_-(\tau)d\tau\right\}}{1 + \frac{3}{2}u(T_1) \int_{T_1}^t \exp\left\{-\int_{T_1}^{\tau} p_-(s)ds\right\}d\tau}.$$

Integrating this inequality from T_1 to t we obtain

$$y(t) \leq -\int_{T_1}^t d\tau \int_T^{\tau} D(s)ds + \ln\left[1 + \frac{3}{2}u(T_1) \int_{T_1}^t \exp\left\{-\int_{T_1}^{\tau} p_-(s)ds\right\}d\tau\right] + y(T_1) + c_T(t - T),$$

$t \geq T_1$. From here it follows (since $p_-(t) \leq 0$, $t \geq t_0$)

$$y(t) \leq - \int_{T_1}^t d\tau \int_T^\tau D(s)ds + \ln \left[1 + \frac{3}{2}u(T_1)(t - T_1) \exp \left\{ - \int_{T_1}^t p_-(s)ds \right\} \right] + y(T_1) + c_T(t - T),$$

$t \geq T_1$. For any $T_2 > T_1$ from here we obtain

$$y(t) \leq - \int_{T_1}^t d\tau \int_T^\tau D(s)ds + \int_{T_1}^t p_-(s)ds + L(t) + \ln(t - T_1) = c_T(t - T), \quad t \geq T_2, \quad (3.13)$$

where $L(t) \equiv \ln \left[1 + \frac{3}{2}u(T_1)(t - T_1) \exp \left\{ - \int_{T_1}^t p_-(s)ds \right\} \right] - \int_{T_1}^t p_-(s)ds - \ln(t - T_1) + y(T_1)$, $t \geq T_2$. We claim that $L(t)$ is bounded on $[T_2, +\infty)$. Indeed, we have

$$\begin{aligned} |L(t)| &= \left| \ln \left[1 + \frac{3}{2}u(T_1)(t - T_1) \exp \left\{ - \int_{T_1}^t p_-(s)ds \right\} \right] - \right. \\ &\quad \left. - \ln \left[\frac{3}{2}u(T_1)(t - T_1) \exp \left\{ - \int_{T_1}^t p_-(s)ds \right\} \right] + \ln \left[\frac{3}{2}u(T_1) \right] + y(T_1) \right| = \\ &= \left| \ln \left[\frac{3}{2}u(T_1) \right] + y(T_1) + \ln \left[1 + \frac{2}{3u(T_1) \exp \left\{ - \int_{T_1}^t p_-(s)ds \right\} (t - T_1)} \right] \right| \leq \\ &\leq \left| \ln \left[\frac{3}{2}u(T_1) \right] \right| + y(T_1) + \ln \left[1 + \frac{2}{3u(T_1)(T_2 - T_1)} \right], \quad t \geq T_2. \end{aligned}$$

Therefore $L(t)$ is bounded on $[T_2, +\infty)$. From (3.13) we get

$$y(t) \leq - \int_{T_2}^t d\tau \int_T^\tau D(s)ds - \int_{T_2}^t p_-(s)ds + J(t), \quad t \geq T_2, \quad (3.14)$$

where

$$J(t) \equiv L(t) - \int_{T_1}^{T_2} d\tau \int_T^\tau D(s)ds - \int_{T_1}^{T_2} p_-(s)ds + \ln(t - T_1) - (t - T_2) \int_T^{T_2} D(s)ds + c_T(t - T), \quad t \geq T_2.$$

Since $L(t)$ is bounded on $[T_2, +\infty)$ we have

$$J(t) = O(t), \quad t \rightarrow +\infty. \quad (3.15)$$

Acting on both sides of (3.14) by the operator $K_{T_2, \alpha}$ we obtain

$$\begin{aligned} \frac{\alpha(\alpha+1)}{t^{\alpha+1}} \int_{T_2}^t (t-\tau)^{\alpha-1} y(\tau) d\tau &\leq -\frac{1}{t^{\alpha+1}} \int_{T_2}^t (t-\tau)^{\alpha+1} D(\tau) d\tau + \\ &+ \frac{\alpha+1}{t^{\alpha+1}} \int_{T_2}^t (t-\tau)^\alpha p_-(\tau) d\tau + \frac{\alpha(\alpha+1)}{t^{\alpha+1}} \int_{T_2}^t (t-\tau)^{\alpha-1} J(\tau) d\tau, \quad t \geq T_2. \end{aligned}$$

Hence,

$$\begin{aligned} \frac{\alpha(\alpha+1)}{t^{\alpha+1}} \int_{T_2}^t (t-\tau)^{\alpha-1} y(\tau) d\tau &\leq -\frac{1}{t^{\alpha+1}} \int_{t_0}^t (t-\tau)^{\alpha+1} D(\tau) d\tau + \\ &+ \frac{\alpha+1}{t^{\alpha+1}} \int_{t_0}^t (t-\tau)^\alpha p_-(\tau) d\tau + \Delta_1(t), \quad t \geq T_2, \end{aligned} \quad (3.16)$$

where

$$\begin{aligned} \Delta_1(t) &\equiv \frac{1}{t^{\alpha+1}} \int_{t_0}^{T_2} (t-\tau)^{\alpha+1} D(\tau) d\tau - \\ &- \frac{\alpha+1}{t^{\alpha+1}} \int_{t_0}^{T_2} (t-\tau)^\alpha p_-(\tau) d\tau - \frac{\alpha(\alpha+1)}{t^{\alpha+1}} \int_{t_0}^{T_2} (t-\tau)^{\alpha-1} J(\tau) d\tau, \quad t \geq T_2. \end{aligned}$$

It follows from (3.15) that $\Delta_1(t)$ is a bounded function on $[T_2, +\infty)$. Then it follows from the condition (d) of the theorem that the right side of (3.16) takes negative values at some points t of the interval $[T_2, +\infty)$, while the left side is always nonnegative. We have obtained a contradiction, which completes the proof of the theorem.

Example 3.1. Let $t_0 = 1$, $p(t) = -Mt^\gamma$, $M > 0$, $\gamma \in \mathbb{R}$, $q(t) \equiv 0$, $r(t) = Nt^\beta - \delta(t)$, $N > 0$, $\beta > -1$, $t \geq 1$, where $\delta(t) = 0$, if $Mt^{2\gamma} - 3\gamma t^{\gamma-1} < 0$ and $\delta(t) = \min\left\{0, \left[\frac{\sqrt{M^2 t^{2\gamma} - 3M\gamma t^{\gamma-1} + Mt^\gamma}}{3}\right]^3 - Mt^\gamma \left[\frac{\sqrt{M^2 t^{2\gamma} - 3M\gamma t^{\gamma-1} + Mt^\gamma}}{3}\right]^2 - M\gamma t^{\gamma-1} \frac{\sqrt{M^2 t^{2\gamma} - 3M\gamma t^{\gamma-1} + Mt^\gamma}}{3}\right\}$,

if $Mt^{2\gamma} - 3\gamma t^{\gamma-1} \geq 0$, $t \geq 1$. Then, obviously, $D(t) = Nt^\beta$, $p_-(t) = -Mt^\gamma$, $t \geq 1$. Therefore, it is easy to check that, for $\gamma-1 < \beta < 2\gamma-1$ all the conditions of Theorem 3.2 are satisfied, but the condition (b) of Theorem 3.1 is not.

Theorem 3.3 Let the condition (a) of Theorem 3.1 and the following conditions be satisfied

(e) $\liminf_{t \rightarrow +\infty} \int_{t_0}^t D(\tau) d\tau > -\infty$,

(f) $p_-(t) = p_{-,1}(t) + p_{-,2}(t)$, $p_{-,j}(t) \leq 0$, $j = 1, 2$, $t \geq t_0$, $\int_{t_0}^{+\infty} |p_{-,1}(t)| dt < +\infty$ and

$p_{-,2}(t)$ is bounded,

(g) for some $\alpha > 1$

$$\limsup_{t \rightarrow +\infty} \frac{1}{t^\alpha} \int_{t_0}^t (t - \tau)^\alpha D(\tau) d\tau = +\infty.$$

Then Eq. (1.1) has oscillatory solutions.

Proof. By the condition (a) to prove the theorem it is enough to show that the relation (3.2) cannot be satisfied. Assume (3.2) is valid. Then (3.3) holds. Rewrite (3.3) in the form

$$y'(t) + \frac{3}{2} \left(y(t) + \frac{p_{-,2}(t)}{3} \right)^2 + p_{-,1}(t)y(t) \leq - \int_T^t D(\tau) d\tau + c_T + \frac{p_{-,2}^2(t)}{9}, \quad t \geq T.$$

It follows from here that

$$y'(t) + p_{-,1}(t)y(t) \leq - \int_T^t D(\tau) d\tau + c_T + \frac{p_{-,2}^2(t)}{9}, \quad t \geq T. \quad (3.17)$$

Since $p_{-,2}(t)$ is bounded it follows from the condition (e) that

$$\int_T^t D(\tau) d\tau - c_T - \frac{p_{-,2}^2(t)}{9} \geq -e_T \quad t \geq T, \quad (3.18)$$

for some $e_T > 0$. We set

$$\lambda_1(t) \equiv - \int_T^t D(\tau) d\tau + c_T + \frac{p_{-,2}^2(t)}{9} - y'(t) - p_{-,1}(t)y(t), \quad t \geq T.$$

It follows from (3.17) that

$$\lambda_1(t) \geq 0, \quad t \geq T. \quad (3.19)$$

Consider the linear equations

$$u' + p_{-,1}(t)u + \lambda_1(t) + \int_T^t D(\tau)d\tau - c_T - \frac{p_{-,2}(t)}{9} = 0, \quad t \geq T. \quad (3.20)$$

$$v' + p_{-,1}(t)v - e_T = 0, \quad t \geq T. \quad (3.21)$$

Obviously $y(t)$ is a solution of Eq. (3.20) on $[T, +\infty)$ and

$$v(t) \equiv \exp\left\{-\int_T^t p_{-,1}(\tau)d\tau\right\} \left[y(T) + e_T \int_T^t \exp\left\{\int_T^\tau p_{-,1}(s)ds\right\} d\tau \right], \quad t \geq T$$

is a solution of Eq. (3.21) on $[T, +\infty)$ with $u(T) = y(T)$. Obviously we can interpret the equations (3.20) and (3.21) as Riccati equations with $\equiv 0$ coefficients of u^2 and v^2 . Then applying Theorem 2.1 to these equations and taking into account (3.18) and (3.19) we conclude that $y(t) \leq v(t)$, $t \geq T$. Hence,

$$p_{-,1}(t)y(t) \geq p_{-,1}(t)v(t), \quad t \geq T \quad (\text{since } y(t) > 0, \quad p_{-,1}(t) \leq 0, \quad t \geq T).$$

This together with (3.17) implies

$$y'(t) \leq -\int_T^t D(\tau)d\tau + f_T(t), \quad t \geq T,$$

where

$$f_T(t) \equiv c_T + \frac{p_{-,2}^2(t)}{9} - p_{-,1}(t)v(t), \quad t \geq T.$$

Integrating the obtained inequality from T to t we obtain

$$y(t) \leq y(T) - \int_T^t d\tau \int_T^\tau D(s)ds + \int_T^t f_T(\tau)d\tau, \quad t \geq T. \quad (3.22)$$

It is not difficult to verify (by integrating by parts) that

$$\int_T^t p_{-,1}(\tau)v(\tau)d\tau = \exp\left\{-\int_T^t p_{-,1}(\tau)d\tau\right\} \left[y(T) + e_T \int_T^t \exp\left\{\int_T^\tau p_{-,1}(s)ds\right\} \right] -$$

$$-y(T) - e_T(t-T), \quad t \geq T.$$

Then, obviously,

$$f_T(t) = O(t), \quad \text{as } t \rightarrow +\infty. \quad (3.23)$$

Without loss of generality we can take that $T > 0$. Then acting on both sides of (3.22) by the operator $K_{T, \alpha-1}$ (here $\alpha > 1$) we obtain

$$\frac{(\alpha-1)\alpha}{t^\alpha} \int_T^t (t-\tau)^{\alpha-1} y(\tau) d\tau \leq \frac{1}{t^\alpha} \int_T^t (t-\tau)^\alpha D(\tau) d\tau + \frac{(\alpha-1)\alpha}{t^\alpha} \int_T^t (t-\tau)^{\alpha-2} d\tau \int_T^\tau f_T(s) ds,$$

$t \geq T$. Hence,

$$\frac{(\alpha-1)\alpha}{t^\alpha} \int_T^t (t-\tau)^{\alpha-1} y(\tau) d\tau \leq \frac{1}{t^\alpha} \int_{t_0}^t (t-\tau)^\alpha D(\tau) d\tau + F_T(t), \quad t \geq T, \quad (3.24)$$

where $F_T(t) \equiv -\frac{1}{t^\alpha} \int_{t_0}^T (t-\tau)^\alpha D(\tau) d\tau + \frac{(\alpha-1)\alpha}{t^\alpha} \int_T^t (t-\tau)^{\alpha-2} d\tau \int_T^\tau f_T(s) ds$, $t \geq T$. It follows from (3.23) that the function $F_T(t)$ is bounded on $[T, +\infty)$. Obviously the left part of (3.24) is nonnegative for all $t \geq T$. However, it follows from the condition (g) of the theorem that the right part of (3.24) takes negative values in some points of $[T, +\infty)$. We have obtained a contradiction, completing the proof of the theorem.

Remark 3.1. *Theorems 3.1 - 3.3 are generalizations of Theorem 1.1.*

Example 3.2. *Assume*

$$p(t) = M \begin{cases} -n^3 \sin^2[n^5 \pi(t-n)], & t \in [n, n + \frac{1}{n^5}]. \\ 0, & t \in (n + \frac{1}{n^5}), \end{cases}$$

$M = \text{const} > 0$, $n = 1, 2, \dots$. Obviously $p(t)$ is continuously differentiable on $[1, +\infty)$ and $p_-(t) = p(t)$, $t \geq 1$. Let $q(t) \equiv 0$, $r(t) = r_0 + \delta_1(t)$, $t \geq 1$, where $r_0 > 0$, $\delta_1(t) = 0$, if $p^2(t) + 3p'(t) < 0$ and $\delta_1(t) = \min\left\{0, \left[\frac{\sqrt{p^2(t)+3p'(t)}-p(t)}{3}\right]^3 + p(t) \left[\frac{\sqrt{p^2(t)+3p'(t)}-p(t)}{3}\right]^2 - p(t) \frac{\sqrt{p^2(t)+3p'(t)}-p(t)}{3}\right\}$, if $p^2(t) + 3p'(t) \geq 0$, $t \geq 1$. Then obviously

$$D(t) \equiv r_0 > 0, \quad t \geq 1 \quad (3.25)$$

and $\int_{t_0}^{+\infty} |p_{-,1}(t)|dt = \int_{t_0}^{+\infty} |p(t)|dt \leq \sum_{n=1}^{+\infty} \frac{1}{n^2} < +\infty$. Therefore all conditions of Theorem 3.3 for this case of coefficients of Eq (1.1) are satisfied. Let us show that for this case the condition (b) of Theorem 3.1 is not satisfied for all $M \geq 12r_0$. We have

$$\int_n^{n+1} p_-^2(t)dt = n^6 \int_n^{n+\frac{1}{n^5}} \sin^4(n^5\pi(t-n))dt = \frac{Mn}{\pi} \int_0^\pi \sin^4\tau d\tau = \frac{3Mn}{8}, \quad n = 1, 2, \dots$$

It follows from here that

$$\int_1^t p_-^2(\tau)d\tau \geq \frac{3M(t-1)t}{16}, \quad t \geq 1.$$

This together with (3.25) implies

$$\int_1^t (t-\tau)^\alpha [(t-\tau)D(\tau) - \frac{\alpha+1}{9}p_-^2(\tau)]d\tau \leq \int_1^t \frac{(t-\tau)^\alpha}{2} [r_0\tau^2 - \frac{M(\alpha+1)}{12}(\tau-1)\tau]d\tau, \quad t \geq 1.$$

Therefore, if $M \geq 12r_0$, then the condition (b) of Theorem 3.1 is not satisfied,

References

1. L. Erbe, Existence of oscillatory solutions and asymptotic behavior for a class of third order linear differential equations. Pacific J. Math., vol. 64, No 2, 1976, pp. 369–385.
2. G. A. Grigorian, Some properties of the solutions of third order linear ordinary differential equations. Rocky J. Math. vol. 46, No 1, 2016, pp. 147–168.
3. G. A. Grigorian, On two comparison tests for second-order linear ordinary differential equations, Diff. Urav., vol 47, (2011), 1225–1240 (in Russian), Diff. Eq., vol. 47, (2011), 1237–1252 (in English).
4. A. C. Lazer, The behavior of solutions of the differential equation $y''' + p(x)t' + q(x)y = 0$. Pacific J. Math., vol. 17, No 3, 1966, pp. 435–466.