

A GENERALIZATION OF THE NORM INEQUALITIES FOR SUMS OF OPERATORS

MOJTABA BAKHERAD

ABSTRACT. Let $\mathbb{B}(\mathcal{H})$ denotes the set of all bounded linear operators on a complex Hilbert space \mathcal{H} . In this paper, we present some norm inequalities for sums of operators which are a generalization some recent results. Among other inequalities, it is shown that if $S, T \in \mathbb{B}(\mathcal{H})$ are normal operators, then

$$\begin{aligned} & \|S + T\| \\ & \leq \frac{1}{2}(\|S\| + \|T\|) + \frac{1}{2} \min_{t>0} \sqrt{(\|S\| - \|T\|)^2 + \left\| \frac{1}{t} f_1(|S|) g_1(|T|) + t f_2(|S|) g_2(|T|) \right\|^2}, \end{aligned}$$

where f_1, f_2, g_1, g_2 are non-negative continuous functions on $[0, \infty)$, in which $f_1(x)f_2(x) = x$ and $g_1(x)g_2(x) = x$ ($x \geq 0$). Moreover, it is shown several inequalities for the numerical radius.

1. INTRODUCTION

Let $\mathbb{B}(\mathcal{H})$ denotes the C^* -algebra of all bounded linear operators on a complex Hilbert space \mathcal{H} with an inner product $\langle \cdot, \cdot \rangle$ and the corresponding norm $\| \cdot \|$. In the case when $\dim \mathcal{H} = n$, we identify $\mathbb{B}(\mathcal{H})$ with the matrix algebra \mathbb{M}_n of all $n \times n$ matrices with entries in the complex field. For $S \in \mathbb{B}(\mathcal{H})$, let $S = \Re(S) + i\Im(S)$ be the Cartesian decomposition of S , where the Hermitian matrices $\Re(S) = \frac{S+S^*}{2}$ and $\Im(S) = \frac{S-S^*}{2i}$ are called the real and imaginary parts of S , respectively. The numerical radius of $S \in \mathbb{B}(\mathcal{H})$ is defined by $w(S) := \sup\{|\langle Sx, x \rangle| : x \in \mathcal{H}, \|x\| = 1\}$. It is well known that $w(\cdot)$ defines a norm on $\mathbb{B}(\mathcal{H})$, which is equivalent to the usual operator norm $\| \cdot \|$. In fact, for any $S \in \mathbb{B}(\mathcal{H})$, $\frac{1}{2}\|S\| \leq w(S) \leq \|S\|$; see . Let $r(\cdot)$ denote the spectral radius. It is well known that for every operator $S \in \mathbb{B}(\mathcal{H})$, we have $r(S) \leq w(S)$. In , the authors showed that $w(S) = \sup_{\theta \in \mathbb{R}} \|\Re(e^{i\theta} S)\| = \sup_{\alpha^2 + \beta^2 = 1} \|\alpha \Re S + \beta \Im S\|$, which is equal to the above definition. For more facts about the numerical radius see and references therein. Let $S, T, X, Y \in \mathbb{B}(\mathcal{H})$.

The operator matrices $\begin{bmatrix} S & 0 \\ 0 & Y \end{bmatrix}$ and $\begin{bmatrix} 0 & T \\ X & 0 \end{bmatrix}$ are called the diagonal and off-diagonal parts of the operator matrix $\begin{bmatrix} S & T \\ X & Y \end{bmatrix}$, respectively.

Date: January 25, 2023.

2010 Mathematics Subject Classification. Primary: 47A30, 47A12; Secondary 47A08, 47L05.

Key words and phrases. Keywords: Operator norm, Positive operator, Norm inequality, Numerical radius.

In [1], it has been shown that if S is an operator in $\mathbb{B}(\mathcal{H})$, then

$$w(S) \leq \frac{1}{2} \left(\|S\| + \|S^2\|^{\frac{1}{2}} \right). \quad (1.1)$$

Several refinements and generalizations of inequality have been given; see [2]. Yamazaki showed that for $S \in \mathbb{B}(\mathcal{H})$ and $t \in [0, 1]$ we have

$$w(S) \leq \frac{1}{2} \left(\|S\| + w(\tilde{S}_t) \right). \quad (1.2)$$

where $S = U|S|$ is the polar decomposition of S and $\tilde{S}_t = |S|^t U |S|^{1-t}$. Horn et al. proved that

$$\|S + T\| \leq \| |S| + |T| \|, \quad (1.3)$$

where $S, T \in \mathbb{B}(\mathcal{H})$ are normal. Davidson and Power proved that if S and T are positive operators in $\mathbb{B}(\mathcal{H})$, then

$$\|S + T\| \leq \max\{\|S\|, \|T\|\} + \|ST\|^{\frac{1}{2}}. \quad (1.4)$$

Inequality has been generalized in [3]. In [4], the author extended this inequality to the form

$$\|S + T\| \leq \max\{\|S\|, \|T\|\} + \frac{1}{2} \left(\| |S|^t |T|^{1-t} \| + \| |S^*|^{1-t} |T^*|^t \| \right), \quad (1.5)$$

in which $S, T \in \mathbb{B}(\mathcal{H})$ and $t \in [0, 1]$. In [5], the authors showed that a generalization of inequality as follows:

$$\|S + T\| \leq \max\{\|S\|, \|T\|\} + \frac{1}{2} \left(\|f(|S|)g(|T|)\| + \|f(|S^*|)g(|T^*|)\| \right),$$

in which $S, T \in \mathbb{B}(\mathcal{H})$ and f, g be two non-negative non-decreasing continuous functions on $[0, \infty)$ such that $f(x)g(x) = x$ ($x \geq 0$). Recently, Shi et al. proved the following inequality

$$\|S + T\| \leq \frac{1}{2}(\|S\| + \|T\|) + \frac{1}{2} \sqrt{(\|S\| - \|T\|)^2 + \left\| \frac{1}{t} |S|^r |T|^s + t |S|^{1-r} |T|^{1-s} \right\|^2} \quad (1.6)$$

for normal operators $S, T \in \mathbb{B}(\mathcal{H})$, $r, s \in [0, 1]$ and $t > 0$.

In this paper, we give several further norm inequalities for sums of bounded linear operators. These inequalities refine and generalize inequalities. Moreover, as another application, we show a new numerical radius inequality which is a generalization [6].

2. MAIN RESULTS

To present our results, we need the following lemmas.

Lemma 2.1. *Let $S, T, X, Y \in \mathbb{B}(\mathcal{H})$. Then*

$$\left\| \begin{bmatrix} S & T \\ X & Y \end{bmatrix} \right\| \leq \left\| \begin{bmatrix} \|S\| & \|T\| \\ \|X\| & \|Y\| \end{bmatrix} \right\|.$$

Lemma 2.2. *If $S, T \in \mathbb{B}(\mathcal{H})$ in which ST is selfadjoint, then*

$$\|ST\| \leq \|\Re(TS)\|.$$

In the first result, a generalization of the inequality is obtained.

Theorem 2.3. *Let $S, T \in \mathbb{B}(\mathcal{H})$ be normal and f_1, f_2, g_1, g_2 be non-negative continuous functions on $[0, \infty)$, in which $f_1(x)f_2(x) = x$ and $g_1(x)g_2(x) = x$ ($x \geq 0$). Then*

$$\begin{aligned} & \|S + T\| \\ & \leq \frac{1}{2}(\|S\| + \|T\|) + \frac{1}{2} \min_{t>0} \sqrt{(\|S\| - \|T\|)^2 + \left\| \frac{1}{t} f_1(|S|)g_1(|T|) + t f_2(|S|)g_2(|T|) \right\|^2}. \end{aligned}$$

Proof. Let S and T be positive operators. Then for all $t > 0$ we have

$$\begin{aligned} & \|S + T\| \\ & = \left\| \begin{bmatrix} S + T & 0 \\ 0 & 0 \end{bmatrix} \right\| \\ & = \left\| \begin{bmatrix} t f_2(S) & g_1(T) \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \frac{1}{t} f_1(S) & 0 \\ g_2(T) & 0 \end{bmatrix} \right\| \\ & \leq \left\| \Re \left(\begin{bmatrix} \frac{1}{t} f_1(S) & 0 \\ g_2(T) & 0 \end{bmatrix} \begin{bmatrix} t f_2(S) & g_1(T) \\ 0 & 0 \end{bmatrix} \right) \right\| \\ & \quad \text{(by Lemma)} \\ & = \left\| \begin{bmatrix} S & \frac{1}{2}(\frac{1}{t} f_1(S)g_1(T) + t f_2(S)g_2(T)) \\ \frac{1}{2}(t f_2(S)g_2(T) + \frac{1}{t} f_1(S)g_1(T)) & T \end{bmatrix} \right\| \\ & \leq \left\| \begin{bmatrix} \|S\| & \frac{1}{2}\|\frac{1}{t} f_1(S)g_1(T) + t f_2(S)g_2(T)\| \\ \frac{1}{2}\|t f_2(S)g_2(T) + \frac{1}{t} f_1(S)g_1(T)\| & \|T\| \end{bmatrix} \right\| \\ & \quad \text{(by Lemma)} \\ & = r \left(\begin{bmatrix} \|S\| & \frac{1}{2}\|\frac{1}{t} f_1(S)g_1(T) + t f_2(S)g_2(T)\| \\ \frac{1}{2}\|\frac{1}{t} f_2(S)g_2(T) + t f_1(S)g_1(T)\| & \|T\| \end{bmatrix} \right) \\ & \quad \text{(since is Hermitian matrix)} \\ & = \frac{1}{2}(\|S\| + \|T\|) + \frac{1}{2} \sqrt{(\|S\| - \|T\|)^2 + \left\| \frac{1}{t} f_1(|S|)g_1(|T|) + t f_2(|S|)g_2(|T|) \right\|^2}, \end{aligned}$$

whence

$$\begin{aligned} & \|S + T\| \\ & \leq \frac{1}{2}(\|S\| + \|T\|) + \frac{1}{2} \min_{t>0} \sqrt{(\|S\| - \|T\|)^2 + \left\| \frac{1}{t} f_1(|S|)g_1(|T|) + t f_2(|S|)g_2(|T|) \right\|^2} \end{aligned}$$

for positive operators S and T . Now, if S and T are normal operators, then by using inequality we have

$$\begin{aligned}
& \|S + T\| \\
& \leq \| |S| + |T| \| \quad (\text{by inequality}) \\
& \leq \frac{1}{2}(\| |S| \| + \| |T| \|) + \frac{1}{2} \min_{t>0} \sqrt{(\| |S| \| - \| |T| \|)^2 + \left\| \frac{1}{t} f_1(|S|)g_1(|T|) + t f_2(|S|)g_2(|T|) \right\|^2} \\
& = \frac{1}{2}(\|S\| + \|T\|) + \frac{1}{2} \min_{t>0} \sqrt{(\|S\| - \|T\|)^2 + \left\| \frac{1}{t} f_1(|S|)g_1(|T|) + t f_2(|S|)g_2(|T|) \right\|^2}.
\end{aligned}$$

□

As a consequence of Theorem we have the following result.

Corollary 2.4. *Let $S, T \in \mathbb{B}(\mathcal{H})$ be normal, f_1, f_2, g_1, g_2 be non-negative continuous functions on $[0, \infty)$ such that $f_1(x)f_2(x) = x$ and $g_1(x)g_2(x) = x$ ($x \geq 0$). Then*

$$\|S + T\| \leq \frac{1}{2}(\|S\| + \|T\|) + \frac{1}{2} \sqrt{(\|S\| - \|T\|)^2 + 4 \|f_1(|S|)g_1(|T|)\| \|f_2(|S|)g_2(|T|)\|}.$$

Proof. If S and T are normal operators, then

$$\begin{aligned}
& \|S + T\| \\
& \leq \frac{1}{2}(\|S\| + \|T\|) + \frac{1}{2} \min_{t>0} \sqrt{(\|S\| - \|T\|)^2 + \left\| \frac{1}{t} f_1(|S|)g_1(|T|) + t f_2(|S|)g_2(|T|) \right\|^2} \\
& \leq \frac{1}{2}(\|S\| + \|T\|) + \frac{1}{2} \min_{t>0} \sqrt{(\|S\| - \|T\|)^2 + \left(\frac{1}{t} \|f_1(|S|)g_1(|T|)\| + t \|f_2(|S|)g_2(|T|)\| \right)^2} \\
& = \frac{1}{2}(\|S\| + \|T\|) + \frac{1}{2} \sqrt{(\|S\| - \|T\|)^2 + 4 \|f_1(|S|)g_1(|T|)\| \|f_2(|S|)g_2(|T|)\|}.
\end{aligned}$$

The last equation follows from $\min_{t>0} h(t) = \min_{t>0} (\frac{1}{t}a + tb) = 2\sqrt{ab}$ ($a, b \geq 0$). □

For invertible normal operators we get the next result.

Corollary 2.5. *Let $S, T \in \mathbb{B}(\mathcal{H})$ be invertible normal and f, g be two positive continuous functions on $[0, \infty)$. Then*

$$\begin{aligned}
& \|S + T\| \\
& \leq \frac{1}{2}(\|S\| + \|T\|) + \frac{1}{2} \min_{t>0} \sqrt{(\|S\| - \|T\|)^2 + \left\| \frac{1}{t} f(|S|)g(|T|) + t|S|(f(|S|))^{-1}|T|(g(|T|))^{-1} \right\|^2}.
\end{aligned}$$

In particular,

$$\|S + T\| \leq \frac{1}{2}(\|S\| + \|T\|) + \frac{1}{2} \min_{t>0} \sqrt{(\|S\| - \|T\|)^2 + \left\| \frac{1}{t} |S|^r |T|^s + t |S|^{1-r} |T|^{1-s} \right\|^2},$$

where $r, s \in \mathbb{R}$.

Proof. For the first inequality, it is enough to put $f_1(t) = f(t)$, $f_2(t) = \frac{t}{f(t)}$, $g_1(t) = g(t)$ and $g_2(t) = \frac{t}{g(t)}$ ($t > 0$) in Theorem . In particular, if $f(t) = t^r$ and $g(t) = t^s$ ($r, s \in \mathbb{R}$) we get the second inequality. \square

Corollary 2.6. *Let $S, T \in \mathbb{B}(\mathcal{H})$ be invertible normal and f, g be two positive continuous functions on $[0, \infty)$. Then*

$$\|S + T\| \leq \max\{\|S\| + \|T\|\} + \|f(|S|)g(|T|)\|^{\frac{1}{2}} \left\| |S|(f(|S|))^{-1}|T|(g(|T|))^{-1} \right\|^{\frac{1}{2}}.$$

In particular,

$$\|S + T\| \leq \max\{\|S\|, \|T\|\} + \left\| |S|^r |T|^{1-s} \right\|^{\frac{1}{2}} \left\| |S|^{1-r} |T|^s \right\|^{\frac{1}{2}}, \quad (2.1)$$

where $r, s \in \mathbb{R}$.

Proof. If $S, T \in \mathbb{B}(\mathcal{H})$ is invertible normal and f, g are two positive continuous functions on $[0, \infty)$, then by using Theorem we have

$$\begin{aligned} & \|S + T\| \\ & \leq \frac{1}{2}(\|S\| + \|T\|) + \frac{1}{2} \sqrt{(\|S\| - \|T\|)^2 + \left\| \frac{1}{t} f(|S|)g(|T|) + t|S|(f(|S|))^{-1}|T|(g(|T|))^{-1} \right\|^2} \\ & \leq \frac{1}{2}(\|S\| + \|T\|) + \frac{1}{2} \left| \|S\| - \|T\| \right| + \frac{1}{2} \left\| \frac{1}{t} f(|S|)g(|T|) + t|S|(f(|S|))^{-1}|T|(g(|T|))^{-1} \right\| \\ & \quad \text{(by the inequality } \sqrt{x^2 + y^2} \leq \sqrt{x^2} + \sqrt{y^2}\text{)} \\ & = \max\{\|S\|, \|T\|\} + \left\| \frac{1}{t} f(|S|)g(|T|) + t|S|(f(|S|))^{-1}|T|(g(|T|))^{-1} \right\| \\ & \leq \max\{\|S\|, \|T\|\} + \frac{1}{2} \left(\frac{1}{t} \|f(|S|)g(|T|)\| + t \left\| |S|(f(|S|))^{-1}|T|(g(|T|))^{-1} \right\| \right). \end{aligned}$$

Now, we take the minimum on t ($t > 0$) and we get

$$\begin{aligned} & \|S + T\| \\ & \leq \max\{\|S\|, \|T\|\} + \frac{1}{2} \min_{t>0} \left(\frac{1}{t} \|f(|S|)g(|T|)\| + t \left\| |S|(f(|S|))^{-1}|T|(g(|T|))^{-1} \right\| \right) \\ & = \max\{\|S\|, \|T\|\} + \|f(|S|)g(|T|)\|^{\frac{1}{2}} \left\| |S|(f(|S|))^{-1}|T|(g(|T|))^{-1} \right\|^{\frac{1}{2}}. \end{aligned}$$

For the second inequality, put $f(t) = t^r$ and $g(t) = t^{1-s}$ ($r, s \in \mathbb{R}$) in the first inequality. \square

Remark 2.7. If $S, T \in \mathbb{B}(\mathcal{H})$ are normal operator, then inequality is a refinement and generalization of inequality for normal operators. In fact, in this case

$$\begin{aligned} \|S + T\| & \leq \max\{\|S\|, \|T\|\} + \left\| |S|^r |T|^{1-s} \right\|^{\frac{1}{2}} \left\| |S|^{1-r} |T|^s \right\|^{\frac{1}{2}} \\ & \leq \max\{\|S\|, \|T\|\} + \frac{1}{2} \left(\left\| |S|^r |T|^{1-s} \right\| + \left\| |S|^{1-r} |T|^s \right\| \right) \\ & \quad \text{(by the arithmetic-geometric inequality),} \end{aligned}$$

where $r, s \in [0, 1]$.

In the next result, we show a generalization of the inequality for arbitrary operators in $S, T \in \mathbb{B}(\mathcal{H})$.

Theorem 2.8. *Let $S, T \in \mathbb{B}(\mathcal{H})$ and f_1, f_2, g_1, g_2 be non-negative continuous functions on $[0, \infty)$, in which $f_1(x)f_2(x) = x$ and $g_1(x)g_2(x) = x$ ($x \geq 0$). Then*

$$\|S + T\| \leq \frac{1}{2}(\|S\| + \|T\|) + \frac{1}{2}\sqrt{(\|S\| - \|T\|)^2 + \max\{\alpha, \beta\}},$$

where $\alpha = \left\| \frac{1}{t}f_1(|S^*|)g_1(|T^*|) + tf_2(|S^*|)g_2(|T^*|) \right\|^2$ and $\beta = \left\| \frac{1}{t}f_1(|S|)g_1(|T|) + tf_2(|S|)g_2(|T|) \right\|^2$.

Proof. Let $\tilde{S} = \begin{bmatrix} 0 & S \\ S^* & 0 \end{bmatrix}$ and $\tilde{T} = \begin{bmatrix} 0 & T \\ T^* & 0 \end{bmatrix}$. Then \tilde{S} and \tilde{T} are normal operators and

$$\|\tilde{S} + \tilde{T}\| = \left\| \begin{bmatrix} 0 & S + T \\ S^* + S^* & 0 \end{bmatrix} \right\| = \|S + T\|.$$

Hence, applying Theorem we get

$$\begin{aligned} \|S + T\| &= \|\tilde{S} + \tilde{T}\| \\ &\leq \frac{1}{2}(\|\tilde{S}\| + \|\tilde{T}\|) + \frac{1}{2}\sqrt{(\|\tilde{S}\| - \|\tilde{T}\|)^2 + \left\| \frac{1}{t}f_1(|\tilde{S}|)g_1(|\tilde{T}|) + tf_2(|\tilde{S}|)g_2(|\tilde{T}|) \right\|^2}. \end{aligned} \quad (2.2)$$

Moreover, it follows from $\|\tilde{S}\| = \|S\|$, $\|\tilde{T}\| = \|T\|$ and

$$f(|\tilde{S}|) = \begin{bmatrix} f(|S^*|) & 0 \\ 0 & f(|S|) \end{bmatrix}, \quad f(|\tilde{T}|) = \begin{bmatrix} f(|T^*|) & 0 \\ 0 & f(|T|) \end{bmatrix}$$

that we get

$$\begin{aligned} &\left\| \frac{1}{t}f_1(|\tilde{S}|)g_1(|\tilde{T}|) + tf_2(|\tilde{S}|)g_2(|\tilde{T}|) \right\| \\ &= \left\| \begin{bmatrix} \frac{1}{t}f_1(|S^*|)g_1(|T^*|) + tf_2(|S^*|)g_2(|T^*|) & 0 \\ 0 & \frac{1}{t}f_1(|S|)g_1(|T|) + tf_2(|S|)g_2(|T|) \end{bmatrix} \right\| \\ &= \max \left\{ \left\| \frac{1}{t}f_1(|S^*|)g_1(|T^*|) + tf_2(|S^*|)g_2(|T^*|) \right\|, \left\| \frac{1}{t}f_1(|S|)g_1(|T|) + tf_2(|S|)g_2(|T|) \right\| \right\}. \end{aligned}$$

Using the recent equality and inequality we reach the desired result. \square

Applying Theorem and a same argument in the proof of Corollary we get the next result.

Corollary 2.9. *Let $S, T \in \mathbb{B}(\mathcal{H})$. Then*

$$\begin{aligned} &\|S + T\| \\ &\leq \max\{\|S\|, \|T\|\} + \max \left\{ \left\| |S^*|^r |T^*|^{1-s} \right\|^{\frac{1}{2}} \left\| |S^*|^{1-r} |T^*|^s \right\|^{\frac{1}{2}}, \left\| |S|^r |T|^{1-s} \right\|^{\frac{1}{2}} \left\| |S|^{1-r} |T|^s \right\|^{\frac{1}{2}} \right\}, \end{aligned}$$

where $r, s \in [0, 1]$. In particular,

$$\|S + T\| \leq \max\{\|S\|, \|T\|\} + \max \left\{ \left\| |S^*|^{\frac{1}{2}} |T^*|^{\frac{1}{2}} \right\|, \left\| |S|^{\frac{1}{2}} |T|^{\frac{1}{2}} \right\| \right\}. \quad (2.3)$$

3. SOME RESULTS FOR THE NUMERICAL RADIUS

In this section, as application of norm inequalities for sums of operators, we present some inequalities for the numerical radius.

Theorem 3.1. *Let $S, T \in \mathbb{B}(\mathcal{H})$. Then*

$$\begin{aligned} w(S - T) &\geq \max\{2w(S), 2w(T)\} - \max\{\|S\|, \|T\|\} \\ &\quad - \max\left\{\| |S^*|^r |T^*|^{1-s} \|^{1/2} \| |S^*|^{1-r} |T^*|^s \|^{1/2}, \| |S|^r |T|^{1-s} \|^{1/2} \| |S|^{1-r} |T|^s \|^{1/2}\right\}, \end{aligned}$$

where $r, s \in [0, 1]$.

Proof. If $S, T \in \mathbb{B}(\mathcal{H})$, then

$$\begin{aligned} 2 \max\{w(S), w(T)\} &= w\left(\begin{bmatrix} S & 0 \\ 0 & T \end{bmatrix}\right) \\ &= w\left(\begin{bmatrix} S+T & 0 \\ 0 & T+S \end{bmatrix}\right) + w\left(\begin{bmatrix} S-T & 0 \\ 0 & T-S \end{bmatrix}\right) \\ &\leq w\left(\begin{bmatrix} S+T & 0 \\ 0 & T+S \end{bmatrix}\right) + w\left(\begin{bmatrix} S-T & 0 \\ 0 & T-S \end{bmatrix}\right) \\ &= w(S+T) + w(S-T), \end{aligned}$$

whence

$$w(S - T) \geq 2 \max\{w(S), w(T)\} - w(S + T).$$

It follows from $w(S + T) \leq \|S + T\|$ and Corollary that

$$\begin{aligned} w(S - T) &\geq \max\{2w(S), 2w(T)\} - w(S + T) \\ &\geq \max\{2w(S), 2w(T)\} - \|S + T\| \\ &\geq \max\{2w(S), 2w(T)\} - \max\{\|S\|, \|T\|\} \\ &\quad - \max\left\{\| |S^*|^r |T^*|^{1-s} \|^{1/2} \| |S^*|^{1-r} |T^*|^s \|^{1/2}, \| |S|^r |T|^{1-s} \|^{1/2} \| |S|^{1-r} |T|^s \|^{1/2}\right\} \\ &\quad \text{(by Corollary),} \end{aligned}$$

where $r, s \in [0, 1]$. □

Remark 3.2. If $S, T \in \mathbb{B}(\mathcal{H})$ are normal operator, then $|S| = |S^*|$, $|T| = |T^*|$, $w(S) = \|S\|$ and $w(T) = \|T\|$. This concludes that Theorem appear as

$$\|S - T\| \geq \max\{\|S\|, \|T\|\} - \| |S|^r |T|^{1-s} \|^{1/2} \| |S|^{1-r} |T|^s \|^{1/2}.$$

In particular, if S and T are positive, then for $r = s = \frac{1}{2}$ we have

$$\|S - T\| \geq \max\{\|S\|, \|T\|\} - \|S^{\frac{1}{2}} T^{\frac{1}{2}}\|.$$

In the next result, we obtain an upper bound for the numerical radius.

Theorem 3.3. Let $S \in \mathbb{B}(\mathcal{H})$ and f_1, f_2, g_1, g_2 be non-negative continuous functions on $[0, \infty)$, in which $f_1(x)f_2(x) = x$ and $g_1(x)g_2(x) = x$ ($x \geq 0$). Then

$$w(S) \leq \frac{1}{2} \|S\| + \frac{1}{4} \max\{\alpha, \beta\},$$

in which $\alpha = \|f_1(|S^*|)g_1(|S|) + f_2(|S^*|)g_2(|S|)\|$ and $\beta = \|f_1(|S|)g_1(|S^*|) + f_2(|S|)g_2(|S^*|)\|$. In particular,

$$w(S) \leq \frac{1}{2} \|S\| + \frac{1}{4} \|f_1(|S^*|)f_2(|S|) + f_2(|S^*|)f_1(|S|)\|.$$

Proof. Using the of $w(\cdot)$ and Theorem for $t = 1$ we have

$$\begin{aligned} w(S) &= \sup_{\theta \in \mathbb{R}} \|\Re(e^{i\theta} T)\| \\ &= \frac{1}{2} \sup_{\theta \in \mathbb{R}} \|e^{i\theta} S + e^{-i\theta} S^*\| \\ &\leq \frac{1}{4} (\|S\| + \|S^*\|) + \frac{1}{4} \max\{\alpha, \beta\} \quad (\text{by Theorem}) \\ &= \frac{1}{2} \|S\| + \frac{1}{4} \max\{\alpha, \beta\}, \end{aligned}$$

in which $\alpha = \|f_1(|S^*|)g_1(|S|) + f_2(|S^*|)g_2(|S|)\|$ and $\beta = \|f_1(|S|)g_1(|S^*|) + f_2(|S|)g_2(|S^*|)\|$. In the special case for $g_1 = f_2$ and $g_2 = f_1$ we get the second inequality. \square

Theorem 3.4. Let $S \in \mathbb{B}(\mathcal{H})$. Then

$$w(S) \leq \max \{ \|\Re S\|, \|\Im S\| \} + \frac{1}{2} \| |\Re S|^{\frac{1}{2}} |\Im S|^{\frac{1}{2}} \|.$$

Proof. Using inequality and the definition $w(\cdot)$ we have

$$\begin{aligned} w(S) &= \sup_{\alpha^2 + \beta^2 = 1} \|\alpha \Re S + \beta \Im S\| \\ &\leq \sup_{\alpha^2 + \beta^2 = 1} \left(\max \{ \|\alpha \Re S\|, \|\beta \Im S\| \} + \max \left\{ \| |\alpha \Re S|^{\frac{1}{2}} |\beta \Im S|^{\frac{1}{2}} \|, \| |\alpha \Re S|^{\frac{1}{2}} |\beta \Im S|^{\frac{1}{2}} \| \right\} \right) \\ &\quad (\text{by inequality}) \\ &\leq \max \{ \|\Re S\|, \|\Im S\| \} + \sup_{\alpha^2 + \beta^2 = 1} \left(\sqrt{|\alpha\beta|} \| |\Re S|^{\frac{1}{2}} |\Im S|^{\frac{1}{2}} \| \right) \\ &\leq \max \{ \|\Re S\|, \|\Im S\| \} + \frac{1}{2} \| |\Re S|^{\frac{1}{2}} |\Im S|^{\frac{1}{2}} \|. \end{aligned}$$

\square

REFERENCES

1. Y. Al-manasrah and F. Kittaneh, A generalization of two refined Young inequalities, Positivity 19 (2015), no. 4, 757–768.
2. M. Bakherad, Some Berezin number inequalities for operator matrices, Czechoslovak Math. J. 68(4)(2018), 997–1009.
3. M. Bakherad, K. Shebrawi Upper bounds for numerical radius inequalities involving off-diagonal operator matrices, Annals of Functional Analysis. J. 9(3)(2018), 297–309.

4. M. Bakherad, R. Lashkaripour, M. Hajmohamadi, and Ulas Yamanci Complete refinements of the Berezin number inequalities, *J. Math. Inequal.* 4 13 (2019), 1117–1128.
5. F.A. Berezin, Covariant and contravariant symbols for operators, *Math. USSR-Izv.* 6(1972), 1117–1151.
6. F.A. Berezin, Quantizations, *Math. USSR-Izv.* 8(1974), 1109–1163.
7. S.S. Dragomir Some inequalities for the norm and the numerical radius of linear operators in Hilbert spaces, *Tamkang Journal of Mathematics.* 39(1) (2008), 1–7.
8. M.T. Garayev, M. Gürdal, S. and Saltan, Hardy Type Inequality For Reproducing Kernel Hilbert Space Operators and Related Problems, *Positivity*, 21 (4) (2017), 1615–1623.
9. M.T. Garayev, M. Gürdal, and A. Okudan, Hardy-Hilbert’s inequality and a power inequality for Berezin numbers for operators, *Math. Inequal. Appl.*, 19 (3) (2016), 883–891.
10. T. Kosem, Inequalities between $\|f(A + B)\|$ and $\|f(A) + f(B)\|$, *Linear Algebra Appl.* 418 (1)(2006), 153–160.
11. U. Yamanci, M. Tapdigoglu, Some results related to the Berezin number inequalities, *Turkish Journal of Mathematics* 43, (4) (2019), 1940–1952.
12. U. Yamanci, T. Remziye, M. Gurdal, Berezin Number, Gruss-Type Inequalities and Their Applications, *Bulletin of the Malaysian Mathematical Sciences Society* (2019) 1–10.

DEPARTMENT OF MATHEMATICS, FACULTY OF MATHEMATICS, UNIVERSITY OF SISTAN AND BALUCHESTAN, ZAHEDAN, IRAN.

Email address: mojtaba.bakherad@yahoo.com