

THE $T1$ THEOREM FOR THE HILBERT TRANSFORM FAILS WHEN $p \neq 2$

MICHEL ALEXIS, JOSE LUIS LUNA-GARCIA, ERIC T. SAWYER, AND IGNACIO URIARTE-TUERO

ABSTRACT. Given $1 < p < \infty$, $p \neq 2$, we show that the $T1$ theorem for the Hilbert transform fails for L^p , despite holding for $p = 2$. More precisely, we construct a pair of locally finite positive Borel measures (σ, ω) that satisfy the two-tailed \mathcal{A}_p condition, and satisfy both of the L^p -testing conditions for the Hilbert transform H , yet $H_\sigma : L^p(\sigma) \not\rightarrow L^p(\omega)$. In the opposite direction, the $T1$ theorem for the Hilbert transform for $p = 2$ was proved a decade ago in the two part paper [LaSaShUr3];[Lac].

1. INTRODUCTION

The $T1$ conjecture of F. Nazarov, S. Treil, and A. Volberg on the boundedness of the Hilbert transform from one weighted space $L^2(\sigma)$ to another $L^2(\omega)$, was settled affirmatively in the two part paper [LaSaShUr3],[Lac], with the assumption of common point masses subsequently removed by Hytönen in [Hyt2]. One might conjecture that the natural extension of this result to $p \neq 2$ holds, namely that the Hilbert transform H is bounded from $L^p(\sigma)$ to $L^p(\omega)$ with general locally finite positive Borel measures σ and ω if and only if the local testing conditions for H ,

$$(1.1) \quad \int_I |H\mathbf{1}_I\sigma|^p d\omega \leq \mathfrak{T}_{H,p}(\sigma, \omega)^p |I|_\sigma \quad \text{and} \quad \int_I |H\mathbf{1}_I\omega|^{p'} d\sigma \leq \mathfrak{T}_{H,p'}(\omega, \sigma)^{p'} |I|_\omega,$$

both hold, along with the two-tailed Muckenhoupt \mathcal{A}_p condition,

$$(1.2) \quad \mathcal{A}_p(\sigma, \omega) \equiv \left(\int_{\mathbb{R}} \frac{|I|^{p-1}}{[|I| + \text{dist}(x, I)]^p} d\omega \right)^{\frac{1}{p}} \left(\int_{\mathbb{R}} \frac{|I|^{p'-1}}{[|I| + \text{dist}(x, I)]^{p'}} d\sigma \right)^{\frac{1}{p'}} < \infty.$$

The purpose of this paper is to dash this hope by constructing a pair of locally finite positive Borel measures (ω, σ_p) as in [LaSaUr2], but with σ_p adjusted according to p , for which (1.1) and (1.2) both hold, yet H_σ is unbounded, i.e. $\int_{\mathbb{R}} |Hf\sigma|^p d\omega = \infty$ and $\int_{\mathbb{R}} |f|^p d\sigma < \infty$.

Remark 1. *On the other hand, T. Hytönen and E. Vuorinen [HyVu] have recently conjectured an extension to weighted L^p spaces that involves quadratic testing and Muckenhoupt conditions, obtained from testing the ℓ^2 -valued extension of the norm inequality for H . In our counterexample here, we derive the unboundedness of the Hilbert transform from the failure of their quadratic Muckenhoupt condition. In the special case of doubling measures, their conjecture was recently proved, even for a large collection of Calderón-Zygmund operators, by the third author and B. Wick [SaWi], with in fact the usual scalar testing conditions in place of the quadratic testing conditions - but with quadratic Muckenhoupt and weak boundedness conditions similar to those in [HyVu].*

Remark 2. *It was pointed out in a seminal paper [NTV, page 911] on the $T1$ theorem in L^2 spaces for well-localized operators, that for general Haar multipliers, the two weight $T1$ theorem fails for all $p \neq 2$, and a Bellman proof can be found in an unpublished note of F. Nazarov [Naz]. However, Vuorinen has proved the ‘quadratic’ $T1$ theorem for dyadic shifts in L^p spaces [Vuo].*

Denote by H the Hilbert transform of a complex Borel measure ν by

$$H\nu(x) \equiv \int_{\mathbb{R}} \frac{1}{y-x} d\nu(y).$$

E. Sawyer is partially supported by a grant from the National Research Council of Canada.

I. Uriarte-Tuero has been partially supported by grant MTM2015-65792-P (MINECO, Spain), and is partially supported by a grant from the National Research Council of Canada.

Here is the *quadratic* \mathcal{A}_p condition of Hytönen and Vuorinen [HyVu],

$$(1.3) \quad \left\| \left(\sum_{i=1}^{\infty} \left(\mathbf{1}_{I_i} \int_{\mathbb{R} \setminus I_i} \frac{f_i(x)}{|x - c_i|} d\sigma(x) \right)^2 \right)^{\frac{1}{2}} \right\|_{L^p(\omega)} \leq \mathcal{A}_p^{\ell^2}(\sigma, \omega) \left\| \left(\sum_{i=1}^{\infty} f_i^2 \right)^{\frac{1}{2}} \right\|_{L^p(\sigma)},$$

taken over all sequences of intervals $\{I_i\}_{i=1}^M$, and all sequences of functions $\{f_i\}_{i=1}^M$, and all $M \in \mathbb{N}$. The constant $\mathcal{A}_p^{\ell^2}(\sigma, \omega)$ is by definition the best possible. As pointed out in [HyVu], this condition is necessary for the norm inequality. We will construct our counterexample (σ, ω) to the $T1$ theorem for H so as to satisfy the \mathcal{A}_p and testing conditions for (σ, ω) , but *fail* the quadratic \mathcal{A}_p condition.

2. CANTOR CONSTRUCTION

We will use analogues of the measure pairs (σ, ω) and $(\dot{\sigma}, \omega)$ constructed in [LaSaUr2] (that were used to demonstrate the failure of necessity of the pivotal condition). Recall the middle-third Cantor set \mathbf{E} and Cantor measure ω on the closed unit interval $I_1^0 = [0, 1]$. At the k^{th} generation in the construction, there is a collection $\{I_j^k\}_{j=1}^{2^k}$ of 2^k pairwise disjoint closed intervals of length $|I_j^k| = \frac{1}{3^k}$. With $K_k = \bigcup_{j=1}^{2^k} I_j^k$, the Cantor set is defined by $\mathbf{E} = \bigcap_{k=1}^{\infty} K_k = \bigcap_{k=1}^{\infty} \left(\bigcup_{j=1}^{2^k} I_j^k \right)$. The Cantor measure ω is the unique probability measure supported in \mathbf{E} with the property that it is equidistributed among the intervals $\{I_j^k\}_{j=1}^{2^k}$ at each scale k , i.e.

$$(2.1) \quad |I_j^k|_{\omega} = 2^{-k}, \quad k \geq 0, 1 \leq j \leq 2^k.$$

Denote the removed open middle third of I_j^k by G_j^k . Let $z_j^k \in G_j^k$ be the center of the interval G_j^k , which is also the center of the interval I_j^k , and for $1 < p < \infty$, define

$$(2.2) \quad \dot{\sigma}_p = \sum_{k,j} s_j^k \delta_{z_j^k},$$

where the sequence of positive numbers s_j^k is chosen to satisfy the following precursor of the \mathcal{A}_p condition:

$$\frac{(s_j^k)^{p-1} |I_j^k|_{\omega}}{|I_j^k|^p} = 1, \quad s_j^k = 2^{-k} \left(\frac{2}{3} \right)^{kp'} \quad k \geq 0, 1 \leq j \leq 2^k.$$

To define the measure σ_p for the counter example, we modify $\dot{\sigma}_p$ as follows, and for convenience in notation we will drop the subscript p from both $\dot{\sigma}_p$ and σ_p . Shift the point masses $s_j^k \delta_{z_j^k}$ in $\dot{\sigma}$ to new positions $s_j^k \delta_{z_j^k}$ in G_j^k where z_j^k is the unique point z in the open interval G_j^k where $H\omega(z)$ vanishes. Such a point z_j^k exists because $H\omega(z)$ is strictly increasing from $-\infty$ to $+\infty$ on the interval G_j^k . Let

$$(2.3) \quad \sigma = \sum_{k,j} s_j^k \delta_{z_j^k}.$$

In the case $p = 2$, it was shown in [LaSaUr2] that the two-tailed Muckenhoupt condition (1.2), and the forward and backward testing conditions (1.1) hold for both of the measure pairs $(\dot{\sigma}, \omega)$ and (σ, ω) . These proofs extend easily to the case $p \neq 2$, and are left for the reader. For example, the self-similarity argument in [LaSaUr2] gives

$$(2.4) \quad \int |H\dot{\sigma}|^p \omega = \frac{2^{p'-1}}{3^{p'}} (1 + \varepsilon) \int |H\dot{\sigma}|^p \omega + \frac{2^{p'-1}}{3^{p'}} (1 + \varepsilon) \int |H\dot{\sigma}|^p \omega + \mathcal{R}_{\varepsilon},$$

where $\mathcal{R}_{\varepsilon} \leq C_{\varepsilon} \mathcal{A}_p(f\dot{\sigma})$, and hence

$$\int |H\dot{\sigma}|^p \omega = \frac{1}{1 - 2 \frac{2^{p'-1}}{3^{p'}} (1 + \varepsilon)} \mathcal{R}_{\varepsilon} \leq C_{\varepsilon} \mathcal{A}_p^2 \left(\int \dot{\sigma} \right),$$

provided $0 < \varepsilon < \left(\frac{3}{2}\right)^{p'} - 1$.

2.1. **Failure of $\mathcal{A}_p^{\ell^2}(\sigma, \omega)$.** It remains to show that $\mathcal{A}_p^{\ell^2}(\sigma, \omega) = \infty$ for $p \neq 2$. We will show that there is $f = \{f_i\}_{i=1}^{\infty}$ satisfying

$$(2.5) \quad \left\| \left(\sum_{i=1}^{\infty} f_i^2 \right)^{\frac{1}{2}} \right\|_{L^p(\sigma)} < \infty \text{ and } \left\| \left(\sum_{i=1}^{\infty} \left(\mathbf{1}_{I_i} \int_{\mathbb{R} \setminus 3I_i} \frac{f_i(x)}{|x - c_i|} d\sigma(x) \right)^2 \right)^{\frac{1}{2}} \right\|_{L^p(\omega)} = \infty.$$

In fact we will take

$$\{I_i\}_{i=1}^{\infty} \equiv \{I_j^k\}_{k,j} \text{ and } f_i = f_{k,j} \equiv b_j^k \mathbf{1}_{\theta I_j^k},$$

where θI_j^k denotes the triadic sibling of I_j^k , and use the estimates

$$\begin{aligned} |I_j^k| &= 3^{-k}, \quad |I_j^k|_{\omega} = 2^{-k}, \quad |I_j^k|_{\sigma} \approx \widehat{s}_j^k [p] = \frac{2^{k(p'-1)}}{3^{kp'}}, \\ \int_{\mathbb{R} \setminus 3I_j^k} \frac{f_{k,j}(x)}{|x - c_i|} d\sigma(x) &\approx b_j^k \int_{\theta I_j^k} \frac{1}{|I_j^k|} d\sigma(x) \approx b_j^k \frac{|I_j^k|_{\sigma}}{|I_j^k|} \approx b_j^k \frac{2^{k(p'-1)}}{3^{k(p'-1)}}. \end{aligned}$$

We define

$$b_j^k = \beta_k = \frac{1}{(k+1)^{\frac{1}{p}} (\ln(k+2))^{\frac{1+\delta}{p}}} \left(\frac{3}{2} \right)^{(p'-1)k} \equiv \frac{1}{a_{k,p,\delta}} \left(\frac{3}{2} \right)^{(p'-1)k},$$

and obtain

$$\left\| \left(\sum_{i=1}^{\infty} f_i^2 \right)^{\frac{1}{2}} \right\|_{L^p(\sigma)}^p = \int_{\mathbb{R}} \left(\sum_{k,j} \left(\beta_k \mathbf{1}_{\theta I_j^k}(y) \right)^2 \right)^{\frac{p}{2}} d\sigma(y).$$

Set $\Omega_k \equiv \bigcup_{j=1}^{2^k} I_j^k = \bigcup_{j=1}^{2^k} \theta I_j^k$. We have the estimate

$$\begin{aligned} \sum_{k,j} \beta_k^2 \mathbf{1}_{\theta I_j^k}(y) &\approx \sum_{k=0}^{\infty} \beta_k^2 \mathbf{1}_{\Omega_k}(y) = \sum_{k=0}^{\infty} \beta_k^2 \sum_{\ell=k}^{\infty} \mathbf{1}_{\Omega_{\ell} \setminus \Omega_{\ell+1}}(y) \\ &= \sum_{\ell=0}^{\infty} \left(\sum_{k=1}^{\ell} \beta_k^2 \right) \mathbf{1}_{\Omega_{\ell} \setminus \Omega_{\ell+1}}(y) \approx \sum_{\ell=0}^{\infty} \beta_{\ell}^2 \mathbf{1}_{\Omega_{\ell} \setminus \Omega_{\ell+1}}(y), \end{aligned}$$

and so changing dummy variables,

$$\begin{aligned} \int_{\mathbb{R}} \left(\sum_{k,j} \left(b_j^k \mathbf{1}_{\theta I_j^k}(y) \right)^2 \right)^{\frac{p}{2}} d\sigma(y) &\approx \int_{\mathbb{R}} \left(\sum_{k=0}^{\infty} \beta_k^2 \mathbf{1}_{\Omega_k \setminus \Omega_{k+1}}(y) \right)^{\frac{p}{2}} d\sigma(y) \\ &\approx \sum_{k=0}^{\infty} \beta_k^p |\Omega_k \setminus \Omega_{k+1}|_{\sigma} \approx \sum_{k=0}^{\infty} \beta_k^p \sum_{j=1}^{2^k} |I_j^k \setminus \Omega_{k+1}|_{\sigma} = \sum_{k=0}^{\infty} \beta_k^p \sum_{j=1}^{2^k} s_j^k \\ &= \sum_{k=0}^{\infty} \beta_k^p \sum_{j=1}^{2^k} \frac{2^{k(p'-1)}}{3^{kp'}} = \sum_{k=0}^{\infty} \left(\frac{1}{a_{k,p,\delta}} \left(\frac{3}{2} \right)^{(p'-1)k} \right)^p \frac{2^{kp'}}{3^{kp'}} = \sum_{k=0}^{\infty} \frac{1}{(k+1) (\ln(k+2))^{1+\delta}} < \infty. \end{aligned}$$

Next we compute

$$\begin{aligned} \left\| \left(\sum_i \left(\mathbf{1}_{I_i} \int_{\mathbb{R} \setminus 3I_i} \frac{f_i(x)}{|x - c_i|} d\sigma(x) \right)^2 \right)^{\frac{1}{2}} \right\|_{L^p(\omega)}^p &\approx \int_{\mathbb{R}} \left(\sum_{k,j} \left(\mathbf{1}_{I_j^k}(y) \beta_k \frac{2^{k(p'-1)}}{3^{k(p'-1)}} \right)^2 \right)^{\frac{p}{2}} d\omega(y) \\ &\approx \int_{\mathbb{R}} \left(\sum_{k,j} \left(\mathbf{1}_{I_j^k}(y) \frac{1}{a_{k,p,\delta}} \left(\frac{3}{2} \right)^{(p'-1)k} \frac{2^{k(p'-1)}}{3^{k(p'-1)}} \right)^2 \right)^{\frac{p}{2}} d\omega(y), \end{aligned}$$

and truncating the sum in k at N we get

$$\sum_{k \leq N, j} \left(\mathbf{1}_{I_j^k}(y) \frac{1}{a_{k,p,\delta}} \right)^2 = \sum_{k=0}^N \frac{1}{(k+1)^{\frac{2}{p}} (\ln(k+2))^{\frac{2}{p}(1+\delta)}} \mathbf{1}_{\Omega_N}(y) \approx \frac{N^{1-\frac{2}{p}}}{(\ln N)^{\frac{2}{p}(1+\delta)}} \mathbf{1}_{\Omega_N}(y) \text{ for } \omega\text{-a.e. } y,$$

and so

$$\int_{\mathbb{R}} \left(\sum_{k \leq N, j} \left(\mathbf{1}_{I_j^k}(y) \frac{1}{a_{k,p,\delta}} \right)^2 \right)^{\frac{p}{2}} d\omega(y) \approx \int_{\mathbb{R}} \left(\frac{N^{1-\frac{2}{p}}}{(\ln N)^{\frac{2}{p}(1+\delta)}} \mathbf{1}_{\Omega_N}(y) \right)^{\frac{p}{2}} d\omega(y) \approx \frac{N^{\frac{p}{2}-1}}{(\ln N)^{(1+\delta)}} |\Omega_N|_{\omega},$$

which blows up to ∞ as $N \nearrow \infty$ provided $p > 2$.

This shows that the $T1$ theorem fails for the Hilbert transform when $p > 2$, and a duality argument shows the same when $1 < p < 2$.

Remark 3. *The arguments outlined above for the Hilbert transform generalize to individual Riesz transforms R_j in higher-dimensions. Indeed, if considering R_1 in \mathbb{R}^n , take ω to the Cantor measure on the segment $[0, 1] \times \{0\}^{n-1}$, and take δ and σ as in (2.2) and (2.3) with $s_j^k \equiv \frac{2^{k(p'-1)}}{3^{nkp'}}$, z_j^k as the centers of the middle-third line segments G_j^k removed in the construction of ω , and z_j^k defined as any one of the zeros of $R_1\omega(x)$ occurring in G_j^k . Finally, to show the failure of $\mathcal{A}_p^{\ell^2}(\sigma, \omega)$, follow the same proof as outlined above, and take $b_j^k = \beta_k = \frac{1}{k^{\frac{1}{p}} (\ln k)^{\frac{1+\delta}{p}} \left(\frac{3^n}{2}\right)^{(p'-1)k}}$. This implies the norm inequality fails for R_1 : indeed, since σ and ω are supported on the line segment $[0, 1] \times \{0\}^{n-1}$, then for this pair of measures we can show that $\mathcal{A}_p^{\ell^2}$ is also necessary for the norm inequality for the Riesz transform R_1 .*

REFERENCES

- [Hyt2] HYTÖNEN, TUOMAS, *The two weight inequality for the Hilbert transform with general measures*, *arXiv:1312.0843v2*.
- [HyVu] TUOMAS HYTÖNEN AND EMIL VUORINEN, *A two weight inequality between $L^p(\ell^2)$ and L^p* , *arXiv:1608.07199v2*.
- [Lac] MICHAEL T. LACEY, *Two weight inequality for the Hilbert transform: A real variable characterization, II*, *Duke Math. J.* Volume **163**, Number 15 (2014), 2821-2840.
- [LaSaShUr3] MICHAEL T. LACEY, ERIC T. SAWYER, CHUN-YEN SHEN, AND IGNACIO URIARTE-TUERO, *Two weight inequality for the Hilbert transform: A real variable characterization I*, *Duke Math. J.*, Volume **163**, Number 15 (2014), 2795-2820.
- [LaSaUr2] LACEY, MICHAEL T., SAWYER, ERIC T., URIARTE-TUERO, IGNACIO, *A Two Weight Inequality for the Hilbert transform assuming an energy hypothesis*, *Journal of Functional Analysis*, Volume **263** (2012), Issue 2, 305-363.
- [Naz] F. NAZAROV, unpublished manuscript.
- [NTV] F. NAZAROV, S. TREIL AND A. VOLBERG, *The Bellman Functions and Two-Weight Inequalities for Haar Multipliers*, *Journal of the American Mathematical Society*, Oct., 1999, Vol. **12**, No. 4 (Oct., 1999), 909-928.
- [SaWi] ERIC T. SAWYER AND BRETT D. WICK, *Two weight Sobolev norm inequalities for smooth Calderon-Zygmund operators and doubling weights*, *arXiv:2207.14684*.
- [Vuo] EMIL VUORINEN, *Two weight L^p -inequalities for dyadic shifts and the dyadic square function*, *Studia Math.* **237** (1) (2017), 25-56.

DEPARTMENT OF MATHEMATICS & STATISTICS, MCMASTER UNIVERSITY, 1280 MAIN STREET WEST, HAMILTON, ONTARIO, CANADA L8S 4K1

Email address: alexism@mcmaster.ca

DEPARTMENT OF MATHEMATICS & STATISTICS, MCMASTER UNIVERSITY, 1280 MAIN STREET WEST, HAMILTON, ONTARIO, CANADA L8S 4K1

Email address: lunagaj@mcmaster.ca

DEPARTMENT OF MATHEMATICS & STATISTICS, MCMASTER UNIVERSITY, 1280 MAIN STREET WEST, HAMILTON, ONTARIO, CANADA L8S 4K1

Email address: Sawyer@mcmaster.ca

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, ROOM 6290, 40 ST. GEORGE STREET, TORONTO, ONTARIO, CANADA M5S 2E4, (ADJUNCT APPOINTMENT), DEPARTMENT OF MATHEMATICS, 619 RED CEDAR RD., ROOM C212, MICHIGAN STATE UNIVERSITY, EAST LANSING, MI 48824 USA

Email address: ignacio.uriartetuero@utoronto.ca