

From valuations on convex bodies to convex functions

Jonas Knoerr and Jacopo Ulivelli

Abstract

We show how the classification of continuous, epi-translation invariant valuations on convex functions of maximal degree of homogeneity established by Cole-santi, Ludwig, and Mussnig can be obtained from a classical result of McMullen by explicitly relating these functionals to valuations on higher dimensional convex bodies. Following this geometric interpretation, we derive a new description of 1-homogeneous, continuous, and epi-translation invariant valuations on convex functions analogous to a classical result by Goodey and Weil.

1 Introduction

For a family \mathcal{S} of subsets of \mathbb{R}^n , a functional $Y : \mathcal{S} \rightarrow \mathbb{R}$ is a real-valued *valuation* if for every $A, B \in \mathcal{S}$,

$$Y(A \cap B) + Y(A \cup B) = Y(A) + Y(B)$$

whenever $A \cup B, A \cap B \in \mathcal{S}$. This concept goes back to Dehn's solution of Hilbert's third problem and has since then played a central role in convex and discrete geometry (see [39, Chapter 6] for a comprehensive exposition of the subject). Valuations on convex bodies of \mathbb{R}^n , that is, valuations on the space \mathcal{K}^n of all non-empty, convex, and compact subsets of \mathbb{R}^n , have been the focus of intense research ever since, with many break-through results in recent years, see [1, 2, 5, 6, 21, 22, 23, 33]. Most of these results concern valuations on convex bodies that are continuous with respect to the Hausdorff metric.

In recent years, these concepts have been extended to families of functions. Here, a functional $Z : \mathcal{F} \rightarrow \mathbb{R}$ defined on a family \mathcal{F} of extended real-valued functions is called a valuation if

$$Z(u \wedge v) + Z(u \vee v) = Z(u) + Z(v)$$

for every $u, v \in \mathcal{F}$ such that the pointwise minimum $u \wedge v$ and maximum $u \vee v$ belong to \mathcal{F} . If \mathcal{F} denotes the family of indicator functions of convex bodies, this recovers the classical notion of valuations on \mathcal{K}^n . In this sense, valuations on functions generalize valuations on sets. There is, however, also the following more geometric interpretation: Assume that these functions are defined on \mathbb{R}^n and consider for $u \in \mathcal{F}$ its epi-graph

$$\text{epi}(u) := \{(x, t) \in \mathbb{R}^n \times \mathbb{R} : u(x) \leq t\} \subset \mathbb{R}^{n+1}.$$

MSC 2020 Classification: 52B45, 26B25.

Keywords: Convex function, Valuation on functions, Convex body.

It is easy to see that $\text{epi}(u \wedge v) = \text{epi}(u) \cup \text{epi}(v)$, $\text{epi}(u \vee v) = \text{epi}(u) \cap \text{epi}(v)$ for $u, v \in \mathcal{F}$. Thus, valuations on functions correspond to valuations on epi-graphs.

Due to their intimate relation with convex bodies, valuations on various spaces related to convexity have been studied, including log-concave and quasi-concave functions [8, 9, 37]. However, there are also many results concerning other classical function spaces, for example, L_p and Orlicz spaces [28, 29, 41, 42], continuous and Lipschitz functions [18, 19], Sobolev spaces and spaces of functions of bounded variation [30, 31, 32, 34, 43], as well as definable functions [4], or general Banach lattices [40].

Valuations on convex functions on \mathbb{R}^n have been one of the most active areas of study [3, 7, 10, 11, 12, 13, 14, 15, 16, 17, 25, 26, 27]. The primary focus is on subspaces of the space $\text{Conv}(\mathbb{R}^n)$ of all convex functions $u : \mathbb{R}^n \rightarrow (-\infty, +\infty]$ that are lower semi-continuous and proper, that is, not identically $+\infty$. We will equip these spaces with the topology induced by epi-convergence (see Section 2.1 for details). In general, the family $\text{Conv}(\mathbb{R}^n)$ seems to be too large to allow for many meaningful valuations; indeed, under very geometric invariance assumptions, there do not exist any non-constant valuations on this space, as observed, for example, in [14, Section 9].

To obtain a sufficiently interesting set of valuations, one has to focus on smaller spaces of convex functions. One of the natural choices is the space of *super coercive* convex functions used in [11, 13, 14]

$$\text{Conv}_{\text{sc}}(\mathbb{R}^n) := \{u \in \text{Conv}(\mathbb{R}^n) : \lim_{|x| \rightarrow +\infty} u(x)/|x| = +\infty\}$$

together with the space of *finite* convex functions

$$\text{Conv}(\mathbb{R}^n, \mathbb{R}) := \{v \in \text{Conv}(\mathbb{R}^n) : v(x) < +\infty \text{ for all } x \in \mathbb{R}^n\}.$$

These two families are in natural bijection using the *Fenchel-Legendre transform* $u \mapsto u^*$, where

$$u^*(y) = \sup_{x \in \mathbb{R}^n} \{x \cdot y - u(x)\}.$$

It will be advantageous to think of elements of $\text{Conv}_{\text{sc}}(\mathbb{R}^n)$ as unbounded convex sets in \mathbb{R}^{n+1} via their epi-graph, whereas we will consider their Fenchel-Legendre transforms (that is, elements of $\text{Conv}(\mathbb{R}^n, \mathbb{R})$) as the associated support functions. This leads to the following notion, introduced by Colesanti, Ludwig, and Mussnig in [14]: A valuation $Z : \text{Conv}_{\text{sc}}(\mathbb{R}^n) \rightarrow \mathbb{R}$ is called *epi-translation invariant* if

$$Z(u(\cdot - x) + c) = Z(u) \quad \text{for all } u \in \text{Conv}_{\text{sc}}(\mathbb{R}^n), x \in \mathbb{R}^n, c \in \mathbb{R},$$

that is, Z is invariant with respect to translations of the epi-graph of u in \mathbb{R}^{n+1} . Let us similarly define *epi-multiplication* by

$$t \cdot u(x) := tu \left(\frac{x}{t} \right) \quad \text{for } u \in \text{Conv}_{\text{sc}}(\mathbb{R}^n), x \in \mathbb{R}^n, t > 0,$$

that is, the epi-graph of $t \cdot u$ is obtained by rescaling $\text{epi}(u)$ by $t > 0$. We call a valuation $Z : \text{Conv}_{\text{sc}}(\mathbb{R}^n) \rightarrow \mathbb{R}$ epi-homogeneous of degree i if

$$Z(t \cdot u) = t^i Z(u) \quad \text{for all } u \in \text{Conv}_{\text{sc}}(\mathbb{R}^n), t > 0.$$

Obviously, continuous epi-translation invariant valuations on $\text{Conv}_{\text{sc}}(\mathbb{R}^n)$ are closely related to continuous translation invariant valuations on convex bodies. As shown by Colesanti, Ludwig, and Mussnig, this relation can be exploited to establish a homogeneous decomposition theorem for this type of valuations, similar to the corresponding decomposition for translation invariant valuations on convex bodies obtained by McMullen [35].

Theorem 1.1 (Colesanti, Ludwig, and Mussnig [14]). *If $Z : \text{Conv}_{\text{sc}}(\mathbb{R}^n) \rightarrow \mathbb{R}$ is a continuous and epi-translation invariant valuation, then there are continuous and epi-translation invariants valuations $Z_0, \dots, Z_n : \text{Conv}_{\text{sc}}(\mathbb{R}^n) \rightarrow \mathbb{R}$ such that Z_i is epi-homogeneous of degree i and $Z = Z_0 + \dots + Z_n$.*

They were furthermore able to provide a complete characterization of valuations of maximal degree of homogeneity. Let $C_c(\mathbb{R}^n)$ denote the family of continuous functions on \mathbb{R}^n with compact support. For a function $u : \mathbb{R}^n \rightarrow (-\infty, +\infty]$ we also define its *domain*

$$\text{dom}(u) := \{x \in \mathbb{R}^n : u(x) < +\infty\}.$$

Theorem 1.2 (Colesanti, Ludwig, and Mussnig [14]). *A functional $Z : \text{Conv}_{\text{sc}}(\mathbb{R}^n) \rightarrow \mathbb{R}$ is a continuous and epi-translation invariant valuation that is epi-homogeneous of degree n , if and only if there exists $\zeta \in C_c(\mathbb{R}^n)$ such that*

$$Z(u) = \int_{\text{dom}(u)} \zeta(\nabla u(x)) \, dx$$

for every $u \in \text{Conv}_{\text{sc}}(\mathbb{R}^n)$.

A similar characterization holds for continuous, translation invariant valuations on \mathcal{K}^n that are homogeneous of degree $n - 1$. Here $Y : \mathcal{K}^n \rightarrow \mathbb{R}$ is called i -homogeneous if

$$Y(tK) = t^i Y(K) \quad \text{for all } K \in \mathcal{K}^n, t \geq 0,$$

and Y is translation invariant if $Y(K + x) = Y(K)$ for every $K \in \mathcal{K}^n$ and $x \in \mathbb{R}^n$.

Theorem 1.3 (McMullen [36]). *A functional $Y : \mathcal{K}^n \rightarrow \mathbb{R}$ is a continuous, translation invariant valuation which is $(n - 1)$ -homogeneous, if and only if there exists a continuous function $\eta : \mathbb{S}^{n-1} \rightarrow \mathbb{R}$ such that*

$$Y(K) = \int_{\mathbb{S}^{n-1}} \eta(\nu) \, dS_{n-1}(K, \nu) \tag{1}$$

for every $K \in \mathcal{K}^n$. The function η is uniquely determined up to the addition of the restriction of a linear function.

Here, $S_{n-1}(K, \cdot)$ denotes the surface area measure of $K \in \mathcal{K}^n$. By considering the epi-graph of $u \in \text{Conv}_{\text{sc}}(\mathbb{R}^n)$ and a suitable change of coordinates, it is easy to see that the representation in Theorem 1.2 can be interpreted as an integral over the boundary of this epi-graph, which we consider as a convex subset of \mathbb{R}^{n+1} . In this sense, Theorem 1.2 and Theorem 1.3 provide a description of n -homogeneous valuations on convex sets in \mathbb{R}^{n+1} . Although the two classes of sets are disjoint, it is not difficult to assign a convex function in $\text{Conv}_{\text{sc}}(\mathbb{R}^n)$ to any $K \in \mathcal{K}^{n+1}$ in a way that is consistent with this geometric interpretation. This naturally leads to the question of whether Theorem 1.2 can actually be deduced from McMullen's Theorem 1.3. Our first result shows that this is indeed the case (see Theorem 3.1 below): We prove that one can recover the representation in Theorem 1.2 from McMullen's Theorem 1.3, including the properties of the function ζ . More precisely, we exploit a link between valuations on \mathcal{K}^{n+1} and valuations on the family

$$\text{Conv}_{\text{cd}}(\mathbb{R}^n) := \{u \in \text{Conv}_{\text{sc}}(\mathbb{R}^n) : u \text{ has compact domain}\},$$

showing that valuations on this space that are continuous with respect to epi-convergence, epi-translation invariant, as well as epi-homogeneous of degree n , admit a representation as in Theorem 1.2. As this space is dense in $\text{Conv}_{\text{sc}}(\mathbb{R}^n)$, this directly provides the representation in Theorem 1.2. We want to stress that this new proof of Theorem 1.2 gives a classification of valuations on $\text{Conv}_{\text{cd}}(\mathbb{R}^n)$. This can be shown with the original approach in [14] as well.

The idea is the following: Any convex body in \mathcal{K}^{n+1} induces a function in $\text{Conv}_{\text{cd}}(\mathbb{R}^n)$ by considering the lower part of its boundary as the graph of a convex function. In this sense, valuations on $\text{Conv}_{\text{cd}}(\mathbb{R}^n)$ and $\text{Conv}_{\text{sc}}(\mathbb{R}^n)$ may be informally regarded as valuations on convex bodies in \mathbb{R}^{n+1} that do not depend on the top part of the convex bodies, only the lower half. In particular, valuations of this type induce continuous, translation invariant valuations on convex bodies that are invariant under modifications of the convex bodies on their top part. This informal interpretation can be formalized by introducing a suitable notion of support for these functionals, which was extensively used by the first named author in [25] and [26] to obtain a description of a certain dense subspace of so-called smooth valuations. The results in these articles are formulated in the *dual setting*; that is, on the space $\text{Conv}(\mathbb{R}^n, \mathbb{R})$, but can be translated to the more geometric setting of [14] and this article using the Fenchel-Legendre transform. As this relation is even more explicit in the setting of this article, we have included a discussion of this construction which is independent of [25, 26].

In the theory of valuations on convex bodies, another important characterization result was established by Goodey and Weil [20].

Theorem 1.4 (Goodey and Weil). *A functional $Y : \mathcal{K}^n \rightarrow \mathbb{R}$ is a continuous, translation invariant valuation which is homogeneous of degree 1, if and only if there are two sequences of convex bodies $(L_j)_j, (W_j)_j$ in \mathcal{K}^n such that*

$$Y(K) = \lim_{j \rightarrow \infty} [V(K, L_j, \dots, L_j) - V(K, W_j, \dots, W_j)] \quad (2)$$

holds uniformly on compact subsets of \mathcal{K}^n .

Here, $V : (\mathcal{K}^n)^n \rightarrow \mathbb{R}$ denotes the so called *mixed volume*, which is a multilinear functional on \mathcal{K}^n with respect to Minkowski addition (compare [39, Theorem 5.1.7]). Equation (2) can equivalently be written (compare [39, Chapter 5.1]) as

$$\begin{aligned} Y(K) &= \lim_{j \rightarrow \infty} [V(K, L_j, \dots, L_j) - V(K, W_j, \dots, W_j)] \\ &= \lim_{j \rightarrow \infty} \frac{1}{n} \int_{\mathbb{S}^{n-1}} h_K dS_n(L_j) - \frac{1}{n} \int_{\mathbb{S}^{n-1}} h_K dS_n(W_j). \end{aligned}$$

Here $h_K(y) := \max_{x \in K} \langle y, x \rangle$ denotes the *support function* of $K \in \mathcal{K}^n$. Following the analogy between the surface area measure and functionals obtained by integrating the gradient of a convex function, as well as the interpretation of the Fenchel-Legendre transform of an element of $\text{Conv}_{\text{sc}}(\mathbb{R}^n)$ as its support function, we establish the following analogous description for valuations on $\text{Conv}_{\text{sc}}(\mathbb{R}^n)$ that are epi-homogeneous of degree 1.

Theorem 1.5. *Every continuous, epi-translation invariant valuation $Z : \text{Conv}_{\text{sc}}(\mathbb{R}^n) \rightarrow \mathbb{R}$ that is epi-homogeneous of degree 1 can be approximated uniformly on compact subsets of $\text{Conv}_{\text{sc}}(\mathbb{R}^n)$ by a sequence $(Z_j)_j$ of valuations on $\text{Conv}_{\text{sc}}(\mathbb{R}^n)$ with the following properties:*

1. Z_j is a continuous, epi-translation invariant valuation for each $j \in \mathbb{N}$.
2. For every $j \in \mathbb{N}$ there exist two functions $\ell_j, w_j \in \text{Conv}_{\text{cd}}(\mathbb{R}^n)$ such that

$$Z_j(u) = \int_{\text{dom}(\ell_j)} u^*(\nabla \ell_j(x)) dx - \int_{\text{dom}(w_j)} u^*(\nabla w_j(x)) dx$$

for all $u \in \text{Conv}_{\text{cd}}(\mathbb{R}^n)$.

Note that this representation only holds on the dense subspace $\text{Conv}_{\text{cd}}(\mathbb{R}^n)$, as the integrals are not convergent for general elements of $\text{Conv}_{\text{sc}}(\mathbb{R}^n)$. This is a technical artifact of this representation, as the contributions of the two integrals cancel outside of a compact subset. More precisely, [26, Proposition 1.2] implies that the support of the valuations Z_j is *uniformly bounded* in the following sense: There exists a compact subset $A \subset \mathbb{R}^n$ such that for all $u, v \in \text{Conv}_{\text{sc}}(\mathbb{R}^n)$ with $u^* = v^*$ in a neighborhood of A we have $Z_j(u) = Z_j(v)$ for all $j \in \mathbb{N}$. In the proof of this result given in Section 4, Z_j has a simple description that is valid for general elements of $\text{Conv}_{\text{sc}}(\mathbb{R}^n)$. However, we have decided to state the version above to emphasize the analogy with the result by Goodey and Weil.

This paper is structured as follows. In Section 2, we recall some facts about convex functions and discuss the relation between $\text{Conv}_{\text{cd}}(\mathbb{R}^n)$ and \mathcal{K}^{n+1} . In particular, we examine how the surface area measure of a convex body in \mathcal{K}^{n+1} is related to the induced convex function. These results are used in Section 2.3 to show how valuations on $\text{Conv}_{\text{cd}}(\mathbb{R}^n)$ may be interpreted as valuations on \mathcal{K}^{n+1} . In Section 3, we use this correspondence to provide the new proof of Theorem 1.2, together with the analogous characterization on $\text{Conv}_{\text{cd}}(\mathbb{R}^n)$. In Section 4, we prove Theorem 1.5 by using a mollification procedure in the dual setting.

2 Preliminaries

We work in the Euclidean space \mathbb{R}^{n+1} , with $n \geq 1$. The vectors e_1, \dots, e_{n+1} denote its standard orthonormal basis, and we identify $H := \text{span}\{e_1, \dots, e_n\} \subset \mathbb{R}^{n+1}$ with \mathbb{R}^n . We use the following convention throughout this paper: Capital letters will be used for vectors in \mathbb{R}^{n+1} while lowercase letters are reserved for elements of \mathbb{R}^n . Both spaces are endowed with the standard Euclidean norm $|\cdot|$ and the usual scalar product $x \cdot y$ for $x, y \in \mathbb{R}^n$. We will denote the ball with radius $R > 0$ centered at $x \in \mathbb{R}^n$ by $B_R(x)$ or $B_R^n(x)$ if we want to emphasize its dimension.

The space of convex bodies, that is, the set of all non-empty, convex, and compact subsets of \mathbb{R}^{n+1} , will be denoted by \mathcal{K}^{n+1} , which is equipped with the Hausdorff metric. We refer to the monograph by Schneider [39] for a comprehensive background on convex bodies.

Given a convex subset $A \subset \mathbb{R}^n$, let $I_A : \mathbb{R}^n \rightarrow (-\infty, +\infty]$ denote the convex *indicator function* of A , that is,

$$I_A(x) = \begin{cases} 0 & x \in A, \\ +\infty & \text{otherwise.} \end{cases}$$

Note that if $K \in \mathcal{K}^n$, then $I_K \in \text{Conv}_{\text{sc}}(\mathbb{R}^n)$. Moreover, $(I_K)^* = h_K$.

2.1 Convex functions

The spaces of functions used in this paper are all subsets of the space

$$\text{Conv}(\mathbb{R}^n) := \{f : \mathbb{R}^n \rightarrow (-\infty, +\infty] : f \text{ is convex, proper and l.s.c.}\}.$$

For a comprehensive source on convex functions, we refer to the monograph by Rockafellar [38]. The space $\text{Conv}(\mathbb{R}^n)$ is equipped with a natural topology induced by epi-convergence, where a sequence $(u_j)_j \subset \text{Conv}(\mathbb{R}^n)$ is called *epi-convergent* to $u \in \text{Conv}(\mathbb{R}^n)$ if for every $x \in \mathbb{R}^n$ the following conditions hold:

1. For every sequence $(x_j)_j$ that converges to x , $u(x) \leq \liminf_{j \rightarrow \infty} u_j(x_j)$.
2. There exists a sequence $(x_j)_j$ converging to x such that $u(x) = \lim_{j \rightarrow \infty} u_j(x_j)$.

In fact, this topology is metrizable, compare [38, Theorem 7.58]. For the space $\text{Conv}_{\text{sc}}(\mathbb{R}^n)$, epi-convergence may be equivalently characterized by Hausdorff convergence of the level sets

$$\{u \leq t\} := \{x \in \mathbb{R}^n : u(x) \leq t\} \quad \text{for } u \in \text{Conv}_{\text{sc}}(\mathbb{R}^n), t \in \mathbb{R}.$$

We will say that $\{u_j \leq t\} \rightarrow \emptyset$ if there exists j_0 such that $\{u_j \leq t\} = \emptyset$ for all $j \geq j_0$.

Lemma 2.1 ([14] Lemma 10). *A sequence of functions $(u_j) \subset \text{Conv}_{\text{sc}}(\mathbb{R}^n)$ epi-converges to $u \in \text{Conv}_{\text{sc}}(\mathbb{R}^n)$ if and only if $\{u_k \leq t\} \rightarrow \{u \leq t\}$ for every $t \in \mathbb{R}$ with $t \neq \min_{x \in \mathbb{R}^n} u(x)$.*

2.2 The space $\text{Conv}_{\text{cd}}(\mathbb{R}^n)$.

Recall that our first result concerns valuations on the space

$$\text{Conv}_{\text{cd}}(\mathbb{R}^n) := \{u \in \text{Conv}_{\text{sc}}(\mathbb{R}^n) : u \text{ has compact domain}\}.$$

These functions can be obtained from convex bodies in \mathbb{R}^{n+1} using the following construction: To every $K \in \mathcal{K}^{n+1}$ we associate the function $\lfloor K \rfloor : \mathbb{R}^n \rightarrow [-\infty, +\infty]$ defined by

$$\lfloor K \rfloor(x) = \inf\{t \in \mathbb{R} : (x, t) \in K\}.$$

In particular, $\lfloor K \rfloor(x) = +\infty$ if and only if $(x, t) \notin K$ for all $t \in \mathbb{R}$.

Lemma 2.2. $\lfloor K \rfloor \in \text{Conv}_{\text{cd}}(\mathbb{R}^n)$ for all $K \in \mathcal{K}^{n+1}$.

Proof. Note first that $\lfloor K \rfloor$ is bounded from below by

$$\inf\{t \in \mathbb{R} : (x, t) \in K \text{ for some } x \in \mathbb{R}^n\},$$

which is finite due to the compactness of K . In particular, $\lfloor K \rfloor(x) \in (-\infty, +\infty]$ for $x \in \mathbb{R}^n$.

Assume that $x \in \mathbb{R}^n$ satisfies $\lfloor K \rfloor(x) < +\infty$. As K is compact, this implies that the infimum is attained, so $(x, \lfloor K \rfloor(x)) \in K$ in this case. This implies that

$$\text{dom}\lfloor K \rfloor = \text{pr}_H(K)$$

where $\text{pr}_H : \mathbb{R}^{n+1} \rightarrow H \cong \mathbb{R}^n$ denotes the natural projection. In particular, $\text{dom}\lfloor K \rfloor$ is compact and non-empty, so $\lfloor K \rfloor$ is proper.

Let us show that $\lfloor K \rfloor$ is lower semi-continuous. If $x \in \text{dom}(\lfloor K \rfloor)$ and $(x_j)_j$ is a sequence in $\text{dom}(\lfloor K \rfloor)$ converging to x , then $(x_j, \lfloor K \rfloor(x_j)) \in K$ for all $j \in \mathbb{N}$. This sequence is bounded in \mathbb{R}^{n+1} , so $t := \liminf_{j \rightarrow \infty} \lfloor K \rfloor(x_j)$ exists and is finite. Thus, (x, t) is a limit point of the sequence $(x_j, \lfloor K \rfloor(x_j))$ and therefore belongs to K . In particular,

$$\lfloor K \rfloor(x) \leq t = \liminf_{j \rightarrow \infty} \lfloor K \rfloor(x_j).$$

On the other hand, $x \in \mathbb{R}^n \setminus \text{dom}(\lfloor K \rfloor)$ implies that $\lfloor K \rfloor$ is equal to $+\infty$ on a neighborhood of x , as the domain is closed. Thus $\lfloor K \rfloor$ is lower semi-continuous.

It is easy to see that $\lfloor K \rfloor$ is convex. Thus, $\lfloor K \rfloor \in \text{Conv}_{\text{cd}}(\mathbb{R}^n)$ for all $K \in \mathcal{K}^{n+1}$. □

Corollary 2.3. $\text{Conv}_{\text{cd}}(\mathbb{R}^n) \subset \text{Conv}_{\text{sc}}(\mathbb{R}^n)$ is dense.

Proof. For $u \in \text{Conv}_{\text{sc}}(\mathbb{R}^n)$, set $u_j := \lfloor \text{epi}(u) \cap (B_j^n(0) \times [-j, j]) \rfloor$. Then $u_j \in \text{Conv}_{\text{cd}}(\mathbb{R}^n)$ for all $j \in \mathbb{N}$ large enough. As u has compact sublevel sets, given $t \in \mathbb{R}$ we have

$$\{u_j \leq t\} = \{u \leq t\} \quad \text{for all } j \in \mathbb{N} \text{ large enough.}$$

Lemma 2.1 thus implies that $(u_j)_j$ converges to u in $\text{Conv}_{\text{sc}}(\mathbb{R}^n)$, which shows the claim. □

Lemma 2.4. *The map $\lfloor \cdot \rfloor : \mathcal{K}^{n+1} \rightarrow \text{Conv}_{\text{cd}}(\mathbb{R}^n)$ is continuous.*

Proof. Consider a sequence $(K_j) \subset \mathcal{K}^{n+1}$ such that $K_j \rightarrow K \in \mathcal{K}^{n+1}$. Then

$$\tilde{K}_j := K_j + [0, e_{n+1}]$$

converges to $\tilde{K} := K + [0, e_{n+1}]$. We may thus choose $R > 0$ such that $\tilde{K}_j, \tilde{K} \subset B_R^n(0) \times [-R, R]$ for all $j \in \mathbb{N}$.

As $\lfloor \tilde{K}_j \rfloor = \lfloor K_j \rfloor$, $\lfloor \tilde{K} \rfloor = \lfloor K \rfloor$ for all $j \in \mathbb{N}$, we obtain

$$\{\lfloor K_j \rfloor \leq t\} = \text{pr}_H(\tilde{K}_j \cap (B_R^n(0) \times [-(R+1), t])).$$

and similarly for the sublevel sets of $\lfloor K \rfloor$. Note that the sets \tilde{K} and $(B_R^n(0) \times [-(R+1), t])$ can not be separated by a hyperplane for $t > \min_{(x,s) \in \tilde{K}} s = \min_{x \in \mathbb{R}^n} \lfloor K \rfloor(x)$. For $t > \min_{x \in \mathbb{R}^n} \lfloor K \rfloor(x)$, [39, Theorem 1.8.10] thus implies $\tilde{K}_j \cap (B_R^n(0) \times [-(R+1), t]) \neq \emptyset$ for almost all $j \in \mathbb{N}$ and

$$\tilde{K}_j \cap (B_R^n(0) \times [-(R+1), t]) \rightarrow \tilde{K} \cap (B_R^n(0) \times [-(R+1), t])$$

for $j \rightarrow \infty$. Applying the natural projection onto H to both sides, we obtain for $t > \min_{x \in \mathbb{R}^n} \lfloor K \rfloor(x)$

$$\{\lfloor K_j \rfloor \leq t\} \rightarrow \{\lfloor K \rfloor \leq t\}.$$

On the other hand, $t < \min_{x \in \mathbb{R}^n} \lfloor K \rfloor(x)$ implies that $\{\lfloor K \rfloor \leq t\} = \emptyset$. Therefore $\{\lfloor K_j \rfloor \leq t\} = \emptyset$ for almost all $j \in \mathbb{N}$, as we may otherwise find a sequence $x_{j_k} \in \mathbb{R}^n$ with

$$(x_{j_k}, \lfloor K_{j_k} \rfloor(x_{j_k})) \in \tilde{K}_{j_k} \cap (B_R^n(0) \times [-(R+1), t]),$$

from which we can construct a limit point $(x, t_0) \in \tilde{K} \cap (B_R^n(0) \times [-(R+1), t])$.

Lemma 2.1 thus implies that $\lfloor K_j \rfloor \rightarrow \lfloor K \rfloor$. As $\text{Conv}_{\text{cd}}(\mathbb{R}^n) \subset \text{Conv}_{\text{sc}}(\mathbb{R}^n)$ is metrizable, this shows that $\lfloor \cdot \rfloor$ is continuous. \square

Conversely, we may associate to any $u \in \text{Conv}_{\text{cd}}(\mathbb{R}^n)$ a convex body in the following way: For $u \in \text{Conv}_{\text{cd}}(\mathbb{R}^n)$, we set $M_u := \max_{x \in \text{dom}(u)} u(x)$, which is finite since the domain of u is compact and u is convex, and define

$$K^u := \text{epi}(u - M_u) \cap R_H(\text{epi}(u - M_u)) + M_u e_{n+1},$$

where R_H is the reflection with respect to H . Obviously, this is a compact and convex set, so $K^u \in \mathcal{K}^{n+1}$. We thus obtain a map

$$\begin{aligned} \text{Conv}_{\text{cd}}(\mathbb{R}^n) &\rightarrow \mathcal{K}^{n+1} \\ u &\mapsto K^u. \end{aligned}$$

By construction $u = \lfloor K^u \rfloor$ for $u \in \text{Conv}_{\text{cd}}(\mathbb{R}^n)$. In particular, $\lfloor \cdot \rfloor : \mathcal{K}^{n+1} \rightarrow \text{Conv}_{\text{cd}}(\mathbb{R}^n)$ is surjective and continuous. It is thus natural to ask whether the map

$$\begin{aligned} \text{Conv}_{\text{cd}}(\mathbb{R}^n) &\rightarrow \mathcal{K}^{n+1} \\ u &\mapsto K^u, \end{aligned}$$

which is a right inverse of $\lfloor \cdot \rfloor$, is also continuous. This is not the case, which can be shown using an example that we are going to exploit in the proof of Theorem 3.1 below (which is in turn taken from [14]).

Informally, the boundary of the epi-graph of $\lfloor K \rfloor$ coincides with the part of the boundary of K that lies above $\text{dom}(\lfloor K \rfloor)$. Let us make this slightly more precise: Consider the lower half-sphere $\mathbb{S}_-^n := \{X \in \mathbb{S}^n : X \cdot e_{n+1} < 0\}$ define the *lower boundary* of K by

$$\partial K_- := \{X \in \partial K : \text{some unit normal to } K \text{ in } X \text{ belongs to } \mathbb{S}_-^n\}.$$

Up to a set of zero n -dimensional Hausdorff measure, we can parametrize ∂K_- using the map

$$\begin{aligned} f_u : \text{dom}(u) &\rightarrow \mathbb{R}^{n+1} \\ x &\mapsto (x, u(x)). \end{aligned}$$

If $\gamma : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$ is bounded and Borel measurable, we can apply the Area formula to obtain

$$\int_{\partial K_-} \gamma(X) d\mathcal{H}(X) = \int_{\text{dom}(u)} \gamma((x, u(x))) \sqrt{1 + |\nabla u(x)|^2} dx, \quad (3)$$

where $\sqrt{1 + |\nabla u(x)|^2}$ is the approximate Jacobian of f_u .

If K is of class C_+^2 , then the *Gauss map* $\nu_K : \partial K \rightarrow \mathbb{S}^n$, that is, the map that associates to $X \in \partial K$ its unique outer normal unit vector establishes a diffeomorphism between ∂K_- and \mathbb{S}_-^n . In this case, we can relate integrals over ∂K_- to integrals with respect to the surface area of K by

$$\int_{\mathbb{S}_-^n} \eta(N) dS_n(K, N) = \int_{\partial K_-} \eta(\nu_K(X)) d\mathcal{H}(X), \quad (4)$$

where $\eta : \mathbb{S}_-^n \rightarrow \mathbb{R}$ is a bounded Borel measurable function, compare (2.61) in [39].

If $u = \lfloor K \rfloor$ for a C_+^2 body K , we may combine the two relations above to obtain for all bounded Borel measurable functions $\eta : \mathbb{S}_-^n \rightarrow \mathbb{R}$

$$\int_{\mathbb{S}_-^n} \eta(N) dS_n(K^u, N) = \int_{\text{dom}(u)} \eta\left(\frac{(\nabla u(x), -1)}{\sqrt{1 + |\nabla u(x)|^2}}\right) \sqrt{1 + |\nabla u(x)|^2} dx. \quad (5)$$

Let us show that this relation holds for arbitrary elements of $\text{Conv}_{\text{cd}}(\mathbb{R}^n)$.

Theorem 2.5. *For every $u \in \text{Conv}_{\text{cd}}(\mathbb{R}^n)$, $\eta : \mathbb{S}_-^n \rightarrow \mathbb{R}$ Borel measurable and bounded*

$$\int_{\mathbb{S}_-^n} \eta(N) dS_n(K^u, N) = \int_{\text{dom}(u)} \eta\left(\frac{(\nabla u(x), -1)}{\sqrt{1 + |\nabla u(x)|^2}}\right) \sqrt{1 + |\nabla u(x)|^2} dx. \quad (6)$$

Proof. Choosing η as the indicator function of a Borel subset of \mathbb{S}_-^n , we may consider both sides of this equation as positive Radon measures on \mathbb{S}_-^n . In particular, it is enough to prove this relation for $\eta \in C_c(\mathbb{S}_-^n)$. If $K \in \mathcal{K}^{n+1}$ is a smooth convex body with positive Gauss curvature, then $u := \lfloor K \rfloor$ satisfies the equation by our previous discussion. If $u \in \text{Conv}_{\text{cd}}(\mathbb{R}^n)$ is an arbitrary function, then we may approximate K^u in the Hausdorff metric by a sequence $(K_j)_j$ of smooth convex bodies with strictly positive Gauss curvature. Let $\eta \in C_c(\mathbb{S}_-^n)$ be given. As the surface area measure is weakly continuous, compare [39, Theorem 4.2.1], we obtain

$$\begin{aligned} \int_{\mathbb{S}_-^n} \eta(N) dS_n(K^u, N) &= \int_{\mathbb{S}_-^n} \eta(N) dS_n(K^u, N) = \lim_{j \rightarrow \infty} \int_{\mathbb{S}_-^n} \eta(N) dS_n(K_j, N) \\ &= \lim_{j \rightarrow \infty} \int_{\mathbb{S}_-^n} \eta(N) dS_n(K_j, N) \\ &= \lim_{j \rightarrow \infty} \int_{\text{dom}(\lfloor K_j \rfloor)} \eta \left(\frac{(\nabla \lfloor K_j \rfloor(x), -1)}{\sqrt{1 + |\nabla \lfloor K_j \rfloor(x)|^2}} \right) \sqrt{1 + |\nabla \lfloor K_j \rfloor(x)|^2} dx. \end{aligned}$$

On the other hand, the map $u \mapsto \int_{\text{dom}(u)} \eta \left(\frac{(\nabla u(x), -1)}{\sqrt{1 + |\nabla u(x)|^2}} \right) \sqrt{1 + |\nabla u(x)|^2} dx$ is continuous with respect to epi-convergence by [14, Proposition 20]. As $\lfloor \cdot \rfloor$ is continuous by Lemma 2.4, we thus obtain

$$\begin{aligned} \int_{\mathbb{S}_-^n} \eta(N) dS_n(K^u, N) &= \lim_{j \rightarrow \infty} \int_{\text{dom}(\lfloor K_j \rfloor)} \eta \left(\frac{(\nabla \lfloor K_j \rfloor(x), -1)}{\sqrt{1 + |\nabla \lfloor K_j \rfloor(x)|^2}} \right) \sqrt{1 + |\nabla \lfloor K_j \rfloor(x)|^2} dx \\ &= \int_{\text{dom}(\lfloor K^u \rfloor)} \eta \left(\frac{(\nabla \lfloor K^u \rfloor(x), -1)}{\sqrt{1 + |\nabla \lfloor K^u \rfloor(x)|^2}} \right) \sqrt{1 + |\nabla \lfloor K^u \rfloor(x)|^2} dx. \end{aligned}$$

As $\lfloor K^u \rfloor = u$, the claim follows. \square

Let us add the following result, which shows that the integrals in Theorem 1.5 are well defined. Note that if we consider the Fenchel-Legendre transform of the function $\lfloor K \rfloor$ associated with $K \in \mathcal{K}^{n+1}$, we obtain

$$h_K(y, -1) = \lfloor K \rfloor^*(y) \quad \text{for } y \in \mathbb{R}^n. \quad (7)$$

Corollary 2.6. *If $u, v \in \text{Conv}_{\text{cd}}(\mathbb{R}^n)$, then $x \mapsto v^*(\nabla u(x))$ is integrable on $\text{dom}(u)$.*

Proof. This follows directly from Theorem 2.5 since support functions are homogeneous of degree 1 and the surface area measure of a convex body is a finite measure. \square

2.3 From valuations on convex functions to valuations on convex bodies

We will use the map $\lfloor \cdot \rfloor : \mathcal{K}^{n+1} \rightarrow \text{Conv}_{\text{cd}}(\mathbb{R}^n)$ to interpret valuations on $\text{Conv}_{\text{cd}}(\mathbb{R}^n)$ as valuations on convex bodies in \mathbb{R}^{n+1} . This is based on the following simple observation.

Lemma 2.7. *If $K, L \in \mathcal{K}^{n+1}$ are such that $K \cup L \in \mathcal{K}^{n+1}$, then*

$$\lfloor K \cap L \rfloor = \lfloor K \rfloor \vee \lfloor L \rfloor, \quad \lfloor K \cup L \rfloor = \lfloor K \rfloor \wedge \lfloor L \rfloor.$$

Proof. By definition

$$\begin{aligned} \lfloor K \cap L \rfloor(x) &= \inf\{t \in \mathbb{R} : (x, t) \in K \cap L\} \\ &\geq \inf\{t \in \mathbb{R} : (x, t) \in K\} \vee \inf\{t \in \mathbb{R} : (x, t) \in L\} = \lfloor K \rfloor(x) \vee \lfloor L \rfloor(x), \\ \lfloor K \cup L \rfloor(x) &= \inf\{t \in \mathbb{R} : (x, t) \in K \cup L\} \\ &\leq \inf\{t \in \mathbb{R} : (x, t) \in K\} \wedge \inf\{t \in \mathbb{R} : (x, t) \in L\} = \lfloor K \rfloor(x) \wedge \lfloor L \rfloor(x). \end{aligned}$$

On the other hand,

$$\text{dom}(\lfloor K \cap L \rfloor) = \text{dom}(\lfloor K \rfloor) \cap \text{dom}(\lfloor L \rfloor), \quad \text{dom}(\lfloor K \cup L \rfloor) = \text{dom}(\lfloor K \rfloor) \cup \text{dom}(\lfloor L \rfloor),$$

as the domains are just the image of the corresponding convex bodies under the natural projection onto $H \cong \mathbb{R}^n$. In particular, both sides of each of the inequalities are finite if and only if one of the two sides is finite. We thus only have to consider points belonging to the corresponding domains. Assume that $\lfloor K \cap L \rfloor(x) < +\infty$. As $\lfloor K \rfloor(x) \vee \lfloor L \rfloor(x) \leq \lfloor K \cap L \rfloor(x) < +\infty$,

$$\begin{aligned} \{(x, t) \in \mathbb{R}^{n+1} : t \in [\lfloor K \rfloor(x), \lfloor K \cap L \rfloor(x)]\} &\subset K, \\ \{(x, t) \in \mathbb{R}^{n+1} : t \in [\lfloor L \rfloor(x), \lfloor K \cap L \rfloor(x)]\} &\subset L \end{aligned}$$

by convexity, as the points corresponding to the boundary points belong to these sets. Thus $(x, \lfloor K \rfloor(x) \vee \lfloor L \rfloor(x)) \in K \cap L$, which implies

$$\lfloor K \cap L \rfloor(x) \leq \lfloor K \rfloor(x) \vee \lfloor L \rfloor(x).$$

Now assume that $\lfloor K \cup L \rfloor(x) < +\infty$. Then $(x, \lfloor K \cup L \rfloor(x)) \in K \cup L$. Without loss of generality, we may assume that $(x, \lfloor K \cup L \rfloor(x)) \in K$, and therefore

$$\lfloor K \rfloor(x) \wedge \lfloor L \rfloor(x) \leq \lfloor K \rfloor(x) \leq \lfloor K \cup L \rfloor(x)$$

by the definition of $\lfloor K \rfloor(x)$. □

Theorem 2.8. For $Z : \text{Conv}_{\text{cd}}(\mathbb{R}^n) \rightarrow \mathbb{R}$ consider $Y : \mathcal{K}^{n+1} \rightarrow \mathbb{R}$ defined by

$$Y(K) = Z(\lfloor K \rfloor).$$

Then Y has the following properties:

1. If Z is a valuation, then so is Y .
2. If Z is continuous, then Y is continuous with respect to the Hausdorff metric.
3. If Z is epi-translation invariant, then Y is translation invariant, that is

$$Y(K + X) = Y(K) \quad \text{for all } K \in \mathcal{K}^{n+1}, X \in \mathbb{R}^{n+1}.$$

4. If Z is epi-homogeneous of degree j , then Y is j -homogeneous, that is,

$$Y(tK) = t^j Y(K) \quad \text{for all } K \in \mathcal{K}^{n+1}, t > 0.$$

Proof. 1. If $K, L \in \mathcal{K}^{n+1}$ satisfy $K \cup L \in \mathcal{K}^{n+1}$, then

$$\lfloor K \cap L \rfloor = \lfloor K \rfloor \vee \lfloor L \rfloor \text{ and } \lfloor K \cup L \rfloor = \lfloor K \rfloor \wedge \lfloor L \rfloor.$$

by Lemma 2.7. Thus

$$\begin{aligned} Y(K \cup L) + Y(K \cap L) &= Z(\lfloor K \cup L \rfloor) + Z(\lfloor K \cap L \rfloor) \\ &= Z(\lfloor K \rfloor \wedge \lfloor L \rfloor) + Z(\lfloor K \rfloor \vee \lfloor L \rfloor) \\ &= Z(\lfloor K \rfloor) + Z(\lfloor L \rfloor) = Y(K) + Y(L). \end{aligned}$$

2. If Z is continuous, then $Y = Z \circ \lfloor \cdot \rfloor$ is continuous due to the continuity of $\lfloor \cdot \rfloor$, compare Lemma 2.4.

3. For $X = (v, c) \in \mathbb{R}^n \times \mathbb{R}$ and $K \in \mathcal{K}^{n+1}$, the definition of $\lfloor K \rfloor$ implies for $x \in \mathbb{R}^n$

$$\begin{aligned} \lfloor K + X \rfloor(x) &= \inf\{s \in \mathbb{R} : (x, s) \in K + X\} = \inf\{s \in \mathbb{R} : (x - v, s - c) \in K\} \\ &= \inf\{s + c : s \in \mathbb{R}, (x - v, s) \in K + X\} \\ &= \lfloor K \rfloor(x - v) + c. \end{aligned}$$

If Z is epi-translation invariant, we obtain

$$Y(K + X) = Z(\lfloor K + X \rfloor) = Z(\lfloor K \rfloor(\cdot - v) + c) = Z(\lfloor K \rfloor) = Y(K).$$

Thus Y is translation invariant.

4. For $t > 0$ we calculate for $x \in \mathbb{R}^n$

$$\begin{aligned} \lfloor tK \rfloor(x) &= \inf\{s \in \mathbb{R} : (x, s) \in tK\} = \inf\left\{s \in \mathbb{R} : \left(\frac{x}{t}, \frac{s}{t}\right) \in K\right\} \\ &= \inf\left\{ts : s \in \mathbb{R}, \left(\frac{x}{t}, s\right) \in K\right\} \\ &= t\lfloor K \rfloor\left(\frac{x}{t}\right). \end{aligned}$$

Thus $\lfloor tK \rfloor = t \cdot \lfloor K \rfloor$, which implies

$$Y(tK) = Z(\lfloor tK \rfloor) = Z(t \cdot \lfloor K \rfloor) = t^j Z(\lfloor K \rfloor) = t^j Y(K)$$

whenever Z is epi-homogeneous of degree j . □

3 New proof of Theorem 1.2

We will deduce the representation formula established in Theorem 1.2 from McMullen's Theorem 1.3. More precisely, we will show that the same representation holds for continuous, epi-translation invariant valuations on $\text{Conv}_{\text{cd}}(\mathbb{R}^n)$ that are epi-homogeneous of degree n . As $\text{Conv}_{\text{cd}}(\mathbb{R}^n) \subset \text{Conv}_{\text{sc}}(\mathbb{R}^n)$ is dense, this directly establishes the representation formula for the corresponding space of valuations on $\text{Conv}_{\text{sc}}(\mathbb{R}^n)$ by continuity.

Theorem 3.1. *Let $Z : \text{Conv}_{\text{cd}}(\mathbb{R}^n) \rightarrow \mathbb{R}$ be a valuation that is continuous, epi-translation invariant, and epi-homogeneous of degree n . Then there exists a unique function $\zeta \in C_c(\mathbb{R}^n)$ such that*

$$Z(u) = \int_{\text{dom}(u)} \zeta(\nabla u(x)) dx \quad (8)$$

for every $u \in \text{Conv}_{\text{cd}}(\mathbb{R}^n)$.

Proof. Given a functional Z with the properties stated above, $Y(K) := Z(\lfloor K \rfloor)$ defines a valuation on \mathcal{K}^{n+1} which is continuous, translation invariant and n -homogeneous by Theorem 2.8. By McMullen's Theorem 1.3 there exists $\eta \in C(\mathbb{S}^n)$ such that

$$Y(K) = \int_{\mathbb{S}^n} \eta(N) dS_n(K, N)$$

for every $K \in \mathcal{K}^{n+1}$. If we define $\tilde{\eta}(N) := [\eta(N) + \eta(R_H N)]/2$, then the valuation

$$\tilde{Y}(K) := \int_{\mathbb{S}^n} \tilde{\eta}(N) dS_n(K, N)$$

thus satisfies

$$Z(u) = Y(K^u) = \tilde{Y}(K^u).$$

We will work with \tilde{Y} and the function $\tilde{\eta} \in C(\mathbb{S}^n)$.

For a convex body K in H and $\ell > 0$, consider the cylinder $C(K, \ell) = K \times [0, \ell] \in \mathcal{K}^{n+1}$. Then by definition $I_K = \lfloor C(K, \ell) \rfloor$, so

$$\begin{aligned} Z(I_K) &= \tilde{Y}(C(K, \ell)) = 2\tilde{\eta}(-e_{n+1})V_n(K) + \ell \int_{\mathbb{S}^n \cap H} \tilde{\eta}(N) dS_n(C(K, \ell), N) \\ &= 2\tilde{\eta}(-e_{n+1})V_n(K) + \ell \int_{\mathbb{S}^{n-1}} \tilde{\eta}(\nu) dS_{n-1}(K, \nu), \end{aligned}$$

where we identify \mathbb{S}^{n-1} and $\mathbb{S}^n \cap H$. As the left-hand side of this equation is independent of $\ell > 0$, we infer that

$$\int_{\mathbb{S}^{n-1}} \tilde{\eta}(\nu) dS_{n-1}(K, \nu) = 0 \quad (9)$$

for every $K \in \mathcal{K}^n$. We may consider the left-hand side of (9) as a valuation on convex bodies in \mathcal{K}^n that is continuous, translation invariant, and $(n-1)$ -homogeneous. As it vanishes identically, McMullen's Theorem 1.3 implies that $\tilde{\eta}|_{\mathbb{S}^n \cap H}$ is the restriction of a linear function to $\mathbb{S}^n \cap H$.

In particular, there exists a linear function $l : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$ such that $\tilde{\eta} + l \equiv 0$ on the equator $\mathbb{S}^n \cap H$, and we set $\hat{\eta} = \tilde{\eta} + \frac{1}{2}[l + l \circ R_H]$. Then $\hat{\eta}$ vanishes on the equator $\mathbb{S}^n \cap H$. Using Theorem 2.5 and the fact that linear functions belong to the kernel of the surface area measure, we obtain for $u \in \text{Conv}_{\text{cd}}(\mathbb{R}^n)$

$$\begin{aligned} Z(u) &= \tilde{Y}(K^u) = \int_{\mathbb{S}^n} \hat{\eta}(N) dS_n(K^u, N) = 2 \int_{\mathbb{S}^n_-} \hat{\eta}(N) dS_n(K^u, N) \\ &= \int_{\text{dom}(u)} 2\hat{\eta} \left(\frac{(\nabla u(x), -1)}{\sqrt{1 + |\nabla u(x)|^2}} \right) \sqrt{1 + |\nabla u(x)|^2} dx, \end{aligned}$$

which for $\zeta(y) := 2\hat{\eta}\left(\frac{(y,-1)}{\sqrt{1+|y|^2}}\right)\sqrt{1+|y|^2}$ gives the desired representation in equation (8). Here we used that $\hat{\eta}$ vanishes on the equator $\mathbb{S}^n \cap H$ and is symmetric with respect to H .

To prove that ζ has compact support, one can use the argument given by Colesanti, Ludwig, and Mussnig in the proof of [14, Proposition 27], which we include for completeness. Suppose by contradiction that the support is not compact. Then we can find a sequence $y_j \in \mathbb{R}^n$ such that $|y_j| \rightarrow \infty$, $\zeta(y_j) \neq 0$ for every $j \in \mathbb{N}$ and

$$\lim_{j \rightarrow \infty} \frac{y_j}{|y_j|} = \nu \in \mathbb{S}^n.$$

Consider the sets

$$B_j := \{x \in y_j^\perp : |x| \leq 1\}, \quad B_\infty := \{x \in \nu^\perp : |x| \leq 1\}$$

and define the cylinders

$$C_j := \left\{ x + t \frac{y_j}{|y_j|} : x \in B_j, t \in \left[0, \frac{1}{|\zeta(y_j)|}\right] \right\}.$$

For $y \in \mathbb{R}^n$ let l_y denote the linear function $x \mapsto x \cdot y$. Consider the sequence

$$u_j = l_{y_j} + I_{C_j}$$

in $\text{Conv}_{\text{cd}}(\mathbb{R}^n)$. By construction, $\min_{x \in \text{dom}(u_j)} u_j(x) = 0$. For $t > 0$, the sublevel sets are given by

$$\{u_j \leq t\} = \left\{ x + t \frac{y_j}{|y_j|} : x \in B_j, s \in \left[0, \min\left\{\frac{t}{|y_j|}, \frac{1}{|\zeta(y_j)|}\right\}\right] \right\},$$

so $\{u_j \leq t\} \rightarrow B_\infty$ in this case. Obviously, the sublevel sets are empty for $t < 0$. Lemma 2.1 thus implies that $(u_j)_j$ converges to I_{B_∞} .

Now note that $S_n(K^{u_j}, \cdot)$ is concentrated on $(H \cap \mathbb{S}^n) \cup \left\{ \frac{(y_j, -1)}{\sqrt{1+|y_j|^2}}, \frac{(y_j, 1)}{\sqrt{1+|y_j|^2}} \right\}$, so

$$\begin{aligned} Z(u_j) &= \int_{\mathbb{S}^n} \hat{\eta}(N) dS_n(K^{u_j}, N) = \left[\hat{\eta}\left(\frac{(y_j, -1)}{\sqrt{1+|y_j|^2}}\right) + \hat{\eta}\left(\frac{(y_j, 1)}{\sqrt{1+|y_j|^2}}\right) \right] \sqrt{1+|y_j|^2} \text{vol}_n(C_j) \\ &= \zeta(y_j) \text{vol}_n(C_j) = \kappa_{n-1}, \end{aligned}$$

because $\hat{\eta}$ is symmetric with respect to H and vanishes on $\mathbb{S}^n \cap H$. Here, vol_n denotes the n -dimensional Lebesgue measure and κ_{n-1} is the volume of the $(n-1)$ -dimensional unit ball. By continuity we obtain

$$Z(I_{B_\infty}) = \lim_{j \rightarrow \infty} Z(u_j) = \kappa_{n-1}.$$

On the other hand, Z is n -homogeneous and B_∞ is a convex body of dimension $n - 1$, so $Z(I_{B_\infty}) = 0$, which is a contradiction. Thus ζ has compact support.

In total, we obtain

$$Z(u) = \int_{\text{dom}(u)} \zeta(\nabla u(x)) dx,$$

where ζ has compact support.

Finally, let us show how one can use McMullen's Theorem 1.3 to see that ζ is uniquely determined by the valuation Z . Let us thus assume that $\zeta, \zeta' \in C_c(\mathbb{R}^n)$ are such that for all $u \in \text{Conv}_{\text{cd}}(\mathbb{R}^n)$

$$Z(u) = \int_{\text{dom}(u)} \zeta(\nabla u(x)) dx = \int_{\text{dom}(u)} \zeta'(\nabla u(x)) dx.$$

Consider the functions $\eta, \eta' \in C(\mathbb{S}_-^n)$ given for $(y, -\sqrt{1 - |y|^2}) \in \mathbb{S}_-^n$, $y \in \{y \in \mathbb{R}^n : |y| < 1\}$, by

$$\begin{aligned} \eta\left(y, -\sqrt{1 - |y|^2}\right) &:= \zeta\left(\frac{y}{\sqrt{1 - |y|^2}}\right) \sqrt{1 - |y|^2}, \\ \eta'\left(y, -\sqrt{1 - |y|^2}\right) &:= \zeta'\left(\frac{y}{\sqrt{1 - |y|^2}}\right) \sqrt{1 - |y|^2}. \end{aligned}$$

As these functions are compactly supported on \mathbb{S}_-^n , we extend them trivially to \mathbb{S}^n . Using Theorem 2.5, we obtain

$$Y(K) = Z(\lfloor K \rfloor) = \int_{\mathbb{S}^n} \eta dS_n(K) = \int_{\mathbb{S}^n} \eta' dS_n(K)$$

for all $K \in \mathcal{K}^{n+1}$. By McMullen's Theorem 1.3, η and η' thus differ by the restriction of a linear function to \mathbb{S}^n . However, they are both equal to 0 on the complement of \mathbb{S}_-^n , and therefore $\eta - \eta'$ vanishes on an open subset. As this difference is the restriction of a linear function, it thus has to vanish identically. In particular, $\eta = \eta'$. \square

Let us add the following observation.

Corollary 3.2. *Let $Z : \text{Conv}_{\text{cd}}(\mathbb{R}^n) \rightarrow \mathbb{R}$ be a continuous and epi-translation invariant valuation that is epi-homogeneous of degree n . Then Z extends uniquely to a continuous valuation on $\text{Conv}_{\text{sc}}(\mathbb{R}^n)$.*

Proof. By Theorem 3.1, any such valuation $Z : \text{Conv}_{\text{cd}}(\mathbb{R}^n) \rightarrow \mathbb{R}$ is given by

$$Z(u) = \int_{\text{dom}(u)} \zeta(\nabla u(x)) dx \quad \text{for } u \in \text{Conv}_{\text{cd}}(\mathbb{R}^n)$$

for some $\zeta \in C_c(\mathbb{R}^n)$. The right-hand side of this equation defines a continuous valuation on $\text{Conv}_{\text{sc}}(\mathbb{R}^n)$ by [14, Proposition 20], which yields the desired continuous extension. As $\text{Conv}_{\text{cd}}(\mathbb{R}^n) \subset \text{Conv}_{\text{sc}}(\mathbb{R}^n)$ is dense, this extension is unique. \square

4 Proof of Theorem 1.5

For the proof of Theorem 1.5, we will switch to the dual setting: Recall that for any functional $Z : \text{Conv}_{\text{sc}}(\mathbb{R}^n) \rightarrow \mathbb{R}$, we may define a functional $Z^* : \text{Conv}(\mathbb{R}^n, \mathbb{R}) \rightarrow \mathbb{R}$ by

$$Z^*(u) := Z(u^*) \quad \text{for } u \in \text{Conv}(\mathbb{R}^n, \mathbb{R}),$$

where u^* denotes the Fenchel-Legendre transform. Then the following holds, compare the discussion in [13, Section 3.1].

- Z is a valuation if and only if Z^* is a valuation.
- Z is continuous if and only if Z^* is continuous.
- Z is epi-translation invariant if and only if Z^* is *dually epi-translation invariant*, that is, invariant with respect to the addition of affine functions to its argument.
- Z is epi-homogeneous of degree i if and only if Z^* is i -homogeneous in the classical sense, that is,

$$Z^*(tu) = t^i Z^*(u) \quad \text{for all } u \in \text{Conv}(\mathbb{R}^n, \mathbb{R}), t \geq 0.$$

Now assume that Z is epi-homogeneous of degree 1. According to [14, Corollary 24], Z^* is then an *additive* valuation, that is,

$$Z^*(u + v) = Z^*(u) + Z^*(v) \quad \text{for all } u, v \in \text{Conv}(\mathbb{R}^n, \mathbb{R}).$$

In [26], this property was used to lift dually epi-translation invariant valuations to *distributions*, that is, continuous linear functionals on the space of smooth functions with compact support. We refer to [24] for a background on distributions.

Theorem 4.1 ([26] Theorem 2). *For every 1-homogeneous, dually epi-translation invariant, continuous valuation $Z : \text{Conv}(\mathbb{R}^n, \mathbb{R}) \rightarrow \mathbb{R}$ there exists a unique distribution $\text{GW}(Z)$ on \mathbb{R}^n with compact support which satisfies*

$$\text{GW}(Z)[u] = Z(u) \quad \text{for all } u \in \text{Conv}(\mathbb{R}^n, \mathbb{R}) \cap C^\infty(\mathbb{R}^n). \quad (10)$$

Note that (10) is well defined due to the compactness of the support. Let us remark that a similar result holds for homogeneous valuations of arbitrary degree of homogeneity, which is based on ideas of Goodey and Weil [20] for translation invariant valuations on convex bodies. We refer to [26] for this more general construction.

For a 1-homogeneous, dually epi-translation invariant, and continuous valuation Z on $\text{Conv}(\mathbb{R}^n, \mathbb{R})$, we define its *support* as

$$\text{supp } Z := \text{supp } \text{GW}(Z).$$

Then this is a compact subset of \mathbb{R}^n which has the property that

$$Z(u) = Z(v) \quad \text{for all } u, v \in \text{Conv}(\mathbb{R}^n, \mathbb{R}) \text{ s.t. } u \equiv v \text{ on a neighborhood of } \text{supp } Z,$$

compare [26, Proposition 6.3]. In particular, the support of such a valuation is a compact subset of \mathbb{R}^n .

The following Lemma can also be deduced from [25, Theorem 1] in combination with Lemma 5.3 of the same article. The proof we give here is self-contained and does not rely on the machinery developed in [25]. It uses a standard convolution argument, which we include for the convenience of the reader.

Lemma 4.2. *Let $Z : \text{Conv}(\mathbb{R}^n, \mathbb{R}) \rightarrow \mathbb{R}$ be a valuation that is continuous, dually epi-translation invariant, and homogeneous of degree 1. If $\text{supp } Z \subset B_R(0)$, then there exists a sequence $(\phi_j)_j$ in $C_c^\infty(\mathbb{R}^n)$ such that*

1. $\text{supp } \phi_j \subset B_{R+1}(0)$ for all $j \in \mathbb{N}$,
2. $\int_{\mathbb{R}^n} \phi_j(x) dx = \int_{\mathbb{R}^n} x_i \phi_j(x) dx = 0$ for all $1 \leq i \leq n$ for all $j \in \mathbb{N}$,

and such that the continuous valuations Z_j given by

$$Z_j(v) := \int_{\mathbb{R}^n} v(x) \phi_j(x) dx$$

are dually epi-translation invariant and converge uniformly to Z on compact subsets of $\text{Conv}(\mathbb{R}^n, \mathbb{R})$.

Proof. Let $T := \text{GW}(Z)$ denote the Goodey-Weil distribution of Z . Fix a non-negative function $\phi \in C_c^\infty(\mathbb{R}^n)$ with $\int_{\mathbb{R}^n} \phi(x) dx = 1$, $\text{supp } \phi \subset B_1(0)$ and consider the convolution $T_j := T * j^n \phi(j \cdot)$ defined by

$$T_j(\psi) = T(\psi * j^n \phi(j \cdot)) = T\left(\int_{\mathbb{R}^n} \psi(x) j^n \phi(j(\cdot - x)) dx\right) \quad \text{for } \psi \in C_c^\infty(\mathbb{R}^n).$$

As T is continuous, we thus obtain

$$T_j(\psi) = \int_{\mathbb{R}^n} \psi(x) j^n T(\phi(j(\cdot - x))) dx \quad \text{for } \psi \in C_c^\infty(\mathbb{R}^n).$$

Let us consider the function

$$\phi_j(x) := j^n T(\phi(j(\cdot - x))).$$

Elementary facts about the convolution of distributions show that $\phi_j \in C_c^\infty(\mathbb{R}^n)$ with $\text{supp } \phi_j \subset B_{R+1}(0)$ for all $j \in \mathbb{N}$. If $l \in C^\infty(\mathbb{R}^n)$ is an affine function, then

$$[l * j^n \phi(j \cdot)](y) = \int_{\mathbb{R}^n} l(y - x) j^n \phi(jx) dx$$

is affine as well. As T has compact support and Z vanishes on affine functions, we obtain

$$\int_{\mathbb{R}^n} l(x) \phi_j(x) dx = T(l * j^n \phi(j \cdot)) = Z(l * j^n \phi(j \cdot)) = 0,$$

which shows the second property. In particular, if we define $Z_j : \text{Conv}(\mathbb{R}^n, \mathbb{R}) \rightarrow \mathbb{R}$ by

$$Z_j(v) := \int_{\mathbb{R}^n} v(x) \phi_j(x) dx,$$

then Z_j is a continuous, dually epi-translation invariant valuation that satisfies $\text{GW}(Z_j) = T_j$. It remains to check that $(Z_j)_j$ converges to Z uniformly on compact subsets. To see this, note that for $v \in \text{Conv}(\mathbb{R}^n, \mathbb{R}) \cap C^\infty(\mathbb{R}^n)$ the function

$$[v * j^n \phi(j \cdot)](y) = \int_{\mathbb{R}^n} v(y-x) j^n \phi(jx) dx,$$

is convex as ϕ is non-negative. Thus, the fact that T has compact support implies that

$$T(v * j^n \phi(j \cdot)) = \int_{\mathbb{R}^n} T(v(\cdot - x)) j^n \phi(jx) dx.$$

In particular, for any $v \in \text{Conv}(\mathbb{R}^n, \mathbb{R}) \cap C^\infty(\mathbb{R}^n)$

$$\begin{aligned} Z_j(v) &= \text{GW}(Z_j)[v] = T_j(v) = \int_{\mathbb{R}^n} T(v(\cdot - x)) j^n \phi(jx) dx \\ &= \int_{\mathbb{R}^n} Z(v(\cdot - x)) j^n \phi(jx) dx. \end{aligned}$$

On $\text{Conv}(\mathbb{R}^n, \mathbb{R})$, the topology induced by epi-convergence coincides with the topology of uniform convergence on compact subsets of \mathbb{R}^n , see [38, Theorem 7.17]. Using this fact, it is easy to see that the map

$$\begin{aligned} \text{Conv}(\mathbb{R}^n, \mathbb{R}) \times \mathbb{R}^n &\rightarrow \text{Conv}(\mathbb{R}^n, \mathbb{R}) \\ (v, x) &\mapsto v(\cdot - x) \end{aligned}$$

is continuous. In particular,

$$v \mapsto \int_{\mathbb{R}^n} Z(v(\cdot - x)) j^n \phi(jx) dx$$

defines a continuous valuation on $\text{Conv}(\mathbb{R}^n, \mathbb{R})$. By continuity, we thus obtain

$$Z_j(v) = \int_{\mathbb{R}^n} Z(v(\cdot - x)) j^n \phi(jx) dx \quad \text{for all } v \in \text{Conv}(\mathbb{R}^n, \mathbb{R}).$$

Let $\epsilon > 0$ be given. Our previous discussion implies

$$\begin{aligned} |Z_j(v) - Z(v)| &= \left| \int_{\mathbb{R}^n} Z(v(\cdot - x)) j^n \phi(jx) dx - \int_{\mathbb{R}^n} Z(v) j^n \phi(jx) dx \right| \\ &\leq \int_{\mathbb{R}^n} |Z(v(\cdot - x)) - Z(v)| j^n \phi(jx) dx. \end{aligned}$$

As the map

$$\begin{aligned} \text{Conv}(\mathbb{R}^n, \mathbb{R}) \times \mathbb{R}^n &\rightarrow \text{Conv}(\mathbb{R}^n, \mathbb{R}) \\ (v, x) &\mapsto v(\cdot - x) \end{aligned}$$

is continuous, it is uniformly continuous on compact subsets. Given a compact subset $K \subset \text{Conv}(\mathbb{R}^n)$, we can thus find $\delta > 0$ such that

$$|Z(v(\cdot - x)) - Z(v)| \leq \epsilon \quad \text{for all } v \in K \text{ and all } x \in \mathbb{R}^n \text{ with } |x| < \delta.$$

As ϕ is supported on $B_1(0)$, $\text{supp } \phi(j \cdot) \subset B_\delta(0)$ for all $j \geq \frac{1}{\delta}$, so

$$|Z_j(v) - Z(v)| \leq \int_{\mathbb{R}^n} |Z(v(\cdot - x)) - Z(v)| j^n \phi(jx) dx \leq \epsilon \quad \text{for all } v \in K \text{ and } j \geq \frac{1}{\delta}.$$

Thus $(Z_j)_j$ converges uniformly to Z on the compact subset $K \subset \text{Conv}(\mathbb{R}^n, \mathbb{R})$, which concludes the proof. \square

Proof of Theorem 1.5. Let $Z : \text{Conv}_{\text{sc}}(\mathbb{R}^n) \rightarrow \mathbb{R}$ be a continuous, epi-translation invariant valuation that is epi-homogeneous of degree 1. Let us thus assume that $\text{supp } Z^* \subset B_R(0)$. By Lemma 4.2, there exists a sequence $\phi_j \in C_c^\infty(\mathbb{R}^n)$ with $\text{supp } \phi_j \subset B_{R+1}(0)$ such that

$$Z_j(u) := \int_{\mathbb{R}^n} u^*(x) \phi_j(x) dx$$

defines a sequence of continuous, epi-translation invariant, and epi-homogeneous valuations of degree 1 that converges uniformly to Z on compact subsets. Here we use that the Legendre transform establishes a homeomorphism between $\text{Conv}_{\text{sc}}(\mathbb{R}^n)$ and $\text{Conv}(\mathbb{R}^n, \mathbb{R})$, so the preimage of any compact subset of $\text{Conv}(\mathbb{R}^n, \mathbb{R})$ under this map is compact.

It remains to see that Z_j has the desired representation on the subspace $\text{Conv}_{\text{cd}}(\mathbb{R}^n)$. Consider the function $b = [B_1^{n+1}(0)] \in \text{Conv}_{\text{cd}}(\mathbb{R}^n)$, that is,

$$b(x) = \begin{cases} 1 - \sqrt{1 - |x|^2} & |x| \leq 1, \\ +\infty & |x| > 1. \end{cases}$$

Then $b^*(x) = \sqrt{1 + |x|^2}$. From a direct calculation one infers that $\det D^2 b^*(x) = (1 + |x|^2)^{-(n/2+1)}$, and using (3) we can thus write for $u \in \text{Conv}_{\text{cd}}(\mathbb{R}^n)$

$$\begin{aligned} Z_j(u) &= \int_{\mathbb{R}^n} u^*(x) \phi_j(x) dx = \int_{\mathbb{R}^n} u^*(x) \phi_j(x) (1 + |x|^2)^{n/2+1} \det D^2 b^*(x) dx \\ &= \int_{\text{dom}(b)} u^*(\nabla b(x)) \phi_j(\nabla b(x)) (1 + |\nabla b(x)|^2)^{n/2+1} dx = \int_{\mathbb{S}_-^n} \frac{u^*(g(N))}{\sqrt{1 + |g(N)|^2}} f_j(N) dN, \end{aligned}$$

where

$$\begin{aligned} g : \mathbb{S}_-^n &\rightarrow H \cong \mathbb{R}^n \\ N &\mapsto \frac{N}{N \cdot e_{n+1}} - e_{n+1} \end{aligned}$$

and $f_j(N) := \phi_j(g(N))(1 + |g(N)|^2)^{n/2+2}$ is a function that is compactly supported on the lower half sphere. We trivially extend f_j to a smooth function on \mathbb{S}^n .

By equation (7), we thus obtain the representation

$$Z_j(u) = \int_{\mathbb{S}^n} h_{K^u}(N) f_j(N) dN.$$

In fact, $Z_j(u) = \int_{\mathbb{S}^n} h_K(N) f_j(N) dN$ for any $K \in \mathcal{K}^{n+1}$ with $h_K = h_{K^u}$ on \mathbb{S}^n_- . As Z_j is epi-translation invariant and epi-homogeneous of degree 1, we thus obtain

$$0 = Z_j(I_{\{v\}} + c) = \int_{\mathbb{S}^n} h_{\{(v,c)\}}(N) f_j(N) dN \quad \text{for all } (v, c) \in \mathbb{R}^{n+1}.$$

As $h_{\{(v,c)\}}(N) = (v, c)^T \cdot N$, the non-negative measure

$$\mu_j(B) := \int_B (1 + \|f_j\|_\infty + f_j) dN \quad \text{for a Borel subset } B \subset \mathbb{S}^n$$

is thus not concentrated on a great sphere and satisfies

$$\int_{\mathbb{S}^n} N d\mu_j(N) = 0.$$

By Minkowski's existence Theorem (see [39, Section 8.2.1]), there thus exists a convex body $L_j \in \mathcal{K}^{n+1}$ such that $\mu_j = S_n(L_j)$. In particular,

$$Z_j(u) = \int_{\mathbb{S}^n} h_{K^u} f_j(N) dN = \int_{\mathbb{S}^n} h_{K^u} dS_n(L_j) - \int_{\mathbb{S}^n} h_{K^u} dS_n(\sqrt[n]{1 + \|f_j\|_\infty} B_1(0)).$$

Here we have used that the surface area measure on \mathbb{R}^{n+1} is n -homogeneous and that $S_n(B_1(0))$ is the spherical Lebesgue measure. Set $W_j := \sqrt[n]{1 + \|f_j\|_\infty} B_1(0)$. By construction, $S_n(L_j)$ and $S_n(W_j)$ are absolutely continuous with respect to the spherical Lebesgue measure, and their densities only differ on the support of f_j , that is, on a compact subset contained in the lower half sphere. Set $\ell_j = \lfloor L_j \rfloor, w_j = \lfloor W_j \rfloor$. Applying Theorem 2.5 again and using that support functions are 1-homogeneous, we thus obtain for $u \in \text{Conv}_{\text{cd}}(\mathbb{R}^n)$

$$\begin{aligned} Z_j(u) &= \int_{\mathbb{S}^n_-} h_{K^u} dS_n(L_j) - \int_{\mathbb{S}^n_-} h_{K^u} dS_n(W_j) \\ &= \int_{\text{dom}(\ell_j)} u^*(\nabla \ell_j(x)) dx - \int_{\text{dom}(w_j)} u^*(\nabla w_j(x)) dx. \end{aligned}$$

Thus Z_j has the desired representation. In particular, Z can be approximated uniformly on compact subsets by valuations of this type. \square

Using the main results of [26], it is possible to deduce Theorem 1.5 from the characterization of 1-homogeneous valuations on convex bodies by Goodey and Weil in Theorem 2 using the ideas in Section 2.3. To avoid unnecessary technicalities, we have decided to present a more direct argument.

References

- [1] S. ALESKER, *Continuous rotation invariant valuations on convex sets*, Ann. of Math. (2), 149 (1999), pp. 977–1005.
- [2] ———, *Description of translation invariant valuations on convex sets with solution of P. McMullen’s conjecture*, Geom. Funct. Anal., 11 (2001), pp. 244–272.
- [3] S. ALESKER, *Valuations on convex functions and convex sets and Monge-Ampère operators*, Adv. Geom., 19 (2019), pp. 313–322.
- [4] Y. BARYSHNIKOV, R. GHRIST, AND M. WRIGHT, *Hadwiger’s Theorem for definable functions*, Adv. Math., 245 (2013), pp. 573–586.
- [5] A. BERNIG AND J. H. G. FU, *Hermitian integral geometry*, Ann. of Math. (2), 173 (2011), pp. 907–945.
- [6] A. BERNIG, J. H. G. FU, AND G. SOLANES, *Integral geometry of complex space forms*, Geom. Funct. Anal., 24 (2014), pp. 403–492.
- [7] L. CAVALLINA AND A. COLESANTI, *Monotone valuations on the space of convex functions*, Anal. Geom. Metr. Spaces, 3 (2015), pp. 167–211.
- [8] A. COLESANTI AND N. LOMBARDI, *Valuations on the space of quasi-concave functions*, in Geometric aspects of functional analysis, vol. 2169 of Lecture Notes in Math., Springer, Cham, 2017, pp. 71–105.
- [9] A. COLESANTI, N. LOMBARDI, AND L. PARAPATITS, *Translation invariant valuations on quasi-concave functions*, Studia Math., 243 (2018), pp. 79–99.
- [10] A. COLESANTI, M. LUDWIG, AND F. MUSSNIG, *Minkowski valuations on convex functions*, Calc. Var. Partial Differential Equations, 56 (2017), pp. Paper No. 162, 29.
- [11] ———, *Valuations on convex functions*, Int. Math. Res. Not. IMRN, (2019), pp. 2384–2410.
- [12] ———, *The Hadwiger theorem on convex functions, I*, arXiv:2009.03702, (2020).
- [13] A. COLESANTI, M. LUDWIG, AND F. MUSSNIG, *Hessian valuations*, Indiana Univ. Math. J., 69 (2020), pp. 1275–1215.
- [14] ———, *A homogeneous decomposition theorem for valuations on convex functions*, Journal of Functional Analysis, 279 (2020), p. 108573.
- [15] A. COLESANTI, M. LUDWIG, AND F. MUSSNIG, *The Hadwiger theorem on convex functions, II*, arXiv:2109.09434, (2021).

- [16] —, *The Hadwiger theorem on convex functions, III: Steiner formulas and mixed Monge-Ampère measures*, Calc. Var. Partial Differential Equations, 61 (2022), pp. Paper No. 181, 37.
- [17] —, *The Hadwiger theorem on convex functions, IV: The Klain approach*, Adv. Math., 413 (2023), p. Paper No. 108832.
- [18] A. COLESANTI, D. PAGNINI, P. TRADACETE, AND I. VILLANUEVA, *A class of invariant valuations on $Lip(S^{n-1})$* , Adv. Math., 366 (2020), pp. 107069, 37.
- [19] —, *Continuous valuations on the space of Lipschitz functions on the sphere*, J. Funct. Anal., 280 (2021), pp. Paper No. 108873, 43.
- [20] P. GOODEY AND W. WEIL, *Distributions and valuations*, Proc. London Math. Soc. (3), 49 (1984), pp. 504–516.
- [21] C. HABERL, *Minkowski valuations intertwining with the special linear group*, J. Eur. Math. Soc. (JEMS), 14 (2012), pp. 1565–1597.
- [22] C. HABERL AND L. PARAPATITS, *The centro-affine Hadwiger theorem*, J. Amer. Math. Soc., 27 (2014), pp. 685–705.
- [23] —, *Moments and valuations*, Amer. J. Math., 138 (2016), pp. 1575–1603.
- [24] L. HÖRMANDER, *The analysis of linear partial differential operators. I*, Classics in Mathematics, Springer-Verlag, Berlin, 2003. Distribution theory and Fourier analysis, Reprint of the second (1990) edition [Springer, Berlin; MR1065993 (91m:35001a)].
- [25] J. KNOERR, *Smooth valuations on convex functions*, To appear in Journal of Differential Geometry, arXiv:2006.12933.
- [26] J. KNOERR, *The support of dually epi-translation invariant valuations on convex functions*, J. Funct. Anal., 281 (2021), pp. Paper No. 109059, 52.
- [27] J. KNOERR, *Singular valuations and the Hadwiger theorem on convex functions*, arXiv:2209.05158, (2022).
- [28] H. KONE, *Valuations on Orlicz spaces and L^ϕ -star sets*, Adv. in Appl. Math., 52 (2014), pp. 82–98.
- [29] J. LI AND D. MA, *Laplace transforms and valuations*, J. Funct. Anal., 272 (2017), pp. 738–758.
- [30] M. LUDWIG, *Fisher information and matrix-valued valuations*, Adv. Math., 226 (2011), pp. 2700–2711.
- [31] M. LUDWIG, *Valuations on function spaces*, Adv. Geom., 11 (2011), pp. 745–756.
- [32] M. LUDWIG, *Valuations on Sobolev spaces*, Amer. J. Math., 134 (2012), pp. 827–842.

- [33] M. LUDWIG AND M. REITZNER, *A classification of $SL(n)$ invariant valuations*, Ann. of Math. (2), 172 (2010), pp. 1219–1267.
- [34] D. MA, *Real-valued valuations on Sobolev spaces*, Sci. China Math., 59 (2016), pp. 921–934.
- [35] P. McMULLEN, *Valuations and Euler-type relations on certain classes of convex polytopes*, Proc. London Math. Soc. (3), 35 (1977), pp. 113–135.
- [36] ———, *Continuous translation invariant valuations on the space of compact convex sets*, Archiv der Mathematik, 34 (1980), pp. 377–384.
- [37] F. MUSSNIG, *Valuations on log-concave functions*, J. Geom. Anal., 31 (2021), pp. 6427–6451.
- [38] R. ROCKAFELLAR AND R. J. B. WETS, *Variational Analysis*, vol. 317 of Grundlehren der Mathematischen Wissenschaften, Springer Berlin, Heidelberg, 1998.
- [39] R. SCHNEIDER, *Convex Bodies: The Brunn–Minkowski Theory*, Encyclopedia of Mathematics and its Applications, Cambridge University Press, 2 ed., 2013.
- [40] P. TRADACETE AND I. VILLANUEVA, *Valuations on Banach lattices*, Int. Math. Res. Not. IMRN, (2020), pp. 287–319.
- [41] A. TSANG, *Valuations on L^p -spaces*, Int. Math. Res. Not. IMRN, (2010), pp. 3993–4023.
- [42] ———, *Minkowski valuations on L^p -spaces*, Trans. Amer. Math. Soc., 364 (2012), pp. 6159–6186.
- [43] T. WANG, *Semi-valuations on $BV(\mathbb{R}^n)$* , Indiana Univ. Math. J., 63 (2014), pp. 1447–1465.

Jonas Knoerr
 Institute of Discrete Mathematics and Geometry
 TU Wien
 Wiedner Hauptstrasse 8-10, 1040 Wien
 Austria
 jonas.knoerr@tuwien.ac.at

Jacopo Ulivelli
 Dipartimento di Matematica
 Sapienza, University of Rome
 Piazzale Aldo Moro 5, 00185 Rome
 Italy
 jacopo.ulivelli@uniroma1.it