

# CONSTRUCTING ALGEBRAIC SOLUTIONS OF PAINLEVE VI EQUATION FROM $p$ -ADIC HODGE THEORY AND LANGLANDS CORRESPONDENCE

JINBANG YANG AND KANG ZUO

ABSTRACT. We construct infinitely many non-isotrivial families of abelian varieties over given four punctured projective lines. These families lead to algebraic solutions of Painleve VI equation. Finally, based on a recent paper by Lin-Sheng-Wang, we prove a complete characterization for the locus of motivic Higgs bundles in the moduli space as fixed points of an “additive” self-map. This is a note based on the lecture given by the second named author on 04.Nov.2022 at Tsinghua University.

## 0. Introduction

Let  $R$  be a commutative ring with identity and  $X$  be a scheme over  $R$ . An abelian scheme  $A$  over  $X$  together with a polarization  $\mu$  is called of *of  $GL_2$ -type*, if there exists a number field  $K$  of degree  $\dim_X A$  such that the ring of integers  $\mathcal{O}_K$  can be embedded into the endomorphism ring  $\text{End}_\mu(A/X)$ . We will call the abelian scheme *of  $GL_2(K)$ -type*, if we want to emphasize the role of  $K$ .

Let  $f: A \rightarrow X$  be an abelian scheme of  $GL_2(K)$ -type. Let  $D$  denote the discriminant locus and let  $X^0$  denote the complement of  $D$  in  $X$ . Let  $\Delta \subset A$  denote the inverse image of  $D$  under the structure morphism  $f$  and let  $A^0$  denote the complement of  $\Delta$  in  $A$ . Then we obtain the smooth abelian scheme  $f^0: A^0 \rightarrow X^0$ .

$$(0.1) \quad \begin{array}{ccccc} A^0 & \longrightarrow & A & \longleftarrow & \Delta \\ \downarrow f^0 & & \downarrow f & & \downarrow \\ X^0 & \longrightarrow & X & \longleftarrow & D \end{array}$$

For  $R = \mathbb{C}$  we consider the Betti-local system

$$\mathbb{V} = R_{\mathbb{B},*}^1 f^0 \mathbb{Z}_{A^0}$$

attached to  $f^0$ , which is a  $\mathbb{Z}$ -local system over the base  $X^0$ . Since  $f$  is of  $GL_2(K)$ -type, the action of  $\mathcal{O}_K$  on  $f$  induces an action of  $K$  on the  $\overline{\mathbb{Q}}$ -local system  $\mathbb{V} \otimes \overline{\mathbb{Q}}$ . Taking the  $K$ -eigen sheaves decomposition

$$\mathbb{V} \otimes \overline{\mathbb{Q}} = \bigoplus_{i=1}^g \mathbb{L}_i.$$

Then there  $\mathbb{L}_i$ 's are of rank-2 over  $X^0$  and defined over the ring of integers of some number field. On the other hand, consider the logarithmic de Rham bundle attached to the abelian scheme  $f$  and denote

$$(V, \nabla) = R_{\text{dR},*}^1 \left( \Omega_{A/X}^*(\log \Delta), d \right).$$

On this de Rham bundle, there is a canonical filtration satisfying Griffiths transversality given by

$$E^{1,0} := R^0 f_* \Omega_{A/X}^1(\log \Delta) \subset V.$$

Taking the grading with respect to this filtration, one gets a logarithmic graded Higgs bundle, which is so-called Kodaira-Spencer map attached to  $f$

$$(0.2) \quad (E, \theta) := (E^{1,0} \oplus E^{0,1}, \theta) := \text{Gr}_{E^{1,0}}(V, \nabla) = \left( R^0 f_* \Omega_{A/X}^1(\log \Delta) \oplus R^1 f_* \mathcal{O}_A, \text{Gr}(\nabla) \right).$$

Since  $f$  is of  $\text{GL}_2(K)$ -type, one also gets a  $K$ -eigen decomposition of the Higgs bundle

$$(0.3) \quad (E, \theta) = \bigoplus_{i=1}^g (E, \theta)_i.$$

Under Hitchin-Simpson's non-abelian Hodge theory, these eigensheaves  $\{(E, \theta)_i\}_{i=1, \dots, g}$  are just those Higgs bundles correspond to the local systems  $\{\mathbb{L}_i\}_{i=1, \dots, g}$ .

Local systems and Higgs bundles arising as subquotient of local systems and Higgs bundles of some families of varieties are called *motivic*. Sometimes they are also called *coming from geometry origin*. Simpson had found a characterization for a rank-2 local system being motivic:

**Theorem 0.1** (Simpson). *A rank-2 local system  $\mathbb{L}$  is motivic if and only if the following two conditions hold:*

- (1)  $\mathbb{L}$  is defined over the ring of integers of some number field, and
- (2) for each element  $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$  the Higgs bundle corresponding to the Galois conjugation  $\mathbb{L}^\sigma$  is again graded.

**Conjecture 0.2** (Simpson). *A rigid local system is motivic.*

Simpson's conjecture has been proved for the case of rank-2 by Corlette-Simpson [CS08] and rank-3 by Langer-Simpson [LS18] for cohomologically rigid local systems. The conjecture predicts that any rigid local system  $\mathbb{L}$  shall enjoy all properties of motivic local systems. For example,

- its corresponding filtered de Rham bundle is isomorphic to the underlying filtered de Rham bundle of some Fontaine-Faltings modules at almost all places, and
- if  $\mathbb{L}$  is in addition cohomologically rigid, then it is defined over the ring of integers of some number field.

Those two properties have been verified by Esnault-Groechenig recently [EG20], [EG18].

In this note, we take  $X$  as the complex projective line  $\mathbb{P}^1$  and  $D$  as the 4 punctures  $\{0, 1, \infty, \lambda\}$ . Our goal is find some motivic rank-2 logarithmic graded Higgs bundles over  $(\mathbb{P}^1, \{0, 1, \infty, \lambda\})$ , which are coming from  $\text{GL}_2$ -type abelian schemes over  $\mathbb{P}^1$  with discriminant locus contained in  $\{0, 1, \infty, \lambda\}$ .

Beauville has shown that there exist exactly 6 non-isotrivial families of elliptic curves over  $\mathbb{P}^1$  with semistable reductions over  $\{0, 1, \infty, \lambda_i\}$  for  $1 \leq i \leq 6$ . All of them are modular curves with respect to certain mixed level structures. Based on Beauville's result, Viehweg-Zuo have shown that there no more non-isotrivial abelian schemes over  $\mathbb{P}^1$  of

$\mathrm{GL}_2(K)$ -type with  $[K : \mathbb{Q}] \geq 2$  and with semistable reductions over  $\{0, 1, \infty, \lambda\}$ . So except Beavuille's example any non-isotrivial smooth abelian schemes over  $\mathbb{P}^1 \setminus \{0, 1, \infty, \lambda\}$  of  $\mathrm{GL}_2(E)$ -type must have non-semistable reduction at some point in  $\{0, 1, \infty, \lambda\}$ . In this case, the some eigenvalues of the local monodromies of motivic local system must be roots of unity other than 1.

In this note, we consider the simplest situation: motivic rank-2 local systems whose local monodromies around  $\{0, 1, \infty, \lambda\}$  are unipotent and around  $\infty$  is quasiunipotent with eigenvalues  $\{-1, -1\}$ . We call such type local monodromy to be *of type-(1/2) $_\infty$* , or just *of type-(1/2)*.

**Theorem 0.3.** *For given  $\lambda \in \mathbb{P}^1 \setminus \{0, 1, \infty\}$ , there exists infinitely many non-isotrivial abelian schemes of  $\mathrm{GL}_2$ -type over  $\mathbb{P}^1 \setminus \{0, 1, \infty, \lambda\}$  with the associated rank-2 eigen local systems being of type-(1/2) $_\infty$ .*

Let  $M_{0,n}$  denote the moduli space of  $n$ -punctured projective lines and let  $S_{0,n}$  denote the total space of the universal family of  $n$ -punctured projective lines with structure morphism

$$p_n: S_{0,n} \rightarrow M_{0,n}.$$

Then  $M_{0,4} \simeq \mathbb{P}^1 \setminus \{0, 1, \infty\}$ , and  $S_{0,4} = \bigcup_{\lambda \in M_{0,4}} (\mathbb{P}^1 \setminus \{0, 1, \infty, \lambda\})$  is an algebraic surface.

Once we vary the parameter  $\lambda$ , Theorem 0.3 implies the following result:

**Theorem 0.4.** *There exist infinitely many non-isotrivial abelian schemes*

$$f: A \rightarrow \tilde{S}_{0,4}$$

*of  $\mathrm{GL}_2$ -type over fiber products of finite étale base changes*

$$\begin{array}{ccc} \tilde{S}_{0,4} := \tilde{M}_{0,4} \times_{M_{0,4}} S_{0,4} & \longrightarrow & S_{0,4} \\ \downarrow & & \downarrow p_4 \\ \tilde{M}_{0,4} & \xrightarrow{\text{finite étale}} & M_{0,4} \end{array}$$

*and such that the local monodromies of  $f$  around  $\{0, 1, \lambda\}$  are unipotent and around  $\{\infty\}$  is quasi-unipotent with all eigenvalues being  $-1$ .*

**Corollary 0.5** (Corollary 4.). *Let  $f: A \rightarrow \tilde{S}_{0,4}$  be a family given in Theorem 0.4. Then all rank-2 eigen local systems associated to the family  $f$  are algebraic solutions of Painlevé VI equation of the type-(1/2) $_\infty$ .*

For given  $\lambda \in \mathbb{P}^1 \setminus \{0, 1, \infty\}$ , any family  $f_\lambda: A_\lambda \rightarrow \mathbb{P}^1$  in Theorem 0.3 has semistable reduction over  $\{0, 1, \lambda\}$  and potentially semistable reduction over  $\infty$ . Thus the eigen Higgs bundles  $(E, \theta)_i$  associated to this family (constructed in 0.3) have the following form

$$(0.4) \quad E_i = \mathcal{O} \oplus \mathcal{O}(-1), \quad \theta_i: \mathcal{O} \xrightarrow{\neq 0} \mathcal{O}(-1) \otimes \Omega_{\mathbb{P}^1}^1(\log\{0, 1, \infty, \lambda\})$$

and are endowed with natural parabolic structures on the punctures  $\{0, 1, \infty, \lambda\}$  of type-(1/2) $_\infty$ . Here type-(1/2) $_\infty$  parabolic structures means that the parabolic structures at 0, 1 and  $\lambda$  are trivial and the parabolic filtration at  $\infty$  is

$$(E_i |_\infty)_\alpha = \begin{cases} E_i |_\infty & 0 \leq \alpha \leq 1/2, \\ 0 & 1/2 < \alpha < 1. \end{cases}$$

Let  $M_{Hig\lambda}^{gr\frac{1}{2}}$  denote the moduli space of rank-2 semi-stable graded Higgs bundles over  $\mathbb{P}^1$  with the parabolic structure on  $\{0, 1, \infty, \lambda\}$  of type- $(1/2)_\infty$ . Then any Higgs bundle  $(E, \theta) \in M_{Hig\lambda}^{gr\frac{1}{2}}$  is parabolic stable and has the form as in (0.4).

In view of  $p$ -adic Hodge theory, a Higgs bundle  $(E, \theta)$  over the Witt ring  $W(\mathbb{F}_q)$  realized by an abelian scheme over  $W(\mathbb{F}_q)$  of  $GL(E)_2$ -type has to be the grading of an  $K$ -eigen sheaf of the Fontaine-Faltings module attached to the abelian scheme. Hence, by Lan-Sheng-Zuo functor  $(E, \theta)$  is *periodic* on  $M_{Hig\lambda}^{gr\frac{1}{2}}$  over  $W(\mathbb{F}_q)$ .

One identifies

$$M_{Hig\lambda}^{gr\frac{1}{2}} = \mathbb{P}^1$$

by sending  $(E, \theta)$  to the zero locus of the Higgs map  $(\theta)_0 \in \mathbb{P}^1$ , and takes then the elliptic curve  $C_\lambda$  of the Weierstrass form  $y^2 = z(z-1)(z-\lambda)$  as the double cover

$$\pi : C_\lambda \rightarrow \mathbb{P}^1$$

ramified on  $\{0, 1, \infty, \lambda\}$ .

**Conjecture 0.6** (Sun-Yang-Zuo [SYZ17]). *The self-map induced by Higgs-de Rham flow on  $M_{Hig\lambda}^{gr\frac{1}{2}}$  over  $\mathbb{F}_q$  comes from multiplication map by  $p$  on the associated elliptic curve over  $\mathbb{F}_q$*

$$\begin{array}{ccc} C_\lambda & \xrightarrow{[p]} & C_\lambda \\ \downarrow & & \downarrow \\ M_{Hig\lambda}^{gr} & \xrightarrow{\phi} & M_{Hig\lambda}^{gr} \end{array}$$

The conjecture implies two things:

- (1) a Higgs bundle  $(E, \theta)$  is periodic if and only if  $\pi^{-1}(\theta)_0$  is a torsion point in  $C_\lambda$  and of order  $p^f - 1$ .
- (2) for a prime  $p > 2$  and assume  $C_\lambda$  is supersingular then  $\phi_\lambda(z) = z^{p^2}$ . Hence, any Higgs bundle  $(E, \theta) \in M_{Hig\lambda}^{gr\frac{1}{2}}(\overline{\mathbb{F}_q})$  is periodic.

The Conjecture has been checked by Sun-Yang-Zuo for  $p < 50$ . Very recently it has been proved by Lin-Sheng-Wang and becomes a theorem.

**Theorem 0.7** (Lin-Sheng-Wang [LSW22]). *Conjecture 0.6 holds true.*

The technique used in Theorem 0.4 combined with Theorem 0.7 lead us to prove the following result on a characterization for motivic Higgs bundles contained in  $M_{Hig\lambda}^{gr\frac{1}{2}}$ .

**Theorem 0.8.** *A Higgs bundle  $(E, \theta) \in M_{Hig\lambda}^{gr\frac{1}{2}}$  is an eigensheaf of the Kodaira-Spencer map attached to an abelian scheme of  $GL_2$ -type if and only if  $\pi^{-1}(\theta)_0$  is a torsion point in  $C_\lambda$ .*

## 1. DISCUSSION ON THEOREM 0.3

The underlying principle behind Theorem 0.3 is very simple, the so-called isomonodromy deformation for motivic local systems over mixed characteristic. To illustrate the idea, let's first look at the situation over complex numbers. We assume, there exists an abelian scheme  $f_{\lambda_0} : A_{\lambda_0} \rightarrow \mathbb{P}^1$  of  $\mathrm{GL}_2(K)$ -type over complex numbers with bad reduction on  $\{0, 1, \lambda_0, \infty\}$  of type-(1/2). Then the filtered logarithmic de Rham bundle decomposes as  $K$ -eigen sheaves

$$(V, \nabla, E^{1,0}) =: (R_{\mathrm{dr}}^1 f_* \Omega_{A_{\lambda_0}/\mathbb{P}^1}^*(\log \Delta), d), R^0 f_* \Omega_{A_{\lambda_0}/\mathbb{P}^1}^1(\log \Delta)) = \bigoplus_{i=1}^g (V, \nabla, E^{1,0})_i,$$

where each eigen sheaf has the form

$$(V, \nabla, E^{1,0})_i \simeq (\mathcal{O} \oplus \mathcal{O}(-1), \nabla_i, \mathcal{O}).$$

Consider a family of 4-punctured projective line

$$(\mathbb{P}^1, \{0, 1, \lambda, \infty\})_{\hat{U}_{\lambda_0}} \rightarrow \hat{U}_{\lambda_0}$$

over a formal neighborhood  $\hat{U}_{\lambda_0} \subset M_{0,4}$  of  $\lambda_0$ . Then by forgetting the Hodge filtration the de Rham bundle extends to a de Rham bundle  $(V, \nabla)_{\hat{U}_{\lambda_0}}$  over  $(\mathbb{P}^1, \{0, 1, \lambda, \infty\})_{\hat{U}_{\lambda_0}}$ . It is known the abelian scheme extends over  $(\mathbb{P}^1, \{0, 1, \lambda, \infty\})_{\hat{U}_{\lambda_0}}$  if and only if the Hodge filtration  $E^{1,0}$  extends to a sub bundle in the de Rham bundle  $(V, \nabla)_{\hat{U}_{\lambda_0}}$ . Using the  $K$ -eigen sheaf decomposition we see that the obstruction for extending the Hodge filtration  $E^{1,0} = \bigoplus_{i=1}^g \mathcal{O}$  lies in  $\bigoplus_{i=1}^g H^1(\mathbb{P}^1, \mathcal{O}(-1)) = 0$ . Hence, the abelian scheme  $f_{\lambda_0}$  extends over the base  $(\mathbb{P}^1, \{0, 1, \lambda, \infty\})_{\hat{U}_{\lambda_0}}$ .

Back to the situation over mixed characteristic, along the diagram below. One

- starts with moduli space  $M_{\mathrm{Hig}\lambda}^{gr\frac{1}{2}}$  of rank-2 stable graded Higgs bundles on  $\mathbb{P}_{\mathbb{Z}_q}^1$  of parabolic type-(1/2) $_{\infty}$  on  $\{0, 1, \lambda, \infty\}$ ,
- finds periodic Higgs bundles over  $\mathbb{P}_{\mathbb{F}_q}^1$  (i.e. fixed points of iterations of the self map on  $M_{\mathrm{Hig}\lambda}^{gr\frac{1}{2}} \otimes \mathbb{F}_q$  induced by Higgs-de Rham flow),
- gets Fontaine-Faltings modules via Lan-Sheng-Zuo functor from those periodic Higgs bundles and lifts these modules to  $\mathbb{P}_{\mathbb{Z}_q}^1$ ,
- obtains rank-2  $\ell$ -adic local systems on  $\mathbb{P}^1 - \{0, 1, \lambda, \infty\}$  over  $\mathbb{F}_q$  by forgetting Hodge filtration in the Fontaine-Faltings module, tensoring with  $\mathbb{Q}_p$ , and applying Deligne's  $p$ - $\ell$  companion conjecture proven by Abe [Abe13],
- finds  $\mathrm{GL}_2$ -type abelian schemes over  $\mathbb{P}_{\mathbb{F}_q}^1$  with bad reductions of type-(1/2) $_{\infty}$  realizing those  $\ell$ -adic local systems via Drinfeld's theorem on Langlands correspondence.
- modifies those  $\mathrm{GL}_2$ -type abelian schemes up to some  $p$ -isogeny and lifts these modified  $\mathrm{GL}_2$ -type abelian schemes to  $\mathbb{P}_{\mathbb{Z}_q}^1$ .

$$\begin{array}{ccccc}
M_{p\text{-adic}}^{\text{cris}}/\chi & \xleftarrow[p\text{-adic RH}]{\text{Fontaine-Faltings}} & M_{\text{dR}}^{FF}/\chi & \xrightarrow[\text{filtration, } \otimes_{\mathbb{Q}_p}]{\text{forgetting Hodge}} & M^{\text{F-isoc}}/\chi & \xleftarrow[\text{companion by Abe}]{\text{Deligne's } p\text{-}\ell} & M_{\ell\text{-adic}} \\
& & \uparrow \text{Higgs-de Rham flow by Lan-Sheng-Zuo} & & & & \\
& & M_{\text{Higg}}^{\text{per}} & \hookrightarrow & M_{\text{Higg}\lambda}^{\text{gr}\frac{1}{2}} & & 
\end{array}$$

And a type of boundedness and rigidity arguments of classifying maps from the log base curve  $(\mathbb{P}^1, \{0, 1, \lambda, \infty\})$  into the fine moduli space of polarized abelian varieties shows that the abelian scheme lifts over complex numbers for any  $\lambda \in M_{0,4}(\mathbb{C})$ .

Below we give more detailed explanations on the technique issue in each step:

**1.1. Constructing Fontaine-Faltings modules over  $\mathbb{Z}_q$  from semistable Higgs bundles over  $\mathbb{F}_q$  via Higgs de Rham flow.** Take a prime  $p$  and an element  $\lambda \in \mathbb{Z}_{q^2}$  such that the modulo  $p$  reduction of  $C_\lambda$  is supersingular. Consider the moduli space  $M_{\text{Hig}\lambda}^{\text{gr}\frac{1}{2}} \otimes \mathbb{F}_{q^2}$  of rank-2 stable logarithmic graded Higgs bundles over  $(\mathbb{P}^1, D)_{\mathbb{F}_{q^2}}$  with parabolic structures on  $D$  of type-(1/2). Identifying  $M_{\text{Hig}\lambda}^{\text{gr}\frac{1}{2}} \otimes \mathbb{F}_{q^2}$  with the projective line  $\mathbb{P}_{\mathbb{F}_{q^2}}^1$ , the self-map induced by Higgs-de Rham flow is given by  $z \mapsto z^{p^2}$ . We show that, for any  $n \geq 1$ , each  $(\bar{E}, \bar{\theta}) \in M_{\text{Hig}\lambda}^{\text{gr}\frac{1}{2}}(\bar{\mathbb{F}}_{q^{2n}})$  is automatically periodic and lifts uniquely to a periodic Higgs bundle  $(E, \theta) \in M_{\text{Hig}\lambda}^{\text{gr}\frac{1}{2}}(\mathbb{Z}_{q^{2n}})$ . Hence, the Lan-Sheng-Zuo functor induces following bijection

$$M_{\text{Hig}\lambda}^{\text{gr}\frac{1}{2}}(\mathbb{F}_{q^{2n}}) \simeq M_{\text{dR}\lambda}^{\text{FF}\frac{1}{2}}(\mathbb{Z}_{q^{2n}})/\chi,$$

where the right hand side is  $M_{\text{dR}\lambda}^{\text{FF}\frac{1}{2}}(\mathbb{Z}_{q^{2n}})$  modulo an equivalent relation. The  $M_{\text{dR}\lambda}^{\text{FF}\frac{1}{2}}(\mathbb{Z}_{q^{2n}})$  is the set of rank-2 Fontaine-Faltings module over  $(\mathbb{P}^1, D)_{\mathbb{Z}_{q^{2n}}}$  such that the monodromy is of type-(1/2), and two Fontaine-Faltings modules in  $M_{\text{dR}\lambda}^{\text{FF}\frac{1}{2}}(\mathbb{Z}_{q^{2n}})$  are called equivalent if they are differed by a constant rank-1 Fontaine-Faltings modules.

**1.2. Deligne's  $p$ - $\ell$  companion conjecture solved by Abe.** Consider the functors defined by forgetting Hodge filtration and tensoring with  $\mathbb{Q}_p$ , which induce an injective map between a set of logarithmic Fontaine-Faltings modules with parabolic structure-( $\frac{1}{2}$ ) and a set of logarithmic  $F$ -isocrystals. These functors keeps the same type parabolic structure, and the result logarithmic  $F$ -isocrystals have a fixed constant determinant, the  $F$ -isocrystal  $\mathcal{E}_{\text{cy}}$  associated to the cyclotomic character. The injectivity follows the fact that the underlying log-parabolic de Rham bundles are parabolic stable mod  $p$ .

We apply furtherly the forgetful functor by sending logarithmic  $F$ -isocrystals to overconvergent  $F$ -isocrystals, which is known injective by work of Kedlaya [Ked07, Proposition 6.3.2]. Hence, by putting every thing together we obtain an injective map

$$M_{\text{Hig}\lambda}^{\text{gr}\frac{1}{2}}(\mathbb{F}_{q^{2n}}) \simeq M_{\text{dR}\lambda}^{\text{FF}\frac{1}{2}}(\mathbb{Z}_{q^{2n}})/\chi \hookrightarrow M_\lambda^{\text{F-iso}\frac{1}{2}\dagger}(\mathbb{Q}_{q^{2n}})/\chi,$$

where  $M_\lambda^{\text{F-iso}\frac{1}{2}\dagger}(\mathbb{Q}_{q^{2n}})$  is the set of overconvergent  $F$ -isocrystals with parabolic structure of type-(1/2) and determinant  $\mathcal{E}_{\text{cy}}$ .

We choose a prime  $\ell \neq p$  and fix an isomorphism  $\phi : \overline{\mathbb{Q}}_p \simeq \overline{\mathbb{Q}}_\ell$ . Given an overconvergent  $F$ -isocrystal  $(V, \nabla, \Phi)^\dagger \in M_\lambda^{F-iso \frac{1}{2}^\dagger}(\mathbb{Q}_{q^{2n}})$ , by applying Deligne's  $p - \ell$  companion proved by Abe to  $(V, \nabla, \Phi)^\dagger$  we find a rank-2  $\ell$ -adic irreducible local system  $\mathbb{L}$  with cyclotomic determinant corresponding to  $(V, \nabla, \Phi)^\dagger$  in the sense the characteristic polynomial of  $\Phi_x$  on  $V_x$  and the characteristic polynomial of  $\sigma_x$  on  $\mathbb{L}_x$  via  $\phi$  at all close point  $x$  in the close fiber. By the compatibility of local-global Langlands correspondence we find the local monodromies of  $\mathbb{L}$  around  $\{0, 1, \lambda\}$  are unipotent and around  $\infty$  is quasi-unipotent with eigenvalues  $\{-1, -1\}$ . Hence, we obtain a bijective functor

$$M_\lambda^{F-iso \frac{1}{2}^\dagger}(\mathbb{Q}_{q^{2n}})/\chi \simeq M_{\ell\text{-adic } \lambda}^{\frac{1}{2}}(\mathbb{F}_{q^{2n}})/\sim,$$

where  $M_{\ell\text{-adic } \lambda}^{\frac{1}{2}}(\mathbb{F}_{q^{2n}})$  is the set of equivalent classes of rank-2  $\ell$ -adic local systems over  $(\mathbb{P}^1 - D)_{\mathbb{F}_q^{2n}}$  with cyclotomic determinant, unipotent local monodromies around  $\{0, 1, \lambda\}$  and quasi-unipotent local monodromy around  $\infty$  of eigenvalues  $\{-1, -1\}$ . The  $\sim$  stands for an equivalent relation, two local systems  $\mathbb{L}$  and  $\mathbb{L}'$  are called equivalent if the restriction of them on geometric fundamental group are isomorphic. As we show that the restriction of  $\mathbb{L}$  to the geometric fundamental group is automatically irreducible, two local systems  $\mathbb{L}$  and  $\mathbb{L}'$  are equivalent if and only if there are differed by a character of the absolute Galois group of  $\mathbb{F}_{q^{2n}}$ .

Composing the above functors together we obtain an injective functor

$$M_{Hig\lambda}^{gr \frac{1}{2}}(\mathbb{F}_{q^{2n}}) \simeq M_{dR\lambda}^{FF \frac{1}{2}}(\mathbb{Z}_{q^{2n}})/\chi \hookrightarrow M_{\ell\text{-adic } \lambda}^{\frac{1}{2}}(\mathbb{F}_{q^{2n}})/\sim.$$

Over any field  $E$  we have

$$M_{Hig\lambda}^{gr \frac{1}{2}} \otimes E \simeq \mathbb{P}_E^1,$$

in particular,

$$\#M_{Hig\lambda}^{gr \frac{1}{2}}(\mathbb{F}_{q^{2n}}) = \#\mathbb{P}^1(\mathbb{F}_{q^{2n}}) = q^{2n} + 1.$$

On the other hand by applying a formula due to Hongjie Yu [Y22], which solved a conjecture by Deligne on counting the number of  $\ell$ -adic local systems on a punctured smooth projective curve  $(C, D)/\mathbb{F}_q$  in terms of the number of semistable parabolic graded Higgs bundles on  $(C, D)/\mathbb{F}_q$ , we find

$$\#M_{\ell\text{-adic } \lambda}^{\frac{1}{2}}(\mathbb{F}_{q^{2n}}) = q^{2n} + 1.$$

This equality implies that the above injective functor is, in fact, bijective

$$(1.1) \quad M_{Hig\lambda}^{gr \frac{1}{2}}(\mathbb{F}_{q^{2n}}) \simeq M_{dR\lambda}^{FF \frac{1}{2}}(\mathbb{Z}_{q^{2n}})/\chi \simeq M_{\ell\text{-adic } \lambda}^{\frac{1}{2}}(\mathbb{F}_{q^{2n}})/\sim.$$

### 1.3. Constructing abelian scheme over $\mathbb{F}_q$ via Langlands correspondence and lifting Hodge filtration of relative differential 1-forms to characteristic zero.

Given a local system  $\mathbb{L} \in M_{\ell\text{-adic } \lambda}^{\frac{1}{2}}(\mathbb{F}_{q^{2n}})$  with cyclotomic determinant. Then one shows that the restriction of  $\mathbb{L}$  to the geometric fundamental group is irreducible with infinite local monodromy at least on one puncture. By applying Drinfeld's theorem [Dri81] to  $\mathbb{L}$  there exists an abelian scheme

$$f : A \rightarrow \mathbb{P}_{\mathbb{F}_{q^{2n}}}^1$$

with the bad reduction  $\Delta$  over  $\{0, 1, \infty, \lambda\}$  and such that:

- (1) The abelian scheme is of  $\mathrm{GL}_2(K)$ -type, where  $K$  is the number field generated by traces of Frobenius on  $(\mathbb{L})_x$  at all close points  $x \in \mathbb{P}^1 - D$ .
- (2) Let

$$\mathbb{V} := R_{\mathrm{et}, f_*}^1 \overline{\mathbb{Q}}_\ell A^0 = \bigoplus_{i=1}^g \mathbb{L}_i$$

be the  $K$ -eigen decomposition of the  $\ell$ -adic local system attached to the family. Then the local system  $\mathbb{L}$  is isomorphic to a  $K$ -eigen sheaf, say  $\mathbb{L}_1$ , and all eigen sheaves have the cyclotomic character. Moreover, the local monodromies matrices of eigen sheaves at any puncture  $x \in \{0, 1, \infty, \lambda\}$  are of the same type.

- (3) Consider the realization of the  $\mathcal{O}_K$ -log Dieudonné crystal attached to  $f$  over  $(\mathbb{P}^1, \{0, 1, \infty, \lambda\})_{\mathbb{Z}_{q^{2n}}}$ , one gets

$$(V, \nabla, \Phi, \mathcal{V}) = \left( R_{\mathrm{cry}, f_*}^1 \mathcal{O}_{A, \mathrm{cryst}} \right) (\mathbb{P}_{\mathbb{Z}_{q^{2n}}}^1).$$

The  $\mathrm{GL}_2$ -structure induces a  $K$ -eigen sheaves decomposition of  $F$ -isocrystals

$$(V, \nabla, \Phi) \otimes \mathbb{Q}_p = \bigoplus_{i=1}^g (V, \nabla, \Phi)_{i, \mathbb{Q}_p}$$

where  $(V, \nabla, \Phi)_{i, \mathbb{Q}_p}$  is a  $\sigma^f$ -log  $F$ -isocrystal with determinant  $\mathcal{E}_{cy}$ . Under  $p$ - $\ell$  companion, the eigen sheaves correspond to local systems with cyclotomic determinant. The parabolic structure induced by the residue of the connection is of type-(1/2).

According the bijection (1.1), each  $(V, \nabla, \Phi)_{i, \mathbb{Q}_p}$  has an integral lattice, which underlies a Fontaine-Faltings module  $(V, \nabla, Fil, \Phi)_i^{FF}$  over  $\mathbb{Z}_{q^{2n}}$ . Consequently the multiplication field  $K$  is unramified at each place above  $p$ . This shows that the above decomposition can be extended over  $\mathbb{Z}_{q^{2n}}$ . In other words, there exists  $(V, \nabla, \Phi)_i$  such that

$$(V, \nabla, \Phi) = \bigoplus_{i=1}^g (V, \nabla, \Phi)_i$$

and

$$(V, \nabla, \Phi)_i \otimes \mathbb{Q}_p \simeq (V, \nabla, \Phi)_i^{FF} \otimes \mathbb{Q}_p.$$

Consider the new  $\mathcal{O}_K$ -lattice

$$(V, \nabla, \Phi)^{FF} := \bigoplus_{i=1}^g (V, \nabla, \Phi)_i^{FF}$$

of  $(V, \nabla, \Phi) \otimes \mathbb{Q}_p$  defined from the Fontaine-Faltings module

$$(V, \nabla, Fil, \Phi)^{FF} := \bigoplus_{i=1}^g (V, \nabla, Fil, \Phi)_i^{FF}.$$

By extending the coefficient, one gets a Verschiebung on  $(V, \nabla, \Phi) \otimes \mathbb{Q}_p$ . By restricting onto the new lattice, one gets a Verschiebung structure  $\mathcal{V}$  on  $(V, \nabla, Fil, \Phi)^{FF}$ .

According equivalence of the Dieudonné functor, from the new modified  $\mathcal{O}_K$ -log Dieudonné crystal  $(V, \nabla, \Phi, \mathcal{V})^{FF}$ , one gets a  $p$ -isogeny  $f' : A' \rightarrow \mathbb{P}^1$  of the original abelian scheme such that  $(V, \nabla, \Phi, \mathcal{V})^{FF}$  is the Dieudonné crystal attached to  $f'$ .



From the family  $f'$ , by taking relative differential 1-forms one gets the natural Hodge filtration on  $(V, \nabla)^{FF} \otimes \mathbb{F}_{q^{2n}}$  given by

$$E^{1,0} := R^0 f'_* \Omega_{A'/\mathbb{P}^1}^1(\log \Delta) \subset (V, \nabla)^{FF} \otimes \mathbb{F}_{q^{2n}} = R_{dR}^1 f'_*(\Omega_{A'/\mathbb{P}^1}^\bullet(\log \Delta), d),$$

which is a rank- $g$  sub bundle. Since the relative Frobenius  $\Phi$  in the Fontaine-Faltings module satisfies the strong  $p$ -divisible condition with respect to the filtration  $Fil$ , the Hodge filtration is coincide with the modulo  $p$  reduction of the filtration in the Fontaine-Faltings module. In other words,  $Fil$  is a filtration lifts the Hodge filtration relative differential 1-forms attached to  $f'$ .

**1.4. Lifting abelian scheme from characteristic  $p$  to characteristic zero by Grothendieck-Messing-Kato logarithmic deformation theorem.** By Zarhin's trick, the fiber product

$$f'^{(4,4)} : A'^{4,4} = (A' \times A'^t)^4 \rightarrow \mathbb{P}^1$$

with the induced  $\mathcal{O}_K^{(4,4)}$ -multiplication carries an  $\mathcal{O}_K^{(4,4)}$ -principle polarization

$$\iota : A'^{4,4} \simeq A'^{4,4t}.$$

This  $\mathcal{O}_K^{4,4}$ -polarization induces an isomorphism between the  $\mathcal{O}_K^{4,4}$ -eigen sheaves of the Dieudonné module and its dual

$$\iota^* : (V, \nabla, \Phi, \mathcal{V})^{FF} = \bigoplus_{i=1}^{8g} (V, \nabla, \Phi, \mathcal{V})_i^{FF} \simeq \bigoplus_{i=1}^{8g} (V, \nabla, \Phi, \mathcal{V})_i^{FF\vee} = (V, \nabla, \Phi, \mathcal{V})^{FF\vee},$$

which carries the Hodge filtration  $Fil$  and  $Fil^\vee$  as liftings of the Hodge bundles  $E_{f'^{4,4}}^{1,0}$  and  $E_{f'^{4,4t}}^{1,0}$ . One checks that  $\text{Gr}_{E_{f'^{4,4}}}^{1,0} (V, \nabla)^{FF} \otimes \mathbb{F}_{q^{2n}}$  and  $\text{Gr}_{E_{f'^{4,4t}}}^{1,0} (V, \nabla)^{FF\vee} \otimes \mathbb{F}_{q^{2n}}$  are parabolic stable with respect to the  $\mathcal{O}_K^{4,4}$ -eigen sheaves decomposition. Hence, we obtain

$$(1.2) \quad \iota^* Fil = Fil^\vee$$

By Faltings-Chai Theorem [FC90] on the fine arithmetic moduli space  $\mathcal{A}_{8g,1,3} =: \mathcal{X}^0$  of principle polarized abelian varieties with level-3 exists over  $\mathbb{Z}[e^{\frac{2i\pi}{3}}, 1/3]$ , which is smooth and carries an universal abelian scheme

$$\mathcal{A}^0 \rightarrow \mathcal{X}^0.$$

Further more, there exists a smooth Toroidal compactification  $\mathcal{X} \supset \mathcal{X}^0$  over  $\mathbb{Z}[e^{\frac{2i\pi}{3}}, 1/3]$  with a smooth compactification of the universal abelian scheme

$$f : \mathcal{A} \rightarrow \mathcal{X}$$

such that  $\mathcal{A} \setminus \mathcal{A}^0$  is a relative normal crossing divisor over  $\mathcal{X} \setminus \mathcal{X}^0 =: \infty$ . For an  $\mathcal{O}_E^{(4,4)}$ -principle polarized abelian scheme  $f'^{(4,4)}$  together with a lifting of the base  $(\mathbb{P}^1, D)_{\mathbb{Z}_{q^{2n}}}$

In order to obtain a period map into the fine moduli space  $\mathcal{X}^0$  for  $f'^{(4,4)}$  we take the base change

$$\chi : (\mathbb{P}^1, D)_\chi \rightarrow (\mathbb{P}^1, D)$$

defined by torsion-6 subgroup of  $f^{s(4,4)}$ . The fiber product of the base change

$$f'_\chi{}^{4,4} : A'_\chi{}^{4,4} \rightarrow \mathbb{P}_\chi^1$$

has semistable reduction on  $D_\chi$  and carries a level-6 structure. Hence,  $f_\chi^{s(\lambda^4, 4)^0}$  induces a log period map

$$\psi : (\mathbb{P}^1, D)_\chi \rightarrow (\mathcal{X}, \infty) \otimes k.$$

By Kato's log deformation theorem [Kat96], the local liftings of  $\psi$  on  $W_n(k)$  defines an obstruction-cocycle in  $H^1(\mathbb{P}_\chi^1 \otimes k, \psi^* \Theta_{\mathcal{X}/k}(\log \infty))$ . The local liftings can be glue if and only if the obstruction class vanishes.

By Faltings-Chai [FC90], the universal Kodaira-Spencer map induces an identification

$$\Theta_{\mathcal{X}}(\log \infty) \simeq S^2 E_{\mathcal{X}}^{0,1}$$

and we may identify the obstruction for lifting of  $\psi$  with the obstruction for lifting the Hodge bundle  $E_{f'^{4,4}}^{1,0}$  satisfying (1.2). As  $E_{f'^{4,4}}^{1,0}$  lifts we show  $\psi$  lifts. Which corresponds to a lifting of the abelian scheme

$$(f'^{4,4} : A_{\mathcal{X}}'^{4,4} \rightarrow \mathbb{P}_{\mathcal{X}}^1)_{\mathbb{Z}_{q^{2n}}}$$

with a lifting of  $\mathcal{O}_K^{4,4}$ -action, as the the Dieudonné crystal with  $Fil$  carries an  $\mathcal{O}_K^{4,4}$ -action.

As  $f'^{4,4}$  over  $\mathbb{P}_{\mathcal{X}}^1/\mathbb{F}_{q^{2n}}$  descends to  $\mathbb{P}^1/\mathbb{F}_{q^{2n}}$  and the Fontaine-Faltings module attached to  $f'^{4,4}$  over  $\mathbb{P}_{\mathcal{X}}^1/W(\mathbb{F}_{q^{2n}})$  descends to  $\mathbb{P}^1/W(\mathbb{F}_{q^{2n}})$  we obtain  $f'^{4,4}$  descends to  $\mathbb{P}^1/W(\mathbb{F}_{q^{2n}})$ .

## 2. Sketch the proof of Theorem 0.8.

Let  $g : C \rightarrow \mathbb{P}^1$  be the Legendre family. Then one may identify the smooth locus of  $g$  with  $M_{0,4}$ , the moduli space of projective line with 4-punctures, which sends  $\lambda$  to the projective line with punctures at  $\{0, 1, \lambda, \infty\}$ . For any  $\lambda \neq 0, 1, \infty$ , the fiber of  $g$  at  $\lambda$  is just the elliptic curve given by the double cover  $\pi_{\lambda} : C_{\lambda} \rightarrow \mathbb{P}^1$  ramified on  $\{0, 1, \lambda, \infty\}$

For a  $\lambda_0 \in M_{0,4}$ , take a Higgs bundle  $(E, \theta)$  in  $M_{Hig}^{gr \frac{1}{2}}_{\lambda_0}$ . Then Theorem 0.8 claims that  $(E, \theta)$  is a motivic Higgs bundle if and only if  $(\theta)_0 \in \mathbb{P}^1$  is a torsion point with respect to  $\lambda_0$  (i.e. the preimage in  $\pi_{\lambda_0}^{-1}(\theta)_0 \in C_{\lambda_0}$  are torsion points). In the following, we give a sketch of the proof of Theorem 0.8.

Assume  $(E, \theta)$  is motivic. Then the modulo  $\mathfrak{p}$  reduction of  $(E, \theta)$  is periodic for almost all places  $\mathfrak{p}$ . According Theorem 0.7, the modulo  $\mathfrak{p}$  reduction of  $(\theta)_0$  is torsion. By a theorem of Pink [Pin04], it itself is torsion.

Conversely, assume  $(\theta)_0$  is a torsion point with order  $m$ , in the following we show  $(E, \theta)$  is motivic. We first show a very special case:

**Case 1.** Assume  $\lambda_0$  takes a value in  $\mathcal{O}_K$ , the ring of integers of some number field, such that  $C_{\lambda_0}$  is an elliptic curve with complex multiplication. Choose a sufficient large place  $\mathfrak{p}$  such that the reduction of  $C_{\lambda_0}$  at  $\mathfrak{p}$  is supersingular and  $\mathfrak{p} \nmid m$ .

Since the modulo  $\mathfrak{p}$  reduction  $(\bar{E}, \bar{\theta})$  of the Higgs bundle  $(E, \theta)$  is also torsion and of order  $m$  with  $\mathfrak{p} \nmid m$ . By Theorem 0.7, the reduction  $(\bar{E}, \bar{\theta})$  is periodic. According the bijection in (1.1), the modulo  $p$  reduction  $(\bar{E}, \bar{\theta})$  lifts to a periodic Higgs bundle  $(E, \theta)^{per}$  on  $(\mathbb{P}^1, \{0, 1, \lambda_0, \infty\})/W(\mathbb{F}_{q_0^{2n}})$ . In Section 1.4 we show that there exists an abelian scheme  $f_{\lambda_0} : A \rightarrow \mathbb{P}^1$  of  $GL_2(K)$ -type over characteristic 0 with bad reduction on  $\{0, 1, \lambda_0, \infty\}$  of type-(1/2) such that  $(E, \theta)^{per}$  is an  $K$ -eigen Higgs bundle attached to  $f_{\lambda_0}$ . Hence,  $(E, \theta)^{per}$  is motivic. In particular  $(\theta^{per})_0$  is also torsion and has the some modulo  $\mathfrak{p}$  reduction as  $\theta_0$ .

We claim that  $(E, \theta) \simeq (E, \theta)^{per}$ . In particular,  $(E, \theta)$  is motivic. Since  $C_{\lambda_0}$  has complex multiplication, the field generated by  $p$ -torsion point must be ramified above  $p$ . In particular, the order of  $(\theta^{per})_0$  does not divided by  $p$ . Together with the fact that  $(\theta)_0 \equiv (\theta^{per})_0 \pmod{\mathfrak{p}}$ , one gets  $(E, \theta) \simeq (E, \theta)^{per}$ .

**Case 2.** Let  $\Sigma_m \subset C$  be the  $m$ -torsion (multiple) section,  $T_m = \pi(\Sigma_m) \subset M_{0,4}$ . Then  $T_m$  is étale over  $M_{0,4}$ . Let  $T'_m$  be the irreducible component of  $T_m$  containing  $(\theta)_0$ .

Recall the fact that the set of elliptic curves with complex multiplication are dense in the moduli space. We can find a subset of infinitely many  $\{\lambda_i\}$  with the some modulo  $\mathfrak{p}$ -reduction such that all  $C_{\lambda_i}$  are elliptic curve with complex multiplication. Choose a point  $z_i$  in the intersection of  $T'_m$  and the fiber above  $\lambda_i$  for each  $i$ .

With the method in case 1, we find abelian schemes  $f_{\lambda_i} : A_{\lambda_i} \rightarrow \mathbb{P}^1$  of  $GL_2(K)$ -type with bad reduction on  $\{0, 1, \lambda_i, \infty\}$ , such that  $(E, \theta)_{z_i}$  is an  $K$ -eigen Higgs bundle attached to  $f_{\lambda_i}$ . One shows that infinitely many of  $\{f_{\lambda_i}\}$  glue together into an abelian scheme  $f : A \rightarrow M_{0,4}$ . Consequently, the subset  $\{z_i\}$ , where  $z_i = (\theta_{z_i})_0$ , glue into a component  $Z_m$  of  $\Sigma_m$  such that for any  $z_\lambda \in Z_m \cap (\mathbb{P}^1, \{0, 1, \lambda, \infty\})$  the Higgs bundle  $(E, \theta)_{z_\lambda}$  with  $(\theta_{z_\lambda})_0 = z_\lambda$  is a motivic Higgs bundle on  $(\mathbb{P}^1, \{0, 1, \lambda, \infty\})$ .

By the construction of  $Z_m$  and  $T'_m$ , Both of them are irreducible, relative dimensional one over the base and their intersect set is infinite. Hence  $Z_m = T'_m$ . In particular  $(E, \theta)$  is motivic.

**Acknowledgement.** *The authors warmly thank Raju Krishnamoorthy, Mao Sheng, Carlos Simpson, Rui-Ran Sun, Hong-Jie Yu and Shing-Tung Yau for helpful discussions.*

## REFERENCES

- [Abe13] Tomoyuki Abe. Langlands correspondence for isocrystals and the existence of crystalline companions for curves J. Amer. Math. Soc. 31 (2018), 921-1057.
- [CS08] Kevin Corlette and Carlos Simpson On the classification of rank-two representations of quasiprojective fundamental groups Compositio Mathematica , Volume 144 , Issue 5 , September 2008 , pp. 1271 - 1331.
- [Dri81] V. G. Drinfeld. The number of two-dimensional irreducible representations of the fundamental group of a curve over a finite field. *Funktsional. Anal. i Prilozhen.*, 15(4):75–76, 1981.
- [EG20] Hélène Esnault and Michael Groechenig. Rigid connections and  $f$ -isocrystals Acta Math. 225 (2020), 103-158.
- [EG18] Hélène Esnault and Michael Groechenig. Cohomologically rigid local systems and integrality November 2018 Selecta Mathematica 24(8)
- [FC90] Gerd Faltings and Ching-Li Chai Degeneration of Abelian Varieties. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge / A Series of Modern Surveys in Mathematics (MATHE3, volume 22)
- [Kat96] Fumiharu Kato. Log smooth deformation theory Tohoku Math. J., Volume 48, 1996, pp. 317-354, doi=10.2748/tmj/1178225336.
- [Ked07] Kiran S. Kedlaya. Semistable reduction for overconvergent  $F$ -isocrystals. I: Unipotence and logarithmic extensions. *Compos. Math.*, 143(5):1164–1212, 2007.
- [LSZ19] Guitang Lan, Mao Sheng, and Kang Zuo. Semistable Higgs bundles, periodic Higgs bundles and representations of algebraic fundamental groups. *J. Eur. Math. Soc. (JEMS)*, 21(10):3053–3112, 2019.
- [LS18] Adrian Langer and Carlos Simpson Rank 3 rigid representations of projective fundamental groups. Compositio Mathematica , Volume 154 , Issue 7 , July 2018 , pp. 1534 - 1570.
- [LSW22] IAojin Lin, Mao Sheng, and Jianping Wang A torsion property of zero of Kodaira-Spencer over  $\mathbb{P}^1$  removing four points. arXiv:2212.02038

- [Pin04] Richard Pink. On the order of the reduction of a point on an abelian variety. *Math. Ann.*, Volume 330, Issue 2, 2004, pp 275-291, doi=10.1007/s00208-004-0548-8.
- [SYZ17] Ruiran Sun, Jinbang Yang, and Kang Zuo. Projective crystalline representations of étale fundamental groups and twisted periodic higgs-de Rham flow, 2017. *J. Eur. Math. Soc.* 24 (2022), no. 6, pp. 1991–2076
- [Y22] Hongjie Yu Rank-2  $\ell$ -adic local systems and Higg bundles over curves. Preprint 2022.

*Email address:* yjb@mail.ustc.edu.cn

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA,  
HEFEI, ANHUI 230026, PR CHINA

*Email address:* zuok@uni-mainz.de

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT MAINZ, MAINZ 55099, GERMANY