

FLAT FLOW SOLUTION TO THE MEAN CURVATURE FLOW WITH VOLUME CONSTRAINT

VESA JULIN

ABSTRACT. In this paper I will revisit the construction of a global weak solution to the volume preserving mean curvature flow via discrete minimizing movement scheme by Mugnai-Seis-Spadaro [19]. This method is based on the gradient flow approach due to Almgren-Taylor-Wang [3] and Luckhaus-Strürzenhecker [14] and my aim is to replace the volume penalization by implementing the volume constraint directly in the discrete scheme, which from practical point of view is perhaps more natural. A technical novelty is the proof of the density estimate which is based on the second variation condition of the energy.

1. INTRODUCTION

A smooth family of set $(E_t)_{t \geq 0}$ is said to evolve according to volume preserving mean curvature flow if the normal velocity V_t is proportional to the mean curvature H_{E_t} as

$$V_t = -(H_{E_t} - \bar{H}_{E_t}) \quad \text{on } \partial E_t, \quad (1.1)$$

where $\bar{H}_{E_t} = \int_{\partial E_t} H_{E_t} d\mathcal{H}^n$. Such a geometric equation has been proposed in the physical literature to model coarsening phenomena, where the system consisting on several subdomains evolve such that it decreases the interfacial area while keeping the total volume unchanged [7, 18]. From purely mathematical point of view the equation (1.1) can be seen as the L^2 -gradient flow of the surface area under the volume constraint [18]. One has to be careful in this interpretation as the Riemannian distance between two sets is in general degenerate [16]. In order to overcome this one may use the idea due to Almgren-Taylor-Wang [3] and Luckhaus-Strürzenhecker [14] and to view (1.1) as the gradient flow of the surface area with respect to a different, non-degenerate, distance. Using the gradient flow structure, one may then construct a discrete-in-time approximation to the solution of (1.1) via the Euler implicit method, also known as the minimizing movements scheme. By letting the time step to zero, one then obtains a candidate for a weak solution of (1.1) called *flat flow*, as the convergence is measured in terms of "flat norm". This method is implemented to the volume preserving setting in [19].

In [19] the authors observe that from technical point of view it is easier to replace the volume constraint of the problem with a volume penalization, as this simplifies certain regularity issues at the level of the discrete approximation. My aim here is to show that one may construct the flat flow

solution to (1.1) by implementing the volume constraint in the minimizing movements scheme directly and thus avoid the volume penalization.

Let me quickly recall the discrete minimizing movements scheme for (1.1). One defines a sequence of sets $(E_k^h)_k$, with fixed time step $h > 0$, iteratively such that $E_0^h = E_0$, where E_0 is the given initial set, and E_{k+1}^h is a minimizer of the functional

$$P(E) + \frac{1}{h} \int_E \bar{d}_{E_k^h} dx \quad \text{under the constraint } |E| = |E_k^h|.$$

Here $P(E)$ denotes the perimeter (generalized surface area) of the set E and \bar{d}_F is the signed distance function of the set F (see next section). One then defined an approximative flat flow solution to (1.1) $(E_t^h)_{t \geq 0}$ from the previous sequence by $E_t^h = E_k^h$ for $t \in [kh, (k+1)h)$. Any cluster point of $(E_t^h)_{t \geq 0}$ is then defined as flat flow solution to (1.1). The advantage is that such a solution is defined for all times and for rough initial data. The main result in the paper is the existence of a flat flow solution.

Theorem 1. *Assume that $E_0 \subset \mathbb{R}^{n+1}$ is an open and bounded set with finite perimeter and let $(E_t^h)_{t \geq 0}$ be an approximative flat flow solution to (1.1) starting from E_0 (see Definition 2.1). Then there exists a family of bounded sets of finite perimeter $(E_t)_{t \geq 0}$ and a subsequence $h_k \rightarrow 0$ such that*

$$\lim_{h_k \rightarrow 0} |E_t^{h_k} \Delta E_t| = 0 \quad \text{for a.e. } t \geq 0$$

and for every $0 < t < s$ it holds $|E_t| = |E_0|$, $P(E_t) \leq P(E_0)$ and

$$|E_t \Delta E_s| \leq C \sqrt{s-t},$$

where C depends on the dimension and on E_0 . Moreover, if the initial set E_0 is $C^{1,1}$ -regular, then any such limit flow $(E_t)_{t \geq 0}$ agrees with the unique classical solution of (1.1) as long as the latter exists.

The above theorem thus provides the existence of a flat flow solution and guarantees that this notion is consistent with the classical solution when the initial set is regular enough. The disadvantage of the flat flow is that it is not clear if it provides a solution to the original equation (1.1) in any weak sense. However, the conditional result in the spirit of Luckhaus-Strürzenhecker [14] holds also in this case.

Theorem 2. *Let $(E_t^h)_{t \geq 0}$ be an approximative flat flow solution to (1.1) and let $(E_t^{h_k})_{t \geq 0}$ be the converging subsequence in Theorem 1. Assume further that it holds*

$$\lim_{h_k \rightarrow 0} P(E_t^{h_k}) = P(E_t) \quad \text{for a.e. } t \geq 0.$$

Then for $n \leq 6$ the flat flow $(E_t)_{t \geq 0}$ is a distributional solution to (1.1) (see Definition 4.5).

One may also try to view the equation (1.1) as a mean curvature flow with forcing, where the forcing term depends on the flow itself. In this way one may try use different methods to construct a solution to the equation see e.g. [5, 6]. I also refer the recent work [13] for a weak-strong uniqueness result related to (1.1).

As I already mentioned, the flat flow is defined for all times and one may study its asymptotical behavior. Indeed, by using the methods from [10, 11] one may deduce the convergence of the flow in low dimensions.

Remark 1.1. Assume that $E_0 \subset \mathbb{R}^{n+1}$, with $n \leq 2$, is as in Theorem 1 and let $(E_t)_{t \geq 0}$ be a limit flat flow. When $n = 1$, the flow E_t converges to a union of disjoint balls exponentially fast and when $n = 2$ the flow converges to a union of disjoint balls up to a possible translation of the components.

The main technical challenge in proving Theorem 1 is to obtain the sharp density estimate for the discrete flow. This is also the main technical novelty of this paper. There are several techniques to deal with the volume constraint in variational problems, e.g. by using the argument from [2] (see also [15, Lemma 17.21]) or from [8] (see also [4, 9]). However, due to the presence of the dissipation term in the energy it is not obvious how to apply these arguments in order to obtain sharp density estimates in terms of the time step h . I will use an argument which is based on the second variation condition of the energy to prove the density estimate in Proposition 3.1. After this the proof of Theorem 1 follows exactly as in [14, 19] and the consistency follows almost directly using the argument in [12]. The proof also provides the dissipation inequality and therefore the results in [10, 11] hold and one obtains the result stated in Remark 1.1. Finally I would like to point out that this article is not self-consistent as several arguments are well-known, in particular, in Section 4.

2. PRELIMINARIES

In this section I will briefly introduce the notation, the definition of the flat flow solution and recall some of its basic properties.

Given a set $E \subset \mathbb{R}^{n+1}$ the distance function $\text{dist}(\cdot, E) : \mathbb{R}^{n+1} \rightarrow [0, \infty)$ is defined, as usual, as $\text{dist}(x, E) := \inf_{y \in E} |x - y|$ and denote the signed distance function by $\bar{d}_E : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$,

$$\bar{d}_E(x) := \begin{cases} -\text{dist}(x, \partial E), & \text{for } x \in E \\ \text{dist}(x, \partial E), & \text{for } x \in \mathbb{R}^{n+1} \setminus E. \end{cases}$$

Then clearly it holds $\text{dist}(\cdot, \partial E) = |\bar{d}_E|$. I denote the ball with radius r centered at x by $B_r(x)$ and by B_r if it is centered at the origin.

For a measurable set $E \subset \mathbb{R}^{n+1}$ the perimeter in an open set $U \subset \mathbb{R}^{n+1}$ is defined by

$$P(E, U) := \sup \left\{ \int_E \text{div } X \, dx : X \in C_0^1(U, \mathbb{R}^{n+1}), \|X\|_{L^\infty} \leq 1 \right\}$$

and write $P(E) = P(E, \mathbb{R}^{n+1})$. If $P(E) < \infty$, then E is called a set of finite perimeter. For an introduction to the topic I refer to [15]. The reduced boundary of a set of finite perimeter E is denoted by $\partial^* E$ and the generalized unit outer normal by ν_E . Note that it holds $P(E, U) = \mathcal{H}^n(\partial^* E \cap U)$ for open sets U . Recall also that if E is regular enough, say with Lipschitz boundary, then $P(E) = \mathcal{H}^n(\partial E)$. For a given vector field $X \in C^1(\mathbb{R}^{n+1}, \mathbb{R}^{n+1})$ and a set of finite perimeter E denote the tangential divergence on ∂E^* as $\text{div}_\tau X = \text{div } X - \langle DX \nu_E, \nu_E \rangle$. The distributional mean curvature $H_E \in L^1(\partial^* E, \mathbb{R})$

is defined via the divergence theorem such that for every test vector field $X \in C_0^1(\mathbb{R}^{n+1}, \mathbb{R}^{n+1})$ it holds

$$\int_{\partial^* E} \operatorname{div}_\tau X \, d\mathcal{H}^n = \int_{\partial^* E} H_E \langle X, \nu_E \rangle \, d\mathcal{H}^n.$$

I will consider a flat flow solution to (1.1) in the spirit of Almgren-Taylor-Wang [3] and Luckhaus-Stürzenhecker [14]. To this aim for a fixed $h \in (0, 1)$ and a given (open) set $F \subset \mathbb{R}^{n+1}$ I define the functional

$$\mathcal{F}_h(E, F) = P(E) + \frac{1}{h} \int_E \bar{d}_F \, dx. \quad (2.1)$$

The flat flow solution is defined analogously as in [19].

Definition 2.1. Let $E_0 \subset \mathbb{R}^{n+1}$ be an open and bounded set of finite perimeter and fix $h \in (0, 1)$. Define the sequence of sets $(E_k^h)_{k=0}^\infty$ iteratively as $E_0^h = E_0$ and E_{k+1}^h is a minimizer of the problem

$$\min \left\{ \mathcal{F}_h(E, E_k^h) : |E| = |E_0| \right\}.$$

Moreover, define an approximative flat flow $(E_t^h)_{t \geq 0}$ for (1.1) starting from E_0 as

$$E_t^h = E_k^h \quad \text{for } t \in [kh, (k+1)h).$$

One has to be careful in the definition of the functional (2.1) if the set F is merely a set of finite perimeter as its value depends on the choice of the representative of F . One may overcome this by choosing a proper representative of the set F . In my case this is not necessary, as the regularity theorem below implies that one may in fact assume the sets E_k^h to be open. The difference in the Definition 2.1 to the scheme in [19] is that here the minimizing problem is under volume constraint. On one hand this makes the minimization problem more natural, but on the other hand, it makes the quantitative density estimates more difficult to prove.

For a given open and bounded set $F \subset \mathbb{R}^{n+1}$ consider the minimization problem

$$\min \left\{ \mathcal{F}_h(E, F) : |E| = |F| \right\}, \quad (2.2)$$

where $\mathcal{F}_h(\cdot, F)$ is defined in (2.1). One may use an argument similar to [9] or [15, Lemma 17.21] to get rid of the volume constraint in (2.2) and deduce that a minimizer of (2.2) is a minimizer also for

$$\min \left\{ \mathcal{F}_h(E, F) + \tilde{\Lambda} \left| |E| - |F| \right| \right\}, \quad (2.3)$$

when $\tilde{\Lambda}$ is chosen large. Note that the constant $\tilde{\Lambda}$ may have unoptimal dependence on E and on h . However, the property (2.3) is enough to deduce qualitative regularity properties since it implies that the minimizer inherits the regularity from the theory of the perimeter minimizers [15]. One may also write the Euler-Lagrange equation and by standard calculations (see e.g. [1]) we have the second variation condition. We state this in the following proposition.

Proposition 2.2. *Let $F \subset \mathbb{R}^{n+1}$ be an open and bounded set, fix $h \in (0, 1)$ and let E be a minimizer of (2.2). Then E can be chosen to be open, which topological boundary is $C^{2,\alpha}$ -regular up to a relatively closed singular set*

which Hausdorff dimension is at most $n - 7$. In fact, the regular part is exactly the reduced boundary $\partial^* E$.

The Euler-Lagrange equation

$$\frac{d_F}{h} = -H_E + \lambda, \quad (2.4)$$

where $\lambda \in \mathbb{R}$ is the Lagrange-multiplier, holds point wise on $\partial^* E$ and in a distributional sense on ∂E . The quadratic form associated with the second variation of the energy is non-negative, i.e., for all $\varphi \in H^1(\partial^* E)$ with $\int_{\partial^* E} \varphi \, d\mathcal{H}^n = 0$ it holds

$$\int_{\partial^* E} |\nabla_{\tau} \varphi|^2 - |B_E|^2 \varphi^2 \, d\mathcal{H}^n + \frac{1}{h} \int_{\partial^* E} \langle \nabla \bar{d}_F, \nu_E \rangle \varphi^2 \, d\mathcal{H}^n \geq 0, \quad (2.5)$$

where $B_E(x)$ denotes the second fundamental form at $x \in \partial^* E$.

Proof. Since the argument is standard, I will only give the outline. As I already mentioned, the minimizer E is also a minimizer of the problem (2.3) for some large constant $\tilde{\Lambda}$, which depend on h and on E itself. This implies that the set E is a Λ -minimizer of the perimeter and thus the reduced boundary $\partial^* E$ is relatively open, $C^{1,\alpha}$ -regular hypersurface and the singular set $\partial E \setminus \partial^* E$ has dimension at most $n - 7$ [15]. The $C^{2,\alpha}$ -regularity then follows from the Euler-Lagrange equation and from standard Schauder-estimates for elliptic PDEs.

One may obtain the second variation condition (2.5) by using the argument from [1]. Indeed, given a function $\varphi \in C_0^1(\partial^* E)$ with $\int_{\partial^* E} \varphi \, d\mathcal{H}^n = 0$, we may construct a family of diffeomorphisms Φ_t such that $\Phi_0 = id$, $|\Phi_t(E)| = |E|$ and $\frac{\partial}{\partial t} \Big|_{t=0} \Phi_t(x) \cdot \nu_E = \varphi$. Then the inequality follows from the minimality of E as

$$\frac{\partial^2}{\partial t^2} \Big|_{t=0} \mathcal{F}_h(\Phi_t(E), F) \geq 0$$

and following the standard calculation of the second variation (see e.g. [1]). Finally one obtains (2.5) for all $\varphi \in H^1(\partial^* E)$ by approximation argument and by the fact that the singular set has zero capacity. \square

3. DENSITY ESTIMATES

This section is the theoretical core of the paper. The aim is to prove the following density estimate.

Proposition 3.1. *Let $F \subset \mathbb{R}^{n+1}$ be an open and bounded set of finite perimeter, fix $h \in (0, 1)$ and let E be a minimizer of (2.2). Then there is a constant $c > 0$, which depends on the dimension n , $|F|$ and on $P(F)$ such that for all $r \leq \sqrt{h}$ and all $x \in \partial E$ it holds*

$$\min \{ |E \cap B_r(x)|, |B_r(x) \setminus E| \} \geq cr^{n+1}$$

and for all $r \leq C_0 \sqrt{h}$, where $C_0 \geq 1$, it holds

$$cr^n \leq P(E, B_r(x)) \leq C_1 r^n,$$

where C_1 depends also on C_0 . Moreover, the following holds

$$\|H_E\|_{L^\infty(\partial^* E)} \leq \frac{1}{c\sqrt{h}} \quad \text{and} \quad \|\bar{d}_F\|_{L^\infty(\partial E)} \leq c^{-1}\sqrt{h}.$$

It is interesting that in [19, Corollary 3.3] the authors obtain similar result for their scheme for a constant which is independent of $P(F)$.

I need several lemmas in order to prove Proposition 3.1 and therefore I postpone its proof to the end of the section. Before proceeding to technical details, I state a useful consequence of Proposition 3.1.

Proposition 3.2. *Let $F, E \subset \mathbb{R}^{n+1}$ be as in Proposition 3.1. Then there are constants $C \geq 1, c > 0$ and $h_0 > 0$, depending on the dimension, $|F|$ and $P(F)$ such that E is (Λ, r) - minimizer of the perimeter for $\Lambda = \frac{C}{\sqrt{h}}$, $r = c\sqrt{h}$ and for $h < h_0$. To be more precise, for sets $G \subset \mathbb{R}^{n+1}$ with $E \Delta G \subset B_{c\sqrt{h}}(x_0)$ it holds*

$$P(E) \leq P(G) + \frac{C}{\sqrt{h}}|E \Delta G|.$$

Proof. The argument is standard but I recall it for the reader's convenience. Let me first show that there is $x \in E$ and $\tilde{c} > 0$ such that for $\rho = \tilde{c}\sqrt{h}$ it holds $B_\rho(x) \subset E$. Fix ρ and apply Besicovich covering theorem to find disjoint balls $\{B_\rho(x_i)\}_{i=1}^N$ such that $x_i \in E$ and

$$\sum_{i=1}^N |B_\rho(x_i)| = N|B_1|\rho^{n+1} \geq c|E|. \quad (3.1)$$

I claim that for some $i = 1, 2, \dots, N$ it holds $B_{\rho/2}(x_i) \subset E$. Indeed, if this is not the case then Proposition 3.1 implies

$$P(E, B_\rho(x_i)) \geq c\rho^n$$

for all i . Since the balls are disjoint, one has by the above and by (3.1)

$$P(E) \geq \sum_{i=1}^N P(E, B_\rho(x_i)) \geq cN\rho^n \geq c\frac{|E|}{\rho} \geq \frac{c}{\sqrt{h}}.$$

This is a contradiction when h is small enough.

Fix $x_0 \in \partial E$ and G as in the claim. Note that in general the set G does not have the same measure as E and one needs to modify it to \tilde{G} with $|\tilde{G}| = |E|$ e.g. by using the argument from [9] as follows. Assume that $|G| < |E|$ (the case $|G| > |E|$ follows from similar argument). Since $B_\rho(x) \subset E$, then by decreasing ρ and r if needed, it holds $B_\rho(x) \subset G$. By continuity there is $z \in \mathbb{R}^{n+1}$ such that $|z - x_0| \geq 2\rho$ and $|G \cup B_\rho(z)| = |E|$. Define $\tilde{G} = G \cup B_\rho(z)$. Then by the minimality of E and Proposition 3.1 it holds

$$P(E) \leq P(\tilde{G}) + \frac{C}{\sqrt{h}}|\tilde{G} \Delta E|.$$

Arguing as in [9] one then deduces

$$\begin{aligned} P(\tilde{G}) - P(G) &\leq \mathcal{H}^n(\partial B_\rho(z) \setminus G) - \mathcal{H}^n(\partial G \cap B_\rho(z)) \\ &\leq \frac{C}{\rho}|B_\rho(z) \setminus G| \leq \frac{C}{\sqrt{h}}|\tilde{G} \Delta E| \end{aligned}$$

and the claim follows as $|\tilde{G} \Delta E| \leq 2|G \Delta E|$. \square

The first technical result which I need is the classical density estimate which can be found e.g. in [20].

Lemma 3.3. *Assume $E \subset \mathbb{R}^{n+1}$ is a set of finite perimeter with distributional mean curvature H_E which satisfies $\|H_E\|_{L^\infty(B_{2R}(x_0))} \leq \Lambda$. Then for all $x \in B_R(x_0)$ and $r \leq \min\{R, \Lambda^{-1}\}$ it holds*

$$P(E, B_r(x)) \geq c_n r^n,$$

for a dimensional constant $c_n > 0$.

For a minimizer of (2.2) it holds the inverse of the isoperimetric inequality.

Lemma 3.4. *Let F and E be as in Proposition 3.1. Then for all $x \in \partial E$ and $r \leq C_0 \sqrt{h}$ it holds*

$$P(E, B_r(x)) \leq \frac{C}{r} \min\{|E \cap B_{2r}(x)|, |B_{2r}(x) \setminus E|\},$$

for a constant which depends on the dimension and on $C_0 > 0$. In, particular it holds $P(E, B_r(x)) \leq Cr^n$.

Proof. Fix $h \in (0, 1)$, x and $r > 0$ as in the claim and without loss of generality assume that $x = 0$. One may also consider only the case $|E \cap B_{2r}| \leq |B_{2r} \setminus E|$ as the other case is similar. In particular, then it holds $|E \cap B_{2r}| \leq \frac{1}{2}|B_{2r}|$.

Since

$$\int_0^{2r} \mathcal{H}^n(\partial B_\rho \cap E) = |E \cap B_{2r}|,$$

there is $\rho \in (r, 2r)$ such that

$$\mathcal{H}^n(\partial B_\rho \cap E) \leq C_n \frac{|E \cap B_{2r}|}{r} \quad \text{and} \quad |B_\rho| \geq \frac{2}{3}|B_{2r}|. \quad (3.2)$$

Consider first the set $E_1 = E \setminus \bar{B}_\rho$. In order to have a competing set with the volume of E , define $\tilde{\rho} < \rho$ to be a radius such that $|B_{\tilde{\rho}}| = |E \cap B_\rho|$ and define $E_2 = E_1 \cup B_{\tilde{\rho}}$. Then it holds by construction that $|E_2| = |E|$. By the minimality of E we have

$$P(E) + \frac{1}{h} \int_E \bar{d}_F dx \leq P(E_2) + \frac{1}{h} \int_{E_2} \bar{d}_F dx.$$

Estimate the perimeter of E_2 using (3.2) as

$$\begin{aligned} P(E_2) &\leq P(E, \mathbb{R}^{n+1} \setminus B_\rho) + \mathcal{H}^n(\partial B_\rho \cap E) + \mathcal{H}^n(\partial B_{\tilde{\rho}}) \\ &\leq P(E, \mathbb{R}^{n+1} \setminus B_\rho) + C_n \frac{|E \cap B_{2r}|}{r}. \end{aligned}$$

Use then $E \Delta E_2 \subset B_{2r}$, $|E| = |E_2|$ and the fact that the signed distance function is 1-Lipschitz to estimate

$$\left| \int_E \bar{d}_F dx - \int_{E_2} \bar{d}_F dx \right| \leq 4r |E \cap B_\rho| \leq 4r |E \cap B_{2r}|.$$

Therefore one obtains by combining the three above inequalities and $r \leq C_0 \sqrt{h}$

$$P(E, B_r) \leq P(E, B_\rho) \leq C_n \frac{|E \cap B_{2r}|}{r} + \frac{4r}{h} |E \cap B_{2r}| \leq C \frac{|E \cap B_{2r}|}{r}.$$

□

By Lemma 3.3 and Lemma 3.4 it is clear that for Proposition 3.1 it is crucial to prove the curvature estimate $\|H_E\|_{L^\infty} \leq \frac{C}{\sqrt{h}}$. The next lemma is a step towards this.

Lemma 3.5. *Let F, E and h be as in Proposition 3.1. Then it holds*

$$\|H_E\|_{L^2(\partial^* E)} \leq \frac{C_1}{\sqrt{h}},$$

where the constant C_1 depends on the dimension and on $|F|$ and $P(F)$.

Proof. The proof relies on the second variation inequality in Proposition 2.2. I would like to point out that in the case of the mean curvature flow, when there is no volume constraint, the proof is considerably easier as one could choose constant function in (2.5). In the volume preserving case I will choose a cut-off function for a test function.

To this aim use first [11, Proposition 2.3] (see also [17, Lemma 2.1]) to find a point $x_0 \in \mathbb{R}^{n+1}$ and a radius $r \in (c, 1)$, where $c = c(n, |F|, P(F))$, such that

$$|E \cap B_r(x_0)| = \frac{1}{2}|B_r|.$$

Note that the minimality of E yields $P(E) \leq P(F) \leq C$. Moreover, by the isoperimetric inequality it holds $P(E) \geq c_n |E|^{\frac{n}{n+1}} = c_n |F|^{\frac{n}{n+1}} \geq c$. These estimates are used repeatedly from now on without mentioning. Without loss of generality assume that $x_0 = 0$. Choose $\rho < r$ such that $|B_r \setminus B_\rho| = \frac{1}{4}|B_r|$. Note that then $r - \rho \geq c_n > 0$ and

$$\frac{3}{4}|B_\rho| \geq |E \cap B_\rho| \geq \frac{1}{4}|B_\rho|. \quad (3.3)$$

Define first a cut-off function $\zeta \in C_0^1(\mathbb{R}^{n+1})$ such that $0 \leq \zeta \leq 1$, $\zeta = 1$ in B_ρ , $\zeta = 0$ outside B_r and $|\nabla \zeta| \leq C_n$. Choose then $\varphi = \zeta - \bar{\zeta}$, where $\bar{\zeta} = \int_{\partial^* E} \zeta \, d\mathcal{H}^n$, as a test function in (2.5), use $|\langle \nabla \bar{d}_F, \nu_E \rangle| \leq 1$ and $|\varphi| \leq 1$, and obtain

$$\int_{\partial^* E} |B_E|^2 (\zeta - \bar{\zeta})^2 \, d\mathcal{H}^n \leq \int_{\partial^* E} |\nabla_\tau \zeta|^2 \, d\mathcal{H}^n + \frac{P(E)}{h}. \quad (3.4)$$

Since $H_E = \text{Trace}(B_E)$, it holds point wise on $\partial^* E$

$$|B_E|^2 \geq \frac{H_E^2}{n}. \quad (3.5)$$

Recall that $0 \leq \zeta \leq 1$. Moreover, by the isoperimetric inequality and by (3.3) it holds $P(E, B_\rho) \geq c_n |E \cap B_\rho|^{\frac{n}{n+1}} \geq c$. Therefore $\bar{\zeta} \geq c$ for $c = c(n, |F|, P(F))$. In particular, it holds $|\zeta(x) - \bar{\zeta}| \geq c$ for $x \in \partial^* E \setminus B_r$. Hence, we have by (3.4)

$$\int_{\partial^* E \setminus B_r} |H_E|^2 \, d\mathcal{H}^n \leq \frac{C}{h} P(F).$$

We repeat the same argument by defining a cut-off function $\zeta \in C^1(\mathbb{R}^{n+1})$ as $\zeta = 0$ in B_r , $\zeta = 1$ outside B_R and $|\nabla \zeta| \leq C_n$, where $R > r$ is such that $|B_R \setminus B_r| = \frac{1}{4}|B_r|$. Using $\varphi = \zeta - \bar{\zeta}$ in (2.5) and arguing as above yields

$$\int_{\partial^* E \cap B_r} |H_E|^2 \, d\mathcal{H}^n \leq \frac{C}{h} P(F)$$

and the claim follows. \square

The last lemma I need is a bound on the Lagrange multiplier in the Euler-Lagrange equation (2.4).

Lemma 3.6. *Let F, E and h be as in Proposition 3.1. Then for the Lagrange multiplier in (2.4), i.e.,*

$$\frac{\bar{d}_F}{h} = -H_E + \lambda \quad \text{on } \partial^* E$$

it holds

$$|\lambda| \leq \frac{C_2}{\sqrt{h}},$$

where the constant C_2 depends on the dimension, on $|F|$ and on $P(F)$.

Proof. Let $\Lambda \geq 0$ be such that $|\lambda| = \frac{\Lambda}{\sqrt{h}}$. Below all the constant depend on $n, |F|$ and $P(F)$. I only treat the case when λ is positive as in the negative case the proof is the similar. Define the set

$$\Sigma = \{x \in \partial^* E : |H_E(x)| < \frac{\hat{C}}{\sqrt{h}}\}.$$

I claim that we may choose $\hat{C} > 2$ such that it depends on $n, |F|, P(F)$ and on C_1 from Lemma 3.5 and it holds

$$\mathcal{H}^n(\Sigma) \geq \frac{P(E)}{2}. \quad (3.6)$$

Indeed, by the Euler-Lagrange equation (2.4), by Lemma 3.5 and by $\hat{C} > 2$ it holds

$$\frac{\hat{C}^2}{h} \mathcal{H}^n(\partial^* E \setminus \Sigma) \leq \int_{\partial^* E \setminus \Sigma} H_E^2 d\mathcal{H}^n \leq \int_{\partial^* E} H_E^2 d\mathcal{H}^n \leq \frac{C_1}{h}.$$

By choosing \hat{C} large enough one then obtains $\mathcal{H}^n(\partial^* E \setminus \Sigma) < P(E)/2$ and (3.6) follows.

By Besikovich covering theorem one finds disjoint balls of radius \sqrt{h} , denote them $\{B_{\sqrt{h}}(x_i)\}_{i=1}^N$, with $x_i \in \Sigma$ such that

$$\sum_{i=1}^N P(E, B_{\sqrt{h}}(x_i)) \geq c_n \mathcal{H}^n(\Sigma) \geq cP(E). \quad (3.7)$$

By the Euler-Lagrange equation (2.4) and by the definition of the set Σ it holds for all $x \in \partial^* E \cap B_{\sqrt{h}}(x_i)$ with $x_i \in \Sigma$ that

$$\begin{aligned} |H_E(x)| &\leq \left| \frac{\bar{d}_F(x)}{h} \lambda \right| \leq \frac{|\bar{d}_F(x) - \bar{d}_F(x_i)|}{h} + \left| \frac{\bar{d}_F(x_i)}{h} - \lambda \right| \\ &\leq \frac{1}{\sqrt{h}} + |H_E(x_i)| \leq \frac{2\hat{C}}{\sqrt{h}}. \end{aligned} \quad (3.8)$$

Therefore by Lemma 3.3 it holds

$$P(E, B_{\sqrt{h}/2}(x_i)) \geq P(E, B_{\sqrt{h}/2\hat{C}}(x_i)) \geq ch^{n/2}.$$

On the other hand applying Lemma 3.4 first with $r = \sqrt{h}/2$ yields

$$|E \cup B_{\sqrt{h}}(x_i)| \geq c\sqrt{h}P(E, B_{\sqrt{h}/2}(x_i))$$

and then with $r = \sqrt{h}$ yields $h^{\frac{n+1}{2}} \geq cP(E, B_{\sqrt{h}}(x_i))$. In conclusion it holds

$$|E \cup B_{\sqrt{h}}(x_i)| \geq c\sqrt{h}P(E, B_{\sqrt{h}}(x_i)) \quad (3.9)$$

for all balls in the cover.

The minimality of E implies

$$\frac{1}{h} \int_{E \setminus F} \bar{d}_F \, dx \leq P(E) + \frac{1}{h} \int_{E \Delta F} |\bar{d}_F| \, dx \leq P(F). \quad (3.10)$$

Note that by (3.8), by $\lambda = \frac{\Lambda}{\sqrt{h}}$ and by the Euler-Lagrange equation (2.4) we have for all $x \in B_{\sqrt{h}}(x_i)$ that

$$\bar{d}_F(x) \geq \lambda h - |H(x)|h \geq (\Lambda - 2\hat{C})\sqrt{h}.$$

Therefore either $\Lambda \leq 4\hat{C}$, in which case the claim follows trivially, or

$$\bar{d}_F(x) \geq \frac{\Lambda}{2}\sqrt{h}.$$

I assume the latter and show that even in this case Λ is bounded. Indeed, by the above discussion the balls $B_{\sqrt{h}}(x_i)$ are in the exterior of F . Therefore we estimate by (3.7) and (3.9) that

$$\begin{aligned} \frac{1}{h} \int_{E \setminus F} \bar{d}_F \, dx &\geq \frac{1}{h} \sum_{i=1}^N \int_{E \cap B_{\sqrt{h}}(x_i)} \bar{d}_F \, dx \\ &\geq \frac{1}{h} \sum_{i=1}^N \left(\frac{\Lambda}{2} \sqrt{h} |E \cap B_{\sqrt{h}}(x_i)| \right) \\ &\geq c\Lambda \sum_{i=1}^N P(E, B_{\sqrt{h}}(x_i)) \\ &\geq c\Lambda \mathcal{H}^n(\Sigma) \geq c\Lambda P(E). \end{aligned}$$

Since $P(E) \geq c_n |E|^{\frac{n}{n+1}} = c_n |F|^{\frac{n}{n+1}}$, the above and (3.10) gives a bound for Λ and the claim follows. \square

Here is the proof of the density estimate.

Proof of Proposition 3.1. By Lemma 3.3, Lemma 3.4, Lemma 3.6 and by the Euler-Lagrange equation (2.4) it is enough to prove

$$\|\bar{d}_F\|_{L^\infty(\partial^* E)} \leq C\sqrt{h}. \quad (3.11)$$

Argue by contradiction and assume that there is $x_0 \in \partial^* E$ such that

$$|\bar{d}_F(x_0)| = \Lambda\sqrt{h}$$

for large $\Lambda \gg 1$. Without loss of generality assume that $x_0 = 0$ and consider only the case $\bar{d}_F(x_0) > 0$ as the other case is similar.

By the Euler-Lagrange equation (2.4), by Lemma 3.3 and Lemma 3.6 it holds for $r_0 = \frac{\sqrt{h}}{2\Lambda}$ that

$$P(E, B_{r_0}) \geq cr_0^n. \quad (3.12)$$

Define radii $r_k = k\sqrt{h} + r_0$ for $k = 0, 1, \dots, n+1$. For every $k = 0, 1, \dots, n+1$ choose a cut-off function $\zeta_k \in C_0^1(\mathbb{R}^{n+1})$ such that $0 \leq \zeta \leq 1$, $\zeta_k = 1$ in B_{r_k} , $\zeta_{k+1} = 0$ in $\mathbb{R}^{n+1} \setminus B_{r_{k+1}}$ and $|\nabla \zeta_k| \leq \frac{2}{\sqrt{h}}$. Choose $\varphi = \zeta_k - \zeta_{k+1}$, where

$\bar{\zeta}_k = \int_{\partial^* E} \zeta_k \, d\mathcal{H}^n$, as a test function in the second variation condition (2.5), use $|\langle \nabla d_F, \nu_E \rangle| \leq 1$ and (3.5), and obtain

$$\frac{1}{n} \int_{\partial^* E} |H_E|^2 (\zeta_k - \bar{\zeta}_k)^2 \, d\mathcal{H}^n \leq \frac{1}{h} \int_{\partial^* E} (\zeta_k - \bar{\zeta}_k)^2 \, d\mathcal{H}^n + \int_{\partial^* E} |\nabla_\tau \zeta_k|^2 \, d\mathcal{H}^n. \quad (3.13)$$

Since $\zeta_k = 0$ outside $B_{r_{k+1}}$, one may estimate

$$\int_{\partial^* E} |\nabla_\tau \zeta_k|^2 \, d\mathcal{H}^n \leq \frac{4}{h} P(E, B_{r_{k+1}}).$$

Moreover, since $0 \leq \zeta_k \leq 1$ it holds

$$\int_{\partial^* E} (\zeta_k - \bar{\zeta}_k)^2 \, d\mathcal{H}^n = \int_{\partial^* E} \zeta_k^2 - \bar{\zeta}_k^2 \, d\mathcal{H}^n \leq \int_{\partial^* E} \zeta_k^2 \, d\mathcal{H}^n \leq P(E, B_{r_{k+1}}).$$

When Λ is large enough, then for all $x \in \partial^* E \cap B_{r_{k+1}}$ and all $k \leq n+1$ it holds $\bar{d}_F(x) \geq \frac{\Lambda}{2} \sqrt{h}$. Then one may deduce from (3.6) that $P(E, B_{r_{k+1}}) \leq \frac{1}{2} P(E)$ when Λ is large enough. This yields

$$0 \leq \bar{\zeta}_k \leq \frac{1}{2}.$$

Also by the Euler-Lagrange equation (2.4), by $\bar{d}_F(x) \geq \frac{\Lambda}{2} \sqrt{h}$, and by Lemma 3.6 it holds $|H_E| \geq \frac{\Lambda}{4\sqrt{h}}$ on $\partial^* E \cap B_{r_k}$, when Λ is large. Therefore it holds

$$\begin{aligned} \frac{1}{n} \int_{\partial^* E} |H_E|^2 (\zeta_k - \bar{\zeta}_k)^2 \, d\mathcal{H}^n &\geq \frac{1}{n} \int_{\partial^* E \cap B_{r_k}} |H_E|^2 (1 - \bar{\zeta}_k)^2 \, d\mathcal{H}^n \\ &\geq c_n \frac{\Lambda^2}{h} P(E, B_{r_k}). \end{aligned}$$

Combining the three above estimates with (3.13) yields

$$c_n \frac{\Lambda^2}{h} P(E, B_{r_k}) \leq \frac{5}{h} P(E, B_{r_{k+1}}).$$

For Λ large enough this implies

$$\Lambda P(E, B_{r_k}) \leq P(E, B_{r_{k+1}}). \quad (3.14)$$

Use (3.14) $(n+1)$ -times from $k=0$ to $k=n$, use then (3.12) and recall that $r_0 = \frac{\sqrt{h}}{2\Lambda}$ and obtain finally that

$$P(E, B_{r_{n+1}}) \geq \Lambda^{n+1} P(E, B_{r_0}) \geq c \Lambda^{n+1} r_0^n = c \Lambda^{n+1} \left(\frac{\sqrt{h}}{2\Lambda} \right)^n = c \Lambda h^{n/2}. \quad (3.15)$$

But now since $r_{n+1} = (n+1)\sqrt{h} + r_0 \leq 2(n+1)\sqrt{h}$, one obtains from Lemma 3.4 with $r = 2(n+1)\sqrt{h}$ that

$$P(E, B_{r_{n+1}}) \leq P(E, B_r) \leq C h^{n/2},$$

which contradicts (3.15) when Λ is large. \square

4. EXISTENCE OF THE FLAT FLOW

Now that the density estimates are proven the proof of Theorem 1 follows from the arguments from [14, 19] without major changes. In this section I consider the approximative flat flow $(E_t^h)_{t \geq 0}$ and the associated sequence $(E_k^h)_{k \geq 0}$ as in the Definition 2.1 starting from an open and bounded set of finite perimeter E_0 . The proof for the following "interpolation" result can be found in [14, Lemma 1.5].

Lemma 4.1. *Let $(E_t^h)_{t \geq 0}$ be an approximative flat flow starting from E_0 and fix $h \in (0, 1)$ and $t > h$. Then for all $l \leq \sqrt{h}$ it holds*

$$|E_t^h \Delta E_{t-h}^h| \leq C \left(l P(E_{t-h}^h) + \frac{1}{l} \int_{E_t^h \Delta E_{t-h}^h} |\bar{d}_{E_{t-h}^h}| dx \right)$$

By the regularity result stated in Proposition 2.2 the Euler-Lagrange equation

$$\frac{\bar{d}_{E_{t-h}^h}}{h} = -H_{E_t^h} + \lambda_{t,h} \quad (4.1)$$

holds point wise on $\partial^* E_t^h$ and in a distributional sense on ∂E_t^h . Here $\lambda_{t,h}$ is the Lagrange multiplier. Using the minimality of E_t^h against the previous set E_{t-h}^h one obtains the important inequality

$$P(E_t^h) + \frac{1}{h} \int_{E_t^h \Delta E_{t-h}^h} |\bar{d}_{E_{t-h}^h}| dx \leq P(E_{t-h}^h). \quad (4.2)$$

Using (4.2) and the argument [14, Lemma 2.1] (see also [19, Lemma 3.6]) one obtains the following dissipation inequality.

Lemma 4.2. *Let $(E_t^h)_{t \geq 0}$ be an approximative flat flow starting from E_0 and fix $h \in (0, 1)$. Then for all $T_2 > T_1 \geq h$ it holds*

$$\int_{T_1}^{T_2} \|H_{E_t^h} - \lambda_{t,h}\|_{L^2(\partial^* E_t^h)}^2 dt \leq C(P(E_{T_1-h}) - P(E_{T_2})).$$

Moreover, it holds

$$\int_{T_1}^{T_2} (\|H_{E_t^h}\|_{L^2(\partial^* E_t^h)}^2 + \lambda_{t,h}^2) dt \leq C(1 + T_2 - T_1).$$

The constant depends on the dimension, $|E_0|$ and $P(E_0)$.

Proof. I will only sketch the proof. Let (E_k^h) be the sequence of sets associated with $(E_t^h)_{t \geq 0}$. For $l \in \mathbb{Z}$ with $2^l \leq 2Ch^{-\frac{1}{2}}$ set

$$K(l) = \{x \in \mathbb{R}^{n+1} : 2^l h < |\bar{d}_{E_{t-h}^h}| \leq 2^{l+1} h\}.$$

Here C is such that $|\bar{d}_{E_{t-h}^h}| \leq Ch$ on ∂E_t^h . Proposition 3.1 yields that for every $x \in \partial E_t^h$

$$|E_t^h \cap B_{2^l h}(x)| \geq c(2^l h)^{n+1} \quad \text{and} \quad \mathcal{H}^n(\partial E_t^h \cap B_{2^l h}(x)) \leq C(2^l h)^n.$$

Therefore for all $x \in \partial E_t^h \cap K(l)$ it holds

$$\begin{aligned} \int_{B_{2^l h}(x) \cap E_t^h \Delta E_{t-h}^h} |\bar{d}_{E_{t-h}^h}| dx &\geq c(2^l h)^{n+2} \quad \text{and} \\ \int_{B_{2^l h}(x) \cap \partial E_t^h} \bar{d}_{E_{t-h}^h}^2 d\mathcal{H}^n &\leq C(2^l h)^{n+2}. \end{aligned}$$

Combing these two yields

$$\int_{B_{2^l h}(x) \cap \partial E_t^h} \bar{d}_{E_{t-h}^h}^2 d\mathcal{H}^n \leq C \int_{B_{2^l h}(x) \cap E_t^h \Delta E_{t-h}^h} |\bar{d}_{E_{t-h}^h}| dx.$$

By applying Besicovitch covering theorem and summing over $l \in \mathbb{Z}$ (see [19, Lemma 3.6] for details) yields

$$\int_{\partial E_t^h} \bar{d}_{E_{t-h}^h}^2 d\mathcal{H}^n \leq \int_{E_t^h \Delta E_{t-h}^h} |\bar{d}_{E_{t-h}^h}| dx$$

which by combining with (4.1) and (4.2) implies

$$h \int_{\partial E_t^h} (H_{E_t^h} - \lambda_{t,h})^2 d\mathcal{H}^n \leq C(P(E_{t-h}^h) - P(E_t^h)).$$

The first inequality then follows by iterating the above.

By [11, Lemma 2.4] it holds

$$|\lambda_{t,h}| \leq C(1 + \|H_{E_t^h} - \lambda_{t,h}\|_{L^1(\partial^* E_t^h)})$$

for a constant that depends on the dimension and on $|E_0|$ and $P(E_0)$. Note that then

$$\|H_{E_t^h}\|_{L^2}^2 + \lambda_{t,h}^2 \leq C(1 + \|H_{E_t^h} - \lambda_{t,h}\|_{L^2}^2).$$

Therefore by the first inequality one obtains

$$\int_{T_1}^{T_2} (\|H_{E_t^h}\|_{L^2}^2 + \lambda_{t,h}^2) dt \leq C \int_{T_1}^{T_2} (1 + \|H_{E_t^h} - \lambda_{t,h}\|_{L^2}^2) dt \leq C(1 + T_2 - T_1).$$

□

The third lemma we need is a quantitative bound on the diameter of the sets (E_t^h) , which is essentially the same as [19, Lemma 3.8].

Lemma 4.3. *Let $(E_t^h)_{t \geq 0}$ be an approximative flat flow starting from E_0 for $h \in (0, 1)$. Then for all $T > 0$ there is R_T , which depends on T , on the dimension and on the diameter of the initial set E_0 , such that $E_t^h \subset B_{R_T}$ for all $t \leq T$.*

Proof. As in [19, Lemma 3.8] define r_t for all $t \leq T$ as

$$r_t := \inf\{r > 0 : E_t^h \subset B_r\}.$$

Arguing as in [19, Lemma 3.8] one deduces that at the point $y \in \partial B_{r_t} \cap \partial E_t^h$ it holds $H_{E_t^h}(y) \geq 0$ and therefore by (4.1)

$$r_t \leq r_{t-h} + h|\lambda_{t,h}|.$$

Iterating this and using Lemma 4.2 yields

$$R_T - R_0 \leq \int_0^T |\lambda_{t,h}| dt \leq \int_0^T (1 + \lambda_{t,h}^2) dt \leq C(1 + T).$$

□

Proof of Theorem 1. Let $(E_t^h)_{t \geq 0}$ be an approximative flat flow starting from E_0 for $h \in (0, 1)$ and fix $T \geq 1$. Then by (4.2) it holds $P(E_t^h) \leq P(E_0)$ and by Lemma 4.3 it holds $E_t^h \subset B_{R_T}$ for all $t \leq T$. I claim that for $0 < t < s$ with $s - t \geq h$ it holds

$$|E_t^h \Delta E_s^h| \leq C\sqrt{t - s}. \quad (4.3)$$

Once (4.3) is obtained, then the convergence of a subsequence $E_t^{h_k} \rightarrow E_t$ in measure follows as in [14, 19].

Let j, k be such that $s \in [jh, (j+1)h)$ and $t \in [(j+k)h, (j+k+1)h)$. Then by applying Lemma 4.1 for $l = \frac{h}{\sqrt{s-t}}$ and by (4.2) one obtains

$$\begin{aligned} |E_t^h \Delta E_s^h| &\leq \sum_{m=j}^{j+k} |E_{mh}^h \Delta E_{(m+1)h}^h| \\ &\leq C \sum_{m=j}^{j+k} \left(\frac{h}{\sqrt{s-t}} P(E_{mh}^h) + \frac{\sqrt{s-t}}{h} \int_{E_{(m+1)h}^h \Delta E_{mh}^h} |\bar{d}_{E_{mh}^h}| dx \right) \\ &\leq C \sum_{m=j}^{j+k} \frac{h}{\sqrt{s-t}} P(E_0) + C\sqrt{s-t} \sum_{m=j}^{j+k} (P(E_{mh}^h) - P(E_{(m+1)h}^h)) \\ &\leq C \frac{kh}{\sqrt{s-t}} P(E_0) + C\sqrt{s-t} P(E_0). \end{aligned}$$

Since $kh \leq 2(s-t)$, one obtains (4.3).

The proof of the consistency principle for $C^{1,1}$ -regular initial sets follows using the arguments in [12]. The volume penalization is used only in [12, Lemma 3.2], but one may overcome this by using the lemma below. \square

Lemma 4.4. *Let $F \subset \mathbb{R}^{n+1}$ be an open and bounded set which satisfies interior and exterior ball condition with radius $r_0 > 0$ and let E be a minimizer of (2.2). There are ρ_0 and h_0 with the property that if G is a set of finite perimeter such that*

$$G \Delta E \subset B_\rho(x) \cap \mathcal{N}_{C_0 h}(\partial F),$$

for $\rho \leq \rho_0$ and $h \leq h_0$ where $\mathcal{N}_{C_0 h}(\partial F) = \{x : \text{dist}(x, \partial F) < C_0 h\}$, then it holds

$$P(E) \leq P(G) + C\rho^{n+1}.$$

Above the constant depends on the dimension, on $r_0, C_0, |F|$ and $P(F)$.

Proof. By approximation one may assume G to be smooth. Since F satisfies interior and exterior ball condition, then by [12, Lemma 3.1] it holds

$$\max_{x \in E \Delta F} \text{dist}(x, \partial F) \leq Ch \tag{4.4}$$

when $h \leq h_0$. As in the proof of Proposition 3.2 the set G does not have the same measure as E and modifies it to \tilde{G} with $|\tilde{G}| = |E|$. Assume again that $|G| < |E|$. Since F satisfies interior ball condition with radius r_0 , there is $y \in G$ such that $B_{r_0/2}(y) \subset G$. By continuity there is $z \in \mathbb{R}^{n+1}$ such that $|z-x| \geq 2\rho_0$ and $|G \cup B_{r_0/2}(z)| = |E|$ when ρ_0 is small. Define $\tilde{G} = G \cup B_{r_0/2}(z)$. Then by the minimality of E and by (4.4) and by the assumption $G \Delta E \subset B_\rho(x) \cap \mathcal{N}_{C_0 h}(\partial F)$ it holds

$$P(E) \leq P(\tilde{G}) + C|\tilde{G} \Delta E| \leq P(\tilde{G}) + C\rho^{n+1}.$$

Arguing as in [9] one then deduces

$$\begin{aligned} P(\tilde{G}) - P(G) &\leq \mathcal{H}^n(\partial B_{r_0/2}(z) \setminus G) - \mathcal{H}^n(\partial G \cap B_{r_0/2}(z)) \\ &\leq \frac{2(n+1)|B_1|}{r_0} |B_{r_0/2}(z) \setminus G| \\ &\leq C|\tilde{G}\Delta E| \leq C\rho^{n+1}. \end{aligned}$$

and the claim follows. \square

The paper concludes with Theorem 2. To this aim I recall the definition of a distributional solution of (1.1) from [14].

Definition 4.5. Family of sets of finite perimeter $(E_t)_{t \geq 0}$ is a distributional solution to (1.1) starting from $E_0 \subset \mathbb{R}^{n+1}$ if the following holds:

- (1) for almost every $t > 0$ the set E_t has mean curvature H_{E_t} in a distributional sense and for every $T > 0$

$$\int_0^T \|H_{E_t}\|_{L^2(\partial^* E_t)}^2 dt < \infty.$$

- (2) There exists $v : \mathbb{R}^{n+1} \times (0, \infty) \rightarrow \mathbb{R}$ with $v \in L^2(0, T; L^2(\partial^* E_t))$ such that for every $\phi \in C_0^1(\mathbb{R}^{n+1} \times [0, \infty))$ it holds

$$\begin{aligned} - \int_0^T \int_{\partial^* E_t} v \phi d\mathcal{H}^n dt &= \int_0^T \int_{\partial^* E_t} (H_{E_t} - \bar{H}_{E_t}) \phi d\mathcal{H}^n dt, \\ \int_0^T \int_{E_t} \partial_t \phi dx dt + \int_{E_0} \phi(\cdot, 0) dx &= - \int_0^T \int_{\partial^* E_t} v \phi d\mathcal{H}^n dt. \end{aligned}$$

Proof of Theorem 2. The proof is exactly the same as [19, Theorem 2.3].

Note that Proposition 3.2 implies that the sets E_t^h are $(Ch^{-1/2}, c\sqrt{h})$ -minimizers of the perimeter, i.e., for every F with $F\Delta E_t^h \subset B_{c\sqrt{h}}(x_0)$ it holds

$$P(E_t^h) \leq P(F) + \frac{C}{\sqrt{h}} |E_t^h \Delta F|. \quad (4.5)$$

\square

ACKNOWLEDGMENTS

The author is supported by the Academy of Finland grants 314227 and 347550.

REFERENCES

- [1] E. ACERBI, N. FUSCO, M. MORINI. *Minimality via second variation for a nonlocal isoperimetric problem.* Comm. Math. Phys. **322** (2013), 515–557.
- [2] F. ALMGREN, *Existence and regularity almost everywhere of solutions to elliptic variational problems with constraints.* Mem. Amer. Math. Soc. **4**(165):viii+199p, (1976).
- [3] F. ALMGREN, J.E. TAYLOR, L. WANG, *Curvature-driven flows: a variational approach.* SIAM J. Control Optim. **31**(2), 387–438 (1993).
- [4] G. ANTONELLI, E. PASQUALETTO, M. POZZETTA, *Isoperimetric sets in spaces with lower bounds on the Ricci curvature.* Nonlinear Anal. **220** (2022), Paper No. 112839, 59 pp.
- [5] G. BELLETTINI, V. CASELLES, A. CHAMBOLLE, M. NOVAGA, *The volume preserving crystalline mean curvature flow of convex sets in \mathbb{R}^N .* J. Math. Pures Appl. (9) **92** (2009), 499–527.

- [6] L. BRONSARD, B. STOTH, *Volume-preserving mean curvature flow as a limit of a nonlocal Ginzburg-Landau equation*. SIAM J. Math. Anal. **28** (1997), 769–807
- [7] W. CARTER, A. ROOSEN, J. CAHN, J. TAYLOR. *Shape evolution by surface diffusion and surface attachment limited kinetics on completely faceted surfaces*. Acta Metallurgica et Materialia **43** (1995), 4309–4323.
- [8] E. GONZALES, U. MASSARI, I. TAMANINI, *On the regularity of boundaries of sets minimizing perimeter with a volume constraint*. Indiana Univ. Math. J. **32** (1983), 25–37.
- [9] M. GRÜTER, *Boundary regularity for solutions of a partitioning problem*. Arch. Rational Mech. Anal. **97** (1987), 261–270.
- [10] V. JULIN, M. MORINI, M. PONSIGLIONE, E. SPADARO, *The Asymptotics of the Area-Preserving Mean Curvature and the Mullins-Sekerka Flow in Two Dimensions*, Math. Ann., (2022), Early online. <https://doi.org/10.1007/s00208-022-02497-3>.
- [11] V. JULIN, J. NIINIKOSKI, *Quantitative Alexandrov Theorem and asymptotic behavior of the volume preserving mean curvature flow*. Preprint 2020.
- [12] V. JULIN, J. NIINIKOSKI, *Consistency of the flat flow solution to the volume preserving mean curvature flow*. Preprint 2022, arXiv:2206.05002.
- [13] T. LAUX, *Weak-strong uniqueness for volume-preserving mean curvature flow*. Preprint arXiv:2205.13040.
- [14] S. LUCKHAUS, T. STÜRZENHECKER, *Implicit time discretization for the mean curvature flow equation*. Calc. Var. Partial. Diff. Eq. **3** (1995), 253–271.
- [15] F. MAGGI, *Sets of finite perimeter and geometric variational problems. An introduction to geometric measure theory*. Cambridge Studies in Advanced Mathematics, 135. Cambridge University Press, Cambridge (2012).
- [16] P.W. MICHOR, D. MUMFORD, *Riemannian geometries on spaces on planar curves*. J. Eur. Mat. Soc. **8**(1), 2006, 1–28.
- [17] M. MORINI, M. PONSIGLIONE, E. SPADARO, *Long time behaviour of discrete volume preserving mean curvature flows*. J. Reine Angew. Math. **784** (2022), 27–51.
- [18] L. MUGNAI, C. SEIS, *On the coarsening rates for attachment-limited kinetics*. SIAM J. Math. Anal. **45** (2013), 324–344.
- [19] L. MUGNAI, C. SEIS, E. SPADARO, *Global solutions to the volume-preserving mean-curvature flow*. Calc. Var. Partial. Diff. Eq. **55** (2016), Art. 18, 23 pp.
- [20] L. SIMON *Introduction to Geometric Measure Theory*. Tsinghua Lectures (2014).

VESA JULIN, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF JYVÄSKYLÄ,
P.O. BOX 35, 40014 JYVÄSKYLÄ, FINLAND
Email address: vesa.julin@jyu.fi