

Square distance functions are Polyak-Łojasiewicz and vice-versa

Guillaume Garrigos

GARRIGOS@LPSM.PARIS

*Université Paris Cité and Sorbonne Université, CNRS
Laboratoire de Probabilités, Statistique et Modélisation
F-75013 Paris, France*

Abstract

This short note gathers known results to state that the squared distance function to a (nonconvex) closed set of an Euclidean space is Polyak-Łojasiewicz. As a fuzzy reciprocal, we also recall that every Polyak-Łojasiewicz function can be bounded from below by the squared distance function to its set of minimizers.

Keywords: Optimization, Polyak-Łojasiewicz.

1. Introduction

The Polyak-Łojasiewicz (PŁ) property (see Definition 1 below) appeared in a work by Polyak (1963, Теорема 4), in parallel to the development of Łojasiewicz’s inequalities (1963, Théorème 4) for the study of analytic gradient flows. The machine learning optimization community recently gained interest for this property (Karimi et al., 2016), because making this hypothesis allows to obtain results in a nonconvex setting which are quantitatively similar to what is usually obtained in a strongly convex setting (Foster et al., 2018; Lei et al., 2019).

While it is generally emphasized that the PŁ property can be derived from strong convexity (or other weakened notions), most works seem to elude the question of what are *nonconvex* Polyak-Łojasiewicz functions, except for a few toy examples. Note that we are considering the original PŁ inequality, which is *global* by nature, and not “local” PŁ inequalities which can be derived for some machine learning models (Hardt and Ma, 2016; Li and Yuan, 2017; Liu et al., 2022).

In this letter we put the emphasis on a class of functions which we show to be PŁ: the squared distance to closed sets (Theorem 9). Not only this provides a class of functions illustrating the PŁ property, which can have an interesting pedagogical purpose, but we also show that every PŁ function can be bounded from below (and above, if smooth) by a squared distance function (Corollary 12). The first result does not seem to appear in the literature, even though it is a simple application of standard results in variational analysis. The second was essentially proved by Bolte et al. (2010), but might have been missed by the machine learning community, and its proof is adapted here to fit our needs.

2. Definitions and notations

Given a closed set $\Omega \subset \mathbb{R}^N$, we note $d_\Omega : \mathbb{R}^N \rightarrow \mathbb{R}$ the *distance function to Ω*

$$(\forall x \in \mathbb{R}^N) \quad d_\Omega(x) := \inf_{p \in \Omega} \|p - x\|,$$

and $\text{proj}_\Omega : \mathbb{R}^N \rightrightarrows \mathbb{R}^N$ the *projection mapping onto Ω*

$$(\forall x \in \mathbb{R}^N) \quad \text{proj}_\Omega(x) := \underset{p \in \Omega}{\text{argmin}} \|p - x\|.$$

Note that we do not assume Ω to be convex, so the projection $\text{proj}_\Omega(x)$ could be a set larger than a singleton. But it is always nonempty, since the function $p \mapsto \|p - x\|$ is coercive and continuous. We recall if needed that d_Ω is Lipschitz continuous (Rockafellar and Wets, 2009, Example 9.6).

Given a function $f : \mathbb{R}^N \rightarrow \mathbb{R}$, and $x \in \mathbb{R}^N$, we note $\partial^F f(x)$ the *Fréchet subdifferential of f at x* defined by

$$x^* \in \partial^F f(x) \Leftrightarrow \liminf_{y \rightarrow x} \frac{f(y) - f(x) - \langle x^*, y - x \rangle}{\|y - x\|} \geq 0.$$

Moreover, we note $\partial^L f(x)$ the *Limiting subdifferential of f at x* , which is composed of vectors $x^* \in \partial^L f(x)$ for which

$$(\exists x_n \rightarrow x)(\exists x_n^* \rightarrow x^*) \quad f(x_n) \rightarrow f(x) \text{ and } x_n^* \in \partial^F f(x_n).$$

It is immediate from the definitions that $\partial^L(\alpha f)(x) = \alpha \partial^L f(x)$ for any $\alpha \geq 0$ and that $\partial^F f(x) \subset \partial^L f(x)$. We recall if needed that if f is continuously differentiable at x , then $\partial^F f(x) = \partial^L f(x) = \{\nabla f(x)\}$ (Rockafellar and Wets, 2009, Exercise 8.8); and that if f is convex then $\partial^F f(x)$ and $\partial^L f(x)$ both coincide with the classical convex subdifferential (Rockafellar and Wets, 2009, Proposition 8.12). Finally, we will say that x is a *Limiting critical point* of f if $0 \in \partial^L f(x)$.

We now turn to the main property of interest in this document. Because square distance functions to nonconvex sets are nonsmooth, we have to consider a nonsmooth version of the PŁ property. We are implicitly making choices when doing so, which we justify in Remark 15.

Definition 1. Let $f : \mathbb{R}^N \rightarrow \mathbb{R}$. We say that f is **Polyak-Łojasiewicz (PŁ)** for short) if there exists $\mu > 0$ such that

$$(\forall x \in \mathbb{R}^N)(\forall x^* \in \partial^L f(x)) \quad f(x) - \inf f \leq \frac{1}{2\mu} \|x^*\|^2.$$

Remark 2 (Invexity). The PŁ property is not related to convexity (see for instance Figure 2 for a nonconvex PŁ function), but shares a strong global property with convex functions, which is that every Limiting critical point must be a global minimizer. See Example 16 on why the word *Limiting* is important here.




Figure 1: Graph of $f(x) = \frac{1}{2}d_{\Omega}(x)^2$, where $\Omega = \{(t, t^2) \in \mathbb{R}^2 \mid t \in \mathbb{R}\}$.

The PL property describes functions having a global quadratic growth with respect to its minimizer set. One can also consider functions having a more general polynomial growth:

Definition 3. Let $f : \mathbb{R}^N \rightarrow \mathbb{R}$, $p > 1$ and $\mu > 0$. We say that f is **globally p -Łojasiewicz** if there exists $\mu > 0$ such that

$$(\forall x \in \mathbb{R}^N)(\forall x^* \in \partial^L f(x)) \quad f(x) - \inf f \leq \frac{1}{q\mu^p} \|x^*\|^q, \quad \text{where } \frac{1}{p} + \frac{1}{q} = 1.$$

Example 4 (Power norms). Let $f(x) = \frac{\mu}{p} \|x\|^p$ for some $p > 1$ and $\mu > 0$. Then it is easy to verify that f is globally p -Łojasiewicz with constant μ .

Example 5 (Uniformly convex functions). Let $f : \mathbb{R}^N \rightarrow \mathbb{R}$ be p -uniformly convex for $p \geq 2$, in the sense that there exists $\mu > 0$ such that for all $x, y \in \mathbb{R}^N$ and all $\alpha \in [0, 1]$,

$$f((1 - \alpha)x + \alpha y) + \alpha(1 - \alpha) \frac{\mu}{p} \|y - x\|^p \leq (1 - \alpha)f(x) + \alpha f(y).$$

Then f is globally p -Łojasiewicz with constant μ (Garrigos et al., 2022, Lemma A.6). When $p = 2$ we recover the well-known fact that strongly convex functions are Polyak-Łojasiewicz.

3. The subdifferential of distance functions

Lemma 6. Let $\Omega \subset \mathbb{R}^N$ be closed and nonempty. Then

1. for every $x \in \Omega$, $0 \in \partial^L d_{\Omega}(x)$;
2. for every $x \notin \Omega$, $\partial^L d_{\Omega}(x) = \frac{x - \text{proj}_{\Omega}(x)}{d_{\Omega}(x)}$;
3. for every $x \in \mathbb{R}^N$, $\partial^L d_{\Omega}(x) \neq \emptyset$.

Proof. For item 1, note that x is a minimizer of d_{Ω} , so the optimality condition $0 \in \partial^L d_{\Omega}(x)$ holds, see (Rockafellar and Wets, 2009, Theorem 10.1). For item 2, the proof can be found in (Rockafellar and Wets, 2009, Example 8.53). Item 3 is a direct consequence of the two previous items, and the fact that the projection on a nonempty closed set is always nonempty. \square

Using a suitable chain rule, we obtain a similar result for powers of distance function.

Lemma 7. *Let $\Omega \subset \mathbb{R}^N$ be closed and nonempty. Let $p > 1$, $\mu > 0$ and let $f(x) := \frac{\mu}{p}d_\Omega(x)^p$. Then*

1. *for all $x \in \Omega$, $\partial^L f(x) = \{0\}$;*
2. *for all $x \notin \Omega$, $\partial^L f(x) = \mu d_\Omega(x)^{p-2}(x - \text{proj}_\Omega(x))$;*
3. *for every $x \in \mathbb{R}^N$, $\partial^L f(x) \neq \emptyset$.*

Proof. Let us write $f = \phi \circ d_\Omega$, where $\phi(t) = \frac{\mu}{p}t^p$. Using the chain rule¹ in (Mordukhovich, 2012, Theorem 1.110.ii), we can write that

$$\partial^L f(x) = \partial^L (\phi'(d_\Omega(x))d_\Omega(\cdot)) (x).$$

Here $\phi'(d_\Omega(x)) = \mu d_\Omega(x)^{p-1}$, so

$$\partial^L f(x) = \partial^L (\mu d_\Omega(x)^{p-1}d_\Omega(\cdot)) (x) = \mu d_\Omega(x)^{p-1} \partial^L d_\Omega(x).$$

Now, use Lemma 6 to see that $\partial^L d_\Omega(x) \neq \emptyset$. So for $x \in \Omega$ we have $d_\Omega(x)^{p-1} = 0$ (because $p > 1$) which implies that $\partial^L f(x) = \{0\}$. Instead, if $x \notin \Omega$, we can use Lemma 6 and the fact that $d_\Omega(x) \neq 0$ to conclude that $\partial^L f(x) = \mu d_\Omega(x)^{p-2}(x - \text{proj}_\Omega(x))$. \square

4. Łojasiewicz inequalities for distance functions

Theorem 8. *Let $\Omega \subset \mathbb{R}^N$ be closed and nonempty. Let $p > 1$, $\mu > 0$ and let $f(x) := \frac{\mu}{p}d_\Omega(x)^p$. Then f is globally p -Łojasiewicz with constant μ .*

Proof. Let $x \in \mathbb{R}^N$ and let us verify that the Łojasiewicz inequality in Definition 3 holds true. Note that here $\inf f = 0$. If $x \in \Omega$ then the inequality is trivial, because $f(x) = 0$ and from Lemma 7 we know that $\partial^L f(x) = \{0\}$. If $x \notin \Omega$, then we use Lemma 7 to write, for all $x^* \in \partial^L f(x)$,

$$\|x^*\|^q = \mu^q d_\Omega(x)^{(p-2)q} d_\Omega(x)^q = \mu^q d_\Omega(x)^p = \mu^{q-1} p f(x),$$

from which we see that Definition 3 is verified. \square

Theorem 9. *Let $\Omega \subset \mathbb{R}^N$ be closed and nonempty. Let $\mu > 0$ and let $f(x) := \frac{\mu}{2}d_\Omega(x)^2$. Then f is Polyak-Łojasiewicz with constant μ .*

Proof. Apply Theorem 8 with $p = q = 2$. \square

Using similar arguments, we can also see that (powers of) distance functions verify two other related properties, which are typically used for describing the geometry of a convex function. See (Bolte et al., 2017; Garrigos et al., 2022) for more details and references on those properties in a convex setting.

Proposition 10. *Let $\Omega \subset \mathbb{R}^N$ be closed and nonempty. Let $p > 1$, $\mu > 0$ and let $f(x) := \frac{\mu}{p}d_\Omega(x)^p$. Then*

1. To help with the notations in Mordukhovich (2012): one must look at (Mordukhovich, 2012, Eq. 1.61) and take $\varphi(x, y) := \phi(y)$ which is strictly differentiable for $p > 1$.

1. f is **globally p -conditioned** with constant μ :

$$(\forall x \in \mathbb{R}^N) \quad \frac{\mu}{p} d_{\operatorname{argmin} f}(x)^p \leq f(x) - \inf f;$$

2. f is **globally p -submetric regular** with constant μ :

$$(\forall x \in \mathbb{R}^N)(\forall x^* \in \partial^L f(x)) \quad \mu d_{\operatorname{argmin} f}(x)^{p-1} \leq \|x^*\|.$$

Proof. The result follows again from Lemma 7. If $x \in \Omega$ the inequalities are trivial, and if $x \notin \Omega$ we use the fact that $d_{\operatorname{argmin} f}(x) = d_{\Omega}(x)$, $f(x) - \inf f = \frac{\mu}{p} d_{\Omega}(x)^p$, and $\|x^*\| = \mu d_{\Omega}(x)^{p-1}$ for all $x^* \in \partial^L f(x)$. \square

5. Distance bounds for globally Łojasiewicz functions

In the previous section we have seen that square distance functions are examples of PL functions. In this section we provide bounds justifying the idea that a PL function is a function growing like the square distance to its minimizers.

Theorem 11. *Let $f : \mathbb{R}^N \rightarrow \mathbb{R}$ be continuous and such that $\operatorname{argmin} f \neq \emptyset$. Let $p > 1$, $\mu > 0$, and let f be globally p -Łojasiewicz with constant μ . Then f is globally p -conditioned with constant $\mu/(p-1)^{p-1}$:*

$$(\forall x \in \mathbb{R}^N) \quad \frac{1}{(p-1)^{p-1}} \frac{\mu}{p} d_{\operatorname{argmin} f}(x)^p \leq f(x) - \inf f.$$

Proof. This result comes from (Bolte et al., 2010, Theorem 24), where the authors need a few extra assumptions (which they do for simplifying their presentation). Since we drop those assumptions, we give here the details of the proof. Let $x \in \mathbb{R}^N \setminus \operatorname{argmin} f$ be fixed, and let us consider a sequence $(x_k)_{k \in \mathbb{N}}$ satisfying $x_0 = x$ and

$$x_{k+1} \in \operatorname{argmin}_{u \in \mathbb{R}^N} f(u) + \frac{1}{2} \|u - x_k\|^2.$$

Note that our assumption that f is bounded from below guarantees that $f + \frac{1}{2} \|\cdot - x_k\|^2$ is coercive. Together with the continuity of f , we conclude that such sequence always exists. Following (Bolte et al., 2010, Lemma 23) and using the definition of our sequence, we write for every $k \in \mathbb{N}$ and every $u \in \mathbb{R}^N$ that

$$\frac{1}{2} \|x_{k+1} - x_k\|^2 \leq \frac{1}{2} \|u - x_k\|^2 + f(u) - f(x_{k+1}).$$

Taking the infimum over the level set $S_{x_{k+1}} := \{u \in \mathbb{R}^N \mid f(u) \leq f(x_{k+1})\}$, and noting that $x_{k+1} \in S_{x_{k+1}}$, we deduce that

$$\|x_{k+1} - x_k\| = d_{S_{x_{k+1}}}(x_k).$$

Now let $\varphi(t) := \frac{p-1}{q^{\frac{1}{q}} \mu^{\frac{1}{p}}} t^{\frac{1}{p}}$ with $\varphi'(t) = \frac{1}{q^{\frac{1}{q}} \mu^{\frac{1}{p}}} t^{\frac{-1}{q}}$ and $\varphi^{-1}(t) = \frac{1}{(p-1)^{p-1}} \frac{\mu}{p} t^p$. It is easy to see that our global p -Łojasiewicz assumption on f can be rewritten as $1 \leq \varphi'(f(x) - \inf f) \|x^*\|$,

for $x^* \in \partial^L f(x)$. Using a chain rule from (Rockafellar and Wets, 2009, Proposition 10.19.b), we know that

$$\partial^L(\varphi \circ (f - \inf f))(x) \subset \varphi'(f(x) - \inf f) \partial^L f(x).$$

We can use now (Drusvyatskiy et al., 2015, Proposition 4.6) to deduce, with the notations of (Bolte et al., 2010), that $|\nabla(\varphi \circ (f - \inf f))|(x) \geq 1$, which means that the assumptions of (Bolte et al., 2010, Corollary 4) are satisfied, and that we can now write

$$\|x_{k+1} - x_k\| = d_{S_{x_{k+1}}}(x_k) \leq \varphi(f(x_k) - \inf f) - \varphi(f(x_{k+1}) - \inf f).$$

Now use the fact that $x_0 = x$, with the triangular inequality, to write

$$\begin{aligned} \|x_k - x\| &\leq \sum_{k=0}^{K-1} \|x_{k+1} - x_k\| \leq \sum_{k=0}^{K-1} \varphi(f(x_k) - \inf f) - \varphi(f(x_{k+1}) - \inf f) \\ &\leq \varphi(f(x_0) - \inf f) - \varphi(f(x_K) - \inf f) \leq \varphi(f(x) - \inf f). \end{aligned}$$

On the one hand, the above inequality implies that the sequence x_k has finite length, which in particular means that it converges to some $x_\infty \in \mathbb{R}^N$, when $k \rightarrow +\infty$. On the second hand, we can pass to the limit on k to obtain

$$\|x_\infty - x\| \leq \varphi(f(x) - \inf f).$$

To conclude the proof, it remains to prove that $x_\infty \in \operatorname{argmin} f$. To see this, apply first-order optimality conditions to the definition of x_{k+1} to write

$$x_{k+1}^* := x_k - x_{k+1} \in \partial^L f(x_{k+1}),$$

and pass to the limit in the global p -Łojasiewicz inequality to obtain

$$f(x_\infty) - \inf f = \lim_{k \rightarrow +\infty} f(x_{k+1}) - \inf f \leq \lim_{k \rightarrow +\infty} \frac{1}{q\mu^p} \|x_{k+1}^*\|^q = 0,$$

where the last equality comes from the fact that $\|x_{k+1}^*\| = \|x_k - x_{k+1}\| \rightarrow 0$. This proves the claim after writing

$$\varphi^{-1}(d_{\operatorname{argmin} f}(x)) \leq \varphi^{-1}(\|x - x_\infty\|) \leq f(x) - \inf f.$$

□

Next corollaries illustrate the idea that PL functions essentially behave like square distance functions.

Corollary 12. *Let $f : \mathbb{R}^N \rightarrow \mathbb{R}$ be continuous and such that $\operatorname{argmin} f \neq \emptyset$. Let f be Polyak-Łojasiewicz with constant $\mu > 0$. Then f is globally 2-conditioned with constant μ :*

$$(\forall x \in \mathbb{R}^N) \quad \frac{\mu}{2} d_{\operatorname{argmin} f}(x)^2 \leq f(x) - \inf f.$$

Proof. Apply Theorem 11 with $p = 2$. □

Corollary 13. *Let $f : \mathbb{R}^N \rightarrow \mathbb{R}$ be differentiable and such that $\operatorname{argmin} f \neq \emptyset$. Let f be Polyak-Łojasiewicz with constant $\mu > 0$, and L -smooth² with constant $L > 0$. Then f behaves like a squared distance function, in the sense that*

$$(\forall x \in \mathbb{R}^N) \quad \frac{\mu}{2} d_{\operatorname{argmin} f}(x)^2 \leq f(x) - \inf f \leq \frac{L}{2} d_{\operatorname{argmin} f}(x)^2.$$

Proof. The first inequality was proved in the previous Corollary. For the second inequality, use the smoothness of f (see e.g. Lemma 1.2.3 in [Nesterov \(2004\)](#)) to write for all $x^* \in \operatorname{argmin} f$ that

$$f(x) - f(x^*) - \langle \nabla f(x^*), x - x^* \rangle \leq \frac{L}{2} \|x - x^*\|^2.$$

The conclusion follows from the fact that $f(x^*) = \inf f$, $\nabla f(x^*) = 0$ and by taking $x^* = \operatorname{proj}_{\operatorname{argmin} f}(x)$. \square

6. Advanced comments

Remark 14 (On the continuity of f). In Theorem 11 we assume that $f : \mathbb{R}^N \rightarrow \mathbb{R}$ is continuous. It is a simple exercise to show that the result is still true if we simply assume that $f : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$ is proper and lower semi-continuous.

Remark 15 (On the choice of the subdifferential). In Definitions 1 and 3 we require the Łojasiewicz inequalities to hold for every $x^* \in \partial^L f(x)$. We could have considered a definition involving the Fréchet subdifferential $\partial^F f(x) \subset \partial^L f(x)$ instead. But the larger the subdifferential, the stronger the result, and we are indeed able to prove our main result in terms of the Limiting subdifferential. One could wonder if the result remains true if we consider yet a larger subdifferential, such as the Clarke subdifferential ([Clarke, 1990](#)). As can be seen in the next counterexample, the answer is no, due to the fact that this Clarke subdifferential can detect sharp local maxima and contain 0 at a point which is not a minimizer (a property incompatible with the PŁ definition). This is not surprising since it is well-known that the Clarke subdifferential is sometimes *too big* (see e.g. the discussion in Section 1.4.4 of [Mordukhovich \(2012\)](#)).

Example 16 (PŁ can fail with Clarke subgradients). Let $\Omega = \mathbb{S}^2$ be the unit circle in \mathbb{R}^2 , and $f(x) = \frac{1}{2} d_{\Omega}(x)^2$ (see Figure 6). At $x = 0$, we compute $f(x) - \inf f = \frac{1}{2}$, and (see Lemma 7) $\partial^L f(x) = x - \operatorname{proj}_{\Omega}(x) = 0 - \mathbb{S} = \mathbb{S}$. Note how $x = 0$ is not a Limiting critical point, even though it is a local maximum! We recover as expected the fact that with $\mu = 1$ we have

$$f(0) - \inf f = \frac{1}{2} \leq \frac{1}{2\mu} \|x^*\|^2 \text{ for all } x^* \in \partial^L f(0) = \mathbb{S}.$$

If we now compute the Clarke subdifferential at $x = 0$, we obtain $\partial^C f(0) = \overline{\operatorname{co}} \partial^L f(0)$ (use Theorem 3.57 from [Mordukhovich \(2012\)](#) with the fact that f is locally Lipschitz continuous). That is, $\partial^C f(0)$ is the unit ball \mathbb{B} , which contains the origin. This means that the inequality

$$f(0) - \inf f = \frac{1}{2} \leq \frac{1}{2\mu} \|x^*\|^2 \text{ for all } x^* \in \partial^C f(0) = \mathbb{B}$$

2. We say that f is L -smooth if ∇f is L -Lipschitz continuous.




Figure 2: Graph of $f(x) = \frac{\mu}{2}d_{\Omega}(x)^2$, where $\mu = 20$ and $\Omega \subset \mathbb{R}^2$ is the unit circle. We can see a (nonsmooth) local maximizer at the origin.

cannot be true when $x^* = 0$, whatever constant $\mu > 0$ we consider. The problem here is that $x = 0$ is a *Clarke* critical point which is not minimizer.

Acknowledgments

The author would like to thank Robert Gower who accepted to proofread this document.

References

- Jérôme Bolte, Aris Daniilidis, Olivier Ley, and Laurent Mazet. Characterizations of Łojasiewicz inequalities: Subgradient flows, talweg, convexity. *Transactions of the American Mathematical Society*, 362(6):3319–3363, 2010. ISSN 0002-9947, 1088-6850. doi: 10.1090/S0002-9947-09-05048-X.
- Jérôme Bolte, Trong Phong Nguyen, Juan Peypouquet, and Bruce W. Suter. From error bounds to the complexity of first-order descent methods for convex functions. *Mathematical Programming*, 165(2):471–507, 2017. ISSN 1436-4646. doi: 10.1007/s10107-016-1091-6.
- F. Clarke. *Optimization and Nonsmooth Analysis*. Classics in Applied Mathematics. Society for Industrial and Applied Mathematics, 1990. ISBN 978-0-89871-256-8.
- D. Drusvyatskiy, A. Ioffe, and A. Lewis. Curves of Descent. *SIAM Journal on Control and Optimization*, 53(1):114–138, 2015. ISSN 0363-0129. doi: 10.1137/130920216.
- Dylan J. Foster, Ayush Sekhari, and Karthik Sridharan. Uniform convergence of gradients for non-convex learning and optimization. *Advances in Neural Information Processing Systems*, 31, 2018.

- Guillaume Garrigos, Lorenzo Rosasco, and Silvia Villa. Convergence of the forward-backward algorithm: Beyond the worst-case with the help of geometry. *Mathematical Programming*, 2022. ISSN 1436-4646. doi: 10.1007/s10107-022-01809-4.
- Moritz Hardt and Tengyu Ma. Identity matters in deep learning. *arXiv preprint arXiv:1611.04231*, 2016.
- Hamed Karimi, Julie Nutini, and Mark Schmidt. Linear convergence of gradient and proximal-gradient methods under the Polyak-Łojasiewicz condition. In *Joint European Conference on Machine Learning and Knowledge Discovery in Databases*, pages 795–811. Springer, 2016.
- Yunwen Lei, Ting Hu, Guiying Li, and Ke Tang. Stochastic Gradient Descent for Nonconvex Learning without Bounded Gradient Assumptions. *arXiv:1902.00908 [cs, math, stat]*, 2019.
- Yuanzhi Li and Yang Yuan. Convergence analysis of two-layer neural networks with relu activation. *Advances in neural information processing systems*, 30, 2017.
- Chaoyue Liu, Libin Zhu, and Mikhail Belkin. Loss landscapes and optimization in over-parameterized non-linear systems and neural networks. *Applied and Computational Harmonic Analysis*, 2022.
- S. Łojasiewicz. Une propriété topologique des sous-ensembles analytiques réels. In *Les Équations Aux Dérivées Partielles (Paris, 1962)*. Éditions du Centre National de la Recherche Scientifique, 1963.
- Boris S. Mordukhovich. *Variational Analysis and Generalized Differentiation I: Basic Theory*. Springer, Berlin ; New York, 1st ed. 2006. corr., 2nd printing 2013 edition, 2012. ISBN 978-3-540-25437-9.
- Yurii Nesterov. *Introductory Lectures on Convex Optimization*, volume 87. Springer Science & Business Media, 2004.
- B.T. Polyak. Gradient methods for the minimisation of functionals. *USSR Computational Mathematics and Mathematical Physics*, 3(4):864–878, 1963. ISSN 00415553. doi: 10.1016/0041-5553(63)90382-3.
- R. Tyrrell Rockafellar and Roger J.-B. Wets. *Variational Analysis*. Springer, New York, 2009. ISBN 978-3-540-62772-2.