

SYMPLECTOMORPHISMS AND SPHERICAL OBJECTS IN THE CONIFOLD SMOOTHING

AILSA KEATING AND IVAN SMITH

ABSTRACT. Let X denote the ‘conifold smoothing’, the symplectic Weinstein manifold which is the complement of a smooth conic in T^*S^3 , or equivalently the plumbing of two copies of T^*S^3 along a Hopf link. Let Y denote the ‘conifold resolution’, by which we mean the complement of a smooth divisor in $\mathcal{O}(-1) \oplus \mathcal{O}(-1) \rightarrow \mathbb{P}^1$. We prove that the compactly supported symplectic mapping class group of X splits off a copy of an infinite rank free group, in particular is infinitely generated; and we classify spherical objects in the bounded derived category $D(Y)$ (the three-dimensional ‘affine A_1 -case’). Our results build on work of Chan-Pomerleano-Ueda and Toda, and both theorems make essential use of working on the ‘other side’ of the mirror.

CONTENTS

1. Introduction	1
2. Wrapped Fukaya categories of Liouville manifolds	3
3. The conifold smoothing	7
4. The conifold resolution	14
5. Proof of Theorem 1.1	17
6. Classification of sphericals	21
7. Miscellanea	29
References	32

1. INTRODUCTION

1.1. **Results.** Consider the open symplectic manifold

$$X = \{u_1v_1 = z - 1, u_2v_2 = z + 1\} \subset \mathbb{C}^4 \times \mathbb{C}^*$$

where $u_i, v_i \in \mathbb{C}$ and $z \in \mathbb{C}^*$, with the restriction ω of the standard exact Kähler form from $\mathbb{C}^4 \times \mathbb{C}^*$. Projection to $z \in \mathbb{C}^*$ defines a Morse-Bott-Lefschetz fibration on X , from which one sees that X is also a plumbing of two copies of T^*S^3 along a Hopf link.

Theorem 1.1. *There is a split injection $\mathbb{Z}^{*\infty} \rightarrow \pi_0 \text{Symp}_c(X, \omega)$, where $\mathbb{Z}^{*\infty}$ denotes the free group on countably infinitely many generators. In particular, the symplectic mapping class group $\pi_0 \text{Symp}_c(X, \omega)$ is infinitely generated.*

The generators of the image $\mathbb{Z}^{*\infty}$ in $\pi_0 \text{Symp}_c(X, \omega)$ will be Dehn twists in certain Lagrangian spheres in X .

Corollary 1.2. *There is a 3-dimensional Stein domain $(X', \partial X')$, with both X' and $\partial X'$ simply connected, and with $\pi_0 \text{Symp}_c(X')$ infinitely generated.*

By contrast, for a simply-connected compact six-manifold X' with simply connected boundary, $\pi_0 \text{Diff}_c(X')$ is always of finite type, cf. [Kup] and Lemma 7.8.

The Milnor fibres of the A_k -singularities for $k > 1$ contain infinitely many exact Lagrangian spheres up to Lagrangian isotopy, but these are obtained from a single sphere under the action of the symplectic mapping class group. We show, in Corollary 7.2, that the action of $\pi_0 \text{Symp}_c(X')$ on the set of isotopy classes of Lagrangian 3-spheres in X' has infinitely many orbits, answering a folk question sometimes attributed to Fukaya.

The mirror Y to X is obtained from the small resolution $\mathcal{O}(-1) \oplus \mathcal{O}(-1) \rightarrow \mathbb{P}^1$ of the 3-fold ordinary double point $xy = (1+w)(1+t) \subset \mathbb{C}^4$ by removing the pullback of the divisor $\{wt = 0\}$. Our methods also resolve the long-standing open problem of classifying all the spherical objects in $D(Y)$ (i.e. classifying sphericals in the ‘three-dimensional affine A_1 -case’). One consequence of this classification is:

Theorem 1.3. *The spherical objects in $D(Y)$ form a single orbit under a natural action of the pure braid group PBr_3 .*

It may be worth highlighting that Theorem 1.1, which is a purely symplectic statement, relies essentially on passing to the other side of mirror symmetry, to exploit a computation of a space of stability conditions accessible only via algebraic geometry; whilst Theorem 1.3, which is a purely algebro-geometric statement, relies essentially on passing back to the symplectic setting, to exploit constraints arising from Nielsen-Thurston theory and symplectic dynamics on surfaces.

Remark 1.4. Propositions 6.4 and 6.16 together imply that if $\phi \in D(Y)$ is a Torelli autoequivalence (i.e. acting trivially on K -theory) which is not a power of a spherical twist, and S, S' are any spherical objects in $D(Y)$, then the total rank of $\text{Ext}^*(\phi^k(S), S')$ grows exponentially in k (moreover the exponential growth rate is an algebraic integer and is at least $\log(2)/12$ [Pen91], etc). This follows from the identification of the ranks of these morphism groups with geometric intersection numbers of curves on a surface, and classical results in surface dynamics. This connects to ideas around categorical entropy, cf. [DHKK14].

1.2. Context. Understanding the symplectic mapping class group is a classical and long-standing problem in symplectic topology. Theorem 1.1 is the first example of a (finite type) symplectic manifold for which the compactly supported symplectic mapping class group is known to be infinitely generated. It complements a result of [SS20], see also [Smi22], which constructed compact symplectic manifolds for which the ‘symplectic Torelli group’ (of mapping classes acting trivially on cohomology) was infinitely generated, and a result of [AS21] which gave a compact smooth manifold with a family of symplectic forms ω_t for which the rank of $\pi_0 \text{Symp}(X, \omega_t)$ grew unbounded with t . In contrast, we expect the symplectic mapping class group of a $K3$ surface, or of (the mirror to) a log Calabi-Yau surface, to be finitely generated.

The classification of spherical objects in either the derived category of an algebraic variety or the Fukaya category of a symplectic manifold is also a well-known and active area of research. In particular, for the Milnor fibres of type A surface singularities a large body of work yielded a complete classification [KS02, IU05, Wu14]. Much less is known on threefolds. For ‘small’ categories associated to flopping contractions, namely the full subcategory of objects supported on the exceptional locus and with trivial push-forward, a classification was

recently obtained in [HWa], see also [SW]. Theorem 1.3 is the first example which treats the ‘big’ category of all complexes supported on the exceptional locus, where one encounters an affine hyperplane arrangement for the associated chamber of flops (or stability conditions) rather than a finite hyperplane arrangement.

1.3. Remarks on the proofs. Homological mirror symmetry for the conifold has been established by Chan, Pomerleano and Ueda in [CPU16], both for the wrapped Fukaya category $\mathcal{W}(X)$ and the compact subcategory generated by the ‘compact core’ of X . In particular, there is a \mathbb{C} -linear equivalence $\mathcal{W}(X) \simeq D(Y)$. Crucially, both categories are also linear over a larger ring $R = SH^0(X) \simeq \Gamma(\mathcal{O}_Y)$. The group of stability conditions on $D(Y)$, and the group of autoequivalences of that category, was analysed by Toda [Tod08, Tod09] (see also [HWb] for closely related results in a more general setting).

The heart of the proof of Theorem 1.1 is to show that the part of the autoequivalence group of $\mathcal{W}(X)$ coming from compactly supported symplectomorphisms surjects to the subgroup of R -linear autoequivalences of $D(Y)$ which moreover act trivially on K -theory. This uses in an essential way the detailed knowledge of $\text{Auteq } D(Y)$ arising from the computation of the space of stability conditions. By contrast, the key ingredient in the proof of Theorem 1.3 is a growth rate estimate for ranks of Floer cohomology under autoequivalences, which is proved using a combination of localisation-type theorems (to reduce Floer theory computations to a two-dimensional surface inside the six-dimensional X) together with classical results of Nielsen and Thurston on dynamics of pseudo-Anosov maps.

The symplectic topology of two-dimensional log Calabi-Yau surfaces has been fruitfully illuminated through mirror symmetry [GHK15, HKa, HKb]. This paper treats one simple three-dimensional example, but many of the techniques and results seem likely to generalise.

Acknowledgements. The authors are grateful to Spencer Dowdall, Paul Hacking, Daniel Pomerleano, Oscar Randal-Williams, Paul Seidel and Michael Wemyss for helpful conversations and correspondence.

This material is based upon work supported by the National Science Foundation under Grant No. 1440140, while the authors were in residence at the Mathematical Sciences Research Institute in Berkeley, California, during the Fall semester 2022. A.K. was partially supported by EPSRC Open Fellowship EP/W001780/1. I.S. is grateful to the Clay Foundation for support at MSRI as a Clay Senior Scholar, and to EPSRC for support through Frontier Research grant EP/X030660/1 (in lieu of an ERC Advanced Grant).

2. WRAPPED FUKAYA CATEGORIES OF LIOUVILLE MANIFOLDS

2.1. Preliminary wrapped Fukaya categories. Let (W, θ) denote a finite type complete Liouville manifold (i.e. the completion of a Liouville domain \mathring{W} by cylindrical ends). A Liouville domain (\mathring{W}, θ) canonically determines a finite type complete Liouville manifold (W, θ) . A Lagrangian is cylindrical in (W, θ) , resp. (\mathring{W}, θ) , if it is invariant under the Liouville vector field outside a compact set, resp. in a collar neighbourhood of the boundary. Note that any cylindrical Lagrangian for (W, θ) is exact Lagrangian isotopic, via the inverse Liouville flow, to a Lagrangian which is the completion of a cylindrical Lagrangian in (\mathring{W}, θ) .

Definition 2.1. The preliminary wrapped Fukaya category $\mathcal{W}_{\text{pr}}(W, \theta)$ is defined as the wrapped Fukaya category set-up as in [GPS, Section 2.4], with the following choices: the category is ungraded, with coefficient ring $\mathbb{Z}/2$; and Lagrangians are exact and cylindrical

(no further brane data is required as there are no gradings). We will write $\mathcal{W}_{\text{pr}}(\overset{\circ}{W}, \theta)$ for $\mathcal{W}_{\text{pr}}(W, \theta)$ when we wish to emphasise the choice of domain.

‘Preliminary’ is intended as in [Sei08b]. We will later restrict to Lagrangians which admit brane structures including a choice of *Spin*-structure and grading, yielding a category $\mathcal{W}(W, \theta)$ which is defined over \mathbb{Z} and \mathbb{Z} -graded.

2.2. Actions of symplectomorphisms. An exact symplectomorphism of $(\overset{\circ}{W}, \theta)$ is a diffeomorphism $\phi : \overset{\circ}{W} \rightarrow \overset{\circ}{W}$ such that $\phi^*\theta = \theta + df$, where f needn’t have support in $\text{Int}(\overset{\circ}{W})$ (and is formally defined on a small open neighbourhood of $\overset{\circ}{W}$ in W). Let $\text{Symp}(\overset{\circ}{W})$ be the group of all symplectomorphisms of $\overset{\circ}{W}$, equipped with the C^∞ topology. Let $\text{Symp}_{\text{ex}}(\overset{\circ}{W}, \theta)$ denote the subgroup of exact symplectomorphisms; the connected component of the identity in $\text{Symp}_{\text{ex}}(\overset{\circ}{W}, \theta)$ is $\text{Ham}(\overset{\circ}{W}, \theta)$ the subgroup of Hamiltonian diffeomorphisms of $(\overset{\circ}{W}, \theta)$.

Lemma 2.2. *The group $\pi_0 \text{Symp}_{\text{ex}}(\overset{\circ}{W}, \theta)$ acts on $\mathcal{W}_{\text{pr}}(W, \theta)$.*

Proof. Fix $\phi \in \text{Symp}_{\text{ex}}(\overset{\circ}{W}, \theta)$. Cylindrical Lagrangians for $(\overset{\circ}{W}, \theta)$ get mapped under ϕ to cylindrical Lagrangians for $(\overset{\circ}{W}, \phi^*\theta)$. This induces a map $\phi : \mathcal{W}_{\text{pr}}(\overset{\circ}{W}, \theta) \rightarrow \mathcal{W}_{\text{pr}}(\overset{\circ}{W}, \phi^*\theta)$ taking a Lagrangian L to $\phi(L)$ and pulling back all Floer data.

On the other hand, as ϕ is an exact symplectomorphism of $\overset{\circ}{W}$, $\phi^*\theta = \theta + df$, for some smooth function f . We claim that $(\overset{\circ}{W}, \theta + tdf)$ is a Liouville domain for all $t \in [0, 1]$. For $t = 1$, this is true tautologically via pullback, with the Liouville flow $Z_{\theta+df}$ for $(\overset{\circ}{W}, \theta + df)$ being the pushforward under ϕ of the Liouville flow Z_θ for $(\overset{\circ}{W}, \theta)$. Now notice, first, that $(\partial\overset{\circ}{W}, \ker(\theta + tdf))$ is a contact manifold for all $t \in [0, 1]$: the positivity one needs comes from interpolating between two volume forms of the same sign. And second, notice that the vector field ω -dual to $\theta + tdf$ is a convex combination of the two outward pointing vector fields Z_θ and $Z_{\theta+df}$, and so is also outward pointing. Thus $\{(\overset{\circ}{W}, \theta + tdf)\}_{t \in [0, 1]}$ is a smooth family of Liouville domains.

The cylindrical completions of $\{(\overset{\circ}{W}, \theta + tdf)\}_{t \in [0, 1]}$ give a smooth family of finite type complete Liouville manifolds, say $\{(W, \theta_t)\}_{t \in [0, 1]}$. This is automatically a convex symplectic deformation in the sense of [SS05, Section 2]. By [SS05, Lemma 5], there is a smooth family of diffeomorphisms $h_t : W \rightarrow W$, with $h_0 = \text{id}$, such that $h_t^*\theta_t = \theta_0 + dg_t$, where $(t, x) \mapsto g_t(x)$ is a compactly supported function on $[0, 1] \times W$. Inspection of the proof shows that the family h_t is canonical up to isotopy through diffeomorphisms with the same properties.

This implies that we have a trivial inclusion of Liouville manifolds $(W, \theta) \hookrightarrow (W, \theta_1)$ in the sense of [GPS20, Definition 2.4]. By [GPS20, Lemma 3.41] (see also [GPS, Lemma 3.4]), there is an induced equivalence $c : \mathcal{W}_{\text{pr}}(W, \theta) \rightarrow \mathcal{W}_{\text{pr}}(W, \theta_1)$. Postcomposing $\phi : \mathcal{W}_{\text{pr}}(\overset{\circ}{W}, \theta) \rightarrow \mathcal{W}_{\text{pr}}(\overset{\circ}{W}, \phi^*\theta)$ with the inverse to c gives an autoequivalence of $\mathcal{W}_{\text{pr}}(\overset{\circ}{W}, \theta)$, which we will also denote ϕ .

To ensure that we obtain a group homomorphism, we require a consistency of continuation maps. This follows from the compatibility of the Liouville inclusions obtained from [SS05, Section 2] with concatenation of families of finite type complete Liouville manifolds.

Finally, if ϕ and ϕ' are in the same connected component of $\text{Symp}_{\text{ex}}(\overset{\circ}{W}, \theta)$, they are related by a Hamiltonian isotopy. Furthermore, if ϕ is Hamiltonian, the maps constructed in the two steps of the argument above are evidently quasi-inverse. This completes the proof. \square

Lemma 2.3. *cf. [BEE12, Lemma 1.1]. The inclusion $\text{Symp}_{\text{ex}}(\mathring{W}, \theta) \hookrightarrow \text{Symp}(\mathring{W})$ is a weak homotopy equivalence.*

Proof. An analogous statement for complete Liouville manifolds is contained in [BEE12, Lemma 1.1]. This shows that given an arbitrary symplectomorphism $\psi : (W, \theta) \rightarrow (W', \theta')$ between finite type complete Liouville manifolds, there exists a symplectic isotopy $\sigma_t : (W, \theta) \rightarrow (W, \theta)$, $t \in [0, 1]$, $\sigma_0 = \text{id}$, such that $\psi \circ \sigma_1$ is exact. Inspecting the proof, we see that the isotopy is canonical up to deformation within the class with the same properties, and that the argument can be carried out for families of symplectomorphisms parametrised by arbitrary compact cell complexes.

We now want to upgrade this to a statement for Liouville domains. We start with the argument for a single map. Suppose $\psi : \mathring{W} \rightarrow \mathring{W}'$ is an arbitrary symplectomorphism. Tautologically, this induces a symplectomorphism from the conical extension of (\mathring{W}, θ) , say (W, θ) , to the conical extension of $(\mathring{W}', \psi^*\theta)$, say (W', θ') . Let us also denote this symplectomorphism by ϕ . By construction, the cylindrical end $E = [0, \infty) \times \partial\mathring{W}$ for (W, θ) is taken to the one for $(W, \psi^*\theta)$, with ψ preserving s -level contact shells, $s \in [0, \infty)$, and $\psi \circ \rho_s = \rho'_s \circ \psi$, where ρ_s is the Liouville flow for (W, θ) and ρ'_s the one for (W', θ') .

We now apply [BEE12, Lemma 1.1] to get a symplectic isotopy σ_t , $t \in [0, 1]$, as above. Inspecting their proof, we see that outside a compact set, σ_t is given by integrating a vector field of the form $e^{-s}\tilde{V}$, where \tilde{V} is independent of s . In particular, for sufficiently large s , the contact level-set $\{s\} \times \partial\mathring{W}$ is displaced by C^∞ -small amounts under σ_t , for any $t \in [0, 1]$. (The statement of their lemma only spells this out for C^0 .) On the other hand, under C^∞ small perturbations, both the contact condition and the condition that a vector field be outward pointing are open. Fix such a sufficiently large s . For any $t \in [0, 1]$, let $\mathring{W}_t := W \setminus \sigma_t^{-1}([s, \infty) \times \partial\mathring{W})$. By construction, (\mathring{W}_t, θ) is a Liouville domain with conical completion W , and is canonically Liouville isomorphic (via σ_t) to $(\mathring{W} \cup ([0, s] \times \partial\mathring{W}), \theta)$.

Altogether, we have deformed ψ to an exact symplectomorphism of $(\mathring{W} \cup ([0, s] \times \partial\mathring{W}), \theta)$. Conjugating by the time s Liouville flow now gives the desired map for ψ . Finally, for a family of maps $\{\psi_b\}_{b \in B}$ indexed by a compact cell complex B , one can choose the parameter s uniformly over B . Together with the observation that the argument in [BEE12] works in families, this completes the proof. \square

Corollary 2.4. *The group $\pi_0 \text{Symp}(\mathring{W})$ acts on $\mathcal{W}_{\text{pr}}(W, \theta)$.*

Proof. This now follows immediately from combining Lemmas 2.2 and 2.3. \square

2.3. Gradings and orientations. We now assume that $2c_1(W, d\theta) = 0 \in H^2(W; \mathbb{Z})$, i.e. the bundle $(\Lambda^{\dim W} T^*W)^{\otimes 2}$ is trivial (here \dim denotes the complex dimension). Then W admits a grading, i.e. a choice of fibrewise universal cover for the Lagrangian Grassmanian bundle $\text{Lag}(W) \rightarrow W$, say $\widehat{\text{Lag}}(W) \rightarrow W$. The isomorphism classes of such covers are in one-to-one correspondence with trivialisations of $(\Lambda^{\dim W} T^*W)^{\otimes 2} \simeq W \times \mathbb{C}$, and form an affine space over $H^1(W; \mathbb{Z})$ (see [Sei00, Lemma 2.2]).

Fix a grading on W . Given a Lagrangian $L \subset W$, the trivialisaton $(\Lambda^{\dim W} T^*W)^{\otimes 2} \simeq W \times \mathbb{C}$ determines a (homotopy class of) map $L \rightarrow \mathbb{C}^*$, and so a class in $H^1(L; \mathbb{Z})$, called the Maslov class of L . There exists a lift of $TL \subset \text{Lag}(W)$ to $\widehat{\text{Lag}}(W)$ precisely when L has Maslov class zero; a choice of such a lift is called a grading on L .

We can now upgrade to the ‘full’ version of the wrapped Fukaya category, which will be used for the rest of this article.

Definition 2.5. The wrapped Fukaya category $\mathcal{W}(W, \theta)$ is defined using the set-up in [GPS, Section 2.4], with the following choices: coefficient ring \mathbb{Z} ; Lagrangians are exact, cylindrical, Maslov zero and spin, and, as objects of $\mathcal{W}(W, \theta)$, equipped with brane data comprising a *Spin*-structure and a grading.

The compact Fukaya category $\mathcal{F}(W, \theta)$ is the subcategory of compact Lagrangian submanifolds.

We will often suppress θ from the notation when it is clear from the context which primitive we are referring to.

Remark 2.6. For homological mirror symmetry statements, we’ll base change the ground ring for $\mathcal{W}(W)$ and $\mathcal{F}(W)$ from \mathbb{Z} to \mathbb{C} ; as this will be clear from context, we will also denote those categories by $\mathcal{W}(W)$ and $\mathcal{F}(W)$.

Suppose that we are given $\phi \in \text{Symp}(W)$. We say that ϕ is gradeable if under pullback by ϕ , our choice of trivialisation of $(\Lambda^{\dim W} T^*W)^{\otimes 2}$ is preserved; a grading for ϕ is then a choice of lift of the pullback map to a bundle automorphism of $\widetilde{\text{Lag}}(W)$. We let $\text{Symp}^{\text{gr}} W$ denote the group of graded symplectomorphisms of W , and $\text{Symp}_{\text{ex}}^{\text{gr}}(W, \theta)$ the subgroup of exact graded symplectomorphisms. From the discussion above there’s an exact sequence¹ (see [Sei00, Lemma 2.4])

$$(2.1) \quad 1 \rightarrow \mathbb{Z} \rightarrow \text{Symp}^{\text{gr}}(W) \rightarrow \text{Symp}(W) \rightarrow H^1(W; \mathbb{Z}).$$

If $\phi \in \text{Symp}(W)$ is gradeable and has compact support, it comes with a preferred grading: the one which fixes $\widetilde{\text{Lag}}(W)$ outside of the support of ϕ . This implies that there is a splitting of the forgetful map from $\text{Symp}_c^{\text{gr}}(W)$ to the group of gradeable compactly supported symplectomorphisms.

The results of the previous subsection readily generalise as follows.

Lemma 2.7. *The group $\pi_0 \text{Symp}_{\text{ex}}^{\text{gr}}(\mathring{W}, \theta)$ acts on $\mathcal{W}(W, \theta)$.*

Proof. Given $\phi \in \text{Symp}_{\text{ex}}^{\text{gr}}(\mathring{W}, \theta)$, we get a map $\mathcal{W}(\mathring{W}, \theta) \rightarrow \mathcal{W}(\mathring{W}, \phi^*\theta)$ by pulling back all Floer data, taking a Lagrangian L to $\phi(L)$, and using the choice of grading for ϕ to determine the grading for $\phi(L)$ (the spin structure is just pulled back).

We can now follow the proof of Lemma 2.2, noting that in any given connected component of $\text{Symp}_{\text{ex}}(\mathring{W}, \theta)$, either all maps are gradeable or none of them are. \square

Lemma 2.8. *The inclusion $\text{Symp}_{\text{ex}}^{\text{gr}}(\mathring{W}, \theta) \hookrightarrow \text{Symp}^{\text{gr}}(\mathring{W})$ is a weak homotopy equivalence.*

Proof. This readily follows from Lemma 2.3, by noticing, first, that for any given connected component of $\text{Symp}(\mathring{W})$, resp. $\text{Symp}_{\text{ex}}(\mathring{W}, \theta)$, either all maps in it are gradeable or none of them are; and that for each such gradeable component there are \mathbb{Z} components in $\text{Symp}^{\text{gr}}(\mathring{W})$, resp. $\text{Symp}_{\text{ex}}^{\text{gr}}(\mathring{W}, \theta)$. \square

We then immediately get the following:

Corollary 2.9. *The group $\pi_0 \text{Symp}^{\text{gr}}(\mathring{W})$ acts on $\mathcal{W}(W, \theta)$.*

¹The final map is not a group homomorphism, but its kernel is nonetheless a subgroup of $\text{Symp}(W)$.

The natural module action of symplectic cohomology on wrapped Floer cohomology shows that $\mathcal{W}(W)$ admits the structure of a category linear over $SH^0(W)$. From the discussion above, we immediately get that graded symplectomorphisms act on $\mathcal{W}(X)$ by $SH^0(W)$ -linear equivalences.

3. THE CONIFOLD SMOOTHING

3.1. Descriptions of the conifold smoothing X . Consider $\mathbb{C}^2 \times \mathbb{C}^2 \times \mathbb{C}^*$ with co-ordinates (u_1, v_1, u_2, v_2, z) , equipped with its standard symplectic form $\omega = d\theta$. The *conifold smoothing* X is the symplectic submanifold defined by the equations

$$(3.1) \quad X = \{u_1v_1 = z - 1, u_2v_2 = z + 1\}.$$

It is the fibre $X = X_{1,-1}$ over $(+1, -1) \in (\mathbb{C}^* \times \mathbb{C}^*) \setminus \Delta$ of a family of symplectic submanifolds

$$(3.2) \quad X_{a,b} = \{u_1v_1 = z - a, u_2v_2 = z - b\}.$$

Let $\pi : X_{a,b} \rightarrow \mathbb{C}^*$ denote the projection to the z co-ordinate. This is a Morse-Bott-Lefschetz fibration, meaning that $d\pi$ has transversely non-degenerate critical submanifolds. The smooth fibres are isomorphic to $\mathbb{C}^* \times \mathbb{C}^* = T^*T^2$, and there are singular fibres $(\mathbb{C} \vee \mathbb{C}) \times \mathbb{C}^*$ respectively $\mathbb{C}^* \times (\mathbb{C} \vee \mathbb{C})$ over a respectively b .

Lemma 3.1. *For fixed a, b , symplectic parallel transport for the fibration $\pi : X_{a,b} \rightarrow \mathbb{C}^*$ is globally well-defined.*

Proof. There is a Hamiltonian T^2 -action on $X_{a,b}$, rotating the (u_i, v_i) -planes, and π is a T^2 -equivariant map. Given any symplectic fibration with a fibrewise Hamiltonian action of a Lie group G , each component of the moment map for the G -action is preserved by symplectic parallel transport (cf. [Aur07, Section 5] or [SW, Lemma 4.1]). In the case at hand, the moment map is fibrewise proper; the result follows. \square

Lemma 3.2. *Any matching path γ between the two critical values in the base \mathbb{C}^* of the Morse-Bott-Lefschetz fibration defines a Lagrangian sphere S_γ in X .*

Proof. Paths emanating from $+1 \in \mathbb{C}^*$, resp. $-1 \in \mathbb{C}^*$ have associated Morse-Bott thimbles $S^1 \times D^2$, respectively $D^2 \times S^1$. These restrict to Lagrangian T^2 s in nearby fibres. The sphere S_γ is then presented via the standard genus one Heegaard splitting of the sphere. \square

Lemma 3.3. *$X_{a,b}$ is naturally a Weinstein manifold, with compact core $S^3 \cup_{\text{Hopf}} S^3$.*

Proof. It suffices to consider $X = X_{1,-1}$. Reverse Liouville flow on each T^*T^2 fibre retracts it to the zero-section $\{|u_i| = |v_i|\}$, and retracts the \mathbb{C}^* base to the unit circle in the z -plane. This shows that X retracts to the union $S_0 \cup S_1$, where $S_i = S_{\gamma_i}$ and the paths γ_0 respectively γ_1 form the arcs of the unit circle in the lower respectively upper half-plane in \mathbb{C}_z^* . \square

It follows that X is the Weinstein completion of an open neighbourhood of this compact core, i.e. of a Weinstein domain obtained by plumbing two copies of the disc cotangent bundle D^*S^3 along a Hopf link.

An affine variety determines a (complete) Weinstein symplectic manifold canonically up to exact symplectomorphism [Sei08a, Section 4b]. We will give two further descriptions of X by giving two further descriptions of the affine variety (3.1).

Recall that if $Z \subset \mathbb{C}^2$ is a Zariski open subset and $f : Z \rightarrow \mathbb{C}$ is a regular function, the *spinning* of f is the 3-fold

$$(3.3) \quad \{(x, y, u, v) \in Z \times \mathbb{C}^2 \mid f(x, y) = uv\}.$$

This is a fibration over Z by affine conics \mathbb{C}^* , with singular fibres $\mathbb{C} \vee \mathbb{C}$ along $f^{-1}(0) \subset Z$.

Lemma 3.4. *X can be obtained by spinning $Z = \mathbb{C}^2 \setminus \{xy = 1\}$ along the map $f(x, y) = xy - 2$.*

Proof. Let $g : \mathbb{C}^2 \rightarrow \mathbb{C}$ be given by $g(x, y) = xy - 1 = z$, so $Z = \mathbb{C}^2 \setminus g^{-1}(0)$ and $g : Z \rightarrow \mathbb{C}^*$ has a Lefschetz singular fibre over $z = -1$. The spinning is then a \mathbb{C}^* -fibration over $\mathbb{C}^2 \setminus \{xy = 1\}$ with singular fibres over $\{xy - 1 = 1\}$. It is globally cut out of $\mathbb{C}^2 \times \mathbb{C}^2 \times \mathbb{C}^*$, with co-ordinates (x, y, u, v, z) , by the equations $xy = z + 1$ and $uv = z - 1$. \square

Z contains an immersed Lagrangian 2-sphere, obtained from parallel transport around the unit circle in the z -plane, cf. [Sei, Section 11]. Viewing this as a union of two Lefschetz thimbles fibred over the upper and lower half-circles with common boundary in the fibre $z = 1$, the immersed 2-sphere naturally ‘spins’ to the compact core $S_0 \cup_{\text{Hopf}} S_1$ of X described above, i.e. the matching 3-spheres are obtained as S^1 -fibrations over the thimbles with fibres degenerating over the boundary of the 2-disc.

Our final description of X is as a log Calabi-Yau 3-fold. We won’t use this directly, but it may be of independent interest. Consider $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, which we view as the trivial $\mathbb{P}^1 \times \mathbb{P}^1$ -bundle over the base \mathbb{P}^1 . We blow up this variety along the (disjoint) curves $\mathbb{P}^1 \times \{pt\} \times \{1\}$ and $\{pt\} \times \mathbb{P}^1 \times \{-1\}$ to obtain a variety \overline{X} with an induced map $\overline{X} \rightarrow X \rightarrow \mathbb{P}^1$. Let D denote the divisor given by the union of two smooth fibres of $\overline{X} \rightarrow \mathbb{P}^1$ together with the subset of the toric boundary comprising its four components isomorphic to the first Hirzebruch surface \mathbb{F}_1 , see Figure 1.




FIGURE 1. The toric compactification of X ; blow up the thickened black edges on the cube for $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$. This slices off wedges of the corresponding moment polytope. Two of the four \mathbb{F}_1 -boundary components of the result have been shaded.

Lemma 3.5. *X is symplectically equivalent to the symplectic completion of $\overline{X} \setminus D$.*

Proof. In an affine chart $\mathbb{C}_p \times \mathbb{C}_q \times \mathbb{C}_z^*$ we blow up $\{p = 1 = z\}$ and $\{q = 1 = -z\}$. Taking $[\lambda, \mu] \in \mathbb{P}^1$ and $[\tilde{\lambda}, \tilde{\mu}] \in \mathbb{P}^1$ to be homogeneous co-ordinates on two different copies of \mathbb{P}^1 , the blow-up is given by

$$(p - 1)/(z - 1) = \lambda/\mu; \quad (q - 1)/(z + 1) = \tilde{\lambda}/\tilde{\mu}.$$

Letting $u_1 = q - 1$, $u_2 = p - 1$, $v_1 = \tilde{\mu}/\tilde{\lambda}$ and $v_2 = \mu/\lambda$ gives

$$u_1 v_1 = z + 1, \quad u_2 v_2 = z - 1;$$

since $z \in \mathbb{C}^*$ none of u_i, v_j can be infinite. Now remove the irreducible divisor from the projective blow-up where any of the u_i, v_j co-ordinates do become infinite. This meets the general fibre in its toric boundary; each special fibre is again toric, isomorphic to $(\mathbb{P}^1 \times \mathbb{P}^1) \cup_{\mathbb{P}^1 \times \{pt\}} (\mathbb{P}^1 \times \mathbb{P}^1)$, and the divisor meets this in a hexagon of lines (one does not remove the entire line of intersection of the two components, but only its intersection with the rest of the boundary). \square

3.2. Exact Lagrangian submanifolds. As remarked previously in Lemma 3.2, any matching path γ between the two critical values in the base of the Morse-Bott-Lefschetz fibration $X \rightarrow \mathbb{C}^*$ defines a Lagrangian sphere S_γ in the total space X . We will later be particularly interested in the collection of matching spheres $\{S_i\}_{i \in \mathbb{Z}}$ corresponding to the matching paths given in Figure 2.




FIGURE 2. Lagrangian matching spheres S_i (with matching paths γ_i) and Lagrangian discs L_i (see Theorem 5.1).

Suppose we are given an embedded $S^1 \subset \mathbb{C}^* \setminus \{-1, 1\}$ in the base of the Morse-Bott-Lefschetz fibration, avoiding the critical values. By similar considerations to those in Lemmas 3.1 and 3.2, parallel transporting the T^2 vanishing cycle about the S^1 gives a Lagrangian T^3 . This T^3 will be exact precisely when the S^1 in the base is Hamiltonian isotopic to the unit circle in \mathbb{C}^* . (Recall here that we are using the standard Kähler form for \mathbb{C}^* as an affine manifold.) For any choice of such circle containing all of $\{-1, 0, 1\}$, call the associated exact torus T .

3.3. Gradings. As X is an affine complete intersection in Euclidean space, we have that $2c_1(X) = 0$, and the isomorphism classes of gradings on X form an affine space over $H^1(X; \mathbb{Z}) \cong \mathbb{Z}$. Our convention is that throughout this paper we work with a fixed, distinguished grading for X : the unique one for which the exact Lagrangian torus T defined above has Maslov class zero (this will also be true of any other tori fibred over S^1 s containing 0 in the base of the Morse-Bott-Lefschetz fibration).

We now want to fix gradings for the S_i . Fix an arbitrary grading for S_0 . For $i \neq 0$, we want to grade S_i as follows: first, note that we can make fibrewise Morse perturbations such that S_0 and S_i intersect transversally in $4|i|$ points. One can check that there is a choice of grading for S_i such that: for $i > 0$, as generators for the Floer complex $CF^*(S_0, S_{-i})$, there are $i + 1$ intersection points with grading zero; $2i$ points with grading one; and $i - 1$ points with grading two; and for $i < 0$, the same is true for $CF^*(S_i, S_0)$.

Remark 3.6. These gradings are chosen with mirror symmetry in mind: S_i will be mirror to $\mathcal{O}_C(-i)$ for a fixed $(-1, -1)$ curve C in a threefold Y , see Section 5.

We fix an arbitrary grading on the Lagrangian torus T .

3.4. Symplectomorphisms of X : first considerations. Let (\mathring{X}, θ) be a Liouville domain such that (X, θ) is its associated completion. Following the notation in Section 2.2, let $\text{Symp}(\mathring{X})$ denote the group of all symplectomorphisms of \mathring{X} equipped with the C^∞ topology, and let $\text{Symp}_{\text{ex}}(\mathring{X}, \theta)$ be the subgroup of exact symplectomorphisms. We define graded versions $\text{Symp}^{\text{gr}}(\mathring{X})$ and $\text{Symp}_{\text{ex}}^{\text{gr}}(\mathring{X})$ similarly.

By Lemma 2.8, the inclusion $\text{Symp}_{\text{ex}}^{\text{gr}}(\mathring{X}, \theta) \hookrightarrow \text{Symp}^{\text{gr}}(\mathring{X})$ is a weak homotopy equivalence. We define $\text{Symp}^{\text{gr}}(X)$ to mean $\text{Symp}^{\text{gr}}(\mathring{X})$, and $\text{Symp}_{\text{ex}}^{\text{gr}}(X)$ to mean $\text{Symp}_{\text{ex}}^{\text{gr}}(\mathring{X}, \theta)$. (Similarly for ungraded versions.) For any other Liouville domain (\mathring{X}', θ) with completion (X, θ) , there is a natural weak homotopy equivalence $\text{Symp}^{\text{gr}}(\mathring{X}) \rightarrow \text{Symp}^{\text{gr}}(\mathring{X}')$, and similarly for exact maps. In particular, the group $\pi_0 \text{Symp}^{\text{gr}}(X)$ is independent of choices. By Corollary 2.9, $\pi_0 \text{Symp}^{\text{gr}}(X)$ acts on the Fukaya category $\mathcal{W}(X)$; this readily restricts to an action on $\mathcal{F}(X)$. Let $\text{Symp}_c(X)$ denote the group of compactly supported symplectomorphisms of X .

Lemma 3.7. *Any element $\phi \in \text{Symp}_c(X)$ is strongly exact: there exists a compactly supported function f such that $\phi^*\theta = \theta + df$, where θ is the Liouville form on X .*

Proof. As ϕ is a symplectomorphism, $\phi^*\theta - \theta$ is a closed one-form; moreover, away from a compact set, ϕ is the identity, and so $\phi^*\theta - \theta$ vanishes there. This means it defines a class in $H_c^1(X; \mathbb{R})$. By Poincaré duality, this is isomorphic to $H_5(X; \mathbb{R})$, which vanishes as X has the homotopy type of a 3-cell complex. Thus $\phi^*\theta - \theta = df$ for some function f , locally constant outside the domain of ϕ . Finally, as the boundary at infinity of X is connected, we see that we can assume that f vanishes outside a compact set. \square

Lemma 3.8. *Any element $\phi \in \text{Symp}_c(X)$ is gradeable. Moreover, it admits a preferred grading; this determines a splitting of the forgetful map $\text{Symp}_c^{\text{gr}}(X) \rightarrow \text{Symp}_c(X)$.*

Proof. See [Sei00, Remark 2.5]. As ϕ has compact support, the obstruction to grading it is a map $h \in [X, \mathbb{C}^*]$ such that $h \equiv 1$ outside a compact set, i.e. a class in $H_c^1(X; \mathbb{Z})$. As this vanishes, any such ϕ is gradeable. Finally, as noted in Section 2.3, any gradeable compactly supported symplectomorphism has a distinguished grading: the one which is the identity on $\widehat{\text{Lag}}(X)$ outside a compact set. \square

It is then immediate that $\pi_0 \text{Symp}_c(X)$ acts on $\mathcal{W}(X)$ and $\mathcal{F}(X)$.

There is a well-known source of compactly supported symplectomorphisms: for any matching path γ in the base of the Morse-Bott-Lefschetz fibration, the Dehn twist $\tau_{S_\gamma} \in \pi_0 \text{Symp}_c X$. By [SW, Lemma 4.13], τ_{S_γ} is the monodromy associated to a full right-handed twist in the path $\gamma \subset \mathbb{C}^*$. More precisely, suppose we parametrise the full right-handed twist as a one-parameter family of compactly supported diffeomorphisms of \mathbb{C}^* , giving a map $\rho : [0, 1] \times \mathbb{C}^* \rightarrow \mathbb{C}^*$; then τ_{S_γ} can be interpreted as the monodromy of the family $X_{\rho(t, -1), \rho(t, 1)}$, $t \in [0, 1]$, with the notation from Equation 3.2. We will revisit this in Lemma 3.10.

3.5. The PBr_3 action. We start with some generalities on the pure braid group PBr_3 . We will usually think of it as a subgroup of the fundamental group of the space of configurations of two points in \mathbb{C}^* , with basepoint $\{-1, 1\} \subset \mathbb{C}^*$; equivalently, it's naturally isomorphic to $\pi_1((\mathbb{C}^*)^2 \setminus \Delta)$.

By [MM09, Theorem 2.3], we have a presentation for PBr_3 with three generators R_1, R_2 and R_3 , given by the full twists described in Figure 3, and the sole relation

$$R_1 R_2 R_3 = R_2 R_3 R_1 = R_3 R_1 R_2.$$

The element $R_1R_2R_3$ generates the centre $Z(\text{PBr}_3) \cong \mathbb{Z}$. Geometrically, it corresponds to an inverse Dehn twist in a boundary S^1 .




FIGURE 3. The generators for PBr_3

Quotienting by the center, we get isomorphisms

$$\text{PBr}_3 \simeq \text{PBr}_3 / Z(\text{PBr}_3) \times Z(\text{PBr}_3) \simeq (\mathbb{Z} * \mathbb{Z}) \times \mathbb{Z}.$$

This decomposition as a direct product of groups is of course not unique: for instance we could have $(\mathbb{Z}\langle R_i \rangle * \mathbb{Z}\langle R_j \rangle) \times \mathbb{Z}\langle R_1R_2R_3 \rangle$ for any $i \neq j$.

Given any matching path γ between -1 and $+1$ in \mathbb{C}^* , consider the full twist $t_\gamma \in \text{PBr}_3$. We want to characterise the subgroup generated by these twists.

Proposition 3.9. *Let $\text{PBr}_3^c \triangleleft \text{PBr}_3$ be the subgroup of PBr_3 generated by the twists t_γ . Then:*
 (i) $\psi \in \text{PBr}_3$ lies in PBr_3^c if and only if the winding number of 1 about 0 and the winding number of -1 about 0 are both zero. In other words,

$$\text{PBr}_3^c = \ker\left\{\text{PBr}_3 \xrightarrow{(\text{forget } -1, \text{forget } 1)} \text{PBr}_2 \times \text{PBr}_2\right\}$$

where PBr_2 is thought of as the fundamental group of the configuration space of one point in \mathbb{C}^* , based at $\{1\}$ for the first factor and at $\{-1\}$ for the second.

(ii) We have an isomorphism $\text{PBr}_3^c \cong \mathbb{Z}^{*\infty}$, with generators the full twists t_i in the matching paths γ_i for S_i from Figure 2.

Proof. Given any γ , it is immediate that $t_\gamma \in \ker\left\{\text{PBr}_3 \xrightarrow{(\text{forget } -1, \text{forget } 1)} \text{PBr}_2 \times \text{PBr}_2\right\}$, and so the inclusion one way is clear.

Now suppose that we're given a pure braid

$$\psi \in K = \ker\left\{\text{PBr}_3 \xrightarrow{(\text{forget } -1, \text{forget } 1)} \text{PBr}_2 \times \text{PBr}_2\right\}.$$

To analyse this, first consider simply forgetting the first strand. This gives a short exact sequence of groups:

$$0 \rightarrow \mathbb{Z} * \mathbb{Z} \rightarrow \text{PBr}_3 \xrightarrow{\text{forget } 1} \mathbb{Z} \rightarrow 0$$

where to see that the kernel is a $\mathbb{Z} * \mathbb{Z}$, think about pulling the first and second strands (corresponding to -1 and 0 with our current choices) taut. The kernel now presents as the subgroup of $\pi_1 \text{Conf}(\mathbb{C}^*, 2)$ generated by pushing the third point around the second point; and pushing the third point around the first point, i.e. R_2 and R_3 in the notation of Figure 3. This is naturally the fundamental group of the wedge of two S^1 s.

Putting things together, we have

$$K = \ker\left\{\mathbb{Z}\langle R_2 \rangle * \mathbb{Z}\langle R_3 \rangle \xrightarrow{\text{forget } 1} \text{PBr}_2\right\}$$

A pure braid $\psi \in \mathbb{Z}\langle R_2 \rangle * \mathbb{Z}\langle R_3 \rangle$ is in K if and only if the power of R_2 in the abelianisation of $\mathbb{Z}\langle R_2 \rangle * \mathbb{Z}\langle R_3 \rangle$ vanishes. To conclude, consider the infinite cyclic cover of the wedge of

two circles with group of deck transformations $\mathbb{Z}\langle R_2 \rangle$. The image of π_1 of this cover in $\mathbb{Z}\langle R_2 \rangle * \mathbb{Z}\langle R_3 \rangle$ is K ; we now see that it is a $\mathbb{Z}^{*\infty}$ with generators $R_2^{-i}R_3R_2^i = t_i$, $i \in \mathbb{Z}$. \square

Mapping $R_2^{-i}R_3R_2^i$ to the Dehn twist τ_{S_i} gives a map $\mathrm{PBr}_3^c \rightarrow \pi_0 \mathrm{Symp}_c X$; this is compatible with the description of τ_{S_i} as the monodromy of the full twist in γ_i from [SW, Lemma 4.13]. We will now upgrade this to a representation of the whole of PBr_3 as symplectomorphisms; in order to do this, we have to enlarge the group we are working in to include non-compactly supported exact symplectomorphisms.

Lemma 3.10. *There is a map*

$$(3.4) \quad \mathrm{PBr}_3 \rightarrow \pi_0 \mathrm{Symp}_{\mathrm{ex}} X$$

extending the composition $\mathrm{PBr}_3^c \rightarrow \pi_0 \mathrm{Symp}_c X \rightarrow \pi_0 \mathrm{Symp}_{\mathrm{ex}} X$. This is compatible with the action of PBr_3 on the collection of matching paths: if γ is a matching path, $R \in \mathrm{PBr}_3$, and $\gamma' = R \cdot \gamma$, then the image of R in $\pi_0 \mathrm{Symp}_{\mathrm{ex}} X$ takes S_γ to $S_{\gamma'}$ (up to Hamiltonian isotopy).

Proof. Consider $\mathbb{C}^4 \times \mathbb{C}^* \times (\mathbb{C}^*)^2$ with coordinates $(u_1, v_1, u_2, v_2, z, a, b)$ and its standard Kähler form. Consider the smooth affine subvariety

$$(3.5) \quad \mathcal{X} = \{u_1 v_1 = z - a, u_2 v_2 = z - b\}.$$

Projection to the final two coordinates (a, b) defines a symplectic fibre bundle $\mathcal{X} \rightarrow (\mathbb{C}^*)^2$. The fibre over $(a, b) \in (\mathbb{C}^*)^2 \setminus \Delta$ is the space $X_{a,b}$ from Equation 3.2, and the family has three-fold ordinary double points along $\Delta = \{a = b\}$.

Now on the one hand, $\pi_1((\mathbb{C}^*)^2 \setminus \Delta)$ is naturally the pure braid group PBr_3 . On the other hand, given any path $\gamma : [0, 1] \rightarrow (\mathbb{C}^*)^2 \setminus \Delta$ in the smooth locus, we can construct parallel transport maps over γ which are globally defined and strongly exact, by the argument of [Kea15, Lemma 2.2]. Symplectic monodromy then defines the representation (3.4). The fact that the fibres over points of Δ have a single ordinary double point implies that the monodromy around a path which is a meridian to Δ (oriented anticlockwise as the boundary of a disc positively transverse to Δ at one point) is a Dehn twist (see also [SW, Lemma 4.13]). This implies that (3.4) extends the previously defined $\mathrm{PBr}_3^c \rightarrow \pi_0 \mathrm{Symp}_c X$. It is then standard, cf. [KS02], that the action is compatible with the natural action on matching paths. \square

Remark 3.11. The proof readily gives a stronger result: we have a natural map $\mathrm{PBr}_3 \rightarrow \pi_0 \mathrm{Symp}_\partial X$, where $\mathrm{Symp}_\partial(X)$ denotes the group of strongly exact symplectomorphisms, i.e. diffeomorphisms ϕ of X such that $\phi^*\theta = \theta + df$ for a compactly supported function f , where θ is the Liouville form on X . We will not need this for our purposes.

Remark 3.12. Let us justify the notation PBr_3^c . Using the Morse-Bott-Lefschetz perspective, given any element of PBr_3 , we could try to build a symplectomorphism of X by first getting fibrewise symplectomorphisms via parallel transport, and then upgrading to a symplectomorphism of the total space. (For technical reasons we use a different set-up in the present work.) For what elements $\psi \in \mathrm{PBr}_3$ does parallel transport induce the identity for fibres near infinity? Consider some arc $\tilde{\gamma}$ from $+\infty$ to 0, avoiding the marked points. We want to know when the monodromy induced by the concatenated path $\tilde{\gamma} \# (-\psi \tilde{\gamma})$ is the identity. Now notice that this is equivalent to asking that $\tilde{\gamma} \# (-\psi \tilde{\gamma})$ have winding number 0 about both -1 and $+1$, which is in turn equivalent to ψ being in the kernel of the two forgetful maps to $\mathrm{PBr}_2 \times \mathrm{PBr}_2$.

Let W^{2n} be a symplectic manifold with $c_1(W) = 0$, and fix a compatible almost complex structure J . A choice of holomorphic volume form $\Theta_J \in \Gamma(\Lambda_{\mathbb{C}}^n T^*W)$ defines a distinguished fibrewise universal cover $\widetilde{\text{Lag}}(W) \rightarrow \text{Lag}(W)$ of the fibre bundle $\text{Lag}(W) \rightarrow W$ whose fibre is the Lagrangian Grassmannian $Gr_{\text{Lag}}(T_w W) \cong U(n)/O(n)$. Explicitly, we have a map

$$\Theta_J : \text{Lag}(W) \rightarrow S^1, \quad (v_1, \dots, v_n) \mapsto \Theta_J(v_1 \wedge \dots \wedge v_n) / |\Theta_J(v_1 \wedge \dots \wedge v_n)|$$

and we consider the bundle $\widetilde{\text{Lag}}(W) \rightarrow \text{Lag}(W)$ with fibre the pairs $(\Lambda, t) \in \text{Lag}(W) \times \mathbb{R}$ with $\Theta_J(\Lambda) = e^{i\pi t}$.

Lemma 3.13. *The map $\text{PBr}_3 \rightarrow \pi_0 \text{Sym}_{\text{ex}}(X)$ can be lifted to graded symplectomorphisms.*

Proof. The form $d \log(z) \wedge d \log(u_1) \wedge d \log(u_2)$ defines a holomorphic volume form on (3.2), and so the family $\mathcal{X} \rightarrow \text{Conf}_2(\mathbb{C}^*)$ over configuration space from (3.5) admits a relative holomorphic volume form. Via the preceding construction, this defines a fibration $\mathcal{X}^\infty \rightarrow \mathcal{X}$ with fibre the universal cover $U(n)/O(n)$. A choice of Ehresmann connection for this fibration enables one to lift (not necessarily uniquely) the parallel transport maps for $\mathcal{X} \rightarrow \text{Conf}_2(\mathbb{C}^*)$ to parallel transport maps for $\mathcal{X}^\infty \rightarrow \text{Conf}_2(\mathbb{C}^*)$, which in particular shows that these symplectomorphisms admit coherent gradings. \square

Choice of gradings for the action of PBr_3 . We now fix a preferred lift $\text{PBr}_3 \rightarrow \pi_0 \text{Sym}_{\text{ex}}^{\text{gr}}(X)$, by specifying the lifts of each of the three generators R_1, R_2 and R_3 . First, the image of R_3 is the Dehn twist $\tau_0 = \tau_{S_0}$; we give it the preferred grading for a compactly supported symplectomorphism. Second, we grade the image of R_2 , say λ , by asking for $\lambda^i S_0 = S_{-i}$ as graded objects. Finally, to choose the grading for the image of R_1 , say ρ , recall that $R_1 R_2 R_3$ generates the center of PBr_3 , and corresponds to the inverse Dehn twist in a boundary parallel circle; in particular, the product $\rho \circ \lambda \circ \tau_{S_0}$ takes S_0 to itself as an ungraded object. Fixing a grading on ρ is the same as fixing the shift of S_0 under $\rho \circ \lambda \circ \tau_{S_0}$; we choose this to be [1].

Remark 3.14. The central map $\rho \circ \lambda \circ \tau_{S_0}$ is isotopic in $\pi_0 \text{Sym}(X)$ to the identity (this follows from the proof of Lemma 3.10), though through maps acting non-trivially on the boundary. From a purely symplectic perspective, the grading of the central element $\rho \circ \lambda \circ \tau_{S_0}$ could have been chosen to be any power of a shift. (In particular, there isn't a 'preferred' shift as we are not in $\pi_0 \text{Sym}_c(X)$.) We will later see that from an HMS perspective it is natural to choose the shift [1].

3.6. K -theoretic actions of symplectomorphism groups. We have that $K(\mathcal{W}(X)) = \mathbb{Z}^2$ generated by two cotangent fibres F_0 and F_1 , by for instance [GPS, Theorem 1.13] or [CRGG]. This is naturally identified with $H_3(X, \partial_\infty X; \mathbb{Z})$, where $\partial_\infty X$ denotes the conical end of X . Moreover, there is a pairing

$$(3.6) \quad K(\mathcal{F}(X)) \times K(\mathcal{W}(X)) \rightarrow \mathbb{Z}, \quad (K, L) \mapsto \chi(HF^*(L, K)).$$

Evaluating this on fibres and zero-sections shows that $\mathbb{Z}^2 = \langle S_0, S_1 \rangle \rightarrow K(\mathcal{F}(X))$ is injective. (Note we don't have an explicit description for $K(\mathcal{F}(X))$.)

Definition 3.15. Let the numerical Grothendieck group of $\mathcal{F}(X)$, $K_{\text{num}}(\mathcal{F}(X))$, be the group $K(\mathcal{F}(X))/\ker$, where \ker is the subgroup generated by compact Lagrangians L s.t. $\chi(HF^*(L, -)) = 0$ for every object $-$ in $\mathcal{W}(X)$.

The pairing 3.6 descends to a non-degenerate pairing $K_{\text{num}}(\mathcal{F}(X)) \times K(\mathcal{W}(X)) \rightarrow \mathbb{Z}$. This implies that $K_{\text{num}}(\mathcal{F}(X)) = \mathbb{Z}^2 = \langle S_0, S_1 \rangle$. This is naturally identified with $H_3(X; \mathbb{Z})$, and

the non-degenerate pairing

$$K_{num}(\mathcal{F}(X)) \times K(\mathcal{W}(X)) \rightarrow \mathbb{Z}$$

is identified with the intersection pairing

$$H_3(X; \mathbb{Z}) \times H_3(X, \partial_\infty X; \mathbb{Z}) \rightarrow \mathbb{Z}.$$

Both $\text{Symp}^{\text{gr}}(X)$ and $\text{Symp}_c(X)$ (with preferred gradings) act on $K(\mathcal{F}(X))$ and $K(\mathcal{W}(X))$ compatibly with this pairing. This implies that there is an induced action on the numerical Grothendieck group. On the other hand, as the intersection pairing of S_0 with both S_0 and S_1 vanishes, and similarly for S_1 , we see that the standard map $H_3(X; \mathbb{Z}) \rightarrow H_3(X, \partial_\infty X; \mathbb{Z})$ has zero image. The following is now immediate:

Corollary 3.16. *Suppose that $f \in \pi_0 \text{Symp}^{\text{gr}}(X)$. Then f_* acts on $K_{num}(\mathcal{F})$. Moreover, f induces the identity on $K(\mathcal{W})$ if and only if it induces the identity on $K_{num}(\mathcal{F})$.*

For arbitrary compactly supported maps, we have the following.

Lemma 3.17. *Compactly supported homeomorphisms act trivially on $H_3(X; \mathbb{Z})$.*

Proof. Let $\phi \in \text{Homeo}_c(X)$ be a compactly supported homeomorphism. Considering the induced action on the long exact sequence for the pair $(X, \partial_\infty X)$, we get a commutative diagram:

$$(3.7) \quad \begin{array}{ccccccc} \dots & \longrightarrow & H_3(\partial_\infty X) & \longrightarrow & H_3(X) & \longrightarrow & H_3(X, \partial_\infty X) & \longrightarrow & \dots \\ & & \phi_* \downarrow & & \phi_* \downarrow & & \phi_* \downarrow & & \\ \dots & \longrightarrow & H_3(\partial_\infty X) & \longrightarrow & H_3(X) & \longrightarrow & H_3(X, \partial_\infty X) & \longrightarrow & \dots \end{array}$$

where all coefficient rings are \mathbb{Z} . By assumption, $\phi_* : H_3(\partial_\infty X) \rightarrow H_3(\partial_\infty X)$ is the identity. On the other hand, recall that in this case the map $H_3(X) \rightarrow H_3(X, \partial_\infty X)$ has zero image. It's then immediate that $\phi_* : H_3(X) \rightarrow H_3(X)$ is the identity. \square

Combined with Corollary 3.16, this implies that any $f \in \pi_0 \text{Symp}_c X$ acts trivially on $K_{num}(\mathcal{F})$ and $K(\mathcal{W})$.

Lemma 3.17 will also imply that there are infinity many orbits of Lagrangian spheres in X ; see Corollary 7.2 for a more general statement. On the other hand, by the proof of Proposition 3.9, we see that all of the S_γ are contained in the orbits of the S_i , $i \in \mathbb{Z}$, under the action of the group generated by Dehn twists in the S_i .

Remark 3.18. We'll see in Section 5 that the subgroup $\mathbb{Z}^{*\infty}$ which maps to $\pi_0 \text{Symp}_c X$ actually split injects into it. This will use mirror symmetry. If one only wants to show that the action of $\mathbb{Z}^{*\infty}$ is faithful, one could do this instead by using a generalisation of Khovanov–Seidel's work on faithful braid group actions, see Proposition 6.4.

4. THE CONIFOLD RESOLUTION

Consider the threefold ordinary double point $\{xy = (1+w)(1+t)\} \subset \mathbb{C}^4$. This has a small resolution, which is the total space of $\mathcal{O}(-1) \oplus \mathcal{O}(-1) \rightarrow \mathbb{P}^1$. Let D be the pullback under the small resolution of the divisor $\{wt = 0\} \subset \mathbb{C}^4$. Let Y be the open subset of $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$ given by the complement of D . Let $C \subset \mathcal{O}(-1) \oplus \mathcal{O}(-1)$ denote the zero-section, which lies in the complement of the divisor D and hence in Y .

A brief calculation shows that $\text{Pic}(Y) \cong \mathbb{Z}$. This is generated by a line bundle \mathcal{L} such that $\mathcal{O}_C(i) \otimes \mathcal{L} \cong \mathcal{O}_C(i+1)$ (by the push-pull formula).

Let $R = \Gamma(\mathcal{O}_Y)$. We have that $\text{Spec } R \cong \{xy = (1+w)(1+t)\} \setminus \{wt = 0\} \subset \mathbb{C}^4$ is the blow-down of Y .

Remark 4.1. Start with $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$; blow up $Z = \mathbb{P}^1 \times \{-1\} \times \{0\} \cup \{-1\} \times \mathbb{P}^1 \times \{\infty\}$, and let F be the proper transform of the toric divisor for $(\mathbb{P}^1)^3$. Then one can check that the variety Y is isomorphic to $\text{Bl}_Z(\mathbb{P}^1)^3 \setminus Z$. In particular, Y is log Calabi-Yau.

We write $D(Y)$ for the bounded derived category of Y .

Definition 4.2. The category $\mathcal{D} \subset D(Y)$ is the full subcategory of $D(Y)$ of complexes whose cohomology sheaves are (set-theoretically) supported on $C \subset Y$.

Remark 4.3. Definition 4.2 is taken from [HWb]. The paper [CPU16] defines \mathcal{D} as the derived category of the abelian subcategory of $\text{Coh}(Y)$ of sheaves (set-theoretically) supported on C . This yields the same category as Definition 4.2 by [Sta18, Lemma 0AEF]. The alternative definition makes it manifest that \mathcal{D} is determined by a formal neighbourhood of C in Y . On the other hand, the formal neighbourhood of a flopping curve is determined by its length [Rei83], where a flopping curve has normal bundle $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$ if and only if it has length one. So the category \mathcal{D} depends only on C being a $(-1, -1)$ -curve.

From Remark 4.3, or directly, one sees that \mathcal{D} is generated by \mathcal{O}_C and $\mathcal{O}_C(-1)$, so the Grothendieck group $K(\mathcal{D})$ of \mathcal{D} is \mathbb{Z}^2 , generated by $[\mathcal{O}_C], [\mathcal{O}_C(-1)]$.

Proposition 4.4. *Suppose ϕ is an autoequivalence of $D(Y)$. Then ϕ is Fourier-Mukai, i.e. induced by an object $P_\phi \in D(Y \times Y)$. Moreover, P_ϕ is unique up to isomorphism.*

Proof. Existence of P_ϕ is in [LO10, Corollary 2.16]. (See also [Bal, Theorem 7.13] and [CS14, Theorem 1.1].) Uniqueness follows from Toën [Toë07, Corollary 8.12], see [LO10, Remark 9.11]. □

The contraction map $Y \rightarrow \text{Spec}(R)$ gives $D(Y)$ the structure of an R -linear category, via pullback. In particular, we can consider those autoequivalences of $D(Y)$ which are R -linear.

Lemma 4.5. *Let ϕ be an R -linear autoequivalence of $D(Y)$ and $x \in Y \setminus C$. Then ϕ preserves the skyscraper sheaf \mathcal{O}_x up to a shift.*

Proof. For any sheaf $\mathcal{E} \in D(Y)$, one can consider the ideal

$$\text{Ann}(\mathcal{E}) := \{r \in R \mid r \cdot \mathcal{E} = 0\}.$$

By R -linearity, $\text{Ann } \mathcal{O}_x = \text{Ann}(\phi \mathcal{O}_x)$. On the other hand, note that $\text{Ann } \mathcal{O}_x$ is simply the maximal ideal of functions vanishing at x . Thus $\phi \mathcal{O}_x$ is also supported at x , and hence by [Huy06, Lemma 4.5] agrees with \mathcal{O}_x up to a shift. □

To simplify notation, we will denote $Y \times_{\text{Spec } R} Y$ by $Y \times_R Y$.

Lemma 4.6. *R -linear autoequivalences of $D(Y)$ preserve \mathcal{D} and are induced by Fourier-Mukai kernels with support on $Y \times_R Y$.*

Proof. Let ϕ be an R -linear autoequivalence of $D(Y)$ with Fourier-Mukai kernel $P_\phi \in D(Y \times Y)$. By Lemma 4.5, for each $x \in Y \setminus C$, the fibre of $\text{Supp } P_\phi \rightarrow Y$ above x is zero dimensional. On the other hand, the fibre of $\text{Supp } P_\phi \rightarrow Y$ above any point is connected [Huy06, Lemma

6.11]. Thus for any $x \in Y \setminus C$, the fibre of $\text{Supp } P_\phi \rightarrow Y$ above x is simply $\{x\}$. By [Huy06, Lemma 3.29], this means that

$$(\text{Supp } P_\phi)|_{Y \setminus C \times Y \setminus C} = \Delta_{Y \setminus C}.$$

Using [Huy06, Lemma 6.11] again, it follows that

$$\text{Supp } P_\phi \subset \Delta_Y \cup C \times C.$$

Both parts of the lemma follow. \square

Let $\text{Stab } \mathcal{D}$ denote the space of Bridgeland stability conditions on \mathcal{D} [Bri07]. We use standard notation: we will write $(Z, \{\mathcal{P}[\theta]\}_{\theta \in \mathbb{R}})$ for the stability condition with central charge $Z : K(\mathcal{D}) \rightarrow \mathbb{C}$ and full additive subcategory of phase θ given by $\mathcal{P}[\theta] \subset \mathcal{D}$. Following [Tod08], let $\text{Stab}_n \mathcal{D}$ denote the subspace of normalised Bridgeland stability conditions, i.e. stability conditions whose central charge Z satisfies $Z([\mathcal{O}_x]) = -1$. $\text{Stab}_n \mathcal{D}$ has an open subset consisting of stability conditions with the standard heart $\mathcal{D} \cap \text{Coh } Y$ and central charges of the form

$$Z_{\beta+i\zeta}(E) := - \int e^{-(\beta+i\zeta)} \text{ch } E \quad \text{for } \beta + i\zeta \in A(\mathcal{D})_{\mathbb{C}}$$

where $A(\mathcal{D})_{\mathbb{C}}$ denotes the complexified ample cone. (See [Tod08, Lemma 4.1]; these correspond to the neighbourhood of the large volume limit.) Following [Tod08, Definition 4.2], let $\text{Stab}_n^\circ \mathcal{D}$ be the connected component of $\text{Stab}_n \mathcal{D}$ containing this open subset.

By [Tod09, Theorem 7.1], $\text{Stab } \mathcal{D}$ is connected. Moreover, we have a rescaling action of $t \in \mathbb{C}$ on $\text{Stab } \mathcal{D}$:

$$(Z, \{\mathcal{P}[\phi]\}_{\phi \in \mathbb{R}}) \mapsto (e^{-i\pi t} Z, \{\mathcal{P}[\phi + \text{Re } t]\}_{\phi \in \mathbb{R}})$$

Assembling all this together, we get a commutative diagram

$$\begin{array}{ccc} \mathbb{C} \times \text{Stab}_n^\circ \mathcal{D} & \xrightarrow{\cong} & \text{Stab } \mathcal{D} \\ \uparrow & & \uparrow \\ 2\mathbb{Z} \times \text{Stab}_n^\circ \mathcal{D} & \xrightarrow{\cong} & \text{Stab}_n \mathcal{D} \end{array}$$

where the top horizontal isomorphism is given by the rescaling action.

Corollary 4.7. *Let ϕ be an R -linear autoequivalence of $D(Y)$ which preserves $\mathcal{O}_x \in D(Y)$ for some $x \in Y \setminus C$ (with no shift). Then the action of ϕ on $\text{Stab } \mathcal{D}$ preserves $\text{Stab}_n^\circ \mathcal{D}$.*

Proof. This follows from the preceding discussion, on noting that the rescaling action of $\mathbb{Z} \subset \mathbb{C}$ agrees with the action of shifts. \square

Definition 4.8. Let $\text{Auteq } \mathcal{D}$ be the group of R -linear autoequivalences of $D(Y)$, and let $\text{Auteq}^\circ \mathcal{D}$ be the subgroup of R -linear autoequivalences preserving $\mathcal{O}_x \in D(Y)$ for some point $x \notin C$.

To justify the notation, we'll see shortly that we could have equally well defined these groups in terms of R -linear autoequivalences of \mathcal{D} .

Theorem 4.9. [Tod08, Tod09]

(i) *We have an isomorphism*

$$\text{Auteq } \mathcal{D} \cong (\mathbb{Z} * \mathbb{Z}) \times \mathbb{Z},$$

where the first factor has generators the spherical twist in $\mathcal{O}_C(-1)$ and $\otimes \mathcal{L}$, where \mathcal{L} generates $\text{Pic}(Y)$, and the final \mathbb{Z} factor is the grading shift.

(ii) We have $\text{Auteq}^\circ \mathcal{D} \cong \mathbb{Z} * \mathbb{Z}$, again with generators the spherical twist in $\mathcal{O}_C(-1)$ and $\otimes \mathcal{L}$. The action of $\text{Auteq}^\circ \mathcal{D}$ on $K(\mathcal{D})$ induces a homomorphism to the stabiliser of $[\mathcal{O}_x]$ in $GL_2(\mathbb{Z}) = \text{Aut } K(\mathcal{D})$, i.e. the infinite dihedral group $\mathbb{Z} \rtimes \mathbb{Z}/2$. Let $\text{Auteq}_{\text{Tor}}^\circ \mathcal{D}$ be its kernel. Then the image of this homomorphism is \mathbb{Z} , and we then obtain a diagram with exact rows

$$(4.1) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \text{Auteq}_{\text{Tor}}^\circ \mathcal{D} & \longrightarrow & \text{Auteq}^\circ \mathcal{D} & \longrightarrow & \mathbb{Z} \longrightarrow 0 \\ & & \uparrow \simeq & & \uparrow \simeq & & \parallel \\ 0 & \longrightarrow & \mathbb{Z}^{*\infty} & \longrightarrow & \mathbb{Z} * \mathbb{Z} & \longrightarrow & \mathbb{Z} \longrightarrow 0 \end{array}$$

The vertical isomorphism maps the standard generators of $\mathbb{Z}^{*\infty}$ to $T_{\mathcal{O}_C(i)}$, $i \in \mathbb{Z}$; and the standard generators of $\mathbb{Z} * \mathbb{Z}$ to $T_{\mathcal{O}_C(-1)}$ and $\otimes \mathcal{L}$.

Proof. For (i), this is [Tod09, Theorem 7.7]. For (ii), the description of $\text{Auteq}^\circ \mathcal{D}$ is clear. To conclude, consider \mathcal{L} a line bundle generating $\text{Pic}(Y) \cong \mathbb{Z}$. Up to possibly replacing \mathcal{L} with its dual line bundle, its action by tensor on $K(\mathcal{D})$, with respect to the basis $\{[\mathcal{O}_C] - [\mathcal{O}_C(-1)], [\mathcal{O}_C]\}$, is given by the matrix $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. As $\otimes \mathcal{L}^i \circ T_{\mathcal{O}_C} \circ \otimes \mathcal{L}^{-i} = T_{\mathcal{O}_C(i)}$, the rest of the claim is then immediate. \square

The proof shows that the \mathbb{Z} quotient on the top line of equation 4.1 is naturally identified with $\text{Pic}(Y)$.

Remark 4.10. For instance comparing with [HWb, Theorem 1.5], we see that $\text{Auteq}_{\text{Tor}}^\circ \mathcal{D}$ could equivalently have been defined as R -linear autoequivalences of the subcategory \mathcal{D} preserving $\text{Stab}_n^\circ \mathcal{D}$.

We'll use the following immediate corollary.

Corollary 4.11. *Suppose that $\phi \in \text{Auteq } \mathcal{D}$ induces the identity on $K(\mathcal{D})$. Then ϕ must lie in the subgroup $\mathbb{Z}^{*\infty} \times 2\mathbb{Z}$, where $\mathbb{Z}^{*\infty} \subset \mathbb{Z} * \mathbb{Z}$ is the subgroup generated by spherical twists in the $\mathcal{O}_C(i)$; and $2\mathbb{Z}$ denotes even shifts.*

5. PROOF OF THEOREM 1.1

For an A_∞ -category \mathcal{C} we write $D(\mathcal{C}) := H^0(\text{Tw}(\mathcal{C}))$ for the cohomological category of the category of twisted complexes; this is a triangulated category in the classical sense. We remark that the group $\text{Auteq } \mathcal{C}$ of A_∞ -autoequivalences of \mathcal{C} admits a natural homomorphism to the group $\text{Auteq}(D(\mathcal{C}))$ of triangulated equivalences of $D(\mathcal{C})$.

The primary mirror symmetric input to the proof of Theorem 1.1 is the following:

Theorem 5.1. [CPU16, Theorem 4.2] *There is a quasi-equivalence of \mathbb{C} -linear triangulated categories*

$$(5.1) \quad \Upsilon : DW(X) \xrightarrow{\sim} D(Y)$$

with $\Upsilon(S_i) = \mathcal{O}_C(-i)$.

It follows that $D(\mathcal{W})$ is in fact split-closed. By construction, the mirror equivalence identifies the subcategory $\langle S_0, S_1 \rangle$ of $DW(X)$ generated by S_0 and S_1 with \mathcal{D} ; passing to K -theory,

the numerical Grothendieck group $K_{\text{num}}\mathcal{F}(X)$ is identified with $K(\mathcal{D})$. Moreover, [CPU16] explicitly identify the objects mirror to \mathcal{O}_Y and $\mathcal{O}_Y(1)$: these are Lagrangian \mathbb{R}^3 s, say L_0 and L_1 , fibred over the base of the Morse-Bott-Lefschetz fibration, see Figure 2, cf. [CPU16, Figure 1.2 and Theorem 1.2].

Corollary 5.2. *The equivalence Υ of (5.1) entwines the $SH^0(X)$ -linear structure of $DW(X)$ and the R -linear structure of $D(Y)$.*

Proof. The existence of an isomorphism of \mathbb{C} -algebras $SH^0(X) \cong \Gamma(\mathcal{O}_Y) = R$ is proved in [CPU16, Appendix]. Both linear structures can be viewed as the general fact that a k -linear dg -category or A_∞ -category \mathcal{C} is also linear over the Hochschild cohomology $HH^0(\mathcal{C})$. At the cohomological level (which is all that we require) this is classical, and a chain-level version is established in [Pom, Section 5.1]. By [GPS20], the wrapped category $\mathcal{W}(X)$ is generated by L_0 and L_1 and is homologically smooth and non-degenerate. This means in particular that the open-closed map $HH_*(\mathcal{W}(X)) \rightarrow SH^*(X)$ is an isomorphism. Since $\mathcal{W}(X)$ is also a Calabi-Yau category [Gan12], the closed-open map $SH^*(X) \rightarrow HH^*(\mathcal{W}(X))$ is also an isomorphism. The action of Υ on HH^* therefore induces a distinguished isomorphism $SH^0(X) \cong R$. The result follows. \square

Recall the exact, Maslov zero torus T introduced in Section 3.2, which was graded in Section 3.3.

Lemma 5.3. *Let $(T, \zeta) \in \text{Ob } \mathcal{W}(X)$ be any Lagrangian brane associated to T . Then under the mirror symmetry isomorphisms above, (T, ζ) is mirror to $\mathcal{O}_x[k]$ for some $x \in Y \setminus C$ and shift $k \in \mathbb{Z}$.*

Proof. Let \mathcal{E} be the mirror to (T, ζ) . By inspection, we can find Hamiltonian isotopies so that the image of T is disjoint from the sphere S_0 , respectively S_1 ; and the Floer theory of T with each of L_0 and L_1 has a single generator, see Figure 2. By considering Floer cohomology with S_0 and S_1 , \mathcal{E} is orthogonal to \mathcal{O}_C and $\mathcal{O}_C(-1)$, and hence the whole of \mathcal{D} . Since \mathcal{E} is orthogonal to \mathcal{O}_p for any $p \in C$, \mathcal{E} has support on $Y \setminus C$.

[CPU16, Theorem 5.3] shows that $\mathcal{O}_Y \oplus \mathcal{O}_Y(1)$ is a tilting object in $D(Y)$. It is shown in the proof of [BCZ17, Theorem 7.2.1] that whenever a smooth variety $Z \rightarrow A$ is projective over A , where A is affine, and \mathcal{E}_Z is a tilting object for Z , and $F \in D(Z)$ has non-proper support, then $\text{Ext}^\ell(\mathcal{E}_Z, F)$ has infinite rank for some ℓ . This implies that in our case \mathcal{E} has proper support.

Finally, the only proper subvarieties of $Y \setminus C$ are finite unions of points. Since (T, ζ) has the self-Floer cohomology of a three-torus, the mirror \mathcal{E} is simple and has no negative self-Exts. By [Huy06, Lemma 4.5], $\mathcal{E} = \mathcal{O}_x[k]$, some $k \in \mathbb{Z}$. \square

Remark 5.4. Varying the local system ζ on T , one expects the family of branes (T, ζ) in X to correspond to a cluster chart $(\mathbb{C}^*)^3$ on the mirror Y (suitably interpreted, to toric SYZ fibres). There are other exact Lagrangian tori in X with vanishing Maslov class, for instance fibred over an S^1 in the base \mathbb{C}^* of the Morse-Bott-Lefschetz fibration which encloses one critical fibre and the puncture. Appropriate such tori do have non-trivial Floer theory with S_0 or S_1 , for certain choices of local systems. In that sense Lemma 5.3 amounts to saying that our choice of T is picking the ‘‘correct’’ cluster chart. (Compare to the closely related two-dimensional situation studied in [Aur07].)

Lemma 4.5 then implies that for any $\phi \in \pi_0 \text{Symp}^{\text{gr}} X$, $\phi(T, \zeta)$ is equal to (T, ζ) up to a shift in $DW(X)$.

Lemma 5.5. *For any $\phi \in \pi_0 \text{Symp}_c X$, equipped with the preferred grading, $\phi(T, \zeta)$ is equal to (T, ζ) in $DW(X)$.*

Proof. Recall from Lemma 3.8 that any compactly supported symplectomorphism of X is gradeable and canonically graded. Such a map therefore acts on the set of graded Lagrangian submanifolds in X . The action of a symplectomorphism on the set of objects $\text{Ob } \mathcal{W}(X)$ of the wrapped category is compatible with the action on graded Lagrangians.

Let α be the angle coordinate for $z \in \mathbb{C}^*$. For all $t \in \mathbb{R}_{\geq 0}$, let σ_t be the time t flow of the symplectic vector field Z_α which is ω -dual to the closed one-form $d\alpha$. (Our convention is that this is inward-pointing with respect to the base \mathbb{C}^* .) We consider the path $\{\sigma_t \circ \phi \circ \sigma_t^{-1}\}_{t \in \mathbb{R}}$ of graded symplectomorphisms. For sufficiently large t , the map $\phi' := \sigma_t \circ \phi \circ \sigma_t^{-1}$ has support in some compact set V disjoint from T ; moreover, we can assume that $X \setminus V$ is path connected (and so contains both T and a conical end of X). As we are using the canonical grading for the map ϕ' , we see that it must therefore fix the grading on (T, ζ) . On the other hand, recall that we already know that ϕ fixes (T, ζ) up to shift. The action of compactly supported symplectomorphisms on the set of graded Lagrangians only depends on the path-component in the group $\text{Symp}_c(X)$ with its C^∞ -topology. Since ϕ lies in the same component of the space of graded compactly supported symplectomorphisms as ϕ' , it must also fix the grading of (T, ζ) . \square

Definition 5.6. Define $\text{Symp}^\circ(X) \subset \text{Symp}^{\text{gr}}(X)$ to be the subgroup of graded symplectomorphisms which take (T, ζ) to itself in $DW(X)$.

Lemma 5.7. *Consider the map $\text{PBr}_3 \rightarrow \pi_0 \text{Symp}^{\text{gr}} X$ defined earlier. Then the intersection of the image with $\pi_0 \text{Symp}^\circ X$ is precisely the subgroup generated by τ_{S_0} and λ , i.e. the image of the subgroup $\mathbb{Z} * \mathbb{Z}$ generated by R_2 and R_3 .*

Proof. By construction, the image of R_3 is τ_{S_0} , with its preferred grading as a compactly supported map; and the image of R_2 is λ , a non-compactly supported map which maps S_i to S_{i+1} for all $i \in \mathbb{Z}$. Now λ induces an R -linear autoequivalence of $\mathcal{W}(X)$, and the mirror autoequivalence in $\text{Auteq } \mathcal{D}$ must take $\mathcal{O}_C(-i)$ to $\mathcal{O}_C(-i-1)$. By Theorem 4.9, the only possibility is for this map to be $\otimes \mathcal{L}$, which fixes \mathcal{O}_x . By Lemma 5.3, we then have $\lambda \in \pi_0 \text{Symp}^\circ X$.

Finally, recall that we choose gradings on the image of PBr_3 so that the generator of the center, i.e. $R_1 R_2 R_3$, maps to a shift by one. The claim then follows. \square

Any element of $\pi_0 \text{Symp}^{\text{gr}}(X)$ induces an $SH^0(X)$ -linear autoequivalence of $\mathcal{W}(X)$. Lemma 4.6, together with the HMS equivalence $\text{Auteq}_{SH^0}(\mathcal{W}(X)) \simeq \text{Auteq}_R(D(Y)) = \text{Auteq } \mathcal{D}$, then implies that the image of $\pi_0 \text{Symp}^{\text{gr}}(X)$ in $\text{Auteq}_{SH^0}(\mathcal{W}(X))$ preserves the subcategory $\langle S_0, S_1 \rangle$ (equivalent under mirror symmetry to \mathcal{D}). Moreover, from our definitions, any element of $\pi_0 \text{Symp}^\circ(X)$ lands not only in $\text{Auteq}(\mathcal{D})$ but in fact in the subgroup $\text{Auteq}^\circ(\mathcal{D})$. Altogether, this gives a diagram:

$$(5.2) \quad \begin{array}{ccc} \pi_0 \mathrm{Symp}_c(X) & \longrightarrow & \pi_0 \mathrm{Symp}^\circ(X) \\ \downarrow & & \downarrow \\ \mathrm{Auteq}_{\mathrm{Tor}}(\langle S_0, S_1 \rangle) & \longrightarrow & \mathrm{Auteq}(\langle S_0, S_1 \rangle) \\ \simeq \downarrow & & \simeq \downarrow \\ \mathrm{Auteq}_{\mathrm{Tor}}^\circ(\mathcal{D}) & \longrightarrow & \mathrm{Auteq}^\circ(\mathcal{D}) \end{array}$$

with the bottom vertical arrows the HMS equivalences.

Recall from Lemma 3.16 that $\pi_0 \mathrm{Symp}^{\mathrm{gr}}(X)$, and in particular $\pi_0 \mathrm{Symp}^\circ(X)$, acts on the numerical K -theory $K_{\mathrm{num}}(\mathcal{F}(X))$.

Passing to the mirror side, consider the image of $\pi_0 \mathrm{Symp}^{\mathrm{gr}}(X)$ in $\mathrm{Auteq}^\circ \mathcal{D}$. As the image of $\mathrm{Auteq}^\circ \mathcal{D}$ in $GL(K(\mathcal{D}))$ lies in a \mathbb{Z} subgroup, it follows that the image of $\pi_0 \mathrm{Symp}^{\mathrm{gr}}(X)$ in $GL(K_{\mathrm{num}} \mathcal{F}(X))$ lies in the mirror \mathbb{Z} subgroup. We are now ready to state a categorical version of our main theorem.

Theorem 5.8. (i) *For graded symplectomorphisms, we have*

$$(5.3) \quad \mathrm{PBr}_3 \hookrightarrow \pi_0 \mathrm{Symp}^{\mathrm{gr}}(X) \twoheadrightarrow \mathrm{Auteq}_{\mathrm{SH}^0} DW(X) \cong \mathrm{Auteq} \mathcal{D} \cong \mathrm{PBr}_3,$$

where under the decomposition $\mathrm{PBr}_3 \cong (\mathbb{Z} * \mathbb{Z}) \times \mathbb{Z}$ with generators R_2, R_3 and $R_1 R_2 R_3$, R_2 maps to λ , R_3 maps to τ_{S_0} , and $R_1 R_3 R_3$ maps to the shift. Moreover, these group homomorphisms compose to give the identity.

(ii) *Exact symplectomorphisms which preserve $(T, \zeta) \in \mathrm{Ob} DW(X)$ fit into the following commutative diagram:*

$$(5.4) \quad \begin{array}{ccccc} \mathbb{Z}^{*\infty} & \longrightarrow & \mathbb{Z} * \mathbb{Z} & \longrightarrow & \mathbb{Z} \\ \downarrow & & \downarrow & & \parallel \\ \pi_0 \mathrm{Symp}_c(X) & \longrightarrow & \pi_0 \mathrm{Symp}^\circ(X) & \longrightarrow & \mathbb{Z} \\ \downarrow & & \downarrow & & \parallel \\ \mathrm{Auteq}_{\mathrm{Tor}}^\circ \mathcal{D} & \longrightarrow & \mathrm{Auteq}^\circ \mathcal{D} & \longrightarrow & \mathbb{Z} \\ \downarrow \simeq & & \downarrow \simeq & & \parallel \\ \mathbb{Z}^{*\infty} & \longrightarrow & \mathbb{Z} * \mathbb{Z} & \longrightarrow & \mathbb{Z} \end{array}$$

where the vertical compositions from the top to bottom lines are the identity. The map $\pi_0 \mathrm{Symp}^\circ(X) \rightarrow \mathbb{Z}$ is given by taking the action on $K_{\mathrm{num}}(\mathcal{F}(X))$, itself naturally identified with $K(\mathcal{D})$, and noting that the image lies in the \mathbb{Z} subgroup.

Proof. Part (i) is immediate from Theorem 4.9 and the preceding construction and discussion of the pure braid group action $\mathrm{PBr}_3 \rightarrow \pi_0 \mathrm{Symp}^{\mathrm{gr}} X$.

For part (ii), the map from the first to the second line is defined by combining the Section 3 constructions with Lemma 5.7. Lemma 3.17 implies that the image of $\pi_0 \mathrm{Symp}_c(X)$ lies in the subgroup $\mathrm{Auteq}_{\mathrm{Tor}}^\circ \mathcal{D}$ of $\mathrm{Auteq}^\circ \mathcal{D}$, which allows us to go from the second line to the third. The isomorphism of the final two rows is given by Theorem 4.9.

To check that composition from the first to the final row is an isomorphism, we proceed as follows. For the first column, this follows from having identified the twists τ_{S_i} and $T_{\mathcal{O}_C(-i)}$, $i \in \mathbb{Z}$ in the homological mirror symmetry isomorphism of Theorem 5.1.

For the second column, recall we have a preferred non-compactly supported symplectomorphism λ , together with a grading choice such that it lives in $\pi_0 \text{Symp}^\circ(X)$. With that choice of grading, λ takes S_i to S_{i+1} , for all $i \in \mathbb{Z}$ (with no shifts). At the level of K -theory, this is the same action as the tensor with our fixed generator \mathcal{L} for $\text{Pic}(Y)$. This implies that $\pi_0 \text{Symp}^\circ(X)$ surjects onto \mathbb{Z} . Surjectivity on the second column then follows from the exactness of the sequence

$$\text{Auteq}_{\text{Tor}}^\circ \mathcal{D} \rightarrow \text{Auteq}^\circ \mathcal{D} \rightarrow \mathbb{Z}.$$

Given that the composition from the first to the final row is an isomorphism, injectivity from the first to the second row, and surjectivity from the second to the third row, are then automatic. \square

Theorem 1.1 immediately follows.

6. CLASSIFICATION OF SPHERICALS

Let \mathcal{S} be the collection of Lagrangian spheres which we know of in X : the images of the S_i , $i \in \mathbb{Z}$, under the action of the group $\mathbb{Z}^{*\infty}$ generated by the Dehn twists in the S_i . In other words, these are all the spheres associated to matching paths between the two critical values in the base of our Morse-Bott-Lefschetz fibration, with all possible choices of gradings.

Definition 6.1. [Sei12, Section (3a)] We say an object S of $DW(X)$ is spherical if $\text{HF}^*(S, S) \cong H^*(S^3; \mathbb{Z})$, and for any object Z of $DW(X)$, $\text{HF}^*(S, Z)$ and $\text{HF}^*(Z, S)$ are of finite total rank and the composition

$$\text{HF}^{3-*}(Z, S) \otimes \text{HF}^*(S, Z) \rightarrow \text{HF}^3(S, S) \cong \mathbb{Z}$$

is a non-degenerate pairing.

Theorem 6.2. *Let S be a spherical object in $DW(X)$. Then S is quasi-isomorphic to an element of \mathcal{S} (as a Lagrangian brane, i.e. together with a choice of grading).*

We will break the proof into a number of stages.

6.1. Twists act trivially on numerical K -theory. Fix a spherical object $S \in DW(X)$ and consider the associated spherical twist T_S . By [Pom, Lemma 5.46] we can assume that the category $DW(X)$ of perfect modules over $\mathcal{W}(X)$ has a strictly $SH^0(X)$ -linear chain model. The total evaluation morphism $\text{Hom}(S, \bullet) \otimes S \rightarrow \bullet$ is then SH^0 -linear, and the cone inherits an $SH^0(X)$ -linear structure. This means that T_S is linear as an autoequivalence. By Theorem 5.8, we know that T_S is an element of $\text{Auteq}_{SH^0} DW(X) \cong \text{Auteq} \mathcal{D} \cong \text{PBr}_3$.

Lemma 6.3. *T_S acts trivially on $K(\mathcal{D}) \cong K_{\text{num}}(\mathcal{F}(X))$.*

Proof. Under the equivalence $\Upsilon : DW(X) \rightarrow D(Y)$ of Theorem 5.1, S corresponds to some complex \mathcal{E}_S of sheaves on Y . As in the proof of Lemma 5.3, the argument in the proof of [BCZ17, Theorem 7.2.1] shows that \mathcal{E}_S must have proper support in Y : otherwise it would have infinite rank morphisms in some degree with the tilting object in $D(Y)$, which violates the definition of being spherical. As \mathcal{E}_S is also indecomposable, it must have connected support, which by the classification of proper subvarieties of Y implies that the support is either C or

a single point. By [Huy06, Lemma 4.5], the support must be C , and so in fact \mathcal{E}_S belongs to the category $\mathcal{D} \subset D(Y)$. It follows that S belongs to the subcategory of $\mathcal{W}(X)$ generated by S_0 and S_1 and in particular belongs to the compact Fukaya category $\mathcal{F}(X) \subset \mathcal{W}(X)$.

The spherical twist T_S acting on $\mathcal{F}(X)$ fits into an exact triangle

$$(6.1) \quad \text{Hom}_{\mathcal{F}}(S, \bullet) \otimes S \longrightarrow \text{id} \longrightarrow T_S$$

and hence acts on (numerical) K -theory by the Picard-Lefschetz type transformation

$$A \mapsto A - \chi(S, A)[S]$$

where $\chi(S, A) \in \mathbb{Z}$ is the Euler characteristic of $\text{Hom}_{\mathcal{F}}(S, A)$. Since the intersection pairing on $K_{num}(\mathcal{F}(X)) = H_3(X)$ vanishes, χ vanishes identically. \square

Lemma 6.3 combined with Corollary 4.11 then implies that T_S lies in $\mathbb{Z}^{*\infty} \times 2\mathbb{Z} \subset (\mathbb{Z} * \mathbb{Z}) \times \mathbb{Z} \cong \text{PBr}_3$.

6.2. A localisation argument. The key input for Theorem 6.2 is the following analogue, in our setting, of a well-known theorem [KS02] due to Khovanov and Seidel, which localises the computation of Floer cohomology to the fixed locus of a finite group action.

For matching paths γ_a, γ_b in the base \mathbb{C}^* of the Morse-Bott-Lefschetz fibration $\pi : X \rightarrow \mathbb{C}^*$, we let $I(\gamma_a, \gamma_b)$ denote the minimal (unsigned, i.e. geometric) interior intersection number between γ_a and γ_b , relative to their end points. Here isotopies of paths are not allowed to cross the puncture; and ‘interior’ means that we’re not including the end points in our count.

Proposition 6.4. *Suppose $S_a, S_b \in \mathcal{S}$ correspond to matching paths $\gamma_a, \gamma_b \subset \mathbb{C}^*$. Then the Floer homology of S_a and S_b with coefficients in $\mathbb{Z}/2$ satisfies*

$$\dim \text{HF}(S_a, S_b; \mathbb{Z}/2) = 4 + 4I(\gamma_a, \gamma_b)$$

The proof of Proposition 6.4 has two ingredients: the existence of a $\mathbb{Z}/2 \times \mathbb{Z}/2$ action on X , preserving the Lagrangians, and with the property that there are no holomorphic discs in the fixed point locus; and an equivariant transversality argument which allows us to compare Floer theory in X with Floer theory in that fixed point locus. We describe the involutions and the reduction to existence of an equivariantly transversal J in this section; the existence of such a J will be broken down into Sections 6.3 through 6.5.

Recall X is given explicitly by

$$X = \{u_1 v_1 = z - 1, u_2 v_2 = z + 1\} \subset \mathbb{C}^4 \times \mathbb{C}^*.$$

X carries two commuting involutions: ι_1 , trading u_1 and v_1 ; and ι_2 , trading u_2 and v_2 . Let $G = \mathbb{Z}/2 \times \mathbb{Z}/2$ be the group generated by the ι_j .

The common fixed point locus of both ι_1 and ι_2 , say X^{inv} , has equation $\{p^2 = z - 1, q^2 = z + 1\} \subset \mathbb{C}^2 \times \mathbb{C}^*$. This is smooth, and Weinstein with the restriction of the form from X . (Topologically, X^{inv} is a sphere with six punctures.)

Projection $X^{\text{inv}} \rightarrow \mathbb{C}^*$ to the z co-ordinate, i.e. to the base of the Morse-Bott-Lefschetz fibration, defines a four-fold cover of \mathbb{C}^* , branched over ± 1 . Note that the branching pattern is as follows: say the sheets are labelled by $(\pm p, \pm q)$. Then above $+1$, the sheets with the same value of q meet; and above -1 , the sheets with the same value of p meet.

For any matching path γ , S_γ meets X^{inv} cleanly in an embedded S^1 , say s_γ (which is itself a matching sphere in X^{inv}). The circle s_γ is a 4-fold cover of γ under projection to z .

Lemma 6.5. *After a Hamiltonian isotopy of X , the matching spheres S_a and S_b meet exactly in the finite set of transverse intersection points of the curves s_a and s_b . In particular, all intersection points lie in the common fixed locus of the involutions ι_1 and ι_2 .*

Proof. Assume that two matching paths γ_a and γ_b intersect transversally in k points apart from the end points. Then by construction, the S_a and S_b intersect cleanly in two S^1 s (above each end point) and k copies of T^2 (above each interior intersection point); and their restrictions to X^{inv} , i.e. the matching spheres s_a and s_b in X^{inv} , intersect transversally in $4 + 4k$ points (with the end points contributing two each, and each ‘interior’ intersection point contributing 4).

There is a symplectic neighbourhood theorem for cleanly intersecting Lagrangian submanifolds, saying that near a component B of the clean intersection locus they are modelled by the conormal bundle of B inside the cotangent bundle of either one. Starting from this, and following the proof of [KS02, Proposition 5.15] (which treats the case of a single involution), there is an (ι_1, ι_2) -equivariant small Hamiltonian perturbation of S_a , fixing the invariant locus, so that S_a and S_b intersect transversally in exactly the intersection points of $s_a \pitchfork s_b$. \square

Any isotopy of γ in the base (rel endpoints) is covered by a Hamiltonian isotopy of S_γ in X and a Hamiltonian isotopy of s_γ in X^{inv} . Assume that after isotopy rel end points, the matching paths γ_a and γ_b intersect minimally rel endpoints, in $I(\gamma_a, \gamma_b)$ points. Then after Hamiltonian isotopy in X , we may assume S_a and S_b intersect transversally in $4 + 4I(\gamma_a, \gamma_b)$ points, all fixed by both ι_j .

Lemma 6.6. *Suppose that there is a compatible $J = \{J_t\}_{t \in [0,1]}$ on X with the properties that*

- (1) *J is G -equivariant;*
- (2) *the moduli spaces of J -holomorphic curves $u : \mathbb{R} \times [0, 1] \rightarrow X$ with boundary conditions on S_a and S_b are regular.*

Then

$$\text{HF}(S_a, S_b; \mathbb{Z}/2) \cong \text{HF}(s_a, s_b; \mathbb{Z}/2)$$

Proof. We have arranged that all intersection points lie in the fixed locus, so the underlying Floer cochain groups agree.

Now suppose that $u : D \rightarrow X$ is a Floer holomorphic disc contributing to the differential and that $\text{Im}(u) \not\subseteq X^{\text{inv}}$. Then notice that each of $u, \iota_1 \circ u, \iota_2 \circ u$ and $\iota_1 \circ \iota_2 \circ u$ give holomorphic discs; moreover, these are either a pair of distinct holomorphic discs (e.g. if $\text{Im}(u) \subset \text{Fix}(\iota_1)$ but $\text{Im}(u) \not\subseteq \text{Fix}(\iota_2)$) or a quadruple (if $\text{Im}(u) \not\subseteq \text{Fix}(\iota_1)$ and $\text{Im}(u) \not\subseteq \text{Fix}(\iota_2)$). Equivariant transversality means that all these discs are regular, so as we are working with $\mathbb{Z}/2$ coefficients, the contribution to the differential is zero.

It follows that the differential on $\text{CF}(S_a, S_b; \mathbb{Z}/2)$ co-incides with that on $\text{CF}(s_a, s_b; \mathbb{Z}/2)$. \square

For computing Floer cohomology of curves on a surface, one can achieve regularity with a time-independent almost complex structure. The computation of $\text{HF}(s_a, s_b; \mathbb{Z}/2)$ is then completed by:

Lemma 6.7. *For any compatible almost complex structure J which makes projection to the base \mathbb{C}^* holomorphic, there are no non-constant holomorphic discs with boundary on s_a and s_b contained wholly in the fixed locus X^{inv} .*

Proof. Suppose we had such a disc $u : D \rightarrow X^{\text{inv}}$; it projects to a disc $\pi \circ u$ in the base (this will in general be immersed). By minimality of intersection, the image of $\pi \circ u$ must contain at least one of the critical values. Consider the auxiliary Riemann surface given by pulling back the cover $X^{\text{inv}} \rightarrow \mathbb{C}^*$ under $\pi \circ u$, say $\Sigma_u \rightarrow D$. This is a four-fold cover with two possible types of critical values / branch points, modelled on the branching patterns for $\pm 1 \in \mathbb{C}^*$. Now notice that for any positive number of such branch points, $\Sigma_u \rightarrow D$ cannot have a section for topological reasons. Thus there are no such holomorphic discs $u \rightarrow X^{\text{inv}}$. \square

Lemma 6.7 immediately implies that for our choices,

$$\text{HF}(s_a, s_b; \mathbb{Z}/2) \cong \text{CF}(s_a, s_b; \mathbb{Z}/2)$$

which has rank $4 + 4I(\gamma_a, \gamma_b)$. Subject to achieving the hypotheses of Lemma 6.6, this completes the proof of Proposition 6.4.

6.3. Equivariant transversality. Let (M, ω) be an exact symplectic manifold with a symplectic involution $\iota : M \rightarrow M$ which setwise preserves a pair of exact Lagrangian submanifolds $L_0, L_1 \subset M$. We write M^ι for the fixed locus, and L_j^ι for the fixed set of $\iota|_{L_j}$, which is Lagrangian in M^ι . We suppose that the L_i intersect transversally, which implies that the L_i^ι intersect transversally in the fixed point locus. Let $n = \dim_{\mathbb{C}}(M)$, $n_\iota = \dim_{\mathbb{C}}(M^\iota)$ and let $n_{\text{anti}} = n - n_\iota$ be the complex codimension of the fixed locus. The normal bundle $\nu_{M^\iota} \rightarrow M^\iota$ is a complex bundle of rank n_{anti} ; this is the (-1) -eigenspace of the action of ι on $T_p M$ for $p \in M^\iota$.

We consider ι -equivariant paths of compatible almost complex structures $J = \{J_t\}_{t \in [0,1]}$ on M . Khovanov and Seidel [KS02, Proposition 5.13] explain how to choose J_t so as to achieve regularity for those curves contributing to $\text{CF}(L_0, L_1)$ which are not contained wholly in the fixed locus M^ι of the involution.

Unfortunately, their work does not immediately apply to our setting: whilst Lemma 6.7 implies that, for suitable data, we have no Floer solutions lying in $X^{\text{inv}} \subset X$ for topological reasons, there may be curves which lie in the fixed loci X^{ι_j} of either one of the two involutions. In general the existence of such curves can obstruct the existence of any ι_j -equivariant regular J , and hence *a fortiori* of a G -equivariant regular J . More precisely, if x_+, x_- are ι_j -invariant intersections of S_a and S_b for which the difference in their Maslov indices, computed in the fixed locus X^{ι_j} , is greater than the difference of the indices computed in the total space, it is impossible to achieve equivariant transversality (because the dimension of the space of solutions in the fixed set is greater than the dimension in the total space).

A sufficient criterion for vanishing of this obstruction was given in [SS10, Section 3.5].

Definition 6.8. A *stable normal trivialization* consists of two pieces of data:

- (1) a stable trivialization $\phi : \nu_{M^\iota} \times \mathbb{C}^r \rightarrow M^\iota \times \mathbb{C}^{n_{\text{anti}}+r}$ of the normal bundle to the fixed point set, which is unitary (respects both the complex and symplectic structures);
- (2) homotopies $h_k : [0; 1] \times L_k^{\text{inv}} \rightarrow \text{Gr}_{\text{Lag}}(n_{\text{anti}} + r)$, for $k = 0, 1$, such that (i) $h_0(0, \cdot) = \phi_*(\nu_{L_0^{\text{inv}}} \times \mathbb{R}^r)$ and $h_0(1, \cdot) = \mathbb{R}^{n_{\text{anti}}+r}$, (ii) $h_1(0, \cdot) = \phi_*(\nu_{L_1^{\text{inv}}} \times i\mathbb{R}^r)$ and $h_1(1, \cdot) = i\mathbb{R}^{n_{\text{anti}}+r}$.

Let $\hat{M} = M \times \mathbb{C}^r$ with $\hat{\iota} = \iota \times (-1)$, then $\hat{M}^\iota = M^\iota$, and if we set $\hat{L}_0 = L_0 \times \mathbb{R}^r$ and $\hat{L}_1 = L_1 \times i\mathbb{R}^r$, then $\hat{L}_i^\iota = L_i^\iota$. We fix the convention that the linear Lagrangian subspaces \mathbb{R} and $i\mathbb{R}$ in \mathbb{C} are graded so the rank one Floer cohomology $\text{HF}(\mathbb{R}, i\mathbb{R}) \cong \mathbb{Z}$ is concentrated in degree zero. There is a canonical isomorphism $\text{CF}(L_0, L_1) \cong \text{CF}(L_0 \times \mathbb{R}^r, L_1 \times i\mathbb{R}^r)$ of Floer

complexes, where the first complex lives inside M and the second inside \hat{M} equipped with a product almost complex structure.

Lemma 6.9. *Suppose (M, L_0, L_1) admit a stable normal trivialization $\Upsilon = (\phi, h_0, h_1)$. There is a Hamiltonian symplectomorphism $\Phi = \Phi_\Upsilon : \hat{M} \rightarrow \hat{M}$ satisfying (i) $\Phi|_{M^\iota} = id$; (ii) $\nu_{\Phi(\hat{L}_0)} = \mathbb{R}^{n_{anti}+r}$; (iii) $\nu_{\Phi(\hat{L}_1)} = i\mathbb{R}^{n_{anti}+r}$.*

Proof. This is proved in the discussion after [SS10, Definition 19]. \square

We have the Floer equation

$$(6.2) \quad \begin{aligned} \partial_s u + J_t(u)(\partial_t u - X_t(u)) &= 0, \\ u(s, 0) &\in L_0, \quad u(s, 1) \in L_1. \end{aligned}$$

where X_t denotes the time-dependent Hamiltonian flow of H_t . The next definition 6.10 and subsequent Lemma 6.11 are taken from [Sei08b, Section 14c].

Definition 6.10. Suppose (M, L_0, L_1) have stably trivial normal structure, and fix a stable normal trivialisation Υ and Hamiltonian symplectomorphism $\Phi = \Phi_\Upsilon$ as above. We define Υ -constrained Floer perturbation data for $(\Phi(\hat{L}_0), \Phi(\hat{L}_1))$ to comprise $(\{\hat{H}_t\}, \{\hat{J}_t\})_{0 \leq t \leq 1}$, where

- (1) \hat{H}_t, \hat{J}_t are \hat{i} -invariant;
- (2) for some extension of the unitary trivialisation ϕ to a map $\bar{\phi} : \mathbb{C}^r \times U(M^\iota) \rightarrow \mathbb{C}^{n_{anti}+r}$, the derivatives $\bar{\partial}_{\hat{J}_t} \bar{\phi}$ vanish to first order along M^ι ;
- (3) for some $h \in (0, \pi/2)$, the flow X_t of \hat{H}_t acts by rotation by e^{iht} on $\nu_{L_0^\iota}$.

Lemma 6.11. *Υ -constrained Floer perturbation data are generically regular for all solutions $u : \mathbb{R} \times [0, 1] \rightarrow M^\iota$ to the equations (6.2) with image inside M^ι .*

Proof. We briefly review the discussion from [Sei08b, Section 14c]. Suppose u satisfies (6.2) for Υ -constrained perturbation data. Using the first order information on \hat{J}_t and \hat{H}_t along M^ι , [Sei08b, Lemma 14.6] shows that, globally over $\mathbb{R} \times [0, 1]$, the anti-invariant part D_{u, \hat{J}_t}^{anti} of the linearisation can be written explicitly as an operator $\partial_s + i\partial_t + h$. By a direct consideration of Fourier coefficients, [Sei08b, Lemma 11.5] shows that this is injective. It follows that invariant curves u are regular in the total space if and only if they are regular in the fixed point set. To prove Lemma 6.11 it then suffices to show that the class of Υ -constrained data is sufficiently large. The \hat{H}_t may be prescribed arbitrarily both on M^ι and outside a small open neighbourhood of M^ι , provided that each is \hat{i} -invariant. In a unitary splitting of $T\hat{M}|_{M^\iota}$, the off-diagonal parts of the almost complex structures \hat{J}_t vanish and the assumptions constrain the component tangential to M^ι , but the normal part may also be prescribed arbitrarily. This gives sufficient flexibility to appeal to standard transversality arguments to see that regular data are generic in this class. \square

Corollary 6.12. *Suppose that (M, L_0, L_1) have stably trivial normal structure. A choice of stable normal trivialisation Υ defines a (non-empty, open) set of \hat{i} -invariant time-dependent almost complex structures on \hat{M} which are simultaneously regular for all moduli spaces of solutions to (6.2) in both \hat{M} and $\hat{M}^\iota = M^\iota$.*

Proof. For generic invariant time-dependent J , regularity holds at all curves $u : \mathbb{R} \times [0, 1] \rightarrow \hat{M}$ which are not entirely contained in M^ι (this is a direct adaptation of the usual regularity theory, and follows from [KS02, Proposition 5.13]). For curves u inside the fixed locus,

after stabilising and applying a Hamiltonian symplectomorphism, Lemma 6.9 reduces us to a product situation, where we may invoke Lemma 6.11 to find an open set of Υ -constrained perturbation data which are regular. \square

The upshot of the discussion at this point is that, if a stable normal trivialisation exists, we can replace (M, L_0, L_1, ι) by $(\hat{M}, \hat{L}_0, \hat{L}_1, \hat{\iota})$ in such a way that the Floer complexes $\text{CF}(\hat{L}_0, \hat{L}_1) = \text{CF}(L_0, L_1)$ are naturally identified (and similarly in the fixed point set M^ι , which never changes); and the complex $\text{CF}(\hat{L}_0, \hat{L}_1)$ admits a regular equivariant J .

6.4. Two involutions. Now suppose that M carries an action of $G = \mathbb{Z}/2 \times \mathbb{Z}/2$, generated by involutions ι_1 and ι_2 , which again preserve the Lagrangians L_j setwise. We assume that the fixed submanifolds M^{ι_1} and M^{ι_2} meet cleanly in the global fixed locus M^G .

Lemma 6.13. *Suppose that*

- (1) M admits an action of $\mathbb{Z}/2 \times \mathbb{Z}/2$ generated by ι_1 and ι_2 ;
- (2) the triple (M, L_0, L_1) admits a stable normal polarisation for ι_1 ;
- (3) the triple $(M^{\iota_1}, L_0^{\iota_1}, L_1^{\iota_1})$ admits a stable normal polarisation for $\iota_2|_{M^{\iota_1}}$;
- (4) the intersection points of L_0 and L_1 all lie in the common fixed locus M^G , so $L_0 \cap L_1 = L_0^G \cap L_1^G$.

Then there is a cochain homotopy equivalence $\text{CF}(L_0, L_1; \mathbb{Z}/2) \simeq \text{CF}(L_0^G, L_1^G; \mathbb{Z}/2)$.

Proof. We apply the involutions iteratively. Existence of a stable normal trivialisation for ι_1 gives an equivariant J which identifies the Floer complexes $\text{CF}(L_0, L_1; \mathbb{Z}/2) = \text{CF}(L_0^{\iota_1}, L_1^{\iota_1}; \mathbb{Z}/2)$, since all intersection points are fixed and all non-fixed discs come in pairs (cf. the proof of Lemma 6.6). Note that, even though this identification of complexes arises from stabilisation (and goes via the intermediate complex $\text{CF}(\hat{L}_0, \hat{L}_1)$), the fixed locus M^{ι_1} for the first involution is never changed. Since the involution $\iota_2|_{M^{\iota_1}}$ admits a stable normal trivialisation, we repeat the argument to identify $\text{CF}(L_0^{\iota_1}, L_1^{\iota_1}; \mathbb{Z}/2)$ with $\text{CF}(L_0^G, L_1^G; \mathbb{Z}/2)$. \square

Remark 6.14. More generally, there is a natural notion of compatibility of stable normal trivialisations for a pair of involutions. The normal bundle satisfies $\nu_{M^G} \cong \nu_{M^{\iota_1}} \oplus \nu_{M^{\iota_2}}$, and ι_2 acts on the normal space $\nu_{M^{\iota_1}}$. This gives an obvious notion of a stable normal trivialisation for ι_1 being ι_2 -equivariant. Since the ordering of the generators ι_j for G was arbitrary, we will say that (M, L_0, L_1) admits *compatible* stable normal trivialisations. By an equivariant version of Moser's theorem, the Hamiltonian symplectomorphism constructed in Lemma 6.9 can be constructed ι_2 -equivariantly (extending ι_2 from M to \hat{M} by the trivial involution on the second factor).

6.5. Constructing stable normal trivialisations. We return to X , its action of $G = \mathbb{Z}/2 \times \mathbb{Z}/2$, and the G -invariant Lagrangian matching spheres S_a and S_b . The following is an analogue of [SS10, Lemma 31].

Lemma 6.15. *The triple (X, S_a, S_b) admits compatible stable normal trivialisations.*

Proof. Consider the map $\mathbb{C}^4 \times \mathbb{C}^* \rightarrow \mathbb{C}^2$ taking $(u_1, v_1, u_2, v_2, z) \mapsto (u_1 v_1 - z, u_2 v_2 - z)$. Then X is the regular fibre over $(-1, 1)$, so we have a short exact sequence of complex vector bundles

$$(6.3) \quad 0 \rightarrow TX \rightarrow T(\mathbb{C}^4 \times \mathbb{C}^*) \rightarrow T\mathbb{C}^2 \rightarrow 0.$$

There is a contractible space of splittings of such a sequence (from a Hermitian metric) which gives a canonical homotopy class of stable trivialisation of TX ; thinking about the family

of regular fibres $X_{c,d}$ over $\text{Conf}_2(\mathbb{C}^*) \subset \mathbb{C}^2$ shows the canonical homotopy class of stable trivialisation is preserved by parallel transport and the PBr_3 -action.

The Lagrangian S_a arises as a vanishing cycle, which means we can find a disc $D \subset \mathbb{C}^2$ normal to the diagonal and a one-parameter family $\mathcal{X}_D \rightarrow D$ with fibre $X \mapsto 1$ and such that $\mathcal{X}_D \supset \Delta_a$ contains a Lagrangian thimble fibred over an arc $[0, 1] \subset D$ and with boundary $\Delta_a \cap X = S_a$.

We have a commutative diagram of vector bundles over S_a ,

$$(6.4) \quad \begin{array}{ccccccc} 0 & \longrightarrow & TS_a \otimes_{\mathbb{R}} \mathbb{C} & \longrightarrow & ((T\Delta_a)|_{S_a}) \otimes_{\mathbb{R}} \mathbb{C} & \longrightarrow & S_a \times \mathbb{C} \longrightarrow 0 \\ & & \downarrow \cong & & \downarrow \cong & & \downarrow = \\ 0 & \longrightarrow & TX|_{S_a} & \longrightarrow & T(\mathcal{X}_D)|_{S_a} & \longrightarrow & S_a \times \mathbb{C} \longrightarrow 0 \end{array}$$

Since Δ_a is contractible, $T\Delta_a$ is trivial, and this induces a stable trivialization of TS_a ; the nullhomotopy of the stabilised Lagrangian Gauss map required for Definition 6.8 thus comes from an extension of that map to Δ_a . The commutativity of (6.4) shows that the complexification of this trivialization is (canonically) homotopic to the trivialization of $TX|_{S_a}$ arising from the previous stable trivialisation of TX from (6.3).

Because of our previous remarks about parallel transport, the same is true for any matching sphere S_b , and furthermore the argument goes through equivariantly. More explicitly, TX has a G -equivariant stable trivialization, compatible with a G -equivariant stable trivialisation of TS_a in the sense that under the natural isomorphism $TS_a \otimes \mathbb{C} \cong TX|_{S_a}$, there is an equivariant homotopy between these two trivializations. Restriction to fixed point sets of either generator of G and to the corresponding anti-invariant directions along those fixed-point loci now yields the desired compatible stable normal trivializations. \square

At this point we have satisfied all of the hypotheses of Lemma 6.13, bearing in mind that in Lemma 6.5 we constructed Hamiltonian isotopies of the matching spheres after which $S_a \cap S_b = s_a \cap s_b$, i.e. all intersection points lie in the common fixed point locus. This completes the construction of the J required for Lemma 6.6, and hence our discussion of the proof of Proposition 6.4.

6.6. Dynamics on surfaces. Having a formula for ranks of Floer cohomologies of matching spheres in terms of intersection numbers on the surface X^{inv} leads to growth rate bounds, using classical arguments from dynamics on surfaces.

Proposition 6.16. *Suppose that an autoequivalence $\phi \in \mathbb{Z}^{*\infty}$ is such that for all $S_i, S_j \in \mathcal{S}$, the rank of the Floer cohomology group $\text{HF}(S_i, \phi^n S_j; \mathbb{Z}/2)$ grows at most linearly with n . Then, up to shifts, ϕ is a power of a spherical twist in an element of \mathcal{S} .*

Proof. The Nielsen-Thurston theorem (see [FLP79]) says that elements of mapping class groups of punctured surfaces are either periodic, reducible, or pseudo-Anosov. Concretely for PBr_3 these cases simplify as follows. The only periodic elements are powers of the full twist (i.e. elements in the centre of PBr_3); the only reducible elements are powers of a full twist in a matching path between marked points; any other element is pseudo-Anosov, which by definition means that it is pseudo-Anosov as a mapping of the four-punctured sphere given by collapsing and removing the boundary of the disc. If ϕ is pseudo-Anosov, its lift $\hat{\phi}$ to the four-fold cover X^{inv} introduced in Proposition 6.4 is still pseudo-Anosov (as it preserves two transverse singular foliations, namely the preimages under the branched cover of the foliations

preserved by ϕ downstairs). Then for any two simple closed curves ℓ, ℓ' on X^{inv} the geometric intersection number $\iota(\ell, \hat{\phi}^k \ell')$ grows exponentially in k [FLP79, Exposé 12, Théorème II]. Taking ℓ, ℓ' to be lifts of arcs γ_i, γ_j downstairs, this geometric intersection number gives a lower bound for the rank of $\text{HF}(S_i, \phi^k S_j; \mathbb{Z}/2)$ by Proposition 6.4.

Suppose, then, that the growth of Floer cohomology is at most linear. Since $\phi \in \mathbb{Z}^{*\infty}$, if it is the image of a power of a full twist in a matching path, that matching path must be associated to some element of \mathcal{S} . If the growth rate is linear for some S_i, S_j , then ϕ is a shift of a non-trivial power of a Dehn twist in a matching sphere; if the growth rate is trivial for all i, j then ϕ is the image of a central element, and acts by a shift. \square

Lemma 6.17. *For a spherical twist $\phi = T_S$, the rank of $\text{HF}(S_i, \phi^n S_j; \mathbb{Z}/2)$ grows at most linearly with n .*

Proof. This follows immediately from the exact triangle for a spherical twist (6.1). \square

6.7. A spherical twist is a power of a Dehn twist. The conclusion of the previous section is that, as an autoequivalence of $DW(X)$, $T_S = \tau_\gamma^l[k]$, where τ_γ denotes the Dehn twist in the matching sphere S_γ associated to a matching path γ , and $k, l \in \mathbb{Z}$ are to be determined.

Lemma 6.18. *We have $k = 0$, so the spherical twist $T_S = \tau_\gamma^l$ is the image of some power of a Dehn twist in a matching path.*

Proof. By conjugating with a suitable element of $\mathbb{Z}^{*\infty}$, we may assume without loss of generality that $S_\gamma = S_1$, mirror to $\mathcal{O}_C(-1)$. In particular, since S_1 is disjoint from the Lagrangian thimble-like object L_0 (see Figure 2), we have that

$$T_S(L_0) = \tau_\gamma^l(L_0)[k] = L_0[k].$$

Suppose for contradiction that $k \neq 0$. Then in the exact triangle

$$\text{HF}^*(S, L_0) \otimes S \longrightarrow L_0 \longrightarrow T_S(L_0) = L_0[k]$$

the second arrow is given by an element of $HW^0(L_0, L_0[k]) = HW^k(L_0, L_0)$. But [CPU16, Remark 6.1] computes that the endomorphisms of L_0 are concentrated in degree zero, so this vanishes. This implies

$$\text{HF}^*(S, L_0) \otimes S \simeq L_0 \oplus L_0[k]$$

Taking self-endomorphisms of this object, since $HW^*(L_0, L_0) = SH^0(X)$ is infinite-dimensional, whilst $\text{HF}^*(S, S)$ has finite rank, we see that $\text{HF}^*(S, L_0)$ must be an infinite-dimensional vector space, which contradicts that S is spherical. We conclude that the shift satisfies $k = 0$. \square

6.8. Reduction to the ‘small category’ \mathcal{C} . Continuing with the same notation, in the course of the previous proof we showed that $T_S(L_0) = L_0$. Strengthening this:

Lemma 6.19. $\text{HF}^*(S, L_0) = 0 = \text{HF}^*(L_0, S)$.

Proof. For any given $n \in \mathbb{Z}$,

$$\text{HF}^*(S, L_0) = \text{HF}^*(S, T_S^n(L_0)) = \text{HF}^*(T_S^{-n} S, L_0) = \text{HF}^*(S[-2n], L_0) = \text{HF}^{*+2n}(S, L_0)$$

where we have used that for any 3-dimensional spherical object \hat{S} , the twist functor acts on the object itself by a non-trivial shift, $T_{\hat{S}}(\hat{S}) = \hat{S}[2]$.

Taking $n \neq 0$ shows that the vector space $HF^*(S, L_0)$ is periodic with some non-trivial grading shift (so infinite rank) or vanishes; again by the definition of S being spherical, it must be the latter. Thus, $HF^*(S, L_0) = 0$. By non-degeneracy of the pairing

$$HF^*(S, L_0) \otimes HF^*(L_0, S) \rightarrow \mathbb{C}$$

(or just by repeating the argument with the order of the branes reversed) we conclude that $HF^*(L_0, S) = 0$ also. \square

In the course of the proof of Lemma 6.3, we showed that the mirror object $\mathcal{E}_S \in D(Y)$ to S belonged to the full subcategory \mathcal{D} of complexes with cohomology supported on the curve C . There is a ‘small’ category, cf. [HWb] for instance,

$$\mathcal{C} = \{F \in D(Y) \mid \text{supp}(F) \subset C, R\pi_*(F) = 0\} \subset \mathcal{D} \subset D(Y)$$

of objects of $D(Y)$ which are both cohomologically supported on C and have trivial push-forward under the small resolution.

Lemma 6.20. *The mirror complex $\mathcal{E}_S \in \mathcal{D}$ belongs to the subcategory \mathcal{C} .*

Proof. Recall that L_0 is mirror to \mathcal{O}_Y , so

$$HF^*(L_0, S) = 0 \Rightarrow \text{Ext}^*(\mathcal{O}_Y, \mathcal{E}_S) = 0.$$

Pushing down under the contraction $\pi : Y \rightarrow \text{Spec}(R) = A$, we obtain

$$\text{Ext}^*(\mathcal{O}_A, R\pi_*\mathcal{E}_S) = 0.$$

However, A is affine and \mathcal{O}_A is a spanning object for the derived category $D(A)$, so this forces $R\pi_*\mathcal{E}_S = 0$. We deduce that $\mathcal{E}_S \in \mathcal{C}$. \square

6.9. Conclusion. To recap, if $S \in DW(X)$ is spherical, it has a mirror \mathcal{E}_S which is a spherical object in $\mathcal{D} \subset D(Y)$. We showed that the twist $T_{\mathcal{E}_S}$ is mirror to τ_{S_γ} for some $S_\gamma \in \mathcal{S}$. Conjugating with a PBr_3 element taking γ to γ_1 , wlog $T_{\mathcal{E}_S}$ is mirror to τ_{S_1} , and we saw above that moreover we get that \mathcal{E}_S is a spherical object in the small category $\mathcal{C} \subset D(Y)$. The classification of spherical objects in \mathcal{C} is standard (cf. [SW, HWb]; in the terminology of [SW], in our case \mathcal{C} corresponds to the one-vertex no-loop quiver). There is a unique spherical object up to shift, given by $\mathcal{O}_C(-1)$. Translating back under the mirror equivalence $DW(X) \simeq D(Y)$, it follows that S is quasi-isomorphic to some shift of S_γ .

This concludes the proof of Theorem 6.2. Passing back through the mirror, this readily implies Theorem 1.3.

7. MISCELLANEA

7.1. Addition of subcritical Weinstein handles. We will need the following generalisation of Lemma 3.17.

Lemma 7.1. *Suppose X' is a Weinstein 6-manifold given by iteratively adding any sequence of Weinstein one- and two-handles to X . Let $f \in \text{Homeo}_c(X')$ be a compactly supported homeomorphism. Then f_* acts trivially on $H_3(X'; \mathbb{Z})$.*

Proof. Take all homology groups to have coefficient ring \mathbb{Z} . Comparing with the proof of Lemma 3.17, it’s enough to show that the map $H_3(X') \rightarrow H_3(X', \partial X')$ has vanishing image. Consider the non-degenerate intersection pairing

$$H_3(X') \times H_3(X', \partial X') \rightarrow \mathbb{Z}.$$

As $X \subset X'$ is given by adding Weinstein one- and two-handles, the inclusion induces an isomorphism $H_3(X) \cong H_3(X')$. In particular, $H_3(X') \cong \mathbb{Z}^2$, generated by S_0 and S_1 . As the intersection pairing of S_i and S_j vanishes for any i, j , we see that for any i the image of S_i in $H_3(X', \partial X')$ vanishes. This completes the proof. \square

Corollary 7.2. *Suppose X' is any Weinstein 6-manifold given by iteratively adding Weinstein one- and two-handles to X . Then under the action of $\pi_0 \text{Symp}_c(X')$, there are infinitely many orbit sets of Lagrangian spheres in X' .*

This answers a folklore question often attributed to Fukaya.

Proof. The spheres S_i belong to pairwise distinct classes in $H_3(X'; \mathbb{Z})$, and any compactly supported symplectomorphism acts trivially on $H_3(X'; \mathbb{Z})$ by Lemma 7.1. \square

Corollary 7.3. *Suppose X' is a Weinstein 6-manifold given by iteratively adding any sequence of Weinstein one- and two-handles to X . Then we have a pair of group homomorphisms*

$$\mathbb{Z}^{*\infty} \hookrightarrow \pi_0 \text{Symp}_c(X') \twoheadrightarrow \mathbb{Z}^{*\infty}$$

which compose to the identity.

In particular, by adding a two-handle, we can find such X' 's which are simply connected.

Proof. We have a natural isomorphism of wrapped Fukaya categories $DW(X) \simeq DW(X')$, compatible with an isomorphism between $SH^0(X)$ and $SH^0(X')$. (By [CRGG, GPS20], the categories $\mathcal{W}(X)$ and $\mathcal{W}(X')$ are non-degenerate and equivalent. The equivalence induces an isomorphism of Hochschild (co)homologies. On the other hand, by [Gan12], the non-degeneracy implies that $HH_*(\mathcal{W}(X)) \rightarrow SH^*(X)$ is an isomorphism. It's then immediate that we get a compatible isomorphism between $SH^0(X)$ and $SH^0(X')$.)

This implies that a compactly supported symplectomorphism $\phi \in \pi_0 \text{Symp}_c(X')$ induces an $SH^0(X)$ -linear autoequivalence of $DW(X)$. As before, combining mirror symmetry for X with Lemmas 4.5, 4.6 and 5.3, we get that ϕ preserves the subcategory $\langle S_0, S_1 \rangle$ (mirror to \mathcal{D}), and fixes the torus (T, ζ) , as an object of $DW(X)$, up to a shift.

By Lemma 7.1, ϕ acts as the identity $H_3(X'; \mathbb{Z}) \cong H_3(X; \mathbb{Z})$, i.e. on the numerical K -theory $K_{\text{num}}\mathcal{F}(X')$. As before, Corollary 4.11 implies that we get a map $\pi_0 \text{Symp}_c(X') \rightarrow \mathbb{Z}^{*\infty} \times 2\mathbb{Z}$. Projecting gives a map $\pi_0 \text{Symp}_c(X') \rightarrow \mathbb{Z}^{*\infty}$, which is surjective by considering the image of the Dehn twists. The claim then follows. \square

7.2. Non-exact deformations. By open-ness of the symplectic condition, for a small neighbourhood U of $(0, 0) \in H^2(X; \mathbb{R}) \cong \mathbb{R}^2$, there is a non-exact symplectic deformation of $(X, d\theta)$ to $(X, \omega_{a,b})$ with $[\omega_{a,b}] = (a, b) \in U$.

We can in fact exhibit non-exact deformations $(X, \omega_{a,b})$ for arbitrary $(a, b) \in \mathbb{R}^2$, via the Morse-Bott-Lefschetz fibration. Let $t_c : T^*S^1 \rightarrow T^*S^1$ be a symplectomorphism of flux c (translating the cylinder). Start with the previously considered Morse-Bott-Lefschetz fibration $\pi : X \rightarrow \mathbb{C}^*$, with Morse-Bott singular fibres at ± 1 . This is a locally Hamiltonian fibration, with monodromies $\text{id} \times \tau_{S^1}$ about -1 and $\tau_{S^1} \times \text{id}$ about 1 . We define $(X, \omega_{a,b})$ to be the total space of the corresponding locally Hamiltonian fibration with local monodromies $t_a \times \tau_{S^1}$ about -1 and $\tau_{S^1} \times t_b$ about 1 .

For any (a, b) , we can still construct Lagrangian matching spheres S_i for the same matching paths as in Figure 2. By inspection, whenever a and b are both non-zero, S_i and S_j are disjoint if $i \neq j$.

Corollary 7.4. *For $a \neq 0$ and $b \neq 0$, there is an injection $\mathbb{Z}^{\oplus\infty} \hookrightarrow \pi_0 \text{Symp}_c(X, \omega_{a,b})$.*

Proof. The Dehn twists in the S_i define a homomorphism $\mathbb{Z}^{\oplus\infty} \rightarrow \pi_0 \text{Symp}_c^{\text{gr}}(X, \omega_{a,b})$. We claim this is injective by considering the action on graded Lagrangian spheres: each S_i admits a \mathbb{Z} -torsor of gradings, and the twist τ_{S_i} shifts the grading on S_i by 2; as they are disjoint, it does not change the grading on any S_j for $i \neq j$. Finally, as the forgetful map from $\text{Symp}_c^{\text{gr}}(X, \omega_{a,b})$ to gradeable compactly supported symplectomorphisms of $(X, \omega_{a,b})$ splits, we get an injection $\mathbb{Z}^{\oplus\infty} \hookrightarrow \pi_0 \text{Symp}_c(X, \omega_{a,b})$ as required. \square

Remark 7.5. By inspection, for our description of the S_i as matching spheres, there is no compact subset of X containing all of the matching spheres S_i : as $|i|$ increases, the S_i have arbitrarily large maximal radial coordinate in the conical end for (the deformation of) X .

7.3. Homeomorphism & diffeomorphism groups.

Lemma 7.6. *The composite $\mathbb{Z}^{*\infty} \rightarrow \pi_0 \text{Symp}_c(X, d\theta) \rightarrow \pi_0 \text{Diff}_c(X)$ factors through $\mathbb{Z}^{\oplus\infty}$.*

Proof. This immediately follows by noting that this is the same map as the composite $\mathbb{Z}^{*\infty} \rightarrow \pi_0 \text{Symp}_c(X, \omega_{a,b}) \rightarrow \pi_0 \text{Diff}_c(X)$ for any deformation $(X, \omega_{a,b})$ of $(X, d\theta)$, and recalling that the Dehn twists in the S_i generate a $\mathbb{Z}^{\oplus\infty}$ in $\pi_0 \text{Symp}_c(X, \omega_{a,b})$. \square

Recall that $\pi_1(X) = \mathbb{Z}$. We view X as the completion of a Weinstein domain $(X, \partial X)$ and let $(X', \partial X')$ denote the Weinstein domain obtained by adding one subcritical two-handle to X so as to kill its fundamental group.

Lemma 7.7. *The boundary $\partial X'$ is diffeomorphic to $(S^2 \times S^3) \# (S^2 \times S^3)$.*

Proof. The boundary ∂X has $\pi_1(\partial X) = \mathbb{Z}$. On the boundary, the subcritical handle addition acts by a surgery, removing $S^1 \times D^4$ and regluing $D^2 \times S^3$, and so $\pi_1(\partial X') = \{1\}$. The long exact sequence for $(X, \partial X)$ shows that $H_2(\partial X) \cong H_2(X) \cong \mathbb{Z}^2$, and hence $H_2(\partial X'; \mathbb{Z}) = \mathbb{Z}^2$ also. In particular, the homology of X' is torsion-free. Since $H^2(X'; \mathbb{Z}) \rightarrow H^2(X; \mathbb{Z})$ is an isomorphism, X' still has vanishing first Chern class, which shows that $\partial X'$ is spin. Simply-connected spin 5-manifolds are classified [Bar65]; when the homology is torsion-free they are given by $(S^2 \times S^3) \#^r$ where r is the second Betti number. \square

Recall that we previously constructed a diagram

$$\begin{array}{ccc} \mathbb{Z}^{*\infty} & \longrightarrow & \pi_0 \text{Symp}_c(X') \\ \downarrow & & \downarrow \\ \mathbb{Z}^{\oplus\infty} & \longrightarrow & \pi_0 \text{Diff}_c(X') \end{array}$$

Lemma 7.8. *The map $\mathbb{Z}^{\oplus\infty} \rightarrow \pi_0 \text{Diff}_c(X')$ factors through a finite rank abelian group.*

Proof. If $(M, \partial M)$ is a simply-connected six-manifold with simply-connected boundary, then Kupers shows [Kup, Theorem A, Remark (iv)] that the mapping class group $\pi_0 \text{Diff}_c(M)$ has finite type in the sense that its classifying space admits a CW structure with only finitely many cells in each dimension. The proof actually yields a stronger result, see [Kup, Theorem 4.2], which is that there is an exact sequence

$$1 \rightarrow \Gamma \rightarrow \pi_0 \text{Diff}_c(M) \rightarrow G \rightarrow 1$$

where Γ is a finitely generated abelian group, and G is an arithmetic group. For any group fitting into such an exact sequence, any abelian subgroup is finitely generated. \square

REFERENCES

- [AS21] Denis Auroux and Ivan Smith. Fukaya categories of surfaces, spherical objects and mapping class groups. *Forum Math. Sigma*, 9:Paper No. e26, 50, 2021. [1.2](#)
- [Aur07] Denis Auroux. Mirror symmetry and T -duality in the complement of an anticanonical divisor. *J. Gökova Geom. Topol. GGT*, 1:51–91, 2007. [3.1](#), [5.4](#)
- [Bal] Matthew Ballard. Sheaves on local Calabi-Yu varieties. arXiv:0801.3499. [4](#)
- [Bar65] D. Barden. Simply connected five-manifolds. *Ann. of Math. (2)*, 82:365–385, 1965. [7.3](#)
- [BCZ17] Arend Bayer, Alastair Craw, and Ziyu Zhang. Nef divisors for moduli spaces of complexes with compact support. *Selecta Math. (N.S.)*, 23(2):1507–1561, 2017. [5](#), [6.1](#)
- [BEE12] Frédéric Bourgeois, Tobias Ekholm, and Yasha Eliashberg. Effect of Legendrian surgery. *Geom. Topol.*, 16(1):301–389, 2012. With an appendix by Sheel Ganatra and Maksim Maydanskiy. [2.3](#), [2.2](#)
- [Bri07] Tom Bridgeland. Stability conditions on triangulated categories. *Ann. of Math. (2)*, 166(2):317–345, 2007. [4](#)
- [CPU16] Kwokwai Chan, Daniel Pomerleano, and Kazushi Ueda. Lagrangian torus fibrations and homological mirror symmetry for the conifold. *Comm. Math. Phys.*, 341(1):135–178, 2016. [1.3](#), [4.3](#), [5.1](#), [5](#), [5](#), [6.7](#)
- [CRGG] Baptiste Chantraine, Georgios Dimitroglou Rizell, Paolo Ghiggini, and Roman Golovko. Geometric generation of the wrapped Fukaya category of Weinstein manifolds and sectors. arXiv:1712.09126. [3.6](#), [7.1](#)
- [CS14] Alberto Canonaco and Paolo Stellari. Fourier-Mukai functors in the supported case. *Compos. Math.*, 150(8):1349–1383, 2014. [4](#)
- [DHKK14] G. Dimitrov, F. Haiden, L. Katzarkov, and M. Kontsevich. Dynamical systems and categories. In *The influence of Solomon Lefschetz in geometry and topology*, volume 621 of *Contemp. Math.*, pages 133–170. Amer. Math. Soc., Providence, RI, 2014. [1.4](#)
- [FLP79] *Travaux de Thurston sur les surfaces*, volume 66 of *Astérisque*. Société Mathématique de France, Paris, 1979. Séminaire Orsay, With an English summary. [6.6](#)
- [Gan12] Sheel Ganatra. *Symplectic Cohomology and Duality for the Wrapped Fukaya Category*. ProQuest LLC, Ann Arbor, MI, 2012. Thesis (Ph.D.)—Massachusetts Institute of Technology. See arXiv:1304.7312. [5](#), [7.1](#)
- [GHK15] Mark Gross, Paul Hacking, and Sean Keel. Mirror symmetry for log Calabi-Yau surfaces I. *Publ. Math. Inst. Hautes Études Sci.*, 122:65–168, 2015. [1.3](#)
- [GPS] Sheel Ganatra, John Pardon, and Vivek Shende. Sectorial descent for wrapped Fukaya categories. arXiv:1809.03427v3. [2.1](#), [2.2](#), [2.5](#), [3.6](#)
- [GPS20] Sheel Ganatra, John Pardon, and Vivek Shende. Covariantly functorial wrapped Floer theory on Liouville sectors. *Publ. Math. Inst. Hautes Études Sci.*, 131:73–200, 2020. [2.2](#), [5](#), [7.1](#)
- [HKa] Paul Hacking and Ailsa Keating. Homological mirror symmetry for log Calabi-Yau surfaces. arXiv:2005.05010. [1.3](#)
- [HKb] Paul Hacking and Ailsa Keating. Symplectomorphisms of mirrors to log Calabi-Yau surfaces. arXiv:2112.06797. [1.3](#)
- [Huy06] D. Huybrechts. *Fourier-Mukai transforms in algebraic geometry*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, Oxford, 2006. [4](#), [4](#), [5](#), [6.1](#)
- [HWa] Wahei Hara and Michael Wemyss. Spherical objects in dimension two and three. arXiv:2205.11552. [1.2](#)
- [HWb] Yuki Hirano and Michael Wemyss. Stability Conditions for 3-fold Flops. arXiv:1907.09742. [1.3](#), [4.3](#), [4.10](#), [6.8](#), [6.9](#)
- [IU05] Akira Ishii and Hokuto Uehara. Autoequivalences of derived categories on the minimal resolutions of A_n -singularities on surfaces. *J. Differential Geom.*, 71(3):385–435, 2005. [1.2](#)
- [Kea15] Ailsa Keating. Lagrangian tori in four-dimensional Milnor fibres. *Geom. Funct. Anal.*, 25(6):1822–1901, 2015. [3.5](#)
- [KS02] Mikhail Khovanov and Paul Seidel. Quivers, Floer cohomology, and braid group actions. *J. Amer. Math. Soc.*, 15(1):203–271, 2002. [1.2](#), [3.5](#), [6.2](#), [6.2](#), [6.3](#), [6.3](#)
- [Kup] Alexander Kupers. Mapping class groups of manifolds with boundary are of finite type. arXiv:2204.01945. [1.1](#), [7.3](#)
- [LO10] Valery A. Lunts and Dmitri O. Orlov. Uniqueness of enhancement for triangulated categories. *J. Amer. Math. Soc.*, 23(3):853–908, 2010. [4](#)

- [MM09] Dan Margalit and Jon McCammond. Geometric presentations for the pure braid group. *J. Knot Theory Ramifications*, 18(1):1–20, 2009. 3.5
- [Pen91] R. C. Penner. Bounds on least dilatations. *Proc. Amer. Math. Soc.*, 113(2):443–450, 1991. 1.4
- [Pom] Dan Pomerleano. Intrinsic mirror symmetry and categorical crepant resolutions. arXiv:2103.01200. 5, 6.1
- [Rei83] Miles Reid. Minimal models of canonical 3-folds. In *Algebraic varieties and analytic varieties (Tokyo, 1981)*, volume 1 of *Adv. Stud. Pure Math.*, pages 131–180. North-Holland, Amsterdam, 1983. 4.3
- [Sei] Paul Seidel. Categorical dynamics and symplectic topology. Lecture notes, available at <https://math.mit.edu/~seidel>. 3.1
- [Sei00] Paul Seidel. Graded Lagrangian submanifolds. *Bull. Soc. Math. France*, 128(1):103–149, 2000. 2.3, 2.3, 3.4
- [Sei08a] Paul Seidel. A biased view of symplectic cohomology. In *Current developments in mathematics, 2006*, pages 211–253. Int. Press, Somerville, MA, 2008. 3.1
- [Sei08b] Paul Seidel. *Fukaya categories and Picard-Lefschetz theory*. Zurich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zürich, 2008. 2.1, 6.3, 6.3
- [Sei12] Paul Seidel. Lagrangian homology spheres in (A_m) Milnor fibres via \mathbb{C}^* -equivariant A_∞ -modules. *Geom. Topol.*, 16(4):2343–2389, 2012. 6.1
- [Smi22] Gleb Smirnov. Symplectic mapping class groups of K3 surfaces and Seiberg-Witten invariants. *Geom. Funct. Anal.*, 32(2):280–301, 2022. 1.2
- [SS05] Paul Seidel and Ivan Smith. The symplectic topology of Ramanujam’s surface. *Comment. Math. Helv.*, 80(4):859–881, 2005. 2.2
- [SS10] Paul Seidel and Ivan Smith. Localization for involutions in Floer cohomology. *Geom. Funct. Anal.*, 20(6):1464–1501, 2010. 6.3, 6.3, 6.5
- [SS20] Nick Sheridan and Ivan Smith. Symplectic topology of K3 surfaces via mirror symmetry. *J. Amer. Math. Soc.*, 33(3):875–915, 2020. 1.2
- [Sta18] The Stacks Project Authors. *Stacks Project*. <https://stacks.math.columbia.edu>, 2018. 4.3
- [SW] Ivan Smith and Michael Wemyss. Double bubble plumbings and two-curve flops. arXiv:2010.10114. 1.2, 3.1, 3.4, 3.5, 3.5, 6.9
- [Tod08] Yukinobu Toda. Stability conditions and crepant small resolutions. *Trans. Amer. Math. Soc.*, 360(11):6149–6178, 2008. 1.3, 4, 4.9
- [Tod09] Yukinobu Toda. Stability conditions and Calabi-Yau fibrations. *J. Algebraic Geom.*, 18(1):101–133, 2009. 1.3, 4, 4.9, 4
- [Toë07] Bertrand Toën. The homotopy theory of dg -categories and derived Morita theory. *Invent. Math.*, 167(3):615–667, 2007. 4
- [Wu14] Weiwei Wu. Exact Lagrangians in A_n -surface singularities. *Math. Ann.*, 359(1-2):153–168, 2014. 1.2

CENTRE FOR MATHEMATICAL SCIENCES, UNIVERSITY OF CAMBRIDGE, WILBERFORCE ROAD, CB3 0WB, UNITED KINGDOM

Email address: amk50@cam.ac.uk

CENTRE FOR MATHEMATICAL SCIENCES, UNIVERSITY OF CAMBRIDGE, WILBERFORCE ROAD, CB3 0WB, UNITED KINGDOM

Email address: is200@cam.ac.uk