

Finite Alphabet Phase Retrieval

Tamir Bendory, Dan Edidin, Ivan Gonzalez

January 26, 2023

Abstract

We consider the finite alphabet phase retrieval problem: recovering a signal whose entries lie in a small alphabet of possible values from its Fourier magnitudes. This problem arises in the celebrated technology of X-ray crystallography to determine the atomic structure of biological molecules. Our main result states that for generic values of the alphabet, two signals have the same Fourier magnitudes if and only if several partitions have the same difference sets. Thus, the finite alphabet phase retrieval problem reduces to the combinatorial problem of determining a signal from those difference sets. Notably, this result holds true when one of the letters of the alphabet is zero, namely, for sparse signals with finite alphabet, which is the situation in X-ray crystallography.

1 Introduction

X-ray crystallography is a leading technology for determining the 3-D atomic structure of biological molecules, such as proteins. Indeed, thousands of new crystal structures are resolved each year, and more than a dozen Nobel Prizes have been awarded for work involving X-ray crystallography. In X-ray crystallography, the crystal—a periodic arrangement of a repeating unit—is illuminated with a beam of X-rays, producing a diffraction pattern, which is equivalent to the magnitude of the Fourier transform of the crystal. The signal to be estimated (the electron density function of the crystal) is supported only at the sparsely-spread positions of atoms [11]. Therefore, the *crystallographic phase retrieval problem* entails recovering a sparse signal from its Fourier magnitudes. The crystallographic phase retrieval is a special case of the *phase retrieval* problem, which refers to all problems that involve recovering a signal from its Fourier magnitudes, see [18, 1, 9, 3] and reference therein. A detailed mathematical model of X-ray crystallography is introduced in [7].

A recent paper by a subset of the authors provides the first rigorous attempt to establish a mathematical theory for the crystallographic phase retrieval problem [2]. In particular, it was conjectured that a generic sparse signal $x \in \mathbb{R}^N$ whose support has size at most K is uniquely determined, up to unavoidable ambiguities, as long as $K \leq N/2$. The conjecture was verified for a small set of parameters; see also [8, 15].

In practice, however, a more accurate model of the crystallographic phase retrieval problem should account for sparse signals whose non-zero entries are taken from a finite (small) alphabet; this alphabet models the relevant type of atoms, such as hydrogen, oxygen, carbon, nitrogen, and so on. In this paper, we make a first step towards this direction. Specifically,

we study the problem of recovering a signal, whose entries are taken from a finite alphabet, from its Fourier magnitudes. We refer to this problem as the *finite alphabet phase retrieval problem*.

In particular, we show that for generic choice of entries in the alphabet, the finite alphabet phase retrieval problem can be reduced to a combinatorial problem involving difference sets. This is similar to the situation for binary phase retrieval (a problem studied before [6]) but new combinatorial subtleties can occur. More specifically, we show that two signals with entries taken from a finite alphabet have the same Fourier magnitudes if and only if the associated partitions have the same difference sets; see Proposition 4.2. Notably, this result remains true when one of the letters of the alphabet is zero, namely, for sparse signals with finite alphabet; see Theorem 4.3. This is the situation in X-ray crystallography where crystals are typically very sparse; the non-zero values occupy only $\sim 1/100$ of the signal's support [6]. Unfortunately, the problem of analyzing if specific difference sets determine a set uniquely (up to unavoidable symmetries) is an extremely difficult combinatorial problem [17, p.350], [16, Section 3]. Therefore, we cannot provide a complete characterization when a finite-alphabet signal can be recovered uniquely, up to unavoidable symmetries, from its Fourier magnitudes.

The rest of the paper is organized as follows. Section 2 provides a necessary background on difference sets, homometric sets, and autocorrelations. Section 3 begins our analysis by studying signals whose entries are taken from two-letters alphabet. Section 4 presents and proves our main results about phase retrieval of signals whose entries are taken from a finite alphabet. Section 5 provides a few examples and introduces the intriguing notion of pseudo-equivalent partitions.

2 Background

We begin by introducing basic definitions about difference sets. For any $i, j \in [0, N - 1]$, we define the *cyclic distance* between i and j by

$$d(i, j) = \min\{N - |i - j|, |i - j|\}. \quad (2.1)$$

We note that $d(i, j) \in [0, \lfloor N/2 \rfloor]$ as illustrated in Figure 1.

Definition 2.1 (Difference sets). Let A, B be two subsets of $[0, N - 1]$. We define the cyclic difference multi-set by

$$A - B = \{d(i, j) \mid i \in A, j \in B\}. \quad (2.2)$$

In particular, the self difference multi-set is given by $A - A = \{d(i, j) \mid i \leq j \in A\}$.

Definition 2.2 (The dihedral group). The dihedral group D_{2N} is the group of symmetries of the regular N -gon. It is a group of order $2N$, which is generated by two elements r (rotation) and s (reflection). The elements r, s satisfy the relations $r^N = s^2 = e$ and $rs = sr^{N-1}$, where e is the identity. The group D_{2N} acts on the set $[0, N - 1]$ as follows. The element $r \in D_{2N}$ acts by cyclic shift; i.e., $r(i) = (i + 1) \bmod N$ and the element s acts by the reflection $s(i) = N - i$.




Figure 1: Illustration of the cyclic distances between the points 0 and 5 and 1 and 5 in $[0, 7]$.

Next, we define equivalent classes. Since the Fourier magnitude is invariant under cyclic shifts and reflection (i.e., under the dihedral group), we can only hope to determine the support of a signal from the signal's Fourier magnitudes up to an action of the dihedral group D_{2N} .

Definition 2.3. Two subsets $A, B \subset [0, N - 1]$ are *equivalent* if there exists an element $\sigma \in D_{2N}$ such that $B = \sigma A$.

Another fundamental definition is that of homometric sets: subsets with the same difference set.

Definition 2.4. Two subsets A, B are *homometric* if $A - A = B - B$.

Lemma 2.5. *If A, B are equivalent then they are homometric.*

Proof. The lemma is an immediate consequence of the fact that the cyclic distance $d(i, j)$ is invariant under cyclic shifts and reflection. \square

A key fact we use about homometric subsets of $[0, N - 1]$ is the following result, originally stated by Patterson [14]. For a modern proof, see [10, Corollary 1] or [5].

Theorem 2.6 (Patterson). *Two sets $A, B \subset [0, N - 1]$ are homometric if and only if their complements are homometric as well.*

We now consider the case of partitions. Let A_1, \dots, A_K and B_1, \dots, B_K be two ordered partitions of $[0, N - 1]$.

Definition 2.7. Two ordered partitions A_1, \dots, A_K and B_1, \dots, B_K are *homometric* if $A_i - A_j = B_i - B_j$ for all pairs $i, j \in \{1, \dots, K\}$. Two ordered partitions A_1, \dots, A_K and B_1, \dots, B_K are *equivalent* if there exists $\sigma \in D_{2N}$ such that $B_i = \sigma A_i$ for all $i \in \{1, \dots, K\}$.

Lemma 2.5 can be directly extended to ordered partitions.

Lemma 2.8. *Equivalent partitions are homometric.*

Before moving to the next section, we remind the reader of a couple of definitions from signal processing.

Definition 2.9 (Power spectrum). The power spectrum of a signal $x \in \mathbb{C}^N$ is the vector $|\hat{x}|^2 \in \mathbb{R}_{\geq 0}^N$, where \hat{x} is the discrete Fourier transform (DFT) of x and the absolute value is taken componentwise. The periodic auto-correlation of x is defined by

$$a_x[\ell] = \sum_{n=0}^{N-1} x[n] \overline{x[\ell+n]}, \quad (2.3)$$

where all indices are taken modulo N .

A key fact, first observed by Patterson [12, 13], is that the DFT of a_x is the power spectrum. The phase retrieval problem is thus equivalent to the problem of recovering a signal from its periodic auto-correlation. In the sequel, we use the terms Fourier magnitudes, power spectrum and autocorrelation interchangeably.

3 Binary and two-alphabet phase retrieval

The binary phase retrieval problem is the problem of recovering a binary signal $x \in \mathbb{R}^N$ from its periodic auto-correlation $a_x \in \mathbb{R}^N$ [6, 4]. Let $S(x)$ denote the support of a signal x . A well known result for binary signals states that $a_x = a_{x'}$ if and only if $S(x) - S(x) = S(x') - S(x')$; i.e., the two supports are homometric (e.g., [2]).

Let us expand upon this result and replace the zeros and ones in the traditional binary phase retrieval with arbitrary scalars α and β . We refer to this problem as the *two-alphabet phase retrieval problem*. When either α or β is zero, then this problem reduces to the binary phase retrieval problem. Otherwise, we do not have a well-defined notion of a support set since neither one of the two letters is assumed to be zero. Instead, we consider the sets

$$S_\alpha(x) = \{i \mid x[i] = \alpha\} \quad \text{and} \quad S_\beta(x) = \{j \mid x[j] = \beta\}.$$

We prove the following result.

Theorem 3.1. *For a generic choice of values of α, β , the following are equivalent for signals $x, x' \in \{\alpha, \beta\}^N$.*

- (i) $a_x = a_{x'}$
- (ii) $S_\alpha(x)$ and $S_\alpha(x')$ are homometric
- (iii) $S_\beta(x)$ and $S_\beta(x')$ are homometric
- (iv) The ordered partitions $(S_\alpha(x), S_\beta(x))$ and $(S_\alpha(x'), S_\beta(x'))$ are homometric.

Proof. Clearly (iv) implies (ii) and (iii). If we show that (ii) implies (iv) then by symmetry we can also conclude that (iii) implies (iv). To see that (ii) implies (iv) we use Patterson's Theorem, Theorem 2.6. Note that $S_\beta(x) = S_\alpha(x)^c$, so by Patterson's Theorem $S_\alpha(x)$ and $S_\alpha(x')$ are homometric if and only if $S_\beta(x)$ and $S_\beta(x')$ are also homometric. To show that the partitions $(S_\alpha(x), S_\beta(x))$ and $(S_\alpha(x'), S_\beta(x'))$ are homometric we must

show that $S_\alpha(x) - S_\beta(x) = S_\alpha(x') - S_\beta(x')$. This follows from the fact that the difference sets $S_\alpha(x) - S_\alpha(x), S_\alpha(x) - S_\beta(x), S_\beta(x) - S_\beta(x)$ (respectively $S_\alpha(x') - S_\alpha(x'), S_\alpha(x') - S_\beta(x'), S_\beta(x') - S_\beta(x')$) form a partition of the multi-set $[0, N - 1] - [0, N - 1]$.

Let $a_x[\ell]$ and $a_{x'}[\ell]$ be the ℓ -th entry of a_x and $a_{x'}$, respectively. Then, $a_x[\ell] = m_\ell\alpha^2 + n_\ell\beta^2 + p_\ell\alpha\beta$, and $a_{x'}[\ell] = m'_\ell\alpha^2 + n'_\ell\beta^2 + p'_\ell\alpha\beta$, where m_ℓ (resp. m'_ℓ) is the multiplicity of ℓ in $S_\alpha(x) - S_\alpha(x)$ (resp. $S_\alpha(x') - S_\alpha(x')$), n_ℓ (resp. n'_ℓ) is the multiplicity of ℓ in $S_\beta(x) - S_\beta(x)$ (resp. $S_\beta(x') - S_\beta(x')$), and p_ℓ (resp. p'_ℓ) is the multiplicity of ℓ in $S_\alpha(x) - S_\beta(x)$ (resp. $S_\alpha(x') - S_\beta(x')$). Hence, if $(S_\alpha(x), S_\beta(x))$ and $(S_\alpha(x'), S_\beta(x'))$ are homometric then $a_x[\ell] = a_{x'}[\ell]$. Thus (iv) \implies (i).

Conversely, suppose that $a_x = a_{x'}$. This means that for each ℓ we have that

$$m_\ell\alpha^2 + n_\ell\beta^2 + p_\ell\alpha\beta = m'_\ell\alpha^2 + n'_\ell\beta^2 + p'_\ell\alpha\beta.$$

Let us define $m = m_\ell - m'_\ell, n = n_\ell - n'_\ell, p = p_\ell - p'_\ell$. This implies that there are integers m, n, p such that $m\alpha^2 + n\beta^2 + p\alpha\beta = 0$. However, for a generic choice of α, β , this is only possible if $m = n = p = 0$; i.e., $m_\ell = m'_\ell, n_\ell = n'_\ell$, and $p_\ell = p'_\ell$, meaning that the multiplicities of ℓ in $S_\alpha(x) - S_\alpha(x), S_\beta(x) - S_\beta(x), S_\alpha(x) - S_\beta(x)$ are equal to the multiplicities of ℓ in $S_\alpha(x') - S_\alpha(x'), S_\beta(x') - S_\beta(x'), S_\alpha(x') - S_\beta(x')$, respectively. Since for generic α, β this is true for every $\ell = 0, \dots, N - 1$ we conclude that the partitions $(S_\alpha(x), S_\beta(x))$ and $(S_\alpha(x'), S_\beta(x'))$ are homometric. Thus, (i) \implies (iv) for generic choice of α, β . □

4 Signals with entries taken from a finite alphabet

We now extend our analysis to account for signals whose entries are taken from a finite alphabet. Let $S = \{\alpha_1, \dots, \alpha_K\}$ be a set K real numbers and let \mathbb{R}_S be the set of all vectors in \mathbb{R}^N whose entries are taken from S . A vector $x \in \mathbb{R}_S$ determines a length K partition $A_1(x), \dots, A_K(x)$ of $[0, N - 1]$, where $A_k(x) = \{n \in [0, N - 1] \mid x[n] = \alpha_k\}$.

Remark 4.1. In this paper we assume that the alphabet is taken from the reals, but the theory is unchanged if we consider complex alphabet entries.

Proposition 4.2. *For a generic choice of $\alpha_1, \dots, \alpha_K$, two vectors $x, x' \in \mathbb{R}_S$ have the same auto-correlation if and only if the associated partitions $\{A_k(x)\}$ and $\{A_k(x')\}$ are homometric.*

Proof. The proof is similar to the proof of the equivalence (i) \iff (iv) in Theorem 3.1.

Since the entries $\alpha_1, \dots, \alpha_K$ are generic we can treat them as indeterminates. By definition, $a_x[\ell] = \sum_{n=0}^{N-1} x[n]x[n+\ell]$ and $a_{x'}[\ell] = \sum_{n=0}^{N-1} x'[n]x'[n+\ell]$, where all indices are taken modulo N . Since the entries of x, x' are taken from the set S , $a_x[\ell]$ and $a_{x'}[\ell]$ are quadratic polynomials in $\alpha_1, \dots, \alpha_K$. The coefficient of $\alpha_i\alpha_j$ in $a_x[\ell]$ is the multiplicity of ℓ in the difference multi-set $A_i(x) - A_j(x)$. Likewise, the coefficient of $\alpha_i\alpha_j$ in $a_{x'}[\ell]$ is the multiplicity of ℓ in the difference multi-set $A_i(x') - A_j(x')$. Hence, $a_x = a_{x'}$ if and only if $A_i(x) - A_j(x) = A_i(x') - A_j(x')$ for all i, j . In other words, $a_x = a_{x'}$ if and only if the corresponding partitions are homometric. □

We are now ready to present our main result, which is motivated by the crystallographic phase retrieval problem of recovering a sparse signal, whose non-zero values are taken from a finite alphabet, from its autocorrelation.

Theorem 4.3 (Sparse signals). *Consider two signals x, x' with entries taken from an alphabet $0, \alpha_2, \dots, \alpha_K$. Then, $a_x = a_{x'}$ if and only if the partitions $A(x)$ and $A(x')$ are homometric.*

Remark 4.4. The significance of this result is that we no longer assume that α_1 is arbitrary. Equivalently, our result states that if we view $a_x, a_{x'}$ as polynomials in the variables $\alpha_1, \dots, \alpha_K$, then $a_x(\alpha_1, \dots, \alpha_K) = a_{x'}(\alpha_1, \dots, \alpha_K)$ if and only if $a_x(0, \alpha_2, \dots, \alpha_K) = a_{x'}(0, \alpha_2, \dots, \alpha_K)$.

Proof. If $a_x = a_{x'}$, then for every ℓ the coefficients of $\alpha_i \alpha_j$ in $a_x[\ell]$ and $a_{x'}[\ell]$ are equal for $i, j \geq 2$. As before, this coefficient is just the multiplicity of ℓ in the difference multi-sets $A_i(x) - A_j(x)$ and $A_i(x') - A_j(x')$ respectively.

We begin with the following lemma.

Lemma 4.5. $A_1[x] - A_1[x] = A_1[x'] - A_1[x']$.

Proof. Let $B[x] = \cup_{i=2}^K A_i[x]$ and $B[x'] = \cup_{i=2}^K A_i[x']$. Then, $\{A_1[x], B[x]\}$ and $\{A_1[x'], B[x']\}$ are length two partitions of $[0, N-1]$. Now, $B[x] - B[x] = \sum_{i,j \geq 2} (A_i[x] - A_j[x]) = B[x'] - B[x']$ ¹. In other words, the subsets $B[x]$ and $B[x']$ are homometric. Since $A_1[x] = B[x]^c$ and $A_1[x'] = B[x']^c$, it follows from Patterson's Theorem that they also have the same difference sets. \square

To complete the proof that the partitions are homometric, we need to show that the difference sets $A_1[x] - A_i[x]$ and $A_1[x'] - A_i[x']$ are equal for all $i > 1$. To do this we can argue inductively. Assume by induction that $A_1[x] - A_i[x] = A_1[x'] - A_i[x']$ for $i < k$ with the initial case $k = 1$ established by Lemma 4.5. Let $B_k[x] = \cup_{j>k} A_k[x]$ and $B_k[x'] = \cup_{j>k} A_k[x']$. We know already that $B_k[x] - B_k[x] = B_k[x'] - B_k[x']$. Hence, by Patterson's Theorem we know that $B_k[x]^c - B_k[x]^c = B_k[x']^c - B_k[x']^c$. But these difference sets are just $\sum_{i,j \leq k} A_i[x] - A_j[x]$ and $\sum_{i,j \leq k} A_i[x'] - A_j[x']$. We a priori know that if $i, j \geq 2$ then $A_i[x] - A_j[x] = A_i[x'] - A_j[x']$. By induction, we know that $A_1[x] - A_j[x] = A_1[x'] - A_j[x']$ for $j < k$. Hence, by the pigeon-hole principle we conclude that $A_1[x] - A_k[x] = A_1[x'] - A_k[x']$ for all k . \square

5 Equivalent, Homometric and pseudo-equivalent partitions

In the conclusion to his 1944 paper [14], Patterson noted that

in very few cases are the atoms of a crystal all of one kind and it seems very probable that the presence of a second kind of atom will often resolve the ambiguities which might occur in the location of the first if taken alone.

In the following example we illustrate this phenomenon for 1-D signals.

¹Here, the notation \sum refers to the sum of multi-sets. If C_1, \dots, C_r are multi-sets, then an element $c \in \sum C_i$ appears with multiplicity equal to the sum of the multiplicities in each of the sets C_i .



Figure 2: The ordered partition $(B_1, B_2) = (\{2, 3\}, \{5, 6\})$ is shown on the left circle with points of B_1 in red and B_2 in blue. The ordered partition $(B'_1, B'_2) = (\{2, 5\}, \{6, 7\})$ is shown on the right circle with the points in B'_1 in red and B'_2 in blue. These partitions are clearly not homometric since the distance between the red points on the left is one but in the circle on the right the distance is three. However, the sets $B_1 \cup B_2 = \{2, 3, 5, 6\}$ and $B'_1 \cup B'_2 = \{2, 5, 6, 7\}$ are homometric; see Example 5.1.

Example 5.1. Consider $A = \{0, 1, 4, 7\}$ and $A' = \{0, 1, 3, 4\}$ as subsets of $[0, 7]$. These sets are homometric but not equivalent. Let $B = A^c = \{2, 3, 5, 6\}$ and $B' = (A')^c = \{2, 5, 6, 7\}$. Direct inspection or Patterson's Theorem implies that the sets B and B' are also homometric but not equivalent. In particular $B - B = B' - B' = \{0^4, 1^2, 2^2, 3, 4\}$ as multi-sets and any two binary signals supported on B, B' have the same autocorrelation.

Now, we consider the decomposition of B and B' into two subsets of size two B_1, B_2 and B'_1, B'_2 , respectively. We can now ask which of the six possible ordered partitions A, B_1, B_2 are homometric with any of the six possible partitions A', B'_1, B'_2 . Of the six possible three-set partitions of the form A, B_1, B_2 , only the partitions $(\{0, 1, 3, 4\}, \{2, 6\}, \{3, 5\})$ are homometric but not equivalent to the partitions $A', B'_1, B'_2 = (\{0, 1, 4, 7\}, \{2, 6\}, \{5, 7\})$. For example the partitions $(\{2, 3\}, \{5, 6\})$ and $(\{2, 5\}, \{6, 7\})$ are not homometric as illustrated in Figure 2.

On the other hand, if we consider partitions of B and B' into three subsets, respectively, then no partition of the form A, B_1, B_2, B_3 is homometric with a partition of the form A, B'_1, B'_2, B'_3 . This reinforces Patterson's intuition that by considering a second kind of atom we increase the likelihood that the ambiguities of the auto-correlation can be resolved.

We next consider pseudo-equivalent partitions. As far as we know, this case was not considered before.

Definition 5.2. Two partitions A_1, \dots, A_K and A'_1, \dots, A'_K of $[0, N-1]$ are *pseudo-equivalent* if there exists elements $\sigma_1, \dots, \sigma_K \in D_{2N}$ such that $A'_i = \sigma_i A_i$

When $K = 2$, any pair of pseudo-equivalent partitions are automatically equivalent because in this case if $\sigma_1 A_1 = A'_1$ then $\sigma_1 A_2 = A'_2$ since $A_2 = A_1^c$ and $A'_2 = A_1'^c$, respectively. However, in general pseudo-equivalent partitions need not be homometric and homometric partitions need not be pseudo-equivalent.

Example 5.3. The ordered partitions $\{0, 1, 4\}, \{7\}, \{3\}, \{2, 5, 6\}$ and $\{0, 1, 4\}, \{3\}, \{7\}, \{2, 5, 6\}$ are pseudo-equivalent and also homometric but not equivalent.

While pseudo equivalent partitions need not be homometric, when the sets in the partitions have approximately equal size the following numerical experiment seems to indicate that non-equivalent homometric partitions are often pseudo-equivalent.

N	Partition Sizes	Equivalent Pairs	Pseudo-equivalent pairs	Total homometric pairs
6	2,2,2,	369	0	369
7	3,2,2	1218	0	1218
8	3,3,2	2005	99	2104
9	3,3,3	813	158	971
10	4,3,3	360	12	372
11	4,4,3	148	1	149
12	4,4,4	62	3	70
13	5,4,4	10	0	10

Table 1: Results of Example 5.4

Example 5.4. We conducted the following experiment. For a given N in the range, $N = 6, \dots, 13$, we considered a set $S(N)$ of partitions where $N_1 = \lceil N/3 \rceil$, $N_2 = \lceil N - N_1/2 \rceil$, $N_3 = N_1 - N_2 - N_3$. When $N = 6, 7$ we considered all partitions and when $N \geq 8$ we took a random sample of size 300. (For $N = 6, 7$ there are fewer than 300 partitions). Each of the $\binom{|S(N)|}{2}$ pairs of partitions in $S(N)$ were tested to see if they were homometric. As indicated in Table 1, except for the case $N = 12$, all homometric pairs found were either equivalent or pseudo-equivalent.

Unfortunately, the problem of analyzing, for a particular partition type, which partitions are homometric but not equivalent seems to be an extremely difficult combinatorial problem. We conclude with the following positive result when each atom in the alphabet appears with multiplicity exactly one.

Proposition 5.5. *For any two ordered partitions of the form $(N - K, \underbrace{1, \dots, 1}_{K \text{ times}})$ are homometric if and only if they are equivalent.*

Proof. We use induction on K . For $K = 1$, any two partitions of type $(N - 1, 1)$ are necessarily equivalent because the dihedral group acts transitively on the set $[0, N - 1]$. Hence, if $(A, \{\alpha\})$ and $(A', \{\alpha'\})$ are two partitions of this form, then there exists $\sigma \in D_{2N}$ such that $\alpha' = \sigma(\alpha)$.

Assume by induction that the statement holds for partitions of the form $(A, \{\alpha_1\}, \dots, \{\alpha_K\})$ with $1 \leq K < N$. This implies that if $(\alpha'_1, \dots, \alpha'_K)$ is any sequence of distinct integers in $[0, N - 1]$ such that for all $1 \leq i < j \leq K$ it has the same difference set as $(\alpha_1, \dots, \alpha_K)$ then there exists $\sigma \in D_{2N}$ such that $\alpha'_i = \sigma\alpha_i$ for $i = 1, \dots, K$. To prove the induction step we must prove that if $(\alpha_1, \dots, \alpha_K, \alpha_{K+1})$ and $(\alpha'_1, \dots, \alpha'_K, \alpha'_{K+1})$ are two sequences with the same difference set, there exists $\sigma \in D_{2N}$ such that $\alpha'_i = \sigma\alpha_i$ for $i = 1, \dots, K+1$. By induction, there exists $\sigma_1 \in D_{2N}$ such that $\sigma_1(\alpha_1, \dots, \alpha_K) = (\alpha'_1, \dots, \alpha'_K)$. In particular, we may assume that $(\alpha'_1, \dots, \alpha'_K, \alpha'_{K+1})$ is equivalent to a sequence of the form $(\alpha_1, \dots, \alpha_K, \alpha'_{K+1})$. Applying a suitable element of D_{2N} we may also assume that $\alpha_1 = 0$. Hence, $d(\alpha_{K+1}, 0) = d(\alpha'_{K+1}, 0)$ which implies that $\alpha'_{K+1} = N - \alpha_{K+1}$ or $\alpha'_{K+1} = \alpha_{K+1}$. In the latter case we are done since the sequences would be equal. If $\alpha'_{K+1} = N - \alpha_{K+1}$ then for $i = 2, \dots, K$ we have that $d(N - \alpha_{K+1}, \alpha_i) = d(\alpha_{K+1}, \alpha_i)$. This implies that either $2\alpha_{K+1} \equiv 0 \pmod{N}$ or $2\alpha_i \equiv 0 \pmod{N}$. Since neither α_{K+1} nor α_i can be zero, we see that in either case N must be even and in the

former case we have that $\alpha_{K+1} = N/2$ so $\alpha'_{K+1} = \alpha_{K+1}$. On the other hand, the integers $\alpha_2, \dots, \alpha_K$ are distinct so we cannot have that $\alpha_i = N/2$ for $2 \leq i \leq K$ unless $K = 2$. In the case of $K = 2$, then we see that our sequences would necessarily be of the form $(0, N/2, \alpha_3)$ and $(0, N/2, N - \alpha_3)$. However, these sequences are also dihedrally equivalent since they are related by the reflection $\alpha \mapsto N - \alpha$. \square

Acknowledgment

The authors were supported by the BSF grant no. 2020159. T.B. is also supported in part by the NSF-BSF grant no. 2019752, and the ISF grant no. 1924/21 and D.E. was also supported by NSF-DMS 1906725 and NSF-DMS 2205626.

References

- [1] Tamir Bendory, Robert Beinert, and Yonina C Eldar. Fourier phase retrieval: Uniqueness and algorithms. In *Compressed Sensing and its Applications*, pages 55–91. Springer, 2017.
- [2] Tamir Bendory and Dan Edidin. Toward a mathematical theory of the crystallographic phase retrieval problem. *SIAM Journal on Mathematics of Data Science*, 2(3):809–839, 2020.
- [3] Tamir Bendory and Dan Edidin. Algebraic theory of phase retrieval. *Notices of the AMS*, 69:3489–3499, 2022.
- [4] Tamir Bendory, Oscar Mickelin, and Amit Singer. Sparse multi-reference alignment: Sample complexity and computational hardness. In *ICASSP 2022-2022 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, pages 8977–8981. IEEE, 2022.
- [5] Martin J. Buerger. Proofs and generalizations of Patterson’s theorems on homometric complementary sets. *Zeitschrift für Kristallographie - Crystalline Materials*, 143(1-6):79–98, 1976.
- [6] Veit Elser. The complexity of bit retrieval. *IEEE Transactions on Information Theory*, 64(1):412–428, 2017.
- [7] Veit Elser, Ti-Yen Lan, and Tamir Bendory. Benchmark problems for phase retrieval. *SIAM Journal on Imaging Sciences*, 11(4):2429–2455, 2018.
- [8] Subhroshekhar Ghosh and Philippe Rigollet. Sparse multi-reference alignment: Phase retrieval, uniform uncertainty principles and the beltway problem. *Foundations of Computational Mathematics*, pages 1–48, 2022.
- [9] Philipp Grohs, Sarah Koppensteiner, and Martin Rathmair. Phase retrieval: uniqueness and stability. *SIAM Review*, 62(2):301–350, 2020.

- [10] Juan E. Iglesias. On Patterson's cyclotomic sets and how to count them. *Z. Krist.*, 156(3-4):187–196, 1981.
- [11] Rick P Millane. Phase retrieval in crystallography and optics. *JOSA A*, 7(3):394–411, 1990.
- [12] A. L. Patterson. A Fourier series method for the determination of the components of interatomic distances in crystals. *Phys. Rev.*, 46:372–376, Sep 1934.
- [13] A. L. Patterson. A direct method for the determination of the components of interatomic distances in crystals. *Zeitschrift für Kristallographie - Crystalline Materials*, 90(1-6):517–542, 1935.
- [14] A. L. Patterson. Ambiguities in the X-ray analysis of crystal structures. *Phys. Rev.*, 65:195–201, Mar 1944.
- [15] Juri Ranieri, Amina Chebira, Yue M Lu, and Martin Vetterli. Phase retrieval for sparse signals: Uniqueness conditions. *arXiv preprint arXiv:1308.3058*, 2013.
- [16] Joseph Rosenblatt. Phase retrieval. *Communications in mathematical physics*, 95(3):317–343, 1984.
- [17] Joseph Rosenblatt and Paul D. Seymour. The structure of homometric sets. *SIAM J. Algebraic Discrete Methods*, 3(3):343–350, 1982.
- [18] Yoav Shechtman, Yonina C Eldar, Oren Cohen, Henry Nicholas Chapman, Jianwei Miao, and Mordechai Segev. Phase retrieval with application to optical imaging: a contemporary overview. *IEEE Signal Processing Magazine*, 32(3):87–109, 2015.