

A HYPERDETERMINANT ON FERMIONIC FOCK SPACE

FRÉDÉRIC HOLWECK AND LUKE OEDING

ABSTRACT. Twenty years ago Cayley’s hyperdeterminant, the degree four invariant of the polynomial ring $\mathbb{C}[\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2]^{\text{SL}_2(\mathbb{C})^{\times 3}}$, was popularized in modern physics as separates genuine entanglement classes in the three qubit Hilbert space and is connected to entropy formulas for special solutions of black holes. In this note we compute the analogous invariant on the fermionic Fock space for $N = 8$, i.e. spin particles with four different locations, and show how this invariant projects to other well-known invariants in quantum information. We also give combinatorial interpretations of these formulas.

1. INTRODUCTION

Arthur Cayley in 1845 [6] established several notions of hyperdeterminant as possible analogues of the classical notion of determinant but for hypermatrices. The most popular hyperdeterminant (see [9]), denoted HDet , arises by generalizing the concept of singular matrix to hypermatrix. Let $A = (a_{ijk})_{i,j,k \in \{0,1\}}$ be a real/complex $2 \times 2 \times 2$ tensor. The hyperdeterminant is:

$$\begin{aligned} \text{HDet}_{222}(A) = & a_{000}^2 a_{111}^2 + a_{010}^2 a_{101}^2 + a_{001}^2 a_{110}^2 + a_{011}^2 a_{100}^2 + 4(a_{000} a_{011} a_{101} a_{110} + a_{001} a_{010} a_{100} a_{111}) \\ & - 2(a_{000} a_{001} a_{110} a_{111} + a_{000} a_{010} a_{101} a_{111} + a_{000} a_{011} a_{100} a_{111} + \\ & a_{001} a_{010} a_{101} a_{110} + a_{001} a_{011} a_{100} a_{110} + a_{010} a_{011} a_{100} a_{101}). \end{aligned}$$

There is a combinatorial picture associated to this polynomial [5]: Consider the cube of Figure 1 labelled by the entries of the $2 \times 2 \times 2$ tensor A , the three groups of $\text{HDet}_{222}(A)$ monomials are deduced from the diagonals, parallelograms and tetrahedra inside the cube.




FIGURE 1. Monomials of Cayley’s Hyperdeterminant from a combinatorial perspective: The four diagonals (red) of the cube provide the first four monomials of degree 4 of type $a_{ijk} a_{\bar{i}\bar{j}\bar{k}}$ (where bar denotes the bit-complement), the six parallelograms (green) provide the six monomials of type $a_{ijk} a_{\bar{i}\bar{j}\bar{k}} a_{i'j'k'} a_{\bar{i}'\bar{j}'\bar{k}'}$ (where ijk and $i'j'k'$ have one bit difference), and the two tetrahedra (blue) give the two monomials of type $a_{ijk} a_{\bar{i}\bar{j}\bar{k}} a_{\bar{i}\bar{j}\bar{k}} a_{\bar{i}\bar{j}\bar{k}}$.

This polynomial gained a lot of attention in the Quantum Information literature in the early 2000's when Miyake [21] showed that this invariant is useful to distinguish the different types of genuine three-qubit entanglement. In the STU model, extremal back holes solutions with 4 electric and 4 magnetic charges have entropy formula that is the square root of Cayley's hyperdeterminant [7]. This surprising analogy was the beginning of several works on the black holes/qubits correspondence [3, 4, 15, 16, 18]. From an algebraic geometry perspective $\text{HDet} = 0$ is the equation of the dual variety X^\vee of the Segre embedding of three copies of \mathbb{P}^1 in \mathbb{P}^7 , i.e. $X = \text{Seg}(\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1) \subset \mathbb{P}(\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2)$. Recall that if $X \subset \mathbb{P}(V)$ is a projective algebraic variety, i.e. the zero set of a collection of homogeneous polynomials, then

$$X^\vee = \overline{\{H \in \mathbb{P}(V^*), \exists x \in X_{\text{smooth}}, T_x X \subset H\}}. \quad (1)$$

In other words, the dual variety of X , is the variety (in the dual projective space) of tangent hyperplanes, i.e. hyperplanes that intersect X tangentially. The bar in **1** refers to the Zariski closure, i.e. the minimal set defined by polynomial equations that contains the set. When X^\vee is a hypersurface, its defining equation, Δ_X , is called the X -discriminant [9]. When $X = \text{Seg}(\mathbb{P}^{d_1} \times \mathbb{P}^{d_2} \times \dots \times \mathbb{P}^{d_n}) \subset \mathbb{P}(\mathbb{C}^{d_1+1} \otimes \mathbb{C}^{d_2+1} \otimes \dots \otimes \mathbb{C}^{d_n+1})$ and $d_i \leq \sum_{j \neq i} d_j$ then the X -discriminant is always a hypersurface called Hyperdeterminant of format $(d_1 + 1) \times (d_2 + 1) \times \dots \times (d_n + 1)$ and denoted by $\text{HDet}_{d_1+1, d_2+1, \dots, d_n+1}$. Cayley's hyperdeterminant is the hyperdeterminant of format $2 \times 2 \times 2$.

In this paper we consider the analogue of Cayley's hyperdeterminant in the \mathcal{F}^+ component of the fermionic Fock spaces \mathcal{F}_{2N} , i.e. $\mathcal{F}_{2N} = \bigwedge^\bullet V = \bigoplus_{k \text{ even}} \bigwedge^k V \oplus \bigoplus_{k \text{ odd}} \bigwedge^k V = \mathcal{F}^+ \oplus \mathcal{F}^-$ where V is a N -dimensional single particle space. We use techniques established by the authors in [13] to compute it on a generic element of \mathcal{F}^+ in the case where the single particle state of the Fock space is 8-dimensional, spin $\frac{1}{2}$ -particle in 4 different locations, i.e. $V = \mathbb{C}^8 = \underbrace{\mathbb{C}^4}_{\text{locations}} \otimes \underbrace{\mathbb{C}^2}_{\text{Spin}}$. We show that this polynomial contains copies of the Cayley analogue for 4 fermions with 8-single particle states, the 4-qubit hyperdeterminant and the 4-bosonic qubit discriminant.

The paper is organized as follows. In Section 2 we recall the principle of the $\text{Spin}(2N)$ action on the fermionic Fock space \mathcal{F}_{2N} and why this action is the natural generalization of the SLOCC group on multiqubit Hilbert space. This allows to connect entanglement classification in fermionic Fock space and classification of spinors as explained in [17, 23]. In Section 3 we compute the equation of the dual variety of $X_{\text{Spin}(16)} \subset \mathbb{P}(\mathcal{F}^+)$ on a generic element of \mathcal{F}^+ . Our computation is based on techniques of [13] and the realization of the exceptional Lie algebra as a \mathbb{Z}_2 -graded algebra $\mathfrak{e}_8 = \mathfrak{so}(16) \oplus \mathcal{F}^+$. This polynomial of degree 240 in 8 variables has a nice combinatorial presentation as planes of a finite geometric cube space. It contains 8 copies of the degree 120 polynomial in 7 variables that measures entanglement in $\mathcal{H} = \bigwedge^4 \mathbb{C}^8$ and several copies of the four-qubit hyperdeterminant. These combinatorial pictures are provided in Section 4. Section 5 is dedicated to concluding remarks.

2. ENTANGLEMENT IN FERMIONIC FOCK SPACE

Here we recall how the fermionic Fock space of a $2n$ -single particle state Hilbert space can be seen as a Hilbert space with an SLOCC action given by the spin group $\text{Spin}(2n, \mathbb{C})$ [17, 23].

Let $\mathcal{H} = \mathbb{C}^N$ denote a Hilbert space for N -single particle states with $N = 2n$ and denote by $\mathcal{F} = \bigwedge^* \mathcal{H}$ the fermionic Fock space obtained from \mathcal{H} . We equip \mathcal{H} with a canonical basis $\{e_I\}$ and we denote by $\{e^J\}$ the basis of the dual space \mathcal{H}^* . Consider the vector space

$\mathcal{V} = \mathcal{H} \oplus \mathcal{H}^*$ where vectors are elements $x = v + \alpha$ with $v \in \mathcal{H}$ and $\alpha \in \mathcal{H}^*$. Let us equip \mathcal{V} with the quadratic form $g = \begin{pmatrix} 0 & I_N \\ I_N & 0 \end{pmatrix}$, which makes \mathcal{H} and \mathcal{H}^* orthogonal complementary subspaces in \mathcal{V} . For all $x \in \mathcal{V}$ one defines an operator acting on $\mathcal{F} = \bigwedge^* \mathcal{H}$ as follows: Consider the operators \hat{e}_I and \hat{e}^J corresponding to the exterior and interior products, i.e.

$$\begin{aligned}
 \hat{e}_I: \mathcal{F} &\rightarrow \mathcal{F} & \hat{e}^J: \mathcal{F} &\rightarrow \mathcal{F} \\
 f &\mapsto \sqrt{2}e_I \wedge f & f &\mapsto \sqrt{2}e_{J \lrcorner} f
 \end{aligned} \quad (2)$$

Then to $x = v^I e_I + \alpha_J e^J \in \mathcal{V}$ (summation over repeated indices) one can associate the operator $\mathcal{O}_x = v^I \hat{e}_I + \alpha_J \hat{e}^J$ that acts on \mathcal{F} . The operators \hat{e}_I and \hat{e}^J are respectively known as creation and annihilation operators and will be denoted from now on by p_I and n_J . Let $|0\rangle$ denote a unit vector of $\mathbb{C} = \bigwedge^0 \mathcal{H}$ defined by the property that $n_J |0\rangle = 0$, $J = 1, \dots, N$. The state $|0\rangle$ is known as the vacuum and corresponds to a state without any excitations. Then an arbitrary state of \mathcal{F} can be expressed as

$$|\Psi\rangle = \sum_{m=0}^N \sum_{I_1, \dots, I_m=1}^N \psi_{I_1, \dots, I_m}^m p_{I_1} \cdots p_{I_m} |0\rangle, \quad (3)$$

where ψ_{I_1, \dots, I_m}^m are skew-symmetric tensors. In the second quantization picture it translates the idea that quantum states may be obtained from the vacuum by excitations.

Note that the creation and annihilation operators, p_I and n_J satisfy the Canonical Anti-commutation Relations (CAR):

$$\{p_I, p_J\} = 0 \quad \{n_I, n_J\} = 0 \quad \{p_I, n_J\} = 2\delta_I^J, \quad (4)$$

where $\{\cdot, \cdot\}$ is the standard anti-commutator. From the CAR one can define a product on \mathcal{V} such that $xy = z$ for $x, y, z \in \mathcal{V}$, where z is such that $\mathcal{O}_x \mathcal{O}_y = \mathcal{O}_z$. One can easily check that given $x \in \mathcal{V}$ one has

$$x^2 = Q(x, x)\mathbf{1}, \quad (5)$$

where Q is the quadratic form on \mathcal{V} defined by g . This shows that \mathcal{V} with this product is the Clifford algebra $\mathcal{C}(\mathcal{V}, Q)$. This algebra acts on \mathcal{F} by sending $x \mapsto \mathcal{O}_x$.

Recall that the existence of g allows us to define the Lie algebra $\mathfrak{so}(\mathcal{V}, Q) = \mathfrak{so}(2N, \mathbb{C})$ as the set of matrices $s \in \mathcal{M}_{2N \times 2N}(\mathbb{C})$ such that,

$$s = \begin{pmatrix} A & B \\ C & -{}^t A \end{pmatrix}, \quad (6)$$

where B and C satisfy $B = -{}^t B$ and $C = -{}^t C$. In other words, s is a skew symmetric matrix, i.e. $\mathfrak{so}(\mathcal{V}, Q) \cong \bigwedge^2 \mathcal{V}$. The Lie algebra $\mathfrak{so}(\mathcal{V}, Q)$ can be embedded into $\mathcal{C}(\mathcal{V}, Q)$ via,

$$\begin{aligned}
 \bigwedge^2 \mathcal{V} &\rightarrow \mathcal{C}(\mathcal{V}, Q) \\
 x \wedge y &\mapsto \frac{1}{4}[x, y].
 \end{aligned} \quad (7)$$

This embedding defines a representation of $\mathfrak{so}(\mathcal{V}, Q)$ acting on \mathcal{F} . Indeed Eq. (7) maps s (from Eq. (6)) to the operator \mathcal{O}_s where

$$\mathcal{O}_s = \frac{1}{2} \sum_{i,j} A_{ij} [p_i, n_j] + B_{ij} p_i p_j + C_{ij} n_i n_j. \quad (8)$$

The action of Eq. (8) preserves the parity of the number of creations and annihilators needed to describe $|\Psi\rangle$ from the vacuum Eq. (3), i.e. one has a decomposition of \mathcal{F}_N in two irreducible representations,

$$\mathcal{F}_N = \mathcal{F}_N^+ \oplus \mathcal{F}_N^-. \quad (9)$$

Here \mathcal{F}_N^+ (resp. \mathcal{F}_N^-) denotes the fermionic states obtained by applying an even (resp. odd) number of creation operators to the vacuum.

The action of $\mathfrak{so}(\mathcal{V}, 2N)$ on \mathcal{F}_N^\pm is known in the mathematics literature as the spin representation [8]. The Lie group with Lie algebra $\mathfrak{so}(\mathcal{V}, 2N)$ acting on \mathcal{F}_N^\pm is the spin group, $\text{Spin}(2N, \mathbb{C})$, i.e. the double covering of $\text{SO}(2N, \mathbb{C})$. The Spin group has a unique closed orbit on $\mathbb{P}(\mathcal{F}_N^\pm)$ known in representation theory as the highest weight orbit or the variety of pure spinors (i.e. obtained from $|0\rangle$ by the action of the spin group). In this setting it is natural to consider as separable fermionic Fock states [23] the ones obtained from the vacuum by reversible operations of the SLOCC group $\text{Spin}(2n, \mathbb{C})$, i.e.

$$X_{\text{sep}} = \mathbb{P}(\text{Spin}(2n, \mathbb{C}) \cdot |0\rangle) \subset \mathbb{P}(\mathcal{F}_N^\pm). \quad (10)$$

In [17] it was shown that for $N = 2n$ fermionic, bosonic and multiqubit systems (n -fermions with $2n$ -single particles states, n -bosonic qubits, n -qubits) can be embedded in many ways in \mathcal{F}_N^\pm . Under these embeddings, the $\text{Spin}(2N, \mathbb{C})$ action on \mathcal{F}_N^\pm boils down to the usual SLOCC groups on the corresponding embedded Hilbert spaces ($\text{SL}(n, \mathbb{C})$, $\text{SL}(2, \mathbb{C})$, $\text{SL}(2, \mathbb{C})^{\times n}$). In this respect it is natural to look for Spin invariant polynomials in order to separate entanglement classes in fermionic Fock space.

3. THE HYPERDETERMINANT ON FERMIONIC FOCK SPACE FOR $N = 8$

We now focus on the expression of the equation of the dual of X_{sep} i.e. the analogue of Cayley's hyperdeterminant for $\text{Spin}(16, \mathbb{C})$ and \mathcal{F}_8^+ . In [13] we showed how dual equations of orbit closures a Lie group G acting on its Lie algebra \mathfrak{g} can be projected to submodules by restriction. Let us recall the principle of our construction. Let \mathfrak{g}_0 denote a subalgebra of \mathfrak{g} acting on a module \mathfrak{g}_1 such that as vector spaces we have

$$\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1. \quad (11)$$

The Lie bracket on \mathfrak{g} is compatible with the bracket on \mathfrak{g}_0 and respects the \mathbb{Z}_2 grading at (11). More specifically, the action of \mathfrak{g}_0 on \mathfrak{g}_1 defines the bracket $[\cdot, \cdot]: \mathfrak{g}_0 \times \mathfrak{g}_1 \rightarrow \mathfrak{g}_1$, and we can insist that it is skew commuting, and there is a consistent restriction of the bracket of \mathfrak{g} to \mathfrak{g}_1 i.e. $[\cdot, \cdot]: \mathfrak{g}_1 \times \mathfrak{g}_1 \rightarrow \mathfrak{g}_0$ (see [14]). Let $X_G \subset \mathbb{P}(\mathfrak{g})$ denote the unique closed G -orbit also known as the adjoint variety for the Lie group G and $Y_{G_0} \subset \mathbb{P}(\mathfrak{g}_1)$ the unique closed G_0 -orbit in $\mathbb{P}(\mathfrak{g}_1)$ known as the highest weight orbit of the \mathfrak{g}_1 -module for the Lie group G_0 . With these notations, in [13, Theorem 3.1] we showed that $Y_{G_0}^\vee \subset X_G^\vee \cap \mathbb{P}(\mathfrak{g}_1)$. In particular if $Y_{G_0}^\vee$ is a hypersurface, its defining equation $\Delta_{Y_{G_0}^\vee}$ will divide the restriction to $\mathbb{P}(\mathfrak{g}_1)$ of Δ_G , the defining equation of X_G^\vee . When the polynomials have the same degree, one gets equality.

Remark 3.1. Theorem 3.1 of [13] is in fact more general and deals with \mathbb{Z}_k -graded Lie algebras. Recently Manivel and Benedetti [20] improved our result by relating the degree relations between the polynomials $\Delta_{Y_{G_0}^\vee}$ and Δ_{X_G} with the grading of \mathfrak{g} .

Let us consider the following realization of the exceptional Lie algebra \mathfrak{e}_8 ,

$$\mathfrak{e}_8 = \mathfrak{so}(16, \mathbb{C}) \oplus \mathcal{F}_8^+. \quad (12)$$

This \mathbb{Z}_2 -branching of \mathfrak{e}_8 with $\mathfrak{g}_1 = \mathcal{F}_8^+$ was used by Antonyan and Elashvili [2] to obtain the orbit classification of spinors in \mathcal{F}_8^\pm . The bracket on $\mathfrak{so}(16, \mathbb{C})$ is the usual bracket on this Lie algebra, while the bracket $[\cdot, \cdot]: \mathfrak{g}_0 \times \mathfrak{g}_1 \rightarrow \mathfrak{g}_1$ is defined by Eq. (8). The bracket $[\cdot, \cdot]: \mathfrak{g}_1 \times \mathfrak{g}_1 \rightarrow \mathfrak{g}_0$ is described in [2, Eqs. (8) and (9)], and essentially reflects the Jordan algebra structure.

Our strategy to find an expression of X_{sep}^\vee will be to restrict an expression of $X_{E_8}^\vee$, which we need to construct. Such an expression exists on a Cartan subalgebra of \mathfrak{e}_8 , i.e. a subspace of semi-simple elements of dimension 8. Recall that the Chevalley's restriction Theorem insures that for a Lie algebra \mathfrak{g} with Cartan algebra \mathfrak{h} then $\mathbb{C}[\mathfrak{g}]^G \simeq \mathbb{C}[\mathfrak{h}]^W$ where W is the Weyl group of type G . Under this restriction, Tevelev [24] showed that when G is a simple Lie group with simply laced Dynkin diagram that

$$\Delta_G \mapsto \prod_{\alpha \in R} \alpha \in \mathbb{C}[\mathfrak{h}]^W,$$

where R is the set of roots of \mathfrak{g} . In other words, the restriction of the G -discriminant of a Lie algebra to semi-simple elements of \mathfrak{g} is the product of the roots. In order to restrict Δ_G to $\mathfrak{g}_1 = \mathcal{F}_8^+$ one needs to identify a Cartan subalgebra of \mathfrak{e}_8 living inside \mathcal{F}_8^+ . Recall that a standard choice [2] for a Cartan of \mathfrak{e}_8 is $\mathfrak{h} = \{\sum_{i=1}^8 x_i e_i \otimes e^i - e_{i+8} \otimes e^{i+8}, x_i \in \mathbb{C}\}$. For this Cartan the 240 roots of \mathfrak{e}_8 are

$$\pm x_i \pm x_j, \frac{1}{2}(\pm x_1 \pm x_2 \pm x_3 \pm x_4 \pm x_5 \pm x_6 \pm x_7 \pm x_8) \quad (13)$$

with an even number of minus signs for the second type of roots. A generic element of a Cartan subalgebra \mathfrak{c} , such that $\mathfrak{c} \subset \mathcal{F}_8^+$ can be described [17] as,

$$|\Psi\rangle = \sum_{i=1}^8 y_i |E_i\rangle, \quad (14)$$

where we have chosen the following basis of \mathfrak{c} :

$$\begin{aligned} |E_1\rangle &= (p_1 p_2 p_3 p_4 + p_{\overline{1}} p_{\overline{2}} p_{\overline{3}} p_{\overline{4}}) |0\rangle & |E_2\rangle &= (p_1 p_2 p_{\overline{3}} p_{\overline{4}} + p_{\overline{1}} p_{\overline{2}} p_3 p_4) |0\rangle \\ |E_3\rangle &= (p_1 p_{\overline{2}} p_3 p_{\overline{4}} + p_{\overline{1}} p_2 p_{\overline{3}} p_4) |0\rangle & |E_4\rangle &= (p_1 p_{\overline{2}} p_{\overline{3}} p_4 + p_{\overline{1}} p_2 p_3 p_{\overline{4}}) |0\rangle \\ |E_5\rangle &= (p_1 p_{\overline{1}} p_4 p_{\overline{4}} + p_2 p_{\overline{2}} p_3 p_{\overline{3}}) |0\rangle & |E_6\rangle &= (p_1 p_{\overline{1}} p_3 p_{\overline{3}} + p_2 p_{\overline{2}} p_4 p_{\overline{4}}) |0\rangle \\ |E_7\rangle &= (p_1 p_{\overline{1}} p_2 p_{\overline{2}} + p_2 p_{\overline{2}} p_4 p_{\overline{4}}) |0\rangle & |E_8\rangle &= (\mathbf{1} + p_1 p_2 p_3 p_4 p_{\overline{1}} p_{\overline{2}} p_{\overline{3}} p_{\overline{4}}) |0\rangle. \end{aligned} \quad (15)$$

In order to obtain the projection of Δ_G to the semi-simple algebra \mathfrak{c} one needs to express the variables x_i in terms of y_j . This can be achieved by considering [17]:

$$\begin{aligned} y_1 &= \frac{1}{2}(x_1 + x_2 + x_3 + x_4 - x_5 - x_6 - x_7 - x_8), & y_2 &= \frac{1}{2}(x_1 + x_2 - x_3 - x_4 - x_5 - x_6 + x_7 + x_8) \\ y_3 &= \frac{1}{2}(x_1 - x_2 + x_3 - x_4 - x_5 + x_6 - x_7 + x_8), & y_4 &= \frac{1}{2}(x_1 - x_2 - x_3 + x_4 - x_5 + x_6 + x_7 - x_8) \\ y_5 &= \frac{1}{2}(x_1 - x_2 - x_3 + x_4 + x_5 - x_6 - x_7 + x_8), & y_6 &= \frac{1}{2}(x_1 - x_2 + x_3 - x_4 + x_5 - x_6 + x_7 - x_8) \\ y_7 &= \frac{1}{2}(x_1 + x_2 - x_3 - x_4 + x_5 + x_6 - x_7 - x_8), & y_8 &= \frac{1}{2}(x_1 + x_2 + x_3 + x_4 + x_5 + x_6 + x_7 + x_8) \end{aligned} \quad (16)$$

The fundamental invariants can be expressed in terms of the y_i 's by inverting the relations above, writing the set of roots R in the y -basis and forming the power sums

$$f_d = \sum_{\alpha \in R} \alpha^d,$$

for d in $\{2, 8, 12, 14, 18, 20, 24, 30\}$. We did this computation in Macaulay2 [11]. Up to symmetry permuting the names of the variables and rescaling the invariant, the monomials that occur in each are listed in the appendix.

The expression of the roots at (13) after the substitution from the inverse of (16) has every root paired with its negative. As such the restriction $\Delta_{E_8|_{\mathfrak{c}}}$, which is the expression of

the $\text{HDet}_{\text{Spin}(16, \mathbb{C})}$ of \mathcal{F}_8^+ on semi-simple elements (Eq. (14)) becomes (after re-scaling)

$$\begin{aligned}
\text{HDet}_{\text{Spin}(16, \mathbb{C})}(\Psi) = & \left(y_8 y_7 y_6 y_5 y_4 y_3 y_2 y_1 \right. \\
& (y_5 - y_6 - y_7 - y_8)(y_5 - y_6 - y_7 + y_8)(y_5 - y_6 + y_7 - y_8)(y_5 - y_6 + y_7 + y_8) \\
& (y_5 + y_6 - y_7 - y_8)(y_5 + y_6 - y_7 + y_8)(y_5 + y_6 + y_7 - y_8)(y_5 + y_6 + y_7 + y_8) \\
& (y_3 - y_4 - y_7 - y_8)(y_3 - y_4 - y_7 + y_8)(y_3 - y_4 + y_7 - y_8)(y_3 - y_4 + y_7 + y_8) \\
& (y_3 - y_4 - y_5 - y_6)(y_3 - y_4 - y_5 + y_6)(y_3 - y_4 + y_5 - y_6)(y_3 - y_4 + y_5 + y_6) \\
& (y_3 + y_4 - y_7 - y_8)(y_3 + y_4 - y_7 + y_8)(y_3 + y_4 + y_7 - y_8)(y_3 + y_4 + y_7 + y_8) \\
& (y_3 + y_4 - y_5 - y_6)(y_3 + y_4 - y_5 + y_6)(y_3 + y_4 + y_5 - y_6)(y_3 + y_4 + y_5 + y_6) \\
& (y_2 - y_4 - y_6 - y_8)(y_2 - y_4 - y_6 + y_8)(y_2 - y_4 + y_6 - y_8)(y_2 - y_4 + y_6 + y_8) \\
& (y_2 - y_4 - y_5 - y_7)(y_2 - y_4 - y_5 + y_7)(y_2 - y_4 + y_5 - y_7)(y_2 - y_4 + y_5 + y_7) \\
& (y_2 + y_4 - y_6 - y_8)(y_2 + y_4 - y_6 + y_8)(y_2 + y_4 + y_6 - y_8)(y_2 + y_4 + y_6 + y_8) \\
& (y_2 + y_4 - y_5 - y_7)(y_2 + y_4 - y_5 + y_7)(y_2 + y_4 + y_5 - y_7)(y_2 + y_4 + y_5 + y_7) \\
& (y_2 - y_3 - y_6 - y_7)(y_2 - y_3 - y_6 + y_7)(y_2 - y_3 + y_6 - y_7)(y_2 - y_3 + y_6 + y_7) \\
& (y_2 - y_3 - y_5 - y_8)(y_2 - y_3 - y_5 + y_8)(y_2 - y_3 + y_5 - y_8)(y_2 - y_3 + y_5 + y_8) \\
& (y_2 + y_3 - y_6 - y_7)(y_2 + y_3 - y_6 + y_7)(y_2 + y_3 + y_6 - y_7)(y_2 + y_3 + y_6 + y_7) \\
& (y_2 + y_3 - y_5 - y_8)(y_2 + y_3 - y_5 + y_8)(y_2 + y_3 + y_5 - y_8)(y_2 + y_3 + y_5 + y_8) \\
& (y_1 - y_4 - y_6 - y_7)(y_1 - y_4 - y_6 + y_7)(y_1 - y_4 + y_6 - y_7)(y_1 - y_4 + y_6 + y_7) \\
& (y_1 - y_4 - y_5 - y_8)(y_1 - y_4 - y_5 + y_8)(y_1 - y_4 + y_5 - y_8)(y_1 - y_4 + y_5 + y_8) \\
& (y_1 + y_4 - y_6 - y_7)(y_1 + y_4 - y_6 + y_7)(y_1 + y_4 + y_6 - y_7)(y_1 + y_4 + y_6 + y_7) \\
& (y_1 + y_4 - y_5 - y_8)(y_1 + y_4 - y_5 + y_8)(y_1 + y_4 + y_5 - y_8)(y_1 + y_4 + y_5 + y_8) \\
& (y_1 - y_3 - y_6 - y_8)(y_1 - y_3 - y_6 + y_8)(y_1 - y_3 + y_6 - y_8)(y_1 - y_3 + y_6 + y_8) \\
& (y_1 - y_3 - y_5 - y_7)(y_1 - y_3 - y_5 + y_7)(y_1 - y_3 + y_5 - y_7)(y_1 - y_3 + y_5 + y_7) \\
& (y_1 + y_3 - y_6 - y_8)(y_1 + y_3 - y_6 + y_8)(y_1 + y_3 + y_6 - y_8)(y_1 + y_3 + y_6 + y_8) \\
& (y_1 + y_3 - y_5 - y_7)(y_1 + y_3 - y_5 + y_7)(y_1 + y_3 + y_5 - y_7)(y_1 + y_3 + y_5 + y_7) \\
& (y_1 - y_2 - y_7 - y_8)(y_1 - y_2 - y_7 + y_8)(y_1 - y_2 + y_7 - y_8)(y_1 - y_2 + y_7 + y_8) \\
& (y_1 - y_2 - y_5 - y_6)(y_1 - y_2 - y_5 + y_6)(y_1 - y_2 + y_5 - y_6)(y_1 - y_2 + y_5 + y_6) \\
& (y_1 - y_2 - y_3 - y_4)(y_1 - y_2 - y_3 + y_4)(y_1 - y_2 + y_3 - y_4)(y_1 - y_2 + y_3 + y_4) \\
& (y_1 + y_2 - y_7 - y_8)(y_1 + y_2 - y_7 + y_8)(y_1 + y_2 + y_7 - y_8)(y_1 + y_2 + y_7 + y_8) \\
& (y_1 + y_2 - y_5 - y_6)(y_1 + y_2 - y_5 + y_6)(y_1 + y_2 + y_5 - y_6)(y_1 + y_2 + y_5 + y_6) \\
& \left. (y_1 + y_2 - y_3 - y_4)(y_1 + y_2 - y_3 + y_4)(y_1 + y_2 + y_3 - y_4)(y_1 + y_2 + y_3 + y_4) \right)^2
\end{aligned} \tag{17}$$

We can restrict this polynomial to $\bigwedge^4 \mathcal{H} = \bigwedge^4 \mathbb{C}^8 \subset \mathcal{F}_8$. Indeed considering the seven-dimensional Cartan subalgebra spanned by $|E_1\rangle, \dots, |E_7\rangle$, the projection of $\text{HDet}_{\text{Spin}(16, \mathbb{C})}$ is obtained by dividing by y_8^2 , simplifying, and then setting $y_8 = 0$ in Eq. (17). One obtains

$$\text{HDet}_{\text{Spin}(16, \mathbb{C})| \bigwedge^4 \mathbb{C}^8}(\Psi) = Q^2 T^4 \tag{18}$$

with

$$\begin{aligned}
Q = & y_7 y_6 y_5 y_4 y_3 y_2 y_1 (y_3 - y_4 - y_5 - y_6)(y_3 - y_4 - y_5 + y_6)(y_3 - y_4 + y_5 - y_6)(y_3 - y_4 + y_5 + y_6) \\
& (y_3 + y_4 - y_5 - y_6)(y_3 + y_4 - y_5 + y_6)(y_3 + y_4 + y_5 - y_6)(y_3 + y_4 + y_5 + y_6)(y_2 - y_4 - y_5 - y_7) \\
& (y_2 - y_4 - y_5 + y_7)(y_2 - y_4 + y_5 - y_7)(y_2 - y_4 + y_5 + y_7)(y_2 + y_4 - y_5 - y_7)(y_2 + y_4 - y_5 + y_7) \\
& (y_2 + y_4 + y_5 - y_7)(y_2 + y_4 + y_5 + y_7)(y_2 - y_3 - y_6 - y_7)(y_2 - y_3 - y_6 + y_7)(y_2 - y_3 + y_6 - y_7) \\
& (y_2 - y_3 + y_6 + y_7)(y_2 + y_3 - y_6 - y_7)(y_2 + y_3 - y_6 + y_7)(y_2 + y_3 + y_6 - y_7)(y_2 + y_3 + y_6 + y_7) \\
& (y_1 - y_4 - y_6 - y_7)(y_1 - y_4 - y_6 + y_7)(y_1 - y_4 + y_6 - y_7)(y_1 - y_4 + y_6 + y_7)(y_1 + y_4 - y_6 - y_7) \\
& (y_1 + y_4 - y_6 + y_7)(y_1 + y_4 + y_6 - y_7)(y_1 + y_4 + y_6 + y_7)(y_1 - y_3 - y_5 - y_7)(y_1 - y_3 - y_5 + y_7) \\
& (y_1 - y_3 + y_5 - y_7)(y_1 - y_3 + y_5 + y_7)(y_1 + y_3 - y_5 - y_7)(y_1 + y_3 - y_5 + y_7)(y_1 + y_3 + y_5 - y_7) \\
& (y_1 + y_3 + y_5 + y_7)(y_1 - y_2 - y_5 - y_6)(y_1 - y_2 - y_5 + y_6)(y_1 - y_2 + y_5 - y_6)(y_1 - y_2 + y_5 + y_6) \\
& (y_1 - y_2 - y_3 - y_4)(y_1 - y_2 - y_3 + y_4)(y_1 - y_2 + y_3 - y_4)(y_1 - y_2 + y_3 + y_4)(y_1 + y_2 - y_5 - y_6) \\
& (y_1 + y_2 - y_5 + y_6)(y_1 + y_2 + y_5 - y_6)(y_1 + y_2 + y_5 + y_6)(y_1 + y_2 - y_3 - y_4)(y_1 + y_2 - y_3 + y_4) \\
& (y_1 + y_2 + y_3 - y_4)(y_1 + y_2 + y_3 + y_4),
\end{aligned}$$

and

$$\begin{aligned}
T = & (y_5 - y_6 - y_7)(y_5 - y_6 + y_7)(y_5 + y_6 - y_7)(y_5 + y_6 + y_7)(y_3 - y_4 - y_7)(y_3 - y_4 + y_7) \\
& (y_3 + y_4 - y_7)(y_3 + y_4 + y_7)(y_2 - y_4 - y_6)(y_2 - y_4 + y_6)(y_2 + y_4 - y_6)(y_2 + y_4 + y_6) \\
& (y_2 - y_3 - y_5)(y_2 - y_3 + y_5)(y_2 + y_3 - y_5)(y_2 + y_3 + y_5)(y_1 - y_4 - y_5)(y_1 - y_4 + y_5) \\
& (y_1 + y_4 - y_5)(y_1 + y_4 + y_5)(y_1 - y_3 - y_6)(y_1 - y_3 + y_6)(y_1 + y_3 - y_6)(y_1 + y_3 + y_6) \\
& (y_1 - y_2 - y_7)(y_1 - y_2 + y_7)(y_1 + y_2 - y_7)(y_1 + y_2 + y_7).
\end{aligned}$$

The factor Q of degree 63 in the expression of $\text{HDet}_{\text{Spin}(16, \mathbb{C})| \Lambda^4 \mathbb{C}^8}$ in Eq. (18) corresponds to the expression $\Delta_{E_7| \Lambda^4 \mathbb{C}^8}$ as computed in [13] from the grading $\mathfrak{e}_7 = \mathfrak{sl}_8 \oplus \Lambda^4 \mathbb{C}^8$ [1, 22]. This last polynomial is the restriction to semi-simple elements of the dual equation of the Grassmannian of four-planes in \mathbb{C}^8 . In the quantum information literature it would be the analogue of Cayley's hyperdeterminant for 4 fermions with 8-single particles states and as such is interesting to study fermionic entanglement. In [13] we also showed how this equation projects to the hyperdeterminant of format $2 \times 2 \times 2 \times 2$. The second factor, which has degree 28 and is named T in Eq. (18), is by construction another $\text{SL}_8(\mathbb{C})$ invariant polynomial for the module $\Lambda^4 \mathbb{C}^8$.

4. A COMBINATORIAL INTERPRETATION

The invariant polynomials $\text{HDet}_{\text{Spin}(16, \mathbb{C})}$ and $\text{HDet}_{\text{Spin}(16, \mathbb{C})| \Lambda^4 \mathbb{C}^8}$ described on semi-simple elements have nice combinatorial interpretations. Consider the cube in $(\mathbb{Z}_2)^3$ with vertices

$$\begin{aligned}
y_1 = & (0, 0, 0), & y_2 = & (0, 1, 0), & y_3 = & (1, 0, 0), & y_4 = & (1, 1, 0), \\
y_5 = & (0, 0, 1), & y_6 = & (0, 1, 1), & y_7 = & (1, 0, 1), & y_8 = & (1, 1, 1).
\end{aligned} \tag{19}$$

The cube in $(\mathbb{Z}_2)^3$ comprises 8 points, 28 lines and 14 planes. The planes are represented in Figure 2. They are planes in the sense that their defining equation is linear. For instance the tetrahedron-plane of Figure 2 is given in the (z_1, z_2, z_3) coordinates of $(\mathbb{Z}_2)^3$ by the equation $z_1 + z_2 + z_3 = 0$. The 14 planes of the cube in \mathbb{Z}_3 generate the $8 \times 14 = 112$ linear forms of Eq. (17) that involve 4 variables. Including the linear forms y_1, \dots, y_8 one gets the product of 120 linear forms (every linear form being squared to make a degree 240 polynomial). Thus we have the following formula parametrized by vertices and planes of the cube in $(\mathbb{Z}_3)^2$:

$$\text{HDet}_{\text{Spin}(16, \mathbb{C})}(\Psi) = \left(\prod_{y_i\text{-vertices}} y_i \prod_{(y_{i_1}, y_{i_2}, y_{i_3}, y_{i_4})\text{-planes}} (y_{i_1} \pm y_{i_2} \pm y_{i_3} \pm y_{i_4}) \right)^2, \tag{20}$$




FIGURE 2. Planes of the cube in $(\mathbb{Z}_2)^3$: 6 planes as the “regular” faces, 6 planes through the “opposite edges” and 2 “tetrahedron”-planes.

with the convention that $i_1 < i_2 < i_3 < i_4$.

The restriction of $\text{HDet}_{\text{Spin}(16, \mathbb{C})}$ to $\bigwedge^4 \mathbb{C}^8$ was obtained by eliminating y_8 in Eq. (17). In our combinatorial picture it could be obtained by projecting the planes of $(\mathbb{Z}_2)^3$ to the projective space $PG(2, 2)$ where the coordinate associated to y_8 is considered as the center. The 2-dimensional projective space over \mathbb{Z}_2 is well-known as the Fano-plane. The planes passing through the center will project to lines while the planes that do not pass through the center will correspond to affine planes in $PG(2, 2)$.




FIGURE 3. The Fano plane obtained by considering the projective space associated to the cube $(\mathbb{Z}_2)^3$ when the point corresponding to variable y_8 is chosen as the center. The planes of $(\mathbb{Z}_2)^3$ passing through y_8 are sent to lines and the planes that do not go through y_8 are mapped to affine plane of the Fano plane.

Therefore one can see Eq. (18) as a product of linear forms parametrized by the vertices, affine planes, and lines in the Fano plane:

$$\text{HDet}_{\text{Spin}(16, \mathbb{C})} | \bigwedge^4 \mathbb{C}^8 (\Psi) = \left(\prod_{y_i\text{-vertices}} y_i \prod_{(y_{i_1}, y_{i_2}, y_{i_3}, y_{i_4})\text{-planes}} (y_{i_1} \pm y_{i_2} \pm y_{i_3} \pm y_{i_4}) \right)^2 \left(\prod_{(y_{j_1}, y_{j_2}, y_{j_3})\text{-lines}} (y_{j_1} \pm y_{j_2} \pm y_{j_3}) \right)^4, \quad (21)$$

with the convention that $i_1 < i_2 < i_3 < i_4$ and $j_1 < j_2 < j_3$. This observation reveals an interesting connection between the Fano plane and the dual of the Grassmannian $\text{Gr}(4, 8)$.

Furthermore one can eliminate three more variables and restrict to a four-dimensional Cartan algebra. Suppose we remove the line $y_5 y_6 y_7$ of Figure (4) and we keep the collinearity relations. One gets the affine plane depicted in Figure 4. Considering the monomials given by the lines of this affine plane one gets the polynomial,

$$\Delta(y_1, y_2, y_3, y_4) = \prod_{(y_i, y_j)\text{-lines}, i < j} (y_i \pm y_j)^2. \quad (22)$$




FIGURE 4. The affine plane obtained by removing the projective line $y_5y_6y_7$.

This last equation is nothing but the expression of the $2 \times 2 \times 2 \times 2$ hyperdeterminant (see [19]) restricted to the semisimple elements of Eq. (14) for $y_5 = y_6 = y_7 = y_8 = 0$. For an understanding of the four-qubit entanglement classification based on the study of Δ one refers to [12] and for other algebraic geometry techniques for this classification see [10].

5. CONCLUSION

In this paper we evaluate for a generic element of the fermionic Fock space, the analogue of the Cayley's hyperdeterminant in the context of the spinor representation of $\text{Spin}(16, \mathbb{C})$. This work was motivated by considerations from quantum information theory as the fermionic Fock space and its spinor representation is a framework to describe many different multipartite Hilbert spaces and their corresponding SLOCC actions. In this setting algebraic invariants are interesting to compute, either to distinguish different classes of entanglement by separating the classes when the polynomials vanish or not, or by measuring some amount of entanglement when we evaluate those invariants.

It is interesting to try to express a given invariant in terms of the fundamental invariants of the corresponding representation. It is known [17] that the ring of invariants $\mathbb{C}[\mathcal{F}^\pm]^{\text{Spin}(16, \mathbb{C})}$ is generated by 8 polynomials of degree 2, 8, 12, 14, 18, 20, 24 and 30. Those polynomials can be evaluated on a generic fermionic Fock states by the same techniques as we used in this paper (See the Appendix). However an expression of $\text{HDet}_{\text{Spin}(16, \mathbb{C})}$ in terms of those invariants sounds quite difficult to obtain due to the number of monomials involved for trying a naive interpolation.

ACKNOWLEDGEMENTS

This work was partially done during the visit of FH to Auburn University during the academic year 2021-2022. FH thanks Pr. Ash Abebe and his staff of the AU Mathematics and Statistics department for making his visiting year in Auburn possible. FH was partially supported by the Conseil Régional de Bourgogne Franche-Comté (GeoQuant project). Both authors acknowledge partial support from the Thomas Jefferson foundation.

APPENDIX: FUNDAMENTAL INVARIANTS ON FERMIONIC FOCK SPACE WITH $N = 8$

One evaluates the fundamental invariants of $\mathbb{C}[\mathcal{F}_8^\pm]^{\text{Spin}(16)}$ on the generic semi-simple element given by Eq. (14). To do so, we restrict to the Cartan of the spin module, the fundamental invariants of $\mathbb{C}[\mathfrak{e}_8]^{E_8}$. Those invariants can be obtained by the Chevalley restriction Theorem (see Section 3) and the observation that the power sum expression

$$f_d = \sum_{\alpha \in R} \alpha^d, \text{ with } R \text{ is the set of roots of } E_8$$

is invariant under the action of the the Weyl group E_8 . A set of fundamental invariants of $\mathbb{C}[\mathcal{F}_8^\pm]^{\text{Spin}(16)}$ consists of homogeneous polynomials of degrees $d = 2, 8, 12, 14, 18, 20, 24, 30$ (one of each). One may obtain a set of invariants by first calculating the expression of the polynomials f_k for $k = 2, 8, 12, 14, 18, 20, 24, 30$ in terms of the x_i 's (See Eq. (13)) and then expressing those polynomials in terms of the y_i 's, i.e. restricting them to a Cartan subalgebra of \mathcal{F}_8^\pm (see Eq. (16)). It turns out that the invariant expressions obtained in this way are non-trivial and generate the invariant ring. Moreover, the fundamental invariants restricted to the y_i 's have an additional \mathfrak{S}_8 -symmetry. Respectively the invariants have 1, 7, 14, 17, 29, 38, 57, and 93 terms up to symmetry that we list below.

f_2 : y_1^2

f_8 : $6\,061\,y_1^8, 7\,196\,y_1^6 y_2^2, 17\,990\,y_1^4 y_2^4, 35\,980\,y_1^4 y_2^2 y_3^2, -28\,560\,y_1^5 y_2 y_3 y_4, -95\,200\,y_1^3 y_2^3 y_3 y_4, 215\,880\,y_1^2 y_2^2 y_3^2 y_4^2$

f_{12} : $1\,407\,661\,y_1^{12}, 270\,402\,y_1^{10} y_2^2, 2\,028\,015\,y_1^8 y_2^4, 3\,785\,628\,y_1^6 y_2^6, 4\,056\,030\,y_1^8 y_2^2 y_3^2, 18\,928\,140\,y_1^6 y_2^4 y_3^2, 47\,320\,350\,y_1^4 y_2^4 y_3^4,$
 $-1\,801\,800\,y_1^9 y_2 y_3 y_4, -21\,621\,600\,y_1^7 y_2^3 y_3 y_4, -45\,405\,360\,y_1^5 y_2^5 y_3 y_4, -151\,351\,200\,y_1^5 y_2^3 y_3^3 y_4, 113\,568\,840\,y_1^6 y_2^2 y_3^2 y_4^2,$
 $283\,922\,100\,y_1^4 y_2^4 y_3^2 y_4^2, -504\,504\,000\,y_1^3 y_2^3 y_3^3 y_4^3$

f_{14} : $1\,723\,681\,y_1^{14}, 114\,695\,y_1^{12} y_2^2, 1\,261\,645\,y_1^{10} y_2^4, 3\,784\,935\,y_1^8 y_2^6, 2\,523\,290\,y_1^{10} y_2^2 y_3^2, 18\,924\,675\,y_1^8 y_2^4 y_3^2, 35\,326\,060\,y_1^6 y_2^6 y_3^2,$
 $88\,315\,150\,y_1^6 y_2^4 y_3^4, -917\,448\,y_1^{11} y_2 y_3 y_4, -16\,819\,880\,y_1^9 y_2^3 y_3 y_4, -60\,551\,568\,y_1^7 y_2^5 y_3 y_4, -201\,838\,560\,y_1^7 y_2^3 y_3^3 y_4,$
 $-423\,860\,976\,y_1^5 y_2^5 y_3^3 y_4, 113\,548\,050\,y_1^8 y_2^2 y_3^2 y_4^2, 529\,890\,900\,y_1^6 y_2^4 y_3^2 y_4^2, 1\,324\,727\,250\,y_1^4 y_2^4 y_3^4 y_4^2, -1\,412\,869\,920\,y_1^5 y_2^3 y_3^3 y_4^3$

f_{18} : $5\,727\,234\,733\,y_1^{18}, 40\,108\,185\,y_1^{16} y_2^2, 802\,163\,700\,y_1^{14} y_2^4, 4\,866\,459\,780\,y_1^{12} y_2^6, 11\,470\,940\,910\,y_1^{10} y_2^8, 1\,604\,327\,400\,y_1^{14} y_2^2 y_3^2,$
 $24\,332\,298\,900\,y_1^{12} y_2^4 y_3^2, 107\,062\,115\,160\,y_1^{10} y_2^6 y_3^2, 172\,064\,113\,650\,y_1^8 y_2^8 y_3^2, 267\,655\,287\,900\,y_1^{10} y_2^4 y_3^4, 802\,965\,863\,700\,y_1^8 y_2^6 y_3^4,$
 $1\,498\,869\,612\,240\,y_1^6 y_2^8 y_3^6, -427\,817\,376\,y_1^{15} y_2 y_3 y_4, -14\,973\,608\,160\,y_1^{13} y_2^3 y_3 y_4, -116\,794\,143\,648\,y_1^{11} y_2^5 y_3 y_4,$
 $-305\,889\,423\,840\,y_1^9 y_2^7 y_3 y_4, -389\,313\,812\,160\,y_1^{11} y_2^3 y_3^3 y_4, -2\,141\,225\,966\,880\,y_1^9 y_2^5 y_3^3 y_4, -3\,670\,673\,086\,080\,y_1^7 y_2^7 y_3^3 y_4,$
 $-7\,708\,413\,480\,768\,y_1^7 y_2^5 y_3^5 y_4, 145\,993\,793\,400\,y_1^{12} y_2^2 y_3^2 y_4^2, 1\,605\,931\,727\,400\,y_1^{10} y_2^4 y_3^2 y_4^2, 4\,817\,795\,182\,200\,y_1^8 y_2^6 y_3^2 y_4^2,$
 $12\,044\,487\,955\,500\,y_1^8 y_2^4 y_3^4 y_4^2, 22\,483\,044\,183\,600\,y_1^6 y_2^6 y_3^4 y_4^2, -7\,137\,419\,889\,600\,y_1^9 y_2^3 y_3^3 y_4^3, -25\,694\,711\,602\,560\,y_1^7 y_2^5 y_3^3 y_4^3,$
 $-53\,958\,894\,365\,376\,y_1^5 y_2^5 y_3^3 y_4^3, 56\,207\,610\,459\,000\,y_1^6 y_2^4 y_3^4 y_4^4$

f_{20} : $91\,628\,415\,661\,y_1^{20}, 199\,229\,630\,y_1^{18} y_2^2, 5\,080\,355\,565\,y_1^{16} y_2^4, 40\,642\,844\,520\,y_1^{14} y_2^6, 132\,089\,244\,690\,y_1^{12} y_2^8,$
 $193\,730\,892\,212\,y_1^{10} y_2^{10}, 10\,160\,711\,130\,y_1^{16} y_2^2 y_3^2, 203\,214\,222\,600\,y_1^{14} y_2^4 y_3^2, 1\,232\,832\,950\,440\,y_1^{12} y_2^6 y_3^2,$
 $2\,905\,963\,383\,180\,y_1^{10} y_2^8 y_3^2, 3\,082\,082\,376\,100\,y_1^{12} y_2^4 y_3^4, 13\,561\,162\,454\,840\,y_1^{10} y_2^6 y_3^4, 21\,794\,725\,373\,850\,y_1^8 y_2^8 y_3^4,$
 $40\,683\,487\,364\,520\,y_1^8 y_2^6 y_3^6, -2\,390\,751\,000\,y_1^{17} y_2 y_3 y_4, -108\,380\,712\,000\,y_1^{15} y_2^3 y_3 y_4, -1\,137\,997\,476\,000\,y_1^{13} y_2^5 y_3 y_4,$
 $-4\,226\,847\,768\,000\,y_1^{11} y_2^7 y_3 y_4, -6\,457\,684\,090\,000\,y_1^9 y_2^9 y_3 y_4, -3\,793\,324\,920\,000\,y_1^{13} y_2^3 y_3^3 y_4,$
 $-29\,587\,934\,376\,000\,y_1^{11} y_2^5 y_3^3 y_4, -77\,492\,209\,080\,000\,y_1^9 y_2^7 y_3^3 y_4, -162\,733\,639\,068\,000\,y_1^9 y_2^5 y_3^5 y_4,$
 $-278\,971\,952\,688\,000\,y_1^7 y_2^7 y_3^5 y_4, 1\,219\,285\,335\,600\,y_1^{14} y_2^2 y_3^2 y_4^2, 18\,492\,494\,256\,600\,y_1^{12} y_2^4 y_3^2 y_4^2,$
 $81\,366\,974\,729\,040\,y_1^{10} y_2^6 y_3^2 y_4^2, 130\,768\,352\,243\,100\,y_1^8 y_2^8 y_3^2 y_4^2, 203\,417\,436\,822\,600\,y_1^{10} y_2^4 y_3^4 y_4^2,$
 $610\,252\,310\,467\,800\,y_1^8 y_2^6 y_3^4 y_4^2, 1\,139\,137\,646\,206\,560\,y_1^6 y_2^8 y_3^4 y_4^2, -98\,626\,447\,920\,000\,y_1^{11} y_2^3 y_3^3 y_4^3,$
 $-542\,445\,463\,560\,000\,y_1^9 y_2^5 y_3^3 y_4^3, -929\,906\,508\,960\,000\,y_1^7 y_2^7 y_3^3 y_4^3, -1\,952\,803\,668\,816\,000\,y_1^7 y_2^5 y_3^5 y_4^3,$
 $1\,525\,630\,776\,169\,500\,y_1^8 y_2^4 y_3^4 y_4^4, 2\,847\,844\,115\,516\,400\,y_1^6 y_2^6 y_3^4 y_4^4, -4\,100\,887\,704\,513\,600\,y_1^5 y_2^5 y_3^5 y_4^5$

$$\begin{aligned}
f_{24}: & 23\ 456\ 287\ 206\ 061\ y_1^{24}, 4\ 630\ 511\ 892\ y_1^{22}y_2^2, 178\ 274\ 707\ 842\ y_1^{20}y_2^4, 2\ 258\ 146\ 299\ 332\ y_1^{18}y_2^6, 12\ 339\ 156\ 564\ 207\ y_1^{16}y_2^8, \\
& 32\ 904\ 417\ 504\ 552\ y_1^{14}y_2^{10}, 45\ 368\ 212\ 013\ 852\ y_1^{12}y_2^{12}, 356\ 549\ 415\ 684\ y_1^{20}y_2^2y_3^2, 11\ 290\ 731\ 496\ 660\ y_1^{18}y_2^4y_3^2, \\
& 115\ 165\ 461\ 265\ 932\ y_1^{16}y_2^6y_3^2, 493\ 566\ 262\ 568\ 280\ y_1^{14}y_2^8y_3^2, 998\ 100\ 664\ 304\ 744\ y_1^{12}y_2^{10}y_3^2, 287\ 913\ 653\ 164\ 830\ y_1^{16}y_2^4y_3^4, \\
& 2\ 303\ 309\ 225\ 318\ 640\ y_1^{14}y_2^6y_3^4, 7\ 485\ 754\ 982\ 285\ 580\ y_1^{12}y_2^8y_3^4, 10\ 979\ 107\ 307\ 352\ 184\ y_1^{10}y_2^{10}y_3^4, \\
& 13\ 973\ 409\ 300\ 266\ 416\ y_1^{12}y_2^6y_3^6, 32\ 937\ 321\ 922\ 056\ 552\ y_1^{10}y_2^8y_3^6, 52\ 934\ 981\ 660\ 448\ 030\ y_1^8y_2^8y_3^6, \\
& -67\ 914\ 166\ 320\ y_1^{21}y_2y_3y_4, -4\ 753\ 991\ 642\ 400\ y_1^{19}y_2^3y_3y_4, -81\ 293\ 257\ 085\ 040\ y_1^{17}y_2^5y_3y_4, \\
& -526\ 470\ 617\ 312\ 640\ y_1^{15}y_2^7y_3y_4, -1\ 535\ 539\ 300\ 495\ 200\ y_1^{13}y_2^9y_3y_4, -2\ 177\ 673\ 917\ 065\ 920\ y_1^{11}y_2^{11}y_3y_4, \\
& -270\ 977\ 523\ 616\ 800\ y_1^{17}y_2^3y_3^3y_4, -3\ 685\ 294\ 321\ 188\ 480\ y_1^{15}y_2^5y_3^3y_4, -18\ 426\ 471\ 605\ 942\ 400\ y_1^{13}y_2^7y_3^3y_4, \\
& -39\ 924\ 021\ 812\ 875\ 200\ y_1^{11}y_2^9y_3^3y_4, -38\ 695\ 590\ 372\ 479\ 040\ y_1^{13}y_2^5y_3^5y_4, -143\ 726\ 478\ 526\ 350\ 720\ y_1^{11}y_2^7y_3^5y_4, \\
& -219\ 582\ 119\ 970\ 813\ 600\ y_1^9y_2^9y_3^3y_4, -376\ 426\ 491\ 378\ 537\ 600\ y_1^9y_2^7y_3^5y_4, 67\ 744\ 388\ 979\ 960\ y_1^{18}y_2^5y_3^2y_4^2, \\
& 1\ 727\ 481\ 918\ 988\ 980\ y_1^{16}y_2^4y_2^2y_4^2, 13\ 819\ 855\ 351\ 911\ 840\ y_1^{14}y_2^6y_2^2y_4^2, 44\ 914\ 529\ 893\ 713\ 480\ y_1^{12}y_2^8y_2^2y_4^2, \\
& 65\ 874\ 643\ 844\ 113\ 104\ y_1^{10}y_2^{10}y_2^2y_4^2, 34\ 549\ 638\ 379\ 779\ 600\ y_1^{14}y_2^4y_3^4y_4^2, 209\ 601\ 139\ 503\ 996\ 240\ y_1^{12}y_2^6y_3^4y_4^2, \\
& 494\ 059\ 828\ 830\ 848\ 280\ y_1^{10}y_2^8y_3^4y_4^2, 922\ 245\ 013\ 817\ 583\ 456\ y_1^{10}y_2^6y_3^6y_4^2, 1\ 482\ 179\ 486\ 492\ 544\ 840\ y_1^8y_2^8y_3^6y_4^2, \\
& -12\ 284\ 314\ 403\ 961\ 600\ y_1^{15}y_2^3y_3^3y_4^3, -128\ 985\ 301\ 241\ 596\ 800\ y_1^{13}y_2^5y_3^3y_4^3, -479\ 088\ 261\ 754\ 502\ 400\ y_1^{11}y_2^7y_3^3y_4^3, \\
& -731\ 940\ 399\ 902\ 712\ 000\ y_1^9y_2^7y_3^3y_4^3, -1\ 006\ 085\ 349\ 684\ 455\ 040\ y_1^{11}y_2^4y_3^4y_4^3, -2\ 634\ 985\ 439\ 649\ 763\ 200\ y_1^{10}y_2^5y_3^5y_4^3, \\
& -4\ 517\ 117\ 896\ 542\ 451\ 200\ y_1^7y_2^7y_3^3y_4^3, 524\ 002\ 848\ 759\ 990\ 600\ y_1^{12}y_2^4y_3^4y_4^3, 2\ 305\ 612\ 534\ 543\ 958\ 640\ y_1^{10}y_2^6y_3^4y_4^3, \\
& 3\ 705\ 448\ 716\ 231\ 362\ 100\ y_1^8y_2^8y_3^4y_4^3, 6\ 916\ 837\ 603\ 631\ 875\ 920\ y_1^8y_2^6y_3^6y_4^3, -5\ 533\ 469\ 423\ 264\ 502\ 720\ y_1^9y_2^5y_3^5y_4^3, \\
& -9\ 485\ 947\ 582\ 739\ 147\ 520\ y_1^7y_2^5y_3^5y_4^3, 12\ 911\ 430\ 193\ 446\ 168\ 384\ y_1^6y_2^6y_3^6y_4^3
\end{aligned}$$

$$\begin{aligned}
f_{30}: & 96\ 076\ 794\ 555\ 968\ 173\ y_1^{30}, 467\ 077\ 693\ 875\ y_1^{28}y_2^2, 29\ 425\ 894\ 714\ 125\ y_1^{26}y_2^4, 637\ 561\ 052\ 139\ 375\ y_1^{24}y_2^6, \\
& 6\ 284\ 530\ 371\ 088\ 125\ y_1^{22}y_2^8, 32\ 260\ 589\ 238\ 252\ 375\ y_1^{20}y_2^{10}, 92\ 871\ 393\ 261\ 635\ 625\ y_1^{18}y_2^{12}, 156\ 146\ 408\ 450\ 881\ 875\ y_1^{16}y_2^{14}, \\
& 58\ 851\ 789\ 428\ 250\ y_1^{26}y_2^6y_3^2, 3\ 187\ 805\ 260\ 696\ 875\ y_1^{24}y_2^4y_3^2, 58\ 655\ 616\ 796\ 822\ 500\ y_1^{22}y_2^2y_3^2, \\
& 483\ 908\ 838\ 573\ 785\ 625\ y_1^{20}y_2^8y_3^2, 2\ 043\ 170\ 651\ 755\ 983\ 750\ y_1^{18}y_2^{10}y_3^2, 4\ 736\ 441\ 056\ 343\ 416\ 875\ y_1^{16}y_2^{12}y_3^2, \\
& 6\ 245\ 856\ 338\ 035\ 275\ 000\ y_1^{14}y_2^{14}y_3^2, 146\ 639\ 041\ 992\ 056\ 250\ y_1^{22}y_2^4y_3^3, 2\ 258\ 241\ 246\ 677\ 666\ 250\ y_1^{20}y_2^6y_3^3, \\
& 15\ 323\ 779\ 888\ 169\ 878\ 125\ y_1^{18}y_2^8y_3^3, 52\ 100\ 851\ 619\ 777\ 585\ 625\ y_1^{16}y_2^{10}y_3^3, 94\ 728\ 821\ 126\ 868\ 337\ 500\ y_1^{14}y_2^{12}y_3^3, \\
& 28\ 604\ 389\ 124\ 583\ 772\ 500\ y_1^{18}y_2^6y_3^6, 156\ 302\ 554\ 859\ 332\ 756\ 875\ y_1^{16}y_2^8y_3^6, 416\ 806\ 812\ 958\ 220\ 685\ 000\ y_1^{14}y_2^{10}y_3^6, \\
& 574\ 688\ 181\ 503\ 001\ 247\ 500\ y_1^{12}y_2^{12}y_3^6, 669\ 868\ 092\ 254\ 283\ 243\ 750\ y_1^{14}y_2^8y_3^8, 1\ 354\ 622\ 142\ 114\ 217\ 226\ 250\ y_1^{12}y_2^{10}y_3^8, \\
& 1\ 986\ 779\ 141\ 767\ 518\ 598\ 500\ y_1^{10}y_2^{10}y_3^{10}, -8\ 718\ 783\ 602\ 760\ y_1^{27}y_2y_3y_4, -1\ 020\ 097\ 681\ 522\ 920\ y_1^{25}y_2^3y_3y_4, \\
& -30\ 602\ 930\ 445\ 687\ 600\ y_1^{23}y_2^5y_3y_4, -368\ 692\ 447\ 750\ 426\ 800\ y_1^{21}y_2^7y_3y_4, -2\ 150\ 705\ 945\ 210\ 823\ 000\ y_1^{19}y_2^9y_3y_4, \\
& -6\ 686\ 740\ 302\ 382\ 740\ 600\ y_1^{17}y_2^{11}y_3y_4, -11\ 658\ 931\ 809\ 282\ 727\ 200\ y_1^{15}y_2^{13}y_3y_4, -102\ 009\ 768\ 152\ 292\ 000\ y_1^{23}y_2^3y_3^3y_4, \\
& -2\ 580\ 847\ 134\ 252\ 987\ 600\ y_1^{21}y_2^5y_3^3y_4, -25\ 808\ 471\ 342\ 529\ 876\ 000\ y_1^{19}y_2^7y_3^3y_4, -122\ 590\ 238\ 877\ 016\ 911\ 000\ y_1^{17}y_2^9y_3^3y_4, \\
& -303\ 132\ 227\ 041\ 350\ 907\ 200\ y_1^{15}y_2^{11}y_3^3y_4, -408\ 062\ 613\ 324\ 895\ 452\ 000\ y_1^{13}y_2^{13}y_3^3y_4, \\
& -54\ 197\ 789\ 819\ 312\ 739\ 600\ y_1^{19}y_2^3y_3^5y_4, -441\ 324\ 859\ 957\ 260\ 879\ 600\ y_1^{17}y_2^5y_3^5y_4, \\
& -1\ 667\ 227\ 248\ 727\ 429\ 989\ 600\ y_1^{15}y_2^9y_3^5y_4, -3\ 182\ 888\ 383\ 934\ 184\ 525\ 600\ y_1^{13}y_2^{11}y_3^5y_4, \\
& -2\ 858\ 103\ 854\ 961\ 308\ 553\ 600\ y_1^{15}y_2^7y_3^7y_4, -8\ 336\ 136\ 243\ 637\ 149\ 948\ 000\ y_1^{13}y_2^9y_3^7y_4, \\
& -11\ 822\ 156\ 854\ 612\ 685\ 380\ 800\ y_1^{11}y_2^{11}y_3^7y_4, -18\ 061\ 628\ 527\ 880\ 491\ 554\ 000\ y_1^{11}y_2^9y_3^9y_4, \\
& 19\ 126\ 831\ 564\ 181\ 250\ y_1^{24}y_2^2y_3^2y_4^2, 879\ 834\ 251\ 952\ 337\ 500\ y_1^{22}y_2^4y_3^2y_4^2, \\
& 13\ 549\ 447\ 480\ 065\ 997\ 500\ y_1^{20}y_2^6y_3^2y_4^2, 91\ 942\ 679\ 329\ 019\ 268\ 750\ y_1^{18}y_2^8y_3^2y_4^2, \\
& 312\ 605\ 109\ 718\ 665\ 513\ 750\ y_1^{16}y_2^{10}y_3^2y_4^2, 568\ 372\ 926\ 761\ 210\ 025\ 000\ y_1^{14}y_2^{12}y_3^2y_4^2, \\
& 33\ 873\ 618\ 700\ 164\ 993\ 750\ y_1^{20}y_2^4y_3^4y_4^2, 429\ 065\ 836\ 868\ 756\ 587\ 500\ y_1^{18}y_2^6y_3^4y_4^2, \\
& 2\ 344\ 538\ 322\ 889\ 991\ 353\ 125\ y_1^{16}y_2^8y_3^4y_4^2, 6\ 252\ 102\ 194\ 373\ 310\ 275\ 000\ y_1^{14}y_2^{10}y_3^4y_4^2, \\
& 8\ 620\ 322\ 722\ 545\ 018\ 712\ 500\ y_1^{12}y_2^{12}y_3^4y_4^2, 4\ 376\ 471\ 536\ 061\ 317\ 192\ 500\ y_1^{16}y_2^6y_3^6y_4^2, \\
& 18\ 756\ 306\ 583\ 119\ 930\ 825\ 000\ y_1^{14}y_2^8y_3^6y_4^2, 37\ 929\ 419\ 979\ 198\ 082\ 335\ 000\ y_1^{12}y_2^{10}y_3^6y_4^2, \\
& 60\ 957\ 996\ 395\ 139\ 775\ 181\ 250\ y_1^{12}y_2^8y_3^8y_4^2, 89\ 405\ 061\ 379\ 538\ 336\ 932\ 500\ y_1^{10}y_2^{10}y_3^8y_4^2, \\
& -8\ 602\ 823\ 780\ 843\ 292\ 000\ y_1^{21}y_2^3y_3^3y_4^3, -180\ 659\ 299\ 397\ 709\ 132\ 000\ y_1^{19}y_2^5y_3^3y_4^3, \\
& -1\ 471\ 082\ 866\ 524\ 202\ 932\ 000\ y_1^{17}y_2^7y_3^3y_4^3, -5\ 557\ 424\ 162\ 424\ 766\ 632\ 000\ y_1^{15}y_2^9y_3^3y_4^3, \\
& -10\ 609\ 627\ 946\ 447\ 281\ 752\ 000\ y_1^{13}y_2^{11}y_3^3y_4^3, -3\ 089\ 274\ 019\ 700\ 826\ 157\ 200\ y_1^{17}y_2^5y_3^5y_4^3, \\
& -20\ 006\ 726\ 984\ 729\ 159\ 875\ 200\ y_1^{15}y_2^7y_3^5y_4^3, -58\ 352\ 953\ 705\ 460\ 049\ 636\ 000\ y_1^{13}y_2^9y_3^5y_4^3, \\
& -82\ 755\ 097\ 982\ 288\ 797\ 665\ 600\ y_1^{11}y_2^{11}y_3^5y_4^3, -100\ 033\ 634\ 923\ 645\ 799\ 376\ 000\ y_1^{13}y_2^7y_3^7y_4^3, \\
& -216\ 739\ 542\ 334\ 565\ 898\ 648\ 000\ y_1^{11}y_2^9y_3^7y_4^3, -331\ 129\ 856\ 344\ 475\ 678\ 490\ 000\ y_1^9y_2^9y_3^9y_4^3, \\
& 1\ 072\ 664\ 592\ 171\ 891\ 468\ 750\ y_1^{18}y_2^4y_3^4y_4^4, 10\ 941\ 178\ 840\ 153\ 292\ 981\ 250\ y_1^{16}y_2^6y_3^4y_4^4, \\
& 46\ 890\ 766\ 457\ 799\ 827\ 062\ 500\ y_1^{14}y_2^8y_3^4y_4^4, 94\ 823\ 549\ 947\ 995\ 205\ 837\ 500\ y_1^{12}y_2^{10}y_3^4y_4^4, \\
& 87\ 529\ 430\ 721\ 226\ 343\ 850\ 000\ y_1^{14}y_2^6y_3^6y_4^4, 284\ 470\ 649\ 843\ 985\ 617\ 512\ 500\ y_1^{12}y_2^8y_3^6y_4^4, \\
& 417\ 223\ 619\ 771\ 178\ 905\ 685\ 000\ y_1^{10}y_2^{10}y_3^6y_4^4, 670\ 537\ 960\ 346\ 537\ 526\ 993\ 750\ y_1^{10}y_2^8y_3^8y_4^4, \\
& -42\ 014\ 126\ 667\ 931\ 235\ 737\ 920\ y_1^{15}y_2^5y_3^5y_4^5, -210\ 070\ 633\ 339\ 656\ 178\ 689\ 600\ y_1^{13}y_2^7y_3^5y_4^5, \\
& -455\ 153\ 038\ 902\ 588\ 387\ 160\ 800\ y_1^{11}y_2^9y_3^5y_4^5, -780\ 262\ 352\ 404\ 437\ 235\ 132\ 800\ y_1^{11}y_2^7y_3^7y_4^5, \\
& -1\ 192\ 067\ 482\ 840\ 112\ 442\ 564\ 000\ y_1^9y_2^9y_3^7y_4^5, 531\ 011\ 879\ 708\ 773\ 152\ 690\ 000\ y_1^{12}y_2^6y_3^6y_4^6, \\
& 1\ 251\ 670\ 859\ 313\ 536\ 717\ 055\ 000\ y_1^{10}y_2^8y_3^6y_4^6, 2\ 011\ 613\ 881\ 039\ 612\ 580\ 981\ 250\ y_1^8y_2^8y_3^6y_4^6, \\
& -2\ 043\ 544\ 256\ 297\ 335\ 615\ 824\ 000\ y_1^9y_2^7y_3^7y_4^6
\end{aligned}$$

REFERENCES

- [1] L. Antonyan, *Classification of four-vectors of an eight-dimensional space*, Trudy Sem. Vektor. Tenzor. Anal **20** (1981), no. 38, 144–161.
- [2] L. Antonyan and A. Elashvili, *Classification of spinors of dimension sixteen*, Trudy Tbiliss. Mat. Inst. Razmadze Akad. Nauk Gruzin. SSR **70** (1982), 5–23.
- [3] L. Borsten, D. Dahanayake, M. J Duff, H. Ebrahim, and W. Rubens, *Black holes, qubits and octonions*, Physics Reports **471** (2009), no. 3-4, 113–219.
- [4] L. Borsten, M. J Duff, and P. Lévy, *The black-hole/qubit correspondence: an up-to-date review*, Classical and Quantum Gravity **29** (2012), no. 22, 224008.
- [5] M. R Bremner, M. G Bickis, and M. Soltanifar, *Cayley’s hyperdeterminant: a combinatorial approach via representation theory*, Linear algebra and its applications **437** (2012), no. 1, 94–112.
- [6] A. Cayley, *On the theory of linear transformations*, Metcalfe and Palmer, 1800.
- [7] M. J Duff, *String triality, black hole entropy, and Cayley’s hyperdeterminant*, Physical Review D **76** (2007), no. 2, 025017.
- [8] W. Fulton and J. Harris, *Representation theory: a first course*, Vol. 129, Springer Science & Business Media, 2013.
- [9] I. Gelfand, M. Kapranov, and A Zelevinsky, *Discriminants, resultants, and multidimensional determinants* (M. Birkhäuser Boston, ed.), 1994.
- [10] M. Gharahi, S. Mancini, and G. Ottaviani, *Fine-structure classification of multiqubit entanglement by algebraic geometry*, Physical Review Research **2** (2020), no. 4, 043003.
- [11] D. R. Grayson and M. E. Stillman, *Macaulay2, a software system for research in algebraic geometry*.
- [12] F. Holweck, J.-G. Luque, and J.-Y. Thibon, *Entanglement of four-qubit systems: a geometric atlas with polynomial compass II (the tame world)*, Journal of Mathematical Physics **58** (2017), no. 2, 022201.
- [13] F. Holweck and L. Oeding, *Hyperdeterminants from the e_8 discriminant*, Journal of Algebra **593** (2022), 622–650.
- [14] F. Holweck and L. Oeding, *Jordan decompositions of tensors*, arXiv preprint arXiv:2206.13662 (2022).
- [15] R. Kallosh and A. Linde, *Strings, black holes, and quantum information*, Physical Review D **73** (2006), no. 10, 104033.
- [16] P. Lévy, *Stringy black holes and the geometry of entanglement*, Physical Review D **74** (2006), no. 2, 024030.
- [17] P. Lévy and F. Holweck, *Embedding qubits into Fermionic Fock space: Peculiarities of the four-qubit case*, Physical Review D **91** (2015), no. 12, 125029.
- [18] P. Lévy, M. Saniga, and P. Vrana, *Three-qubit operators, the split Cayley hexagon of order two, and black holes*, Physical Review D **78** (2008), no. 12, 124022.
- [19] J.-G. Luque and J.-Y. Thibon, *Polynomial invariants of four qubits*, Physical Review A **67** (2003), no. 4, 042303.
- [20] L. Manivel and V. Benedetti, *On the automorphisms of hyperplane sections of generalized Grassmannians*, Transformation Groups (2022), 1–45.
- [21] A. Miyake, *Classification of multipartite entangled states by multidimensional determinants*, Physical Review A **67** (2003), no. 1, 012108.
- [22] L. Oeding, *A Translation of “Classification of four-vectors of an 8-dimensional space,” by Antonyan, L. V., with an appendix by the translator*, to appear volume 83 (2022) of “Trudy Moskovskogo matematicheskogo obshchestva” (2022). arXiv:2205.09741.
- [23] G. Sárosi and P. Lévy, *Entanglement in Fermionic Fock space*, Journal of Physics A: Mathematical and Theoretical **47** (2014), no. 11, 115304.
- [24] E. A. Tevelev, *Projectively dual varieties*, Journal of Mathematical Sciences **117** (2003), no. 6, 4585–4732.

Email address: frederic.holweck@utbm.fr

LABORATOIRE INTERDISCIPLINAIRE CARNOT DE BOURGOGNE, ICB/UTBM, UMR 6303 CNRS, UNIVERSITÉ BOURGOGNE FRANCHE-COMTÉ, 90010 BELFORT CEDEX, FRANCE

Email address: oeding@auburn.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, AUBURN UNIVERSITY, AUBURN, AL, USA