

# BIRKHOFF GENERIC POINTS ON CURVES IN HOROSPHERES

OMRI SOLAN AND ANDREAS WIESER

ABSTRACT. Let  $\{a_t : t \in \mathbb{R}\} < \mathrm{SL}_d(\mathbb{R})$  be a diagonalizable subgroup whose expanding horospherical subgroup  $U < \mathrm{SL}_d(\mathbb{R})$  is abelian. By the Birkhoff ergodic theorem, for any  $x \in \mathrm{SL}_d(\mathbb{R})/\mathrm{SL}_d(\mathbb{Z})$  and for almost every point  $u \in U$  the point  $ux$  is Birkhoff generic for  $a_t$  when  $t \rightarrow \infty$ . We prove that the same is true when  $U$  is replaced by any non-degenerate analytic curve in  $U$ .

This Birkhoff genericity result has various applications in Diophantine approximation. For instance, we obtain density estimates for Dirichlet improvability along typical points on a curve in Euclidean space. Other applications address approximations by algebraic numbers and best approximations (in the sense of Lagarias).

## CONTENTS

|  |    |
|--|----|
| 1. Introduction  | 2  |
| 1.1. Dirichlet improvability and densities                 | 2  |
| 1.2. Best approximations of points on curves               | 3  |
| 1.3. Expanded translates of analytic curves in horospheres | 4  |
| 1.4. Birkhoff genericity and the main result               | 5  |
| 1.5. Ingredients of the proof                              | 6  |
| 1.6. Further research                                      | 7  |
| 1.7. Structure of the article                              | 8  |
| 2. Notation and preliminaries                              | 8  |
| 2.1. $(C, \alpha)$ -good functions                         | 10 |
| 3. Long intervals  | 12 |
| 3.1. Contraction for long intervals with short vectors     | 12 |
| 3.2. Atypicality of long intervals with short vectors      | 14 |
| 4. The Short Intervals Lemma                               | 17 |
| 5. Quantitative non-divergence                             | 20 |
| 6. Unipotent invariance                                    | 23 |
| 7. Avoiding singular sets and a proof of Theorem 1.10      | 25 |
| 7.1. The linearization technique                           | 25 |
| 8. Proof of the applications                               | 28 |
| 8.1. Proof of Theorem 1.1                                  | 28 |
| 8.2. Proof of Theorem 1.4                                  | 28 |
| 8.3. Proof of Theorem 1.6                                  | 33 |
| References   | 35 |

---

*Date:* January 26, 2023.

The first named author is funded by the generous donation of Dr. Arthur A. Kaselemas. The second named author is supported by the ERC grant HomDyn, ID 833423.

## 1. INTRODUCTION

**1.1. Dirichlet improvability and densities.** Let  $\xi = (\xi_1, \dots, \xi_d) \in \mathbb{R}^d$  be a vector. By Dirichlet's theorem on simultaneous approximations, the set of inequalities

$$|p_1 + q_1\xi_1 + \dots + q_d\xi_d| \leq N^{-d}, \quad \max_i |q_i| \leq N$$

has a non-zero solution  $(p, q_1, \dots, q_d) \in \mathbb{Z}^{d+1}$  for all  $N \in \mathbb{N}$ . We say that  $\xi$  is not  $\mu$ -Dirichlet improvable for  $\mu \in (0, 1)$  if there exist infinitely many  $N \in \mathbb{N}$  for which the inequalities

$$(1.1) \quad |p_1 + q_1\xi_1 + \dots + q_d\xi_d| \leq \mu N^{-d}, \quad \max_i |q_i| \leq \mu N$$

have no non-zero solution  $(p, q_1, \dots, q_d) \in \mathbb{Z}^{d+1}$ . Davenport and Schmidt [9] proved that almost every  $\xi \in \mathbb{R}^d$  is not  $\mu$ -Dirichlet improvable for any  $\mu < 1$  (in the language of homogeneous dynamics, this also follows from the Dani correspondence [7] and recurrence).

Now let  $\phi : [0, 1] \rightarrow \mathbb{R}^d$  be a curve. If the image of  $\phi$  is not contained in an affine hyperplane and  $\phi$  is analytic, Shah [32] proved that for almost every  $s$  the vector  $\phi(s)$  is not  $\mu$ -Dirichlet improvable for any  $\mu < 1$ . Among other results, we propose here a strengthening of Shah's result describing the logarithmic density of exceptions to Dirichlet improvability. Define for any  $\xi \in \mathbb{R}^d$  and  $\mu \in (0, 1)$

$$\begin{aligned} \overline{\mathcal{D}}_\xi(\mu) &= \limsup_{N' \rightarrow \infty} \frac{1}{\log(N')} \sum_{\substack{N \leq N', \\ (1.1) \text{ holds for } N, \xi}} \frac{1}{N}, \\ \underline{\mathcal{D}}_\xi(\mu) &= \liminf_{N' \rightarrow \infty} \frac{1}{\log(N')} \sum_{\substack{N \leq N', \\ (1.1) \text{ holds for } N, \xi}} \frac{1}{N}. \end{aligned}$$

**Theorem 1.1** (Density in Dirichlet improvability). *There exists an explicit continuous strictly increasing function  $f : [0, 1] \rightarrow [0, 1]$  with  $f(0) = 0$  and  $f(1) = 1$  with the following properties. Suppose that  $\phi : [0, 1] \rightarrow \mathbb{R}^d$  is an analytic curve not contained in any affine hyperplane. Then for every  $\mu \in (0, 1)$  and for almost every  $s \in [0, 1]$  we have*

$$\overline{\mathcal{D}}_{\phi(s)}(\mu) = \underline{\mathcal{D}}_{\phi(s)}(\mu) = f(\mu).$$

*Remark 1.2.* It is a simple consequence of the Birkhoff ergodic theorem that there is a function  $f$  as above so that for every  $\mu \in (0, 1)$  and for almost every  $\xi \in \mathbb{R}^d$  we have  $\overline{\mathcal{D}}_\xi(\mu) = \underline{\mathcal{D}}_\xi(\mu) = f(\mu)$ . This function  $f$  coincides with the one in Theorem 1.1 and is given by the measure of a parametrized family of neighborhoods of the cusp in  $\mathrm{SL}_{d+1}(\mathbb{R})/\mathrm{SL}_{d+1}(\mathbb{Z})$ . The speed of convergence  $f(\mu) \rightarrow 1$  for  $\mu \rightarrow 1$  has been studied in recent papers by Kleinbock and Yu [19] for  $d = 1$  and Kleinbock, Strömbergsson, Yu [18] for  $d > 1$ . The decay of  $f(\mu)$  for  $\mu \rightarrow 0$  is easily established via Siegel's formula [39].

*Remark 1.3.* The methods in Theorem 1.1 yield similar results for Diophantine approximation of matrices (see e.g. [29, 41]).

We turn now to algebraic approximations: The *polynomial height*  $\|\alpha\|$  of an algebraic number  $\alpha \in \mathbb{R}$  is defined to be the height  $\mathrm{ht}(\cdot)$  of its minimal polynomial, that is, the maximal absolute value of the coefficients for the minimal integer multiple

of the minimal polynomial. From Theorem 1.1, one obtains the logarithmic density of solutions to

$$P(s) \leq \mu N^{-d}, \quad P \in \mathbb{Z}[x] \text{ of degree } d \text{ with } \text{ht}(P) \leq \mu N$$

for almost every  $s \in [0, 1]$ .

Consider now the following problem. Given  $s \in \mathbb{R}$ ,  $\mu > 0$ , and  $N \in \mathbb{N}$  does there exist an algebraic number  $s' \in \mathbb{R}$  of degree  $d$  with

$$(1.2) \quad |s' - s| \leq \mu N^{-(d+1)} \text{ and } \|s'\| \leq N?$$

Define

$$\begin{aligned} \overline{\mathcal{A}}_s(\mu) &= \limsup_{N' \rightarrow \infty} \frac{1}{\log(N')} \sum_{\substack{N \leq N', \\ (1.2) \text{ holds for } N, s}} \frac{1}{N}, \\ \underline{\mathcal{A}}_s(\mu) &= \liminf_{N' \rightarrow \infty} \frac{1}{\log(N')} \sum_{\substack{N \leq N', \\ (1.2) \text{ holds for } N, s}} \frac{1}{N}. \end{aligned}$$

**Theorem 1.4** (Algebraic approximations). *There is an explicit continuous function  $f : \mathbb{R}_{>0} \times \mathbb{R} \rightarrow (0, 1)$  which is strictly increasing in the first coordinate with the following properties. Then for every  $\mu \in (0, 1)$  and for almost every  $s \in \mathbb{R}$  we have*

$$\overline{\mathcal{A}}_s(\mu) = \underline{\mathcal{A}}_s(\mu) = f(\mu, s).$$

**1.2. Best approximations of points on curves.** Fix a norm  $\|\cdot\|$  on  $\mathbb{R}^d$  for  $d \geq 2$  and let  $\xi \in \mathbb{R}^d$ . Following Lagarias [21, 22] we say that a vector  $(p, q) \in \mathbb{Z}^d \times \mathbb{N}$  is a best approximation of  $\xi$  if

$$\|p - q\xi\| < \min_{q' < q} \min_{p' \in \mathbb{Z}^d} \|p' - q'\xi\|.$$

If  $\xi$  has an irrational coordinate, one obtains by recursion an infinite sequence  $v_k = (p_k, q_k)$  of best approximations of  $\xi$ . In a recent article [36], Shapira and Weiss obtain various new results about statistical properties of the best approximations of Lebesgue almost every  $\xi$ , some of which we now recall.

Define for any  $v = (p, q) \in \mathbb{Z}^d \times \mathbb{N}$

$$\text{disp}(\xi, v) = q^{\frac{1}{d}}(p - q\xi).$$

By Dirichlet's theorem, there exists a constant  $C > 0$  depending only on the norm  $\|\cdot\|$  so that  $\|\text{disp}(\xi, v)\| \leq C$  whenever  $v$  is a best approximation of  $\xi$ . Moreover, we may associate a certain sequence of lattices with the best approximations. To that end, we identify  $\mathbb{R}^d \simeq \{x \in \mathbb{R}^{d+1} : x_{d+1} = 0\}$  and denote for any  $v \in \mathbb{R}^{d+1} \setminus \mathbb{R}^d$  by  $\pi^v : \mathbb{R}^{d+1} \rightarrow \mathbb{R}^d$  the projection parallel to  $v$ . For  $v \in \mathbb{Z}^d \times \mathbb{N}$  the projection  $\Lambda_v = \pi^v(\mathbb{Z}^{d+1})$  defines a lattice in  $\mathbb{R}^d$  and hence a point

$$[\Lambda_v] \in X_d = \text{SL}_d(\mathbb{R})/\text{SL}_d(\mathbb{Z})$$

after dilation (not to be confused with the shape of the lattice  $\Lambda_v$ ).

**Theorem 1.5** (Shapira, Weiss [36, Thm. 1.1]). *For any norm  $\|\cdot\|$  on  $\mathbb{R}^d$  there exists a Borel probability measure  $\mu_{d, \|\cdot\|}$  on  $X_d \times \mathbb{R}^d$  such that for Lebesgue almost  $\xi \in \mathbb{R}^d$  the following holds. Let  $(v_k)$  be the sequence of best approximations of  $\xi$ . Then the sequence*

$$\{([\Lambda_{v_k}], \text{disp}(\xi, v_k))\} \subset X_d \times \mathbb{R}^d$$

is equidistributed with respect to  $\mu_{d,\|\cdot\|}$ .

The measure  $\mu_{d,\|\cdot\|}$  is further characterized in [36] and is notably not a product measure. The projection of  $\mu_{d,\|\cdot\|}$  to  $\mathbb{R}^d$  is boundedly supported and absolutely continuous with respect to Lebesgue with a non-trivial density. We refer to [36] for a more precise description of the measure  $\mu_{d,\|\cdot\|}$  and for various extensions of the above theorem. We remark that similar questions to Theorem 1.5 had been studied previously – cf. [36, §3] for a discussion.

As a consequence of the Birkhoff genericity results in the current article, we obtain results for the best approximations of points on curves.

**Theorem 1.6** (Best approximations of points on curves). *Let  $\phi : [0, 1] \rightarrow \mathbb{R}^d$  be an analytic curve not contained in any affine hyperplane. Then for almost every  $s \in [0, 1]$  the following holds. Let  $(v_k)$  be the sequence of best approximations of  $\phi(s)$ . Then the sequence*

$$\{([\Lambda_{v_k}], \text{disp}(\phi(s), v_k))\} \subset X_d \times \mathbb{R}^d$$

is equidistributed with respect to  $\mu_{d,\|\cdot\|}$ .

A similar result can be obtained for local properties of the best approximations (e.g. the statistics of  $v_k \bmod M$  for any  $M \geq 1$ ).

**1.3. Expanded translates of analytic curves in horospheres.** Before turning to our main results, we briefly recall existing work on expanding translates of curves in horospheres. For concreteness, let  $G = \text{SL}_d(\mathbb{R})$  and let  $\{a_t\} < G$  be a diagonalizable subgroup. The expanding horospherical subgroup is then given by

$$G^+ = \{g \in G : a_t g a_{-t} \rightarrow 1 \text{ as } t \rightarrow -\infty\}.$$

A long line of research has established variants of equidistribution of expanded pieces of  $G^+$ -orbits in  $G/\Gamma$  for  $\Gamma < G$  a lattice. Here, equidistribution may for example be deduced by Margulis' thickening technique as well as mixing and can be made effective; this idea originated in Margulis' thesis [23] (see also [15, Prop. 2.4.8]). Motivated by this, one may ask under what conditions translates of a curve

$$\varphi : [0, 1] \rightarrow G^+$$

(or more generally a submanifold) would equidistribute [34]. In his influential work [32], Shah resolved this question when  $a_t = \text{diag}(e^{(d-1)t}, e^{-t}, \dots, e^{-t})$  and when the curve  $\varphi$  is analytic and is not contained in any proper subgroup of  $G^+$  (see also [33, 31]). Subsequently, Shah and L. Yang [29] and finally P. Yang [41] generalized Shah's result to all diagonalizable subgroups. Given the objectives of the current article, we shall state P. Yang's result [41] here in the case

$$(1.3) \quad a_t = \text{diag}(e^{t/m}, \dots, e^{t/m}, e^{-t/n}, \dots, e^{-t/n}) \in G$$

where  $m + n = d$ . In this case, obstructions to equidistribution may be described via constraining pencils introduced by Aka, Breuillard, de Saxcé, and Rosenzweig [1]. Let  $P < G$  be the parabolic group given by

$$(1.4) \quad P = \{g \in G : \lim_{t \rightarrow \infty} a_t g a_{-t} \text{ exists}\}$$

and note that we may identify  $P \backslash G \simeq \text{Gr}(m, d)$ .

**Definition 1.7.** For an integer  $r \leq m$  and a proper subspace  $W \subset \mathbb{R}^d$  the pencil  $\mathfrak{P}_{W,r}$  is

$$\mathfrak{P}_{W,r} = \{V \in \text{Gr}(m, d) : \dim(V \cap W) \geq r\}.$$

The pencil is constraining if  $\frac{\dim(W)}{r} < \frac{d}{m}$  and weakly constraining if  $\frac{\dim(W)}{r} \leq \frac{d}{m}$ .

For example, when  $m = 1$  the pencil  $\mathfrak{P}_{W,1}$  consists of all lines contained in  $W$  and is always constraining. If  $m = n = 2$  the pencil  $\mathfrak{P}_{W,r}$  is weakly constraining if  $\dim(W) = 2$  and  $r = 1$ , constraining if  $\dim(W) \in \{1, 2, 3\}$  and  $r = 2$ , and not constraining if  $\dim(W) = 3$  and  $r = 1$ .

**Definition 1.8.** A partial flag subvariety of  $P \backslash G$  (for  $\{a_t\}$ ) is a subvariety of the form  $PHg$  where  $H < G$  is reductive group containing  $\{a_t\}$  and where  $g \in G$ .

**Theorem 1.9** (P. Yang [41]). *Let  $\Gamma < G$  be a lattice. Let  $\{a_t : t \in \mathbb{R}\}$  be as in (1.3) and  $\varphi : [0, 1] \rightarrow G$  be an analytic curve whose image in  $P \backslash G$  is not contained in a weakly constraining pencil or in a partial flag subvariety. Then for any  $x \in G/\Gamma$  and any  $f \in C_c(G/\Gamma)$*

$$\int_0^1 f(a_t \varphi(s)x) ds \rightarrow \int_{G/\Gamma} f d\mu_{G/\Gamma} \quad (t \rightarrow \infty).$$

The same conclusion was obtained in earlier work by Shah and L. Yang [29] under a stronger assumption called supergenericity. The conditions on  $\varphi$  are not to be considered sharp: Suppose that  $m = 1$  in which case  $G^+ \simeq \mathbb{R}^{d-1}$  and the above conditions for  $\varphi : [0, 1] \rightarrow G^+$  translate to the image of  $\varphi$  not being contained in any affine hyperplane; Theorem 1.9 in this case is due to Shah [32]. For  $d = 3$  work by Shi and Weiss [38] (with an additional average over  $t$ ) and Kleinbock, de Saxcé, Shah, and P. Yang [16] addresses conditions under which translates of planar lines equidistribute. Similar results exist for  $d > 3$  [30]. In the case  $m = 1$ , the analyticity assumption has been weakened recently by Shah and P. Yang [35].

**1.4. Birkhoff genericity and the main result.** In this article, we establish that almost every point on a curve in an expanding horosphere is Birkhoff generic for the diagonalizable flow under suitable conditions.

In view of Birkhoff's ergodic theorem, one may ask whether for an ergodic measure preserving system  $X = (X, \mathcal{B}, \mu, T)$  the ergodic averages  $\frac{1}{N} \sum_{n=1}^N f(T^n x)$  converge for almost every point  $x$  when the point  $x$  is sampled with respect to a different measure. Assuming that  $X$  is locally compact  $\sigma$ -compact, we say that a point  $x$  is Birkhoff generic (with respect to  $\mu$  and  $T$ ) if

$$\frac{1}{N} \sum_{n=1}^N f(T^n x) \rightarrow \int f d\mu$$

for any  $f \in C_c(X)$ . By the Birkhoff ergodic theorem,  $\mu$ -almost every point is Birkhoff generic with respect to  $\mu$ . The above question for  $X = \mathbb{T} = \mathbb{R}/\mathbb{Z}$  has been studied by various authors including Host [13] and Hochman and Shmerkin [12]. Host established that  $\nu$ -almost every point is Birkhoff generic for  $\times 2$  when  $\nu$  is a  $\times 3$  invariant and ergodic measure  $\nu$  on  $\mathbb{T}$  of positive entropy. For example, almost any point on the middle-third Cantor set is Birkhoff generic for  $\times 2$  i.e. normal in base 2 [28, 5].

Now let  $\{a_t : t \in \mathbb{R}\}$  be as in (1.3) and let  $G^+$  be the expanding horospherical subgroup. Birkhoff genericity for fractal measures on  $G^+$  is known in many cases – see [10, 40, 25, 26]. For certain subgroups of  $G^+$  (and for general diagonalizable subgroups), Shi [37] established Birkhoff genericity. Moreover, Fraczek, Shi, and Ulcigrai [11] and Zhang [42] proved Birkhoff genericity for certain submanifolds of the full horosphere in  $\mathrm{SL}_d(\mathbb{R}) \times (\mathbb{R}^d)^k$  which project surjectively onto the full horosphere in  $\mathrm{SL}_d(\mathbb{R})$ . All of the works [37, 11, 42] are based on previous work of Chaika and Eskin [6].

In the current article, we establish the following Birkhoff genericity result.

**Theorem 1.10** (Birkhoff genericity along curves). *Let  $G = \mathrm{SL}_d(\mathbb{R})$ ,  $\Gamma = \mathrm{SL}_d(\mathbb{Z})$ ,  $X = G/\Gamma$ ,  $\{a_t : t \in \mathbb{R}\}$  as in (1.3), and  $P$  the associated parabolic subgroup as in (1.4). Let  $\varphi : [0, 1] \rightarrow G$  be an analytic curve whose image in  $P \backslash G$  is not contained in any weakly constraining pencil or in any partial flag subvariety. Then for any  $x \in X$  and for almost every  $s \in [0, 1]$  the point  $\varphi(s)x$  is Birkhoff generic with respect to the Haar probability measure on  $X$  and the flow  $\{a_t : t > 0\}$  i.e. for every  $f \in C_c(X)$*

$$\frac{1}{T} \int_0^T f(a_t \varphi(s)x) dt \rightarrow \int_X f dm_X.$$

*Remark 1.11.* The non-degeneracy assumptions on the curve  $\varphi$  in the case  $m = 1$  in either of the above results translates to assuming that the image of  $\varphi$  in  $G/P \simeq \mathbb{P}^d$  is not contained in any proper subspace.

*Remark 1.12.* We do not expect the conditions in either of the above theorems to be sharp and refer to §1.6 for a discussion. In the upcoming work of Shah and P. Yang Theorem 1.10 for degenerate curves is established in some important cases.

*Remark 1.13.* The lattice  $\mathrm{SL}_d(\mathbb{Z})$  may be replaced by a congruence lattice  $\Gamma < \mathrm{SL}_d(\mathbb{Z})$  in the above without difficulty.

**Acknowledgments:** The authors would like to thank Barak Weiss for suggesting this beautiful topic. We are grateful towards Elon Lindenstrauss for various conversations around this topic and for so generously sharing his ideas. Last but not least, we thank Roland Prohaska and Pengyu Yang for related discussions.

**1.5. Ingredients of the proof.** The proof consists of roughly four main ingredients, the first three not being novel:

- **Establishing unipotent invariance:** This is a general tool, first introduced in [6, Prop. 3.1] and explored more in [37, Prop. 2.2]. It shows that limits of Birkhoff averages starting at a suitable expanding set are invariant under unipotent directions, which are determined from local information of the starting point. Both [6, 37] use effective estimates of correlations to prove this unipotent invariance. We rephrase the proof with martingales, as this is a recurring theme in the paper.
- **Linearization:** Provided that a measure  $\mu$  is invariant under a unipotent subgroup, Ratner’s measure classification [27] implies that the ergodic components are of algebraic nature. Linearization is a general technique, introduced by Dani and Margulis [8], transforming the problem of showing that  $\mu$  is the Haar measure to countably many algebraic-like problems. These problems can be summarized as showing that “some vectors do not

spend a lot of time in a certain set, with high probability.” For the purposes of this overview, we omit discussing here the appearance of the (related) quantitative non-divergence techniques of Kleinbock and Margulis [14].

- **Linear Dynamics:** The linearization technique requires that one rules out (among other things) the existence of a nonzero vector  $v \in \mathbb{R}^d$  such that  $\sup_{s \in [0,1]} \|a_t \varphi(s)v\| \rightarrow 0$ . Statements of this flavor have been established in the works of Shah [32], Shah, L. Yang [29] and Yang [41] (usually called ‘basic lemma’) in various instances.

**Interlude, Combinatorial structure of the problems:** For the sake of this introduction, we provide a combinatorial structure of the problems the linearization technique yields. Let  $\rho : \mathrm{SL}_d \rightarrow \mathrm{SL}_N$  be a representation defined over  $\mathbb{Q}$ . For each time  $t$  we have a collection  $A_t$  of disjoint intervals  $I \subset [0, 1]$ , each of which is a connected component of a set of the form  $\{s \in [0, 1] : \|\rho(a_t \varphi(s))v_I\| \leq 1\}$  for  $v_I \in \mathbb{Z}^N$  nonzero. For  $\varepsilon > 0$  let

$$\mathrm{Bad}_t(\varepsilon) = \bigcup_{I \in A_t} \{s \in I : \|\rho(a_t \varphi(s))v_I\| \leq \varepsilon\}$$

The goal is to bound the amount of  $t \in [0, T]$  for which  $s \in \mathrm{Bad}_t(\varepsilon)$  in terms of  $\varepsilon$  almost surely for  $s \in [0, 1]$ .

- **Interval combinatorics:** We first employ an argument by Kleinbock and Margulis [14] in their work on quantitative non-divergence, and obtain a bound on the total volume of exceptionally long intervals, that is,

$$\left| \bigcup_{\substack{I \in A_t \\ |I| > e^{-rt+t'}}} I \right| \ll e^{-\delta t'},$$

for some  $\delta > 0$  and  $t' < \delta t$ . By the Borel–Cantelli lemma one may hence ignore the intervals  $\{I \in A_t : |I| > e^{-rt+\sqrt{t}}\}$  (from now on these are called *long intervals*). The Short Intervals Lemma 4.1 is used to bound the rest (these are called *short intervals*). The Short Intervals Lemma takes the place of Margulis functions, which are used for instance in [6, 37].

**1.6. Further research.** One can try to extend the setting of the problem into a more general one:

**Problem 1.1.** *Extend Theorems 1.10 to non-abelian horospheres (cf. Yang [41]).*

One obstruction in using our technique occurs when ruling out long intervals (Proposition 3.4). For instance, when  $G^+$  is abelian, a curve  $s \mapsto a_t \varphi(s)v$  through a vector  $v \in \mathbb{R}^d$  can be described in a small neighborhood of a point  $s_0 \in [0, 1]$  through a unipotent curve  $\exp(xY)w$  with  $Y$  nilpotent and  $w \in \mathbb{R}^d$ . The same is not true when the horosphere is non-abelian.

Similarly, one may ask for the following extension:

**Problem 1.2.** *Extend Theorems 1.10 to smooth curves (cf. Shah and Yang [35]).*

In the spirit of Shi’s work [37] (see also [32, Thm. 1.10]) we pose the following problem:

**Problem 1.3.** Let  $\rho : G = \mathrm{SL}_d(\mathbb{R}) \rightarrow L$  be a representation of Lie groups and let  $\Gamma < L$  be a lattice. Let  $\varphi : [0, 1] \rightarrow G$  be an analytic curve whose image in  $P \backslash G$  is not contained in any weakly constraining pencil or in any partial flag subvariety. Let  $x \in L/\Gamma$ . Show that for almost every  $s \in [0, 1]$  and any  $f \in C_c(L/\Gamma)$

$$\frac{1}{T} \int_0^T f(\rho(a_t \varphi(s))x) dt \rightarrow \int f dm_{\overline{\rho(G)x}}$$

where  $m_{\overline{\rho(G)x}}$  is the invariant probability measure on the orbit closure of  $x$ .

In the spirit of [10, 40, 25, 26] one may ask the following:

**Problem 1.4.** Can one prove Theorem 1.10 for friendly measures (see [17]), in place of the uniform measure on a curve?

One could also consider non-conventional ergodic averages (see e.g. the work of Bourgain [3, 4] for polynomial averages):

**Problem 1.5.** Let  $\{t_n\}$  be a sequence of integers. Determine sufficient conditions on the curve  $\varphi$  and the sequence  $\{t_n\}$  so that for all  $x \in X$  and for almost every  $s \in [0, 1]$

$$\frac{1}{N} \sum_{n=1}^N f(a_{t_n} \varphi(s)x) \rightarrow \int_X f dm_X$$

for all  $f \in C_c(X)$ .

A different type of questions lies in changing the methods of our proof:

**Problem 1.6.** Can one extend the argument given in §6 to prove invariance in more directions, not only the tangent one?

This would nullify the need to use the linearization technique, though one would still have to prove non-escape of mass.

**1.7. Structure of the article.** In §2 we define the notations that we use and introduce preliminary tools on  $(C, \alpha)$ -good functions. In §3 we use elementary algebraic manipulations combined with quantitative non-divergence [14] and results in linear dynamics [32, 41] to control the ‘long intervals’ in the sense of §1.5. In §4 we prove a completely probabilistic lemma which will help us control the ‘short intervals’. In §5 we combine the previous tools and give a demonstration of the method, proving a (partially quantitative) nondivergence result. In §6 establish unipotent invariance of limits the Birkhoff averages. It is independent of the previous sections, and it uses probabilistic methods similar to those in §4. In §7 we combine the previous results with Ratner’s measure classification [27] and the linearization technique [8] to prove Theorem 1.10. Finally, in §8 we establish the applications in §1.1, 1.2.

## 2. NOTATION AND PRELIMINARIES

As in Theorem 1.10,  $\{a_t : t \in \mathbb{R}\}$  shall be the (real) diagonalizable subgroup of  $G = \mathrm{SL}_d(\mathbb{R})$ ,  $d \geq 3$  defined in (1.3). Let  $P$  be the parabolic subgroup in (1.4) and let  $\mathfrak{p}$  be its Lie algebra.

Let  $Z = Z(\{a_t\}) < P$  be the centralizer subgroup of  $\{a_t\}$ . Note that  $Z$  acts by conjugation on the expanded horospherical subgroup

$$G^+ = \{g \in G : a_t g a_{-t} \rightarrow \mathrm{id} \text{ as } t \rightarrow -\infty\}.$$



For  $\{a_t\}$  as above, one may identify  $G^+ \simeq \text{Lie}(G^+) = \mathfrak{g}^+ \simeq \text{Mat}_{mn}(\mathbb{R})$  and

$$\begin{aligned} Z &\simeq (\text{GL}_m(\mathbb{R}) \times \text{GL}_n(\mathbb{R}))^1 \\ &:= \{(g_1, g_2) \in \text{GL}_m(\mathbb{R}) \times \text{GL}_n(\mathbb{R}) : \det(g_1) \det(g_2) = 1\} \end{aligned}$$

where the action of  $Z$  on  $G^+$  corresponds to the action of  $(\text{GL}_m(\mathbb{R}) \times \text{GL}_n(\mathbb{R}))^1$  on  $\text{Mat}_{mn}(\mathbb{R})$  via  $(g_1, g_2).X = g_1 X g_2^{-1}$ .

**Example 2.1.** If  $a_t = (e^{(d-1)t}, e^{-t}, \dots, e^{-t})$  then  $G^+ \simeq \mathbb{R}^{d-1}$ ,  $Z \simeq \text{GL}_{d-1}(\mathbb{R})$ , and the action of  $Z$  on  $G^+$  corresponds to the (non-standard) action of  $\text{GL}_{d-1}(\mathbb{R})$  on  $\mathbb{R}^{d-1}$  via  $v \mapsto g.v = \det(g)^{-1}(A^{-1})^t v$ .

**Lemma 2.2.** *There are real subvarieties  $V_0 \subset V_1 \subset \dots \subset V_{\min\{m,n\}} = \mathfrak{g}^+$  so that each  $V_k$  is invariant under  $Z$  and the action of  $Z$  on  $V_k \setminus V_{k-1}$  has at most two orbits both of which are open in  $V_k$  (for the Hausdorff topology).*

*Proof.* Under the identification  $\mathfrak{g}^+ \simeq \text{Mat}_{mn}(\mathbb{R})$  the action of  $Z$  preserves the rank. For each  $k \leq \min\{m, n\}$  let  $V_k \subset \mathfrak{g}^+$  be the subvariety of matrices of rank at most  $k$ . Fix a rank  $k \leq \min\{m, n\}$  and consider the action of  $Z$  on  $V_k \setminus V_{k-1}$ ; we prove that it has at most two orbits<sup>1</sup>. A section for this action is given by

$$S_k = \left\{ \begin{pmatrix} A & 0_{k,n-k} \\ 0_{m-k,k} & 0_{m-k,n-k} \end{pmatrix} : A \in \text{GL}_k(\mathbb{R}) \right\} \subset V_k$$

where a subgroup of  $Z$  identifiable with  $(\text{GL}_k(\mathbb{R}) \times \text{GL}_k(\mathbb{R}))^1$  still acts on  $S_k$  (in the natural manner). This new action is transitive on matrices  $A$  whose determinants share the same sign.  $\square$

We let  $\varphi : [0, 1] \rightarrow G$  be an analytic curve. We shall often assume that the image of  $\varphi$  in  $P \setminus G$  is not contained in any weakly constraining pencil. Equivalently, the image of  $\varphi$  in  $P \setminus G$  is not contained in any weakly unstable Schubert variety [41, Props. 6.4, 6.5].

In the following we shall always write  $r = \frac{1}{m} + \frac{1}{n} > 0$  for the expansion rate of  $G^+$  (that is, the eigenvalue of  $\text{Ad}(a_t)$  acting on  $\mathfrak{g}^+$  is  $e^{rt}$ ). We define  $\Phi^+(s, \xi) \in \mathfrak{g}^+$  and  $\Phi^{0-}(s, \xi) \in \mathfrak{p}$  for  $s \in [0, 1]$  and  $\xi$  sufficiently small through

$$(2.1) \quad \varphi(s + \xi)\varphi(s)^{-1} = \exp(\Phi^{0-}(s, \xi)) \exp(\Phi^+(s, \xi)).$$

We remark that  $\Phi^{0-}(\cdot, \cdot)$  and  $\Phi^+(\cdot, \cdot)$  are analytic in both variables. Moreover, we have  $\Phi^{0-}(s, \xi) = O(\xi)$  and

$$(2.2) \quad \Phi^+(s, \xi) = \xi Y_s + O(\xi^2)$$

for some analytic curve  $s \in [0, 1] \rightarrow Y_s$ .

**Lemma 2.3.** *If the image of  $\varphi$  in  $P \setminus G$  is non-constant, the curve  $s \mapsto Y_s$  is non-zero at all but finitely many points.*

*Proof.* As  $s \mapsto Y_s$  is analytic, we may assume by contradiction that  $Y_s = 0$  for all  $s \in [0, 1]$ . Then for  $\xi$  small the distance between  $P\varphi(s)$  and  $P\varphi(s + \xi)$  is of order  $O(\xi^2)$  as

$$P\varphi(s + \xi) = P(I + O(\xi^2))\varphi(s) = P\varphi(s)(I + O(\xi^2))$$

by (2.2). In particular, the derivative of  $s \mapsto P\varphi(s)$  is zero everywhere and hence  $s \mapsto P\varphi(s)$  is constant.  $\square$

<sup>1</sup>A more careful argument show that the action of  $Z$  on  $V_k \setminus V_{k-1}$  is transitive if  $k < \max\{m, n\}$  and has two orbits if  $k = m = n$ .

The following lemma is a direct consequence of (2.1) and (2.2).

**Lemma 2.4.** *For any  $\xi \in \mathbb{R}$  sufficiently small and all  $t > 0$  we have*

$$a_t \varphi(s + \xi) = (I + O(\xi + \xi^2 e^{rt})) \exp(\xi e^{rt} Y_s) a_t \varphi(s).$$

*In particular, the distance between  $a_t \varphi(s + \xi)$  and  $a_t \varphi(s)$  is of order  $O(\xi e^{rt})$ .*

Let  $k \leq \min\{m, n\}$  be minimal such that  $V_k \cap \{Y_s : s \in [0, 1]\}$  is infinite. By Lemma 2.3, we have  $k > 0$ . By analyticity, we have  $Y_s \in V_k$  for all  $s \in [0, 1]$ . Since  $V_{k-1} \cap \{Y_s : s \in [0, 1]\}$  is finite by construction, we may assume without loss of generality that  $Y_s \notin V_{k-1}$  for all  $s \in [0, 1]$  by a covering argument. In particular, by Lemma 2.2 the points  $\{Y_s : s \in [0, 1]\}$  lies on the same  $Z$  orbit. Hence there exists an analytic curve  $s \mapsto z(s) \in Z$  such that

$$(2.3) \quad z(s) \cdot Y_s = Y_0$$

for all  $s \in [0, 1]$ . We note that after the above reduction  $Y_s \neq 0$  for all  $s \in [0, 1]$ .

*Remark 2.5.* It is not generally necessary (cf. [41]) to map  $Y_s$  to, say,  $Y_{s'}$  within the centralizer of the flow and indeed it is only possible when the horosphere is abelian (or equivalently minimal). In the context of this article, it is however convenient.

**2.1.  $(C, \alpha)$ -good functions.** We recall the following (by now standard) definition from [14]:

**Definition 2.6** ( $(C, \alpha)$ -good functions). Let  $I \subset \mathbb{R}$  be an interval and let  $f : I \rightarrow \mathbb{R}$ . Given  $C > 0$  and  $\alpha > 0$  we say that  $f$  is  $(C, \alpha)$ -good if for any subinterval  $J \subset I$  and any  $\varepsilon > 0$  we have

$$|\{x \in J : |f(x)| < \varepsilon\}| \leq C \left( \frac{\varepsilon}{\sup_{x \in J} |f(x)|} \right)^\alpha |J|.$$

Here (and henceforth),  $|A|$  for a subset  $A \subset \mathbb{R}$  indicates the Lebesgue measure of  $A$ .

For example, polynomials of degree at most  $n$  are  $(C_n, \frac{1}{n})$ -good for some  $C_n > 0$  (cf. [8, Lemma 4.1], [14, Prop. 3.2]).

For any  $\gamma > 0$  and  $J = [a - b, a + b] \subset \mathbb{R}$  an interval define  $\gamma \cdot J = [a - \gamma b, a + \gamma b]$  the centered  $\gamma$ -dilation of  $J$ .

**Lemma 2.7.** *Let  $f : [0, 1] \rightarrow \mathbb{R}$  be  $(C, \alpha)$ -good for some  $(C, \alpha)$  and let  $I \subset [0, 1]$  be a maximal subinterval with  $\sup_{s \in I} |f(s)| \leq 1$ . If  $J \subset I$  is an interval with  $\sup_{s \in J} |f(s)| \leq \varepsilon$  then  $\frac{1}{C\varepsilon^\alpha} \cdot J \cap [0, 1] \subset I$ .*

*Proof.* Write  $I = [a, b]$  and  $J = [s_0 - c, s_0 + c]$ . By the maximality condition of  $I$ , we get that either  $|f(a)| = 1$  or  $a = 0$ . Similarly, either  $|f(b)| = 1$  or  $b = 1$ . To show the desired containment, it is sufficient to show that

$$\left[ s_0 - \frac{1}{C\varepsilon^\alpha} c, s_0 \right] \cap [0, 1] \subseteq I \quad \text{and} \quad \left[ s_0, s_0 + \frac{1}{C\varepsilon^\alpha} c \right] \cap [0, 1] \subseteq I.$$

We will show the first containment, and the second will follow similarly. If  $a = 0$  the containment follows since  $s_0 \in I$ . Otherwise,  $|f(a)| = 1$ . Applying the  $(C, \alpha)$ -property on  $I^- = [a, s_0]$  yields that the measure of  $J \cap I^\pm$  relative to  $I^\pm$  is at most  $C\varepsilon^\alpha$  which implies the desired dilation statement.  $\square$

The following local to global property for  $(C, \alpha)$ -good functions is already mentioned in [33, §2.1], we include a full proof for lack of reference.

**Lemma 2.8** (Locality of the  $(C, \alpha)$ -goodness property.). *Let  $I$  be an interval and  $f : I \rightarrow \mathbb{R}$  be a function. Suppose that  $I = \bigcup_{i=1}^k I_i$  is a union of intervals such that  $f$  is  $(C, \alpha)$  good on each interval  $I_i$  for  $C > 1$  and  $\alpha > 0$ , and for every  $i, j \in \{1, \dots, k\}$  the intersection  $I_{i_1} \cap I_{i_2}$  is either empty or has length  $|I_{i_1} \cap I_{i_2}| \geq \eta$ . Then  $f$  is  $(C^k k (\frac{|I|}{\eta})^{k-1}, \alpha)$ -good.*

*Proof.* Let  $J \subset I$  be a subinterval. We will prove the goodness property with respect to this interval. For any  $i \in \{1, \dots, k\}$  set  $J_i = I_i \cap J$ . In the following we shall call to indices  $i, j \in \{1, \dots, k\}$  paired if  $I_i \cap I_j \neq \emptyset$ . Whenever  $i, j$  are paired, we have either  $J_i \cap J_j = I_i \cap I_j$  or one of the interval  $J_i, J_j$  may be discarded in the induced cover of  $J$ . Indeed, if the intersection  $I_i \cap I_j$  does not intersect  $\bar{J}$  one of the intervals  $J_i, J_j$  is empty and if it contains a boundary point of  $J$  then either  $J_i \subset J_j$  or  $J_j \subset J_i$ . After discarding intervals, we may assume without loss of generality that  $I_i \cap I_j = J_i \cap J_j$  for all  $i, j$ .

Now suppose that  $i, j$  are paired. We claim that

$$(2.4) \quad \sup_{x \in J_i} |f(x)| \geq \tilde{\eta} \sup_{x \in J_j} |f(x)|$$

where  $\tilde{\eta} = (\eta/(C|I|))^{1/\alpha}$ . Indeed, as  $f$  is  $(C, \alpha)$ -good on  $I_j$  we have

$$\eta \leq |J_i \cap J_j| \leq |\{x \in J_j : |f(x)| \leq \sup_{x \in J_i} |f(x)|\}| \leq C \left( \frac{\sup_{x \in J_i} |f(x)|}{\sup_{x \in J_j} |f(x)|} \right)^\alpha |I|$$

which implies (2.4).

Note that  $\sup_{x \in J} |f(x)| = \sup_{x \in J_{i_0}} |f(x)|$  for some  $i_0$ . Consider the graph whose vertices are indices  $i = 1, \dots, k$  and whose edges are  $(i, j)$  where  $i, j$  are paired. It is connected since the intervals  $J_i$  cover the connected set  $J$ . Iterating (2.4) over a path between  $i_0$  to any other vertex  $i$  yields that for every  $i = 1, \dots, k$  we have  $\sup_{x \in J_i} |f(x)| \geq \tilde{\eta}^{k-1} \sup_{x \in J} |f(x)|$ . Finally, for every  $\varepsilon > 0$  we have

$$\begin{aligned} |\{x \in J : |f(x)| \leq \varepsilon\}| &\leq \sum_{i=1}^k |\{x \in J_i : |f(x)| \leq \varepsilon\}| \\ &\leq C \sum_{i=1}^k \left( \frac{\varepsilon}{\sup_{x \in J_i} |f(x)|} \right)^\alpha |J_i| \\ &\leq C \tilde{\eta}^{-\alpha(k-1)} \sum_{i=1}^k \left( \frac{\varepsilon}{\sup_{x \in J} |f(x)|} \right)^\alpha |J_i| \\ &\leq C k \tilde{\eta}^{-\alpha(k-1)} \left( \frac{\varepsilon}{\sup_{x \in J} |f(x)|} \right)^\alpha |J| \end{aligned}$$

as desired.  $\square$

The following lemma in particular establishes uniform  $(C, \alpha)$ -goodness for certain families of analytic functions.

**Lemma 2.9.** *Suppose that  $\varphi : [0, 1] \rightarrow M_{N \times N}(\mathbb{R})$  is analytic. Then there exist constants  $C > 0$ ,  $\alpha > 0$ , and  $M > 0$  such that for any  $v \in \mathbb{R}^N$  and  $g \in M_{N \times N}(\mathbb{R})$  the following hold:*

- *The function  $s \mapsto \|g\varphi(s)v\|$  is  $(C, \alpha)$ -good.*

- For any  $x > 0$  the set

$$\{s \in [0, 1] : \|g\varphi(s)v\| < x\}$$

has at most  $M$  connected components.

*Proof.* Since  $f$  is  $(C, \alpha)$ -good if and only if  $f^2$  is  $(C, \alpha/2)$ -good, we may prove the lemma for  $s \mapsto \|g\varphi(s)v\|^2$  instead. Let  $\mathcal{F}$  be the linear span of all products of two entry functions of  $\varphi$ . The space  $\mathcal{F}$  is a finite-dimensional vector space as it is generated by  $N^2(N^2 + 1)/2$  elements. The map  $s \mapsto \|g\varphi(s)v\|^2$  is in  $\mathcal{F}$  for every  $t \in \mathbb{R}$ ,  $v \in \mathbb{R}^N$  and  $g \in M_{N \times N}(\mathbb{R})$ , hence we will prove the conclusions of the lemma for all  $f \in \mathcal{F}$ .

Let  $S_{\mathcal{F}}$  be a unit sphere in  $\mathcal{F}$  (with respect to any norm). Normalizing functions to  $S_{\mathcal{F}}$  we may prove the lemma only for  $f \in S_{\mathcal{F}}$ .

For every  $p = (f, s) \in S_{\mathcal{F}} \times [0, 1]$  there is  $k = k_p \in \mathbb{N}$  such that  $|f^{(k)}(s)| > 0$  by analyticity of  $f$ . We can find a neighborhood  $U_p \times I_p$  of  $p = (f, s) \in S_{\mathcal{F}} \times [0, 1]$  and  $\varepsilon_p > 0$  such that for every  $p_0 = (f_0, s_0) \in U_p \times I_p$  we have  $|f_0^{(k)}(s_0)| > \varepsilon_p$ . Moreover, we may assume that  $I_p$  is always an interval. Since  $S_{\mathcal{F}} \times [0, 1]$  is compact we deduce that there is a finite collection  $P \subset S_{\mathcal{F}} \times [0, 1]$  such that  $\{U_p \times I_p : p \in P\}$  covers  $S_{\mathcal{F}} \times [0, 1]$ .

By the mean value theorem if  $f \in U_p$  then  $\{s \in I_p : |f(s)| < x\}$  has at most  $k_p$  connected components. The second part of the lemma follows from the finiteness of  $P$ . For the first part of the lemma we may apply [14, Lemma 3.3] and obtain that for any  $f \in U_p$  the restriction  $f|_{I_p}$  is  $(C_p, \alpha_p)$ -good for some  $C_p, \alpha_p > 0$  (depending on  $k_p$ ). Thus, the first part of the lemma follows also from the finiteness of  $P$  and Lemma 2.8.  $\square$

### 3. LONG INTERVALS

#### 3.1. Contraction for long intervals with short vectors.

**Claim 3.1.** *Let  $P(x) = a_0x^n + a_1x^{n-1} + a_2x^{n-2} + \dots + a_n$  be a polynomial. Then for every  $s > 0$  we have*

$$\max_{x \in [0, s]} |P(x)| \asymp_n \max_{k=0, \dots, n} |a_k s^{n-k}|.$$

*Proof.* Using rescaling it is sufficient to prove the lemma for  $s = 1$  then the claim follows from the fact that both sides are norms in the coefficient vector.  $\square$

**Claim 3.2.** *Let  $h \in \mathfrak{sl}_N(\mathbb{R}) \setminus \{0\}$  be a nilpotent matrix. Then there is a diagonalizable flow  $t \mapsto b_t = \exp(b't) \in \mathrm{SL}_N(\mathbb{R})$  and a constant  $\delta \geq \frac{1}{N}$  such that for every  $t_0 \geq 0$ , vector  $v \in \mathbb{R}^N$  and interval  $I \subseteq \mathbb{R}$  with  $|I| \geq e^{t_0}$  we have*

$$\max_{s \in I} \|b_{t_0} \exp(sh)v\| \asymp_h e^{-\delta t_0} \max_{s \in I} \|\exp(sh)v\|.$$

*Proof.* First, we will prove the claim whenever  $h$  is a nontrivial Jordan block,

$$h = \begin{pmatrix} 0 & 1 & & \\ & 0 & \ddots & \\ & & \ddots & 1 \\ & & & 0 \end{pmatrix}.$$

Let  $b(t) = \exp(t \operatorname{diag}(\frac{1-N}{2}, \frac{3-N}{2}, \dots, \frac{N-1}{2}))$ . We may shrink  $I$  and assume that  $|I| = e^{t_0}$ . By changing  $v$  we may assume that  $I = [0, e^{t_0}]$ . Now if  $v = (v_i)_{i=0}^{N-1}$  then by Claim 3.1 the maximal  $k$ -th coordinate satisfies

$$\max_{s \in I} |(\exp(sh)v)_k| \asymp_N \max_{i=0, \dots, N-1-k} e^{it_0} |v_{i+k}|.$$

It follows that

$$\max_{s \in I} \|\exp(sh)v\| \asymp_N \max_{i=0, \dots, N-1} e^{it_0} |v_i|.$$

Similarly

$$\begin{aligned} \max_{s \in I} \|b(t_0) \exp(sh)v\| &\asymp_N \max_{k=0, \dots, N-1} (e^{(1+2k-N)t_0/2} \max_{i=0, \dots, N-1-k} e^{it_0} |v_{i+k}|) \\ &\asymp_N e^{(1-N)t_0/2} \max_{i=0, \dots, N-1} e^{it_0} |v_i|. \end{aligned}$$

The result follows for  $\delta = \frac{N-1}{2}$ . Now, whenever  $h$  of Jordan normal form, say,  $h = \operatorname{diag}(h_0, \dots, h_k)$  where  $h_i \in \mathfrak{sl}_{N_i}(\mathbb{R})$  is a Jordan block. Let  $b_0(t), \dots, b_k(t)$  be the corresponding homomorphisms and  $\delta_i = \frac{N_i-1}{2}$  for  $i = 0, \dots, k$ . Let  $\delta = \frac{1}{N} \sum_{i=0}^k N_i \delta_i$  be the weighted average of the  $\delta_i$ . It satisfies that  $\sum_{i=0}^k N_i (\delta - \delta_i) = 0$ . Then  $b(t) = \operatorname{diag}(e^{-(\delta - \delta_i)t} b_i(t) : i = 0, \dots, k)$  satisfies the desired result. Note that if  $h$  is nontrivial, then one of the  $N_i$  is greater or equal 2, and hence  $\delta \geq \frac{1}{N}$ . Since every  $h \in \mathfrak{sl}_N(\mathbb{R})$  is conjugate to a Jordan normal form, the result follows.  $\square$

Recall the definition of  $z(\cdot)$  from (2.3).

**Lemma 3.3.** *For every nontrivial  $\rho : G \rightarrow \operatorname{SL}_N(\mathbb{R})$  there exists diagonalizable flow  $t \in \mathbb{R} \mapsto b_t \in \operatorname{SL}_N(\mathbb{R})$  and constants  $\delta_1, \delta_2 > 0$  such that for every  $t_1 < \delta_1 t_0$ , we have the following: For every interval  $J \subseteq [0, 1]$  with  $|J| \geq e^{-rt_0+t_1}$  and for every  $v \in \mathbb{R}^N$  we have*

$$(3.1) \quad \max_{s \in J} \|b_{t_1} \rho(a_{t_0} z(s) \varphi(s)) v\| \asymp e^{-\delta_2 t_1} \max_{s \in J} \|\rho(a_{t_0} \varphi(s)) v\|.$$

*Proof.* Since for every  $L < |J|$  and a function  $f : J \rightarrow \mathbb{R}$  we have

$$\max_{s \in J} f(s) = \max_{\substack{J' \subset J \\ |J'|=L}} \max_{s \in J'} f(s),$$

it is sufficient to prove the claim for any subinterval of  $J$  with length  $e^{-rt_0+t_1}$ , hence we may assume that  $|J| = e^{-rt_0+t_1}$  and write  $J = [s_0, s_0 + e^{-rt_0+t_1}]$ . Set  $s = s_0 + \varepsilon$  for some  $\varepsilon \in [0, e^{-rt_0+t_1}]$ . Lemma 2.4 implies that

$$a_{t_0} \varphi(s) = (I + O(\varepsilon + \varepsilon^2 e^{rt_0})) \exp(\varepsilon e^{rt_0} Y_{s_0}) a_{t_0} \varphi(s_0).$$

Hence,

$$\begin{aligned} a_{t_0} z(s) \varphi(s) &= z(s) a_{t_0} \varphi(s) \\ &= (I + O(\varepsilon)) z(s_0) (I + O(\varepsilon + \varepsilon^2 e^{rt_0})) \exp(\varepsilon e^{rt_0} Y_{s_0}) a_{t_0} \varphi(s_0) \\ &= A_{s_0}(\varepsilon) \exp(\varepsilon e^{rt_0} Y_0) z(s_0) a_{t_0} \varphi(s_0) \end{aligned}$$

where  $A_{s_0}(\varepsilon) \in \operatorname{SL}_d(\mathbb{R})$  satisfies  $A_{s_0}(\varepsilon) = I + O(\varepsilon + \varepsilon^2 e^{rt_0})$ . The range for  $\varepsilon$  implies that  $\varepsilon + \varepsilon^2 e^{rt_0} \ll e^{-rt_0+2t_1}$ .

The homomorphism  $x \mapsto \rho(\exp(xY_0))$  is unipotent, and is of the form  $x \mapsto \exp(h_0 x)$  for some nonzero nilpotent  $h_0 \in \mathfrak{sl}_N(\mathbb{R})$  (explicitly,  $h_0 = \operatorname{D}\rho(Y_0)$ ). Let

$t \mapsto b(t)$  and  $\delta_0 > 0$  be the diagonalizable flow and the constant from Claim 3.2 for this  $h_0$  respectively. For every  $v \in \mathbb{R}^N$  we have

$$\begin{aligned}
(3.2) \quad & \max_{s \in J} \|b(t_1)\rho(a_{t_0}z(s)\varphi(s))v\| \\
&= \max_{\varepsilon \in [0, e^{-rt_0+t_1}]} \|b(t_1)\rho(A_{s_0}(\varepsilon)\exp(\varepsilon e^{rt_0}Y_0)z(s_0)a_{t_0}\varphi(s_0))v\| \\
&= \max_{\varepsilon \in [0, e^{-rt_0+t_1}]} \|\rho(A_{s_0}(\varepsilon))^{b(t_1)}b(t_1)\exp(\varepsilon e^{rt_0}h_0)v'\|,
\end{aligned}$$

where  $\rho(A_{s_0}(\varepsilon))^{b(t_1)}$  is the conjugation of  $\rho(A_{s_0}(\varepsilon))$  by  $b(t_1)$  and

$$v' = \rho(z(s_0)a_{t_0}\varphi(s_0))v.$$

Since  $A_{s_0}(\varepsilon)$  is of distance  $O(e^{-rt_0+2t_1})$  from the identity matrix, it follows that  $\rho(A_{s_0}(\varepsilon))$  is of distance  $O(e^{-rt_0+2t_1})$  from the identity matrix as well. Since the eigenvalues of  $b(t_1), b(t_1)^{-1}$  are bounded by  $Rt_1$  for some  $R > 0$  depending only on  $b$ , and hence only on  $\rho$ , it follows that  $\rho(A_{s_0}(\varepsilon))^{b(t_1)}$  is of distance  $e^{-rt_0+(2R+2)t_1}$  of the identity element. Provided that  $t_1 < \frac{r}{2R+2}t_0$  we get that  $\rho(A_{s_0}(\varepsilon))^{b(t_1)}$  is bounded and hence we may omit this term in Eq. (3.2), that is, it is proportional to

$$(3.3) \quad \max_{x \in [0, e^{t_1}]} \|b(t_1)\exp(xh_0)v'\|.$$

Now Claim 3.2 implies that

$$(3.4) \quad \max_{x \in [0, e^{t_1}]} \|b(t_1)\exp(xh_0)v'\| \asymp e^{-\delta t_1} \max_{x \in [0, e^{t_1}]} \|\exp(xh_0)v'\|.$$

We have seen that the left-hand side of (3.4) is asymptotic to the left-hand side of (3.1). Moreover, we obtain in a similar manner

$$\max_{x \in [0, e^{t_1}]} \|\exp(xh_0)v'\| \asymp \max_{s \in J} \|\rho(z(s)a_{t_0}\varphi(s))v\| \asymp \max_{s \in J} \|\rho(a_{t_0}\varphi(s))v\|.$$

using that  $s \mapsto z(s)$  is bounded, which implies that  $s \mapsto \rho(z(s))$  is bounded as well. The result thus follows from (3.4).  $\square$

**3.2. Atypicality of long intervals with short vectors.** Lemma 3.3 together with the quantitative non-divergence results by Kleinbock and Margulis [14] implies the following result, which roughly states that in any representation the set of points  $s$  contained in an unexpectedly long interval with a short vector is small (unless a natural obstruction occurs).

**Proposition 3.4** (Avoiding exceptional intervals). *Let  $\varphi : [0, 1] \rightarrow G$  be an analytic curve whose image in  $P \backslash G$  is not contained in any weakly constraining pencil or in any partial flag subvariety with respect to  $\{a_t\}$ . Let  $\rho : \mathrm{SL}_d \rightarrow \mathrm{SL}_N$  be a representation defined over  $\mathbb{Q}$  and let  $\mathcal{U} \subset \mathbb{Z}^N$  be a set of non-zero vectors. There exist constants  $\delta > 0$  and  $C_1 > 0$  depending on  $\rho$  with the following property.*

*Let  $\psi : (0, \infty) \rightarrow (0, \infty)$  be a function with  $\psi(t) < \delta t$  for all  $t > 0$ . Then one of the following is true for all  $g \in \mathrm{SL}_d(\mathbb{R})$ ,  $R > 0$ , and  $t > 0$  sufficiently large (depending on  $g, R$ ):*

- (1) *Let  $\mathcal{VB}_t$  be the union of all intervals  $I \subset [0, 1]$  with  $|I| \geq e^{-rt+\psi(t)}$  and with*

$$I \subset \{s \in [0, 1] : \|\rho(a_t\varphi(s)g)v\| < R\}$$

*for some  $v \in \mathcal{U}$ . Then  $|\mathcal{VB}_t| \leq C_2 e^{-\delta\psi(t)}$  for a constant  $C_2 = C_2(R, g)$ .*

- (2) *There exists a  $\rho(\mathrm{SL}_d(\mathbb{R}))$ -invariant vector  $v \in \mathcal{U}$  with  $\|v\| \leq C_1 R$ .*

We shall heavily use the following result from linear dynamics (often referred to as 'basic lemma') – see also Shah's basic lemma in [32, §4] as well as [29, §4].

**Theorem 3.5** (Yang [41]). *Let  $\varphi : [0, 1] \rightarrow G$  be an analytic curve whose image in  $P \backslash G$  is not contained in any weakly constraining pencil. Let  $\rho : G \rightarrow \mathrm{SL}_N(\mathbb{R})$  be a representation, let  $V^+ \subset \mathbb{R}^N$  be the sum of all eigenspaces of eigenvalue  $> 1$  for  $\rho(a_t)$ ,  $t \geq 0$ , and write  $\pi_+ : \mathbb{R}^N \rightarrow V^+$  for the projection with respect to the weight decomposition. Let  $v \in \mathbb{R}^N$  be a nonzero vector such that for all  $s \in [0, 1]$*

$$(3.5) \quad \pi_+(\rho(\varphi(s))v) = 0.$$

*Then  $\rho(G)v$  is closed. Moreover, if the image  $\varphi$  in  $P \backslash G$  is not contained in any partial flag subvariety with respect to  $\{a_t\}$ , then  $v$  is  $G$ -invariant.*

*Proof.* We merely point out how to combine the proven statements in [41]. If  $\rho(G)v$  were not closed, [41, Props. 2.7, 6.5] would imply that the image of  $\varphi$  in  $P \backslash G$  would be contained in a weakly constraining pencil yielding a contradiction. So  $\rho(G)v$  is closed. We now proceed as in [41, Props. 5.4, 5.6]. In particular, by (3.5) there exists for every  $s \in [0, 1]$  an element  $\xi(s) \in G$  with

$$\lim_{t \rightarrow \infty} \rho(a_t \varphi(s))v = \rho(\xi(s)\varphi(s))v =: w_s$$

Fix  $s \in [0, 1]$  and let  $F_s < G$  be the stabilizer subgroup of  $w_s$ . By Matsushima's criterion,  $F_s$  is reductive. It is easy to see that  $F_s$  contains  $\{a_t\}$ .

We claim that we may assume  $\xi(s) \in P$ . Indeed, by definition of  $w_s$  there exists for every  $t > 0$  some  $\varepsilon_t$  in a small neighborhood of the identity of  $G$  such that  $w_s = a_t \xi(s) a_t^{-1} \varepsilon_t w_s$ . In particular,  $a_t \xi(s)^{-1} a_t^{-1}$  is bounded in  $G/F$  which implies that  $\xi(s) \in PF$  by a local calculation (indeed,  $\xi(s) a_t^{-1}$  is in a neighborhood of the identity coset of  $F \backslash G$  for  $t$  large enough). This proves the claim.

Now let  $g = \xi(0)\varphi(0)$  and  $F = F_0$ . As  $\xi(0) \in P$  and  $a_t \varphi(s)v$  converges,  $a_t \xi(0)\varphi(s)g^{-1}F$  converges in  $G/F$ . Moreover,  $\xi(0)\varphi(s)g^{-1}$  is in a neighborhood of the identity for  $s$  close to 0 and a Lie algebra calculation shows that  $\xi(0)\varphi(s)g^{-1} \in PF$ . Hence,  $\varphi(s) \in PFg$  for all  $s \in [0, 1]$  where  $PFg$  is a proper flag subvariety of  $P \backslash G$  unless  $F = G$  i.e.  $v$  is  $G$ -invariant.  $\square$

*Proof of Proposition 3.4.* Suppose for simplicity first that  $\rho$  is irreducible (and nonzero). Let  $z(\cdot)$  be as in (2.3), let  $\delta > 0$  to be determined, and let  $t \mapsto b_t$  be as in Lemma 3.3. If  $\delta$  is sufficiently small, then for any interval  $J \subset [0, 1]$  with  $|J| \geq e^{-rt+\psi(t)}$  and any  $v \in \mathbb{R}^N$  with

$$\max_{s \in J} \|\rho(a_t \varphi(s))v\| \leq R$$

we have for some constant  $c = c(\rho) > 0$

$$(3.6) \quad \max_{s \in J} \|b_{\psi(t)} \rho(z(s) a_t \varphi(s))v\| \leq cR e^{-\delta \psi(t)}.$$

We now invoke the quantitative non-divergence results by Kleinbock and Margulis [14, Thm. 5.2]. To that end, note first that for any  $v \in \mathbb{R}^N$  the map

$$s \mapsto \|b_{\psi(t)} \rho(a_t z(s) \varphi(s))v\|$$

is  $(C, \alpha)$ -good for some  $(C, \alpha)$  depending on  $\rho$  and  $\varphi$ . Indeed, we may apply Lemma 2.9 for  $s \mapsto \rho(z(s)\varphi(s))$  and  $g = b_{\psi(t)}\rho(a_t)$ . In particular, [14, Thm. 5.2] implies that for some sufficiently small  $\kappa > 0$  one of the following options holds for every  $t > 0$ :

(a) The measure of the set of  $s \in [0, 1]$  so that

$$\|b_{\psi(t)}\rho(a_t z(s)\varphi(s)g)v\| \leq ce^{-\delta\psi(t)}R$$

for some nonzero  $v \in \mathbb{Z}^N$  is  $\ll e^{-\kappa\delta\psi(t)}$ .

(b) There exists a rational subspace  $U_t \subset \mathbb{Q}^N$  such that

$$\sup_{s \in [0, 1]} \|b_{\psi(t)}\rho(a_t z(s)\varphi(s)g)U_t(\mathbb{Z})\| \leq ce^{-\frac{\delta}{2}\psi(t)}R.$$

We prove that option (b) can only occur in a bounded set of times (depending on  $g, R$ ); this concludes the special case when  $\rho$  is irreducible in view of (3.6). Suppose option (b) holds for a sequence  $t_j \rightarrow \infty$ . Restricting to a subsequence we may assume that the subspaces  $U_j = U_{t_j}$  have equal dimension  $m$ . Also, we may assume that if  $u_j \in \bigwedge^m \mathbb{Z}^N$  is a pure integral vector corresponding to  $U_j(\mathbb{Z})$  then  $\frac{u_j}{\|u_j\|} \rightarrow u$  for some (possibly irrational) pure vector  $u \in \bigwedge^m W_i$  of norm one. Note that (b) implies

$$(3.7) \quad \sup_{s \in [0, 1]} \|\rho(a_{t_j}\varphi(s)g)u_j\| \ll e^{\star\psi(t_j)}R.$$

Let  $\pi_+ : W = \bigwedge^m \mathbb{R}^N \rightarrow W^+$  be the projection onto the  $a_t$ -expanded subspace  $W^+$ . By definition,  $W^+$  is the direct sum of all  $a_t$ -eigenspaces with eigenvalue  $e^{\lambda t}$  for some  $\lambda > 0$  and the projection is with respect to the eigenspace decomposition. Note that there exists  $\alpha > 0$  (depending on  $\rho$ ) such that for any  $w \in W$

$$\|\rho(a_t)\pi_+(w)\| \gg e^{\alpha t}\|\pi_+(w)\|.$$

In particular, for any  $s \in [0, 1]$  by discreteness of  $W(\mathbb{Z})$

$$\begin{aligned} \|\rho(a_{t_j}\varphi(s)g)u_j\| &\gg \|\pi_+(\rho(a_{t_j}\varphi(s)g)u_j)\| \gg e^{\alpha t_j}\|\pi_+(\rho(\varphi(s)g)u_j)\| \\ &\gg e^{\alpha t_j}\|u_j\|\|\pi_+(\rho(\varphi(s)g)u)\| \gg e^{\alpha t_j}\|\pi_+(\rho(\varphi(s)g)u)\| \end{aligned}$$

which implies together with (3.7) that

$$\sup_{s \in [0, 1]} \|\pi_+(\rho(\varphi(s)g)u)\| \ll e^{-\alpha t_j + \star\psi(t_j)}R.$$

For  $j \rightarrow \infty$  this implies that  $\pi_+(\rho(\varphi(s)g)u) = 0$ . By Theorem 3.5 we obtain that  $u$  is  $\rho(G)$ -invariant. But  $u$  is pure and hence the corresponding subspace is also  $\rho(G)$ -invariant. By irreducibility of  $\rho$ , this is a contradiction, which concludes this special case.

Now suppose that  $\rho$  is general. Write  $\mathbb{R}^N$  as a direct sum of subspaces

$$(3.8) \quad \mathbb{R}^N = \bigoplus_{i=0}^k W_i$$

where  $W_0$  is the subspace of  $G = \mathrm{SL}_d(\mathbb{R})$ -invariant vectors and where  $W_i$  for  $i > 0$  are irreducible subspaces (all defined over  $\mathbb{Q}$ ). This is possible because  $\rho$  is defined over  $\mathbb{Q}$ . We may replace  $R$  by a constant multiple and assume that the norm  $\|\cdot\|$  is given by  $\|v\| = \max_i \|\pi_i(v)\|$  where  $\pi_i : \mathbb{R}^N \rightarrow W_i$  denotes the  $G$ -invariant projection relative to (3.8). We let  $M > 0$  be a sufficiently large integer so that  $M\pi_i(\mathbb{Z}^N) \subset W_i(\mathbb{Z})$ .

Let  $I \subset [0, 1]$  be an interval with  $|I| \geq e^{-rt + \psi(t)}$  and with

$$I \subset \{s \in [0, 1] : \|\rho(a_t\varphi(s)g)v\| < R\}$$



for some  $v \in V(\mathbb{Z}) \setminus \{0\}$ . Then for every  $i = 0, \dots, k$ , the interval  $I$  is contained in

$$\{s \in [0, 1] : \|\rho(a_t \varphi(s)g)\pi_i(v)\| < R\}.$$

By the already proven special case, we know that for all  $i > 0$ , the union of all connected components of the above sets of length at least  $e^{-rt+\psi(t)}$  and with  $\pi_i(v) \neq 0$  has measure  $\ll MR e^{-\kappa\delta\psi(t)}$  for some  $\delta, \kappa > 0$  depending only on  $N$ . Then either (1) holds or there exists an interval  $I$  and a vector  $v \in \mathcal{U}$  as above with the additional property that  $\pi_i(v) = 0$  for all  $i > 0$ . But then  $v \in W_0(\mathbb{Z})$  as well as  $\|v\| \leq R$  so (2) holds.  $\square$

#### 4. THE SHORT INTERVALS LEMMA

For every interval  $I = [a - b, a + b] \subset \mathbb{R}$  denote by  $|I| := 2b$  the length and by  $\gamma \cdot I = [a - \gamma b, a + \gamma b]$  for all  $\gamma > 0$  the centered  $\gamma$ -dilation as in §2.1.

**Lemma 4.1** (Short Intervals). *Fix  $0 < \beta < 1 < \gamma$  and a function  $f : \mathbb{N} \rightarrow \mathbb{N}$  such that  $\lim_{n \rightarrow \infty} \frac{f(n)}{n/\log \log n} = 0$ . Let  $(A_n)_{n=0}^\infty$  be a sequence of sets of intervals in  $[-1, 1]$  such that for every  $I \in A_n$  we have  $\beta^{n+f(n)} \leq |I| \leq \beta^{n-f(n)}$  and the intervals  $\{\gamma \cdot I : I \in A_n\}$  are disjoint. Let  $B_n = \bigcup_{I \in A_n} I$ . Then*

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \#\{1 \leq n \leq N : x \in B_n\} \leq \frac{1}{\gamma} \text{ a.s. for } x \in [-1, 1]$$

Before proving the lemma we mention an important tool, introduced by Azuma, [2, Thm. 1]. The form we use here is not identical to the stated by Azuma, and we will provide a reduction from [2, Thm. 1] to Lemma 4.3.

**Definition 4.2.** Let  $(M_n)_{n=0}^\infty$  be a sequence of random variables such that  $M_n$  is  $\Sigma_n$  integrable, where  $\Sigma_n$  are a filtration of a probability space  $(\Sigma, \mu)$ . The sequence  $(M_n)_{n=0}^\infty$  is called a *Martingale* if  $\mathbb{E}(M_n | \Sigma_{n-1}) = M_{n-1}$ . It is called *sup-martingale* if  $\mathbb{E}(M_n | \Sigma_{n-1}) \leq M_{n-1}$ .

**Lemma 4.3** (Azuma's inequality). *For every sup-martingale  $(M_n)$  and a sequence of positive numbers  $c_n$  such that  $|M_n - M_{n-1}| \leq c_n$  for every  $n \geq 1$  one has*

$$(4.1) \quad \limsup_{n \rightarrow \infty} \frac{M_n}{\sqrt{2(c_1^2 + \dots + c_n^2) \log n}} \leq 1.$$

*Reduction to [2, Thm. 1]:* First, whenever  $(M_n)_{n=0}^\infty$  is a martingale, (4.1) follows from [2, Eq. (3.1)] where  $x_n = (M_n - M_{n-1})/c_n$  and  $a_{nk} = c_k$ . If  $M_n$  is only a supmartingale, we will find a martingale  $(M'_n)_{n=0}^\infty$  such that  $M_n \leq M'_n$  for all  $n \geq 0$ . Indeed, denote the filtration of the martingale by  $(\Sigma_n)_{n=0}^\infty$ . We will find a martingale  $(M'_n)_{n=0}^\infty$ , defined on the same filtration, such that  $M'_n \geq M_n$ . Then, using the lemma of  $M'_n$  we will get the desired result. We will construct  $(M'_n)_{n=0}^\infty$ , by setting  $M'_0 = M_0$  and recursively choosing  $M'_n$  by demanding  $M'_n - M'_{n-1} \geq M_n - M_{n-1}$  and  $\mathbb{E}(M'_n - M'_{n-1} | \Sigma_{n-1}) = 0$ . To this end, we show how to construct for every random variable  $X$  with  $\mathbb{E}(X) \leq 0$  another random variable  $X'$  with

- $\mathbb{E}(X') = 0$ ,
- $X' \geq X$  a.s., and
- if  $X \in [-c, c]$  almost surely then so is  $X'$ .

Then we could use this construction to define  $M'_n - M'_{n-1}$  from  $M_n - M_{n-1}$  conditioned on  $\Sigma_{n-1}$ . If  $X$  is the constant random variable, define  $X' = 0$ . If  $X \leq 0$  a.s., define  $X' = X - \mathbb{E}(X)$ . Otherwise, denote by  $\max(X)$  the minimal value such that

$\mathbb{P}(X \leq \max(X)) = 1$ . By the assumption,  $\max(X) > 0$  and  $\max(X) > \mathbb{E}(X)$ . Then we choose  $X' = \frac{-\mathbb{E}(X)}{\max(X) - \mathbb{E}(X)} \max(X) + \frac{\max(X)}{\max(X) - \mathbb{E}(X)} X$ , this is a convex combination between  $X$  and  $\max(X)$ , and it can be seen to satisfy the conditions above.  $\square$

**Claim 4.4.** *Let  $A$  be a collection of intervals such that  $\gamma I$  are disjoint for  $I \in A$ . Let  $J$  be an interval in  $\mathbb{R}$ . Then the total length of  $J \cap \bigcup_{I \in A} I$  is at most  $|J|/\gamma + 2 \max_{I \in A} |I|$ .*

*Proof.* Let  $a = \max_{I \in A} |I|$ . Without loss of generality assume all  $I \in A$  intersect  $J$  non-trivially. Then

$$\left| J \cap \bigcup_{I \in A} I \right| \leq \sum_{I \in A} |I|.$$

On the other hand, if we enlarge  $J$  by  $\gamma a$  on each side it would contain all intervals  $\gamma I$  for  $I \in A$ . Hence  $|J| + 2\gamma a \geq \sum_{I \in A} \gamma |I|$ . The result follows.  $\square$

*Proof of Lemma 4.1.* We first perform a few reductions. Fix  $\delta > 0$ . Let  $f'(n) := \left\lceil \delta \frac{n}{\log \log n} \right\rceil$ , for all but finitely many  $n \in \mathbb{N}$  we have  $f'(n) \geq f(n)$ . Without changing the correctness of the claim, replace  $A_n$  with an empty set whenever  $f'(n) < f(n)$ . Hence it is sufficient to show that if for every  $I \in A_n$  we have  $\beta^{n+f'(n)} \leq |I| \leq \beta^{n-f'(n)}$  then

$$(4.2) \quad \limsup_{N \rightarrow \infty} \frac{1}{N} \#\{1 \leq n \leq N : x \in B_n\} \leq \frac{1}{\gamma} + o_\delta(1) \text{ a.s. for } x \in [-1, 1].$$

Note that the sequence  $(n + f'(n))_{n=1}^\infty$  goes over all integers once, up to a negligible proportion. Whenever possible, replace  $A_n$  by  $A'_n = A_{n'}$  where  $n'$  satisfies  $n = n' + f(n')$  (and by the empty set otherwise). With this choice  $\beta^n \leq |I| \leq \beta^{n-2f'(n)}$  for every  $I \in A_n$ . Replacing  $\delta$  with  $2\delta$  allows us to assume that  $\beta^n \leq |I| \leq \beta^{n-f'(n)}$  for every  $I \in A_n$ .

Fix  $a > 0$  to be determined later. Fix a sequence of positive integers  $n_0 < n_1 < n_2 < \dots$  that satisfy  $n_{i+1} = n_i + f'(n_{i+1}) + 2a$ . Denote

$$I_m = \{n_{2m-1}, n_{2m-1} + 1, \dots, n_{2m} - 1\}$$

and let  $T_m = n_{2m} + a$ . Note that for every  $n \in I_m$  we have  $n \leq T_m - a$  and  $n - f'(n) \geq T_{m-1} + a$ .

Let  $J_m = \bigcup_{k=1}^m I_k$ . It is enough to show that

$$(4.3) \quad \limsup_{m \rightarrow \infty} \frac{1}{\#J_m} \sum_{i \in J_m} \mathbb{1}(x \in B_i) = \limsup_{m \rightarrow \infty} \frac{1}{\#J_m} \sum_{k=1}^m \sum_{i \in I_k} \mathbb{1}(x \in B_i) \leq \frac{1}{\gamma} + o_{\delta,a}(1) \text{ a.s. for } x.$$

This inequality together with the similar inequality for the shifted sequence  $n'_i = n_{i+1}$  proves the bound of Eq. (4.2) for  $N \in (n_i)_{i=0}^\infty$ . Since  $\frac{n_i - n_{i-1}}{n_i} \xrightarrow{i \rightarrow \infty} 0$  this sequence is dense enough to imply Eq. (4.2).

Consider a sequence of i.i.d. uniform  $x_n \in [-1, 1]$  and some  $c_0 \in [-1, 1]$ . Define  $y_m = c_0 + \sum_{i=0}^{m-1} x_i \beta^{T_i}$  for all  $m \in \mathbb{N} \cup \{\infty\}$ . Since the law of  $y_\infty$  is continuous with

positive density on  $[c_0 - \sum_{i=0}^{\infty} \beta^{T_i}, c_0 + \sum_{i=0}^{\infty} \beta^{T_i}]$ , it is sufficient to bound from above

$$\limsup_{m \rightarrow \infty} \frac{1}{\#J_m} \sum_{k=1}^m \sum_{i \in I_k} \mathbb{1}(y_\infty \in B_i).$$

for all  $c_0 \in \mathbb{R}$ . Since  $|y_\infty - y_m| < \beta^{T_m} \frac{1}{1-\beta}$  we can relate the condition of  $y_\infty$  being inside an interval with a similar condition on  $y_m$ . For every  $n \in I_m$  any interval  $I \in A_n$  is of length  $|I| \geq \beta^n$ . If  $y_\infty$  lies in  $I$  then  $y_m$  lies in  $(1 + \varepsilon)I$  for  $\varepsilon = 2\beta^a \frac{1}{1-\beta} > \beta^{T_m-n} \frac{1}{1-\beta}$ .

Hence it is sufficient to bound

$$\limsup_{m \rightarrow \infty} \frac{1}{\#J_m} \sum_{k=1}^m \sum_{n \in I_k} \mathbb{1}(y_k \in B'_n),$$

for  $B'_n = \bigcup_{I \in A_n} (1 + \varepsilon)I$ .

Condition on  $y_m$ . For every  $n \in I_{m+1}$ , the conditioned probability that  $y_{m+1} \in B'_n$  is at most  $p = \frac{1+\varepsilon}{\gamma} + (1+\varepsilon)\beta^a$ , by Claim 4.4 applied to  $J = [y_m - \beta^{T_m}, y_m + \beta^{T_m}]$ ,  $A = \{(1 + \varepsilon)I : I \in A_n\}$  and  $\gamma/(1 + \varepsilon)$  (instead of  $\gamma$ ). Hence the conditional expectation satisfies  $\mathbb{E}(\sum_{n \in I_{m+1}} \mathbb{1}(y_{m+1} \in B'_n) | y_m) \leq p \#I_{m+1}$ . Hence

$$M_m = \sum_{k=0}^m \sum_{i \in I_k} (\mathbb{1}(y_\infty \in B_i) - p),$$

is a sup-martingale, with differences  $|M_m - M_{m-1}| < \#I_m$ . Azuma's Inequality 4.3, implies that

$$(4.4) \quad \limsup_{m \rightarrow \infty} \frac{M_m}{\sqrt{2(\sum_{k=0}^m (\#I_k)^2) \log m}} \leq 1.$$

We will show that

$$(4.5) \quad \limsup_{m \rightarrow \infty} \frac{\sqrt{2(\sum_{k=0}^m (\#I_k)^2) \log m}}{\sum_{k=0}^m \#I_k} = o_\delta(1),$$

as the product of Eqs. (4.4) and (4.5) is the desired inequality. Indeed,  $\sum_{k=0}^m (\#I_k)^2 \leq \#I_m (\sum_{k=0}^m \#I_k)$ , hence it is sufficient to have

$$\limsup_{m \rightarrow \infty} \frac{\#I_m \log m}{\sum_{k=0}^m \#I_k} = o_\delta(1).$$

Denote by  $M = n_{2m}$ . Note that  $\lim_{m \rightarrow \infty} \frac{\sum_{k=0}^m \#I_k}{M} = 1/2$ . Since  $m \leq \sum_{n=1}^M \frac{1}{f'(n)}$ , So we need to have

$$\limsup_{M \rightarrow \infty} \frac{f'(M) \log \left( \sum_{n=1}^M \frac{1}{f'(n)} \right)}{M} = o_\delta(1).$$

We may replace  $f'(M)$  by  $\delta M / \log \log M$  and obtain

$$\frac{\delta \log \left( \sum_{n=1}^M \frac{\log \log n}{\delta n} \right)}{\log \log M} \leq \frac{\delta \log (\log M \log \log M) - \delta \log \delta}{\log \log M} + o_M(1) \xrightarrow{M \rightarrow \infty} \delta.$$

Altogether, using  $\varepsilon = 2\beta^a \frac{1}{1-\beta}$  and  $p = (1 + \varepsilon)/\gamma + (1 + \varepsilon)\beta^a$

$$\begin{aligned} \limsup_{m \rightarrow \infty} \frac{1}{\#J_m} \sum_{k=0}^m \sum_{i \in I_k} \mathbb{1}(y_k \in B'_i) &\leq p + \limsup_{m \rightarrow \infty} \left( \frac{M_m}{\#J_m} \right) \\ &\leq \frac{1 + \frac{2\beta^a}{1-\beta}}{\gamma} + \left( 1 + \frac{2\beta^a}{1-\beta} \right) \beta^a + 2\sqrt{\delta} \\ &= \frac{1}{\gamma} + O(\beta^a + \sqrt{\delta}) \end{aligned}$$

This implies Eq. (4.3), as desired.  $\square$

## 5. QUANTITATIVE NON-DIVERGENCE

We write

$$\text{ht} : X \rightarrow \mathbb{R}_{>0}, \quad g\Gamma \mapsto \inf\{\|gv\| : 0 \neq v \in \mathbb{Z}^d\}^{-1}$$

for the standard height function on  $X$ . This is a proper and continuous function on  $X$ . In this section, we prove the following result.

**Theorem 5.1** (Quantitative non-divergence). *Let  $\varphi : [0, 1] \rightarrow G$  be an analytic curve so that the image of  $\varphi$  in  $P \backslash G$  is not contained in any weakly constraining pencil or in any proper flag subvariety with respect to  $\{a_t\}$ . Let  $x \in X$ , and let  $\mu_{s,T} = \frac{1}{T} \int_0^T \delta_{a_t \varphi(s)x} dt$  for any  $T > 0$ . There is a constant  $\kappa > 0$  so that for almost every  $s \in [0, 1]$*

$$\limsup_{T \rightarrow \infty} \mu_{s,T}(\{y \in X : \text{ht}(y) > \varepsilon^{-1}\}) \ll \varepsilon^\kappa.$$

The authors suspect that the assumptions on  $\varphi$  can be weakened (see e.g. the difference in assumptions between [41, Thms. 1.1, 1.3]); the current proof however requires the strong assumptions given above. The proof uses ingredients from the quantitative non-divergence results of Kleinbock and Margulis [14], a construction of which we summarize in the following lemma.

**Lemma 5.2.** *For any  $t > 0$  there exist a finite collection  $\mathcal{P}_t$  consisting of sets of intervals with the following properties:*

- $|\mathcal{P}_t| \leq 1 + 3 + 3^2 + \dots + 3^{d-1}$ .
- Any  $P \in \mathcal{P}_t$  is a set of disjoint intervals.
- For any  $P \in \mathcal{P}_t$  and any  $J \in P$  there exists a primitive sublattice  $\Lambda_J < \mathbb{Z}^d$  so that  $J$  is a connected component of

$$\{s \in [0, 1] : \|a_t \varphi(s) g \Lambda_J\| \leq 1\}.$$

- For any  $s \in [0, 1]$  with  $\text{ht}(a_t \varphi(s)x) > \frac{1}{\varepsilon}$  there exists  $P \in \mathcal{P}_t$  and  $J \in P$  with  $\|a_t \varphi(s) \Lambda_J\| \leq \varepsilon$ .

Here,  $\|\Lambda\|$  denotes the covolume of a lattice  $\Lambda \subset \mathbb{R}^d$  in the subspace it spans. As mentioned, a proof of Lemma 5.2 is (at least implicitly) contained in [14]. We give here direct proof for the reader's convenience.

*Proof.* Write  $x = g\Gamma$ . We first construct a finite tree of tuples  $(I, \mathcal{F})$  where  $I \subset [0, 1]$  is a closed interval and  $\mathcal{F}$  is a partial flag of primitive sublattices of  $\mathbb{Z}^d$ . Here,  $(I, \mathcal{F}) \preceq (I', \mathcal{F}')$  if  $I' \subseteq I$  and  $\mathcal{F} \subset \mathcal{F}'$  and any element  $(I, \mathcal{F})$  in the tree satisfies

- For all  $\Lambda \in \mathcal{F}$

$$\sup_{s \in I} \|a_t \varphi(s) g \Lambda\| \leq 1.$$

- For all  $\Lambda \notin \mathcal{F}$  compatible<sup>2</sup> to  $\mathcal{F}$

$$\sup_{s \in I} \|a_t \varphi(s) g \Lambda\| > 1.$$

Moreover, the root of the tree is the tuple  $([0, 1], \{\{0\}, \mathbb{Z}^d\})$ .

To construct the tree, we may construct the children of a tuple  $(I, \mathcal{F})$  already in the tree. Consider all intervals  $J \subset I$  for which there exists  $\Lambda_J \notin \mathcal{F}$  compatible to  $\mathcal{F}$  so that  $J$  is a connected component of  $\{s \in I : \|a_t \varphi(s) g \Lambda_J\| \leq 1\}$ . If  $\mathcal{J}$  is the collection of all such intervals we take a minimal subcollection  $\mathcal{J}' \subset \mathcal{J}$  so that  $\bigcup_{J \in \mathcal{J}'} J = \bigcup_{J \in \mathcal{J}} J$  and  $\sum_{J \in \mathcal{J}'} 1_J \leq 2$ . Note that by assumption on  $(I, \mathcal{F})$  all intervals  $J \in \mathcal{J}$  are proper subintervals of  $I$ . The children of  $(I, \mathcal{F})$  are then  $(J, \mathcal{F} \cup \{\Lambda_J\})$  for  $J \in \mathcal{J}'$ ; the necessary properties are easily verified.

For any member  $(I, \mathcal{F})$  of the tree and  $(I', \mathcal{F}')$  its parent we denote by  $\Lambda_{I, \mathcal{F}}$  the primitive lattice used to construct  $\mathcal{F}$  from  $\mathcal{F}'$  (so that  $\{\Lambda_{I, \mathcal{F}}\} = \mathcal{F} \setminus \mathcal{F}'$ ). Furthermore, we let  $J(I, \mathcal{F})$  be the connected component of  $\{s \in [0, 1] : \|a_t \varphi(s) g \Lambda_{I, \mathcal{F}}\| \leq 1\}$  containing  $I$ .

We now construct a partition  $\mathcal{P}_t$  of the tree. In the following, the level  $\ell$  of the tree consists of the flags of length  $\ell+2$  (i.e.  $\ell$  is the number of lattices  $\neq \{0\}, \mathbb{Z}^d$ ); we partition each level of the tree and combine. The level zero is partitioned trivially. Now, we enumerate each level according to the left endpoint of the underlying interval. We partition the first level according to progressions of 3. For each progression of the first level, we partition the children (members of the second level) according to progressions of 3. We proceed in this manner at every level. The construction is such that whenever two members of the tree belong to the same partition, so do their parents. Also, members of the same partition are disjoint.

The following is the crucial property of this partition: Let  $P \in \mathcal{P}_t$ , let  $(I_1, \mathcal{F}_1), (I_2, \mathcal{F}_2) \in P$ , and let  $J_i = J(I_i, \mathcal{F}_i)$  for  $i = 1, 2$ . We claim that  $J_1, J_2$  are disjoint. Suppose without loss of generality that  $I_1$  is to the left of  $I_2$ . If  $(I_1, \mathcal{F}_1)$  and  $(I_2, \mathcal{F}_2)$  have the same parent  $(I, \mathcal{F})$ , then the right endpoint of  $I_1$  and the left endpoint of  $I_2$  are in  $I^\circ$ . By construction, these are the corresponding endpoints of  $J_1, J_2$  respectively and the claim follows in this case. Suppose now that  $(I_1, \mathcal{F}_1)$  and  $(I_2, \mathcal{F}_2)$  have different parents  $(I'_1, \mathcal{F}'_1)$  and  $(I'_2, \mathcal{F}'_2)$ . If the right endpoint of  $I_1$  is in  $(I'_1)^\circ$  and the left endpoint of  $I_2$  is in  $(I'_2)^\circ$ , we are done. By construction of the partition, it remains to treat the case where the right endpoint of  $I_1$  is in  $(I'_1)^\circ$  and the left endpoint of  $I_2$  coincides with the left endpoint of  $I'_2$ . Let  $(I, \mathcal{F})$  be the first common ancestor of  $(I_1, \mathcal{F}_1)$  and  $(I_2, \mathcal{F}_2)$  and let  $(I''_1, \mathcal{F}''_1) \preceq (I_1, \mathcal{F}_1)$  be the corresponding child of  $(I, \mathcal{F})$ . If  $J_1 \cap J_2 \neq \emptyset$  we also have  $I_1 \cap J_2 \neq \emptyset$  and hence  $I''_1 \cap J_2 \neq \emptyset$ . But this is impossible: there are two members of the tree that are to the right of  $(I''_1, \mathcal{F}''_1)$ ; one of these would be rendered obsolete by  $J_2$  contradicting minimality in the construction of the tree.

Given a member  $(I, \mathcal{F})$  of the tree (the tree depends on  $t$ ) we write

$$\text{Bad}_t(I, \mathcal{F}; \varepsilon) = \{s \in I : \|a_t \varphi(s) g \Lambda_{I, \mathcal{F}}\| \leq \varepsilon\}.$$

<sup>2</sup>For a partial flag  $\mathcal{F}$  of primitive lattices in  $\mathbb{Z}^d$  and a primitive lattice  $\Lambda \subset \mathbb{Z}^d$  we say that  $\Lambda$  is compatible to  $\mathcal{F}$  if  $\mathcal{F} \cup \{\Lambda\}$  is flag.

Furthermore, we write for  $P \in \mathcal{P}_t$

$$(5.1) \quad \begin{aligned} \text{Bad}_{t,P}(\varepsilon) &:= \bigcup_{(I,\mathcal{F}) \in P} \text{Bad}_t(I,\mathcal{F};\varepsilon) \\ &\subset \bigcup_{(I,\mathcal{F}) \in P} \{s \in J(I,\mathcal{F}) : \|a_t \varphi(s) g \Lambda_{I,\mathcal{F}}\| \leq \varepsilon\} =: \text{Bad}'_{t,P}(\varepsilon) \end{aligned}$$

where the latter union is still disjoint as established above. Lastly, we set  $\text{Bad}_t(\varepsilon) = \bigcup_{P \in \mathcal{P}_t} \text{Bad}_{t,P}(\varepsilon)$ .

Now let  $s \in [0, 1]$  with  $\|a_t \varphi(s)v\| \leq \varepsilon$  for some  $v \in \mathbb{Z}^d$  primitive. We claim that  $s \in \text{Bad}_t(\varepsilon)$ . The proof proceeds by induction. To begin, observe that  $s \in I_1$  for some  $(I_1, \mathcal{F}_1)$  of level one in the tree. Writing  $\mathcal{F}_1 = \{\{0\}, \Lambda_1, \mathbb{Z}^d\}$  we are done if  $\mathbb{Z}v = \Lambda_1$  or if  $\|a_t \varphi(s) g \Lambda_1\| \leq \varepsilon$ . Otherwise, there exists a child  $(I_2, \mathcal{F}_2)$  of  $(I_1, \mathcal{F}_1)$  with  $s \in I_2$ . Indeed, either  $v \in \Lambda_1$  (in which case one may add  $\mathbb{Z}v$  to the flag) or  $\Lambda_1$  and  $v$  generate a lattice  $\Lambda$  with  $\|a_t \varphi(s) g \Lambda\| \leq \varepsilon$  (and the same estimate is true for the primitive lattice  $\mathbb{Q}\Lambda \cap \mathbb{Z}^d$ ). Suppose now that  $s \in I$  for some  $(I, \mathcal{F})$  in the tree where the flag  $\mathcal{F}$  is given by  $\Lambda_0 = \{0\} \subset \Lambda_1 \subset \dots \subset \Lambda_i \subset \mathbb{Z}^d$  and where  $\|a_t \varphi(s) g \Lambda_j\| > \varepsilon$  for all  $j \leq i$ ,  $j \neq j_0$ . Here,  $\Lambda_{j_0}$  is the primitive lattice added to the flag of the parent of  $(I, \mathcal{F})$  to obtain  $\mathcal{F}$ . If  $\|a_t \varphi(s) g \Lambda_{j_0}\| \leq \varepsilon$ , we are done. Otherwise, let  $j \in \{0, \dots, i\}$  be the largest index with  $v \notin \Lambda_j$ . The primitive lattice  $\Lambda$  generated by  $\Lambda_j$  and  $v$  is compatible with the flag and satisfies  $\|a_t \varphi(s) g \Lambda\| \leq \varepsilon$ . In particular,  $\mathcal{F} \cup \{\Lambda\} \supsetneq \mathcal{F}$  concluding the induction step. This also concludes the lemma.  $\square$

*Proof of Theorem 5.1.* Write  $x = g\Gamma$ . Throughout the argument we may assume that  $\varepsilon > 0$  is sufficiently small. We begin first with  $t \in \mathbb{N}$  and choose an arbitrary enumeration of  $\mathcal{P}_t = \{P_{t,1}, \dots, P_{t,k}\}$  as constructed in Lemma 5.2 where  $k \leq 1 + 3 + \dots + 3^{d-1} = k_d$ . We may assume that  $k = k_d$  by setting  $P_{t,j} = \emptyset$  for  $k < j \leq k_d$ . Moreover, we set

$$\text{Bad}_{t,j}(\varepsilon) = \bigcup_{J \in P_{t,j}} \{s \in J : \|a_t \varphi(s) \Lambda_J\| \leq \varepsilon\}.$$

We show that for every  $k$  and for almost every  $s$

$$(5.2) \quad \limsup_{T \rightarrow \infty} \frac{1}{T} \#\{1 \leq t \leq T : s \in \text{Bad}_{t,j}(\varepsilon)\} \ll \varepsilon^*.$$

Applying Proposition 3.4 to all exterior products of the standard representation and the variety of pure vectors therein, we obtain (as the standard representation is irreducible) that the measure of the union of all intervals in  $P_{t,j}$  of length at least  $e^{-rt+\sqrt{t}}$  is  $\ll e^{-*\sqrt{t}}$ . Let  $\text{Bad}'_{t,j}(\varepsilon) \subset \text{Bad}_{t,j}(\varepsilon)$  be the set of points not contained in that union. The Borel Cantelli lemma implies that (5.2) is equivalent to

$$(5.3) \quad \limsup_{T \rightarrow \infty} \frac{1}{T} \#\{1 \leq t \leq T : s \in \text{Bad}'_{t,j}(\varepsilon)\} \ll \varepsilon^*$$

for almost every  $s$ .

We use Lemma 4.1 to establish (5.3) after further partitioning. By Lemma 2.9 for any  $J$  the number of connected components of  $\{s \in J : \|a_t \varphi(s) g \Lambda_J\| \leq \varepsilon\}$  is at most  $M$  for some integer  $M$  depending only on  $\varphi$ . We may hence partition  $\text{Bad}'_{t,j}(\varepsilon)$  into at most  $M$  disjoint subsets, denoted for simplicity by  $B_t^i$ , so that for

any  $J \in P_{t,j}$

$$B_t^i \cap J \subset \{s \in J : \|a_t \varphi(s) g \Lambda_J\| \leq \varepsilon\}$$

is a connected component. At this point, one would like to invoke Lemma 4.1, but observe that the lower bound  $|B_t^i \cap J| \geq e^{-rt - \sqrt{t}}$  for  $t$  large enough is not necessarily satisfied. To remedy that, we replace  $B_t^i \cap J$  by the connected component of  $\{s \in J : \|a_t \varphi(s) g \Lambda_J\| \leq 2\varepsilon\}$  containing it. Note that by Lipschitz continuity of  $\varphi$  this connected component has size  $\gg e^{-rt}$ , and we continue with these intervals. Recall that by Lemma 2.9 the function  $s \mapsto \|a_t \varphi(s) \Lambda_J\|$  is  $(C, \alpha)$ -good for any  $J$  where  $C$  and  $\alpha$  depend only on  $\varphi$ . Hence, by Lemma 2.7 the conditions of Lemma 4.1 are satisfied for  $\gamma = \frac{1}{C(2\varepsilon)^\alpha}$ . This yields that for almost every  $s$

$$\limsup_{T \rightarrow \infty} \frac{1}{T} \#\{1 \leq t \leq T : s \in B_t^i\} \ll \varepsilon^\alpha.$$

Summing over  $i$  the estimates (5.2), (5.3) follow and after summing over  $1 \leq j \leq k_d$  for almost every  $s \in [0, 1]$

$$\limsup_{T \rightarrow \infty} \frac{1}{T} \sum_{t=1}^T \mathbf{1}_{\{y \in X : \text{ht}(y) > \varepsilon^{-1}\}}(a_t \varphi(s)x) \ll \varepsilon^\alpha.$$

For any  $\delta \in (0, 1)$  and any point  $y \in X$  we have  $\text{ht}(a_\delta y) \asymp \text{ht}(y)$  and hence the continuous time result in the theorem follows.  $\square$

## 6. UNIPOTENT INVARIANCE

The proof we present here for unipotent invariance is a variant of the proof given at [6, Prop. 3.1], [37, Prop. 2.2]. Instead of the estimate of correlation they use, we use Martingales directly.

For every locally compact separable metric space  $X$ , denote by  $\Delta(X)$  the set of Borel probability measures on  $X$ . Note that there is a metric  $d(\cdot, \cdot)$  inducing the weak\*-topology on  $\Delta(X)$ .

**Definition 6.1.** For every probability space  $(\Omega, \Sigma, \mathbb{P})$  and a measurable space  $X$ , denote by  $RV(X; \Sigma)$  the set of  $X$  valued  $\Sigma$ -measurable random variables.

**Lemma 6.2.** *Let  $X$  be a locally compact metric space and let  $(\Sigma_n)_{n=0}^\infty$  be a filtration of a probability space  $(\Omega, \Sigma, \mathbb{P})$  by countably generated  $\sigma$ -algebras. Let  $(p_n)_{n=0}^\infty$  be a sequence of random points in  $X$ , where  $p_n \in RV(X; \Sigma_n)$ . Let  $M \in RV(\text{Hom}_{\text{cont}}(X, \Delta(X)); \Sigma)$  be a random continuous function  $x \mapsto M_x$ . Assume that the conditional distribution  $\text{Law}(p_{n+1} | \Sigma_n) \in RV(\Delta(X), \Sigma_n) \subseteq RV(\Delta(X), \Sigma)$  satisfy that with probability 1 we have*

$$d(\text{Law}(p_{n+1} | \Sigma_n), M_{p_n}) \xrightarrow{n \rightarrow \infty} 0.$$

*Then with probability 1 every partial limit  $\mu$  of  $\mu_n = \frac{1}{n} \sum_{k=1}^n \delta_{p_k} \in RV(\Delta(X); \Sigma_n)$  is invariant under convolution with  $M$ , that is:*

$$\mu = \int_X M_x d\mu(x).$$

*Proof.* To show that almost surely every partial limit  $\mu$  of  $\mu_n$  satisfies  $\mu = \int_X M_x d\mu(x)$ , we should show that for a countable dense set of functions  $F \subset C_c(X)$  and for every  $f \in F$  we have  $\int_X M_x(f) d\mu(x) = \int_X f d\mu$ . Here  $M_x(f) = \int_X f dM_x$ . To show

that it is sufficient to prove that for every  $f \in F$ , almost surely  $\int_X M_x(f) d\mu_n(x) - \int_X f d\mu_n \xrightarrow{n \rightarrow \infty} 0$ . By assumption,

$$(6.1) \quad \int_X M_x(f) d\mu_n(x) - \frac{1}{n} \sum_{k=1}^n \text{Law}(p_{k+1} | \Sigma_k)(f) \xrightarrow{n \rightarrow \infty} 0.$$

On the other hand, define

$$B_n := \sum_{k=1}^n \text{Law}(p_{k+1} | \Sigma_k)(f) - \sum_{k=1}^n f(p_{k+1}) \in RV(\mathbb{R}, \Sigma_n).$$

The process  $B_n$  is a martingale, as the step  $n$  increment is a random variable minus its conditional expectation. Since its increments are bounded Azuma's inequality [2] implies that  $\frac{B_n}{n} \xrightarrow{n \rightarrow \infty} 0$ . This, together with Eq. (6.1) implies the desired result.  $\square$

For every  $T, s$  denote by  $\mu_{s,T} = \frac{1}{T}(t \mapsto a_t \varphi(s) \Lambda)_* m_{[0,T]}$  the pushforward of the uniform measure on  $[0, T]$  where  $\Lambda \in X_d = \text{SL}_d(\mathbb{R}) / \text{SL}_d(\mathbb{Z})$  is fixed. Also, recall the definition of  $Y_s \in \mathfrak{g}^+$  from Lemma 2.4.

**Theorem 6.3** (Unipotent invariance). *For almost every  $s$  every partial limit of  $\mu_{s,T}$  is invariant under the unipotent flow  $x \mapsto \exp(xY_s)$ .*

*Proof.* Fix  $0 < \lambda < 1$ , and let  $x_0, x_1, \dots \in [0, 1]$  be uniform independent random variables. Define the sigma-algebras  $\Sigma_n = \sigma(x_0, \dots, x_n)$  for  $n \geq 1$  and  $\Sigma = \sigma((x_i)_{i=0}^\infty)$ . Let  $s = (1 - \lambda) \sum_{i=0}^\infty \lambda^i x_i \in RV([0, 1], \Sigma)$ . Then  $s$  is chosen at random via a continuous distribution with support  $[0, 1]$ , hence it is sufficient to show that every partial limit of  $\mu_{s,T}$  is invariant under  $x \mapsto \exp(xY_s)$  for almost every  $s$  chosen in this way. Let  $s_n = (1 - \lambda) \sum_{i=0}^n \lambda^i x_i \in RV([0, 1], \Sigma_n)$ ,  $T_0 = |\log \lambda|/r \in \mathbb{R}$  and for every  $n$  consider the random point  $p_n = a_{nT_0} \varphi(s_n) \Lambda \in RV(X, \Sigma_n)$ , and the random measures  $\mu_n = \frac{1}{n} \sum_{i=0}^{n-1} \delta_{p_i} \in RV(\Delta(X), \Sigma_n)$  and  $\mu'_n = \frac{1}{n} \sum_{i=0}^{n-1} \delta_{a_{iT_0} \varphi(s) \Lambda} \in RV(\Delta(X), \Sigma)$ . Since  $|s - s_n| = O(\lambda^{n+1})$  it follows that  $p_i$  and  $a_{iT_0} \varphi(s) \Lambda$  are  $O(\lambda)$ -close (cf. Lemma 2.4). Similarly,  $a_{-t} p_i$  and  $a_{-t} a_{iT_0} \varphi(s) \Lambda$  are  $O(e^{-rt} \lambda)$ -close whenever  $t < iT_0$ . Hence  $(a_{-t})_* \mu_n$  is  $O(e^{-rt} \lambda + \frac{1}{\sqrt{n}})$  close to  $(a_{-t})_* \mu'_n$  for  $t < \sqrt{n} T_0 / 3$ .

Suppose we show that every partial limit of  $\mu_n$  is  $\exp(xY_s)$ -invariant. Let  $\mu'$  be a partial limit of the measures  $\mu'_n$  with respect to some subsequence  $n_k$ . Without loss of generality, assume that  $\mu_{n_k}$  converge to a measure  $\mu$ , which by assumption is  $\exp(xY_s)$ -invariant. In particular, also  $(a_{-t})_* \mu$  is  $\exp(xY_s)$ -invariant for every  $t \geq 0$ . Moreover,  $(a_{-t})_* \mu'$  is  $O(e^{-rt} \lambda)$  close to  $(a_{-t})_* \mu$  for all  $t$ . In addition, it is clear that  $\mu'$  is invariant under  $(a_{-t})_*$  for  $t = T_0$ . These facts imply that  $\mu'$  is invariant under  $\exp(xY_s)$ . Since every partial limit of  $\mu_{s,T}$  is a convolution of  $\mu'$  with a uniform distribution of  $a_{[0,T_0]}$ , for  $T = nT_0$ , this would imply that every partial limit of  $\mu_{s,T}$  is  $\exp(xY_s)$ -invariant. Hence the proof is reduced to proving the following claim:

**Claim 6.4.** *Every partial limit of  $\mu_n$  is invariant  $x \mapsto \exp(xY_s)$ .*

To prove the claim we introduce a new operator, and we show that every partial limit of  $\mu_n$  is invariant under the new operator. Let  $\nu_s \in RV(\Delta(G), \Sigma)$  denote the following random measure on  $G$ : choose  $y$  uniformly on  $[0, \lambda(1 - \lambda)]$  and set  $\nu_s = \text{Law}(a_{T_0} \exp(yY_s))$ . Alternatively, if  $\nu'_s = \text{Law}(\exp(yY_s)) \in RV(\Delta(G^+), \Sigma)$



then  $\nu_s = \delta_{a_{T_0}} * \nu'_s$ . Now, the law  $Law(p_{n+1}|\Sigma_n)$  can be computed as follows. We know  $s_n$ , hence by Lemma 2.4

$$\begin{aligned} p_{n+1} &= a_{(n+1)T_0} \varphi(s_{n+1}) \Lambda \\ &= a_{T_0} (I + O(\lambda^{n+1})) \exp((1-\lambda)\lambda x_{n+1} Y_{s_n}) p_n \\ &= (I + O(\lambda^n)) a_{T_0} \exp((1-\lambda)\lambda x_{n+1} Y_{s_n}) p_n \\ &= (I + O(\lambda^n)) a_{T_0} \exp((1-\lambda)\lambda x_{n+1} Y_s) p_n. \end{aligned}$$

We can see that  $d(Law(p_{n+1}|\Sigma_n), \nu_s * \delta_{p_n}) \xrightarrow{n \rightarrow \infty} 0$ . Now Lemma 6.2 shows that every partial limit  $\mu$  of  $\mu_n$  is invariant to convolution with  $\nu_s$ . The theorem is now reduced to the following claim:

**Claim 6.5.** *Every measure  $\mu$  for which  $\nu_s * \mu = \mu$  is invariant under  $x \mapsto \exp(xY_s)$ .*

Inductively,  $\mu = \nu_s^{*n} * \mu$ . Note that as  $\nu_s = \delta_{a_{T_0}} * \nu'_s$  and  $\nu'_s = Law(\exp(yY_s))$  for  $y \sim Unif([0, \lambda(1-\lambda)])$  we get that  $Law(\exp(\kappa y Y_s))$  convolution-commutes with  $\nu'_s$  for all  $\kappa > 0$  and

$$\delta_{a_{T_0}} * Law(\exp(AyY_s)) = Law(\exp(\lambda^{-1}AyY_s)) * \delta_{a_{T_0}},$$

for every  $A \in \mathbb{R}$ . Hence

$$\begin{aligned} \mu &= \nu_s^{*n} * \mu = \nu_s^{*(n-1)} * \delta_{a_{T_0}} * Law(\exp(yY_s)) * \mu \\ &= \dots = Law(\exp(\lambda^{-n}yY_s)) * \nu_s^{*(n-1)} * \delta_{a_{T_0}} * \mu. \end{aligned}$$

Hence,  $\mu$  is the convolution of  $Law(\exp(\lambda^{-n}yY_s))$  with some other measure for every  $n$ . This implies that  $\mu$  is  $\exp(xY_s)$ -invariant for all  $x \in \mathbb{R}$ .  $\square$

## 7. AVOIDING SINGULAR SETS AND A PROOF OF THEOREM 1.10

In this section, we combine the linearization technique developed by Dani and Margulis [8] with the methods from the previous sections to establish that for almost every point on the curve the trajectory under the diagonal flow does not accumulate on a singular set.

**7.1. The linearization technique.** Consider the one-parameter unipotent subgroup  $W = \{\exp(xY_0) : x \in \mathbb{R}\} < G^+$ . By Theorem 6.3, for almost every  $s$  any limit of the measures

$$\nu_{s,T} = \frac{1}{T} \int_0^T \delta_{z(s)a_t \varphi(s)x} dt = z(s) * \mu_{s,T}$$

is  $W$ -invariant. Here,  $x \in X$  will remain fixed.

Let  $\mathcal{H}$  be the collection of all closed connected subgroups of  $G = \mathrm{SL}_d(\mathbb{R})$  such that  $H \cap \Gamma < H$  is a lattice and such that the subgroup of  $H$  generated by all unipotent one-parameter subgroups acts ergodically on  $H\Gamma/\Gamma$ . It is easy to see from the Borel density theorem that  $\mathcal{H}$  is countable (in fact, this would also be true if  $\Gamma$  were non-arithmetic [8, Prop. 2.1]). We define for  $W$  as above and  $H \in \mathcal{H}$

$$N(W, H) = \{g \in G : W \subset gHg^{-1}\},$$

$$S(W, H) = \bigcup_{\mathcal{H} \ni H' \subsetneq H} N(W, H').$$

The following is a consequence of Ratner's measure classification [27] (see also [24, Thm. 2.2]):

**Theorem 7.1.** *Let  $\mu$  be a  $W$ -invariant probability measure on  $X = G/\Gamma$  with  $\mu(N(W, H)\Gamma) = 0$  for all proper subgroups  $H \in \mathcal{H}$  of  $G$ . Then  $\mu$  is the  $G$ -invariant probability measure on  $X$ .*

By Theorems 5.1 and 6.3, the following proposition thus implies Theorem 1.10.

**Proposition 7.2.** *For almost every  $s \in [0, 1]$  and any weak\*-limit  $\nu$  of the measures  $\nu_{s,T}$  (as  $T \rightarrow \infty$ ) we have  $\nu(N(H, W)\Gamma/\Gamma) = 0$ .*

As mentioned we apply the linearization technique to prove Proposition 7.2. For any  $H \in \mathcal{H}$  we consider the  $G$ -space (via the adjoint representation)

$$V_H = \bigwedge^{\dim(H)} \mathfrak{g}$$

and a unit vector  $v_H \in \bigwedge^{\dim(H)} \mathfrak{h}$ . Whenever not implicitly clear, we write  $\rho_H : G \rightarrow \mathrm{SL}(V_H)$  for the representation and  $\eta_H : G \rightarrow V_H$ ,  $g \mapsto g.v_H$  for the orbit map at  $v_H$ . Observe that  $N(W, H) = \eta_H^{-1}(A_H)$  where

$$A_H = \{v \in V_H : w \wedge v = 0\}.$$

For any  $R > 0$  we write  $B_R = \{v \in V_H : \|v\| \leq R\}$ . Finally, we define

$$S_{H,R} = \{g \in G : \exists \gamma \in \Gamma \text{ with } \eta_H(g\gamma) \neq \pm \eta_H(g) \text{ and } \eta_H(g\gamma), \eta_H(g) \in B_R \cap A_H\}.$$

The following proposition is a consequence of the fact that

$$G/\mathrm{Stab}_\Gamma(v_H) \rightarrow G/\Gamma \times V_H, \quad g \mapsto (g\Gamma, \eta_H(g))$$

is proper.

**Proposition 7.3** ([24, Prop. 3.2]). *Let  $R > 0$ . Then  $S_{H,R}$  is a subset of  $S(W, H)$  with  $S_{H,R}\Gamma \subset X$  closed. For any compact subset  $K \subset X \setminus S_{H,R}\Gamma$  there exists a neighborhood  $\mathcal{O}$  of  $B_R \cap A_H \subset V_H$  such that any coset  $\eta_H^{-1}(\mathcal{O})\Gamma \cap K$  has a unique representative in  $\eta_H^{-1}(\mathcal{O})$ .*

**Proposition 7.4.** *Let  $r > 0$ ,  $H \in \mathcal{H}$ , and let  $\varepsilon > 0$ . Then there exists a closed subset  $\mathcal{S} \subset S(W, H)$  so that for any compact subset  $K \subset X \setminus \mathcal{S}\Gamma$  there exists an open neighborhood  $\mathcal{O}'$  of  $B_r \cap A_H$  with the following property. For almost every  $s \in [0, 1]$  and  $T$  sufficiently large*

$$\nu_{s,T}(K \cap \pi(\eta_H^{-1}(\mathcal{O}')) \leq \varepsilon.$$

*Proof.* In the following, we first take times  $t \in \mathbb{N}$ . Let  $\delta \in (0, 1)$  (to be chosen polynomial in  $\varepsilon$  later) and  $R = r/\delta$ . Set  $\mathcal{S} = S_{H,R}$ , let  $K \subset X \setminus \mathcal{S}\Gamma$  be compact, and choose  $\mathcal{O}$  as in Proposition 7.3. Let  $\beta > 0$  so that  $\mathcal{O}$  contains

$$\{v \in V_H : \|v\| < R + \beta, \|w \wedge v\| < \beta\}.$$

By replacing  $\mathcal{O}$  we may assume that it is equal to the above displayed set. We will show that the open neighborhood

$$\mathcal{O}' = \{v \in V_H : \|v\| < \delta(R + \beta), \|w \wedge v\| < \delta\beta\} = \delta\mathcal{O}.$$

of  $B_r \cap A_H$  satisfies the requirements of the proposition.

Consider the set of points

$$J_t = \{s \in [0, 1] : z(s)a_t\varphi(s)x \in K \cap \pi(\eta_H^{-1}(\mathcal{O}'))\}$$

Note that for any  $s \in J_t$  there exists a unique vector  $v_s \in \eta_H(x)$  (up to signs) such that  $z(s)a_t\varphi(s).v_s \in \mathcal{O}$ . We let  $I_t(s)$  be the largest interval containing  $s$  such that

$z(s')a_t\varphi(s').v_s \in \mathcal{O}$  for all  $s' \in I_t(s)$ . By construction, these intervals are either equal or disjoint.

We will say that  $s$  is special at time  $t \in \mathbb{N}$  if  $s \in J_t$  and  $I_t(s)$  has length at least  $e^{-rt+\sqrt{t}}$  and is special if it is special for infinitely many times  $t \in \mathbb{N}$ . We claim that the set of special points has zero measure. As  $s \mapsto z(s)$  is bounded, there exists a constant  $C_1 > 0$  so that for any  $t > 0$

$$\begin{aligned} I_t(s) &\subset \{s \in [0, 1] : \|z(s)a_t\varphi(s).v_s\| \leq R + \beta\} \\ &\subset \{s \in [0, 1] : \|a_t\varphi(s).v_s\| \leq C_1(R + \beta)\}. \end{aligned}$$

Also, note that no nontrivial pure vector in  $V_H$  is  $G$ -invariant as  $\mathfrak{g}$  is simple. Hence, Proposition 3.4 applied to the variety of pure vectors in  $V_H$  implies that the measure of the set of special points at time  $t$  is  $\ll e^{-\star\sqrt{t}}$  where the implicit constant depends on  $R$  (and hence on  $\delta$ ) and  $x$ . The claim thus follows from the Borel-Cantelli lemma.

Let  $I$  be one of the non-special intervals  $I_t(s)$ . Lemma 2.9 shows that the functions  $s' \mapsto \|a_t\varphi(s').v_s\|$  and  $s' \mapsto \|w \wedge a_t\varphi(s').v_s\|$  are  $(C, \alpha)$ -good where  $C$  and  $\alpha$  only depend on  $\dim(H)$  and  $\varphi$ . In particular,

$$f_s : s' \mapsto \max\{(R + \beta)^{-1}\|a_t\varphi(s').v_s\|, \beta^{-1}\|w \wedge a_t\varphi(s').v_s\|\}$$

is  $(C, \alpha)$ -good. Moreover, any set of the form  $\{s' : |f_s(s')| < \kappa\}$  has at most  $M^2$  connected components where  $M$  is as in Lemma 2.9 for the representation  $\rho_H$ . We define for each  $k \leq M^2$  a collection of intervals  $\mathcal{I}_{t,k}$  by picking for each interval  $I_t(s)$  the  $k$ -th connected component of  $\{s' \in I_t(s) : |f_s(s')| < \delta\}$  (if it exists). By Lemma 2.7 we have that for any  $I' \in \mathcal{I}_{t,k}$  the dilation  $\frac{1}{C\delta^\alpha}.I'$  is contained in one of the intervals  $I_t(s)$ . We let  $B_{t,k}$  be the union of all intervals in  $\mathcal{I}_{t,k}$ . By Lemma 4.1, we have for almost every  $s \in J$

$$\limsup_{T \rightarrow \infty} \frac{1}{T} \#\{1 \leq t \leq T : s \in B_{t,k}\} \ll \delta^\alpha.$$

Summing over  $k$ , for almost every  $s \in [0, 1]$

$$\limsup_{T \rightarrow \infty} \frac{1}{T} \#\{1 \leq t \leq T : z(s)a_t\varphi(s).x \in K \cap \pi(\eta_H^{-1}(\mathcal{O}'))\} \ll \delta^\alpha,$$

which concludes the discrete-time case of the lemma for a suitable choice of  $\delta = \delta(\varepsilon)$ . To obtain the lemma for continuous time, it suffices to observe that by Lipschitz continuity there exists a constant  $C_2 \geq 1$  (depending on  $z(\cdot)$ ) so that  $z(s)a_{\lfloor t \rfloor}\varphi(s) \in \mathcal{O}'$  implies  $z(s)a_t\varphi(s) \in C_2\mathcal{O}'$ .  $\square$

*Proof of Proposition 7.2.* For any  $H \in \mathcal{H}$  write  $\text{Bad}_H \subset [0, 1]$  for the set of points  $s \in [0, 1]$  for which the sequence of measures  $\nu_{s,T}$  has a weak\*-limit  $\nu$  satisfying  $\nu(N(W, H)\Gamma/\Gamma) > 0$ . For any compact subset  $K' \subset N(W, H)$  and any  $\kappa > 0$  the sets  $\text{Bad}_H(K', \kappa) \subset \text{Bad}_H$  are similarly defined with the inequality  $\nu(K'\Gamma/\Gamma) > \kappa$ . These are Borel measurable subsets. Note that  $\text{Bad}_H = \bigcup_{K', \kappa} \text{Bad}_H(K', \kappa)$ . Let  $H \in \mathcal{H}$  be minimal (with respect to inclusion) so that  $\text{Bad}_H$  has a positive measure. In particular,  $\text{Bad}_{H'}$  is a nullset for any  $\mathcal{H} \ni H' \subsetneq H$  and for almost every  $s \in [0, 1]$  any limit  $\nu$  of the measure  $\nu_{s,T}$  satisfies  $\nu(S(W, H)\Gamma) = 0$ . We suppose by contradiction that  $H \neq G$ . Choose  $K' \subset N(W, H) \setminus S(W, H)$  compact and  $\kappa > 0$  such that  $\text{Bad}_H(K', \kappa)$  has positive measure. Let  $r > 0$  be such that  $\eta_H(K') \subset B_r$ . Let  $\mathcal{S}$  be as in Proposition 7.4 for  $\varepsilon = \frac{\kappa}{2}$  and choose a compact neighborhood

$K \subset X \setminus \mathcal{S}\Gamma$  of  $K'\Gamma$ . By Proposition 7.4 there exists an open neighborhood  $\mathcal{O}$  of  $\eta_H(K')$  so that for almost every  $s$  and  $T$  sufficiently large

$$\nu_{s,T}(K \cap \eta_H^{-1}(\mathcal{O})\Gamma) \leq \frac{\kappa}{2}.$$

For  $T \rightarrow \infty$  this implies that any weak\*-limit  $\nu$  of  $\nu_{s,T}$  satisfies  $\nu(K'\Gamma) \leq \frac{\kappa}{2}$ . This is a contradiction to  $\text{Bad}(K', \kappa)$  having a positive measure and the proposition follows.  $\square$

## 8. PROOF OF THE APPLICATIONS

**8.1. Proof of Theorem 1.1.** Let  $X = \text{SL}_{d+1}(\mathbb{R})/\text{SL}_{d+1}(\mathbb{Z})$ . Write  $\|\cdot\|_\infty$  for the supremum norm on  $\mathbb{R}^d$  and set for  $\mu \in (0, 1]$

$$X(\mu) = \{\Lambda \in X : \exists \lambda \in \Lambda \setminus \{0\} \text{ with } \|\lambda\|_\infty \leq \mu\}$$

Note that  $X(1) = X$  and that  $X(\mu)$  for  $\mu < 1$  is a subset of (Haar) measure in  $(0, 1)$ . We write  $f(\mu)$  for that measure.

Define  $a_t = \text{diag}(e^{dt}, e^{-t}, \dots, e^{-t})$  for  $t \in \mathbb{R}$  and for  $\xi \in \mathbb{R}^d$

$$u(\xi) = \begin{pmatrix} 1 & \xi_1 & \cdots & \xi_d \\ & 1 & & \\ & & \ddots & \\ & & & 1 \end{pmatrix} \in \text{SL}_{d+1}(\mathbb{R}).$$

By the Dani correspondence, (1.1) for  $\xi$  and  $N$  has a non-zero solution if and only if  $a_{\log(N)}u(\xi)\mathbb{Z}^{d+1} \in X(\mu)$ . As  $X(\mu)$  is Jordan-measurable (with respect to the Haar measure), Theorem 1.10 implies that for almost every  $s$  and for every  $\mu > 0$

$$(8.1) \quad \begin{aligned} & \frac{1}{\log(N)} \int_1^N \frac{1_{X(\mu)}(a_{\log(n)}u(\phi(s))\mathbb{Z}^{d+1})}{n} dn \\ &= \frac{1}{\log(N)} \sum_{n=1}^N \frac{1_{X(\mu)}(a_{\log(n)}u(\phi(s))\mathbb{Z}^{d+1})}{n} + o(1) \end{aligned}$$

converges to  $f(\mu)$ . This together with Dani correspondence implies Theorem 1.1.

**8.2. Proof of Theorem 1.4.** Let  $s \in \mathbb{R}$ . Define

$$\begin{aligned} C_s &= \left\{ (c_i)_{i=1}^d \in \mathbb{R}^d : |c_i| \leq 1 \text{ for } i = 1, \dots, d, \left| \sum_{i=1}^d c_i s^i \right| \leq 1 \right\} \subset \mathbb{R}^d, \\ M_{s,\mu} &= \left\{ (\eta, (c_i)_{i=1}^d) \in \mathbb{R} \times \mathbb{R}^d : (c_i)_{i=1}^d \in C_s, \mu \left| \sum_{i=1}^d i c_i s^{i-1} \right| \geq |\eta| \right\}, \\ f(s, \mu) &= m_{X_{d+1}}(\{\Lambda \in X : \Lambda \cap M_X \neq \{0\}\}). \end{aligned}$$

We will prove Theorem 1.4 with this function  $f$ . The sets  $C_s$  and  $M_{s,\mu}$  are illustrated respectively in Figures 1 and 2.

Let  $a_t$  be as in (1.3) for  $n = d, m = 1$ , and set for any  $s \in \mathbb{R}$

$$\varphi(s) = \begin{pmatrix} 1 & s & \cdots & s^d \\ & 1 & & \\ & & \ddots & \\ & & & 1 \end{pmatrix}.$$




FIGURE 1. The set  $C_s$  for  $d = 2, s = \sqrt{2}$ .




FIGURE 2. The set  $M_{s,\mu}$  for  $d = 2, s = \sqrt{2}, \mu = 1/4$ .

Let  $\Lambda_{s,t} := a_t \varphi(s) \mathbb{Z}^{d+1}$ . For every  $\varepsilon > 0$  denote by  $M_{s,\mu}^\varepsilon$  the  $\varepsilon$ -neighborhoods of  $M_{s,\mu}$ , and  $M_{s,\mu}^{-\varepsilon}$  the set of  $v \in M_{s,\mu}$  such that the  $\varepsilon$ -ball around  $v$  is contained in  $M_{s,\mu}$ .

**Claim 8.1.** *For every  $s \in \mathbb{R}, \mu > 0, \varepsilon > 0$  and  $t > t_0(s, \mu, \varepsilon)$  the following occur:*

- (i) Let  $s' \in \mathbb{R}$  be a real number with  $|s' - s| < \mu e^{-t(1+1/d)}$  and  $\sum_{i=0}^d b_i (s')^i = 0$  for some  $(b_i)_{i=0}^d \in \mathbb{Z}^{d+1}$  with  $|b_i| < e^{t/d}$  for all  $i = 0, \dots, d$ . Then the vector  $a_t \varphi(s) ((b_i)_{i=0}^d)$  lies in  $\Lambda_{s,t} \cap M_{s,\mu}^\varepsilon$ .
- (ii) Let  $(\eta, (c_i)_{i=1}^d) \in \Lambda_{s,t} \cap M_{s,\mu}^{-\varepsilon}$ . Then the vector

$$(b_i)_{i=0}^d = (a_t \varphi(s))^{-1} (\eta, (c_i)_{i=1}^d) \in \mathbb{Z}^{d+1}$$

satisfies that  $|b_i| < e^{t/d}$  for all  $i = 0, \dots, d$  and  $\sum_{i=0}^d b_i x^i$  has a real root  $s'$  with  $|s' - s| < \mu e^{-t(1+1/d)}$ .

The proof of Claim 8.1 is a straightforward but lengthy computation, and we leave it to the end of the subsection. We are now ready to prove Theorem 1.4.

*Proof of upper bound in Theorem 1.4.* For an upper bound of  $\overline{\mathcal{A}}_s(\mu)$ , as in (8.1), it is sufficient to bound

$$(*)_T = \frac{1}{T} \int_0^T \mathbb{1}((1.2) \text{ holds for } N' = e^{t/d}) dt.$$

By Claim 8.1,

$$(*)_T \leq \frac{1}{T} \int_0^T \mathbb{1}(\Lambda_{s,t} \cap M_{s,\mu}^\varepsilon \neq \{0\}) dt + \frac{t_0(s, \mu, \varepsilon)}{T}$$

Apply Theorem 1.10 to  $\varphi, \mathbb{Z}^{d+1}, a_t$ . This is possible as the image of  $\varphi$  is not contained in any affine hyperplane of the expanding horosphere of  $a_t$ . This implies that  $\lim_{T \rightarrow \infty} (*)_T \leq m_X(\Lambda \in X : \Lambda \cap M_{s,\mu}^\varepsilon \neq \{0\})$  almost surely. Since  $\bigcap_{\varepsilon > 0} M_{s,\mu}^\varepsilon = M_{s,\mu}$  it follows that

$$m_X(\Lambda \in X : \Lambda \cap M_{s,\mu}^\varepsilon \neq \{0\}) \xrightarrow{\varepsilon \rightarrow 0} f(s, \mu).$$

□

The lower bound is slightly more complicated, as we need to guarantee that the algebraic approximations are of degree  $d$  and not less. One possible argument for the irreducibility replaces the space  $X_{d+1}$  with the congruence cover  $\mathrm{SL}_{d+1}(\mathbb{R})/(\ker(\mathrm{SL}_{d+1}(\mathbb{Z}) \rightarrow \mathrm{SL}_{d+1}(\mathbb{Z}/M\mathbb{Z})))$  where  $M = p_1 p_2 \cdots p_k$  is a product of distinct primes and obtaining that the resulting polynomials are random independent nonzero polynomials modulo  $p_i$ . Since random polynomials  $b_0 + b_1 x + \cdots + b_d x^d \in \mathfrak{F}_p[x]$  are irreducible with probability  $\sim \frac{1}{d}$ , we get that the integral polynomial is irreducible with high probability. We choose a more straightforward proof, relying on the fact that if  $s'$  is an algebraic number of degree  $d' < d$  then it is too good an approximation with this degree, and hence occurs with 0 frequency. To control the size of the minimal polynomial of  $s'$  provided that it satisfies a degree  $d$  polynomial with certain coefficients, we introduce the following lemma:

**Lemma 8.2.** *Let  $Q(x)|P(x)$  integer polynomials of degree  $d' < d$ . Then  $\mathrm{ht}(Q) \leq c(d', d) \mathrm{ht}(P)$ , where  $c(d', d)$  is a constant depending only on  $d, d'$ .*

*Proof.* For every polynomial  $P \in \mathbb{R}[x]$  denote by  $\max\mathrm{coef}(P)$  the maximal absolute value of a coefficient of  $P$ . It is sufficient to show that for every  $k, l \geq 0$  there exists a constant  $C(k, l) > 0$  such that for every  $Q_1, Q_2 \in \mathbb{R}[x]$  with  $\deg(Q_1) = k, \deg(Q_2) = l$  we have

$$\max\mathrm{coef}(Q_1) \max\mathrm{coef}(Q_2) \leq C(k, l) \max\mathrm{coef}(Q_1 Q_2).$$

This holds with

$$C(k, l) = \left( \min_{\substack{\max\text{coef}(Q_1)=1 \\ \max\text{coef}(Q_2)=1}} \max\text{coef}(Q_1 Q_2) \right)^{-1}.$$

□

*Proof of lower bound in Theorem 1.4.* Fix  $s \in \mathbb{R}$ . We say that  $t$  is *almost  $d$ -algebraic  $\mu$ -approximable* if it satisfies (1.2) for  $N = e^{t/d}, \mu$  with  $s'$  an algebraic number which is a root of a polynomial  $P(x) = b_0 + b_1x + b_2x^2 + \cdots + b_dx^d$ , which is not necessarily irreducible, with  $|b_i| < e^{t/d}$ . If  $s'$  is not of degree  $d$ , there is an integer irreducible polynomial  $Q(x)|P(x)$  of degree  $d' < d$  such that  $Q(s') = 0$ . To bound the maximal coefficient of  $Q$ , we use the Lemma 8.2. Thus, if  $t$  is almost  $d$ -algebraic  $\mu$ -approximable, with an approximation  $s'$  of degree  $d' < d$ , then  $s'$  is the root of an integral irreducible polynomial  $Q(x)$  of degree  $d'$  and coefficients less than  $c(d', d)e^{t/d}$  in absolute value. By the upper bound of Theorem 1.4, we get that for almost every  $s$ , approximations of  $s$  with  $s'$  algebraic of degree  $d' < d$  with  $\|s'\| < e^{t/d}c(d', d)$ , then for proportion 1 of  $t$ , such  $s'$  has  $s - s' = \omega((e^{t/d}c(d', d))^{-d'-1})$ . Since this approximation is too bad to make  $t$  almost  $d$ -algebraic  $\mu$ -approximable, it follows that it is enough to bound from below

$$(**)_T = \frac{1}{T} \int_0^T \mathbb{1}(t \text{ is almost } d\text{-algebraic } \mu\text{-approximable}) dt.$$

As in the proof of the upper bound,

$$(**)_T \geq \frac{1}{T} \int_0^T \mathbb{1}(\Lambda_{s,t} \cap M_{s,\mu}^{-\varepsilon} \neq \{0\}) dt \xrightarrow{T \rightarrow \infty} m_X(\Lambda \in X : \Lambda \cap M_{s,\mu}^{-\varepsilon} \neq \{0\}),$$

for every  $\varepsilon > 0$ . Now,  $m_X(\Lambda \in X : \Lambda \cap M_{s,\mu}^{-\varepsilon} \neq \{0\}) \xrightarrow{\varepsilon \rightarrow 0} m_X(\Lambda \in X : \Lambda \cap M_{s,\mu}^{\circ} \neq \{0\})$ , where  $M_{s,\mu}^{\circ}$  is the interior of  $M_{s,\mu}$ . Since measure of the boundary  $\partial M_{s,\mu}$  is zero, Siegel's formula [39] implies that  $\Lambda \cap \partial M_{s,\mu} = \{0\}$  for almost every  $\Lambda \in X$ . Hence  $m_X(\Lambda \in X : \Lambda \cap M_{s,\mu}^{\circ} \neq \{0\}) = f(s, \mu)$ . The desired claim follows. □

*Proof of Claim 8.1.* Let  $(b_i)_{i=0}^d \in \mathbb{Z}^{d+1}$ . Let

$$(8.2) \quad (\eta, (c_i)_{i=1}^d) = a_t \varphi(s) (b_i)_{i=0}^d \in \Lambda_{s,t}.$$

Let  $P(x) = \sum_{i=0}^d b_i x^i$ , and  $s'$  be a root of  $P(s)$ . In particular,  $\eta = e^t P(s)$  and  $c_i = e^{-t/d} b_i$  so that

$$\sum_{i=1}^d c_i s^i = e^{-t/d} (P(s) - b_0), \quad \sum_{i=1}^d i c_i s^{i-1} = e^{-t/d} P'(s).$$

By Taylor's Theorem (see [20, §20.3]),

$$(8.3) \quad P(s') = P(s) + (s' - s)P'(s) + \frac{1}{2}P''(s'')(s' - s)^2$$

for some  $s'' \in [s, s']$ . Assume we are in Case (i) of the claim. We get that  $P''(s'') = O(e^{t/d})$ , and hence  $P(s') = 0 = P(s) + (s' - s)P'(s) + O(e^{t/d}(s' - s)^2)$ . This implies

that

$$\begin{aligned} \left| \sum_{i=1}^d ic_i s^{i-1} \right| e^{-t} \mu = |P'(s)| e^{-t(1+1/d)} \mu > |P'(s)| |s' - s| = \left| P(s) + O(e^{-t(2+1/d)}) \right| \\ = \left| e^{-t} \eta + O(e^{-t(2+1/d)}) \right|, \end{aligned}$$

that is,  $|\sum_{i=1}^d ic_i s^{i-1}| \mu > |\eta + O(e^{-t(1+2/d)})|$ . Consequently,  $(\eta, (c_i)_{i=1}^d)$  satisfies the second condition of  $M_{s,\mu}$  up to an  $o_t(1)$  error. In addition, we get that  $\eta = O(1)$ . For the first condition, we have that  $|c_i| < 1$  because  $|b_i| < e^{t/d}$ . Now  $\sum_{i=1}^d c_i s^i = e^{-t/d}(P(s) - b_0) = e^{-t(1+1/d)} \eta - b_0 e^{-t/d}$ . Since  $|b_0| < e^{t/d}$ , we deduce that  $|\sum_{i=1}^d c_i s^i| < 1 + O(e^{-t(1+1/d)})$ , and hence  $(c_i)_{i=1}^d \in C_s$  up to an  $o_t(1)$  error.

This implies that  $(\eta, (c_i)_{i=1}^d) \in M_{s,\mu}^\varepsilon$  for all  $t$  sufficiently large.

Assume now the assumptions of Case (ii). Then  $|b_i| = e^{t/d} |c_i| < e^{t/d}$  for every  $i = 1, \dots, d$ , by Eq. (8.2). As for  $b_0 = e^{-t} \eta - e^{t/d} \sum_{i=1}^d s^i c_i$ , from the definition of  $M_{s,\mu}^{-\varepsilon}$  we get that  $|\sum_{i=1}^d s^i c_i| < 1 - \delta(s)\varepsilon$ , where  $\delta(s) = (s^2 + s^4 + \dots + s^{2d})^{-1/2}$ . Also,  $\eta = O_{s,\mu}(1)$  follows. This implies that  $|b_0| < e^{t/d}$  for all  $t$  large enough.

Now it remains to find a root  $s'$  of  $P(x)$  which is close to  $s$ . For all  $s''$  within distance 1 of  $s$  we have  $P''(s'') = O(e^{t/d})$ , and hence (8.3) implies that for all  $s' \in [s-1, s+1]$

$$(8.4) \quad P(s') = P(s) + (s' - s)P'(s) + O(e^{t/d}(s' - s)^2)$$

Note that by the definition of  $M_{s,\mu}$ , if  $\sum_{i=1}^d ic_i s^{i-1} = 0$  then also  $\eta = 0$ , and hence  $M_{s,\mu} \cap \{(\eta, (c_i)_{i=1}^d) : \sum_{i=1}^d ic_i s^{i-1} = 0\}$  has trivial interior in  $\{(\eta, (c_i)_{i=1}^d) : \sum_{i=1}^d ic_i s^{i-1} = 0\}$ . This implies that

$$\left| \sum_{i=1}^d ic_i s^{i-1} \right| \geq \delta(s)\varepsilon,$$

and hence  $|P'(s)| > \delta(s)\varepsilon e^{t/d}$ . Also, recall that  $\eta = O(1)$  so that  $P(s) = O(e^{-t})$ . Let  $s_0$  be chosen with  $s_0 - s = 2 \frac{P(s)}{P'(s)}$  so that in particular  $|s_0 - s| = O(e^{-(1+1/d)t})$ . Then by (8.4)

$$\begin{aligned} P(s_0) &= -P(s) + O\left(e^{t/d} \left(\frac{P(s)}{P'(s)}\right)^2\right) \\ &= -P(s) + O\left(P(s)e^{-(1+1/d)t}\right). \end{aligned}$$

Consequently,  $P(s_0)$  and  $P(s)$  have opposite signs (or they both vanish) and hence there is  $s'$  between  $s$  and  $s_0$  with  $P(s') = 0$ . Moreover, by (8.4), this  $s'$  satisfy that

$$\begin{aligned} -\frac{P(s)}{P'(s)} &= (s' - s) + O(e^{t/d}(s' - s)^2) \\ &= (s' - s) + O(e^{-(2+1/d)t}) \end{aligned}$$

and hence

$$s' - s = -e^{-(1+1/d)t} \frac{\eta}{\sum_{i=1}^d ic_i s^{i-1}} + O(e^{-2(1+1/d)t}).$$



By assumption,  $(\eta \pm \delta, (c_i)_{i=1}^d) \in M_{s,\mu}$  for any  $\delta < \varepsilon$  which can be seen to imply  $|s' - s| < \mu e^{-(1+1/d)t}$ .  $\square$

**8.3. Proof of Theorem 1.6.** The result follows directly from the methods of [36] and Theorem 1.10, here we merely point out the specific ingredients we use and provide an overview. Write  $X = X_{d+1} = \mathrm{SL}_{d+1}(\mathbb{R})/\mathrm{SL}_{d+1}(\mathbb{Z})$ ,  $a_t = \mathrm{diag}(e^t, \dots, e^t, e^{-dt})$  for  $t \in \mathbb{R}$  and for  $\xi \in \mathbb{R}^d$

$$u(\xi) = \begin{pmatrix} 1 & & & \xi_1 \\ & 1 & & \vdots \\ & & \ddots & \xi_d \\ & & & 1 \end{pmatrix} \in \mathrm{SL}_{d+1}(\mathbb{R}).$$

We now recall the construction of a section for the diagonal flow  $a_t$  from [36] (following mostly their notation). For  $v \in \mathbb{R}^{d+1}$  we let  $\bar{v} = (v_1, \dots, v_d)^t$ . For  $r > 0$  define a cylindrical set and its top by

$$C_r = \{v \in \mathbb{R}^{d+1} : \|\bar{v}\| \leq r, |v_{d+1}| \leq 1\},$$

$$D_r = \{v \in \mathbb{R}^{d+1} : \|\bar{v}\| \leq r, v_{d+1} = 1\}.$$

Now let  $\mathcal{S}_r \subset X$  be the set of lattices with at least one primitive vector in  $D_r$  and  $\mathcal{S}_r^\# \subset \mathcal{S}_r$  the set of lattices with exactly one primitive vector in  $D_r$ . Choose  $r_0 > 0$  sufficiently large such that  $C_{r_0}$  has Lebesgue measure at least  $2^d$ . By [36, Lemma 8.4],  $\mathcal{S}_{r_0}$  is a cross section in the sense of [36, Def. 4.2] (with respect to the Haar measure  $m_X$  on  $X$ ). By work of Ambrose and Kakutani this induces a finite measure  $\mu_{\mathcal{S}_{r_0}}$  on  $\mathcal{S}_{r_0}$  with natural properties [36, Thm. 4.4] and with  $\mu_{\mathcal{S}_{r_0}}(\mathcal{S}_{r_0} \setminus \mathcal{S}_{r_0}^\#) = 0$ . Moreover, the cross-section  $\mathcal{S}_{r_0}$  is  $m_X$ -reasonable (see [36, Def. 5.3, Thm. 8.6]) which allows us to relate notions of Birkhoff genericity (for  $a_t$  and for returns to the cross-section).

The significance of the cross section  $\mathcal{S}_{r_0}$  and its subset  $\mathcal{S}_{r_0}^\#$  can be seen as follows: let  $\xi \in \mathbb{R}^d$  and suppose that  $t > 0$  is such that  $a_t u(\xi) \mathbb{Z}^{d+1} \in \mathcal{S}_{r_0}^\#$ . Then there exists a unique  $(p, q) \in \mathbb{Z}^{d+1}$  primitive such that  $a_t u(\xi)(p, q)^t \in D_{r_0}$ , that is,  $q = e^{dt}$  and  $\|q^{\frac{1}{d}}(p - q\xi)\| \leq r_0$ . It is however not guaranteed that the so obtained vector  $(p, q)$  is a best approximation. To that end, for any  $\Lambda \in \mathcal{S}_{r_0}^\#$  write  $v(\Lambda) = (v_\Lambda, 1)$  for the unique primitive vector in  $D_{r_0} \cap \Lambda$  and set  $r(\Lambda) = \|v_\Lambda\|$ . Let  $\mathcal{B} \subset \mathcal{S}_{r_0}^\#$  be the set of lattices  $\Lambda$  with

$$\{\lambda \in \Lambda \cap C_{r(\Lambda)} : \lambda \text{ primitive}\} = \{\pm v(\Lambda)\}.$$

Now whenever  $a_t u(\xi) \mathbb{Z}^{d+1} \in \mathcal{B}$  it is easy to see that the so obtained vector  $(p, q)$  as above is a best approximation. In fact, all but finitely many best approximations arise in this manner [36, Prop. 10.4].

We also remark that the set  $\mathcal{B}$  is a well-behaved subset of  $\mathcal{S}_{r_0}$ : it is a positive measure Jordan measurable subset with respect to  $\mu_{\mathcal{S}_{r_0}}$  and it is tempered [36, Lemma 9.2, Prop. 9.8]. This implies that whenever  $x \in X$  is an  $a_t$ -generic point (with respect to  $m_X$ ) the sequence of visits  $\{a_t x : a_t x \in \mathcal{B}\}$  is equidistributed with respect to  $\nu := \frac{1}{\mu_{\mathcal{S}_{r_0}}(\mathcal{B})} \mu_{\mathcal{S}_{r_0}}|_{\mathcal{B}}$  – see [36, Thm. 5.11]. By Theorem 1.10 we may apply this to the point  $u(\phi(s)) \mathbb{Z}^{d+1}$  for almost every  $s \in [0, 1]$ .

It remains to deduce Theorem 1.6 from the above equidistribution result. The map (see [36, §8.2])

$$\Lambda \in \mathcal{S}_{r_0}^\# \mapsto (\pi^{e_{d+1}}(u(-v_\Lambda)\Lambda), v_\Lambda) \in X_d \times \overline{B_{r_0}}$$

is continuous and surjective. Moreover, it is easy to check that it associates to any  $\xi \in \mathbb{R}^d$  and  $t > 0$  with  $a_t u(\xi)\mathbb{Z}^{d+1} \in \mathcal{B}$  the projection lattice and the displacement vector of the corresponding best approximation. The measure  $\mu_{d, \|\cdot\|}$  in Theorem 1.5 is the pushforward of  $\nu$  under the above map and hence the theorem follows. Note that the properties of the measure  $\nu$  are studied in [36, §11].

## REFERENCES

- [1] Menny Aka, Emmanuel Breuillard, Lior Rosenzweig, and Nicolas de Saxcé, *Diophantine approximation on matrices and Lie groups*, *Geom. Funct. Anal.* **28** (2018), no. 1, 1–57. MR 3777412
- [2] K. Azuma, *Weighted sums of certain dependent random variables*, *Tohoku Mathematical Journal, Second Series* **19** (1967), no. 3, 357–367.
- [3] J. Bourgain, *On the maximal ergodic theorem for certain subsets of the integers*, *Israel J. Math.* **61** (1988), no. 1, 39–72. MR 937581
- [4] ———, *On the pointwise ergodic theorem on  $L^p$  for arithmetic sets*, *Israel J. Math.* **61** (1988), no. 1, 73–84. MR 937582
- [5] J. W. S. Cassels, *On a problem of Steinhaus about normal numbers*, *Colloq. Math.* **7** (1959), 95–101. MR 113863
- [6] J. Chaika and A. Eskin, *Every flat surface is Birkhoff and Oseledets generic in almost every direction*, *J. Mod. Dyn.* **9** (2015), no. 01, 1–23.
- [7] S. G. Dani, *Divergent trajectories of flows on homogeneous spaces and Diophantine approximation*, *J. Reine Angew. Math.* **359** (1985), 55–89. MR 794799
- [8] S.G. Dani and G.A. Margulis, *Limit distributions of orbits of unipotent flows and values of quadratic forms*, *Advances in Soviet Mathematics* **16** (1993), no. 1, 91–137.
- [9] H. Davenport and W. M. Schmidt, *Dirichlet’s theorem on diophantine approximation. II*, *Acta Arith.* **16** (1969/70), 413–424. MR 279040
- [10] Manfred Einsiedler, Lior Fishman, and Uri Shapira, *Diophantine approximations on fractals*, *Geom. Funct. Anal.* **21** (2011), no. 1, 14–35. MR 2773102
- [11] Frączek, Krzysztof and Shi, Ronggang and Ulcigrai, Corinna, *Genericity on curves and applications: pseudo-integrable billiards, Eaton lenses and gap distributions*, *J. Mod. Dyn.* **12** (2018), 55–122. MR 3808209
- [12] Michael Hochman and Pablo Shmerkin, *Equidistribution from fractal measures*, *Invent. Math.* **202** (2015), no. 1, 427–479. MR 3402802
- [13] Bernard Host, *Nombres normaux, entropie, translations*, *Israel J. Math.* **91** (1995), no. 1-3, 419–428. MR 1348326
- [14] D. Kleinbock and G.A. Margulis, *Flows on homogeneous spaces and Diophantine approximation on manifolds*, *Ann. Math.* **148** (1998), 339–360.
- [15] D. Y. Kleinbock and G. A. Margulis, *Bounded orbits of nonquasiunipotent flows on homogeneous spaces*, *Sinai’s Moscow Seminar on Dynamical Systems*, *Amer. Math. Soc. Transl. Ser. 2*, vol. 171, Amer. Math. Soc., Providence, RI, 1996, pp. 141–172. MR 1359098
- [16] Dmitry Kleinbock, Nicolas de Saxcé, Nimish A. Shah, and Pengyu Yang, *Equidistribution in the space of 3-lattices and dirichlet-improvable vectors on planar lines*, arXiv preprint (2021).
- [17] Dmitry Kleinbock, Elon Lindenstrauss, and Barak Weiss, *On fractal measures and Diophantine approximation*, *Selecta Math. (N.S.)* **10** (2004), no. 4, 479–523. MR 2134453
- [18] Dmitry Kleinbock, Andreas Strömbergsson, and Shucheng Yu, *A measure estimate in geometry of numbers and improvements to Dirichlet’s theorem*, *Proc. Lond. Math. Soc. (3)* **125** (2022), no. 4, 778–824. MR 4500198
- [19] Dmitry Kleinbock and Shucheng Yu, *A dynamical Borel-Cantelli lemma via improvements to Dirichlet’s theorem*, *Mosc. J. Comb. Number Theory* **9** (2020), no. 2, 101–122. MR 4096116
- [20] M. Kline, *Calculus: an intuitive and physical approach*, Courier Corporation, 1998.
- [21] J. C. Lagarias, *Best simultaneous Diophantine approximations. I. Growth rates of best approximation denominators*, *Trans. Amer. Math. Soc.* **272** (1982), no. 2, 545–554. MR 662052
- [22] ———, *Best simultaneous Diophantine approximations. II. Behavior of consecutive best approximations*, *Pacific J. Math.* **102** (1982), no. 1, 61–88. MR 682045
- [23] Grigoriy A. Margulis, *On some aspects of the theory of Anosov systems*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2004, With a survey by Richard Sharp: Periodic orbits of hyperbolic flows, Translated from the Russian by Valentina Vladimirovna Szulikowska. MR 2035655
- [24] S. Mozes and N. Shah, *On the space of ergodic invariant measures of unipotent flows*, *Ergodic Theory Dynam. Systems* **15** (1995), no. 1, 149–159.
- [25] Roland Prohaska and Cagri Sert, *Markov random walks on homogeneous spaces and Diophantine approximation on fractals*, *Trans. Amer. Math. Soc.* **373** (2020), no. 11, 8163–8196. MR 4169685

- [26] Roland Prohaska, Cagri Sert, and Ronggang Shi, *Expanding measures: Random walks and rigidity on homogeneous spaces*, (2021).
- [27] M. Ratner, *On Raghunathan's measure conjecture*, Ann. Math. **134** (1991), no. 3, 545–607.
- [28] Wolfgang M. Schmidt, *On normal numbers*, Pacific J. Math. **10** (1960), 661–672. MR 117212
- [29] Nimish Shah and Lei Yang, *Equidistribution of curves in homogeneous spaces and Dirichlet's approximation theorem for matrices*, Discrete Contin. Dyn. Syst. **40** (2020), no. 9, 5247–5287. MR 4128307
- [30] Nimish Shah and Pengyu Yang, *Equidistribution of expanding degenerate manifolds in the space of lattices*, arXiv preprint (2021).
- [31] Nimish A. Shah, *Asymptotic evolution of smooth curves under geodesic flow on hyperbolic manifolds*, Duke Math. J. **148** (2009), no. 2, 281–304. MR 2524497
- [32] ———, *Equidistribution of expanding translates of curves and Dirichlet's theorem on Diophantine approximation*, Invent. Math. **177** (2009), no. 3, 509–532. MR 2534098
- [33] ———, *Limiting distributions of curves under geodesic flow on hyperbolic manifolds*, Duke Math. J. **148** (2009), no. 2, 251–279. MR 2524496
- [34] ———, *Equidistribution of translates of curves on homogeneous spaces and Dirichlet's approximation*, Proceedings of the International Congress of Mathematicians. Volume III, Hindustan Book Agency, New Delhi, 2010, pp. 1332–1343. MR 2827843
- [35] Nimish A. Shah and Pengyu Yang, *Equidistribution of non-uniformly stretching translates of shrinking smooth curves and weighted dirichlet approximation*, arXiv preprint (2022).
- [36] Uri Shapira and Barak Weiss, *Geometric and arithmetic aspects of approximation vectors*, arXiv preprint (2022).
- [37] R. Shi, *Pointwise equidistribution for one parameter diagonalizable group action on homogeneous space*, Transactions of the American Mathematical Society **373** (2020), no. 6, 4189–4221.
- [38] Ronggang Shi and Barak Weiss, *Invariant measures for solvable groups and Diophantine approximation*, Israel J. Math. **219** (2017), no. 1, 479–505. MR 3642031
- [39] Carl Ludwig Siegel, *A mean value theorem in geometry of numbers*, Ann. of Math. (2) **46** (1945), 340–347. MR 12093
- [40] David Simmons and Barak Weiss, *Random walks on homogeneous spaces and Diophantine approximation on fractals*, Invent. Math. **216** (2019), no. 2, 337–394. MR 3953505
- [41] Pengyu Yang, *Equidistribution of expanding translates of curves and Diophantine approximation on matrices*, Invent. Math. **220** (2020), no. 3, 909–948. MR 4094972
- [42] Han Zhang, *Genericity on submanifolds and application to universal hitting time statistics*, to appear in Pure Appl. Math. Q. (2021).

EINSTEIN INSTITUTE OF MATHEMATICS, HEBREW UNIVERSITY, ISRAEL

EINSTEIN INSTITUTE OF MATHEMATICS, HEBREW UNIVERSITY, ISRAEL