

SCHOENBERG CORRESPONDENCE FOR k -(SUPER)POSITIVE MAPS ON MATRIX ALGEBRAS

B.V.RAJARAMA BHAT, PURBAYAN CHAKRABORTY, AND UWE FRANZ

ABSTRACT. We prove a Schoenberg-type correspondence for non-unital semigroups which generalizes an analogous result for unital semigroup proved by Michael Schürmann [Sch85]. It characterizes the generators of semigroups of linear maps on $M_n(\mathbb{C})$ which are k -positive, k -superpositive, or k -entanglement breaking. As a corollary we reprove Lindblad, Gorini, Kossakowski, Sudarshan's theorem[Lind76] [GKS76]. We present some concrete examples of semigroup of operators and study how their positivity properties can improve with time.

CONTENTS

1. Introduction	1
2. Preliminaries	2
2.1. Cones	2
2.2. The Algebra $\text{Lin}(M_n, M_n)$	3
2.3. The cones of k -positive, k -superpositive and k -entanglement breaking maps on M_n	4
2.4. Examples	6
2.5. Non-unital semigroups	9
2.6. Examples of k -positive semigroups	9
3. A Schoenberg correspondence for general non-unital semigroups	10
4. Application to the Semigroups of k -(super)positive, k -entanglement breaking Linear Maps on M_n	11
5. Positive Semigroups on M_2	14
6. A general 4-parameter family of semigroups	15
6.1. The depolarising channel	15
6.2. Transposition	16
References	17

1. INTRODUCTION

We characterise the generators of semigroups of linear maps acting on the matrix algebras M_n which forms various cones relevant to quantum information. In their famous papers Lindblad[Lind76], Gorini, Kosakowski, Sudarshan[GKS76] characterised the generators of semigroups of completely positive identity preserving maps. The question of semigroups of other quantum maps (e.g. k -positive or

P.C. and U.F. were supported by the ANR Project No. ANR-19-CE40-0002. Bhat is supported by the J C Bose Fellowship JBR/2021/000024 of SERB(India) .

k -superpositive etc.) arose naturally. In this paper we take this question of characterising generators of semigroups of general class of quantum maps and in particular k -(super)positive case.

Let C be a cone in \mathbb{R}^n and denote by C° the cone dual to C [cf. Def.2.1]. Schneider and Vidyasagar [SV70] in their paper showed that if ϕ is an operator on \mathbb{R}^n then the semigroup $(\Phi)_{t \geq 0} = \exp(t\phi)_{t \geq 0}$ leaves the cone C invariant if and only if $\langle \phi x | y \rangle \geq 0$ for any $x \in C$ and $y \in C^\circ$ satisfying $\langle x | y \rangle \geq 0$. This already establishes an important key result for semigroups of operators and its generators in \mathbb{R}^n . This kind of correspondence is known as ‘Schoenberg correspondence’. Michael Shürmann extends this results further to the case of unital Banach algebras, coalgebras, Hopf algebras. We can take advantage of these results to establish the correspondence between a general semigroup of k -(super)positive maps and its generator but the problem is that in all of the previous considerations the semigroup starts at the identity i.e. at time $t = 0$, $\Phi_t = \text{Id}$. As we know the identity map though completely positive, is not a k -super positive map in general. So to apply Schoenberg type correspondence for a general k -super positive semigroup we need to start at some point other than identity which should in principle lie inside the cone of k -super positive maps. The natural choice is an idempotent. We replace the identity at $t = 0$ with an idempotent and extend the Schoenberg correspondence that Schürmann proved in his paper and this constitute the key result of this paper. We can of course deduce the non-counital analogue of version for semigroups in coalgebra but that discussion is out of scope here.

In section 2, we discuss the preliminary notion of convex cone, dual cone and some of their properties without going into details. Then we shall briefly discuss the cones of k -positive maps, completely positive maps, k -superpositive maps and k -entanglement breaking maps in the space $\text{Lin}(M_n(\mathbb{C}), M_n(\mathbb{C}))$. As an example, a general four parameter map of depolarizing channel, transposition, diagonal and identity map has been studied. We end the section 2 by recalling the idea of non-unital semigroup and giving an example of a non-unital semigroup not arising as conditional expectation onto some unital $*$ -subalgebra of $M_n(\mathbb{C})$. Section 3 contains the main result of this paper i.e. the non-unital version of Schürmann’s result on Schoenberg correspondence for semigroup of operators on unital Banach algebra. In section 4, we apply this result for different cones discussed in section 2. Furthermore, we characterise the generator of semigroup of k -positive maps and finally retrieve the Linblad, Gorini, Kossakowski, Sudarsan’s theorem. We discuss a characterisation of positive semigroup on $M_2(\mathbb{C})$ in section 5. We end our paper with section 6 where we describe the time evolution of the semigroup generated by the four parameter family of operators as discussed in section 2.

2. PRELIMINARIES

2.1. Cones. We recall some definitions and basic facts on cones.

Definition 2.1. A subset C of a topological vector space V over \mathbb{R} is called a cone if for any two elements $x, y \in C$ and $\alpha \geq 0$ we have $\alpha x + y \in C$. Furthermore, we call a cone solid if it has nonempty interior and it is pointed if $C \cap (-C) = \{0\}$.

Definition 2.2. For a cone $C \subseteq V$ we define its dual cone C° ,

$$C^\circ := \{z \in V'; \langle z, x \rangle \geq 0, x \in C\}$$

where V' is the topological dual space of V .

The following results are well known so we mention without proof.[see Lemma 3.2, Corollary 3.3 [AT07]]

Proposition 2.3. *Let C be a closed convex cone in \mathbb{R}^n . Then the following statements are equivalent:*

- a. C is pointed i.e. $C \cap (-C) = \{0\}$.
- b. $C^\circ - C^\circ = \mathbb{R}^n$.
- c. C° has non-empty interior.
- d. $\text{span}(C^\circ) = \mathbb{R}^n$.

If we assume C is closed (which is the case for all cones we give as examples) the same results hold if we replace C by C° in the above proposition via the Bipolar theorem $(C^\circ)^\circ = C$ [see Theorem 5.5, [Sim11]].

In the following discussion we will often use Dirac's bra-ket notation to denote rank-one operators. To define it in short, we understand $|x\rangle$ to be a vector in some Hilbert space \mathbb{C}^n and $\langle x|$ to be its dual vector. Then for two vectors $x, y \in \mathbb{C}^n$ we define the rank one operator $|x\rangle\langle y|$ by-

$$|x\rangle\langle y|(z) := \langle y|z\rangle x$$

for any $z \in \mathbb{C}^n$. If $\{e_i\}_{i=1}^n$ is the standard basis of \mathbb{C}^n and $\{E_{ij}; 1 \leq i, j \leq n\}$ is the standard matrix units (i.e. 1 at ij -th position and zero everywhere else) then one can easily verify that $E_{ij} = |e_i\rangle\langle e_j|$.

2.2. The Algebra $\text{Lin}(M_n, M_n)$. We will consider the $*$ -algebra M_n as a Hilbert space with the Hilbert-Schmidt inner product

$$\langle A, B \rangle = \text{Tr}(A^* B), \quad A, B \in M_n,$$

and we will also consider the Hilbert-Schmidt inner product

$$\langle R, S \rangle = \text{Tr}(R^* S), \quad R, S \in \text{Lin}(M_n, M_n),$$

on $\text{Lin}(M_n, M_n)$.

In addition to the involution $*$ on $\text{Lin}(M_n, M_n)$ defined w.r.t. the inner product on M_n , where $S \in \text{Lin}(M_n, M_n)$ gets mapped to the unique S^* satisfying the condition

$$\langle S^*(A), B \rangle = \langle A, S(B) \rangle, \quad \forall A, B \in M_n,$$

we will also consider the *multiplicative* involution $\# : \text{Lin}(M_n, M_n) \rightarrow \text{Lin}(M_n, M_n)$ defined by

$$S^\#(A) = S(A^*)^*$$

for $S \in \text{Lin}(M_n, M_n)$ and $A \in M_n$. It is multiplicative in the sense that $(S_1 \circ S_2)^\# = S_1^\# \circ S_2^\#$ for $S_1, S_2 \in \text{Lin}(M_n, M_n)$.

For a pair of matrices $A, B \in M_n$ we define a linear map $T_{A,B} : M_n \rightarrow M_n$ by

$$T_{A,B}(X) = AXB, \quad \text{for } X \in M_n.$$

Denote by M_n^{op} the matrix algebra M_n equipped with the opposite multiplication, $A \cdot_{\text{op}} B = BA$ for $a, b \in M_n$.

Proposition 2.4. *The map $M_n \times M_n \ni (A, B) \mapsto T_{A,B} \in \text{Lin}(M_n, M_n)$ extends to a unique isomorphism of $*$ -algebras $T : M_n \otimes M_n^{\text{op}} \ni A \otimes B \rightarrow T_{A \otimes B} \in \text{Lin}(M_n, M_n)$, which is also an isomorphism of Hilbert spaces.*

If we define an involution $\dagger : M_n \otimes M_n^{\text{op}} \rightarrow M_n \otimes M_n^{\text{op}}$ on simple tensors $A \otimes B \in M_n \otimes M_n^{\text{op}}$ by $(A \otimes B)^\dagger = B^* \otimes A^*$ and extend conjugate-linearly, then we have

$$T(M^\dagger) = T(M)^\#, \quad M \in M_n \otimes M_n^{\text{op}}.$$

2.3. The cones of k -positive, k -superpositive and k -entanglement breaking maps on M_n . For linear maps from M_n to M_n we can define the notions of k -positivity and k -superpositivity, see [SSZ09] and the references therein.

Definition 2.5. A map $S \in \text{Lin}(M_n, M_n)$ is called *positive*, if it maps positive matrices to positive matrices. It is called *k -positive*, if $S \otimes \text{id}_{M_k} : M_n \otimes M_k \rightarrow M_n \otimes M_k$ is positive.

A linear map $S \in \text{Lin}(M_n, M_n)$ is called *completely positive*, if it is k -positive for all $k \in \mathbb{N}$. It is known that a linear map is completely positive if and only if it is n -positive and in this case it admits a Kraus representation

$$S(X) = \sum_{i=1}^p L_i^* X L_i, \quad X \in M_n,$$

with certain matrices $L_1, \dots, L_p \in M_n$.

Definition 2.6. A map $S \in \text{Lin}(M_n, M_n)$ is called *k -superpositive*, if it admits a Kraus representation $S(X) = \sum_{i=1}^p L_i^* X L_i$ where all the matrices L_1, \dots, L_p have rank less than or equal to k .

Definition 2.7. A map $S \in \text{Lin}(M_n, M_n)$ is called *entanglement breaking* if for any $X \in (M_k \otimes M_n)^+$ and $k \geq 1$, $(\text{id}_k \otimes S)X$ is separable. We denote the set of all entanglement breaking maps on M_n by \mathcal{EB} .

It can be shown that the cone of 1-super positive maps coincides with the set of entanglement breaking maps.

Proposition 2.8. $\mathcal{S}_1 = \mathcal{EB}$.

Proof. [HSR03]. □

In the terminology of [CMW19] we can define k -entanglement breaking maps-

Definition 2.9. A linear map $S \in \text{Lin}(M_n, M_n)$ is called *k -entanglement breaking* if it is k -positive and $(\text{Id}_k \otimes S)X$ is separable whenever $X \in (M_k \otimes M_n)^+$. We denote the cone of k -entanglement breaking maps by \mathcal{EB}_k .

We define the following cones in $\text{Lin}(M_n, M_n)$,

$$\begin{aligned} \mathcal{P}_k &= \{S \in \text{Lin}(M_n, M_n); S \text{ is } k\text{-positive}\}, \\ \mathcal{CP} &= \{S \in \text{Lin}(M_n, M_n); S \text{ is completely positive}\} \\ \mathcal{S}_k &= \{S \in \text{Lin}(M_n, M_n); S \text{ is } k\text{-superpositive}\}, \\ \mathcal{EB}_k &= \{S \in \text{Lin}(M_n, M_n); S \text{ is } k\text{-entanglement breaking}\}. \end{aligned}$$

With the cones defined above we have the following chain of inclusions

- (i) $\mathcal{EB} = \mathcal{S}_1 \subset \mathcal{S}_k \subset \mathcal{S}_n = \mathcal{CP} = \mathcal{P}_n \subset \mathcal{P}_k \subset \mathcal{P}_1$
- (ii) $\mathcal{EB} = \mathcal{EB}_n \subset \mathcal{EB}_k \subset \mathcal{P}_k \subset \mathcal{P}_1$
- (iii) $\mathcal{EB} = \mathcal{S}_1 \subset \mathcal{S}_k \subset \mathcal{EB}_k^\circ \subset \mathcal{P}_1$

It is easy to see that \mathcal{P}_k is a closed cone and the cone \mathcal{EB}_k is also closed (see Theorem 3.12,[DMS21]). Furthermore we have the duality relations that $\mathcal{P}_k^\circ = \mathcal{S}_k$ for $k = 1, \dots, n$. In summary,

$$\mathcal{P}_k^\circ = \mathcal{S}_k, \quad \mathcal{S}_k^\circ = \mathcal{P}_k \quad \text{and} \quad (\mathcal{EB}_k^\circ)^\circ = \mathcal{EB}_k$$

for $k=1, \dots, n$.

Proposition 2.10. *The cones $\mathcal{P}_k, \mathcal{S}_k, \mathcal{EB}_k, \mathcal{EB}_k^\circ \subseteq \text{Lin}(M_n, M_n)^{\text{her}}$, $k = 1, \dots, n$, are closed, pointed and solid. Furthermore they are stable under composition.*

Stability under composition is easy to check for k -positivite maps and therefore for CP maps. For k -superpositive maps, if $S, T \in \mathcal{S}_k$ have Kraus representations

$$S(X) = \sum_{i=1}^q L_i^* X L_i \quad \text{and} \quad T(X) = \sum_{j=1}^q K_j^* X K_j,$$

with Kraus operators of rank less than or equal to k , then so does their composition $S \circ T$,

$$S \circ T(X) = \sum_{i=1}^p \sum_{j=1}^q S_i K_j X (S_i K_j)^*,$$

since $\text{rank}(L_i K_j) \leq \min \text{rank}(L_i), \text{rank}(K_j) \leq k$. For the stability of composition of k -entanglement breaking maps see [DMS21].

To show that they are solid (i.e., $C + (-C) = \text{Lin}(M_n, M_n)$), or, equivalently, they have non-empty interior) and pointed (i.e., $C \cap (-C) = \{0\}$), we will show below in Lemma 2.11 that the depolarising channel $P : M_n \rightarrow M_n$, $P(X) = \text{Tr}(X) \frac{1}{n} I_n$ is in the interior of $\mathcal{S}_1 = \mathcal{EB}$. Since all other cones contain \mathcal{EB} , they are also solid. Which implies that they are pointed, as duals of solid cones.

Lemma 2.11. *The depolarising channel $P : M_n \rightarrow M_n$, $P(X) = \text{Tr}(X) \frac{1}{n} I_n$ belongs to the interior of the cone $\mathcal{S}_1 = \mathcal{EB}$ of entanglement breaching maps.*

Proof. We will use the duality between $\mathcal{EB} = \mathcal{S}_1$ and $\mathcal{PM} = \mathcal{P}_1$. To show that P belongs to the interior of \mathcal{EB} , it is sufficient to show that

$$\forall T \in \mathcal{PM} = \mathcal{EB}^\circ, T \neq 0 \quad \Rightarrow \quad \langle T, P \rangle > 0.$$

Let $T \in \mathcal{PM}$ such that $\langle T, P \rangle = 0$. Choose an orthonormal bases $(u_i)_{i=1, \dots, n}$ of \mathbb{C}^n . Then the set of rank 1 operators $(|u_i\rangle\langle u_j|)_{i,j=1, \dots, n}$ is an orthonormal basis of M_n and we have

$$0 = \langle T, P \rangle = \frac{1}{n^2} \sum_{i,j=1}^m \left\langle T(|u_i\rangle\langle u_j|), \underbrace{P(|u_i\rangle\langle u_j|)}_{= \frac{\delta_{ij}}{n} I_n} \right\rangle = \frac{1}{n^3} \sum_{i=1}^n \text{Tr} \left(T(|u_i\rangle\langle u_i|)^* \right)$$

Since T is positive map, each term in this sum is positive, and therefore has to vanish. Since the orthonormal basis $(u_i)_{i=1, \dots, n}$ is arbitrary, this implies

$$\forall u \in \mathbb{C}^n, \quad \text{Tr} \left(T(|u\rangle\langle u|) \right) = 0.$$

and therefore $T = 0$. □

2.4. Examples. We take the linear combinations of some well known maps e.g. depolarizing channel, transpose, or map which is conditional expectation onto diagonal and discuss the criterion of positivity, k -positivity or complete positivity of such combinations. Later we can use these examples to generate identity preserving semigroups, which are of our interest. We have already defined *depolarising channel* in lemma 2.11. as a linear map $P : M_n \rightarrow M_n$ such that,

$$P(X) = \frac{1}{n} \text{Tr}(X) I_n$$

We have

$$P(X) = \frac{1}{n} \sum_{j,k=1}^n \langle e_j, X e_j \rangle |e_k\rangle\langle e_k| = \sum_{j,k=1}^n \frac{1}{\sqrt{n}} |e_k\rangle\langle e_j| X \left(\frac{1}{\sqrt{n}} |e_k\rangle\langle e_j| \right)^*$$

which shows that $P \in \mathcal{S}_1 = \mathcal{EB}$. The Choi-Jamiołkowski matrix of P is

$$C_P = \sum_{j,k=1}^n E_{jk} \otimes P(E_{jk}) = \frac{1}{n} \sum_{j,k=1}^n E_{jj} \otimes E_{kk} = \frac{1}{n} I_n \otimes I_n.$$

We consider also the *transposition map*, $T(X) = X^T$. It is known that T is positive, but not 2-positive, i.e.

$$T \in \mathcal{P}_1, \quad T \notin \mathcal{P}_2,$$

for $n \geq 2$.

The Choi-Jamiołkowski matrix of T is

$$C_T = \sum_{j,k=1}^n E_{jk} \otimes T(E_{jk}) = \sum_{j,k=1}^n E_{jk} \otimes E_{kj}.$$

Conditional expectation onto the the diagonal: Consider the linear map $D : M_n \rightarrow M_n$, $D(X) = (\delta_{jk} x_{jk})_{1 \leq j,k \leq n}$ for $X = (x_{jk})_{1 \leq j,k \leq n} \in M_n$. This map is the conditional expectation onto the *-subalgebra of diagonal matrices (w.r.t. the standard basis).

We have

$$D(X) = \sum_{j=1}^n |e_j\rangle\langle e_j| X |e_j\rangle\langle e_j|,$$

which shows that $D \in \mathcal{S}_1 = \mathcal{EB}$.

We can furthermore show that D belongs to the boundary of \mathcal{CP} , and therefore also to the boundary of \mathcal{EB} . Indeed, denote by $C \in M_n$ that matrix that cyclically permutes the vectors of the standard basis,

$$C e_j = e_{j \oplus 1} = \begin{cases} e_{j+1} & \text{if } 1 \leq j \leq n-1, \\ e_1 & \text{if } j = n, \end{cases}$$

where we use \oplus to denote the addition modulo n in $\{1, \dots, n\}$. Then the completely positive map T_C with $T_C(X) = C X C^*$ acts as

$$T_C(E_{jk}) = E_{j \oplus 1, k \oplus 1}.$$

Thus we have

$$\langle T_C, D \rangle = \sum_{j,k=1}^n \text{Tr}(T_C(E_{jk})^* D(E_{jk})) = \sum_{j=1}^n \text{Tr}(E_{j \oplus 1, j \oplus 1} E_{jj}) = 0.$$

The Choi-Jamiołkowski matrix of D is

$$C_D = \sum_{j,k=1}^n E_{jk} \otimes D(E_{jk}) = \sum_{j=1}^n E_{jj} \otimes E_{jj}.$$

We have

$$C_{Id} = \sum_{j,k=1}^n E_{jk} \otimes E_{jk}.$$

We are interested in the 4 parameter family of linear map- $\alpha P + \beta D + \gamma T + \delta \text{Id}$. These are exactly the linear maps on $M_n(\mathbb{C})$ that are invariant under the action of the hyperoctahedral group as signed permutations and have been considered from that perspective in [JPPY23]. We wish to decide to which cone it belongs depending on the real values of the coefficients. We can easily check that the Choi-Jamiołkowski matrices $C_P, C_D, C_T, C_{Id} \in M_n \otimes M_n$ commute. Therefore, we can simultaneously diagonalise these four matrices. The minimal polynomials of C_D, C_T, C_{Id} have degree two, and these matrices have two distinct eigenvalues. Computing the traces, we also get the multiplicities.

Eigenvalues of C_D	$\rho = 0$	$\rho = 1$
$\dim \ker(C_D - \rho I_n \otimes I_n)$	$n^2 - n$	n
Eigenvalues of C_T	$\sigma = -1$	$\sigma = 1$
$\dim \ker(C_T - \sigma I_n \otimes I_n)$	$\frac{1}{2}(n^2 - n)$	$\frac{1}{2}(n^2 + n)$
Eigenvalues of C_{Id}	$\tau = 0$	$\tau = n$
$\dim \ker(C_{Id} - \tau I_n \otimes I_n)$	$n^2 - 1$	1

In particular, C_{Id} is a multiple of the orthogonal projection onto $\Omega = \sum_{j=1}^n e_j \otimes e_j$, which is also an eigenvector for the other matrices.

Denote by

$$V(\rho, \sigma, \tau) = \{v \in \mathbb{C}^n \otimes \mathbb{C}^n : C_D v = \rho v, C_T v = \sigma v, C_{Id} v = \tau v\}$$

the joint eigenspaces of C_D, C_T, C_{Id} . We can verify that, $V(\rho, \sigma, \tau)$ is non-null corresponding to the four triples $(\rho, \sigma, \tau) = (0, -1, 0), (0, 1, 0), (1, 1, 0), (1, 1, n)$.

Proposition 2.12. *Let $\alpha, \beta, \gamma, \delta \in \mathbb{R}$ and set*

$$\Phi(\alpha, \beta, \gamma, \delta) = \alpha P + \beta D + \gamma T + \delta \text{Id}.$$

Then Φ is completely positive if and only if $\alpha, \beta, \gamma, \delta$ satisfy the inequality

$$\alpha \geq \max\{n|\gamma|, -n(\beta + \gamma), -n(\beta + \gamma) - n^2\delta\}.$$

Proof. We check that the Choi-Jamiołkowski matrix $C_{\Phi(\alpha, \beta, \gamma, \delta)} = \alpha C_P + \beta C_D + \gamma C_T + \delta C_{Id}$ of $C_{\Phi(\alpha, \beta, \gamma, \delta)}$ has eigenvalues

$$\frac{\alpha}{n} - \gamma, \quad \frac{\alpha}{n} + \gamma, \quad \frac{\alpha}{n} + \beta + \gamma, \quad \frac{\alpha}{n} + \beta + \gamma + n\delta,$$

corresponding to the four triples $(\rho, \sigma, \tau) = (0, -1, 0), (0, 1, 0), (1, 1, 0), (1, 1, n)$ with non-trivial eigenspaces. \square

Restricting to equivariant maps as in the terminology of [BCS20] (see also [COS18, Theorem 2.2]) we can also characterize the k -positivity criterion of the above mentioned family.

2.4.1. *(1,0)-unitarily equivariant case - linear combinations of identity and depolarising channel.* The identity map and the depolarising channel are $(1,0)$ -unitarily equivariant (in the terminology of [BCS20, Definition 1.1 (iii)]). For $\Phi_{\alpha,\delta} = \alpha P + \delta \text{Id}$, $\Phi_{\alpha,\delta}(X) = \delta X + \frac{\alpha}{n} \text{Tr}(X) I_n$ we

$$\Phi_{\alpha,\delta}(U X U^*) = U \Phi_{\alpha,\delta}(X) U^*,$$

for all $U, X \in M_n$ with U unitary.

Let $1 \leq k \leq n$. By [BCS20, Theorem 2.4], $\Phi_{\alpha,\delta}$ is k -positive, if and only if

$$C_{\Phi_{\alpha,\delta}}^{(k)} = \sum_{i,j=1}^k E_{ij} \otimes \Phi_{\alpha,\delta}(E_{ij}) = \sum_{i,j=1}^k E_{ij} \otimes \left(\frac{\alpha}{n} I_n^{(k)} \otimes I_n + \delta \sum_{i,j=1}^k E_{ij} \otimes E_{ij} \right) \in M_n \otimes M_n$$

is positive, where $I_n^{(k)} = \sum_{i=1}^k E_{ii} \in M_n$.

Note that $C_{\Phi_{0,1}}^{(k)} = \sum_{i,j=1}^k E_{ij} \otimes E_{ij}$ commutes with $I_n^{(k)} \otimes I_n$, satisfies $\left(C_{\Phi_{0,1}}^{(k)} \right)^2 = k C_{\Phi_{0,1}}^{(k)}$, and is a multiple of the orthogonal projection onto $\Omega_k = \sum_{i=1}^k e_i \otimes e_i$. One can show that the eigenvalues of $C_{\Phi_{\alpha,\delta}}^{(k)} = \frac{\alpha}{n} I_n^{(k)} \otimes I_n + \delta C_{\Phi_{0,1}}^{(k)}$ are given by

$$\text{spec}(C_{\Phi_{0,1}}^{(k)}) = \left\{ 0, \frac{\alpha}{n}, \frac{\alpha}{n} + k\delta \right\}.$$

We summarize our results in the following Lemma.

Lemma 2.13. ([Tom85, Theorem 2]) *Let $\alpha, \delta \in \mathbb{R}$. The linear map $\Phi_{\alpha,\delta}$ is k -positive iff the matrix $C_{\Phi_{\alpha,\delta}}^{(k)}$ is positive iff α, δ satisfy the following two inequalities*

$$\alpha \geq 0 \quad \text{and} \quad \delta \geq -\frac{\alpha}{kn}.$$

2.4.2. *(0,1)-unitarily equivariant case - linear combinations of transposition and depolarising channel.* The transposition and the depolarising channel are $(0,1)$ -unitarily equivariant (in the terminology of [BCS20, Definition 1.1 (iii)]). For $\Psi_{\alpha,\gamma} = \alpha P + \gamma T$, $\Psi_{\alpha,\gamma}(X) = \gamma X^T + \frac{\alpha}{n} \text{Tr}(X) I_n$, we have

$$\Psi_{\alpha,\gamma}(U X U^*) = \bar{U} \Psi_{\alpha,\gamma}(X) U^T,$$

for all $U, X \in M_n$ with U unitary.

By [BCS20, Theorem 2.4], $\Psi_{\alpha,\gamma}$ is k -positive, if and only if

$$C_{\Psi_{\alpha,\gamma}}^{(k)} = \sum_{i,j=1}^k E_{ij} \otimes \Psi_{\alpha,\gamma}(E_{ij}) = \frac{\alpha}{n} I_n^{(k)} \otimes I_n + \gamma \sum_{i,j=1}^k E_{ij} \otimes E_{ji} \in M_n \otimes M_n$$

is positive, where $I_n^{(k)} = \sum_{i=1}^k E_{ii} \in M_n$.

Note that $C_{\Psi_{0,1}}^{(k)} = \sum_{i,j=1}^k E_{ij} \otimes E_{ji}$ satisfies

$$I_n^{(k)} \otimes I_n C_{\Psi_{0,1}}^{(k)} = C_{\Psi_{0,1}}^{(k)} = C_{\Psi_{0,1}}^{(k)} I_n^{(k)} \otimes I_n,$$

and $(C_{\Psi_{0,1}}^{(k)})^2 = \sum_{i,j=1}^k E_{ii} \otimes E_{jj} = I_n^{(k)} \otimes I_n^{(k)}$. which implies

$$\text{spec}(C_{\Psi_{0,1}}^{(1)}) = \{0, 1\} \quad \text{and} \quad \text{spec}(C_{\Psi_{0,1}}^{(k)}) = \{-1, 0, 1\} \text{ for } k \geq 2.$$

We see that the eigenvalues of $C_{\Psi_{\alpha,\gamma}}^{(k)} = \frac{\alpha}{n} I_n^{(k)} \otimes I_n + \gamma C_{\Psi_{0,1}}^{(k)}$ are

$$\text{spec}(C_{\Psi_{\alpha,\gamma}}^{(1)}) = \left\{ 0, \frac{\alpha}{n}, \frac{\alpha}{n} + \gamma \right\} \quad \text{and} \quad \text{spec}(C_{\Psi_{\alpha,\gamma}}^{(k)}) = \left\{ 0, \frac{\alpha}{n}, \frac{\alpha}{n} + \gamma, \frac{\alpha}{n} - \gamma \right\}$$

for $k \geq 2$.

Therefore we have the following Lemma.

Lemma 2.14. ([Tom85, Theorem 3]) *Let $\alpha, \gamma \in \mathbb{R}$. The linear map $\Psi_{\alpha, \gamma}$ is*

- i. 1-positive iff $\alpha \geq 0$ and $\alpha \geq -n\gamma$.*
- ii. k -positive for $k \geq 2$, iff $\alpha \geq n|\gamma|$.*

2.5. Non-unital semigroups. We will be interested in semigroups of linear operators $(T_t)_{t \geq 0}$ (on some Banach space V), which do not start from the identity, i.e. we have $T_s T_t = T_{s+t}$ for all $s, t \geq 0$, but not necessarily $T_0 = \text{id}$. We still want $t \mapsto T_t$ to be continuous.

The semigroup property implies

$$T_0^2 = T_0$$

i.e., T_0 is idempotent. Then we can decompose V as $V = \text{Im}(T_0) \oplus \text{Ker}(T_0)$, where $\text{Im}(T_0) = T_0(V) = \text{Ker}(\text{id}_V - T_0)$, and $\text{Ker}(T_0) = \text{Im}(\text{id}_V - T_0) = (\text{id}_V - T_0)(V)$. We assume that T_0 is bounded, so both subspaces are closed. With respect to this decomposition T_0 has the form

$$T_0 = \begin{pmatrix} \text{id}_{\text{Im}(T_0)} & 0 \\ 0 & 0 \end{pmatrix}$$

Furthermore, the semigroup property implies $T_0 T_t = T_t T_0 = T_t$ for all $t \geq 0$. Therefore

$$\text{Ker}(T_0) \subseteq \text{Ker}(T_t) \quad \text{and} \quad \text{Im}(T_t) \subseteq \text{Im}(T_0).$$

W.r.t. the decomposition $V = \text{Im}(T_0) \oplus \text{Ker}(T_0)$ we can write the T_t as

$$T_t = \begin{pmatrix} \tilde{T}_t & 0 \\ 0 & 0 \end{pmatrix}$$

with some linear operators $\tilde{T}_t \in B(\text{Im}(T_0))$, which form a continuous semigroup $(\tilde{T}_t)_{t \geq 0}$ with initial value $\tilde{T}_0 = \text{Id}_{\text{Im}(T_0)}$. This allows to extend classical result on unital semigroups to the non-unital case.

In our examples, if T_0 is a conditional expectation onto some unital $*$ -subalgebra, then we are lead to study semigroups $(\tilde{T}_t)_{t \geq 0}$ that preserve the corresponding cones of $T_0(M_n) \subseteq M_n$.

But in general T_0 need not be a conditional expectation, as the example in the following subsection shows.

2.6. Examples of k -positive semigroups. Since the identity map $\text{id} : M_n \rightarrow M_n$ is not k -superpositive for $k < n$, there exist no k -superpositive semigroups $(T_t : M_n \rightarrow M_n)_{t \geq 0}$ with $T_0 = \text{id}$. But there do exist semigroups of k -superpositive linear maps on M_n that start with an idempotent k -superpositive map T_0 . Very simple examples are given by $T_0(X) = PXP$ with P a k -dimensional orthogonal projection.

Another class of examples are semigroups $T_t : M_2 \otimes M_n \rightarrow M_2 \otimes M_n$ of the form

$$T_t \left(\begin{pmatrix} A & B \\ C & D \end{pmatrix} \right) = \begin{pmatrix} S_t(A) & 0 \\ 0 & \alpha(S_t(A)) \end{pmatrix}$$

with $(S_t)_{t \geq 0}$ a k -superpositive semigroup acting on M_n and $\alpha : M_n \rightarrow M_n$ any k -superpositive linear map. If α is not a $*$ -homomorphism, then T_0 is a k -superpositive idempotent whose image is not a $*$ -subalgebra, and so T_0 is not a conditional expectation.

3. A SCHOENBERG CORRESPONDENCE FOR GENERAL NON-UNITAL SEMIGROUPS

We give a non-unital version of [Sch85, Lemma 2.1].

For X a Banach space, we will denote by $B_{X'}$ and $S_{X'}$ the unit ball and the unit sphere of the dual space X' . If $C \subseteq X$ is a cone in X then C° denotes the dual cone in X' , i.e.

$$C^\circ = \{\varphi \in X'; \forall v \in C, \varphi(v) \geq 0\}.$$

Note that we have (for C a closed convex cone)

$$(C^\circ)^\circ = C.$$

Theorem 3.1. *Let A be a real Banach algebra with a closed convex cone $C \subseteq A$ with non-empty interior. Let $a_0 \in C$ be an idempotent such that for any $c \in C$, we have $a_0 c a_0 \in C$.*

We assume furthermore that for any $c \in C$ we have $c^n \in C$ for $n \geq 1$.

Then, for any $b \in A$ such that $ba_0 = a_0 b = b$, the following statements are equivalent.

- (i): *b is a_0 -conditionally positive on C° , i.e., $\varphi(b) \geq 0$ for all $\varphi \in C^\circ$ with $\varphi(a_0) = 0$.*
- (ii): *$\exp_{a_0}(tb) := \lim_{n \rightarrow \infty} (a_0 + \frac{tb}{n})^n \in C$ for all $t \geq 0$.*

Proof. (ii) \Rightarrow (i): This follows by differentiation at $t = 0$. If $\varphi \in C^\circ$ is such that $\varphi(a_0) = 0$, then

$$\varphi(b) = \lim_{\substack{t \rightarrow 0 \\ t > 0}} \varphi \left(\frac{\exp_{a_0}(tb) - \exp_{a_0}(0)}{t} \right) = \lim_{\substack{t \rightarrow 0 \\ t > 0}} \frac{1}{t} \varphi(\exp_{a_0}(tb)) \geq 0,$$

(i) \Rightarrow (ii): We have

$$(3.1) \quad \exp_{a_0}(tb) := \lim_{n \rightarrow \infty} \left(a_0 + \frac{tb}{n} \right)^n = a_0 + \sum_{n=1}^{\infty} \frac{(tb)^n}{n!}$$

and we want to show that this quantity is positive for $t \geq 0$ if b satisfies condition (i). Without loss of generality we can take $t = 1$.

We will prove in four steps that $\exp_{a_0}(b)$ is positive.

Step I: For any interior point $c \in C$ there exists $\delta > 0$ such that

$$\forall \varphi \in C^\circ, \quad \varphi(c) \geq \delta \|\varphi\|.$$

Indeed, let $c \in C$ be an interior point. Then there exists $\delta > 0$ such that $c + \delta B_A \subseteq C$, where B_A is the unit ball in A . Therefore for $v \in B_A$,

$$\varphi(c \pm \delta v) \geq 0,$$

and

$$\varphi(c) \geq \delta \sup_{v \in B_A} |\varphi(v)| = \delta \|\varphi\|.$$

Step II: for any $\rho > 0$ there exists $\eta > 0$ such that

$$\forall \varphi \in C^\circ \cap B_{A'}, \quad \varphi(a_0) < \eta \Rightarrow \varphi(b) > -\rho.$$

Indeed, fix $\rho > 0$ and set

$$V_n(\rho) = \left\{ \varphi \in C^\circ \cap B_{A'}; \varphi(a_0) \leq \frac{1}{n} \text{ and } \varphi(b) \leq -\rho \right\}$$

By the a_0 -conditional positivity of b w.r.t. C° , we have

$$\bigcap_{n \geq 1} V_n(\rho) = \emptyset.$$

Since $C^\circ \cap B_{A'}$ is compact by the Banach-Alaoglu theorem in the weak-* topology, and since the $V_n(\rho) \subseteq C^\circ \cap B_{A'}$ are weak-* closed, there exists $n_0 \in \mathbb{N}$ such that $V_{n_0}(\rho) = \emptyset$. Take $\eta = \frac{1}{n_0}$.

Step III: For any $\varepsilon > 0$ and $c \in C$ an interior point, there exists $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$,

$$\forall \varphi \in C^\circ \cap S_{A'}, \quad \varphi \left(a_0 + \frac{b + \varepsilon c}{n} \right) \geq 0,$$

where $S_{A'}$ denotes the unit sphere in A' . Let $\delta > 0$ be the real number guaranteed by Step I such that $\varphi(c) \geq \delta \|\varphi\|$ for all $\varphi \in C^\circ$. Let $\eta > 0$ be the real number guaranteed by Step II such that for all $\varphi \in C^\circ \cap B_{A'}$ with $\varphi(a_0) < \eta$ we have $\varphi(b) \geq -\varepsilon\delta$.

Let $\varphi \in C^\circ \cap S_{A'}$. We distinguish two cases, according to the value of φ on a_0 .

Case $\varphi(a_0) < \eta$: in this case we have

$$\varphi \left(a_0 + \frac{b + \varepsilon c}{n} \right) = \underbrace{\varphi(a_0)}_{\geq 0} + \frac{1}{n} \left(\underbrace{\varphi(b)}_{\geq -\varepsilon\delta} + \varepsilon \underbrace{\varphi(c)}_{\geq \delta} \right) \geq 0.$$

Case $\varphi(a_0) \geq \eta$: now we get

$$\varphi \left(a_0 + \frac{b + \varepsilon c}{n} \right) = \underbrace{\varphi(a_0)}_{\geq \eta} + \frac{1}{n} \varphi(b + \varepsilon c) \geq \eta - \frac{\|b + \varepsilon c\|}{n},$$

which is positive as soon as $n \geq \frac{\|b + \varepsilon c\|}{\eta}$.

Step IV: By the Bipolar theorem, this means that for any $\varepsilon > 0$ and $c \in C$ an interior point there exists an $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$, $a_0 + \frac{b + \varepsilon c}{n} \in C$. Then we have $a_0 \left(a_0 + \frac{b + \varepsilon c}{n} \right) a_0 = a_0 + \frac{b + \varepsilon a_0 c a_0}{n} \in C$.

Since C is stable under taking powers and closed, we get

$$\exp_{a_0}(b + \varepsilon a_0 c a_0) = \lim_{n \rightarrow \infty} \left(a_0 + \frac{b + \varepsilon a_0 c a_0}{n} \right)^n \in C.$$

To conclude the proof we let $\varepsilon \searrow 0$. □

4. APPLICATION TO THE SEMIGROUPS OF k -(SUPER)POSITIVE, k -ENTANGLEMENT BREAKING LINEAR MAPS ON M_n

Now we apply Theorem 3.1 to the algebra $L(M_n, M_n)^{\text{her}}$ of hermitianity preserving linear maps from M_n to M_n .

Theorem 4.1. *Let*

$$A = L(M_n, M_n)^{\text{her}} = \{T \in L(M_n, M_n); T \circ * = * \circ T\}$$

and let $C \subseteq A$ be one of the cones $\mathcal{P}_1 = \mathcal{PM}, \mathcal{P}_2, \dots, \mathcal{P}_{n-1}, \mathcal{P}_n = \mathcal{CP} = \mathcal{S}_n, \mathcal{S}_{n-1}, \dots, \mathcal{S}_2, \mathcal{S}_1 = \mathcal{EB}, \mathcal{EB}_{n-1}, \dots, \mathcal{EB}_1$.

Fix an idempotent map $\Phi_0 \in C$. Then for $\Psi \in A$ with $\Psi \circ \Phi_0 = \Phi_0 \circ \Psi = \Psi$ the following are equivalent.

(i) We have $\exp_{\Phi_0}(t\Psi) = \Phi_0 + \sum_{n=1}^{\infty} \frac{t^n \Psi \circ \Phi_0^n}{n!} \in C$ for all $t \geq 0$;

(ii) Ψ is Φ_0 -conditionally positive on C° , i.e., we have

$$\forall v \in C^\circ, \quad \langle v, \Phi_0 \rangle = 0 \quad \Rightarrow \quad \langle v, \Psi \rangle \geq 0.$$

Proof. Proposition 2.10 ensures that all the cones are convex, solid, pointed and stable under composition. Thus the result follows from Theorem 3.1. \square

Corollary 4.2. *Take $C = \mathcal{P}_k$ with $k \in \{1, \dots, n\}$ and $\Phi_0 = \text{Id} = T_{I \otimes I}$. Then we know that*

$$C^\circ = \mathcal{S}_k = \text{convex hull of } \{T_{V \otimes V^*}; V \in M_n, \text{rank}(V) \leq k\}.$$

Since

$$\langle T_{V \otimes V^*}, \text{Id} \rangle = \text{Tr}(V^* \otimes V) = |\text{Tr}(V)|^2,$$

we see that $\Psi = T_S$ with $S = \sum A_i \otimes B_i \in (M_n \otimes M_n^{op})^{\text{sa}}$ generates a semigroup $T_t = \exp(t\Psi)$, $t \geq 0$, of k -positive maps if and only if

$$\forall V \in M_n, \quad (\text{rank}(V) \leq k \text{ and } \text{Tr}(V) = 0) \quad \Rightarrow \quad \sum \text{Tr}(A_i V^*) \text{Tr}(B_i V) \geq 0.$$

Let $\{B_j\}_{j=1, \dots, n^2}$ be an orthonormal basis of M_n such that $B_1 = I_n$ and $\text{Tr}(B_j) = 0$ for $2 \leq j \leq n^2$. Taking $\{B_i \otimes B_j\}_{i,j=1}^{n^2}$ as a basis of $M_n \otimes M_n^{op}$, or equivalently $\{B_i \otimes B_j^*\}_{i,j=1}^{n^2}$ of $M_n \otimes M_n^*$ further we write $S = \sum_{i,j=1}^{n^2} D_{ij} B_i \otimes B_j^*$, where D_{ij} is a $n^2 \times n^2$ matrix in \mathbb{C} , so that the map T_S is given by

$$(4.1) \quad \Psi(X) = T_S(X) = \sum_{i,j=1}^{n^2} D_{ij} B_i X B_j^*$$

for all $X \in M_n$. The semigroup $\exp(t\Psi)$ is Hermiticity preserving if and only if the generator T_S is Hermiticity perserving if and only if the matrix (D_{ij}) is Hermitian.

Proposition 4.3. *A semigroup $T_t = \exp t\Psi$ is completely positive if and only if the matrix $(D_{ij})_{i,j=1}^{n^2}$ is Hermitian and for any $v = (0, v_2, v_3, \dots, v_{n^2}) \in \mathbb{C}^{n^2}$ we have*

$$\langle v | (D_{ij})_{i,j=1}^{n^2} v \rangle \geq 0,$$

Proof. Let V be a $n \times n$ matrix with the basis decomposition $V = \sum_{j=1}^{n^2} v_j B_j$. The trace conditions on B_j 's implies that $\text{Tr}(V) = 0$ if and only if $v_1 = 0$. By a direct application of Corollary 4.2 we see that the map T_S with $S = \sum_{i,j=1}^{n^2} B_i \otimes B_j^*$ generates a completely positive semigroup (i.e. n positive) if and only if for any $V \in M_n$ with $\text{Tr}(V) = 0$, we have $\sum_{i,j=1}^{n^2} D_{ij} \text{Tr}(B_i V^*) \text{Tr}(B_j^* V) \geq 0$. Writing the basis decomposition of V in this condition, we obtain

$$(\forall (0, v_2, v_3, \dots, v_{n^2}) \in \mathbb{C}^{n^2}) \quad \Rightarrow \quad \sum_{i,j=2}^{n^2} \sum_{k,l=2}^{n^2} D_{ij} \bar{v}_k v_l \text{Tr}(B_i B_k^*) \text{Tr}(B_j^* B_l) \geq 0.$$

Because of the orthonormality of the basis B_i 's, $\text{Tr}(B_i B_k^*) = \delta_{ik}$ and $\text{Tr}(B_j^* B_l) = \delta_{jl}$. Thus the above condition becomes

$$\sum_{i,j=2}^{n^2} \sum_{k,l=2}^{n^2} D_{ij} \bar{v}_k v_l \delta_{ik} \delta_{jl} = \sum_{i,j=2}^{n^2} D_{ij} \bar{v}_i v_j \geq 0,$$

which is the desired result. \square

From this result we can re-derive the Lindblad[Lind76], Gorini, Kossakowski, Sudarshan's theorem [GKS76] on generator of CP semigroup.

Theorem 4.4. *An identity preserving semigroup $\exp(t\Psi)$ will be completely positive for all time $t \geq 0$ if and only if the generator Ψ has the form*

$$\Psi(X) = i[H, X] + \sum_{j=1}^k \{V_j^* X V_j - \frac{1}{2}(V_j^* V_j X - X V_j^* V_j)\}$$

for all $X \in M_n$, where H is an Hermitian matrix and $V \in M_n$.

Proof. We use the basis decomposition 4.1 of the generator Ψ

$$\Psi(X) = D_{11}X + \sum_{i=2}^{n^2} D_{i1}B_i X + \sum_{j=2}^{n^2} D_{1j}X B_j^* + \sum_{i,j=2}^{n^2} D_{ij}B_i X B_j^*$$

As (D_{ij}) is Hermitian, if we denote $W := \sum_{i=2}^{n^2} D_{i1}B_i$ and $\kappa := D_{11}$ then $W^* = \sum_{j=2}^{n^2} D_{1j}B_j^*$ and κ is a real number. From the previous proposition 4.3 we know that $(D_{ij})_{i,j=2}^{n^2}$ is a positive matrix. Therefore there exists $A_1, A_2, \dots, A_k \in M_{n^2-1}$ such that $(D_{ij})_{i,j=2}^{n^2} = A_1 A_1^* + A_2 A_2^* + \dots + A_k A_k^*$. If we write $A_r = (a_r(p, q))$ then in terms of the coefficients of the matrix A_r we have

$$D_{ij} = \sum_{r=1}^k \sum_{p=2}^{n^2} a_r(i, p) \overline{a_r(j, p)}.$$

Substituting these in the above expression of Ψ and regrouping the terms we obtain

$$\begin{aligned} \Psi(X) &= \kappa X + WX + XW^* + \sum_{r=1}^k \sum_{p=2}^{n^2} \left(\sum_{i=2}^{n^2} a_r(i, p) B_i \right) X \left(\sum_{j=2}^{n^2} a_r(j, p) B_j \right)^* \\ &= \kappa X + WX + XW^* + \sum_{p=2}^{n^2} \sum_{r=1}^k V_{p,r} X V_{p,r}^* \end{aligned}$$

where $V_{p,r} := \sum_{i=2}^{n^2} a_r(i, p) B_i$. That the semigroup preserves the identity is equivalent to the generator mapping it to zero i.e. $\Psi(I_n) = 0$. Plugging this condition we get

$$0 = \kappa + W + W^* + \sum_p \sum_r V_{p,r} V_{p,r}^*.$$

So we can set $W = iH - \frac{1}{2}\kappa - \frac{1}{2} \sum_p \sum_r V_{p,r} V_{p,r}^*$, where H is a Hermitian matrix and substituting it in the above expression of $\Psi(X)$, we have

$$\begin{aligned} \Psi(X) &= \kappa X + \left(iH - \frac{1}{2}\kappa - \frac{1}{2} \sum_p \sum_r V_{p,r} V_{p,r}^* \right) X + X \left(iH - \frac{1}{2}\kappa - \frac{1}{2} \sum_p \sum_r V_{p,r}^* V_{p,r} \right) \\ &\quad + \sum_p \sum_r V_{p,r} X V_{p,r}^* \\ &= i[H, X] + \sum_p \sum_r \left\{ V_{p,r} X V_{p,r}^* - \frac{1}{2} \left(V_{p,r} V_{p,r}^* X - X V_{p,r}^* V_{p,r} \right) \right\} \end{aligned}$$

□

5. POSITIVE SEMIGROUPS ON M_2

In [Car04] positive semigroup has been characterized in terms of the generator of the semigroup. Here we give another characterization of the same, following the discussion above. The Pauli matrices are unitary matrices

$$\sigma_0 = \frac{1}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad \sigma_1 = \frac{1}{2} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad \sigma_2 = \frac{1}{2} \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \quad \sigma_3 = \frac{1}{2} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix},$$

which form an orthogonal basis of M_2 with respect to Hilbert-Schmidt inner product. Moreover, they satisfy the following relations

$$\sigma_p^2 = \sigma_0^2, \quad \sigma_p \sigma_q = -\sigma_q \sigma_p, \quad \sigma_p \sigma_q = i \sigma_r \sigma_0$$

if $(p, q, r) \in \{(1, 2, 3), (2, 3, 1), (3, 1, 2)\}$. Decomposing a matrix V with respect to these basis i.e $V = \sum_{p=0}^3 v_p \sigma_p$, we observe that V has the form

$$\frac{1}{2} \begin{bmatrix} v_0 + v_3 & v_1 - i v_2 \\ v_1 + i v_2 & v_0 - v_3 \end{bmatrix}.$$

It is easy to see that

- (i) V has trace zero if and only if $v_0 = 0$,
- (ii) it has rank one if and only if $\det(V) = 0$ i.e. $v_0^2 = v_1^2 + v_2^2 + v_3^2$.

For a semigroup $\exp(t\Phi)$ on M_2 we can write the generator again in terms of Pauli basis as in 4.1-

$$\Phi = T_S, \quad \text{where } S = \sum_{p,q=0}^3 D_{p,q} \sigma_p \otimes \sigma_q.$$

Proposition 5.1. *A semigroup $\exp(t\Phi)$ is positive if and only if the matrix $(D_{pq})_{p,q=0}^3$ is Hermitian and for all $v = (v_1, v_2, v_3) \in \mathbb{C}^3$ with $v_1^2 + v_2^2 + v_3^2 = 0$ we have*

$$(5.1) \quad \langle v | (D_{pq})_{p,q=1}^3 v \rangle \geq 0$$

where $(D_{pq})_{p,q=1}^3$ is the 3×3 submatrix of the matrix $(D_{ij})_{i,j=0}^3$, defined above. Moreover, it is identity preserving if and only if the following relations are satisfied,

$$\sum_{p=0}^3 D_{pp} = 0, \quad \text{and } (D_{p0} + D_{0p}) + i(D_{qr} - D_{rq}) = 0$$

for $(p, q, r) \in \{(1, 2, 3), (2, 3, 1), (3, 1, 2)\}$.

Proof. As a direct application of corollary 4.2 it follows that the semigroup $\exp(t\Phi)$ is positive if and only if for any $V \in M_2$ with $\text{rank}(V) = 1$ and $\text{Tr}(V) = 0$

$$(5.2) \quad \sum_{p,q=0}^3 D_{pq} \text{Tr}(\sigma_p V^*) \text{Tr}(\sigma_q V) \geq 0$$

As we observed, expanding V in the Pauli basis $V = \sum_{j=0}^3 v_j \sigma_j$ the rank and trace conditions translate into $v_0 = 0$ and $\sum_{j=1}^3 v_j^2 = 0$. Plugging this decomposition of

V in the expression 5.2 it becomes

$$\begin{aligned} \sum_{p,q=0}^3 \sum_{j,k=0}^3 D_{pq} \bar{v}_j v_k \text{Tr}(\sigma_p \sigma_j) \text{Tr}(\sigma_q \sigma_k) &= \sum_{p,q=0}^3 \sum_{j,k=0}^3 \bar{v}_j v_k \delta_{pj} \delta_{qk} \\ &= \frac{1}{4} \sum_{p,q=1}^3 D_{pq} \bar{v}_p v_q \geq 0, \end{aligned}$$

which gives the desired inequality.

To obtain the conditions of hermiticity and identity preserving property, it is easy to see that-

- (i) $\exp(t\Phi)$ is hermiticity preserving if and only if the generator $\Phi = T_S$ has the same property, which is equivalent to the matrix $(D_{pq})_{p,q=0}^3$ being hermitian.
- (ii) The semigroup preserves identity if and only if the generator Φ takes identity to zero i.e. $T_S(\sigma_0) = 0$.

□

6. A GENERAL 4-PARAMETER FAMILY OF SEMIGROUPS

Now we discuss as an example the semigroup generated by 4 parameter family $L = \alpha P + \beta D + \gamma T + \delta \text{Id}$ and time evolution for specific cases in M_2 . As all the operators P, D, T, Id commute, the semigroup generated is given by

$$\begin{aligned} \exp(tL) &= e^{t(\beta+\gamma+\delta)}(e^{t\alpha} - 1)P + e^{t(\gamma+\delta)}(e^{t\beta} - 1)D + e^{t\delta} \frac{(e^{t\gamma} - e^{-t\gamma})}{2} T \\ &\quad + e^{t\delta} \frac{(e^{t\gamma} + e^{-t\gamma})}{2} \text{Id} \end{aligned}$$

6.1. The depolarising channel. We consider the semigroup $T_{\alpha,\delta} = \exp t(\alpha P + \delta \text{Id})$. This semigroup is again a linear combination of the operator P and identity,

$$T_{\alpha,\delta}(t) = \exp(t(\alpha P + \delta \text{Id})) = \underbrace{e^{\delta t}(e^{\alpha t} - 1)}_{=: \alpha(t)} P + \underbrace{e^{\delta t}}_{=: \delta(t)} \text{Id},$$

which, by Lemma 2.13, is k -positive iff

$$\alpha(t) \geq 0 \quad \text{and} \quad \delta(t) \geq -\frac{\alpha(t)}{kn}$$

This condition is satisfied for all $t \geq 0$ iff $e^{\alpha t} \geq 1$ for all $t \geq 0$ iff $\alpha \geq 0$.

If we consider the identity preserving semigroup generated by the operator P and the identity operator i.e. the semigroup $T_{\alpha,-\alpha}$. we can see that it is a convex combination of P and the identity operator

$$T_{\alpha,-\alpha} = \exp t\alpha(P - \text{Id}) = (1 - e^{-t\alpha})P + e^{-t\alpha}\text{Id}.$$

which is completely positive for all $t \geq 0$ iff $\alpha \geq 0$. What is more, in this case the semigroup converges to P as $t \rightarrow \infty$, which we know by Lemma 2.11 to be an interior point of the cone \mathcal{EB} or 1-superpositive. So if $\alpha \geq 0$, the semigroup $T_{\alpha,-\alpha}$

enters the cone \mathcal{EB} in finite time $t_1 > 0$. The Choi matrix of the semigroup is given by

$$C_{T_{\alpha,-\alpha}} = \begin{bmatrix} \frac{1}{2}(1 + e^{-t\alpha}) & 0 & 0 & e^{-t\alpha} \\ 0 & \frac{1}{2}(1 - e^{-t\alpha}) & 0 & 0 \\ 0 & 0 & \frac{1}{2}(1 - e^{-t\alpha}) & 0 \\ e^{-t\alpha} & 0 & 0 & \frac{1}{2}(1 + e^{-t\alpha}) \end{bmatrix}.$$

To decide when the semigroup enters \mathcal{EB} we use Peres-Horodeci (or PPT) criterion. If T is the transpose map then

$$(\text{Id} \otimes T)C_{\exp(tL)} = \begin{bmatrix} \frac{1}{2}(1 + e^{-t\alpha}) & 0 & 0 & 0 \\ 0 & \frac{1}{2}(1 - e^{-t\alpha}) & e^{-t\alpha} & 0 \\ 0 & e^{-t\alpha} & \frac{1}{2}(1 - e^{-t\alpha}) & 0 \\ 0 & 0 & 0 & \frac{1}{2}(1 + e^{-t\alpha}) \end{bmatrix}.$$

So $T_{\alpha,-\alpha}$ is in \mathcal{EB} if and only if $C_{T_{\alpha,-\alpha}}$ is separable if and only if $(\text{Id} \otimes T)C_{T_{\alpha,-\alpha}}$ is positive (by PPT criterion, see [Per96],[HHH96]). We can easily check that the above matrix is positive iff the determinant

$$\begin{vmatrix} \frac{1}{2}(1 - e^{-t\alpha}) & e^{-t\alpha} \\ e^{-t\alpha} & \frac{1}{2}(1 - e^{-t\alpha}) \end{vmatrix} = -\frac{3}{4}e^{-2t\alpha} - 2e^{-t\alpha} + 1.$$

is positive. Replacing $x = e^{-t\alpha}$ we observe that the polynomial $-\frac{3}{4}x^2 - 2x + 1$ has the positive root $x_0 := -\frac{4}{3} + 2\frac{\sqrt{7}}{3}$ in the interval $[0, 1]$. The polynomial is positive on the interval $[0, x_0]$ and negative on $[x_0, 1]$. Thus the semigroup enters the cone \mathcal{S}_1 at the time

$$t_1 = -\frac{1}{\alpha} \ln \left(-\frac{4}{3} + \frac{2\sqrt{7}}{3} \right).$$

6.2. Transposition. We consider the identity preserving semigroup generated by the transposition and the depolarizing channel i.e. by the generator $L = \alpha P + \gamma T - (\alpha + \gamma)\text{Id}$, for $\alpha, \gamma \in \mathbb{R}$. Since all the quantities T, P, Id commutes we have

$$\begin{aligned} \exp(tL) &= \exp(t\gamma(T - \text{Id})) \exp(t\alpha(P - \text{Id})) \\ &= e^{-t\alpha} \frac{1 + e^{-2t\gamma}}{2} \text{Id} + e^{-t\alpha} \frac{1 - e^{-2t\gamma}}{2} T + (1 - e^{-t\alpha})P, \end{aligned}$$

which is a convex combination of the Id, T and P .

If we take $\alpha, \gamma > 0$, then in general it is a positive semigroup but not necessarily completely positive.

If the parameter $\alpha = 0$ then the semigroup is just convex combination of Id and T , which converges to $\frac{1}{2}(\text{Id} + T)$ as $t \rightarrow \infty$.

If $\alpha > 0$, then the semigroup eventually becomes completely positive, even 1-superpositive as it converges to P .

We compute the Choi matrix of the semigroup $\exp(tL)$,

$$(6.1) \quad C_{\exp(tL)} = \rho_t C_{\text{Id}} + \mu_t C_T + \nu_t C_P,$$

where $\rho_t = e^{-t\alpha} \frac{1 + e^{-2t\gamma}}{2}$, $\mu_t = e^{-t\alpha} \frac{1 - e^{-2t\gamma}}{2}$ and $\nu_t = (1 - e^{-t\alpha})$. Using the same arguments as in Proposition 2.12, the Choi matrix is positive iff the eigenvalues $\frac{\nu_t}{n} - \mu_t$ and $\frac{\nu_t}{n} + \mu_t$ are positive. Combining these two conditions, we conclude that the semigroup $\exp(tL)$ becomes CP at time t iff

$$\frac{\nu_t}{n} \geq |\mu_t| \quad \text{i.e.} \quad 2(e^{\alpha t} - 1) \geq n|1 - e^{-2t\gamma}|.$$

The above inequality shows that even if $\gamma \gg \alpha > 0$ it is possible that the semigroup not CP for certain time but ultimately becomes CP and then superpositive after finite time. If $\gamma = \alpha > 0$ then substituting $x = e^{\alpha t}$ in the above inequality gives the following criterion,

$$2x^3 - (2+n)x^2 + n \geq 0.$$

The polynomial $p(x) = 2x^3 - (2+n)x^2 + n$ has two positive roots- 1 and $\frac{n+\sqrt{n^2+8n}}{4}$, and $p(x) \geq 0$ for $x \geq \frac{n+\sqrt{n^2+8n}}{4}$. The root 1 corresponds to the time $t = 0$. So the semigroup becomes CP at time $t_1 = \frac{1}{\alpha} \ln \frac{n+\sqrt{n^2+8n}}{4}$.

We can find the time when the semigroup becomes 1-superpositive using again PPT criterion for the case $M_2(\mathbb{C})$.

$$(\text{Id} \otimes T)C_{\exp tL} = \begin{bmatrix} \rho_t + \mu_t + \frac{1}{2}\nu_t & 0 & 0 & \mu_t \\ 0 & \frac{1}{2}\nu_t & \rho_t & 0 \\ 0 & \rho_t & \frac{1}{2}\nu_t & 0 \\ \mu_t & 0 & 0 & \rho_t + \mu_t + \frac{1}{2}\nu_t \end{bmatrix}.$$

The above matrix is positive iff the determinant $\begin{vmatrix} \frac{1}{2}\nu_t & \rho_t \\ \rho_t & \frac{1}{2}\nu_t \end{vmatrix} = \frac{1}{4}\nu_t^2 - \rho_t^2$ is positive, which gives the condition

$$(6.2) \quad 2 \leq e^{t\alpha} - e^{-2t\gamma}.$$

If $\gamma = \alpha > 0$ then substituting $x = e^{t\alpha}$ in the above inequality gives that

$$x^3 - 2x^2 - 1 \geq 0.$$

If ξ is the positive root of the polynomial then we see that at the time $t_2 = \frac{1}{\alpha} \ln \xi$ the semigroup becomes 1-superpositive.

REFERENCES

- [AT07] Charalambos Dionisios Aliprantis, Rabee Tourky, Cones and duality, Graduate studies in Mathematics vol. 84, American Mathematical Society, 2007.
- [BCS20] Ivan Bardet, Benoît Collins, Gunjan Sapra, Characterization of equivariant maps and application to entanglement detection, Ann. Henri Poincaré 21 (2020), 3385-3406
- [Car04] Raffaella Carbone, Optimal log-Sobolev inequality and hypercontractivity for positive semigroups on $M_2(\mathbb{C})$. Infinite Dimensional Analysis. Quantum Probability and Related Topics, vol. 7, No. 3, 317-335.
- [CMW19] Matthias Christandl, Alexander Müller-Hermes, Michael M. Wolf, When do composed maps become entanglement breaking?, Ann. Henri Poincaré. 20(7):20295-20322, 2019
- [COS18] Benoît Collins, H. Osaka, G. Sapra, On a family of linear maps from $M_n(\mathbb{C})$ to $M_{n^2}(\mathbb{C})$. Linear Algebra Appl. 555, 398-411 (2018)
- [DMS21] Repana Devendra, Nirupama Mallik, Kappil Sumesh, Mapping cone of k-entanglement breaking maps. <https://arxiv.org/abs/2105.14991v2>
- [FS00] Uwe Franz and Michael Schürmann, Lévy processes on quantum hypergroups. In: Heyer, Herbert (ed.) et al., Infinite dimensional harmonic analysis. Transactions of the 2nd Japanese-German symposium, University of Kyoto, Japan, September 20-24, 1999. p. 93-114 (2000).
- [GKS76] Vittorio Gorini, Andrzej Kossakowski, E.C. George Sudarshan, Completely positive dynamical semigroups of N-level systems. Journal of Mathematical Physics. 17, 821 (1976)
- [HHH96] Michal Horodecki, Pawel Horodecki, Ryszard Horodecki, Separability of mixed states: necessary and sufficient conditions, vol. 223, 1-8, 1996.
- [HSR03] Michael Horodecki, Peter W. Shor, Mary Beth Ruskai, Entanglement breaking channels, Reviews in Mathematical Physics. vol. 15, No. 6, 629-641, 2003.

- [JPPY23] Yeong-Gwang Jung, Jeongeun Park, Sang-Jung Park, and Sang-Gyun Youn, A universal framework for entanglement detection under group symmetry, arXiv: 2301.03849v1
- [Lind76] Göran Lindblad, On the generators of quantum dynamical semigroups. *Communications in Mathematical Physics*. 48, 119-130 (1976).
- [Per96] Asher Peres, Separability criterion for density matrices, *Phys. Rev. Lett.* 77, 1413-1415, 1996.
- [Sch85] Michael Schürmann, Positive and conditionally positive linear functionals on coalgebras. *Quantum probability and applications II*, Proc. 2nd Workshop, Heidelberg/Ger. 1984, Lect. Notes Math. 1136, 475-492 (1985).
- [Sim11] Barry Simon, *Convexity: An analytical viewpoint*, Cambridge Tracts in Mathematics 187. Cambridge: Cambridge University Press 2011.
- [SSZ09] Lukasz Skowronek, Erling Størmer, and Karol Życzkowski, Cones of positive maps and their duality relations. *J. Math. Phys.* 50, No. 6, 062106 (2009).
- [SV70] Hans Schneider and Mathukumalli Vidyasagar, Cross-positive matrices. *SIAM J. numer. Anal.* 7, 508-519 (1970).
- [Tom85] Jun Tomiyama, On the geometry of positive maps in matrix algebras. II. *Linear Algebra Appl.* 69, 169-177 (1985)

INDIAN STATISTICAL INSTITUTE, STAT-MATH. UNIT, R V COLLEGE POST, BENGALURU 560059, INDIA

Email address: bhat@isibang.ac.in

LABORATOIRE DE MATHÉMATIQUES DE BESANÇON, UNIVERSITÉ DE FRANCHE-COMTÉ, 16, ROUTE DE GRAY, 25 030 BESANÇON CEDEX, FRANCE

Email address: purbayan.chakraborty@univ-fcomte.fr

U.F., DÉPARTEMENT DE MATHÉMATIQUES DE BESANÇON, UNIVERSITÉ DE FRANCHE-COMTÉ, 16, ROUTE DE GRAY, 25 030 BESANÇON CEDEX, FRANCE

Email address: uwe.franz@univ-fcomte.fr

URL: <http://lmb.univ-fcomte.fr/uwe-franz>