

# Testing quantum satisfiability

Ashley Montanaro<sup>\*1,2</sup>, Changpeng Shao<sup>†1</sup>, and Dominic Verdon<sup>‡1</sup>

<sup>1</sup>School of Mathematics, University of Bristol, UK

<sup>2</sup>Phasecraft Ltd., UK

January 26, 2023

## Abstract

Quantum  $k$ -SAT (the problem of determining whether a  $k$ -local Hamiltonian is frustration-free) is known to be  $\text{QMA}_1$ -complete for  $k \geq 3$ , and hence likely hard for quantum computers to solve. Building on a classical result of Alon and Shapira, we show that quantum  $k$ -SAT can be solved in randomised polynomial time given the ‘property testing’ promise that the instance is either satisfiable (by any state) or far from satisfiable by a product state; by ‘far from satisfiable by a product state’ we mean that  $\epsilon n^k$  constraints must be removed before a product state solution exists, for some fixed  $\epsilon > 0$ . The proof has two steps: we first show that for a satisfiable instance of quantum  $k$ -SAT, most subproblems on a constant number of qubits are satisfiable by a product state. We then show that for an instance of quantum  $k$ -SAT which is far from satisfiable by a product state, most subproblems are unsatisfiable by a product state. Given the promise, quantum  $k$ -SAT may therefore be solved by checking satisfiability by a product state on randomly chosen subsystems of constant size.

## 1 Introduction

**Property testing.** In computer science, one often needs to determine whether a structure defined by a large quantity of data possesses a certain property which is *global*, in the sense that it is defined with reference to all the data. Property testing algorithms aim to determine with high probability whether a structure possesses a global property by performing *local* checks; that is, checking properties of randomly chosen small subsets of the data defining the structure. Since the time complexity of a local check is smaller than that of a global check, and one only inspects a small subset of the data, this approach, when feasible, often leads to algorithms with excellent time and query complexity.

Of course, in general the structure may not possess the global property, but be close enough to possessing it that one is highly unlikely to perform a local check which witnesses this fact. For this reason property testing algorithms usually require a promise that the structure either possesses the property or is far from possessing the property, where distance from possessing the property is defined by a measure specific to the problem and is usually quantified by some constant  $\epsilon$ .

In [AS03], Alon and Shapira showed that the NP-hard problem of classical  $k$ -SAT (satisfiability of Boolean functions on  $n$  variables, where each function depends on exactly  $k$  variables) is amenable to a property testing approach. Specifically, with the promise that an instance of classical  $k$ -SAT is either satisfiable or far from satisfiable, satisfiability may be determined in randomised constant time, by choosing constant-sized subsets of variables and checking satisfiability of those functions which depend only on variables in those subsets. Because of the promise, their result is limited to dense instances of  $k$ -SAT, where there are  $\Omega(n^k)$  functions.

---

\*ashley.montanaro@bristol.ac.uk

†changpeng.shao@bristol.ac.uk

‡dominic.verdon@bristol.ac.uk

**Testing quantum satisfiability.** In [Bra11] Bravyi defined a quantum analogue of classical  $k$ -SAT, which we will here call quantum  $k$ -SAT. An instance of this problem is defined by  $n$  Hilbert spaces of dimension 2 (qubits) and, for every subset  $s$  of  $k$  qubits, a projector  $\Pi_s$  on the Hilbert space of those qubits; we say that the instance is satisfiable if there is a state of all  $n$  qubits which is in the kernel of all of the projections. While classical  $k$ -SAT is NP-complete whenever  $k \geq 3$ , quantum  $k$ -SAT is QMA<sub>1</sub>-complete whenever  $k \geq 3$ , where QMA<sub>1</sub> is the quantum analogue of the class MA with one-sided error. In the terminology of physics quantum  $k$ -SAT is precisely the problem of determining whether a  $k$ -local Hamiltonian on  $n$  qubits is frustration free.

Since the problem seems very difficult in general, previous work on algorithms for quantum  $k$ -SAT has focused on tractable special cases. There are many interesting results on random quantum  $k$ -SAT, for instance [LLM<sup>+</sup>10, LMSS10, BMR10, HLL<sup>+</sup>13]. It has been shown that quantum 2-SAT can be solved in linear time [dBG16, ASSZ18]; moreover, in this case there is always a satisfying state which is a product of one- and two-qubit states [CCD<sup>+</sup>11]. The case where there are relatively few nontrivial projectors all of rank 1 is also often tractable [AKS12, AdGS21].

In this work we also treat a special case: the dense case, where the interaction hypergraph (i.e. the  $k$ -regular hypergraph on  $n$  vertices whose edges are subsets of  $k$  qubits whose associated projection is nontrivial) contains  $\Omega(n^k)$  edges. We show that the results of [AS03] extend to the quantum setting. Given the promise that the instance is either satisfiable or far from satisfiable *by a product state*, quantum  $k$ -SAT may be solved in randomised polynomial time in  $n$  by checking satisfiability by a product state on constant-sized subsets of qubits. (The polynomial dependence on  $n$  is due to the fact that, in the data specifying an instance of quantum  $k$ -SAT, the projectors are given to  $\text{poly}(n)$  precision, so all computations involving them will pick up a  $\text{poly}(n)$  factor.)

**Some comments on our results.** Before stating our results formally, we will make two comments. Firstly, as we emphasised in the last paragraph, the promise is that the instance, if unsatisfiable, is far from satisfiable *by a product state*. Obviously, this is a less stringent requirement than being far from satisfiable by any state. The reason we are able to weaken the promise in this way is that an instance of quantum  $k$ -SAT which is satisfiable by any state is locally satisfiable by a product state with high probability. This result (Theorem 1.5) is essentially a fact about quantum states which may be of independent interest; there are some connections with monogamy of entanglement [KW04, CW04, BCY11] and de Finetti theorems [BH16, BH13] which we discuss briefly in Remark 2.12.

Secondly, our proofs are combinatorial in nature; we make no use of the norm. In particular, we do not require the polynomial lower bound on the ground state energy in the unsatisfiable case that is usually given as a promise in the definition of quantum  $k$ -SAT; our testing algorithm therefore solves a harder problem than quantum  $k$ -SAT as it is usually defined.

## 1.1 Results

We will first state the three basic definitions we will use in this work. Throughout we write  $\sigma_k(S)$  for the set of  $k$ -element subsets of a set  $S$  and  $[n]$  for the set  $\{1, \dots, n\}$ . For any subset  $x \in \sigma_k(S)$  we use the notation  $\bar{x} := (S - x)$  for the complement.

**Definition 1.1** (Quantum  $k$ -SAT). An instance of quantum  $k$ -SAT (without a lower bound on the ground state energy) is defined by the following data.

- A number of qubits  $n$ .
- For every  $k$ -element subset  $s \in \sigma_k([n])$ , a projector  $\Pi_s$  on  $(\mathbb{C}^2)^{\otimes n}$  which acts non-trivially only on the qubits in  $s$ . These projectors are given as matrices whose entries have precision polynomial in  $n$  (i.e. they are specified by  $\text{poly}(n)$  bits).

We write the instance as  $([n], \{\Pi_s\})$ . The problem is to determine whether there exists some state  $|x\rangle \in (\mathbb{C}^2)^{\otimes n}$  such that the following equation holds:

$$\sum_{s \in \sigma_k([n])} \Pi_s |x\rangle = 0. \quad (1)$$

If such a state exists we say that the instance is *satisfiable*; if it does not we say that the instance is *unsatisfiable*. If there is a product state  $|\phi\rangle$  satisfying (1) we say that the instance is *satisfiable by a product state*.

The usual definition of quantum  $k$ -SAT includes a promise that the ground-state energy has a polynomial lower bound if the instance is unsatisfiable [Bra11, §1]; Definition 1.1 is therefore a strictly harder problem.

**Definition 1.2** ( $\epsilon$ -far from satisfiable by a product state). Let  $\Sigma \subset \sigma_k([n])$  be such that, for some product state  $|\phi\rangle \in (\mathbb{C}^2)^{\otimes n}$ :

$$\sum_{s \in (\sigma_k([n]) - \Sigma)} \Pi_s |\phi\rangle = 0. \quad (2)$$

We say that an instance  $([n], \{\Pi_s\})$  of quantum  $k$ -SAT is  $\epsilon$ -far from satisfiable by a product state if this implies that  $|\Sigma| \geq \epsilon n^k$ .

**Remark 1.3.** Our definition of  $\epsilon$ -far differs slightly from that in [AS03]; whereas they remove individual clauses (which in our case correspond to rank-one projections) we remove the whole projection for a given  $k$ -subset. This does not make much difference to the overall theory, since defining  $\epsilon$ -far in terms of the number of rank-one projections removed would only alter  $\epsilon$  by a constant factor.

**Definition 1.4** (Local satisfiability). Let  $([n], \{\Pi_s\}_{s \in \sigma_k([n])})$  be an instance of quantum  $k$ -SAT. Let  $C \subset [n]$ .

We say that the *restriction* of the instance to  $C$  is the instance of quantum  $k$ -SAT defined by the data  $(C, \{\Pi_s\}_{s \in \sigma_k(C)})$ . We say that  $([n], \{\Pi_s\})$  is *locally satisfiable by a product state at  $C$*  if  $(C, \{\Pi_s\}_{s \in \sigma_k(C)})$  is satisfiable by a product state.

Our result is based on the following two theorems, whose proofs are given in the remainder of the paper.

**Theorem 1.5.** *Let  $([n], \{\Pi_s\})$  be a satisfiable instance of quantum  $k$ -SAT, and let  $c \in \mathbb{N}$  be fixed, where  $c \geq 3$ . Let  $C \in \sigma_c([n])$  be a subset chosen uniformly at random. The probability that the instance is locally satisfiable by a product state at  $C$  is greater than  $p \in (0, 1)$  whenever  $n > \Psi(p, c)$ , where  $\Psi(p, c) = O(2^{6c} / \log(1/p))$ .*

*Equivalently, let  $|\psi\rangle \in (\mathbb{C}^2)^{\otimes n}$  be any state and let  $C \in \sigma_c([n])$  be a subset chosen uniformly at random. The probability that the subspace  $\text{supp}(\text{Tr}_{\bar{C}}(|\psi\rangle\langle\psi|)) \subseteq (\mathbb{C}^2)^{\otimes c}$  contains a product state is greater than  $p \in (0, 1)$  whenever  $n > \Psi(p, c)$ .*

**Remark 1.6.** The theorem applies to the case where  $c \geq 3$ ; a similar result holds for  $c = 2$ , but the result for  $c = 2$  is much simpler and stronger and is treated separately in Proposition 2.11. We give a precise bound on  $\Psi(p, c)$  in the proof. The constant factors in this bound are far from optimal, since we made many approximations to simplify the proof; if required, the reader could easily improve the bound by tightening the analysis without changing the structure of the proof. However, we would be surprised if the exponential dependence on  $c$  could be removed without a different approach.

For the second theorem, we assume that  $n$  is large enough (but still finite).

**Theorem 1.7.** *There exists a constant  $c(k, \epsilon)$  (i.e. not depending on  $n$ ) such that the following holds. Let  $([n], \{\Pi_s\})$  be any instance of quantum  $k$ -SAT which is  $\epsilon$ -far from satisfiable by a product state. Then, for a randomly chosen subset  $C \in \sigma_{c(k, \epsilon)}([n])$ , the instance is locally unsatisfiable by a product state at  $C$  with probability  $p > 0.75$ .*

**Remark 1.8.** The main weakness of Theorem 1.7 is that we do not have a good upper bound for the constant  $c(k, \epsilon)$ . Our proof in Section 3 does yield an upper bound, but it is extremely large (bigger than  $((3!)!) \cdots !$ , where the chain of factorials has length  $(5/\epsilon)4^{k-1}$ ). The bound has this form because the degree of the  $m$ -qubit Segre variety is the factorial  $m!$  (Lemma 3.6). It is an open question whether the upper bound can be made small enough for the testing result to be practically useful rather than merely theoretically interesting. Improvements on this scale are not unknown in classical property testing (see e.g. the discussion in [AS03, § 1.1]).

**Remark 1.9.** Together, Theorems 1.5 and 1.7 imply that, for large enough  $n$ , an instance  $([n], \{\Pi_s\})$  of quantum  $k$ -SAT cannot be both satisfiable and far from satisfiable by a product state.

The consequent result about testability of quantum  $k$ -SAT is as follows.

**Corollary 1.10.** *With a promise that the instance is either satisfiable or  $\epsilon$ -far from satisfiable by a product state, quantum  $k$ -SAT may be solved in randomised polynomial time in  $n$ .*

*Proof.* Let  $([n], \{\Pi_s\})$  be an instance of quantum  $k$ -SAT, and let  $c$  be some constant. Given a subset  $C \in \sigma_c([n])$ , we can check whether or not  $(n, \{\Pi_s\})$  is locally satisfiable by a product state at  $C$  in time polynomial in  $n$  (and exponential in  $c$ ) using Gröbner basis methods. Indeed, let  $\{\Pi_1, \dots, \Pi_N\}$  be the set of nontrivial projectors on  $C$ . Let  $\Pi_i = \sum_{j=1}^{r_i} |a_{ij}\rangle \langle a_{ij}|$ , where  $r_i \leq 2^k$  is the rank of  $\pi_i$  and  $|a_{ij}\rangle \in (\mathbb{C}^2)^{\otimes k}$ . Let  $|\phi\rangle = |\phi_1\rangle \otimes \cdots \otimes |\phi_c\rangle$  be a unknown product state defined on  $C$ . We set

$$|\phi_i\rangle = x_{i0}|0\rangle + x_{i1}|1\rangle, \quad i \in \{1, 2, \dots, c\}.$$

Let  $s_i \in \sigma_k(C)$  be the set of qubits on which  $\Pi_i$  acts. For each  $s_i$ , we write  $|\phi_{s_i}\rangle := \bigotimes_{v \in s_i} |\phi_v\rangle$ . Then (1) implies the following equations:

$$\langle a_{ij} | \phi_{s_i} \rangle = 0, \quad i \in \{1, 2, \dots, N\}, j \in \{1, 2, \dots, r_i\}. \quad (3)$$

We also have that  $|x_{i0}|^2 + |x_{i1}|^2 = 1$ . As we can rescale (3) arbitrarily, we can relax this to  $|x_{i0}|^2 + |x_{i1}|^2 \neq 0$ ; equivalently,  $x_{ij} \neq 0$  for at least one  $j$ . To ensure this we introduce some new variables  $\{y_{ij}\}$  with the same index sets as the  $\{x_{ij}\}$ , and some new equations:

$$(x_{i0}y_{i0} - 1)(x_{i1}y_{i1} - 1) = 0, \quad i \in \{1, \dots, c\}. \quad (4)$$

In total we obtain  $c + \sum_{i=1}^N r_i$  equations. Each equation has degree at most  $k$  or 2, and there are  $4c$  variables over the field  $\mathbb{C}$ . We want to determine if there is a solution for this polynomial system over  $\mathbb{C}$ . It is known that this polynomial system has no zeros in  $\mathbb{C}$  if and only if its Gröbner basis is  $\{1\}$ . This is Hilbert's Nullstellensatz, which is an EXPSpace-complete problem [VZGG13, Chapter 21.7]. By [BFS15, Proposition 1], an upper bound on the time complexity in this case is

$$D(c + \sum_{i=1}^N r_i) \binom{4c + D}{D}^\omega \text{poly}(n),$$

where  $D \leq 1 + 3c + (k-1) \sum_{i=1}^N r_i$  is the maximal degree of the elements in the reduced Gröbner basis, and  $\omega < 2.373$  is the exponent of matrix multiplication. The  $\text{poly}(n)$  factor is due to the polynomial precision of the matrix entries specifying the quantum  $k$ -SAT instance (Definition 1.1).

We can therefore straightforwardly check satisfiability of an instance  $([n], \{\Pi_s\})$  of quantum  $k$ -SAT with the testing promise. Since we are only interested in scaling w.r.t.  $n$ , we may assume that  $n$  is large enough that (by Theorem 1.5) if the instance is satisfiable then it will be locally satisfiable by a product state on  $c(k, \epsilon)$ -subsets with probability  $p > 0.75$ , and (by Theorem 1.7) if the instance is  $\epsilon$ -far from satisfiable by a product state then it will be locally unsatisfiable by a product state on  $c(k, \epsilon)$ -subsets with probability  $p > 0.75$ .

We choose subsets  $\{C_i \in \sigma_{c(k,\epsilon)}([n])\}_{i=1}^m$  at random and check whether the instance is locally satisfiable by a product state on these subsets. If the majority are locally satisfiable by a product state we conclude that the instance is satisfiable; otherwise we conclude that the instance is  $\epsilon$ -far from satisfiable by a product state. By the Chernoff bound for the binomial distribution the probability that the conclusion is incorrect tends exponentially to zero as  $m$  increases.  $\square$

## 1.2 Acknowledgements

We thank Niel de Beaudrap and Aram Harrow for useful discussions. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 817581). We acknowledge support from EPSRC grant EP/T001062/1.

## 2 Proof of Theorem 1.5

We will make use of some combinatorial lemmas, some of which are well-known. For the reader's convenience we prove all these lemmas, but to avoid a long digression we have relegated the proofs to Appendix A. Throughout we consider the binomial coefficient  $\binom{x}{k}$  to be the polynomial  $x(x-1)\cdots(x-k+1)/k!$ . It is defined for every real number  $x$ , and is positive and increasing for  $x \geq k-1$ .

**Lemma 2.1** (Scaling binomial coefficients). *Let  $\theta, x \in \mathbb{R}_{\geq 0}$  and  $c \in \mathbb{N}$ , where  $0 \leq \theta \leq 1$  and  $x \geq c$ . Then, for some  $\alpha \in [0, 1)$ , we have:*

$$\theta \binom{x}{c} = \binom{\theta^{1/c}x}{c} + \alpha \binom{\theta^{1/c}x}{c-1} \leq \binom{\theta^{1/c}x + 1}{c}. \quad (5)$$

**Definition 2.2.** Let  $G$  be an  $l$ -regular hypergraph on  $n$  vertices, and let  $k \leq l$ . We define the  $k$ -shadow  $G|_k$  of  $G$  to be the  $k$ -regular hypergraph on  $n$  vertices whose edges are precisely those  $k$ -subsets which are contained in an edge of  $G$ .

**Definition 2.3** (c.f. [BE15, Fit18]). Let  $G$  be an  $l$ -regular hypergraph on  $n$  vertices, let  $k \leq l$ , and let  $\omega \in [0, 1]$ . We say that a  $k$ -regular hypergraph on  $n$  vertices  $G|_{k,\omega}$  is a *partial  $(k, \omega)$ -shadow* of  $G$  if it can be obtained by the following construction. For each edge  $E$  of  $G$ , choose a subset  $K_E \subset \sigma_k(E)$  of size  $|K_E| = \lfloor \omega \binom{l}{k} \rfloor$ . The edge set of  $G|_{k,\omega}$  is then defined as  $\bigcup_{E \in G} K_E$ .

Obviously there are many different partial  $(k, \omega)$ -shadows of an  $l$ -regular hypergraph  $G$ ; they depend on the choice made of the set  $K_E$  for each edge  $E \in G$ . Recall that the *edge density*  $0 \leq \theta \leq 1$  of a  $k$ -regular hypergraph on  $n$  vertices with edge set  $E$  is defined by  $|E| = \theta \binom{n}{k}$ .

**Lemma 2.4** (Edge density of shadows [Kee08, Thm. 1]). *If  $G$  is an  $l$ -regular hypergraph with edge density  $\theta$ , then the  $k$ -shadow  $G|_k$  has edge density  $\theta|_k \geq (\theta^{1/l} - 2/n)^k$ .*

**Lemma 2.5** (Edge density of partial shadows). *Let  $n \in \mathbb{N}$  and let  $c \in \mathbb{N}$ . Let  $l := n/2^c + \delta$  and  $k := l/2 + \epsilon$ , where  $-3 \leq \delta < -1$  and  $-1/2 \leq \epsilon \leq 0$ , be natural numbers. Let  $G$  be an  $l$ -regular hypergraph on  $n$  vertices with edge density  $\theta \in [0, 1]$ .*

*Suppose that  $n \geq 3(2^c)$  and  $n \geq 3(2^{c+1}) \left( \frac{1}{2} + \frac{1}{\ln(A)} \left( \frac{A^{1/3}}{4} - \ln(\theta) \right) \right)$  for some constant  $A > 1$ . Then the edge density  $\theta|_{k,1/2}$  of a partial  $(k, 1/2)$ -shadow of  $G$  has the following lower bound:*

$$\theta|_{k,1/2} \geq \frac{\theta}{K_A} := \frac{\theta}{2\sqrt{A \exp(A^{1/3}/2)}}. \quad (6)$$

**Lemma 2.6** (Edge density for a certain construction). *Let  $n \in \mathbb{N}$  be odd. We construct an  $(n-1)/2$ -regular hypergraph  $G$  as follows. For every subset  $X \in \sigma_{(n-1)/2}([n])$ , we do precisely one of the following:*

1. Add  $X$  as an edge to  $G$ .
2. Add all the subsets in  $\sigma_{(n-1)/2}(\overline{X})$  as edges to  $G$ .

The edge density  $\theta$  of a hypergraph  $G$  constructed in this way satisfies  $\theta \geq 1/2$ .

Recall that the *degree* of a vertex in a hypergraph is the number of edges containing it.

**Lemma 2.7** (Degree of a random vertex). *Let  $G$  be a  $k$ -regular hypergraph on  $n$  vertices with edge density  $\theta$ , where  $k = \beta n$  for some  $\beta \in [0, 1]$ . Let  $N$  be the degree of a vertex picked uniformly at random, and let  $\tau := N/\binom{n-1}{k-1}$ . Then, for any  $\epsilon > 0$  and  $\alpha \in (0, 1)$ :*

$$\Pr[|\theta - \tau| \geq \min(\Delta(\alpha, \beta, n), (1 - \theta)(1 - \alpha))] \leq \alpha, \quad (7)$$

where

$$\Delta(\alpha, \beta, n) = \frac{5 \exp\left(\frac{1}{12\alpha(1-\alpha)n} + \frac{1}{12\beta(1-\beta)n}\right)}{\sqrt{2\pi\alpha(1-\alpha)\beta(1-\beta)n}} =: \frac{5(L_{\alpha,\beta})^{1/n}}{M_{\alpha,\beta}\sqrt{n}}. \quad (8)$$

The final ingredients for the proof are two facts about entangled subspaces. Recall that a subspace  $L \subset \mathbb{C}^2 \otimes \mathbb{C}^m$  is *completely entangled* if there is no product state in  $L$ .

**Lemma 2.8** ([ATL11, Lem. 1 and Lem. 2]). *Let  $L \subset \mathbb{C}^2 \otimes \mathbb{C}^m$  be a completely entangled subspace, for any  $m \geq 2$ . Then the orthogonal complement  $L^\perp \subset \mathbb{C}^2 \otimes \mathbb{C}^m$  contains a product state.*

**Corollary 2.9.** *Let  $m \in \mathbb{N}$ , and let  $V \subset (\mathbb{C}^2)^{\otimes m}$  be a subspace of dimension  $\geq 2$ . Let  $L \subset \mathbb{C}^2 \otimes (\mathbb{C}^2)^{\otimes m}$  be a subspace such that:*

1.  $\mathbb{C}^2 \otimes V^\perp \subset L$ .
2.  $(\mathbb{C}^2 \otimes V) \cap L$  is a completely entangled subspace of  $\mathbb{C}^2 \otimes V$ .

Then  $L^\perp \subset \mathbb{C}^2 \otimes \mathbb{C}^m$  contains a product state.

*Proof.* Since  $\mathbb{C}^2 \otimes V^\perp \subset L$ , we have  $L = (\mathbb{C}^2 \otimes V^\perp) \oplus ((\mathbb{C}^2 \otimes V^\perp)^\perp \cap L) = (\mathbb{C}^2 \otimes V^\perp) \oplus ((\mathbb{C}^2 \otimes V) \cap L)$ . Then  $L^\perp = (\mathbb{C}^2 \otimes V) \cap ((\mathbb{C}^2 \otimes V) \cap L)^\perp = ((\mathbb{C}^2 \otimes V) \cap L)^\perp$ , where  $\perp$  is the orthogonal complement in  $\mathbb{C}^2 \otimes V$ . But  $(\mathbb{C}^2 \otimes V) \cap L$  is a completely entangled subspace of  $\mathbb{C}^2 \otimes V$ , so by Lemma 2.8 we know that  $((\mathbb{C}^2 \otimes V) \cap L)^\perp$  contains a product state; therefore  $L^\perp$  contains a product state.  $\square$

**Lemma 2.10.** *Let  $|\psi\rangle \in (\mathbb{C}^2)^{\otimes n}$  be a state and let  $x \in [n]$ . Let  $S_1, S_2 \subset ([n] - \{x\})$  be two subsets such that  $S_1 \cap S_2 = \emptyset$  and  $S_1 \sqcup S_2 = ([n] - \{x\})$ . Then at least one of the following is true:*

1. The subspace  $\text{supp}(\text{Tr}_{S_2}(|\psi\rangle\langle\psi|))$  contains a product state  $|\phi_x\rangle \otimes |\phi_{S_1}\rangle$ .
2. The subspace  $\text{supp}(\text{Tr}_{S_1}(|\psi\rangle\langle\psi|))$  contains a product state  $|\phi_x\rangle \otimes |\phi_{S_2}\rangle$ .

*Proof.* To clarify our notation, we first recall the definition of supports and kernels. By definition, for any subset  $S \subset [n]$ , we have  $\text{supp}(\text{Tr}_{\overline{S}}(|\psi\rangle\langle\psi|)) = \ker(\text{Tr}_{\overline{S}}(|\psi\rangle\langle\psi|))^\perp$ , where the orthogonal complement is taken in the space  $(\mathbb{C}^2)^{\otimes |S|}$ . If we write  $(\mathbb{C}^2)^{\otimes n} = (\mathbb{C}^2)^{\otimes |S|} \otimes (\mathbb{C}^2)^{\otimes (n-|S|)}$ , where the first factor corresponds to those qubits in the set  $S$ , then:

$$\ker(\text{Tr}_{\overline{S}}(|\psi\rangle\langle\psi|)) = \text{span}\{|v\rangle \in (\mathbb{C}^2)^{\otimes |S|} : \langle v| \otimes \mathbb{1} |\psi\rangle = 0\}, \quad (9)$$

$$\text{supp}(\text{Tr}_S(|\psi\rangle\langle\psi|)) = \text{span}\{(\langle v| \otimes \mathbb{1}) |\psi\rangle : |v\rangle \in (\mathbb{C}^2)^{|S|}\}. \quad (10)$$

We now observe that if any of (i)  $\dim(\text{supp}(\text{Tr}_{\overline{S}}(|\psi\rangle\langle\psi|)))$ , (ii)  $\dim(\text{supp}(\text{Tr}_{\overline{S_1}}(|\psi\rangle\langle\psi|)))$  or (iii)  $\dim(\text{supp}(\text{Tr}_{\overline{S_2}}(|\psi\rangle\langle\psi|)))$  equal 1, then at least one of the two claims in this lemma is true. Indeed, for (i) observe that if  $\dim(\text{supp}(\text{Tr}_{\overline{S}}(|\psi\rangle\langle\psi|))) = 1$ , then  $|\psi\rangle = |\psi_x\rangle \otimes |\psi_{[n]-\{x\}}\rangle$ . To see this, observe that there exists  $|w\rangle \in \mathbb{C}^2$  such that  $(\langle w| \otimes \mathbb{1}) |\psi\rangle = 0$ ;

it follows by the Schmidt decomposition that  $|\psi\rangle$  is a product state. The argument for (ii) and (iii) is similar.

We therefore now assume that those three dimensions are greater than 1. We consider the subspace

$$\begin{aligned} \ker(\mathrm{Tr}_{\{x\} \sqcup S_1}(|\psi\rangle\langle\psi|)) \cap (\mathrm{supp}(\mathrm{Tr}_{\{x\}}(|\psi\rangle\langle\psi|)) \otimes \mathrm{supp}(\mathrm{Tr}_{S_1}(|\psi\rangle\langle\psi|))) \\ \subseteq \mathrm{supp}(\mathrm{Tr}_{\{x\}}(|\psi\rangle\langle\psi|)) \otimes \mathrm{supp}(\mathrm{Tr}_{S_1}(|\psi\rangle\langle\psi|)). \end{aligned}$$

We have a dichotomy:

- (a) The subspace  $\ker(\mathrm{Tr}_{\{x\} \sqcup S_1}(|\psi\rangle\langle\psi|)) \cap (\mathrm{supp}(\mathrm{Tr}_{\{x\}}(|\psi\rangle\langle\psi|)) \otimes \mathrm{supp}(\mathrm{Tr}_{S_1}(|\psi\rangle\langle\psi|)))$  is a completely entangled subspace of  $\mathrm{supp}(\mathrm{Tr}_{\{x\}}(|\psi\rangle\langle\psi|)) \otimes \mathrm{supp}(\mathrm{Tr}_{S_1}(|\psi\rangle\langle\psi|))$ .
- (b) The subspace  $\ker(\mathrm{Tr}_{\{x\} \sqcup S_1}(|\psi\rangle\langle\psi|)) \cap (\mathrm{supp}(\mathrm{Tr}_{\{x\}}(|\psi\rangle\langle\psi|)) \otimes \mathrm{supp}(\mathrm{Tr}_{S_1}(|\psi\rangle\langle\psi|)))$  contains a product state.

In case (a), we are precisely in the situation of Corollary 2.9, where  $V = \mathrm{supp}(\mathrm{Tr}_{S_1}(|\psi\rangle\langle\psi|))$  and  $L = \ker(\mathrm{Tr}_{\{x\} \sqcup S_1}(|\psi\rangle\langle\psi|))$ ; therefore claim 1 is true. On the other hand, in case (b) there exists a product state  $|v\rangle \otimes |w\rangle \in \mathrm{supp}(\mathrm{Tr}_{\{x\}}(|\psi\rangle\langle\psi|)) \otimes \mathrm{supp}(\mathrm{Tr}_{S_1}(|\psi\rangle\langle\psi|))$  such that  $\langle v| \otimes \langle w| \otimes \mathbb{1} |\psi\rangle = 0$ . Consider the state  $|\psi'\rangle := (\mathbb{1} \otimes \langle w| \otimes \mathbb{1}) |\psi\rangle \in \mathrm{supp}(\mathrm{Tr}_{S_1}(|\psi\rangle\langle\psi|))$ . Since  $\langle v| \otimes \mathbb{1} |\psi'\rangle = 0$ , we have that  $\dim(\ker(\mathrm{Tr}_{\{x\}}(|\psi'\rangle\langle\psi'|))) = 1$ ; by the Schmidt decomposition this implies that  $|\psi'\rangle = |\psi'_x\rangle \otimes |\psi'_{S_2}\rangle$ , and so claim 2 is true.  $\square$

We can now prove the theorem. To warm up, we prove the case  $c = 2$  first.

**Proposition 2.11.** *Let  $([n], \{\Pi_s\})$  be a satisfiable instance of quantum  $k$ -SAT, and let  $x_1 \in [n]$  be any qubit. Then there is at most one qubit  $x_2 \in ([n] - \{x_1\})$  such that the instance is not locally satisfiable by a product state at  $\{x_1, x_2\}$ .*

*Equivalently, let  $|\psi\rangle \in (\mathbb{C}^2)^{\otimes n}$  be any state, and let  $x_1 \in [n]$  be any qubit. Then there is at most one qubit  $x_2 \in ([n] - \{x_1\})$  such that the subspace  $\mathrm{supp}(\mathrm{Tr}_{\{x_1, x_2\}}(|\psi\rangle\langle\psi|)) \subseteq (\mathbb{C}^2)^{\otimes 2}$  does not contain a product state.*

*Proof.* The equivalence of the two statements will be shown in the first paragraph of the proof of Theorem 1.5 below. We therefore need only prove the second statement. Let  $x_2 \in ([n] - \{x_1\})$  be any qubit. We now use Lemma 2.10; in the notation of that lemma, let  $x := x_1$ ,  $S_1 := \{x_2\}$  and  $S_2 := ([n] - \{x_1, x_2\})$ . Then either  $\mathrm{supp}(\mathrm{Tr}_{\{x_1, x_2\}}(|\psi\rangle\langle\psi|))$  contains a product state or  $\mathrm{supp}(\mathrm{Tr}_{\{x_1, x'\}}(|\psi\rangle\langle\psi|))$  contains a product state for any  $x' \in ([n] - \{x_1, x_2\})$ .  $\square$

We now move onto the case  $c \geq 3$ .

**Theorem** (Restatement of Theorem 1.5). *Let  $([n], \{\Pi_s\})$  be a satisfiable instance of quantum  $k$ -SAT, and let  $c \in \mathbb{N}$  be some constant, where  $c \geq 3$ . Let  $C \in \sigma_c([n])$  be a subset chosen uniformly at random. The probability that the instance is locally satisfiable by a product state at  $C$  is greater than  $p \in (0, 1)$  whenever:*

$$n > \Psi(p, c) := \max \left( \frac{(2^c - 1)(c - 1) \ln(4)}{\ln(1/p)} + (2^c - 2), \quad e^2 \left( \frac{180(c - 2)(2)^{3(c-2)}}{\sqrt{2\pi}(3 - 1/(2)^{c-2})} \right)^2, \quad 6(2^{c+1})(c - 1) \right).$$

*Equivalently, let  $|\psi\rangle \in (\mathbb{C}^2)^{\otimes n}$  be any state and let  $C \in \sigma_c([n])$  be a subset chosen uniformly at random. The probability that the subspace  $\mathrm{supp}(\mathrm{Tr}_C(|\psi\rangle\langle\psi|)) \subseteq (\mathbb{C}^2)^{\otimes c}$  contains a product state is greater than  $p \in (0, 1)$  whenever  $n > \Psi(p, c)$ .*

**Remark 2.12.** Before giving the proof, we remark that the following weaker fact follows quite straightforwardly from known results:

Let  $([n], \{\Pi_s\})$  be a satisfiable instance of quantum  $k$ -SAT, let  $c \in \mathbb{N}$  be some constant, and let  $\epsilon > 0$ . Let  $C \in \sigma_c([n])$  be a subset chosen uniformly at random. The probability that there exists a product state  $|\phi\rangle \in (\mathbb{C}^2)^{\otimes c}$  such that  $\sum_{s \in \sigma_k(C)} \langle \phi | \Pi_s | \phi \rangle < \epsilon$  is greater than  $p$  whenever  $p > O(n^{k-c-1/4}) \binom{c}{k} / \epsilon$ .

To prove this, one can use [BH16, Eq. 91], which implies that there is a product state  $|\phi\rangle \in (\mathbb{C}^2)^{\otimes n}$  such that  $\sum_{s \in \sigma_k([n])} \langle \phi | \Pi_s | \phi \rangle \leq O(n^{k-1/4})$ . We can write

$$\sum_{s \in \sigma_k([n])} \langle \phi | \Pi_s | \phi \rangle = \sum_{C \in \sigma_c([n])} \sum_{s \in \sigma_k(C)} \frac{1}{\binom{c}{k}} \langle \phi | \Pi_s | \phi \rangle = \sum_{C \in \sigma_c([n])} S_C,$$

where  $S_C := \sum_{s \in \sigma_k(C)} \binom{c}{k}^{-1} \langle \phi | \Pi_s | \phi \rangle$ . Since there are  $O(n^c)$   $c$ -subsets, the expected value of  $S_C$  for a randomly chosen  $c$ -subset is  $O(n^{(k-c)-1/4})$ ; the result then follows by Markov's inequality.

An alternative approach to proving this fact, which has some issues but nevertheless reveals something about the nature of the problem, is to use monogamy of the squashed entanglement [KW04, Thm. 8]. For this, let  $|\psi\rangle$  be a state satisfying  $([n], \{\Pi_s\})$ , and let  $l_1, l_2$  be constants. It follows straightforwardly from monogamy that, for a randomly chosen  $C \in \sigma_c([n])$ , the expected entanglement across every bipartition of  $C$  is  $O(1/n)$ . Then by [BCY11, Thm. §II],  $\text{Tr}_{\overline{C}}(|\psi\rangle\langle\psi|)$  is close to a biseparable state across every bipartition of  $C$ . If we could conclude that this implied closeness to a fully separable state, the fact would follow.

Theorem 1.5 is a stronger, exact analogue, where we demand exact local satisfiability by a product state rather than approximate satisfiability. We were unable to derive Theorem 1.5 from the non-exact result, and the proof we are about to provide follows a different approach, which does not make use of any continuous entanglement measure.

*Proof of Theorem 1.5.* We first observe the equivalence of the two statements in this theorem. If  $([n], \{\Pi_s\})$  is satisfiable, let  $|\psi\rangle \in (\mathbb{C}^2)^{\otimes n}$  be a satisfying state; clearly any state  $|\psi_C\rangle \in \text{supp}(\text{Tr}_{\overline{C}}(|\psi\rangle\langle\psi|))$  is a local solution to  $([n], \{\Pi_s\})$  at  $C$ , so the second statement implies the first. In the other direction, let  $|\psi\rangle \in (\mathbb{C}^2)^{\otimes n}$  be a state; then for all  $C \in \sigma_c([n])$  let  $\Pi_C$  be the projector onto  $\ker(\text{Tr}_{\overline{C}}(|\psi\rangle\langle\psi|))$ , and consider the instance  $([n], \{\Pi_C\}_{C \in \sigma_c([n])})$  of quantum  $c$ -SAT, which is of course satisfied by the state  $|\psi\rangle$ .

Given this equivalence it is sufficient to show that for any state  $|\psi\rangle \in (\mathbb{C}^2)^{\otimes n}$  and a randomly chosen subset  $C \in \sigma_c([n])$ , there exists a product state in  $\text{supp}(\text{Tr}_{\overline{C}}(|\psi\rangle\langle\psi|))$  with high probability.

We will choose qubits  $\{x_1, \dots, x_c\} =: C$  one by one at random. Let us pick the first qubit  $x_1$ . There are two possibilities:

- (a)  $n - 1$  is even.
- (b)  $n - 1$  is odd.

In case (a), let  $\mathcal{B}([n] - \{x_1\})$  be the set of all *equal bipartitions* of the set  $([n] - \{x_1\})$ , where an equal bipartition is a pair of sets  $B_1, B_2 \subset ([n] - \{x_1\})$  of size  $|B_1| = |B_2| = (n - 1)/2$  such that  $B_1 \cup B_2 = ([n] - \{x_1\})$ . We write such a bipartition as  $B_1^i | B_2^i$ , where  $i \in I$  is some index for the bipartitions. We observe that, for any finite set  $S$  with even cardinality,  $|\mathcal{B}(S)| = \binom{|S|}{|S|/2} / 2$ . Now, for any equal bipartition  $B_1 | B_2 \in \mathcal{B}([n] - \{x_1\})$ , we know by Lemma 2.10 that at least one of the following holds:

- (i) The subspace  $\text{supp}(\text{Tr}_{B_2}(|\psi\rangle\langle\psi|))$  contains a product state  $|\phi_{x_1}\rangle \otimes |\phi_{B_1}\rangle$ .
- (ii) The subspace  $\text{supp}(\text{Tr}_{B_1}(|\psi\rangle\langle\psi|))$  contains a product state  $|\phi_{x_1}\rangle \otimes |\phi_{B_2}\rangle$ .

We define an  $(n - 1)/2$ -regular hypergraph  $G_1$  on the set  $([n] - \{x_1\})$  as follows. For each equal bipartition  $B_1^i | B_2^i \in \mathcal{B}([n] - \{x_1\})$ , if  $\text{supp}(\text{Tr}_{B_j^i}(|\psi\rangle\langle\psi|))$  contains a product state then add the set  $B_j^i$  as an edge to the graph  $G_1$ . Clearly, the edge density  $\theta_1$  of  $G_1$  has the lower bound  $\theta_1 \geq 1/2$ . The physical significance of this hypergraph  $G_1$  is that,



for any edge  $E$  of  $G_1$ , we know that  $\text{supp}(\text{Tr}_{\overline{\{x_1\} \sqcup E}}(|\psi\rangle\langle\psi|))$  contains a product state, which we will write as  $|\psi_1\rangle \otimes |\psi_E\rangle$ .

This definition of  $G_1$  depended on the fact that  $n - 1$  was even. In case (b), we make a slightly different definition. For any subset  $X \in \sigma_{n/2-1}([n] - \{x_1\})$ , at least one of the following holds, by Lemma 2.10:

- (i) The subspace  $\text{supp}(\text{Tr}_{\overline{\{x_1\} \sqcup X}}(|\psi\rangle\langle\psi|))$  contains a product state  $|\phi_{x_1}\rangle \otimes |\phi_X\rangle$ .
- (ii) The subspace  $\text{supp}(\text{Tr}_X(|\psi\rangle\langle\psi|))$  contains a product state  $|\phi_{x_1}\rangle \otimes |\phi_{\overline{\{x_1\} \sqcup X}}\rangle$ .

We now define an  $(n/2 - 1)$ -regular hypergraph  $G_1$  on the set  $([n] - \{x_1\})$  as follows. For every  $X \in \sigma_{n/2-1}([n] - \{x_1\})$ , if  $\text{supp}(\text{Tr}_{\overline{\{x_1\} \sqcup X}}(|\psi\rangle\langle\psi|))$  contains a product state then we add  $X$  as an edge to  $G_1$ . On the other hand, if  $\text{supp}(\text{Tr}_X(|\psi\rangle\langle\psi|))$  contains a product state then we add all the subsets in  $\sigma_{n/2-1}(\overline{\{x_1\} \sqcup X})$  as edges to  $G_1$ . By Lemma 2.6 the edge density  $\theta_1$  of  $G$  satisfies  $\theta_1 \geq 1/2$ . Again, the physical significance of this hypergraph  $G_1$  is that, for any edge  $E$  of  $G_1$ , we know that  $\text{supp}(\text{Tr}_{\overline{\{x_1\} \sqcup E}}(|\psi\rangle\langle\psi|))$  contains a product state, which we will write as  $|\psi_1\rangle \otimes |\psi_E\rangle$ .

Having obtained the hypergraph  $G_1$ , we now pick the next qubit  $x_2 \in ([n] - \{x_1\})$  at random. We define a new hypergraph  $W_2$  ( $W$  for ‘working’, since this graph is an intermediate step towards obtaining  $G_2$ ) on the set  $([n] - \{x_1, x_2\})$  with the edges  $\{(E - \{x_2\}) \mid E \text{ is an edge of } G_1 \text{ and } x_2 \in E\}$ . This hypergraph  $W_2$  is  $k_2$ -regular, where  $k_2 = ((n - 3)/2)$  if  $n - 1$  was even, and  $k_2 = (n/2 - 2)$  if  $n - 1$  was odd. Now we define the graph  $G_2$ ; again the definition will depend on whether  $k_2$  is even or odd. If  $k_2$  is even then, for every edge  $E$  of  $W_2$  and every  $X \in \sigma_{k_2/2}(E)$ , we again have the following dichotomy, by Lemma 2.10:

- (i) The subspace  $\text{supp}(\text{Tr}_{\overline{\{x_1, x_2\} \sqcup X}}(|\psi_1\rangle \otimes |\psi_E\rangle))$  contains a product state  $|\psi_1\rangle \otimes |\psi_2\rangle \otimes |\psi_X\rangle$ .
- (ii) The subspace  $\text{supp}(\text{Tr}_X(|\psi_1\rangle \otimes |\psi_E\rangle))$  contains a product state  $|\psi_1\rangle \otimes |\psi_2\rangle \otimes |\psi_{E - (X \sqcup x_2)}\rangle$ .

We then define the  $(k_2/2)$ -regular hypergraph  $G_2$  as follows. For every edge  $E$  in  $W_2$ , and every  $X \in \sigma_{k_2/2}(E)$ , if  $\text{supp}(\text{Tr}_{\overline{\{x_1, x_2\} \sqcup X}}(|\psi_1\rangle \otimes |\psi_E\rangle))$  contains a product state then we add  $X$  as an edge to  $G_2$ . On the other hand, if  $\text{supp}(\text{Tr}_X(|\psi_1\rangle \otimes |\psi_E\rangle))$  contains a product state then we add  $\overline{X}$  as an edge to  $G_2$ . This graph contains a partial  $(k_2/2, 1/2)$ -shadow of  $W_2$ .

If  $k_2$  is odd then we define the  $(k_2 - 1)/2$ -regular hypergraph  $G_2$  as follows: for every edge  $E$  of  $W_2$  and every  $X \in \sigma_{(k_2-1)/2}(E)$ , if  $\text{supp}(\text{Tr}_{\overline{\{x_1, x_2\} \sqcup X}}(|\psi_1\rangle \otimes |\psi_E\rangle))$  contains a product state then we add  $X$  as an edge to  $G_2$ . On the other hand, if  $\text{supp}(\text{Tr}_X(|\psi_1\rangle \otimes |\psi_E\rangle))$  contains a product state then we add all the subsets in  $\sigma_{(k_2-1)/2}(\overline{X})$  as edges to  $G_2$ . By Lemma 2.6 applied to each edge  $E$ , this graph contains a partial  $((k_2 - 1)/2, 1/2)$ -shadow of  $W_2$ .

The physical significance of the hypergraph  $G_2$  is that for any edge  $E$  of  $G_2$ , we have that  $\text{supp}(\text{Tr}_{\overline{\{x_1, x_2\} \sqcup E}}(|\psi\rangle\langle\psi|))$  contains a product state over the qubit  $x_1$ , the qubit  $x_2$  and the qubits in  $E$ , which we will write as  $|\psi_1\rangle \otimes |\psi_2\rangle \otimes |\psi_E\rangle$ .

We now iterate this construction for the qubits  $\{x_3, \dots, x_{c-1}\}$  and obtain a  $k_{c-1}$ -regular hypergraph  $G_{c-1}$  on the set  $([n] - \{x_1, \dots, x_{c-1}\})$ ; for any edge  $E$  of  $G_{c-1}$ , we know that  $\text{supp}(\text{Tr}_{\overline{\{x_1, \dots, x_{c-1}\} \sqcup E}}(|\psi\rangle\langle\psi|))$  contains a product state, which we write as  $|\psi_1\rangle \otimes \dots \otimes |\psi_{c-1}\rangle \otimes |\psi_E\rangle$ .

Let us obtain bounds on the edge size  $k_{c-1}$  and edge density  $\theta_{c-1}$  of the hypergraph  $G_{c-1}$ . We determine the edge size first. The hypergraph  $G_1$  was  $k_1$ -regular, where either  $k_1 = (n - 1)/2$  or  $k_1 = (n/2 - 1)$ , and had edge density  $\theta_1 \geq 1/2$ . To get  $G_2$  from  $G_1$  we defined the graph  $W_2$  by picking a single qubit; this graph had edges of size  $l_2 = k_1 - 1$ . Iterating this construction, we see that in general the graph  $G_{i+1}$  will have edges of size  $k_{i+1} = l_{i+1}/2$  or  $k_{i+1} = l_{i+1}/2 - 1/2$ , where  $l_{i+1} = k_i - 1$ . It follows that  $k_i/2 > k_{i+1} \geq k_i/2 - 1$ , and  $k_0 = n$ . Solving the linear recurrence, we obtain:

$$n/2^i > k_i \geq n/2^i + 1/2^{i-1} - 2 > n/2^i - 2,$$

$$n/2^i - 1 > l_{i+1} \geq n/2^i + 1/2^{i-1} - 3 > n/2^i - 3. \quad (11)$$

In particular,

$$n/2^{c-1} > k_{c-1} \geq n/2^{c-1} + 1/2^{c-2} - 2. \quad (12)$$

We now obtain a lower bound on the edge density  $\theta_{c-1}$ . To simplify the calculation we will make numerous approximations. By (11), Lemma 2.5 will always hold when we go from  $W_i$  to  $G_i$ , as long as  $n$  is big enough; we at least need  $n \geq 3(2^{c-1})$  for the step from  $W_{c-1}$  to  $G_{c-1}$ , so we assume this from now on. There is a free constant  $A$  in Lemma 2.5; we arbitrarily set it to  $A = 216W(2^{2/3}/6)^3 \simeq 2.107$ , where  $W(x)$  is the Lambert  $W$ -function; this implies that  $K_A = 4$ .

On the other hand, Lemma 2.7 gives us a bound on the change in edge density when we go from  $G_i$  to  $W_{i+1}$ . The bound is a minimum over two functions; we will just consider the function (8). Let  $\beta_i$  be the value of  $\beta$  when we apply Lemma 2.7 to go from  $G_i$  to  $W_i$ ; by (11), we always have  $1/2 > 1/2^i > \beta_i \geq 1/2^i - 2/n \geq 1/2^i - 2/3(2^{c-2}) \geq 1/3(2^{c-2})$ . Let  $\bar{\beta} := 1/3(2^{c-2})$ . We also fix some  $\bar{\alpha} \in (0, 1)$ , which we will determine later. By the union bound, Lemma 2.5 and Lemma 2.7 we then have

$$\Pr[\theta_{c-1} \geq \Phi] \geq 1 - (c-2)(1 - \bar{\alpha}), \quad (13)$$

where  $\Phi := \frac{1}{2(4)^{c-2}} - (c-2)\Delta(\bar{\alpha}, \bar{\beta}, n)$ .

Finally, we pick the qubit  $x_c$ . As long as it is contained in an edge of  $G_{c-1}$ , there will be a product state in  $\text{supp}(\text{Tr}_{\{x_1, \dots, x_{c-1}, x_c\}}(|\psi\rangle\langle\psi|))$ . The worst case is that the edges of  $G_{c-1}$  form a complete hypergraph on a subset  $C \subset ([n] - \{x_1, \dots, x_{c-1}\})$ , with some overspill onto a single vertex not in  $C$ . Let  $|C| = \sigma(n - c + 1)$  for some  $\sigma \in [0, 1]$ . By (12) and Lemma 2.1, we have:

$$\sigma \geq (\theta_{c-1})^{\frac{2^{c-1}}{n+2-2^c}} \geq (\Phi)^{\frac{2^{c-1}}{n+2-2^c}}. \quad (14)$$

Let  $p$  be the probability that  $\text{supp}(\text{Tr}_{\bar{C}}(|\psi\rangle\langle\psi|))$  contains a product state; by (13) and (14) we have that

$$p \geq (1 - (c-2)(1 - \bar{\alpha}))(\Phi)^{\frac{2^{c-1}}{n+2-2^c}}. \quad (15)$$

We will assume that  $p = \bar{p}$  is given and find values for the as yet undefined variables  $\bar{\alpha}, n$  which satisfy (15) while minimising  $n$ . (Again, we will make approximations, so the minimisation will not be tight.) Assuming  $1 - (c-2)(1 - \bar{\alpha}) > 0$ , we need:

$$\left( \frac{\bar{p}}{(1 - (c-2)(1 - \bar{\alpha}))} \right)^{\frac{n+2-2^c}{2^{c-1}}} \leq \frac{1}{2(4)^{c-2}} - (c-2) \frac{5(L_{\bar{\alpha}, \bar{\beta}})^{1/n}}{M_{\bar{\alpha}, \bar{\beta}} \sqrt{n}}. \quad (16)$$

We make  $n$  big enough that

$$(c-2) \frac{5(L_{\bar{\alpha}, \bar{\beta}})^{1/n}}{M_{\bar{\alpha}, \bar{\beta}} \sqrt{n}} \leq \frac{1}{2t(4)^{c-2}} \quad (17)$$

for some  $t > 1$ ; this is to say that

$$\frac{1}{n} \ln(L_{\bar{\alpha}, \bar{\beta}}) - \frac{1}{2} \ln(n) \leq \ln \left( \frac{M_{\bar{\alpha}, \bar{\beta}}}{10(c-2)(4)^{c-2}t} \right). \quad (18)$$

Let us assume that  $\bar{\alpha} \geq 1 - 1/3(2^{c-2})$ ; then

$$L_{\bar{\alpha}, \bar{\beta}} \leq \exp \left( \frac{1}{6\bar{\alpha}(1 - \bar{\alpha})} \right), \quad M_{\bar{\alpha}, \bar{\beta}} \geq \sqrt{2\pi\bar{\alpha}}(1 - \bar{\alpha}),$$

and the equation (18) reduces to

$$\frac{1}{6n\bar{\alpha}(1 - \bar{\alpha})} - \frac{1}{2} \ln(n) \leq \ln \left( \frac{\sqrt{2\pi\bar{\alpha}}(1 - \bar{\alpha})}{10(c-2)(4)^{c-2}t} \right). \quad (19)$$

Now assume that

$$\ln(n) \geq 2 \left( \ln \left( \frac{10(c-2)(4)^{c-2}t}{\sqrt{2\pi\bar{\alpha}}(1-\bar{\alpha})} \right) + 1 \right) \quad \Leftrightarrow \quad n \geq e^2 \left( \frac{10t(c-2)(4)^{c-2}}{\sqrt{2\pi\bar{\alpha}}(1-\bar{\alpha})} \right)^2, \quad (20)$$

so the equation (19) reduces to:

$$\frac{1}{6n\bar{\alpha}(1-\bar{\alpha})} \leq 1 \quad \Leftrightarrow \quad n \geq \frac{1}{6\bar{\alpha}(1-\bar{\alpha})}$$

which is already implied by (20). By (17), the equation (16) then reduces to:

$$\left( \frac{\bar{p}}{1-(c-2)(1-\bar{\alpha})} \right)^{\frac{n+2-2^c}{2^c-1}} \leq \frac{t-1}{2t(4)^{c-2}}. \quad (21)$$

Clearly a larger  $\bar{\alpha}$  here will reduce the required size of  $n$  in (21), but it will worsen it in (20); so we set  $\bar{\alpha} = 1 - 1/3(2^{c-2})$ . We then assume that:

$$\frac{\bar{p}}{1-(c-2)(1-\bar{\alpha})} < 1 \quad \Leftrightarrow \quad \bar{p} < 1 - \frac{c-2}{3(2^{c-2})}.$$

We also arbitrarily set  $t = 2$ . Given these assumptions, (21) reduces to

$$n \geq \frac{(2^c - 1)(c - 1) \ln(4)}{\ln(1 - (c - 2)/3(2^{c-2})) - \ln(\bar{p})} + (2^c - 2),$$

which is implied by the simpler

$$n \geq \frac{(2^c - 1)(c - 1) \ln(4)}{\ln(1/\bar{p})} + (2^c - 2). \quad (22)$$

In addition to the lower bounds (20) and (22), there is also a lower bound on  $n$  coming from the second lower bound on  $n$  in Lemma 2.5. By (13) and (17) we always have  $\theta_i \geq \Phi \geq \frac{1}{4(4)^{c-2}}$ , so using  $\ln(A) \geq \ln(2)$  we obtain the following bound:

$$n \geq 6(2^{c+1})(c - 1).$$

Substituting the chosen values of  $\bar{\alpha}$  and  $t$  into (20), we obtain the theorem.  $\square$

### 3 Proof of Theorem 1.7

We first remark that one idea for a proof of this theorem is to add the usual polynomial ground state energy bound to the definition of quantum  $k$ -SAT, then use the standard approach with a  $\delta$ -net to reduce the problem of satisfiability by a product state to an instance of classical  $k'$ -SAT. It is straightforwardly seen that this instance of  $k'$ -SAT is also  $\epsilon$ -far from satisfiable, so the result from [AS03] can be directly applied to show local unsatisfiability. The problem is that the constant  $\delta$  determining the precision of the net, and therefore also the number  $k'$ , depends polynomially on  $n$ ; this implies that the size of the subsets to be checked also depends at least polynomially on  $n$ . Since the time taken to check for a product state solution is exponential in the size of the subset, as we saw in the proof of Corollary 1.10, we end up with time exponential in  $n$ .

Instead, we give a proof which is heavily inspired by Alon and Shapira's proof of the classical analogue (to the point where we were able to lift most of the constants from their work), and which is in any case more general than the above approach since it does not require any lower bound on the ground state energy when an instance of quantum  $k$ -SAT is unsatisfiable. Firstly, we consider the problem of extending a local assignment, which in our setting is a product subspace. We define the notion of a bad qubit; that is, a qubit to which either the local assignment cannot be extended, or such that extension to that qubit greatly reduces the dimension of possible extensions to the other qubits.

If, in the course of constructing a local assignment by extension, one extends through  $5(4)^{k-1}/\epsilon$  bad qubits, then the assignment can no longer be extended (Lemma 3.4). We show that if an instance of quantum  $k$ -SAT is  $\epsilon$ -far from satisfiable by a product state, then there are always more than  $\epsilon n/5$  bad qubits with respect to any local assignment (Lemma 3.5).

At this point, Alon and Shapira [AS03, Claim 2.4] built a binary tree of possible assignments to a set of randomly picked variables and showed that each branch of the tree will run into more than  $5(4)^{k-1}/\epsilon$  bad variables with high probability; they then used the union bound to conclude that there is no local assignment to those variables with high probability. This approach cannot be applied directly in the quantum setting because there are in general continuously many assignments to each variable (the space of possible assignments is a Hilbert space and not a binary set) and it is therefore not possible to build a finite tree. Instead, we use a backtracking approach which is summarised in the first paragraph of the proof of the theorem at the end of this section. We still need to discretise somehow; for this we use Lemma 3.6, which implies that we need only eliminate a finite number of possible solutions.

**Notation 3.1.** Throughout this section we write  $|H| := \dim(H)$  for the dimension of a Hilbert space  $H$  and  $|X|$  for the cardinality of a set  $X$ . When the context does not adequately distinguish these two meanings we explicitly use  $\dim(H)$  for the dimension and  $\#(X)$  for the cardinality.

Let  $V, W$  be two sets. We write  $\sigma_k(\underline{V} \sqcup W)$  for the set of  $k$ -element subsets of  $V \sqcup W$  containing all the elements of  $V$ ; that is, underlining one of the sets in the union means all its elements must be contained in the  $k$ -element subsets.

For a subset  $X \subseteq [n]$  we write  $H_X := (\mathbb{C}^2)^{\otimes |X|}$  for the Hilbert space of the qubits in  $X$ .

**Definition 3.2.** Fix a subset  $S \subset [n]$ , and let  $H_S := (\mathbb{C}^2)^{\otimes |S|}$  be the Hilbert space of the qubits in  $S$ . We say that an *assignment* to  $S$  is a product subspace  $P_S \subseteq H_S$  such that every state in the subspace is a local solution for  $([n], \{\Pi_s\})$  at  $S$ ; that is, a subspace  $P_S = V_1 \otimes \cdots \otimes V_{|S|} \subseteq H_S$  such that every state  $|\phi_S\rangle \in P_S$  satisfies  $\sum_{s \in \sigma_k(S)} \Pi_s |\phi_S\rangle = 0$ . For every  $\{x_1, \dots, x_j\} \in \sigma_j([n] - S)$ , where  $1 \leq j \leq k-1$ , we define  $L_{S, P_S}(x_1, \dots, x_j) \subseteq H_{x_1, \dots, x_j}$  to be the space of all states  $|v_{x_1, \dots, x_j}\rangle \in H_{x_1, \dots, x_j}$  such that

$$\sum_{s \in \sigma_k(\{\underline{x_1, \dots, x_j}\} \sqcup S)} \Pi_s (|v_{x_1, \dots, x_j}\rangle \otimes |\phi_S\rangle) = 0 \quad \forall |\phi_S\rangle \in P_S. \quad (23)$$

We define  $L_{S, P_S, j} := \bigoplus_{\{x_1, \dots, x_j\} \in \sigma_j([n] - S)} L_{S, P_S}(x_1, \dots, x_j)$ . In words,  $L_{S, P_S}(x_1, \dots, x_j)$  is the space of possible assignments  $|v_{x_1, \dots, x_j}\rangle$  on the qubits  $x_1, \dots, x_j$  that are compatible with the assignment  $(S, P_S)$ , in the sense that  $|v_{x_1, \dots, x_j}\rangle \otimes P_S$  is in the kernel of every projector on the qubits  $\{x_1, \dots, x_j\} \sqcup S$  which acts on all of the qubits  $x_1, \dots, x_j$ . The space  $L_{S, P_S, j}$  is the direct sum of all these possible  $j$ -qubit ‘extensions’ of the assignment  $(S, P_S)$ , and its dimension measures how many possible  $j$ -qubit extensions there are to the local assignment  $(S, P_S)$ .

In what follows we will consider taking a qubit  $x \notin S$  and extending the assignment  $P_S$  on  $S$  to an assignment  $V_x \otimes P_S$  on  $\{x\} \sqcup S$ , where  $V_x \subseteq L_{S, P_S}(x)$ . We define

$$\delta_{S, P_S, x, V_x, j} := \sum_{\{x_1, \dots, x_j\} \in \sigma_j([n] - (\{x\} \sqcup S))} |L_{S, P_S}(x_1, \dots, x_j)| - |L_{\{x\} \sqcup S, V_x \otimes P_S}(x_1, \dots, x_j)|$$

and  $\delta_{S, P_S, x, V_x} := \sum_{j=1}^{k-1} \delta_{S, P_S, x, V_x, j} n^{k-j-1}$ . In words,  $\delta_{S, P_S, x, V_x}$  measures how much the dimension of the space of possible extensions has been reduced by extending the assignment  $(S, P_S)$  to  $(\{x\} \sqcup S, V_x \otimes P_S)$ .

We say that a qubit  $x \in ([n] - S)$  *conflicts* with respect to (w.r.t.)  $(S, P_S)$  if  $L_{S, P_S}(x) = \emptyset$ . If a qubit  $x \in ([n] - S)$  does not conflict w.r.t.  $(S, P_S)$ , we say that it is *heavy* w.r.t.  $(S, P_S)$  if  $\min_{V_x \subseteq L_{S, P_S}(x)} \delta_{S, P_S, x, V_x} > \epsilon n^{k-1}/5$ . We say that a qubit  $x \in ([n] - S)$  is *bad* w.r.t.  $(S, P_S)$  if it is either heavy or conflicting.

Finally, in the proof of Lemma 3.5 we will define a set of subsets  $\Sigma \subset \sigma_k([n])$  and only consider projectors on subsets not in  $\Sigma$ . For a local assignment  $(S, P_S)$  and every  $\{x_1, \dots, x_j\} \in \sigma_j([n] - S)$  we therefore define  $L_{S, P_S}^\Sigma(x_1, \dots, x_j) \subset H_{x_1, \dots, x_j}$  to be the space of all states  $|v_{x_1, \dots, x_j}\rangle \in H_{x_1, \dots, x_j}$  such that

$$\sum_{s \in \sigma_k(\{x_1, \dots, x_j\} \sqcup S) - \Sigma} \Pi_s(|v_{x_1, \dots, x_j}\rangle \otimes |\phi_S\rangle) = 0 \quad \forall |\phi_S\rangle \in P_S.$$

This is identical to (3.2), but with the projectors in  $\Sigma$  removed. The other definitions above can be similarly generalised.

We first show that it is impossible to extend an assignment through more than  $5(d^2)^{k-1}/\epsilon$  heavy qubits.

**Definition 3.3.** We define a *chain* of length  $m$  to be a local assignment  $(\{y_1, \dots, y_m\}, P_1 \otimes \dots \otimes P_m)$  such that, for each  $2 \leq l \leq m$ , each qubit  $y_l$  is heavy with respect to the assignment  $(\{y_1, \dots, y_{l-1}\}, P_1 \otimes \dots \otimes P_{l-1})$ , and  $y_1$  is heavy with respect to the empty assignment  $(\emptyset, \emptyset)$ .

**Lemma 3.4.** Let  $(\{y_1, \dots, y_\gamma\}, P_1 \otimes \dots \otimes P_\gamma)$  be a chain of length  $\gamma(k, \epsilon) := 5(4)^{k-1}/\epsilon$ . Then every qubit  $x \in ([n] - \{y_1, \dots, y_\gamma\})$  is conflicting w.r.t. the chain.

*Proof.* Let us construct the chain  $(\{y_1, \dots, y_\gamma\}, P_1 \otimes \dots \otimes P_\gamma)$  starting from the empty assignment  $(\emptyset, \emptyset)$ . We define

$$W_i := \sum_{j=1}^{k-1} |L_{\{y_1, \dots, y_i\}, P_1 \otimes \dots \otimes P_{i,j}}| n^{k-j-1}.$$

The initial size of each  $|L_{\emptyset, \emptyset, j}|$  is at most  $\sum_{\{v_1, \dots, v_j\} \in \sigma_j([n])} 2^j = \binom{n}{j} 2^j \leq \frac{2^j}{j!} n^j$ . Therefore the initial value of  $W_0 = \sum_{j=1}^{k-1} |L_{\emptyset, \emptyset, j}| n^{k-j-1}$  is at most  $\sum_{j=1}^{k-1} \frac{2^j}{j!} n^j n^{k-j-1} \leq 2^{k-1} n^{k-1}$ .

When we extend to the qubit  $y_i$  from the assignment  $(\{y_1, \dots, y_{i-1}\}, P_1 \otimes \dots \otimes P_{i-1})$  we have

$$\sum_{j=1}^{k-1} \delta_{\{y_1, \dots, y_{i-1}\}, P_1 \otimes \dots \otimes P_{i-1, y_i, P_{i,j}}} n^{k-j-1} = \delta_{\{y_1, \dots, y_{i-1}\}, P_1 \otimes \dots \otimes P_{i-1, y_i, P_i}} > \epsilon n^{k-1}/5$$

by definition of heaviness, and therefore  $W_i \leq W_{i-1} - \epsilon n^{k-1}/5$ . So by the time we have made  $5d^{k-1}/\epsilon$  extensions we must have  $W = 0$ . In particular  $L_{\{y_1, \dots, y_\gamma\}, P_1 \otimes \dots \otimes P_\gamma}(x) = 0$  for all  $x \in ([n] - \{y_1, \dots, y_\gamma\})$ .  $\square$

We now show that the  $\epsilon$ -far condition implies that there will always be many bad qubits with respect to any local assignment.

**Lemma 3.5.** Let  $([n], \{\Pi_s\})$  be an instance of quantum  $k$ -SAT which is  $\epsilon$ -far from satisfiable by a product state. Let  $S \subset [n]$  and let  $P_S$  be some local assignment to  $S$ . Then there are at least  $\epsilon n/5$  bad qubits w.r.t.  $(S, P_S)$ .

*Proof.* Suppose that there are fewer than  $\epsilon n/5$  bad qubits w.r.t.  $(S, P_S)$ . We will define a subset  $\Sigma \subset \sigma_k([n])$  satisfying (2) and such that  $|\Sigma| < \epsilon n^k$ , contradicting the assumption.

This is a two-step process. Let  $X_{nb} \subset ([n] - S)$  be the set of qubits which are not bad w.r.t.  $(S, P_S)$ . In the first step we will extend the assignment  $(S, P_S)$  to the qubits in  $X_{nb}$  one at a time, while adding  $k$ -subsets to  $\Sigma$  according to the following prescription. Let  $x \in X_{nb}$  be the first qubit to which we extend; we choose some  $V_x \subseteq L_{S, P_S}(x)$  minimising  $\delta_{S, P_S, x, V_x}$ , and extend the assignment to  $(\{x\} \sqcup S, V_x \otimes P_S)$ . Let us consider in turn each  $\{x_1, \dots, x_j\} \in \sigma_j([n] - (\{x\} \sqcup S))$ , where  $1 \leq j \leq k-1$ . Clearly  $L_{\{x\} \sqcup S, V_x \otimes P_S}(x_1, \dots, x_j) \subseteq L_{S, P_S}(x_1, \dots, x_j)$ . If this inclusion is an equality we add no  $k$ -subsets to  $\Sigma$ . However, if the inclusion is strict then we add all the subsets

$s \in \sigma_k(\{x_1, \dots, x_j\} \sqcup \{x\} \sqcup S)$  to  $\Sigma$ . Then for any states  $|v_{x_1, \dots, x_j}\rangle \in L_{S, P_S}(x_1, \dots, x_j)$ ,  $|v_x\rangle \in V_x$  and  $|\phi_S\rangle \in P_S$  we have:

$$\begin{aligned}
& \sum_{s \in \sigma_k(\{x_1, \dots, x_j\} \sqcup \{x\} \sqcup S) - \Sigma} \Pi_s(|v_{x_1, \dots, x_j}\rangle \otimes |v_x\rangle \otimes |\phi_S\rangle) \\
= & \sum_{s \in \sigma_k(\{x_1, \dots, x_j\} \sqcup S)} \Pi_s(|v_{x_1, \dots, x_j}\rangle \otimes |v_x\rangle \otimes |\phi_S\rangle) \\
& + \sum_{s \in \sigma_k(\{x_1, \dots, x_j\} \sqcup \{x\} \sqcup S)} \Pi_s(|v_{x_1, \dots, x_j}\rangle \otimes |v_x\rangle \otimes |\phi_S\rangle) \\
& - \sum_{s \in \sigma_k(\{x_1, \dots, x_j\} \sqcup \{x\} \sqcup S)} \Pi_s(|v_{x_1, \dots, x_j}\rangle \otimes |v_x\rangle \otimes |\phi_S\rangle) \\
= & 0.
\end{aligned}$$

Here the second equality is by definition of  $L_{S, P_S}(x_1, \dots, x_j)$ . After adding these subsets to  $\Sigma$ , we therefore have an equality  $L_{\{x\} \sqcup S, V_x \otimes P_S}^\Sigma(x_1, \dots, x_j) = L_{S, P_S}(x_1, \dots, x_j)$ . After doing this for all the  $\{x_1, \dots, x_j\}$ , we enlarge the assignment to  $(\{x\} \sqcup S, V_x \otimes P_S)$ . We then repeat this process until we have extended to an assignment on  $X_{nb} \sqcup S$ .

Importantly, we claim that each qubit in  $X_{nb}$  remains not bad throughout this process, even as the assignment is extended. Indeed, it is clear that none of these qubits will ever conflict, since we add subsets to  $\Sigma$  so that  $L^\Sigma(x)$  is fixed throughout. It remains to show that none of these qubits will become heavy. To see this, suppose that we extend the assignment to  $(\{y_1, \dots, y_m\} \sqcup S, V_{y_1} \otimes \dots \otimes V_{y_m} \otimes P_S)$ , removing the relevant subsets as above, and then extend to  $x \in (X_{nb} - (\{y_1, \dots, y_m\} \sqcup S))$  with subspace  $V_x \in L_{\{y_1, \dots, y_m\} \sqcup S, V_{y_1} \otimes \dots \otimes V_{y_m} \otimes P_S}^\Sigma(x) = L_{S, P_S}(x)$  (the equality is due to the addition of subsets to  $\Sigma$  detailed above). Alternatively, we could have extended to  $x$  directly from  $(S, P_S)$ , without extending through  $\{y_1, \dots, y_m\}$  first. It is sufficient to show that, for any  $\{x_1, \dots, x_j\} \in [n] - (\{x, y_1, \dots, y_m\} \sqcup S)$ , there is an inclusion:

$$L_{\{x\} \sqcup S, V_x \otimes P_S}(x_1, \dots, x_j) \subseteq L_{\{x, y_1, \dots, y_m\} \sqcup S, V_x \otimes V_{y_1} \otimes \dots \otimes V_{y_m} \otimes P_S}(x_1, \dots, x_j). \quad (24)$$

Here the subspaces are taken prior to the removal of subsets following the extension to qubit  $x$ ; but on the RHS we have removed all the relevant subsets following extension to  $\{y_1, \dots, y_m\}$ .

We will now prove the inclusion. Let  $|v_{x_1, \dots, x_j}\rangle \in L_{\{x\} \sqcup S, V_x \otimes P_S}(x_1, \dots, x_j)$ . We will consider two cases. The first is where  $j = k - 1$ . In the following equations  $\Sigma$  is defined after extension to  $\{y_1, \dots, y_m\}$ . For all  $|v_x\rangle \in V_x$ ,  $|v_{y_i}\rangle \in V_{y_i}$  and  $|\phi_S\rangle \in P_S$ , we have:

$$\begin{aligned}
& \sum_{s \in \{x_1, \dots, x_{k-1}\} \sqcup \{x\} \sqcup \{y_1, \dots, y_m\} \sqcup S - \Sigma} \Pi_s(|v_{x_1, \dots, x_{k-1}}\rangle \otimes |v_x\rangle \otimes |v_{y_1}\rangle \otimes \dots \otimes |v_{y_m}\rangle \otimes |\phi_S\rangle) \\
= & \sum_{s \in \{x_1, \dots, x_{k-1}\} \sqcup \{y_1, \dots, y_m\} \sqcup S - \Sigma} \Pi_s(|v_{x_1, \dots, x_{k-1}}\rangle \otimes |v_x\rangle \otimes |v_{y_1}\rangle \otimes \dots \otimes |v_{y_m}\rangle \otimes |\phi_S\rangle) \\
& + \Pi_{\{x_1, \dots, x_{k-1}, x\}}(|v_{x_1, \dots, x_{k-1}}\rangle \otimes |v_x\rangle \otimes |v_{y_1}\rangle \otimes \dots \otimes |v_{y_m}\rangle \otimes |\phi_S\rangle) \\
= & 0.
\end{aligned}$$

Here the first term on the second line is zero since  $\Sigma$  is defined so that

$$L_{\{y_1, \dots, y_m\} \sqcup S, V_{y_1} \otimes \dots \otimes V_{y_m} \otimes P_S}^\Sigma(x_1, \dots, x_{k-1}) = L_{S, P_S}(x_1, \dots, x_{k-1});$$

the second term is zero by definition of  $L_{\{x\} \sqcup S, V_x \otimes P_S}(x_1, \dots, x_{k-1})$ . Therefore  $|v_{x_1, \dots, x_j}\rangle \in L_{\{x, y_1, \dots, y_m\} \sqcup S, V_x \otimes V_{y_1} \otimes \dots \otimes V_{y_m} \otimes P_S}(x_1, \dots, x_j)$  and we obtain the inclusion (24).

On the other hand, suppose that  $j < k - 1$ . Then for all  $|v_x\rangle \in V_x$ ,  $|v_{y_i}\rangle \in V_{y_i}$  and  $|\phi_S\rangle \in P_S$  we have the following equation:

$$\sum_{s \in \{x_1, \dots, x_{k-1}\} \sqcup \{x\} \sqcup \{y_1, \dots, y_m\} \sqcup S - \Sigma} \Pi_s(|v_{x_1, \dots, x_{k-1}}\rangle \otimes |v_x\rangle \otimes |v_{y_1}\rangle \otimes \dots \otimes |v_{y_m}\rangle \otimes |\phi_S\rangle)$$

$$\begin{aligned}
&= \sum_{s \in \{\underline{x_1, \dots, x_{k-1}}\} \sqcup \{y_1, \dots, y_m\} \sqcup S - \Sigma} \Pi_s(|v_{x_1, \dots, x_{k-1}}\rangle \otimes |v_x\rangle \otimes |v_{y_1}\rangle \otimes \dots \otimes |v_{y_m}\rangle \otimes |\phi_S\rangle) \\
&+ \sum_{s \in \{\underline{x_1, \dots, x_{k-1}, x}\} \sqcup \{y_1, \dots, y_m\} \sqcup S - \Sigma} \Pi_s(|v_{x_1, \dots, x_{k-1}}\rangle \otimes |v_x\rangle \otimes |v_{y_1}\rangle \otimes \dots \otimes |v_{y_m}\rangle \otimes |\phi_S\rangle) \\
&= 0.
\end{aligned}$$

Here the terms in the second line are both zero since  $\Sigma$  is defined so that:

$$\begin{aligned}
L_{\{y_1, \dots, y_m\} \sqcup S, V_{y_1} \otimes \dots \otimes V_{y_m} \otimes P_S}(x_1, \dots, x_{k-1}) &= L_{S, P_S}(x_1, \dots, x_{k-1}), \\
L_{\{y_1, \dots, y_m\} \sqcup S, V_{y_1} \otimes \dots \otimes V_{y_m} \otimes P_S}(x_1, \dots, x_{k-1}, x) &= L_{S, P_S}(x_1, \dots, x_{k-1}, x).
\end{aligned}$$

Therefore  $|v_{x_1, \dots, x_j}\rangle \in L_{\{x, y_1, \dots, y_m\} \sqcup S, V_x \otimes V_{y_1} \otimes \dots \otimes V_{y_m} \otimes P_S}(x_1, \dots, x_j)$  and we obtain the inclusion (24). We have shown that a qubit in  $X_{nb}$  cannot become bad upon extension.

The first step ends when we have extended to an assignment on  $V_{nb} \sqcup S$ . In the second step we add to  $\Sigma$  all subsets containing any qubit in  $[n] - (V_{nb} \sqcup S)$  and then extend the assignment to the whole of  $[n]$ . We thus obtain a product state  $|\phi\rangle \in (\mathbb{C}^2)^{\otimes n}$  satisfying (2).

To finish we must show that we added less than  $\epsilon n^k$  subsets to  $\Sigma$ . In the first step, after extending to a qubit  $x$  we added all the subsets  $s \in \sigma_k(\{\underline{x_1, \dots, x_j}\} \sqcup \underline{x} \sqcup S)$  each time that  $|L_{\{x\} \sqcup S, V_x \otimes P_S}(x_1, \dots, x_j)|$  was smaller than  $|L_{S, P_S}(x_1, \dots, x_j)|$ . This is at most  $\binom{n}{k-j-1} \leq n^{k-j-1}$  subsets each time, since  $|S| \leq n$ . Since the qubit  $x$  was never heavy, we therefore added at most  $\sum_{j=1}^{k-1} \delta_{S, P_S, x, V_x, j} n^{k-j-1} = \delta_{S, P_S, x, V_x} \leq \epsilon n^{k-1}/5$  projectors every time we extended. As there are at most  $n$  qubits in  $X_{nb}$ , in the first step we added less than  $\epsilon n^k/5$  subsets to  $\Sigma$ .

In the second step, we added all the subsets which contained a qubit not in  $S \sqcup V_{nb}$ . Since by assumption there were  $\leq \epsilon n/5$  bad qubits to begin with, we therefore added at most  $(\epsilon n/5) \binom{n}{k-1} \leq \epsilon n^k/5$  subsets in the second step. Altogether we added less than  $\epsilon n^k$  subsets, as claimed.  $\square$

For discretisation we will also need the following lemma. For a state  $|\phi\rangle \in \mathbb{C}^2$ , we write  $\langle|\phi\rangle\rangle \in \mathbb{P}^1$  for the corresponding point on the complex projective line (a.k.a. the Bloch sphere).

**Lemma 3.6.** *Let  $L \subseteq (\mathbb{C}^2)^{\otimes m}$  be a subspace. We define*

$$X := \{x \in \mathbb{P}^1 : \exists \{|\phi_i\rangle \in \mathbb{C}^2\}_{i=1}^m \text{ s.t. } |\phi_1\rangle |\phi_2\rangle \dots |\phi_m\rangle \in L \text{ and } \langle|\phi_1\rangle\rangle = x\}.$$

*Then either (i)  $X = \mathbb{P}^1$ , or (ii)  $\#(X) \leq m!$ . The bound in case (ii) is tight.*

*Proof.* It is straightforward to show that  $X$  is either finite or covers the whole space. Indeed, the subspace  $L$  corresponds to a projective linear subspace  $\mathbb{P}(L) \subseteq \mathbb{P}^{2^m-1}$ . By the Segre embedding the product states form a projective subvariety  $\Sigma_m := \mathbb{P}^1 \times \dots \times \mathbb{P}^1 \subseteq \mathbb{P}^{2^m-1}$ . But it is an elementary result in algebraic geometry that the projection map  $\pi_1 : \mathbb{P}^1 \times \dots \times \mathbb{P}^1 \rightarrow \mathbb{P}^1$  is closed in the Zariski topology, since every projective variety is complete [Har13, §2 Thm. 4.9]. The intersection  $\mathbb{P}(L) \cap \Sigma_m$  is an algebraic set in  $\Sigma_m$ , and therefore by closure of the projection map, the image under the projection will be an algebraic set in  $\mathbb{P}^1$ . This set will either be zero-dimensional, in which case it is a finite set of points; or one-dimensional, in which case it must be the whole of  $\mathbb{P}^1$ .

The only thing left is to bound the number of points in the case where the image is zero-dimensional. We first note that for any variety  $S \subset \Sigma_m$ , the image  $\pi_1(S)$  is irreducible; indeed, were it not irreducible then by continuity of  $\pi_1$  the fibres would give a nontrivial decomposition of  $S$ . If  $\pi_1(S) \neq \mathbb{P}^1$ , it must therefore be a single point. It follows that all we need to do is bound the number of irreducible components of  $\mathbb{P}(L) \cap \Sigma_m$ . We know that  $\mathbb{P}(L)$  is an intersection of hyperplanes  $\{P_i\}_{i=1}^{2^m-|L|}$ . We use [Har13, §1 Thm. 7.7], which yields the following statement in our special case. For

any subvariety  $V \subset \mathbb{P}^{2^m-1}$  of dimension  $\geq 1$ , and for any hyperplane  $P \subset \mathbb{P}^{2^m-1}$  not containing  $V$ , let  $Z_1, \dots, Z_s$  be the irreducible components of  $V \cap P$ ; then:

$$\sum_{j=1}^s \deg(Z_j) \leq \deg(V). \quad (25)$$

Here  $\deg(Z_j), \deg(V) \in \mathbb{N}_{>0}$  are the degrees of the varieties in  $\mathbb{P}^{2^m-1}$ . If  $P$  contains  $V$  then the intersection is just  $V$  again. We observe that the irreducible components of  $\mathbb{P}(L) \cap \Sigma_m$  can be obtained by the following process: take the intersection  $\Sigma_m \cap P_1$ ; then for each of the irreducible components of that intersection take the intersection with  $P_2$ ; then for each of the irreducible components of that intersection take the intersection with  $P_3$ ; etc. Since  $\deg(\Sigma_m) = m!$ , by (25) we finish with at most  $m!$  irreducible components (which would all have degree 1). The bound follows.

To see that the bound is tight, observe that by definition of the degree, the variety  $\Sigma_m$  intersects a generic linear subvariety  $\mathbb{P}(V) \subset \mathbb{P}^{2^m-1}$  of dimension  $2^m - (m+1)$  in precisely  $m!$  points, and generically these points will be mapped to  $m!$  different points in  $\mathbb{P}^1$  by the projection  $\pi_1$ .  $\square$

We can now prove the theorem.

**Definition 3.7.** Let  $(S, P_S)$  be some local assignment. For any  $X \subseteq ([n] - S)$  we define  $\overline{L}_{S, P_S}(X) \subseteq H_X$  to be the space of all states  $|v_X\rangle \in H_X$  such that

$$\sum_{s \in \sigma_k(X \sqcup S)} \Pi_s(|v_X\rangle \otimes |\phi_S\rangle) = 0 \quad \forall |\phi_S\rangle \in P_S. \quad (26)$$

(The difference between this and (23) is that all the subsets of  $X \sqcup S$  are included, not just those containing every element of  $X$ .)

**Theorem** (Restatement of Theorem 1.7). *There exists a constant  $c(k, \epsilon)$  (i.e. not depending on  $n$ ) such that the following holds. Let  $([n], \{\Pi_s\})$  be any instance of quantum  $k$ -SAT which is  $\epsilon$ -far from satisfiable by a product state. Then, for a randomly chosen subset  $C \in \sigma_{c(k, \epsilon)}([n])$ , the instance is locally unsatisfiable by a product state at  $C$  with probability  $p > 0.75$ .*

*Proof of Theorem 1.7.* The idea of the proof is as follows. We will build a randomly chosen subset  $C \subseteq [n]$  of qubits by picking qubits at random from  $[n]$ , one by one. For any local assignment on some subset of the qubits which have already been picked, we know by Lemma 3.5 that the next qubit we pick has an  $\epsilon/5$  chance of being bad w.r.t. that assignment. By varying the local assignment we consider before picking each qubit, we will show that, after enough qubits have been picked, there can be no local assignment to all the qubits in  $C$  with high probability. This is essentially a backtracking argument; we know by Lemma 3.4 that we cannot extend a chain through more than  $\gamma(k, \epsilon)$  heavy variables, so we simply build such a maximal chain and gradually eliminate all the possible assignments to that chain. We have not endeavoured to optimise this procedure and it can certainly be improved, although whether one can bring the constant  $c(k, \epsilon)$  down to the level in the classical setting [AS03, Thm. 1.1] using such a backtracking approach is by no means clear.

The key to our backtracking argument is the ability to eliminate possible assignments to a chain. We formalise this as follows, assuming for the time being that we can pick bad qubits each time; of course this will not be the case when we pick qubits at random, but in the end we will simply ignore those we pick that are not bad. Suppose that, given any chain  $(\{y_1, \dots, y_m\}, P_1 \otimes \dots \otimes P_m)$  of length  $m$ , there exists an *elimination procedure*, which is defined by a rule in the following way. The procedure will define a set  $\Gamma \subseteq ([n] - \{y_1, \dots, y_m\})$  of picked qubits; at the beginning of the procedure  $\Gamma = \emptyset$ . At each step of the procedure, we:

- Choose some local assignment  $(S, P_S)$  on a subset  $S \subset \{y_1, \dots, y_m\} \sqcup \Gamma$ , according to the rule.



- Pick at random a qubit  $x \in ([n] - (\{y_1, \dots, y_m\} \sqcup \Gamma))$  which is bad w.r.t. the assignment  $(S, P_S)$ .
- Add the qubit  $x$  to  $\Gamma$ .

The rule defines an elimination procedure with constant  $\phi(m) \in \mathbb{N}$  if, while  $|\Gamma| \leq \phi(m)$ , we can stop the procedure and conclude that there is no local assignment to  $\{y_1, \dots, y_m\} \sqcup \Gamma$  which assigns  $P_1 \otimes \dots \otimes P_{m-1}$  to the qubits  $\{y_1, \dots, y_{m-1}\}$ .

We claim that if an elimination procedure exists for chains of length  $m$  with constant  $\phi(m)$ , then there also exists an elimination procedure for chains of length  $m-1$  with constant  $\phi(m-1) = ((\phi(m) + 2)! + 1)(\phi(m) + 1)$ . The procedure is defined as follows. Let  $(\{y_1, \dots, y_{m-1}\}, P_1 \otimes \dots \otimes P_{m-1})$  be a chain of length  $m-1$ , and let  $|p_{m-1}\rangle \in P_{m-1}$  be any state. We pick at random a bad qubit  $y_m^0 \in ([n] - \{y_1, \dots, y_{m-1}\})$  with respect to the local assignment  $(\{y_1, \dots, y_{m-1}\}, P_1 \otimes \dots \otimes P_{m-2} \otimes |p_{m-1}\rangle)$  and add it to  $\Gamma$ . If it is conflicting, then the procedure is complete. If it is heavy, then we perform the length  $m$  elimination procedure on the chain  $(\{y_1, \dots, y_{m-1}, y_m^0\}, P_1 \otimes \dots \otimes P_{m-2} \otimes |p_{m-1}\rangle \otimes \bar{L}_{\{y_1, \dots, y_{m-1}\}, P_1 \otimes \dots \otimes P_{m-2} \otimes |p_{m-1}\rangle}(y_m^0))$ . This gives us a set  $\Gamma_m^0 \subseteq ([n] - \{y_1, \dots, y_m^0\})$  of qubits, where  $|\Gamma_m^0| \leq \phi(m)$ , such that there is no local assignment to  $\{y_1, \dots, y_m^0\} \sqcup \Gamma_m^0$  which assigns  $P_1 \otimes \dots \otimes P_{m-2} \otimes |p_{m-1}\rangle$  to the qubits  $\{y_1, \dots, y_{m-1}\}$ . We add the qubits in  $\Gamma_m^0$  to  $\Gamma$ . Now consider the following subspace:

$$L := \bar{L}_{\{y_1, \dots, y_{m-2}\}, P_1 \otimes \dots \otimes P_{m-2}}(y_{m-1}, y_m^0, \Gamma_m^0) \subseteq H_{y_{m-1}} \otimes H_{y_m^0} \otimes H_{\Gamma_m^0} \cong \mathbb{C}^2 \otimes \mathbb{C}^{2^{|\Gamma_m^0|+1}}.$$

Let  $X = \{|\phi_0\rangle\} \in \mathbb{P}^1 : |\phi_0\rangle \otimes |\phi_1\rangle \otimes \dots \otimes |\phi_{|\Gamma_m^0|+1}\rangle \in L$ . Since there is no local assignment to  $\{y_1, \dots, y_m^0\} \sqcup \Gamma_m^0$  which assigns  $P_1 \otimes \dots \otimes P_{m-2} \otimes |p_{m-1}\rangle$  to the qubits  $\{y_1, \dots, y_{m-1}\}$ , we know that  $X \neq \mathbb{P}^1$ , since  $|\phi_{m-1}\rangle \notin X$ . But then, by Lemma 3.6, we have  $\#(X) \leq (|\Gamma_m^0| + 2)!$ . The worst case is where this is an equality: in this case let the corresponding states be  $\{|p_{m-1}^i\rangle \in H_{y_{m-1}}\}_{i=1}^{(|\Gamma_m^0|+2)!}$ . Now for each  $1 \leq i \leq (|\Gamma_m^0| + 2)!$  in turn we pick at random a qubit  $y_m^i \in ([n] - \Gamma)$  which is bad with respect to the local assignment  $(\{y_1, \dots, y_{m-1}\}, P_1 \otimes \dots \otimes P_{m-2} \otimes |p_{m-1}^i\rangle)$  and add this qubit to  $\Gamma$ . These qubits  $\{y_m^i\}_{i=1}^{(|\Gamma_m^0|+2)!}$  are either conflicting or heavy with respect to their corresponding local assignments; in the worst case they are all heavy. In this case, we consider each of the following chains in turn:

$$(\{y_1, \dots, y_{m-1}, y_m\}, P_1 \otimes \dots \otimes P_{m-2} \otimes |p_{m-1}^i\rangle) \otimes \bar{L}_{\{y_1, \dots, y_{m-1}\}, P_1 \otimes \dots \otimes P_{m-2} \otimes |p_{m-1}^i\rangle}(y_m^i).$$

For each of these chains in turn we perform the length  $m$  elimination procedure, thereby obtaining a set  $\Gamma_m^i \subseteq ([n] - \Gamma)$  of qubits, where  $|\Gamma_m^i| \leq \phi(m)$ , which we add to  $\Gamma$ . By definition of the length  $m$  elimination procedure, for each  $\Gamma_i$  there is no local assignment to  $\{y_1, \dots, y_m^i\} \sqcup \Gamma_m^i$  which assigns  $P_1 \otimes \dots \otimes P_{m-2} \otimes |p_{m-1}^i\rangle$  to the qubits  $\{y_1, \dots, y_{m-1}\}$ . We now stop the procedure and conclude that there is no local assignment to  $\{y_1, \dots, y_{m-1}\} \sqcup \Gamma$  which assigns  $P_1 \otimes \dots \otimes P_{m-2}$  to the qubits  $\{y_1, \dots, y_{m-2}\}$ . Since  $\Gamma = \bigsqcup_{i=0}^{(|\Gamma_m^0|+2)!} (\Gamma_m^i \sqcup \{y_m^i\})$ , we see that  $|\Gamma| \leq ((\phi(m) + 2)! + 1)(\phi(m) + 1) = \phi(m-1)$ .

We now observe that an elimination procedure exists for chains of length  $\gamma := \gamma(k, \epsilon)$ , with constant  $\phi(\gamma) = 1$ . Indeed, for any such chain  $(\{y_1, \dots, y_\gamma\}, P_1 \otimes \dots \otimes P_\gamma)$ , as soon as we add any qubit to  $\Gamma$  we can conclude by Lemma 3.4 that there is no local assignment to  $\{y_1, \dots, y_\gamma\} \sqcup \Gamma$  which assigns  $P_1 \otimes \dots \otimes P_{\gamma-1}$  to the qubits  $\{y_1, \dots, y_{\gamma-1}\}$ . It follows that an elimination procedure exists for every  $1 \leq m \leq \gamma$ , with constant  $\phi(m)$  defined by the recurrence relation  $\phi(m-1) = ((\phi(m) + 2)! + 1)(\phi(m) + 1)$ .

Finally, we observe that this gives rise to an elimination procedure for the empty assignment  $(\emptyset, \emptyset)$ , with constant  $\phi(0) = \phi(1) + 1$ . This procedure is as follows: pick a qubit  $x \in [n]$  which is bad w.r.t.  $(\emptyset, \emptyset)$  and add it to  $\Gamma$ . If it conflicts then there is no local assignment on  $\Gamma = \{x\}$ . If it is heavy, then use the length 1 elimination procedure for the chain  $(\{x\}, L_{\emptyset, \emptyset}(x))$  to obtain a set  $\Gamma_1 \subseteq ([n] - \{x\})$ , where  $|\Gamma_1| \leq \phi(1)$ , such that there is no local assignment on  $\{x\} \sqcup \Gamma_1$ .

The proof of the theorem now reduces to showing that if we pick qubits  $C = \{x_1, \dots, x_c\} \subseteq [n]$  at random, one at a time, then they will implement the elimination procedure for the empty assignment with high probability. Indeed, we recall from

Lemma 3.5 that, with respect to any local assignment  $(S, P_S)$ , there are more than  $\epsilon n/5$  bad qubits in  $([n] - S)$ . For large enough  $n$ , every time we pick a new qubit there is therefore a roughly  $\epsilon/5$  chance of it being bad w.r.t. any local assignment. (The actual value will generally be slightly lower than  $\epsilon/5$  because we cannot pick any of the bad qubits that lie in  $([n] - S) \cap \tilde{C}$ , where  $\tilde{C}$  is the set of qubits which have already been picked; this is why we need  $n$  to be large enough.) The elimination procedure for the empty assignment requires us to pick a bad qubit w.r.t. some local assignment  $\phi(0)$  times. Using the Chernoff bound for the binomial distribution, we find that the probability that we have not picked  $\phi(0)$  bad qubits after  $c(k, \epsilon) := 5\phi(0)/\epsilon + 1$  qubits have been picked is less than 0.25.  $\square$

## References

- [AdGS21] Marco Aldi, Niel de Beaudrap, Sevag Gharibian, and Seyran Saeedi. On efficiently solvable cases of quantum k-SAT. *Communications in Mathematical Physics*, 381(1):209–256, January 2021. [arXiv:1712.09617](#), doi:10.1007/s00220-020-03843-9.
- [AKS12] Andris Ambainis, Julia Kempe, and Or Sattath. A quantum Lovász local lemma. *Journal of the ACM*, 59(5):1–24, 2012. [arXiv:0911.1696](#), doi:10.1145/2371656.2371659.
- [AS03] Noga Alon and Asaf Shapira. Testing satisfiability. *Journal of Algorithms*, 47(2):87–103, 2003. URL: <https://www.tau.ac.il/~nogaa/PDFS/asafsodaproc2.pdf>, doi:10.1016/S0196-6774(03)00019-1.
- [ASSZ18] Itai Arad, Miklos Santha, Aarthi Sundaram, and Shengyu Zhang. Linear-time algorithm for quantum 2SAT. *Theory of Computing*, 14(1):1–27, 2018. [arXiv:1508.06340](#), doi:10.4086/toc.2018.v014a001.
- [ATL11] R Augusiak, J Tura, and M Lewenstein. A note on the optimality of decomposable entanglement witnesses and completely entangled subspaces. *Journal of Physics A: Mathematical and Theoretical*, 44(21):212001, April 2011. [arXiv:1012.3786](#), doi:10.1088/1751-8113/44/21/212001.
- [BCY11] Fernando G.S.L. Brandao, Matthias Christandl, and Jon Yard. Faithful squashed entanglement. *Communications in Mathematical Physics*, 306(3):805–830, 2011. [arXiv:1010.1750](#), doi:10.1007/s00220-011-1302-1.
- [BE15] Béla Bollobás and Tom Eccles. Partial shadows of set systems. *Combinatorics, Probability and Computing*, 24(5):825–828, 2015. doi:10.1017/S0963548314000790.
- [BFS15] Magali Bardet, Jean-Charles Faugere, and Bruno Salvy. On the complexity of the F5 Gröbner basis algorithm. *Journal of Symbolic Computation*, 70:49–70, 2015. [arXiv:1312.1655](#), doi:10.1016/j.jsc.2014.09.025.
- [BH13] Fernando G.S.L. Brandao and Aram W. Harrow. Quantum de Finetti theorems under local measurements with applications. In *Proceedings of the Forty-Fifth Annual ACM Symposium on Theory of Computing*, STOC '13, page 861–870, New York, NY, USA, 2013. Association for Computing Machinery. [arXiv:1210.6367](#), doi:10.1145/2488608.2488718.
- [BH16] Fernando G.S.L. Brandao and Aram W Harrow. Product-state approximations to quantum states. *Communications in Mathematical Physics*, 342(1):47–80, 2016. [arXiv:1310.0017](#), doi:10.1007/s00220-016-2575-1.

- [BMR10] Sergey Bravyi, Cristopher Moore, and Alexander Russell. Bounds on the quantum satisfiability threshold. In Andrew Chi-Chih Yao, editor, *Innovations in Computer Science 2010*, pages 482–489. Tsinghua University Press, 2010. [arXiv:0907.1297](#).
- [Bra11] Sergey Bravyi. Efficient algorithm for a quantum analogue of 2-SAT. In K. Mahdavi, D. Koslover, and L.L. Brown, editors, *Cross Disciplinary Advances in Quantum Computing*, volume 536 of *Contemporary Mathematics*. American Mathematical Society, 2011. [arXiv:quant-ph/0602108](#).
- [CCD<sup>+</sup>11] Jianxin Chen, Xie Chen, Runyao Duan, Zhengfeng Ji, and Bei Zeng. No-go theorem for one-way quantum computing on naturally occurring two-level systems. *Phys. Rev. A*, 83:050301, May 2011. [arXiv:1004.3787](#), doi:10.1103/PhysRevA.83.050301.
- [CW04] Matthias Christandl and Andreas Winter. Squashed entanglement: an additive entanglement measure. *Journal of Mathematical Physics*, 45(3):829–840, 2004. [arXiv:quant-ph/0308088](#), doi:10.1063/1.1643788.
- [dBG16] Niel de Beaudrap and Sevag Gharibian. A Linear Time Algorithm for Quantum 2-SAT. In Ran Raz, editor, *31st Conference on Computational Complexity (CCC 2016)*, volume 50 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 27:1–27:21, Dagstuhl, Germany, 2016. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik. [arXiv:1508.07338](#), doi:10.4230/LIPIcs.CCC.2016.27.
- [Fit18] Matthew Fitch. Kruskal-Katona type problem. 2018. [arXiv:1805.00340](#).
- [Har13] R. Hartshorne. *Algebraic Geometry*. Graduate Texts in Mathematics. Springer New York, 2013.
- [HLL<sup>+</sup>13] B. Hsu, C. R. Laumann, A. M. Läuchli, R. Moessner, and S. L. Sondhi. Approximating random quantum optimization problems. *Phys. Rev. A*, 87:062334, Jun 2013. [arXiv:1304.2837](#), doi:10.1103/PhysRevA.87.062334.
- [Kee08] Peter Keevash. Shadows and intersections: Stability and new proofs. *Advances in Mathematics*, 218(5):1685–1703, 2008. [arXiv:0806.2023](#), doi:<https://doi.org/10.1016/j.aim.2008.03.023>.
- [KW04] Masato Koashi and Andreas Winter. Monogamy of quantum entanglement and other correlations. *Physical Review A*, 69(2):022309, 2004. [arXiv:quant-ph/0310037](#), doi:10.1103/PhysRevA.69.022309.
- [LLM<sup>+</sup>10] Christopher R Laumann, AM Läuchli, R Moessner, A Scardicchio, and Shivaji Lal Sondhi. Product, generic, and random generic quantum satisfiability. *Physical Review A*, 81(6):062345, 2010. [arXiv:0910.2058](#), doi:10.1103/PhysRevA.81.062345.
- [LMSS10] C. R. Laumann, R. Moessner, A. Scardicchio, and S. L. Sondhi. Random quantum satisfiability. *Quantum Information and Computation*, 10(1):1–15, Jan 2010.
- [VZGG13] Joachim Von Zur Gathen and Jürgen Gerhard. *Modern Computer Algebra*. Cambridge University Press, Cambridge, 2013.

## A Proofs of combinatorial lemmas

**Lemma** (Restatement of Lemma 2.1). *Let  $\theta, x \in \mathbb{R}_{\geq 0}$  and  $c \in \mathbb{N}$ , where  $0 \leq \theta \leq 1$  and  $x \geq c$ . Then, for some  $\alpha \in [0, 1)$ , we have:*

$$\theta \binom{x}{c} = \binom{\theta^{1/c} x}{c} + \alpha \binom{\theta^{1/c} x}{c-1} < \binom{\theta^{1/c} x + 1}{c}. \quad (27)$$

*Proof.* Setting  $m = x - l$  for  $l < x$ , we have:

$$\frac{\theta \binom{x}{c} - \binom{m}{c}}{\binom{m}{c-1}} = \frac{\theta x \cdots (x - c + 1)}{cm \cdots (m - c + 2)} - \frac{m - c + 1}{c}.$$

Now, if we set  $m = \theta^{1/c} x$ , we obtain:

$$\begin{aligned} \alpha &:= \frac{\theta \binom{x}{c} - \binom{\theta^{1/c} x}{c}}{\binom{\theta^{1/c} x}{c-1}} = \frac{1}{c} \left( \frac{\theta x \cdots (x - c + 1)}{\theta^{1/c} x \cdots (\theta^{1/c} x - c + 2)} - (\theta^{1/c} x - c + 1) \right) \\ &= \frac{1}{c} \left( \frac{\theta^{1/c} x \cdots (x - c + 1)}{x(x - \theta^{-1/c}) \cdots (x - \theta^{-1/c}(c - 2))} - (\theta^{1/c} x - c + 1) \right) \\ &\leq \frac{1}{c} (\theta^{1/c} (x - c + 1) - (\theta^{1/c} x - c + 1)) \\ &= \frac{(c - 1)(1 - \theta^{1/c})}{c} < 1. \end{aligned}$$

The inequality then follows from the standard binomial recurrence relation  $\binom{x}{c} + \binom{x}{c-1} = \binom{x+1}{c}$ .  $\square$

**Lemma** (Restatement of Lemma 2.4). *If  $G$  is an  $l$ -regular hypergraph with edge density  $\theta$ , then the  $k$ -shadow  $G|_k$  has edge density  $\theta|_k \geq (\theta^{1/l} - 2/n)^k$ .*

*Proof.* The number of edges in  $G|_k$  is the minimum number of  $k$ -edges in a  $k$ -regular hypergraph such that all the edges in  $G$  appear as complete  $l$ -vertex subgraphs of that graph; for a given number of  $k$ -edges we may therefore ask what is the optimal arrangement maximising the number of complete  $l$ -vertex subgraphs. It follows from [Kee08, Thm. 1] that a complete  $k$ -regular hypergraph on a subset  $S \subseteq [n]$  is an optimal arrangement. It then follows from Lemma 2.1 that  $G|_k$  has at least  $\binom{\lfloor \theta^{1/l} n \rfloor}{k}$  edges. Observing that  $\lfloor \theta^{1/l} n \rfloor \geq \theta^{1/l} n - 1 = (\theta^{1/l} - 2/n)n + 1$  and using the inequality in (5) gives the result.  $\square$

**Lemma** (Restatement of Lemma 2.5). *Let  $n, c \in \mathbb{N}$  be such that  $l := n/2^c + \delta$  and  $k := l/2 + \epsilon$ , where  $-3 \leq \delta < -1$  and  $-1/2 \leq \epsilon \leq 0$ , are natural numbers. Let  $G$  be an  $l$ -regular hypergraph on  $n$  vertices with edge density  $\theta \in [0, 1]$ .*

*Suppose that  $n \geq 3(2^c)$  and  $n \geq 3(2^{c+1}) \left( \frac{1}{2} + \frac{1}{\ln(A)} \left( \frac{A^{1/3}}{4} - \ln(\theta) \right) \right)$  for some constant  $A > 1$ . Then the edge density  $\theta|_{k, 1/2}$  of a partial  $(k, 1/2)$ -shadow of  $G$  has the following lower bound:*

$$\theta|_{k, 1/2} \geq \frac{\theta}{K_A} := \frac{\theta}{2\sqrt{A \exp(A^{1/3}/2)}}$$

*Proof.* We first fix some notation; let  $\omega = 1/2$ ,  $\alpha = 1/2^c$ , and  $\beta = 1/2^{c+1}$ ; then  $l = (\alpha + \delta/n)n$  and  $k = (\beta + \epsilon/n)n$ . Let  $\theta'$  be the edge density of a  $k$ -regular hypergraph  $H'$ . We will find the largest edge density  $\theta$  of an  $l$ -regular hypergraph  $H$  which has  $H'$  as a  $(k, \omega)$ -shadow. We observe that we can add an  $l$ -edge to  $H$  precisely when at least  $\omega \binom{l}{k}$  of the  $k$ -subsets of that  $l$ -edge are  $k$ -edges of  $H'$ . We thus have the following equation for the maximum number of edges in  $H$ . Let  $S(H') \subseteq [n]$  be the set of vertices which are contained in some  $k$ -edge of  $H'$ . For  $0 \leq i \leq |S(H')|$ , let  $N_i$  be the number of subsets of  $S(H')$  of size  $i$  such that the induced subgraph on that subset has  $\geq \omega \binom{l}{k}$  edges. Then:

$$\theta \binom{n}{l} = \sum_{i=0}^l N_i \binom{n - |S(H')|}{l - i}. \quad (28)$$

We now observe that  $N_i = 0$  for all  $i \neq l$ . Indeed, consider  $N_{l-1}$ . The required density of edges in a subset of  $H'$  of size  $l-1$  which contains  $\omega \binom{l}{k}$  edges is  $\omega \binom{l}{k} / \binom{l-1}{k} = \omega l / (l-k) = \frac{\omega(\alpha+\delta/n)}{\alpha+\delta/n-(\beta+\epsilon/n)}$ , which is greater than 1 for all choices of  $-3 \leq \delta < -1$ ,  $-1/2 \leq \epsilon \leq 0$ . We therefore have  $N_{l-1} = 0$ . The sum (28) therefore reduces to a much simpler equation:

$$\theta = \frac{N_l}{\binom{n}{l}}. \quad (29)$$

We immediately observe that as soon as  $\theta' > \omega$  then we can obtain  $\theta = 1$  by distributing the edges uniformly over the  $n$  vertices, giving  $N_l = \binom{n}{l}$ .

Assuming from now on that  $\theta' < \omega$ , we make some further observations which will enable us to bound  $\theta$ . To maximise  $N_l$  the best distribution of  $k$ -edges is a uniform distribution over as many vertices as possible, with density  $\omega$ . If we multiplied the edge density by  $\frac{1}{\omega}$ , we could form the complete hypergraph on these vertices, and so we see by Lemma 2.1 that the maximum number of vertices is upper bounded by  $\lceil (\theta'/\omega)^{1/k} n \rceil$ . We therefore obtain the following bound:

$$N_l \leq \binom{\lceil (\theta'/\omega)^{1/k} n \rceil}{l} \leq \left( (\theta'/\omega)^{1/k} + \frac{1}{n} \right)^l \binom{n}{l} = \left( (\theta'/\omega)^{\frac{1}{\beta n + \epsilon}} + \frac{1}{n} \right)^{\alpha n + \delta} \binom{n}{l}.$$

We now claim that, if  $n > 3(2^c)$  and furthermore  $n \geq \frac{3(2^{c+1})}{\ln(A)} \ln(1/(2\theta'))$  for some constant  $A > 1$ , then:

$$\frac{N_l}{\binom{n}{l}} \leq A(2\theta')^2 \exp\left(\alpha A^{1/3}\right). \quad (30)$$

Indeed, note that  $0 < \theta' < \omega < 1$ ; thus, since  $n > 3(2^c)$  implies that  $\alpha n + \delta > 0$ :

$$\begin{aligned} \left( (\theta'/\omega)^{\frac{1}{\beta n + \epsilon}} + \frac{1}{n} \right)^{\alpha n + \delta} &\leq \left( (\theta'/\omega)^{\frac{1}{\beta n}} + \frac{1}{n} \right)^{\alpha n + \delta} \\ &= (\theta'/\omega)^{\frac{\alpha n + \delta}{\beta n}} \left( 1 + \frac{1}{n} (\omega/\theta')^{\frac{1}{\beta n}} \right)^{\alpha n + \delta} \\ &\leq (\theta'/\omega)^{\frac{\alpha n - 3}{\beta n}} \left( 1 + \frac{1}{n} (\omega/\theta')^{\frac{1}{\beta n}} \right)^{\alpha n} \\ &\leq (\theta'/\omega)^{\frac{\alpha n - 3}{\beta n}} \exp\left(\alpha (\omega/\theta')^{\frac{1}{\beta n}}\right) \\ &= (\theta'/\omega)^{\frac{\alpha}{\beta}} (\omega/\theta')^{\frac{3}{\beta n}} \exp\left(\alpha (\omega/\theta')^{\frac{1}{\beta n}}\right). \end{aligned}$$

Substituting (30) into (29) and inverting the inequality, we obtain the statement in the theorem.  $\square$

**Lemma** (Restatement of Lemma 2.6). *Let  $n \in \mathbb{N}$  be odd. We construct an  $(n-1)/2$ -regular hypergraph  $G$  as follows. For every subset  $X \in \sigma_{(n-1)/2}([n])$ , we do precisely one of the following:*

1. Add  $X$  as an edge to  $G$ .
2. Add all the subsets in  $\sigma_{(n-1)/2}(\overline{X})$  as edges to  $G$ .

The edge density  $\theta$  of a hypergraph  $G$  constructed in this way satisfies  $\theta \geq 1/2$ .

*Proof.* We call a choice of **1** or **2** for each  $X \in \sigma_{(n-1)/2}([n])$  a *configuration*. For a given configuration, let  $S_1 \subset \sigma_{(n-1)/2}([n])$  be the subsets for which option **1** is chosen, and let  $S_2 \subset \sigma_{(n-1)/2}([n])$  be the subsets for which option **2** is chosen. We observe that there is always a configuration minimising the number of edges in  $G$  (a *minimal configuration*) which satisfies the property that, for every  $X \in S_2$ ,  $\sigma_{(n-1)/2}(\overline{X}) \subset S_1$ . Indeed, suppose this is not the case, and there is some  $Y \in \sigma_{(n-1)/2}(\overline{X}) \subset S_2$ . But then one can change the configuration by moving  $Y$  to  $S_1$  without increasing the number of edges in  $G$ . If we

start with a minimal configuration, by doing this for all  $X \in S_2$ , we obtain a minimal configuration satisfying the desired property.

Let us calculate the minimal number of edges when such a configuration is chosen. Let  $C$  be the  $(n-1)/2$ -regular hypergraph on those vertices of  $G$  contained in an edge of  $S_1$ , whose edges are precisely the subsets in  $S_1$ . Every  $Y \in S_2$  corresponds to a induced complete hypergraph on  $(n+1)/2$  vertices of  $C$ . By Lemma 2.4, the arrangement of  $(n-1)/2$ -edges maximising the number of induced complete hypergraphs is the complete hypergraph on  $C$ . We have  $\bar{Y} \subset C$  for every  $Y \in S_2$ , which implies that  $\bar{C} \subset Y$ . We therefore obtain the following inequality giving a lower bound on the size of  $C$ :

$$\binom{|C|}{(n-1)/2} + \binom{|C|}{(n+1)/2 - (n-|C|)} \geq \binom{n}{(n-1)/2}.$$

Setting  $|C| = n-2$ , we find the inequality is violated; but setting  $|C| = n-1$ , we find the equality is satisfied. The minimal number of edges in  $G$  is therefore  $\binom{n-1}{(n-1)/2} = \frac{n+1}{2n} \binom{n}{(n-1)/2}$ , and the result follows.  $\square$

**Lemma** (Restatement of Lemma 2.7). *Let  $G$  be a  $k$ -regular hypergraph on  $n$  vertices with edge density  $\theta$ , where  $k = \beta n$  for some  $\beta \in [0, 1]$ . Let  $N$  be the degree of a vertex picked uniformly at random, and let  $\tau := N/\binom{n-1}{k-1}$ . Then, for any  $\epsilon > 0$  and  $\alpha \in (0, 1)$ :*

$$\Pr[|\theta - \tau| \geq \min(\Delta(\alpha, \beta, n), (1-\theta)(1-\alpha))] \leq \alpha,$$

where

$$\Delta(\alpha, \beta, n) = \frac{5 \exp\left(\frac{1}{12\alpha(1-\alpha)n} + \frac{1}{12\beta(1-\beta)n}\right)}{\sqrt{2\pi\alpha(1-\alpha)\beta(1-\beta)n}} =: \frac{5(L_{\alpha,\beta})^{1/n}}{M_{\alpha,\beta}\sqrt{n}}.$$

*Proof.* We prove the case where  $\tau < \theta$ ; the case  $\tau > \theta$  follows by considering the complement of the hypergraph. For any hypergraph  $G$  with edge density  $\theta$ , let  $C \subset [n]$  be the subset of vertices whose degree is less than or equal to  $\tilde{\tau}\binom{n-1}{k-1}$ , where  $\tilde{\tau} < \theta$ ; clearly  $\Pr\left(\tilde{N} \leq \tilde{\tau}\binom{n-1}{k-1}\right) = |C|/n$ , where  $\tilde{N}$  is the degree of a vertex of  $G$  picked uniformly at random. Let  $S := \sum_{v \in C} d_v$ , where  $d_v$  is the degree of the vertex  $v \in C$ ; we must have  $S \leq |C|\tilde{\tau}\binom{n-1}{k-1}$ . We will obtain a lower bound for  $\tilde{\tau}$  in terms of  $\theta$  and  $|C|$  by defining a configuration of edges which minimises  $S$ . Since we are interested in how the lower bound changes as  $n$  increases, we will consider  $|C|$  to be some function of  $n$ ; but  $\theta$  will remain constant. The configuration is defined as follows: we first place all the edges contained entirely in  $(n-|C|)$ , which does not increase  $S$ . Then we place all the edges which only contain one vertex in  $C$ , at the cost  $S \mapsto S+1$ . Then we can place all the edges which only contain two vertices of  $C$ , at the cost  $S \mapsto S+2$ , etc. We will stop when we have placed all  $\theta\binom{n}{k}$  edges. Let  $r+1$  be the number of vertices of  $C$  in the final edge we place. The following inequality encodes the fact that we have placed  $\theta\binom{n}{k}$  edges:

$$\begin{aligned} \sum_{i=0}^r \binom{n-|C|}{k-i} \binom{|C|}{i} &\leq \theta \binom{n}{k} \leq \sum_{i=0}^{r+1} \binom{n-|C|}{k-i} \binom{|C|}{i} \\ \Rightarrow P(X_{n-|C|,|C|,k} \leq r) &\leq \theta \leq P(X_{n-|C|,|C|,k} \leq r+1). \end{aligned} \quad (31)$$

Here  $X_{n-|C|,|C|,k}$  is a random variable distributed according to the classical hypergeometric distribution with parameters  $n, |C|, k$ . The following inequality encodes the fact that  $S \leq |C|\tilde{\tau}\binom{n-1}{k-1}$ :

$$\begin{aligned} \sum_{i=0}^r \frac{i}{|C|} \binom{n-|C|}{k-i} \binom{|C|}{i} &\leq \tilde{\tau} \binom{n-1}{k-1} \leq \sum_{i=0}^{r+1} \frac{i}{|C|} \binom{n-|C|}{k-i} \binom{|C|}{i} \\ \Rightarrow P(X_{n-|C|,|C|-1,k-1} \leq r-1) &\leq \tilde{\tau} \leq P(X_{n-|C|,|C|-1,k-1} \leq r). \end{aligned} \quad (32)$$

Here  $X_{n-|C|,|C|-1,k-1}$  is a random variable distributed according to the classical hypergeometric distribution with parameters  $n-1, |C|-1, k-1$ . It follows that:

$$\begin{aligned} P(X_{n-|C|,|C|,k} \leq r) - P(X_{n-|C|,|C|-1,k-1} \leq r) \\ \leq \theta - \tilde{\tau} \\ \leq P(X_{n-|C|,|C|,k} \leq r+1) - P(X_{n-|C|,|C|-1,k-1} \leq r-1). \end{aligned} \quad (33)$$

For some intuition, we recall the standard model for the classical hypergeometric distribution with parameters  $n, |C|, k$ : there are  $n$  balls,  $|C|$  of which are white and  $(n-|C|)$  of which are black, and we pick  $k$  of them at random without replacement; the hypergeometric distribution counts the number of white balls picked. We observe that

$$P(X_{a,b,c} \leq r+1) = \frac{a+b-c}{a+b} P(X_{a,b-1,c} \leq r+1) + \frac{c}{a+b} P(X_{a,b-1,c-1} \leq r). \quad (34)$$

This equation is obtained by conditioning on whether a specific white ball is picked. We further observe:

$$P(X_{a,b,c} \leq r) = P(X_{a,b,c-1} \leq r-1) + \frac{a+r-(c-1)}{a+b-(c-1)} P(X_{a,b,c-1} = r). \quad (35)$$

This equation is obtained by conditioning on how many of the first  $c-1$  balls picked are white. Recall  $k = \beta n$ ; let  $|C| = \alpha n$ . Then, substituting (34) and (35) into (33), we obtain:

$$\begin{aligned} \theta - \tilde{\tau} &= P(X_{n-|C|,|C|-1,k-1} = r) + \frac{n-|C|+r+2-k}{n} P(X_{n-|C|,|C|-1,k-1} = r+1) \\ &\leq P(X_{n-|C|,|C|-1,k-1} = r) + (4-\alpha-\beta) P(X_{n-|C|,|C|-1,k-1} = r+1). \end{aligned}$$

For the inequality we used that  $r < n$ . The mode of  $X_{n-|C|,|C|-1,k-1}$  is  $M := \lfloor \frac{k|C|}{n+1} \rfloor$ , so by Stirling's approximation we have:

$$\begin{aligned} \theta - \tilde{\tau} &\leq 5 \frac{\binom{n-|C|}{k-M} \binom{|C|}{M}}{\binom{n}{k}} \leq \frac{5e^{f(n)}}{\sqrt{2\pi\alpha\beta(1-\alpha)(1-\beta)n}} \\ &\leq \frac{5 \exp\left(\frac{1}{12\alpha(1-\alpha)n} + \frac{1}{12\beta(1-\beta)n}\right)}{\sqrt{2\pi\alpha(1-\alpha)\beta(1-\beta)n}}, \end{aligned} \quad (36)$$

where

$$\begin{aligned} f(n) &= \frac{1}{12\alpha n} + \frac{1}{12\beta n} + \frac{1}{12(1-\alpha)n} + \frac{1}{12(1-\beta)n} - \frac{1}{12n+1} - \frac{1}{12\alpha\beta n+1} \\ &\quad - \frac{1}{12(1-\alpha)\beta n+1} - \frac{1}{12(1-\alpha)(1-\beta)n+1} - \frac{1}{12\alpha(1-\beta)n+1}. \end{aligned}$$

The free variables are  $\alpha$  and  $\tilde{\tau}$ , so we fix  $\alpha$  and determine the minimal  $\tilde{\tau}$ . Since

$$\Pr\left(\tilde{N} \leq \tilde{\tau} \binom{n-1}{k-1}\right) = \frac{|C|}{n} = \alpha,$$

and, by taking the sum of the degrees of all the vertices,

$$\theta \leq (1-\alpha) + \alpha\tilde{\tau} \quad \Rightarrow \quad \theta - \tilde{\tau} \leq (1-\theta)(1-\alpha),$$

we obtain the bound in the statement of the lemma.  $\square$