

UNIQUENESS OF CONICAL SINGULARITIES FOR MEAN CURVATURE FLOWS

TANG-KAI LEE AND XINRUI ZHAO

ABSTRACT. In this paper, we prove the uniqueness of asymptotically conical tangent flows in all codimensions. This is based on an early work of Chodosh-Schulze, who proved the uniqueness in the hypersurface case.

0. Introduction

Mean curvature flow (MCF) is a family of immersions $\mathbf{x}: M^n \times I \rightarrow \mathbb{R}^{n+k}$ such that $\partial_t \mathbf{x} = \mathbf{H}$ where \mathbf{H} is the mean curvature vector of the immersed submanifold $M_t = \mathbf{x}(M, t)$. To understand the structure of a MCF, studying its singularities is the most important problem since they are unavoidable in general. For example, every MCF starting from a closed hypersurface develops a finite-time singularity. The singularities are usually investigated via a blowup technique. This leaves the most fundamental question about the *uniqueness* of these singularity models.

To elaborate, given a MCF M_t with a spacetime singularity at $(\mathbf{0}, 0) \in \mathbb{R}^{n+k} \times \mathbb{R}$, we can define a sequence of MCFs $M_t^i := \lambda_i M_{\lambda_i^{-2}t}$ by rescaling using any sequence $\lambda_i \rightarrow \infty$. Based on Huisken's monotonicity [Hui90] and Brakke's compactness [Bra78], White [Whi94] and Ilmanen [Ilm95] proved that M_t^i weakly converges to a self-similar limiting flow, called a *tangent flow*, after replacing λ_i 's with a subsequence. When the limiting flow is smooth, it is generated by a *self-shrinker*, a submanifold satisfying $\mathbf{H} = -\frac{x^\perp}{2}$. In general, this limiting flow depends on the sequence we choose to rescale the flow. That is to say, choosing different subsequences may result in different tangent flows at the same singularity. Determining the uniqueness of tangent flows is a crucial question if we would like to understand the singularity structure of the flow. See e.g., [Sch14, CM15, CM19, Zhu20, CS21, LSS22]. A similar problem also plays an important role in the study of minimal submanifolds, that is, the uniqueness of tangent cones. For related references, one can see [AA81, Sim83, Har97, Sim08, CM14].

In this paper, we will focus on asymptotically conical self-shrinkers. An *asymptotically conical* self-shrinker is a self-shrinker that is (smoothly) asymptotic to a cone at infinity. Such kinds of self-shrinkers are important in understanding two-dimensional self-shrinkers, see for example [Wan14]. Asymptotically conical self-shrinkers have other interesting properties. For example, given any cone in \mathbb{R}^{n+k} , there is at most one asymptotically conical self-shrinker asymptotic to it even in the case that the codimension is higher than one [Wan14, Kha20]. Also, asymptotically conical self-shrinkers as tangent flows of hypersurface MCFs are unique, proven by Chodosh-Schulze [CS21].

Date: February 3, 2023.

Following their framework and combining with an extension theorem that originated from White's curvature estimate, we can prove a higher-codimensional analogue of Chodosh-Schulze's result.

Theorem 0.1 (Uniqueness of asymptotically conical tangent flows). Let $\Sigma^n \subseteq \mathbb{R}^{n+k}$ be a multiplicity one asymptotically conical self-shrinker and $(M_t^n)_{t \in (-T, 0)}$ a MCF in \mathbb{R}^{n+k} such that a tangent flow at $(\mathbf{0}, 0)$ is the MCF generated by Σ . Then it is the unique tangent flow of $(M_t^n)_{t \in (-T, 0)}$ at $(\mathbf{0}, 0)$.

Uniqueness properties of tangent flows for geometric flows usually follow from a Łojasiewicz-type inequality. For geometric applications, these kinds of inequalities were first proven by Simon [Sim83] by using the Lyapunov-Schmidt reduction and applying the original inequality invented by Łojasiewicz [Loj65] for real analytic functions. As a consequence, Simon used these infinite dimensional Łojasiewicz inequalities, now often called Łojasiewicz-Simon inequalities, to prove the uniqueness of tangent cones with smooth cross sections of minimal surfaces. Simon's framework was later applied to the study of compact MCFs by Schulze [Sch14]. By setting up appropriate function spaces on asymptotically self-shrinking hypersurfaces, Chodosh-Schulze [CS21] successfully obtained Łojasiewicz-Simon inequalities on them. It is worth mentioning that the Łojasiewicz inequalities obtained by Colding-Minicozzi [CM15, CM19] and Zhu [Zhu20] were proved without using Simon's theory, and they were able to get explicit Łojasiewicz inequalities by hand.

The proof of Theorem 0.1 is based on the ideas of Chodosh-Schulze [CS21]. The first main idea is, as mentioned above, to construct some function spaces to replace the ordinary Hölder spaces so that we can establish the desired elliptic estimates, based on which Simon's techniques [Sim83] can be applied. Elements in these new function spaces are entire vector fields with appropriate decay rates. Establishing good mapping properties among this new function spaces leads to the Simon-Łojasiewicz inequality for entire vector fields with sufficiently small norms (in the new function spaces). Nevertheless, in a blowup sequence, the locally smooth convergence only gives us closeness on (probably large) compact parts of the flows. As a result, we need to show that some Łojasiewicz-type inequalities can be derived for vector fields defined only over parts of the self-shrinker. To deal with this localization process, we need to say that the *rough conical scale* (Definition 3.7 (2)) improves whenever the flow can be written as the graph of a (small enough) vector field on a large enough compact set. Based on this, we can extend the graphicality up to the *shrinker scale* (Definition 3.7 (1)). This allows us to obtain the desired localized Łojasiewicz-Simon inequality, leading to the uniqueness of tangent flows.

The main goal of this note is to implement the plan above for higher-codimensional MCFs. It is known that extending results in the hypersurface setting to the higher-codimensional one is not always direct. The main difficulties include the complexity of curvature tensors and the lack of certain maximum principles. In our case, most of the work in [CS21] can be generalized to the higher-codimensional setting by standard extensions. However, there are still some places where we need new arguments, and they are all related to the use of certain maximum principles in the hypersurface case. For example, when proving the extension result in the hypersurface case, Chodosh-Schulze used a barrier principle to show a curvature estimate. This, in general, does not apply to a MCF of submanifolds of higher codimensions. To deal with this, we combine the asymptotic conicality and White's curvature estimate to obtain the desired curvature estimate (cf.

Proposition 4.1). This finally leads to the extension step that is necessary for us to utilize the localized Łojasiewicz-Simon inequality.

This paper is organized as follows. In Section 1 and 2, we give some preliminary notations and results for asymptotically conical self-shrinkers and their normal bundles. In Section 3, we talk about entire Łojasiewicz-Simon inequalities and localized Łojasiewicz-Simon inequalities. In Section 4, the uniqueness of higher-codimensional asymptotically conical tangent flow of MCF is proven.

ACKNOWLEDGEMENT

The authors are very grateful to Prof. Tobias Colding and Prof. William Minicozzi for their constant support and several inspiring conversations. They would also like to thank Prof. Jonathan Zhu for clarifying some results in his paper, thank him, Prof. Otis Chodosh, Prof. Yng-Ing Lee and Prof. Mu-Tao Wang for their interests, and thank Zhihan Wang for pointing out some imprecise statements in an early version. During the project, Lee was partially supported by NSF Grant DMS 2005345 and Zhao was supported by NSF Grant DMS 1812142 and NSF Grant DMS 1811267.

1. Geometric and analytic preliminaries

Most of the properties and definitions in this section are similar to the ones in [CS21]. We mention the parts that we will need or that require some slight modifications in the higher-codimensional case.

1.1. Estimate on ends. Consider an asymptotic conical self-shrinker $\Sigma^n \subseteq \mathbb{R}^{n+k}$ and assume that the asymptotic cone

$$\mathcal{C} = \lim_{t \rightarrow 0} \sqrt{-t} \Sigma$$

is the cone over $\Gamma^{n-1} \subseteq S^{n+k-1}$. The metric on \mathcal{C} is $g_{\mathcal{C}} = dr \otimes dr + r^2 g_{\Gamma}$. Then from the smooth convergence we know the following decay of curvatures (cf. [Wan14, CS21]).

Lemma 1.1. There exist $C > 0$ and $R_1 \geq 0$ such that for $x \in \Sigma \setminus B_{R_1}$ and $i \geq 0$,

$$|\nabla^i A(x)| \leq C|x|^{-i-1}.$$

From this lemma and the self-shrinker equation $\mathbf{H} = \frac{x^\perp}{2}$, we have an improved estimate

$$|\nabla^i x^\perp| \leq C|x|^{-i-1}.$$

We list some other estimates we will use later. Their proofs are similar to those in the hypersurface case.

Lemma 1.2. For $R > 0$ sufficiently large, there is $W \in \Gamma(N(\mathcal{C} \setminus B_R(0)))$ so that

$$\text{graph } W := \{p + W(p) : p \in \mathcal{C} \setminus B_R(0)\} \subseteq \Sigma$$

parametrizes Σ outside a compact set. The section W satisfies

$$(1.3) \quad |W| = O(r^{-1}) \text{ and } |\nabla^{(j)} W| = O(r^{-1-j+\eta})$$

as $r \rightarrow \infty$ for any $\eta > 0$ and $j \geq 1$. Moreover, the radial derivatives satisfy sharper relations $|\nabla_{\partial_r}^j W| = O(r^{-1-j})$.

Corollary 1.4. For large $R > 0$ and by denoting the corresponding parametrization of $\Sigma \setminus B_R(0)$ using \mathcal{C} as F , we have the following improved estimates on the induced metric

$$F^*g_\Sigma = dr \otimes dr + r^2g_\Gamma + h$$

where h is a symmetric $(0, 2)$ -tensor on $\mathcal{C} \setminus B_R(0)$ satisfying $|h| = O(r^{-2})$ and $|\nabla^{(j)}h| = O(r^{-2-j+\eta})$ as $r \rightarrow \infty$, for all $j \geq 1$ and $\eta > 0$.

The idea of Corollary 1.4 is the same as the one in [CS21]. If we write $F : \mathcal{C} \setminus B_R(0) \rightarrow \Sigma$, $F(p) = p + W(p)$ and $W = W^\beta \mathbf{n}_\beta$ where \mathbf{n}_β 's form a local orthonormal basis of $N\mathcal{C}$, then we know that

$$r^{-1}(\partial_{\omega_i} W)(p) = r^{-1}(\partial_{\omega_i} W^\beta) \mathbf{n}_\beta + r^{-1}W^\beta \partial_{\omega_i} \mathbf{n}_\beta.$$

Using a rotation as in Lemma 1.2, we can compute (using the fact that $A_{\mathcal{C}}(\partial_r, \cdot) = 0$)

$$\partial_r F = \partial_r + (\partial_r W)(p) \text{ and } r^{-1}\partial_{\omega_i} F = r^{-1}\partial_{\omega_i} + r^{-1}(\partial_{\omega_i} W^\beta) \mathbf{n}_\beta + r^{-1}W^\beta \partial_{\omega_i} \mathbf{n}_\beta.$$

Since we have $|W| = O(r^{-1})$, $|\partial_r W| = O(r^{-2})$, $r^{-1}|\partial_{\omega_i} W| = O(r^{-2+\eta})$, $\langle r^{-1}\partial_{\omega_i} \mathbf{n}_\beta, \partial_r \rangle$, and $|\langle r^{-1}\partial_{\omega_i} \mathbf{n}_\beta, \nu_{\mathcal{C},k} \rangle| \leq C$, we have $|h| = O(r^{-2})$. The higher derivative estimates follow from interpolation, as in Lemma 1.2. We also have the following estimates for the normal bundle of Σ .

Lemma 1.5. We have $|\nabla^{(j)}(r\partial_r W - W)|(p) = O(r^{-1-j})$ for any $j \geq 0$,

$$\begin{aligned} A_\Sigma(\partial_r F, \partial_r F) &= O(r^{-3}), \\ A_\Sigma(\partial_r F, r^{-1}\partial_{\omega_i} F) &= O(r^{-3}), \text{ and} \\ A_\Sigma(r^{-1}\partial_{\omega_i} F, r^{-1}\partial_{\omega_j} F) &= A_{\mathcal{C}}(r^{-1}\partial_{\omega_i}, r^{-1}\partial_{\omega_j}) + O(r^{-3+\eta}) \end{aligned}$$

as $r \rightarrow \infty$, and $|\nabla_{\mathcal{C}}^{(k)}(A_\Sigma \circ F - A_{\mathcal{C}})| = O(r^{-3-k+\eta})$ for any $\eta > 0$ and $k \geq 1$. Moreover, the vector field $V := \mathbf{proj}_{T\Sigma} F(p) - r\partial_r F$ is tangent to Σ and satisfies $|V| = O(r^{-1})$, $|\nabla^{(k)}V| = O(r^{-1-k+\eta})$ for $\eta > 0$. And the radial derivative satisfies

$$\vec{x} \cdot \nabla_\Sigma f = r(\partial_r f)^\perp + \alpha_3 \cdot \nabla_{g_{\mathcal{C}}} f,$$

where $|\alpha_3| = O(r^{-1})$ and $|\nabla^{(j)}\alpha_3| = O(r^{-1-j+\eta})$ for $\eta > 0$ and $j \geq 1$, as $r \rightarrow \infty$.

1.2. Mapping properties between weighted Hölder spaces. Next, we define some relevant weighted Hölder spaces of normal sections as in [CS21]. The spaces defined in [CS21] were constructed based on the ones in [KKM18]. We will then mention a Schauder-type estimate for the stability operator for the self-shrinker equation. We will follow our notations in the previous subsection.

From now on, we fix a cutoff function $\chi : [0, \infty) \rightarrow [0, 1]$ so that $\text{supp}\chi \subseteq [R, \infty)$, $\chi \equiv 1$ in $[2R, \infty)$, and $|\nabla^j \chi| \leq CR^{-j}$ for $j \geq 1$ and C independent of R sufficiently large. This allows us to define the primary Hölder spaces.

Definition 1.6. Let $\gamma \in \mathbb{R}$ and $\alpha \in (0, 1)$.

(1) (Homogeneously weighted Hölder spaces) For a normal vector field V over Σ , we define a norm

$$\|V\|_{0;-\gamma}^{\text{hom}} := \sup_{x \in \Sigma} \tilde{r}(x)^\gamma |V(x)|$$

and a semi-norm

$$[V]_{\alpha;-\gamma-\alpha}^{\text{hom}} := \sup_{x,y \in \Sigma} \frac{1}{\tilde{r}(x)^{-\gamma-\alpha} + \tilde{r}(y)^{-\gamma-\alpha}} \frac{|V(x) - \mathbf{proj}_{N_x \Sigma} V(y)|}{|x-y|^\alpha}.$$

Then we let $C_{\text{hom},-\gamma}^{0,\alpha}(\Sigma)$ to be the set of normal vector fields V such that

$$\|V\|_{0,\alpha;-\gamma}^{\text{hom}} := \|V\|_{0;-\gamma}^{\text{hom}} + [V]_{\alpha;-\gamma-\alpha}^{\text{hom}} < \infty.$$

Similarly, we define $C_{\text{hom},-\gamma}^{2,\alpha}(\Sigma)$ to be the set of V such that

$$\|V\|_{2,\alpha;-\gamma}^{\text{hom}} = \sum_{j=0}^2 \|(\nabla_\Sigma)^{(j)} V\|_{0,\alpha;-\gamma}^{\text{hom}} < \infty.$$

(2) (Anisotropically weighted Hölder spaces) We define $C_{\text{an},-1}^{2,\alpha}(\Sigma)$ to be the space of $V \in C_{\text{hom},-1}^{2,\alpha}(\Sigma)$ such that

$$\|V\|_{2,\alpha;-1}^{\text{an}} := \|V\|_{2,\alpha;-1}^{\text{hom}} + \|\vec{x} \cdot \nabla_\Sigma V\|_{0,\alpha;-1}^{\text{hom}} < \infty.$$

(3) (Cone Hölder spaces) We define $\mathcal{CS}_{-1}^{0,\alpha}(\Sigma) := C_{\text{hom},-1}^{0,\alpha}(\Sigma)$ and

$$\mathcal{CS}_{-1}^{2,\alpha}(\Sigma) := C^{2,\alpha}(\Gamma) \times C_{\text{an},-1}^{2,\alpha}(\Sigma).$$

An element $(v, V) \in \mathcal{CS}_{-1}^{2,\alpha}(\Sigma)$ will be considered as a section on Σ given by

$$U = U_{(v,V)}(r, \omega) = \chi(r)(v(\omega))^\perp r + V(r, \omega)$$

for $r \geq R$, and $u = V$ otherwise. We will write $U = U_{(v,V)}$ if everything is clear. Finally, we define the norm

$$\|U\|_{\mathcal{CS}_{-1}^{2,\alpha}(\Sigma)} := \inf_{U = \chi(r)(v(\omega))^\perp r + V(r, \omega)} \|v\|_{C^{2,\alpha}(\Gamma)} + \|V\|_{2,\alpha;-1}^{\text{an}}.$$

Next, we will prove a Schauder-type estimate for the operator $\mathcal{L}_{\frac{1}{2}}^\perp$, defined by

$$\mathcal{L}_{\frac{1}{2}}^\perp V := \Delta_\Sigma^\perp V - \frac{1}{2} \cdot \nabla_{x^T}^\perp V + \frac{1}{2} V.$$

An important ingredient is a vector-valued version of Brandt's non-standard interior Schauder estimate.

Proposition 1.7. Suppose $B_2 \in \mathbb{R}^n$ and there are $C^{0,\alpha}$ functions $a_{ij}, b_i^{kl}, c^{kl}, u^k, f: B_2 \times [-2, 0] \rightarrow \mathbb{R}$ such that

$$\frac{\partial}{\partial t} u^k = a_{ij} D_{ij}^2 u^k + b_i^{kl} D_i u^l + c^{kl} u^l + f$$

for all $k = 1, \dots, n$. If the system is uniformly parabolic in the sense that

$$a_{ij} \xi_i \xi_j \geq \lambda |\xi|^2$$

for some $\lambda > 0$ and

$$\sup_{t \in [-2, 0]} (\|a_{ij}(\cdot, t)\|_{C^{0,\alpha}(B_1)} + \|b_i^{kl}(\cdot, t)\|_{C^{0,\alpha}(B_1)} + \|c^{kl}(\cdot, t)\|_{C^{0,\alpha}(B_1)}) \leq \Lambda$$

for some $\Lambda > 0$, then for $T \in (-1, 0]$,

$$\sup_{t \in [-1, T]} \|u(\cdot, t)\|_{C^{2,\alpha}(B_1)} \leq C \sup_{t \in [-2, T]} (\|u(\cdot, t)\|_{C^{0,\alpha}(B_2)} + \|f(\cdot, t)\|_{C^{0,\alpha}(B_2)})$$

for some $C = C(n, \alpha, \lambda, \Lambda)$ where $u = (u^1, \dots, u^m)$.

Proof. By the non-standard Schauder estimate for parabolic equations established in [Bra69] (see also [CS21, Theorem 3.6]), for any k and $t \in [-2, 0]$, we have

$$\|u^k(\cdot, t)\|_{C^{2,\alpha}(B_1)} \leq \|u^k(\cdot, t)\|_{C^{0,\alpha}(B_2)} + \|b_i^{kl}(\cdot, t)D_i u^l(\cdot, t) + c^{kl}(\cdot, t)u^l(\cdot, t) + f(\cdot, t)\|_{C^{0,\alpha}(B_2)}.$$

Then the proposition follows by rearrangement after combining this and the standard estimate

$$\|h\|_{C^2(B_1)} \leq \varepsilon [D^2 h]_{\alpha, B_2} + C_1(\alpha) \varepsilon^{-C_2(\alpha)} \|h\|_{C^0(B_2)}$$

for any $\varepsilon > 0$ and any $C^{2,\alpha}$ function $h: B_2 \rightarrow \mathbb{R}$. \square

We would like to use this estimate and the diffeomorphism generated by the position vector field to obtain the Schauder-type estimate we would like. Similar as in [CS21], we consider the time dependent vector field $X_t = \frac{1}{-2t}x^T$ and define $\Phi_t: \Sigma \rightarrow \Sigma$ as $\frac{\partial}{\partial t}\Phi_t = X_t \circ \Phi_t$ with $\Phi_{-1} = Id$ with $\hat{g}_t = (-t)\Phi_t^*g_\Sigma$. Then we know that if $F: \Sigma \rightarrow \mathbb{R}^{n+k}$ is the embedding of Σ in \mathbb{R}^{n+k} , $\hat{F}_t = \sqrt{-t}(F \circ \Phi_t): \Sigma \rightarrow \mathbb{R}^{n+1}$ is a MCF and $\hat{g}_t = \hat{F}_t^*g_{\mathbb{R}^{n+k}}$.

We can parameterize the end of Σ using $\mathcal{C} \setminus B_r(0)$ as we did before. We denote the parametrization as $F: \mathcal{C} \setminus B_r(0), p \rightarrow p + W(p)$. Then we define

$$\tilde{\Phi}_t = F^{-1} \circ \Phi_t \circ F$$

and

$$\phi_t: (R, \infty) \times \Gamma \rightarrow (R, \infty) \times \Gamma, (r, \omega) \mapsto ((-t)^{-1/2}r, \omega) \text{ for } t \in [-1, 0).$$

Similarly as in [CS21], we know that for $t \in [-1, 0)$ and r sufficiently large,

$$d_{g_C}(\tilde{\Phi}_t(t, \theta), \phi_t(r, \theta)) \lesssim \frac{1}{\sqrt{-t} \cdot r},$$

and in the flat cylindrical metric $dr^2 + g_\Gamma$, we have

$$|D^{(j)}(\tilde{\Phi}_t - \phi_t)|(r, \theta) \lesssim \frac{1}{\sqrt{-t} \cdot r^{1+j-\eta}}.$$

In the higher-codimensional case, the stability operator for the self-shrinker equation is¹

$$L = \Delta_\Sigma^\perp - \frac{1}{2}\nabla_{x^T}^\perp + \frac{1}{2} + \sum_{k,\ell} \langle \cdot, A_{k\ell} \rangle A_{k\ell} = \mathcal{L}_{\frac{1}{2}}^\perp + \sum_{k,\ell} \langle \cdot, A_{k\ell} \rangle A_{k\ell}.$$

¹One can see [ALW14], [AS13] or [LL15] for the higher-codimensional generalization of the one defined in [CM12].

Then for a normal section V such that $\mathcal{L}_{\frac{1}{2}}^\perp V + \sum_{k,\ell} \langle V, A_{k\ell} \rangle A_{k\ell} = E$, if we define $a: N\Sigma \rightarrow N\Sigma$ by $a(V) = \sum_{k,\ell} \langle V, A_{k\ell} \rangle A_{k\ell}$, we would know that $\|a\|_{C^{0,\alpha}(B_1(x))} = O(|x|^{-2})$. Following this, consider

$$\hat{V}(x, t) = \sqrt{-t} V(\Phi_t(x)), \quad \hat{E}(x, t) = \frac{1}{\sqrt{-t}} E(\Phi_t(x)) \text{ and } \hat{a}(x, t) = \frac{1}{-t} a(\Phi_t(x)).$$

Then we have

$$\frac{\partial \hat{V}}{\partial t} - \Delta_{\hat{g}_t}^\perp \hat{V} - \hat{a}(\hat{V}) = \hat{E}.$$

Using these, we have the corresponding claim in higher codimension.

Proposition 1.8. Consider $a: N\Sigma \rightarrow N\Sigma$ with $\|a\|_{C^{0,\alpha}(B_1(x))} = O(|x|^{-2})$ for $x \in \Sigma$ with $|x| \rightarrow \infty$, i.e., $a \in C_{\text{hom};-2}^{0,\alpha}(\Sigma)$. There exists $C = C(\Sigma, a)$ such that if $V \in C_{\text{loc}}^{2,\alpha}(\Sigma) \cap C_{\text{hom};+1}^0(\Sigma)$ satisfies $\mathcal{L}_{\frac{1}{2}}^\perp V + aV \in \mathcal{CS}_{-1}^{0,\alpha}(\Sigma)$, then $V \in \mathcal{CS}_{-1}^{2,\alpha}(\Sigma)$ and we have the estimate

$$\|V\|_{\mathcal{CS}_{-1}^{2,\alpha}(\Sigma)} \leq C \left(\|V\|_{C_{\text{hom};+1}^0(\Sigma)} + \|\mathcal{L}_{\frac{1}{2}}^\perp V + aV\|_{\mathcal{CS}_{-1}^{0,\alpha}(\Sigma)} \right).$$

The proof is essentially the same as in the codimension one case based on the discussion above and Proposition 1.7. Note that in the higher-codimensional case, we set

$$W := r(\partial_r V)^\perp - V = -2E + 2\Delta^\perp V + 2a(V) - (\vec{x} \cdot \nabla_\Sigma^\perp V - r(\partial_r V)^\perp)$$

and the estimate comes from Lemma 1.5.

2. Estimates on weighted spaces

In this section, we establish some linear estimates on weighted Sobolev spaces, extending those in the hypersurface case in [CS21] to the higher-codimensional case.

Let $\rho(x) := (4\pi)^{-n/2} e^{-|x|^2/4}$ be the n -dimensional backward heat kernel. For any normal vector field or any tensor V over Σ , we define

$$\|V\|_{L_W^2(\Sigma)}^2 := \int_\Sigma |V|_g^2 \cdot \rho \, dx,$$

and let L_W^2 be the space of normal vector field V over Σ such that $\|V\|_{L_W^2(\Sigma)}^2 < \infty$. Using this, we consider

$$\|V\|_{H_W^k(\Sigma)}^2 := \sum_{i=0}^k \|(\nabla^\perp)^i V\|_{L_W^2}^2,$$

and let $H_W^k(\Sigma)$ be the associated Sobolev space of normal vector fields. When it is clear, we will omit the symbol Σ and just write $\|V\|_{L_W^2}$ and $\|V\|_{H_W^k}$.

The following Sobolev inequality was due to Ecker [Eck00] in the hypersurface case.

Proposition 2.1. For $V \in H_W^1(\Sigma)$, we have

$$\int_\Sigma |V|^2 \cdot |x|^2 \rho \, dx \leq 4 \int_\Sigma (n|V|^2 + 4|\nabla^\perp V|^2) \rho \, dx.$$

Proof. Consider a smooth normal vector field V over Σ with compact support. Then the first variation formula implies

$$\begin{aligned} - \int_{\Sigma} \langle |V|^2 \rho(x) x, \mathbf{H} \rangle dx &= \int_{\Sigma} \operatorname{div} (|V|^2 \rho(x) x) dx \\ &= \int_{\Sigma} \left(n|V|^2 + \langle \nabla |V|^2, x \rangle - \frac{1}{2} |V|^2 \cdot |x^T|^2 \right) \rho dx. \end{aligned}$$

Since Σ is a self-shrinker (that is, $\mathbf{H} = -\frac{x^\perp}{2}$), we derive

$$\begin{aligned} \frac{1}{2} \int_{\Sigma} |V|^2 \cdot |x|^2 \rho dx &= \int_{\Sigma} (n|V|^2 + 2|V| \langle \nabla |V|, x \rangle) \rho dx \\ &\leq \int_{\Sigma} \left(n|V|^2 + 4|\nabla^\perp V|^2 + \frac{1}{4}|x|^2 |V|^2 \right) \rho dx \end{aligned}$$

by the Cauchy-Schwarz inequality and the Kato inequality. Then the inequality for V follows after rearrangement. This implies the inequality for $V \in H_W^1(\Sigma)$ using an approximation argument. \square

Lemma 2.2. For $V \in H_W^2(\Sigma)$, we have $\|\nabla^\perp V\|_{L_W^2}^2 \leq \|\mathcal{L}^\perp V\|_{L_W^2} \|V\|_{L_W^2}$. In particular, the inclusion $H_W^1(\Sigma) \hookrightarrow L_W^2(\Sigma)$ is compact.

Proof. As above, it suffices to prove the inequality for smooth V with compact support. Using integration by parts, we have

$$\begin{aligned} 0 &= \int_{\Sigma} \mathcal{L} |V|^2 \rho dx \\ &= 2 \int_{\Sigma} \left(\langle \Delta^\perp V, V \rangle + |\nabla^\perp V|^2 - \frac{1}{2} \langle \nabla_{x^T}^\perp V, V \rangle \right) \rho dx \\ &= 2 \int_{\Sigma} (\langle \mathcal{L}^\perp V, V \rangle + |\nabla^\perp V|^2) \rho dx. \end{aligned}$$

As a result, the first conclusion follows from the Cauchy-Schwarz inequality. The same argument in [CS21, lemma 3.12] implies the second conclusion. \square

Proposition 2.3. For $V \in H_W^2(\Sigma)$, we have

$$\|V\|_{H_W^2}^2 \leq C \left(\|\mathcal{L}^\perp V\|_{L_W^2}^2 + \|V\|_{L_W^2}^2 \right)$$

for some $C = C(\Sigma)$.

Proof. As above, it suffices to prove the inequality for smooth V with compact support. Around a point $p \in \Sigma$, take normal coordinates e_1, \dots, e_n such that $\nabla_{e_i} e_j(p) = 0$. At p , we have

$$\begin{aligned}
\Delta |\nabla^\perp V|^2 &= \Delta \left(\sum_i \langle \nabla_{e_i}^\perp V, \nabla_{e_i}^\perp V \rangle \right) \\
&= 2 \sum_{i,k} \langle \nabla_{e_k}^\perp \nabla_{e_k}^\perp \nabla_{e_i}^\perp V, \nabla_{e_i}^\perp V \rangle + 2 \sum_{i,k} \langle \nabla_{e_k}^\perp \nabla_{e_i}^\perp V, \nabla_{e_k}^\perp \nabla_{e_i}^\perp V \rangle \\
&= 2 \sum_{i,k} \langle \nabla_{e_k}^\perp \nabla_{e_i}^\perp \nabla_{e_k}^\perp V + \nabla_{e_k} (R^\perp(e_i, e_k)V), \nabla_{e_i}^\perp V \rangle + 2|(\nabla^\perp)^2 V|^2 \\
&= 2 \sum_{i,k} \langle \nabla_{e_i}^\perp \nabla_{e_k}^\perp \nabla_{e_k}^\perp V + R^\perp(e_i, e_k) \nabla_{e_k}^\perp V + \nabla_{e_k} (R^\perp(e_i, e_k)V), \nabla_{e_i}^\perp V \rangle + 2|(\nabla^\perp)^2 V|^2 \\
&= 2 \langle \nabla^\perp \Delta^\perp V, \nabla^\perp V \rangle + 2|(\nabla^\perp)^2 V|^2 + 4 \langle R^\perp(e_i, e_k) \nabla_{e_k}^\perp V, \nabla_{e_i}^\perp V \rangle + 2 \langle (\nabla_{e_k} R^\perp)(e_i, e_k)V, \nabla_{e_i}^\perp V \rangle,
\end{aligned}$$

where the normal curvature

$$R^\perp(X, Y)W = \nabla_Y^\perp \nabla_X^\perp W - \nabla_X^\perp \nabla_Y^\perp W + \nabla_{[X, Y]}^\perp W$$

is defined for any tangent vectors X, Y and any normal vector W . Therefore, using Lemma 1.1 and the Ricci equation

$$\langle R^\perp(X, Y)W, Z \rangle = \sum_{k=1}^n (\langle A(X, e_k), W \rangle \langle A(Y, e_k), Z \rangle - \langle A(Y, e_k), W \rangle \langle A(X, e_k), Z \rangle),$$

we can estimate

$$\Delta |\nabla^\perp V|^2 = 2 \langle \nabla^\perp \Delta^\perp V, \nabla^\perp V \rangle + 2|(\nabla^\perp)^2 V|^2 + O(|V|^2 + |\nabla^\perp V|^2).$$

Therefore, we have

$$\begin{aligned}
&\mathcal{L} |\nabla^\perp V|^2 \\
&= 2 \langle \nabla^\perp \mathcal{L}^\perp V, \nabla^\perp V \rangle + 2|(\nabla^\perp)^2 V|^2 - \frac{1}{2} \nabla_{x^T} |\nabla^\perp V|^2 + \langle \nabla^\perp \nabla_{x^T}^\perp V, \nabla^\perp V \rangle + O(|V|^2 + |\nabla^\perp V|^2) \\
&= 2 \langle \nabla^\perp \mathcal{L}^\perp V, \nabla^\perp V \rangle + 2|(\nabla^\perp)^2 V|^2 + \sum_i (-\langle \nabla_{x^T}^\perp \nabla_{e_i}^\perp V, \nabla_{e_i}^\perp V \rangle + \langle \nabla_{e_i}^\perp \nabla_{x^T}^\perp V, \nabla^\perp e_i V \rangle) \\
&+ O(|V|^2 + |\nabla^\perp V|^2) \\
&= 2 \langle \nabla^\perp \mathcal{L}^\perp V, \nabla^\perp V \rangle + 2|(\nabla^\perp)^2 V|^2 + \sum_i \langle R^\perp(x^T, e_i)V, \nabla_{e_i}^\perp V \rangle + O(|V|^2 + |\nabla^\perp V|^2) \\
&= 2 \langle \nabla^\perp \mathcal{L}^\perp V, \nabla^\perp V \rangle + 2|(\nabla^\perp)^2 V|^2 + O(|V|^2 + |\nabla^\perp V|^2)
\end{aligned}$$

using Lemma 2.2 and the Ricci equation again. Hence, after integrating over Σ and using integration by parts, we get

$$0 = -2 \|\mathcal{L}^\perp V\|_{L_W^2}^2 + 2 \|(\nabla^\perp)^2 V\|_{L_W^2}^2 + O\left(\|V\|_{L_W^2}^2 + \|\nabla^\perp V\|_{L_W^2}^2\right).$$

The proposition then follows after combining this with lemma 2.2. \square

Now we can derive the existence of weak solutions to the stability operator L . To this end, recall that we define

$$LV := \mathcal{L}^\perp V + \sum_{i,j} \langle V, A_{ij} \rangle A_{ij} + V = \Delta^\perp V - \frac{1}{2} \nabla_{x^T}^\perp V + \sum_{i,j} \langle V, A_{ij} \rangle A_{ij} + V$$

for $V \in N\Sigma$, and

$$B_\gamma(U, V) = \int_\Sigma \left(\langle \nabla^\perp U, \nabla^\perp V \rangle - (\gamma + 1) \langle U, V \rangle - \sum_{i,j} \langle U, A_{ij} \rangle \langle V, A_{ij} \rangle \right) \rho \, dx,$$

which is the bounded bilinear form associated to $-(L + \gamma)$. By taking large enough γ such that $\gamma \geq \sup_\Sigma |A|^2 + 2$, we have

$$\|V\|_{H_W^2}^2 \leq |B_\gamma(V, V)|.$$

Using this estimate, the standard Fredholm alternative, the Lax-Milgram theorem, Lemma 2.2 and Proposition 2.3, we obtain the following result.

Theorem 2.4. The space $\text{Ker}L$ of the weak solutions to $LV = 0$ is finite-dimensional. Moreover, for $U \in L_W^2(\Sigma)$, $LV = U$ has a weak solution in $H_W^1(\Sigma)$ if and only if U is L_W^2 -orthogonal to $\text{Ker}L$, and if V is orthogonal to $\text{Ker}L$ also, then $\|V\|_{H_W^2} \lesssim \|U\|_{L_W^2}$.

Finally we generalize [CS21, Lemma 3.15] to sections of the normal bundle.

Proposition 2.5. There exists $C > 0$ such that for any section $X \in L_W^2(\Sigma) \cap C^0(\Sigma)$ of the normal bundle, if $V \in H_W^1(\Sigma)$ satisfies $LV = X$, then $V \in C_{\text{hom};+1}^0(\Sigma)$ and for R sufficiently large,

$$\|V\|_{C_{\text{hom};+1}^0(\Sigma)} \leq C (\|X\|_{C^0(\Sigma)} + \|V\|_{C^0(\Sigma \cap B_R(0))}).$$

Proof. Define $\varphi(x) = \alpha|x|$ with α to be specified later, and consider the section $U = \frac{V}{\varphi}$. We have that

$$LU = L\left(\frac{V}{\varphi}\right) = \frac{LV}{\varphi} + 2\nabla_{\frac{1}{\varphi}}^\perp V + \left(\Delta\left(\frac{1}{\varphi}\right) - \frac{1}{2}\nabla_{x^T}\left(\frac{1}{\varphi}\right)\right)V.$$

Since we know

$$\nabla\left(\frac{1}{\varphi}\right) = -\frac{\alpha x^T}{\varphi^2|x|}$$

and

$$\Delta\left(\frac{1}{\varphi}\right) = -\alpha\left(\frac{n - \langle x^\perp, \mathbf{H} \rangle}{\varphi^2|x|}\right) + \frac{2\alpha^2|x^T|^2}{|x|^2\varphi^3} + \frac{\alpha|x^T|^2}{\varphi^2|x|^3},$$

we can derive

$$\Delta\left(\frac{1}{\varphi}\right) - \frac{1}{2}\nabla_{x^T}\left(\frac{1}{\varphi}\right) = \frac{\alpha|x|}{2\varphi^2} - \frac{n\alpha}{\varphi^2|x|} + \frac{2\alpha^2|x^T|^2}{|x|^2\varphi^3} + \frac{\alpha|x^T|^2}{\varphi^2|x|^3} = \frac{1}{2\alpha|x|} - \frac{n}{\alpha|x|^3} + \frac{3|x^T|^2}{|x|^5\alpha}.$$

Combining these with $|A| = O(r^{-1})$, we can estimate

$$\begin{aligned}
\langle \mathcal{L}_{\frac{1}{2}}U, U \rangle &= \langle LU, U \rangle - |AU|^2 = \left\langle \frac{X}{\varphi} + 2\nabla_{\frac{\nabla^\perp}{\varphi}}(U\varphi) + \frac{1}{2}U, U \right\rangle + O(r^{-2})|U|^2 \\
(2.6) \quad &= \frac{1}{2}|U|^2 + \left\langle \frac{X}{\varphi}, U \right\rangle - \frac{2|U|^2|\nabla\varphi|^2}{\varphi^2} + 2\varphi\nabla^\perp U \left(U, \nabla \frac{1}{\varphi} \right) + O(r^{-2})|U|^2.
\end{aligned}$$

Next, we take a cutoff function $\eta : [0, +\infty) \rightarrow \mathbf{R}$ such that $\eta' \geq 0$ and $\eta(x) = \begin{cases} 0, & 0 < x < \frac{1}{2} \\ 1, & x > 1 \end{cases}$.

Then we have

$$(2.7) \quad \nabla^\perp U(U, \nabla \eta^2(|U|)) = 2\eta\eta'\nabla^\perp U(U, \nabla|U|) = 2\eta\eta'\nabla^\perp U(U, \frac{\nabla^\perp U(U)}{|U|}) = 2\eta\eta' \frac{|\nabla^\perp U(U)|^2}{|U|} \geq 0.$$

We set

$$(2.8) \quad \alpha = 2R^{-1} \sup_{\Sigma \cap B_R(0)} |V| + 8R^{-1} \sup_{\Sigma \cap B_R(0)} |X|.$$

This implies $\alpha R \geq 2 \sup_{\Sigma \cap B_R(0)} |V|$, and hence, we have

$$\begin{aligned}
\int_{\Sigma \setminus B_R(0)} \eta^2(|U|) \langle \mathcal{L}_{\frac{1}{2}}U, U \rangle \rho dx &= \int_{\Sigma \setminus B_R(0)} \left(\eta^2(|U|) \left(-|\nabla^\perp U|^2 + \frac{1}{2}|U|^2 \right) - \nabla^\perp U(U, \nabla \eta^2(|U|)) \right) \rho dx \\
(2.9) \quad &\leq \int_{\Sigma \setminus B_R(0)} \eta^2(|U|) \left(-|\nabla^\perp U|^2 + \frac{1}{2}|U|^2 \right)
\end{aligned}$$

by (2.7). On the other hand, as $|A| = O(r^{-1})$ and $\alpha R \geq 8 \sup_{\Sigma} |X|$, we have

$$\begin{aligned}
\eta^2(|U|) \langle \mathcal{L}_{\frac{1}{2}}U, U \rangle &= \eta^2(|U|) \left(\frac{1}{2}|U|^2 + \left\langle \frac{X}{\varphi}, U \right\rangle + 2\varphi\nabla^\perp U \left(U, \nabla \frac{1}{\varphi} \right) + O(R^{-2})|U|^2 \right) \\
(2.10) \quad &\geq \eta^2(|U|) \left(\frac{1}{8}|U|^2 + O(R^{-2})|\nabla^\perp U|^2 + O(R^{-2})|U|^2 \right).
\end{aligned}$$

Note that if we apply the Sobolev inequality (Proposition 2.1) to $\eta(|U|)U$, we have

$$\begin{aligned}
&R^2 \int_{\Sigma \setminus B_R} \eta^2|U|^2 \rho dx \\
&\leq 4n \int_{\Sigma \setminus B_R} \eta^2|U|^2 \rho dx + 16 \int_{\Sigma \setminus B_R} |\nabla^\perp(\eta U)|^2 \rho dx \\
&\leq 4n \int_{\Sigma \setminus B_R} \eta^2|U|^2 \rho dx + 32 \int_{\Sigma \setminus B_R} \eta^2 |\nabla^\perp U|^2 \rho dx + 32 \int_{\Sigma \setminus B_R} (\eta'(|U|))^2 |\nabla^\perp U| \cdot |U|^2 \rho dx \\
&\leq 4n \int_{\Sigma \setminus B_R} \eta^2|U|^2 \rho dx + 32 \int_{\Sigma \setminus B_R} \eta^2 |\nabla^\perp U|^2 \rho dx + 32 \sup \eta' \cdot \int_{\Sigma \setminus B_R} |\nabla^\perp U|^2 \rho dx
\end{aligned}$$

since $\eta' \neq 0$ only when $|U| \in (1/2, 1)$. Consequently, combining (2.9), (2.10) and Proposition 2.1, we get that

$$\begin{aligned} & \frac{R^2 - 4n}{32} \int_{\Sigma \setminus B_R} (1 + O(R^{-2})) \eta^2 |U|^2 \rho \, dx + \int_{\Sigma \setminus B_R} \left(\frac{1}{2} + O(R^{-2}) \right) \eta^2 |U|^2 \rho \, dx \\ & \leq (4 + O(R^{-2})) \int_{\Sigma \setminus B_R} |\nabla^\perp U|^2 \rho \, dx. \end{aligned}$$

Thus, if we take R sufficiently large, we can derive that $\eta^2(|U|)|U|^2 = 0$, which implies that $|U| \leq 1$. As $U = \frac{V}{\alpha|x|}$, this is equivalent to that

$$(2.11) \quad \|V\|_{C_{\text{hom};+1}^0(\Sigma)} \lesssim \|X\|_{C^0(\Sigma)} + \|V\|_{C^0(\Sigma \cap B_R(0))}.$$

This proves the desired estimate. \square

Combining Proposition 2.5 with the Schauder-type estimate (Proposition 1.8), we obtain the following corollary, which is essential when applying Simon's theory to obtain a Łojasiewicz inequality.

Corollary 2.12. For $U \in \mathcal{CS}_{-1}^{0,\alpha}(\Sigma)$, there exists a solution $V \in \mathcal{CS}_{-1}^{2,\alpha}(\Sigma)$ to $LV = U$ if and only if U is L_W^2 -orthogonal to $\text{Ker}L \subseteq H_W^1(\Sigma)$.

3. Łojasiewicz-Simon inequalities

In this section, we will set up the frameworks that enable us to apply Simon's theory to establish a Łojasiewicz-Simon inequality for entire vector fields over an asymptotically conical self-shrinker, and localize it to the form we would like to apply later. We will follow the guidelines of [CS21] and [Sim83].

Assume M is a normal graph of $V \in N\Sigma$ over Σ . Recall that the Gaussian area (or the F -functional) and the entropy of M are defined to be

$$F(M) := \int_M \rho \, dx \text{ and } \lambda(M) := \sup_{x_0 \in \mathbb{R}^{n+k}, t_0 > 0} F(t_0 M + x_0),$$

where we recall that $\rho(x) := (4\pi)^{-n/2} e^{-|x|^2/4}$. By [Sch14, Section 3], we know that given a variation W of V ,

$$\delta_W F(M) = - \int_\Sigma \left\langle \Pi_{N\Sigma} \left(\mathbf{H}_M + \frac{x^\perp}{2} \right) \Big|_{x=y+V(y)}, W(y) \right\rangle J(y, V, \nabla^\Sigma V) \rho(y + V(y)) \, dy$$

where $\Pi_{N\Sigma}$ is the projection onto $N\Sigma$ and $J(y, V, \nabla^\Sigma V) = \text{Jac}(D \exp_y(V))$ is the area element. Thus, the linearization of F is

$$\mathcal{M}(V) = J(y, V, \nabla^\Sigma V) \rho(y + V(y)) \rho(y)^{-1} \cdot \Pi_{N\Sigma} \left(\mathbf{H}_M + \frac{x^\perp}{2} \right) \Big|_{x=y+V(y)}.$$

To exploit the ideas in [Sim83], we need some good properties of the map \mathcal{M} . The proofs are parallel to the ones in [CS21] with some standard modifications.

Lemma 3.1. For β small enough depending on Σ , the map

$$\mathcal{M}: \{u \in \mathcal{CS}_{-1}^{2,\alpha}(\Sigma) : \|u\|_{\mathcal{CS}_{-1}^{2,\alpha}(\Sigma)} < \beta\} \rightarrow \mathcal{CS}_{-1}^{0,\alpha}(\Sigma)$$

is Fréchet differentiable with its derivative at 0 being L .

Proof. First we prove the continuity of \mathcal{M} . For $V \in \mathcal{CS}_{-1}^{2,\alpha}(\Sigma)$, note that

$$J(y, V, \nabla^\Sigma V) \rho(y + V(y)) \rho(y)^{-1} = J(y, V, \nabla^\Sigma V) \cdot e^{-\frac{2\langle V(y), y \rangle + |V(y)|^2}{4}},$$

which is uniformly bounded in $C^{0,\alpha}(\Sigma)$ as $y \rightarrow \infty$.

For the last term, at a point $y \in \Sigma$, if we let F be a parametrization of Σ near y , then we write \tilde{F} to be the parametrization of M near $y + V(y)$, given by

$$(3.2) \quad \tilde{F}(y) = F(y) + V(y) = F(y) + V^\beta(y) \mathbf{n}_\beta(y)$$

where $\{\mathbf{n}_1, \dots, \mathbf{n}_k\}$ is a local orthonormal basis of $N\Sigma$ near p . Then we have

$$\partial_i \tilde{F} = \partial_i F + (\partial_i V^\beta) \mathbf{n}_\beta - V^\beta A_{ik}^\beta \partial_k F$$

where $A_{ij} = A_{ij}^\beta \mathbf{n}_\beta$ is the second fundamental form of Σ . Therefore, by direct calculations (similar to those used to derive [Wan14, (2.27)]), we can see that if we let $\Phi := \langle V, A \rangle = (\langle V, A_{ij} \rangle)_{ij}$, then

$$N_\beta := -(\text{id} - \Phi)^{-1} \nabla^\Sigma V^\beta + \mathbf{n}_\beta$$

forms a local orthogonal basis ($\beta = 1, \dots, k$) of NM near $\tilde{F}(y)$. Combining this with the self-shrinker equation and the curvature decay (Lemma 1.1), we have

$$\left| \left\langle x, \frac{N_\beta}{|N_\beta|} \right\rangle \right| = \frac{1}{|N_\beta|} (V^\beta - \langle y, \nabla^\Sigma V^\beta \rangle) + O(|y|^{-1})$$

at $x = y + V(y)$. This shows that \mathcal{M} is a bounded map, and its differentiability follows from a similar standard argument. \square

By Theorem 2.4, $\text{Ker} L \subseteq H_W^1(\Sigma)$ is finite-dimensional. Define $\Pi: L_W^2(\Sigma) \rightarrow L_W^2(\Sigma) \cap \mathcal{CS}_{-1}^{2,\alpha}(\Sigma)$ to be the projection onto $\text{Ker} L$ and $\mathcal{N} := \mathcal{M} + \Pi$. By Lemma 3.1, \mathcal{N} is Fréchet differentiable with its derivative at 0 being $L + \Pi$, which is an isomorphism from $\mathcal{CS}_{-1}^{2,\alpha}(\Sigma)$ to $\mathcal{CS}_{-1}^{0,\alpha}(\Sigma)$. Therefore, the inverse function theorem implies that there exists open neighborhoods $W_1 \subseteq \mathcal{CS}_{-1}^{2,\alpha}(\Sigma)$ and $W_2 \subseteq \mathcal{CS}_{-1}^{0,\alpha}(\Sigma)$ of 0 such that $\mathcal{N}: W_1 \rightarrow W_2$ is bijective with the inverse $\Psi: W_2 \rightarrow W_1$. We next prove that these maps behave nicely when restricted to these neighborhoods.

Lemma 3.3. After shrinking W_1 and W_2 if necessary, we can find $C > 0$ such that

$$\|\mathcal{M}(V_1) - \mathcal{M}(V_2)\|_{L_W^2} \leq C \|V_1 - V_2\|_{H_W^2}$$

for $V_1, V_2 \in W_1$ and

$$\|\Psi(U_1) - \Psi(U_2)\|_{H_W^2} \leq C \|U_1 - U_2\|_{L_W^2}$$

for $U_1, U_2 \in W_2$.

Proof. At a point $y \in \Sigma$, if we let F be a parametrization of Σ near y , then we write \tilde{F} to be the parametrization of M near $y + V(y)$, given by (3.2). We may assume $g_{ij} = \langle \partial_i F, \partial_j F \rangle = \delta_{ij}$ at y . Then using the calculations in the proof of [Kha20, lemma 3.18], we have

$$\tilde{g}_{ij} := \langle \partial_i \tilde{F}, \partial_j \tilde{F} \rangle = \delta_{ij} + \langle \partial_i V, \partial_j V \rangle - 2 \langle V, A_{ij} \rangle + V^\beta V^\gamma A_{ik}^\beta A_{jk}^\gamma$$

at $y + V(y)$ where we write $V = V^\beta \mathbf{n}_\beta$ and $A_{ij} = A_{ij}^\beta \mathbf{n}_\beta$ in a local orthonormal basis $\{\mathbf{n}_\beta\}_{\beta=1, \dots, k}$. Using this, we can derive

$$(3.4) \quad \tilde{g}^{ij} = \delta^{ij} - \langle \partial_i V, \partial_j V \rangle + 2 \langle V, A_{ij} \rangle + Q_{ij}(y, V, DV)$$

and

$$(3.5) \quad \tilde{A}_{ij} = A_{ij} + (\partial_{ij}^2 V^\beta) \mathbf{n}_\beta - A_{ij}^\beta (\partial_k V^\beta) (\partial_k F) + Q(y, V, DV)$$

with $|Q_{ij}(y, V, DV)|, |Q(y, V, DV)| \lesssim |y|^{-2} (|V|^2 + |DV|^2)$. Using (3.4) and (3.5), we can get the expression of the mean curvature \tilde{H} of M , and use the same arguments as in [Sim96] to derive

$$\mathcal{M}(V_1) - \mathcal{M}(V_2) = L(V_1 - V_2) + A \cdot (\nabla^\perp)^2 (V_1 - V_2) + B \cdot \nabla^\perp (V_1 - V_2) + C(V_1 - V_2)$$

with

$$\sup_{\Sigma} (|A| + |B| + |C|) \lesssim \|V_1\|_{\mathcal{CS}_{-1}^{2,\alpha}(\Sigma)} + \|V_2\|_{\mathcal{CS}_{-1}^{2,\alpha}(\Sigma)}.$$

Combining this with Lemma 2.2, the first conclusion follows. The second conclusion follows from the same arguments as in [Sim96, Proof of Lemma 1 in 3.12] with the estimates above in place of those there. \square

At this point, we can follow the same arguments as in [Sim83] to derive a Łojasiewicz-Simon inequality for M . For example, we use Theorem 2.4 and Corollary 2.12 to replace [Sim83, Equation (2.5)] and the preceding existence result, and Lemma 3.3 in place of [Sim83, Equation (2.10)]. Then, the arguments there will produce the following Łojasiewicz-Simon inequality for entire vector fields.

Theorem 3.6 (Entire Łojasiewicz-Simon inequality). There exists $\beta_0 > 0$, $\theta \in (0, 1/2)$ and $C > 0$ such that if M is the graph over Σ of a vector field $U \in \mathcal{CS}_{-1}^{2,\alpha}(\Sigma)$ with $\|U\|_{\mathcal{CS}_{-1}^{2,\alpha}(\Sigma)} \leq \beta_0$, then

$$|F(M) - F(\Sigma)|^{1-\theta} \leq C \|\mathcal{M}(U)\|_{L_W^2(\Sigma)} \leq C \left(\int_M |\varphi|^2 \rho \, dx \right)^{1/2}.$$

Next, we need to localize the Simon-Łojasiewicz inequality obtained above, since in a blowup sequence of MCF, we can only write the flow as the graph of a vector field on a large compact set. To this end, we will introduce some scales related to the concerned self-shrinkers. For θ given by Theorem 3.6, define

$$\Theta := \left(\frac{1 - \frac{\theta}{2}}{1 - \theta} \right)^{\frac{1}{4}} \in \left(1, \left(\frac{3}{2} \right)^{\frac{1}{4}} \right].$$

Definition 3.7. For a submanifold $M^n \subseteq \mathbb{R}^{n+k}$, let $\varphi = \mathbf{H} + \frac{x^\perp}{2}$, and we define the following scales of M .

(1) (Shrinker scale)² The shrinker scale $\mathbf{R}(M)$ of M is defined by

$$e^{-\frac{\mathbf{R}(M)^2}{4}} = \int_M |\varphi|^2 dx.$$

(2) (Rough conical scale) For $l \in \mathbb{N}$ and $C_l > 0$ (to be chosen) fixed, the rough conical scale $\tilde{\mathbf{r}}_l(M)$ of M is the largest radius \tilde{r} such that $M \cap B_{\tilde{r}}$ is smooth and

$$|(\nabla^\perp)^k A_M(x)| \leq \frac{C_l}{(1+r(x))^{1+k}}$$

for all $k = 0, \dots, l-1$ and $x \in M \cap B_{\tilde{r}}$.

(3) (Conical scale) For a fixed asymptotically conical self-shrinker Σ^n and $\beta_0 = \beta_0(\Sigma)$ given by the entire Łojasiewicz inequality (Theorem 3.6), the conical scale $\mathbf{r}_l(M)$ of M is the largest radius $r \leq \tilde{\mathbf{r}}_l(M)$ such that there exists $U \in \mathcal{CS}_{-1}^{2,\alpha}(\Sigma)$ with $\|U\|_{\mathcal{CS}_{-1}^{2,\alpha}(\Sigma)} \leq \beta_0$ such that

$$\text{graph } U|_{\Sigma \cap B_r} \subseteq M \text{ and } M \cap B_{r-1} \subseteq \text{graph } U.$$

(4) (Core graphical condition) We say M satisfies the core graphical condition $(*_b, \underline{r})$ if $\tilde{\mathbf{r}}_l(M) \geq r$ and there exists a C^{l+1} vector field U over $\Sigma \cap B_{\underline{r}}$ with $\|U\|_{C^{l+1}} \leq b$ such that

$$\text{graph } U \subseteq M \text{ and } M \cap B_{\underline{r}-1} \subseteq \text{graph } U.$$

(5) (Roughly conical approximate shrinker) For $R \geq \underline{r}$, we say that $M^n \subseteq \mathbb{R}^{n+k}$ is a roughly conical approximate shrinker up to scale R if $\Theta R \leq \tilde{r}_\ell(M)$, M satisfies the core graphical hypothesis $(*_b, \underline{r})$, and $|\phi| + (1+|x|)|\nabla^\perp \phi| \leq s(1+|x|)^{-1}$ on $M \cap B_{\Theta R}$.

At this point, we are able to obtain a higher-codimensional version of the graphical estimate as in [CS21, Proposition 7.2]. The proof is similar, and as in Proposition 1.8, we need to take the normal projections of the corresponding vector fields.

Proposition 3.8. Taking ℓ sufficiently large, there are constants $b, s > 0$ sufficiently small, depending on Σ , β_0 , C_ℓ , and λ_0 such that if $M^n \subseteq \mathbb{R}^{n+k}$ has $\lambda(M) \leq \lambda_0$ and is a roughly conical approximate shrinker up to scale R , then there exists $U : \Sigma \rightarrow \mathbb{R}^k$ with

$$\text{graph } U|_{\Sigma \cap B_R(0)} \subseteq M \text{ and } M \cap B_{R-1}(0) \subseteq \text{graph } U$$

and $\|U\|_{\mathcal{CS}_{-1}^{2,\alpha}(\Sigma)} \leq \beta_0$. That is to say, the conical scale satisfies $r_\ell(M) \geq R$.

Using Proposition 3.8, we can get the final version of the Łojasiewicz-Simon inequality we would like. This can be done based on the same arguments in the proofs of Theorem 6.1 and Theorem 8.1 in [CS21].

²Here we define the shrinker scale following [CS21], and it is different from the shrinker scale defined in [CM15] and [CM19].

Theorem 3.9 (Localized Lojasiewicz-Simon inequality). Let $M \subseteq \mathbb{R}^N$ with $\lambda(M) \leq \lambda_0$. If $\mathbf{R}(M)$ is sufficiently large with $\mathbf{R}(M) \leq \tilde{\mathbf{r}}_l(M) - 1$ and M satisfies the core graphical condition $(*_b, \underline{r})$, then there exists $C = C(\Sigma, \lambda_0)$ such that

$$|F(M) - F(\Sigma)| \leq C \left(\int_M |\varphi|^2 \rho \, dx \right)^{\frac{1}{2(1-\theta/3)}}$$

where θ is the one given by Theorem 3.6.

4. Uniqueness for asymptotically conical self-shrinkers

Now we are in a position to prove Theorem 0.1. Fix some sufficiently large \underline{r} and sufficiently small ε (which will be chosen in the proof of Proposition 4.1). Assume a tangent flow of a MCF \widetilde{M}_t is modelled on Σ . That means that after rescaling, we can obtain a rescaled MCF M_τ ($\tau \in [-2, \infty)$) so that there is a map

$$U: (\Sigma \cap B_{\varepsilon^{-1}}) \times [-2, \varepsilon^{-2}] \rightarrow \mathbb{R}^k$$

such that for all $\tau \in [-2, \varepsilon^{-2}]$, we have

$$\text{graph } U(\cdot, \tau) \subseteq M_\tau, M_\tau \cap B_{\varepsilon^{-1}-1} \subseteq \text{graph } U(\cdot, \tau),$$

$\|U\|_{C^{l+1}(\Sigma \cap B_{\varepsilon^{-1}})} \leq \varepsilon$, and $F(M_\tau) - F(\Sigma) \leq \varepsilon$. As in [CS21], we define the graphical time to be

$$\bar{\tau} := \sup\{t \in [-2, \infty) : M_\tau \text{ satisfies } (*_{b, \underline{r}}) \text{ for all } \tau \in [-2, t]\}.$$

We are going to establish the following extension result.

Proposition 4.1. There exists $\underline{r}_0 = \underline{r}_0(\Sigma, M_{-2})$ sufficiently large such that for $\underline{r} \geq \underline{r}_0$, $\varepsilon = \varepsilon(\Sigma, \underline{r})$ sufficiently small and $C_l = C_l(\Sigma, \underline{r})$ sufficiently large, we have

$$(4.2) \quad \tilde{\mathbf{r}}_l \geq \frac{1}{2} e^{\frac{\tau}{2}} \underline{r}$$

for all $\tau \in [0, \bar{\tau})$ such that $\mathbf{R}(M_\tau) > \tilde{\mathbf{r}}_l(M_\tau)$.

In the proof of proposition 4.1, the key ingredient is White's Brakke estimate.³

Theorem 4.3 ([Whi05]). For $\varepsilon > 0$ small enough, there exists a constant $C(\varepsilon) > 0$ such that the following holds. If $\hat{M}_t^k \subseteq \mathbb{R}^{n+k}$ ($t < 0$) flow by the MCF with $\lambda(\hat{M}_s) \leq \lambda_0$, and

$$(-4\pi t_0)^{-\frac{n}{2}} \int_{\hat{M}_{t_0}} e^{\frac{|x-x_0|^2}{4t_0}} \leq \varepsilon$$

for some $x_0 \in \mathbb{R}^{n+k}$ and $t_0 < 0$, then

$$\sup_{\hat{M}_t \cap B_{\sqrt{-t_0}/2}(x_0)} |A|^2 \leq \frac{C(\varepsilon)}{-t_0}$$

for all $t \in [\frac{t_0}{4}, 0)$. Also, the constant $C(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$.

³The version we apply here is the one used in [CM15].

Proof of Proposition 4.1. Let $\tau_0 \in [0, \bar{\tau})$. Consider the flow $\hat{M}_t := \sqrt{-t}M_{\tau_0 - \log(-t)}$, which is an ordinary MCF (depending on τ_0). Then $\hat{M}_{-1} = M_{\tau_0}$. For $\delta_0 > 0$, we take $\varepsilon = \varepsilon(\delta_0) > 0$ so small that the corresponding constant $C(\varepsilon)$ in theorem 4.3 satisfies $C(\varepsilon) \leq 4\delta_0$. Based on [CIM15, Lemma 2.4] and the entropy bound λ_0 , we can take $R_0 = R_0(\delta_0)$ large enough such that

$$(4.4) \quad (16\pi)^{-\frac{n}{2}} \int_{\hat{M}_{-4} \setminus B_{R_0}(x_0)} e^{-\frac{|x-x_0|^2}{16}} \leq \frac{\varepsilon}{2}$$

for any $x_0 \in \mathbb{R}^{n+k}$. Next, by taking \underline{r} large and b small, using Lemma 1.1 and $(*_b, \underline{r})$, we have

$$|A_{\hat{M}_{-4}}|^2 \leq \sigma_0 \text{ on } \hat{M}_{-4} \cap (\overline{B_{\underline{r}}} \setminus B_{\underline{r}/2})$$

for any given $\sigma_0 > 0$. We may also assume $\underline{r} \geq 4R_0$. By definition, if σ_0 is sufficiently small, we have

$$(4.5) \quad \lambda(\hat{M}_{-4} \cap B_{R_0}(x_0)) \leq 1 + \frac{\varepsilon}{2}$$

for $B_{R_0}(x_0) \subseteq \overline{B_{\underline{r}}} \setminus B_{\underline{r}/2}$. Combining (4.4) and (4.5), we obtain

$$(4.6) \quad (16\pi)^{-\frac{n}{2}} \int_{\hat{M}_{-4}} e^{-\frac{|x-x_0|^2}{16}} \leq 1 + \varepsilon.$$

Therefore, Theorem 4.3 implies

$$\sup_{\hat{M}_t \cap B_1(x_0)} |A|^2 \leq \frac{C(\varepsilon)}{4} \leq \delta_0$$

for $t \in [-1, 0]$ and $x_0 \in \overline{B_{\underline{r}-R_0}} \setminus B_{\underline{r}/2+R_0}$. As a result, we can take a universal constant $\delta_0 = \delta_0(n)$ (such that, for example, $n\delta_0 \leq \frac{1}{4}$) and assume $\hat{M}_{-1} \cap (\overline{B_{\underline{r}-R_0-1}} \setminus B_{\underline{r}/2+R_0+1})$ stays in $\overline{B_{\underline{r}-R_0-1}} \setminus B_{\underline{r}/2+R_0+1}$ under the MCF in the time interval $[-1, 0)$. Thus, by applying the Shi-type estimate [AB10, Lemma 3], we derive

$$(4.7) \quad |\nabla^k A_{\hat{M}_t}| \leq C(M_{-2}, \lambda_0, w) \text{ on } \hat{M}_t \cap \overline{B_{\underline{r}-R_0-1}}$$

for $t \in [-1+w, 0)$ and $k = 0, \dots, l-1$.

Now we follow the ideas in [CS21] to prove the proposition. We may assume (4.2) is true for $\tau \in [-2, 1]$ by taking ε small enough. On the other hand, given $\tau \in [1, \bar{\tau})$ and

$$x \in M_\tau \cap \left(\overline{B_{e^{\frac{\tau}{2}}(x-R_0-1)}} \setminus B_{\underline{r}} \right),$$

consider

$$\tau_0 := \tau + 2 \log \frac{\underline{r} - R_0 - 1}{|x|} \in [0, \tau).$$

Then,

$$t := -e^{\tau_0 - \tau} = -\frac{(\underline{r} - R_0 - 1)^2}{|x|^2} \in [-1+w, 0)$$

where $w := 1 - \left(1 - \frac{R_0-1}{\underline{r}}\right)^2 \in (0, 1)$. After rescaling, we get

$$\hat{x} := \sqrt{-t}x \in \hat{M}_t \cap \partial B_{\underline{r}-R_0-1}.$$

Consequently, the curvature estimate (4.7) implies

$$|\hat{x}|^{-1-k}|\hat{x}|^{1+k}|\nabla^k A_{\hat{M}_t}(x)| = |\nabla^k A_{\hat{M}_t}(x)| \leq C = C(M_{-2}, \lambda_0, w).$$

Rescaling back, we have

$$|x|^{1+k}|\nabla^k A_{M_\tau}(x)| \leq C(\underline{r} - R_0 + 1)^{1+k}.$$

Finally, choosing $C_l = (M_{-2}, \lambda_0, w)$ sufficiently large in (2) of definition 3.7, we have

$$\tilde{\mathbf{r}}_l(M_\tau) \geq e^{\frac{\tau}{2}}(\underline{r} - R_0 - 1) \geq \frac{1}{2}e^{\frac{\tau}{2}}$$

by our choices of R_0 . □

With the Lojasiewicz-Simon inequality (Theorem 3.9) and the extension result (Proposition 4.1), we are able to prove the higher-codimensional version of the theorem proven by Chodosh-Schulze.

Proof of Theorem 0.1. After establishing Theorem 3.9 and Proposition 4.1, we can follow the arguments in [CS21, Section 9], which are standard after having a Lojasiewicz-type inequality, verbatim to get the uniqueness of asymptotically conical tangent flows. □

REFERENCES

- [AA81] W. K. Allard and F. J. Almgren Jr., *On the radial behavior of minimal surfaces and the uniqueness of their tangent cones*, Ann. of Math. (2) **113** (1981), no. 2, 215–265. MR607893
- [AB10] B. Andrews and C. Baker, *Mean curvature flow of pinched submanifolds to spheres*, J. Differential Geom. **85** (2010), no. 3, 357–395. MR2739807
- [ALW14] B. Andrews, H. Li, and Y. Wei, *\mathcal{F} -stability for self-shrinking solutions to mean curvature flow*, Asian J. Math. **18** (2014), no. 5, 757–777. MR3287002
- [AS13] C. Arezzo and J. Sun, *Self-shrinkers for the mean curvature flow in arbitrary codimension*, Math. Z. **274** (2013), no. 3-4, 993–1027. MR3078255
- [Bra78] K. A. Brakke, *The motion of a surface by its mean curvature*, Mathematical Notes, vol. 20, Princeton University Press, Princeton, N.J., 1978. MR485012
- [Bra69] A. Brandt, *Interior Schauder estimates for parabolic differential- (or difference-) equations via the maximum principle*, Israel J. Math. **7** (1969), 254–262. MR249803
- [CHS84] L. Caffarelli, R. Hardt, and L. Simon, *Minimal surfaces with isolated singularities*, Manuscripta Math. **48** (1984), no. 1-3, 1–18, DOI 10.1007/BF01168999. MR753722
- [CS21] O. Chodosh and F. Schulze, *Uniqueness of asymptotically conical tangent flows*, Duke Math. J. **170** (2021), no. 16, 3601–3657. MR4332673
- [CIM15] T. H. Colding, T. Ilmanen, and W. P. Minicozzi II, *Rigidity of generic singularities of mean curvature flow*, Publ. Math. Inst. Hautes Études Sci. **121** (2015), 363–382. MR3349836
- [CM12] T. H. Colding and W. P. Minicozzi II, *Generic mean curvature flow I: generic singularities*, Ann. of Math. (2) **175** (2012), no. 2, 755–833. MR2993752
- [CM14] ———, *On uniqueness of tangent cones for Einstein manifolds*, Invent. Math. **196** (2014), no. 3, 515–588. MR3211041
- [CM15] ———, *Uniqueness of blowups and Lojasiewicz inequalities*, Ann. of Math. (2) **182** (2015), no. 1, 221–285. MR3374960
- [CM19] ———, *Regularity of elliptic and parabolic systems* (2019). Preprint available at <https://arxiv.org/abs/1905.00085>.
- [CM20] ———, *Complexity of parabolic systems*, Publ. Math. Inst. Hautes Études Sci. **132** (2020), 83–135. MR4179832

- [CM21] ———, *Singularities of Ricci flow and diffeomorphisms* (2021). Preprint available at <https://arxiv.org/abs/2109.06240>.
- [Eck00] K. Ecker, *Logarithmic Sobolev inequalities on submanifolds of Euclidean space*, J. Reine Angew. Math. **522** (2000), 105–118. MR1758578
- [Har97] R. M. Hardt, *Singularities of harmonic maps*, Bull. Amer. Math. Soc. (N.S.) **34** (1997), no. 1, 15–34. MR1397098
- [Hui84] G. Huisken, *Flow by mean curvature of convex surfaces into spheres*, J. Differential Geom. **20** (1984), no. 1, 237–266. MR772132
- [Hui90] ———, *Asymptotic behavior for singularities of the mean curvature flow*, J. Differential Geom. **31** (1990), no. 1, 285–299. MR1030675
- [Ilm95] T. Ilmanen, *Singularities of Mean Curvature Flow of Surfaces* (1995). Preprint available at <http://www.math.ethz.ch/~ilmanen/papers/pub.html>.
- [KKM18] N. Kapouleas, S. J. Kleene, and N. M. Møller, *Mean curvature self-shrinkers of high genus: non-compact examples*, J. Reine Angew. Math. **739** (2018), 1–39. MR3808256
- [Kha20] I. Khan, *Uniqueness of Asymptotically Conical Higher Codimension Self-Shrinkers and Self-Expanders* (2020). Preprint available at <https://arxiv.org/abs/2005.07611>.
- [LL15] Y.-I. Lee and Y.-K. Lue, *The stability of self-shrinkers of mean curvature flow in higher co-dimension*, Trans. Amer. Math. Soc. **367** (2015), no. 4, 2411–2435. MR3301868
- [Loj65] S. Lojasiewicz, *Ensembles semi-analytiques* (1965). Lectures Notes, Institut des Hautes Études Scientifiques.
- [LSS22] J. D. Lotay, F. Schulze, and G. Székelyhidi, *Neck pinches along the Lagrangian mean curvature flow of surfaces* (2022). Preprint available at <https://arxiv.org/abs/2208.11054>.
- [Sch14] F. Schulze, *Uniqueness of compact tangent flows in mean curvature flow*, J. Reine Angew. Math. **690** (2014), 163–172. MR3200339
- [Sim83] L. Simon, *Asymptotics for a class of nonlinear evolution equations, with applications to geometric problems*, Ann. of Math. (2) **118** (1983), no. 3, 525–571. MR727703
- [Sim96] ———, *Theorems on regularity and singularity of energy minimizing maps*, Lectures in Mathematics ETH Zürich, Birkhäuser Verlag, Basel, 1996. Based on lecture notes by Norbert Hungerbühler. MR1399562
- [Sim08] ———, *A general asymptotic decay lemma for elliptic problems*, Handbook of geometric analysis. No. 1, Adv. Lect. Math. (ALM), vol. 7, Int. Press, Somerville, MA, 2008, pp. 381–411. MR2483370
- [Wan14] L. Wang, *Uniqueness of self-similar shrinkers with asymptotically conical ends*, J. Amer. Math. Soc. **27** (2014), no. 3, 613–638. MR3194490
- [Whi94] B. White, *Partial regularity of mean-convex hypersurfaces flowing by mean curvature*, Internat. Math. Res. Notices **4** (1994), 185–192. MR1266114
- [Whi05] ———, *A local regularity theorem for mean curvature flow*, Ann. of Math. (2) **161** (2005), no. 3, 1487–1519. MR2180405
- [Zhu20] J. J. Zhu, *Lojasiewicz inequalities, uniqueness and rigidity for cylindrical self-shrinkers* (2020). Preprint available at <https://arxiv.org/abs/2011.01633>.

MIT, DEPT. OF MATH., 77 MASSACHUSETTS AVENUE, CAMBRIDGE, MA 02139-4307.

Email address: tangkai@mit.edu and xrzhao@mit.edu