

MOLLIFIED MOMENTS OF CUBIC DIRICHLET L-FUNCTIONS OVER THE EISENSTEIN FIELD

AHMET M. GÜLOĞLU AND HAMZA YESILYURT

ABSTRACT. We prove, assuming the generalized Riemann Hypothesis (GRH) that the density of the L -functions associated with primitive cubic Dirichlet characters over the Eisenstein field that do not vanish at the central point $s = 1/2$ is positive. This is achieved by computing the first mollified moment assuming a subconvexity bound, and obtaining a sharp upper bound for the higher mollified moments for these L -functions under GRH. The proportion of non-vanishing is explicit, but extremely small.

1. INTRODUCTION

S. Chowla conjectured in [4] that, for any real non-principal Dirichlet character χ , $L(1/2, \chi) \neq 0$. Li [18] showed that over the rational function field, there are infinitely many quadratic Dirichlet L -functions such that $L(1/2, \chi) = 0$. It is believed, however, that the number of such characters should be of density zero among all quadratic characters. For quadratic Dirichlet L -functions, Özlük and Snyder [20] showed, under the Generalized Riemann Hypothesis (GRH), that at least $15/16$ of the L -functions attached to quadratic characters do not vanish, by computing the one-level density for the low-lying zeroes in the family. The conjectures of Katz and Sarnak [16] imply that $L(1/2, \chi) \neq 0$ for almost all quadratic Dirichlet L -functions. By computing the first two mollified moments Soundararajan [25] proved without assuming GRH that at least 87.5% of the quadratic Dirichlet L -functions do not vanish at $s = 1/2$. It is well-known that using the first two (non-mollified) moments does not lead to a positive proportion of non-vanishing, as they grow too fast (see Conjecture 1.5.3 in [5] and the work of Jutila [15]). In the function field case, Bui and Florea [2] obtained a proportion of non-vanishing of at least 94% for quadratic Dirichlet L -functions by computing the one-level density.

In this paper, we consider the family of L -functions attached to primitive cubic Dirichlet characters defined over the Eisenstein field $K = \mathbb{Q}(\omega)$, where $\omega = e^{2\pi i/3}$. Many new conceptual and technical difficulties appear when considering cubic characters, and the results in the literature are fewer and weaker. Several authors studied the one level density for families of cubic characters over both \mathbb{Q} and K for test functions whose Fourier transforms are supported in $(-1, 1)$. These families have unitary symmetry. For a family with unitary symmetry, one needs to go beyond $(-1, 1)$ to obtain a positive proportion of non-vanishing. This was achieved for a thin subfamily of cubic characters over K by David and the first author [9], where the test functions are supported in $(-13/11, 13/11)$, which in turn gives $2/13$ as

Key words and phrases. Non-vanishing, Mollified Moments, Cubic Dirichlet characters, Cubic Gauss sums, Hecke L -functions.

The first author is supported by TÜBİTAK Research Grant no. 119F413.

the proportion of non-vanishing. These results are obtained under GRH. As for the moments, the first moment of $L(1/2, \chi)$, where χ is a primitive cubic character, was computed by Baier and Young [1] over \mathbb{Q} , by Luo [17] for the same thin sub-family as in [9] over K , and by David, Florea, and Lalin [6] over function fields. In these three papers, the authors obtained lower bounds for the number of non-vanishing cubic twists, but not positive proportions, by using upper bounds on higher moments. The lower bound in [6] was later improved by Ellenberg, Li, and Shusterman in [10]. Obtaining an asymptotic for the second moment for cubic Dirichlet L -functions is still an open question over functions fields or number fields. Recently, David, Florea and Lalin [7] proved using mollified moments of L -functions that there is a positive proportion of non-vanishing for cubic Dirichlet L -functions at $s = 1/2$ over function fields in the non-Kummer case. This is obtained by computing the first mollified moment, and by finding a sharp upper bound for the second mollified moment. Using the same approach, we prove the following result.

Theorem 1.1. *Let \mathcal{F} be the family of primitive cubic Dirichlet characters defined in (4). Then, assuming GRH for the L -functions in \mathcal{F} , for $X \rightarrow \infty$,*

$$\lim_{X \rightarrow \infty} \frac{\#\{\chi \in \mathcal{F} : L(1/2, \chi) \neq 0, n_\chi \leq X\}}{\#\{\chi \in \mathcal{F} : n_\chi \leq X\}} \geq \frac{3e^{-e^{101.3}}}{(\sqrt{3}-1)^2},$$

where n_χ denotes the norm of the conductor of χ .

As in [7], our positive proportion of non-vanishing is extremely small, since the implied constant in the upper bound for the second moment is too large.

1.1. Statement of Results. In what follows, we shall write $\mathcal{F}(X)$ to denote characters $\chi \in \mathcal{F}$ with $n_\chi \leq X$. Our first result gives an upper bound for the k^{th} mollified moment of the L -functions for any real $k > 0$.

Theorem 1.2. *Let $k, \kappa > 0$ be real numbers such that $k\kappa$ is an integer bounded by an absolute constant. Let \mathcal{F} be the family of primitive cubic Dirichlet characters defined in (4). Then, assuming GRH for the L -functions in \mathcal{F} , the bound*

$$(1) \quad \sum_{\chi \in \mathcal{F}(X)} |L(1/2, \chi)|^k |M(\chi, \kappa)|^{k\kappa} \ll_{k, \kappa} |\mathcal{F}(X)|$$

holds for X sufficiently large in terms of the parameters related to the mollifiers $M(\chi, \kappa)$ defined in section 3.1.

The proof of this theorem is given in section 4.1. As in [7], we start with an upper bound for $\log |L(s, \chi)|$ given in Lemma 3.1. The main contribution to (1) comes from prime ideals and their squares that appear in this upper bound. Most of section 4 deals with the former, which also leads to the definition of the mollifiers given in section 3.1. The contribution of squares of primes is dealt with separately in section 4.7.

Our next result gives an asymptotic formula for the first mollified moment of the family.

Theorem 1.3. *Let \mathcal{F} be the family defined in (4). Assume that for some ς with $0 \leq \varsigma < \frac{1}{14}$, the subconvexity bound*

$$(2) \quad L(1/2 + it, \Psi) \ll_\varepsilon ((1 + t^2)n_\Psi)^{\varsigma/2 + \varepsilon}$$

holds for $\Psi = \psi\chi$ and $\Psi = \psi$ for all $\chi \in \mathcal{F}$ and all ray class characters ψ modulo 9. Then,

$$(3) \quad \sum_{\chi \in \mathcal{F}(X)} L(1/2, \chi)M(\chi, 1) = C_X X \log X + O(X),$$

where C_X given in (79) converges to a constant as $X \rightarrow \infty$.

Using the definition of the mollifier $M(\chi, 1)$ in (3) leads to the twisted sum (57) of $L(1/2, \chi)$ over the family of characters χ . Then, using approximate functional equation for each L -function $L(s, \chi)$ gives a character sum over the family and a dual sum of shifted cubic Gauss sums given by (59). The former is split into two sums S_1 that runs over the cubes and the sum S_2 of remaining terms. The first sum S_1 gives the main term in (3) and is handled by using Lemma 2.8 in section 5.3. The second sum S_2 is estimated using Lemma 2.9, while the third sum S_3 is estimated in Lemma 5.4, where we use the deep work of Kubota and Patterson on the average of cubic Gauss sums (see Lemma 2.5).

2. PRELIMINARY

2.1. Notation. For $n \geq 2$, we put $\log_n X = \log(\log_{n-1} X)$, where $\log_1 X = \log X$. We use $N\mathfrak{a}$ for the norm of the ideal \mathfrak{a} , and $N(a)$ for the norm of the ideal $a\mathbb{Z}[\omega]$. We use $e(z) = \exp(2\pi iz)$ and put $\text{Tr}(z) = z + \bar{z}$. We use $\mathfrak{p}, \mathfrak{q}$ for prime ideals, and write $\mathfrak{a} \in \mathcal{I}$ to mean that $\forall \mathfrak{p}, \mathfrak{p} \mid \mathfrak{a} \Rightarrow N\mathfrak{p} \in \mathcal{I}$ for an interval \mathcal{I} .

The following lemma is used to find optimal bounds in the proofs of several results throughout the paper.

Lemma 2.1 ([11, Lemma 2.4]). *Suppose that*

$$L(H) = \sum_{i=1}^m A_i H^{a_i} + \sum_{j=1}^n B_j H^{-b_j},$$

where A_i, B_j, a_i, b_j are positive, and that $H_1 \leq H_2$. Then, there is some H with $H_1 \leq H \leq H_2$ such that

$$L(H) \ll \sum_{i=1}^m \sum_{j=1}^n \left(A_i^{b_j} B_j^{a_i} \right)^{1/(a_i+b_j)} + \sum_{i=1}^m A_i H_1^{a_i} + \sum_{j=1}^n B_j H_2^{-b_j},$$

where the implied constant depends only on m and n .

2.2. Cubic Characters. The ring of integers $\mathbb{Z}[\omega]$ of K has class number one and six units $\{\pm 1, \pm\omega, \pm\omega^2\}$. Each non-trivial principal ideal \mathfrak{n} co-prime to 3 has a unique generator $n \equiv 1 \pmod{3}$.

The cubic Dirichlet characters on $\mathbb{Z}[\omega]$ are given by the cubic residue symbols. For each prime $\pi \in \mathbb{Z}[\omega]$ with $\pi \equiv 1 \pmod{3}$, there are two primitive characters of conductor (π) ; the cubic residue character χ_π satisfying

$$\chi_\pi(\alpha) = \left(\frac{\alpha}{\pi} \right)_3 \equiv \alpha^{(N(\pi)-1)/3} \pmod{\pi\mathbb{Z}[\omega]},$$

and its conjugate $\bar{\chi}_\pi = \chi_\pi^2$.

In general, for $n \in \mathbb{Z}[\omega]$ with $n \equiv 1 \pmod{3}$, the cubic residue symbol χ_n is defined multiplicatively using the characters of prime conductor by

$$\chi_n(\alpha) = \left(\frac{\alpha}{n} \right)_3 = \prod_{\pi^{v_\pi} \parallel n} \chi_\pi(\alpha)^{v_\pi}.$$

Such a character χ_n is primitive when it is a product of characters of distinct prime conductors, i.e. either χ_π or $\overline{\chi}_\pi = \chi_\pi^2 = \chi_{\pi^2}$. Moreover, χ_n is a (cubic) Hecke character of conductor $n\mathbb{Z}[\omega]$ if $\chi_n(\omega) = 1$. Since

$$\left(\frac{\omega}{n}\right)_3 = \prod_{\pi|n} \omega^{v_\pi(n)(N(\pi)-1)/3} = \omega^{\sum_{\pi|n} v_\pi(n)(N(\pi)-1)/3} = \omega^{(N(n)-1)/3},$$

we conclude that a given Dirichlet character χ is a primitive cubic Hecke character provided that $\chi = \chi_n$, where

- (1) $n = n_1 n_2^2$, where n_1, n_2 are square-free and co-prime, and
- (2) $N(n) \equiv 1 \pmod{9}$, or equivalently, $N(n_1) \equiv N(n_2) \pmod{9}$.

In this case, χ has conductor $n_1 n_2 \mathbb{Z}[\omega]$.

We recall the cubic reciprocity theorem for cubic characters.

Lemma 2.2 ([13, page 114, Theorem 1]). *Let $m, n \in \mathbb{Z}[\omega]$, $m, n \equiv \pm 1 \pmod{3}$. Then,*

$$\left(\frac{m}{n}\right)_3 = \left(\frac{n}{m}\right)_3.$$

2.3. The family \mathcal{F} of cubic Dirichlet characters. We shall consider the family \mathcal{F} of cubic characters $\chi_{c_2 \overline{\chi}_{c_1}}$ given by cubic residue symbols, where $c_1, c_2 \equiv 1 \pmod{3} \in \mathbb{Z}[\omega]$ are square-free and co-prime, and $c_2 c_1^2 \equiv 1 \pmod{9}$. We naturally exclude the case $c_1 = c_2 = 1$. By a slight abuse of notation (dropping the letter χ), we shall write

$$(4) \quad \begin{aligned} \mathcal{F} &= \left\{ c_2 c_1^2 \in \mathbb{Z}[\omega] \setminus \{1\} : \begin{array}{l} c_1, c_2 \equiv 1 \pmod{3} \text{ both square-free,} \\ (c_1, c_2) = 1, c_2 c_1^2 \equiv 1 \pmod{9} \end{array} \right\} \\ &= \left\{ qd \in \mathbb{Z}[\omega] \setminus \{1\} : \begin{array}{l} q, d \equiv 1 \pmod{3}, q \text{ square-free,} \\ d \mid q, qd \equiv 1 \pmod{9} \end{array} \right\}. \end{aligned}$$

Note that for $c = qd \in \mathcal{F}$, the conductor of the character χ_{qd} is $q\mathbb{Z}[\omega]$. We shall write n_c in short, instead of n_{χ_c} , and $c \in \mathcal{F}(X)$ to mean that $c \in \mathcal{F}$ and $n_c \leq X$.

2.4. Cubic Gauss Sums. For any $n \equiv 1 \pmod{3}$, the shifted cubic Gauss sum is defined by

$$g(r, n) = \sum_{\alpha \pmod{n}} \chi_n(\alpha) e(\text{Tr}(r\alpha/n)).$$

The following two lemmas are classical results about cubic Gauss sums which can be found in [12], or easily checked, and we include them without proof.

Lemma 2.3. *Let $n, n_1, n_2 \equiv 1 \pmod{3}$ and s, r be elements of $\mathbb{Z}[\omega]$.*

If $(s, n) = 1$,

$$g(rs, n) = \overline{\chi_n}(s) g(r, n).$$

If $(n_1, n_2) = 1$,

$$g(r, n_1 n_2) = \overline{\chi_{n_2}(n_1)} g(r, n_1) g(r, n_2) = g(r n_1, n_2) g(r, n_1).$$

Lemma 2.4. *Let $\pi \equiv 1 \pmod{3}$ be a prime, $(\pi, r) = 1$, where $r \equiv 1 \pmod{3}$. Let k, j be integers with $k > 0$ and $j \geq 0$.*

If $k = j + 1$,

$$g(r\pi^j, \pi^k) = N(\pi^j) \times \begin{cases} -1 & \text{if } 3 \mid k \\ g(r, \pi) & \text{if } k \equiv 1 \pmod{3} \\ \overline{g(r, \pi)} & \text{if } k \equiv 2 \pmod{3}. \end{cases}$$

If $k \neq j + 1$,

$$g(r\pi^j, \pi^k) = \begin{cases} \varphi_K(\pi^k) & \text{if } 3 \mid k, k \leq j \\ 0 & \text{otherwise.} \end{cases}$$

For any character λ , we put $g_\lambda(r, b) = \lambda(b)g(r, b)$ and define

$$h(r, \lambda, s) = \sum_{\substack{b \equiv 1 \pmod{3} \\ (b, r) = 1}} g_\lambda(r, b) N(b)^{-s},$$

The following lemma contains the analytic information on the generating function $h(r, \lambda, s)$.

Lemma 2.5 ([9][Lemma 6.1]). *Write $r = r_1 r_2^2 r_3^3$, where r_1, r_2 are square-free and co-prime, and $r_i \equiv 1 \pmod{3}$. Then, $h(r, \lambda, s + 1/2)$ can be meromorphically continued to the whole complex plane; it is entire for $\Re(s) > 1/2$ except possibly for a simple pole at $s = 5/6$ with residue*

$$\operatorname{Res}_{s=5/6} h(r, \lambda, s + \frac{1}{2}) \ll N(r_1)^{-1/6} \log 2N(r_1) \log 2N(r_3^*)$$

when $r_2 = 1$ and λ^3 is not the principal character, and zero otherwise.

Let $\varepsilon > 0$ and $\sigma_1 = 1 + \varepsilon$. For $s = \sigma + it$ satisfying $\sigma_1 - 1/2 \leq \sigma \leq \sigma_1$ and $|s - 5/6| > 1/12$,

$$h(r, \lambda, s + \frac{1}{2}) \ll N(r_1 r_2^2)^{\frac{1}{2}(\sigma_1 - \sigma)} N(r_1 r_3^*)^\varepsilon (1 + t^2)^{\sigma_1 - \sigma}.$$

2.5. Hecke L -functions. Let χ be a primitive Hecke character to some modulus $\mathfrak{m} = m\mathbb{Z}[\omega]$ of trivial infinite type. The Hecke L -function associated with χ is given by

$$L(s, \chi) = \sum_{\mathfrak{a}} \chi(\mathfrak{a}) (N\mathfrak{a})^{-s}$$

where \mathfrak{a} runs over the integral ideals of $\mathbb{Z}[\omega]$ and we put $\chi(\mathfrak{a}) = 0$ whenever \mathfrak{a} is not coprime to \mathfrak{m} . The completed Hecke L -series is then defined by

$$\Lambda(s, \chi) = (|d_K| N(m))^{s/2} (2\pi)^{-s} \Gamma(s) L(s, \chi),$$

where $d_K = -3$ is the discriminant of K .

Proposition 2.6 ([19, VII. Cor. 8.6]). *The completed L -series $\Lambda(s, \chi)$ is entire, provided χ is primitive and $\mathfrak{m} \neq \mathbb{Z}[\omega]$. Furthermore, it satisfies the functional equation*

$$\Lambda(s, \chi) = W(\chi) (Nm)^{-1/2} \Lambda(1 - s, \overline{\chi}),$$

where

$$W(\chi) = \sum_{\substack{x \pmod{\mathfrak{m}} \\ (x, \mathfrak{m}) = 1}} \chi(x) e(\operatorname{Tr}(x/m\sqrt{-3}))$$

is the associated cubic Gauss sum, and x varies over a system of representatives of $(\mathbb{Z}[\omega]/\mathfrak{m})^\times$.

2.6. Dedekind Zeta Function and the prime ideal theorem. Note that

$$\zeta_K(s) = \zeta(s) \prod_{p \neq 3} \left(1 - \left(\frac{-3}{p}\right) p^{-s}\right)^{-1} = \zeta(s) L(s, \left(\frac{-3}{\cdot}\right)).$$

We know by the class number formula of Dirichlet that

$$(5) \quad L(1, \left(\frac{-3}{\cdot}\right)) = \frac{2\pi h(-3)}{\omega_K \sqrt{|\mathfrak{d}_K|}} = \frac{\pi}{3\sqrt{3}},$$

which gives the residue of $\zeta_K(s)$ at $s = 1$.

By [24, Theorem 2] and [26] it follows that there exist positive absolute constants c_1, c_2 such that

$$(6) \quad \pi_K(x) = \text{Li}(x) + O(x \exp(-c_1 \sqrt{\log x}))$$

for $x \geq c_2$, where $\pi_K(x)$ denotes the number of prime ideals of $\mathbb{Z}[\omega]$ whose norm does not exceed x .

2.7. Family Related Results.

Lemma 2.7 ([8, Chapter 17, 1st Lemma]). *Let*

$$\delta(y) = \begin{cases} 0 & \text{if } 0 < y < 1 \\ 1/2 & \text{if } y = 1 \\ 1 & \text{if } y > 1. \end{cases}$$

For any $y > 0, a > 0, T > 0$,

$$\left| \frac{1}{2\pi i} \int_{a-iT}^{a+iT} \frac{y^s}{s} ds - \delta(y) \right| < \begin{cases} y^a \min\{1, T^{-1} |\log y|^{-1}\} & \text{if } y \neq 1 \\ aT^{-1} & \text{if } y = 1. \end{cases}$$

Lemma 2.8. *Assume that the subconvexity bound*

$$(7) \quad L(1/2 + it, \psi) \ll_\varepsilon ((1 + t^2)n_\psi)^{\varsigma/2+\varepsilon}$$

holds for some $0 \leq \varsigma \leq 1/2$, and for all ray class characters ψ modulo 9. Then, for any $n \in \mathbb{Z}[\omega]$,

$$\sum_{\substack{c \in \mathcal{F}(X) \\ (c, n) = 1}} 1 = C_1(n) X \log X + C_2(n) X + O_\varepsilon \left((X^{\frac{4\varsigma+1}{4\varsigma+2}+5\varepsilon} + X^{1/2+5\varepsilon}) (N(n))^\varepsilon \right),$$

where

$$(8) \quad \begin{aligned} C_1(n) &= \frac{4\pi^2 F_{\psi_0}(1; n)}{2187} \\ C_2(n) &= \frac{4F_{\psi_0}(1; n)}{81} \left\{ \frac{\pi^2 \log(3/e)}{27} + \lim_{s \rightarrow 1} ((s-1)^2 \zeta_K^2(s))' \right\} + \frac{4\pi^2}{2187} F'_{\psi_0}(1; n) \end{aligned}$$

and $F_\psi(s; n)$ is defined in (10).

Proof. Using Lemma 2.7 and proceeding similar to [8, Chapter 17], we can show for $T \geq 1$ and $\rho = 1 + 1/(2 \log X)$ that

$$\sum_{\substack{c \in \mathcal{F}(X) \\ (c, n) = 1}} 1 = \frac{1}{2\pi i} \int_{\rho-iT}^{\rho+iT} X^s \left(\sum_{\substack{c \in \mathcal{F} \cup \{1\} \\ (c, n) = 1}} \frac{1}{n_c^s} \right) \frac{ds}{s} + O(X^\varepsilon + X^{1+2\varepsilon} T^{-1}).$$

Note that by the definition of the family \mathcal{F} (see equation (4)) we have

$$\begin{aligned} \sum_{\substack{c \in \mathcal{F} \cup \{1\} \\ (c,n)=1}} \frac{1}{n^s} &= \sum_{\substack{q \equiv 1 \pmod{3} \\ q \text{ square-free} \\ (q,n)=1}} \frac{1}{N(q)^s} \sum_{\substack{d \equiv 1 \pmod{3} \\ d|q \\ qd \equiv 1 \pmod{9}}} 1 \\ &= \frac{1}{h_{(9)}} \sum_{\psi \pmod{9}} G_\psi(s; n), \end{aligned}$$

where ψ runs over the ray class characters modulo 9, $h_{(9)} = 9$, and for any ray class character Ψ of conductor not co-prime to 3, and for any $\alpha \in \mathbb{Z}[\omega]$,

$$(9) \quad G_\Psi(s; \alpha) = \prod_{\substack{\pi \equiv 1 \pmod{3} \\ \pi \nmid \alpha}} \left(1 + \frac{\Psi(\pi)}{N(\pi)^s} (1 + \Psi(\pi)) \right) = L(s, \Psi) L(s, \Psi^2) F_\Psi(s; \alpha),$$

where

$$L(s, \Psi) = \prod_{\pi \equiv 1 \pmod{3}} \left(1 - \frac{\Psi(\pi)}{N(\pi)^s} \right)^{-1} = \sum_{\mathfrak{a}} \frac{\Psi(\mathfrak{a})}{(N\mathfrak{a})^s} \quad (\text{since } \Psi(1 - \omega) = 0)$$

is the Hecke L-function associated with the ray class character Ψ , and

$$(10) \quad \begin{aligned} F_\Psi(s; \alpha) &= \prod_{\substack{\pi \equiv 1 \pmod{3} \\ \pi \nmid \alpha}} \left(1 - \frac{\Psi^2(\pi) + \Psi^3(\pi) + \Psi^4(\pi)}{N(\pi)^{2s}} + \frac{\Psi^4(\pi) + \Psi^5(\pi)}{N(\pi)^{3s}} \right) \\ &\quad \cdot \prod_{\substack{\pi \equiv 1 \pmod{3} \\ \pi | \alpha}} \left(1 - \frac{\Psi(\pi)}{N(\pi)^s} \right) \left(1 - \frac{\Psi^2(\pi)}{N(\pi)^s} \right). \end{aligned}$$

Now, the integral above can be written as

$$\frac{1}{2\pi i h_{(9)}} \sum_{\psi \pmod{9}} \int_{\varrho - iT}^{\varrho + iT} X^s G_\psi(s; n) \frac{ds}{s}.$$

Note that since

$$|F_\Psi(s; \alpha)| \leq \prod_{\substack{\pi \equiv 1 \pmod{3} \\ \pi | \alpha}} \left(1 + \frac{1}{N(\pi)^\sigma} \right)^2 \exp \left(\sum_{\pi \equiv 1 \pmod{3}} \left(\frac{3}{N(\pi)^{2\sigma}} + \frac{2}{N(\pi)^{3\sigma}} \right) \right),$$

$F_\Psi(s; \alpha)$ converges absolutely for any $\alpha \in \mathbb{Z}[\omega]$ in the region $\text{Re } s \geq 1/2 + \varepsilon$, uniformly for any character Ψ , whereas $L(s, \Psi)$ and $L(s, \Psi^2)$ both have analytic continuation to entire functions except for a simple pole at $s = 1$ if the characters are principal. Therefore, we see that $G_\psi(s; \alpha)$ is analytic in the region $\text{Re } s \geq 1/2 + \varepsilon$, except for a double pole at $s = 1$ when ψ is principal, and a simple pole if ψ has order 2. Since $h_{(9)} = 9$, there is no character of order 2. Therefore, shifting the contour to $\text{Re } s = 1/2 + \varepsilon$, we pick up the residue at $s = 1$ when $\psi = \psi_0$, and using the crude estimate $F_\psi(s; n) \ll (N(n))^\varepsilon$ together with (7), we deduce that

$$\begin{aligned} \int_{1/2+\varepsilon}^{\varrho} X^s G_\psi(s; n) \frac{ds}{s} &\ll (X^{1/2+\varepsilon} T^{2\varsigma-1+4\varepsilon} + X T^{4\varepsilon-1}) (N(n))^\varepsilon \\ \int_{1/2+\varepsilon-iT}^{1/2+\varepsilon+iT} X^s G_\psi(s; n) \frac{ds}{s} &\ll X^{1/2+\varepsilon} T^{2\varsigma+4\varepsilon} (N(n))^\varepsilon. \end{aligned}$$

Combining all the estimates and applying Lemma 2.1 with $T \in [1, X]$, we get the claimed error term.

Since $L(s, \psi_0)L(s, \psi_0^2) = (1 - 3^{-s})^2 \zeta_K^2(s)$ has a double pole at $s = 1$, the residue is given by

$$\begin{aligned} & \lim_{s \rightarrow 1} (X^s s^{-1} F_{\psi_0}(s; \mathbf{m})(1 - 3^{-s})^2 (s - 1)^2 \zeta_K^2(s))' \\ &= \frac{4}{9} F_{\psi_0}(1; \mathbf{m}) (\operatorname{Res}_{s=1} \zeta_K(s))^2 X \log X \\ & \quad + \frac{4}{9} X \left(\log(3/e) F_{\psi_0}(1; \mathbf{m}) (\operatorname{Res}_{s=1} \zeta_K(s))^2 \right. \\ & \quad \left. + F'_{\psi_0}(1; \mathbf{m}) (\operatorname{Res}_{s=1} \zeta_K(s))^2 + F_{\psi_0}(1; \mathbf{m}) \lim_{s \rightarrow 1} ((s - 1)^2 \zeta_K^2(s))' \right). \end{aligned}$$

The desired result follows upon using (5). \square

Lemma 2.9. *Let χ be a non-principal cubic ray class character of conductor $\mathbf{m} = m\mathbb{Z}[\omega]$ with $(m, 3) = 1$. Assume that the sub-convexity bound (7) holds for some $0 \leq \varsigma \leq 1/2$, and each for $L(s, \psi\chi)$, where ψ varies over the ray class characters modulo 9. Then,*

$$\sum_{c \in \mathcal{F}(X)} \chi(c) \ll_{\varepsilon} X^{\frac{4\varsigma+1}{4\varsigma+2}+5\varepsilon} (n_{\chi})^{\frac{\varsigma}{2\varsigma+1}+3\varepsilon} + X^{1/2+\varepsilon} (n_{\chi})^{\varsigma+3\varepsilon}.$$

Proof. We only give a sketch of the proof as it is similar to that of Lemma 2.8. Using Perron's formula, we have as before for $T \geq 1$ that

$$\sum_{c \in \mathcal{F}(X)} \chi(c) = \frac{1}{18\pi i} \sum_{\psi \bmod 9} \int_{\varrho-iT}^{\varrho+iT} X^s G_{\psi\chi}(s; \mathbf{m}) \frac{ds}{s} + O\left(X^{\varepsilon} + X^{1+2\varepsilon} T^{-1}\right),$$

where $G_{\Psi}(s; \mathbf{m})$ was defined in (9). We see that $G_{\psi\chi}(s; \mathbf{m})$ is analytic in the region $\operatorname{Re} s \geq 1/2 + \varepsilon$ for each ψ , since χ is not principal and the conductors of ψ and χ are coprime. We shift the contour to $\operatorname{Re} s = 1/2 + \varepsilon$. Using the estimate $F_{\Psi}(s; \mathbf{m}) \ll (\mathbf{N}\mathbf{m})^{\varepsilon}$ together with (7), we deduce that

$$\begin{aligned} & \int_{1/2+\varepsilon}^{\varrho} X^s G_{\psi\chi}(s; \mathbf{m}) \frac{ds}{s} \ll X^{1/2+\varepsilon} T^{2\varsigma-1+4\varepsilon} (n_{\psi\chi})^{\varsigma+3\varepsilon} + XT^{4\varepsilon-1} (n_{\psi\chi})^{3\varepsilon} \\ & \int_{1/2+\varepsilon-iT}^{1/2+\varepsilon+iT} X^s G_{\psi\chi}(s; \mathbf{m}) \frac{ds}{s} \ll X^{1/2+\varepsilon} T^{2\varsigma+4\varepsilon} (n_{\psi\chi})^{\varsigma+3\varepsilon}. \end{aligned}$$

Combining all the estimates and applying Lemma 2.1 with $T \in [1, X]$, we get the claimed error term. \square

3. DEFINITION OF THE MOLLIFIERS

To define the mollifiers we shall need the following result first, the proof of which will be skipped as it is very similar to [3, Theorem 2.1].

Lemma 3.1. *Let $\lambda_0 \approx 0.491225$ be the unique number satisfying $e^{-\lambda_0} = \lambda_0 + \lambda_0^2/2$. Assuming GRH for $L(s, \chi)$, the inequality*

$$\begin{aligned} \log |L(1/2 + it, \chi)| \leq & \operatorname{Re} \sum_{1 < \mathbf{N}\mathbf{a} \leq x} \frac{\Lambda(\mathbf{a})\chi(\mathbf{a})}{\mathbf{N}(\mathbf{a})^{1/2+\lambda/\log x+it} \log \mathbf{N}\mathbf{a}} \frac{\log(x/\mathbf{N}\mathbf{a})}{\log x} \\ & + \frac{1+\lambda}{2\log x} \log \left(\frac{3n_\chi |1/2 + \lambda/\log x + it|^2}{4\pi^2} \right) \\ & + \frac{2e^{-\lambda}}{x^{1/2} \log^2 x |1/2 + \lambda/\log x + it|^2} \end{aligned}$$

holds for any $\lambda \geq \lambda_0$, any t which does not coincide with the ordinates of zeros of $L(s, \chi)$, and any $x \geq 3$.

3.1. The Mollifier. Let $k, \kappa > 0$ be real numbers such that $k\kappa$ is an integer bounded by an absolute constant. Put $k_0 = \max\{3, 4k^2\}$.

Definition 3.2. Let $\alpha > 1$ and $\Theta, \beta < 1$ be fixed real numbers satisfying

$$(11) \quad \alpha\beta > \frac{2k\kappa + 3}{2}.$$

Assume that X satisfies

$$(12) \quad (\log_2 X)^{\alpha\beta} > 2, \quad \alpha \log_3 X \geq 1 - \log \Theta.$$

Let $J = \lfloor \log(\Theta(\log_2 X)^\alpha) \rfloor$, and for each $0 \leq j \leq J$, put

$$\mathcal{I}_j = (X^{\theta_{j-1}}, X^{\theta_j}], \quad \ell_j = 2 \lfloor \theta_j^{-\beta} \rfloor,$$

where

$$\theta_{-1} = \frac{\log k_0}{\log X}, \quad \theta_j = \frac{e^j}{(\log_2 X)^\alpha} \quad (j \geq 0).$$

Note that ℓ_j is a positive even integer since each $\theta_j < 1$.

Lemma 3.3 ([21, Section 3, Lemma 1]). *For any integer $\ell \geq 0$, put*

$$E_\ell(x) = \sum_{0 \leq n \leq \ell} \frac{x^n}{n!}.$$

Then, $E_{2\ell}(x) > 0$ and is concave up. Furthermore,

$$\begin{aligned} e^x &< E_{2\ell}(x) & (x \leq 0) \\ e^x &< (1 + e^{-2\ell})E_{2\ell}(x) & (x \leq 2\ell/e^2). \end{aligned}$$

Remark 3.4. Note that for $x \geq \ell$,

$$E_\ell(x) \leq (\ell + 1) \frac{x^\ell}{\ell!}$$

as $x^n/n!$ is increasing for $0 \leq n \leq \ell$.

Definition 3.5. For $0 \leq j \leq J$, let

$$(13) \quad D_{j,k}(c) = \prod_{r=0}^j (1 + e^{-\ell_r}) E_{\ell_r}(k \operatorname{Re} F_r(c, j)),$$

and

$$S_{j,k}(c) = \exp\left(k \operatorname{Re} \sum_{k_0 < \mathfrak{N}\mathfrak{p} \leq X^{\theta_j/2}} \frac{\chi_c(\mathfrak{p}^2)}{2\mathfrak{N}(\mathfrak{p})^{1+2/(\theta_j \log X)}} \left(1 - \frac{2 \log \mathfrak{N}\mathfrak{p}}{\theta_j \log X}\right)\right),$$

where

$$(14) \quad F_r(c, j) = \sum_{\mathfrak{p} \in \mathcal{I}_r} \frac{\chi_c(\mathfrak{p}) f(\mathfrak{p}, j)}{\mathfrak{N}(\mathfrak{p})^{1/2}}, \quad f(\mathfrak{p}, j) = \frac{1}{(\mathfrak{N}\mathfrak{p})^{1/(\theta_j \log X)}} \left(1 - \frac{\log \mathfrak{N}\mathfrak{p}}{\theta_j \log X}\right).$$

Definition 3.6. For each $r = 0, \dots, J$, let

$$(15) \quad \mathcal{T}_r = \left\{c \in \mathcal{F}(X) : \max_{r \leq j \leq J} \operatorname{Re} F_r(c, j) \leq \frac{\ell_r}{ke^2}\right\}.$$

Lemma 3.7. For each $c \in \mathcal{F}(X)$, either $c \notin \mathcal{T}_0$ or

$$\begin{aligned} |L(1/2, \chi_c)|^k &\leq \sum_{\substack{0 \leq j < J \\ j < u \leq J}} \exp(k/\theta_j + O(kk_0)) D_{j,k}(c) S_{j,k}(c) \left(\frac{e^2 k \operatorname{Re} F_{j+1}(c, u)}{\ell_{j+1}}\right)^{s_{j+1}} \\ &\quad + \exp(k/\theta_J + O(kk_0)) D_{J,k}(c) S_{J,k}(c) \end{aligned}$$

for any positive even integers s_1, \dots, s_J .

Proof. Taking $t = 0$ and $\lambda = 1$ in Lemma 3.1 we have for any $x \geq 3$ and $c \in \mathcal{F}$ that

$$\begin{aligned} \log |L(1/2, \chi_c)| &< \operatorname{Re} \sum_{1 < \mathfrak{N}\mathfrak{a} \leq x} \frac{\Lambda(\mathfrak{a}) \chi_c(\mathfrak{a})}{\mathfrak{N}(\mathfrak{a})^{1/2+1/\log x} \log \mathfrak{N}\mathfrak{a}} \frac{\log(x/\mathfrak{N}\mathfrak{a})}{\log x} \\ &\quad + \frac{1}{\log x} \log\left(\frac{3n_c(1/2 + 1/\log 3)^2}{4\pi^2}\right) + \frac{8}{ex^{1/2} \log^2 x}. \end{aligned}$$

The contribution of powers of prime ideals \mathfrak{p}^n with $n \geq 3$ is

$$\leq \sum_{n \geq 3, (\mathfrak{N}\mathfrak{p})^n \leq x} \frac{1}{n(\mathfrak{N}\mathfrak{p})^{n(1/2+1/\log x)}} \frac{\log(x/(\mathfrak{N}\mathfrak{p})^n)}{\log x} < \sum_{\mathfrak{p}} (\mathfrak{N}\mathfrak{p})^{-3/2}.$$

Moreover, removing the terms, if any, with $3 \leq \mathfrak{N}\mathfrak{p} \leq k_0$ we end up with the inequality

$$(16) \quad \begin{aligned} \log |L(1/2, \chi_c)| &< \operatorname{Re} \left(\sum_{k_0 < \mathfrak{N}\mathfrak{p} \leq x} \frac{\chi_c(\mathfrak{p})}{\mathfrak{N}(\mathfrak{p})^{1/2+1/\log x}} \frac{\log(x/\mathfrak{N}\mathfrak{p})}{\log x} \right) \\ &\quad + \operatorname{Re} \left(\sum_{k_0 < \mathfrak{N}\mathfrak{p} \leq \sqrt{x}} \frac{\chi_c(\mathfrak{p}^2)}{2\mathfrak{N}(\mathfrak{p})^{1+2/\log x}} \frac{\log(x/\mathfrak{N}\mathfrak{p}^2)}{\log x} \right) + \frac{\log n_c}{\log x} + O(k_0). \end{aligned}$$

Now, assume that $c \in \mathcal{T}_0$. If $c \in \mathcal{T}_r$ for every $0 \leq r \leq J$, then we choose $x = X^{\theta_j}$ in (16), multiply both sides by k and exponentiate, then apply Lemma 3.3. This gives the last term in the claimed upper bound above. Otherwise, choose $x = X^{\theta_j}$ in (16), where $j+1 \geq 1$ is the smallest r for which $c \notin \mathcal{T}_r$. Note that $k \operatorname{Re} F_r(c, j) \leq \ell_r/e^2$ for each $0 \leq r \leq j$ since $c \in \mathcal{T}_r$ for each such r , and we can apply Lemma 3.3. Furthermore, since $c \notin \mathcal{T}_{j+1}$, we have $(e^2 k/\ell_{j+1}) \operatorname{Re} F_{j+1}(c, u) > 1$ for some $u = u(j) > j$. This gives the first term in the claim. \square

Definition 3.8 (The Mollifier). *For each $c \in \mathcal{F}(X)$, we define the mollifier by*

$$(17) \quad \begin{aligned} M(c, \kappa) &= \prod_{j=0}^J M_j(c, \kappa) \\ M_j(c, \kappa) &= E_{\ell_j} \left(-\frac{1}{\kappa} F_j(c, J) \right). \end{aligned}$$

Using the definition of $E_\ell(x)$, we see that

$$\begin{aligned} M_j(c, \kappa) &= \sum_{n=0}^{\ell_j} \frac{1}{n!} \left(-\frac{1}{\kappa} \sum_{\mathbf{p} \in \mathcal{I}_j} \frac{\chi_c(\mathbf{p})}{N(\mathbf{p})^{1/2}} f(\mathbf{p}, J) \right)^n \\ &= \sum_{n=0}^{\ell_j} \frac{1}{n!} \left(-\frac{1}{\kappa} \right)^n \sum_{\mathbf{p}_1, \dots, \mathbf{p}_n \in \mathcal{I}_j} \frac{\chi_c(\mathbf{p}_1 \cdots \mathbf{p}_n)}{N(\mathbf{p}_1 \cdots \mathbf{p}_n)^{1/2}} f(\mathbf{p}_1 \cdots \mathbf{p}_n, J) \\ &= \sum_{n=0}^{\ell_j} \left(-\frac{1}{\kappa} \right)^n \sum_{\substack{\mathbf{a} \in \mathcal{I}_j \\ \Omega(\mathbf{a})=n}} \frac{\chi_c(\mathbf{a})}{N(\mathbf{a})^{1/2}} f(\mathbf{a}, J) \nu(\mathbf{a}) \\ &= \sum_{\substack{\mathbf{a} \in \mathcal{I}_j \\ 0 \leq \Omega(\mathbf{a}) \leq \ell_j}} \frac{\chi_c(\mathbf{a}) \lambda(\mathbf{a}) f(\mathbf{a}, J) \nu(\mathbf{a})}{N(\mathbf{a})^{1/2} \kappa^{\Omega(\mathbf{a})}}, \end{aligned}$$

where we extended the definition of f to be a completely multiplicative function using (14), and

$$\lambda(\mathbf{a}) = (-1)^{\Omega(\mathbf{a})}, \quad \nu(\mathbf{p}_1^{r_1} \cdots \mathbf{p}_s^{r_s}) = \frac{1}{r_1! \cdots r_s!}.$$

Note that ν is multiplicative and λ is completely multiplicative. Furthermore, for any integer $n \geq 0$,

$$M_j(c, \kappa)^n = \sum_{\mathbf{a} \in \mathcal{I}_j} \frac{\chi_c(\mathbf{a}) \lambda(\mathbf{a}) f(\mathbf{a}, J)}{N(\mathbf{a})^{1/2} \kappa^{\Omega(\mathbf{a})}} \nu_n(\mathbf{a}; \ell_j)$$

where

$$\nu_n(\mathbf{a}; \ell_j) = \sum_{\substack{\mathbf{a}_1, \dots, \mathbf{a}_n \\ \mathbf{a}_1 \cdots \mathbf{a}_n = \mathbf{a} \\ 0 \leq \Omega(\mathbf{a}_i) \leq \ell_j}} \nu(\mathbf{a}_1) \cdots \nu(\mathbf{a}_n).$$

Remark 3.9. The inequality $\nu_n(\mathbf{a}; \ell_j) \leq \nu_n(\mathbf{a})$ holds with equality when $\Omega(\mathbf{a}) \leq \ell_j$, where $\nu_n(\cdot)$ is the n -fold convolution of ν . Furthermore,

$$\nu_n(\mathbf{p}^m) = \frac{n^m}{m!}.$$

4. UPPER BOUND FOR MOLLIFIED MOMENTS

We start with a simple preliminary result.

Lemma 4.1. *There exists a number $X_0(\Theta) > 0$ such that for $X \geq X_0$,*

$$\begin{aligned} \sum_{N\mathfrak{p} > k_0} \frac{1}{(N\mathfrak{p})^{3/2}} &< 1, \\ \sum_{\mathfrak{p} \in \mathcal{I}_r} \frac{1}{(N\mathfrak{p})^\sigma} &< \frac{2}{(\sigma-1) \lfloor X^{\theta_{r-1}} \rfloor^{\sigma-1}} \quad (\sigma > 1), \\ \sum_{\mathfrak{p} \in \mathcal{I}_0} \frac{1}{N\mathfrak{p}} &< 2 \log \log X^{\theta_0}, \\ \sum_{X^{\theta_j} < N\mathfrak{p} \leq X^{\theta_j}} \frac{1}{N\mathfrak{p}} &= J - j + O\left(\frac{1}{\log X^{\theta_j}}\right) \quad 0 \leq j \leq J-1, \\ \sum_{k_0 < N\mathfrak{p} \leq X^{\theta_j}} \frac{\log^2 N\mathfrak{p}}{N\mathfrak{p}} &< 2(\log X^{\theta_j})^2 \quad 0 \leq j \leq J. \end{aligned}$$

Proof.

$$\begin{aligned} \sum_{N\mathfrak{p} > k_0} \frac{1}{N(\mathfrak{p})^{3/2}} &< \sum_{3 < N\mathfrak{p} \leq 31} \frac{1}{N(\mathfrak{p})^{3/2}} + \sum_{N\mathfrak{p} > 31} \frac{1}{N(\mathfrak{p})^{3/2}} \\ &< \frac{1}{2^3} + \frac{1}{5^3} + \frac{2}{7^{3/2}} + \frac{1}{11^3} + \frac{2}{13^{3/2}} + \frac{1}{17^3} \\ &\quad + \frac{2}{19^{3/2}} + \frac{1}{23^3} + \frac{1}{29^3} + \frac{2}{31^{3/2}} + \frac{2}{3} < 1 \end{aligned}$$

since

$$\sum_{N\mathfrak{p} > 31} \frac{1}{N(\mathfrak{p})^{3/2}} < \sum_{p > 36} \frac{2}{p^{3/2}} < \sum_{n > 36} \frac{2}{n^{3/2}} < 2 \int_{36}^{\infty} x^{-3/2} dx = \frac{2}{3}.$$

For the second claim we have

$$\sum_{\mathfrak{p} \in \mathcal{I}_r} \frac{1}{(N\mathfrak{p})^\sigma} < 2 \sum_{n > \lfloor X^{\theta_{r-1}} \rfloor} \frac{1}{n^\sigma} < 2 \int_{\lfloor X^{\theta_{r-1}} \rfloor}^{\infty} x^{-\sigma} dx = \frac{2}{(\sigma-1) \lfloor X^{\theta_{r-1}} \rfloor^{\sigma-1}}.$$

Let $z = \sqrt{\log X^{\theta_0}} > k_0$. Then, for X large enough, using (6) implies

$$\begin{aligned} \sum_{\mathfrak{p} \in \mathcal{I}_0} \frac{1}{N\mathfrak{p}} &< \sum_{k_0 < p \leq z} \frac{2}{p} + \sum_{z < N\mathfrak{p} \leq X^{\theta_0}} \frac{1}{N\mathfrak{p}} < 2 \int_1^z \frac{dx}{x} + \log \left(\frac{\log X^{\theta_0}}{\log z} \right) + O\left(\frac{1}{\log z}\right) \\ &= 2 \log z + \log \left(\frac{\log X^{\theta_0}}{\log z} \right) + O\left(\frac{1}{\log z}\right) < 2 \log_2 X^{\theta_0}. \end{aligned}$$

The next claim follows using definition 3.2 and (6). For the last claim, choose $z = \exp((2^{-1} \log^2 X^{\theta_j})^{1/3})$. Then, it follows from (6) that for X large enough,

$$\begin{aligned} \sum_{k_0 < N\mathfrak{p} \leq X^{\theta_j}} \frac{\log^2 N\mathfrak{p}}{N\mathfrak{p}} &\leq 2 \sum_{k_0 < n \leq z} \frac{\log^2 n}{n} + \sum_{z < N\mathfrak{p} \leq X^{\theta_j}} \frac{\log^2 N\mathfrak{p}}{N\mathfrak{p}} \\ &\leq 2 \log^2 z \int_1^z \frac{dx}{x} + \log^2 X^{\theta_j} - \log^2 z + O\left(\frac{1}{\log^3 z}\right) \\ &< 2 \log^2 X^{\theta_j}. \end{aligned}$$

□

4.1. **Proof of Theorem 1.2.** We split the sum in (1) into the sums

$$(18) \quad \sum_{c \in \mathcal{F}(X) \cap \mathcal{T}_0} |L(1/2, \chi_c)|^k |M(c, \kappa)|^{k\kappa}$$

and

$$(19) \quad \sum_{c \in \mathcal{F}(X) \setminus \mathcal{T}_0} |L(1/2, \chi_c)|^k |M(c, \kappa)|^{k\kappa}.$$

For each $c \in \mathcal{F}(X) \setminus \mathcal{T}_0$ (see (15)), there exists $0 \leq u = u(c) \leq J$ such that $\operatorname{Re} F_0(c, u) > \ell_0/(ke^2)$. Therefore, applying Cauchy-Schwarz inequality shows that the inequality

$$(19)^2 \leq (J+1) \sum_{c \in \mathcal{F}(X)} |L(1/2, \chi_c)|^{2k} \sum_{u=0}^J \sum_{c \in \mathcal{F}(X)} |M(c, \kappa)|^{2k\kappa} \left(\frac{ke^2 \operatorname{Re} F_0(c, u)}{\ell_0} \right)^{2s_0}$$

holds for any positive *even* integer s_0 . We choose $s_0 = (2k\kappa + 3)\ell_0$. Then, applying Lemma 4.2 with $j = -1$ together with Lemma 4.1 shows that the inequality

$$\begin{aligned} & \sum_{c \in \mathcal{F}(X)} (\operatorname{Re} F_0(c, u))^{2s_0} |M(c, \kappa)|^{2k\kappa} \\ & < \frac{|\mathcal{F}(X)|(2s_0)!(\log_2 X^{\theta_0})^{s_0}}{2^{s_0}(2\ell_0)!} \left(\frac{5}{3} \right)^{2\ell_0} \exp(0.006k^3 + 2k + 2k^2) \\ & \quad + (2e^{1/4}k^2 + 4ke^{1/(4k)}) \log_2 X^{\theta_0} + \frac{0.024k^3 + k^2 + k^4}{4[k_0]^2} \\ & \quad + 4(e-2)k^2(J + o_X(1)) + o(4^{s_0} X^{(1/2+\varepsilon)\eta_0}) \end{aligned}$$

holds, where η_0 is defined in (34). The last o -term can be omitted provided that

$$\eta_0 = 1 + (4k\kappa + 6)\ell_0\theta_0 + 2k\kappa \sum_{0 \leq r \leq J} \theta_r \ell_r \leq \frac{1}{1/2 + \varepsilon} \left(1 - \frac{s_0 \log 4}{\log X} \right).$$

Recalling that $\theta_r = e^r (\log_2 X)^{-\alpha}$, we see that

$$\eta_0 < 1 + 2(4k\kappa + 6)\theta_0^{1-\beta} + 4k\kappa \frac{e^{(1-\beta)}\theta_J^{1-\beta}}{e^{1-\beta} - 1}.$$

Since $\theta_0 < \theta_J \leq \Theta$, it is sufficient to assume that

$$(20) \quad \Theta \leq \left(\frac{1-2\varepsilon - o(1)}{1+2\varepsilon} \left(8k\kappa + 12 + \frac{4k\kappa e^{1-\beta}}{e^{1-\beta} - 1} \right)^{-1} \right)^{1/(1-\beta)}.$$

Using the inequalities (cf. [22, Eq. (2)])

$$(21) \quad \left(\frac{n}{e} \right)^n \sqrt{2\pi n} \exp(1/(12n+1)) < n! < \left(\frac{n}{e} \right)^n \sqrt{2\pi n} \exp(1/(12n))$$

it follows that

$$\frac{(2s_0)!}{(2\ell_0)!} \left(\frac{5}{3} \right)^{2\ell_0} \left(\frac{ke^2}{\sqrt{2\ell_0}} \right)^{2s_0} < \left(ke(4k\kappa + 6) \left(\frac{5e}{6\ell_0} \right)^{2/(4k\kappa+6)} \right)^{(4k\kappa+6)\ell_0}.$$

Since $\ell_0 > \theta_0^{-\beta}$ for $(\log_2 X)^{\alpha\beta} > 2$, which we assumed in (12), we deduce that

$$\begin{aligned} & \sum_{u=0}^J \sum_{c \in \mathcal{F}(X)} |M(c, \kappa)|^{2k\kappa} \left(\frac{ke^2 \operatorname{Re} F_0(c, u)}{\ell_0} \right)^{2s_0} \\ & \ll J |\mathcal{F}(X)| \exp \left((2e^{1/4}k^2 + 4ke^{1/(4k)}) \log_2 X + 4(e-2)k^2 (J + o_X(1)) + O(k) \right. \\ & \quad \left. + (4k\kappa + 6)\ell_0 \log \left(ke(4k\kappa + 6) \left(\frac{5e}{6} \right)^{1/(2k\kappa+3)} (\log_2 X)^{1/2-\alpha\beta/(2k\kappa+3)} \right) \right). \end{aligned}$$

Combining this estimate with the result of Proposition 4.4 and recalling that $J < \alpha \log_3 X$, it follows that

$$\begin{aligned} (19) & \ll |\mathcal{F}(X)| \exp \left(k(\log_2 X)^2 + \log(\alpha \log_3 X) \right. \\ & \quad \left. + (4k\kappa + 6)(\log_2 X)^{\alpha\beta} \log \left(ke(4k\kappa + 6) \left(\frac{5e}{6} \right)^{1/(2k\kappa+3)} (\log_2 X)^{1/2-\alpha\beta/(2k\kappa+3)} \right) \right. \\ & \quad \left. + (e^{1/4}k^2 + 2ke^{1/(4k)}) \log_2 X + 2k^2 \alpha \log_3 X \right) \\ & = o(|\mathcal{F}(X)|), \end{aligned}$$

as $X \rightarrow \infty$, provided that (11) and (20) hold.

As for the sum (18) we use the upper bound for $|L(1/2, \chi_c)|^k$ in Lemma 3.7 to get

$$\begin{aligned} (18) & \leq \exp(k/\theta_J + O(kk_0)) \sum_{c \in \mathcal{F}(X)} D_{J,k}(c) S_{J,k}(c) |M(c, \kappa)|^{k\kappa} \\ & \quad + \sum_{\substack{0 \leq j < J \\ j < u \leq J}} \exp \left(\frac{k}{\theta_j} + O(kk_0) \right) \\ & \quad \cdot \sum_{c \in \mathcal{F}(X)} D_{j,k}(c) S_{j,k}(c) \left(\frac{e^2 k \operatorname{Re} F_{j+1}(c, u)}{\ell_{j+1}} \right)^{s_{j+1}} |M(c, \kappa)|^{k\kappa}, \end{aligned}$$

which holds for any positive even integers s_1, \dots, s_J . To estimate the sums

$$(22) \quad \sum_{c \in \mathcal{F}(X)} D_{J,k}(c) S_{J,k}(c) |M(c, \kappa)|^{k\kappa},$$

$$(23) \quad \sum_{c \in \mathcal{F}(X)} D_{j,k}(c) S_{j,k}(c) \left(\frac{e^2 k \operatorname{Re} F_{j+1}(c, u)}{\ell_{j+1}} \right)^{s_{j+1}} |M(c, \kappa)|^{k\kappa},$$

we use Cauchy-Schwarz inequality and deduce that

$$(22)^2 \leq \sum_{c \in \mathcal{F}(X)} (D_{J,k}(c))^2 |M(c, \kappa)|^{2k\kappa} \sum_{c \in \mathcal{F}(X)} (S_{J,k}(c))^2,$$

and

$$(23)^2 \leq \sum_{c \in \mathcal{F}(X)} (D_{j,k}(c))^2 \left(\frac{e^2 k \operatorname{Re} F_{j+1}(c, u)}{\ell_{j+1}} \right)^{2s_{j+1}} |M(c, \kappa)|^{2k\kappa} \sum_{c \in \mathcal{F}(X)} (S_{j,k}(c))^2.$$

Then, we use (33) in Lemma 4.2 together with (49) to conclude that

$$(22)^2 \ll \mathcal{S}_k |\mathcal{F}(X)| \left(|\mathcal{F}(X)| \exp(8k^2) \prod_{r=0}^J (1 + e^{-\ell_r})^2 (1 + 2^{-\ell_r/2}) + o\left(X^{(1/2+\varepsilon)\eta_J}\right) \right),$$

where \mathcal{S}_k is given by (50) and

$$\eta_J = 1 + (2k\kappa + 1) \sum_{0 \leq r \leq J} \theta_r \ell_r \leq 1 + 2(2k\kappa + 1) \frac{e^{(1-\beta)} \theta_J^{1-\beta}}{e^{1-\beta} - 1}.$$

As before the o -term is $o(|\mathcal{F}(X)|)$ and can be omitted under the assumption that

$$(24) \quad \Theta \leq \left(\frac{(e^{1-\beta} - 1)(1 - 2\varepsilon)}{2e^{1-\beta}(2k\kappa + 1)(1 + 2\varepsilon)} \right)^{1/(1-\beta)}.$$

Finally for (23), we choose, for each $0 \leq j < J$, $s_{j+1} = 2 \lfloor 1/(4a\theta_{j+1}) \rfloor$ for some $a > 1$ to be determined. Note that $s_{j+1} \geq 2k\kappa\ell_{j+1} + 2$ holds for each j as long as

$$(25) \quad \Theta \leq \left(\frac{1}{8a(k\kappa + 1)} \right)^{1/(1-\beta)}.$$

With these choices, we can then use (32) in Lemma 4.2 together with (49) and Lemma 4.1 to get for each $0 \leq j < J$ that

$$(23)^2 < \mathcal{S}_k |\mathcal{F}(X)|^2 (1 + o(1)) \exp(8k^2) \prod_{r=0}^j (1 + e^{-\ell_r})^2 (1 + 2^{-\ell_r/2}) \\ \cdot \left(\frac{e^2 k}{2\ell_{j+1}} \right)^{2s_{j+1}} \frac{(2s_{j+1})!}{\lfloor \frac{2s_{j+1} - 4k\kappa\ell_{j+1}}{3} \rfloor!} \left(\frac{5}{3} \right)^{\lfloor \frac{2s_{j+1} - 4k\kappa\ell_{j+1}}{3} \rfloor} \left(1 + O\left(\frac{1}{\log X^{\theta_j}} \right) \right)^{s_{j+1}} \\ \cdot \exp\left(e^{1/4} k^2 + 2ke^{1/(4k)} + 4(e-2)k^2(J-j-1) + O_k\left(\frac{1}{\log X^{\theta_j}} \right) \right),$$

where we replaced the o -term in (32) by $o(|\mathcal{F}(X)|)$ imposing the condition that

$$\eta_j \leq (1/2 + \varepsilon)^{-1} \left(1 - \frac{\log 4}{2a\theta_{j+1} \log X} \right),$$

and this holds provided that

$$(26) \quad \Theta \leq \left(\frac{e^{1-\beta} - 1}{2(2k\kappa + 1)e^{1-\beta}} \left(\frac{1 - 2\varepsilon - o(1)}{1 + 2\varepsilon} - \frac{1}{a} \right) \right)^{1/(1-\beta)}.$$

Now, write $2s_{j+1} = A$, and $\lfloor (2s_{j+1} - 4k\kappa\ell_{j+1})/3 \rfloor = B$. Then, $B/A < 1/3$ and

$$(27) \quad \beta + B/A - 1 > \beta - \frac{2}{3} - \frac{2a(8k\kappa + 1)\Theta^{1-\beta}}{3}.$$

Thus, $\beta + B/A - 1 > 0$ provided that

$$(28) \quad \beta > \frac{2}{3}, \quad \Theta < \left(\frac{3\beta - 2}{2a(8k\kappa + 3)} \right)^{1/(1-\beta)}.$$

Assuming (28) holds, it follows using (21) that

$$\begin{aligned}
(29) \quad & \left(\frac{5}{3}\right)^B \left(\frac{e^2 k}{2\ell_{j+1}}\right)^A \frac{A!}{B!} \\
& < \left(\left(\frac{5}{3}\right)^{B/A} \frac{e^2 k}{2\theta_{j+1}^{-\beta}}\right)^A A^{A-B} \exp\left(\log 2 + \frac{1}{12A} + B \log(4e) - A\right) \\
& < \left(\left(\frac{5}{3}\right)^{1/3} \frac{e^2 k \theta_{j+1}^{\beta+B/A-1}}{2a^{1-B/A}}\right)^A \\
& < \exp\left(\frac{1}{2a\theta_{j+1}} \log\left(\left(\frac{5}{3a^2}\right)^{1/3} \frac{e^2 k \theta_{j+1}^{\beta+B/A-1}}{2}\right)\right),
\end{aligned}$$

where we used $\ell_{j+1} > \theta_{j+1}^{-\beta}$ for the first inequality, which holds assuming (28), the second inequality follows since $B/A < 1/3$, $A \leq 1/(a\theta_{j+1})$, and $A \geq 4$ implies $1/(12A) + \log 2 + B \log(4e) < A$, and the third inequality follows since $B/A < 1/3$ and $A > 1/(2a\theta_{j+1})$ assuming that

$$(30) \quad \Theta < \left(\frac{2}{e^2 k} \left(\frac{3a^2}{5}\right)^{1/3}\right)^{1/(\beta-2/3)}.$$

Therefore, using (29) we see that

$$\begin{aligned}
(23)^2 & < |\mathcal{F}(X)|^2 (1 + o(1)) \mathcal{S}_k \exp(8k^2 + e^{1/4}k^2 + 2ke^{1/(4k)}) \prod_{r=0}^j (1 + e^{-\ell_r})^2 (1 + 2^{-\ell_r/2}) \\
& \cdot \exp\left(\frac{1}{2a\theta_{j+1}} \log\left(\left(\frac{5}{3a^2}\right)^{1/3} \frac{e^2 k \theta_{j+1}^{\beta+B/A-1}}{2}\right) + O\left(\frac{1}{\theta_{j+1} \log X^{\theta_j}}\right)\right) \\
& + 4(e-2)k^2(J-j-1) + O_k\left(\frac{1}{\log X^{\theta_j}}\right).
\end{aligned}$$

Combining the estimates for (22) and (23) we conclude that

$$(18) < |\mathcal{F}(X)|(1 + o_X(1)) \mathcal{D}_k \left(C_J \exp(e^{1/4}k^2/2 + ke^{1/(4k)}) + \exp(ek/\Theta)\right)$$

where

$$\mathcal{D}_k = \mathcal{S}_k^{1/2} \exp(4k^2 + O(kk_0)) \prod_{r=0}^J (1 + e^{-\ell_r})(1 + 2^{-\ell_r/2})^{1/2},$$

and

$$\begin{aligned}
C_J &= \sum_{\substack{0 \leq j < J \\ j < u \leq J}} \exp\left(\frac{k}{\theta_j} + \frac{1}{4a\theta_{j+1}} \log\left(\left(\frac{5}{3a^2}\right)^{1/3} \frac{e^2 k \theta_{j+1}^{\beta+B/A-1}}{2}\right)\right) + O\left(\frac{1}{\theta_{j+1} \log X^{\theta_j}}\right) \\
&\quad + 2(e-2)k^2(J-j-1) \\
&= \sum_{0 \leq j \leq J-1} (J-j) \exp\left(\frac{ke^{J-j}}{\theta_J} + \frac{e^{J-j-1}}{4a\theta_J} \log\left(\left(\frac{5}{3a^2}\right)^{1/3} \frac{e^2 k (e^{j+1-J} \theta_J)^{\beta+B/A-1}}{2}\right)\right) \\
&\quad + O\left(\frac{1}{e^{j+1-J} e^{j-J} \theta_J^2 \log X}\right) + 2(e-2)k^2(J-1-j) \\
&= \sum_{0 \leq r \leq J-1} (r+1) \exp\left(\frac{ke^{r+1}}{\theta_J} + \frac{e^r}{4a\theta_J} \log\left(\left(\frac{5}{3a^2}\right)^{1/3} \frac{e^2 k (e^{-r} \theta_J)^{\beta+B/A-1}}{2}\right)\right) \\
&\quad + O\left(\frac{e^{2r}}{\theta_J^2 \log X}\right) + 2(e-2)k^2 r \\
&< \sum_{0 \leq r \leq J-1} (r+1) \exp\left(f(r) + O\left(\frac{(\log_2 X)^{2\alpha}}{\log X}\right)\right).
\end{aligned}$$

Here,

$$\begin{aligned}
(31) \quad f(r) &= R_1 e^r - R_2 r e^r + 2k^2 r, \\
R_1 &= \frac{e}{\Theta} \left(ke + \frac{1}{4a} \log\left(\left(\frac{5}{3a^2}\right)^{1/3} \frac{e^2 k}{2}\right) \right. \\
&\quad \left. + \frac{\log \Theta}{4a} \left(\beta - \frac{2}{3} - \frac{2a(8k\kappa + 1)\Theta^{1-\beta}}{3} \right) \right), \\
R_2 &= \frac{e}{4a\Theta} \left(\beta - \frac{2}{3} - \frac{2a(8k\kappa + 1)\Theta^{1-\beta}}{3} \right).
\end{aligned}$$

To finish the proof of Theorem 1.2, we observe that it is possible to choose the parameters $\varepsilon, a, \beta, \alpha$ in the given order (for example, one possible choice, is to take $\varepsilon = 1/64, a = 6, \beta = 3/4, \alpha = 3 + 2k\kappa$), and then choose Θ small enough in such a way that $R_1 < 0, R_2 > 0$, and (11), (20), (24), (25), (26), (27), (28), (30), and (51) are all satisfied. This shows that $C_J \ll 1$ and for X sufficiently large in terms of the chosen parameters, (1) holds. This finishes the proof.

4.2. Behind the Scenes. In this part, we provide the technical results used in the proof of Theorem 1.2.

Lemma 4.2. *Let j, u be integers satisfying $-1 \leq j \leq J-1$ and $j+1 \leq u \leq J$. For any positive integer s_{j+1} satisfying $s_{j+1} \geq 2k\kappa\ell_{j+1} + 2$,*

$$\begin{aligned}
(32) \quad & \sum_{c \in \mathcal{F}(X)} D_{j,k}(c)^2 (\operatorname{Re} F_{j+1}(c, u))^{2s_{j+1}} |M(c, \kappa)|^{2k\kappa} \\
& < \exp(8k^2) \prod_{r=0}^j (1 + e^{-\ell_r})^2 (1 + 2^{-\ell_r/2}) \\
& \cdot \frac{|\mathcal{F}(X)|(2s_{j+1})!}{4^{s_{j+1}} \left\lfloor \frac{2s_{j+1} - 4k\kappa\ell_{j+1}}{3} \right\rfloor!} \left(\sum_{\mathfrak{p} \in \mathcal{I}_{j+1}} \frac{1}{N\mathfrak{p}} \right)^{s_{j+1}} \left(\frac{5}{3} \right)^{\left\lfloor \frac{2s_{j+1} - 4k\kappa\ell_{j+1}}{3} \right\rfloor} \\
& \cdot \exp\left(\frac{0.024k^3 + k^2 + k^4}{4[X\theta_j]^2} + \sum_{\mathfrak{p} \in \mathcal{I}_{j+1}} \frac{0.006k^3 + 2k + 2k^2}{(N\mathfrak{p})^{3/2}} + \frac{e^{1/4}k^2 + 2ke^{1/(4k)}}{N\mathfrak{p}} \right) \\
& + \sum_{X^{\theta_{j+1}} < N\mathfrak{p} \leq X^{\theta_j}} \frac{4(e-2)k^2}{N\mathfrak{p}} + o\left(4^{s_{j+1}} X^{(1/2+\varepsilon)\eta_j}\right)
\end{aligned}$$

where the first line of the upper bound is to be omitted for $j = -1$. Furthermore,

$$\begin{aligned}
(33) \quad & \sum_{c \in \mathcal{F}(X)} (D_{J,k}(c))^2 |M(c, \kappa)|^{2k\kappa} \\
& < |\mathcal{F}(X)| \exp(8k^2) \prod_{r=0}^J (1 + e^{-\ell_r})^2 (1 + 2^{-\ell_r/2}) + o\left(X^{(1/2+\varepsilon)\eta_J}\right)
\end{aligned}$$

where

$$(34) \quad \eta_j = 1 + 2s_{j+1}\theta_{j+1} + 2k\kappa \sum_{0 \leq r \leq J} \theta_r \ell_r + \sum_{0 \leq r \leq j} \theta_r \ell_r.$$

and the second term $2s_{j+1}\theta_{j+1}$ is to be omitted when $j = J$.

Proof. Using the definitions (13) and (17), we see that

$$\sum_{c \in \mathcal{F}(X)} D_{j,k}(c)^2 (\operatorname{Re} F_{j+1}(c, u))^{2s_{j+1}} |M(c, \kappa)|^{2k\kappa} = \prod_{r=0}^j (1 + e^{-\ell_r})^2 \sum_{c \in \mathcal{F}(X)} \prod_{r=0}^J \mathcal{E}(r, c),$$

where, for any $-1 \leq j \leq J$,

$$\mathcal{E}(r, c) = \begin{cases} (E_{\ell_r}(k \operatorname{Re} F_r(c, j)))^2 |M_r(c, \kappa)|^{2k\kappa} & \text{if } 0 \leq r \leq j \\ (\operatorname{Re} F_r(c, u))^{2s_r} |M_r(c, \kappa)|^{2k\kappa} & \text{if } r = j+1 \\ |M_r(c, \kappa)|^{2k\kappa} & \text{if } j+1 < r \leq J. \end{cases}$$

Now writing $\operatorname{Re} z = \frac{1}{2}(z + \bar{z})$, we obtain for any $n \geq 0$ that

$$\begin{aligned}
(35) \quad & (\operatorname{Re} F_r(c, u))^n = \frac{n!}{2^n} \sum_{\substack{\mathfrak{a}, \mathfrak{b} \in \mathcal{I}_r \\ \Omega(\mathfrak{a}\mathfrak{b})=n}} \frac{f(\mathfrak{a}\mathfrak{b}, u)\nu(\mathfrak{a})\nu(\mathfrak{b})}{N(\mathfrak{a}\mathfrak{b})^{1/2}} \chi_c(\mathfrak{a}\mathfrak{b}^2) \\
& E_{\ell_r}(k \operatorname{Re} F_r(c, j)) = \sum_{\substack{\mathfrak{a}, \mathfrak{b} \in \mathcal{I}_r \\ 0 \leq \Omega(\mathfrak{a}\mathfrak{b}) \leq \ell_r}} (k/2)^{\Omega(\mathfrak{a}\mathfrak{b})} \frac{f(\mathfrak{a}\mathfrak{b}, j)\nu(\mathfrak{a})\nu(\mathfrak{b})}{N(\mathfrak{a}\mathfrak{b})^{1/2}} \chi_c(\mathfrak{a}\mathfrak{b}^2).
\end{aligned}$$

Therefore,

$$(36) \quad D_{j,k}(c)^2 = \prod_{r=0}^j (1 + e^{-\ell_r})^2 \sum_{\substack{\mathbf{a}_{1r}, \mathbf{b}_{1r}, \mathbf{a}_{2r}, \mathbf{b}_{2r} \in \mathcal{I}_r \\ 0 \leq \Omega(\mathbf{a}_{1r} \mathbf{b}_{1r}) \leq \ell_r \\ 0 \leq \Omega(\mathbf{a}_{2r} \mathbf{b}_{2r}) \leq \ell_r}} (k/2)^{\Omega(\mathbf{a}_{1r} \mathbf{b}_{1r} \mathbf{a}_{2r} \mathbf{b}_{2r})} \\ \cdot \frac{f(\mathbf{a}_{1r} \mathbf{b}_{1r} \mathbf{a}_{2r} \mathbf{b}_{2r}, j) \nu(\mathbf{a}_{1r}) \nu(\mathbf{a}_{2r}) \nu(\mathbf{b}_{1r}) \nu(\mathbf{b}_{2r})}{\mathbf{N}(\mathbf{a}_{1r} \mathbf{b}_{1r} \mathbf{a}_{2r} \mathbf{b}_{2r})^{1/2}} \chi_c(\mathbf{a}_{1r} \mathbf{b}_{1r}^2 \mathbf{a}_{2r} \mathbf{b}_{2r}^2).$$

Moreover, since $2k\kappa$ is a positive even integer,

$$(37) \quad |M_r(c, \kappa)|^{2k\kappa} = \sum_{\mathbf{a}, \mathbf{b} \in \mathcal{I}_r} \frac{\chi_c(\mathbf{a} \mathbf{b}^2) \lambda(\mathbf{a} \mathbf{b}) f(\mathbf{a} \mathbf{b}, J)}{\mathbf{N}(\mathbf{a} \mathbf{b})^{1/2} \kappa^{\Omega(\mathbf{a} \mathbf{b})}} \nu_{k\kappa}(\mathbf{a}; \ell_r) \nu_{k\kappa}(\mathbf{b}; \ell_r).$$

Using (35), (36) and (37) we see that for $0 \leq r \leq j$,

$$\mathcal{E}(r, c) = \sum_{\substack{\mathbf{a}_{1r}, \mathbf{b}_{1r}, \mathbf{a}_{2r}, \mathbf{b}_{2r}, \mathbf{a}_{3r}, \mathbf{b}_{3r} \in \mathcal{I}_r \\ 0 \leq \Omega(\mathbf{a}_{1r} \mathbf{b}_{1r}) \leq \ell_r \\ 0 \leq \Omega(\mathbf{a}_{2r} \mathbf{b}_{2r}) \leq \ell_r \\ (c, \mathbf{m}_r) = 1}} \dots \sum_{\substack{\mathbf{a}_{3r}, \mathbf{b}_{3r} \in \mathcal{I}_r \\ 0 \leq \Omega(\mathbf{a}_{3r} \mathbf{b}_{3r}) \leq \ell_r \\ (c, \mathbf{m}_r) = 1}} \frac{\lambda(\mathbf{a}_{3r} \mathbf{b}_{3r}) f(\mathbf{a}_{1r} \mathbf{b}_{1r} \mathbf{a}_{2r} \mathbf{b}_{2r}, j) f(\mathbf{a}_{3r} \mathbf{b}_{3r}, J)}{\mathbf{N}(\mathbf{a}_{1r} \mathbf{b}_{1r} \mathbf{a}_{2r} \mathbf{b}_{2r} \mathbf{a}_{3r} \mathbf{b}_{3r})^{1/2} \kappa^{\Omega(\mathbf{a}_{3r} \mathbf{b}_{3r})}} \\ \cdot \left(\frac{k}{2}\right)^{\Omega(\mathbf{a}_{1r} \mathbf{b}_{1r} \mathbf{a}_{2r} \mathbf{b}_{2r})} \nu(\mathbf{a}_{1r}) \nu(\mathbf{b}_{1r}) \nu(\mathbf{a}_{2r}) \nu(\mathbf{b}_{2r}) \nu_{k\kappa}(\mathbf{a}_{3r}; \ell_r) \nu_{k\kappa}(\mathbf{b}_{3r}; \ell_r) \chi_c(\mathbf{m}_r),$$

for $r = j + 1$,

$$\mathcal{E}(r, c) = \frac{(2s_r)!}{4^{s_r}} \sum_{\substack{\mathbf{a}_{1r}, \mathbf{b}_{1r}, \mathbf{a}_{2r}, \mathbf{b}_{2r} \in \mathcal{I}_r \\ \Omega(\mathbf{a}_{1r} \mathbf{b}_{1r}) = 2s_r \\ (c, \mathbf{m}_r) = 1}} \dots \sum_{\substack{\mathbf{a}_{2r}, \mathbf{b}_{2r} \in \mathcal{I}_r \\ \Omega(\mathbf{a}_{2r} \mathbf{b}_{2r}) = 2s_r \\ (c, \mathbf{m}_r) = 1}} \frac{\lambda(\mathbf{a}_{2r} \mathbf{b}_{2r}) f(\mathbf{a}_{1r} \mathbf{b}_{1r}, u) f(\mathbf{a}_{2r} \mathbf{b}_{2r}, J)}{\mathbf{N}(\mathbf{a}_{1r} \mathbf{b}_{1r} \mathbf{a}_{2r} \mathbf{b}_{2r})^{1/2} \kappa^{\Omega(\mathbf{a}_{2r} \mathbf{b}_{2r})}} \\ \cdot \nu(\mathbf{a}_{1r}) \nu(\mathbf{b}_{1r}) \nu_{k\kappa}(\mathbf{a}_{2r}; \ell_r) \nu_{k\kappa}(\mathbf{b}_{2r}; \ell_r) \chi_c(\mathbf{m}_r),$$

and for $j + 1 < r \leq J$,

$$\mathcal{E}(r, c) = \sum_{\substack{\mathbf{a}_{1r}, \mathbf{b}_{1r} \in \mathcal{I}_r \\ (c, \mathbf{m}_r) = 1}} \frac{\lambda(\mathbf{a}_{1r} \mathbf{b}_{1r}) f(\mathbf{a}_{1r} \mathbf{b}_{1r}, J)}{\mathbf{N}(\mathbf{a}_{1r} \mathbf{b}_{1r})^{1/2} \kappa^{\Omega(\mathbf{a}_{1r} \mathbf{b}_{1r})}} \nu_{k\kappa}(\mathbf{a}_{1r}; \ell_r) \nu_{k\kappa}(\mathbf{b}_{1r}; \ell_r) \chi_c(\mathbf{m}_r).$$

Here,

$$\mathbf{m}_r = \begin{cases} \mathbf{a}_{1r} \mathbf{b}_{1r}^2 \mathbf{a}_{2r} \mathbf{b}_{2r}^2 \mathbf{a}_{3r} \mathbf{b}_{3r}^2 & \text{if } 0 \leq r \leq j \\ \mathbf{a}_{1r} \mathbf{b}_{1r}^2 \mathbf{a}_{2r} \mathbf{b}_{2r}^2 & \text{if } r = j + 1 \\ \mathbf{a}_{1r} \mathbf{b}_{1r}^2 & \text{if } j + 1 < r \leq J. \end{cases}$$

Note that \mathbf{m}_r 's are pairwise co-prime since the intervals \mathcal{I}_r are disjoint. For each $r = 0, \dots, J$, let $\tilde{\mathcal{E}}(r, c)$ denote the sum of the terms of $\mathcal{E}(r, c)$ for which $\mathbf{m}_r = \mathfrak{A}$, and put

$$\mathbf{m} = \prod_{r=0}^J \mathbf{m}_r.$$

Notice that $\mathbf{m} = \mathfrak{A}$ if and only if $\mathbf{m}_r = \mathfrak{A}$ for each r . To estimate

$$\sum_{c \in \mathcal{F}(X)} \left(\prod_{r=0}^J \mathcal{E}(r, c) - \prod_{r=0}^J \tilde{\mathcal{E}}(r, c) \right)$$

we interchange the summation over c after expanding the products and taking the difference so that the innermost sum is

$$\sum_{\substack{c \in \mathcal{F}(X) \\ (c, \mathbf{m})=1}} \chi_c(\mathbf{m}).$$

Then, applying Lemma 2.9 with $\varsigma = 0$ gives the bound $X^{1/2+\varepsilon}(\mathbf{N}\mathbf{m})^\varepsilon$ for this sum, which in turn yields that

$$\begin{aligned} & \prod_{r=0}^j (1 + e^{-\ell_r})^2 \sum_{c \in \mathcal{F}(X)} \left(\prod_{r=0}^J \mathcal{E}(r, c) - \prod_{r=0}^J \tilde{\mathcal{E}}(r, c) \right) \\ (38) \quad & \ll X^{1/2+\varepsilon} \prod_{r=0}^j (1 + e^{-\ell_r})^2 \left(E_{\ell_r} \left(k \sum_{\mathbf{p} \in \mathcal{I}_r} \frac{f(\mathbf{p}, j)}{\mathbf{N}(\mathbf{p})^{1/2-\varepsilon}} \right) \right)^2 \\ & \cdot \left(\sum_{\mathbf{p} \in \mathcal{I}_{j+1}} \frac{f(\mathbf{p}, u)}{\mathbf{N}(\mathbf{p})^{1/2-\varepsilon}} \right)^{2s_{j+1}} \left(\prod_{r=0}^J E_{\ell_r} \left(\frac{1}{\kappa} \sum_{\mathbf{p} \in \mathcal{I}_r} \frac{f(\mathbf{p}, J)}{\mathbf{N}(\mathbf{p})^{1/2-\varepsilon}} \right) \right)^{2k\kappa} \end{aligned}$$

where the sum over the interval \mathcal{I}_{j+1} is to be omitted when $j = J$. Since

$$\sum_{\mathbf{p} \in \mathcal{I}_r} f(\mathbf{p}, j) (\mathbf{N}\mathbf{p})^{\varepsilon-1/2} < 2X^{\theta_r(1/2+\varepsilon)}$$

for any $0 \leq r, j \leq J$, and

$$2kX^{\theta_r(1/2+\varepsilon)} > \ell_r, \quad 2\kappa^{-1}X^{\theta_r(1/2+\varepsilon)} > \ell_r$$

for sufficiently large X depending on the choice of α and β , it follows from Remark 3.4 that

$$(39) \quad (38) \ll 4^{s_{j+1}} X^{(1/2+\varepsilon)\eta_j} \prod_{r=0}^j \left(\frac{(2k)^{\ell_r} (\ell_r + 1) (1 + e^{-\ell_r})}{(\ell_r)!} \right)^2 \prod_{r=0}^J \left(\frac{2^{\ell_r} (\ell_r + 1)}{\kappa^{\ell_r} (\ell_r)!} \right)^{2k\kappa},$$

where η_j is given by (34). Using Stirling's formula and the definition of ℓ_r , it follows that (39) = $o_X(4^{s_{j+1}} X^{(1/2+\varepsilon)\eta_j})$.

We have shown so far that

$$\begin{aligned} \prod_{r=0}^j (1 + e^{-\ell_r})^2 \sum_{c \in \mathcal{F}(X)} \prod_{r=0}^J \mathcal{E}(r, c) &= \prod_{r=0}^j (1 + e^{-\ell_r})^2 \sum_{c \in \mathcal{F}(X)} \prod_{r=0}^J \tilde{\mathcal{E}}(r, c) \\ &\quad + o\left(4^{s_{j+1}} X^{(1/2+\varepsilon)\eta_j}\right) \end{aligned}$$

for any $-1 \leq j \leq J$. Combining the estimates for each $\tilde{\mathcal{E}}(r, c)$ that will be derived in the next section in (42), (44) and (45) gives the claimed bound in (32). Similarly, using (42) yields (33). \square

4.3. **Upper Bound for $\tilde{\mathcal{E}}(r, c)$ for $0 \leq r \leq j$.** Using Remark 3.9 we can write $\tilde{\mathcal{E}}(r, c) = \tilde{\mathcal{E}}_1(r, c) + \tilde{\mathcal{E}}_2(r, c)$, where

$$\begin{aligned} \tilde{\mathcal{E}}_1(r, c) &= \sum_{\substack{\mathbf{a}_{1r}, \mathbf{b}_{1r}, \mathbf{a}_{2r}, \mathbf{b}_{2r}, \mathbf{a}_{3r}, \mathbf{b}_{3r} \in \mathcal{I}_r \\ \mathfrak{m}_r = \mathfrak{c}_r \\ (c, \mathfrak{m}_r) = 1}} \dots \dots \sum \frac{\lambda(\mathbf{a}_{3r}, \mathbf{b}_{3r}) f(\mathbf{a}_{1r}, \mathbf{b}_{1r}, \mathbf{a}_{2r}, \mathbf{b}_{2r}, j) f(\mathbf{a}_{3r}, \mathbf{b}_{3r}, J)}{\mathbf{N}(\mathbf{a}_{1r}, \mathbf{b}_{1r}, \mathbf{a}_{2r}, \mathbf{b}_{2r}, \mathbf{a}_{3r}, \mathbf{b}_{3r})^{1/2} \kappa^{\Omega(\mathbf{a}_{3r}, \mathbf{b}_{3r})}} \\ &\cdot \left(\frac{k}{2}\right)^{\Omega(\mathbf{a}_{1r}, \mathbf{b}_{1r}, \mathbf{a}_{2r}, \mathbf{b}_{2r})} \nu(\mathbf{a}_{1r}) \nu(\mathbf{b}_{1r}) \nu(\mathbf{a}_{2r}) \nu(\mathbf{b}_{2r}) \nu_{k\kappa}(\mathbf{a}_{3r}) \nu_{k\kappa}(\mathbf{b}_{3r}) \end{aligned}$$

and

$$\begin{aligned} \tilde{\mathcal{E}}_2(r, c) &= \sum_{\substack{\mathbf{a}_{1r}, \mathbf{b}_{1r}, \mathbf{a}_{2r}, \mathbf{b}_{2r}, \mathbf{a}_{3r}, \mathbf{b}_{3r} \in \mathcal{I}_r \\ 0 \leq \Omega(\mathbf{a}_{1r}, \mathbf{b}_{1r}) \leq \ell_r \\ 0 \leq \Omega(\mathbf{a}_{2r}, \mathbf{b}_{2r}) \leq \ell_r \\ \max\{\Omega(\mathbf{a}_{3r}), \Omega(\mathbf{b}_{3r})\} > \ell_r \\ \mathfrak{m}_r = \mathfrak{c}_r, (c, \mathfrak{m}_r) = 1}} \dots \dots \sum \frac{\lambda(\mathbf{a}_{3r}, \mathbf{b}_{3r}) f(\mathbf{a}_{1r}, \mathbf{b}_{1r}, \mathbf{a}_{2r}, \mathbf{b}_{2r}, j) f(\mathbf{a}_{3r}, \mathbf{b}_{3r}, J)}{\mathbf{N}(\mathbf{a}_{1r}, \mathbf{b}_{1r}, \mathbf{a}_{2r}, \mathbf{b}_{2r}, \mathbf{a}_{3r}, \mathbf{b}_{3r})^{1/2} \kappa^{\Omega(\mathbf{a}_{3r}, \mathbf{b}_{3r})}} \\ &\cdot \left(\frac{k}{2}\right)^{\Omega(\mathbf{a}_{1r}, \mathbf{b}_{1r}, \mathbf{a}_{2r}, \mathbf{b}_{2r})} \nu(\mathbf{a}_{1r}) \nu(\mathbf{b}_{1r}) \nu(\mathbf{a}_{2r}) \nu(\mathbf{b}_{2r}) \nu_{k\kappa}(\mathbf{a}_{3r}; \ell_r) \nu_{k\kappa}(\mathbf{b}_{3r}; \ell_r) \\ &- \sum_{\substack{\mathbf{a}_{1r}, \mathbf{b}_{1r}, \mathbf{a}_{2r}, \mathbf{b}_{2r}, \mathbf{a}_{3r}, \mathbf{b}_{3r} \in \mathcal{I}_r \\ \max\{\Omega(\mathbf{a}_{1r}, \mathbf{b}_{1r}), \Omega(\mathbf{a}_{2r}, \mathbf{b}_{2r})\} > \ell_r \\ \Omega(\mathbf{a}_{3r}), \Omega(\mathbf{b}_{3r}) > \ell_r \\ \mathfrak{m}_r = \mathfrak{c}_r \\ (c, \mathfrak{m}_r) = 1}} \dots \dots \sum \frac{\lambda(\mathbf{a}_{3r}, \mathbf{b}_{3r}) f(\mathbf{a}_{1r}, \mathbf{b}_{1r}, \mathbf{a}_{2r}, \mathbf{b}_{2r}, j) f(\mathbf{a}_{3r}, \mathbf{b}_{3r}, J)}{\mathbf{N}(\mathbf{a}_{1r}, \mathbf{b}_{1r}, \mathbf{a}_{2r}, \mathbf{b}_{2r}, \mathbf{a}_{3r}, \mathbf{b}_{3r})^{1/2} \kappa^{\Omega(\mathbf{a}_{3r}, \mathbf{b}_{3r})}} \\ &\cdot \left(\frac{k}{2}\right)^{\Omega(\mathbf{a}_{1r}, \mathbf{b}_{1r}, \mathbf{a}_{2r}, \mathbf{b}_{2r})} \nu(\mathbf{a}_{1r}) \nu(\mathbf{b}_{1r}) \nu(\mathbf{a}_{2r}) \nu(\mathbf{b}_{2r}) \nu_{k\kappa}(\mathbf{a}_{3r}) \nu_{k\kappa}(\mathbf{b}_{3r}). \end{aligned}$$

4.3.1. *Estimate of $\tilde{\mathcal{E}}_1(r, c)$.* We can write

$$\begin{aligned} \tilde{\mathcal{E}}_1(r, c) &= \sum_{\substack{\mathbf{a}_{1r}, \mathbf{b}_{1r}, \mathbf{a}_{2r}, \mathbf{b}_{2r} \in \mathcal{I}_r \\ \mathbf{a}_{1r}, \mathbf{a}_{2r}, \mathbf{b}_{1r}^2, \mathbf{b}_{2r}^2 = \mathfrak{c}_r}} \dots \dots \sum \left(\frac{k}{2}\right)^{\Omega(\mathbf{a}_{1r}, \mathbf{b}_{1r})} \frac{\lambda(\mathbf{a}_{2r}, \mathbf{b}_{2r}) f(\mathbf{a}_{1r}, \mathbf{b}_{1r}, j) f(\mathbf{a}_{2r}, \mathbf{b}_{2r}, J)}{\mathbf{N}(\mathbf{a}_{1r}, \mathbf{b}_{1r}, \mathbf{a}_{2r}, \mathbf{b}_{2r})^{1/2} \kappa^{\Omega(\mathbf{a}_{2r}, \mathbf{b}_{2r})}} \\ &\cdot \nu_2(\mathbf{a}_{1r}) \nu_2(\mathbf{b}_{1r}) \nu_{k\kappa}(\mathbf{a}_{2r}) \nu_{k\kappa}(\mathbf{b}_{2r}) \\ &= \sum_{\substack{\mathbf{a}_r, \mathbf{b}_r \in \mathcal{I}_r \\ \mathbf{a}_r, \mathbf{b}_r^2 = \mathfrak{c}_r \\ (c, \mathbf{a}_r, \mathbf{b}_r) = 1}} \sum \frac{W(\mathbf{a}_r) W(\mathbf{b}_r)}{\mathbf{N}(\mathbf{a}_r, \mathbf{b}_r)^{1/2}} = \sum_{\substack{\mathbf{c}_r \in \mathcal{I}_r \\ (c, \mathbf{c}_r) = 1}} \frac{1}{\mathbf{N}(\mathbf{c}_r)^{1/2}} \sum_{\substack{\mathbf{a}_r, \mathbf{b}_r \in \mathcal{I}_r \\ \mathbf{a}_r, \mathbf{b}_r = \mathbf{c}_r \\ \mathbf{c}_r, \mathbf{b}_r = \mathfrak{c}_r}} W(\mathbf{a}_r) W(\mathbf{b}_r) \end{aligned}$$

where $W = U * V$ is the convolution of U and V , and

$$U(\mathbf{a}) = (k/2)^{\Omega(\mathbf{a})} f(\mathbf{a}, j) \nu_2(\mathbf{a}), \quad V(\mathbf{a}) = \frac{\lambda(\mathbf{a}) f(\mathbf{a}, J) \nu_{k\kappa}(\mathbf{a})}{\kappa^{\Omega(\mathbf{a})}}.$$

Since all factors are multiplicative in \mathbf{c}_r , we obtain

$$\tilde{\mathcal{E}}_1(r, c) = \prod_{\substack{\mathfrak{p} \in \mathcal{I}_r \\ \mathfrak{p} \nmid c}} \left(1 + \sum_{n \geq 1} \frac{1}{\mathbf{N}(\mathfrak{p})^{n/2}} \sum_{\substack{\mathbf{a}_r, \mathbf{b}_r \in \mathcal{I}_r \\ \mathbf{a}_r, \mathbf{b}_r = \mathfrak{p}^n \\ \mathfrak{p}^n, \mathbf{b}_r = \mathfrak{c}_r}} W(\mathbf{a}_r) W(\mathbf{b}_r) \right).$$

Noting that

$$W(\mathfrak{p}^i) = \frac{k^i}{i!} \sum_{0 \leq a \leq i} \binom{i}{a} f(\mathfrak{p}, j)^a (-f(\mathfrak{p}, J))^{i-a} = \frac{k^i}{i!} (f(\mathfrak{p}, j) - f(\mathfrak{p}, J))^i$$

we have that

$$\sum_{\substack{\mathbf{a}_r, \mathbf{b}_r \in \mathcal{I}_r \\ \mathbf{a}, \mathbf{b}_r = \mathbf{p}^n \\ \mathbf{p}^n \nmid \mathbf{b}_r}} W(\mathbf{a}_r)W(\mathbf{b}_r) = (-1)^n b_n(\mathbf{p}),$$

where

$$b_n(\mathbf{p}) = \sum_{\substack{0 \leq m \leq n \\ 3|n+m}} W(\mathbf{p}^{n-m})W(\mathbf{p}^m) = k^n (f(\mathbf{p}, J) - f(\mathbf{p}, j))^n \sum_{\substack{0 \leq m \leq n \\ 3|n+m}} \frac{1}{m!(n-m)!}.$$

Since $b_1(\mathbf{p}) = 0$, we conclude that

$$\tilde{\mathcal{E}}_1(r, c) = \prod_{\substack{\mathbf{p} \in \mathcal{I}_r \\ \mathbf{p} \nmid c}} \left(1 + \sum_{n \geq 2} \frac{(-1)^n b_n(\mathbf{p})}{N(\mathbf{p})^{n/2}} \right).$$

We will show that the sum over n is an alternating series and is thus bounded above by $b_2(\mathbf{p})/N\mathbf{p}$. First note that if $2 \mid n$, then

$$\begin{aligned} \sum_{\substack{0 \leq m \leq n+1 \\ 3|n+1+m}} \frac{1}{m!(n+1-m)!} &= \sum_{\substack{0 \leq m \leq n/2-1 \\ 3|n+1+m}} \frac{2}{m!(n+1-m)!} \\ &< \sum_{\substack{1 \leq m \leq n/2 \\ 3|n+m}} \frac{1}{m!(n-m)!} \leq \sum_{\substack{0 \leq m \leq n \\ 3|n+m}} \frac{1}{m!(n-m)!}. \end{aligned}$$

If $2 \nmid n$, then $n \geq 3$ and we have

$$\begin{aligned} \sum_{\substack{0 \leq m \leq n+1 \\ 3|n+1+m}} \frac{1}{m!(n+1-m)!} &= \sum_{\substack{0 \leq m \leq (n-5)/2 \\ 3|n+1+m}} \frac{2}{m!(n+1-m)!} + \frac{1}{\left(\frac{n+1}{2}\right)! \left(\frac{n+1}{2}\right)!} \\ &< \sum_{\substack{0 \leq m \leq (n-3)/2 \\ 3|n+m}} \frac{2}{m!(n-m)!} \leq \sum_{\substack{0 \leq m \leq n \\ 3|n+m}} \frac{1}{m!(n-m)!}. \end{aligned}$$

Furthermore, since $e^{-x} > 1 - x$ for $x \neq 0$, we have

$$\begin{aligned} 0 < f(\mathbf{p}, J) - f(\mathbf{p}, j) &= f(\mathbf{p}, J) - \frac{1}{(N\mathbf{p})^{1/(\theta_j \log X)}} \left(1 - \frac{\log N\mathbf{p}}{\theta_j \log X} \right) \\ &< 1 - \frac{1}{(N\mathbf{p})^{1/(\theta_j \log X)}} \left(1 - \frac{\log N\mathbf{p}}{\theta_j \log X} \right) < 1 - \left(1 - \frac{\log N\mathbf{p}}{\theta_j \log X} \right)^2 \\ &< \min \left\{ 1, \frac{2 \log N\mathbf{p}}{\theta_j \log X} \right\}. \end{aligned}$$

Hence, $0 < b_n(\mathbf{p}) < (2k)^n/n! \rightarrow 0$ as $n \rightarrow \infty$ and $b_n(\mathbf{p})/(N\mathbf{p})^{n/2}$ is decreasing since

$$\frac{b_{n+1}(\mathbf{p})(N\mathbf{p})^{n/2}}{b_n(\mathbf{p})(N\mathbf{p})^{(n+1)/2}} < \frac{k(f(\mathbf{p}, J) - f(\mathbf{p}, j))}{N(\mathbf{p})^{1/2}} < \frac{k}{\sqrt{N\mathbf{p}}} < 1$$

for all $\mathfrak{p} \in \mathcal{I}_r$. This shows that we have an alternating series. Therefore,

$$(40) \quad \begin{aligned} \tilde{\mathcal{E}}_1(r, c) &< \prod_{\substack{\mathfrak{p} \in \mathcal{I}_r \\ \mathfrak{p} \nmid c}} \left(1 + \frac{b_2(\mathfrak{p})}{N\mathfrak{p}}\right) < \prod_{\mathfrak{p} \in \mathcal{I}_r} \left(1 + \frac{k^2(f(\mathfrak{p}, J) - f(\mathfrak{p}, j))^2}{N(\mathfrak{p})}\right) \\ &< \exp\left(k^2 \sum_{\mathfrak{p} \in \mathcal{I}_r} \frac{1}{N\mathfrak{p}} \min\left\{1, \frac{4 \log^2 N\mathfrak{p}}{(\theta_j \log X)^2}\right\}\right). \end{aligned}$$

4.3.2. *Estimate of $\tilde{\mathcal{E}}_2(r, c)$.* Using Remark 3.9, the fact that

$$\max\{\Omega(\mathfrak{a}_{1r}\mathfrak{b}_{1r}), \Omega(\mathfrak{a}_{2r}\mathfrak{b}_{2r}), \Omega(\mathfrak{a}_{3r}), \Omega(\mathfrak{b}_{3r})\} > \ell_r$$

and the inequality $f(\mathfrak{a}, \cdot) \leq 1$, we see that

$$\begin{aligned} |\tilde{\mathcal{E}}_2(r, c)| &< \frac{2}{2^{\ell_r}} \sum_{\substack{\mathfrak{a}_{1r}, \mathfrak{b}_{1r}, \mathfrak{a}_{2r}, \\ \mathfrak{b}_{2r}, \mathfrak{a}_{3r}, \mathfrak{b}_{3r} \in \mathcal{I}_r \\ \mathfrak{m}_r = \overline{\mathfrak{a}\mathfrak{b}}}^{\cdot} \frac{k^{\Omega(\mathfrak{a}_{1r}\mathfrak{b}_{1r}\mathfrak{a}_{2r}\mathfrak{b}_{2r})} 2^{\Omega(\mathfrak{a}_{3r}\mathfrak{b}_{3r})}}{N(\mathfrak{a}_{1r}\mathfrak{b}_{1r}\mathfrak{a}_{2r}\mathfrak{b}_{2r}\mathfrak{a}_{3r}\mathfrak{b}_{3r})^{1/2} k^{\Omega(\mathfrak{a}_{3r}\mathfrak{b}_{3r})}} \\ &\quad \cdot \nu(\mathfrak{a}_{1r})\nu(\mathfrak{b}_{1r})\nu(\mathfrak{a}_{2r})\nu(\mathfrak{b}_{2r})\nu_{k\kappa}(\mathfrak{a}_{3r})\nu_{k\kappa}(\mathfrak{b}_{3r}) \\ &= \frac{2}{2^{\ell_r}} \sum_{\substack{\mathfrak{a}, \mathfrak{b}, \mathfrak{a}_{3r}, \mathfrak{b}_{3r} \in \mathcal{I}_r \\ \mathfrak{a}\mathfrak{b}^2\mathfrak{a}_{3r}\mathfrak{b}_{3r}^2 = \overline{\mathfrak{a}\mathfrak{b}}}^{\cdot} \frac{k^{\Omega(\mathfrak{a}\mathfrak{b})} 2^{\Omega(\mathfrak{a}_{3r}\mathfrak{b}_{3r})} \nu_2(\mathfrak{a})\nu_2(\mathfrak{b})\nu_{k\kappa}(\mathfrak{a}_{3r})\nu_{k\kappa}(\mathfrak{b}_{3r})}{N(\mathfrak{a}\mathfrak{b}\mathfrak{a}_{3r}\mathfrak{b}_{3r})^{1/2} k^{\Omega(\mathfrak{a}_{3r}\mathfrak{b}_{3r})}} \\ &= \frac{2}{2^{\ell_r}} \sum_{\substack{\mathfrak{A}, \mathfrak{B} \in \mathcal{I}_r \\ \mathfrak{A}\mathfrak{B}^2 = \overline{\mathfrak{a}\mathfrak{b}}}^{\cdot} \frac{H(\mathfrak{A})H(\mathfrak{B})}{N(\mathfrak{A}\mathfrak{B})^{1/2}} = \frac{2}{2^{\ell_r}} \sum_{\mathfrak{C} \in \mathcal{I}_r} \frac{1}{N(\mathfrak{C})^{1/2}} \sum_{\substack{\mathfrak{A}, \mathfrak{B} \in \mathcal{I}_r \\ \mathfrak{A}\mathfrak{B} = \mathfrak{C} \\ \mathfrak{A}\mathfrak{B}^2 = \overline{\mathfrak{a}\mathfrak{b}}}^{\cdot} H(\mathfrak{A})H(\mathfrak{B}). \end{aligned}$$

where

$$H(\mathfrak{A}) = \sum_{\substack{\mathfrak{a}, \mathfrak{a}_{3r} \\ \mathfrak{a}\mathfrak{a}_{3r} = \mathfrak{A}}} \nu_2(\mathfrak{a}) k^{\Omega(\mathfrak{a})} \nu_{k\kappa}(\mathfrak{a}_{3r}) (2/\kappa)^{\Omega(\mathfrak{a}_{3r})}.$$

This gives

$$\begin{aligned} 2^{\ell_r-1} |\tilde{\mathcal{E}}_2(r, c)| &< \prod_{\mathfrak{p} \in \mathcal{I}_r} \left(1 + \sum_{n \geq 2} \frac{1}{N(\mathfrak{p})^{n/2}} \sum_{\substack{0 \leq m \leq n \\ 3|m+n}} H(\mathfrak{p}^{n-m}) H(\mathfrak{p}^m)\right) \\ &< \prod_{\mathfrak{p} \in \mathcal{I}_r} \left(1 + \sum_{n \geq 2} \frac{(8k)^n}{n! N(\mathfrak{p})^{n/2}}\right) \end{aligned}$$

since

$$H(\mathfrak{p}^m) = \sum_{0 \leq i \leq m} \nu_2(\mathfrak{p}^i) k^{\Omega(\mathfrak{p}^i)} \nu_{k\kappa}(\mathfrak{p}^{m-i}) (2/\kappa)^{\Omega(\mathfrak{p}^{m-i})} = \frac{(4k)^m}{m!}.$$

Therefore, since $N\mathfrak{p} > k_0 \geq (2k)^2$,

$$(41) \quad |\tilde{\mathcal{E}}_2(r, c)| < 2^{1-\ell_r} \exp\left(4k^2(e^4 - 5) \sum_{\mathfrak{p} \in \mathcal{I}_r} \frac{1}{N\mathfrak{p}}\right).$$

Combining (41) with (40) and using Lemma 4.1 it follows that for sufficiently large X ,

$$(42) \quad \prod_{r=0}^j |\tilde{\mathcal{E}}(r, c)| < \exp(8k^2) \prod_{r=0}^j (1 + 2^{-\ell_r/2}).$$

4.4. **Bound for $\tilde{\mathcal{E}}(j+1)$.** Recall that for $r = j+1$,

$$\begin{aligned} \tilde{\mathcal{E}}(r) &= \frac{(2s_r)!}{4^{s_r}} \sum_{\substack{\mathbf{a}_{2r}, \mathbf{b}_{2r} \in \mathcal{I}_r \\ 0 \leq \Omega(\mathbf{a}_{2r}), \Omega(\mathbf{b}_{2r}) \leq k\kappa\ell_r}} \frac{\lambda(\mathbf{a}_{2r}\mathbf{b}_{2r})f(\mathbf{a}_{2r}\mathbf{b}_{2r}, J)\nu_{k\kappa}(\mathbf{a}_{2r}; \ell_r)\nu_{k\kappa}(\mathbf{b}_{2r}; \ell_r)}{N(\mathbf{a}_{2r}\mathbf{b}_{2r})^{1/2}\kappa^{\Omega(\mathbf{a}_{2r}, \mathbf{b}_{2r})}} \\ &\cdot \sum_{\substack{\mathbf{a}_{1r}, \mathbf{b}_{1r} \in \mathcal{I}_r \\ \Omega(\mathbf{a}_{1r}\mathbf{b}_{1r})=2s_r \\ \mathbf{a}_{1r}\mathbf{b}_{1r}^2, \mathbf{a}_{2r}\mathbf{b}_{2r}^2 = \square \\ (c, \prod_i \mathbf{a}_{ir}, \mathbf{b}_{ir})=1}} \frac{f(\mathbf{a}_{1r}\mathbf{b}_{1r}, u)\nu(\mathbf{a}_{1r})\nu(\mathbf{b}_{1r})}{N(\mathbf{a}_{1r}\mathbf{b}_{1r})^{1/2}}. \end{aligned}$$

We first replace $\mathbf{a}_{ir}, \mathbf{b}_{ir}$ with $\mathbf{r}\mathbf{a}_{1r}, \mathbf{r}\mathbf{b}_{1r}$ and $\mathbf{X}\mathbf{a}_{2r}, \mathbf{X}\mathbf{b}_{2r}$, respectively, with $(\mathbf{a}_{ir}, \mathbf{b}_{ir}) = 1$ for $i = 1, 2$, use the inequalities $|\lambda|, f \leq 1$, $\nu_n(\mathbf{a}; \cdot) \leq \nu_n(\mathbf{a})$, and then remove the condition that $(c, \prod_i \mathbf{a}_{ir}, \mathbf{b}_{ir}) = 1$. This shows that $4^{s_r}|\tilde{\mathcal{E}}(r)|/((2s_r)!)$ is bounded by

$$\sum_{\substack{\mathbf{a}_{2r}, \mathbf{b}_{2r}, \mathbf{X} \in \mathcal{I}_r \\ (\mathbf{a}_{2r}, \mathbf{b}_{2r})=1 \\ 0 \leq \Omega(\mathbf{X}\mathbf{a}_{2r}), \Omega(\mathbf{X}\mathbf{b}_{2r}) \leq k\kappa\ell_r}} \frac{\nu_{k\kappa}(\mathbf{X}\mathbf{a}_{2r})\nu_{k\kappa}(\mathbf{X}\mathbf{b}_{2r})}{N(\mathbf{X}^2\mathbf{a}_{2r}\mathbf{b}_{2r})^{1/2}\kappa^{\Omega(\mathbf{X}^2\mathbf{a}_{2r}, \mathbf{b}_{2r})}} \sum_{\substack{\mathbf{a}_{1r}, \mathbf{b}_{1r}, \mathbf{r} \in \mathcal{I}_r \\ (\mathbf{a}_{1r}, \mathbf{b}_{1r})=1 \\ \Omega(\mathbf{r}^2\mathbf{a}_{1r}\mathbf{b}_{1r})=2s_r \\ \mathbf{a}_{1r}\mathbf{b}_{1r}^2, \mathbf{a}_{2r}\mathbf{b}_{2r}^2 = \square}} \frac{\nu(\mathbf{r}\mathbf{a}_{1r})\nu(\mathbf{r}\mathbf{b}_{1r})}{N(\mathbf{r}^2\mathbf{a}_{1r}\mathbf{b}_{1r})^{1/2}}.$$

Next, replacing $\mathbf{a}_{1r}, \mathbf{b}_{2r}$ with $\mathbf{\eta}\mathbf{a}_{1r}, \mathbf{\eta}\mathbf{b}_{2r}$, and $\mathbf{a}_{2r}, \mathbf{b}_{1r}$ with $\mathbf{\mathfrak{Y}}\mathbf{a}_{2r}, \mathbf{\mathfrak{Y}}\mathbf{b}_{1r}$, where $(\mathbf{a}_{ir}, \mathbf{b}_{3-i,r}) = 1$ for $i = 1, 2$ gives

$$\begin{aligned} |\tilde{\mathcal{E}}(r)| &\leq \frac{(2s_r)!}{4^{s_r}} \sum_{\substack{\mathbf{a}_{2r}, \mathbf{b}_{2r}, \mathbf{X}, \mathbf{\eta}, \mathbf{\mathfrak{Y}} \in \mathcal{I}_r \\ (\mathbf{\mathfrak{Y}}\mathbf{a}_{2r}, \mathbf{\eta}\mathbf{b}_{2r})=1 \\ 0 \leq \Omega(\mathbf{X}\mathbf{\mathfrak{Y}}\mathbf{a}_{2r}), \Omega(\mathbf{\eta}\mathbf{X}\mathbf{b}_{2r}) \leq k\kappa\ell_r}} \frac{\nu_{k\kappa}(\mathbf{X}\mathbf{\mathfrak{Y}}\mathbf{a}_{2r})\nu_{k\kappa}(\mathbf{X}\mathbf{\eta}\mathbf{b}_{2r})}{N(\mathbf{X}^2\mathbf{\eta}^2\mathbf{\mathfrak{Y}}^2\mathbf{a}_{2r}\mathbf{b}_{2r})^{1/2}\kappa^{\Omega(\mathbf{X}^2\mathbf{\eta}\mathbf{\mathfrak{Y}}\mathbf{a}_{2r}, \mathbf{b}_{2r})}} \\ &\cdot \sum_{\substack{\mathbf{a}_{1r}, \mathbf{b}_{1r}, \mathbf{r} \in \mathcal{I}_r \\ (\mathbf{\eta}\mathbf{a}_{1r}, \mathbf{\mathfrak{Y}}\mathbf{b}_{1r})=1 \\ (\mathbf{a}_{2r}, \mathbf{b}_{1r})=1=(\mathbf{a}_{1r}, \mathbf{b}_{2r}) \\ \Omega(\mathbf{r}^2\mathbf{a}_{1r}\mathbf{b}_{1r})=2s_r - \Omega(\mathbf{\eta}\mathbf{\mathfrak{Y}}) \\ \mathbf{a}_{1r}\mathbf{b}_{1r}^2, \mathbf{a}_{2r}\mathbf{b}_{2r}^2 = \square}} \frac{\nu(\mathbf{r}\mathbf{\eta}\mathbf{a}_{1r})\nu(\mathbf{r}\mathbf{\mathfrak{Y}}\mathbf{b}_{1r})}{N(\mathbf{r}^2\mathbf{a}_{1r}\mathbf{b}_{1r})^{1/2}}. \end{aligned}$$

Now, replace $\mathbf{a}_{1r}, \mathbf{a}_{2r}$ with $\mathbf{z}\mathbf{a}_{1r}, \mathbf{z}\mathbf{a}_{2r}$ and $\mathbf{b}_{1r}, \mathbf{b}_{2r}$ with $\mathbf{z}\mathbf{b}_{1r}, \mathbf{z}\mathbf{b}_{2r}$, where $(\mathbf{a}_{1r}, \mathbf{a}_{2r}) = 1 = (\mathbf{b}_{1r}, \mathbf{b}_{2r})$ to get

$$\begin{aligned} |\tilde{\mathcal{E}}(r)| &\leq \frac{(2s_r)!}{4^{s_r}} \sum_{\substack{\mathbf{a}_{2r}, \mathbf{b}_{2r}, \mathbf{X}, \mathbf{\eta}, \mathbf{\mathfrak{Y}}, \mathbf{z}, \mathbf{z} \in \mathcal{I}_r \\ (\mathbf{\mathfrak{Y}}\mathbf{z}\mathbf{a}_{2r}, \mathbf{\eta}\mathbf{z}\mathbf{b}_{2r})=1 \\ 0 \leq \Omega(\mathbf{X}\mathbf{\mathfrak{Y}}\mathbf{z}\mathbf{a}_{2r}), \Omega(\mathbf{X}\mathbf{\eta}\mathbf{z}\mathbf{b}_{2r}) \leq k\kappa\ell_r}} \frac{\nu_{k\kappa}(\mathbf{X}\mathbf{\mathfrak{Y}}\mathbf{z}\mathbf{a}_{2r})\nu_{k\kappa}(\mathbf{X}\mathbf{\eta}\mathbf{z}\mathbf{b}_{2r})}{N(\mathbf{X}^2\mathbf{\eta}^2\mathbf{\mathfrak{Y}}^2\mathbf{z}\mathbf{z}\mathbf{a}_{2r}\mathbf{b}_{2r})^{1/2}\kappa^{\Omega(\mathbf{X}^2\mathbf{\eta}\mathbf{\mathfrak{Y}}\mathbf{z}\mathbf{z}\mathbf{a}_{2r}, \mathbf{b}_{2r})}} \\ &\cdot \sum_{\substack{\mathbf{a}_{1r}, \mathbf{b}_{1r}, \mathbf{r} \in \mathcal{I}_r \\ (\mathbf{a}_{1r}, \mathbf{a}_{2r})=1=(\mathbf{b}_{1r}, \mathbf{b}_{2r}) \\ (\mathbf{\eta}\mathbf{z}\mathbf{a}_{1r}, \mathbf{\mathfrak{Y}}\mathbf{z}\mathbf{b}_{1r})=1 \\ (\mathbf{z}\mathbf{a}_{2r}, \mathbf{z}\mathbf{b}_{1r})=1=(\mathbf{z}\mathbf{a}_{1r}, \mathbf{z}\mathbf{b}_{2r}) \\ \Omega(\mathbf{r}^2\mathbf{a}_{1r}\mathbf{b}_{1r})=2s_r - \Omega(\mathbf{\eta}\mathbf{\mathfrak{Y}}\mathbf{z}\mathbf{z}) \\ \mathbf{z}^2\mathbf{a}_{1r}\mathbf{a}_{2r}, \mathbf{z}\mathbf{b}_{1r}^2, \mathbf{z}\mathbf{b}_{2r}^2 = \square}} \frac{\nu(\mathbf{r}\mathbf{\eta}\mathbf{z}\mathbf{a}_{1r})\nu(\mathbf{r}\mathbf{\mathfrak{Y}}\mathbf{z}\mathbf{b}_{1r})}{N(\mathbf{r}^2\mathbf{z}\mathbf{z}\mathbf{a}_{1r}\mathbf{b}_{1r})^{1/2}}. \end{aligned}$$

Finally writing $\mathbf{z} = \mathbf{a}\mathbf{b}^2\mathbf{c}^3$ and $\mathbf{z} = \mathbf{A}\mathbf{B}^2\mathbf{C}^3$, where $(\mathbf{a}, \mathbf{b}) = 1 = (\mathbf{A}, \mathbf{B})$, all square-free, shows that $\mathbf{z}^2\mathbf{a}_{1r}\mathbf{a}_{2r} = \mathbf{a}^2\mathbf{b}^4\mathbf{c}^6\mathbf{a}_{1r}\mathbf{a}_{2r} = \square$ if and only if $\mathbf{a}^2\mathbf{b}\mathbf{a}_{1r}\mathbf{a}_{2r} = \square$ if $\mathbf{a}_{1r} = \mathbf{a}_1\mathbf{b}_1^2\mathbf{c}_1^3$ and $\mathbf{a}_{2r} = \mathbf{a}_2\mathbf{b}_2^2\mathbf{c}_2^3$, where $\mathbf{a} = \mathbf{a}_1\mathbf{a}_2$ and $\mathbf{b} = \mathbf{b}_1\mathbf{b}_2$. Similarly, we get $\mathbf{z}\mathbf{b}_{1r}^2, \mathbf{z}\mathbf{b}_{2r}^2 = \mathbf{A}\mathbf{B}^2\mathbf{C}^3\mathbf{b}_{1r}^2, \mathbf{b}_{2r}^2 = \square$ if $\mathbf{b}_{1r} = \mathbf{A}_1\mathbf{B}_1^2\mathbf{C}_1^3$ and $\mathbf{b}_{2r} = \mathbf{A}_2\mathbf{B}_2^2\mathbf{C}_2^3$ where $\mathbf{A} = \mathbf{A}_1\mathbf{A}_2$

and $\mathfrak{B} = \mathfrak{B}_1 \mathfrak{B}_2$. Making these changes leads to the bound

$$\begin{aligned}
|\tilde{\mathcal{E}}(r)| &\leq \frac{(2s_r)!}{4^{s_r}} \sum_{\mathfrak{c}_2, \mathfrak{c}_2, \mathfrak{x}, \mathfrak{y}, \mathfrak{z}, \mathfrak{c}, \mathfrak{c} \in \mathcal{I}_r} \frac{\nu(\mathfrak{h}\mathfrak{c}^3)\nu(\mathfrak{y}\mathfrak{c}^3)\nu_{k\kappa}(\mathfrak{x}\mathfrak{y}\mathfrak{c}^3\mathfrak{c}_2^3)\nu_{k\kappa}(\mathfrak{x}\mathfrak{y}\mathfrak{c}^3\mathfrak{c}_2^3)}{\mathsf{N}(\mathfrak{c}^6\mathfrak{c}^6\mathfrak{x}^2\mathfrak{y}^2\mathfrak{z}^2\mathfrak{c}_2^3\mathfrak{c}_2^3)^{1/2}\kappa^\Omega(\mathfrak{x}^2\mathfrak{y}\mathfrak{z}\mathfrak{c}^3\mathfrak{c}_2^3\mathfrak{c}_2^3)} \\
&\cdot \sum_{\substack{\mathfrak{a} \in \mathcal{I}_r \\ \mathfrak{a} \text{ sq-free} \\ \mathfrak{a} = \mathfrak{a}_1 \mathfrak{a}_2}} \frac{\nu(\mathfrak{a}\mathfrak{a}_1)\nu_{k\kappa}(\mathfrak{a}\mathfrak{a}_2)}{\mathsf{N}(\mathfrak{a})^{3/2}\kappa^\Omega(\mathfrak{a}\mathfrak{a}_2)} \sum_{\substack{\mathfrak{A} \in \mathcal{I}_r \\ \mathfrak{A} \text{ sq-free} \\ \mathfrak{A} = \mathfrak{A}_1 \mathfrak{A}_2}} \frac{\nu(\mathfrak{A}\mathfrak{A}_1)\nu_{k\kappa}(\mathfrak{A}\mathfrak{A}_2)}{\mathsf{N}(\mathfrak{A})^{3/2}\kappa^\Omega(\mathfrak{A}\mathfrak{A}_2)} \sum_{\substack{\mathfrak{b} \in \mathcal{I}_r \\ \mathfrak{b} \text{ sq-free} \\ \mathfrak{b} = \mathfrak{b}_1 \mathfrak{b}_2}} \frac{\nu(\mathfrak{b}^2\mathfrak{b}_1^2)\nu_{k\kappa}(\mathfrak{b}^2\mathfrak{b}_2^2)}{\mathsf{N}(\mathfrak{b})^3\kappa^\Omega(\mathfrak{b}^2\mathfrak{b}_2^2)} \\
&\cdot \sum_{\substack{\mathfrak{B} \in \mathcal{I}_r \\ \mathfrak{B} \text{ sq-free}, \mathfrak{B} = \mathfrak{B}_1 \mathfrak{B}_2 \\ 0 \leq \Omega(\mathfrak{x}\mathfrak{y}\mathfrak{a}\mathfrak{b}^2\mathfrak{c}^3\mathfrak{a}_2\mathfrak{b}_2^3\mathfrak{c}_2^3) \leq k\kappa\ell_r \\ 0 \leq \Omega(\mathfrak{x}\mathfrak{y}\mathfrak{A}\mathfrak{B}^2\mathfrak{c}^3\mathfrak{A}_2\mathfrak{B}_2^3\mathfrak{c}_2^3) \leq k\kappa\ell_r}} \frac{\nu(\mathfrak{B}^2\mathfrak{B}_1^2)\nu_{k\kappa}(\mathfrak{B}^2\mathfrak{B}_2^2)}{\mathsf{N}(\mathfrak{B})^3\kappa^\Omega(\mathfrak{B}^2\mathfrak{B}_2^2)} \sum_{\substack{\mathfrak{c}_1, \mathfrak{c}_1, \mathfrak{r} \in \mathcal{I}_r \\ \Omega(\mathfrak{r}^2\mathfrak{c}_1^3\mathfrak{c}_1^3) = N}} \frac{\nu(\mathfrak{r}\mathfrak{c}_1^3)\nu(\mathfrak{r}\mathfrak{c}_1^3)}{\mathsf{N}(\mathfrak{r}^2\mathfrak{c}_1^3\mathfrak{c}_1^3)^{1/2}},
\end{aligned} \tag{43}$$

where

$$N = 2s_r - \Omega(\mathfrak{y}\mathfrak{z}\mathfrak{a}\mathfrak{b}^2\mathfrak{c}^3\mathfrak{A}\mathfrak{B}^2\mathfrak{c}^3\mathfrak{a}_1\mathfrak{b}_1^2\mathfrak{A}_1\mathfrak{B}_1^2).$$

Since

$$\Omega(\mathfrak{y}\mathfrak{z}\mathfrak{a}\mathfrak{b}^2\mathfrak{c}^3\mathfrak{A}\mathfrak{B}^2\mathfrak{c}^3\mathfrak{a}_1\mathfrak{b}_1^2\mathfrak{A}_1\mathfrak{B}_1^2) \leq \Omega(\mathfrak{y}\mathfrak{z})(\mathfrak{a}\mathfrak{b}^2)^2(\mathfrak{A}\mathfrak{B}^2)^2\mathfrak{c}^3\mathfrak{c}^3 \leq 4k\kappa\ell_r$$

we see that $N \geq 2s_r - 4k\kappa\ell_r$. Writing $N = 3q + R$ with $0 \leq R < 3$, it follows that

$$\begin{aligned}
&\sum_{\substack{\mathfrak{c}_1, \mathfrak{c}_1, \mathfrak{r} \in \mathcal{I}_r \\ \Omega(\mathfrak{r}^2\mathfrak{c}_1^3\mathfrak{c}_1^3) = N}} \frac{\nu(\mathfrak{r}\mathfrak{c}_1^3)\nu(\mathfrak{r}\mathfrak{c}_1^3)}{\mathsf{N}(\mathfrak{r}^2\mathfrak{c}_1^3\mathfrak{c}_1^3)^{1/2}} \\
&\leq \sum_{\substack{a, b, c \geq 0 \\ 2a+3b+3c=N}} \sum_{\substack{\mathfrak{r} \in \mathcal{I}_r \\ \Omega(\mathfrak{r})=a}} \frac{\nu(\mathfrak{r})}{\mathsf{N}(\mathfrak{r})} \sum_{\substack{\mathfrak{c}_1 \in \mathcal{I}_r \\ \Omega(\mathfrak{c}_1)=b}} \frac{\nu(\mathfrak{c}_1)}{3^b\mathsf{N}(\mathfrak{c}_1)^{3/2}} \sum_{\substack{\mathfrak{c}_1 \in \mathcal{I}_r \\ \Omega(\mathfrak{c}_1)=c}} \frac{\nu(\mathfrak{c}_1)}{3^c\mathsf{N}(\mathfrak{c}_1)^{3/2}} \\
&= \sum_{\substack{a, b, c \geq 0 \\ 2a+3b+3c=N}} \frac{1}{a!b!c!} \left(\sum_{\mathfrak{p} \in \mathcal{I}_r} \frac{1}{\mathsf{N}\mathfrak{p}} \right)^a \left(\sum_{\mathfrak{p} \in \mathcal{I}_r} \frac{1}{3(\mathsf{N}\mathfrak{p})^{3/2}} \right)^{b+c}.
\end{aligned}$$

Using Lemma 4.1 we have

$$\begin{aligned}
\sum_{\substack{\mathfrak{c}_1, \mathfrak{c}_1, \mathfrak{r} \in \mathcal{I}_r \\ \Omega(\mathfrak{r}^2\mathfrak{c}_1^3\mathfrak{c}_1^3) = N}} \frac{\nu(\mathfrak{r}\mathfrak{c}_1^3)\nu(\mathfrak{r}\mathfrak{c}_1^3)}{\mathsf{N}(\mathfrak{r}^2\mathfrak{c}_1^3\mathfrak{c}_1^3)^{1/2}} &< \sum_{\substack{0 \leq 2a \leq 3q+R \\ 3|R-2a}} \frac{1}{a!(q - \frac{2a-R}{3})!} \left(\sum_{\mathfrak{p} \in \mathcal{I}_r} \frac{1}{\mathsf{N}\mathfrak{p}} \right)^a \left(\frac{2}{3} \right)^{(N-2a)/3} \\
&= \sum_{\substack{0 \leq m \leq q \\ 2|3m+R}} \frac{(2/3)^{q-m}}{\left(\frac{3m+R}{2} \right)!(q-m)!} \left(\sum_{\mathfrak{p} \in \mathcal{I}_r} \frac{1}{\mathsf{N}\mathfrak{p}} \right)^{(3m+R)/2} \\
&\leq \left(\sum_{\mathfrak{p} \in \mathcal{I}_r} \frac{1}{\mathsf{N}\mathfrak{p}} \right)^{s_r} \frac{1}{q!} \sum_{\substack{0 \leq m \leq q \\ 2|3m+R}} \binom{q}{m} \left(\frac{2}{3} \right)^{q-m} \\
&\leq \frac{1}{\lfloor \frac{2s_r-4k\kappa\ell_r}{3} \rfloor!} \left(\sum_{\mathfrak{p} \in \mathcal{I}_r} \frac{1}{\mathsf{N}\mathfrak{p}} \right)^{s_r} \left(\frac{5}{3} \right)^{\lfloor \frac{2s_r-4k\kappa\ell_r}{3} \rfloor}.
\end{aligned}$$

where in the last inequality we used the fact that $(5/3)^q/q!$ is decreasing for $q \geq 1$ and in our case $q = \lfloor N/3 \rfloor \geq 1$ by the choice of s_r . Since the last result is

independent of N , we can bound the remaining terms in (43) independent of the innermost sum. The sum over \mathfrak{B} is bounded by

$$\prod_{\mathfrak{p} \in \mathcal{I}_r} \left(1 + \frac{\nu(\mathfrak{p}^2) \nu_{k\kappa}(\mathfrak{p}^2)}{N(\mathfrak{p})^3 \kappa^{\Omega(\mathfrak{p}^2)}} \sum_{\substack{\mathfrak{B}_1, \mathfrak{B}_2 \\ \mathfrak{p} = \mathfrak{B}_1 \mathfrak{B}_2}} \frac{\nu(\mathfrak{B}_1^2) \nu_{k\kappa}(\mathfrak{B}_2^2)}{\kappa^{\Omega(\mathfrak{B}_2^2)}} \right) = \prod_{\mathfrak{p} \in \mathcal{I}_r} \left(1 + \frac{k^2 + k^4}{8(N\mathfrak{p})^3} \right).$$

The same bound holds for the sum over \mathfrak{b} . As for the sum over \mathfrak{a} , similarly for \mathfrak{A} , we have

$$\begin{aligned} \sum_{\substack{\mathfrak{a} \in \mathcal{I}_r \\ \mathfrak{a} \text{ sq-free} \\ \mathfrak{a} = \mathfrak{a}_1 \mathfrak{a}_2}} \frac{\nu(\mathfrak{a}\mathfrak{a}_1) \nu_{k\kappa}(\mathfrak{a}\mathfrak{a}_2)}{N(\mathfrak{a})^{3/2} \kappa^{\Omega(\mathfrak{a}\mathfrak{a}_2)}} &\leq \sum_{\substack{\mathfrak{a} \in \mathcal{I}_r \\ \mathfrak{a} \text{ sq-free}}} \frac{\nu(\mathfrak{a}) \nu_{k\kappa}(\mathfrak{a})}{N(\mathfrak{a})^{3/2} \kappa^{\Omega(\mathfrak{a})}} \sum_{\substack{\mathfrak{a}_1, \mathfrak{a}_2 \\ \mathfrak{a} = \mathfrak{a}_1 \mathfrak{a}_2}} \frac{\nu(\mathfrak{a}_1) \nu_{k\kappa}(\mathfrak{a}_2)}{\kappa^{\Omega(\mathfrak{a}_2)}} \\ &= \prod_{\mathfrak{p} \in \mathcal{I}_r} \left(1 + \frac{k + k^2}{(N\mathfrak{p})^{3/2}} \right) < \exp \left(\sum_{\mathfrak{p} \in \mathcal{I}_r} \frac{k(1+k)}{N(\mathfrak{p})^{3/2}} \right). \end{aligned}$$

Finally we see that

$$\begin{aligned} &\sum_{\mathfrak{c}_2, \mathfrak{c}_3, \mathfrak{x}, \mathfrak{y}, \mathfrak{z}, \mathfrak{c} \in \mathcal{I}_r} \dots \sum \frac{\nu(\mathfrak{y}\mathfrak{c}^3) \nu(\mathfrak{z}\mathfrak{c}^3) \nu_{k\kappa}(\mathfrak{x}\mathfrak{z}\mathfrak{c}_2^3 \mathfrak{c}_3^3) \nu_{k\kappa}(\mathfrak{x}\mathfrak{y}\mathfrak{c}^3 \mathfrak{c}_3^3)}{N(\mathfrak{c}_2^6 \mathfrak{c}_3^6 \mathfrak{x}^2 \mathfrak{y}^2 \mathfrak{z}^2 \mathfrak{c}_2^3 \mathfrak{c}_3^3)^{1/2} \kappa^{\Omega(\mathfrak{x}^2 \mathfrak{y} \mathfrak{z} \mathfrak{c}^3 \mathfrak{c}_2^3 \mathfrak{c}_3^3)}} \\ &\leq \left(\sum_{\mathfrak{y} \in \mathcal{I}_r} \frac{\nu(\mathfrak{y}) \nu_{k\kappa}(\mathfrak{y})}{N(\mathfrak{y}) \kappa^{\Omega(\mathfrak{y})}} \sum_{\mathfrak{c}_2 \in \mathcal{I}_r} \frac{\nu_{k\kappa}(\mathfrak{c}_2^3)}{N(\mathfrak{c}_2)^{3/2} \kappa^{\Omega(\mathfrak{c}_2^3)}} \sum_{\mathfrak{c} \in \mathcal{I}_r} \frac{\nu(\mathfrak{c}^3) \nu_{k\kappa}(\mathfrak{c}^3)}{N(\mathfrak{c})^3 \kappa^{\Omega(\mathfrak{c}^3)}} \right)^2 \sum_{\mathfrak{x} \in \mathcal{I}_r} \frac{\nu_{k\kappa}(\mathfrak{x}) \nu_{k\kappa}(\mathfrak{x})}{N(\mathfrak{x}) \kappa^{\Omega(\mathfrak{x}^2)}}. \end{aligned}$$

The sum over \mathfrak{x} is

$$\begin{aligned} \sum_{\mathfrak{x} \in \mathcal{I}_r} \frac{\nu_{k\kappa}(\mathfrak{x}) \nu_{k\kappa}(\mathfrak{x})}{N(\mathfrak{x}) \kappa^{\Omega(\mathfrak{x}^2)}} &= \prod_{\mathfrak{p} \in \mathcal{I}_r} \left(1 + \sum_{n \geq 1} \frac{\nu_{k\kappa}(\mathfrak{p}^n) \nu_{k\kappa}(\mathfrak{p}^n)}{N(\mathfrak{p}^n) \kappa^{\Omega(\mathfrak{p}^{2n})}} \right) \\ &= \prod_{\mathfrak{p} \in \mathcal{I}_r} \left(1 + \sum_{n \geq 1} \frac{k^{2n}}{(n!)^2 N(\mathfrak{p})^n} \right) < \exp \left(e^{1/4} k^2 \sum_{\mathfrak{p} \in \mathcal{I}_r} \frac{1}{N(\mathfrak{p})} \right) \end{aligned}$$

since $k^2/N\mathfrak{p} < 1/4$ and for $0 \leq x \leq 1/4$,

$$\sum_{n \geq 1} \frac{x^n}{(n!)^2} \leq e^{1/4} x.$$

Similarly, we see that

$$\begin{aligned} \sum_{\mathfrak{c}_2 \in \mathcal{I}_r} \frac{\nu_{k\kappa}(\mathfrak{c}_2^3)}{N(\mathfrak{c}_2)^{3/2} \kappa^{\Omega(\mathfrak{c}_2^3)}} &= \prod_{\mathfrak{p} \in \mathcal{I}_r} \left(1 + \sum_{n \geq 1} \frac{k^{3n}}{(3n)! N(\mathfrak{p})^{3n/2}} \right) < \exp \left(\sum_{\mathfrak{p} \in \mathcal{I}_r} \frac{0.003k^3}{N(\mathfrak{p})^{3/2}} \right) \\ \sum_{\mathfrak{y} \in \mathcal{I}_r} \frac{\nu(\mathfrak{y}) \nu_{k\kappa}(\mathfrak{y})}{N(\mathfrak{y}) \kappa^{\Omega(\mathfrak{y})}} &= \prod_{\mathfrak{p} \in \mathcal{I}_r} \left(1 + \sum_{n \geq 1} \frac{k^n}{(n!)^2 (N\mathfrak{p})^n} \right) < \exp \left(\sum_{\mathfrak{p} \in \mathcal{I}_r} \frac{ke^{1/(4k)}}{N\mathfrak{p}} \right) \\ \sum_{\mathfrak{c} \in \mathcal{I}_r} \frac{\nu(\mathfrak{c}^3) \nu_{k\kappa}(\mathfrak{c}^3)}{N(\mathfrak{c})^3 \kappa^{\Omega(\mathfrak{c}^3)}} &= \prod_{\mathfrak{p} \in \mathcal{I}_r} \left(1 + \sum_{n \geq 1} \frac{k^{3n}}{((3n)!)^2 N(\mathfrak{p})^{3n}} \right) < \exp \left(\sum_{\mathfrak{p} \in \mathcal{I}_r} \frac{0.003k^3}{N(\mathfrak{p})^3} \right), \end{aligned}$$

by using the inequality

$$\sum_{n \geq 1} \frac{x^n}{(3n)!} < x \sum_{n \geq 1} \frac{1}{(3n)! 8^{n-1}} \quad (0 < x \leq 1/8)$$

and the fact that since $\omega^2 + \omega + 1 = 0$,

$$\begin{aligned} 8 \sum_{n \geq 1} \frac{1}{(3n)! 8^n} &= \frac{8}{3} \left(\sum_{n \geq 1} \frac{(\omega/8)^n}{n!} + \sum_{n \geq 1} \frac{(\omega^2/8)^n}{n!} + \sum_{n \geq 1} \frac{1}{n! 8^n} \right) \\ &= \frac{8}{3} (e^{1/8} + e^{\omega/8} + e^{\omega^2/8} - 3) \approx 0.002604. \end{aligned}$$

Combining all these estimates and using Lemma 4.1 we conclude that

$$(44) \quad |\tilde{\mathcal{E}}(r, c)| < \frac{(2s_r)!}{4^{s_r} \lfloor \frac{2s_r - 4k\kappa\ell_r}{3} \rfloor!} \left(\sum_{\mathfrak{p} \in \mathcal{I}_r} \frac{1}{N\mathfrak{p}} \right)^{s_r} \left(\frac{5}{3} \right)^{\lfloor \frac{2s_r - 4k\kappa\ell_r}{3} \rfloor} \\ \cdot \exp \left(\frac{0.024k^3 + k^2 + k^4}{4 \lfloor X^{\theta_{r-1}} \rfloor^2} + \sum_{\mathfrak{p} \in \mathcal{I}_r} \frac{0.006k^3 + 2k + 2k^2}{(N\mathfrak{p})^{3/2}} + \frac{e^{1/4}k^2 + 2ke^{1/(4k)}}{N\mathfrak{p}} \right).$$

4.5. Bound for $\tilde{\mathcal{E}}(r, c)$ for $j + 1 < r \leq J$. Recall that for $j + 1 < r \leq J$,

$$\tilde{\mathcal{E}}(r, c) = \sum_{\substack{\mathbf{a}_{1r}, \mathbf{b}_{1r} \in \mathcal{I}_r \\ \mathbf{a}_{1r}, \mathbf{b}_{1r} = \overline{\mathfrak{a}\mathfrak{b}} \\ (c, \mathbf{a}_{1r}, \mathbf{b}_{1r}) = 1}} \frac{\lambda(\mathbf{a}_{1r}, \mathbf{b}_{1r}) f(\mathbf{a}_{1r}, \mathbf{b}_{1r}, J)}{N(\mathbf{a}_{1r}, \mathbf{b}_{1r})^{1/2} \kappa^{\Omega(\mathbf{a}_{1r}, \mathbf{b}_{1r})}} \nu_{k\kappa}(\mathbf{a}_{1r}; \ell_r) \nu_{k\kappa}(\mathbf{b}_{1r}; \ell_r).$$

Taking absolute values we have as before

$$(45) \quad \begin{aligned} |\tilde{\mathcal{E}}(r, c)| &\leq \sum_{\substack{\mathbf{a}_{1r}, \mathbf{b}_{1r} \in \mathcal{I}_r \\ \mathbf{a}_{1r}, \mathbf{b}_{1r} = \overline{\mathfrak{a}\mathfrak{b}}}} \frac{\nu_{k\kappa}(\mathbf{a}_{1r}) \nu_{k\kappa}(\mathbf{b}_{1r})}{N(\mathbf{a}_{1r}, \mathbf{b}_{1r})^{1/2} \kappa^{\Omega(\mathbf{a}_{1r}, \mathbf{b}_{1r})}} \\ &= \sum_{\mathfrak{a} \in \mathcal{I}_r} \frac{1}{N(\mathfrak{a})^{1/2} \kappa^{\Omega(\mathfrak{a})}} \sum_{\substack{\mathbf{a}_{1r}, \mathbf{b}_{1r} \in \mathcal{I}_r \\ \mathfrak{a} = \mathbf{a}_{1r} \mathbf{b}_{1r} \\ \mathfrak{a} \mathbf{b}_{1r} = \overline{\mathfrak{a}\mathfrak{b}}}} \nu_{k\kappa}(\mathbf{a}_{1r}) \nu_{k\kappa}(\mathbf{b}_{1r}) \\ &= \prod_{\mathfrak{p} \in \mathcal{I}_r} \left(1 + \sum_{n \geq 1} \frac{1}{N(\mathfrak{p})^{n/2} \kappa^n} \sum_{\substack{0 \leq m \leq n \\ 3|m+n}} \nu_{k\kappa}(\mathfrak{p}^{n-m}) \nu_{k\kappa}(\mathfrak{p}^m) \right) \\ &< \prod_{\mathfrak{p} \in \mathcal{I}_r} \left(1 + \sum_{n \geq 2} \frac{(2k)^n}{n! N(\mathfrak{p})^{n/2}} \right) \leq \exp \left(4(e-2)k^2 \sum_{\mathfrak{p} \in \mathcal{I}_r} \frac{1}{N\mathfrak{p}} \right). \end{aligned}$$

4.6. Moments of $L(s, \chi_c)$.

Lemma 4.3. *For any complex numbers $z_{\mathfrak{p}}$ with $|z_{\mathfrak{p}}| \leq 1$, any positive integer $l = 3q + r$ with $0 \leq r < 3$, and any $\varepsilon > 0$,*

$$\begin{aligned} \sum_{c \in \mathcal{F}(X)} \left| \sum_{3 < N\mathfrak{p} \leq y} \frac{\chi_c(\mathfrak{p}) z_{\mathfrak{p}}}{N(\mathfrak{p})^{1/2}} \right|^{2l} &< |\mathcal{F}(X)| \left(\max \left\{ 1, \sum_{3 < N\mathfrak{p} \leq y} \frac{|z_{\mathfrak{p}}|^2}{N\mathfrak{p}} \right\} \right)^l \frac{(l!)^2 (19/9)^{2(l-r)/3}}{(2(l-r)/3)!} \\ &+ O \left(X^{1/2+\varepsilon} \left(\sum_{3 < N\mathfrak{p} \leq y} \frac{|z_{\mathfrak{p}}|}{N(\mathfrak{p})^{1/2-\varepsilon}} \right)^{2l} \right). \end{aligned}$$

Proof. Put $\mathcal{I} = (3, y]$. Then,

$$\left| \sum_{\mathfrak{p} \in \mathcal{I}} \frac{\chi_c(\mathfrak{p}) z_{\mathfrak{p}}}{N(\mathfrak{p})^{1/2}} \right|^{2l} = (l!)^2 \sum_{\substack{\mathbf{a}_1, \mathbf{a}_2 \in \mathcal{I} \\ \Omega(\mathbf{a}_i) = l}} \frac{\chi_c(\mathbf{a}_1 \mathbf{a}_2^2)}{N(\mathbf{a}_1 \mathbf{a}_2)^{1/2}} \nu(\mathbf{a}_1) \nu(\mathbf{a}_2) z_{\mathbf{a}_1} \overline{z_{\mathbf{a}_2}},$$

and summing over $c \in \mathcal{F}(X)$ we get

$$(|l|)^2 \sum_{\substack{\mathbf{a}_1, \mathbf{a}_2 \in \mathcal{I} \\ \Omega(\mathbf{a}_i) = l}} \frac{\nu(\mathbf{a}_1)\nu(\mathbf{a}_2)z_{\mathbf{a}_1}\overline{z_{\mathbf{a}_2}}}{N(\mathbf{a}_1\mathbf{a}_2)^{1/2}} \sum_{\substack{c \in \mathcal{F}(X) \\ (c, \mathbf{a}_1\mathbf{a}_2) = 1}} \chi_c(\mathbf{a}_1\mathbf{a}_2^2).$$

When $\mathbf{a}_1\mathbf{a}_2^2 = \square$, this sum is trivially bounded by

$$|\mathcal{F}(X)|(|l|)^2 \sum_{\substack{\mathbf{a}_1, \mathbf{a}_2 \in \mathcal{I} \\ \Omega(\mathbf{a}_i) = l \\ \mathbf{a}_1\mathbf{a}_2^2 = \square}} \frac{\nu(\mathbf{a}_1)\nu(\mathbf{a}_2)|z_{\mathbf{a}_1}||z_{\mathbf{a}_2}|}{N(\mathbf{a}_1\mathbf{a}_2)^{1/2}}.$$

Otherwise, applying Lemma 2.9 with $\varsigma = 0$ gives

$$\sum_{\substack{c \in \mathcal{F}(X) \\ (c, \mathbf{a}_1\mathbf{a}_2) = 1}} \chi_c(\mathbf{a}_1\mathbf{a}_2^2) \ll X^{1/2+\varepsilon} N(\mathbf{a}_1\mathbf{a}_2)^\varepsilon,$$

which then leads to the claimed error term. For the case of cubes, replacing $\mathbf{a}_1, \mathbf{a}_2$ by $\mathbf{b}\mathbf{a}_1, \mathbf{b}\mathbf{a}_2$ with $(\mathbf{a}_1, \mathbf{a}_2) = 1$ we see that $\mathbf{a}_1\mathbf{b}\mathbf{a}_2^2\mathbf{b}^2 = \square$ if and only if $\mathbf{a}_1, \mathbf{a}_2$ are cubes. Thus,

$$\begin{aligned} \sum_{\substack{\mathbf{a}_1, \mathbf{a}_2 \in \mathcal{I} \\ \Omega(\mathbf{a}_i) = l \\ \mathbf{a}_1\mathbf{a}_2^2 = \square}} \frac{\nu(\mathbf{a}_1)\nu(\mathbf{a}_2)|z_{\mathbf{a}_1}||z_{\mathbf{a}_2}|}{N(\mathbf{a}_1\mathbf{a}_2)^{1/2}} &\leq \sum_{\substack{\mathbf{b} \in \mathcal{I} \\ 0 \leq \Omega(\mathbf{b}) \leq l \\ 3|l - \Omega(\mathbf{b})}} \frac{|z_{\mathbf{b}}|^2 \nu(\mathbf{b})}{N(\mathbf{b})} \left(\sum_{\substack{\mathbf{a} \in \mathcal{I} \\ \Omega(\mathbf{b}\mathbf{a}^3) = l}} \frac{\nu(\mathbf{a}^3)|z_{\mathbf{a}}|^3}{N(\mathbf{a})^{3/2}} \right)^2 \\ &< \sum_{\substack{\mathbf{b} \in \mathcal{I} \\ 0 \leq \Omega(\mathbf{b}) \leq l \\ 3|l - \Omega(\mathbf{b})}} \frac{|z_{\mathbf{b}}|^2 \nu(\mathbf{b})}{9^{(l - \Omega(\mathbf{b}))/3} \left(\left(\frac{l - \Omega(\mathbf{b})}{3} \right)! \right)^2 N(\mathbf{b})} \end{aligned}$$

since

$$\begin{aligned} \sum_{\substack{\mathbf{a} \in \mathcal{I} \\ \Omega(\mathbf{b}\mathbf{a}^3) = l}} \frac{\nu(\mathbf{a}^3)|z_{\mathbf{a}}|^3}{N(\mathbf{a})^{3/2}} &\leq 3^{-(l - \Omega(\mathbf{b}))/3} \sum_{\substack{\mathbf{a} \in \mathcal{I} \\ \Omega(\mathbf{a}) = (l - \Omega(\mathbf{b}))/3}} \frac{\nu(\mathbf{a})}{N(\mathbf{a})^{3/2}} \\ &= \frac{1}{3^{(l - \Omega(\mathbf{b}))/3} \left(\frac{l - \Omega(\mathbf{b})}{3} \right)!} \left(\sum_{\mathbf{p} \in \mathcal{I}} \frac{1}{N(\mathbf{p})^{3/2}} \right)^{(l - \Omega(\mathbf{b}))/3} \end{aligned}$$

and by Lemma 4.1 the last sum over primes is less than 1. Here, we also used the inequalities $\nu(ab) \leq \nu(a)\nu(b)$, $\nu(b)^2 \leq \nu(b)$, $\nu(a^3) \leq 3^{-\Omega(a)}\nu(a)$, and ignored the condition that $(a_1, a_2) = 1$. Writing $l = 3q + r$ with $0 \leq r < 3$, the sum over b becomes

$$\begin{aligned} &\sum_{0 \leq j \leq q} \frac{1}{9^{q-j} ((q-j)!)^2} \sum_{\substack{\mathbf{b} \in \mathcal{I} \\ \Omega(\mathbf{b}) = 3j+r}} \frac{|z_{\mathbf{b}}|^2 \nu(\mathbf{b})}{N(\mathbf{b})} \\ &= \sum_{0 \leq j \leq q} \frac{1}{9^{q-j} ((q-j)!)^2 (3j+r)!} \left(\sum_{3 < N\mathbf{p} \leq y} \frac{|z_{\mathbf{p}}|^2}{N\mathbf{p}} \right)^{3j+r} \\ &\leq \left(\max \left\{ 1, \sum_{3 < N\mathbf{p} \leq y} \frac{|z_{\mathbf{p}}|^2}{N\mathbf{p}} \right\} \right)^l \frac{(1+1+1/9)^{2(l-r)/3}}{(2(l-r)/3)!}. \end{aligned}$$

This gives the desired result. \square

The next result used in the proof of Theorem 1.2 is not optimal, but is sufficient for our purposes in this paper.

Proposition 4.4. *For a real number $k > 0$,*

$$\sum_{c \in \mathcal{F}(X)} |L(1/2, \chi_c)|^{2k} \ll |\mathcal{F}(X)| \exp(2k(\log \log X)^2).$$

Proof. Put

$$N(v) = |\{c \in \mathcal{F}(X) : L(1/2, \chi_c) \neq 0 \wedge \log |L(1/2, \chi_c)| \geq v\}|.$$

Then, since trivially $N(v) \leq |\mathcal{F}(X)|$,

$$\begin{aligned} \sum_{c \in \mathcal{F}(X)} |L(1/2, \chi_c)|^{2k} &= \int_{-\infty}^{-\infty} e^{2kv} dN(v) = 2k \int_{-\infty}^{\infty} e^{2kv} N(v) dv \\ (46) \quad &= 2k \int_z^{\infty} e^{2kv} N(v) dv + O(e^{2kz} |\mathcal{F}(X)|) \end{aligned}$$

where $z = (\log \log X)^2$.

Taking $\lambda = 1$ and $t = 0$ in Lemma 3.1 shows that for any $x \geq 3$,

$$\begin{aligned} \log |L(1/2, \chi_c)| &< \operatorname{Re} \sum_{1 < \mathbf{N}\mathbf{a} \leq x} \frac{\Lambda(\mathbf{a}) \chi_c(\mathbf{a})}{\mathbf{N}(\mathbf{a})^{1/2+1/\log x} \log \mathbf{N}\mathbf{a}} \frac{\log(x/\mathbf{N}\mathbf{a})}{\log x} \\ (47) \quad &+ \frac{\log n_c}{\log x} + \frac{3}{x^{1/2} \log^2 x}. \end{aligned}$$

If $L(1/2, \chi_c) \neq 0$ and $\log |L(1/2, \chi_c)| \geq v$ for some $c \in \mathcal{F}(X)$, then taking $x = 3$ yields

$$v \leq \log |L(1/2, \chi_c)| < \frac{\log n_c}{\log 3} + \frac{3^{1/2}}{\log^2 3} < 1.44 + \log X.$$

Thus, we shall assume from now on that $z \leq v < \log X + 2$.

Choose x such that

$$\log x = \frac{A}{v} \log X, \quad A = A(v) = \log \log v.$$

Note that

$$\log x > \frac{\log \log(2 + \log X)}{2 + \log X} \log X \gg \log \log \log X$$

as $A(v)/v$ is decreasing for $v \geq z$.

Since

$$\sum_{\substack{(\mathbf{N}\mathbf{p})^k \leq x \\ k \geq 3}} \frac{1}{k \mathbf{N}\mathbf{p}^{k/2}} \leq \frac{1}{3(1 - 3^{-1/2})} \sum_{\mathbf{N}\mathbf{p} \geq 3} \frac{1}{(\mathbf{N}\mathbf{p})^{3/2}} < 1,$$

it follows from (71) that

$$\begin{aligned} v \leq \log |L(1/2, \chi)| &< \operatorname{Re} \sum_{3 < \mathbf{N}\mathbf{p} \leq x} \frac{\chi(\mathbf{p})}{\mathbf{N}(\mathbf{p})^{1/2+1/\log x}} \frac{\log(x/\mathbf{N}\mathbf{p})}{\log x} \\ &+ \operatorname{Re} \sum_{3 < \mathbf{N}\mathbf{p} \leq \sqrt{x}} \frac{\chi(\mathbf{p}^2)}{2\mathbf{N}(\mathbf{p})^{1+2/\log x}} \frac{\log(x/\mathbf{N}\mathbf{p}^2)}{\log x} + \frac{v}{A} + O(1). \end{aligned}$$

Furthermore, as $\sqrt{x} \geq 3$ is satisfied with our choice of x , we have

$$\sum_{3 < N\mathbf{p} \leq \sqrt{x}} \frac{\chi(\mathbf{p}^2)}{2N(\mathbf{p})^{1+2/\log x}} \frac{\log(x/N\mathbf{p}^2)}{\log x} \leq \sum_{p \leq \sqrt{x}} \frac{1}{p} = \log \log x + O(1),$$

which implies that

$$\operatorname{Re} \sum_{3 < N\mathbf{p} \leq x} \frac{\chi_c(\mathbf{p})}{N(\mathbf{p})^{1/2+1/\log x}} \frac{\log(x/N\mathbf{p})}{\log x} > v - \frac{v}{A} - \log \log x + O(1) > v - \frac{2v}{A} > 0$$

for $v \geq z$ since

$$\log \log x < \log \log X < \frac{z}{\log \log z} \leq \frac{v}{A}$$

for sufficiently large X ($X > 181$ works). Applying Lemma 4.3 with

$$z_{\mathbf{p}} = \frac{\log(x/N\mathbf{p})}{\log x N(\mathbf{p})^{1/\log x}}$$

we derive the estimate

$$\begin{aligned} N(v) &< v_1^{-2l} \sum_{c \in \mathcal{F}(X)} \left| \sum_{3 < N\mathbf{p} \leq x} \frac{\chi_c(\mathbf{p}) z_{\mathbf{p}}}{N(\mathbf{p})^{1/2}} \right|^{2l} \\ &\ll v_1^{-2l} |\mathcal{F}(X)| (2 \log \log x)^l \frac{(l!)^2 (19/9)^{2(l-r)/3}}{(2(l-r)/3)!} + v_1^{-2l} X^{1/2+\varepsilon} x^{2l(1/2+\varepsilon)}, \end{aligned}$$

where $v_1 = v - 2v/A$. Using (21) gives for $l \geq 3$ and $3 \mid l$ that

$$\frac{(l!)^2 (19/9)^{2(l-r)/3}}{(2(l-r)/3)!} < \left(\frac{19}{6e^2} \right)^{2l/3} l^{4l/3} \sqrt{3\pi l} \exp\left(\frac{1}{6l} - \frac{1}{8l+1} \right) < l^{4l/3}.$$

Thus, choosing $l = 3 \lfloor v/(9A) \rfloor$, we deduce for $\varepsilon < 1/(12)$ that

$$\begin{aligned} (48) \quad N(v) &\ll |\mathcal{F}(X)| \exp\left(l(\log(2 \log \log x) + \frac{4}{3} \log l - 2 \log v_1) \right) \\ &\quad + X \exp\left(2l(1/2 + \varepsilon) \log x - 2l \log v_1 - (1/2 - \varepsilon) \log X \right) \\ &\ll |\mathcal{F}(X)| \exp\left(-\frac{v \log v}{22A} \right) + X \exp\left(-\frac{v \log v}{2A} \right) \end{aligned}$$

since using the inequality $-\log(1-x) < x/(1-x)$ we have that

$$\begin{aligned} \log(2 \log \log x) + \frac{4}{3} \log l - 2 \log v_1 &< \log 2 + \frac{1}{2} \log z - \frac{2}{3} \log v + \frac{4}{A-2} \\ &\leq -\frac{1}{6} \log v + \frac{4}{\log \log z - 2} < -\frac{1}{7} \log v, \end{aligned}$$

and

$$\begin{aligned} &2l(1/2 + \varepsilon) \log x - 2l \log v_1 - (1/2 - \varepsilon) \log X \\ &< \frac{2v}{3A} (1/2 + \varepsilon) \frac{A}{v} \log X - 2l \log v_1 - (1/2 - \varepsilon) \log X \\ &< (2\varepsilon - 1/6) \log X - 6 \lfloor v/(9A) \rfloor \log v_1 < -6 \lfloor v/(9A) \rfloor \left(\log v - \frac{2}{A-2} \right) \\ &< -5(v/(9A) - 1) \log v < -\frac{v \log v}{2A}. \end{aligned}$$

Inserting (48) in (46) we get the claimed result. \square

4.7. **Squares of Primes.** In this part we prove the following result.

Lemma 4.5. For $0 \leq j \leq J$ and X satisfying (55), the estimate

$$(49) \quad \sum_{c \in \mathcal{F}(X)} (S_{j,k}(c))^2 \ll \mathcal{S}_k |\mathcal{F}(X)|$$

holds, where

$$(50) \quad \mathcal{S}_k = 4 \cdot 5^{2kD} (2kD)! \exp(4k(1+D)),$$

with D given in (53), provided that

$$(51) \quad \Theta < \frac{1}{8kD + 4 \log(16\sqrt{5})}.$$

Remark 4.6. For $k = 2, \kappa = 1$, one can take $D = 1$, in which case

$$\mathcal{S}_2 \cong 5.3316663123 \times 10^{11}.$$

This follows by using the inequality (see [23, Section 3, Theorem 5])

$$\sum_{x < p \leq 5x} \frac{1}{p} < \log \log(5x) - \log \log x + \frac{1}{\log^2 x},$$

which holds for $x \geq 286$ (where we take $x = 8 \cdot 5^{n-1}$ and $n \geq 3$), and the fact that $n \sum_{p \in \mathcal{I}_n} 1/p$ for $n = 1, 2$ are 0.416533, 0.353, respectively.

Proof. Recall that $S_{j,k}(c) = \exp(k \operatorname{Re} K(c, j))$, where

$$K(c, j) = \sum_{\mathfrak{p} \in \mathcal{I}} \frac{\chi_c(\mathfrak{p}^2) B_j(\mathfrak{p})}{N\mathfrak{p}}, \quad B_j(\mathfrak{p}) = \frac{1}{2N(\mathfrak{p})^{2/(\theta_j \log X)}} \left(1 - \frac{2 \log N\mathfrak{p}}{\theta_j \log X}\right),$$

and $\mathcal{I} = (k_0, X^{\theta_j/2}]$. Note that $B_j(\mathfrak{p}) \leq 1/2$. We extend B_j to be a completely multiplicative function below.

Define the intervals

$$\mathcal{I}_m = (k_0 5^{m-1}, k_0 5^m] \cap \mathcal{I}$$

for $m = 1, \dots, M$ with

$$M = \left\lceil \frac{\frac{1}{2} \theta_j \log X - \log k_0}{\log 5} \right\rceil.$$

Let

$$Q_m = \{c \in \mathcal{F}(X) : |\operatorname{Re} K_m(c, j)| > 2^{-m}, |\operatorname{Re} K_n(c, j)| \leq 2^{-n} \text{ for } m < n \leq M\}$$

where

$$K_n(c, j) = \sum_{\mathfrak{p} \in \mathcal{I}_n} \frac{\chi_c(\mathfrak{p}^2) B_j(\mathfrak{p})}{N\mathfrak{p}}.$$

Then,

$$\sum_{c \in \mathcal{F}(X)} (S_{j,k}(c))^2 \leq \sum_{c \in \mathcal{F}(X) \setminus \cup_m Q_m} (S_{j,k}(c))^2 + \sum_{m=1}^M \sum_{c \in Q_m} (S_{j,k}(c))^2.$$

For the first sum on the right, we have

$$(52) \quad \sum_{c \in \mathcal{F}(X) \setminus \cup_m Q_m} (S_{j,k}(c))^2 \leq e^{2k} |\mathcal{F}(X)|.$$

If $c \in Q_m$ for some m , then we have

$$\operatorname{Re} K(c, j) = \sum_{n=1}^M \operatorname{Re} K_n(c, j) < \sum_{n=1}^m \frac{D}{n} + \sum_{n>m} 2^{-n} < D(2 + \log m) + \frac{1}{2^m},$$

where we used the estimate

$$(53) \quad \sum_{\mathfrak{p} \in \mathcal{I}_n} \frac{B_j(\mathfrak{p})}{N\mathfrak{p}} \leq \sum_{p \in \mathcal{I}_n} \frac{1}{p} < \frac{D}{n}$$

for some positive integer D . Therefore,

$$\sum_{c \in Q_m} (S_{j,k}(c))^2 \leq \sum_{c \in \mathcal{F}(X)} \exp\left(2k\left(\frac{1}{2^m} + D(2 + \log m)\right)\right) (2^m \operatorname{Re} K_m(c, j))^4.$$

Noting that

$$(\operatorname{Re} K_m(c, j))^4 = \frac{4!}{2^4} \sum_{\substack{\mathfrak{a}, \mathfrak{b} \in \mathcal{I}_m \\ \Omega(\mathfrak{ab})=4}} \frac{\nu(\mathfrak{a})\nu(\mathfrak{b})B_j(\mathfrak{ab})}{N(\mathfrak{ab})} \chi_c(\mathfrak{ab}^2),$$

and using Lemma 2.9 with $\varsigma = 0$, we deduce that

$$\begin{aligned} \sum_{c \in \mathcal{F}(X)} (\operatorname{Re} K_m(c, j))^4 &= \frac{3}{2} \sum_{\substack{\mathfrak{a}, \mathfrak{b} \in \mathcal{I}_m \\ \Omega(\mathfrak{ab})=4}} \frac{\nu(\mathfrak{a})\nu(\mathfrak{b})B_j(\mathfrak{ab})}{N(\mathfrak{ab})} \sum_{c \in \mathcal{F}(X)} \chi_c(\mathfrak{ab}^2) \\ &\ll |\mathcal{F}(X)| \sum_{\substack{\mathfrak{a}, \mathfrak{b} \in \mathcal{I}_m \\ \Omega(\mathfrak{ab})=4 \\ \mathfrak{ab}^2 = \square}} \frac{\nu(\mathfrak{a})\nu(\mathfrak{b})B_j(\mathfrak{ab})}{N(\mathfrak{ab})} + X^{1/2+\varepsilon} \left(\sum_{\mathfrak{p} \in \mathcal{I}_m} \frac{B_j(\mathfrak{p})}{(N\mathfrak{p})^{1-\varepsilon}} \right)^4. \end{aligned}$$

Now replacing $\mathfrak{a}, \mathfrak{b}$ by $\mathfrak{ca}^3, \mathfrak{cb}^3$ with $(\mathfrak{a}, \mathfrak{b}) = 1$, using $B_j(\mathfrak{p}) \leq 1/2$, we see that

$$\begin{aligned} \sum_{\substack{\mathfrak{a}, \mathfrak{b} \in \mathcal{I}_m \\ \Omega(\mathfrak{ab})=4 \\ \mathfrak{ab}^2 = \square}} \frac{\nu(\mathfrak{a})\nu(\mathfrak{b})B_j(\mathfrak{ab})}{N(\mathfrak{ab})} &\leq \sum_{\substack{\mathfrak{c} \in \mathcal{I}_m \\ \Omega(\mathfrak{c})=2}} \frac{\nu(\mathfrak{c})B_j(\mathfrak{c}^2)}{(N\mathfrak{c})^2} = \frac{1}{2} \left(\sum_{\mathfrak{p} \in \mathcal{I}_m} \frac{B_j(\mathfrak{p}^2)}{N(\mathfrak{p}^2)} \right)^2 \\ &\leq \frac{1}{8} \left(\sum_{p > 5^{m-1}k_0} \frac{1}{p^2} \right)^2 \leq \frac{1}{8} \left(\int_{3 \cdot 5^{m-1}}^{\infty} x^{-2} dx \right)^2 = \frac{1}{72 \cdot 5^{2m-2}}. \end{aligned}$$

Using this bound together with the estimate

$$\sum_{\mathfrak{p} \in \mathcal{I}_m} \frac{B_j(\mathfrak{p})}{(N\mathfrak{p})^{1-\varepsilon}} \leq \sum_{p \in \mathcal{I}_m} \frac{1}{p^{1-\varepsilon}} \leq \int_1^{k_0 5^m} x^{\varepsilon-1} dx < \frac{1}{\varepsilon} (k_0 5^m)^\varepsilon,$$

we conclude that

$$\begin{aligned} \sum_{1 \leq m \leq M} \sum_{c \in Q_m} (S_{j,k}(c))^2 &\ll |\mathcal{F}(X)| \sum_{1 \leq m \leq M} \exp\left(2k\left(\frac{1}{2^m} + D(2 + \log m)\right)\right) (4/5)^m \\ &\quad + X^{1/2+\varepsilon} \sum_{1 \leq m \leq M} \exp\left(2k\left(\frac{1}{2^m} + D(\log m + 2)\right)\right) 2^{4m} \left(\frac{1}{\varepsilon} (k_0 5^m)^\varepsilon\right)^4. \end{aligned}$$

The first term on the right is bounded by

$$(54) \quad 4 \cdot 5^{2kD} (2kD)! \exp(2k + 4kD) |\mathcal{F}(X)|,$$

where we used the inequality

$$\sum_{m \geq 1} m^A x^m \leq \frac{A!x}{(1-x)^{A+1}}$$

which holds for $x < 1$.

The second term is bounded by

$$\exp\left((1/2 + \varepsilon) \log X + 2k(1 + D(2 + \log M)) + \log \frac{\varepsilon^{-4} k_0^{4\varepsilon}}{2^4 5^{4\varepsilon} - 1} + (M+1) \log(2^4 5^{4\varepsilon})\right).$$

We choose $\varepsilon = 1/8$. Then, for X satisfying

$$(55) \quad 2k + 4kD + \log \frac{8^4 k_0^{1/2}}{16\sqrt{5} - 1} < \frac{1}{8} \log X,$$

the second term is $\ll X$ provided that (51) holds. Combining (52) and (54) the claim in (49) follows. \square

5. FIRST MOLLIFIED MOMENT

5.1. Approximate Functional Equation.

Lemma 5.1. *Let χ be a primitive ray class character, $G(u)$ be any even function, holomorphic and bounded in the region $|\operatorname{Re}(u)| < 3$ with $G(0) = 1$, and $Y > 0$ be a constant real number. Then, for any complex number s with $\operatorname{Re}(s) \in [0, 1]$,*

$$\begin{aligned} L(s, \chi) &= \sum_{\mathfrak{b}} \chi(\mathfrak{b})(N\mathfrak{b})^{-s} V_s \left(\frac{N\mathfrak{b}}{Y\sqrt{3n_\chi}} \right) \\ &\quad + (2\pi)^{2s-1} (3n_\chi)^{(1/2-s)} \frac{W(\chi)}{\sqrt{n_\chi}} \sum_{\mathfrak{b}} \frac{\overline{\chi(\mathfrak{b})}}{(N\mathfrak{b})^{1-s}} V_{1-s} \left(\frac{Y N\mathfrak{b}}{\sqrt{3n_\chi}} \right), \end{aligned}$$

where

$$V_s(y) = \frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} G(u) (2\pi y)^{-u} \frac{\Gamma(s+u)}{\Gamma(s)} \frac{du}{u}.$$

Remark 5.2. For $G(u) = 1$, $s = 1/2$, $0 < \alpha \leq 1/2 - \varepsilon$ with $\varepsilon < 1/2$, and $A > 0$, it follows from Stirling's formula for the Gamma function (see, for example, [14, Eqn. 5.112]) by shifting the contour to $\operatorname{Re} u = -\alpha$ and to $\operatorname{Re} u = A$, respectively, that

$$(56) \quad \begin{aligned} y^a V_{1/2}^{(a)}(y) &= \delta_{0,a} + O_{\varepsilon,\alpha}(y^\alpha) \\ y^a V_{1/2}^{(a)}(y) &\ll_A y^{-A} \end{aligned}$$

for any integer $a \geq 0$, where $\delta_{0,a}$ is the Kronecker delta function.

Proof. By the residue theorem,

$$\Lambda(s, \chi) = \frac{1}{2\pi i} \int_R G(u) \Lambda(s+u, \chi) Y^u \frac{du}{u},$$

where we use counterclockwise orientation over the rectangle R with vertices $\pm 2 \pm iT$ and $T > 0$ does not correspond to the ordinate of a zero of $G(u) \Lambda(s+u, \chi)$. By Stirling's formula for $\Gamma(s)$ and the classical convexity bound, it follows that the horizontal integrals vanish as $T \rightarrow \infty$. For the vertical integral at $\operatorname{Re}(u) = 2$, we have

$$\frac{1}{2\pi i} \int_{(2)} G(u)(2\pi)^{-s-u}(3n_\chi)^{(s+u)/2} \Gamma(s+u) \left(\sum_{\mathfrak{b}} \frac{\chi(\mathfrak{b})}{N(\mathfrak{b})^{s+u}} \right) \frac{Y^u}{u} du$$

while for the vertical integral at $\operatorname{Re}(u) = -2$ we change u to $-u$ and use the functional equation to get

$$\begin{aligned} \frac{1}{2\pi i} \int_{(2)} G(u)\Lambda(s-u, \chi)Y^{-u} \frac{du}{u} &= \frac{W(\chi)}{\sqrt{n_\chi}} \frac{1}{2\pi i} \int_{(2)} G(u)\Lambda(1-s+u, \bar{\chi})Y^{-u} \frac{du}{u} \\ &= \frac{W(\chi)}{\sqrt{n_\chi}} \frac{1}{2\pi i} \int_{(2)} G(u)(2\pi)^{s-u-1}(3n_\chi)^{(1-s+u)/2} \Gamma(1-s+u) \\ &\quad \cdot \left(\sum_{\mathfrak{b}} \frac{\overline{\chi(\mathfrak{b})}}{N(\mathfrak{b})^{1-s+u}} \right) Y^{-u} \frac{du}{u}. \end{aligned}$$

The result follows by combining the two integrals and dividing both sides by $(3n_\chi)^{s/2}(2\pi)^{-s}\Gamma(s)$. \square

5.2. First Mollified Moment. Using the definition of the mollifier in (17) we see that

$$M(c, \kappa) = \prod_{j=0}^J \sum_{\substack{\mathfrak{a} \in \mathcal{I}_j \\ 0 \leq \Omega(\mathfrak{a}) \leq \ell_j}} \frac{\chi_c(\mathfrak{a})\lambda(\mathfrak{a})f(\mathfrak{a}, J)\nu(\mathfrak{a})}{N(\mathfrak{a})^{1/2}\kappa^{\Omega(\mathfrak{a})}} = \sum_{\mathfrak{a}} \frac{\chi_c(\mathfrak{a})\lambda(\mathfrak{a})f(\mathfrak{a}, J)v_J(\mathfrak{a})}{N(\mathfrak{a})^{1/2}\kappa^{\Omega(\mathfrak{a})}},$$

where

$$v_J(\mathfrak{a}) = \sum_{\substack{\mathfrak{a}_j \in \mathcal{I}_j \text{ for } 0 \leq j \leq J \\ \mathfrak{a} = \mathfrak{a}_0 \cdots \mathfrak{a}_J \\ 0 \leq \Omega(\mathfrak{a}_j) \leq \ell_j}} \nu(\mathfrak{a}_0) \cdots \nu(\mathfrak{a}_J).$$

The first mollified moment we shall study is given by

$$(57) \quad \sum_{c \in \mathcal{F}(X)} L(1/2, \chi_c) M(c, 1) = \sum_{\mathfrak{a}} \frac{\lambda(\mathfrak{a})f(\mathfrak{a}, J)v_J(\mathfrak{a})}{N(\mathfrak{a})^{1/2}} \sum_{c \in \mathcal{F}(X)} \chi_c(\mathfrak{a}) L(1/2, \chi_c).$$

In Lemma 5.1 taking $s = 1/2$, $\chi = \chi_c$, $G(u) \equiv 1$, replacing Y by $Y(3n_\chi)^{-1/2}$ and factoring each ideal \mathfrak{b} as $\mathfrak{b} = (1-\omega)^{r_2} b \mathbb{Z}[\omega]$ with $b \equiv 1 \pmod{3}$ and $r_2 \geq 0$ gives

$$(58) \quad \begin{aligned} L(1/2, \chi_c) &= \sum_{r_2 \geq 0} 3^{-r_2/2} \sum_{b \equiv 1 \pmod{3}} \frac{\chi_c(b)}{N(b)^{1/2}} V\left(\frac{3^{r_2} N(b)}{Y}\right) \\ &\quad + \frac{W(\chi_c)}{\sqrt{n_c}} \sum_{r_2 \geq 0} 3^{-r_2/2} \sum_{b \equiv 1 \pmod{3}} \frac{\overline{\chi_c(b)}}{N(b)^{1/2}} V\left(\frac{3^{r_2} N(b) Y}{3n_c}\right), \end{aligned}$$

where

$$V(y) = V_{1/2}(y) = \frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} (2\pi y)^{-u} \frac{\Gamma(1/2+u)}{\Gamma(1/2)} \frac{du}{u}.$$

Here we used the fact that $\chi_c(1-\omega) = 1$ whenever $c \equiv 1 \pmod{9}$ for $c \in \mathcal{F}(X)$, which follows from [13][Chapter 9, Theorem 1], Supplement to the Cubic Reciprocity Law.

In what follows, we shall write $\mathfrak{a} = a\mathbb{Z}[\omega]$ with $a \equiv 1 \pmod{3}$. Hence, $\chi_c(\mathfrak{a}\mathfrak{b}) = \chi_c(ab)$. By the cubic reciprocity law, $\chi_c(\mathfrak{a}\mathfrak{b}) = \chi_c(ab) = \chi_{ab}(c)$ unless $ab = 1$.

Substituting (58) into (57) we can write

$$\sum_{c \in \mathcal{F}(X)} \chi_c(a) L(1/2, \chi_c) = S_1 + S_2 + S_3,$$

where

$$\begin{aligned}
S_1 &= \sum_{r_2 \geq 0} 3^{-r_2/2} \sum_{\substack{b \equiv 1 \pmod{3} \\ ab = \square}} \frac{1}{N(b)^{1/2}} V\left(\frac{3^{r_2} N(b)}{Y}\right) \sum_{\substack{c \in \mathcal{F}(X) \\ (c, ab) = 1}} 1 \\
(59) \quad S_2 &= \sum_{r_2 \geq 0} 3^{-r_2/2} \sum_{\substack{b \equiv 1 \pmod{3} \\ ab \neq \square}} \frac{1}{N(b)^{1/2}} V\left(\frac{3^{r_2} N(b)}{Y}\right) \sum_{\substack{c \in \mathcal{F}(X) \\ (c, ab) = 1}} \chi_{ab}(c) \\
S_3 &= \sum_{r_2 \geq 0} 3^{-r_2/2} \sum_{b \equiv 1 \pmod{3}} \frac{1}{N(b)^{1/2}} \sum_{c \in \mathcal{F}(X)} \frac{\chi_c(ab^2) W(\chi_c)}{\sqrt{n_c}} V\left(\frac{3^{r_2} Y N(b)}{3n_c}\right).
\end{aligned}$$

5.3. Estimate of S_1 and S_2 .

Remark 5.3. One can define a ray class character χ modulo $3ab\mathbb{Z}[\omega]$, for any given fractional ideal \mathfrak{a} coprime to $3ab$ by $\chi(\mathfrak{a}) = \chi_{ab}(\alpha)\chi_{ab}(\beta)^{-1}$, where $\mathfrak{a} = \alpha\beta^{-1}\mathbb{Z}[\omega]$ with $\alpha, \beta \equiv 1 \pmod{3}$. It agrees with χ_{ab} on \mathcal{F} . We denote this character by χ_{ab} below.

When $ab = \square$, using Lemma 2.8 with $n = ab$ gives

$$\sum_{\substack{c \in \mathcal{F}(X) \\ (c, ab) = 1}} 1 = C_1(ab)X \log X + C_2(ab)X + O_\varepsilon\left(\left(X^{\frac{4\varsigma+1}{4\varsigma+2}+5\varepsilon} + X^{1/2+5\varepsilon}\right)(N(ab))^\varepsilon\right),$$

provided that (7) holds for some $\varsigma \leq 1/2$. Therefore, using the definition of $V(y)$, we have that

$$\begin{aligned}
(60) \quad S_1 &= \frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} \left(\frac{Y}{2\pi}\right)^u \frac{\Gamma(1/2+u)}{u\Gamma(1/2)} H(u) du \\
&+ O\left(\left(X^{\frac{4\varsigma+1}{4\varsigma+2}+5\varepsilon} + X^{1/2+5\varepsilon}\right) \sum_{r_2 \geq 0} 3^{-r_2/2} \sum_{b \equiv 1 \pmod{3}} \frac{(N(ab))^\varepsilon}{N(b)^{1/2}} \left|V\left(\frac{3^{r_2} N(b)}{Y}\right)\right|\right),
\end{aligned}$$

where

$$H(u) = \sum_{r_2 \geq 0} 3^{-r_2(1/2+u)} \sum_{\substack{b \equiv 1 \pmod{3} \\ ab = \square}} \frac{C_1(ab)X \log X + C_2(ab)X}{N(b)^{1/2+u}},$$

and C_1, C_2 were defined in (8). Since ab is a cube, writing $a = a_1 a_2^2 a_3^3$, where a_1, a_2 are square-free and co-prime, we can replace b above by $a_1^2 a_2 b^3$ (with a different b) to get

$$(61) \quad H(u) = \sum_{r_2 \geq 0} 3^{-r_2(1/2+u)} \sum_{b \equiv 1 \pmod{3}} \frac{C_1(a_1 a_2 a_3 b)X \log X + C_2(a_1 a_2 a_3 b)X}{N(a_1^2 a_2 b^3)^{1/2+u}}.$$

Using (10) we have for any $\alpha \in \mathbb{Z}[\omega]$ that

$$\frac{F'_{\psi_0}(1; \alpha)}{F_{\psi_0}(1; \alpha)} = \sum_{\substack{\pi \equiv 1 \pmod{3} \\ \pi \nmid \alpha}} \frac{6 \log N(\pi) \left(1 - \frac{1}{N(\pi)}\right)}{N(\pi)^2 \left(1 - \frac{3}{N(\pi)^2} + \frac{2}{N(\pi)^3}\right)} + 2 \sum_{\substack{\pi \equiv 1 \pmod{3} \\ \pi | \alpha}} \frac{\log N(\pi)}{N(\pi) - 1}.$$

Furthermore, $F_{\psi_0}(1; \alpha) \leq 1$, and

$$F_\psi(1; \alpha) \ll \prod_{\substack{\pi \equiv 1 \pmod{3} \\ \pi | \alpha}} \left(1 + \frac{1}{N\pi}\right)^2$$

where ψ is any character modulo 9 of order 2. Therefore, for any $\varepsilon > 0$,

$$(62) \quad F_{\psi_0}(1; \alpha) \leq 1, \quad F'_{\psi_0}(1; \alpha), F_{\psi}(1; \alpha) \ll (\mathbf{N}(\alpha))^\varepsilon.$$

This shows that $H(u)$ converges absolutely and uniformly in the region $\operatorname{Re} u \geq -1/6 + \varepsilon$ for any $\varepsilon > 0$. Therefore, moving the contour to $\operatorname{Re} s = -1/6 + \varepsilon$ in (60) and using (62) with $\varepsilon/2$, where $\varepsilon < 1/3$, we pick up the residue $H(0)$ at $s = 0$ and conclude that the integral in (60) equals

$$\begin{aligned} H(0) - \frac{1}{2\pi i} \int_{-1/6+\varepsilon+i\infty}^{-1/6+\varepsilon-i\infty} \left(\frac{Y}{2\pi}\right)^u \frac{\Gamma(1/2+u)}{u\Gamma(1/2)} H(u) du \\ = H(0) + O\left(\frac{X \log X + X(\mathbf{N}(a_3))^{\varepsilon/2}}{Y^{1/6-\varepsilon} \mathbf{N}(a_1^2 a_2)^{1/3+\varepsilon/2}}\right). \end{aligned}$$

As for S_2 , using Lemma 2.9 gives the bound

$$(63) \quad S_2 \ll \sum_{r_2 \geq 0} 3^{-r_2/2} \sum_{\substack{b \equiv 1 \pmod{3} \\ ab \neq 1}} \frac{1}{\mathbf{N}(b)^{1/2}} \left| V\left(\frac{3^{r_2} \mathbf{N}(b)}{Y}\right) \right| \cdot \left(X^{\frac{4s+1}{4s+2}+5\varepsilon} (\mathbf{N}(ab))^{\frac{s}{2s+1}+3\varepsilon} + X^{1/2+\varepsilon} (\mathbf{N}(ab))^{s+3\varepsilon} \right).$$

Finally, using (56) for the error terms in (60) and (63), we conclude that

$$(64) \quad \begin{aligned} S_1 + S_2 - H(0) \ll & (XY)^{\frac{4s+1}{4s+2}+\varepsilon} (\mathbf{N}(a))^{\frac{s}{2s+1}+\varepsilon} + X^{1/2+\varepsilon} Y^{1/2+s+\varepsilon} (\mathbf{N}(a))^{s+\varepsilon} \\ & + \frac{X^{1+\varepsilon} (\mathbf{N}(a_3))^\varepsilon}{Y^{1/6-\varepsilon} \mathbf{N}(a_1^2 a_2)^{1/3+\varepsilon/2}} \end{aligned}$$

upon re-defining ε .

5.4. Estimate of S_3 .

Lemma 5.4. *For small $\varepsilon > 0$, the inequality*

$$(65) \quad S_3 \ll \mathbf{N}(a)^{2/3} \frac{X^{3/2+2\varepsilon}}{Y^{5/6+\varepsilon}} + \mathbf{N}(a)^{2/7+2\varepsilon} \frac{X^{3/2+3\varepsilon}}{Y^{9/14+2\varepsilon}}$$

holds unconditionally.

Proof. We will first show for $3 \leq y \leq X$ and small $\varepsilon > 0$ that

$$(66) \quad \sum_{c \in \mathcal{F}(y)} \chi_c(ab^2) \frac{W(\chi_c)}{\sqrt{n_c}} \ll y^{2/3+\varepsilon} \mathbf{N}(a^2 b)^{1/3} + y^{6/7+\varepsilon} \mathbf{N}(a^2 b)^{1/7+\varepsilon}.$$

Applying Lemma 2.7 and proceeding as in the proof of Lemma 2.9 we see that

$$\begin{aligned} \sum_{c \in \mathcal{F}(y) \cup \{1\}} \chi_c(ab^2) \frac{W(\chi_c)}{\sqrt{n_c}} &= \frac{1}{2\pi i} \int_{\varrho-iT}^{\varrho+iT} y^s \left(\sum_{c \in \mathcal{F} \cup \{1\}} \frac{\chi_c(ab^2) W(\chi_c)}{n_c^{s+1/2}} \right) \frac{ds}{s} \\ &+ O(y^{1+\varepsilon} T^{-1}), \end{aligned}$$

where $\varrho = 1 + 1/(2 \log y)$, and $T \geq 1$. Next, we look at the integral. First note that for $c = c_1^2 c_2 \in \mathcal{F}$,

$$W(\chi_c) = \chi_{c_1^2}(c_2) \chi_{c_2}(c_1) \overline{W(\chi_{c_1})} W(\chi_{c_2}) = \overline{W(\chi_{c_1})} W(\chi_{c_2}),$$

where the second equality follows by cubic reciprocity law. Note also that

$$W(\chi_{c_i}) = \chi_{c_i}(\sqrt{-3}) g(1, c_i).$$

Hence,

$$W(\chi_c) = \chi_c(\sqrt{-3})\overline{g(1, c_1)}g(1, c_2) = \overline{g(1, c_1)}g(1, c_2),$$

since $\chi_c(\sqrt{-3}) = \chi_c(w(1 - \omega)) = 1$ whenever $c \equiv 1 \pmod{9}$. It follows that

$$\begin{aligned} \sum_{c \in \mathcal{F}} \chi_c(ab^2) \frac{W(\chi_c)}{(n_c)^{1/2+s}} &= \sum_{\substack{c_1 \equiv 1 \pmod{3} \\ c_1 \text{ square-free} \\ (c_1, ab)=1}} \sum_{\substack{c_2 \equiv 1 \pmod{3} \\ c_2 \text{ square-free} \\ c_1^2 c_2 \equiv 1 \pmod{9} \\ (c_2, abc_1)=1}} \chi_{c_1^2 c_2}(ab^2) \frac{\overline{g(1, c_1)}g(1, c_2)}{N(c_1 c_2)^{1/2+s}} \\ &= \sum_{\substack{c_1 \equiv 1 \pmod{3} \\ (c_1, ab)=1}} \chi_{c_1^2}(ab^2) \frac{\overline{g(1, c_1)}}{N(c_1)^{1/2+s}} \sum_{\substack{c_2 \equiv 1 \pmod{3} \\ c_1^2 c_2 \equiv 1 \pmod{9} \\ (c_2, abc_1)=1}} \frac{g(a^2 b, c_2)}{N(c_2)^{1/2+s}} \\ &= \frac{1}{h_{(9)}} \sum_{\psi \pmod{9}} \sum_{\substack{c_1 \equiv 1 \pmod{3} \\ (c_1, ab)=1}} (\chi_{ab^2} \psi)(c_1^2) \frac{\overline{g(1, c_1)}}{N(c_1)^{1/2+s}} \sum_{\substack{c_2 \equiv 1 \pmod{3} \\ (c_2, abc_1)=1}} \psi(c_2) \frac{g(a^2 b, c_2)}{N(c_2)^{1/2+s}}, \end{aligned}$$

where in the second equality, we used Lemma 2.4 to replace $\chi_{c_2}(ab^2)g(1, c_2)$ by $g(a^2 b, c_2)$ and remove the condition that c_1, c_2 be square-free. Next we remove the condition that $(c_1, c_2) = 1$ to get

$$\begin{aligned} &\sum_{d \equiv 1 \pmod{3}} \mu_K(d) \sum_{\substack{c_1 \equiv 1 \pmod{3} \\ (c_1, ab)=1 \\ d|c_1}} (\chi_{ab^2} \psi)(c_1^2) \frac{\overline{g(1, c_1)}}{N(c_1)^{1/2+s}} \sum_{\substack{c_2 \equiv 1 \pmod{3} \\ (c_2, ab)=1 \\ d|c_2}} \psi(c_2) \frac{g(a^2 b, c_2)}{N(c_2)^{1/2+s}} \\ &= \sum_{d \equiv 1 \pmod{3}} \mu_K(d) \frac{\psi(d)}{N(d)^{1/2+s}} \sum_{\substack{c_1 \equiv 1 \pmod{3} \\ (c_1, ab)=1 \\ d|c_1}} (\chi_{ab^2} \psi)(c_1^2) \frac{\overline{g(1, c_1)}}{N(c_1)^{1/2+s}} \sum_{\substack{c_2 \equiv 1 \pmod{3} \\ (dc_2, ab)=1}} \psi(c_2) \frac{g(a^2 b, dc_2)}{N(c_2)^{1/2+s}} \\ &= \sum_{\substack{d \equiv 1 \pmod{3} \\ (d, ab)=1}} \mu_K(d) \frac{\psi(d^3) \chi_{ab^2}(d^2) \overline{g(1, d)} g(a^2 b, d)}{N(d)^{1+2s}} \\ &\quad \cdot \sum_{\substack{c_1 \equiv 1 \pmod{3} \\ (c_1, abd)=1}} (\chi_{ab^2} \psi)(c_1^2) \frac{\overline{g(d, c_1)}}{N(c_1)^{1/2+s}} \sum_{\substack{c_2 \equiv 1 \pmod{3} \\ (c_2, abd)=1}} \psi(c_2) \frac{g(a^2 bd, c_2)}{N(c_2)^{1/2+s}} \\ &= \sum_{\substack{d \equiv 1 \pmod{3} \\ (d, ab)=1}} \mu_K(d) \frac{\psi(d^3)}{N(d)^{2s}} \sum_{\substack{c_1 \equiv 1 \pmod{3} \\ (c_1, abd)=1}} \psi(c_1^2) \frac{\overline{g(a^2 bd, c_1)}}{N(c_1)^{1/2+s}} \sum_{\substack{c_2 \equiv 1 \pmod{3} \\ (c_2, abd)=1}} \psi(c_2) \frac{g(a^2 bd, c_2)}{N(c_2)^{1/2+s}}, \end{aligned}$$

where in the last equality we used Lemma 2.3 to get $\chi_{ab^2}(c_1^2) \overline{g(d, c_1)} = \overline{g(a^2 bd, c_1)}$, and $\chi_{ab^2}(d^2) \overline{g(1, d)} = \overline{g(a^2 b, d)}$, and that $|g(a^2 b, d)|^2 = N(d)$ as $(d, ab) = 1$. Now, using definition (2.5) we can write the last expression as

$$H_\psi(s; a, b) = \sum_{\substack{d \equiv 1 \pmod{3} \\ (d, ab)=1}} f(d, s) \overline{h(a^2 bd, \overline{\psi^2}, 1/2 + \overline{s})} h(a^2 bd, \psi, 1/2 + s),$$

where

$$f(d, s) = \frac{\mu_K(d) \psi(d^3)}{N(d)^{2s}}.$$

Hence, we have shown so far that

$$\sum_{c \in \mathcal{F} \cup \{1\}} \frac{\chi_c(ab^2)W(\chi_c)}{n_c^{s+1/2}} = \frac{1}{h_{(9)}^\psi} \sum_{\psi \bmod 9} H_\psi(s; a, b).$$

It follows from Lemma 2.5 that for $\sigma \geq 2/3 + \varepsilon$, the estimate

$$(67) \quad \begin{aligned} H_\psi(s; a, b) &\ll (1+t^2)^{2(1-\sigma+\varepsilon)} \mathbf{N}(a^2b)^{1-\sigma+\varepsilon} \sum_{d \equiv 1 \pmod 3} \mathbf{N}(d)^{-2\sigma} \mathbf{N}(d)^{1-\sigma+\varepsilon} \\ &\ll (1+t^2)^{2(1-\sigma+\varepsilon)} \mathbf{N}(a^2b)^{1-\sigma+\varepsilon} \end{aligned}$$

holds uniformly for any ψ , provided $|s - 5/6| > 1/12$. Now,

$$\frac{1}{2\pi i} \int_{\varrho-iT}^{\varrho+iT} y^s \left(\sum_{c \in \mathcal{F} \cup \{1\}} \frac{\chi_c(ab^2)W(\chi_c)}{n_c^{s+1/2}} \right) \frac{ds}{s} = \frac{1}{h_{(9)}^\psi} \sum_{\psi \bmod 9} \frac{1}{2\pi i} \int_{\varrho-iT}^{\varrho+iT} y^s H_\psi(s; a, b) \frac{ds}{s},$$

and moving the contour to $\operatorname{Re} s = 2/3 + \varepsilon$ the last integral equals

$$\begin{aligned} \frac{1}{2\pi i} \int_{\varrho-iT}^{\varrho+iT} y^s H_\psi(s; a, b) \frac{ds}{s} &= -\frac{1}{2\pi i} \int_{2/3+\varepsilon-iT}^{2/3+\varepsilon-iT} y^s H_\psi(s; a, b) \frac{ds}{s} \\ &\quad - \frac{1}{2\pi i} \left(\int_{\varrho+iT}^{2/3+\varepsilon+iT} + \int_{2/3+\varepsilon-iT}^{\varrho-iT} \right) y^s H_\psi(s; a, b) \frac{ds}{s} \\ &\quad + \operatorname{Res}_{s=5/6} (y^s s^{-1} H_\psi(s; a, b)). \end{aligned}$$

Using (67) with $\varepsilon < 1/12$ we see that for $T \geq 1$, the horizontal integrals are

$$(68) \quad \begin{aligned} T^{-1} \int_{2/3+\varepsilon}^{\varrho} y^\sigma (T^4 \mathbf{N}(a^2b))^{1-\sigma+\varepsilon} d\sigma \\ \ll T^{-1} (T^4 \mathbf{N}(a^2b))^{1+\varepsilon} \left(\left(\frac{y}{T^4 \mathbf{N}(a^2b)} \right)^{2/3+\varepsilon} + \left(\frac{y}{T^4 \mathbf{N}(a^2b)} \right)^\varrho \right), \end{aligned}$$

and the vertical integral is

$$(69) \quad \ll y^{2/3+\varepsilon} \mathbf{N}(a^2b)^{1/3} T^{4/3}.$$

Combining the previous estimates with (68) and (69) we have

$$(70) \quad \begin{aligned} \sum_{c \in \mathcal{F}(y) \cup \{1\}} \chi_c(ab^2) \frac{W(\chi_c)}{\sqrt{n_c}} &= \operatorname{Res}_{s=5/6} (y^s s^{-1} H_\psi(s; a, b)) \\ &\quad + O\left(y^\varepsilon + y^{1+2\varepsilon} T^{-1} + y T^{-1+4\varepsilon} (\mathbf{N}(a^2b))^\varepsilon + y^{2/3+\varepsilon} \mathbf{N}(a^2b)^{1/3} T^{4/3}\right). \end{aligned}$$

Using Lemma 2.1 for $T \in [1, y]$, the O -term is

$$\ll (y \mathbf{N}(a^2b))^\varepsilon + y^{2/3+\varepsilon} \mathbf{N}(a^2b)^{1/3} + y^{6/7+\varepsilon} \mathbf{N}(a^2b)^{1/7+\varepsilon}.$$

According to Lemma 2.5, the possible residue, which may exist when $r_2 = 1$ (when $a^2bd = r_1 r_2^2 r_3^3$) and ψ^3 or ψ^6 is not principal, is given by

$$\lim_{s \rightarrow 5/6} (y^s s^{-1} (s - 5/6)^2 H_\psi(s; a, b))',$$

whenever $H_\psi(s; a, b)$ has a double pole at $s = 5/6$, and equals

$$\frac{6}{5} y^{5/6} \lim_{s \rightarrow 5/6} (s - 5/6) H_\psi(s; a, b)$$

when $H_\psi(s; a, b)$ has a single pole.

When $h(a^2bd, \lambda, s + 1/2)$, where λ is either ψ or $\overline{\psi^2}$, has no pole at $s = 5/6$, we have by Lemma 2.5 with $r = 1/(12) + \varepsilon$ and $\varepsilon \leq 1/(24)$ that

$$(71) \quad h(a^2bd, \lambda, 5/6 + 1/2) = \frac{1}{2\pi i} \int_{|z-5/6|=r} \frac{h(a^2bd, \lambda, z + 1/2)}{z - 5/6} dz \ll N(a^2bd)^{\frac{1}{8} + \varepsilon},$$

and when it does have a simple pole then we have the same bound for

$$\lim_{s \rightarrow 5/6} ((s - 5/6)h(a^2bd, \lambda, s + 1/2))',$$

which will be used later below. Hence, when $H_\psi(s; a, b)$ has a simple pole at $s = 5/6$, it follows from Lemma 2.5 that

$$(72) \quad \begin{aligned} \operatorname{Res}_{s=5/6} (y^s s^{-1} H_\psi(s; a, b)) &\ll y^{5/6} N(a^2b)^{\frac{1}{8} + \varepsilon} \sum_{\substack{d \equiv 1 \pmod{3} \\ (d, ab) = 1}} N(d)^{1/8 - 1/6 - 5/3 + \varepsilon} \\ &\ll y^{5/6} N(a^2b)^{\frac{1}{8} + \varepsilon}. \end{aligned}$$

In the latter case, where $H_\psi(s; a, b)$ has a double pole at $s = 5/6$, the residue is given by

$$(73) \quad \begin{aligned} &\lim_{s \rightarrow 5/6} (s^{-1} y^s (s - 5/6)^2 H_\psi(s; a, b))' \\ &= \left(-\frac{1}{(5/6)^2} y^{5/6} + \frac{6}{5} y^{5/6} \log y \right) \lim_{s \rightarrow 5/6} ((s - 5/6)^2 H_\psi(s; a, b)) \\ &\quad + \frac{6}{5} y^{5/6} \lim_{s \rightarrow 5/6} ((s - 5/6)^2 H_\psi(s; a, b))' \end{aligned}$$

By Lemma 2.5,

$$(74) \quad \begin{aligned} &\lim_{s \rightarrow 5/6} (s - 5/6)^2 H_\psi(s; a, b) \\ &= \sum_{\substack{d \equiv 1 \pmod{3} \\ (d, ab) = 1}} f(d, 5/6) \operatorname{Res}_{s=5/6} \overline{h(a^2bd, \overline{\psi^2}, 1/2 + s)} \operatorname{Res}_{s=5/6} h(a^2bd, \psi, 1/2 + s) \\ &\ll N(ab)^\varepsilon \sum_{\substack{d \equiv 1 \pmod{3} \\ (d, ab) = 1}} N(d)^{-5/3 - 1/3 + \varepsilon} \ll N(ab)^\varepsilon, \end{aligned}$$

and

$$\begin{aligned} &\lim_{s \rightarrow 5/6} ((s - 5/6)^2 H_\psi(s; a, b))' \\ &= \lim_{s \rightarrow 5/6} \sum_{\substack{d \equiv 1 \pmod{3} \\ (d, ab) = 1}} (f(d, s) (s - 5/6) \overline{h(a^2bd, \overline{\psi^2}, 1/2 + s)} (s - 5/6) h(a^2bd, \psi, 1/2 + s))' \\ &= \sum_{\substack{d \equiv 1 \pmod{3} \\ (d, ab) = 1}} f'(d, 5/6) \operatorname{Res}_{s=5/6} \overline{h(a^2bd, \overline{\psi^2}, 1/2 + s)} \operatorname{Res}_{s=5/6} h(a^2bd, \psi, 1/2 + s) \\ &\quad + \sum_{\substack{d \equiv 1 \pmod{3} \\ (d, ab) = 1}} f(d, 5/6) \lim_{s \rightarrow 5/6} ((s - 5/6) \overline{h(a^2bd, \overline{\psi^2}, 1/2 + s)})' \operatorname{Res}_{s=5/6} h(a^2bd, \psi, 1/2 + s) \\ &\quad + \sum_{\substack{d \equiv 1 \pmod{3} \\ (d, ab) = 1}} f(d, 5/6) \lim_{s \rightarrow 5/6} ((s - 5/6) h(a^2bd, \psi, 1/2 + s))' \operatorname{Res}_{s=5/6} \overline{h(a^2bd, \overline{\psi^2}, 1/2 + s)}. \end{aligned}$$

Using (71) and Lemma 2.5 we obtain

$$(75) \quad \begin{aligned} \lim_{s \rightarrow 5/6} ((s - 5/6)^2 H_\psi(s; a, b))' &\ll N(ab)^\varepsilon \sum_{\substack{d \equiv 1 \pmod{3} \\ (d, ab) = 1}} N(d)^{-5/3 - 1/3 + \varepsilon} \\ &+ N(a^2 b)^{\frac{1}{8} + \varepsilon} \sum_{\substack{d \equiv 1 \pmod{3} \\ (d, ab) = 1}} N(d)^{-5/3 - 1/6 + 1/8 + \varepsilon} \\ &\ll N(a^2 b)^{\frac{1}{8} + \varepsilon}. \end{aligned}$$

Combining (70), (72), (73), (74) and (75) we obtain the claimed estimate in (66).

Using partial integration (noting that $n_c > 3$) together with the bound in (56) we deduce that

$$\begin{aligned} \sum_{c \in \mathcal{F}(X)} \chi_c(ab^2) \frac{W(\chi_c)}{\sqrt{n_c}} V\left(\frac{3^{r_2} Y N(b)}{3n_c}\right) &= \int_3^X V\left(\frac{3^{r_2} Y N(b)}{3y}\right) d \sum_{c \in \mathcal{F}(y)} \chi_c(ab^2) \frac{W(\chi_c)}{\sqrt{n_c}} \\ &= V\left(\frac{3^{r_2} Y N(b)}{3X}\right) \sum_{c \in \mathcal{F}(X)} \chi_c(ab^2) \frac{W(\chi_c)}{\sqrt{n_c}} + \int_3^X \left(\sum_{c \in \mathcal{F}(y)} \chi_c(ab^2) \frac{W(\chi_c)}{\sqrt{n_c}} \right) V'(A/y) A y^{-2} dy \\ &\ll \frac{X^{3/2+2\varepsilon} N(a^2 b)^{1/3}}{(3^{r_2} Y N(b))^{5/6+\varepsilon}} + \frac{X^{3/2+3\varepsilon} N(a^2 b)^{1/7+\varepsilon}}{(3^{r_2} Y N(b))^{9/(14)+2\varepsilon}}. \end{aligned}$$

Using this bound in the definition of S_3 , the desired estimate in (65) follows. \square

Proof of Theorem 1.3. Put

$$\mathcal{E}(X, a) = \sum_{c \in \mathcal{F}(X)} \chi_c(a) L(1/2, \chi_c) - H(0),$$

where (see the definition of $H(u)$ in (61))

$$\begin{aligned} H(0) &= \frac{\sqrt{3}}{\sqrt{3}-1} \sum_{b \equiv 1 \pmod{3}} \frac{1}{N(a_1^2 a_2 b^3)^{1/2}} \left(\frac{4}{81} F_{\psi_0}(1; a_1 a_2 a_3 b) \right. \\ &\quad \cdot \left(\frac{\pi^2 X \log X}{27} + \left\{ \frac{\pi^2 \log(3/e)}{27} + \lim_{s \rightarrow 1} ((s-1)^2 \zeta_K^2(s))' \right\} X \right) \\ &\quad \left. + \frac{4\pi^2 X}{2187} F'_{\psi_0}(1; a_1 a_2 a_3 b) \right). \end{aligned}$$

Combining (64) with (65), we see that

$$\begin{aligned} \mathcal{E}(X, a) &\ll (XY)^{\frac{4s+1}{4s+2} + \varepsilon} N(a)^{\frac{s}{2s+1} + \varepsilon} + X^{1/2+\varepsilon} Y^{1/2+s+\varepsilon} N(a)^{s+\varepsilon} \\ &\quad + \frac{X^{1+\varepsilon} N(a_3)^\varepsilon}{Y^{1/6-\varepsilon} N(a_1^2 a_2)^{1/3+\varepsilon/2}} + N(a)^{2/3} \frac{X^{3/2+2\varepsilon}}{Y^{5/6+\varepsilon}} + N(a)^{2/7+2\varepsilon} \frac{X^{3/2+3\varepsilon}}{Y^{9/14+2\varepsilon}}. \end{aligned}$$

By Lemma 2.1 with $Y \in [1, X]$ we conclude that

$$\begin{aligned} \mathcal{E}(X, a) &\ll X^{\frac{4\varsigma+1}{4\varsigma+2}+\varepsilon} \mathbf{N}(a)^{\frac{\varsigma}{2\varsigma+1}+\varepsilon} + X^{1/2+\varepsilon} (\mathbf{N}(a))^{\varsigma+\varepsilon} \\ &+ \frac{X^{5/6+\varepsilon} (\mathbf{N}(a_3))^\varepsilon}{\mathbf{N}(a_1^2 a_2)^{1/3+\varepsilon/2}} + \mathbf{N}(a)^{2/3} X^{2/3+\varepsilon} + \mathbf{N}(a)^{2/7+2\varepsilon} X^{6/7+\varepsilon} \\ &+ X^{\frac{60\varsigma+15}{46\varsigma+16}+\varepsilon} \mathbf{N}(a)^{\frac{17\varsigma+2}{46\varsigma+16}+2\varepsilon} + X^{\frac{21\varsigma+15}{14\varsigma+16}+\varepsilon} \mathbf{N}(a)^{\frac{13\varsigma+2}{14\varsigma+16}+2\varepsilon} \\ &+ X^{\frac{28\varsigma+7}{22\varsigma+8}+\varepsilon} \mathbf{N}(a)^{\frac{13\varsigma+2}{2(11\varsigma+4)}+\varepsilon} + X^{\frac{9\varsigma+7}{6\varsigma+8}+\varepsilon} \mathbf{N}(a)^{\frac{9\varsigma+2}{6\varsigma+8}+\varepsilon} \\ &+ X^{\frac{12\varsigma+7}{12\varsigma+8}+\varepsilon} \mathbf{N}(a_1)^{-\frac{1}{2}-\varepsilon} \mathbf{N}(a_2)^{-\frac{2+\varepsilon(6\varsigma-1)}{4(3\varsigma+2)}} \mathbf{N}(a_3)^{\frac{3\varsigma+6\varepsilon(\varsigma+1)}{6\varsigma+4}} \\ &+ X^{\frac{28\varsigma+7}{28\varsigma+8}+\varepsilon} \mathbf{N}(a_1)^{-\frac{7\varsigma+2+\varepsilon(2+10\varsigma)}{2(7\varsigma+2)}} \mathbf{N}(a_2)^{-\frac{4\varsigma+2+\varepsilon(4\varsigma-1)}{28\varsigma+8}} \mathbf{N}(a_3)^{\frac{3\varsigma+\varepsilon(18\varsigma+6)}{2(7\varsigma+2)}}. \end{aligned}$$

For each $a = a_1 a_2^2 a_3^3$, let

$$M(X, a) = \frac{4\sqrt{3}\pi^2 X \log X}{2187(\sqrt{3}-1)} \sum_{\substack{b \equiv 1 \pmod{3} \\ ab = \square}} \frac{F_{\psi_0}(1; ab)}{\mathbf{N}(b)^{1/2}}$$

be the main term of $H(0)$. Then, using (62) with $\varepsilon < 1/2$ we have that

$$H(0) - M(x, a) \ll X \frac{\mathbf{N}(a_1 a_2 a_3)^\varepsilon}{\mathbf{N}(a_1^2 a_2)^{1/2}}.$$

Thus, summing over \mathbf{a} , we see that the contribution of these terms is

$$\begin{aligned} &\ll X \sum_{\substack{a_1, a_2, a_3 \equiv 1 \pmod{3} \\ (a_1, a_2) = 1 \\ a_1, a_2 \text{ square-free}}} \frac{\nu_J(a_1 a_2^2 a_3^3)}{\mathbf{N}(a_1 a_2 a_3)^{3/2-\varepsilon}} = X \prod_{0 \leq r \leq J} \sum_{\substack{a_{1r}, a_{2r}, a_{3r} \in \mathcal{I}_r \\ (a_{1r}, a_{2r}) = 1 \\ a_{1r}, a_{2r} \text{ square-free} \\ 0 \leq \Omega(a_{1r} a_{2r}^2 a_{3r}^3) \leq \ell_r}} \frac{\nu(a_{1r} a_{2r}^2 a_{3r}^3)}{\mathbf{N}(a_{1r} a_{2r} a_{3r})^{3/2-\varepsilon}} \\ &< X \prod_{0 \leq r \leq J} \sum_{\substack{a_{1r} \in \mathcal{I}_r \\ a_{1r} \text{ square-free}}} \frac{\nu(a_{1r})}{\mathbf{N}(a_{1r})^{3/2-\varepsilon}} \sum_{\substack{a_{2r} \in \mathcal{I}_r \\ a_{2r} \text{ square-free}}} \frac{\nu(a_{2r}^2)}{\mathbf{N}(a_{2r})^{3/2-\varepsilon}} \sum_{a_{3r} \in \mathcal{I}_r} \frac{\nu(a_{3r}^3)}{\mathbf{N}(a_{3r})^{3/2-\varepsilon}} \\ &< X \exp\left(\sum_{k_0 < \mathbf{N}\mathfrak{p} \leq X^{\theta_J}} \left(\frac{1}{(\mathbf{N}\mathfrak{p})^{3/2-\varepsilon}} + \frac{1}{2(\mathbf{N}\mathfrak{p})^{3/2-\varepsilon}} + \sum_{n \geq 1} \frac{1}{(3n)!(\mathbf{N}\mathfrak{p})^{(3/2-\varepsilon)n}} \right) \right) \ll X. \end{aligned}$$

Next, summing $\mathcal{E}(X, a)$ over a , we see that

$$\begin{aligned} &\sum_{\mathbf{a}} \frac{\lambda(\mathbf{a}) f(\mathbf{a}, J) \nu_J(\mathbf{a})}{\mathbf{N}(\mathbf{a})^{1/2}} \mathcal{E}(X, a) \\ &\ll \sum_{\mathbf{a}} \frac{\nu_J(\mathbf{a})}{\mathbf{N}(\mathbf{a})^{1/2}} \left(\frac{X^{5/6+\varepsilon} (\mathbf{N}(a_3))^\varepsilon}{\mathbf{N}(a_1^2 a_2)^{1/3+\varepsilon/2}} + X^{\frac{12\varsigma+7}{12\varsigma+8}+\varepsilon} \frac{\mathbf{N}(a_3)^{\frac{3\varsigma+6\varepsilon(\varsigma+1)}{6\varsigma+4}}}{\mathbf{N}(a_1)^{\frac{1}{2}+\varepsilon} \mathbf{N}(a_2)^{\frac{2+\varepsilon(6\varsigma-1)}{4(3\varsigma+2)}}} \right. \\ (76) \quad &\left. + X^{\frac{28\varsigma+7}{28\varsigma+8}+\varepsilon} \frac{\mathbf{N}(a_3)^{\frac{3\varsigma+\varepsilon(18\varsigma+6)}{2(7\varsigma+2)}}}{\mathbf{N}(a_1)^{\frac{7\varsigma+2+\varepsilon(2+10\varsigma)}{2(7\varsigma+2)}} \mathbf{N}(a_2)^{\frac{4\varsigma+2+\varepsilon(4\varsigma-1)}{28\varsigma+8}}} \right) \\ &+ \sum_{i=1}^8 X^{u_i} \prod_{0 \leq r \leq J} E_{\ell_r} \left(\sum_{\mathfrak{p} \in \mathcal{I}_r} (\mathbf{N}\mathfrak{p})^{v_i-1/2} \right) \end{aligned}$$

where (u_i, v_i) run over the set

$$\left\{ \left(\frac{4\varsigma+1}{4\varsigma+2} + \varepsilon, \frac{\varsigma}{2\varsigma+1} + \varepsilon \right), (1/2 + \varepsilon, \varsigma + \varepsilon), (2/3 + \varepsilon, 2/3), (6/7 + \varepsilon, 2/7 + 2\varepsilon) \right. \\ \left. \left(\frac{60\varsigma+15}{46\varsigma+16} + \varepsilon, \frac{17\varsigma+2}{46\varsigma+16} + 2\varepsilon \right), \left(\frac{21\varsigma+15}{14\varsigma+16} + \varepsilon, \frac{13\varsigma+2}{14\varsigma+16} + 2\varepsilon \right) \right. \\ \left. \left(\frac{28\varsigma+7}{22\varsigma+8} + \varepsilon, \frac{13\varsigma+2}{2(11\varsigma+4)} + \varepsilon \right), \left(\frac{9\varsigma+7}{6\varsigma+8} + \varepsilon, \frac{9\varsigma+2}{6\varsigma+8} + \varepsilon \right) \right\}$$

for $1 \leq i \leq 8$. Since

$$\sum_{\mathfrak{p} \in \mathcal{I}_r} (\mathbb{N}\mathfrak{p})^{\gamma-1/2} < 2X^{\theta_r(\gamma+1/2)} \quad \text{and} \quad 2X^{\theta_r(\gamma+1/2)} > \ell_r$$

for sufficiently large X depending on the choice of α and β , we see that

$$E_{\ell_r} \left(\sum_{\mathfrak{p} \in \mathcal{I}_r} (\mathbb{N}\mathfrak{p})^{\gamma-1/2} \right) \leq \left(\sum_{\mathfrak{p} \in \mathcal{I}_r} (\mathbb{N}\mathfrak{p})^{\gamma-1/2} \right)^{\ell_r} < (2X^{\theta_r(\gamma+1/2)})^{\ell_r}.$$

Therefore, the last eight terms of (76) are bounded by

$$2^{\sum_r \ell_r} \sum_{i=1}^8 X^{u_i+(1/2+v_i)\eta},$$

where

$$\eta = \sum_{0 \leq r \leq J} \theta_r \ell_r \leq \frac{2e^{1-\beta}}{e^{1-\beta} - 1} \Theta^{1-\beta}.$$

Since $J \leq \alpha \log_3 X$,

$$2^{\sum_r \ell_r} \leq X^{(\alpha \log_3 X + 1)(\log_2 X)^{\alpha\beta} \log 4 / \log X} \leq X^\varepsilon,$$

and we see that these error terms are $O(X)$ provided that

$$(77) \quad \Theta \leq \left(\frac{(e^{1-\beta} - 1)(1 - u_i - \varepsilon)}{2e^{1-\beta}(1/2 + v_i)} \right)^{\frac{1}{1-\beta}} \quad i = 1, \dots, 8,$$

$\varsigma < \frac{1}{14}$ is fixed, which leads to the sub-convexity assumption in (2), and ε is sufficiently small in terms of ς . On the other hand, the first three terms in (76) are already $O(X)$.

Finally the main term is given by

$$(78) \quad \sum_{\mathfrak{a}} \frac{\lambda(\mathfrak{a})f(\mathfrak{a}, J)v_J(\mathfrak{a})}{\mathbb{N}(\mathfrak{a})^{1/2}} M(X, \mathfrak{a}) = C_X X \log X$$

where

$$(79) \quad C_X = \frac{4\sqrt{3}\pi^2}{81h_{(9)}(\sqrt{3}-1)} \sum_{\mathfrak{a}} \frac{\lambda(\mathfrak{a})f(\mathfrak{a}, J)v_J(\mathfrak{a})}{\mathbb{N}(\mathfrak{a})^{1/2}} \sum_{\substack{b \equiv 1 \pmod{3} \\ ab = \square}} \frac{F_{\psi_0}(1; ab)}{\mathbb{N}(b)^{1/2}}.$$

Note that C_X is absolutely bounded independent of X since

$$C_X \ll \sum_{a_1, a_2, a_3} \frac{v_J(a_1 a_2^2 a_3^3)}{\mathbb{N}(a_1 a_2^2 a_3^3)^{1/2}} \sum_{b \equiv 1 \pmod{3}} \frac{1}{\mathbb{N}(a_1^2 a_2 b^3)^{1/2}} \\ < \prod_{\substack{\mathfrak{p} \in \mathcal{I}_r \\ 0 \leq r \leq J}} \left(1 + \frac{1}{(\mathbb{N}\mathfrak{p})^{3/2}} \right) \left(1 + \frac{1}{2(\mathbb{N}\mathfrak{p})^{3/2}} \right) \left(1 + \sum_{n \geq 1} \frac{1}{(3n)!(\mathbb{N}\mathfrak{p})^{3n/2}} \right) \ll 1.$$

Combining (78) with the estimates above and inserting these in (57) proves the claimed result in Theorem 1.3. \square

5.5. Lower Bound for the main term of the first moment. Write

$$F_\psi(s; \mathbf{m}) = g_\psi(s)h_\psi(s; \mathbf{m}),$$

where

$$\begin{aligned} h_\psi(s; \alpha) &= \prod_{\substack{\pi \equiv 1 \pmod{3} \\ \pi | \alpha}} \left(1 - \frac{\psi(\pi)}{N(\pi)^s}\right) \left(1 - \frac{\psi^2(\pi)}{N(\pi)^s}\right) \\ &\quad \cdot \left(1 - \frac{\psi^2(\pi) + \psi^3(\pi) + \psi^4(\pi)}{N(\pi)^{2s}} + \frac{\psi^4(\pi) + \psi^5(\pi)}{N(\pi)^{3s}}\right)^{-1} \\ g_\psi(s) &= \prod_{\pi \equiv 1 \pmod{3}} \left(1 - \frac{\psi^2(\pi) + \psi^3(\pi) + \psi^4(\pi)}{N(\pi)^{2s}} + \frac{\psi^4(\pi) + \psi^5(\pi)}{N(\pi)^{3s}}\right). \end{aligned}$$

For $\psi = \psi_0$, the principal character modulo 9, we see that

$$h_{\psi_0}(s; \alpha) = \prod_{\substack{\pi \equiv 1 \pmod{3} \\ \pi | \alpha}} \left(1 - \frac{2}{N(\pi)^s + 2}\right) = \prod_{\substack{\pi \equiv 1 \pmod{3} \\ \pi | \alpha}} \left(1 + \frac{2}{N(\pi)^s}\right)^{-1},$$

Thus,

$$\sum_{\substack{b \equiv 1 \pmod{3} \\ ab = \square}} \frac{F_{\psi_0}(1; ab)}{N(b)^{1/2}} = g_{\psi_0}(1) \sum_{\substack{b \equiv 1 \pmod{3} \\ ab = \square}} \frac{1}{N(b)^{1/2}} \prod_{\substack{\pi \equiv 1 \pmod{3} \\ \pi | ab}} \frac{N(\pi)}{N(\pi) + 2}.$$

Inserting this in (79) gives

$$\begin{aligned} C_X &= c_0 \sum_{\mathbf{a}} \frac{\lambda(\mathbf{a})f(\mathbf{a}, J)v_J(\mathbf{a})}{N(\mathbf{a})^{1/2}} \sum_{\substack{b \equiv 1 \pmod{3} \\ ab = \square}} \frac{1}{N(b)^{1/2}} \prod_{\substack{\pi \equiv 1 \pmod{3} \\ \pi | ab}} \frac{N(\pi)}{N(\pi) + 2} \\ &= c_0 \sum_{\mathbf{c}} \frac{1}{N(\mathbf{c})^{3/2}} \prod_{\substack{\pi \equiv 1 \pmod{3} \\ \pi | \mathbf{c}}} \frac{N(\pi)}{N(\pi) + 2} \sum_{\substack{\mathbf{a}, b \equiv 1 \pmod{3} \\ ab = \mathbf{c}^3}} \lambda(\mathbf{a})f(\mathbf{a}, J)v_J(\mathbf{a}) \\ &= c_0 \prod_{\substack{\xi \equiv 1 \pmod{3} \\ \xi \text{ prime}}} \left(1 + \frac{N(\xi)}{N(\xi) + 2} \sum_{n \geq 1} \frac{1}{N(\xi)^{3n/2}} \sum_{\substack{\mathbf{a}, b \equiv 1 \pmod{3} \\ ab = \xi^{3n}}} \lambda(\mathbf{a})f(\mathbf{a}, J)v_J(\mathbf{a})\right), \end{aligned}$$

where

$$c_0 = \frac{4\sqrt{3}\pi^2 g_{\psi_0}(1)}{2187(\sqrt{3} - 1)}.$$

For $\xi \notin \mathcal{I}_r$ for any $r = 0, \dots, J$, $ab = \xi^{3n}$ implies that $\mathbf{a} = 1$. Thus, the contribution of these primes is

$$(80) \quad \prod_{\substack{\xi \equiv 1 \pmod{3} \\ N\xi > X^{\theta_J}}} \left(1 + \frac{N(\xi)}{(N(\xi) + 2)(N(\xi)^{3/2} - 1)}\right) > 1.$$

For the rest of the primes ξ , we have

$$(81) \quad \prod_{\substack{0 \leq r \leq J \\ \xi \in \mathcal{I}_r}} \left(1 + \frac{N(\xi)}{N(\xi) + 2} \sum_{n \geq 1} \frac{1}{N(\xi)^{3n/2}} \sum_{\substack{\mathbf{a}, b \equiv 1 \pmod{3} \\ ab = \xi^{3n}}} \lambda(\mathbf{a}) f(\mathbf{a}, J) \nu_J(\mathbf{a}) \right)$$

For $\xi \in \mathcal{I}_r$, we have

$$\sum_{n \geq 1} \frac{1}{N(\xi)^{3n/2}} \sum_{\substack{0 \leq m \leq 3n \\ 0 \leq m \leq \ell_r}} \lambda(\xi^m) f(\xi^m, J) \nu(\xi^m) = \frac{1}{(1 - N(\xi)^{-3/2})} \sum_{0 \leq m \leq \ell_r} (-1)^m b_m,$$

where

$$b_m = \frac{f(\xi, J)^m}{m! N(\xi)^{3 \max(1, \lceil m/3 \rceil)/2}} > 0.$$

Since

$$\frac{b_{m+1}}{b_m} = \frac{f(\xi, J) N(\xi)^{3 \max(1, \lceil m/3 \rceil)/2}}{(m+1) N(\xi)^{3 \max(1, \lceil (m+1)/3 \rceil)/2}} < 1,$$

we see that

$$\sum_{0 \leq m \leq \ell_r} (-1)^m b_m > b_0 - b_1 = \frac{1 - f(\xi, J)}{N(\xi)^{3/2}} > 0.$$

Combining (80) and (81) we conclude that

$$(82) \quad C_X > c_0 = \frac{4\pi^2}{2187(1 - 1/\sqrt{3})} \prod_{\pi \equiv 1 \pmod{3}} \left(1 - \frac{3}{N(\pi)^2} + \frac{2}{N(\pi)^3} \right).$$

Proof of Theorem 1.1. Assuming Lindelöf Hypothesis (taking $\varsigma = 0$), the condition in (77) is satisfied if

$$(83) \quad \Theta^{1-\beta} \leq \frac{1-\beta}{2e} \min \left\{ \frac{2-12\varepsilon}{7}, \frac{2-28\varepsilon}{(11+28\varepsilon)}, \frac{1-32\varepsilon}{(10+32\varepsilon)}, \frac{1-16\varepsilon}{(6+8\varepsilon)}, \frac{1-4\varepsilon}{2+2\varepsilon} \right\} \\ = \frac{(1-32\varepsilon)(1-\beta)}{(10+32\varepsilon)2e}$$

with $\varepsilon < 1/32$. We take

$$k = 2, \quad \kappa = 1, \quad a = 1.0299, \quad \beta = 0.916, \quad \epsilon = 10^{-5}, \quad \alpha = 7,$$

and

$$\Theta = 5.8025935515 \times 10^{-44}.$$

Then, the conditions in (11), (20), (24), (25), (26), (27), (28), (30), (51) and (83) are all satisfied. Furthermore, we have

$$R_1 \approx -4.7107876828 \times 10^{-40}, \quad R_2 \approx 2.8043085602 \times 10^{42},$$

which were defined in (31). Since $e^r \geq r$, we have for $r \geq 0$ that

$$f(r) = e^r (R_1 - R_2 r) + 8r \leq r(R_1 - R_2 r + 8) \leq (R_1 + 8)r.$$

Thus, we see that

$$\sum_{0 \leq r \leq J-1} (r+1) \exp \left(f(r) + O \left(\frac{(\log_2 X)^{2\alpha}}{\log X} \right) \right) \\ < \sum_{r \geq 0} (r+1) e^{r(R_1+8)} = \frac{1}{(1 - e^{R_1+8})^2}$$

for sufficiently large X . Moreover, with our choices,

$$\mathcal{D}_2 \leq 2.6176409874 \times 10^{15}.$$

This leads to the inequality

$$(18) < |\mathcal{F}(X)|(1 + o_X(1))2.6176409874 \times 10^{15} \left(2 \exp(2e^{1/4} + 2e^{1/8}) + \exp(2e/\Theta) \right) \\ = |\mathcal{F}(X)|(1 + o_X(1)) \left(6.5837089699 \times 10^{17} + \exp(\exp(101.248586291)) \right).$$

We conclude that the estimate

$$\sum_{c \in \mathcal{F}(X) \cap \mathcal{T}_0} |L(1/2, \chi_c)|^2 |M(c, 1)|^2 < e^{e^{101.3}} |\mathcal{F}(X)|(1 + o_X(1))$$

holds for sufficiently large X . Furthermore, it follows from Lemma 2.8 with $n = 1$ and $\varsigma = 0$ that

$$|\mathcal{F}(X)| \leq (|C_1(1)| + o_X(1))X \log X, \quad |C_1(1)| \leq \frac{4\pi^2 g_{\psi_0}(1)}{2187} = (1 - 1/\sqrt{3})c_0.$$

Combining this result with the asymptotic formula for the first moment (82) and using Cauchy-Schwarz inequality it follows that

$$\frac{1}{|\mathcal{F}(X)|} \sum_{\substack{c \in \mathcal{F}(X) \\ L(1/2, \chi_c) \neq 0}} 1 > \frac{c_0^2 + O(1/\log X)}{e^{e^{101.3}}(1 + o_X(1))((1 - 1/\sqrt{3})c_0 + o_X(1))^2}.$$

The claim of Theorem 1.1 follows upon letting X tend to infinity. \square

REFERENCES

1. Baier, S.; Young, M. P. *Mean values with cubic characters*. J. Number Theory 130 (2010), no. 4, 879 - 903.
2. Bui, H.; Florea, A. *Zeros of quadratic Dirichlet L-functions in the hyperelliptic ensemble*, Transactions of the AMS (2018), no. 11, 8013 - 8045.
3. Chandee, V. *Explicit upper bounds for L-functions on the critical line*. Proc. Am. Math. Soc. 137(12), 4049-4063 (2009)
4. Chowla, S. *The Riemann hypothesis and Hilbert's tenth problem*, Norske Vid. Selsk. Forh. (Trondheim) 38 (1965) 62-64. MR0186643.
5. Conrey, J. B., Farmer, D. W., Keating, J. P., Rubinstein, M. O., Snaith, N. C. *Integral moments of L-functions*, Proc. London Math. Soc. (3) 91 (2005), no. 1, 33 - 104. MR 2149530
6. David, C.; Florea, A.; Lalin, M. *Mean values of cubic L-functions over function fields*, Algebra Number Theory 16 (2022), no. 5, 1259 - 1326.
7. David, C., Florea, A., Lalin, M. (2021). *Nonvanishing for cubic L-functions*. Forum of Mathematics, Sigma, 9, E69. doi:10.1017/fms.2021.62
8. Davenport H. *Multiplicative Number Theory*. Second edition. Revised by Hugh L. Montgomery. Graduate Texts in Mathematics, 74. Springer-Verlag, New York-Berlin, 1980.
9. David, C., Gülöglu, A. M. *One-Level Density and Non-Vanishing for Cubic L-Functions Over the Eisenstein Field*. Int. Math. Res. Not. IMRN 2022, no. 23, 18833-18873.
10. Ellenberg, J. S.; Li, W.; Shusterman, M. *Nonvanishing of hyperelliptic zeta functions over finite fields*, Algebra Number Theory 14 (2020), no. 7, 1895 - 1909.
11. Graham, S. W.; Kolesnik, G. *van der Corput's method of exponential sums*, London Mathematical Society Lecture Note Series, 126. Cambridge University Press, Cambridge, 1991. vi+120 pp. ISBN: 0-521-33927-8
12. Heath-Brown, D. R.; Patterson, S. J. *The distribution of Kummer sums at prime arguments*, J. Reine Angew. Math., 310 (1979), 111 - 130.
13. Ireland, K.; Rosen, M. *A classical introduction to modern number theory*. Second edition. Graduate Texts in Mathematics, 84. Springer-Verlag, New York, 1990. xiv+389 pp. ISBN: 0-387-97329-X

14. Iwaniec, H.; Kowalski, E. *Analytic number theory*. American Mathematical Society Colloquium Publications, 53. American Mathematical Society, Providence, RI, 2004.
15. Jutila, M. *On the mean value of $L(1/2, \chi)$ for real characters*, Analysis 1 (1981), no. 2, 149 - 161. MR 632705
16. Katz, N.; Sarnak, P. *Random Matrices, Frobenius Eigenvalues and Monodromy*, AMS Colloq. Publ. 45 (1999).
17. Luo, W. *On Hecke L -series associated with cubic characters*, Compos. Math. 140 (2004), no. 5, 1191 - 1196.
18. Li, W. *Vanishing of Hyperelliptic L -functions at the Central Point*, J. Number Theory 191 (2018), 85-103.
19. Neukirch, J. *Algebraic Number Theory*. Translated from the 1992 German original and with a note by Norbert Schappacher. With a foreword by G. Harder. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 322. Springer-Verlag, Berlin, 1999.
20. Özlük, A. E.; Snyder, C. *On the distribution of the nontrivial zeros of quadratic L -functions close to the real axis*, Acta Arith. 91 (1999), no. 3, 209 - 228.
21. Radziwiłł, M., Soundararajan, K. *Moments and distribution of central L -values of quadratic twists of elliptic curves*. Invent. Math. 202(3), 1029 - 1068 (2015).
22. Robbins, H. (1955). *A Remark on Stirling's Formula*. The American Mathematical Monthly, 62(1), 26 - 29. <https://doi.org/10.2307/2308012>
23. J.B. Rosser and L. Schoenfeld. *Approximate formulas for some functions of prime numbers*. Illinois J. Math. 6 (1962), 64 - 94.
Acta Arith. 143 (2010), no. 1, 81–99.
24. Serre, J.P., *Quelques applications du théorème de densité de Chebotarev*, Inst. Hautes Etudes Sci. Publ. Math., no. 54, pp. 323 - 401, 1981. CHECK THIS
25. Soundararajan, K. *Nonvanishing of quadratic Dirichlet L -functions at $s = 1/2$* , Ann. of Math. (2) 152 (2000), no. 2, 447 - 488. MR 1804529
26. Stark, H. *Some effective cases of the brauer-siegel theorem*, Inventiones mathematicae, vol. 23, pp. 135-152, 1974.

DEPARTMENT OF MATHEMATICS, BILKENT UNIVERSITY, ANKARA, TURKEY
Email address: guloglu@fen.bilkent.edu.tr

DEPARTMENT OF MATHEMATICS, BILKENT UNIVERSITY, ANKARA, TURKEY
Email address: hamza@fen.bilkent.edu.tr