

Beyond scalar quasi-arithmetic means: Quasi-arithmetic averages and quasi-arithmetic mixtures in information geometry*

Frank Nielsen

Sony Computer Science Laboratories Inc
Tokyo, Japan

Abstract

We generalize quasi-arithmetic means beyond scalars by considering the gradient map of a Legendre type real-valued function. The gradient map of a Legendre type function is proven strictly comonotone with a global inverse. It thus yields a generalization of strictly monotone and differentiable functions generating scalar quasi-arithmetic means. Furthermore, the Legendre transformation gives rise to pairs of dual quasi-arithmetic averages via the convex duality. We study the invariance and equivariance properties under affine transformations of quasi-arithmetic averages via the lens of dually flat spaces of information geometry. We show how these quasi-arithmetic averages are used to express points on dual geodesics and sided barycenters in the dual affine coordinate systems. We then consider quasi-arithmetic mixtures and describe several parametric and non-parametric statistical models which are closed under the quasi-arithmetic mixture operation.

Keywords: quasi-arithmetic mean ; Legendre-type function ; information geometry ; affine Legendre invariance ; Jensen divergence ; comparative convexity ; Jensen-Shannon divergence

1 Introduction

We first start by generalizing the notion of quasi-arithmetic means [22] (Definition 1) which relies on strictly monotone and differentiable functions to other non-scalar types such as vectors and matrices in Section 2: Namely, we show how the gradient of a strictly convex and differentiable function of Legendre type [48] (Definition 2) is co-monotone (Proposition 1) and admits a global inverse. Legendre functions bring the counterpart notion of quasi-arithmetic mean generators to non-scalar types: Quasi-arithmetic averages (Definition 3). In Section 3, we show how quasi-arithmetic averages occur in the dually flat manifolds of information geometry [3, 5]: Quasi-arithmetic averages are used to parameterize the points on dual geodesics (§3.1) and to characterize the dual barycenters with respect to the canonical divergence [3] (§3.2). We explain the dualities between steep exponential families [9], regular Bregman divergences [8], and quasi-arithmetic averages in Section 3.3. The invariance and equivariance properties of quasi-arithmetic averages are studied in Section 4 under the information-geometric lens: The invariance and equivariance of quasi-arithmetic averages under affine transformations (Proposition 2) generalizes the invariance property of quasi-arithmetic means (Property 1) and bring new insights from information geometry. Finally, in Section 5, we define quasi-arithmetic mixtures (Definition 4), show their role in defining a generalization of Jensen-Shannon divergence [36], and discusses the underlying information geometry of parametric and non-parametric statistical models closed under the operation of quasi-arithmetic mixing. We propose a geometric generalization of the Jensen-Shannon divergence (Definition 20) based on affine connections [5] which recovers the ordinary Jensen-Shannon divergence and the geometric Jensen-Shannon divergence when the affine connections are chosen as the mixture connection and the exponential connection, respectively.

*Check the latest technical report: <http://arxiv.org/abs/2301.10980>

2 Quasi-arithmetic averages and information geometry

2.1 Scalar quasi-arithmetic means

Let $\Delta_{n-1} = \{(w_1, \dots, w_n) : w_i \geq 0, \sum_i w_i = 1\} \subset \mathbb{R}^d$ denotes the closed $(d-1)$ -dimensional standard simplex sitting in \mathbb{R}^d and $\Delta_{n-1}^\circ = \partial\Delta_{n-1}$ the open standard simplex where ∂ denotes the set boundary operator. Weighted quasi-arithmetic means [22] generalize the weighted arithmetic mean $A(x_1, \dots, x_n; w) = \sum_i w_i x_i$ as follows:

Definition 1 (Weighted quasi-arithmetic mean) *Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be a strictly monotone and differentiable real-valued function. The weighted quasi-arithmetic mean (QAM) $M_f(x_1, \dots, x_n; w)$ between n scalars $x_1, \dots, x_n \in I \subset \mathbb{R}$ with respect to a normalized weight vector $w \in \Delta_{n-1}$, is defined by*

$$M_f(x_1, \dots, x_n; w) := f^{-1} \left(\sum_{i=1}^n w_i f(x_i) \right).$$

The notion of quasi-arithmetic means and its properties were studied by Knopp [24], Jessen [23], Kolmogorov [25], Nagumo [30] and De Finetti [18] in the late 1920's-early 1930's. Let us write for short $M_f(x_1, \dots, x_n) := M_f(x_1, \dots, x_n; \frac{1}{n}, \dots, \frac{1}{n})$ the quasi-arithmetic mean, and $M_{f,\alpha}(x, y) := M_f(x, y; \alpha, 1-\alpha)$, the weighted bivariate mean. Mean $M_f(x_1, \dots, x_n)$ is called a quasi-arithmetic mean because we have:

$$f(M_f(x_1, \dots, x_n)) = \frac{1}{n} \sum_i f(x_i) = A(f(x_1), \dots, f(x_n)),$$

the arithmetic mean with respect to the f -representation [53] of scalars. A QAM has also been called a f -mean [1] in the literature to emphasize its underlying generator f . A QAM like any other generic mean [14] satisfies the in-betweenness property:

$$\min\{x_1, \dots, x_n\} \leq M_f(x_1, \dots, x_n; w) \leq \max\{x_1, \dots, x_n\}.$$

See also the recent works on aggregators [16]. QAMs have been used in machine learning (e.g., [26]) and statistics (e.g., [7]).

We have the following invariance property of QAMs:

Property 1 (Invariance of quasi-arithmetic mean [32]) $M_g(x, y) = M_f(x, y)$ if and only if $g(t) = \lambda f(t) + c$ for $\lambda \in \mathbb{R} \setminus \{0\}$ and $c \in \mathbb{R}$.

See [6, 28] for the more general case of invariance of weighted quasi-arithmetic means with weights defined by functions.

Let $\mathcal{CM}(a, b)$ denotes the class of continuous strictly monotone functions on $[a, b]$, and \sim the equivalence relation $f \sim g$ if and only if $M_f = M_g$. Then the quasi-arithmetic mean induced by f is $M_{[f]}$ where $[f]$ denotes the equivalence class of functions in $\mathcal{CM}(a, b)$ which contains f . When $f(t) = t$, we recover the arithmetic mean A : $M_{\text{id}}(x, y) = A(x, y)$ where $\text{id}(x) = x$ is the scalar identity function.

The power means $M_p(x, y) := M_{f_p}(x, y)$, also called Hölder means [46, 54], are obtained for the following continuous family of QAM generators index by $p \in \mathbb{R}$:

$$f_p(t) = \begin{cases} \frac{t^p - 1}{p}, & p \in \mathbb{R} \setminus \{0\}, \\ \log(t), & p = 0. \end{cases}, \quad f_p^{-1}(t) = \begin{cases} (1 + tp)^{\frac{1}{p}}, & p \in \mathbb{R} \setminus \{0\}, \\ \exp(t), & p = 0. \end{cases},$$

Special cases of the power means are the harmonic mean ($H = M_{-1}$), the geometric mean ($G = M_0$), the arithmetic mean ($A = M_1$), and the quadratic mean ($Q = M_2$). A QAM is said positively homogeneous if and only if $M_f(\lambda x, \lambda y) = \lambda M_f(x, y)$ for all $\lambda > 0$. The power means M_p are the only positively homogeneous QAMs [22]. QAMs provide a versatile way to construct means [14] by specifying a functional generator $f \in$

$\mathcal{CM}(I)$. For example, the log-sum-exp mean¹ is obtained for the QAM generator $f_{\text{LSE}}(u) = \exp(u) = f_0^{-1}(u)$ with $f_{\text{LSE}}^{-1}(u) = \log u = f_0(u)$ (these functions are the inverse of the geometric mean functions):

$$\text{LSE}(x, y) = \log \left(\frac{\exp^x + \exp^y}{2} \right) = M_{f_{\text{LSE}}}(x, y).$$

2.2 Quasi-arithmetic averages

To generalize scalar QAMs to other non-scalar types such as vectors or matrices, we face two difficulties:

1. we need to ensure that the generator $G : \mathbb{X} \rightarrow \mathbb{R}$ admits a global inverse G^{-1} , and
2. we would like the smooth function G to bear a generalization of monotonicity of univariate functions.

Indeed, the inverse function theorem [17] in multivariable calculus states only the existence locally of an inverse continuously differentiable function G^{-1} for a multivariate function G provided that the Jacobian matrix of G is not singular (non-zero determinant).

We shall thus consider a well-behaved class \mathcal{F} of non-scalar functions G (i.e., vector or matrix functions) which admits global inverse functions G^{-1} belonging to the same class \mathcal{F} : Namely, we consider the gradient maps of Legendre-type functions where Legendre-type functions are defined as follows:

Definition 2 (Legendre type function [48]) (Θ, F) is of Legendre type if the function $F : \Theta \subset \mathbb{X} \rightarrow \mathbb{R}$ is strictly convex and differentiable with $\Theta \neq \emptyset$ and

$$\lim_{\lambda \rightarrow 0} \frac{d}{d\lambda} F(\lambda\theta + (1-\lambda)\bar{\theta}) = -\infty, \quad \forall \theta \in \Theta, \forall \bar{\theta} \in \partial\Theta. \quad (1)$$

The condition of Eq. 1 is related to the notion of steepness in exponential families [9].

Legendre-type functions $F(\Theta)$ admits a convex conjugate $F^*(\eta)$ of Legendre type via the Legendre transform (Theorem 1 [48]):

$$F^*(\eta) = \langle \nabla F^{-1}(\eta), \eta \rangle - F(\nabla F^{-1}(\eta)),$$

where $\langle \theta, \eta \rangle = \theta^\top \eta$ denotes the inner product in \mathbb{X} (e.g., Euclidean inner product $\langle \theta, \eta \rangle = \theta^\top \eta$ for $\mathbb{X} = \mathbb{R}^d$, the Hilbert-Schmidt inner product $\langle A, B \rangle := \text{tr}(AB^\top)$ where $\text{tr}(\cdot)$ denotes the matrix trace for $\mathbb{X} = \text{Mat}_{d,d}(\mathbb{R})$, etc.), and $\eta \in H$ with H the image of the gradient map $\nabla F : \Theta \rightarrow H$. Moreover, we have $\nabla F^* = \nabla F^{-1}$.

The gradient of a strictly convex function of Legendre type can also be interpreted as a generalization the notion of monotonicity of a univariate function: A function $G : \mathbb{X} \rightarrow \mathbb{R}$ is said strictly increasing co-monotone if

$$\forall \theta_1, \theta_2 \in \mathbb{X}, \theta_1 \neq \theta_2, \quad \langle \theta_1 - \theta_2, G(\theta_1) - G(\theta_2) \rangle > 0.$$

and strictly decreasing co-monotone if $-G$ is strictly increasing co-monotone.

Proposition 1 (Gradient co-monotonicity) *The gradient functions $\nabla F(\theta)$ and $\nabla F^*(\eta)$ of the Legendre-type convex conjugates F and F^* in \mathcal{F} are strictly increasing co-monotone functions.*

Proof: We have to prove that

$$\langle \theta_2 - \theta_1, \nabla F(\theta_2) - \nabla F(\theta_1) \rangle > 0, \quad \forall \theta_1 \neq \theta_2 \in \Theta \quad (2)$$

$$\langle \eta_2 - \eta_1, \nabla F^*(\eta_2) - \nabla F^*(\eta_1) \rangle > 0, \quad \forall \eta_1 \neq \eta_2 \in H \quad (3)$$

The inequalities follow by interpreting the terms of the left-hand-side of Eq. 2 and Eq. 3 as Jeffreys-symmetrization [36] of the dual Bregman divergences [12]:

$$\begin{aligned} B_F(\theta_1 : \theta_2) &= F(\theta_1) - F(\theta_2) - \langle \theta_1 - \theta_2, \nabla F(\theta_2) \rangle \geq 0, \\ B_{F^*}(\eta_1 : \eta_2) &= F^*(\eta_1) - F^*(\eta_2) - \langle \eta_1 - \eta_2, \nabla F(\theta_2) \rangle \geq 0, \end{aligned}$$

¹Also called the exponential mean [14] since it is a f -mean for the exponential function.

where the first equality holds if and only if $\theta_1 = \theta_2$ and the second inequality holds iff $\eta_1 = \eta_2$. Indeed, we have the following Jeffreys-symmetrization of the dual Bregman divergences:

$$\begin{aligned} B_F(\theta_1 : \theta_2) + B_F(\theta_2 : \theta_1) &= \langle \theta_2 - \theta_1, \nabla F(\theta_2) - \nabla F(\theta_1) \rangle > 0, \quad \forall \theta_1 \neq \theta_2 \\ B_{F^*}(\eta_1 : \eta_2) + B_{F^*}(\eta_2 : \eta_1) &= \langle \eta_2 - \eta_1, \nabla F^*(\eta_2) - \nabla F^*(\eta_1) \rangle > 0, \quad \forall \eta_1 \neq \eta_2 \end{aligned}$$

The symmetric divergences $\text{JB}_F(\theta_1, \theta_2) := B_F(\theta_1 : \theta_2) + B_F(\theta_2 : \theta_1)$ and $\text{JB}_{F^*}(\eta_1, \eta_2) := B_{F^*}(\eta_1 : \eta_2) + B_{F^*}(\eta_2 : \eta_1)$ are called Jeffreys-Bregman divergences in [39]. \square

Remark 1 *Co-monotonicity can be interpreted as a multivariate generalization of monotone univariate functions: A smooth univariate strictly increasing monotone function f is such that $f'(x) > 0$. Since $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$, a strictly monotone function is such that $(x+h-x)(f(x+h) - f(x)) > 0$ for small enough $h > 0$.*

Let us now define the quasi-arithmetic averages (QAAs) as follows:

Definition 3 (Quasi-arithmetic averages) *Let $F : \Theta \rightarrow \mathbb{R}$ be a strictly convex and smooth real-valued function of Legendre-type in \mathcal{F} . The weighted quasi-arithmetic average of $\theta_1, \dots, \theta_n$ and $w \in \Delta_{n-1}$ is defined by the gradient map ∇F as follows:*

$$M_{\nabla F}(\theta_1, \dots, \theta_n; w) := \nabla F^* \left(\sum_i w_i \nabla F(\theta_i) \right), \quad (4)$$

where ∇F^* is the gradient map of the Legendre transform F^* of F .

We recover the usual definition of scalar QAMs M_f (Definition 1) when $F(t) = \int_a^t f(u)du$ for a strictly increasing or strictly decreasing and continuous function f : $M_f = M_{F'}$ (with $f^{-1} = (F')^{-1}$). Notice that we only need to consider F to be strictly convex or strictly concave and smooth to define a multivariate QAM since $M_{\nabla F} = M_{-\nabla F}$.

Let us now give some examples of vector and matrix quasi-arithmetic averages:

Example 1 (Separable quasi-arithmetic average) *When the strictly convex d -variate real-valued function $F(\theta)$ is separable, i.e., $F(\theta) = \sum_{i=1}^d f_i(\theta_i)$ with $f_i : I_i \rightarrow \mathbb{R}$ for strictly convex and differen-*

tiable univariate functions $f_i(\theta_i) \in \mathcal{CM}(I_i)$, the global gradient maps are $\nabla F(\theta) = \begin{bmatrix} f'_1(\theta_1) \\ \vdots \\ f'_d(\theta_d) \end{bmatrix}$ and

$\nabla F^(\eta) = \begin{bmatrix} f_1'^{-1}(\eta_1) \\ \vdots \\ f_d'^{-1}(\eta_d) \end{bmatrix} = \nabla F^{-1}(\eta)$ so that we have $M_{\nabla F}(\theta, \theta') = \begin{bmatrix} M_{f'_1}(\theta_1, \theta'_1) \\ \vdots \\ M_{f'_d}(\theta_d, \theta'_d) \end{bmatrix}$, the componentwise*

quasi-arithmetic scalar means.

Example 2 (Non-separable quasi-arithmetic average) *Consider the non-separable d -variate real-valued function $F(\theta) = \log(1 + \exp_{i=1}^d e^{\theta_i}) = \text{LSE}(0, \theta_1, \dots, \theta_2)$. This function called $\text{LSE}_0^+(\theta_1, \dots, \theta_2) = \text{LSE}(0, \theta_1, \dots, \theta_2)$ is strictly convex and differentiable of Legendre type [40], with the reciprocal gradient maps*

$\nabla F(\theta) = \begin{bmatrix} \frac{e^{\theta_1}}{1 + \sum_{j=1}^d e^{\theta_j}} \\ \vdots \\ \frac{e^{\theta_d}}{1 + \sum_{j=1}^d e^{\theta_j}} \end{bmatrix}$ and $\nabla F^(\eta) = \begin{bmatrix} \log \frac{\eta_1}{1 - \sum_{j=1}^d \eta_j} \\ \vdots \\ \log \frac{\eta_d}{1 - \sum_{j=1}^d \eta_j} \end{bmatrix}$. We shall call this quasi-arithmetic average the*

categorical mean as it is induced by the cumulant function of the family of categorical distributions (see §3).

Example 3 (Matrix example) Consider the strictly convex function [52, 11]:

$$\begin{aligned} F & : \text{Sym}_{++}(d) \rightarrow \mathbb{R} \\ \theta & \mapsto -\log \det(\theta), \end{aligned}$$

where $\det(\cdot)$ denotes the matrix determinant. Function $F(\theta)$ is strictly convex and differentiable [11] on the domain of d -dimensional symmetric positive-definite matrices $\text{Sym}_{++}(d)$ (open convex cone). We have $F(\theta) = -\log \det(\theta)$, $\nabla F(\theta) = -\theta^{-1} =: \eta(\theta)$, $\nabla F^{-1}(\eta) = -\eta^{-1} =: \theta(\eta)$, and $F^*(\eta) = \langle \theta(\eta), \eta \rangle - F(\theta(\eta)) = -d - \log \det(-\eta)$, where the dual parameter η belongs to the d -dimensional negative-definite matrix domain, and the inner matrix product is the Hilbert-Schmidt inner product $\langle A, B \rangle := \text{tr}(AB^\top)$, where $\text{tr}(\cdot)$ denotes the matrix trace. It follows that $M_{\nabla F}(\theta_1, \theta_2) = 2(\theta_1^{-1} + \theta_2^{-1})^{-1}$ is the matrix harmonic mean [2] generalizing the scalar harmonic mean $H(a, b) = \frac{2ab}{a+b}$ for $a, b > 0$. Other examples of matrix means are reported in [10].

3 Use of quasi-arithmetic averages in dually flat manifolds

In this section, we shall elicit the roles of quasi-arithmetic averages in information geometry [5], and report the invariance and equivariance properties of quasi-arithmetic averages with respect to affine transformations from the lens of information geometry.

Let (M, g, ∇, ∇^*) be a dually flat space (DFS) where ∇ and ∇^* are the dual torsion-free flat affine connections such that $\frac{\nabla + \nabla^*}{2}$ is the Riemannian metric Levi-Civita connection ∇^g induced by g (we have $\nabla^* = 2\nabla^g - \nabla$ and $\nabla^{**} = \nabla$). Let $F(\theta)$ and $F^*(\eta)$ denotes the Legendre-type potential functions with θ denoting the ∇ -affine coordinate system and η denoting the ∇^* -affine coordinate system. A point P in a DFS can thus be represented either by the coordinates $\theta(P)$ or by the coordinates $\eta(P)$. Let us denote this duality of coordinates by $P \begin{bmatrix} \theta(P) \\ \eta(P) \end{bmatrix}$. In a DFS, the dual canonical divergences [5] $D_{\nabla, \nabla^*}(P : Q)$ and $D_{\nabla^*, \nabla}^*(P : Q) = D_{\nabla^*, \nabla}(P : Q)$ between two points P and Q of M can be expressed using the coordinate systems as dual Bregman divergences. We have the following identities:

$$D_{\nabla, \nabla^*}(P : Q) = B_F(\theta(P) : \theta(Q)) = B_{F^*}(\eta(Q) : \eta(P)) = D_{\nabla^*, \nabla}(Q : P).$$

3.1 Quasi-arithmetic averages in dual parameterizations of dual geodesics




Figure 1: The points on dual geodesics in a dually flat spaces have dual coordinates expressed with quasi-arithmetic averages.

In a DFS $(M, g, \nabla, \nabla^*) = \text{DFS}(F, \theta \in \Theta; F^*, \eta \in H)$, the primal geodesics $\gamma_{\nabla}(P, Q; t)$ are obtained as line segments in the θ -coordinate system (because the Christoffel symbols of the connection ∇ vanishes in the θ -coordinate system) while the dual geodesics $\gamma_{\nabla^*}(P, Q; t)$ are line segments in the η -coordinate system (because the Christoffel symbols of the dual connection ∇^* vanishes in the η -coordinate system). The dual geodesics define interpolation schemes $(PQ)^{\nabla}(t) = \gamma_{\nabla}(P, Q; t)$ and $(PQ)^{\nabla^*}(t) = \gamma_{\nabla^*}(P, Q; t)$ between input

points P and Q with $P = \gamma_{\nabla}(P, Q; 0) = \gamma_{\nabla^*}(P, Q; 0)$ and $Q = \gamma_{\nabla}(P, Q; 1) = \gamma_{\nabla^*}(P, Q; 1)$ when t ranges in $[0, 1]$. We express the coordinates of the interpolated points on γ_{∇} and γ_{∇^*} using quasi-arithmetic averages as follows (Figure 1):

$$(PQ)^{\nabla}(t) = \gamma_{\nabla}(P, Q; t) = \begin{bmatrix} M_{\text{id}}(\theta(P), \theta(Q); 1-t, t) \\ M_{\nabla F^*}(\eta(P), \eta(Q); 1-t, t) \end{bmatrix}, \quad (5)$$

$$(PQ)^{\nabla^*}(t) = \gamma_{\nabla^*}(P, Q; t) = \begin{bmatrix} M_{\nabla F}(\theta(P), \theta(Q); 1-t, t) \\ M_{\text{id}}(\eta(P), \eta(Q); 1-t, t) \end{bmatrix}. \quad (6)$$

Quasi-arithmetic averages were used by a geodesic bisection algorithm to approximate the circumcenter of the minimum enclosing balls with respect to the canonical divergence in a DFS in [45].

3.2 Quasi-arithmetic average coordinates of dual barycenters with respect to the canonical divergence

Consider a finite set of n points P_1, \dots, P_n on the DFS (M, g, ∇, ∇^*) . The points P_i $\begin{bmatrix} \theta_i \\ \eta_i \end{bmatrix}$ can be expressed in the dual coordinate systems either as $\theta(P_i) = \theta_i$ or $\eta(P_i) = \eta_i$. The right centroid point $\bar{C}_R \in M$ defined by $\bar{C}_R = \arg \min_{P \in M} D_{\nabla, \nabla^*}(P_i : P)$ (or equivalently as a right-sided Bregman centroid [8] $\bar{\theta}_R = \arg \min_{\theta} B_F(\theta_i : \theta)$) has dual coordinates

$$\bar{\theta}_R = \theta(\bar{C}_R) = \frac{1}{n} \sum_{i=1}^n \theta_i = M_{\text{id}}(\theta_1, \dots, \theta_n), \quad (7)$$

$$\bar{\eta}_R = \nabla F(\bar{\theta}_R) = M_{\nabla F^*}(\eta_1, \dots, \eta_n). \quad (8)$$

Similarly, the left centroid point $\bar{C}_L \in M$ defined by $\bar{C}_L = \arg \min_{P \in M} D_{\nabla, \nabla^*}(P : P_i)$ (or equivalently as a left-sided Bregman centroid [41] $\bar{\theta}_L = \arg \min_{\theta} B_F(\theta : \theta_i)$) has coordinates

$$\bar{\theta}_L = M_{\nabla F}(\theta_1, \dots, \theta_n), \quad (9)$$

$$\bar{\eta}_L = \nabla F(\bar{\theta}_L) = M_{\text{id}}(\eta_1, \dots, \eta_n). \quad (10)$$

Thus we have the two dual sided centroids \bar{C}_R and \bar{C}_L (reference duality [53]) on the dually flat manifold M expressed using the dual coordinates as

$$\bar{C}_R \left[\begin{array}{l} \bar{\theta}_L = M_{\nabla F}(\theta_1, \dots, \theta_n) \\ \bar{\eta}_L = \nabla F(\bar{\theta}_L) = M_{\text{id}}(\eta_1, \dots, \eta_n) \end{array} \right], \quad \bar{C}_L \left[\begin{array}{l} \bar{\theta}_L = M_{\nabla F}(\theta_1, \dots, \theta_n) \\ \bar{\eta}_L = \nabla F(\bar{\theta}_L) = M_{\text{id}}(\eta_1, \dots, \eta_n) \end{array} \right]$$

Let $\underline{\theta} = \frac{1}{n} \sum_{i=1}^n \theta_i$ and $\underline{\eta} = \frac{1}{n} \sum_{i=1}^n \eta_i$. Then we have $\underline{\theta} = \nabla F^*(M_{\nabla F^*}(\eta_1, \dots, \eta_n))$ and $\underline{\eta} = \nabla F(M_{\nabla F}(\theta_1, \dots, \theta_n))$. The dual DFS centroids [47] were studied as Bregman sided centroids and expressed as quasi-arithmetic averages in [41].

Figure 2 illustrates the dual centroids expressed using the quasi-arithmetic averages.

We may consider the barycenter of n weighted points P_1, \dots, P_n (weight vector $w \in \Delta_{n-1}$) with respect to a Jensen divergence [39] J_F defined as the minimization of $\sum_{i=1}^n w_i J_F(\theta, \theta_i)$. In [39], the following iterative algorithm was proposed: Let $\theta^{(0)} = \sum_{i=1}^n w_i \theta_i$, and iteratively update $\theta^{(t+1)} = M_{\nabla F} \left(\frac{\theta^{(t)} + \theta_1}{2}, \dots, \frac{\theta^{(t)} + \theta_n}{2}; w \right)$.

3.3 Tripartite duality of densities/divergences/means and dual quasi-arithmetic averages

Banerjee et al. [8] proved a bijection between natural regular exponential families $\mathcal{E}_F = \{p_{\theta}(x) = \exp(x \cdot \theta - F(\theta))\}$ with cumulant functions F and ‘‘regular’’ dual Bregman divergences B_{F^*} by rewriting the densities as $p_{\theta}(x) = p_{\eta}(x) = \exp(-B_{F^*}(x : \eta) + F^*(x))$ with $\eta = \nabla F(\theta)$ (using the Young equality $F(\theta) + F^*(\eta) - \langle \theta, \eta \rangle =$




Figure 2: Dual centroids in a dually flat spaces have dual coordinates expressed with quasi-arithmetic averages.




Figure 3: Overview of the bijections between regular exponential families, Bregman divergences of Legendre-type, and quasi-arithmetic averages.

0). Furthermore, a bijection between Bregman divergences B_F and quasi-arithmetic averages $M_{\nabla F}$ was informally mentioned in [45]. Using these bijections, we can cast the maximum likelihood estimator (MLE) of an exponential family \mathcal{E}_F as a dual right-sided Bregman centroid problem [34]. Figure 3 summarizes the dualities between convex conjugate functions, exponential families and Bregman divergences, and maximum likelihood estimator, Bregman centroid expressed as multivariate QAMs.

The categorical mean of Example 2 is induced by the gradient map of the cumulant function of the exponential family of categorical distributions [5] (the family of discrete distributions on a finite alphabet).

A Legendre-type function F induces two dual quasi-arithmetic weighted averages $M_{\nabla F}$ and $M_{\nabla F^*}$ by the gradient maps of the convex conjugates F and F^* .

When $\nabla F = \nabla F^* = \nabla F^{-1}$ (meaning that the convex conjugate gradients are reciprocal to each others), we have $M_{\nabla F} = M_{\nabla F^*}$. This holds for example for the scalar and matrix harmonic means which are self-dual means with $\nabla F(x) = x^{-1} = \nabla F^*(x)$.

Consider the Mahalanobis divergence Δ^2 (i.e., the squared Mahalanobis distance Δ) as a Bregman divergence obtained for the quadratic form generator $F_Q(\theta) = \frac{1}{2} \theta^\top Q \theta + c\theta + \kappa$ for a symmetric positive-

definite $d \times d$ matrix Q , $c \in \mathbb{R}^d$ and $\kappa \in \mathbb{R}$. We have:

$$\Delta^2(\theta_1, \theta_2) = B_{F_Q}(\theta_1 : \theta_2) = \frac{1}{2}(\theta_2 - \theta_1)^\top Q(\theta_2 - \theta_1).$$

When $Q = I$, the identity matrix, the Mahalanobis divergence² coincides with the Euclidean divergence² (i.e., the squared Euclidean distance). The Legendre convex conjugate is $F^*(\eta) = \frac{1}{2}\eta^\top Q^{-1}\eta = F_{Q^{-1}}(\eta)$, and we have $\eta = \nabla F_Q(\theta) = Q\theta$ and $\theta = \nabla F_Q^*(\eta) = Q^{-1}\eta$. Thus we get the following dual quasi-arithmetic averages:

$$M_{\nabla F_Q}(\theta_1, \dots, \theta_n; w) = Q^{-1} \left(\sum_{i=1}^n w_i Q \theta_i \right) = \sum_{i=1}^n w_i \theta_i = M_{\text{id}}(\theta_1, \dots, \theta_n; w), \quad (11)$$

$$M_{\nabla F_Q^*}(\eta_1, \dots, \eta_n; w) = Q \left(\sum_{i=1}^n w_i Q^{-1} \eta_i \right) = M_{\text{id}}(\eta_1, \dots, \eta_n; w). \quad (12)$$

The dual quasi-arithmetic average functions $M_{\nabla F_Q}$ and $M_{\nabla F_Q^*}$ induced by a Mahalanobis Bregman generator F_Q coincide since $M_{\nabla F_Q} = M_{\nabla F_Q^*} = M_{\text{id}}$. This means geometrically that the left-sided and right-sided centroids of the underlying canonical divergences match. The average $M_{\nabla F_Q}(\theta_1, \dots, \theta_n; w)$ expresses the centroid $C = \bar{C}_R = \bar{C}_L$ in the θ -coordinate system ($\theta(C) = \underline{\theta}$) and the average $M_{\nabla F_Q^*}(\eta_1, \dots, \eta_n; w)$ expresses the same centroid in the η -coordinate system ($\eta(C) = \underline{\eta}$). In that case of self-dual flat Euclidean geometry, there is an affine transformation relating the θ - and η -coordinate systems: $\eta = Q\theta$ and $\theta = Q^{-1}\eta$. As we shall see this is because the underlying geometry is self-dual Euclidean flat space ($(M, g_{\text{Euclidean}}, \nabla_{\text{Euclidean}}, \nabla_{\text{Euclidean}}^* = \nabla_{\text{Euclidean}})$) and that both dual connections coincide with the Euclidean connection (i.e., the Levi-Civita connection of the Euclidean metric). In this particular case, the dual coordinate systems are just related by affine transformations of one to another.

4 Invariance and equivariance properties of quasi-arithmetic averages

Recall that a dually flat manifold [5] (M, g, ∇, ∇^*) has a canonical divergence [3] D_{∇, ∇^*} which can be expressed either as a primal Bregman divergence in the ∇ -affine coordinate system θ (using the convex potential function $F(\theta)$) or as a dual Bregman divergence in the ∇^* -affine coordinate system η (using the convex conjugate potential function $F^*(\eta)$), or as dual Fenchel-Young divergences [38] using the mixed coordinate systems θ and η . The dually flat manifold (M, g, ∇, ∇^*) (a particular case of Hessian manifolds [49] which admit a global coordinate system) is characterized by $(\theta, F(\theta); \eta, F^*(\eta))$ which we shall denote by $(M, g, \nabla, \nabla^*) \leftarrow \text{DFS}(\theta, F(\theta); \eta, F^*(\eta))$ (or in short $(M, g, \nabla, \nabla^*) \leftarrow (\Theta, F(\theta))$). However, the choices of parameters θ and η and potential functions F and F^* are not unique since they can be chosen up to affine reparameterizations and additive affine terms [5]: $(M, g, \nabla, \nabla^*) \leftarrow \text{DFS}([\theta, F(\theta); \eta, F^*(\eta)])$ where $[\cdot]$ denotes the equivalence class that has been called purposely the affine Legendre invariance in [31]:

- First, consider changing the potential function $F(\theta)$ by adding an affine term: $\bar{F}(\theta) = F(\theta) + \langle c, \theta \rangle + d$. We have $\nabla \bar{F}(\theta) = \nabla F(\theta) + c = \bar{\eta}$. Inverting $\nabla \bar{F}(x) = \nabla F(x) + c = y$, we get $\nabla \bar{F}^{-1}(y) = \nabla F(y - c)$. We check that $B_F(\theta_1 : \theta_2) = B_{\bar{F}}(\theta_1 : \theta_2) = D_{\nabla, \nabla^*}(P_1 : P_2)$ with $\theta(P_1) =: \theta_1$ and $\theta(P_2) =: \theta_2$. It is indeed well-known that Bregman divergences modulo affine terms coincide [8]. For the quasi-arithmetic averages $M_{\nabla \bar{F}}$ and $M_{\nabla F}$, we thus obtain the following invariance property: $M_{\nabla \bar{F}}(\theta_1, \dots, \theta_n; w) = M_{\nabla F}(\theta_1, \dots, \theta_n; w)$.
- Second, consider an affine change of coordinates $\bar{\theta} = A\theta + b$ for $A \in \text{GL}(d)$ and $b \in \mathbb{R}^d$, and define the potential function $\bar{F}(\bar{\theta})$ such that $\bar{F}(\bar{\theta}) = F(\theta)$. We have $\theta = A^{-1}(\bar{\theta} - b)$ and $\bar{F}(x) = F(A^{-1}(x - b))$.

²The squared Euclidean/Mahalanobis divergence are not metric distances since they fail the triangle inequality.

It follows that $\nabla\bar{F}(x) = (A^{-1})^\top \nabla F(A^{-1}(x - b))$, and we check that $B_{\bar{F}(\bar{\theta}_1:\bar{\theta}_2)} = B_F(\theta_1 : \theta_2)$:

$$\begin{aligned} B_{\bar{F}(\bar{\theta}_1:\bar{\theta}_2)} &= \bar{F}(\bar{\theta}_1) + \bar{F}(\bar{\theta}_2) - \langle \bar{\theta}_1 - \bar{\theta}_2, \nabla\bar{F}(\bar{\theta}_2) \rangle, \\ &= F(\theta_1) - F(\theta_2) - (A(\theta_1 - \theta_2))^\top (A^{-1})^\top \nabla F(\theta_2), \\ &= F(\theta_1) - F(\theta_2) - (\theta_1 - \theta_2)^\top \underbrace{A^\top (A^{-1})^\top}_{(A^{-1}A)^\top = I} \nabla F(\theta_2) = B_F(\theta_1 : \theta_2). \end{aligned}$$

This highlights the invariance that $D_{\nabla, \nabla^*}(P_1 : P_2) = B_F(\theta_1 : \theta_2) = B_{\bar{F}(\bar{\theta}_1:\bar{\theta}_2)}$, i.e., the canonical divergence does not change under a reparameterization of the ∇ -affine coordinate system. For the induced quasi-arithmetic averages $M_{\nabla\bar{F}}$ and $M_{\nabla F}$, we have $\nabla\bar{F}(x) = (A^{-1})^\top \nabla F(A^{-1}(x - b)) = y$, we calculate $x = \nabla\bar{F}(x)^1(y) = A \nabla\bar{F}^{-1}(((A^{-1})^\top)^{-1}y) + b$, and we have

$$\begin{aligned} M_{\nabla\bar{F}}(\bar{\theta}_1, \dots, \bar{\theta}_n; w) &:= \nabla\bar{F}^{-1}\left(\sum_i w_i \nabla\bar{F}(\bar{\theta}_i)\right), \\ &= (\nabla\bar{F})^{-1}\left((A^{-1})^\top \sum_i w_i \nabla F(\theta_i)\right), \\ &= A \nabla F^{-1}\left(\underbrace{((A^{-1})^\top)^{-1} (A^{-1})^\top}_{=I} \sum_i w_i \nabla F(\theta_i)\right), \\ M_{\nabla\bar{F}}(\bar{\theta}_1, \dots, \bar{\theta}_n; w) &= A M_{\nabla F}(\theta_1, \dots, \theta_n; w) + b \end{aligned}$$

More generally, we may define $\bar{F}(\bar{\theta}) = F(A\theta + b) + \langle c, \theta \rangle + d$ and get via Legendre transformation $\bar{F}^*(\bar{\eta}) = F^*(A^*\eta + b^*) + \langle c^*, \eta \rangle + d^*$ (with A^*, b^*, c^* and d^* expressed using A, b, c and d since these parameters are linked by the Legendre transformation).

- Third, the canonical divergences should be considered relative divergences (and not absolute divergences), and defined according to a prescribed arbitrary “unit” $\lambda > 0$. Thus we can scale the canonical divergence by $\lambda > 0$, i.e., $D_{\lambda, \nabla, \nabla^*} := \lambda D_{\nabla, \nabla^*}$. We have $D_{\lambda, \nabla, \nabla^*}(P_1 : P_2) = \lambda B_F(\theta_1 : \theta_2) = \lambda B_{F^*}(\eta_1 : \eta_2) = \lambda Y_F(\theta_1 : \theta_2) = \lambda Y_{F^*}(\eta_1 : \eta_2)$, and $\lambda B_F(\theta_1 : \theta_2) = B_{\lambda F}(\theta_1 : \theta_2)$ (and $\nabla \lambda F = \lambda \nabla F$). We check the scale invariance of quasi-arithmetic averages: $M_{\lambda \nabla F} = M_{\nabla F}$.

Thus we end up with the following invariance and equivariance properties of the quasi-arithmetic averages which have been obtained from an information-geometric viewpoint:

Proposition 2 (Invariance and equivariance of quasi-arithmetic averages) *Let $F(\theta)$ be a function of Legendre type. Then $\bar{F}(\bar{\theta}) := \lambda(F(A\theta + b) + \langle c, \theta \rangle + d)$ for $A \in \text{GL}(d)$, $b, c \in \mathbb{R}^d$, $d \in \mathbb{R}^d$ and $\lambda \in \mathbb{R}_{>0}$ is a Legendre-type function, and we have $M_{\nabla\bar{F}} = A M_{\nabla F} + b$.*

This proposition generalizes Property 1 of scalar QAMs, and untangles the role of scale $\lambda > 0$ from the other invariance roles brought by the Legendre transformation.

5 Quasi-arithmetic mixtures and information geometry

Consider a quasi-arithmetic mean M_f . We consider n probability distributions P_1, \dots, P_n all dominated by a measure μ , and denote by $p_1 = \frac{dP_1}{d\mu}, \dots, p_n = \frac{dP_n}{d\mu}$ their Radon-Nikodym derivatives. Let us define statistical M_f -mixtures of p_1, \dots, p_n :

Definition 4 *The M_f -mixture of n densities p_1, \dots, p_n weighted by $w \in \Delta_n^\circ$ is defined by*

$$(p_1, \dots, p_n; w)^{M_f}(x) := \frac{M_f(p_1(x), \dots, p_n(x); w)}{\int M_f(p_1(x), \dots, p_n(x); w) d\mu(x)}.$$

The quasi-arithmetic mixture (QAMIX for short) $(p_1, \dots, p_n; w)^{M_f}$ generalizes the ordinary statistical mixture $\sum_{i=1}^d w_i p_i(x)$ when $f(t) = t$ and $M_f = A$ is the arithmetic mean. A statistical M_f -mixture can be interpreted as the M_f -integration of its weighted component densities, the densities p_i 's. The power mixtures $(p_1, \dots, p_n; w)^{M_p}$ (including the ordinary and geometric mixtures) are called α -mixtures in [4] with $\alpha(p) = 1 - 2p$ (or $p = \frac{1-\alpha}{2}$). A nice characterization of the α -mixtures is that these mixtures are the density centroids of the weighted mixture components with respect to the α -divergences [4] (proven by calculus of variation):

$$(p_1, \dots, p_n; w)^{M_\alpha} = \arg \min_p w_i D_\alpha(p_i, p),$$

where D_α denotes the α -divergences [5, 44]. M_f -mixtures have also been used to define a generalization of the Jensen-Shannon divergence [36] between densities p and q as follows:

$$D_{\text{JS}}^{M_f}(p, q) := \frac{1}{2} (D_{\text{KL}}(p : (pq)^{M_f}) + D_{\text{KL}}(q : (pq)^{M_f})) \geq 0, \quad (13)$$

where $D_{\text{KL}}(p : q) = \int p(x) \log \frac{p(x)}{q(x)} d\mu(x)$ is the Kullback-Leibler divergence, and $(pq)^{M_f} := (p, q; \frac{1}{2}, \frac{1}{2})^{M_f}$. The ordinary JSD is recovered when $f(t) = t$ and $M_f = A$:

$$D_{\text{JS}}(p, q) = \frac{1}{2} \left(D_{\text{KL}} \left(p : \frac{p+q}{2} \right) + D_{\text{KL}} \left(q : \frac{p+q}{2} \right) \right).$$

Quasi-arithmetic mixtures of two components have also been used to upper bound the probability of error in Bayesian hypothesis testing [35].

Let us give some examples of parametric families of probability distributions that are closed under quasi-arithmetic mixing:

- Consider a natural exponential family [9] $\mathcal{E} = \{p_\theta = \exp(\theta \cdot x - F(\theta)) : \theta \in \Theta\}$. Function $F(\theta) = \log \int \exp(\theta \cdot x) d\mu$ is called the cumulant function (the log-normalizer function called log-partition in statistical physics), and is of Legendre type when the exponential family is steep [9]. Regular exponential families with full exponential families with open natural parameter spaces Θ are steep. We have Shannon entropy of density $p_\theta \in \mathcal{E}$ expressed using the negative convex conjugate [42] which is concave: $H(p_\theta) = -F^*(\eta)$ with $\eta = \nabla F(\theta)$. Since exponential families are closed under geometric mixtures, i.e. $(p_{\theta_1} p_{\theta_2})^G = p_{\frac{\theta_1 + \theta_2}{2}}$, we have Shannon entropy which can be expressed using the convex conjugate:

$$H((p_{\theta_1} p_{\theta_2})^G) = -F^* \left(\nabla F \left(\frac{\theta_1 + \theta_2}{2} \right) \right).$$

We can rewrite $\nabla F \left(\frac{\theta_1 + \theta_2}{2} \right)$ as the quasi-arithmetic average $M_{\nabla F^*}(\eta_1, \eta_2)$. More generally, the geometric mixtures of n densities of an exponential family belongs to that exponential family:

$$(p_{\theta_1}, \dots, p_{\theta_n}; w)^G \propto \prod_{i=1}^n p_{\theta_i}^{w_i} = p_{\sum_{i=1}^n w_i \theta_i}.$$

That is the normalization constant of $\prod_{i=1}^n p_{\theta_i}^{w_i}$ is $\exp(F(\sum_i w_i \theta_i)) - \sum w_i F(\theta_i) = \exp(-J_F(\theta_1, \dots, \theta_n; w))$, where J_F is called the Jensen diversity index [15, 39].

We may also build an exponential family by considering $n + 1$ probability distributions P_0, \dots, P_n mutually absolutely continuous and all dominated by a reference measure μ . Let p_0, \dots, p_n denote their Radon-Nikodym densities such that $\log \frac{p_1}{p_0}, \dots, \log \frac{p_n}{p_0}$ are linear independent. Then consider the geometric mixture family:

$$\mathcal{G} = \{(p_0, \dots, p_n; w)^G = : w \in \Delta_n^\circ\}.$$

We have $(p_0, p_1, \dots, p_n; w)^G \propto \exp \left(\sum_{i=1}^n w_i \log \frac{p_i}{p_0} \right) p_0(x)$. We let the natural parameter be $\theta = (w_1, \dots, w_n) \in \Delta_n^\circ$. When $n = 1$, Grünwald [21] called \mathcal{G} a likelihood ratio exponential family (LREF).

Uni-order LREFs ($n = 1$) have been studied in [13, 37]: It has the advantage of considering a non-parametric statistical model [53] as a 1D exponential family model which yields a convenient framework for studying the statistical model \mathcal{G} under the lens of well-studied exponential families [9].

Remark 2 We may consider another equivalent definition of the ordinary JSD [50, 27] given by

$$D_{\text{JS}}(p, q) = H\left(\frac{p+q}{2}\right) - \frac{H(p) + H(q)}{2} \geq 0, \quad (14)$$

where $H(p) = -\int p(x) \log p(x) d\mu(x)$ is the strictly concave Shannon entropy. Thus we may consider the following generalization of the JSD:

$$H_{\text{JS}}^{M_f, M_g}(p, q) = H((pq)^{M_f}) - M_g(H(p), H(q)), \quad (15)$$

where M_f and M_g are two quasi-arithmetic means. The first QAM is used to build a quasi-arithmetic mixture while the second QAM is used to average scalars. When $f(t) = g(t) = t$, we recover the ordinary JSD with $M_f = M_g = A$. Let us introduce the (M, N) -Jensen divergences [43] according to two generic symmetric bivariate means M and N :

$$J_F^{M, N}(\theta_1, \theta_2) = M(F(\theta_1), F(\theta_2)) - F(N(\theta_1, \theta_2)). \quad (16)$$

We recover the ordinary Jensen divergence [15, 39] when $M = N = A$:

$$J_F(\theta_1, \theta_2) = J_F^{A, A}(\theta_1, \theta_2) = \frac{F(\theta_1) + F(\theta_2)}{2} - F\left(\frac{\theta_1 + \theta_2}{2}\right).$$

Jensen divergences are non-negative and equal to zero only when $\theta_1 = \theta_2$ when F is a strictly convex function. Similarly, by definition, $J_F^{M, N}(\theta_1, \theta_2) \geq 0$ with equality only when $\theta_1 = \theta_2$ when F is said (M, N) -convex [33] and [32] (Appendix A). A (G, A) -convex function is said log-convex and a (G, G) -convex function is said multiplicative convex. We can test whether a function g is strictly (M_{f_1}, M_{f_2}) -convex by checking whether the function $f_2 \circ g \circ f_1$ is strictly convex or not (see the correspondence Lemma A.2.2 in [32]). For densities p_{θ_1} and p_{θ_2} belonging to a same exponential family \mathcal{E} , we have

$$\begin{aligned} H_{\text{JS}}^{G, A}(p_{\theta_1}, p_{\theta_2}) &= H((p_{\theta_1} p_{\theta_2})^G) - A(H(p_{\theta_1}), H(p_{\theta_2})), \\ &= -F^*(M_{\nabla F^*}(\eta_1, \eta_2)) + \frac{F^*(\eta_1) + F^*(\eta_2)}{2} = J_{F^*}^{\nabla F^*, A}(\eta_1, \eta_2), \end{aligned}$$

where $J^{\nabla F^*, A}$ is the $(\nabla F^*, A)$ -Jensen divergence defined according to two means. Thus we have $H_{\text{JS}}^{G, A}(p_{\theta_1}, p_{\theta_2}) \geq 0$ iff F^* is $(\nabla F^*, A)$ -convex.

To see how $H_{\text{JS}}^{G, A}$ differs from D_{JS}^G defined in [36], let us introduce the cross-entropy between p and q : $H^\times(p : q) = -\int p(x) \log q(x) d\mu(x)$. Then $D_{\text{KL}}(p : q) = H^\times(p : q) - H(p)$ with $H(p) = H^\times(p : p)$, and we have

$$D_{\text{JS}}^G(p : q) = \frac{1}{2} (D_{\text{KL}}(p : (pq)^G) + D_{\text{KL}}(q : (pq)^G)) \geq 0, \quad (17)$$

$$= \frac{1}{2} (H^\times(p : (pq)^G) - H(p) + H^\times(q : (pq)^G) - H(q)) \geq 0, \quad (18)$$

$$= H^\times((pq)^A : (pq)^G) - \frac{H(p) + H(q)}{2} \geq 0. \quad (19)$$

However, we have $H_{\text{JS}}^{G, A}(p : q) = H^\times((pq)^G : (pq)^G) - \frac{H(p) + H(q)}{2}$. Therefore, the dissimilarity $H_{\text{JS}}^{G, A}(p : q)$ can be potentially negative when $H^\times((pq)^G : (pq)^G) \leq \frac{H(p) + H(q)}{2}$.

- Let p_0, p_1, \dots, p_n denotes $n+1$ linearly independent densities, and consider their (arithmetic/standard) mixture family [5]: $\mathcal{M} = \{m_\theta(x) = \sum_{i=0}^n w_i p_i(x) : w \in \Delta_n^\circ\}$ with $\theta = (w_1, \dots, w_n) \in \Delta_{n-1}^\circ$ (and $w_0 = 1 - \sum_{i=1}^n w_i$). The Shannon negentropy $F(\theta) = -H(m_\theta)$ is a Legendre type function [40]. Since the mixture of two densities of a mixture family \mathcal{M} belongs to \mathcal{M} (i.e., $\frac{m_{\theta_1} + m_{\theta_2}}{2} = m_{\frac{\theta_1 + \theta_2}{2}}$), we have $H\left(\frac{m_{\theta_1} + m_{\theta_2}}{2}\right) = H\left(m_{\frac{\theta_1 + \theta_2}{2}}\right) = -F\left(\frac{\theta_1 + \theta_2}{2}\right)$. It follows that $D_{\text{JS}}(m_{\theta_1} : m_{\theta_2}) = J_F(\theta_1 : \theta_2) \geq 0$.
- Consider the family of scale Cauchy distributions $\mathcal{C} = \left\{ p_s(x) = \frac{1}{\pi s} \frac{1}{1 + \left(\frac{x}{s}\right)^2} : s \in \mathbb{R}_{>0} \right\}$. The harmonic mixture $(p_{s_1} p_{s_2})^H$ of two Cauchy distributions is a Cauchy distribution $p_{s_{12}}$ [35] with parameter $s_{12} = \sqrt{\frac{s_1 s_2^2 + s_2 s_1^2}{s_1 + s_2}}$. More generally, the harmonic mixtures of n scale Cauchy distributions is a Cauchy distribution.
- The power mixture of central multivariate t -distributions is a central multivariate t -distribution [35].




Figure 4: Statistical models in the quasi-mixture family are parametrized by a vector in the open standard simplex.

In general, we may consider quasi-arithmetic paths between densities on the space \mathcal{P} of probability density functions with a common support all dominated by a reference measure. On \mathcal{P} , we can build a parametric statistical model called a M_f -mixture family of order n as follows:

$$\mathcal{F}_{p_0, p_1, \dots, p_n}^{M_f} := \{(p_0, p_1, \dots, p_n; (\theta, 1))^{M_f} : \theta \in \Delta_n^\circ\}.$$

In particular, power q -paths have been investigated in [29] with applications in annealing importance sampling and other Monte Carlo methods. The information geometry of such a density space with quasi-arithmetic paths has been investigated in [19] by considering quasi-arithmetic means with respect to a monotone increasing and concave function. See also [51, 20].

We conclude by giving a geometric definition of a generalization of the Jensen-Shannon divergence on \mathcal{P} according to an arbitrary affine connection [5, 53] ∇ :

Definition 5 (Affine connection-based ∇ -Jensen-Shannon divergence) *Let ∇ be an affine connection on the space of densities \mathcal{P} , and $\gamma_\nabla(p, q; t)$ the geodesic linking density $p = \gamma_\nabla(p, q; 0)$ to density $q = \gamma_\nabla(p, q; 1)$. Then the ∇ -Jensen-Shannon divergence is defined by:*

$$D_{\nabla}^{\text{JS}}(p, q) := \frac{1}{2} \left(D_{\text{KL}} \left(p : \gamma_\nabla \left(p, q; \frac{1}{2} \right) \right) + D_{\text{KL}} \left(q : \gamma_\nabla \left(p, q; \frac{1}{2} \right) \right) \right). \quad (20)$$

When $\nabla = \nabla^m$ is chosen as the mixture connection [5], we end up with the ordinary Jensen-Shannon divergence since $\gamma_{\nabla^m}(p, q; \frac{1}{2}) = \frac{p+q}{2}$. When $\nabla = \nabla^e$, the exponential connection, we get the geometric Jensen-Shannon divergence [36] since $\gamma_{\nabla^e}(p, q; \frac{1}{2}) = (pq)^G$ is a statistical geometric mixture.

When the space of densities \mathcal{P} is an exponential family or a mixture family with carrying a dually flat structure $(\mathcal{P}, g_{\text{Fisher}}, \nabla^m, \nabla^e)$ where g_{Fisher} denotes the Riemannian Fisher information metric [5], we have

the Kullback-Leibler divergence D_{KL} which can be expressed using the canonical divergence D_{∇^m, ∇^e} , and the Jensen-Shannon divergence can be written geometrically as

$$D_{\text{JS}}(p : q) = D_{\nabla}^{\text{JS}}(P, Q) := \frac{1}{2} \left(D_{\nabla^m, \nabla^e} \left(p : \gamma_{\nabla^m}(p, q; \frac{1}{2}) \right) + D_{\nabla^m, \nabla^e} \left(q : \gamma_{\nabla^m}(p, q; \frac{1}{2}) \right) \right),$$

where P and Q denote the points on \mathcal{P} representing the densities p and q .

Furthermore, we may consider the α -connections [5] ∇^α of parametric or non-parametric statistical models, and skew the geometric Jensen-Shannon divergence to define the β -skewed ∇^α -JSD:

$$D_{\nabla^\alpha, \beta}^{\text{JS}}(p, q) = \beta D_{\text{KL}}(p : \gamma_{\nabla^\alpha}(p, q; \beta)) + (1 - \beta) D_{\text{KL}}(q : \gamma_{\nabla^\alpha}(p, q; \beta)).$$

6 Concluding remarks

In this paper, we presented two generalizations of the scalar quasi-arithmetic means [22] through the lens of information geometry and discussed some of their applications:

- The first generalization of scalar quasi-arithmetic means consisted in defining pairs of *quasi-arithmetic averages* induced by the gradient maps of pairs of Legendre-type functions. These dual quasi-arithmetic averages are used in information geometry to express points on dual geodesics and sided barycenters in the dually affine θ - and η -coordinate systems. Furthermore, we proved that $M_{\nabla F} = M_{\nabla \bar{F}} = M_{[\nabla F]}$ where $[\nabla \bar{F}]$ denotes the equivalence class of Legendre type functions such that $\bar{F}(\bar{\theta}) = \lambda F(\theta + b) + \langle c, \theta \rangle + d \sim F(\theta)$. This property generalizes the well-known fact that quasi-arithmetic means $M_f = M_g$ iff $g = \lambda f + c$ and distinguishes the scaling invariance by $\lambda > 0$ with the Legendre invariance by c .
- The second generalization of quasi-arithmetic means defined *statistical quasi-arithmetic mixtures* by normalizing quasi-arithmetic means of their densities: In particular, we showed how exponential families are closed under geometric mixtures, and described a generic way to build exponential families of order n from geometric mixtures of $n + 1$ linear independent log ratio densities $\log \frac{p_i}{p_0}$. The statistical geometric mixture family construction holds similarly for other quasi-arithmetic mixture families. Last, we gave a generic geometric definition of the Jensen-Shannon divergence based on affine connections which generalizes both the ordinary Jensen-Shannon divergence [27] and the geometric Jensen-Shannon divergence [36]. This demonstrates the rich interplay of divergences with information geometry.

References

- [1] S Abramovich, M Klaričić Bakula, M Matić, and J Pečarić. A variant of Jensen–Steffensen’s inequality and quasi-arithmetic means. *Journal of mathematical analysis and applications*, 307(1):370–386, 2005.
- [2] M Alić, B Mond, J Pečarić, and V Volenec. The arithmetic-geometric-harmonic-mean and related matrix inequalities. *Linear Algebra and its Applications*, 264:55–62, 1997.
- [3] Shun-ichi Amari. Differential-geometrical methods in statistics. *Lecture Notes on Statistics*, 28:1, 1985.
- [4] Shun-ichi Amari. Integration of stochastic models by minimizing α -divergence. *Neural computation*, 19(10):2780–2796, 2007.
- [5] Shun-ichi Amari. *Information Geometry and Its Applications*. Applied Mathematical Sciences. Springer Japan, 2016.
- [6] Mahmut Bajraktarević. *Sur une équation fonctionnelle aux valeurs moyennes*. 1958.
- [7] Miguel A Ballester and José Luis García-Lapresta. Using quasiarithmetic means in a sequential decision procedure. In *EUSFLAT Conf.(2)*, pages 479–484, 2007.

- [8] Arindam Banerjee, Srujana Merugu, Inderjit S Dhillon, Joydeep Ghosh, and John Lafferty. Clustering with Bregman divergences. *Journal of machine learning research*, 6(10), 2005.
- [9] Ole Barndorff-Nielsen. *Information and exponential families: in statistical theory*. John Wiley & Sons, 2014.
- [10] Rajendra Bhatia, Stephane Gaubert, and Tanvi Jain. Matrix versions of the Hellinger distance. *Letters in Mathematical Physics*, 109(8):1777–1804, 2019.
- [11] Stephen Boyd, Stephen P Boyd, and Lieven Vandenberghe. *Convex optimization*. Cambridge university press, 2004.
- [12] Lev M. Bregman. The relaxation method of finding the common point of convex sets and its application to the solution of problems in convex programming. *USSR computational mathematics and mathematical physics*, 7(3):200–217, 1967.
- [13] Rob Breckelmanns, Frank Nielsen, Alireza Makhzani, Aram Galstyan, and Greg Ver Steeg. Likelihood ratio exponential families. *arXiv preprint arXiv:2012.15480*, 2020.
- [14] Peter S Bullen, Dragoslav S Mitrinovic, and Means Vasic. *Means and their inequalities*, volume 31. Springer Science & Business Media, 2013.
- [15] Jacob Burbea and C Rao. On the convexity of some divergence measures based on entropy functions. *IEEE Transactions on Information Theory*, 28(3):489–495, 1982.
- [16] Tomasa Calvo, Gaspar Mayor, and Radko Mesiar. *Aggregation operators: new trends and applications*, volume 97. Springer Science & Business Media, 2002.
- [17] Francis Clarke. On the inverse function theorem. *Pacific Journal of Mathematics*, 64(1):97–102, 1976.
- [18] Bruno De Finetti. *Sul concetto di media*. Istituto italiano degli attuari, 1931.
- [19] Shinto Eguchi and Osamu Komori. Path connectedness on a space of probability density functions. In *Geometric Science of Information: Second International Conference, GSI 2015, Palaiseau, France, October 28-30, 2015, Proceedings 2*, pages 615–624. Springer, 2015.
- [20] Shinto Eguchi, Osamu Komori, and Atsumi Ohara. Information geometry associated with generalized means. In *Information Geometry and Its Applications: On the Occasion of Shun-ichi Amari's 80th Birthday, IGAI IV Liblice, Czech Republic, June 2016*, pages 279–295. Springer, 2018.
- [21] Peter D Grünwald. *The minimum description length principle*. MIT press, 2007.
- [22] Godfrey Harold Hardy, John Edensor Littlewood, George Pólya, and György Pólya. *Inequalities*. Cambridge university press, 1952.
- [23] Børge Jessen. Bemærkninger om konvekse Funktioner og Uligheder imellem Middelværdier. I. *Matematisk tidsskrift. B*, pages 17–28, 1931.
- [24] Konrad Knopp. Über reihen mit positiven gliedern. *Journal of the London Mathematical Society*, 1(3):205–211, 1928.
- [25] Andreï Nikolaevich Kolmogorov. Sur la notion de la moyenne. *G. Bardi, tip. della R. Accad. dei Lincei*, 12:388–391, 1930.
- [26] Osamu Komori and Shinto Eguchi. A unified formulation of k -Means, fuzzy c -Means and Gaussian mixture model by the Kolmogorov–Nagumo average. *Entropy*, 23(5):518, 2021.
- [27] Jianhua Lin. Divergence measures based on the Shannon entropy. *IEEE Transactions on Information theory*, 37(1):145–151, 1991.

- [28] László Losonczi. Equality of two variable weighted means: reduction to differential equations. *aequationes mathematicae*, 58(3):223–241, 1999.
- [29] Vaden Masrani, Rob Brekelmans, Thang Bui, Frank Nielsen, Aram Galstyan, Greg Ver Steeg, and Frank Wood. q -paths: Generalizing the geometric annealing path using power means. In *Uncertainty in Artificial Intelligence*, pages 1938–1947. PMLR, 2021.
- [30] Mitio Nagumo. Über eine Klasse der Mittelwerte. In *Japanese journal of mathematics: transactions and abstracts*, volume 7, pages 71–79. The Mathematical Society of Japan, 1930.
- [31] Naomichi Nakajima and Toru Ohmoto. The dually flat structure for singular models. *Information Geometry*, 4(1):31–64, 2021.
- [32] Constantin Niculescu and Lars-Erik Persson. *Convex functions and their applications*, volume 23. Springer, 2006.
- [33] Constantin P Niculescu. Convexity according to means. *Mathematical Inequalities and Applications*, 6:571–580, 2003.
- [34] Frank Nielsen. k -MLE: A fast algorithm for learning statistical mixture models. In *2012 IEEE international conference on acoustics, speech and signal processing (ICASSP)*, pages 869–872. IEEE, 2012.
- [35] Frank Nielsen. Generalized Bhattacharyya and Chernoff upper bounds on Bayes error using quasi-arithmetic means. *Pattern Recognition Letters*, 42:25–34, 2014.
- [36] Frank Nielsen. On the Jensen–Shannon symmetrization of distances relying on abstract means. *Entropy*, 21(5):485, 2019.
- [37] Frank Nielsen. Revisiting Chernoff Information with Likelihood Ratio Exponential Families. *Entropy*, 24(10):1400, 2022.
- [38] Frank Nielsen. Statistical Divergences between Densities of Truncated Exponential Families with Nested Supports: Duo Bregman and Duo Jensen Divergences. *Entropy*, 24(3):421, 2022.
- [39] Frank Nielsen and Sylvain Boltz. The Burbea-Rao and Bhattacharyya centroids. *IEEE Transactions on Information Theory*, 57(8):5455–5466, 2011.
- [40] Frank Nielsen and Gaëtan Haderes. Monte Carlo information-geometric structures. In *Geometric Structures of Information*, pages 69–103. Springer, 2019.
- [41] Frank Nielsen and Richard Nock. Sided and symmetrized Bregman centroids. *IEEE Transactions on Information Theory*, 55(6):2882–2904, 2009.
- [42] Frank Nielsen and Richard Nock. Entropies and cross-entropies of exponential families. In *2010 IEEE International Conference on Image Processing*, pages 3621–3624. IEEE, 2010.
- [43] Frank Nielsen and Richard Nock. Generalizing skew Jensen divergences and Bregman divergences with comparative convexity. *IEEE Signal Processing Letters*, 24(8):1123–1127, 2017.
- [44] Frank Nielsen, Richard Nock, and Shun-ichi Amari. On clustering histograms with k -means by using mixed α -divergences. *Entropy*, 16(6):3273–3301, 2014.
- [45] Richard Nock and Frank Nielsen. Fitting the smallest enclosing Bregman ball. In *European Conference on Machine Learning*, pages 649–656. Springer, 2005.
- [46] Zsolt Páles and Lars-Erik Persson. Hardy-type inequalities for means. *Bulletin of the Australian Mathematical Society*, 70(3):521–528, 2004.

- [47] Bruno Pelletier. Informative barycentres in statistics. *Annals of the Institute of Statistical Mathematics*, 57(4):767–780, 2005.
- [48] Ralph Tyrrell Rockafellar. Conjugates and Legendre transforms of convex functions. *Canadian Journal of Mathematics*, 19:200–205, 1967.
- [49] Hirohiko Shima and Katsumi Yagi. Geometry of Hessian manifolds. *Differential geometry and its applications*, 7(3):277–290, 1997.
- [50] Robin Sibson. Information radius. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*, 14(2):149–160, 1969.
- [51] Rui F Vigelis, Luiza HF de Andrade, and Charles C Cavalcante. On the existence of paths connecting probability distributions. In *Geometric Science of Information: Third International Conference, GSI 2017, Paris, France, November 7-9, 2017, Proceedings 3*, pages 801–808. Springer, 2017.
- [52] Roger Webster et al. *Convexity*. Oxford University Press, 1994.
- [53] Jun Zhang. Nonparametric information geometry: From divergence function to referential-representational biduality on statistical manifolds. *Entropy*, 15(12):5384–5418, 2013.
- [54] XH Zhang, GD Wang, and YM Chu. Convexity with respect to Hölder mean involving zero-balanced hypergeometric functions. *J. Math. Anal. Appl.*, 353(1):256–259, 2009.