

TRANSFORMATION AND SYMMETRIES FOR THE ANDREWS-GARVAN CRANK FUNCTION

RISHABH SARMA

ABSTRACT. Let $R(z, q)$ be the two-variable generating function of Dyson's rank function. In a recent joint work with Frank Garvan, we investigated the transformation of the elements of the p -dissection of $R(\zeta_p, q)$, where ζ_p is a primitive p -th root of unity, under special congruence subgroups of $SL_2(\mathbb{Z})$, leading us to interesting symmetry observations. In this work, we derive analogous transformation results for the two-variable crank generating function $C(z, q)$ in terms of generalized eta products. We consider the action of the group $\Gamma_0(p)$ on the elements of the p -dissection of $C(z, q)$, leading us to new symmetries for the crank function. As an application, we give a new proof of the crank theorem predicted by Dyson in 1944 and resolved by Andrews and Garvan in 1988. Furthermore, we present identities expressing the elements of the crank dissection in terms of generalized eta products for primes $p = 11, 13, 17$ and 19 .

1. INTRODUCTION

Let $p(n)$ denote the number of partitions of n . The following are Ramanujan's famous partition congruences :

$$\begin{aligned} p(5n + 4) &\equiv 0 \pmod{5}, \\ p(7n + 5) &\equiv 0 \pmod{7}, \\ p(11n + 6) &\equiv 0 \pmod{11}. \end{aligned}$$

In 1944, Dyson, in search of a combinatorial explanation of the mod 5 and 7 congruences of Ramanujan gave the definition of rank of a partition [10]. It is defined as the largest part minus the number of parts in a partition. In particular, the rank statistic decomposes the partitions of $5n + 4$ and $7n + 5$ into five and seven equinumerous classes respectively, a conjecture that was proven 10 years later by Atkin and Swinnerton-Dyer. He also prophesied the existence of a "crank" statistic for the mod 11 congruence, and it wasn't until 1988 that the elusive crank was found by Andrews and Garvan [1]. The crank of a partition is defined as the largest part if there are no ones in the partition and otherwise the number of parts larger than the number of ones minus the number of ones. Let $M(r, t, n)$ denote the number of partitions of n with crank congruent to $r \pmod{t}$. Then they show that

$$(1.1) \quad M(r, 5, 5n + 4) = \frac{1}{5}p(5n + 4), \quad 0 \leq r \leq 4,$$

Date: January 27, 2023.

2010 Mathematics Subject Classification. 05A19, 11B65, 11F11, 11F37, 11P82, 11P83, 33D15.

Key words and phrases. Crank function, Dyson's rank function, modular forms, partitions.

$$(1.2) \quad M(r, 7, 7n + 5) = \frac{1}{7}p(7n + 5), \quad 0 \leq r \leq 6,$$

$$(1.3) \quad M(r, 11, 11n + 6) = \frac{1}{11}p(11n + 6), \quad 0 \leq r \leq 10.$$

Let $M(m, n)$ denote the number of partitions of n with crank m . Then the two variable generating function for the crank function, due to Andrews and Garvan (see [1]) is given by

$$\sum_{n=0}^{\infty} \sum_m M(m, n) z^m q^n = (1 - z)q + \frac{(q; q)_{\infty}}{(zq; q)_{\infty}(z^{-1}q; q)_{\infty}}.$$

Equivalently we define

$$C(z, q) := \frac{(q; q)_{\infty}}{(zq; q)_{\infty}(z^{-1}q; q)_{\infty}} = 1 + (-1 + z + z^{-1})q + \sum_{n=1}^{\infty} \sum_m M(m, n) z^m q^n.$$

In this paper, we look at the transformation and symmetry of this function, when z is a primitive p -th root of unity. We find that the crank function satisfies the same type of symmetries as the rank generating function, which was the subject of our interest in [16].

Let $N(m, n)$ denote the number of partitions of n with rank m and let $R(z, q)$ denote the two-variable generating function for the Dyson rank function so that

$$R(z, q) = \sum_{n=0}^{\infty} \sum_m N(m, n) z^m q^n.$$

The following two identities (1.4) and (1.5) (refer [12] and [13]) originally found in Ramanujan's Lost Notebook [22] give the 5-dissection of the q -series expansion of $C(\zeta_5, q)$ and $R(\zeta_5, q)$ respectively.

$$F(q) = \frac{(1 - q)(1 - q^2)(1 - q^3) \cdots}{(1 - 2q \cos \frac{2n\pi}{5} + q^2)(1 - 2q^2 \cos \frac{2n\pi}{5} + q^4) \cdots},$$

$$f(q) = 1 + \frac{q}{(1 - 2q \cos \frac{2n\pi}{5} + q^2)} \frac{q^4}{(1 - 2q \cos \frac{2n\pi}{5} + q^2)(1 - 2q^2 \cos \frac{2n\pi}{5} + q^4)} + \cdots,$$

$$(1.4) \quad F(q^{\frac{1}{5}}) = A(q) - 4q^{\frac{1}{5}} \cos^2 \frac{2n\pi}{5} B(q) + 2q^{\frac{2}{5}} \cos \frac{4n\pi}{5} C(q) - 2q^{\frac{3}{5}} \cos \frac{2n\pi}{5} D(q),$$

$$(1.5) \quad f(q^{\frac{1}{5}}) = \left\{ A(q) - 4 \sin^2 \frac{n\pi}{5} \phi(q) \right\} + q^{\frac{1}{5}} B(q) + 2q^{\frac{2}{5}} \cos \frac{2n\pi}{5} C(q) - 2q^{\frac{3}{5}} \cos \frac{2n\pi}{5} \left\{ D(q) + 4 \sin^2 \frac{2n\pi}{5} \frac{\psi(q)}{q} \right\},$$

where

$$A(q) = \frac{1 - q^2 - q^3 + q^9 + \cdots}{(1 - q)^2(1 - q^4)^2(1 - q^6)^2 \cdots},$$

$$\begin{aligned}
B(q) &= \frac{(1-q^5)(1-q^{10})(1-q^{15})\cdots}{(1-q)(1-q^4)(1-q^6)\cdots}, \\
C(q) &= \frac{(1-q^5)(1-q^{10})(1-q^{15})\cdots}{(1-q^2)(1-q^3)(1-q^7)\cdots}, \\
D(q) &= \frac{1-q-q^4+q^7+\cdots}{(1-q^2)^2(1-q^3)^2(1-q^7)^2\cdots},
\end{aligned}$$

and

$$\begin{aligned}
\phi(q) &= -1 + \left\{ \frac{1}{1-q} + \frac{q^5}{(1-q)(1-q^4)(1-q^6)} + \frac{q^{20}}{(1-q)(1-q^4)(1-q^6)(1-q^9)(1-q^{11})} + \cdots \right\}, \\
\psi(q) &= -1 + \left\{ \frac{1}{1-q^2} + \frac{q^5}{(1-q^2)(1-q^3)(1-q^7)} + \frac{q^{20}}{(1-q^2)(1-q^3)(1-q^7)(1-q^8)(1-q^{12})} + \cdots \right\}.
\end{aligned}$$

We rewrite Ramanujan's identities (1.4) and (1.5) in terms of generalized eta products to get

$$\begin{aligned}
(1.6) \quad q^{-\frac{1}{24}}C(\zeta_5, q) &= \frac{\eta(25z)\eta_{5,2}(5z)}{\eta_{5,1}(5z)^2} + (\zeta_5 + \zeta_5^4 - 1) \frac{\eta(25z)}{\eta_{5,1}(5z)} \\
&\quad - (\zeta_5 + \zeta_5^4 + 1) \frac{\eta(25z)}{\eta_{5,2}(5z)} - (\zeta_5 + \zeta_5^4) \frac{\eta(25z)\eta_{5,1}(5z)}{\eta_{5,2}(5z)^2},
\end{aligned}$$

$$\begin{aligned}
(1.7) \quad q^{-\frac{1}{24}}(R(\zeta_5, q) - (\zeta_5 + \zeta_5^4 - 2)\phi(q^5) + (1 + 2\zeta_5 + 2\zeta_5^4)q^{-2}\psi(q^5)) \\
= \frac{\eta(25z)\eta_{5,2}(5z)}{\eta_{5,1}(5z)^2} + \frac{\eta(25z)}{\eta_{5,1}(5z)} + (\zeta_5 + \zeta_5^4) \frac{\eta(25z)}{\eta_{5,2}(5z)} - (\zeta_5 + \zeta_5^4) \frac{\eta(25z)\eta_{5,1}(5z)}{\eta_{5,2}(5z)^2}.
\end{aligned}$$

One can observe that the RHS of the above two equations are weakly holomorphic modular forms of weight $\frac{1}{2}$ on the group $\Gamma_0(25) \cap \Gamma_1(5)$. The modularity of the rank generating function $R(z, q)$, where z is replaced by a primitive p -th root of unity for a general prime p was first studied by Bringmann and Ono [6]. Building on the groundbreaking work of Zwegers [24] and [25] around the turn of the century, who realised how Ramanujan's mock theta functions occur naturally as the holomorphic parts of certain real analytic modular forms, Bringmann and Ono showed that $q^{-1/24}R(\zeta_p, q)$ is the holomorphic part a weak Maass form of weight $\frac{1}{2}$ on $\Gamma_1(576.p^4)$, where the non-holomorphic part is a Mordell integral of a theta function. Later, together with Rhoades [7], they used this result to prove that the rank sum modulo p is a weakly holomorphic modular form of weight $\frac{1}{2}$ on $\Gamma_1(576.p^6)$. Garvan, in 2019, observed that the introduction of a simple multiplier and a generalized correction series factor reduces the modularity of the rank function to larger and more natural congruence subgroups of $\mathrm{SL}_2(\mathbb{Z})$, and he obtained a generalized analogue of Ramanujan's result (1.5).

Theorem 1.1 (Garvan [15, Theorem 1.2, p.202]). *For $p > 3$ prime and $1 \leq a \leq \frac{1}{2}(p-1)$ define*

$$(1.8) \quad \Phi_{p,a}(q) := \begin{cases} \sum_{n=0}^{\infty} \frac{q^{pn^2}}{(q^a; q^p)_{n+1} (q^{p-a}; q^p)_n}, & \text{if } 0 < 6a < p, \\ -1 + \sum_{n=0}^{\infty} \frac{q^{pn^2}}{(q^a; q^p)_{n+1} (q^{p-a}; q^p)_n}, & \text{if } p < 6a < 3p, \end{cases}$$

and

$$(1.9) \quad \mathcal{R}_p(\zeta_p, z) := q^{-\frac{1}{24}} R(\zeta_p, q) - \chi_{12}(p) \sum_{a=1}^{\frac{1}{2}(p-1)} (-1)^a \left(\zeta_p^{3a+\frac{1}{2}(p+1)} + \zeta_p^{-3a-\frac{1}{2}(p+1)} - \zeta_p^{3a+\frac{1}{2}(p-1)} - \zeta_p^{-3a-\frac{1}{2}(p-1)} \right) q^{\frac{a}{2}(p-3a)-\frac{p^2}{24}} \Phi_{p,a}(q^p),$$

where

$$(1.10) \quad \chi_{12}(n) := \left(\frac{12}{n} \right) = \begin{cases} 1 & \text{if } n \equiv \pm 1 \pmod{12}, \\ -1 & \text{if } n \equiv \pm 5 \pmod{12}, \\ 0 & \text{otherwise,} \end{cases}$$

and $q = \exp(2\pi iz)$ with $\Im(z) > 0$. Then the function

$$\eta(p^2 z) \mathcal{R}_p(z)$$

is a weakly holomorphic modular form of weight 1 on the group $\Gamma_0(p^2) \cap \Gamma_1(p)$.

In [15], Garvan also considered the modularity of each element of the p -dissection of $\eta(p^2 z) \mathcal{R}_p(z)$ and obtained the following.

Theorem 1.2 (Garvan [15, Corollary 1.3, p.202]). *Let $p > 3$ be prime and $s_p = \frac{1}{24}(p^2 - 1)$. Then the function*

$$\prod_{n=1}^{\infty} (1 - q^{pn}) \sum_{n=\lceil \frac{1}{p}(s_p) \rceil}^{\infty} \left(\sum_{k=0}^{p-1} N(k, p, pn - s_p) \zeta_p^k \right) q^n$$

is a weakly holomorphic modular form of weight 1 on the group $\Gamma_1(p)$.

In a subsequent joint work with Garvan [16], we generalized and improved these results in his paper [15] on transformations for Dyson's rank function leading us to exciting observations of symmetry among the elements of dissection of the rank generating function. In this paper, we try to implement our ideas in [16] and those of Garvan [15] to obtain analogous results for the crank generating function.

Here and throughout this paper we use the standard q -notation:

$$(a; q)_{\infty} = \prod_{k=0}^{\infty} (1 - aq^k),$$

$$(a; q)_n = \frac{(a; q)_\infty}{(aq^n; q)_\infty},$$

$$(a_1, a_2, \dots, a_j; q)_\infty = (a_1; q)_\infty (a_2; q)_\infty \dots (a_j; q)_\infty,$$

$$(a_1, a_2, \dots, a_j; q)_n = (a_1; q)_n (a_2; q)_n \dots (a_j; q)_n.$$

We also define a class of generalized eta products and functions that we will use frequently in expressions for our crank identities and to study their transformations in the subsequent sections.

Definition 1.3. Following Biagioli (see [4]), define

$$(1.11) \quad f_{N,\rho}(z) := (-1)^{\lfloor \rho/N \rfloor} q^{(N-2\rho)^2/(8N)} (q^\rho, q^{N-\rho}, q^N; q^N)_\infty.$$

Then, for a vector $\vec{n} = (n_0, n_1, n_2, \dots, n_{\frac{p-1}{2}}) \in \mathbb{Z}^{\frac{1}{2}(p+1)}$, define

$$(1.12) \quad j(z) = j(p, \vec{n}, z) = \eta(pz)^{n_0} \prod_{k=1}^{\frac{p-1}{2}} f_{p,k}(z)^{n_k}.$$

We note that

$$(1.13) \quad f_{N,\rho}(z) = f_{N,N+\rho}(z) = f_{N,-\rho}(z),$$

and

$$(1.14) \quad f_{N,\rho}(z) = \eta(Nz) \eta_{N,\rho}(z),$$

where

$$(1.15) \quad \eta(z) := q^{\frac{1}{24}} \prod_{n=1}^{\infty} (1 - q^n),$$

$$(1.16) \quad \eta_{N,k}(z) = q^{\frac{N}{2}P_2(k/N)} \prod_{\substack{m>0 \\ m \equiv \pm k \pmod{N}}} (1 - q^m),$$

where $z \in \mathfrak{h}$, $P_2(t) = \{t\}^2 - \{t\} + \frac{1}{6}$ is the second periodic Bernoulli polynomial, and $\{t\} = t - [t]$ is the fractional part of t .

We also need transformations for $\eta(z)$ due to Knopp and the theta function $f_{N,\rho}(z)$ due to Biagioli.

Theorem 1.4. [18, Theorem 2, p.51] For $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{Z})$ we have

$$\eta(z) \Big| [A]_{\frac{1}{2}} = \nu_\eta(A) \eta(z),$$

where

$$\nu_\eta(A) = \begin{cases} \left(\frac{d}{c}\right)^* \exp\left(\frac{\pi i}{12}((a+d)c - bd(c^2 - 1) - 3c)\right) & \text{if } c \text{ is odd,} \\ \left(\frac{c}{d}\right)_* \exp\left(\frac{\pi i}{12}((a+d)c - bd(c^2 - 1) + 3d - 3 - 3cd)\right) & \text{if } d \text{ is odd,} \end{cases}$$

$$\left(\frac{d}{c}\right)^* = \left(\frac{c}{|d|}\right), \quad \left(\frac{c}{d}\right)_* = \left(\frac{c}{|d|}\right) (-1)^{(\text{sgnc}-1)(\text{sgnd}-1)/4},$$

and (\cdot) is the Jacobi symbol.

Theorem 1.5. [4, Lemma 2.1, p.278] *Let $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(N)$. Then*

$$f_{N,\rho}(z) \Big| [A]_{\frac{1}{2}} = (-1)^{\rho b + \lfloor \rho a/N \rfloor + \lfloor \rho/N \rfloor} \exp\left(\frac{\pi i a b}{N} \rho^2\right) \nu_{\theta_1}({}^N A) f_{N,\rho a}(z),$$

where

$${}^N A = \begin{pmatrix} a & bN \\ c/N & d \end{pmatrix} \in SL_2(\mathbb{Z}),$$

and $\nu_{\theta_1} = \nu_{\eta}^3$ is the theta-multiplier.

Since its discovery, it has been well established that the crank statistic plays a central role in the theory of partitions. Perhaps the most important breakthrough came in 2005 when Mahlburg showed that the crank itself satisfies Ramanujan-type congruences using Hecke's theory of modular forms. In his seminal paper, Mahlburg establishes congruences for infinitely many non-nested arithmetic progressions modulo powers of primes by tying the crank generating function to the Dedekind eta function and Klein forms. Rewriting his functions in the framework of generalized eta products, we have the following result :

Theorem 1.6 (Mahlburg [19, Corollary 3.3]). *Let $p > 3$ be prime, suppose $1 \leq \ell \leq (p-1)$, and define*

$$\mathcal{F}_1\left(\frac{\ell}{p}; z\right) := \frac{\eta(z)^2}{E_{0,\ell}(z)},$$

where, following Yang [23], we consider the generalized eta product

$$E_{0,\ell}(z) = q^{\frac{1}{12}} \prod_{m=1}^{\infty} (1 - \zeta_p^{\ell} q^{m-1})(1 - \zeta_p^{-\ell} q^m).$$

Then $\mathcal{F}_1\left(\frac{\ell}{p}; z\right)$ is a weakly holomorphic modular form of weight 1 on the congruence subgroup $\Gamma_1(2p^2)$.

Mahlburg further goes on to establish a rescaled form of the crank generating function as the sum of a weakly holomorphic function on $\Gamma_1(2p^2)$ and a modular form on $\Gamma_0(p)$ of integral weights ([19], Section 4). We show that after multiplying with an appropriate eta product and a power of q , we can deduce the modularity of the crank generating function to a simpler congruence subgroup of $SL_2(\mathbb{Z})$, resembling the result for the rank generating function in Theorem 1.1. Hence Theorem 1.7 below, where we generalize Ramanujan's result (1.4) for the crank dissection modulo 5.

Theorem 1.7. *Let $p > 3$ be prime. Then the function*

$$\eta(p^2 z) \mathcal{C}_p(z)$$

is a weakly holomorphic modular form of weight 1 on the group $\Gamma_0(p^2) \cap \Gamma_1(p)$, where $\mathcal{C}_p(z) = q^{\frac{1}{24}} C(\zeta_p^{\ell}, q)$.

This result is analogous to the modularity of the rank generating function that was studied in [15]. There we also considered the modularity and transformation of elements of the p -dissection of the rank generating function defined as follows. We note that in the definition for the rank dissection elements $\mathcal{K}_{p,m}^{(R)}(\zeta_p^d; z)$, we have two cases pertaining to whether $-24m$ is a quadratic residue or non-residue modulo p . Throughout [16], we refer to the rank dissection elements by $\mathcal{K}_{p,m}(\zeta_p^d; z)$. We have added the superscript (R) here to distinguish from our crank dissection elements which we define later.

Definition 1.8. ([16], Definition 1.4) For $p > 3$ prime, $0 \leq m \leq p - 1$ and $1 \leq d \leq p - 1$ define $\mathcal{K}_{p,m}^{(R)}(\zeta_p^d; z)$ as follows :

(i) For $m = 0$ or $\left(\frac{-24m}{p}\right) = -1$ define

$$(1.17) \quad \mathcal{K}_{p,m}^{(R)}(\zeta_p^d; z) := q^{m/p} \prod_{n=1}^{\infty} (1 - q^{pn}) \sum_{n=\lceil \frac{1}{p}(s_p-m) \rceil}^{\infty} \left(\sum_{k=0}^{p-1} N(k, p, pn + m - s_p) \zeta_p^{kd} \right) q^n,$$

where $s_p = \frac{1}{24}(p^2 - 1)$, and $q = \exp(2\pi iz)$.

(ii) For $\left(\frac{-24m}{p}\right) = 1$ define

(1.18)

$$\begin{aligned} \mathcal{K}_{p,m}^{(R)}(\zeta_p^d; z) := & q^{m/p} \prod_{n=1}^{\infty} (1 - q^{pn}) \left(\sum_{n=\lceil \frac{1}{p}(s_p-m) \rceil}^{\infty} \left(\sum_{k=0}^{p-1} N(k, p, pn + m - s_p) \zeta_p^{kd} \right) q^n \right. \\ & \left. - 4\chi_{12}(p) (-1)^{a+d+1} \sin\left(\frac{d\pi}{p}\right) \sin\left(\frac{6ad\pi}{p}\right) q^{\frac{1}{p}(\frac{a}{2}(p-3a)-m)} \Phi_{p,a}(q) \right), \end{aligned}$$

where $1 \leq a \leq \frac{1}{2}(p - 1)$ has been chosen so that

$$-24m \equiv (6a)^2 \pmod{p}.$$

In case of the crank generating function, we have the following analogous expression for elements of the dissection and subsequent modularity results. These dissection elements do not feature the correction factor involving the series $\Phi_{p,a}(q)$ as in the quadratic residue case of the rank in the previous definition. Our transformation and symmetry results for elements of the the crank generating function are also analogous to that of the rank generating function in [16].

Definition 1.9. For $p > 3$ prime, $0 \leq m \leq p - 1$ and $1 \leq \ell \leq p - 1$, define

$$\mathcal{K}_{p,m}^{(C)}(\zeta_p^\ell; z) := q^{m/p} \prod_{n=1}^{\infty} (1 - q^{pn}) \sum_{n=\lceil \frac{1}{p}(s_p-m) \rceil}^{\infty} \left(\sum_{k=0}^{p-1} M(k, p, pn + m - s_p) \zeta_p^{k\ell} \right) q^n.$$

Theorem 1.10. Let $p > 3$ be prime and suppose $0 \leq m \leq p - 1$. Then

(i) $\mathcal{K}_{p,0}^{(C)}(\zeta_p; z)$ is a weakly holomorphic modular form of weight 1 on $\Gamma_1(p)$.

(ii) If $1 \leq m \leq (p-1)$ then $\mathcal{K}_{p,m}^{(C)}(\zeta_p, z)$ is a weakly holomorphic modular form of weight 1 on $\Gamma(p)$. In particular,

$$\mathcal{K}_{p,m}^{(C)}(\zeta_p, z) | [A]_1 = \exp\left(\frac{2\pi ibm}{p}\right) \mathcal{K}_{p,m}^{(C)}(\zeta_p, z),$$

$$\text{for } A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_1(p).$$

Analogous to the rank, the action of the congruence subgroup $\Gamma_0(p)$ on $\mathcal{K}_{p,m}^{(C)}(\zeta_p, z)$ leads to our other two important results of the paper, the first one of which illustrates the symmetry among the elements of the crank dissection.

Theorem 1.11. *Let $p > 3$ be prime, $0 \leq m \leq p-1$, and $1 \leq d \leq p-1$. Then*

$$(1.19) \quad \mathcal{K}_{p,m}^{(C)}(\zeta_p, z) | [A]_1 = \frac{\sin(\pi/p)}{\sin(d\pi/p)} (-1)^{d+1} \zeta_p^{mak} \mathcal{K}_{p,ma^2}^{(C)}(\zeta_p^d, z),$$

assuming $1 \leq a, d \leq (p-1)$ and

$$A = \begin{pmatrix} a & k \\ p & d \end{pmatrix} \in \Gamma_0(p).$$

Note : When m is a quadratic residue/non-residue modulo p , the same is correspondingly true for ma^2 . In light of this fact and Theorem 1.11, we have the following corollary.

Corollary 1.12. *Let $p > 3$ be prime. The identities for $\mathcal{K}_{p,m}^{(C)}(\zeta_p, z)$ in terms of generalized eta functions are completely determined by three particular ones namely*

$$\mathcal{K}_{p,0}^{(C)}(\zeta_p, z), \mathcal{K}_{p,m^+}^{(C)}(\zeta_p, z), \mathcal{K}_{p,m^-}^{(C)}(\zeta_p, z),$$

where $\left(\frac{m^+}{p}\right) = 1$, $\left(\frac{m^-}{p}\right) = -1$.

We illustrate our symmetry Theorem 1.11 with an example. The following are the elements of the 11-dissection of $\eta(p^2z) \mathcal{C}_p(\zeta_p, z)$ for $p = 11$ in terms of generalized eta functions. A similar form of this identity can also be found in the works of Garvan [14] and Hirschhorn [17].

$$\mathcal{K}_{11,0}^{(C)}(\zeta_{11}, z) = 0,$$

$$\mathcal{K}_{11,1}^{(C)}(\zeta_{11}, z) = -(1 + \zeta_{11}^2 + \zeta_{11}^4 + \zeta_{11}^5 + \zeta_{11}^6 + \zeta_{11}^8 + \zeta_{11}^9) \frac{\eta(11z)^2}{\eta_{11,3}(z)},$$

$$\mathcal{K}_{11,2}^{(C)}(\zeta_{11}, z) = -(1 + \zeta_{11}^2 + \zeta_{11}^5 + \zeta_{11}^6 + \zeta_{11}^9) \frac{\eta(11z)^2 \eta_{11,1}(z)}{\eta_{11,4}(z) \eta_{11,5}(z)},$$

$$\mathcal{K}_{11,3}^{(C)}(\zeta_{11}, z) = -(1 + \zeta_{11}^4 + \zeta_{11}^7) \frac{\eta(11z)^2}{\eta_{11,4}(z)},$$

$$\mathcal{K}_{11,4}^{(C)}(\zeta_{11}, z) = -(\zeta_{11}^3 + \zeta_{11}^8) \frac{\eta(11z)^2}{\eta_{11,5}(z)},$$

$$\mathcal{K}_{11,5}^{(C)}(\zeta_{11}, z) = \frac{\eta(11z)^2}{\eta_{11,1}(z)},$$

$$\mathcal{K}_{11,6}^{(C)}(\zeta_{11}, z) = -(2 + \zeta_{11}^2 + \zeta_{11}^3 + \zeta_{11}^4 + \zeta_{11}^5 + \zeta_{11}^6 + \zeta_{11}^7 + \zeta_{11}^8 + \zeta_{11}^9) \frac{\eta(11z)^2 \eta_{11,5}(z)}{\eta_{11,2}(z) \eta_{11,3}(z)},$$

$$\mathcal{K}_{11,7}^{(C)}(\zeta_{11}, z) = (\zeta_{11}^2 + \zeta_{11}^9) \frac{\eta(11z)^2 \eta_{11,3}(z)}{\eta_{11,1}(z) \eta_{11,4}(z)},$$

$$\mathcal{K}_{11,8}^{(C)}(\zeta_{11}, z) = (1 + \zeta_{11}^3 + \zeta_{11}^8) \frac{\eta(11z)^2 \eta_{11,2}(z)}{\eta_{11,1}(z) \eta_{11,3}(z)},$$

$$\mathcal{K}_{11,9}^{(C)}(\zeta_{11}, z) = (1 + \zeta_{11}^2 + \zeta_{11}^4 + \zeta_{11}^7 + \zeta_{11}^9) \frac{\eta(11z)^2}{\eta_{11,2}(z)},$$

$$\mathcal{K}_{11,10}^{(C)}(\zeta_{11}, z) = -(\zeta_{11}^2 + \zeta_{11}^4 + \zeta_{11}^7 + \zeta_{11}^9) \frac{\eta(11z)^2 \eta_{11,4}(z)}{\eta_{11,2}(z) \eta_{11,5}(z)}.$$

We consider the $m = 7$ element of the dissection

$$\mathcal{K}_{11,7}^{(C)}(\zeta_{11}, z) = (\zeta_{11}^2 + \zeta_{11}^9) \frac{\eta(11z)^2 \eta_{11,3}(z)}{\eta_{11,1}(z) \eta_{11,4}(z)},$$

and find $\mathcal{K}_{11,7}^{(C)}(\zeta_{11}, z)|[A]_1$ for $A = \begin{pmatrix} 3 & 1 \\ 11 & 4 \end{pmatrix} \in \Gamma_0(11)$.

Using the transformations for $\eta(z)$ and the theta function $f_{N,\rho}(z)$ (Theorems 1.4 and 1.5) and (1.13) and (1.14), we get

$$\begin{aligned} \mathcal{K}_{11,7}^{(C)}(\zeta_{11}, z)|[A]_1 &= (\zeta_{11}^2 + \zeta_{11}^9) \frac{\eta(11z)^3 f_{11,3}(z)}{f_{11,1}(z) f_{11,4}(z)}|[A]_1 \\ &= (\zeta_{11}^2 + \zeta_{11}^9) \frac{\eta(11z)^3 f_{11,9}(z)}{f_{11,3}(z) f_{11,12}(z)} \frac{e^{\frac{27\pi i}{11}}}{e^{\frac{3\pi i}{11}} \cdot e^{\frac{48\pi i}{11}}} \\ &= -\frac{\sin(\frac{\pi}{11})}{\sin(\frac{4\pi}{11})} \exp\left(\frac{9\pi i}{11}\right) (1 + \zeta_{11}^3 + \zeta_{11}^8) \frac{\eta(11z)^2 \eta_{11,2}(z)}{\eta_{11,3}(z) \eta_{11,1}(z)} \\ &= -\frac{\sin(\frac{\pi}{11})}{\sin(\frac{4\pi}{11})} \exp\left(\frac{9\pi i}{11}\right) \mathcal{K}_{11,8}(\zeta_{11}^4, z), \end{aligned}$$

which agrees with our symmetry result. We also make note of the fact that $m = 7$ and 8 are both quadratic non-residues modulo 11 .

Similarly, the $m = 0$ element of the dissection is also of special interest in regards to the fact that there also lies beautiful symmetry among the zeta-coefficients in the identity for $\mathcal{K}_{p,0}^{(C)}(\zeta_p, z)$ expressed in terms of generalized eta-functions.

The following is an identity for $\mathcal{K}_{13,0}^{(C)}(\zeta_{13}, z)$ in terms of generalized eta-functions :

$$(1.20) \quad \mathcal{K}_{13,0}^{(C)}(\zeta_{13}, z) = (q^{13}; q^{13})_{\infty} \sum_{n=1}^{\infty} \left(\sum_{k=0}^{12} M(k, 13, 13n-7) \zeta_{13}^k \right) q^n$$

$$= \sum_{r=1}^6 c_{13,r} j(13, \pi_r(\vec{n}_1), z),$$

where

$$\vec{n}_1 = (15, -1, -3, -2, -2, -2, -3),$$

and the coefficients are :

$$\begin{aligned} c_{13,6} &= 2 \zeta_{13}^{11} + \zeta_{13}^9 + \zeta_{13}^8 + \zeta_{13}^7 + \zeta_{13}^6 + \zeta_{13}^5 + \zeta_{13}^4 + 2 \zeta_{13}^2 + 2, \\ c_{13,1} &= -(\zeta_{13}^{11} + \zeta_{13}^{10} + \zeta_{13}^7 + \zeta_{13}^6 + \zeta_{13}^3 + \zeta_{13}^2), \\ c_{13,5} &= \zeta_{13}^{11} + \zeta_{13}^{10} + \zeta_{13}^9 + \zeta_{13}^8 + \zeta_{13}^5 + \zeta_{13}^4 + \zeta_{13}^3 + \zeta_{13}^2 + 1, \\ c_{13,2} &= \zeta_{13}^{10} - \zeta_{13}^9 + \zeta_{13}^8 + \zeta_{13}^5 - \zeta_{13}^4 + \zeta_{13}^3 + 1, \\ c_{13,4} &= -(\zeta_{13}^{10} + \zeta_{13}^9 + \zeta_{13}^7 + \zeta_{13}^6 + \zeta_{13}^4 + \zeta_{13}^3 + 2), \\ c_{13,3} &= -(\zeta_{13}^8 + \zeta_{13}^5). \end{aligned}$$

We provide a proof of this identity in the last section, but meanwhile we reveal an underlying hidden symmetry among the coefficients in the above identity. Rewriting them, we find that

$$\begin{aligned} c_{13,1} &= 2 + 2(\zeta_{13}^2 + \zeta_{13}^{11}) + (\zeta_{13}^4 + \zeta_{13}^9) + (\zeta_{13}^5 + \zeta_{13}^8) + (\zeta_{13}^6 + \zeta_{13}^7), \\ c_{13,2} &= \frac{\sin(\pi/13)}{\sin(2\pi/13)} (2 + 2(\zeta_{13}^4 + \zeta_{13}^9) + (\zeta_{13}^8 + \zeta_{13}^5) + (\zeta_{13}^{10} + \zeta_{13}^3) + (\zeta_{13}^{12} + \zeta_{13})), \\ c_{13,3} &= -\frac{\sin(\pi/13)}{\sin(3\pi/13)} (2 + 2(\zeta_{13}^6 + \zeta_{13}^7) + (\zeta_{13}^{12} + \zeta_{13}) + (\zeta_{13}^2 + \zeta_{13}^{11}) + (\zeta_{13}^5 + \zeta_{13}^8)), \\ c_{13,4} &= -\frac{\sin(\pi/13)}{\sin(4\pi/13)} (2 + 2(\zeta_{13}^8 + \zeta_{13}^5) + (\zeta_{13}^3 + \zeta_{13}^{10}) + (\zeta_{13}^7 + \zeta_{13}^6) + (\zeta_{13}^{11} + \zeta_{13}^2)), \\ c_{13,5} &= \frac{\sin(\pi/13)}{\sin(5\pi/13)} (2 + 2(\zeta_{13}^{10} + \zeta_{13}^3) + (\zeta_{13}^7 + \zeta_{13}^6) + (\zeta_{13}^{12} + \zeta_{13}) + (\zeta_{13}^4 + \zeta_{13}^9)), \\ c_{13,6} &= \frac{\sin(\pi/13)}{\sin(6\pi/13)} (2 + 2(\zeta_{13}^{12} + \zeta_{13}^1) + (\zeta_{13}^{11} + \zeta_{13}^2) + (\zeta_{13}^4 + \zeta_{13}^9) + (\zeta_{13}^{10} + \zeta_{13}^3)). \end{aligned}$$

Hence our other major symmetry result :

Theorem 1.13. *Let $p > 3$ be prime and the t vectors \vec{n}_ℓ , $1 \leq \ell \leq t$ and $j(z)$ be defined as in Definition 1.3. Suppose*

$$\mathcal{K}_{p,0}^{(C)}(\zeta_p, z) = \sum_{\ell=1}^t \sum_{r=1}^{\frac{1}{2}(p-1)} c_{p,r,\ell}(\zeta_p) j(p, \pi_r(\vec{n}_\ell), z),$$

where for $1 \leq r \leq \frac{p-1}{2}$, π_r is permutation on $\{1, 2, \dots, \frac{1}{2}(p-1)\}$ defined as $\pi_r(i) = i'$ where $ri' \equiv \pm i \pmod{p}$ and the functions $j(p, \pi_r(\vec{n}_\ell), z)$ are linearly independent (over \mathbb{Q}). Then

$$c_{p,r,\ell}(\zeta_p) = \frac{\sin(\pi/p)}{\sin(r\pi/p)} w(r, p) c_{p,1,\ell}(\zeta_p^r),$$

where $w(r, p) = \pm 1$.

The paper is organized as follows. In Section 2 we build the framework of our transformation results and express the crank generating function in terms of eta functions and determine its modularity under special congruence subgroups of $SL_2(\mathbb{Z})$. We also state the conditions for modularity of the eta-quotients which are the building blocks in the expressions for $\mathcal{K}_{p,m}^{(C)}(\zeta_p, z)$ in the last section. In Section 3, we state and illustrate our two main results on the symmetry of crank function. Section 4 is devoted to calculating lower bounds for the orders of $\mathcal{K}_{p,m}^{(C)}(\zeta_p, z)$ at the cusps of $\Gamma_1(p)$ which we utilize to prove identities in the subsequent section. Finally, in Section 5, we give a new proof of the crank theorem and also state identities for $\mathcal{K}_{p,m}^{(C)}(\zeta_p, z)$ when $p = 13, 17$ and 19 in terms of generalized eta-functions, proving two such identities mod 13 using the Valence formula, the modularity conditions and the orders at cusps determined in the previous sections.

2. PRELIMINARY DEFINITIONS AND RESULTS

In Section 1, we defined $M(r, t, n)$ to be the number of partitions of n with crank congruent to $r \pmod{t}$, and let $\zeta_p = \exp(2\pi i/p)$. Then we have

$$C(\zeta_p, q) = \sum_{n=0}^{\infty} \left(\sum_{k=0}^{p-1} M(k, p, n) \zeta_p^k \right) q^n.$$

We introduce a class of generalized eta products and its transformation under matrices in $SL_2(\mathbb{Z})$ due to Yifan Yang. We later show how it relates to the crank function.

Theorem 2.1 (Yang [23]). *Let p be prime and g, h be arbitrary real numbers not simultaneously congruent to 0 modulo p . For $z \in \mathbb{H}$, we define the generalized Dedekind eta functions $E_{g,h}(z)$ by*

$$E_{g,h}(z) = q^{B(g/p)/2} \prod_{m=1}^{\infty} \left(1 - \zeta_p^h q^{m-1+\frac{g}{p}} \right) \left(1 - \zeta_p^{-h} q^{m-\frac{g}{p}} \right),$$

where $B(x) = x^2 - x + 1/6$. Then the following transformations hold :

- i) $E_{g+p,h} = E_{-g,-h} = -\zeta_p^{-h} E_{g,h}$.
- ii) $E_{g,h+p} = E_{g,h}$.

Moreover for $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{Z})$, we have

- iii) $E_{g,h}(z+b) = e^{\pi i b B(g/p)} E_{g,bg+h}(z)$ for $c = 0$.

iv) $E_{g,h}(\gamma z) = \varepsilon(a, b, c, d)e^{\pi i \delta} E_{g',h'}(z)$ for $c \neq 0$ where

$$\varepsilon(a, b, c, d) = \begin{cases} e^{\pi i (bd(1-c^2) + c(a+d-3))/6} & \text{if } c \text{ is odd,} \\ -ie^{\pi i (ac(1-d^2) + d(b-c+3))/6} & \text{if } d \text{ is odd,} \end{cases}$$

$$\delta = \frac{g^2 ab + 2ghbc + h^2 cd}{p^2} - \frac{gb + h(d-1)}{p}, \text{ and}$$

$$(g' \ h') = (g \ h) \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

2.1. Modularity and transformation results.

Theorem 2.2. *Let $p > 3$ be prime, suppose $1 \leq \ell \leq (p-1)$, and define*

$$\mathcal{F}_1 \left(\frac{\ell}{p}; z \right) := \frac{\eta(z)^2}{E_{0,\ell}(z)},$$

where, following Yang [23], we consider the generalized eta product

$$E_{0,\ell}(z) = q^{\frac{1}{12}} \prod_{m=1}^{\infty} (1 - \zeta_p^\ell q^{m-1})(1 - \zeta_p^{-\ell} q^m).$$

Then

$$\mathcal{F}_1 \left(\frac{\ell}{p}; z \right) | [A]_1 = \mu(A, \ell) \mathcal{F}_1 \left(\frac{\bar{d}\ell}{p}; z \right),$$

where

$$\mu(A, \ell) = \exp \left(\pi i \left(\frac{\ell(d-1)}{p} + \frac{cd\ell^2}{p^2} - \frac{c\ell}{p} \right) \right), \text{ and } A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(p).$$

Here \bar{m} is the least nonnegative residue of $m \pmod{p}$.

Proof. Using Theorems 2.1 and 1.4, we have

$$\mathcal{F}_1 \left(\frac{\ell}{p}; z \right) | [A]_1 = \begin{cases} \frac{e^{\frac{2\pi i b}{12}} \eta(z)^2}{e^{\frac{\pi i b}{6}} E_{0,\ell}(z)} & \text{if } c = 0, \\ \frac{e^{\frac{\pi i}{6} [(a+d)c - bd(c^2-1) - 3c]} \eta(z)^2}{e^{\frac{\pi i}{6} [bd(1-c^2) + c(a+d-3)]} e^{\pi i \left[\frac{\ell^2 cd}{p^2} - \frac{\ell(d-1)}{p} \right]} E_{\ell c, \ell d}(z)} & \text{if } c \neq 0 \text{ and odd,} \\ \frac{e^{\frac{\pi i}{6} [(a+d)c - bd(c^2-1) - 3c]} \eta(z)^2}{-ie^{\frac{\pi i}{6} [bd(1-c^2) + c(a+d-3)]} e^{\pi i \left[\frac{\ell^2 cd}{p^2} - \frac{\ell(d-1)}{p} \right]} E_{\ell c, \ell d}(z)} & \text{if } c \neq 0 \text{ and even,} \end{cases}$$

$$= e^{\pi i \left[\frac{\ell(d-1)}{p} - \frac{\ell^2 cd}{p^2} \right]} \frac{\eta(z)^2}{E_{\ell c, \ell d}(z)}.$$

Now, using Theorem 2.1 we have $E_{\ell c, \ell d}(z) = (-\zeta_p^{-\ell d})^{\frac{\ell c}{p}} E_{0, \ell d}(z) = e^{\pi i \left[\frac{c\ell}{p} - \frac{2\ell^2 cd}{p^2} \right]} E_{0, \overline{d\ell}}(z)$.

Then,

$$\mathcal{F}_1 \left(\frac{\ell}{p}; z \right) | [A]_1 = e^{\pi i \left[\frac{\ell(d-1)}{p} + \frac{cd\ell^2}{p^2} - \frac{c\ell}{p} \right]} \frac{\eta(z)^2}{E_{0, \overline{d\ell}}(z)}$$

and we have our result. □

Corollary 2.3. *Let $p > 3$ be prime and suppose $1 \leq \ell \leq \frac{1}{2}(p-1)$. Then*

$$\mathcal{F}_1 \left(\frac{\ell}{p}; z \right) | [A]_1 = \mathcal{F}_1 \left(\frac{\ell}{p}; z \right),$$

where $A \in \Gamma_0(p^2) \cap \Gamma_1(p)$.

Proof. Let $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(p^2) \cap \Gamma_1(p)$. Then $c \equiv 0 \pmod{p^2}$ and $a \equiv d \equiv 1 \pmod{p}$.

So

- i) $p\ell(d-1) + cd\ell^2 - c\ell p \equiv 0 \pmod{p^2}$,
- ii) $p\ell(d-1) + cd\ell^2 - c\ell p \equiv d\ell + \ell + cd\ell + c\ell \equiv \ell(c+1)(d+1) \equiv 0 \pmod{2}$.

Therefore,

$$\mu(A, \ell) = \exp \left(\pi i \left(\frac{\ell(d-1)}{p} + \frac{cd\ell^2}{p^2} - \frac{c\ell}{p} \right) \right) = 1$$

and we have our result. □

Let $\mathcal{F}_1^* \left(\frac{\ell}{p}; z \right) = \frac{\eta(p^2 z)}{\eta(z)} \mathcal{F}_1 \left(\frac{\ell}{p}; z \right)$. The generating function of the crank due to Andrews and Garvan is given by

$$C(z, q) = \sum_{n=0}^{\infty} \sum_m M(m, n) z^m q^n = \prod_{n=1}^{\infty} \frac{1 - q^n}{(1 - zq^n)(1 - z^{-1}q^n)}.$$

Replacing z by ζ_p^ℓ in the above expression and multiplying both sides by $\eta(p^2 z)$, we can express the crank generating function in terms of our eta functions as

$$(2.1) \quad \eta(p^2 z) \mathcal{C}_p(z) = (1 - \zeta_p^\ell) \mathcal{F}_1^* \left(\frac{\ell}{p}; z \right), \text{ where } \mathcal{C}_p(z) = q^{\frac{-1}{24}} C(\zeta_p^\ell, q).$$

This leads to our first main result stated in the introduction section. We restate it here.

Theorem 2.4. *Let $p > 3$ be prime. Then the function*

$$\eta(p^2 z) \mathcal{C}_p(z)$$

is a weakly holomorphic modular form of weight 1 on the group $\Gamma_0(p^2) \cap \Gamma_1(p)$.

Proof. In the light of the above expression, the proof of the theorem is equivalent to showing that $\mathcal{F}_1^* \left(\frac{\ell}{p}; z \right)$ is a weakly holomorphic modular form of weight 1 on the group $\Gamma_0(p^2) \cap \Gamma_1(p)$. The transformation condition holds using Corollary 2.3 and the well-known result that $\frac{\eta(p^2 z)}{\eta(z)}$ is a modular function on $\Gamma_0(p^2)$ when $p > 3$ is prime. That is

$$\mathcal{F}_1^* \left(\frac{\ell}{p}; z \right) | [A]_1 = \mathcal{F}_1^* \left(\frac{\ell}{p}; z \right)$$

for $A \in \Gamma_0(p^2) \cap \Gamma_1(p)$.

We next check the cusp conditions and show that $\mathcal{F}_1^* \left(\frac{\ell}{p}; z \right) | [A]_1(z)$ when expanded as a series in $q_{p^2} = \exp\left(\frac{2\pi iz}{p^2}\right)$ has only finitely many terms with negative exponents for $A \in \text{SL}_2(\mathbb{Z})$. By Theorems 2.1 and 1.4, we have

$$\mathcal{F}_1^* \left(\frac{\ell}{p}; z \right) | [A]_1 = \frac{\eta(p^2 z)}{\eta(z)} \frac{\eta(z)^2}{E_{0,\ell}(z)} | [A]_1 = \frac{\eta(p^2 z)}{\eta(z)} | [A]_1 \cdot \varepsilon_A \frac{\eta(z)^2}{E_{\ell c, \ell d}(z)},$$

where ε_A is a root of unity.

Using the fact that $\frac{\eta(p^2 z)}{\eta(z)}$ is a modular function on $\Gamma_0(p^2)$ and the definition of $\eta(z)$ and $E_{g,h}(z)$, we find that the series has only finitely many terms with negative exponents. □

Next, we define the (weight k) Atkin U_p operator by

$$(2.2) \quad F | [U_p]_k := \frac{1}{p} \sum_{r=0}^{p-1} F \left(\frac{z+r}{p} \right) = p^{\frac{k}{2}-1} \sum_{n=0}^{p-1} F | [T_r]_k,$$

where

$$T_r = \begin{pmatrix} 1 & r \\ 0 & p \end{pmatrix},$$

and the more general $U_{p,m}$ is defined as

$$(2.3) \quad F | [U_{p,m}]_k := \frac{1}{p} \sum_{r=0}^{p-1} \exp\left(-\frac{2\pi i r m}{p}\right) F \left(\frac{z+r}{p} \right) = p^{\frac{k}{2}-1} \sum_{r=0}^{p-1} \exp\left(-\frac{2\pi i r m}{p}\right) F | [T_r]_k.$$

We note that $U_p = U_{p,0}$. In addition, if

$$F(z) = \sum_n a(n) q^n = \sum_n a(n) \exp(2\pi i z n),$$

then

$$F | [U_{p,m}]_k = q^{m/p} \sum_n a(pn + m) q^n = \exp(2\pi i m z / p) \sum_n a(pn + m) \exp(2\pi i n z).$$

Combining Equation (2.1) and Definition 1.9, we have

Proposition 2.5. For $p > 3$ be a prime and $0 \leq m \leq p - 1$ we have

$$(2.4) \quad \mathcal{K}_{p,m}^{(C)}(\zeta_p^\ell, z) = (1 - \zeta_p^\ell) \mathcal{F}_1^* \left(\frac{\ell}{p}; z \right) \mid [U_{p,m}]_1$$

We are now ready to prove the modularity result of the crank function stated in the introduction section. We restate the theorem here and note that the proof utilizes the same matrix transformations as it did for the rank dissection elements $\mathcal{K}_{p,m}^{(R)}(z)$ [15, Theorem 6.3, p.234]. We include the proof here for the sake of completion.

Theorem 2.6. Let $p > 3$ be prime and suppose $0 \leq m \leq p - 1$. Then

- (i) $\mathcal{K}_{p,0}^{(C)}(z)$ is a weakly holomorphic modular form of weight 1 on $\Gamma_1(p)$.
- (ii) If $1 \leq m \leq (p - 1)$ then $\mathcal{K}_{p,m}^{(C)}(z)$ is a weakly holomorphic modular form of weight 1 on $\Gamma(p)$. In particular,

$$\mathcal{K}_{p,m}^{(C)}(z) \mid [A]_1 = \exp \left(\frac{2\pi ibm}{p} \right) \mathcal{K}_{p,m}^{(C)}(z),$$

$$\text{for } A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_1(p).$$

Proof. We let

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_1(p)$$

so that $a \equiv d \equiv 1 \pmod{p}$ and $c \equiv 0 \pmod{p}$. Let $0 \leq k \leq p - 1$. We take $k' \equiv b + k \pmod{p}$ so that

$$T_k A = B_k T_{k'},$$

and

$$B_k = T_k A T_{k'}^{-1} = \begin{pmatrix} a + ck & \frac{1}{p}(-k'(a + kc) + b + kd) \\ pc & d - k'c \end{pmatrix} \in \Gamma_0(p^2) \cap \Gamma_1(p).$$

Thus,

$$\begin{aligned} & \mathcal{K}_{p,m}^{(C)}(z) \mid [A]_1 \\ &= \frac{1}{\sqrt{p}} (1 - \zeta_p) \sum_{k=0}^{p-1} \exp \left(-\frac{2\pi ikm}{p} \right) \mathcal{F}_1^* \left(\frac{1}{p}; z \right) \mid [(T_k A)]_1 \\ &= \frac{1}{\sqrt{p}} (1 - \zeta_p) \sum_{k=0}^{p-1} \exp \left(-\frac{2\pi i(k' - b)m}{p} \right) \mathcal{F}_1^* \left(\frac{1}{p}; z \right) \mid [(B_k T_{k'})]_1 \\ &= \frac{1}{\sqrt{p}} (1 - \zeta_p) \exp \left(\frac{2\pi ibm}{p} \right) \sum_{k'=0}^{p-1} \exp \left(-\frac{2\pi ik'm}{p} \right) \mathcal{F}_1^* \left(\frac{1}{p}; z \right) \mid [T_{k'}]_1 \end{aligned}$$

(since $\frac{\eta(p^2 z)}{\eta(z)}$ is a modular function on $\Gamma_0(p^2)$ and Corollary 2.3 since $B_k \in \Gamma_0(p^2) \cap \Gamma_1(p)$)

$$= \exp \left(\frac{2\pi ibm}{p} \right) \mathcal{K}_{p,m}^{(C)}(z),$$

as required. Thus each function $\mathcal{K}_{p,m}^{(C)}(z)$ has the desired transformation property. It is clear that each $\mathcal{K}_{p,m}^{(C)}(z)$ is holomorphic on \mathfrak{h} . The cusp conditions follow by a standard argument. We examine orders at each cusp in more detail in a later section. \square

2.2. Modularity conditions for eta-quotients. In [16], we derived conditions for an eta-quotient $j(p, \vec{n}, z)$ defined in 1.3 to be a weakly holomorphic modular form of weight 1 on $\Gamma(p)$. In later sections, these are used in the proof of one our symmetry results and to derive and prove identities for $\mathcal{K}_{p,m}(\zeta_p, z)$, where $p = 13, 17$ and 19 and $0 \leq m \leq p - 1$.

Theorem 2.7 ([16], Theorem 3.1). *Let p, \vec{n} and $j(p, \vec{n}, z)$ be as defined in 1.3. Then $j(p, \vec{n}, z)$ is a weakly holomorphic modular form of weight 1 on $\Gamma(p)$ satisfying the modularity condition $j(p, \vec{n}, z) | [A]_1 = \exp(\frac{2\pi i b m}{p}) j(p, \vec{n}, z)$ for $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_1(p)$ provided the following conditions are met :*

$$(1) \quad n_0 + \sum_{k=1}^{\frac{p-1}{2}} n_k = 2,$$

$$(2) \quad n_0 + 3 \sum_{k=1}^{\frac{p-1}{2}} n_k \equiv 0 \pmod{24},$$

$$(3) \quad \sum_{k=1}^{\frac{p-1}{2}} k^2 n_k \equiv 2m \pmod{p}.$$

3. TWO NEW SYMMETRY RESULTS ANALOGOUS TO THE RANK

Theorem 3.1. *Suppose $p > 3$ prime, $0 \leq m \leq p - 1$, and $1 \leq d \leq p - 1$. Then*

$$(3.1) \quad \mathcal{K}_{p,m}^{(C)}(\zeta_p, z) | [A]_1 = \frac{\sin(\pi/p)}{\sin(d\pi/p)} (-1)^{d+1} \zeta_p^{mak} \mathcal{K}_{p,ma^2}^{(C)}(\zeta_p^d, z),$$

assuming $1 \leq a, d \leq (p - 1)$ and

$$A = \begin{pmatrix} a & k \\ p & d \end{pmatrix} \in \Gamma_0(p).$$

Proof. We undergo the same matrix transformations as we did in the case of $\mathcal{K}_{p,m}^{(R)}(\zeta_p, z)$ ([16], Proposition 4.7), and for $0 \leq r \leq p - 1$ let $T_r = \begin{pmatrix} 1 & r \\ 0 & p \end{pmatrix}$, and $B_r = \begin{pmatrix} a + pr & \frac{1}{p}(k + rd - r'(a + pr)) \\ p^2 & d - r'p \end{pmatrix}$, where $0 \leq r' \leq p - 1$ is chosen so that $r' \equiv rd^2 + dk \pmod{p}$. Then

$$T_r A = B_r T_{r'}, \quad r \equiv r'a^2 - ak \pmod{p}, \quad \text{and} \quad B_r \in \Gamma_0(p^2).$$

Using the above matrix transformation and applying the transformation for $\mathcal{F}_1\left(\frac{1}{p}; z\right)$ (Theorem 2.2) under matrices in $\Gamma_0(p)$ and the well-known result that $\frac{\eta(p^2z)}{\eta(z)}$ is a modular function on $\Gamma_0(p^2)$ when $p > 3$ is prime, we arrive at

$$\begin{aligned}
\mathcal{K}_{p,m}^{(C)}(\zeta_p, z) \mid [A]_1 &= (1 - \zeta_p) \mathcal{F}_1^* \left(\frac{1}{p}; z \right) \mid [U_{p,m}]_1 \mid [A]_1 \\
&= \frac{1 - \zeta_p}{\sqrt{p}} \sum_{r=0}^{p-1} \zeta_p^{-rm} \frac{\eta(p^2z)}{\eta(z)} \mathcal{F}_1 \left(\frac{1}{p}; z \right) \mid [T_r]_1 \mid [A]_1 \\
&= \frac{1 - \zeta_p}{\sqrt{p}} \sum_{r=0}^{p-1} \zeta_p^{-rm} \frac{\eta(p^2z)}{\eta(z)} \mathcal{F}_1 \left(\frac{1}{p}; z \right) \mid [B_r]_1 \mid [T_{r'}]_1 \\
&= \frac{1 - \zeta_p}{\sqrt{p}} \sum_{r=0}^{p-1} \zeta_p^{-rm} \frac{\eta(p^2z)}{\eta(z)} \mu(B_r, 1) \mathcal{F}_1 \left(\frac{d}{p}; z \right) \mid [T_{r'}]_1 \\
&= \frac{(1 - \zeta_p) \zeta_p^{\frac{d-1}{2}}}{\sqrt{p}} (-1)^d \zeta_p^{mak} \sum_{r'=0}^{p-1} \zeta_p^{-r'ma^2} \frac{\eta(p^2z)}{\eta(z)} \mathcal{F}_1 \left(\frac{d}{p}; z \right) \mid [T_{r'}]_1, \\
&= \frac{\sin(\pi/p)}{\sin(d\pi/p)} (-1)^d \zeta_p^{mak} \frac{(1 - \zeta_p^d)}{\sqrt{p}} \sum_{r'=0}^{p-1} \zeta_p^{-r'ma^2} \frac{\eta(p^2z)}{\eta(z)} \mathcal{F}_1 \left(\frac{d}{p}; z \right) \mid [T_{r'}]_1,
\end{aligned}$$

since

$$\begin{aligned}
\mu(B_r, 1) &= \exp \left(\pi i \left(\frac{(d - r'p - 1)}{p} + \frac{p^2(d - r'p)}{p^2} - p \right) \right) \\
&= (-1)^{d+1} \exp \left(\pi i \left(\frac{d-1}{p} \right) \right) (-1)^{-r'(p+1)}, \\
&= (-1)^{d+1} \zeta_p^{\frac{d-1}{2}},
\end{aligned}$$

$$\frac{(1 - \zeta_p) \zeta_p^{\frac{d-1}{2}}}{(1 - \zeta_p^d)} = \frac{\sin(\pi/p)}{\sin(d\pi/p)},$$

and

$$\zeta_p^{-rm} = \zeta_p^{m(-r'a^2+ak)} = \zeta_p^{mak} \zeta_p^{-mr'a^2},$$

and as r runs through a complete residue system mod p so does r' . The result follows. \square

Our other symmetry result is between the coefficients of the identity for the $\mathcal{K}_{p,0}^{(C)}(\zeta_p, z)$ element of the dissection expressed in terms of generalized eta functions $j(p, \vec{n}, z)$.

Let $\vec{n} = (n_0, n_1, n_2, \dots, n_{\frac{1}{2}(p-1)}) \in \mathbb{Z}^{\frac{1}{2}(p+1)}$. Then, for $1 \leq r \leq \frac{1}{2}(p-1)$, the action of the permutation $\pi_r : [\frac{1}{2}(p-1)] \rightarrow [\frac{1}{2}(p-1)]$, where $[\frac{1}{2}(p-1)] = \{1, 2, \dots, \frac{1}{2}(p-1)\}$

defined by $\pi_r(i) = i'$ where $ri' \equiv \pm i \pmod{p}$, on the generalized eta function is given by ([16], Lemma 4.12)

$$j(p, \pi_r(\vec{n}), z) = \eta(pz)^{n_0} \prod_{k=1}^{\frac{p-1}{2}} f_{p,rk}(z)^{n_k}.$$

Using Theorem 3.1 and the conditions for modularity of the eta quotients in Theorem 2.7, we determine the transformation of $\mathcal{K}_{p,0}^{(C)}(\zeta_p, z)$. The symmetry is exactly the same for the rank element $\mathcal{K}_{p,0}^{(R)}(\zeta_p, z)$. We give the statement of the result and omit the proof here (see [16], Theorem 5.1 for details).

Theorem 3.2. *Let $p > 3$ be prime. Suppose there are t vectors $\vec{n}_1, \vec{n}_2, \dots, \vec{n}_t \in \mathbb{Z}^{\frac{1}{2}(p+1)}$ such that the set of functions $j(p, \pi_r(\vec{n}_k), z), 1 \leq k \leq t, 1 \leq r \leq \frac{1}{2}(p-1)$ are linearly independent (over \mathbb{Q}) and*

$$\mathcal{K}_{p,0}^{(C)}(\zeta_p, z) = \sum_{k=1}^t \sum_{r=1}^{\frac{1}{2}(p-1)} c_{p,r,k}(\zeta_p) j(p, \pi_r(\vec{n}_k), z),$$

where $c_{p,r,k}(\zeta_p) \in \mathbb{Q}[\zeta_p]$. Then for $1 \leq d \leq \frac{p-1}{2}$, and $\vec{n} = (n_0, n_1, n_2, \dots, n_{\frac{p-1}{2}})$, we have

$$c_{p,d,k}(\zeta_p) = \frac{\sin(\pi/p)}{\sin(d\pi/p)} (-1)^{d+1+L(\vec{n},d)} c_{p,1,k}(\zeta_p^d)$$

where

$$L(\vec{n}, d) = L(\vec{n}, a, b, d, p) = bd(1+a) \sum_{k=1}^{\frac{p-1}{2}} kn_k + \sum_{k=1}^{\frac{p-1}{2}} \left(\left\lfloor \frac{dka}{p} \right\rfloor + \left\lfloor \frac{dk}{p} \right\rfloor \right) n_k,$$

and a, b are chosen so that $A = \begin{pmatrix} a & b \\ p & d \end{pmatrix} \in \Gamma_0(p)$.

4. LOWER BOUNDS FOR ORDER OF AT CUSPS

In this section, we calculate lower bounds for the orders of $\mathcal{K}_{p,m}^{(C)}(\zeta_p, z)$ at the cusps of $\Gamma_1(p)$, which we use in proving the $\mathcal{K}_{p,m}^{(C)}(\zeta_p, z)$ identities in the subsequent section.

We first establish a few other useful transformation results.

Let $p > 3$ be prime, $1 \leq \ell \leq (p-1)$, and define $\mathcal{F}_2\left(\frac{\ell}{p}; z\right) := \frac{\eta(z)^2}{E_{\ell,0}(z)}$, where $E_{\ell,0}(z) = q^{B(\ell/p)/2} \prod_{m=1}^{\infty} \left(1 - q^{m-1+\frac{\ell}{p}}\right) \left(1 - q^{m-\frac{\ell}{p}}\right)$.

Theorem 4.1. (1) $E_{0,\ell}(z) | [S]_1 = -i\zeta_{2p}^{\ell} E_{\ell,0}(z)$.

(2) $\mathcal{F}_2\left(\frac{\ell}{p}; z\right) | [S]_1 = \zeta_{2p}^{-\ell} \mathcal{F}_1\left(\frac{-\ell}{p}; z\right)$.

$$(3) E_{0,\ell}(z) | [T]_1 = e^{\pi i/6} E_{0,\ell}(z).$$

Proof. Proofs of (1) and (3) follow easily using the definition of $E_{g,h}(z)$. (2) follows from (1) and the transformation of $\eta(z)$ under S i.e. $\eta(z) | [S]_1 = \exp(\frac{-\pi i}{4}) \eta(z)$ (Theorem 1.4). \square

$$\text{Let } \mathfrak{Y}_p := \left\{ \mathcal{F}_1 \left(\frac{\ell}{p}; z \right), \mathcal{F}_2 \left(\frac{\ell}{p}; z \right) : 0 < a < p \right\}.$$

The action of $\text{SL}_2(\mathbb{Z})$ on each element can be given explicitly using Theorems 1.4 and 4.1. Also each $\mathcal{F} \in \mathfrak{Y}_p$ has a q -expansion

$$\mathcal{F}(z) = \sum_{m \geq m_0} a(m) \exp \left(2\pi i z \frac{m}{24p^2} \right),$$

where $a(m_0) \neq 0$. We define

$$\text{ord}(\mathcal{G}; \infty) := \frac{m_0}{24p^2}.$$

For any cusp $\frac{a}{c}$ with $(a, c) = 1$ we define

$$\text{ord} \left(\mathcal{F}; \frac{a}{c} \right) := \text{ord}(\mathcal{F} | [A]_1; \infty),$$

where $A \in \text{SL}_2(\mathbb{Z})$ and $A\infty = \frac{a}{c}$. We note that $\mathcal{F} | [A]_1 \in \mathfrak{Y}_p$. We also note that this definition coincides with the definition of invariant order at a cusp [8, p.2319], [4, p.275]. The order of each function $\mathcal{F}(z) \in \mathfrak{Y}_p$ is defined in the natural way i.e.

$$\text{ord} \left(\mathcal{F}; \frac{a}{c} \right) := 2 \text{ord} \left(\eta(z); \frac{a}{c} \right) + \text{ord} \left(1/E_{*,*}; \frac{a}{c} \right),$$

where $\text{ord} \left(\eta(z); \frac{a}{c} \right)$ is the usual invariant order of $\eta(z)$ at the cusp $\frac{a}{c}$ [20, p.34].

We determine $\text{ord}(\mathcal{F}; \infty)$ for each $\mathcal{F}(z) \in \mathfrak{Y}_p$. After some calculation we find

Proposition 4.2. *Let $p > 3$ be prime. Then*

$$\begin{aligned} \text{ord} \left(\mathcal{F}_1 \left(\frac{a}{p}; z \right); \infty \right) &= 0, \\ \text{ord} \left(\mathcal{F}_2 \left(\frac{a}{p}; z \right); \infty \right) &= \frac{a}{2p} - \frac{a^2}{2p^2}. \end{aligned}$$

We also need [20, Corollary 2.2].

Proposition 4.3. *Let $N \geq 1$ and let*

$$F(z) = \prod_{m|N} \eta(mz)^{r_m},$$

where each $r_m \in \mathbb{Z}$. Then for $(a, c) = 1$,

$$\text{ord} \left(F(z); \frac{a}{c} \right) = \sum_{m|N} \frac{(m, c)^2 r_m}{24m}.$$

From [9, Corollary 4, p.930] we have

Proposition 4.4. *Let $p > 3$ be prime. Then a set of inequivalent cusps \mathcal{S}_p for $\Gamma_1(p)$ is given by*

$$i\infty, 0, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{\frac{1}{2}(p-1)}, \frac{2}{p}, \frac{3}{p}, \dots, \frac{\frac{1}{2}(p-1)}{p}.$$

We next calculate lower bounds of the invariant order of $\mathcal{K}_{p,m}^{(C)}(\zeta_p, z)$ at each cusp of $\Gamma_1(p)$.

Theorem 4.5. *Let $p > 3$ be prime, and suppose $0 \leq m \leq p-1$. Then*

(i)

$$\text{ord} \left(\mathcal{K}_{p,m}^{(C)}(\zeta_p, z); 0 \right) \begin{cases} \geq 0 & \text{if } p = 11, \\ = -\frac{1}{24p}(p-1)(p-11) & \text{otherwise;} \end{cases}$$

(ii)

$$\text{ord} \left(\mathcal{K}_{p,m}^{(C)}(\zeta_p, z); \frac{1}{n} \right) \begin{cases} = -\frac{1}{2p} \left[\frac{(p-1)(p+1)}{12} + n(n-p) \right] & \text{if } 2 \leq n < \frac{p}{2} - \frac{1}{2} \sqrt{\frac{2p^2+1}{3}}, \\ \geq 0 & \text{otherwise;} \end{cases}$$

and

(iii)

$$\text{ord} \left(\mathcal{K}_{p,m}^{(C)}(\zeta_p, z); \frac{n}{p} \right) \geq \left(\frac{p^2-1}{24p} \right).$$

Proof. We derive lower bounds for $\text{ord} \left(\mathcal{K}_{p,m}^{(C)}(\zeta_p, z); \zeta \right)$ for each cusp ζ of $\Gamma_1(p)$ not equivalent to $i\infty$.

We have

$$\mathcal{K}_{p,m}^{(C)}(\zeta_p, z) := q^{\frac{1}{24}}(1-\zeta_p) \mathcal{F}_1^* \left(\frac{1}{p}; z \right) \mid [U_{p,m}]_1 = \frac{1}{\sqrt{p}}(1-\zeta_p) \sum_{k=0}^{p-1} \zeta_p^{-km} \mathcal{F}_1^* \left(\frac{1}{p}; z \right) \mid [T_k]_1$$

where

$$(4.1) \quad \mathcal{F}_1^* \left(\frac{1}{p}; z \right) = \frac{\eta(p^2z)}{\eta(z)} \mathcal{F}_1 \left(\frac{1}{p}; z \right).$$

We calculate

$$\mathcal{F}_1^* \left(\frac{1}{p}; z \right) \mid [T_k A]_1,$$

for each $0 \leq k \leq p-1$ and each $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z})$.

Case 1. $a + kc \not\equiv 0 \pmod{p}$. Choose $0 \leq k' \leq p-1$ such that

$$(a + kc)k' \equiv (b + kd) \pmod{p}.$$

Then

$$T_k A = C_k T_{k'},$$

where

$$C_k = T_k A T_{k'}^{-1} = \begin{pmatrix} a + ck & \frac{1}{p}(-k'(a + kc) + b + kd) \\ pc & d - k'c \end{pmatrix} \in \Gamma_0(p).$$

From Theorem 2.2 we have

$$(4.2) \quad \begin{aligned} \mathcal{F}_1^* \left(\frac{1}{p}; z \right) | [T_k A]_1 &= \mathcal{F}_1^* \left(\frac{1}{p}; z \right) | [C_k T_{k'}]_1 \\ &= (F_p(z) | [C_k T_{k'}]_0) \left(\mu(C_k, 1) \mathcal{F}_1 \left(\frac{\overline{d - k'c}}{p}; z \right) | [T_{k'}]_1 \right), \end{aligned}$$

where

$$(4.3) \quad F_p(z) = \frac{\eta(p^2 z)}{\eta(z)}.$$

Case 2. $a + kc \equiv 0 \pmod{p}$. In this case we find that

$$T_k A = D_k P,$$

where

$$P = \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix},$$

$$D_k = \begin{pmatrix} \frac{1}{p}(a + kc) & b + kd \\ c & pd \end{pmatrix} \in \mathbf{SL}_2(\mathbb{Z}),$$

and

$$E_k = D_k S = \begin{pmatrix} b + kd & \frac{-1}{p}(a + kc) \\ pd & -c \end{pmatrix} \in \Gamma_0(p).$$

From Theorem 2.2 we have

$$\mathcal{F}_1 \left(\frac{1}{p}; z \right) | [E_k]_1 = \mu(E_k, 1) \mathcal{F}_1 \left(\frac{\overline{-c}}{p}; z \right).$$

By Theorem 4.1(2) we have

$$\mathcal{F}_1 \left(\frac{1}{p}; z \right) | [D_k]_1 = \mu(E_k, 1) \mathcal{F}_1 \left(\frac{\overline{-c}}{p}; z \right) | [S^{-1}]_1 = \mu(E_k, 1) \zeta_{2p} \mathcal{F}_2 \left(\frac{\overline{c}}{p}; z \right),$$

and

$$\mathcal{F}_1 \left(\frac{1}{p}; z \right) | [T_k A]_1 = \mu(E_k, 1) \zeta_{2p} \mathcal{F}_2 \left(\frac{\overline{c}}{p}; z \right) | [P]_1,$$

so that

$$(4.4) \quad \mathcal{F}_1^* \left(\frac{1}{p}; z \right) | [T_k A]_1 = (F_p(z) | [D_k P]_0) \left(\mu(E_k, 1) \zeta_{2p} \mathcal{F}_2 \left(\frac{\overline{c}}{p}; z \right) | [P]_1 \right).$$

Now we are ready to exam each cusp ζ of $\Gamma_1(p)$. We choose

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathbf{SL}_2(\mathbb{Z}), \quad \text{so that } A(\infty) = \frac{a}{c} = \zeta.$$

(i) $\zeta = 0$. Here $a = 0, c = 1$ and we assume $0 \leq k \leq p-1$. If $k \neq 0$ then applying (4.2) we have

$$\begin{aligned} \text{ord} \left(\mathcal{F}_1^* \left(\frac{1}{p}; \frac{z+k}{p} \right); 0 \right) &= \frac{1}{p} \text{ord} \left(F_p(z); \frac{k}{p} \right) + \frac{1}{p} \text{ord} \left(\mathcal{F}_1 \left(\frac{\overline{d-k'c}}{p}; z \right); i\infty \right) \\ &= 0 + 0 = 0, \end{aligned}$$

by Propositions 4.3 and 4.2. Now applying (4.4) with $k = 0$ we have

$$\begin{aligned} \text{ord} \left(\mathcal{F}_1^* \left(\frac{1}{p}; \frac{z}{p} \right); 0 \right) &= p \text{ord} \left(F_p(z); 0 \right) + p \text{ord} \left(\mathcal{F}_2 \left(\frac{\bar{c}}{p}; z \right); i\infty \right) \\ &= -\frac{1}{24p}(p^2 - 1) + p \left(\frac{1}{2p} - \frac{1}{2p^2} \right) \\ &= -\frac{1}{24p}(p-1)(p-11) \end{aligned}$$

again by Propositions 4.3 and 4.2. The result (i) follows since

$$\text{ord} \left(\mathcal{K}_{p,m}^{(C)}(z); 0 \right) \geq \min_{0 \leq k \leq p-1} \text{ord} \left(\mathcal{F}_1^* \left(\frac{1}{p}; \frac{z+k}{p} \right); 0 \right).$$

(ii) $\zeta = \frac{1}{n}$, where $2 \leq n \leq \frac{1}{2}(p-1)$. Let $A = \begin{pmatrix} 1 & 0 \\ n & 1 \end{pmatrix}$ so that $A(\infty) = 1/n$. If $kn \not\equiv -1 \pmod{p}$ we apply (4.2) with $C_k = \begin{pmatrix} 1+kn & * \\ pn & 1-k'n \end{pmatrix}$, and find that

$$\begin{aligned} \text{ord}_{\text{holo}} \left(\mathcal{F}_1^* \left(\frac{1}{p}; \frac{z+k}{p} \right); \frac{1}{n} \right) &= \frac{1}{p} \text{ord} \left(F_p(z); \frac{1+kn}{pn} \right) + \frac{1}{p} \text{ord} \left(\mathcal{F}_1 \left(\frac{\overline{1-kn}}{p}; z \right); i\infty \right) \\ &= 0 + 0 = 0. \end{aligned}$$

Now we assume $kn \equiv -1 \pmod{p}$ and we will apply (4.4). We have

$$\begin{aligned} \text{ord}_{\text{holo}} \left(\mathcal{F}_1^* \left(\frac{1}{p}; \frac{z+k}{p} \right); \frac{1}{n} \right) &= p \text{ord} \left(F_p(z); \frac{(1+kn)/p}{n} \right) + p \text{ord} \left(\mathcal{F}_2 \left(\frac{n}{p}; z \right); i\infty \right) \\ &= -\frac{1}{24p}(p^2 - 1) + p \left(\frac{n\ell}{2p} - \frac{n^2\ell^2}{2p^2} \right) \\ &\begin{cases} = -\frac{1}{2p} \left[\frac{(p-1)(p+1)}{12} - n(p-n) \right] & \text{if } 2 \leq n < \frac{p}{2} - \frac{1}{2}\sqrt{\frac{2p^2+1}{3}}, \\ \geq 0 & \text{otherwise,} \end{cases} \end{aligned}$$

and the result (ii) follows.

(iii) $\zeta = \frac{n}{p}$, where $2 \leq n \leq \frac{1}{2}(p-1)$. Choose b, d so that $A = \begin{pmatrix} n & b \\ p & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$ and $A(\infty) = n/p$. Since $n \not\equiv 0 \pmod{p}$ we may apply (4.2) for each k . We find that $C_k = \begin{pmatrix} n+kp & * \\ p^2 & d-k'p \end{pmatrix}$, and

$$\begin{aligned} \mathrm{ord} \left(\mathcal{F}_1^* \left(\frac{1}{p}; \frac{z+k}{p} \right); \frac{n}{p} \right) &= \frac{1}{p} \mathrm{ord} \left(F_p(z); \frac{n+kp}{p^2} \right) + \frac{1}{p} \mathrm{ord} \left(\mathcal{F}_1 \left(\frac{\overline{d-k'p}}{p}; z \right); i\infty \right) \\ &= \frac{p^2-1}{24p} + 0 = \frac{p^2-1}{24p}. \end{aligned}$$

The result (iii) follows. □

5. CRANK THEOREM AND CRANK MOD p IDENTITIES

5.1. Dyson's rank conjectures and the crank theorem. 10 years after they were proposed by Dyson [10], Atkin and Swinnerton Dyer [3] came up with a proof of the Dyson's rank conjectures stated below.

Dyson's Rank Conjecture 5.1 (1944). *For all nonnegative integers n ,*

$$(5.1) \quad N(0, 5, 5n+4) = N(1, 5, 5n+4) = \cdots = N(4, 5, 5n+4) = \frac{1}{5}p(5n+4),$$

$$(5.2) \quad N(0, 7, 7n+5) = N(1, 7, 7n+5) = \cdots = N(6, 7, 7n+5) = \frac{1}{7}p(7n+5).$$

Garvan, in [15] notes that their proof involved finding and proving identities for basic hypergeometric functions, theta-functions and Lerch-type series using the theory of elliptic functions. It also involved identifying the generating functions for rank differences $N(0, p, pn+r) - N(k, p, pn+r)$ for $p = 5, 7$ for each $1 \leq k \leq \frac{1}{2}(p-1)$ and each $r = 0, 1, \dots, p-1$. Subsequently in his paper ([15], Section 6.3), Garvan gives a new proof of these conjectures using the theory of modular forms, by calculating the orders of $\mathcal{K}_{p,0}^{(R)}(\zeta_p, z)$ at the cusps of $\Gamma_1(p)$ for $p = 5$ and 7 and making use of the Valence formula.

Prior to Andrews and Garvan's discovery of the actual crank statistic [1] predicted by Dyson, Garvan in his thesis [11] introduced the vector crank statistic defined below, which simultaneously gives a combinatorial interpretation of the distribution of the partitions of $5n+4, 7n+5$ and $11n+6$ into five, seven and eleven equinumerous classes respectively. His proof of the vector crank theorem mod 11 involves an elegant utilization of Jacobi's Triple Product identity and Winquist's Identity to show that the coefficient of q^{11n+6} in the two variable generating function of the vector crank modulo 11 is zero (see [11] and [12] for details).

In this section, we employ Garvan's idea in [15] to give a new proof of the crank theorem modulo 11 (Equation (1.3)) using the Valence formula and the orders of $\mathcal{K}_{11,0}^{(C)}(\zeta_{11}, z)$ at the cusps of $\Gamma_1(11)$ calculated in the previous section. It is easy to observe that in our setup,

the crank theorem is equivalent to showing

$$(5.3) \quad \mathcal{K}_{11,0}^{(C)}(\zeta_{11}, z) = 0.$$

To prove (5.3) we use

Theorem 5.2 (The Valence Formula [21](p.98)). *Let $f \neq 0$ be a modular form of weight k with respect to a subgroup Γ of finite index in $\Gamma(1) = SL_2(\mathbb{Z})$. Then*

$$(5.4) \quad ORD(f, \Gamma) = \frac{1}{12} \mu k,$$

where μ is index $\widehat{\Gamma}$ in $\widehat{\Gamma(1)}$,

$$ORD(f, \Gamma) := \sum_{\zeta \in R^*} ORD(f, \zeta, \Gamma),$$

R^* is a fundamental region for Γ , and

$$ORD(f; \zeta; \Gamma) = n(\Gamma; \zeta) ord(f; \zeta),$$

for a cusp ζ and $n(\Gamma; \zeta)$ denotes the fan width of the cusp $\zeta \pmod{\Gamma}$.

Remark. For $\zeta \in \mathfrak{h}$, $ORD(f; \zeta; \Gamma)$ is defined in terms of the invariant order $ord(f; \zeta)$, which is interpreted in the usual sense. See [21, p.91] for details of this and the notation used.

In the following table, we give the order of $\mathcal{K}_{11,0}^{(C)}(\zeta_{11}, z)$ at the cusps using Theorem 4.5. Here $2 \leq m \leq \frac{p-1}{2}$.

cusps ζ	$n(\Gamma_1(11); \zeta)$	$ord(\mathcal{K}_{11,0}^{(C)}(\zeta_{11}, z); \zeta)$	$ORD(\mathcal{K}_{11,0}^{(C)}(\zeta_{11}, z), \Gamma_1(11), \zeta)$
$i\infty$	1	≥ 1	≥ 1
0	11	≥ 0	≥ 0
$\frac{1}{11}$	11	> 0	> 0
$\frac{m}{11}$	1	$\geq \frac{5}{11}$	≥ 1

An easy calculation by hand shows that $M(0, 11, 6) = \dots = M(10, 11, 6)$. Expanding $\mathcal{K}_{11,0}^{(C)}(\zeta_{11}, z)$ we see that this implies the coefficient of q^1 is 0 and thus we have that $ORD(\mathcal{K}_{11,0}^{(C)}(\zeta_{11}, z), \Gamma_1(11), i\infty) \geq 2$. Hence $ORD(\mathcal{K}_{11,0}^{(C)}(\zeta_{11}, z); \Gamma_1(11)) \geq 6$. But $\mu k = \frac{60}{12} = 5$. The Valence Formula implies that $\mathcal{K}_{11,0}^{(C)}(\zeta_{11}, z)$ is identically zero which proves the crank theorem.

Note : The same method could be applied to prove the mod 5 and 7 and cases of the crank theorem i.e. Equations (1.1) and (1.2).

5.2. Crank mod p identities. In this section, we give identities for $\mathcal{K}_{p,m}^{(C)}(\zeta_p, z)$ when $p = 13, 17$ and 19 in terms of generalized eta-functions given in Definition 1.3. Crank mod 13 identities were first studied by Bilgici in [5]. In [16], we gave such identities for elements of the the rank generating function. In general, these identities are of the form

$$(5.5) \quad \mathcal{K}_{p,m}^{(C)}(\zeta_p, z) = \sum_{k=1}^r c_{p,m,k} j_{p,m,k}(z)$$

where $j_{p,m,k}(z)$ are eta-quotients comprising of generalized eta-functions, and $c_{p,m,k}$ are cyclotomic coefficients. From Theorem 2.6, we know that $\mathcal{K}_{p,m}^{(C)}(\zeta_p, z)$ is a weakly holomorphic modular form of weight 1 on $\Gamma(p)$. The proof of the identities primarily involve establishing the equality using the Valence formula and showing that the RHS is also a weakly holomorphic modular form of weight 1 on $\Gamma(p)$.

In Section 7 of [16], we developed an algorithm for proving rank mod p identities that utilizes the Valence Formula. Here, we follow the steps of the algorithm to prove crank identities modulo 13.

For the cases $p = 17$ and 19 , we simply give the form of the identities omitting the values of the coefficients, for each of the cases $m = 0$ and when m is a quadratic residue and a non-residue modulo p . Their proofs follow the same technique using the algorithm as it does for $p = 13$.

5.3. Crank mod 13 identities.

5.3.1. **Identity for $\mathcal{K}_{13,0}^{(C)}$.** The following is an identity for $\mathcal{K}_{13,0}^{(C)}(\zeta_{13}, z)$ in terms of generalized eta-functions :

$$(5.6) \quad \begin{aligned} \mathcal{K}_{13,0}^{(C)}(\zeta_{13}, z) &= (q^{13}; q^{13})_{\infty} \sum_{n=1}^{\infty} \left(\sum_{k=0}^{12} M(k, 13, 13n-7) \zeta_{13}^k \right) q^n \\ &= \sum_{r=1}^6 c_{13,r} j(13, \pi_r(\vec{n}_1), z), \end{aligned}$$

where

$$\vec{n}_1 = (15, -1, -3, -2, -2, -2, -3),$$

and the coefficients are :

$$\begin{aligned} c_{13,6} &= 2\zeta_{13}^{11} + \zeta_{13}^9 + \zeta_{13}^8 + \zeta_{13}^7 + \zeta_{13}^6 + \zeta_{13}^5 + \zeta_{13}^4 + 2\zeta_{13}^2 + 2, \\ c_{13,1} &= -(\zeta_{13}^{11} + \zeta_{13}^{10} + \zeta_{13}^7 + \zeta_{13}^6 + \zeta_{13}^3 + \zeta_{13}^2), \\ c_{13,5} &= \zeta_{13}^{11} + \zeta_{13}^{10} + \zeta_{13}^9 + \zeta_{13}^8 + \zeta_{13}^5 + \zeta_{13}^4 + \zeta_{13}^3 + \zeta_{13}^2 + 1, \\ c_{13,2} &= \zeta_{13}^{10} - \zeta_{13}^9 + \zeta_{13}^8 + \zeta_{13}^5 - \zeta_{13}^4 + \zeta_{13}^3 + 1, \\ c_{13,4} &= -(\zeta_{13}^{10} + \zeta_{13}^9 + \zeta_{13}^7 + \zeta_{13}^6 + \zeta_{13}^4 + \zeta_{13}^3 + 2), \\ c_{13,3} &= -(\zeta_{13}^8 + \zeta_{13}^5). \end{aligned}$$

We follow the steps of the aforementioned algorithm in the process of proving the identity.

Step 1 We check the conditions for modularity as in Theorem 2.7 for $j(13, \pi_r(\vec{n}_1), z)$, $1 \leq r \leq 6$ involved in (5.6). Here, $p = 13, m = 0, n_0 = 15$. In [16], Theorem 3.6 tells us that it suffices to check modularity for $r = 1$. With $\vec{n}_1 = (15, -1, -3, -2, -2, -2, -3)$, we easily see that

$$n_0 + \sum_{k=1}^5 n_k = 2,$$

$$n_0 + 3 \sum_{k=1}^5 n_k = -24 \equiv 0 \pmod{24},$$

$$\sum_{k=1}^5 k^2 n_k = -221 \equiv 0 \pmod{13},$$

as required.

Since $m = 0$, we skip Step 2 in accordance with our algorithm.

Step 3 Using [15, Theorem 6.7, p.237] and [15, Theorem 6.13, p.243], we calculate the orders of each of the six functions f at each cusp of $\Gamma_1(13)$.

ORD($f, \zeta, \Gamma_1(13)$)

f	cusp ζ								
	$i\infty$	0	$1/n$	$2/13$	$3/13$	$4/13$	$5/13$	$6/13$	
$j(13, \pi_1(\vec{n}_1), z)$	3	-1	-1	2	3	2	2	1	
$j(13, \pi_2(\vec{n}_1), z)$	2	-1	-1	2	1	2	3	3	
$j(13, \pi_3(\vec{n}_1), z)$	3	-1	-1	1	2	3	2	2	
$j(13, \pi_4(\vec{n}_1), z)$	2	-1	-1	2	3	3	1	2	
$j(13, \pi_5(\vec{n}_1), z)$	2	-1	-1	3	2	1	3	2	
$j(13, \pi_6(\vec{n}_1), z)$	1	-1	-1	3	2	2	2	3	

where $2 \leq n \leq 6$.

Step 4 Now we calculate lower bounds $\lambda(13, 0, \zeta)$ of orders $Ord(\mathcal{K}_{13,0}(\zeta_{13}, z), \zeta, \Gamma_1(13))$ at cusps ζ of the LHS of equation (5.6).

cusp ζ	$\lambda(13, 0, \zeta)$
$i\infty$	
0	-1
$1/n$	0
$n/13$	7/13

where $2 \leq n \leq 6$.

Step 5 Next we compile the lower bounds of the LHS and RHS of equation (5.6) and calculate B .

cusp ζ	$n(\Gamma_1(13); \zeta)$	$\text{ORD}(LHS; \zeta)$	$\text{ORD}(RHS; \zeta)$	$\text{ORD}(LHS - RHS; \zeta)$
$i\infty$	1			
0	13	≥ -1	≥ -1	≥ -1
$1/n$	13	≥ 0	≥ -1	≥ -1
$n/13$	1	≥ 1	1	≥ 1

where $2 \leq n \leq 6$.

Thus the constant B is -1 .

Step 6 The LHS and RHS are weakly holomorphic modular forms of weight 1 on $\Gamma_1(13)$. So in the Valence Formula, $\frac{\mu k}{12} = 7$. The result follows provided we can show that $\text{ORD}(LHS - RHS, i\infty, \Gamma_1(13)) \geq 9$. This is easily verified using MAPLE.

5.3.2. **A quadratic residue case.** The following is an identity for $\mathcal{K}_{13,2}^{(C)}(\zeta_{13}, z)$ in terms of generalized eta-functions :

$$(5.7) \quad \mathcal{K}_{13,2}^{(C)}(\zeta_{13}, z) = q^{\frac{2}{13}}(q^{13}; q^{13})_{\infty} \left(\sum_{n=1}^{\infty} \left(\sum_{k=0}^{12} M(k, 13, 13n-5) \zeta_{13}^k \right) q^n \right)$$

$$= \frac{f_{13,1}(z)}{f_{13,6}(z)} \sum_{k=-1}^0 \sum_{r=1}^6 \left(\frac{\eta(13z)}{\eta(z)} \right)^{2k} c_{13,r,k} j(13, \pi_r(\vec{n}_1), z),$$

where

$$\vec{n}_1 = (15, -1, -3, -2, -2, -2, -3),$$

and the coefficients are :

$$\begin{aligned} c_{13,1,0} &= -(\zeta_{13}^{11} + \zeta_{13}^9 + \zeta_{13}^7 + \zeta_{13}^6 + \zeta_{13}^4 + \zeta_{13}^2 + 1), \\ c_{13,2,0} &= 13(\zeta_{13}^{11} - \zeta_{13}^9 - \zeta_{13}^8 - \zeta_{13}^5 - \zeta_{13}^4 + \zeta_{13}^2 - 1), \\ c_{13,3,0} &= -13(\zeta_{13}^9 + \zeta_{13}^4), \\ c_{13,4,0} &= \zeta_{13}^{11} + \zeta_{13}^{10} + \zeta_{13}^8 + \zeta_{13}^5 + \zeta_{13}^3 + \zeta_{13}^2 + 2, \\ c_{13,5,0} &= -(\zeta_{13}^{10} + \zeta_{13}^7 + \zeta_{13}^6 + \zeta_{13}^3), \\ c_{13,6,0} &= -(\zeta_{13}^{11} + \zeta_{13}^{10} + \zeta_{13}^9 + 2\zeta_{13}^8 + 2\zeta_{13}^7 + 2\zeta_{13}^6 + 2\zeta_{13}^5 + \zeta_{13}^4 + \zeta_{13}^3 + \zeta_{13}^2), \\ c_{13,1,-1} &= 0, \\ c_{13,2,-1} &= 0, \end{aligned}$$

$$\begin{aligned}
c_{13,3,-1} &= 0, \\
c_{13,4,-1} &= 0, \\
c_{13,5,-1} &= 0, \\
c_{13,6,-1} &= -(\zeta_{13}^{10} + \zeta_{13}^7 + \zeta_{13}^6 + \zeta_{13}^3).
\end{aligned}$$

We follow the steps of the algorithm in the process of proving the above identity.

Step 1 We first make note of two aspects of our functions $\frac{f_{13,1}(z)}{f_{13,6}(z)} j(13, \pi_r(\vec{n}_1), z)$ and $\frac{f_{13,1}(z)}{f_{13,6}(z)} \left(\frac{\eta(13z)}{\eta(z)}\right)^{-2} j(13, \pi_r(\vec{n}_1), z)$ involved in (5.7). First, from Apostol's [2, Theorem 4.9, p.87], we observe that the eta quotient $\left(\frac{\eta(13z)}{\eta(z)}\right)^2$ is modular under the congruence subgroup $\Gamma_0(13)$. Also, writing the theta quotient $\frac{f_{13,1}(z)}{f_{13,6}(z)}$ in its equivalent vector notation $\vec{n}' = (n'_0, n'_1, n'_2, \dots, n'_6) = (0, 1, 0, 0, 0, 0, -1)$ in accordance with our Definition 1.3, we observe that it doesn't contribute to the weight and

$$\begin{aligned}
n'_0 + 3 \sum_{k=1}^6 n'_k &= 0 \equiv 0 \pmod{24}, \\
\sum_{k=1}^6 k^2 n'_k &= -35 \equiv 4 \pmod{13}.
\end{aligned}$$

Thus it suffices to check the first step for the functions $j(13, \pi_r(\vec{n}_1), z)$, $1 \leq r \leq 6$, which was achieved in Step 1 of the proof for the previous identity $\mathcal{K}_{13,0}^{(C)}(\zeta_{13}, z)$.

Step 2 Next, we calculate the orders of the generalized eta-functions at $i\infty$ and considering the identity with zero coefficients removed, we find that $k_0 = 1$. Thus we divide each generalized eta-function by $j_0 = \frac{f_{13,1}(z)}{f_{13,6}(z)} \left(\frac{\eta(13z)}{\eta(z)}\right)^{-2} j(13, \pi_6(\vec{n}_1), z)$, which has the lowest order at $i\infty$.

Step 3 Using [15, Theorem 6.7, p.237] and [15, Theorem 6.13, p.243], we calculate the orders of each of the six functions f at each cusp of $\Gamma_1(13)$.

$$\text{ORD}(f, \zeta, \Gamma_1(13))$$

f	$i\infty$	0	$1/n$	cusp ζ					
				$2/13$	$3/13$	$4/13$	$5/13$	$6/13$	
$\frac{f_{13,1}(z)}{f_{13,6}(z)}j(13, \pi_1(\vec{n}_1), z)/j_0$	3	-1	-1	0	2	1	1	-1	
$\frac{f_{13,1}(z)}{f_{13,6}(z)}j(13, \pi_2(\vec{n}_1), z)/j_0$	2	-1	-1	0	0	1	2	1	
$\frac{f_{13,1}(z)}{f_{13,6}(z)}j(13, \pi_3(\vec{n}_1), z)/j_0$	3	-1	-1	-1	1	2	1	0	
$\frac{f_{13,1}(z)}{f_{13,6}(z)}j(13, \pi_4(\vec{n}_1), z)/j_0$	2	-1	-1	0	2	2	0	0	
$\frac{f_{13,1}(z)}{f_{13,6}(z)}j(13, \pi_5(\vec{n}_1), z)/j_0$	2	-1	-1	1	1	0	2	0	
$\frac{f_{13,1}(z)}{f_{13,6}(z)}j(13, \pi_6(\vec{n}_1), z)/j_0$	1	-1	-1	1	1	1	1	1	
$\frac{f_{13,1}(z)}{f_{13,6}(z)}\left(\frac{\eta(13z)}{\eta(z)}\right)^{-2}j(13, \pi_6(\vec{n}_1), z)/j_0$	0	0	0	0	0	0	0	0	

where $2 \leq n \leq 6$.

Step 4 Now we calculate lower bounds $L(13, 2, \zeta)$ of orders at cusps ζ of the LHS of equation (5.7) divided by j_0 .

cusp ζ	$Ord\left(\frac{\mathcal{K}_{13,2}^{(C)}(\zeta_{13,z})}{j_0}, \zeta, \Gamma_1(13)\right) \geq$
$i\infty$	
0	-1
$1/n$	0
$2/13$	-14/13
$3/13$	-11/13
$4/13$	1/13
$5/13$	-4/13
$6/13$	-1

where $2 \leq n \leq 6$.

Step 5 Next we compile the lower bounds of the LHS and RHS of equation (5.7) divided by j_0 and deduce B .

cusp ζ	$n(\Gamma_1(13); \zeta)$	$ORD(LHS; \zeta)$	$ORD(RHS; \zeta)$	$ORD(LHS - RHS; \zeta)$
$i\infty$	1			
0	13	≥ -1	≥ -1	≥ -1
$1/n$	13	0	≥ -1	≥ -1
$2/13$	1	≥ -1	-2	≥ -2
$3/13$	1	≥ 0	-1	≥ -1
$4/13$	1	≥ 1	-1	≥ -1
$5/13$	1	≥ 0	-1	≥ -1
$6/13$	1	≥ -1	-2	≥ -2

where $2 \leq n \leq 6$.

Thus the constant B is -13 .

Step 6 The LHS and RHS are weakly holomorphic modular forms of weight 0 on $\Gamma_1(13)$. So in the Valence Formula, $\frac{\mu_k}{12} = 0$. The result follows provided we can show that $\text{ORD}(LHS - RHS, i\infty, \Gamma_1(13)) \geq 14$. This is easily verified using MAPLE.

5.3.3. **A quadratic non-residue case.** The following is an identity for $\mathcal{K}_{13,1}^{(C)}(\zeta_{13}, z)$ in terms of generalized eta-functions :

$$(5.8) \quad \mathcal{K}_{13,1}^{(C)}(\zeta_{13}, z) = q^{\frac{1}{13}}(q^{13}; q^{13})_{\infty} \sum_{n=1}^{\infty} \left(\sum_{k=0}^{12} M(k, 13, 13n-6) \zeta_{13}^k \right) q^n$$

$$= \frac{f_{13,1}(z)}{f_{13,5}(z)} \sum_{k=-1}^0 \sum_{r=1}^6 \left(\frac{\eta(13z)}{\eta(z)} \right)^{2k} c_{13,r,k} j(13, \pi_r(\vec{n}_1), z),$$

where

$$\vec{n}_1 = (15, -1, -3, -2, -2, -2, -3),$$

and the coefficients are :

$$\begin{aligned} c_{13,1,0} &= \zeta_{13}^{11} + \zeta_{13}^9 + \zeta_{13}^7 + \zeta_{13}^6 + \zeta_{13}^4 + \zeta_{13}^2 + 2, \\ c_{13,2,0} &= -(\zeta_{13}^9 + \zeta_{13}^8 + \zeta_{13}^5 + \zeta_{13}^4), \\ c_{13,3,0} &= \zeta_{13}^{11} + \zeta_{13}^{10} + \zeta_{13}^9 + \zeta_{13}^4 + \zeta_{13}^3 + \zeta_{13}^2, \\ c_{13,4,0} &= -(\zeta_{13}^{11} + \zeta_{13}^{10} + \zeta_{13}^9 + \zeta_{13}^8 + \zeta_{13}^7 + \zeta_{13}^6 + \zeta_{13}^5 + \zeta_{13}^4 + \zeta_{13}^3 + \zeta_{13}^2 + 1), \\ c_{13,5,0} &= -\zeta_{13}^{11} + \zeta_{13}^{10} - \zeta_{13}^8 - \zeta_{13}^5 + \zeta_{13}^3 - \zeta_{13}^2 + 1, \\ c_{13,6,0} &= -(\zeta_{13}^{10} + \zeta_{13}^9 + \zeta_{13}^8 + 2\zeta_{13}^7 + 2\zeta_{13}^6 + \zeta_{13}^5 + \zeta_{13}^4 + \zeta_{13}^3), \\ c_{13,1,-1} &= 0, \\ c_{13,2,-1} &= 0, \\ c_{13,3,-1} &= 0, \\ c_{13,4,-1} &= 0, \\ c_{13,5,-1} &= 0, \\ c_{13,6,-1} &= -(\zeta_{13}^{11} + \zeta_{13}^{10} + \zeta_{13}^9 + \zeta_{13}^8 + \zeta_{13}^7 + \zeta_{13}^6 + \zeta_{13}^5 + \zeta_{13}^4 + \zeta_{13}^3 + \zeta_{13}^2 + 1). \end{aligned}$$

Utilizing the same algorithm, the proof follows similar to the identity for the quadratic residue case $\mathcal{K}_{13,2}^{(C)}(\zeta_{13}, z)$ done previously. We omit the details here.

Below, we present one identity for each of the quadratic residue, quadratic non-residue and $m = 0$ cases for $p = 17$ and $p = 19$. These are new and donot seem to appear in the literature elsewhere.

5.4. Crank mod 17 identities.

5.4.1. **Identity for $\mathcal{K}_{17,0}^{(C)}$.** The following is an identity for $\mathcal{K}_{17,0}^{(C)}(\zeta_{17}, z)$ in terms of generalized eta-functions :

(5.9)

$$\begin{aligned} \mathcal{K}_{17,0}^{(C)}(\zeta_{17}, z) &= (q^{17}; q^{17})_{\infty} \sum_{n=1}^{\infty} \left(\sum_{k=0}^{16} M(k, 17, 17n - 12) \zeta_{17}^k \right) q^n \\ &= \sum_{k=0}^1 \sum_{r=1}^8 \left(\frac{\eta(17z)}{\eta(z)} \right)^{3k} (c_{17,r,k} j(17, \pi_r(\vec{n}_1), z) + d_{17,r,k} j(17, \pi_r(\vec{n}_2), z)), \end{aligned}$$

where

$$\begin{aligned} \vec{n}_1 &= (15, -3, -1, -2, -1, -2, -1, -2, -1) \\ \vec{n}_2 &= (27, -2, -2, -3, -2, -4, -4, -4, -4), \end{aligned}$$

and the coefficients $c_{17,r,k}, d_{17,r,k}$, $1 \leq r \leq 8, 0 \leq k \leq 1$ are linear combinations of cyclotomic integers like the mod 11 and 13 identities found previously. We donot include them here. We make note of the fact that the same set of theta functions j and vectors \vec{n}_1 and \vec{n}_2 were also involved in the corresponding identity $\mathcal{K}_{17,0}^{(R)}$ for the rank ([16], Equation 7.10). However, there are eight fewer functions in the identity for the crank.

5.4.2. **A quadratic residue case.** The following is an identity for $\mathcal{K}_{17,12}^{(C)}(\zeta_{17}, z)$ in terms of generalized eta-functions :

(5.10)

$$\begin{aligned} \mathcal{K}_{17,12}^{(C)}(\zeta_{17}, z) &= q^{\frac{12}{17}} (q^{17}; q^{17})_{\infty} \left(\sum_{n=1}^{\infty} \left(\sum_{k=0}^{16} M(k, 17, 17n) \zeta_{17}^k \right) q^n \right) \\ &= \frac{f_{17,7}(z)}{f_{17,5}(z)} \sum_{k=-1}^1 \sum_{r=1}^8 \left(\frac{\eta(17z)}{\eta(z)} \right)^{3k} (c_{17,r,k} j(17, \pi_r(\vec{n}_1), z) + d_{17,r,k} j(17, \pi_r(\vec{n}_2), z)), \end{aligned}$$

where

$$\vec{n}_1 = (15, -3, -1, -2, -1, -2, -1, -2, -1),$$

$$\vec{n}_2 = (27, -2, -2, -3, -2, -4, -4, -4, -4),$$

and the coefficients $c_{17,r,k}, d_{17,r,k}$, $1 \leq r \leq 8, -1 \leq k \leq 1$ are linear combinations of cyclotomic integers like the identities found previously. We donot include them here. Several of these coefficients are zero. We make note of the fact that the same set of theta functions j and vectors \vec{n}_1 and \vec{n}_2 were also involved in the corresponding identity $\mathcal{K}_{17,12}^{(R)}$ for the rank ([16], Equation 7.11). However, there are much fewer functions in the identity for the crank.

5.4.3. A quadratic non-residue case. The following is an identity for $\mathcal{K}_{17,1}^{(C)}(\zeta_{17}, z)$ in terms of generalized eta-functions :

(5.11)

$$\begin{aligned} \mathcal{K}_{17,1}^{(C)}(\zeta_{17}, z) &= q^{\frac{1}{17}}(q^{17}; q^{17})_{\infty} \sum_{n=1}^{\infty} \left(\sum_{k=0}^{16} M(k, 17, 17n - 11) \zeta_{17}^k \right) q^n \\ &= \frac{f_{17,7}(z)}{f_{17,8}(z)} \sum_{k=-1}^1 \sum_{r=1}^8 \left(\frac{\eta(17z)}{\eta(z)} \right)^{3k} (c_{17,r,k} j(17, \pi_r(\vec{n}_1), z) + d_{17,r,k} j(17, \pi_r(\vec{n}_2), z)), \end{aligned}$$

where

$$\begin{aligned} \vec{n}_1 &= (15, -3, -1, -2, -1, -2, -1, -2, -1), \\ \vec{n}_2 &= (27, -2, -2, -3, -2, -4, -4, -4, -4), \end{aligned}$$

and the coefficients $c_{17,r,k}, d_{17,r,k}$, $1 \leq r \leq 8, -1 \leq k \leq 1$ are linear combinations cyclotomic integers like the identities found previously. We donot include them here. Several of these coefficients are zero. We make note of the fact that the same set of theta functions j and vectors \vec{n}_1 and \vec{n}_2 were also involved in the corresponding identity $\mathcal{K}_{17,1}^{(R)}$ for the rank ([16], Equation 7.12). However, there are much fewer functions in the identity for the crank.

5.5. Crank mod 19 identities.

5.5.1. Identity for $\mathcal{K}_{19,0}^{(C)}$. The following is an identity for $\mathcal{K}_{19,0}^{(C)}(\zeta_{19}, z)$ in terms of generalized eta-functions :

(5.12)

$$\mathcal{K}_{19,0}^{(C)}(\zeta_{19}, z) = (q^{19}; q^{19})_{\infty} \sum_{n=1}^{\infty} \left(\sum_{k=0}^{18} M(k, 19, 19n - 15) \zeta_{19}^k \right) q^n$$

$$= \sum_{k=0}^1 \sum_{r=1}^9 \left(\frac{\eta(19z)}{\eta(z)} \right)^{4k} \left(c_{19,r,k} j(19, \pi_r(\vec{n}_1), z) + d_{19,r,k} j(19, \pi_r(\vec{n}_2), z) + e_{19,r,k} j(19, \pi_r(\vec{n}_3), z) \right),$$

where

$$\begin{aligned} \vec{n}_1 &= (27, -3, -2, -4, -4, -3, -3, -2, -3, -1), \\ \vec{n}_2 &= (39, -5, -2, -5, -5, -3, -5, -2, -5, -5), \\ \vec{n}_3 &= (39, -5, -4, -3, -4, -5, -4, -4, -5, -3), \end{aligned}$$

and the coefficients $c_{19,r,k}, d_{19,r,k}, e_{19,r,k}, 1 \leq r \leq 9, 0 \leq k \leq 1$ are linear combinations of cyclotomic integers like the mod 11, 13 and 17 identities found previously. We donot include them here. We make note of the fact that the same set of theta functions j and vectors \vec{n}_1, \vec{n}_2 and \vec{n}_3 were also involved in the corresponding identity $\mathcal{K}_{19,0}^{(R)}$ for the rank ([16], Equation 7.13). However, there are nine fewer functions in the identity for the crank.

5.5.2. A quadratic residue case. The following is an identity for $\mathcal{K}_{19,15}^{(C)}(\zeta_{19}, z)$ in terms of generalized eta-functions :

(5.13)

$$\begin{aligned} \mathcal{K}_{19,15}^{(C)}(\zeta_{19}, z) &= q^{\frac{15}{19}}(q^{19}; q^{19})_{\infty} \left(\sum_{n=1}^{\infty} \left(\sum_{k=0}^{18} M(k, 19, 19n) \zeta_{19}^k \right) q^n \right) \\ &= \frac{f_{19,6}(z)}{f_{19,5}(z)} \left(\sum_{k=-1}^1 \sum_{r=1}^9 \left(\frac{\eta(19z)}{\eta(z)} \right)^{4k} \left(c_{19,r,k} j(19, \pi_r(\vec{n}_1), z) + d_{19,r,k} j(19, \pi_r(\vec{n}_2), z) \right. \right. \\ &\quad \left. \left. + e_{19,r,k} j(19, \pi_r(\vec{n}_3), z) \right) \right), \end{aligned}$$

where

$$\begin{aligned} \vec{n}_1 &= (39, -5, -5, -4, -5, -4, -5, -3, -5, -1), \\ \vec{n}_2 &= (39, -3, -5, -5, -5, -3, -2, -4, -5, -5), \\ \vec{n}_3 &= (39, -4, -5, -4, -3, -3, -4, -5, -5, -4), \end{aligned}$$

and the coefficients $c_{19,r,k}, d_{19,r,k}, e_{19,r,k}, 1 \leq r \leq 9, -1 \leq k \leq 1$ are linear combinations of cyclotomic integers like the identities found previously. We donot include them here. Several of these coefficients are zero. We make note of the fact that the same set of theta functions

j and vectors \vec{n}_1 and \vec{n}_2 were also involved in the corresponding identity $\mathcal{K}_{19,15}^{(R)}$ for the rank ([16], Equation 7.14). However, there are much fewer functions in the identity for the crank.

5.5.3. A quadratic non-residue case. The following is an identity for $\mathcal{K}_{19,1}^{(C)}(\zeta_{19}, z)$ in terms of generalized eta-functions :

(5.14)

$$\begin{aligned} \mathcal{K}_{19,1}^{(C)}(\zeta_{19}, z) &= q^{\frac{1}{19}}(q^{19}; q^{19})_{\infty} \sum_{n=1}^{\infty} \left(\sum_{k=0}^{18} M(k, 19, 19n - 14) \zeta_{19}^k \right) q^n \\ &= \frac{f_{19,8}(z)}{f_{19,9}(z)} \left(\sum_{k=-1}^1 \sum_{r=1}^9 \left(\frac{\eta(19z)}{\eta(z)} \right)^{4k} \left(c_{19,r,k} j(19, \pi_r(\vec{n}_1), z) + d_{19,r,k} j(19, \pi_r(\vec{n}_2), z) \right. \right. \\ &\quad \left. \left. + e_{19,r,k} j(19, \pi_r(\vec{n}_3), z) \right) \right), \end{aligned}$$

where

$$\begin{aligned} \vec{n}_1 &= (39, -5, -5, -4, -5, -4, -5, -3, -5, -1), \\ \vec{n}_2 &= (39, -3, -5, -5, -5, -3, -2, -4, -5, -5), \\ \vec{n}_3 &= (39, -4, -5, -4, -3, -3, -4, -5, -5, -4), \end{aligned}$$

and the coefficients $c_{19,r,k}, d_{19,r,k}, e_{19,r,k}, 1 \leq r \leq 9, -1 \leq k \leq 1$ are linear combinations of cyclotomic integers like the identities found previously. We donot include them here. Several of these coefficients are zero. We make note of the fact that the same set of theta functions j and vectors \vec{n}_1 and \vec{n}_2 were also involved in the corresponding identity $\mathcal{K}_{19,1}^{(R)}$ for the rank ([16], Equation 7.16). However, there are much fewer functions in the identity for the crank.

6. CONCLUDING REMARKS

Building on our work on symmetries of the rank generating function [16], we have found analogous results for the crank generating function in this paper. We restate the fact that the transformation and modularity properties as well as the p -dissection identities are similar for the rank and crank functions. It would be interesting to examine the underlying theory behind this resemblance more closely. Two other natural problems to consider are finding crank dissection identities for higher primes $p \geq 23$ and also exploring such symmetries for other partition statistics and also their overpartition analogues.

7. ACKNOWLEDGMENTS

The author would like to thank his advisor Frank Garvan for suggesting this problem and his immense support and guidance throughout the course of this project.

REFERENCES

1. G.E. Andrews and F. G. Garvan, *Dyson's crank of a partition*, Bull. Amer. Math. Soc. (N.S.) **18** (1988), 167–171.
2. T.M. Apostol, *Modular functions and Dirichlet series in number theory*, Graduate Texts in Mathematics, No. 41, Springer-Verlag, New York-Heidelberg, 1976.
3. A. O. L. Atkin and P. Swinnerton-Dyer, *Some properties of partitions*, Proc. London Math. Soc. (3) **4** (1954), 84–106.
4. A. J. F. Biagioli, *A proof of some identities of Ramanujan using modular forms*, Glasgow Math. J. **31** (1989), 271–295.
5. G. Bilgici, *The crank of partitions mod 13*, Ramanujan J. **30**(3) (2013), 403–424.
6. K. Bringmann and K. Ono, *Dyson's ranks and Maass forms*, Ann. of Math. (2) **171** (2010), 419–449.
7. K. Bringmann, K. Ono, and R. C. Rhoades, *Eulerian series as modular forms*, J. Amer. Math. Soc. **21** (2008), 1085–1104.
8. K. Bringmann and H. Swisher, *On a conjecture of Koike on identities between Thompson series and Rogers-Ramanujan functions*, Proc. Amer. Math. Soc. **135** (2007), 2317–2326.
9. B. Cho, J. K. Koo, Y. K. Park, *Arithmetic of the Ramanujan-Göllnitz-Gordon continued fraction*, J. Number Theory **129** (2009), 922–947.
10. F. J. Dyson, *Some guesses in the theory of partitions*, Eureka (Cambridge) **8** (1944), 10–15.
11. F. G. Garvan, *Generalizations of Dyson's rank*, Thesis (Ph.D.), The Pennsylvania State University (1986).
12. F. G. Garvan, *New combinatorial interpretations of Ramanujan's partition congruences mod 5, 7 and 11*, Trans. Amer. Math. Soc. **305** (1988), 47–77.
13. F. G. Garvan, *Combinatorial interpretations of Ramanujan's partition congruences* in “Ramanujan Revisited: Proc. of the Centenary Conference, Univ. of Illinois at Urbana-Champaign, June 1-5, 1987”, Acad. Press, San Diego, 1988.
14. F. G. Garvan, *The crank of partitions mod 8, 9 and 10*, Trans. Amer. Math. Soc. **322** (1990), 79–94.
15. F. G. Garvan, *Transformation properties for Dyson's rank function*, Trans. Amer. Math. Soc. **371** (2019), 199–248.
16. F. G. Garvan, and R. Sarma, *New symmetries for Dyson's rank function*, submitted, preprint arXiv:2301.08960, 2023.
17. M. D. Hirschhorn, *A birthday present for Ramanujan*, Amer. Math. Monthly **97** (1990), 398–400.
18. M. I. Knopp, “Modular Functions in Analytic Number Theory,” Markham Publishing Co., Chicago, Illinois, 1970.
19. K. Mahlburg, *Partition congruences and the Andrews-Garvan-Dyson crank*, Proc. Natl. Acad. Sci. USA **102** (2005), 15373–15376.
20. G. Köhler, *Eta Products and Theta Series Identities*, Springer Monographs in Mathematics, Springer Heidelberg, 2011.
21. R. A. Rankin, “Modular Forms and Functions,” Cambridge University Press, 1977.
22. S. Ramanujan, “The Lost Notebook and Other Unpublished Papers,” Springer-Verlag, Berlin, 1988.
23. Y. Yang, *Rank generating functions as weakly holomorphic modular forms*, Bull. London Math. Soc. **36** (2004), 671–682.
24. S. P. Zwegers, *Mock θ -functions and real analytic modular forms*, in “ q -Series with Applications to Combinatorics, Number Theory, and Physics,” Contemp. Math. **291** (2001), 269–277.
25. S. P. Zwegers, “Mock Theta Functions,” Ph.D. thesis, Universiteit Utrecht, 2002, 96 pp.