

ON THE NEUMANN (p, q) -EIGENVALUE PROBLEM IN HÖLDER SINGULAR DOMAINS

PRASHANTA GARAIN, VALERII PCHELINTSEV, ALEXANDER UKHLOV

ABSTRACT. In the article we study the Neumann (p, q) -eigenvalue problems in bounded Hölder γ -singular domains $\Omega_\gamma \subset \mathbb{R}^n$. In the case $1 < p < \infty$ and $1 < q < p_\gamma^*$ we prove solvability of this eigenvalue problem and existence of the minimizer of the associated variational problem. In addition, we establish some regularity results of the eigenfunctions and some estimates of (p, q) -eigenvalues.

1. INTRODUCTION

In this article, we investigate the following Neumann (p, q) -eigenvalue problem:

$$(1.1) \quad -\Delta_p u := -\operatorname{div}(|\nabla u|^{p-2} \nabla u) = \lambda \|u\|_{L^q(\Omega_\gamma)}^{p-q} |u|^{q-2} u \text{ in } \Omega_\gamma, \quad \frac{\partial u}{\partial \nu} = 0 \text{ on } \partial\Omega_\gamma,$$

in bounded domains with anisotropic Hölder γ -singularities

$$\Omega_\gamma = \{x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n : 0 < x_n < 1, 0 < x_i < g_i(x_n), i = 1, 2, \dots, n-1\},$$

where $g_i(t) = t^{\gamma_i}$, $\gamma_i \geq 1$, $0 < t < 1$, are Hölder functions and we denote by $\gamma = \log(g_1(t) \cdot \dots \cdot g_{n-1}(t))(\log t)^{-1} + 1$. It is evident that $\gamma \geq n$. In the case $g_1 = g_2 = \dots = g_{n-1}$ we will say that domain Ω_γ is a domain with σ -Hölder singularity, $\sigma = (\gamma - 1)/(n - 1)$. Note that the main results of the article include the case of Lipschitz domains $\Omega \subset \mathbb{R}^n$, in this case we can put $\gamma = n$. We assume that $1 < p < \gamma$ and $1 < q < p_\gamma^*$, where $p_\gamma^* = \gamma p / (\gamma - p)$.

The associated Dirichlet eigenvalue problem of (1.1), i.e.,

$$(1.2) \quad -\Delta_p u = \lambda \|u\|_{L^q(\Omega)}^{p-q} |u|^{q-2} u \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega,$$

is widely studied in the literature concerning the existence, regularity among other qualitative properties of eigenfunctions for $p = q$ as well as for $p \neq q$. When $p = q$ in (1.2), we refer to Lindqvist [22], García Azorero-Peral Alonso [11], Anane-Tousli [1, 2], Drábek-Kufner-Nicolosi [6] and the references therein. When $p \neq q$ in (1.2), we refer to Franzina-Lamberti [9], Ercole [7], Garain-Ukhlov [10] and the references therein.

In contrast to the Dirichlet problem (1.2), the Neumann problem (1.1) is less understood in the literature. In this concern, for $p = q$ in (1.1), i.e., for the equation

$$(1.3) \quad -\Delta_p u = \lambda |u|^{p-2} u \text{ in } \Omega, \quad \frac{\partial u}{\partial \nu} = 0 \text{ on } \partial\Omega,$$

in Lipschitz domains $\Omega \subset \mathbb{R}^n$ we refer to Greco-Lucia [18] for the linear case $p = q = 2$. In the nonlinear setting, Lê [21] proved existence, regularity among other

⁰**Key words and phrases:** p -Laplacian, Neumann eigenvalue problem, existence, regularity, quasiconformal mappings.

⁰**2010 Mathematics Subject Classification:** 35P15, 35P30, 35A01, 35J92, 46E35, 30C65.

qualitative properties of the eigenfunctions of (1.3). For $p \neq q$ in (1.1) under the hypothesis that $\Omega = B_1 \cup B_2$ for two disjoint balls B_1 and B_2 , Croce-Henrot-Pisante [4] proved existence of a bounded eigenfunction of (1.1) in the space $W_{\text{per}}^{1,p}(\Omega)$, which stands for the Sobolev space of functions in $W^{1,p}(\Omega)$ taking constant boundary values, see also Nazarov [25] for related results.

Estimates of Neumann eigenvalues of the p -Laplace operator in non-convex domains is a long-standing complicated problem [20, 26, 27]. This problem was partially solved on the base of composition operators on Sobolev spaces, see for example, [13, 15, 16]. In this article we prove that the first non-trivial Neumann (p, q) -eigenvalues can be characterized by the the Min-Max Principle

$$\lambda_{p,q}(\Omega_\gamma) = \min \left\{ \frac{\|\nabla u\|_{L^p(\Omega_\gamma)}^p}{\|u\|_{L^q(\Omega_\gamma)}^p} : u \in W^{1,p}(\Omega_\gamma) \setminus \{0\}, \int_{\Omega_\gamma} |u|^{q-2} u \, dx = 0 \right\}.$$

By using this characterization we give lower estimates of Neumann (p, q) -eigenvalues in bounded Hölder singular domains $\Omega_\gamma \subset \mathbb{R}^n$. In particular, we give the following lower estimate of the first nontrivial Neumann (p, q) -eigenvalue

$$\lambda_{3,2}(\Omega_\gamma) \geq \left(12\pi \sqrt{(\gamma_1 - 1)^2 + (\gamma_2 - 1)^2 + 3} \right)^{-3},$$

where $\Omega_\gamma \subset \mathbb{R}^3$ is a domain with anisotropic Hölder γ -singularities, $\gamma = \gamma_1 + \gamma_2 + 1$, $3 < \gamma < 6$.

So, this example gives, in particular, in the frameworks the conjecture by V. Maz'ya [23], that exact Sobolev embeddings in anisotropic Hölder singular domains depends not on $\gamma = \gamma_1 + \dots + \gamma_{n-1} + 1$ only.

In addition, in any bounded Hölder singular domain Ω_γ , for $q = 2$, we prove existence of eigenfunction of (1.1) in $W^{1,p}(\Omega_\gamma)$ (see Theorem 4.1-4.2) with zero mean, but not necessarily takes the constant boundary values as in [4]. Further, we study the associated minimizing problem of (1.1) (see Theorem 4.2) and prove the boundedness of the eigenfunctions among other qualitative properties (see Theorem 4.3). To this end, we follow the approach from Ercole [7].

This article is organized as follows: In Section 2, we prove Min-Max Principle for the first non-trivial Neumann (p, q) -eigenvalue. In Section 3, we give estimates of Neumann (p, q) -eigenvalues in Hölder singular domains. Finally, in Section 4, we mention the functional setting related to the problem (1.1) and further state and prove our regularity results.

2. NEUMANN EIGENVALUE PROBLEM

2.1. Sobolev spaces. Let $\Omega \subset \mathbb{R}^n$ be an open set. Then the Sobolev space $W^{1,p}(\Omega)$, $1 \leq p \leq \infty$, is defined as a Banach space of locally integrable weakly differentiable functions $u : \Omega \rightarrow \mathbb{R}$ equipped with the following norm:

$$\|u\|_{W^{1,p}(\Omega)} := \|u\|_{L^p(\Omega)} + \|\nabla u\|_{L^p(\Omega)},$$

where $L^p(\Omega)$ is the Lebesgue space with the standard norm. In accordance with the non-linear potential theory [24] we consider elements of Sobolev spaces $W^{1,p}(\Omega)$ as equivalence classes up to a set of p -capacity zero [23].

The Sobolev space $W_{\text{loc}}^{1,p}(\Omega)$ is defined as follows: $u \in W_{\text{loc}}^{1,p}(\Omega)$ if and only if $u \in W^{1,p}(U)$ for every open and bounded set $U \subset \Omega$ such that $\bar{U} \subset \Omega$, where \bar{U} is the closure of the set U .

We consider domains with anisotropic Hölder γ -singularities

$\Omega_\gamma = \{x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n : 0 < x_n < 1, 0 < x_i < g_i(x_n), i = 1, 2, \dots, n-1\}$, where $g_i(t) = t^{\gamma_i}$, $\gamma_i \geq 1$, $0 < t < 1$ are Hölder functions and we denote by $\gamma = \log(g_1(t) \cdot \dots \cdot g_{n-1}(t))(\log t)^{-1} + 1$.

Let us recall the Sobolev embedding theorem in Hölder singular domains [12, 14].

Theorem 2.1. *Let $\Omega_\gamma \subset \mathbb{R}^n$ be a domain with anisotropic Hölder γ -singularities. Suppose $1 < p < \gamma$. Then the embedding operator*

$$W^{1,p}(\Omega_\gamma) \hookrightarrow L^r(\Omega_\gamma)$$

is compact for any $1 < r < p_\gamma^*$, where $p_\gamma^* = \gamma p / (\gamma - p)$.

Throughout Section 2-3, we assume that $1 < p < \gamma$ and $1 < q < p_\gamma^*$ unless otherwise mentioned. Now for $\lambda \in \mathbb{R}$, we consider the following Neumann (p, q) -eigenvalue problem

$$(2.1) \quad -\Delta_p u := -\operatorname{div}(|\nabla u|^{p-2} \nabla u) = \lambda \|u\|_{L^q(\Omega_\gamma)}^{p-q} u \text{ in } \Omega_\gamma, \quad \frac{\partial u}{\partial \nu} = 0 \text{ on } \partial\Omega_\gamma,$$

where $\Omega_\gamma \subset \mathbb{R}^n$ and ν is the outward unit normal to $\partial\Omega_\gamma$ in the weak formulation:

The pair $(\lambda, u) \in \mathbb{R} \times W^{1,p}(\Omega_\gamma) \setminus \{0\}$ is an eigenpair of (1.1) if for every $\phi \in W^{1,p}(\Omega_\gamma)$, we have

$$(2.2) \quad \int_{\Omega_\gamma} |\nabla u|^{p-2} \nabla u \nabla \phi \, dx = \lambda \|u\|_{L^q(\Omega_\gamma)}^{p-q} \int_{\Omega_\gamma} u \phi \, dx.$$

We refer to λ as an eigenvalue and u as the eigenfunction corresponding to λ .

Now we prove the Min-Max Principle for the first non-trivial Neumann (p, q) -eigenvalue. We establish Theorem 2.3 following the proof of [4, Lemma 2]. To this end, first we obtain the following auxiliary result.

Lemma 2.2. *Let $v \in W^{1,p}(\Omega_\gamma) \setminus \{0\}$ be such that $\int_{\Omega_\gamma} |v|^{q-2} v \, dx = 0$. Then there exists a constant $C = C(\Omega_\gamma) > 0$ such that*

$$\|v\|_{L^p(\Omega_\gamma)} \leq C \|\nabla v\|_{L^p(\Omega_\gamma)}.$$

Proof. By contradiction, suppose for every $n \in \mathbb{N}$, there exists $v_n \in W^{1,p}(\Omega_\gamma) \setminus \{0\}$ such that $\int_{\Omega_\gamma} |v_n|^{q-2} v_n \, dx = 0$ and

$$(2.3) \quad \|v_n\|_{L^p(\Omega_\gamma)} > n \|\nabla v_n\|_{L^p(\Omega_\gamma)}.$$

Without loss of generality, let us assume that $\|v_n\|_{L^p(\Omega_\gamma)} = 1$. If not, we define

$$u_n = \frac{v_n}{\|v_n\|_{L^p(\Omega_\gamma)}},$$

then $\|u_n\|_{L^p(\Omega_\gamma)} = 1$ and (2.3) holds for u_n and also $\int_{\Omega_\gamma} |u_n|^{q-2} u_n \, dx = 0$. By (2.3), since $\|\nabla v_n\|_{L^p(\Omega_\gamma)} \rightarrow 0$ as $n \rightarrow \infty$, we have that the sequence of functions $\{v_n\}, n \in \mathbb{N}$, is uniformly bounded in $W^{1,p}(\Omega_\gamma)$. By Theorem 2.1, since the embedding operator

$$W^{1,p}(\Omega_\gamma) \hookrightarrow L^q(\Omega_\gamma), \quad 1 < q < p_\gamma^*,$$

is compact, there exists $v \in W^{1,p}(\Omega_\gamma)$ such that

$$v_n \rightharpoonup v \text{ weakly in } W^{1,p}(\Omega_\gamma), \quad v_n \rightarrow v \text{ strongly in } L^q(\Omega_\gamma), \quad \forall 1 < q < p_\gamma^*,$$

and $g \in L^q(\Omega_\gamma)$ such that

$$|v_n| \leq g \text{ a.e. in } \Omega_\gamma, \quad \nabla v_n \rightharpoonup \nabla v \text{ weakly in } L^p(\Omega_\gamma).$$

Since $\|\nabla v_n\|_{L^p(\Omega_\gamma)} \rightarrow 0$ as $n \rightarrow \infty$, we have $\nabla v_n \rightarrow 0$ weakly in $L^p(\Omega_\gamma)$, hence $\nabla v = 0$ a.e. in Ω_γ , which gives that $v = \text{constant}$ a.e. in Ω_γ . This combined with the fact that

$$0 = \lim_{n \rightarrow \infty} \int_{\Omega_\gamma} |v_n|^{q-2} v_n dx = \int_{\Omega_\gamma} |v|^{q-2} v dx$$

gives that $v = 0$ a.e. in Ω_γ . This contradicts the hypothesis that $\|v_n\|_{L^p(\Omega_\gamma)} = 1$. This completes the proof. \square

Theorem 2.3. *There exists $u \in W^{1,p}((\Omega_\gamma) \setminus \{0\})$ such that $\int_{\Omega_\gamma} |u|^{q-2} u dx = 0$. Moreover,*

$$\lambda_{p,q}(\Omega_\gamma) = \inf \left\{ \frac{\|\nabla u\|_{L^p(\Omega_\gamma)}^p}{\|u\|_{L^q(\Omega_\gamma)}^p} : u \in W^{1,p}(\Omega_\gamma) \setminus \{0\}, \int_{\Omega_\gamma} |u|^{q-2} u dx = 0 \right\} = \frac{\|\nabla u\|_{L^p((\Omega_\gamma))}^p}{\|u\|_{L^q((\Omega_\gamma))}^p}.$$

Further, $\lambda_{p,q}(\Omega_\gamma) > 0$.

Proof. Let $n \in \mathbb{N}$ and define the functionals $G : W^{1,p}(\Omega_\gamma) \rightarrow \mathbb{R}$ by

$$G(v) = \int_{\Omega_\gamma} |v|^{q-2} v dx,$$

and $H_{\frac{1}{n}} : W^{1,p}(\Omega_\gamma) \rightarrow \mathbb{R}$ by

$$H_{\frac{1}{n}}(v) = \|\nabla v\|_{L^p(\Omega_\gamma)}^p - (\lambda_{p,q}^p + \frac{1}{n}) \|v\|_{L^q(\Omega_\gamma)}^p,$$

where we denoted $\lambda_{p,q}(\Omega_\gamma)$ by $\lambda_{p,q}$. By the definition of infimum, for every $n \in \mathbb{N}$, there exists $u_n \in W^{1,p}(\Omega_\gamma) \setminus \{0\}$ such that $\int_{\Omega_\gamma} |u_n|^{q-2} u_n dx = 0$ and $H_{\frac{1}{n}}(u_n) < 0$. Without loss of generality, let us assume that $\|\nabla u_n\|_{L^p(\Omega_\gamma)} = 1$. By Lemma 2.2, the sequence $\{u_n\}$ is uniformly bounded in $W^{1,p}(\Omega_\gamma)$. Hence, by Theorem 2.1, because the embedding operator

$$W^{1,p}(\Omega_\gamma) \hookrightarrow L^q(\Omega_\gamma), \quad 1 < q < p_\gamma^*,$$

is compact, there exists $u \in W^{1,p}(\Omega_\gamma)$ such that $u_n \rightharpoonup u$ weakly in $W^{1,p}(\Omega_\gamma)$ and $u_n \rightarrow u$ strongly in $L^q(\Omega_\gamma)$ and there exists $g \in L^q(\Omega_\gamma)$ such that $|u_n| \leq g$ a.e. in Ω_γ and $\nabla u_n \rightharpoonup \nabla u$ weakly in $L^p(\Omega_\gamma)$.

Since $|u_n| \leq g$ a.e. in Ω_γ and $u_n \rightarrow u$ a.e. in Ω_γ . Then,

$$\| |u_n|^{q-2} u_n \| \leq \| |u_n|^{q-1} \| \leq \| |g|^{q-1} \| \in L^{q'}(\Omega_\gamma).$$

So, by the Lebesgue Dominated Convergence Theorem (see, for example, [8]), it follows that

$$0 = \lim_{n \rightarrow \infty} \int_{\Omega_\gamma} |u_n|^{q-2} u_n dx = \int_{\Omega_\gamma} |u|^{q-2} u dx.$$

So, $\int_{\Omega_\gamma} |u|^{q-2} u dx = 0$. Due to $H_{\frac{1}{n}}(u_n) < 0$, we obtain

$$(2.4) \quad \|\nabla u_n\|_{L^p(\Omega_\gamma)}^p - (\lambda_{p,q}^p + \frac{1}{n}) \|u_n\|_{L^q(\Omega_\gamma)}^p < 0.$$

Since $\nabla u_n \rightharpoonup \nabla u$ weakly in $L^p(\Omega_\gamma)$, by the weak lower semicontinuity of norm, we have

$$\|\nabla u\|_{L^p(\Omega_\gamma)} \leq \liminf_{n \rightarrow \infty} \|\nabla u_n\|_{L^p(\Omega_\gamma)} = 1.$$

So, by passing to the limit in (2.4), we get

$$\lambda_{p,q} \geq \frac{\|\nabla u\|_{L^p(\Omega_\gamma)}^p}{\|u\|_{L^q(\Omega_\gamma)}^p}.$$

Therefore, by the definition of $\lambda_{p,q}$, we obtain

$$\lambda_{p,q} = \frac{\|\nabla u\|_{L^p(\Omega_\gamma)}^p}{\|u\|_{L^q(\Omega_\gamma)}^p}.$$

Now, since $H_\perp^n(u_n) < 0$ and $\|\nabla u_n\|_{L^p(\Omega_\gamma)} = 1$, we have

$$1 - (\lambda_{p,q}^p + \frac{1}{n}) \|u_n\|_{L^q(\Omega_\gamma)}^p < 0.$$

Letting $n \rightarrow \infty$, we get

$$\|u\|_{L^q(\Omega_\gamma)} \lambda_{p,q} \geq 1,$$

which gives $\lambda_{p,q} > 0$ and $u \neq 0$ a.e. in Ω_γ . \square

3. ESTIMATES OF NEUMANN (p, q) -EIGENVALUES

In this section we give estimates of Neumann (p, q) -eigenvalues in Hölder singular domains. The suggested method is based on Theorem 2.3 and on the composition operators theory on Sobolev spaces [29, 31, 32, 33].

3.1. Composition operators on Sobolev spaces. The seminormed Sobolev space $L^{1,p}(\Omega)$ in a domain $\Omega \subset \mathbb{R}^n$ is the space of all locally integrable weakly differentiable functions with the following seminorm:

$$\|f\|_{L^{1,p}(\Omega)} = \left(\int_\Omega |\nabla f(x)|^p dx \right)^{\frac{1}{p}}.$$

Let Ω and $\tilde{\Omega}$ be domains in the Euclidean space $\subset \mathbb{R}^n$. Then a homeomorphism $\varphi : \Omega \rightarrow \tilde{\Omega}$ belongs to the Sobolev class $W_{\text{loc}}^{1,p}(\Omega)$, $1 \leq p \leq \infty$, if its coordinate functions φ_j belong to $W_{\text{loc}}^{1,p}(\Omega)$, $j = 1, \dots, n$. In this case the formal Jacobi matrix $D\varphi(x) = \left(\frac{\partial \varphi_i}{\partial x_j}(x) \right)$, $i, j = 1, \dots, n$, and its determinant (Jacobian) $J(x, \varphi) = \det D\varphi(x)$ are well defined at almost all points $x \in \Omega$. The norm $|D\varphi(x)|$ of the matrix $D\varphi(x)$ is the norm of the corresponding linear operator $D\varphi(x) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ defined by the matrix $D\varphi(x)$.

Recall that a Sobolev homeomorphism $\varphi : \Omega \rightarrow \tilde{\Omega}$ of the class $W_{\text{loc}}^{1,1}(\Omega)$ has finite distortion [30] if

$$D\varphi(x) = 0 \text{ a.e. on the set } Z = \{x \in \Omega : |J(x, \varphi)| = 0\}.$$

The result of [29] gives the analytic description of composition operators on seminormed Sobolev spaces (see, also [31]) and asserts that

Theorem 3.1. [29] *Let $\varphi : \Omega \rightarrow \tilde{\Omega}$ be a homeomorphism between two domains Ω and $\tilde{\Omega} \subset \mathbb{R}^n$. Then φ induces by the composition rule $\varphi^*(f) = f \circ \varphi$ a bounded composition operator*

$$\varphi^* : L^{1,p}(\tilde{\Omega}) \rightarrow L^{1,s}(\Omega), \quad 1 \leq s < p < \infty,$$

if and only if $\varphi \in W_{\text{loc}}^{1,1}(\Omega)$, has finite distortion, and

$$(3.1) \quad K_{p,s}(\varphi; \Omega) = \left(\int_{\Omega} \left(\frac{|D\varphi(x)|^p}{|J(x, \varphi)|} \right)^{\frac{s}{p-s}} dx \right)^{\frac{p-s}{ps}} < \infty.$$

The homeomorphisms which satisfy conditions of Theorem 3.1 are called weak (p, s) -quasiconformal mappings [34].

3.2. Spectral estimates. Let $\Omega \subset \mathbb{R}^n$ be a bounded domain. Then Ω is called an (r, s) -Sobolev-Poincaré domain, $1 \leq r, s \leq \infty$, if there exists a constant $C < \infty$, such that for any function $f \in L^{1,s}(\Omega)$, the (r, s) -Sobolev-Poincaré inequality

$$\inf_{c \in \mathbb{R}} \|f - c\|_{L^r(\Omega)} \leq C \|\nabla f\|_{L^s(\Omega)}$$

holds. We denote by $B_{r,s}(\Omega)$ the best constant in this inequality.

The weak quasiconformal mappings permits us to "transfer" the Sobolev-Poincaré inequalities from one domain to another. In the work [17], the authors obtained the following result.

Theorem 3.2. *Let a bounded domain $\Omega \subset \mathbb{R}^n$ be a (r, s) -Sobolev-Poincaré domain, $1 < s \leq r < \infty$, and there exists a weak (p, s) -quasiconformal homeomorphism $\varphi : \Omega \rightarrow \tilde{\Omega}$ of a domain Ω onto a bounded domain $\tilde{\Omega}$, possesses the Luzin N -property (an image of a set of measure zero has measure zero) and such that*

$$M_{r,q}(\Omega) = \left(\int_{\Omega} |J(x, \varphi)|^{\frac{r}{r-q}} dx \right)^{\frac{r-q}{rq}} < \infty$$

for some $1 \leq q < r$. Then in the domain $\tilde{\Omega}$ the (q, p) -Sobolev-Poincaré inequality

$$\inf_{c \in \mathbb{R}} \left(\int_{\tilde{\Omega}} |f(y) - c|^q dy \right)^{\frac{1}{q}} \leq B_{q,p}(\tilde{\Omega}) \left(\int_{\tilde{\Omega}} |\nabla f(y)|^p dy \right)^{\frac{1}{p}}, \quad f \in W^{1,p}(\tilde{\Omega}),$$

holds and for $1 < s < p$, we have

$$B_{q,p}(\tilde{\Omega}) \leq K_{p,s}(\varphi; \Omega) M_{r,q}(\Omega) B_{r,s}(\Omega).$$

Here $B_{r,s}(\Omega)$ is the best constant in the (r, s) -Sobolev-Poincaré inequality in the domain Ω and $K_{p,s}(\varphi; \Omega)$ is as defined in (3.1).

Recall that Ω_γ are bounded domains with anisotropic Hölder γ -singularities (introduced in [12]):

$$\Omega_\gamma = \{x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n : 0 < x_n < 1, 0 < x_i < g_i(x_n), i = 1, 2, \dots, n-1\},$$

where $g_i(t) = t^{\gamma_i}$, $\gamma_i \geq 1$, $0 < t < 1$ are Hölder functions and for the function

$$G = \prod_{i=1}^{n-1} g_i \text{ denote by}$$

$$\gamma = \frac{\log G(t)}{\log t} + 1, \quad 0 < t < 1.$$

It is evident that $\gamma \geq n$. In the case $g_1 = g_2 = \dots = g_{n-1}$ we will say that domain Ω_γ is a domain with σ -Hölder singularity, $\sigma = (\gamma - 1)/(n - 1)$. For $g_1(t) = g_2(t) = \dots = g_{n-1}(t) = t$ we will use notation Ω_1 instead of Ω_γ .

Define the mapping $\varphi_a : \Omega_1 \rightarrow \Omega_\gamma$, $a > 0$, by

$$\varphi_a(x) = \left(\frac{x_1}{x_n} g_1^a(x_n), \dots, \frac{x_{n-1}}{x_n} g_{n-1}^a(x_n), x_n^a \right).$$

Let us formulate the following result from the work [17] in the refined form because of the very useful remark by Charles Fefferman.

Theorem 3.3. *Let $(n - p)/(\gamma - p) < a < p(n - s)/s(\gamma - p)$. Then the mapping $\varphi_a : \Omega_1 \rightarrow \Omega_\gamma$ be a weak (p, s) -quasiconformal mapping, $1 < s < p < \gamma$, from the Lipschitz convex domain Ω_1 onto the ‘‘cusp’’ domain Ω_γ with*

$$K_{p,s}(\varphi_a; \Omega_1) \leq a^{-\frac{1}{p}} \sqrt[n-1]{\sum_{i=1}^{n-1} (a\gamma_i - 1)^2 + n - 1 + a^2}.$$

By Theorem 2.1 the embedding operator

$$W^{1,p}(\Omega_\gamma) \hookrightarrow L^q(\Omega_\gamma)$$

is compact for any $1 < q < p_\gamma^*$ and by Theorem 2.3 the first non-trivial Neumann eigenvalue $\lambda_{p,q}(\Omega_\gamma)$ can be characterized as

$$\lambda_{p,q}(\Omega_\gamma) = \inf \left\{ \frac{\|\nabla u\|_{L^p(\Omega_\gamma)}^p}{\|u\|_{L^q(\Omega_\gamma)}^p} : u \in W^{1,p}(\Omega_\gamma) \setminus \{0\}, \int_{\Omega_\gamma} |u|^{q-2} u \, dx = 0 \right\}.$$

Furthermore, $\lambda_{p,q}(\Omega_\gamma)^{-\frac{1}{p}}$ is equal to the best constant $B_{q,p}(\Omega_\gamma)$ in the (q, p) -Sobolev-Poincaré inequality

$$\inf_{c \in \mathbb{R}} \left(\int_{\Omega_\gamma} |f(x) - c|^q \, dx \right)^{\frac{1}{q}} \leq B_{q,p}(\Omega_\gamma) \left(\int_{\Omega_\gamma} |\nabla f(x)|^p \, dx \right)^{\frac{1}{p}}, \quad f \in W^{1,p}(\Omega_\gamma).$$

We are ready to prove spectral estimates in cusp domains.

Theorem 3.4. *Let*

$$\Omega_\gamma := \{x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n : n \geq 3, 0 < x_n < 1, 0 < x_i < x_n^{\gamma_i}, i = 1, 2, \dots, n - 1\}$$

$\gamma_i \geq 1$, $\gamma := 1 + \sum_{i=1}^{n-1} \gamma_i$, be domains with anisotropic Hölder γ -singularities.

Then for $1 < s < p < \gamma$ and $1 < q < p_\gamma^*$, we have

$$\frac{1}{\lambda_{p,q}(\Omega_\gamma)} \leq \inf_{a \in I_a} a^{\frac{p}{q}-1} \left(\sum_{i=1}^{n-1} (a\gamma_i - 1)^2 + n - 1 + a^2 \right)^{\frac{p}{2}} B_{r,s}^p(\Omega_1),$$

where $I_a = (n/\gamma, p(n - s)/s(\gamma - p))$ and $s < n$ is chosen such that $p_\gamma^* < \frac{ns}{n-s}$. $B_{r,s}(\Omega_1)$ is the best constant in the (r, s) -Sobolev-Poincaré inequality in the domain Ω_1 , $1 < q < r < \infty$.

Proof. By Theorem 3.3 there exists a locally Lipschitz homeomorphism $\varphi_a : \Omega_1 \rightarrow \Omega_\gamma$, $(n - p)/(\gamma - p) < a < p(n - s)/s(\gamma - p)$,

$$\varphi_a(x) = \left(\frac{x_1}{x_n} g_1^a(x_n), \dots, \frac{x_{n-1}}{x_n} g_{n-1}^a(x_n), x_n^a \right)$$

which maps the convex Lipschitz domain Ω_1 onto the cusp domain Ω_γ , possesses the Luzin N -property, and it is a weak (p, s) -quasiconformal mapping, $1 < s < p < \gamma$.

Let us check conditions of Theorem 3.2. Since φ_a is a weak (p, s) -quasiconformal mapping, by Theorem 3.3, the constant $K_{p,s}(\varphi_a; \Omega_1)$ is finite and satisfy the estimate

$$(3.2) \quad K_{p,s}(\varphi_a; \Omega_1) \leq a^{-\frac{1}{p}} \sqrt[n-1]{\sum_{i=1}^{n-1} (a\gamma_i - 1)^2 + n - 1 + a^2}.$$

The domain Ω_1 is a Lipschitz domain and so is a (r, s) -Sobolev-Poincaré domain, i.e. $B_{r,s}(\Omega_1) < \infty$.

Let $1 < q < r < \infty$. Next we estimate the quantity $M_{r,q}(\Omega_1)$.

$$\begin{aligned} M_{r,q}(\Omega_1) &= \left(\int_{\Omega_1} |J(x, \varphi_a)|^{\frac{r}{r-q}} dx \right)^{\frac{r-q}{rq}} = a^{\frac{1}{q}} \left(\int_{\Omega_1} (x_n^{a\gamma-n})^{\frac{r}{r-q}} dx \right)^{\frac{r-q}{rq}} \\ &= a^{\frac{1}{q}} \left(\int_0^1 (x_n^{a\gamma-n})^{\frac{r}{r-q}} \left(\int_0^{x_n} dx_1 \dots \int_0^{x_n} dx_{n-1} \right) dx_n \right)^{\frac{r-q}{rq}} \\ &= a^{\frac{1}{q}} \left(\int_0^1 (x_n^{a\gamma-n})^{\frac{r}{r-q}} \cdot x_n^{n-1} dx_n \right)^{\frac{r-q}{rq}} < \infty, \end{aligned}$$

if

$$\frac{(a\gamma - n)r}{r - q} + n - 1 > -1, \text{ i.e. } a > \frac{nq}{\gamma r}.$$

Recall that $r > q$. Therefore, since $0 < x_n < 1$, we obtain

$$(3.3) \quad M_{r,q}(\Omega_1) \leq a^{\frac{1}{q}} \text{ if } a > \frac{n}{\gamma} > \frac{nq}{\gamma r}.$$

So, the conditions of Theorem 3.2 are fulfilled. Therefore, noting that $1 < q < p_\gamma^*$, we have $\lambda_{p,q}^{-1}(\Omega_\gamma) = B_{q,p}^p(\Omega_\gamma)$ and by Theorem 3.2, we obtain for $1 < s < p < \gamma$ the following estimate

$$\begin{aligned} \frac{1}{\lambda_{p,q}(\Omega_\gamma)} &\leq K_{p,s}^p(\varphi_a; \Omega_1) M_{r,q}^p(\Omega_1) B_{r,s}^p(\Omega_1) \\ &\leq a^{\frac{p}{q}-1} \left(\sum_{i=1}^{n-1} (a\gamma_i - 1)^2 + n - 1 + a^2 \right)^{\frac{p}{2}} B_{r,s}^p(\Omega_1), \end{aligned}$$

where $\max\{(n-p)/(\gamma-p), n/\gamma\} < a < p(n-s)/s(\gamma-p)$ and $B_{r,s}(\Omega_1)$ is the best constant in the (r, s) -Sobolev-Poincaré inequality in the domain Ω_1 . To deduce the last inequality above, we have also used the estimates (3.2) and (3.3). Finally remark, that $(n-p)/(\gamma-p) < n/\gamma$ and so $\max\{(n-p)/(\gamma-p), n/\gamma\} = n/\gamma$. \square

Remark 3.5. The estimate of the constant in the (r, s) -Sobolev-Poincaré inequality in the domain Ω_1 was obtained in [16]:

$$B_{r,s}(\Omega_1) \leq n \left(\frac{1-\delta}{1/n-\delta} \right)^{1-\delta} \omega_n^{1-\frac{1}{n}} \left(\frac{1}{(n+1)!} \right)^{\frac{1}{n}-\delta}, \quad \delta = \frac{1}{s} - \frac{1}{r} \geq 0.$$

The problem of exact value of constants in the (r, s) -Sobolev-Poincaré inequality in the case $s \neq r$ is a complicated open problem even in the case of the unit disk $\mathbb{D} \subset \mathbb{R}^2$.

Corollary 3.6. *Let us consider an application of Theorem 3.4 to the following spectral problem*

$$(3.4) \quad -\operatorname{div}(|\nabla u| \nabla u) = \lambda \|u\|_{L^2(\Omega_\gamma)} \cdot u \quad \text{in } \Omega_\gamma \subset \mathbb{R}^3,$$

with the Neumann boundary condition in an anisotropic Hölder γ -singular domains Ω_γ , where

$$\Omega_\gamma := \{x = (x_1, x_2, x_3) \in \mathbb{R}^3 : 0 < x_3 < 1, 0 < x_i < x_3^{\gamma_i}, i = 1, 2\},$$

$\gamma_i \geq 1$, $\gamma := 1 + \gamma_1 + \gamma_2$. Then the first non-trivial Neumann $(3, 2)$ -eigenvalue of the spectral problem (3.4) satisfies the estimate:

$$\frac{1}{\lambda_{3,2}(\Omega_\gamma)} \leq \inf_{\frac{3}{\gamma} < a < \frac{3(3-s)}{s(\gamma-3)}} \sqrt{a} \left((a\gamma_1 - 1)^2 + (a\gamma_2 - 1)^2 + 2 + a^2 \right)^{\frac{3}{2}} B_{r,s}^3(\Omega_1).$$

Let us take $r = 5/2$, $s = 3/2$ then by [16]

$$B_{\frac{5}{2}, \frac{3}{2}}^3(\Omega_1) \leq 3 \cdot 11^{\frac{11}{15}} \left(\frac{4\pi}{3} \right)^{\frac{2}{3}} \left(\frac{1}{24} \right)^{\frac{1}{15}} \approx 12\pi.$$

If $3 < \gamma < 6$ we can take $a = 1$ and so in this case

$$\frac{1}{\lambda_{3,2}(\Omega_\gamma)} \leq (12\pi)^3 \left((\gamma_1 - 1)^2 + (\gamma_2 - 1)^2 + 3 \right)^{\frac{3}{2}}.$$

4. EXISTENCE AND REGULARITY RESULTS

Throughout this section, we assume that $1 < p < \gamma$ and $q = 2$ unless otherwise stated. Let

$$X := \left\{ u \in W^{1,p}(\Omega_\gamma) : \int_{\Omega_\gamma} u \, dx = 0 \right\},$$

By Theorem 2.1, we endow the norm $\|\cdot\|_X$ on X defined by

$$(4.1) \quad \|u\|_X = \left(\int_{\Omega_\gamma} |\nabla u|^p \, dx \right)^{\frac{1}{p}}.$$

The Lebesgue space $Y := L^2(\Omega_\gamma)$ be endowed with the norm

$$(4.2) \quad \|u\|_Y := \left(\int_{\Omega_\gamma} |u|^2 \, dx \right)^{\frac{1}{2}}.$$

4.1. Regularity results. Let us formulate the regularity results for the Neumann (p, q) -eigenvalue problem.

Theorem 4.1. *Let $1 < p < \gamma$. Then the following properties hold:*

(a) *There exists a sequence $\{w_n\}_{n \in \mathbb{N}} \subset X \cap Y$ such that $\|w_n\|_Y = 1$ and for every $v \in X$, we have*

$$(4.3) \quad \int_{\Omega_\gamma} |\nabla w_{n+1}|^{p-2} \nabla w_{n+1} \nabla v \, dx = \mu_n \int_{\Omega_\gamma} w_n v \, dx,$$

where

$$\mu_n \geq \lambda := \inf \left\{ \int_{\Omega_\gamma} |\nabla u|^p \, dx : u \in X \cap Y, \|u\|_Y = 1 \right\}.$$

(b) Moreover, the sequences $\{\mu_n\}_{n \in \mathbb{N}}$ and $\{\|w_{n+1}\|_X^p\}_{n \in \mathbb{N}}$ given by (4.3) are non-increasing and converge to the same limit μ , which is bounded below by λ . Further, there exists a subsequence $\{n_j\}_{j \in \mathbb{N}}$ such that both $\{w_{n_j}\}_{j \in \mathbb{N}}$ and $\{w_{n_{j+1}}\}_{j \in \mathbb{N}}$ converges in X to the same limit $w \in X \cap Y$ with $\|w\|_Y = 1$ and (μ, w) is an eigenpair of (2.1).

Theorem 4.2. *Let $1 < p < \gamma$ and $q = 2$. Suppose $\{u_n\}_{n \in \mathbb{N}} \subset X \cap S_Y$ is a minimizing sequence for λ , that is $\|u_n\|_Y = 1$ and $\|u_n\|_X^p \rightarrow \lambda$. Then there exists a subsequence $\{u_{n_j}\}_{j \in \mathbb{N}}$ which converges weakly in X to $u \in X \cap S_Y$ such that $\lambda = \|u\|_X^p$. Moreover, u is an eigenfunction of (1.1) corresponding to λ and its associated eigenfunctions are precisely the scalar multiple of those vectors at which λ is reached.*

Moreover, we have the following regularity results.

Theorem 4.3. *Let $1 < p < \gamma$ and $q = 2$. Assume that $\lambda > 0$ is an eigenvalue of the problem (1.1) and $u \in X \setminus \{0\}$ is a corresponding eigenfunction. Then*

(i) $u \in L^\infty(\Omega_\gamma)$.

(ii) Moreover, if $u \in X \setminus \{0\}$ is nonnegative in Ω_γ , then $u > 0$ in Ω_γ . Further, for every $\omega \Subset \Omega_\gamma$ there exists a positive constant c depending on ω such that $u \geq c > 0$ in ω .

Remark 4.4. Theorems 4.1–4.3 are correct in the case of Lipschitz domains $\Omega \subset \mathbb{R}^n$. In this case $p_n^* = np/(n-p)$, where $1 < p < n$.

4.2. Operators associated to eigenvalue problems. Let us denote by X^* and Y^* denotes the dual of X and Y respectively. Let us define the operator $A : X \rightarrow X^*$ by

$$(4.4) \quad \langle A(v), w \rangle = \int_{\Omega_\gamma} |\nabla v|^{p-2} \nabla v \nabla w \, dx, \quad \forall v, w \in X$$

and $B : Y \rightarrow Y^*$ by

$$(4.5) \quad \langle B(v), w \rangle = \int_{\Omega_\gamma} vw \, dx, \quad \forall v, w \in Y.$$

First we state some useful results. The following result from [3, Theorem 9.14] will be useful for us.

Theorem 4.5. *Let V be a real separable reflexive Banach space and V^* be the dual of V . Assume that $A : V \rightarrow V^*$ is a bounded, continuous, coercive and monotone operator. Then A is surjective, i.e., given any $f \in V^*$, there exists $u \in V$ such that $A(u) = f$. If A is strictly monotone, then A is also injective.*

Moreover, for the following algebraic inequality, see [5, Lemma 2.1].

Lemma 4.6. *Let $1 < p < \infty$. Then for any $a, b \in \mathbb{R}^N$, there exists a positive constant $C = C(p)$ such that*

$$(4.6) \quad (|a|^{p-2}a - |b|^{p-2}b, a - b) \geq C(|a| + |b|)^{p-2}|a - b|^2.$$

Next, we prove the following result.

Lemma 4.7. (i) *The operators A defined by (4.4) and B defined by (4.5) are continuous. (ii) Moreover, A is bounded, coercive and monotone.*

Proof. (i) **Continuity:** We only prove the continuity of A , since the continuity of B would follow similarly. To this end, suppose $v_n \in X$ such that $v_n \rightarrow v$ in the norm of X . Thus, up to a subsequence $\nabla v_n \rightarrow \nabla v$ in Ω_γ . We observe that

$$(4.7) \quad \|\nabla v_n\|_{L^{\frac{p}{p-1}}(\Omega_\gamma)}^{p-1} \leq c,$$

for some constant $c > 0$, which is independent of n . Thus, up to a subsequence, we have

$$(4.8) \quad |\nabla v_n|^{p-2} \nabla v_n \rightharpoonup |\nabla v|^{p-2} \nabla v \text{ weakly in } L^{p'}(\Omega_\gamma).$$

Since, the weak limit is independent of the choice of the subsequence, as a consequence of (4.8), we have

$$\lim_{n \rightarrow \infty} \langle Av_n, w \rangle = \langle Av, w \rangle$$

for every $w \in X$. Thus A is continuous.

(ii) **Boundedness:** Using the estimate (4.9), we have

$$\|Av\|_{X^*} = \sup_{\|w\|_X \leq 1} |\langle Av, w \rangle| \leq \|v\|_X^{p-1} \|w\|_X \leq \|v\|_X^{p-1}.$$

Thus, A is bounded.

Coercivity: We observe that

$$\langle Av, v \rangle = \|v\|_X^p.$$

Since $p > 1$, we have A is coercive.

Monotonicity: Using Lemma 4.6, it follows that there exists a constant $C = C(p) > 0$ such that for every $v, w \in X$, we have

$$\begin{aligned} \langle Av - Aw, v - w \rangle &= \int_{\Omega_\gamma} (|\nabla v|^{p-2} \nabla v - |\nabla w|^{p-2} \nabla w, \nabla(v - w)) \, dx \\ &= \int_{\Omega_\gamma} (|\nabla v|^{p-2} \nabla v - |\nabla w|^{p-2} \nabla w, \nabla v - \nabla w) \, dx \\ &\geq C(p) \int_{\Omega_\gamma} (|\nabla v| + |\nabla w|)^{p-2} |\nabla v - \nabla w|^2 \, dx \geq 0. \end{aligned}$$

Thus, A is a monotone operator. \square

Lemma 4.8. *The operators A defined by (4.4) and B defined by (4.5) satisfy the following properties:*

$$(H_1) \quad A(tv) = |t|^{p-2} t A(v) \quad \forall t \in \mathbb{R} \quad \text{and} \quad \forall v \in X.$$

$$(H_2) \quad B(tv) = tB(v) \quad \forall t \in \mathbb{R} \quad \text{and} \quad \forall v \in Y.$$

$$(H_3) \quad \langle A(v), w \rangle \leq \|v\|_X^{p-1} \|w\|_X \text{ for all } v, w \in X, \text{ where the equality holds if and only if } v = 0 \text{ or } w = 0 \text{ or } v = tw \text{ for some } t > 0.$$

$$(H_4) \quad \langle B(v), w \rangle \leq \|v\|_Y^{p-1} \|w\|_Y \text{ for all } v, w \in Y, \text{ where the equality holds if and only if } v = 0 \text{ or } w = 0 \text{ or } v = tw \text{ for some } t \geq 0.$$

(H₅) *For every $w \in Y \setminus \{0\}$ there exists $u \in X \setminus \{0\}$ such that*

$$\langle A(u), v \rangle = \langle B(w), v \rangle \quad \forall v \in X.$$

Proof. (H_1) Follows by the definition of A in (4.4).

(H_2) Follows by the definition of B in (4.5).

(H_3) First using Cauchy-Schwartz inequality and then by Hölder's inequality with exponents $\frac{p}{p-1}$ and p , for every $v, w \in X$, we obtain

$$(4.9) \quad \begin{aligned} \langle Av, w \rangle &= \int_{\Omega_\gamma} |\nabla v|^{p-2} \nabla v \nabla w \, dx \leq \int_{\Omega_\gamma} |\nabla v|^{p-1} |\nabla w| \, dx \\ &\leq \left(\int_{\Omega_\gamma} |\nabla v|^p \, dx \right)^{\frac{p-1}{p}} \left(\int_{\Omega_\gamma} |\nabla w|^p \, dx \right)^{\frac{1}{p}} = \|v\|_X^{p-1} \|w\|_X. \end{aligned}$$

Let the equality

$$(4.10) \quad \langle A(v), w \rangle = \|v\|_X^{p-1} \|w\|_X$$

holds for every $v, w \in X$. We claim that either $v = 0$ or $w = 0$ or $v = tw$ for some constant $t > 0$. Indeed, if $v = 0$ or $w = 0$, this is trivial. Therefore, we assume $v \neq 0$ and $w \neq 0$ and prove that $v = tw$ for some constant $t > 0$. By the estimate (4.9) if the equality (4.10) holds, then we have

$$(4.11) \quad \langle A(v), w \rangle = \int_{\Omega_\gamma} |\nabla v|^{p-1} |\nabla w| \, dx,$$

which gives us

$$(4.12) \quad \int_{\Omega_\gamma} f(x) \, dx = 0,$$

where

$$f(x) = |\nabla v|^{p-1} |\nabla w| - |\nabla v|^{p-2} \nabla v \nabla w.$$

By Cauchy-Schwartz inequality, we have $f \geq 0$ in Ω_γ . Hence using this fact in (4.12), we have $f = 0$ in Ω_γ , which reduces to

$$(4.13) \quad |\nabla v|^{p-1} \nabla w = |\nabla v|^{p-2} \nabla v \nabla w \text{ in } \Omega_\gamma,$$

which gives $\nabla v(x) = c(x) \nabla w(x)$ for some $c(x) \geq 0$.

On the otherhand, if the equality (4.10) holds, then by the estimate (4.9) we have

$$(4.14) \quad f_1 = f_2,$$

where

$$f_1 = \int_{\Omega_\gamma} |\nabla v|^{p-1} \nabla w \, dx, \quad f_2 = \left(\int_{\Omega_\gamma} |\nabla v|^p \, dx \right)^{\frac{p-1}{p}} \left(\int_{\Omega_\gamma} |\nabla w|^p \, dx \right)^{\frac{1}{p}}.$$

Thus, we have $|\nabla v|(x) = d|\nabla w|(x)$ a.e. $x \in \Omega_\gamma$ for some constant $d > 0$. Thus, we obtain $c(x) = d$. As a consequence, we get $\nabla v = d\nabla w$ a.e. in Ω_γ and therefore, we deduce that $\|v - dw\|_X = 0$, which gives $v = dw$ a.e. in Ω_γ . Hence, the property (H_3) is verified.

(H_4) The hypothesis (H_4) can be verified similarly.

(H_5) We observe that X is a separable and reflexive Banach space. By Lemma 4.7, the operator $A : X \rightarrow X^*$ is bounded, coercive and monotone.

By the Sobolev embedding theorem, we have X is continuously embedded in Y . Therefore, $B(w) \in X^*$ for every $w \in Y \setminus \{0\}$.

Hence, by Theorem 4.5, for every $w \in Y \setminus \{0\}$, there exists $u \in X \setminus \{0\}$ such that

$$\langle A(u), v \rangle = \langle B(w), v \rangle \quad \forall v \in X.$$

Hence the property (H_5) holds. This completes the proof. \square

4.3. Proof of the regularity results: Proof of Theorem 4.1:

(a) First we recall the definition of the operators $A : X \rightarrow X^*$ from (4.4) and $B : Y \rightarrow Y^*$ from (4.5) respectively. Then, noting the property (H_5) from Lemma 4.8 and proceeding along the lines of the proof in [7, page 579 and pages 584–585], the result follows.

(b) We observe that X is uniformly convex Banach space and by the Sobolev embedding theorem [12, 14], X is compactly embedded in Y . Next, using Lemma 4.7-(i), the operators $A : X \rightarrow X^*$ and $B : Y \rightarrow Y^*$ are continuous and by Lemma 4.8, the properties $(H_1) - (H_5)$ holds. Noting these facts, the result follows from [7, page 579, Theorem 1]. \square

Proof of Theorem 4.2: The proof follows due to the same reasoning as in the proof of Theorem 4.1-(b) except that here we apply [7, page 583, Proposition 2] in place of [7, page 579, Theorem 1].

Proof of Theorem 4.3:

(i) Due to the homogeneity of the equation (1.1), without loss of generality, we assume that $\|u\|_Y = 1$. Let $k \geq 1$ and set $A(k) := \{x \in \Omega_\gamma : u(x) > k\}$. Choosing $v = (u - k)^+$ as a test function in (2.2), we obtain

$$(4.15) \quad \int_{A(k)} |\nabla u|^p dx = \lambda \int_{A(k)} u(u - k) dx \\ \leq \lambda \int_{A(k)} |u|(u - k) dx \leq \lambda \int_{A(k)} |u|(u - k) dx.$$

Noting (4.15) and proceeding along the lines of the proof of [10, Theorem 3.4, Pages 7–8], the result follows.

(ii) By [28, Theorem 1.2], the result follows.

Acknowledgements.

The authors thank Pier Domenico Lamberti for very useful and fruitful discussions and remarks on the topic.

The second author was supported by RSF Grant No. 23-21-00080.

REFERENCES

- [1] A. Anane, N. Tsouli, On the second eigenvalue of the p -Laplacian, In Nonlinear partial differential equations (Fès, 1994), volume 343 of Pitman Res. Notes Math. Ser., pages 1–9. Longman, Harlow, 1996.
- [2] A. Anane, Simplicité et isolation de la première valeur propre du p -laplacien avec poids, C. R. Acad. Sci. Paris Sér. I Math., 305(16) (1987), 725–728.
- [3] P. G. Ciarlet, Linear and nonlinear functional analysis with applications, Society for Industrial and Applied Mathematics, Philadelphia, PA, 2013.
- [4] G. Croce, A. Henrot, G. Pisante, Corrigendum to "An isoperimetric inequality for a nonlinear eigenvalue problem" [Ann. I. H. Poincaré - AN 29(1) (2012) 21–34]. Ann. Inst. H. Poincaré Anal. Non Linéaire, 32(2) (2015), 485–487.

- [5] L. Damascelli, Comparison theorems for some quasilinear degenerate elliptic operators and applications to symmetry and monotonicity results, *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 15(4) (1998), 493–516.
- [6] P. Drábek, A. Kufner, F. Nicolosi, Quasilinear elliptic equations with degenerations and singularities, volume 5 of *De Gruyter Series in Nonlinear Analysis and Applications*. Walter de Gruyter & Co., Berlin, 1997.
- [7] G. Ercole, Solving an abstract nonlinear eigenvalue problem by the inverse iteration method, *Bull. Braz. Math. Soc. (N.S.)*, 49(3) (2018), 577–591.
- [8] H. Federer, *Geometric measure theory*, Springer Verlag, Berlin, (1969).
- [9] G. Franzina, P. D. Lamberti, Existence and uniqueness for a p -Laplacian nonlinear eigenvalue problem, *Electron. J. Differential Equations*, 10 (2010), 10pp.
- [10] P. Garain, A. Ukhlov, On (p, q) -eigenvalues of subelliptic operators on nilpotent Lie groups. *Trans. A. Razmadze Math. Inst.*, 176 (2022), 207–216.
- [11] J. P. García Azorero and I. Peral Alonso. Existence and nonuniqueness for the p -Laplacian: nonlinear eigenvalues. *Comm. Partial Differential Equations*, 12(12) (1987), 1389–1430.
- [12] V. Gol'dshtein, L. Gurov, Applications of change of variables operators for exact embedding theorems, *Integral Equ. Oper. Theory*, 19 (1994), 1–24.
- [13] V. Gol'dshtein, V. Pchelintsev, A. Ukhlov, On the first eigenvalue of the degenerate p -Laplace operator in non-convex domains, *Integral Equ. Oper. Theory*, 90 (2018), 21 pp.
- [14] V. Gol'dshtein, A. Ukhlov, Weighted Sobolev spaces and embedding theorems, *Trans. Amer. Math. Soc.*, 361(7) (2009), 3829–3850.
- [15] V. Gol'dshtein, A. Ukhlov, On the first Eigenvalues of Free Vibrating Membranes in Conformal Regular Domains. *Arch. Rational Mech. Anal.* 221(2) (2016), 893–915.
- [16] V. Gol'dshtein, A. Ukhlov, The spectral estimates for the Neumann-Laplace operator in space domains. *Adv. in Math.*, 315 (2017), 166–193.
- [17] V. Gol'dshtein, A. Ukhlov, Composition Operators on Sobolev Spaces and Neumann Eigenvalues. *Complex Anal. Oper. Theory*, 13 (2019), 2781–2798.
- [18] A. Greco, M. Lucia, Laplacian eigenvalues for mean zero functions with constant Dirichlet data. *Forum Math.*, 20(5) (2008), 763–782.
- [19] J. Heinonen, T. Kilpelinen, O. Martio, *Nonlinear potential theory of degenerate elliptic equations*, Clarendon Press. Oxford, New York, Tokio (1993).
- [20] L. Esposito, C. Nitsch, C. Trombetti, Best constants in Poincaré inequalities for convex domains, *J. Convex Anal.*, 20 (2013), 253–264.
- [21] An Lê, Eigenvalue problems for the p -Laplacian, *Nonlinear Anal.*, 64(5) (2006), 1057–1099.
- [22] P. Lindqvist, Addendum: "On the equation $\operatorname{div}(|\nabla u|^{p-2}\nabla u) + \lambda|u|^{p-2}u = 0$ ", [*Proc. Amer. Math. Soc.* 109(1) (1990), 157–164], *Proc. Amer. Math. Soc.*, 116(2) (1992), 583–584.
- [23] V. Maz'ya, *Sobolev Spaces with applications to elliptic partial differential equations*, Springer, Berlin, 2011.
- [24] V. G. Maz'ya, V. P. Havin, Non-linear potential theory, *Russian Math. Surveys*, 27 (1972), 71–148.
- [25] A. I. Nazarov, On symmetry and asymmetry in a problem of shape optimization, [arXiv:1208.3640](https://arxiv.org/abs/1208.3640).
- [26] L. E. Payne, H. F. Weinberger, An optimal Poincaré inequality for convex domains, *Arch. Rat. Mech. Anal.*, 5 (1960), 286–292.
- [27] G. Pólya, G. Szegő, *Isoperimetric Inequalities in Mathematical Physics*, Princeton University Press, 1951.
- [28] N. S. Trudinger, On Harnack type inequalities and their application to quasilinear elliptic equations, *Comm. Pure Appl. Math.*, 20 (1967), 721–747.
- [29] A. Ukhlov, On mappings, which induce embeddings of Sobolev spaces, *Siberian Math. J.* 34 (1993), 185–192.
- [30] S. K. Vodop'yanov, V. M. Gol'dstein, Yu. G. Reshetnyak, On geometric properties of functions with generalized first derivatives, *Uspekhi Mat. Nauk*, 34 (1979), 17–65.
- [31] S. K. Vodop'yanov, A. D. Ukhlov, Superposition operators in Sobolev spaces, *Russian Mathematics: Izvestiya VUZ* 46 (2002), 11–33.
- [32] S. K. Vodop'yanov, A. D. Ukhlov, Set functions and its applications in the theory of Lebesgue and Sobolev spaces. *Siberian Adv. Math.*, 14 (2004), 1–48.
- [33] S. K. Vodop'yanov, A. D. Ukhlov, Set functions and their applications in the theory of Lebesgue and Sobolev spaces, *Siberian Adv. in Math*, 15 (2005), 91–125.

- [34] S. K. Vodop'yanov, A. D. Ukhlov, Sobolev spaces and (P, Q) -quasiconformal mappings of Carnot groups, *Siberian Math. J.* 39 (1998), 665–682.

Department of Mathematics, Indian Institute of Technology Indore, Khandwa Road, Simrol, Indore 453552, India
E-mail address: pgarain92@gmail.com

Regional Scientific and Educational Mathematical Center, Tomsk State University, 634050 Tomsk, Lenin Ave. 36, Russia
E-mail address: vpchelintsev@vtomske.ru

Department of Mathematics, Ben-Gurion University of the Negev, P.O.Box 653, Beer Sheva, 8410501, Israel
E-mail address: ukhlov@math.bgu.ac.il