

Fourier transform from the symmetric square representation of PGL_2 and SL_2

G rard Laumon

Universit  Paris-Sud et CNRS
gerard.laumon@u-psud.fr

Emmanuel Letellier

Universit  Paris Cit , IMJ-PRG, CNRS
emmanuel.letellier@imj-prg.fr

January 27, 2023

Abstract

Let G be a connected reductive group over $\overline{\mathbb{F}}_q$ and let $\rho^\vee : G^\vee \rightarrow \mathrm{GL}_n$ be an algebraic representation of the dual group G^\vee . Assuming that G and ρ^\vee are defined over \mathbb{F}_q , Braverman and Kazhdan defined an operator on the space $C(G(\mathbb{F}_q))$ of complex valued functions on $G(\mathbb{F}_q)$. In this paper we are interested in the case where G is either SL_2 or PGL_2 and ρ^\vee is the symmetric square representation of G^\vee . We construct a natural $G \times G$ -equivariant embedding $G \hookrightarrow \mathcal{G} = \mathcal{G}_\rho$ and an involutive operator (Fourier transform) $\mathcal{F}^\mathcal{G}$ on the space of functions $C(\mathcal{G}(\mathbb{F}_q))$ that extends Braverman-Kazhdan’s operator.

Contents

| | | |
|----------|--|----------|
| 1 | Introduction | 2 |
| 1.1 | Motivations | 2 |
| 1.2 | Braverman-Kazhdan-Fourier operators on G | 3 |
| 1.3 | The problem | 5 |
| 1.4 | The main results | 5 |
| | | |
| 2 | Braverman-Kazhdan kernels | 7 |
| 2.1 | Generalities for tori | 7 |
| 2.2 | The symmetric square representations | 7 |
| 2.3 | Explicit computation : The case $G = \mathrm{SL}_2$ | 9 |
| 2.4 | Explicit computation : The case $G = \mathrm{PGL}_2$ | 11 |
| 2.5 | Explicit computation : The case $G = \mathrm{GL}_2$ | 12 |

| | | |
|----------|---|-----------|
| 3 | Extending the BK-Fourier kernel: The case $G = \mathrm{SL}_2, \mathrm{PGL}_2$ | 14 |
| 3.1 | Preliminaries | 14 |
| 3.2 | The tori case | 18 |
| 3.3 | Main result | 20 |
| 3.4 | Fourier transforms on quadratic spaces | 21 |
| 3.5 | Proof of Theorem 3.3.1 | 24 |
| 3.6 | Remarks on the torus case | 25 |
| 4 | Extending the BK-Fourier kernel: The case $G = \mathrm{GL}_2$ | 26 |
| 4.1 | The stack \mathcal{G} | 27 |
| 4.2 | The BK-Fourier kernels $\phi^{\mathcal{T}}, \phi^{\mathcal{T}\sigma}$ | 27 |
| 4.3 | The BK-Fourier kernel $\phi^{\mathcal{G}}$ | 29 |
| 5 | The characteristic 2 case | 30 |
| 5.1 | BK-Fourier operator on G | 30 |
| 5.2 | Extending the BK-Fourier operator | 31 |

1 Introduction

1.1 Motivations

Let \mathbb{A} be the ring of adèles of a global field F and let $\psi : \mathbb{A}/F \rightarrow \mathbb{C}^\times$ be a non-trivial additive character. We can define the Fourier transform of a locally constant function $f : \mathfrak{gl}_n(\mathbb{A}) \rightarrow \mathbb{C}$ with compact support as

$$\mathcal{F}^{\mathfrak{gl}_n}(f)(x) = \int_{\mathfrak{gl}_n(\mathbb{A})} \psi(\mathrm{Tr}(xy))f(y)dy.$$

Tate, for $n = 1$ [10], and Godement-Jacquet for arbitrary n [5] proved that this Fourier transform provides functional equations between L-functions. The involutivity property of $\mathcal{F}^{\mathfrak{gl}_n}$ is an important ingredient.

Given now a connected reductive group G together with a representation $\rho^\vee : G^\vee \rightarrow \mathrm{GL}_n$, Braverman-Kazhdan [1], Lafforgue [6] and more recently Ngô, considered a Fourier operator (defined spectrally) on a certain space of functions on $G(\mathbb{A})$. This Fourier operator is not involutive and one of the challenge would be to extend this operator to an (involutive) Fourier transform in order to extend Godement-Jacquet's work to L-functions defined from ρ^\vee .

The problem of defining a Fourier transform relatively to ρ^\vee makes also sense over finite fields and the aim of this paper is to construct such a Fourier transform in the simplest non-trivial

case, namely when ρ^\vee is the symmetric power representation of SL_2 or PGL_2 . We now explain this in details.

Let G be a connected reductive algebraic group with an \mathbb{F}_q -structure and associated geometric Frobenius $F : G \rightarrow G$. In this paper, all geometric Frobenius will be denoted by F and relative to q . Denote by G^\vee the dual group of G (in the sense of Deligne-Lusztig), a connected reductive algebraic group defined over \mathbb{F}_q . Lusztig has defined a partition of the set $\widehat{G^F}$ of all irreducible $\overline{\mathbb{Q}_\ell}$ -characters of G^F (with ℓ a prime not dividing q) whose parts, called *Lusztig series*, are indexed by the set of F -stable semisimple conjugacy classes of G^\vee . We then denote by $\mathrm{LS}(G)$ the set of Lusztig series of G^F .

Let us equip GL_n with its standard Frobenius and assume given a morphism $\rho^\vee : G^\vee \rightarrow \mathrm{GL}_n$ which commutes with Frobenius. It induces a map between sets of F -stable semisimple conjugacy classes and so a transfert map

$$t_\rho : \mathrm{LS}(G) \rightarrow \mathrm{LS}(\mathrm{GL}_n)$$

between Lusztig series (see [8, §3.6] for more details). This is the analogue of Langlands functoriality for finite fields.

1.2 Braverman-Kazhdan-Fourier operators on G

For a finite set X , let $C(X)$ denotes the $\overline{\mathbb{Q}_\ell}$ -space of all functions $X \rightarrow \overline{\mathbb{Q}_\ell}$.

Standard Fourier operator on $C(\mathrm{GL}_n^F)$

Fix a non-trivial additive character $\psi : \mathbb{F}_q \rightarrow \overline{\mathbb{Q}_\ell}^\times$ and denote by Tr the trace map on matrices. We have a natural Fourier transform $\mathcal{F}^{\mathrm{gl}_n} : C(\mathrm{gl}_n^F) \rightarrow C(\mathrm{gl}_n^F)$ defined by

$$\mathcal{F}^{\mathrm{gl}_n}(f)(y) = \sum_{x \in \mathrm{gl}_n^F} \psi(\mathrm{Tr}(yx))f(x),$$

for any $f \in C(\mathrm{gl}_n^F)$. The above Fourier transform restricts to an operator

$$\mathcal{F}^{\mathrm{GL}_n} : C(\mathrm{GL}_n^F) \rightarrow C(\mathrm{GL}_n^F)$$

(which unlike $\mathcal{F}^{\mathrm{gl}_n}$ is not involutive) defined by

$$\mathcal{F}^{\mathrm{GL}_n}(f)(h) = \sum_{g \in \mathrm{GL}_n^F} \psi(\mathrm{Tr}(hg))f(g).$$

The above operator defines a function $\gamma^{\mathrm{GL}_n} : \widehat{\mathrm{GL}_n^F} \rightarrow \overline{\mathbb{Q}_\ell}$ which is constant on Lusztig series and which satisfies

$$\mathcal{F}^{\mathrm{GL}_n}(\chi) = \gamma^{\mathrm{GL}_n}(\chi)\chi^*$$

for any irreducible character χ of GL_n^F where χ^* denotes the dual character of χ (see [8, §3.1] for more details).

BK-Fourier operator on $C(G^F)$

We transfer the gamma function γ^{GL_n} to a gamma function $\gamma_\rho^G : \widehat{G^F} \rightarrow \overline{\mathbb{Q}_\ell}$ by the formula

$$\gamma_\rho^G := c_G c_{\mathrm{GL}_n}^{-1} \gamma^{\mathrm{GL}_n} \circ \mathfrak{t}_\rho, \quad (1.1)$$

with

$$c_G := (-1)^{\mathbb{F}_q - \mathrm{rank}(G)} q^{\nu(G)}$$

where $\nu(G)$ is the number of positive roots of G .

Following Braverman-Kazhdan [2], define a central function $\phi_\rho^G : G^F \rightarrow \overline{\mathbb{Q}_\ell}$, called BK-Fourier kernel, by the formula

$$\phi_\rho^G := \sum_{\chi \in \widehat{G^F}} \chi(1) \gamma_\rho^G(\chi) \chi^*.$$

The BK-Fourier operator $\mathcal{F}_\rho^G : C(G^F) \rightarrow C(G^F)$ is defined by

$$\mathcal{F}_\rho^G(f)(h) = \sum_{g \in G^F} \phi_\rho^G(hg) f(g).$$

Notice that we can recover the gamma function γ_ρ^G by the formula

$$\mathcal{F}_\rho^G(\pi) = \gamma_\rho^G(\pi) \pi^*$$

for any irreducible character π (see [8, §3.5] for more details).

This is the spectral definition of \mathcal{F}_ρ^G due to Braverman-Kazhdan. They also gave an explicit conjectural formula for ϕ_ρ^G which they prove for $G = \mathrm{GL}_m$ and which is proved in full generality in [8] (see also [4] [3] for a more geometrical proof).

1.3 The problem

A crucial property of Fourier transforms is the involutivity property. While $\mathcal{F}^{\text{GL}_n}$ arises as the restriction of the Fourier transform $\mathcal{F}^{\text{gl}_n}$, we can ask ourselves whether the operator \mathcal{F}_ρ^G is also the restriction of an involutive operator. More precisely,

Question : is there an open $G \times G$ -equivariant embedding $G \hookrightarrow \mathcal{G}_\rho$ defined over \mathbb{F}_q (where $G \times G$ acts by right and left multiplication on G) and an involutive operator $C(\mathcal{G}_\rho^F) \rightarrow C(\mathcal{G}_\rho^F)$ restricting to \mathcal{F}_ρ^G on functions on G^F ?

This question has a positive answer in some simple cases. For instance, if $\rho^\vee : \text{SL}_n \hookrightarrow \text{GL}_n$ is the inclusion then there exists a dual morphism $\rho : \text{GL}_n \rightarrow \text{PGL}_n$ which is the natural quotient [8, §3.5]. The BK-Fourier kernel ϕ_ρ^G is then just $\rho_!(\psi \circ \text{Tr})$. We can then take the quotient stack $[\text{gl}_n/\text{Ker}(\rho)] = [\text{gl}_n/\text{GL}_1]$ as \mathcal{G}_ρ and consider on it the kernel $\bar{\rho}_!(\psi \circ \text{Tr})$ where $\bar{\rho} : \text{gl}_n \rightarrow [\text{gl}_n/\text{GL}_1]$ is the quotient. This defines an involutive Fourier transform on the space of functions $C(\mathcal{G}_\rho^F)$ which extends \mathcal{F}_ρ^G .

We will see in this paper, that in fact \mathcal{G}_ρ may not be not unique (see §3.6).

1.4 The main results

Here we mention only the results in odd characteristic (the characteristic 2 case is much simpler and is treated independently in §5). Let G be either SL_2 or PGL_2 equipped with the standard Frobenius F and consider the symmetric square representation $\rho^\vee : G^\vee \rightarrow \text{GL}_3$ (see §2.2).

We will use the following identification for G

$$\begin{aligned} G &= \{(x, \alpha, b) \in \text{GL}_2 \times \text{GL}_1 \times \text{GL}_1 \mid \det(x) = \alpha^2\} / \text{GL}_1 \times \text{GL}_1 && \text{if } G = \text{SL}_2 \\ G &= \{(x, \alpha, b) \in \text{GL}_2 \times \text{GL}_1 \times \text{GL}_1 \mid \det(x) = \alpha^2\} / \text{GL}_1 \times \mu_2 \times \text{GL}_1 && \text{if } G = \text{PGL}_2 \end{aligned}$$

where μ_2 acts on the second coordinate by $\alpha \mapsto -\alpha$ and $\text{GL}_1 \times \text{GL}_1$ acts by $(s, t) \cdot (x, \alpha, b) = (sx, s\alpha, tb)$.

Consider the stack $\mathcal{G} = \mathcal{G}_\rho$ defined as follows

$$\begin{aligned} \mathcal{G} &:= [\{(x, \alpha, b) \in \text{gl}_2 \times \text{gl}_1 \times \text{gl}_1 \mid \det(x) = \alpha^2\} / \text{GL}_1 \times \text{GL}_1] && \text{if } G = \text{SL}_2 \\ \mathcal{G} &:= [\{(x, \alpha, b) \in \text{gl}_2 \times \text{gl}_1 \times \text{gl}_1 \mid \det(x) = \alpha^2\} / \text{GL}_1 \times \mu_2 \times \text{GL}_1] && \text{if } G = \text{PGL}_2 \end{aligned}$$

Notice that the inclusion $i : G \hookrightarrow \mathcal{G}$ is a $G \times G$ -equivariant open embedding (where $G \times G$ acts by left and right multiplication).

Consider the kernel $\phi^{\mathcal{G}} : \mathcal{G}^F \rightarrow \overline{\mathbb{Q}}_\ell$ defined as follows (see §3.3):

If $G = \mathrm{SL}_2$,

$$\begin{aligned} \phi^{\mathcal{G}}([x, \alpha, b]) &= \sum_{s, t \in \mathbb{F}_q^\times} \psi(s(\mathrm{Tr}(x) + 2\alpha) + tb) \\ &= 1 - q\delta_{\mathrm{Tr}(x) + 2\alpha = 0}. \end{aligned}$$

If $G = \mathrm{PGL}_2$,

$$\phi^{\mathcal{G}}([x, \alpha, b]) = \begin{cases} \sum_{s, t \in \mathbb{F}_q^\times, \varepsilon \in \mu_2} \psi(s(\mathrm{Tr}(x) + 2\varepsilon\alpha) + tb) & \text{if } \alpha \in \mathfrak{gl}_1^F, \\ 0 & \text{if } \alpha \in \mathfrak{gl}_1^{F'}. \end{cases}$$

where $F' : \mathfrak{gl}_1 \rightarrow \mathfrak{gl}_1, x \mapsto -x^q$.

Our first main theorem is the following one (see §3.3).

Theorem 1.4.1. (i) *The Fourier transform $\mathcal{F}^{\mathcal{G}} : C(\mathcal{G}^F) \rightarrow C(\mathcal{G}^F)$ defined from the kernel $\phi^{\mathcal{G}}$ is involutive, more precisely*

$$\mathcal{F}^{\mathcal{G}} \circ \mathcal{F}^{\mathcal{G}}(f) = q^5 f$$

for any $f \in C(\mathcal{G}^F)$.

(ii) $\mathcal{F}^{\mathcal{G}}$ extends the BK-Fourier operator $\mathcal{F}^G = \mathcal{F}_\rho^G$, i.e.

$$\mathcal{F}^G = i^* \circ \mathcal{F}^{\mathcal{G}} \circ i_!$$

In the course of proving the above theorem we need the following unsatisfactory trick

$$\sum_{s \in \mathbb{F}_q^\times} \psi(s\tau) = \sum_{s \in \mathbb{F}_q^\times} \psi(s\tau^2)$$

as an essential ingredient.

Finally we will discuss the case of the symmetric square representation of $G = \mathrm{GL}_2$. The BK-Fourier kernel turns out to be more complicated. However there is still a natural stack \mathcal{G}_ρ and a natural Fourier operator on the space of functions on \mathbb{F}_q -points but it is not involutive.

2 Braverman-Kazhdan kernels

We choose a finite field \mathbb{F}_q and a prime ℓ which does not divide q . In sections 2.3 2.4 2.5 the characteristic of \mathbb{F}_q will be assumed to be odd. Throughout this paper $\alpha_o : \mathbb{F}_q^\times \rightarrow \overline{\mathbb{Q}}_\ell^\times$ will denote the character that takes the value 1 at squares and the value -1 at non-squares. We also fix a non-trivial additive linear character $\psi : \mathbb{F}_q \rightarrow \overline{\mathbb{Q}}_\ell^\times$ and denote by Tr the trace map on matrices. The maximal torus of GL_n of diagonal matrices is denoted by T_n .

2.1 Generalities for tori

Let S, S^\vee be two tori defined over \mathbb{F}_q with geometric Frobenius F . Recall that (S, F) and (S^\vee, F) are dual if there exists an isomorphism $X(S) \simeq Y(S^\vee)$ compatible with Galois action where $X(S)$ denotes the character group of S and $Y(S^\vee)$ the co-character group of S^\vee .

Given a morphism $\rho^\vee : S^\vee \rightarrow S'^\vee$ defined over \mathbb{F}_q , we get a *dual* morphism $\rho : S' \rightarrow S$ (see for instance [8, §3.5]). More concretely, if

$$\rho^\vee : S^\vee \rightarrow T_n^\vee = T_n, \quad s \mapsto (\alpha_1(s), \dots, \alpha_n(s))$$

with $\alpha_1, \dots, \alpha_n \in X(S^\vee)$, then

$$\rho : T_n \rightarrow S, (t_1, \dots, t_n) \mapsto \alpha_1(t_1) \cdots \alpha_n(t_n)$$

where $\alpha_1, \dots, \alpha_n$ are regarded as co-characters of S under the isomorphism $X(S^\vee) \simeq Y(S)$.

If T_n is equipped with the standard Frobenius that raises matrix coefficients to their q -th power, the BK-Fourier kernel ϕ_ρ^S is then (see [8, Lemma 5.2.1])

$$\phi_\rho^S = c_S c_{T_n}^{-1} \rho_!^F(\psi \circ \text{Tr}|_{T_n}).$$

where

$$c_S := (-1)^{\mathbb{F}_q - \text{rank}(S)}. \tag{2.1}$$

2.2 The symmetric square representations

Let G be denote either $\text{PGL}_2, \text{SL}_2$ or GL_2 over $\overline{\mathbb{F}}_q$ equipped with standard Frobenius $F : G \rightarrow G$ that raises matrix coefficients to their q -th power. Denote by (G^\vee, F) the dual of (G, F) (in the sense of Deligne-Lusztig) so that $\text{PGL}_2^\vee = \text{SL}_2, \text{SL}_2^\vee = \text{PGL}_2$ and $\text{GL}_2^\vee = \text{GL}_2$ equipped with standard Frobenius.

Denote by

$$\rho^\vee : G^\vee \rightarrow \text{GL}_3$$

the *symmetric square representation* of G^\vee , namely if we let $\text{Sym}^2 : \text{GL}_2 \rightarrow \text{GL}_3$ be defined by

$$\text{Sym}^2 \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} a^2 & ab & b^2 \\ 2ac & ad + bc & 2bd \\ c^2 & cd & d^2 \end{pmatrix},$$

then

$$\begin{aligned} \rho^\vee &= \text{Sym}^2 \text{ if } G = \text{GL}_2 \\ \rho^\vee &= \text{Sym}^2|_{\text{SL}_2} \text{ if } G = \text{PGL}_2 \\ \rho^\vee &= \text{Sym}^2 \cdot \det^{-1} \text{ if } G = \text{SL}_2. \end{aligned}$$

Remark 2.2.1. Notice that we have factorizations

$$\begin{array}{ccccc} \text{SL}_2 & \xrightarrow{\pi^\vee} & \text{GL}_2 & \xrightarrow{\text{Sym}^2} & \text{GL}_3 \\ & \searrow f^\vee & & \nearrow \text{Sym}^2 \cdot \det^{-1} & \\ & & \text{PGL}_2 & & \end{array} \quad (2.2)$$

where f^\vee is the canonical quotient and π^\vee the canonical inclusion.

The morphism ρ^\vee commutes thus with the Frobenius morphisms. We denote by T the maximal torus of G consisting of diagonal matrices, and we denote by $W = \{1, \sigma\} \simeq S_2$ the Weyl group of G with respect to T . The morphism ρ^\vee induces a morphism $W \rightarrow S_3$ mapping the non-trivial element σ of W to the transposition $(13) \in S_3$ which we also denote by σ .

We choose an element $g \in G$ (resp. $g \in \text{GL}_3$) such that

$$g^{-1}F(g) = \sigma.$$

We then denote by T_σ (resp. $T_{3,\sigma}$) the F -stable maximal torus gTg^{-1} (resp. gT_3g^{-1}).

Under conjugation by g^{-1} we have

$$(T_\sigma, F) \simeq (T, \sigma F), \quad (T_{3,\sigma}, F) \simeq (T_3, \sigma F).$$

In particular, $T_\sigma^F \simeq T^{\sigma F}$ and $T_{3,\sigma}^F \simeq T_3^{\sigma F}$. It will be more convenient to represent elements of T_σ^F in $T^{\sigma F}$.

The morphism ρ^\vee restricts to a W -equivariant morphism $T^\vee \rightarrow T_3$ inducing (see [8, Section 3.5]) a W -equivariant morphism $\rho : T_3 \rightarrow T$ given as follows

$$\begin{aligned} \rho(a, b, c) &= (a^2b, bc^2) & \text{if } G = \text{GL}_2 \\ \rho(a, b, c) &= (a/c, c/a) & \text{if } G = \text{SL}_2 \\ \rho(a, b, c) &= [a^2b, bc^2] & \text{if } G = \text{PGL}_2 \end{aligned}$$

where $[a^2b, bc^2]$ denotes the class of (a^2b, bc^2) in PGL_2 .

We thus get morphisms $\rho^F : T_3^F \rightarrow T^F$ and $\rho^{\sigma F} : T_3^{\sigma F} \rightarrow T^{\sigma F}$, and the Braverman-Kazhdan kernels (BK-Fourier kernels) ϕ^T and $\phi^{T\sigma}$ on T^F and $T^{\sigma F}$ respectively are given by (see §2.1)

$$\phi^T = c_{T_3} c_T^{-1} \rho_!^F(\psi \circ \mathrm{Tr}), \quad \phi^{T\sigma} = c_{T_{3,\sigma}} c_{T_\sigma}^{-1} \rho_!^{\sigma F}(\psi \circ \mathrm{Tr}). \quad (2.3)$$

(we omit ρ from the notation of the BK-Fourier kernels as the context will be always clear). The BK-Fourier kernel on G^F is then given by the formula [8, Theorem 5.2.3]

$$\phi^G = \frac{1}{2} (R_T^G(\phi^T) + R_{T_\sigma}^G(\phi^{T\sigma})), \quad (2.4)$$

where R_T^G and $R_{T_\sigma}^G$ denote Deligne-Lusztig induction.

The right hand side of (2.4) can be computed explicitly. For $G = \mathrm{GL}_2$ we get :

| | | | | | |
|----------|---|--|--|---|-------|
| | $g = \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}$ $a \in \mathbb{F}_q^\times$ | $g = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$ $a \neq b \in \mathbb{F}_q^\times$ | $g = \begin{pmatrix} x & 0 \\ 0 & x^q \end{pmatrix}$ $x \neq x^q \in \mathbb{F}_{q^2}^\times$ | $g = \begin{pmatrix} a & 1 \\ 0 & a \end{pmatrix}$ $a \in \mathbb{F}_q^\times$ | |
| ϕ^G | $\frac{1}{2} ((q+1)\phi^T(g) + (1-q)\phi^{T\sigma}(g))$ | $\phi^T(g)$ | $\phi^{T\sigma}(g)$ | $\frac{1}{2} (\phi^T(g_s) + \phi^{T\sigma}(g_s))$ | (2.5) |

where g_s denotes the semisimple part of g .

Remark 2.2.2. Notice that the map $f^\vee : \mathrm{SL}_2 \rightarrow \mathrm{PGL}_2$ in (2.2) is a *normal* morphism (i.e. its image is a normal subgroup) with dual morphism $f = f^\vee : \mathrm{SL}_2 \rightarrow \mathrm{PGL}_2$ (see [8, §3.5]). From the transitivity of the transfert map between Lusztig series (see [8, §3.6]), we get from [8, Lemma 4.1.1] that

$$\phi^{\mathrm{PGL}_2} = \begin{cases} f_!^F(\phi^{\mathrm{SL}_2}) \\ -q^{-1} \pi_!^F(\phi^{\mathrm{GL}_2}). \end{cases} \quad (2.6)$$

Notice that $c_{\mathrm{PGL}_2} c_{\mathrm{GL}_2}^{-1} = -1$ and $c_{\mathrm{PGL}_2} c_{\mathrm{SL}_2}^{-1} = 1$ (see definition of the gamma function (1.1)).

2.3 Explicit computation : The case $G = \mathrm{SL}_2$

Recall that in this case

$$\begin{aligned} \rho^F : T_3^F &\rightarrow T^F, & (a, b, c) &\mapsto (a/c, c/a), \\ \rho^{\sigma F} : T_3^{\sigma F} &\rightarrow T^{\sigma F}, & (a', b, a'^q) &\mapsto (a'^{1-q}, a'^{q-1}). \end{aligned}$$

Notice that $c_{T_3} c_T^{-1} = c_{T_{3,\sigma}} c_{T_\sigma}^{-1} = 1$ and

$$\mathrm{Ker}(\rho^F) = \{(s, t, s) \mid s, t \in \mathbb{F}_q^\times\} = \mathrm{Ker}(\rho^{\sigma F}).$$

Therefore

$$\begin{aligned}
\phi^T(a/c, c/a) &= \rho_1^F(\psi \circ \text{Tr})(a/c, c/a) \\
&= \sum_{s,t} \psi(s(a+c) + bt) \\
&= - \sum_s \psi(s(a+c)) \\
\phi^{T\sigma}(a'^{1-q}, a'^{q-1}) &= \rho_1^{\sigma F}(\psi \circ \text{Tr})(a'^{1-q}, a'^{q-1}) \\
&= \sum_{s,t} \psi(s(a' + a'^q) + bt) \\
&= - \sum_s \psi(s(a' + a'^q))
\end{aligned}$$

where $a' \in \mathbb{F}_q^\times$.

We now use the identification

$$\text{SL}_2 \simeq \{(x, \alpha) \in \text{GL}_2 \times \text{GL}_1 \mid \det(x) = \alpha^2\} / \text{GL}_1$$

where GL_1 acts by $s \cdot (x, \alpha) = (sx, s\alpha)$, and we write the elements of SL_2 in the form $[x, \alpha]$ with $\det(x) = \alpha^2$.

Notice that

$$\text{SL}_2^F = \{(x, \alpha) \in \text{GL}_2^F \times \text{GL}_1^F \mid \det(x) = \alpha^2\} / \text{GL}_1^F.$$

The morphism ρ^F and $\rho^{\sigma F}$ becomes

$$\rho^F(a, b, c) = [(a^2, c^2), ac], \quad \rho^{\sigma F}(a', b, a'^q) = [(a'^2, a'^{2q}), a'^{q+1}].$$

The crucial ingredient is the following obvious identity

$$\sum_{s \in \mathbb{F}_q^\times} \psi(s\tau) = \sum_{s \in \mathbb{F}_q^\times} \psi(s\tau^2) = \begin{cases} q-1 & \text{if } \tau = 0, \\ -1 & \text{otherwise.} \end{cases}$$

Indeed we get

$$\phi^T([(a^2, c^2), ac]) = - \sum_{s \in \mathbb{F}_q^\times} \psi(s(a+c)) = - \sum_{s \in \mathbb{F}_q^\times} \psi(s(a+c)^2) = - \sum_{s \in \mathbb{F}_q^\times} \psi(s(a^2 + c^2 + 2ac)).$$

We thus have for $[x, \alpha] \in T^F$

$$\phi^T([x, \alpha]) = - \sum_{s \in \mathbb{F}_q^\times} \psi(s(\text{Tr}(x) + 2\alpha)),$$

and similarly for $[x', \alpha] \in T_\sigma^F$, we have

$$\phi^{T_\sigma}([x', \alpha]) = - \sum_{s \in \mathbb{F}_q^\times} \psi(s(\text{Tr}(x') + 2\alpha)).$$

We thus deduce from Formula (2.4) the following one.

Proposition 2.3.1. *For $[x, \alpha] \in \text{SL}_2^F$ we have*

$$\phi^G([x, \alpha]) = - \sum_{s \in \mathbb{F}_q^\times} \psi(s(\text{Tr}(x) + 2\alpha)) = \begin{cases} -(q-1) & \text{if } \text{Tr}(x) + 2\alpha = 0, \\ 1 & \text{otherwise.} \end{cases}$$

Remark 2.3.2. It will be useful to get rid of the sign by writing SL_2 in the form

$$\text{SL}_2 \simeq [\{(x, \alpha, b) \in \text{GL}_2 \times \text{GL}_1 \times \text{GL}_1 \mid \det(x) = \alpha\} / \text{GL}_1 \times \text{GL}_1]$$

where $\text{GL}_1 \times \text{GL}_1$ acts as $(s, t) \cdot (x, \alpha, b) = (sx, s\alpha, tb)$. Indeed, under this identification

$$\phi^G([x, \alpha, b]) = \sum_{s, t \in \mathbb{F}_q^\times} \psi(s(\text{Tr}(x) + 2\alpha) + bt).$$

2.4 Explicit computation : The case $G = \text{PGL}_2$

Recall that

$$\rho : T_3 \rightarrow T, \quad (a, b, c) \mapsto [a^2b, bc^2] = [a/c, c/a].$$

and so

$$\text{Ker}(\rho^F) = \{(s, t, \varepsilon s) \mid s, t \in \mathbb{F}_q^\times, \varepsilon \in \mu_2\}, \quad \text{Ker}(\rho^{\sigma F}) = \{(u, t, u^q) \mid t \in \mathbb{F}_q^\times, u^q = \pm u\}.$$

We thus have

$$\begin{aligned} \phi^T([a/c, c/a]) &= \sum_{s, t \in \mathbb{F}_q^\times, \varepsilon \in \mu_2} \psi(s(a + \varepsilon c) + t) \\ &= - \sum_{s, \varepsilon} \psi(s(a + \varepsilon c)^2) \\ &= - \sum_{s, \varepsilon} \psi(s(a^2 + c^2 + 2\varepsilon ac)) \\ \phi^{T_\sigma}([a'^{1-q}, a'^{q-1}]) &= - \sum_{s \in \mathbb{F}_q^\times, \varepsilon \in \mu_2} \psi(s(a'^2 + a'^{2q} + 2\varepsilon a'^{q+1})). \end{aligned}$$

Let us identify

$$\mathrm{PGL}_2 = \mathrm{SL}_2/\mu_2 = \{(x, \alpha) \in \mathrm{GL}_2 \times \mathrm{GL}_1 \mid \det(x) = \alpha^2\} / \mathrm{GL}_1 \times \mu_2$$

where μ_2 acts by multiplication on the second coordinate. Notice that

$$\mathrm{PGL}_2^F = \{(x, \alpha) \in \mathrm{GL}_2^F \times (\mathrm{GL}_1^F \sqcup \mathrm{GL}_1^{F'}) \mid \det(x) = \alpha^2\} / \mathrm{GL}_1^F \times \mu_2$$

where $F' : \mathrm{GL}_1 \rightarrow \mathrm{GL}_1, x \mapsto -x^q$ is the standard Frobenius twisted by the non-trivial element of $H^1(F, \mu_2) = \mu_2$.

Using Formula (2.4) we deduce the following one.

Proposition 2.4.1. *For $[x, \alpha] \in \mathrm{PGL}_2^F$ we have*

$$\phi^G([x, \alpha]) = \begin{cases} -\sum_{s \in \mathbb{F}_q^\times, \varepsilon \in \mu_2} \psi(s(\mathrm{Tr}(x) + 2\varepsilon\alpha)) & \text{if } \alpha \in \mathrm{GL}_1^F, \\ 0 & \text{if } \alpha \in \mathrm{GL}_1^{F'}. \end{cases}$$

Notice that the result could have been also deduced from Formula (2.6) and Proposition 2.3.1.

Remark 2.4.2. From Proposition 2.3.1 and Proposition 2.4.1, we see that the kernel ϕ^G descends to the GIT quotient $T//W$ when G is SL_2 or PGL_2 . We will see in the next section that this is not true when $G = \mathrm{GL}_2$.

2.5 Explicit computation : The case $G = \mathrm{GL}_2$

We have

$$\rho : T_3 \rightarrow T, \quad (a, b, c) \mapsto (a^2b, bc^2).$$

and so

$$\mathrm{Ker}(\rho^F) = \{(s, s^{-2}, \varepsilon s) \mid s \in \mathbb{F}_q^\times, \varepsilon \in \mu_2\}, \quad \mathrm{Ker}(\rho^{\sigma F}) = \{(s, s^{-2}, s^q) \mid s^q = \pm s, s \neq 0\}.$$

For $b \in \mathbb{F}_q$, put

$$\kappa(b) := \sum_{s \in \mathbb{F}_q^\times} \psi(s^{-2}b) \quad \text{and} \quad \kappa'(b) := \sum_{\{s \neq 0, s^q = -s\}} \psi(s^{-2}b).$$

Notice that if $r \in \mathbb{F}_q^\times$ is a non-square, then

$$\kappa(rb) = \kappa'(b).$$

Since $c_{T_3, T} = c_{T_3, \sigma, T_\sigma} = -1$, we have (for $a, b, c \in \mathbb{F}_q^\times$)

$$\begin{aligned} \phi^T(a^2b, bc^2) &= - \sum_{s \in \mathbb{F}_q^\times, \varepsilon \in \mu_2} \psi(s(a + \varepsilon c) + s^{-2}b) \\ &= \begin{cases} - \sum_{s, \varepsilon} \psi(s + s^{-2}(a + \varepsilon c)^2b) & \text{if } a^2 \neq c^2, \\ -(\kappa(b) + \sum_s \psi(s + s^{-2}(a + \varepsilon c)^2b)) & \text{if } a = \varepsilon c, \end{cases} \end{aligned}$$

using the variable change $s(a + \varepsilon c) \leftrightarrow s$ when $a + \varepsilon c \neq 0$.

We get the formula for ϕ^{T_σ} as follows (where $s_o \in \mathbb{F}_{q^2}$ satisfies $s_o^q = -s_o$)

$$\begin{aligned} \phi^{T_\sigma}(a'^2b, ba'^{2q}) &= - \sum_{s^q=s} \psi(s(a' + a'^q) + s^{-2}b) - \sum_{s^q=-s} \psi(s(a' - a'^q) + s^{-2}b) \\ &= - \sum_{s^q=s} \psi(s(a' + a'^q) + s^{-2}b) - \sum_{s^q=s} \psi(ss_o(a' - a'^q) + s_o^{-2}s^{-2}b) \\ &= \begin{cases} - \sum_{s \in \mathbb{F}_q^\times, \varepsilon \in \mu_2} \psi(s + s^{-2}(a' + \varepsilon a'^q)^2b) & \text{if } a'^q \neq \pm a', \\ - \sum_{s \in \mathbb{F}_q^\times} \psi(s + s^{-2}(a' + a'^q)^2b) - \kappa'(b) & \text{if } a'^q = a', \\ - \sum_{s \in \mathbb{F}_q^\times} \psi(s + s^{-2}(a' - a'^q)^2b) - \kappa(b) & \text{if } a'^q = -a'. \end{cases} \end{aligned}$$

using the variable changes $ss_o(a' - a'^q) \leftrightarrow s$ if $a'^q \neq a'$ and $s(a' + a'^q) \leftrightarrow s$ if $a'^q \neq -a'$.

Remark 2.5.1. The case $a'^q = -a'$ is not essential as we can write (a'^2b, ba'^{2q}) in the form (a^2d, da^2) with $a, d \in \mathbb{F}_q^\times$ by putting $(a, d) := (s_o a', s_o^{-2}b)$.

We now consider the identification

$$\mathrm{GL}_2 \simeq \{(x, \alpha, b) \in \mathrm{GL}_2 \times \mathrm{GL}_1 \times \mathrm{GL}_1 \mid \det(x) = \alpha^2\} / \mathrm{GL}_1 \times \mu_2 \quad (2.7)$$

where GL_1 acts as $t \cdot [x, \alpha, b] = [tx, t\alpha, t^{-1}b]$ and μ_2 acts by multiplication on the second coordinate (under this identification, the element $[x, \alpha, b]$ corresponds to $xb \in \mathrm{GL}_2$).

We have

$$\mathrm{GL}_2^F = \{(x, \alpha, b) \in \mathrm{GL}_2^F \times (\mathrm{GL}_1^F \sqcup \mathrm{GL}_1^{F'}) \times \mathrm{GL}_1^F \mid \det(x) = \alpha^2\} / \mathrm{GL}_1^F \times \mu_2.$$

In the following table we write the elements of GL_2^F in the form $[x, \alpha, b]$ and we put

$$\Phi([x, \alpha, b]) := \sum_{s \in \mathbb{F}_q^\times, \varepsilon \in \mu_2} \psi(s + s^{-2}b(\mathrm{Tr}(x) + 2\varepsilon\alpha)).$$

| | | | | |
|------------------|---|--|--|--|
| | $\left[\begin{pmatrix} a^2 & 0 \\ 0 & a^2 \end{pmatrix}, a^2, b \right]$ $a, b \in \mathbb{F}_q^\times$ | $\left[\begin{pmatrix} a^2 & 0 \\ 0 & c^2 \end{pmatrix}, ac, b \right]$ $a, c \in \mathbb{F}_q^\times, a^2 \neq c^2$ | $\left[\begin{pmatrix} a'^2 & 0 \\ 0 & a'^{2q} \end{pmatrix}, a'^{q+1}, b \right]$ $a'^2 \neq a'^{2q}$ | |
| ϕ^T | $-\kappa(b) - 1 - \Phi([x, \alpha, b])$ | $-\Phi([x, \alpha, b])$ | \times | |
| $\phi^{T\sigma}$ | $-\kappa'(b) - 1 - \Phi([x, \alpha, b])$ | \times | $-\Phi([x, \alpha, b])$ | |

(2.8)

Notice that for $b \neq 0$

$$\kappa(b) + \kappa'(b) = -2, \quad \frac{1}{2}(\kappa(b) - \kappa'(b)) = \sum_{t \in \mathbb{F}_q^\times} \alpha_o(t) \psi(tb) =: S(\alpha_o, \psi_b)$$

where recall that α_o is the non-trivial square root of the identity character of \mathbb{F}_q^\times (it takes the value 1 at squares and -1 at non-squares).

Using Table (2.5) and the above table we get the following proposition.

Proposition 2.5.2. *For $[x, \alpha, b] \in \mathrm{GL}_2^F$ we have $\phi^G([x, \alpha, b]) = 0$ if $\alpha \in \mathrm{GL}_1^{F'}$ and otherwise we have*

| | | | | | |
|----------|---|--|--|---|--|
| | $\left[\begin{pmatrix} a^2 & 0 \\ 0 & a^2 \end{pmatrix}, a^2, b \right]$ $a, b \in \mathbb{F}_q^\times$ | $\left[\begin{pmatrix} a^2 & 0 \\ 0 & c^2 \end{pmatrix}, ac, b \right]$ $a, c \in \mathbb{F}_q^\times, a^2 \neq c^2$ | $\left[\begin{pmatrix} a'^2 & 0 \\ 0 & a'^{2q} \end{pmatrix}, a'^{q+1}, b \right]$ $a'^2 \neq a'^{2q}$ | $\left[\begin{pmatrix} a^2 & 1 \\ 0 & a^2 \end{pmatrix}, a^2, b \right]$ $a, b \in \mathbb{F}_q^\times$ | |
| ϕ^G | $-qS(\alpha_o, \psi_b) - \Phi([x, \alpha, b])$ | $-\Phi([x, \alpha, b])$ | $-\Phi([x, \alpha, b])$ | $-\Phi([x, \alpha, b])$ | |

(2.9)

This can be re-written for $g \in \mathrm{GL}_2^F$ as

$$\phi^G(g) = \begin{cases} 0 & \text{if } \det(g) \notin (\mathbb{F}_q^\times)^2, \\ -\sum_{s, \varepsilon} \psi \left(s + s^{-2} \left(\mathrm{Tr}(g) + 2\varepsilon \sqrt{\det(g)} \right) \right) & \text{if } \det(g) \in (\mathbb{F}_q^\times)^2 \text{ and } g \text{ not central,} \\ -\sum_{s, \varepsilon} \psi \left(s + s^{-2} \left(\mathrm{Tr}(g) + 2\varepsilon \sqrt{\det(g)} \right) \right) - qS(\alpha_o, \psi_\lambda) & \text{if } g = \lambda \cdot I_2. \end{cases}$$

Notice that in the last case, $\phi^G(g) = 1 - \sum_s \psi(s + 4s^{-2}\lambda) - qS(\alpha_o, \psi_\lambda)$.

We see that ϕ^G does not descend to $T//W$ as the values of ϕ^G depends on the unipotent part.

3 Extending the BK-Fourier kernel: The case $G = \mathrm{SL}_2, \mathrm{PGL}_2$

3.1 Preliminaries

By a finite groupoid, we shall mean a groupoid with a finite number of isomorphism classes and whose automorphism groups are finite. For a finite groupoid X we denote by \overline{X} the set of

isomorphism classes of X . By a $\overline{\mathbb{Q}_\ell}$ -valued function $X \rightarrow \overline{\mathbb{Q}_\ell}$ on X we shall mean a $\overline{\mathbb{Q}_\ell}$ -valued function on \overline{X} and we denote by $C(X)$ the $\overline{\mathbb{Q}_\ell}$ -vector space of functions on X . Given a morphism $\varphi : X \rightarrow Y$ of finite groupoids, we have the operator $\varphi^* : C(Y) \rightarrow C(X)$ defined by composing with φ and the operator $\varphi_! : C(X) \rightarrow C(Y)$ defined by

$$\varphi_!(f)(y) = \sum_{x \in X_y} \frac{1}{|\text{Aut}(x)|} y_X^*(f)(x)$$

where \bullet is a point, $X_y = \bullet \times_Y X$,

$$\begin{array}{ccc} X_y & \xrightarrow{y_X} & X \\ \downarrow & & \downarrow \varphi \\ \bullet & \xrightarrow{y} & Y \end{array}$$

If G is a finite group acting on a finite set X , we denote by $[X/G]$ the groupoid of G -equivariant maps $G \rightarrow X$ and by $\pi : X \rightarrow [X/G]$ the quotient map. Then for $f \in C(X)$ we have

$$\pi_!(f)(\pi(x)) = \sum_{g \in G} f(g \cdot x)$$

for any $x \in X$.

Given a function $K \in C(X \times X)$ we can define an operator

$$\mathcal{F} : C(X) \rightarrow C(X)$$

with kernel K by

$$\mathcal{F}(f)(y) = \sum_{x \in \overline{X}} \frac{1}{|\text{Aut}(x)|} K(y, x) f(x)$$

for $f \in C(X)$ and $y \in \overline{X}$.

We have the following straightforward lemma.

Lemma 3.1.1. *The operator \mathcal{F} is involutive, i.e. $\mathcal{F} \circ \mathcal{F} = 1$, if and only if for all $y, z \in \overline{X}$ we have*

$$\Delta(z, x) := \sum_{y \in \overline{X}} \frac{1}{|\text{Aut}(y)|} K(z, y) K(y, x) = \begin{cases} |\text{Aut}(z)| & \text{if } z = x, \\ 0 & \text{otherwise.} \end{cases} \quad (3.1)$$

Let Z be a finite set endowed with a right action of a finite group H .

Lemma 3.1.2. *Let $K^Z \in C(Z \times Z)$ be H -invariant for the diagonal action of H on $Z \times Z$ and define $K^{[Z/H]} \in C([Z/H] \times [Z/H])$ by*

$$K^{[Z/H]}(\bar{x}, \bar{y}) = \sum_{h \in H} K^Z(h \cdot x, y) = \sum_{h \in H} K^Z(x, h \cdot y)$$

where $x, y \in Z$ maps to $\bar{x} \in [Z/H]$ and $\bar{y} \in [Z/H]$ respectively. If K^Z satisfies (3.1) then so does $K^{[Z/H]}$.

Proof. Suppose that K^Z satisfies (3.1).

$$\begin{aligned} \sum_{\bar{y} \in [Z/H]} \frac{1}{|C_H(y)|} K^{[Z/H]}(\bar{x}, \bar{y}) K^{[Z/H]}(\bar{y}, \bar{z}) &= \sum_{\bar{y} \in [Z/H]} \frac{1}{|C_H(y)|} \sum_{h \in H} K^Z(h \cdot x, y) \sum_{h' \in H} K^Z(y, h' \cdot z) \\ &= \frac{1}{|H|} \sum_{y \in Z} \sum_{h \in H} K^Z(h \cdot x, y) \sum_{h' \in H} K^Z(y, h' \cdot z) \\ &= \frac{1}{|H|} \sum_{h, h' \in H} \sum_{y \in Z} K^Z(h \cdot x, y) K^Z(y, h' \cdot z) \\ &= \frac{1}{|H|} \sum_{h, h' \in H} \delta_{hx, h'x} \\ &= |C_H(x)| \delta_{\bar{x}, \bar{z}}. \end{aligned}$$

□

Let X be an $\overline{\mathbb{F}}_q$ -scheme on which an $\overline{\mathbb{F}}_q$ -algebraic group G acts on the right. We denote by $[X/G]$ the associated quotient stack. We assume that X , G and the action of G on X are defined over \mathbb{F}_q and we denote by F the associated geometric Frobenius on X , G and $[X/G]$.

Recall that the isomorphism classes of G -torsors over $\text{Spec}(\mathbb{F}_q)$ are parametrized by the set $H^1(F, G) = H^1(F, G/G^o)$ of F -conjugacy classes on G and so the groupoid $[X/G]^F$ of \mathbb{F}_q -points of $[X/G]$ decomposes as

$$[X/G]^F = \coprod_{\bar{h} \in H^1(F, G)} [X^{F \circ h} / G^{F \circ h}]$$

where $h \in G$ denotes a representative of \bar{h} and $G^{F \circ h} = \{g \in G \mid F(h^{-1}gh) = g\}$.

If G is connected then $H^1(F, G)$ is trivial and so $[X/G]^F = [X^F / G^F]$.

Let

$$\pi : [X/G^o] \rightarrow [X/G]$$

be the natural G/G^o -torsor.

Notice that the induced map

$$\pi^F : [X/G^o]^F \rightarrow [X/G]^F$$

is not surjective. But, for any $h \in G$, it induces a surjective map

$$[X/G^o]^{Foh} = [X^{Foh}/G^{o(Foh)}] \rightarrow [X^{Foh}/G^{Foh}]$$

with fibers isomorphic to $(G/G^o)^{Foh} = G^{Foh}/G^{o(Foh)}$.

Let $S : [X/G^o] \times [X/G^o] \rightarrow \overline{\mathbb{F}}_q$ be a G/G^o -invariant algebraic map for the diagonal action of G/G^o . Then

$$S(F(y), F(x)) = S(F \circ h(y), F \circ h(x)) = S(y, x)^q$$

for all $x, y \in [X/G^o]$ and $h \in G$, and so S induces maps

$$S^{Foh} : [X/G^o]^{Foh} \times [X/G^o]^{Foh} \rightarrow \mathbb{F}_q.$$

For all $\bar{h} \in H^1(F, G)$, we get a kernel

$$K_{\bar{h}}^{[X/G^o]} : [X/G^o]^{Foh} \times [X/G^o]^{Foh} \rightarrow \overline{\mathbb{Q}}_\ell, \quad (x, y) \mapsto \psi(S(x, y)).$$

and so a kernel

$$K_{\bar{h}}^{[X/G]} : [X^{Foh}/G^{Foh}] \times [X^{Foh}/G^{Foh}] \rightarrow \overline{\mathbb{Q}}_\ell,$$

defined by

$$K_{\bar{h}}^{[X/G]}(\bar{x}, \bar{y}) = \sum_{w \in (G/G^o)^{Foh}} \psi(S(w \cdot x, y)),$$

for $x, y \in [X/G^o]^{Foh}$ with images denoted respectively by \bar{x}, \bar{y} in the quotient $[X^{Foh}/G^{Foh}]$.

We define a kernel

$$K^{[X/G]} : [X/G]^F \times [X/G]^F \rightarrow \overline{\mathbb{Q}}_\ell$$

by

$$K^{[X/G]}(\bar{x}, \bar{y}) = \begin{cases} K_{\bar{h}}^{[X/G]}(\bar{x}, \bar{y}) & \text{if } \bar{x}, \bar{y} \text{ are both in } [X^{Foh}/G^{Foh}] \text{ for some } \bar{h} \in H^1(F, G), \\ 0 & \text{otherwise.} \end{cases}$$

Proposition 3.1.3. *The Fourier transform $\mathcal{F}^{[X/G]} : C([X/G]^F) \rightarrow C([X/G]^F)$ defined from the kernel $K^{[X/G]}$ is involutive if and only if all the kernels $K_{\bar{h}}^{[X/G^o]}$, with $\bar{h} \in H^1(F, G)$, satisfy (3.1).*

Remark 3.1.4. The above construction of the kernel $K^{[X/G]}$ has the following geometrical counterpart. The geometric analogue of ψ is the Artin-Schreier sheaf \mathcal{L}_ψ on \mathbb{A}^1 and so $K^{[X/G]}$ corresponds to

$$\mathcal{K}^{[X/G]} := S^*(\mathcal{L}_\psi).$$

The sheaf $\mathcal{K}^{[X/G]}$ is G/G^o -equivariant for the diagonal action of G/G^o (see for instance [7, §2.3] for the definition of equivariance).

We consider

$$[X/G^o] \times [X/G^o] \xrightarrow{\pi \times 1} [X/G] \times [X/G^o] \xrightarrow{1 \times \pi} [X/G] \times [X/G]$$

Then the complex $(\pi \times 1)_! \mathcal{K}^{[X/G^o]}$ is naturally G/G^o -equivariant and so by [8, Lemma 2.3.4] it descends to a complex $\mathcal{K}^{[X/G]}$ on $[X/G] \times [X/G]$, i.e. $\mathcal{K}^{[X/G]}$ is the unique complex (up to a unique isomorphism) such that

$$(1 \times \pi)^* \mathcal{K}^{[X/G]} = (\pi \times 1)_! \mathcal{K}^{[X/G^o]}.$$

Then $\mathcal{K}^{[X/G]}$ is the geometrical counterpart of $K^{[X/G]}$, i.e. if we introduce Frobenius action, then $K^{[X/G]}$ is the characteristic function of $\mathcal{K}^{[X/G]}$ equipped with its natural Frobenius (see [8, §6.1]).

Then the geometrical counterpart of Proposition 3.1.3 says that the geometric Fourier transform $\mathcal{F}^{[X/G^o]}$ defined from $\mathcal{K}^{[X/G^o]}$ is involutive if and only if the geometric Fourier transform $\mathcal{F}^{[X/G]}$ defined from $\mathcal{K}^{[X/G]}$ is.

3.2 The tori case

Let S be a torus defined over \mathbb{F}_q and assume given a surjective morphism

$$\rho : T_n \rightarrow S$$

defined over \mathbb{F}_q (the morphism ρ may not be surjective on \mathbb{F}_q -points).

Consider the BK-Fourier kernel

$$\phi_\rho^S = \rho_!^F(\psi \circ \text{Tr}|_{T_n})$$

and consider the Fourier operator $\mathcal{F}_\rho^S : C(S^F) \rightarrow C(S^F)$ defined from ϕ_ρ^S .

We wish to extend this operator to an involutive Fourier transform.

Put

$$H := \text{Ker}(\rho), \quad \mathfrak{t}_n := \text{Lie}(T_n).$$

Consider the natural open embedding

$$\iota : S \simeq T_n / \text{Ker}(\rho) \hookrightarrow [\mathfrak{t}_n / H].$$

On $[\mathfrak{t}_n / H]^F$ define the kernel

$$\phi^{[\mathfrak{t}_n / H]} := q_!^F(\psi \circ \text{Tr})$$

where $q : \mathfrak{t}_n \rightarrow [\mathfrak{t}_n / H]$.

Concretely

$$\phi^{[\mathfrak{t}_n / H]}(y) = \begin{cases} \sum_{t \in H^F} \psi(\text{Tr}(tx)) & \text{if } y = q^F(x), \\ 0 & \text{if } x \notin \text{Im}(q^F). \end{cases}$$

Notice that q^F is not surjective if H is not connected.

The products $\mathfrak{t}_n \times \mathfrak{t}_n \rightarrow \mathfrak{t}_n$ and $H \times H \rightarrow H$ induce a map

$$m : [\mathfrak{t}_n / H] \times [\mathfrak{t}_n / H] \rightarrow [\mathfrak{t}_n / H]$$

which is compatible with \mathbb{F}_q -structures. We thus get a morphism of groupoids

$$m^F : [\mathfrak{t}_n / H]^F \times [\mathfrak{t}_n / H]^F \rightarrow [\mathfrak{t}_n / H]^F.$$

We then define a two-variable kernel $K^{[\mathfrak{t}_n / H]} : [\mathfrak{t}_n / H]^F \times [\mathfrak{t}_n / H]^F \rightarrow \overline{\mathbb{Q}}_\ell$ by

$$K^{[\mathfrak{t}_n / H]} = (m^F)^*(\phi^{[\mathfrak{t}_n / H]}).$$

Proposition 3.2.1. (1) The Fourier transform $\mathcal{F}^{[\mathfrak{t}_n / H]} : C([\mathfrak{t}_n / H]^F) \rightarrow C([\mathfrak{t}_n / H]^F)$ defined from the kernel $K^{[\mathfrak{t}_n / H]}$ is involutive, i.e.

$$\mathcal{F}^{[\mathfrak{t}_n / H]} \circ \mathcal{F}^{[\mathfrak{t}_n / H]}(f) = q^n f^-$$

for any $f \in C([\mathfrak{t}_n / H]^F)$ where $f^- \in C([\mathfrak{t}_n / H]^F)$ is defined as $f^-([x]) := f([-x])$.

(2) It extends the Fourier operator $\mathcal{F}_\rho^S : C(S^F) \rightarrow C(S^F)$, i.e.

$$\mathcal{F}_\rho^S = \iota^* \circ \mathcal{F}^{[\mathfrak{t}_n / H]} \circ \iota_!$$

Proof. The assertion (1) follows from the discussion in §3.1 as the Fourier transform on $C(\mathfrak{t}_n^F)$ defined from the kernel $\psi \circ \text{Tr}$ is involutive. The second assertion is straightforward. \square

3.3 Main result

In this section the characteristic of the base field \mathbb{F}_q is assumed to be odd, G is SL_2 or PGL_2 and $\rho^\vee : G \rightarrow \mathrm{GL}_3$ is as in §2.2. We consider the stack \mathcal{G} defined as follows :

$$\begin{aligned} \mathcal{G} &:= [\{(x, \alpha, b) \in \mathfrak{gl}_2 \times \mathfrak{gl}_1 \times \mathfrak{gl}_1 \mid \det(x) = \alpha^2\} / \mathrm{GL}_1 \times \mathrm{GL}_1] && \text{if } G = \mathrm{SL}_2, \\ \mathcal{G} &:= [\{(x, \alpha, b) \in \mathfrak{gl}_2 \times \mathfrak{gl}_1 \times \mathfrak{gl}_1 \mid \det(x) = \alpha^2\} / \mathrm{GL}_1 \times \mu_2 \times \mathrm{GL}_1] && \text{if } G = \mathrm{PGL}_2, \end{aligned}$$

where μ_2 acts on the second coordinate and $\mathrm{GL}_1 \times \mathrm{GL}_1$ as $(s, t) \cdot (x, \alpha) = (sx, s\alpha, tb)$.

The natural inclusion $G \hookrightarrow \mathcal{G}, g \mapsto [g, 1, 1]$ is open and $G \times G$ -equivariant for the left and right multiplication of G on \mathcal{G} .

We have

$$\begin{aligned} \mathcal{G}^F &= [\{(x, \alpha, b) \in \mathfrak{gl}_2^F \times \mathfrak{gl}_1^F \times \mathfrak{gl}_1^F \mid \det(x) = \alpha^2\} / \mathrm{GL}_1^F \times \mathrm{GL}_1^F] && \text{if } G = \mathrm{SL}_2, \\ \mathcal{G}^F &= [\{(x, \alpha, b) \in \mathfrak{gl}_2^F \times (\mathfrak{gl}_1^F \sqcup \mathfrak{gl}_1^{F'}) \times \mathfrak{gl}_1 \mid \det(x) = \alpha^2\} / \mathrm{GL}_1 \times \mathrm{GL}_1 \times \mu_2] && \text{if } G = \mathrm{PGL}_2, \end{aligned}$$

where $F' : \mathfrak{gl}_1 \rightarrow \mathfrak{gl}_1, x \mapsto -x^q$.

We extend the kernel ϕ^G to a kernel $\phi^{\mathcal{G}}$ on \mathcal{G}^F as follows (see Remark 2.3.2).

If $G = \mathrm{SL}_2$, define $\phi^{\mathcal{G}}$ by

$$\phi^{\mathcal{G}}([x, \alpha, b]) = \sum_{s, t \in \mathbb{F}_q^\times} \psi(s(\mathrm{Tr}(x) + 2\alpha) + tb).$$

If $G = \mathrm{PGL}_2$, define $\phi^{\mathcal{G}}$ by

$$\phi^{\mathcal{G}}([x, \alpha, b]) = \begin{cases} \sum_{s, t \in \mathbb{F}_q^\times, \varepsilon \in \mu_2} \psi(s(\mathrm{Tr}(x) + 2\varepsilon\alpha) + tb) & \text{if } \alpha \in \mathfrak{gl}_1^F, \\ 0 & \text{if } \alpha \in \mathfrak{gl}_1^{F'}. \end{cases}$$

Consider

$$\mathcal{F}^{\mathcal{G}} : C(\mathcal{G}^F) \rightarrow C(\mathcal{G}^F)$$

defined by

$$\mathcal{F}^{\mathcal{G}}(f)([x', \alpha', b']) = \sum_{[x, \alpha, b] \in \mathcal{G}^F} \frac{1}{|\mathrm{Aut}([x, \alpha, b])|} \phi^{\mathcal{G}}([x'x, \alpha'\alpha, b'b]) f([x, \alpha, b])$$

for any function $f \in C(\mathcal{G}^F)$. Here is the main result of our paper.

Theorem 3.3.1. *The Fourier transform $\mathcal{F}^{\mathcal{G}}$ is involutive, namely*

$$\mathcal{F}^{\mathcal{G}} \circ \mathcal{F}^{\mathcal{G}}(f) = q^5 f$$

for any $f \in C(\mathcal{G}^F)$.

3.4 Fourier transforms on quadratic spaces

Assume given a vector space V of dimension $n = 2m + 1$ over \mathbb{F}_q and a non-degenerate quadratic form $Q : V \rightarrow \mathbb{F}_q$ with associated bilinear form

$$B(v', v) = Q(v' + v) - Q(v') - Q(v)$$

for all $v', v \in V$.

From the classification of quadratic forms over finite fields, we can find a basis of V in which Q has the form

$$Q(x_1, \dots, x_{2m+1}) = x_1 x_{2m+1} + \dots + x_m x_{m+2} + c x_{m+1}^2$$

where $c \in \mathbb{F}_q^\times$ is well-determined by Q (up to a multiplication by a square).

Consider the groupoid

$$X = [\{v \in V \mid Q(v) = 0\} / \mathbb{F}_q^\times].$$

Define the kernel

$$K : X \times X \rightarrow \overline{\mathbb{Q}}_\ell, \quad ([v'], [v]) \mapsto \sum_{s \in \mathbb{F}_q^\times} \psi(B(sv', v)) = \sum_{s \in \mathbb{F}_q^\times} \psi(B(v', sv))$$

and the corresponding operator $\mathcal{F}^X : C(X) \rightarrow C(X)$ on the space of functions defined by

$$\mathcal{F}^X(f)([v']) = \sum_{[v] \in X} \frac{1}{|\text{Aut}([v])|} K([v'], [v]) f([v])$$

for all $f \in C(X)$.

The aim of this section is to prove the following result.

Theorem 3.4.1. *The Fourier operator \mathcal{F}^X is involutive, i.e.*

$$\mathcal{F}^X \circ \mathcal{F}^X(f) = q^{2m} f.$$

Remark 3.4.2. The theorem is not true anymore if the dimension of V is even.

To prove Theorem 3.4.1, we first evaluate the sum

$$f(v') = \sum_{v \in V, Q(v)=0} \psi(B(v', v))$$

or equivalently the sum

$$f(v') = \frac{1}{q} \sum_{v \in V, \lambda \in \mathbb{F}_q^\times} \psi(\lambda Q(v) + B(v', v)).$$

We have

$$\begin{aligned} f(v') &= \frac{1}{q} \sum_{v \in V} \psi(B(v', v)) + \frac{1}{q} \sum_{v \in V, \lambda \in \mathbb{F}_q^\times} \psi(\lambda(Q(v) + B(v'/\lambda, v))) \\ &= q^{2m} \delta_0(v') + \frac{1}{q} \sum_{v \in V, \lambda \in \mathbb{F}_q^\times} \psi\left(\lambda\left(Q\left(\frac{v'}{\lambda} + v\right) - Q\left(\frac{v'}{\lambda}\right)\right)\right) \\ &= q^{n-1} \delta_0(v') + \frac{1}{q} \sum_{\lambda_1 \in \mathbb{F}_q^\times} \psi(-\lambda_1 Q(v')) \sum_{v_1 \in V} \psi(\lambda_1^{-1} Q(v_1)) \end{aligned}$$

by the variables change $v_1 = \frac{v'}{\lambda} + v$ and $\lambda_1 = 1/\lambda$.

A straightforward calculation shows that

$$\sum_{v \in V} \psi(\lambda Q(v)) = q^m \alpha_o(c\lambda) S(\alpha_o, \psi).$$

Therefore, setting $\alpha_o(0) = 0$ and noticing that

$$\sum_{x \in \mathbb{F}_q} \psi(yx) \alpha_o(x) = \alpha_o(y) S(\alpha_o, \psi)$$

we get

$$\begin{aligned} f(v') &= q^{2m} \delta_0(v') + q^{m-1} S(\alpha_o, \psi) \alpha_o(c) \sum_{\lambda_1 \in \mathbb{F}_q} \psi(-\lambda_1 Q(v')) \alpha_o(\lambda_1) \\ &= q^{2m} \delta_0(v') + q^{m-1} S(\alpha_o, \psi)^2 \alpha_o(c) \alpha_o(-Q(v')) \end{aligned}$$

and using that $S(\alpha_o, \psi)^2 = \alpha_o(-1)q$ we end up with

$$f(v') = q^{2m} \delta_0(v') + q^m \alpha_o(c) \alpha_o(Q(v')). \quad (3.2)$$

We now prove Theorem 3.4.1.

Proof of Theorem 3.4.1. By Lemma 3.1.1 we need to prove that for $[v], [v''] \in X$ we have

$$\Delta([v''], [v]) = \sum_{[v'] \in X} \frac{1}{|\text{Aut}([v'])|} K([v''], [v']) K([v'], [v]) = \begin{cases} q^{2m} |\text{Aut}([v])| & \text{if } [v''] = [v], \\ 0 & \text{otherwise.} \end{cases}$$

We have

$$\begin{aligned} \Delta([v''], [v]) &= \sum_{s, s'' \in \mathbb{F}_q^\times} \sum_{[v'] \in X} \frac{1}{|\text{Aut}([v'])|} \psi(B(s''v'', v')) \psi(B(v', sv)) \\ &= \sum_{s, s''} \sum_{[v']} \frac{1}{|\text{Aut}([v'])|} \psi(B(sv + s''v'', v')) \end{aligned}$$

Notice that

$$\begin{aligned} \sum_{s \in \mathbb{F}_q^\times} \sum_{v', Q(v')=0} \psi(B(sX, v')) &= \sum_s \left(1 + \sum_{[v'] \neq 0} \sum_{\sigma \in \mathbb{F}_q^\times} \psi(B(sX, \sigma v')) \right) \\ &= (q-1) + \sum_{[v'] \neq 0} \sum_{\sigma} \sum_s \psi(B(sX, \sigma v')) \\ &= (q-1) + \sum_{[v'] \neq 0} \sum_{\sigma} \sum_s \psi(B(sX, v')) \\ &= \sum_s \left(1 + (q-1) \sum_{[v'] \neq 0} \psi(B(sX, v')) \right) \\ &= (q-1) \sum_s \sum_{[v'] \in X} \frac{1}{|\text{Aut}([v'])|} \psi(B(sX, v')) \end{aligned}$$

Therefore

$$\begin{aligned} \Delta([v''], [v]) &= \sum_{s, s''} \sum_{[v'] \in X} \frac{1}{|\text{Aut}([v'])|} \psi(B(sv + s''v'', v')) \\ &= \frac{1}{q-1} \sum_{s, s''} \sum_{v', Q(v')=0} \psi(B(sv + s''v'', v')) \end{aligned}$$

From Formula (3.2) we get

$$\Delta([v''], [v]) = \frac{1}{q-1} \left(q^{2m} \sum_{s, s''} \delta_o(sv + s''v'') + q^m \alpha_o(c) \sum_{s, s''} \alpha_o(Q(sv + s''v'')) \right)$$

The first sum equals $(q-1)|\text{Aut}([v])|$ if $[v] = [v'']$ and 0 otherwise. Since

$$Q(sv + s''v'') = B(sv, s''v'') + Q(sv) + Q(s''v'') = ss''B(v, v'')$$

we get that

$$\sum_{s, s''} \alpha_o(Q(sv + s''v'')) = \sum_{s, s''} \alpha_o(ss''B(v, v'')) = 0$$

because α_o is a non-trivial character on \mathbb{F}_q^\times and $\alpha_o(0) = 0$.

We deduce that

$$\Delta([v''], [v]) = \begin{cases} q^{2m} |\text{Aut}([v])| & \text{if } [v] = [v''], \\ 0 & \text{otherwise.} \end{cases}$$

□

3.5 Proof of Theorem 3.3.1

3.5.1. Assume first that $G = \text{SL}_2$.

In this case we have

$$\mathcal{G} = \left[\{(x, \alpha) \in \mathfrak{gl}_2 \times \mathfrak{gl}_1 \mid \det(x) = \alpha^2\} / \text{GL}_1 \right] \times [\mathfrak{gl}_1 / \text{GL}_1].$$

Put

$$\mathcal{G}_1 := \left[\{(x, \alpha) \in \mathfrak{gl}_2 \times \mathfrak{gl}_1 \mid \det(x) = \alpha^2\} / \text{GL}_1 \right], \quad \mathcal{G}_2 := [\mathfrak{gl}_1 / \text{GL}_1].$$

Then

$$\phi^{\mathcal{G}} = \phi^{\mathcal{G}_1} \boxtimes \phi^{\mathcal{G}_2}$$

where

$$\phi^{\mathcal{G}_1}([x, \alpha]) := \sum_{s \in \mathbb{F}_q^\times} \psi(s(\text{Tr}(x) + 2\alpha)), \quad \phi^{\mathcal{G}_2}([z]) = \sum_{s \in \mathbb{F}_q^\times} \psi(sz).$$

Since the Fourier transform $\mathcal{F}^{\mathcal{G}_2}$ with respect to the kernel $\phi^{\mathcal{G}_2}$ is clearly involutive, it remains to see that the Fourier transform $\mathcal{F}^{\mathcal{G}_1}$ with kernel $\phi^{\mathcal{G}_1}$ is involutive.

Consider

$$Q : (\mathfrak{gl}_2 \times \mathfrak{gl}_1)^F \rightarrow \mathbb{F}_q^\times, \quad (x, \alpha) \mapsto \det(x) - \alpha^2.$$

The associated bilinear form is given by

$$B((y, \beta), (x, \alpha)) = \text{Tr}(\iota(y)x) - 2\beta\alpha$$

where $\iota : \mathfrak{gl}_2 \rightarrow \mathfrak{gl}_2$ is the involution

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$$

sending a matrix to the transpose of its co-factor matrix.

By Theorem 3.4.1, the Fourier transform $\hat{\mathcal{F}}^{\mathcal{G}_1} : C(\mathcal{G}_1^F) \rightarrow C(\mathcal{G}_1^F)$ defined by

$$\hat{\mathcal{F}}^{\mathcal{G}_1}(f)([y, \beta]) = \sum_{[x, \alpha]} \frac{1}{|\text{Aut}([x, \alpha])|} \phi^{\mathcal{G}_1}([\iota(y)x, \alpha\beta]) f([x, \alpha])$$

is involutive from which we deduce the involutivity of $\mathcal{F}^{\mathcal{G}_1}$ as $\text{Tr}(\iota(y)\iota(x)) = \text{Tr}(yx)$ for all $x, y \in \mathfrak{gl}_2$.

3.5.2. Let us now assume that $G = \text{PGL}_2$ in which case we have

$$\mathcal{G} = \left[\left\{ (x, \alpha, b) \in \mathfrak{gl}_2 \times \mathfrak{gl}_1 \times \mathfrak{gl}_1 \mid \det(x) = \alpha^2 \right\} / \text{GL}_1 \times \text{GL}_1 \times \mu_2 \right].$$

If we put $X = \{(x, \alpha, b) \in \mathfrak{gl}_2 \times \mathfrak{gl}_1 \times \mathfrak{gl}_1 \mid \det(x) = \alpha^2\}$, then we get a μ_2 -torsor

$$\tilde{f} : [X/\text{GL}_1 \times \text{GL}_1] \longrightarrow [X/\text{GL}_1 \times \text{GL}_1 \times \mu_2] = \mathcal{G}.$$

which extends the natural quotient $f : \text{SL}_2 \rightarrow \text{PGL}_2$ (notice that $[X/\text{GL}_1 \times \text{GL}_1]$ is the stack \mathcal{G} for $G = \text{SL}_2$). If we denote by ϕ (instead of $\phi^{\mathcal{G}}$ to avoid any confusion) the kernel on $[X/\text{GL}_1 \times \text{GL}_1]^F$ studied in the previous section §3.5.1 we get that

$$\phi^{\mathcal{G}} = \tilde{f}_1^F(\phi)$$

extending the formula (2.6). Therefore, by §3.5.1 and the result of §3.1 we get Theorem 3.3.1 for $G = \text{PGL}_2$.

3.6 Remarks on the torus case

Recall, see Formula (2.3), that the BK-Fourier kernel on T^F is given by

$$\phi^T = \rho_1^F(\psi \circ \text{Tr})$$

where $\rho : T_3 \rightarrow T$ is given by

$$\begin{aligned} \rho(a, b, c) &= (a/c, c/a) & \text{if } G = \text{SL}_2, \\ \rho(a, b, c) &= [a^2b, bc^2] & \text{if } G = \text{PGL}_2. \end{aligned}$$

The following morphism induces a bijection between isomorphism classes of objects

$$\tilde{\rho} : [t_3/\text{Ker}(\rho)] \rightarrow \mathcal{T}, \quad [a, b, c] \mapsto [(a^2, c^2), ac, b]$$

where

$$\begin{aligned} \mathcal{T} &= \left[\left\{ (x, \alpha, b) \in t_2 \times \mathfrak{gl}_1 \times \mathfrak{gl}_1 \mid \det(x) = \alpha^2 \right\} / \text{GL}_1 \times \text{GL}_1 \right] && \text{if } G = \text{SL}_2, \\ \mathcal{T} &= \left[\left\{ (x, \alpha, b) \in t_2 \times \mathfrak{gl}_1 \times \mathfrak{gl}_1 \mid \det(x) = \alpha^2 \right\} / \text{GL}_1 \times \mu_2 \times \text{GL}_1 \right] && \text{if } G = \text{PGL}_2. \end{aligned}$$

Moreover, the size of the stabilizers are the same on both sides of $\tilde{\rho}$. Therefore, since the kernel $\phi^{[t_3/\text{Ker}(\rho)]}$ from §3.2 defines an involutive Fourier transforms $\mathcal{F}^{[t_3/\text{Ker}(\rho)]}$, the kernel

$$\phi^{\mathcal{T}} := \tilde{\rho}_!^F(\phi^{[t_3/\text{Ker}(\rho)]})$$

defines also an involutive Fourier transforms $\mathcal{F}^{\mathcal{T}}$ on $C(\mathcal{T}^F)$.

Notice that even if $\tilde{\rho}$ is not an isomorphism of stacks (as $[t_3/\text{Ker}(\rho)]$ is smooth and \mathcal{T} is singular), both embeddings

$$T \simeq T_3/\text{Ker}(\rho) \hookrightarrow [t_3/\text{Ker}(\rho)], \quad T \hookrightarrow \mathcal{T}$$

provide a good space of functions with an involutive Fourier transform on it that extends the BK-Fourier operator $\mathcal{F}^T = \mathcal{F}_\rho^T : C(T^F) \rightarrow C(T^F)$. However, the first one does not have an analogue for the group G while the second one does.

Following [9], we can extend the definition of Deligne-Lusztig induction R_T^G and $R_{T_\sigma}^G$ to operators

$$R_{\mathcal{T}}^{\mathcal{G}} : C(\mathcal{T}^F) \rightarrow C(\mathcal{G}^F), \quad R_{\mathcal{T}_\sigma}^{\mathcal{G}} : C(\mathcal{T}_\sigma^F) \rightarrow C(\mathcal{G}^F).$$

Then we can verify that

$$\phi^{\mathcal{G}} = \frac{1}{2} \left(R_{\mathcal{T}}^{\mathcal{G}}(\phi^{\mathcal{T}}) + R_{\mathcal{T}_\sigma}^{\mathcal{G}}(\phi^{\mathcal{T}_\sigma}) \right) \tag{3.3}$$

where $\phi^{\mathcal{T}_\sigma}$ is obtained by pushing $\psi \circ \text{Tr}$ along the map $t_3^{\sigma F} \rightarrow \mathcal{T}^{\sigma F}$.

4 Extending the BK-Fourier kernel: The case $G = \text{GL}_2$

In this section q is assumed to be odd.

4.1 The stack \mathcal{G}

Assume that $G = \mathrm{GL}_2$ with $\rho^\vee = \mathrm{Sym}^2$. By analogy with the SL_2 and PGL_2 cases, the results of §2.5 suggest to consider the stack

$$\mathcal{G} := \left[\left\{ (x, \alpha, b) \in \mathfrak{gl}_2 \times \mathfrak{gl}_1 \times \mathfrak{gl}_1 \mid \det(x) = \alpha^2 \right\} / \mathrm{GL}_1 \times \mu_2 \right]$$

where GL_1 acts as $t \cdot [x, \alpha, b] = [tx, t\alpha, t^{-1}b]$ and μ_2 acts on the second coordinate as $\alpha \mapsto -\alpha$.

Notice that the open substack of $[x, \alpha, b]$ with $\alpha, b \neq 0$ is isomorphic to GL_2 via the projection

$$(x, \alpha, b) \mapsto bx.$$

We thus have a $G \times G$ -equivariant embedding

$$G \hookrightarrow \mathcal{G}$$

for the right and left multiplication.

Consider the toric analogue of \mathcal{G}

$$\mathcal{T} := \left[\left\{ (x, \alpha, b) \in \mathfrak{t}_2 \times \mathfrak{gl}_1 \times \mathfrak{gl}_1 \mid \det(x) = \alpha^2 \right\} / \mathrm{GL}_1 \times \mu_2 \right]$$

Then $\rho : T_3 \rightarrow T$, $(a, b, c) \mapsto (a^2b, bc^2)$ extends to a morphism of stacks

$$\tilde{\rho} : [t_3/\mathrm{Ker}(\rho)] \rightarrow \mathcal{T}, \quad [a, b, c] \mapsto [(a^2, c^2), ac, b]. \quad (4.1)$$

Notice that $\tilde{\rho}$ is an isomorphism on the open subset $\{(a, c) \neq (0, 0)\}$ but not globally as \mathcal{T} is singular.

Above $\{[0, b, 0]\}$ with $b \neq 0$, the morphism becomes

$$\left([b \neq 0] / \mathrm{GL}_1 \right) \rightarrow [b \neq 0] / \mathrm{GL}_1 = \mathrm{Spec}(\overline{\mathbb{F}}_q) \times \mathrm{B}(\mu_2)$$

where $s \in \mathrm{GL}_1$ acts by s^2 on the source and by s on the target.

Therefore the morphism of stacks (4.1) is not a bijection between isomorphism classes of objects (unlike the cases of SL_2 and PGL_2).

4.2 The BK-Fourier kernels $\phi^{\mathcal{T}}, \phi^{\mathcal{T}\sigma}$

Recall that

$$\mathrm{Ker}(\rho) = \{(s, s^{-2}, \varepsilon s) \mid \varepsilon \in \mu_2, s \in \mathrm{GL}_1\}.$$

Let $q : t_3 \rightarrow [t_3/\mathrm{Ker}(\rho)]$ be the quotient map.

Consider on $[t_3/\mathrm{Ker}(\rho)]^F$ and on $[t_3/\mathrm{Ker}(\rho)]^{\sigma F}$ the kernels

$$\phi^F = q_1^F(\psi \circ \text{Tr}), \quad \phi^{\sigma F} = q_1^{\sigma F}(\psi \circ \text{Tr})$$

i.e.

$$\phi^F([a, b, c]) = \sum_{s \in \mathbb{F}_q^\times, \varepsilon \in \mu_2} \psi(s(a + \varepsilon c) + s^{-2}b), \quad \phi^{\sigma F}([a', b, a'^q]) = \sum_{s^q = \pm s} \psi(s(a' + \varepsilon a'^q) + b s^{-2})$$

considered in §3.2.

Analogously to the cases $G = \text{SL}_2, \text{PGL}_2$ (see §3.6), we consider the kernels

$$\phi^{\mathcal{T}} := -(\tilde{\rho}^F)_!(\phi^F), \quad \text{and} \quad \phi^{\mathcal{T}\sigma} := -(\tilde{\rho}^{\sigma F})_!(\phi^{\sigma F}).$$

They are given by the formulas

$$\begin{aligned} \phi^{\mathcal{T}}([(a^2, c^2), ac, b]) &= - \sum_{s \in \mathbb{F}_q^\times, \varepsilon \in \mu_2} \psi(s(a + \varepsilon c) + s^{-2}b) \\ \phi^{\mathcal{T}\sigma}([a'^2, a'^{2q}, a'^{q+1}, b]) &= - \sum_{s^q = \pm s} \psi(s(a' + \varepsilon a'^q) + b s^{-2}) \end{aligned}$$

By operating the variable change $s(a + \varepsilon c) \leftrightarrow s'$ when $a + \varepsilon c \neq 0$ we get

$$\phi^{\mathcal{T}}([(a^2, c^2), ac, b]) = \begin{cases} - \sum_{s, \varepsilon} \psi(s + s^{-2}b(a^2 + c^2 + 2\varepsilon ac)) & \text{if } a^2 \neq c^2 \\ -\kappa(b) - \sum_s \psi(s + 4s^{-2}ba^2) & \text{if } a = \varepsilon c \neq 0 \\ -2\kappa(b) & \text{if } a = c = 0 \end{cases}$$

We thus get the values on the elements $[x, \alpha, b]$ as follows

| | $\left[\begin{pmatrix} a^2 & 0 \\ 0 & a^2 \end{pmatrix}, a^2, b \right]$ $a, b \in \mathbb{F}_q$ | $\left[\begin{pmatrix} a^2 & 0 \\ 0 & c^2 \end{pmatrix}, ac, b \right]$ $a, c \in \mathbb{F}_q, a^2 \neq c^2$ | $\left[\begin{pmatrix} a'^2 & 0 \\ 0 & a'^{2q} \end{pmatrix}, a'^{q+1}, b \right]$ $a'^2 \neq a'^{2q}$ | |
|----------------------------|---|---|--|-------|
| $\phi^{\mathcal{T}}$ | $\begin{cases} -\kappa(b) - 1 - \Phi([x, \alpha, b]) & \text{if } a \neq 0 \\ -2\kappa(b) & \text{if } a = 0 \end{cases}$ | $-\Phi([x, \alpha, b])$ | \times | (4.2) |
| $\phi^{\mathcal{T}\sigma}$ | $\begin{cases} -\kappa'(b) - 1 - \Phi([x, \alpha, b]) & \text{if } a \neq 0 \\ -2\kappa'(b) & \text{if } a = 0 \end{cases}$ | \times | $-\Phi([x, \alpha, b])$ | |

where for $[x, \alpha, b] \in \mathcal{G}^F$ we put

$$\Phi([x, \alpha, b]) := \sum_{s \in \mathbb{F}_q^\times, \varepsilon \in \mu_2} \psi(s + s^{-2}b(\text{Tr}(x) + 2\varepsilon\alpha)).$$

To avoid any confusion put

$$\overline{\mathcal{T}} := \left[\left\{ (x, \alpha, b) \in \mathfrak{t}_2 \times \mathfrak{gl}_1 \times \mathfrak{gl}_1 \mid \det(x) = \alpha^2 \right\} / \mathrm{GL}_1 \times \mu_2 \times \mathrm{GL}_1 \right]$$

for the toric stack \mathcal{T} is the $G = \mathrm{PGL}_2$ case (see §3.6).

Denote by $\tilde{\pi} : \mathcal{T} \rightarrow \overline{\mathcal{T}}$ the canonical GL_1 -torsor that extends the quotient $\pi : T_2 \rightarrow T_2/\mathrm{GL}_1$.

Proposition 4.2.1. *We have*

$$\tilde{\pi}_1^F(\phi^{\mathcal{T}}) = -\phi^{\overline{\mathcal{T}}}, \quad \tilde{\pi}_1^{\sigma F}(\phi^{\mathcal{T}\sigma}) = -\phi^{\overline{\mathcal{T}\sigma}}.$$

Proof. This is clear from the diagram

$$\begin{array}{ccc} & \mathfrak{t}_3 & \\ \tilde{\rho} \circ q \swarrow & & \searrow \\ \mathcal{T} & \xrightarrow{\tilde{\pi}} & \overline{\mathcal{T}} \end{array}$$

as $\phi^{\mathcal{T}}$ and $\phi^{\overline{\mathcal{T}}}$ are the pushforward of $\psi \circ \mathrm{Tr}$ along the diagonal arrows. \square

Let $\mathcal{F}^{\mathcal{T}} : C(\mathcal{T}^F) \rightarrow C(\overline{\mathcal{T}}^F)$ be the Fourier operator defined by the kernel $\phi^{\mathcal{T}}$. From the above proposition we thus have the following commutative diagram

$$\begin{array}{ccc} C(\overline{\mathcal{T}}^F) & \xrightarrow{\tilde{\pi}^*} & C(\mathcal{T}^F) \\ \mathcal{F}^{\overline{\mathcal{T}}} \uparrow & & \uparrow -\mathcal{F}^{\mathcal{T}} \\ C(\overline{\mathcal{T}}^F) & \xrightarrow{\tilde{\pi}^*} & C(\mathcal{T}^F) \end{array} \quad (4.3)$$

Remark 4.2.2. (Non-involutivity) The Fourier operator $\mathcal{F}^{\mathcal{T}}$ defined from $\phi^{\mathcal{T}} = \tilde{\rho}_1^F(\phi^F)$ is not involutive. This follows from Lemma 3.1.1 as a direct calculation shows that

$$\Delta([(1, 0), 0, 1], [(0, 1), 0, 1]) = 4\alpha_o(-1)q \neq 0.$$

4.3 The BK-Fourier kernel $\phi^{\mathcal{G}}$

We define the BK-Fourier kernel $\phi^{\mathcal{G}}$ on \mathcal{G}^F by the formula (3.3).

We compute the values of the BK-Fourier kernel $\phi^{\mathcal{G}}$ on \mathcal{G}^F : we have $\phi^{\mathcal{G}}([x, \alpha, b]) = 0$ if

$\alpha \in \mathfrak{gl}_1^{F'}$ and otherwise it is given by the following table

| | | | | |
|----------------------|---|--|---|---|
| | $\left[\begin{pmatrix} a^2 & 0 \\ 0 & a^2 \end{pmatrix}, a^2, b \right]$ $a, b \in \mathbb{F}_q$ | $\left[\begin{pmatrix} a^2 & 0 \\ 0 & c^2 \end{pmatrix}, ac, b \right]$ $a, b, c \in \mathbb{F}_q, a^2 \neq c^2$ | $\left[\begin{pmatrix} x^2 & 0 \\ 0 & x^{2q} \end{pmatrix}, x^{q+1}, b \right]$ $x^2 \neq x^{2q}, b \in \mathbb{F}_q$ | $\left[\begin{pmatrix} a^2 & 1 \\ 0 & a^2 \end{pmatrix}, a^2, b \right]$ $a, b \in \mathbb{F}_q$ |
| $\phi^{\mathcal{G}}$ | $\begin{cases} -(qS(\alpha_o, \psi_b)\delta_{b \neq 0} + q\delta_{b=0} + \Phi([x, \alpha, b])) & \text{if } a \neq 0 \\ -2(qS(\alpha_o, \psi_b)\delta_{b \neq 0} + q\delta_{b=0} - 1) & \text{if } a = 0 \end{cases}$ | $-\Phi([x, \alpha, b])$ | $-\Phi([x, \alpha, b])$ | $\begin{cases} -(q\delta_{b=0} + \Phi([x, \alpha, b])) & \text{if } a \neq 0 \\ -2(q\delta_{b=0} - 1) & \text{if } a = 0 \end{cases}$ |

(4.4)

We can see by a direct calculation that the operator $\mathcal{F}^{\mathcal{G}}$ on $C(\mathcal{G}^F)$ defined from $\phi^{\mathcal{G}}$ is not involutive (see Remark 4.2.2).

As in the toric case (see Diagram (4.3)) we have the following commutative diagram

$$\begin{array}{ccc}
 C(\overline{\mathcal{G}}^F) & \xrightarrow{\tilde{\pi}^*} & C(\mathcal{G}^F) \\
 \mathcal{F}^{\overline{\mathcal{G}}} \uparrow & & \uparrow -\mathcal{F}^{\mathcal{G}} \\
 C(\overline{\mathcal{G}}^F) & \xrightarrow{\tilde{\pi}^*} & C(\mathcal{G}^F)
 \end{array} \tag{4.5}$$

where $\overline{\mathcal{G}}$ is the stack \mathcal{G} in the $G = \mathrm{PGL}_2$ case and $\tilde{\pi} : \mathcal{G} \rightarrow \overline{\mathcal{G}}$ is the canonical GL_1 -torsor that extends $\pi : \mathrm{GL}_2 \rightarrow \mathrm{PGL}_2$.

5 The characteristic 2 case

In this section we assume that \mathbb{F}_q is a finite field of characteristic 2 (every element of \mathbb{F}_q is then a square in \mathbb{F}_q).

We consider only the case $G = \mathrm{GL}_2$ (the cases of SL_2 and PGL_2 are similar).

5.1 BK-Fourier operator on G

First of all, notice that since the characteristic is 2 we have

$$\mathrm{Ker}(\rho^F) = \mathrm{Ker}(\rho^{\sigma F}) = \{(s, s^{-2}, s) \mid s \in \mathbb{F}_q^\times\}.$$

Following the calculation in §2.5 we find

$$\phi^T([(a^2, c^2), ac, b]) = \begin{cases} -\sum_{s \in \mathbb{F}_q^\times} \psi(s + s^{-2}(a^2 + c^2)b) & \text{if } a^2 \neq c^2 \text{ i.e. } a \neq c, \\ -\kappa(b) & \text{if } a^2 = c^2. \end{cases}$$

and

$$\phi^{T_\sigma}([(a'^2, a'^{2q}), a'^{q+1}, b]) = \begin{cases} -\sum_{s \in \mathbb{F}_q^\times} \psi(s + s^{-2}(a'^2 + a'^{2q})b) & \text{if } a'^2 \neq a'^{2q} \text{ i.e. } a'^q \neq a', \\ -\kappa(b) & \text{otherwise.} \end{cases}$$

Since the characteristic is 2, the identification (2.7) becomes

$$G \simeq \{(x, \alpha, b) \in \mathrm{GL}_2 \times \mathrm{GL}_1 \times \mathrm{GL}_1 \mid \det(x) = \alpha^2\} / \mathrm{GL}_1$$

Writing the elements of G as $[x, \alpha, b]$ we get the values of the BK-Fourier kernel as follows

| | | | | | |
|----------|--|--|---|--|-------|
| | $\left[\begin{pmatrix} a^2 & 0 \\ 0 & a^2 \end{pmatrix}, a^2, b \right]$ $a, b \in \mathbb{F}_q$ | $\left[\begin{pmatrix} a^2 & 0 \\ 0 & c^2 \end{pmatrix}, ac, b \right]$ $a, b, c \in \mathbb{F}_q, a^2 \neq c^2$ | $\left[\begin{pmatrix} x^2 & 0 \\ 0 & x^{2q} \end{pmatrix}, x^{q+1}, b \right]$ $x^2 \neq x^{2q}, b \in \mathbb{F}_q$ | $\left[\begin{pmatrix} a^2 & 1 \\ 0 & a^2 \end{pmatrix}, a^2, b \right]$ $a, b \in \mathbb{F}_q$ | |
| ϕ^G | $-\kappa(b)$ | $-\sum_{s \in \mathbb{F}_q^\times} \psi(s + s^{-2}b \mathrm{Tr}(x))$ | $-\sum_{s \in \mathbb{F}_q^\times} \psi(s + s^{-2}b \mathrm{Tr}(x))$ | $-\kappa(b)$ | (5.1) |

5.2 Extending the BK-Fourier operator

Since the characteristic is 2, the morphism

$$\rho : T_3 \rightarrow T, \quad (a, b, c) \mapsto (a^2b, bc^2)$$

extends to a group morphism

$$\rho : \mathrm{GL}_2 \times \mathrm{GL}_1 \rightarrow \mathrm{GL}_2, \quad (x', b) \mapsto \iota_2(x')b$$

where $\iota_2 : \mathfrak{gl}_2 \rightarrow \mathfrak{gl}_2$ is the algebra isomorphism defined by

$$\iota_2 \begin{pmatrix} a & b \\ c & d \end{pmatrix} := \begin{pmatrix} a^2 & b^2 \\ c^2 & d^2 \end{pmatrix}.$$

Remark 5.2.1. Notice that

$$\phi^G = -\rho_!(\psi \circ \mathrm{Tr})$$

where $\mathrm{Tr}(x', b) = \mathrm{Tr}(x') + b$ for $(x', b) \in \mathfrak{gl}_2 \times \mathfrak{gl}_1$.

The kernel of ρ is

$$\mathrm{Ker}(\rho) = \left\{ \left(\begin{pmatrix} s & 0 \\ 0 & s \end{pmatrix}, s^{-2} \right) \mid s \in \mathrm{GL}_1 \right\}.$$

We consider the $G \times G$ -equivariant open embedding

$$G \simeq (\mathrm{GL}_2 \times \mathrm{GL}_1)/\mathrm{Ker}(\rho) \hookrightarrow \mathcal{G} := [(\mathfrak{gl}_2 \times \mathfrak{gl}_1)/\mathrm{Ker}(\rho)]. \quad (5.2)$$

We then consider on \mathcal{G}^F the kernel

$$\phi^{\mathcal{G}}([x', b]) = - \sum_{s \in \overline{\mathbb{F}}_q^\times} \psi(s \mathrm{Tr}(x') + s^{-2}b)$$

Consider the two-variable kernel $K^{\mathcal{G}} : \mathcal{G}^F \times \mathcal{G}^F \rightarrow \overline{\mathbb{Q}}_\ell$ defined by

$$K^{\mathcal{G}}([x'_1, b_1], [x'_2, b_2]) = \phi^{\mathcal{G}}([x'_1 x'_2, b_1 b_2]).$$

Theorem 5.2.2. (1) The Fourier operator $\mathcal{F}^{\mathcal{G}} : C(\mathcal{G}^F) \rightarrow C(\mathcal{G}^F)$ defined from $K^{\mathcal{G}}$ is involutive. (2) $\mathcal{F}^{\mathcal{G}}$ extends the BK-Fourier operator \mathcal{F}^G .

Proof. The first assertion follows from Lemma 3.1.2 and the fact that the Fourier transform on $\mathfrak{gl}_2 \times \mathfrak{gl}_1$ defined from the kernel

$$(x', b) \mapsto \psi(\mathrm{Tr}(x') + b)$$

is involutive. For the second assertion we need to see that the restriction of $\phi^{\mathcal{G}}$ to G along the embedding (5.2) is the kernel ϕ^G whose values are given by the Table (5.1). To see that we use (when $\mathrm{Tr}(x') \neq 0$) the variable change $s \mathrm{Tr}(x') \leftrightarrow s$ which allows us to pass from

$$\sum_s \psi(s \mathrm{Tr}(x') + s^{-2}b) \quad \text{to} \quad \sum_s \psi(s + s^{-2}b \mathrm{Tr}(x))$$

with $x = \iota_2(x')$.

□

References

- [1] BRAVERMAN, A. AND KAZHDAN, D.: *γ -functions of representations and lifting* (with an appendix by V. Vologodsky), GAFA 2000, Geom. Funct. Anal. 2000, Special Volume, Part I, 237–278.
- [2] BRAVERMAN, A. AND KAZHDAN, D.: *γ -sheaves on reductive groups*, Studies in memory of Issai Schur (Chevaleret/Rehovot, 2000), Progr. Math. **210**, Birkhäuser Boston, Boston, MA (2003), 22–47.
- [3] CHEN, T.-H.: *On a conjecture of Braverman-Kazhdan*, J. Amer. Math. Soc., to appear (arXiv:1909.05467v2).

- [4] CHENG, S. AND NGÔ, B. C.: *On a conjecture of Braverman and Kazhdan*, Int. Math. Res. Not. IMRN, **20** (2018), 6177–6200.
- [5] GODEMENT, R. AND JACQUET, H.: *Zeta functions of simple algebras*, Lecture Notes in Math. **260**, Springer-Verlag (1972).
- [6] LAFFORGUE, L.: *Du transfert automorphe de Langlands aux formules de Poisson non linéaires*, Ann. Inst. Fourier **66** (2016), 899–1012.
- [7] LAUMON, G. AND LETELLIER, E.: *On derived Lusztig correspondence*, arXiv:2011.13768.
- [8] LAUMON, G. AND LETELLIER, E.: *Note on a conjecture of Braverman-Kazhdan*, arXiv:1906.07476.
- [9] LETELLIER, E.: *Fourier transforms of invariant functions on finite reductive Lie algebras*, *Lecture Notes in Mathematics, 1859*. Springer-Verlag, Berlin, 2005. xii+165 pp.
- [10] TATE, J.: *Fourier Analysis in Number Fields and Hecke's Zeta-Functions*, Ph.D. thesis, Princeton University, Princeton, NJ, 1950.