

**COMBINATORIAL RELATIONS AMONG RELATIONS FOR
LEVEL 2 STANDARD $C_n^{(1)}$ -MODULES**

MIRKO PRIMC AND TOMISLAV ŠIKIĆ

ABSTRACT. For an affine Lie algebra $\hat{\mathfrak{g}}$ the coefficients of certain vertex operators which annihilate level k standard $\hat{\mathfrak{g}}$ -modules are the defining relations for level k standard modules. In this paper we study a combinatorial structure of the leading terms of these relations for level $k = 2$ standard $\hat{\mathfrak{g}}$ -modules for affine Lie algebras of type $C_n^{(1)}$ and the main result is a construction of combinatorially parameterized relations among the coefficients of annihilating fields. It is believed that the constructed relations among relations will play a key role in a construction of Groebner-like basis of the maximal ideal of the universal vertex operator algebra $V_{\hat{\mathfrak{g}}}^k$ for $k = 2$.

1. INTRODUCTION

Let \mathfrak{g} be a simple complex Lie algebra and $\hat{\mathfrak{g}}$ the corresponding affine Kac-Moody Lie algebra (cf. [K]). In [MP1] combinatorial bases of all standard (i.e. integrable highest weight) $\hat{\mathfrak{g}}$ -modules $L(\Lambda)$ were constructed for $\mathfrak{g} = \mathfrak{sl}_2$. A part of that construction can be carried through for the vacuum $\hat{\mathfrak{g}}$ -modules $L(k\Lambda_0)$ for all simple \mathfrak{g} and all positive integer k . The key ingredient is the vertex operator algebra structure¹ on the generalized Verma $\hat{\mathfrak{g}}$ -module $N(k\Lambda_0)$ generated by fields

$$x(z) = \sum_{m \in \mathbb{Z}} x(m)z^{-m-1} = Y(x(-1)\mathbf{1}, z), \quad x \in \mathfrak{g},$$

where $x(m) = x_m = x \otimes t^m$ is the usual notation for the elements of the affine Lie algebra and $Y(v, z)$ is the usual notation for vertex operators. We also write $\mathfrak{g} = \mathfrak{g} \otimes t^0 \subset \hat{\mathfrak{g}}$ and $\hat{\mathfrak{g}}_{<0} = \coprod_{m < 0} \mathfrak{g} \otimes t^m$.

Let R be the finite dimensional \mathfrak{g} -module generated by the singular vector in $N(k\Lambda_0)$, i.e.

$$R = U(\mathfrak{g}) \cdot x_{\theta}(-1)^{k+1} \mathbf{1} \cong L_{\mathfrak{g}}((k+1)\theta),$$

where x_{θ} is a root vector for the maximal root θ with respect to a chosen Cartan decomposition of \mathfrak{g} . Then the coefficients $r(m)$, $r \in R$, $m \in \mathbb{Z}$, of vertex operators

$$Y(r, z) = \sum_{m \in \mathbb{Z}} r(m)z^{-m-k-1}$$

span a loop $\hat{\mathfrak{g}}$ -module \bar{R} . Since $\bar{R}N(k\Lambda_0) \subset N(k\Lambda_0)$ is the maximal submodule of the generalized Verma module (cf. [MP1], [PŠ1], [PŠ2] and the references therein) we have

$$(1.1) \quad L(k\Lambda_0) = N(k\Lambda_0)/\bar{R}N(k\Lambda_0) \quad \text{and} \quad \bar{R}|_{L(k\Lambda_0)} = 0,$$

^{2000 Mathematics Subject Classification.} Primary 17B67; Secondary 17B69, 05A19.

¹ $N(k\Lambda_0)$ is often denoted as $V^k(\mathfrak{g})$ or $V_{\hat{\mathfrak{g}}}(k, 0)$ (cf. [LL]), but we shall use the notation from [PŠ1] and [PŠ2].

and for this reason we call the elements of \bar{R} *the relations* for level k standard (vacuum) $\hat{\mathfrak{g}}$ -modules. Note that (1.1) is the consequence of representation theory of affine Kac-Moody Lie algebras and the associated vertex operator algebras.

One can use the relations \bar{R} to construct a combinatorial bases of $L(k\Lambda_0)$ —the basic idea is to reduce the PBW spanning set of $L(k\Lambda_0)$ to a basis \mathcal{B} by using relations $r|_{L(k\Lambda_0)} = 0$, and to parameterize the monomial vectors

$$u(\pi)\mathbf{1} \in \mathcal{B} \subset L(k\Lambda_0) = U(\hat{\mathfrak{g}})\mathbf{1}$$

with monomials π in the symmetric algebra $S(\hat{\mathfrak{g}})$. This is done in steps:

- Choose an (appropriately) totally ordered weight basis B of \mathfrak{g} and the corresponding basis $\bar{B} = \bar{B} \cup \{c\}$ of $\hat{\mathfrak{g}}$, where c is the canonical central element in $\hat{\mathfrak{g}}$ and

$$\bar{B} = \{b(m) \mid b \in B, m \in \mathbb{Z}\}.$$

We extend the strict order $<$ on B to \bar{B} so that $m < m'$ implies $b(m) < b'(m')$. Since $b(m)\mathbf{1} = 0$ for $m \geq 0$, in some arguments it is enough to consider basis elements in

$$\bar{B}_{<0} = \{b(m) \mid b \in B, m < 0\} = \bar{B} \cap \hat{\mathfrak{g}}_{<0}.$$

- Denote by \mathcal{P} the set of monomials

$$\pi = \prod_{b(j) \in \bar{B}} b(j)^{n_{b(j)}} \in S(\hat{\mathfrak{g}})$$

and by $\mathcal{P}_{<0} = \mathcal{P} \cap S(\hat{\mathfrak{g}}_{<0})$. Sometimes we say that $\pi \in \mathcal{P}_{<0}$ is a colored partition of length $\ell(\pi)$, degree $|\pi|$ and support $\text{supp } \pi$,

$$\ell(\pi) = \sum_{b(j) \in \bar{B}} n_{b(j)}, \quad |\pi| = \sum_{b(j) \in \bar{B}} n_{b(j)} \cdot j, \quad \text{supp } \pi = \{b(j) \in \bar{B}_{<0} \mid n_{b(j)} > 0\},$$

with colored parts $b(j) \in \text{supp } \pi$ of degree $j < 0$ and color $b \in B$, appearing in the partition $n_{b(j)}$ times. For $\ell \geq 0$ and $m \in \mathbb{Z}$ set

$$\mathcal{P}^\ell = \{\pi \in \mathcal{P} \mid \ell(\pi) = \ell\}, \quad \mathcal{P}^\ell(m) = \{\pi \in \mathcal{P} \mid \ell(\pi) = \ell, |\pi| = m\}.$$

Instead of the product $\mu\nu \in S(\hat{\mathfrak{g}}_{<0})$ we shall write $\mu \cup \nu \in \mathcal{P}_{<0}$ saying that $\mu \cup \nu$ is the partition having all the parts of μ and ν together. Likewise we write

$$\rho \subset \pi \quad \text{if} \quad \pi = \rho\kappa$$

for some κ , saying that $\rho \subset \pi$ is an *embedding of ρ into π* . We extend the total order $<$ on \bar{B} to \mathcal{P} so that $\mu < \mu'$ implies $\mu\kappa < \mu'\kappa$. For arguments by induction we also need that $<$ on $\mathcal{P}_{<0}$ is a well order and that $\ell(\pi) > \ell(\pi')$ or $|\pi| < |\pi'|$ implies $\pi < \pi'$.

- For the filtration $U_\ell(\hat{\mathfrak{g}})$, $\ell \geq 0$, let

$$p_\ell: U_\ell(\hat{\mathfrak{g}}) \rightarrow S^\ell(\hat{\mathfrak{g}}) \cong U_\ell(\hat{\mathfrak{g}})/U_{\ell-1}(\hat{\mathfrak{g}})$$

be the canonical projection. For each $\pi \in \mathcal{P}^\ell$ choose $u(\pi) \in U_\ell(\hat{\mathfrak{g}})$ such that $p_\ell(u(\pi)) = \pi$. Then we have a PBW basis

$$\{u(\pi)\mathbf{1} \mid \pi \in \mathcal{P}_{<0}\}$$

of $N(k\Lambda_0)$ and the corresponding spanning set on the quotient $L(k\Lambda_0)$.

- Determine the set of leading terms $\ell(\bar{R}) \subset \mathcal{P}$ of relations $r(m) \in \bar{R} \setminus \{0\}$:

$$\rho = \ell(r(m)) \quad \text{if} \quad r(m) = c_\rho u(\rho) + \sum_{\rho < \kappa} c_\kappa u(\kappa), \quad c_\rho \neq 0.$$

Then we can parameterize a basis of the loop module \bar{R} by its leading terms, i.e. there is a basis $\{r(\rho) \mid \rho \in \ell(\bar{R})\}$ of \bar{R} such that $\ell(r(\rho)) = \rho$, i.e.

$$(1.2) \quad r(\rho) = u(\rho) + \sum_{\rho < \kappa} c_\kappa u(\kappa).$$

- By using relations $r(\rho)|_{L(k\Lambda_0)} = 0$ reduce the spanning set $\{u(\pi)\mathbf{1} \mid \pi \in \mathcal{P}_{<0}\}$ of $L(k\Lambda_0)$ to (hopefully²) a basis

$$\mathcal{B} = \{u(\pi) \cdot \mathbf{1} \mid \pi \in \mathcal{P}_{<0} \setminus (\ell(\bar{R}))\}.$$

Here $\pi \in \mathcal{P}_{<0} \setminus (\ell(\bar{R}))$ denotes monomials π which are not in the ideal (\bar{R}) generated by relations \bar{R} .

If we think of monomials π as colored partitions, then the spanning set of monomial vectors $\mathcal{B} \subset L(k\Lambda_0)$ is parameterized by partitions which do not contain any subpartition $\rho \in \ell(\bar{R})$ —this is some sort of combinatorial “difference conditions” on parts of the partition π .

Note that the last two steps in constructing combinatorial basis depend on a choice of order $<$ on \mathcal{P} . In some cases the spanning set \mathcal{B} is a basis and we have the corresponding combinatorial description of the character of $L(k\Lambda_0)$, but to prove linear independence of \mathcal{B} might be a difficult task. One possible way to prove that \mathcal{B} is a basis is to have the “correct” combinatorial character formula, obtained by some combinatorial method (like in [DK] where the related conjecture in [CMPP] is proved for $k = 1$), or any other way. Here we pursue the idea that the correct character formula for $L(k\Lambda_0)$ can be obtained by constructing a combinatorial basis of $\bar{R}N(k\Lambda_0)$ because the character formula for $N(k\Lambda_0) \cong S(\hat{\mathfrak{g}}_{<0})$ is obvious, and

$$\text{ch } L(k\Lambda_0) = \text{ch } N(k\Lambda_0) - \text{ch } \bar{R}N(k\Lambda_0).$$

In order to describe a combinatorial basis of $\bar{R}N(k\Lambda_0)$ we set

$$u(\rho \subset \pi) = r(\rho)u(\kappa)$$

for an embedding $\rho \subset \pi$ such that $\pi = \rho\kappa$, $\rho \in \ell(\bar{R})$. Note that

$$r(\rho)u(\kappa) = u(\pi) + \sum_{\pi < \tau} c_\tau u(\tau),$$

so we have $\ell(u(\rho \subset \pi)) = \pi$. With this notation we can write the spanning set of $\bar{R}N(k\Lambda_0)$ as

$$(1.3) \quad u(\rho \subset \pi)\mathbf{1}, \quad \rho \in \ell(\bar{R}), \pi \in (\ell(\bar{R})) \cap \mathcal{P}_{<0}.$$

If for any two embeddings $\rho_1 \subset \pi$ and $\rho_2 \subset \pi$ we have a relation

$$(1.4) \quad u(\rho_1 \subset \pi)\mathbf{1} - u(\rho_2 \subset \pi)\mathbf{1} = \sum_{\pi < \pi', \rho' \subset \pi'} c_{\rho' \subset \pi'} u(\rho' \subset \pi')\mathbf{1},$$

then we can reduce the spanning set (1.3) by using (1.4), and for each π we may take just one embedding $\rho(\pi) \subset \pi$, $\rho(\pi) \in \ell(\bar{R})$, and the corresponding vector for the reduced spanning set of $\bar{R}N(k\Lambda_0)$

$$(1.5) \quad u(\rho(\pi) \subset \pi)\mathbf{1}, \quad \pi \in (\ell(\bar{R})) \cap \mathcal{P}_{<0}.$$

²The obtained spanning set \mathcal{B} is not necessarily a basis—see [MP2].

Such a spanning set is obviously linearly independent since the leading terms are the elements of the PBW basis of $N(k\Lambda_0)$, i.e.

$$\ell(u(\rho(\pi) \subset \pi)\mathbf{1}) = u(\pi)\mathbf{1}.$$

The reasoning above applies to all untwisted affine Lie algebras, so *relations among relations (1.4)* (would) imply linear independence of the spanning set \mathcal{B} of $L(k\Lambda_0)$.

Remark 1.1. If $\pi = \rho_1\rho_2\kappa$, $\rho_1, \rho_2 \in \ell(\bar{R})$, then we have two embeddings $\rho_1 \subset \pi$ and $\rho_2 \subset \pi$ and $\ell(\pi) \geq 2k + 2$. From (1.2) we have

$$\begin{aligned} r(\rho_1)r(\rho_2)u(\kappa) &= (u(\rho_1) + \sum_{\rho_1 \prec \kappa_1} c_{\kappa_1} u(\kappa_1))r(\rho_2)u(\kappa) \\ &= r(\rho_1)(u(\rho_2) + \sum_{\rho_2 \prec \kappa_2} c_{\kappa_2} u(\kappa_1))u(\kappa) \end{aligned}$$

and (1.4) easily follows. Hence the problem is to check (1.4) “only” for

$$k + 2 \leq \ell(\pi) \leq 2k + 1.$$

For $k = 1$ we have $k + 2 = 2k + 1 = 3$, i.e. we have to check (1.4) only for $\ell(\pi) = 3$, and this was done in [PŠ1].

On the other hand, for $k = 2$ we have to check (1.4) for $4 \leq \ell(\pi) \leq 5$. The main result of this paper—the theorem below—gives (1.4) only for $\ell(\pi) = 4$, and it is hoped that the case $\ell(\pi) = 5$ will somehow follow from that³.

Theorem 1.2. *For any two embeddings $\rho_1 \subset \pi$ and $\rho_2 \subset \pi$ in $\pi \in \mathcal{P}^4(m)$, where $\rho_1, \rho_2 \in \ell(\bar{R})$, we have a level 2 relation for $C_n^{(1)}$*

$$(1.6) \quad u(\rho_1 \subset \pi) - u(\rho_2 \subset \pi) = \sum_{\pi \prec \pi', \rho \subset \pi'} c_{\rho \subset \pi'} u(\rho \subset \pi').$$

2. THE ARRAY OF NEGATIVE ROOT VECTORS OF $C_n^{(1)}$

We fix a simple Lie algebra \mathfrak{g} of type C_n , $n \geq 2$, i.e. $\mathfrak{g} = \mathfrak{sp}_{2n}(\mathbb{C})$. For a given Cartan subalgebra \mathfrak{h} and the corresponding root system Δ we can write (as in [B])

$$\Delta = \{\pm(\varepsilon_i \pm \varepsilon_j) \mid i, j = 1, \dots, n\} \setminus \{0\}$$

with simple roots $\alpha_1 = \varepsilon_1 - \varepsilon_2$, $\alpha_2 = \varepsilon_2 - \varepsilon_3$, \dots , $\alpha_{n-1} = \varepsilon_{n-1} - \varepsilon_n$, $\alpha_n = 2\varepsilon_n$. Then $\theta = 2\varepsilon_1$. For each $\alpha \in \Delta$ we choose a root vector X_α such that $[X_\alpha, X_{-\alpha}] = \alpha^\vee$. For root vectors X_α we shall use the following notation:

$$\begin{aligned} X_{ij} &\text{ or just } ij && \text{if } \alpha = \varepsilon_i + \varepsilon_j, \quad i \leq j, \\ X_{\underline{ij}} &\text{ or just } \underline{ij} && \text{if } \alpha = -\varepsilon_i - \varepsilon_j, \quad i \geq j, \\ X_{i\bar{j}} &\text{ or just } i\bar{j} && \text{if } \alpha = \varepsilon_i - \varepsilon_j, \quad i \neq j. \end{aligned}$$

With the previous notation $x_\theta = X_{11}$. We also write for $i = 1, \dots, n$

$$X_{i\bar{i}} = \alpha_i^\vee \text{ or just } i\bar{i}.$$

These vectors X_{ab} form a basis B of \mathfrak{g} which we shall write in a triangular scheme. For example, for $n = 3$ the basis B is

³All necessary relations among relations in [MP1] for all levels follow from Lemma 9.2.




Figure 1

In order to simplify counting of embeddings of leading terms of relations, we will use the usual matrix notation for the basis B , i.e. we will use $i = 1, \dots, 2n$ for the first index for rows and $j = 1, \dots, 2n$ for the second index for columns/diagonals. For example, for $n = 3$ the basis B is




Figure 2

Moreover, we shall write $\bar{B}_{<0} = \coprod_{j>0} B \otimes t^{-j}$ in the following scheme




Figure 3

which we call *the array of negative root vectors of $C_n^{(1)}$* .

By the main theorem in [PŠ2], *the monomial*

$$(2.1) \quad \rho = \prod_{\beta \in \mathcal{B}} X_{\beta}(-j-1)^{m_{\beta,j+1}} \prod_{\alpha \in \mathcal{A}} X_{\alpha}(-j)^{m_{\alpha,j}}$$

is the leading term of the relation $r(\rho) \in \bar{R}$ for level k standard modules of affine Lie algebra of the type $C_n^{(1)}$, where

$$(2.2) \quad \sum_{\beta \in \mathcal{B}} m_{\beta,j+1} = b \quad , \quad \sum_{\alpha \in \mathcal{A}} m_{\alpha,j} = a \quad , \quad a + b = k + 1$$

and \mathcal{B} and \mathcal{A} are the admissible pair of cascades in degree $-j-1$ and $-j$.

For a general rank we may visualize the admissible pair of cascades \mathcal{B} and \mathcal{A} as on Figure 4 (see Figure 1 in [PŠ2]).




Figure 4

With our new way of writing $\bar{B}_{<0} = \coprod_{m<0} B \otimes t^m$ on Figure 3, we may reinterpret the pair of admissible cascades $(\mathcal{B}, \mathcal{A})$ on Figure 4 as the downward zig-zag line on Figure 5.




Figure 5

Note that Figure 5 consists of the triangle Δ on Figure 4 together with its mirror image Δ' with respect to the hypotenuse, but translated along a pair of parallel sides of Δ and Δ' so that the other two sides coincide. Of course, the obtained parallelogram is rotated for $\pi/4$.

On $\bar{B}_{<0}$, written as on Figure 3, we introduce a partial order \trianglelefteq by

$$(2.3) \quad X_{i,j} \trianglelefteq X_{p,r} \quad \text{if} \quad i \in \{1, \dots, p\} \quad \text{and} \quad j \in \{r, p+r-i\}.$$

Anotherwords, $b = X_{i,j}$ lies in the cone bellow the vertex $a = X_{p,r}$, as depicted on Figure 6 below:




Figure 6

With this conventions we can restate the description of leading terms ρ of relations $r(\rho) \in \bar{R}$ in [PŠ2] (cf. [P]) as: *for any zig-zag downward line*

$$a_1 \triangleright a_2 \triangleright \cdots \triangleright a_r$$

of r points in $\bar{B}_{<0}$, $1 \leq r \leq k+1$, the monomial

$$(2.4) \quad \rho = a_1^{m_1} a_2^{m_2} \cdots a_r^{m_r}, \quad m_1 + m_2 + \cdots + m_r = k+1,$$

is the leading term of the relation $r(\rho) \in \bar{R}$ for level k standard modules of affine Lie algebra of the type $C_n^{(1)}$. Moreover, these are all leading terms of \bar{R} in $\mathcal{P}_{<0}$.

In the next section we shall count the number of embeddings $\rho \subset \pi$ for $\text{supp } \pi$ in three successive triangles, i.e. in the trapezoid⁴ T on Figure 6.

3. COLORED PARTITIONS ALLOWING AT LEAST TWO EMBEDDINGS

For a colored partition

$$(3.1) \quad \pi = \prod_{a \in \bar{B}_{<0}} a^{\pi(a)}$$

we have $|\text{supp } \pi| \leq \ell(\pi)$. The basis $\bar{B}_{<0}$ is written as the array of $2n+1$ rows and $a_1 \triangleright \cdots \triangleright a_r$ implies $r \leq 2n+1$.

Lemma 3.1. *Let $\ell(\pi) = k+2$ and assume that π allows two embeddings of leading terms of relations for level k standard modules. Then $\text{supp } \pi$ is one of the following types (for $r, s \in \mathbb{N}$ and δ in the set of two symbols $|$ and $||$):*

- (A_r) $\text{supp } \pi = \{a_1, \dots, a_r\}$, $r \geq 2$, $a_1 \triangleright \cdots \triangleright a_r$.
- ($B_{r\delta}$) $\text{supp } \pi = \{a_1, \dots, a_r, b, c\}$, $r \geq 1$, $a_1 \triangleright \cdots \triangleright a_r$, $a_r \triangleright b$, $a_r \triangleright c$ and b and c are not comparable. We set δ to be $|$ if b and c are in the same row, and $||$ otherwise.
- ($C_{\delta r}$) $\text{supp } \pi = \{b, c, a_1, \dots, a_r\}$, $r \geq 1$, $a_1 \triangleright \cdots \triangleright a_r$, $b \triangleright a_1$, $c \triangleright a_1$ and b and c are not comparable. We set δ to be $|$ if b and c are in the same row, and $||$ otherwise.
- ($D_{r\delta s}$) $\text{supp } \pi = \{a_1, \dots, a_r, b, c, d_1, \dots, d_s\}$, $r, s \geq 1$, $a_1 \triangleright \cdots \triangleright a_r$, $a_r \triangleright b \triangleright d_1$, $a_r \triangleright c \triangleright d_1$, $d_1 \triangleright \cdots \triangleright d_s$, and b and c are not comparable. We set δ to be $|$ if b and c are in the same row, and $||$ otherwise.

Proof. Let $\pi = a_1^{f_1} \cdots a_r^{f_r}$, $f_1 \cdots f_r > 0$, $f_1 + \cdots + f_r = k+2$, $a_1 \triangleright \cdots \triangleright a_r$. If in π we erase one factor a_j , $j \in \{1, \dots, r\}$, we get r leading terms of relations

$$\rho_1 = a_1^{f_1-1} \cdots a_r^{f_r}, \quad \dots \quad \rho_r = a_1^{f_1} \cdots a_r^{f_r-1}$$

and we have r embeddings $\rho_1 \subset \pi$, \dots , $\rho_r \subset \pi$. In particular, for $r \geq 2$ we have at least two embeddings in π .

Now let $\ell(\pi) = k+2$ and let $\rho_1 \subset \pi$ and $\rho_2 \subset \pi$ be two different embeddings of leading terms of relations, $\ell(\rho_1) = \ell(\rho_2) = k+1$. Then there are parts c_1 and c_2 of π such that

$$\pi = c_1 \rho_1 = c_2 \rho_2 = c_1 c_2 (\rho_1 \cap \rho_2),$$

⁴Note that the position of the trapezoid in Figure 6 is in accord with Figure 3 only when the middle triangle is $B \otimes t^j$ for j even, and for j odd the figure should be flipped. However, in our arguments this will make no difference because the flipped zig-zag downward line is again a zig-zag downward line.

and $\rho_1 \neq \rho_2$ implies $c_1 \neq c_2$. Hence

$$\rho_1 = c_2(\rho_1 \cap \rho_2), \quad \rho_2 = c_1(\rho_1 \cap \rho_2).$$

Since ρ_1 and ρ_2 are the leading terms of relations, (2.4) implies that the partition $\rho_1 \cup \rho_2$ can be written as

$$a_1 \triangleright \cdots \triangleright a_t$$

and ρ_1 and ρ_2 as

$$(3.2) \quad \begin{aligned} \rho_1: & \quad a_1 \triangleright \cdots \triangleright a_p \triangleright c_2 \triangleright a_{p+1} \triangleright \cdots \triangleright a_t, \\ \rho_2: & \quad a_1 \triangleright \cdots \triangleright a_q \triangleright c_1 \triangleright a_{q+1} \triangleright \cdots \triangleright a_t. \end{aligned}$$

If $q < p$, then

$$a_q \triangleright c_1 \triangleright a_{q+1} \triangleright \cdots \triangleright c_2$$

and $\text{supp } \pi$ is of the type A_r for some $r \geq 2$.

If $q = p$ and c_1 and c_2 are comparable, say $c_2 \triangleright c_1$, then $a_p \triangleright c_2 \triangleright a_{p+1}$ and $a_p \triangleright c_1 \triangleright a_{p+1}$ implies

$$a_p \triangleright c_2 \triangleright c_1 \triangleright a_{p+1},$$

and again $\text{supp } \pi$ is of the type A_r for some $r \geq 2$. Hence the lemma follows and $\text{supp } \pi$ is of the type $B_{r\delta}$, $C_{\delta r}$ or $D_{r\delta s}$, depending on the position of incomparable parts c_1 and c_2 in ρ_1 and ρ_2 written as (3.2) for $p = q$.

The distinction, and the notation of relative position of parts c_1 and c_2 , i.e. $\delta = \parallel$ if c_1 and c_2 are in different rows, and $\delta = |$ otherwise, will be convenient later on. \square

Lemma 3.2. (i) *Let $2 \leq r \leq \min\{2n+1, k+2\}$. Then for any choice of $a_1 \triangleright \cdots \triangleright a_r$ there is*

$$\binom{k+1}{r-1}$$

partitions π of the form

$$(3.3) \quad \pi = a_1^{f_1} \cdots a_r^{f_r}, \quad f_1 \cdots f_r > 0, \quad f_1 + \cdots + f_r = k+2$$

and for each such π there are r embeddings $\rho_1 \subset \pi, \dots, \rho_r \subset \pi$ for

$$\rho_1 = a_1^{f_1-1} \cdots a_r^{f_r}, \quad \dots \quad \rho_r = a_1^{f_1} \cdots a_r^{f_r-1}.$$

(ii) *Let $r, s \geq 0$, $3 \leq r+2+s \leq k+2$, $3 \leq r+a+s \leq 2n+1$ with $a = 1$ if δ is $|$ and $a = 2$ if δ is \parallel . For any choice of $\{a_1, \dots, a_r, c, d, b_1, \dots, b_s\}$ of the type $B_{r\delta}$ (for $s = 0$), $C_{\delta s}$ (for $r = 0$) or $(D_{r\delta s})$ there is*

$$\binom{k-1}{s+r-1}$$

partitions π of the form

$$(3.4) \quad \pi = a_1^{f_1} \cdots a_r^{f_r} c d b_1^{g_1} \cdots b_s^{g_s}, \quad f_1 \cdots f_r g_1 \cdots g_s > 0, \quad f_1 + \cdots + f_r + g_1 + \cdots + g_s = k,$$

and for each such π there are 2 embeddings $\rho_1 \subset \pi$ and $\rho_2 \subset \pi$ for

$$\rho_1 = a_1^{f_1} \cdots a_r^{f_r} c b_1^{g_1} \cdots b_s^{g_s}, \quad \rho_2 = a_1^{f_1} \cdots a_r^{f_r} d b_1^{g_1} \cdots b_s^{g_s}.$$

Proof. Since there is $\binom{k+1}{r-1}$ ways of writing $f_1 + \cdots + f_r = k+2$, $f_1 \cdots f_r > 0$, the first statement is clear, and the second statement is obvious since $\ell(\rho) = k+1$. This proves (i), and the proof of (ii) is similar. \square

As in [PŠ1] for a colored partition π set

$$N(\pi) = \max\{\#\mathcal{E}(\pi) - 1, 0\} \quad \mathcal{E}(\pi) = \{\rho \in \ell(\bar{R}) \mid \rho \subset \pi\}.$$

Let T be a trapezoid consisting of three consecutive triangles in the array $\bar{B}_{<0}$. Set

$$\Sigma_T(A_r) = \sum_{S \subset T, S \text{ is of the type } A_r} 1, \quad N_T(A_r) = \sum_{\pi, \text{supp } \pi \subset T, \text{supp } \pi \text{ is of the type } A_r} N(\pi),$$

and likewise for types $B_{r\delta}$, $C_{\delta r}$ and $D_{r\delta s}$.

Note that two different orientations for trapezoids are possible: with a long base down, say T , and with the long base up, say T' . However we have an ‘‘up-down symmetry’’ so that $\Sigma_T(A_r) = \Sigma_{T'}(A_r)$ and

$$\Sigma_T(B_{r\delta}) = \Sigma_{T'}(C_{\delta r}), \quad \Sigma_T(C_{\delta r}) = \Sigma_{T'}(B_{r\delta}), \quad \Sigma_T(D_{r\delta s}) = \Sigma_{T'}(D_{s\delta r}),$$

so we shall consider only a trapezoid oriented as T . Clearly Lemma 3.2 implies:

Lemma 3.3. *For r and s as in Lemma 3.2 we have:*

- (1) $N_T(A_r) = (r-1) \binom{k+1}{r-1} \Sigma_T(A_r)$,
- (2) $N_T(B_{r\delta}) = \binom{k-1}{r-1} \Sigma_T(B_{r\delta})$,
- (3) $N_T(C_{\delta r}) = \binom{k-1}{r-1} \Sigma_T(C_{\delta r})$,
- (4) $N_T(D_{r\delta s}) = \binom{k-1}{s+r-1} \Sigma_T(D_{r\delta s})$.

Lemma 3.4. *For $2 \leq r \leq 2n+1$ we have:*


$$\begin{aligned} \Sigma_T(A_r) &= \sum_{i_1=r}^{2n+1} \sum_{j_1=1}^{4n+1-i_1} \sum_{i_2=r-1}^{i_1-1} \sum_{j_2=j_1}^{j_1+i_1-i_2} \cdots \sum_{i_p=r-p+1}^{i_{p-1}-1} \sum_{j_p=j_{p-1}}^{j_{p-1}+i_{p-1}-i_p} \cdots \sum_{i_r=1}^{i_{r-1}-1} \sum_{j_r=j_{r-1}}^{j_{r-1}+i_{r-1}-i_r} 1 \\ &= \sum_{i_1=r}^{2n+1} \sum_{i_2=r-1}^{i_1-1} \cdots \sum_{i_p=r-p+1}^{i_{p-1}-1} \cdots \sum_{i_r=1}^{i_{r-1}-1} \pi(i_1, \dots, i_r), \\ \pi(i_1, \dots, i_r) &= (4n+1-i_1)(i_1-i_2+1) \cdots (i_{p-1}-i_p+1) \cdots (i_{r-1}-i_r+1). \end{aligned}$$

Proof. We want to count all

$$S = \{a_1, \dots, a_r\} \subset T, \quad a_1 \triangleright \cdots \triangleright a_r.$$

The points $a_p = (i_p, j_p) \in T$ are in distinct rows $i_1 > \cdots > i_r$. For the first point a_1 in the row i_1 we may choose $j_1 \in \{1, \dots, 4n+1-i_1\}$, altogether $4n+1-i_1$ points. For the second point $a_2 \triangleright a_1$ in the row i_2 we may choose $j_2 \in \{j_1, \dots, j_1+i_1-i_2\}$, altogether i_1-i_2+1 points. In this way we continue till the point $a_r \in T$. \square

Remark 3.5. The types $B_{r\delta}$, $C_{\delta s}$ and $D_{r\delta s}$ differ from the type A_r by having two incomparable points b and c . In the case when b and c are on the same line ℓ (i.e. δ is \mid) in our counting procedure for Σ_T we first determine all possible distances d between b and c , and then all possible lines ℓ through b and c —we sketch the three corresponding configurations as:



After that we argue in a manner very similar to the proof of Lemma 3.4.

Lemma 3.6. *For $1 \leq r \leq 2n$ we have:*

$$\begin{aligned} \Sigma_T(B_r) &= \sum_{d=1}^{2n+1-r} \sum_{\ell=1}^{2n+2-d-r} \sum_{i_1=\ell+d+r-1}^{2n+1} \sum_{i_2=\ell+d+r-2}^{i_1-1} \dots \sum_{i_r=\ell+d}^{i_{r-1}-1} \pi(d, \ell; i_1, \dots, i_r), \\ \pi(d, \ell; i_1, \dots, i_r) &= (4n+1-i_1)(i_1-i_2+1) \dots (i_{r-1}-i_r+1)(i_r-\ell-d+1). \end{aligned}$$

Proof. We should determine the number of all possible $\{a_1, \dots, a_r, b, c\} \subset T$, $r \geq 1$, such that $a_1 \triangleright \dots \triangleright a_r$, $a_r \triangleright b$, $a_r \triangleright c$ and $b \neq c$ are on the same line ℓ . The distance $d = j_2 - j_1$ between the points $b = (\ell, j_1)$ and $c = (\ell, j_2)$, $j_2 > j_1$, may be 1 because for $\ell = 1$ we still can place $r \leq 2n$ points $a_1 \triangleright \dots \triangleright a_r$ above both b and c . In general for $d \geq 1$ and b and c on the line $1 \leq \ell \leq 2n+1$ the point a_r is on the line $i_r \geq \ell+d$ and we need at least $r-1$ lines above to place the remaining $r-1$ points $a_1 \triangleright \dots \triangleright a_{r-1}$. Hence

$$(3.5) \quad \ell + d + r - 1 \leq 2n + 1,$$

and for $\ell = 1$ this implies

$$1 \leq d \leq 2n + 1 - r.$$

Once we have fixed d , by (3.5) we should take

$$1 \leq \ell \leq 2n + 2 - d - r.$$

Now we proceed as in the proof of Lemma 3.4, the only difference being that for the vertex x in the equilateral triangle $\Delta(b, c, x)$ there is

$$i_r - \ell - d + 1$$

places on the line $\ell + d$ which are in the cone below a_r . \square

Lemma 3.7. *For $1 \leq r \leq 2n$ we have:*

$$\begin{aligned} \Sigma_T(C_r) &= 1 \sum_{d=1}^{2n+1-r} \sum_{\ell=d+r}^{2n+1} \sum_{i_1=r}^{\ell-d} \sum_{i_2=r-1}^{i_1-1} \dots \sum_{i_r=1}^{i_{r-1}-1} \pi(d, \ell; i_1, \dots, i_r), \\ \pi(d, \ell; i_1, \dots, i_r) &= (4n+1-\ell-d)(\ell-d-i_1+1)(i_1-i_2+1) \dots (i_{r-1}-i_r+1). \end{aligned}$$

Proof. We should determine the number of all possible $\{b, c, a_1, \dots, a_r\} \subset T$, $r \geq 1$, such that $a_1 \triangleright \dots \triangleright a_r$, $b \triangleright a_1$, $c \triangleright a_1$ and $b \neq c$ are on the same line ℓ . The distance $d = j_2 - j_1$ between the points $b = (\ell, j_1)$ and $c = (\ell, j_2)$, $j_2 > j_1$, may be 1 because on the “smallest” top row of trapezoid T we have at least 2 points. For r points a_1, \dots, a_r we need r distinct rows, and ℓ -th row is above them at the distance d . Hence

$$d + r \leq 2n + 1.$$

For fixed d we clearly have

$$d + r \leq \ell \leq 2n + 1.$$

There are $4n+1-\ell-d$ ways to put two points b and c on the line ℓ at distance d . For fixed d , and b and c on the line $1 \leq \ell \leq 2n+1$, the point a_1 is on the line $i_1 \leq \ell-d$. Now we proceed as in the proof of Lemma 3.4, the only difference being that for the vertex y in the equilateral triangle $\Delta(b, y, c)$ there is

$$\ell - d - i_1 + 1$$

places on the line i_1 which are in the cone below y . \square

Lemma 3.8. For $2 \leq r + s \leq 2n$ we have:

$$\Sigma_T(D_{r|s}) = \sum_{d=1}^{\lfloor \frac{2n+2-r-s}{2} \rfloor} \sum_{\ell=s+d}^{2n+2-r-d} \sum_{i_1=\ell+d-r+1}^{2n+1} \sum_{i_2=\ell+d-r+2}^{i_1-1} \cdots \sum_{i_r=\ell+d}^{i_{r-1}-1} \sum_{l_1=s}^{\ell-d} \sum_{l_2=s-1}^{l_1-1} \cdots \sum_{l_s=1}^{l_{s-1}-1} \pi(d, \ell; i_1, \dots, i_r; l_1, \dots, l_s),$$

$$\pi(d, \ell; i_1, \dots, i_r; l_1, \dots, l_s) = (4n+1-i_1)(i_1-i_2+1) \cdots (i_{r-1}-i_r+1)(i_r-\ell-d+1) \\ (\ell-d-l_1+1)(l_1-l_2+1) \cdots (l_{s-1}-l_s+1).$$

Proof. We should determine the number of all possible $\{a_1, \dots, a_r, b, c, d_1, \dots, d_s\}$, $r, s \geq 1$, $a_1 \triangleright \cdots \triangleright a_r$, $a_r \triangleright b \triangleright d_1$, $a_r \triangleright c \triangleright d_1$, $d_1 \triangleright \cdots \triangleright d_s$, and $b \neq c$ are on the same line ℓ . We need to place r distinct rows above ℓ at the distance d , and s distinct rows below ℓ at the distance d . Hence $r-1+s-1+2d \leq 2n+1$, and for positive integer d we have


$$1 \leq d \leq \lfloor \frac{2n+3-r-s}{2} \rfloor.$$

For fixed d we have $s+d \leq \ell \leq 2n+1-r-d$, and we proceed as in the proof of lemmas above: the factors in $\pi(d, \ell; i_1, \dots, i_r; l_1, \dots, l_s)$ count the number of points we can place in one row when a point in the row above is chosen. In particular, the factor $(i_r - \ell - d + 1)$ counts the number of places the vertex x of the rhombus $R(b, y, c, x)$ can take on the line $\ell + d$ in the cone below a_r , and $(\ell - d - l_1 + 1)$ is the number of places on the line l_1 which are in the cone below the vertex y . \square

Remark 3.9. The types $B_{r||}$, $C_{||s}$ and $D_{r||s}$ have two incomparable points b and c in different lines in two possible configurations



Due to the left-right symmetry it is enough to consider Σ_T for only the first one counted twice. As before, we first determine all possible distances $d = j_2 - j_1 \geq 1$ between $b = (\ell - h, j_1)$ and $c = (j_2, \ell)$, and then all possible h and lines ℓ through c —we sketch the three corresponding configurations as:



Lemma 3.10. For $1 \leq r \leq 2n-1$ we have:

$$\Sigma_T(B_{r||}) = 2 \sum_{d=1}^{2n-r} \sum_{h=1}^{2n+1-d-r} \sum_{\ell=h+1}^{2n+2-d-r} \sum_{i_1=\ell+d+r-1}^{2n+1} \sum_{i_2=\ell+d+r-2}^{i_1-1} \cdots \sum_{i_r=\ell+d}^{i_{r-1}-1} \pi(d, \ell; i_1, \dots, i_r),$$

$$\pi(d, \ell; i_1, \dots, i_r) = (4n+1-i_1)(i_1-i_2+1) \cdots (i_{r-1}-i_r+1)(i_r-\ell-d+1).$$

Proof. We argue as in the proof of Lemma 3.6: for $d \geq 1$, $h \geq 1$ and c on the line $1 \leq \ell \leq 2n+1$ the point a_r is on the line $i_r \geq \ell + d$ and we need at least $r-1$ lines above to place the remaining $r-1$ points $a_1 \triangleright \cdots \triangleright a_{r-1}$. Hence

$$(3.6) \quad \ell + d + r - 1 \leq 2n + 1,$$

and for $\ell = h + 1$ this implies $d \leq 2n + 1 - r - h$. For $h = 1$ we determine the bounds for d , for chosen d we determine the bounds for h , and by using (3.6) we determine the bounds for ℓ :

$$1 \leq d \leq 2n - r, \quad 1 \leq h \leq 2n + 1 - d - r, \quad h + 1 \leq \ell \leq 2n + 2 - d - r.$$

Now we proceed as in the proof of Lemma 3.6. \square

Lemma 3.11. *For $1 \leq r \leq 2n - 1$ we have:*

$$\begin{aligned} \Sigma_T(C_{\parallel r}) &= 2 \sum_{d=1}^{2n-r} \sum_{h=1}^{2n+1-d-r} \sum_{\ell=d+h+r}^{2n+1} \sum_{i_1=r}^{\ell-d-h} \sum_{i_2=r-1}^{i_1-1} \cdots \sum_{i_r=1}^{i_{r-1}-1} \pi(d, h, \ell; i_1, \dots, i_r), \\ \pi(d, h, \ell; i_1, \dots, i_r) &= (4n + 1 - \ell - d)(\ell - d - h - i_1 + 1)(i_1 - i_2 + 1) \cdots (i_{r-1} - i_r + 1). \end{aligned}$$

Proof. For r points a_1, \dots, a_r we need r distinct rows, and ℓ -th row is above them at the distance $d + h$. Hence

$$d + h + r \leq 2n + 1.$$

For $h = 1$ we determine the bounds for d , for chosen d we determine the bounds for h , and for chosen h we determine the bounds for ℓ :

$$1 \leq d \leq 2n - r, \quad 1 \leq h \leq 2n + 1 - r - d, \quad r + d + h \leq \ell \leq 2n + 1.$$

There are $4n + 1 - \ell - d$ ways to put c on the line ℓ at distance d from b . For fixed d, h and c on the line $1 \leq \ell \leq 2n + 1$, the point a_1 is on the line $i_1 \leq \ell - d - h$. Now we proceed as in the proof of Lemma 3.4, the only difference being that for the vertex y in the triangle $\Delta(b, y, c)$ there is

$$\ell - d - h - i_1 + 1$$

places on the line i_1 which are in the cone below y . \square

Lemma 3.12. *For $2 \leq r + s \leq 2n - 1$ we have:*

$$\begin{aligned} \Sigma_T(D_{r\parallel s}) &= 2 \sum_{d=1}^{\lfloor \frac{2n+1-r-s}{2} \rfloor} \sum_{h=1}^{2n+2-2d-r-s} \sum_{\ell=s+d+h}^{2n+2-r-d} \sum_{i_1=\ell-r+d+1}^{2n+1} \sum_{i_2=\ell-r+d+2}^{i_1-1} \cdots \sum_{i_r=\ell+d}^{i_{r-1}-1} \\ &\quad \sum_{l_1=s}^{\ell-d-h} \sum_{l_2=s-1}^{l_1-1} \cdots \sum_{l_s=1}^{l_{s-1}-1} \pi(d, \ell; i_1, \dots, i_r; l_1, \dots, l_s), \\ \pi(d, \ell; i_1, \dots, i_r; l_1, \dots, l_s) &= (4n + 1 - i_1)(i_1 - i_2 + 1) \cdots (i_{r-1} - i_r + 1)(i_r - \ell - d + 1) \\ &\quad (\ell - d - h - l_1 + 1)(l_1 - l_2 + 1) \cdots (l_{s-1} - l_s + 1). \end{aligned}$$

Proof. We need to place r distinct rows above ℓ at the distance d , and s distinct rows below $\ell - h$ at the distance d . Hence $r - 1 + s - 1 + 2d + h \leq 2n + 1$. For $h = 1$ we determine the bounds for d , for chosen d we determine the bounds for h , and for chosen h we determine the bounds for ℓ :

$$1 \leq d \leq \lfloor \frac{2n+2-r-s}{2} \rfloor, \quad 1 \leq h \leq 2n + 3 - 2d - r - s, \quad s + d + h \leq \ell \leq 2n + 1 - r - d.$$

Now we proceed as in the proof of lemmas above: the factors in $\pi(d, \ell; i_1, \dots, i_r; l_1, \dots, l_s)$ count the number of points we can place in one row when a point in the row above is chosen. In particular, the factor $(i_r - \ell - d + 1)$ counts the number of places the vertex x of the parallelogram $P(b, y, c, x)$ can take on the line $\ell + d$ in the cone below a_r , and $(\ell - d - h - l_1 + 1)$ is the number of places on the line l_1 which are in the cone below the vertex y . \square

4. LEVEL 2 RELATIONS FOR $C_n^{(1)}$ FOR TWO EMBEDDINGS IN π , $\ell(\pi) = 4$

In this section we prove our main result:

Theorem 4.1. *For any two embeddings $\rho_1 \subset \pi$ and $\rho_2 \subset \pi$ in $\pi \in \mathcal{P}^4(m)$, where $\rho_1, \rho_2 \in \ell(\bar{R})$, we have a level 2 relation for $C_n^{(1)}$*

$$(4.1) \quad u(\rho_1 \subset \pi) - u(\rho_2 \subset \pi) = \sum_{\pi < \pi', \rho \subset \pi'} c_{\rho \subset \pi'} u(\rho \subset \pi').$$

For a colored partition (3.1) we define *the shape of π* as the ordinary partition of $|\pi| \in -\mathbb{N}$

$$\text{sh } \pi = \prod_{a \in \bar{B}_{<0}} |a|^{\pi(a)}$$

with parts $|a| \in -\mathbb{N}$ for $a \in \text{supp } \pi = \{a \in \bar{B}_{<0} \mid \pi(a) > 0\}$. Note that $|\text{sh } \pi| = |\pi| = \sum_{a \in \bar{B}_{<0}} |a| \cdot \pi(a)$. We shall depict the ordinary partition $p = (-3)^2(-2)$ with its Young tableaux

$$\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \\ \hline \end{array} .$$

To prove the theorem, for a fixed ordinary partition p of length 4 we need to count “the number of two-embeddings for $\text{sh } \pi = p$ ”

$$\sum_{\text{sh } \pi = p, \pi \in \mathcal{P}^4(m)} N(\pi).$$

It turns out it is enough, but much easier, to count for a trapezoid T the number

$$N_T = \sum_{\pi, \ell(\pi)=4, \text{supp } \pi \subset T} N(\pi).$$

By Lemma 3.3 we have,

$$\begin{aligned} N_T(A_4) &= 3 \Sigma_T(A_4), & N_T(A_3) &= 6 \Sigma_T(A_3), & N_T(A_2) &= 3 \Sigma_T(A_2), \\ N_T(B_{2\parallel}) &= \Sigma_T(B_{2\parallel}), & N_T(B_{2\mid}) &= \Sigma_T(B_{2\mid}), \\ N_T(B_{1\parallel}) &= \Sigma_T(B_{1\parallel}), & N_T(B_{1\mid}) &= \Sigma_T(B_{1\mid}), \\ N_T(C_{\parallel 2}) &= \Sigma_T(C_{\parallel 2}), & N_T(C_{\mid 2}) &= \Sigma_T(C_{\mid 2}), \\ N_T(C_{\parallel 1}) &= \Sigma_T(C_{\parallel 1}), & N_T(C_{\mid 12}) &= \Sigma_T(C_{\mid 1}), \\ N_T(D_{1\parallel 1}) &= \Sigma_T(D_{1\parallel 1}), & N_T(D_{1\mid 1}) &= \Sigma_T(D_{1\mid 1}), \end{aligned}$$

so by using the software package *Mathematica* for summations in Lemmas 3.4–3.12 we get:

Lemma 4.2. *Let T be a trapezoid consisting of three consecutive triangles in the array $\bar{B}_{<0}$ and $k = 2$. Then*

$$\begin{aligned} N_T(A_4) &= \frac{48n^8 + 672n^7 + 2296n^6 - 4613n^4 + 798n^3 + 1009n^2 - 210n}{280}, \\ N_T(A_3) &= \frac{56n^6 + 420n^5 + 590n^4 - 225n^3 - 151n^2 + 30n}{15}, \\ N_T(A_2) &= \frac{20n^4 + 60n^3 + 19n^2 - 3n}{2}, \end{aligned}$$

$$\begin{aligned}
N_T(B_{2|}) &= \frac{128n^7 + 952n^6 + 1652n^5 + 490n^4 - 553n^3 - 182n^2 + 33n}{630}, \\
N_T(B_{1|}) &= \frac{48n^5 + 140n^4 + 120n^3 + 25n^2 - 3n}{30}, \\
N_T(B_{2||}) &= \frac{144n^8 + 992n^7 + 840n^6 - 1456n^5 - 1239n^4 + 518n^3 + 255n^2 - 54n}{1260}, \\
N_T(B_{1||}) &= \frac{56n^6 + 132n^5 + 50n^4 - 45n^3 - 16n^2 + 3n}{45}, \\
N_T(C_{2|}) &= \frac{16n^7 + 112n^6 + 160n^5 - 20n^4 - 101n^3 - 2n^2 + 15n}{90}, \\
N_T(C_{1|}) &= \frac{8n^5 + 20n^4 + 10n^3 - 5n^2 - 3n}{6}, \\
N_T(C_{2||}) &= \frac{32n^8 + 208n^7 + 112n^6 - 392n^5 - 182n^4 + 217n^3 + 38n^2 - 33n}{315}, \\
N_T(C_{1||}) &= \frac{16n^6 + 32n^5 - 20n^3 - n^2 + 3n}{15}, \\
N_T(D_{1|1}) &= \frac{256n^7 + 1512n^6 + 2884n^5 + 1575n^4 - 686n^3 - 567n^2 + 66n}{2520}, \\
N_T(D_{1||1}) &= \frac{72n^8 + 368n^7 + 448n^6 - 322n^5 - 707n^4 - 28n^3 + 187n^2 - 18n}{1260}.
\end{aligned}$$

Lemma 4.3. *Let T be a trapezoid consisting of three consecutive triangles in the array $\bar{B}_{<0}$ and $k = 2$. Then*

$$N_T = \sum_{r=2}^4 N_T(A_r) + \sum_{r=1}^2 \sum_{\delta=|,||} (N_T(B_{r\delta}) + N_T(C_{\delta r}) + N_T(D_{1\delta 1})) = \frac{7(10n-1)}{4} \binom{2n+6}{7}.$$

Proof. Lemma 4.2 gives

$$\begin{aligned}
N_T &= \frac{1120n^8 + 11648n^7 + 47824n^6 + 9800n^5 + 103390n^4 + 50372n^3 + 6426n^2 - 1260n}{2520} \\
&= \frac{(10n-1)n(n+1)(n+2)(n+3)(2n+1)(2n+3)(2n+5)}{180} \\
&= \frac{(10n-1)2n(2n+2)(2n+4)(2n+6)(2n+1)(2n+3)(2n+5)}{180 \cdot 16} \\
&= \frac{7(10n-1)}{4} \binom{2n+6}{7}.
\end{aligned}$$

□

Proof of Theorem 4.1. By Theorem 9.2 and Proposition 10.1 in [PŠ1] for $\pi \in \mathcal{P}^{k+2}(m)$ we have a space of relations for annihilating fields

$$(4.2) \quad Q_{k+2}(m) = U(\mathfrak{g})q_{(k+1)\theta}(m) \oplus U(\mathfrak{g})q_{(k+2)\theta}(m) \oplus U(\mathfrak{g})q_{(k+2)\theta-\alpha^*}(m).$$

Especially, for level $k = 2$ and each degree $m \leq -k - 2$ from the above equation, it follows

$$(4.3) \quad Q_4(m) = U(\mathfrak{g})q_{3\theta}(m) \oplus U(\mathfrak{g})q_{4\theta}(m) \oplus U(\mathfrak{g})q_{4\theta-\alpha^*}(m)$$

where $\alpha^* = \alpha_1 = \varepsilon_1 - \varepsilon_2$. Moreover, again thanks to the mentioned Theorem 9.2, the proof of Theorem 4.1 is reduced to the verification of equality

$$(4.4) \quad \sum_{\pi \in \mathcal{P}^4(m)} N(\pi) = \dim Q_4(m) .$$

The Weyl dimension formula in the case of symplectic Lie algebra $\mathfrak{g} = \mathfrak{sp}_{2n}$ (with the corresponding $\rho = n\varepsilon_1 + (n-1)\varepsilon_2 + \dots + 2\varepsilon_{n-1} + \varepsilon_n$) gives

$$(4.5) \quad \dim L(s\theta) = \binom{2n+2s-1}{2s},$$

$$(4.6) \quad \dim L(4\theta - \alpha^*) = \frac{(2n+7)(n-1)}{4} \binom{2n+5}{6}.$$

Hence from (4.5) and (4.6) we have

$$(4.7) \quad \dim Q_4(m) = \dim L(3\theta) + \dim L(4\theta) + \dim L(4\theta - \alpha^*) = 2n \binom{2n+6}{7}.$$

Since all technical Lemmas 3.1 - 3.12 were proved for three successive triangles (i.e. for the trapezoid) the equality (4.4) can be replaced with the equivalent one

$$(4.8) \quad \sum_{m=4}^{12} \sum_{\pi \in \mathcal{P}^4(m)} N(\pi) = 9 \times 2n \binom{2n+6}{7} - 2 \times \dim L(4\theta) = \frac{7(10n-1)}{4} \binom{2n+6}{7}.$$

Indeed, using Young tableaux we can list the following cases




Figure 7

It is important to note that Young tableaux from $m = -4$ to $m = -12$ covers all cases which can occur in the trapezoid scheme of three successive triangles. Now, from Lemma 4.3 and (4.8) it is obvious that equality (4.4) holds. \square

Remark 4.4. From Lemmas 3.4–3.12 and Faulhaber’s formula (cf. [O]) we see that N_T is a polynomial in n of degree at most 8. So it is enough to check Lemma 4.3 for 9 values of n , say for $n = 1, \dots, 9$. The value of N_T for $n = 1$ equals 126 and this corresponds to the “second simplest case” in [MP1]. The value of N_T for $n = 9$ equals 53 905 698—this number illustrates the degree of complexity when passing from $C_1^{(1)} = A_1^{(1)}$ to $C_n^{(1)}$ for higher ranks n (of course, this discussion goes only for level $k = 2$).

ACKNOWLEDGEMENT

This work is partially supported by the QuantiXLie Centre of Excellence, a project cofinanced by the Croatian Government and European Union through the European Regional Development Fund—the Competitiveness and Cohesion Operational Programme (Grant KK.01.1.1.01.0004).

REFERENCES

- [B] N. Bourbaki, *Algèbre commutative*, Hermann, Paris, 1961.
- [CMPP] S. Capparelli, A. Meurman, A. Primc and M. Primc, *New partition identities from $C_\ell^{(1)}$ -modules*, Glas. Mat. Ser. III **57(77)** (2022), 161–184.
- [DK] J. Dousse and I. Konan, *Characters of level 1 standard modules of $C_n^{(1)}$ as generating functions for generalised partitions*, arXiv:2212.12728.
- [K] V. G. Kac, *Infinite-dimensional Lie algebras* 3rd ed, Cambridge Univ. Press, Cambridge, 1990.
- [LL] J. Lepowsky and H.-S. Li, *Introduction to Vertex Operator Algebras and Their Representations*, Progress in Math. Vol. 227, Birkhäuser, Boston, 2003.
- [MP1] A. Meurman and M. Primc, *Annihilating fields of standard modules of $\mathfrak{sl}(2, \mathbb{C})^-$ and combinatorial identities*, Memoirs of the Amer. Math. Soc. **137**, No. 652 (1999).
- [MP2] A. Meurman and M. Primc, *A basis of the basic $\mathfrak{sl}(3, \mathbb{C})^-$ -module*, Commun. Contemp. Math. **3** (2001), 593–614.
- [O] G. Orosi, *A simple derivation of Faulhaber’s formula*, Applied Math. E-Notes **18** (2018), 124–126.
- [P] M. Primc, *Some Combinatorial Coincidences for Standard Representations of Affine Lie Algebras*, part of the book D. Adamović, P. Papi, *Affine, Vertex and W-algebras*, Springer INdAM Series (SINDAMS, Vol. **37**) (2019) 203–218.
- [PŠ1] M. Primc and T. Šikić, *Combinatorial Bases of Basic Modules for Affine Lie Algebras $C_n^{(1)}$* , J. Math. Phys. **57**(9) (2016), 1–19.
- [PŠ2] M. Primc and T. Šikić, *Leading terms of relations for standard modules of affine Lie algebras $C_n^{(1)}$* , Ramanujan J. **48** (2019), 509–543.

MIRKO PRIMC, UNIVERSITY OF ZAGREB, FACULTY OF SCIENCE, BIJENIČKA 30, 10000 ZAGREB, CROATIA

Email address: primc@math.hr

TOMISLAV ŠIKIĆ, UNIVERSITY OF ZAGREB, FACULTY OF ELECTRICAL ENGINEERING AND COMPUTING, UNSKA 3, 10000 ZAGREB, CROATIA

Email address: tomlslav.sikic@fer.hr