

On the Convergence of No-Regret Learning Dynamics in Time-Varying Games

Ioannis Anagnostides¹, Ioannis Panageas², Gabriele Farina³, and Tuomas Sandholm⁴

^{1,3,4}Carnegie Mellon University

²University of California Irvine

⁴Strategy Robot, Inc.

⁴Optimized Markets, Inc.

⁴Strategic Machine, Inc.

{ianagnos,gfarina,sandholm}@cs.cmu.edu, and ipanagea@ics.uci.edu

March 24, 2023

Abstract

Most of the literature on learning in games has focused on the restrictive setting where the underlying repeated game does not change over time. Much less is known about the convergence of no-regret learning algorithms in dynamic multiagent settings. In this paper, we characterize the convergence of *optimistic gradient descent (OGD)* in time-varying games. Our framework yields sharp convergence bounds for the equilibrium gap of OGD in zero-sum games parameterized on natural variation measures of the sequence of games, subsuming known results for static games. Furthermore, we establish improved *second-order* variation bounds under strong convexity-concavity, as long as each game is repeated multiple times. Our results also apply to time-varying *general-sum* multi-player games via a bilinear formulation of correlated equilibria, which has novel implications for meta-learning and for obtaining refined variation-dependent regret bounds, addressing questions left open in prior papers. Finally, we leverage our framework to also provide new insights on dynamic regret guarantees in static games.

1 Introduction

Most of the classical results in the literature on learning in games—exemplified by, among others, the work of Hart and Mas-Colell [HM00], Foster and Vohra [FV97], and Freund and Schapire [FS99]—rest on the assumption that the underlying repeated game remains invariant throughout the learning process. Yet, in many learning environments that is unrealistic [Duv+22; Zha+22; Car+19; MS21]. One such class is settings where the underlying game is actually changing, such as routing problems on the internet [MO11], online advertising auctions [LST16], and dynamic mechanism design [Pap+22; DMZ21]. Another such class consists of settings in which many similar games need to be solved [Har+22]. For example, one may want to solve variations of a game for the purpose of sensitivity analysis with respect to the modeling assumptions used to construct the game model. Another example is solving multiple versions of a game any one of which might be faced in the future.

Despite the considerable interest in such dynamic multiagent environments, much less is known about the convergence of *no-regret learning* algorithms in time-varying games. No-regret dynamics are natural learning algorithms that have desirable convergence properties in static settings. Also, the state-of-the-art algorithms for finding minimax equilibria in two-player zero-sum games are based on advanced forms of no-regret dynamics [FKS21; BS19a]. Indeed, all the superhuman milestones in poker have used them in the equilibrium-finding module of their architectures [Bow+15; BS17; BS19b].

In this paper, we seek to fill this knowledge gap by understanding properties of no-regret dynamics in time-varying games. In particular, we primarily investigate the convergence of *optimistic gradient descent (OGD)* [Chi+12; RS13] in time-varying games. Unlike traditional no-regret learning algorithms, such as (online) gradient descent, OGD has been recently shown to exhibit *last-iterate* convergence in *static* (two-player) zero-sum games [Das+18; GPD20; COZ22; GTG22]. For the more challenging scenario where the underlying game can vary in every round, a fundamental question arises: *Under what conditions on the sequence of games does OGD (with high probability) approximate the sequence of Nash equilibria?*

1.1 Our Results

In this paper, we develop a framework that enables us to characterize the convergence of OGD in time-varying games. Specifically, building on the work of Zhang et al. [Zha+22], our first contribution is to identify natural variation measures on the sequence of games whose sublinear growth guarantees that *almost all* iterates of OGD are (approximate) Nash equilibria in time-varying (two-player) zero-sum games (Corollary 3.6). More precisely, in Theorem 3.5 we derive a sharp non-asymptotic characterization of the equilibrium gap of OGD as a function of the *minimal first-order* variation of the Nash equilibria and the *second-order* variation of the payoff matrices. It is a compelling property, in light of the multiplicity of Nash equilibria, that the variation of the Nash equilibria is measured in terms of the most favorable—*i.e.*, one that minimizes the variation—such sequence. Additionally, we show that our convergence bounds can be further improved by considering a variation measure that depends on the deviation of *approximate* Nash equilibria of the games, a measure that could be arbitrarily smaller than the one based on (even the least varying) sequence of exact Nash equilibria (Proposition 3.3).

From a technical standpoint, our analysis revolves around a connection between the convergence of OGD in time-varying games and *dynamic regret*. In particular, building on the work of Zhang et al. [Zha+22], the first key observation is that *dynamic regret is always nonnegative under any sequence of Nash equilibria* (Property 3.2). By combining that property with a dynamic *RVU*

bound [Zha+22] (Lemmas 3.1 and A.1), we obtain in Theorem 3.4 a variation-dependent bound for the second-order path length of OGD in time-varying games. In turn, this leads to our main result, Theorem 3.5, discussed above. As such, we extend the regret-based framework of Anagnostides et al. [Ana+22b] from static to time-varying games. In the special case of static games, our result reduces to a tight $T^{-1/2}$ rate. It is worth stressing that Property 3.2 is in fact more general, being intricately tied to the admission of a *minimax theorem* (Property A.3), and applies even under a sequence of approximate Nash equilibria—with slackness that gently degrades with the approximation thereof.

Moreover, for strongly convex-concave time-varying games, we obtain a refined *second-order* variation bound on the sequence of Nash equilibria, as long as each game is repeated multiple times (Theorem 3.8); this is inspired by an improved second-order bound for dynamic regret under analogous conditions due to Zhang et al. [Zha+17]. As a byproduct of our techniques, we point out that *any* no-regret learners are approaching a Nash equilibrium under strong convexity-concavity (Proposition 3.9). Those results apply even in non-strongly convex-concave settings by suitably trading-off the magnitude of a regularizer that makes the game strongly convex-concave. This offers significant gains in the *meta-learning setting* as well, wherein each game is repeated multiple times.

Next, we extend our results to time-varying general-sum multi-player games via a bilinear formulation of *correlated equilibria*. As such, we recover similar convergence bounds parameterized on the variation of the correlated equilibria (Theorem 3.12). To illustrate the power of our framework, we immediately recover natural and algorithm-independent similarity measures for the meta-learning setting (Proposition A.13) even in general games (Corollary A.22), thereby addressing an open question of Harris et al. [Har+22]. Our techniques also imply new per-player regret bounds in zero-sum and general-sum games (Corollaries 3.7 and A.23), the latter addressing a question left open by Zhang et al. [Zha+22]. We further parameterize the convergence of (vanilla) gradient descent in time-varying *potential* games in terms of the deviation of the potential functions (Theorem 3.10).

Finally, building on our techniques in time-varying games, we investigate the best dynamic-regret guarantees possible in *static* games. Although this is a basic question, it has apparently eluded prior research. We first show that instances of optimistic mirror descent guarantee $O(\sqrt{T})$ dynamic per-player regret (Proposition 3.13), matching the known rate of (online) gradient descent but for the significantly weaker notion of *external* regret. We further point out that $O(\log T)$ dynamic regret is attainable, but in a stronger two-point feedback model. In stark contrast, even obtaining sublinear dynamic regret for each player is precluded in general-sum games (Proposition 3.15). This motivates studying a relaxation of dynamic regret that constrains the number of switches in the comparator, for which we derive accelerated rates in general games (Theorem 3.16) by leveraging the techniques of Syrgkanis et al. [Syr+15] in conjunction with our dynamic RVU bound (Lemma 3.1).

1.2 Further Related Work

Even in static (two-player) zero-sum games, the pointwise convergence of no-regret learning algorithms is a tenuous affair. Indeed, traditional learning dynamics within the no-regret framework, such as (online) mirror descent, may even diverge away from the equilibrium; *e.g.*, see [SAF02; MPP18; Vla+20; GVM21]. Notwithstanding, the *empirical frequency* of no-regret learners is well-known to approach the set of Nash equilibria in zero-sum games [FS99], and the set of coarse correlated equilibria in general-sum games [HM00]—a standard relaxation of the Nash equilibrium [MV78; Aum74]. Unfortunately, those classical results are of little use beyond static games, thereby offering a crucial impetus for investigating iterate-convergence in games with a time-varying component—a ubiquitous theme in many practical scenarios of interest [DMZ21; Pap+22; Ven21; Gar17; Van10;

RG22; PKB22; YH15; RJW21].

Indeed, there has been a considerable effort endeavoring to extend the scope of traditional game-theoretic results to the time-varying setting, approached from a variety of different standpoints [LST16; Zha+22; Car+19; MO11; MS21; Duv+22]. In particular, our techniques in Section 3.1 are strongly related to the ones used by Zhang et al. [Zha+22], but our primary focus is different: Zhang et al. [Zha+22] were mainly interested in obtaining variation-dependent regret bounds, while our results revolve around iterate-convergence to Nash equilibria. We stress again that minimizing regret and approaching Nash equilibria are two inherently distinct problems, although connections have emerged [Ana+22b], and are further cultivated in this paper.

Another closely related direction is on *meta-learning* in games [Har+22], wherein each game can be repeated for multiple iterations. Such considerations are motivated in part by a number of use-cases in which many “similar” games—or multiple game variations—ought to be solved [BS16], such as Poker with different stack-sizes. While the meta-learning problem is a special case of our general setting, our results are strong enough to have new implications for meta-learning in games, even though the algorithms considered herein are not tailored to operate in that setting.

Finally, although our focus is on the convergence of OGD in time-varying games, some of our results—namely, the ones formalized in Appendix A.1.7—can be viewed as part of an ongoing effort to characterize the class of variational inequalities (VIs) that are amenable to efficient algorithms; see [DDJ21; CZ22; Azi+20; BMW21; CP04; DL15; MV21; MRS20; Nou+19; Son+20; YKH20; Das22], and references therein. We also highlight that the techniques used to establish last-iterate convergence even in monotone (time-invariant) settings are particularly involved [GPD20; COZ22; GTG22]; the simplicity of our framework, therefore, in the more challenging time-varying regime is a compelling aspect of this paper.

2 Preliminaries

Notation We let $\mathbb{N} := \{1, 2, \dots\}$ be the set of natural numbers. For a number $p \in \mathbb{N}$, we let $\llbracket p \rrbracket := \{1, \dots, p\}$. For a vector $\mathbf{w} \in \mathbb{R}^d$, we use $\|\mathbf{w}\|_2$ to represent its Euclidean norm; we also overload that notation so that $\|\cdot\|_2$ denotes the spectral norm when the argument is a matrix.

For a two-player zero-sum game, we denote by $\mathcal{X} \subseteq \mathbb{R}^{d_x}$ and $\mathcal{Y} \subseteq \mathbb{R}^{d_y}$ the strategy sets of the two players—namely, Player x and Player y , respectively—where $d_x, d_y \in \mathbb{N}$ represent the corresponding dimensions. It is assumed that \mathcal{X} and \mathcal{Y} are nonempty convex and compact sets. For example, in the special case where $\mathcal{X} := \Delta^{d_x}$ and $\mathcal{Y} := \Delta^{d_y}$ —each set corresponds to a probability simplex—the game is said to be in *normal form*. Further, we denote by $D_{\mathcal{X}}$ the ℓ_2 -diameter of \mathcal{X} , and by $\|\mathcal{X}\|_2$ the maximum ℓ_2 -norm attained by a point in \mathcal{X} . We will always assume that the strategy sets remain invariant, while the payoff matrix can change in each round. For notational convenience, we will denote by $\mathbf{z} := (\mathbf{x}, \mathbf{y})$ the concatenation of \mathbf{x} and \mathbf{y} , and by $\mathcal{Z} := \mathcal{X} \times \mathcal{Y}$ the Cartesian product of \mathcal{X} and \mathcal{Y} . In general n -player games, we instead use subscripts indexed by $i \in \llbracket n \rrbracket$ to specify quantities related to a player. Superscripts are typically reserved to identify the time index. Finally, to simplify the exposition, we use the $O(\cdot)$ notation to suppress time-independent parameters of the problem; precise statements are given in Appendix A.

Dynamic regret We operate in the usual online learning setting under full feedback. Namely, at every time $t \in \mathbb{N}$ the learner decides on a strategy $\mathbf{x}^{(t)} \in \mathcal{X}$, and then observes a utility $\mathbf{x} \mapsto \langle \mathbf{x}, \mathbf{u}_x^{(t)} \rangle$, for $\mathbf{u}_x^{(t)} \in \mathbb{R}^{d_x}$. Following Daskalakis, Deckelbaum, and Kim [DDK11], we will insist on allowing only

$O(1)$ previous utilities to be stored; this will preclude trivial exploration protocols when learning in games.

A strong performance benchmark in this online setting is *dynamic regret*, defined for a time horizon $T \in \mathbb{N}$ as follows:

$$\text{DReg}_x^{(T)}(s_x^{(T)}) := \sum_{t=1}^T \langle \mathbf{x}^{(t,*)} - \mathbf{x}^{(t)}, \mathbf{u}_x^{(t)} \rangle, \quad (1)$$

where $s_x^{(T)} := (\mathbf{x}^{(1,*)}, \dots, \mathbf{x}^{(T,*)}) \in \mathcal{X}^T$ is the sequence of *comparators*; setting $\mathbf{x}^{(1,*)} = \mathbf{x}^{(2,*)} = \dots = \mathbf{x}^{(T,*)}$ in (1) we recover the standard notion of (*external*) *regret* (denoted simply by $\text{Reg}_x^{(T)}$), which is commonly used to establish convergence of the time-average strategies in static two-player zero-sum games [FS99]. On the other hand, the more general notion of dynamic regret, introduced in (1), has been extensively used in more dynamic environments; e.g., [Zha+20; Zha+17; Jad+15; Ces+12; HS09; Kal+21; Mok+16]. We also let $\text{DReg}_x^{(T)} := \max_{s_x^{(T)} \in \mathcal{X}^T} \text{DReg}_x^{(T)}(s_x^{(T)})$. While ensuring $o(T)$ dynamic regret is clearly hopeless in a truly adversarial environment, Section 3.4 reveals that non-trivial guarantees are possible when learning in zero-sum games.

Optimistic gradient descent *Optimistic gradient descent (OGD)* [Chi+12; RS13] is a no-regret algorithm defined with the following update rule:

$$\begin{aligned} \mathbf{x}^{(t)} &:= \Pi_{\mathcal{X}} \left(\hat{\mathbf{x}}^{(t)} + \eta \mathbf{m}_x^{(t)} \right), \\ \hat{\mathbf{x}}^{(t+1)} &:= \Pi_{\mathcal{X}} \left(\hat{\mathbf{x}}^{(t)} + \eta \mathbf{u}_x^{(t)} \right). \end{aligned} \quad (\text{OGD})$$

Here, $\eta > 0$ is the *learning rate*; $\hat{\mathbf{x}}^{(1)} := \arg \min_{\hat{\mathbf{x}} \in \mathcal{X}} \|\hat{\mathbf{x}}\|_2^2$ represents the initialization of OGD; $\mathbf{m}_x^{(t)} \in \mathbb{R}^{d_x}$ is the *prediction vector* at time t , and it is set as $\mathbf{m}_x^{(t)} := \mathbf{u}_x^{(t-1)}$ when $t \geq 2$, and $\mathbf{m}_x^{(1)} := \mathbf{0}_{d_x}$; and finally, $\Pi_{\mathcal{X}}(\cdot)$ represents the Euclidean projection to the set \mathcal{X} , which is well-defined, and can be further computed efficiently for structured sets, such as the probability simplex. For our purposes, we will posit access to a projection oracle for the set \mathcal{X} , in which case the update rule (OGD) is efficiently implementable.

In a multi-player n -player game, each Player $i \in \llbracket n \rrbracket$ is associated with a utility function $u_i : \times_{i=1}^n \mathcal{X}_i \rightarrow \mathbb{R}$. We recall the following fundamental definition [Nas50].

Definition 2.1 (Approximate Nash equilibrium). A joint strategy profile $(\mathbf{x}_1^*, \dots, \mathbf{x}_n^*) \in \times_{i=1}^n \mathcal{X}_i$ is an ϵ -approximate Nash equilibrium (NE), for an $\epsilon \geq 0$, if for any Player $i \in \llbracket n \rrbracket$ and any possible deviation $\mathbf{x}'_i \in \mathcal{X}_i$,

$$u_i(\mathbf{x}_1^*, \dots, \mathbf{x}_i^*, \dots, \mathbf{x}_n^*) \geq u_i(\mathbf{x}_1^*, \dots, \mathbf{x}'_i, \dots, \mathbf{x}_n^*) - \epsilon.$$

3 Convergence in Time-Varying Games

In this section, we formalize our results regarding convergence in time-varying games. We organize this section as follows: First, in Section 3.1, we build the foundations of our framework by studying the convergence of OGD in time-varying bilinear saddle-point problems, culminating in the non-asymptotic characterization of Theorem 3.5; Section 3.2 formalizes our improvements under strong convexity-concavity; we then extend our results (in Section 3.3) to time-varying multi-player general-sum and potential games; and finally, Section 3.4 concerns dynamic regret guarantees in static games.

3.1 Bilinear Saddle-Point Problems

We first study an online learning setting wherein two players interact in a sequence of time-varying bilinear saddle-point problems. More precisely, we assume that in every repetition $t \in \llbracket T \rrbracket$ the players select a pair of strategies $(\mathbf{x}^{(t)}, \mathbf{y}^{(t)}) \in \mathcal{X} \times \mathcal{Y}$. Then, Player x receives the utility $\mathbf{u}_x^{(t)} := -\mathbf{A}^{(t)}\mathbf{y}^{(t)} \in \mathbb{R}^{d_x}$, where $\mathbf{A}^{(t)} \in \mathbb{R}^{d_x \times d_y}$ represents the payoff matrix at the t -th repetition; similarly, Player y receives the utility $\mathbf{u}_y^{(t)} := (\mathbf{A}^{(t)})^\top \mathbf{x}^{(t)} \in \mathbb{R}^{d_y}$. The proofs of this subsection are included in Appendix A.1.

Dynamic RVU bound The first key ingredient that we need is the property of *regret bounded by variation in utilities (RVU)*, in the sense of Syrgkanis et al. [Syr+15], but with respect to dynamic regret; such a bound is established below, and is analogous to that obtained by Zhang et al. [Zha+22].

Lemma 3.1 (RVU bound for dynamic regret). *Consider any sequence of utilities $(\mathbf{u}_x^{(1)}, \dots, \mathbf{u}_x^{(T)})$ up to time $T \in \mathbb{N}$. The dynamic regret (1) of OGD with respect to any sequence of comparators $(\mathbf{x}^{(1,\star)}, \dots, \mathbf{x}^{(T,\star)}) \in \mathcal{X}^T$ can be bounded by*

$$\begin{aligned} \frac{D_{\mathcal{X}}^2}{2\eta} + \frac{D_{\mathcal{X}}}{\eta} \sum_{t=1}^{T-1} \|\mathbf{x}^{(t+1,\star)} - \mathbf{x}^{(t,\star)}\|_2 + \eta \sum_{t=1}^T \|\mathbf{u}_x^{(t)} - \mathbf{m}_x^{(t)}\|_2^2 \\ - \frac{1}{2\eta} \sum_{t=1}^T \left(\|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t)}\|_2^2 + \|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t+1)}\|_2^2 \right). \end{aligned} \quad (2)$$

In the special case of external regret— $\mathbf{x}^{(1,\star)} = \mathbf{x}^{(2,\star)} = \dots = \mathbf{x}^{(T,\star)}$ —(2) recovers the bound for OGD of Syrgkanis et al. [Syr+15]. The key takeaway from Lemma 3.1 is that the overhead of dynamic regret in (2) grows with the *first-order* variation of the sequence of comparators. In Lemma A.1 we also articulate an extension of Lemma 3.1 for the more general *optimistic mirror descent (OMD)* algorithm under a certain class of Bregman divergences.

Having established Lemma 3.1, we next point out a crucial property (previously also observed by Zhang et al. [Zha+22]): by selecting a sequence of Nash equilibria (recall Definition 2.1) as the comparators, the *sum* of the players' dynamic regrets is always nonnegative:

Property 3.2. *Suppose that $\mathcal{Z} \ni \mathbf{z}^{(t,\star)} = (\mathbf{x}^{(t,\star)}, \mathbf{y}^{(t,\star)})$ is an $\epsilon^{(t)}$ -approximate Nash equilibrium of the t -th game. Then, for $s_x^{(T)} = (\mathbf{x}^{(t,\star)})_{1 \leq t \leq T}$ and $s_y^{(T)} = (\mathbf{y}^{(t,\star)})_{1 \leq t \leq T}$,*

$$\text{DReg}_x^{(T)}(s_x^{(T)}) + \text{DReg}_y^{(T)}(s_y^{(T)}) \geq -2 \sum_{t=1}^T \epsilon^{(t)}.$$

In particular, if $\epsilon^{(t)} = 0$ for all $t \in \llbracket T \rrbracket$, we have

$$\text{DReg}_x^{(T)}(s_x^{(T)}) + \text{DReg}_y^{(T)}(s_y^{(T)}) \geq 0. \quad (3)$$

In fact, as we show in Property A.3, Property 3.2 applies even in certain (time-varying) nonconvex-nonconcave min-max optimization problems, and it is a consequence of the *minimax theorem*; Property 3.2 also holds for time-varying variational inequalities (VIs) that satisfy the so-called *MVI property* (see Remark A.4). For comparison, it is evident that under a *static* sequence of two-player zero-sum games, it holds that $\text{Reg}_x^{(T)} + \text{Reg}_y^{(T)} \geq 0$.

Next, building on the work of Zhang et al. [Zha+22], let us introduce some natural measures of the games’ variation. First, the first-order variation of the Nash equilibria is defined for $T \geq 2$ as

$$\mathcal{V}_{\text{NE}}^{(T)} := \inf_{\mathbf{z}^{(t,\star)} \in \mathcal{Z}^{(t,\star)}, \forall t \in \llbracket T \rrbracket} \sum_{t=1}^{T-1} \|\mathbf{z}^{(t+1,\star)} - \mathbf{z}^{(t,\star)}\|_2, \quad (4)$$

where $\mathcal{Z}^{(t,\star)}$ is the (nonempty) set of Nash equilibria of the t -th game. We recall that there can be a multiplicity of Nash equilibria [van91]; as such, a compelling feature of the variation measure (4) is that it depends on the most favorable sequence of Nash equilibria—one that minimizes the first-order variation.

It is also important to point out the well-known fact that Nash equilibria can change abruptly even under a “small” perturbation in the payoff matrix (see Example A.5), which is a caveat of the variation (4). To address this, and in accordance with Property 3.2, we consider a more favorable variation measure, defined as

$$\mathcal{V}_{\epsilon\text{-NE}}^{(T)} := \inf \left\{ \sum_{t=1}^{T-1} \|\mathbf{z}^{(t+1,\star)} - \mathbf{z}^{(t,\star)}\|_2 + C \sum_{t=1}^T \epsilon^{(t)} \right\},$$

for a sufficiently large parameter $C > 0$; the infimum above is subject to $\epsilon^{(t)} \in \mathbb{R}_{\geq 0}$ and $\mathbf{z}^{(t,\star)} \in \mathcal{Z}_{\epsilon^{(t)}}^{(t,\star)}$ for all $t \in \llbracket T \rrbracket$, where we denote by $\mathcal{Z}_{\epsilon^{(t)}}^{(t,\star)}$ the set of $\epsilon^{(t)}$ -approximate NE. It is evident that $\mathcal{V}_{\epsilon\text{-NE}}^{(T)} \leq \mathcal{V}_{\text{NE}}^{(T)}$ since one can take $\epsilon^{(1)} = \dots = \epsilon^{(T)} = 0$; in fact, $\mathcal{V}_{\epsilon\text{-NE}}^{(T)}$ can be arbitrarily smaller:

Proposition 3.3. *For any $T \geq 4$, there is a sequence of T games such that $\mathcal{V}_{\text{NE}}^{(T)} \geq \frac{T}{2}$ while $\mathcal{V}_{\epsilon\text{-NE}}^{(T)} \leq \delta$, for any $\delta > 0$.*

Moreover, we also introduce a quantity that captures the variation of the payoff matrices:

$$\mathcal{V}_{\mathbf{A}}^{(T)} := \sum_{t=1}^{T-1} \|\mathbf{A}^{(t+1)} - \mathbf{A}^{(t)}\|_2^2, \quad (5)$$

where we recall that here $\|\cdot\|_2$ denotes the spectral norm. Unlike (4), the variation measure (5) depends on the *second-order* variation (of the payoff matrices), which could translate to a lower-order impact compared to (4) (see, *e.g.*, Corollary A.8). We stress that while our convergence bounds will be parameterized based on (4) and (5), the underlying algorithm—namely OGD—will remain oblivious to those variation measures.

We are ready now to establish a refined bound on the second-order path-length of OGD in time-varying zero-sum games.

Theorem 3.4 (Detailed version in Theorem A.6). *Suppose that both players employ OGD with learning rate $\eta \leq \frac{1}{4L}$ in a time-varying bilinear saddle-point problem, where $L := \max_{t \in \llbracket T \rrbracket} \|\mathbf{A}^{(t)}\|_2$. Then, for any $T \in \mathbb{N}$, the second-order path length $\sum_{t=1}^T (\|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t)}\|_2^2 + \|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t+1)}\|_2^2)$ can be bounded by*

$$O\left(1 + \mathcal{V}_{\epsilon\text{-NE}}^{(T)} + \mathcal{V}_{\mathbf{A}}^{(T)}\right). \quad (6)$$

It is worth noting that when the deviation of the payoff matrices is controlled by the deviation of the players' strategies, in the sense that $\sum_{t=1}^{T-1} \|\mathbf{A}^{(t+1)} - \mathbf{A}^{(t)}\|_2^2 \leq W^2 \sum_{t=1}^{T-1} \|\mathbf{z}^{(t+1)} - \mathbf{z}^{(t)}\|_2^2$ for some parameter $W \in \mathbb{R}_{>0}$, the variation measure $\mathcal{V}_{\mathbf{A}}^{(T)}$ in (6)—and in the subsequent convergence bounds—can be entirely eliminated; see Corollary A.8. The same, in fact, applies under an improved prediction mechanism (Remark A.12), but that prediction is not implementable in our online learning setting.

Armed with Theorem 3.4, we are ready to establish Theorem 3.5. The key observation is that the Nash equilibrium gap at the t -th game can be bounded in terms of the quantity $\|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t)}\|_2 + \|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t+1)}\|_2$, which in turn allows us to use (6) to bound the cumulative (squared) Nash equilibrium gaps across the sequence of games; the aforeclaimed property was established in [Ana+22b] for static games (Claim A.9), but readily extends to our setting as well, and in fact applies to any member of OMD under a *smooth regularizer*. Below, we use the notation $\text{EQGAP}^{(t)}(\mathbf{z}^{(t)}) \in \mathbb{R}_{\geq 0}$ to represent the Nash equilibrium gap of the joint strategy profile $\mathbf{z}^{(t)} \in \mathcal{Z}$ at the t -th game.

Theorem 3.5 (Main result; detailed version in Theorem A.10). *Suppose that both players employ OGD with learning rate $\eta = \frac{1}{4L}$ in a time-varying bilinear saddle-point problem, where $L := \max_{t \in [T]} \|\mathbf{A}^{(t)}\|_2$. Then,*

$$\sum_{t=1}^T \left(\text{EQGAP}^{(t)}(\mathbf{z}^{(t)}) \right)^2 = O \left(1 + \mathcal{V}_{\epsilon-NE}^{(T)} + \mathcal{V}_{\mathbf{A}}^{(T)} \right), \quad (7)$$

where $(\mathbf{z}^{(t)})_{1 \leq t \leq T}$ is the sequence of joint strategy profiles produced by OGD.

We next state some immediate consequences of this result. (Item 2 below follows from (7) by Jensen's inequality.)

Corollary 3.6. *In the setting of Theorem 3.5,*

1. *If at least a δ -fraction of the iterates of OGD have at least $\epsilon > 0$ Nash equilibrium gap, then $\epsilon^2 \delta \leq O \left(\frac{1}{T} \left(\mathcal{V}_{\epsilon-NE}^{(T)} + \mathcal{V}_{\mathbf{A}}^{(T)} + 1 \right) \right)$;*
2. *The average Nash equilibrium gap of OGD is bounded as $O \left(\sqrt{\frac{1}{T} \left(\mathcal{V}_{\epsilon-NE}^{(T)} + \mathcal{V}_{\mathbf{A}}^{(T)} + 1 \right)} \right)$.*

In particular, in terms of asymptotic implications, if $\lim_{T \rightarrow +\infty} \frac{\mathcal{V}_{\epsilon-NE}^{(T)}}{T}, \lim_{T \rightarrow +\infty} \frac{\mathcal{V}_{\mathbf{A}}^{(T)}}{T} = 0$, then (i) for any $\epsilon > 0$ the fraction of iterates of OGD with at least an ϵ Nash equilibrium gap converges to 0; and (ii) the average Nash equilibrium gap of the iterates of OGD converges to 0.

In the special case where $\mathcal{V}_{\epsilon-NE}^{(T)}, \mathcal{V}_{\mathbf{A}}^{(T)} = O(1)$, Theorem 3.5 recovers the $T^{-1/2}$ rate of OGD in static bilinear saddle-point problems. It is also worth pointing out that Theorem 3.5 readily extends to more general time-varying variational inequality problems as well (Remark A.4).

We also state below another interesting consequence of Theorem 3.4, which bounds each player's *individual regret* parameterized based on the variation measures.

Corollary 3.7 (Detailed version in Corollary A.11). *In the setup of Theorem 3.4, it holds that $\text{Reg}_x^{(T)}, \text{Reg}_y^{(T)} = O \left(\frac{1}{\eta} + \eta(\mathcal{V}_{NE}^{(T)} + \mathcal{V}_{\mathbf{A}}^{(T)}) \right)$.*

The $O(\cdot)$ notation here is considered in the regime $\eta \ll 1$. Hence, selecting optimally the learning rate gives an $O(\sqrt{\mathcal{V}_{\text{NE}}^{(T)} + \mathcal{V}_{\mathbf{A}}^{(T)}}$) bound on the *individual* regret of each player; while that optimal value depends on the variation measures, which are not known to the learners, there are techniques that would allow bypassing this [Zha+22]. Corollary 3.7 can also be readily parameterized in terms of the improved variation measure $\mathcal{V}_{\epsilon\text{-NE}}^{(T)}$. Finally, in Appendix A.1.7 we highlight certain implications of our framework on solving (static) general VIs.

Meta-Learning Our results also have immediate applications in the *meta-learning* setting [Har+22]. More precisely, meta-learning in games is a special case of time-varying games which consists of a sequence of $H \in \mathbb{N}$ separate games, each of which is repeated for $m \in \mathbb{N}$ consecutive rounds, so that $T := m \times H$. The central goal in meta-learning is to obtain convergence bounds parameterized by the *similarity* of the games; identifying suitable similarity metrics is a central question in that line of work.

In this context, we highlight that Theorem 3.5 readily provides a meta-learning guarantee parameterized by the following notion of similarity between the Nash equilibria:

$$\inf_{\mathbf{z}^{(h,\star)} \in \mathcal{Z}^{(h,\star)}, \forall h \in \llbracket H \rrbracket} \sum_{h=1}^{H-1} \|\mathbf{z}^{(h+1,\star)} - \mathbf{z}^{(h,\star)}\|_2, \quad (8)$$

where $\mathcal{Z}^{(h,\star)}$ is the set of Nash equilibria of the h -th game in the meta-learning sequence,¹ as well as the similarity of the payoff matrices—corresponding to the term $\mathcal{V}_{\mathbf{A}}^{(T)}$ in (7). In fact, under a suitable prediction—the one used by Harris et al. [Har+22]—the dependence on $\mathcal{V}_{\mathbf{A}}^{(T)}$ can be entirely removed; see Proposition A.13 for our formal result. A compelling aspect of our meta-learning guarantee is that the considered algorithm is oblivious to the boundaries of the meta-learning. We further provide some novel results on meta-learning in general-sum games in Section 3.3.

3.2 Strongly Convex-Concave Games

In this subsection, we show that under additional structure we can significantly improve the variation measures established in Theorem 3.4. More precisely, we first assume that each objective function $f(\mathbf{x}, \mathbf{y})$ is μ -strongly convex with respect to \mathbf{x} and μ -strongly concave with respect to \mathbf{y} . Our second assumption is that each game is played for *multiple* rounds $m \in \mathbb{N}$, instead of only a single round; this is akin to the meta-learning setting. The key insight is that, as long as m is large enough, $m = \Omega(1/\mu)$, those two assumptions suffice to obtain a *second-order* variation bound in terms of the sequence of Nash equilibria,

$$\mathcal{S}_{\text{NE}}^{(H)} := \sum_{h=1}^{H-1} \|\mathbf{z}^{(h+1,\star)} - \mathbf{z}^{(h,\star)}\|_2^2, \quad (9)$$

where $\mathbf{z}^{(h,\star)}$ is a Nash equilibrium of the h -th game. This significantly refines the result of Theorem 3.4, and is inspired by the improved dynamic regret bounds obtained by Zhang et al. [Zha+17]. Below we sketch the key ideas of the improvement; the proofs are included in Appendix A.2.

In this setting, it is assumed that Player x obtains the utility $\mathbf{u}_x^{(t)} := -\nabla_{\mathbf{x}} f^{(t)}(\mathbf{x}^{(t)}, \mathbf{y}^{(t)})$ at every time $t \in \llbracket T \rrbracket$, while its regret will be denoted by $\text{Reg}_{\mathcal{L}, y}^{(T)}$; similar notation applies for Player y . The

¹In accordance to Theorem 3.5, (8) can be refined using a sequence of approximate Nash equilibria.

first observation is that, focusing on a single (static) game, under strong convexity-concavity the sum of the players' regrets are *strongly nonnegative* (Lemma A.15):

$$\text{Reg}_{\mathcal{L},x}^{(m)}(\mathbf{x}^*) + \text{Reg}_{\mathcal{L},y}^{(m)}(\mathbf{y}^*) \geq \frac{\mu}{2} \sum_{t=1}^m \|\mathbf{z}^{(t)} - \mathbf{z}^*\|_2^2, \quad (10)$$

for any Nash equilibrium $\mathbf{z}^* \in \mathcal{Z}$ of the game. In turn, this can be cast in terms of dynamic regret over the sequence of the h games (Lemma A.16). Next, combining those dynamic-regret lower bounds with a suitable RVU-type property leads to a refined second-order path length bound as long as that $m = \Omega(1/\mu)$, which in turn leads to our main result below. Before we present its statement, let us introduce the following measure of variation of the gradients:

$$\mathcal{V}_{\nabla f}^{(H)} := \sum_{h=1}^{H-1} \max_{\mathbf{z} \in \mathcal{Z}} \|F^{(h+1)}(\mathbf{z}) - F^{(h)}(\mathbf{z})\|_2^2, \quad (11)$$

where let $F : \mathbf{z} := (\mathbf{x}, \mathbf{y}) \mapsto (\nabla_{\mathbf{x}} f(\mathbf{x}, \mathbf{y}), -\nabla_{\mathbf{y}} f(\mathbf{x}, \mathbf{y}))$. This variation measure is analogous to $\mathcal{V}_{\mathbf{A}}^{(T)}$ we introduced in (5) for time-varying bilinear saddle-point problems.

Theorem 3.8 (Detailed version in Theorem A.18). *Let $f^{(h)} : \mathcal{X} \times \mathcal{Y}$ be a μ -strongly convex-concave and L -smooth function, for $h \in \llbracket H \rrbracket$. Suppose that both players employ OGD with learning rate $\eta = \min \left\{ \frac{1}{8L}, \frac{1}{2\mu} \right\}$ for T repetitions, where $T = m \times H$ and $m \geq \frac{2}{\eta\mu}$. Then,*

$$\sum_{t=1}^T \left(\text{EQGAP}^{(t)}(\mathbf{z}^{(t)}) \right)^2 = O(1 + \mathcal{S}_{NE}^{(H)} + \mathcal{V}_{\nabla f}^{(H)}),$$

where $\mathcal{S}_{NE}^{(H)}$ and $\mathcal{V}_{\nabla f}^{(H)}$ are defined in (9) and (11).

Our techniques also imply improved regret bounds in this setting, as we formalize in Corollary A.19.

There is another immediate but important implication of (10): *any* no-regret algorithm in a (static) strongly convex-concave setting ought to be approaching the Nash equilibrium; in contrast, this property is spectacularly false in (general) monotone settings [MPP18].

Proposition 3.9. *Let $f : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}$ be a μ -strongly convex-concave function. If players incur regrets such that $\text{Reg}_{\mathcal{L},x}^{(T)} + \text{Reg}_{\mathcal{L},y}^{(T)} \leq CT^{1-\omega}$, for some parameters $C > 0$ and $\omega \in (0, 1]$, then for any $\epsilon > 0$ and $T > \left(\frac{2C}{\mu\epsilon^2} \right)^{1/\omega}$ there is a pair of strategies $\mathbf{z}^{(t)} \in \mathcal{Z}$ such that $\|\mathbf{z}^{(t)} - \mathbf{z}^*\|_2 \leq \epsilon$, where \mathbf{z}^* is a Nash equilibrium.*

The insights of this subsection are also of interest in general monotone settings by incorporating a strongly convex regularizer; tuning its magnitude allows us to trade-off between a better approximation and the benefits of strong convexity-concavity revealed in this subsection.

3.3 General-Sum Multi-player Games

Next, we turn our attention to general-sum multi-player games. For simplicity, in this subsection we posit that the game is represented in normal form, so that each Player $i \in \llbracket n \rrbracket$ has a finite set of available actions \mathcal{A}_i , and $\mathcal{X}_i := \Delta(\mathcal{A}_i)$. The proofs of this subsection are included in Appendix A.3.

Potential Games First, we study the convergence of (online) gradient descent (GD) in time-varying *potential games* (see Definition A.20 for the formal description).² In our time-varying setup, it is assumed that each round $t \in \llbracket T \rrbracket$ corresponds to a different potential game described with a potential function $\Phi^{(t)}$. We further let $d : (\Phi, \Phi') \mapsto \max_{\mathbf{z} \in \times_{i=1}^n \mathcal{X}_i} (\Phi(\mathbf{z}) - \Phi'(\mathbf{z}))$, so that $\mathcal{V}_\Phi^{(T)} := \sum_{t=1}^{T-1} d(\Phi^{(t)}, \Phi^{(t+1)})$; we emphasize the fact that $d(\cdot, \cdot)$ is not symmetric. Analogously to Theorem 3.5, we use $\text{EQGAP}^{(t)}(\mathbf{z}^{(t)}) \in \mathbb{R}_{\geq 0}$ to represent the NE gap of the joint strategy profile $\mathbf{z}^{(t)} := (\mathbf{x}_1^{(t)}, \dots, \mathbf{x}_n^{(t)})$ at the t -th game.

Theorem 3.10. *Suppose that each player employs (online) GD with a sufficiently small learning rate. Then,*

$$\sum_{t=1}^T \left(\text{EQGAP}^{(t)}(\mathbf{z}^{(t)}) \right)^2 = O(\Phi_{\max} + \mathcal{V}_\Phi^{(T)}),$$

where Φ_{\max} is such that $|\Phi^{(t)}(\cdot)| \leq \Phi_{\max}$ for any $t \in \llbracket T \rrbracket$.

We refer to Appendix B for some illustrative experiments.

General games Unfortunately, unlike the settings considered thus far, computing Nash equilibria in general games is computationally hard [DGP08; CDT09]. Instead, learning algorithms are known to converge to relaxations of the Nash equilibrium, known as (*coarse*) *correlated equilibria*. For our purposes, we will employ a bilinear formulation of (coarse) correlated equilibria, which dates back to the seminal work of Hart and Schmeidler [HS89]. This will allow us to translate the results of Section 3.1 to general multi-player games.

Specifically, correlated equilibria³ can be expressed via a game between the n players and a *mediator*. Intuitively, the mediator is endeavoring to identify a correlated strategy $\boldsymbol{\mu} \in \Xi := \Delta(\times_{i=1}^n \mathcal{A}_i)$ for which no player has an incentive to deviate from the recommendation. In contrast, the players are trying to optimally deviate so as to maximize their own utility. More precisely, there exist matrices $\mathbf{A}_1, \dots, \mathbf{A}_n$, with each matrix \mathbf{A}_i depending solely on the utility of Player i , for which the bilinear problem can be expressed as

$$\min_{\boldsymbol{\mu} \in \Xi} \max_{(\bar{\mathbf{x}}_1, \dots, \bar{\mathbf{x}}_n) \in \times_{i=1}^n \bar{\mathcal{X}}_i} \sum_{i=1}^n \boldsymbol{\mu}^\top \mathbf{A}_i \bar{\mathbf{x}}_i, \quad (12)$$

where $\bar{\mathcal{X}}_i := \text{conv}(\mathcal{X}_i, \mathbf{0})$; incorporating the $\mathbf{0}$ vector will be useful for our purposes. This zero-sum game has the property that there exists a strategy $\boldsymbol{\mu}^* \in \Xi$ such that $\max_{\bar{\mathbf{x}}_i \in \bar{\mathcal{X}}_i} (\boldsymbol{\mu}^*)^\top \mathbf{A}_i \bar{\mathbf{x}}_i \leq 0$, for any Player $i \in \llbracket n \rrbracket$, which corresponds to a correlated equilibrium.

Before we proceed, it is important to note that the learning paradigm considered here deviates from the traditional one in that there is an additional learning agent, resulting in a less decentralized protocol. Yet, the dynamics induced by solving (12) via online algorithms remain *uncoupled* [HM00], in the sense that each player obtains feedback—corresponding to the deviation benefit—that depends solely on its own utility.

Now in the time-varying setting, the matrices $\mathbf{A}_1, \dots, \mathbf{A}_n$ that capture the players' utilities can change in each repetition. Crucially, we show that the structure of the induced bilinear problem (12)

²Unlike two-player zero-sum games, gradient descent is known to approach Nash equilibria in potential games.

³The following bilinear formulation applies to coarse correlated equilibria as well (with different payoff matrices), but we will focus solely on the stronger variant (CE) for the sake of exposition.

is such that there is a sequence of correlated equilibria that guarantee nonnegative dynamic regret; this refines Property 3.2 in that only one player’s strategies suffice to guarantee nonnegativity, even if the strategies of the other player remain invariant. Below, we denote by $\text{DReg}_\mu^{(T)}$ the dynamic regret of the min player in (12), and by $\text{Reg}_i^{(T)}$ the regret of each Player i up to time $T \in \mathbb{N}$, so that the regret of the max player in (12) can be expressed as $\sum_{i=1}^n \text{Reg}_i^{(T)}$.

Property 3.11. *Suppose that $\Xi \ni \boldsymbol{\mu}^{(t, \star)}$ is a correlated equilibrium of the game at any time $t \in \llbracket T \rrbracket$. Then,*

$$\text{DReg}_\mu^{(T)}(\boldsymbol{\mu}^{(1, \star)}, \dots, \boldsymbol{\mu}^{(T, \star)}) + \sum_{i=1}^n \text{Reg}_i^{(T)} \geq 0.$$

As a result, this enables us to apply Theorem 3.5 parameterized on (i) the variation of the CE

$$\mathcal{V}_{\text{CE}}^{(T)} := \inf_{\boldsymbol{\mu}^{(t, \star)} \in \Xi^{(t, \star)}, \forall t \in \llbracket T \rrbracket} \sum_{t=1}^{T-1} \|\boldsymbol{\mu}^{(t+1, \star)} - \boldsymbol{\mu}^{(t, \star)}\|_2,$$

where $\Xi^{(t, \star)}$ denotes the set of CE of the t -th game, and (ii) the variation in the players’ utilities $\mathcal{V}_{\mathbf{A}}^{(T)} \leq \sum_{i=1}^n \sum_{t=1}^{T-1} \|\mathbf{A}_i^{(t+1)} - \mathbf{A}_i^{(t)}\|_2^2$; below, we denote by $\text{CEGAP}^{(t)}(\boldsymbol{\mu}^{(t)})$ the CE gap of $\boldsymbol{\mu}^{(t)} \in \Xi$ at the t -th game.

Theorem 3.12. *Suppose that each player employs OGD in (12) with a suitable learning rate. Then,*

$$\sum_{t=1}^T \left(\text{CEGAP}^{(t)}(\boldsymbol{\mu}^{(t)}) \right)^2 = O(1 + \mathcal{V}_{\text{CE}}^{(T)} + \mathcal{V}_{\mathbf{A}}^{(T)}).$$

There are further interesting implications of our framework that are worth highlighting. First, we obtain meta-learning guarantees for general games that depend on the (algorithm-independent) similarity of the correlated equilibria (Corollary A.22); that was left as an open question by Harris et al. [Har+22], where instead algorithm-dependent similarity metrics were derived. Further, by applying Corollary 3.7, we derive natural variation-dependent per-player regret bounds in general games (Corollary A.23), addressing a question left by Zhang et al. [Zha+22]; we suspect that obtaining such results—parameterized on the variation of the CE—are not possible without the presence of the additional player.

3.4 Dynamic Regret Bounds in Static Games

Finally, in this subsection we switch gears by investigating dynamic regret guarantees when learning in static games. The proofs of this subsection are included in Appendix A.4.

First, we point out that while traditional no-regret learning algorithms guarantee $O(\sqrt{T})$ external regret, instances of OMD—a generalization of OGD; see (13) in Appendix A—in fact guarantee $O(\sqrt{T})$ dynamic regret in two-player zero-sum games, which is a much stronger performance measure:

Proposition 3.13. *Suppose that both players in a (static) two-player zero-sum game employ OMD with a smooth regularizer. Then, $\text{DReg}_x^{(T)}, \text{DReg}_y^{(T)} = O(\sqrt{T})$.*

In proof, the dynamic regret for each player under OMD with a smooth regularizer can be bounded by the *first-order* path length of that player’s strategies, which in turn can be bounded by $O(\sqrt{T})$ given that the second-order path length is $O(1)$ (Theorem 3.4). In fact, Theorem 3.4 readily extends Proposition 3.13 to time-varying zero-sum games as well, implying that $\text{DReg}_x^{(T)}, \text{DReg}_y^{(T)} = O\left(\sqrt{T(1 + \mathcal{V}_{\epsilon\text{-NE}}^{(T)} + \mathcal{V}_{\mathbf{A}}^{(T)})}\right)$.

A question that arises from Proposition 3.13 is whether the $O(\sqrt{T})$ guarantee for dynamic regret of OMD can be improved in the online learning setting. Below, we point out a significant improvement to $O(\log T)$, but under a stronger two-point feedback model;⁴ namely, we posit that in every round each player can select an additional auxiliary strategy, and each player then gets to additionally observe the utility corresponding to the auxiliary strategies. Notably, this is akin to how the *extra-gradient method* works [Hsi+19] (also cf. [RS13, Section 4.2] for multi-point feedback models in the bandit setting).

Observation 3.14. *Under two-point feedback, there exist learning algorithms that guarantee $\text{DReg}_x^{(T)}, \text{DReg}_y^{(T)} = O(\log T)$ in two-player zero-sum games.*

In particular, it suffices for each player to employ OMD, but with the twist that the first strategy in each round is the *time-average* of OMD; the auxiliary strategy is the standard output of OMD. Then, the dynamic regret of each player will grow as $O\left(\sum_{t=1}^T \frac{1}{t}\right) = O(\log T)$ since the duality gap of the average strategies is decreasing with a rate of T^{-1} [RS13]. It is an interesting question whether Observation 3.14 can be improved to $O(1)$.

General-sum games In contrast, no (efficient) sublinear dynamic-regret guarantees are possible in general games:

Proposition 3.15. *Unless $\text{PPAD} \subseteq \text{P}$, any polynomial-time algorithm incurs $\sum_{i=1}^n \text{DReg}_i^{(T)} = \Omega(T)$, even if $n = 2$, where $\Omega(\cdot)$ here hides polynomial factors.*

Indeed, this follows since computing a Nash equilibrium to $(1/\text{poly})$ accuracy in two-player games is PPAD-hard [CDT09]. In fact, Proposition 3.15 applies beyond the online learning setting. This motivates considering a relaxation of dynamic regret, wherein the sequence of comparators is subject to the constraint $\sum_{t=1}^{T-1} \mathbb{1}\{\mathbf{x}^{(t+1,*)} \neq \mathbf{x}^{(t,*)}\} \leq K - 1$, for some parameter $K \in \mathbb{N}$; this will be referred to as $K\text{-DReg}_x^{(T)}$. Naturally, external regret coincides with $K\text{-DReg}_x^{(T)}$ under $K = 1$. In this context, we employ Lemma 3.1 to bound $K\text{-DReg}_x^{(T)}$ under OGD:

Theorem 3.16 (Detailed version in Theorem A.24). *Suppose that all n players employ OGD in an L -smooth game. Then, for any $K \in \mathbb{N}$,*

1. $\sum_{i=1}^n K\text{-DReg}_i^{(T)} = O(K\sqrt{n}L)$;
2. $K\text{-DReg}_i^{(T)} = O(K^{3/4}T^{1/4}n^{1/4}\sqrt{L})$, for $i \in \llbracket n \rrbracket$.

One question that arises here is whether the per-player bound of $O(K^{3/4}T^{1/4})$ (Item 2) can be improved to $\tilde{O}(K)$, where $\tilde{O}(\cdot)$ hides logarithmic factors. The main challenge is that, even for

⁴The same asymptotic bound was independently established by Cai and Zheng [CZ23] but in the standard feedback model.

$K = 1$, all known methods that obtain $\tilde{O}(1)$ [DFG21; PSS21; Ana+22a; Far+22] rely on non-smooth regularizers that violate the preconditions of Lemma A.1—our dynamic RVU bound that generalizes Lemma 3.1 beyond (squared) Euclidean regularization. It would also be interesting to give a natural game-theoretic interpretation to the limit point of no-regret learners with K -DReg = $o(T)$, even for a fixed $K \in \mathbb{N}$; for $K = 1$, it corresponds to the fundamental coarse correlated equilibrium. At a superficial level, it seems to be related to the variant considered by Harrow, Natarajan, and Wu [HNW16].

4 Conclusions and Future Work

In this paper, we developed a framework for characterizing iterate-convergence of no-regret learning algorithms—primarily optimistic gradient descent (OGD)—in time-varying games. There are many promising avenues for future research. Besides closing the obvious gaps we highlighted in Section 3.4, it is important to characterize the behavior of no-regret learning algorithms in other fundamental multiagent settings, such as Stackelberg (security) games [Bal+15]. Moreover, our results operate in the full-feedback model where each player receives feedback on all possible actions. Extending the scope of our framework to capture partial-feedback models as well is another interesting direction for future work.

Acknowledgements

We are grateful to Vince Conitzer and Caspar Oesterheld for helpful feedback. We also thank anonymous referees for emphasizing a number of similarities with the work of Zhang et al. [Zha+22] that we previously missed. This material is based on work supported by the National Science Foundation under grants IIS-1901403 and CCF-1733556 and by the ARO under award W911NF2210266.

References

- [Ana+22a] Ioannis Anagnostides, Gabriele Farina, Christian Kroer, Chung-Wei Lee, Haipeng Luo, and Tuomas Sandholm. “Uncoupled Learning Dynamics with $O(\log T)$ Swap Regret in Multiplayer Games”. In: *NeurIPS 2022*. 2022.
- [Ana+22b] Ioannis Anagnostides, Ioannis Panageas, Gabriele Farina, and Tuomas Sandholm. “On Last-Iterate Convergence Beyond Zero-Sum Games”. In: *International Conference on Machine Learning, ICML 2022*. Vol. 162. Proceedings of Machine Learning Research. PMLR, 2022, pp. 536–581.
- [Aum74] Robert Aumann. “Subjectivity and Correlation in Randomized Strategies”. In: *Journal of Mathematical Economics* 1 (1974), pp. 67–96.
- [Azi+20] Wäiss Azizian, Ioannis Mitliagkas, Simon Lacoste-Julien, and Gauthier Gidel. “A Tight and Unified Analysis of Gradient-Based Methods for a Whole Spectrum of Differentiable Games”. In: *The 23rd International Conference on Artificial Intelligence and Statistics, AISTATS 2020*. Vol. 108. Proceedings of Machine Learning Research. PMLR, 2020, pp. 2863–2873.

- [Bal+15] Maria-Florina Balcan, Avrim Blum, Nika Haghtalab, and Ariel D. Procaccia. “Commitment Without Regrets: Online Learning in Stackelberg Security Games”. In: *Proceedings of the Sixteenth ACM Conference on Economics and Computation, EC '15*. ACM, 2015, pp. 61–78.
- [BMW21] Heinz H. Bauschke, Walaa M. Moursi, and Xianfu Wang. “Generalized monotone operators and their averaged resolvents”. In: *Math. Program.* 189.1 (2021), pp. 55–74.
- [Bow+15] Michael Bowling, Neil Burch, Michael Johanson, and Oskari Tammelin. “Heads-up Limit Hold’em Poker is Solved”. In: *Science* 347.6218 (Jan. 2015).
- [BS16] Noam Brown and Tuomas Sandholm. “Strategy-Based Warm Starting for Regret Minimization in Games”. In: *AAAI Conference on Artificial Intelligence (AAAI)*. 2016.
- [BS17] Noam Brown and Tuomas Sandholm. “Superhuman AI for heads-up no-limit poker: Libratus beats top professionals”. In: *Science* (Dec. 2017), eaao1733.
- [BS19a] Noam Brown and Tuomas Sandholm. “Solving imperfect-information games via discounted regret minimization”. In: *AAAI Conference on Artificial Intelligence (AAAI)*. 2019.
- [BS19b] Noam Brown and Tuomas Sandholm. “Superhuman AI for multiplayer poker”. In: *Science* 365.6456 (2019), pp. 885–890.
- [Car+19] Adrian Rivera Cardoso, Jacob D. Abernethy, He Wang, and Huan Xu. “Competing Against Nash Equilibria in Adversarially Changing Zero-Sum Games”. In: *Proceedings of the 36th International Conference on Machine Learning, ICML 2019*. Vol. 97. Proceedings of Machine Learning Research. PMLR, 2019, pp. 921–930.
- [CDT09] Xi Chen, Xiaotie Deng, and Shang-Hua Teng. “Settling the Complexity of Computing Two-Player Nash Equilibria”. In: *Journal of the ACM* (2009).
- [Ces+12] Nicolò Cesa-Bianchi, Pierre Gaillard, Gábor Lugosi, and Gilles Stoltz. “Mirror Descent Meets Fixed Share (and feels no regret)”. In: *Advances in Neural Information Processing Systems 25: 26th Annual Conference on Neural Information Processing Systems 2012*. 2012, pp. 989–997.
- [Chi+12] Chao-Kai Chiang, Tianbao Yang, Chia-Jung Lee, Mehrdad Mahdavi, Chi-Jen Lu, Rong Jin, and Shenghuo Zhu. “Online optimization with gradual variations”. In: *Conference on Learning Theory*. 2012, pp. 6–1.
- [COZ22] Yang Cai, Argyris Oikonomou, and Weiqiang Zheng. “Tight Last-Iterate Convergence of the Extragradient Method for Constrained Monotone Variational Inequalities”. In: *CoRR* abs/2204.09228 (2022).
- [CP04] Patrick L. Combettes and Teemu Pennanen. “Proximal Methods for Cohypomonotone Operators”. In: *SIAM J. Control. Optim.* (2004), pp. 731–742.
- [CZ22] Yang Cai and Weiqiang Zheng. “Accelerated Single-Call Methods for Constrained Min-Max Optimization”. In: *CoRR* abs/2210.03096 (2022).
- [CZ23] Yang Cai and Weiqiang Zheng. “Doubly Optimal No-Regret Learning in Monotone Games”. In: *CoRR* abs/2301.13120 (2023).

- [Das+18] Constantinos Daskalakis, Andrew Ilyas, Vasilis Syrgkanis, and Haoyang Zeng. “Training GANs with Optimism”. In: *6th International Conference on Learning Representations, ICLR 2018*. OpenReview.net, 2018.
- [Das22] Constantinos Daskalakis. *Non-Concave Games: A Challenge for Game Theory’s Next 100 Years*. 2022.
- [DDJ21] Jelena Diakonikolas, Constantinos Daskalakis, and Michael I. Jordan. “Efficient Methods for Structured Nonconvex-Nonconcave Min-Max Optimization”. In: *The 24th International Conference on Artificial Intelligence and Statistics, AISTATS 2021*. Vol. 130. Proceedings of Machine Learning Research. PMLR, 2021, pp. 2746–2754.
- [DDK11] Constantinos Daskalakis, Alan Deckelbaum, and Anthony Kim. “Near-optimal no-regret algorithms for zero-sum games”. In: *Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*. 2011.
- [DFG21] Constantinos Daskalakis, Maxwell Fishelson, and Noah Golowich. “Near-Optimal No-Regret Learning in General Games”. In: *Advances in Neural Information Processing Systems 34: Annual Conference on Neural Information Processing Systems 2021, NeurIPS 2021*. 2021, pp. 27604–27616.
- [DGP08] Constantinos Daskalakis, Paul Goldberg, and Christos Papadimitriou. “The Complexity of Computing a Nash Equilibrium”. In: *SIAM Journal on Computing* (2008).
- [DL15] Cong D. Dang and Guanghui Lan. “On the convergence properties of non-Euclidean extragradient methods for variational inequalities with generalized monotone operators”. In: *Comput. Optim. Appl.* (2015), pp. 277–310.
- [DMZ21] Yuan Deng, Vahab Mirrokni, and Song Zuo. “Non-Clairvoyant Dynamic Mechanism Design with Budget Constraints and Beyond”. In: *Proceedings of the 22nd ACM Conference on Economics and Computation*. EC ’21. New York, NY, USA: Association for Computing Machinery, 2021, p. 369.
- [Duv+22] Benoit Duvocelle, Panayotis Mertikopoulos, Mathias Staudigl, and Dries Vermeulen. “Multiagent online learning in time-varying games”. In: *Mathematics of Operations Research* (2022).
- [Far+22] Gabriele Farina, Ioannis Anagnostides, Haipeng Luo, Chung-Wei Lee, Christian Kroer, and Tuomas Sandholm. “Near-Optimal No-Regret Learning for General Convex Games”. In: *NeurIPS 2022*. 2022.
- [FKS21] Gabriele Farina, Christian Kroer, and Tuomas Sandholm. “Faster Game Solving via Predictive Blackwell Approachability: Connecting Regret Matching and Mirror Descent”. In: *Proceedings of the AAAI Conference on Artificial Intelligence*. 2021.
- [FS99] Yoav Freund and Robert Schapire. “Adaptive game playing using multiplicative weights”. In: *Games and Economic Behavior* 29 (1999), pp. 79–103.
- [FV97] Dean Foster and Rakesh Vohra. “Calibrated Learning and Correlated Equilibrium”. In: *Games and Economic Behavior* 21 (1997), pp. 40–55.
- [Gar17] Daniel F. Garrett. “Dynamic mechanism design: Dynamic arrivals and changing values”. In: *Games and Economic Behavior* 104 (2017), pp. 595–612.

- [GPD20] Noah Golowich, Sarath Pattathil, and Constantinos Daskalakis. “Tight last-iterate convergence rates for no-regret learning in multi-player games”. In: *Advances in Neural Information Processing Systems 33: Annual Conference on Neural Information Processing Systems 2020, NeurIPS 2020*. 2020.
- [GTG22] Eduard Gorbunov, Adrien Taylor, and Gauthier Gidel. *Last-Iterate Convergence of Optimistic Gradient Method for Monotone Variational Inequalities*. 2022.
- [GVM21] Angeliki Giannou, Emmanouil-Vasileios Vlatakis-Gkaragkounis, and Panayotis Mertikopoulos. “Survival of the strictest: Stable and unstable equilibria under regularized learning with partial information”. In: *Conference on Learning Theory, COLT 2021*. Vol. 134. Proceedings of Machine Learning Research. PMLR, 2021, pp. 2147–2148.
- [Har+22] Keegan Harris, Ioannis Anagnostides, Gabriele Farina, Mikhail Khodak, Zhiwei Steven Wu, and Tuomas Sandholm. “Meta-Learning in Games”. In: *CoRR* abs/2209.14110 (2022).
- [HM00] Sergiu Hart and Andreu Mas-Colell. “A Simple Adaptive Procedure Leading to Correlated Equilibrium”. In: *Econometrica* 68 (2000), pp. 1127–1150.
- [HNW16] Aram W. Harrow, Anand Natarajan, and Xiaodi Wu. “Tight SoS-Degree Bounds for Approximate Nash Equilibria”. In: *31st Conference on Computational Complexity, CCC 2016*. Ed. by Ran Raz. Vol. 50. LIPIcs. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2016, 22:1–22:25.
- [HS09] Elad Hazan and C. Seshadhri. “Efficient Learning Algorithms for Changing Environments”. In: *Proceedings of the 26th Annual International Conference on Machine Learning*. ICML ’09. Association for Computing Machinery, 2009, pp. 393–400.
- [HS89] Sergiu Hart and David Schmeidler. “Existence of Correlated Equilibria”. In: *Mathematics of Operations Research* 14.1 (1989), pp. 18–25.
- [Hsi+19] Yu-Guan Hsieh, Franck Iutzeler, Jérôme Malick, and Panayotis Mertikopoulos. “On the convergence of single-call stochastic extra-gradient methods”. In: *Advances in Neural Information Processing Systems 32: Annual Conference on Neural Information Processing Systems 2019, NeurIPS 2019*. 2019, pp. 6936–6946.
- [Jad+15] Ali Jadbabaie, Alexander Rakhlin, Shahin Shahrampour, and Karthik Sridharan. “Online Optimization : Competing with Dynamic Comparators”. In: *Proceedings of the Eighteenth International Conference on Artificial Intelligence and Statistics*. 2015, pp. 398–406.
- [Kal+21] Deepak S. Kalhan, Amrit Singh Bedi, Alec Koppel, Ketan Rajawat, Hamed Hassani, Abhishek K. Gupta, and Adrish Banerjee. “Dynamic Online Learning via Frank-Wolfe Algorithm”. In: *IEEE Trans. Signal Process.* 69 (2021), pp. 932–947.
- [LST16] Thodoris Lykouris, Vasilis Syrgkanis, and Éva Tardos. “Learning and Efficiency in Games with Dynamic Population”. In: *Proceedings of the Twenty-Seventh Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2016*. SIAM, 2016, pp. 120–129.
- [MO11] Ishai Menache and Asuman E. Ozdaglar. *Network Games: Theory, Models, and Dynamics*. Synthesis Lectures on Communication Networks. Morgan & Claypool Publishers, 2011.

- [Mok+16] Aryan Mokhtari, Shahin Shahrampour, Ali Jadbabaie, and Alejandro Ribeiro. “Online optimization in dynamic environments: Improved regret rates for strongly convex problems”. In: *55th IEEE Conference on Decision and Control, CDC 2016*. IEEE, 2016, pp. 7195–7201.
- [MPP18] Panayotis Mertikopoulos, Christos H. Papadimitriou, and Georgios Piliouras. “Cycles in Adversarial Regularized Learning”. In: *Proceedings of the Twenty-Ninth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2018*. SIAM, 2018, pp. 2703–2717.
- [MRS20] Eric Mazumdar, Lillian J. Ratliff, and S. Shankar Sastry. “On Gradient-Based Learning in Continuous Games”. In: *SIAM J. Math. Data Sci.* (2020), pp. 103–131.
- [MS21] Panayotis Mertikopoulos and Mathias Staudigl. “Equilibrium Tracking and Convergence in Dynamic Games”. In: *2021 60th IEEE Conference on Decision and Control (CDC)*. IEEE, 2021, pp. 930–935.
- [MV21] Oren Mangoubi and Nisheeth K. Vishnoi. “Greedy adversarial equilibrium: an efficient alternative to nonconvex-nonconcave min-max optimization”. In: *STOC ’21: 53rd Annual ACM SIGACT Symposium on Theory of Computing, 2021*. ACM, 2021, pp. 896–909.
- [MV78] H. Moulin and J.-P. Vial. “Strategically zero-sum games: The class of games whose completely mixed equilibria cannot be improved upon”. In: *International Journal of Game Theory* 7.3-4 (1978), pp. 201–221.
- [Nas50] John Nash. “Equilibrium points in N-person games”. In: *Proceedings of the National Academy of Sciences* 36 (1950), pp. 48–49.
- [Nou+19] Maher Nouiehed, Maziar Sanjabi, Tianjian Huang, Jason D. Lee, and Meisam Razaviyayn. “Solving a Class of Non-Convex Min-Max Games Using Iterative First Order Methods”. In: *Advances in Neural Information Processing Systems 32: Annual Conference on Neural Information Processing Systems 2019*. 2019, pp. 14905–14916.
- [Pap+22] Christos H. Papadimitriou, George Pierrakos, Alexandros Psomas, and Aviad Rubinfeld. “On the complexity of dynamic mechanism design”. In: *Games and Economic Behavior* 134 (2022), pp. 399–427.
- [PKB22] Jorge I. Poveda, Miroslav Krstic, and Tamer Basar. “Fixed-Time Seeking and Tracking of Time-Varying Nash Equilibria in Noncooperative Games”. In: *American Control Conference, ACC 2022, Atlanta, GA, USA, June 8-10, 2022*. IEEE, 2022, pp. 794–799.
- [PSS21] Georgios Piliouras, Ryann Sim, and Stratis Skoulakis. “Optimal No-Regret Learning in General Games: Bounded Regret with Unbounded Step-Sizes via Clairvoyant MWU”. In: *arXiv preprint arXiv:2111.14737* (2021).
- [RG22] Aitazaz Ali Raja and Sergio Grammatico. “Payoff Distribution in Robust Coalitional Games on Time-Varying Networks”. In: *IEEE Trans. Control. Netw. Syst.* 9.1 (2022), pp. 511–520.
- [RJW21] Jad Rahme, Samy Jelassi, and S. Matthew Weinberg. “Auction Learning as a Two-Player Game”. In: *9th International Conference on Learning Representations, ICLR 2021*. 2021.

- [RS13] Alexander Rakhlin and Karthik Sridharan. “Optimization, learning, and games with predictable sequences”. In: *Advances in Neural Information Processing Systems*. 2013, pp. 3066–3074.
- [SAF02] Yuzuru Sato, Eizo Akiyama, and J. Doyne Farmer. “Chaos in learning a simple two-person game”. In: *Proceedings of the National Academy of Sciences* 99.7 (2002), pp. 4748–4751.
- [Sha12] Shai Shalev-Shwartz. “Online Learning and Online Convex Optimization”. In: *Foundations and Trends in Machine Learning* 4 (2012).
- [Son+20] Chaobing Song, Zhengyuan Zhou, Yichao Zhou, Yong Jiang, and Yi Ma. “Optimistic Dual Extrapolation for Coherent Non-monotone Variational Inequalities”. In: *Advances in Neural Information Processing Systems 33: Annual Conference on Neural Information Processing Systems 2020*. 2020.
- [Syr+15] Vasilis Syrgkanis, Alekh Agarwal, Haipeng Luo, and Robert E Schapire. “Fast convergence of regularized learning in games”. In: *Advances in Neural Information Processing Systems*. 2015, pp. 2989–2997.
- [Van10] Ngo Van Long. *A survey of dynamic games in economics*. Vol. 1. World Scientific, 2010.
- [van91] Eric van Damme. *Stability and perfection of Nash equilibria*. Vol. 339. Springer, 1991.
- [Ven21] Xavier Venel. “Regularity of dynamic opinion games”. In: *Games and Economic Behavior* 126 (2021), pp. 305–334.
- [Vla+20] Emmanouil-Vasileios Vlatakis-Gkaragkounis, Lampros Flokas, Thanasis Liancas, Panayotis Mertikopoulos, and Georgios Piliouras. “No-Regret Learning and Mixed Nash Equilibria: They Do Not Mix”. In: *Advances in Neural Information Processing Systems 33: Annual Conference on Neural Information Processing Systems 2020, NeurIPS 2020*. 2020.
- [YH15] Maojiao Ye and Guoqiang Hu. “Distributed Seeking of Time-Varying Nash Equilibrium for Non-Cooperative Games”. In: *IEEE Trans. Autom. Control*. 60.11 (2015), pp. 3000–3005.
- [YKH20] Junchi Yang, Negar Kiyavash, and Niao He. “Global Convergence and Variance-Reduced Optimization for a Class of Nonconvex-Nonconcave Minimax Problems”. In: *CoRR* abs/2002.09621 (2020).
- [YM22] Yuepeng Yang and Cong Ma. “ $O(T^{-1})$ Convergence of Optimistic-Follow-the-Regularized-Leader in Two-Player Zero-Sum Markov Games”. In: *arXiv preprint arXiv:2209.12430* (2022).
- [Zha+17] Lijun Zhang, Tianbao Yang, Jinfeng Yi, Rong Jin, and Zhi-Hua Zhou. “Improved Dynamic Regret for Non-degenerate Functions”. In: *Advances in Neural Information Processing Systems 30: Annual Conference on Neural Information Processing Systems 2017*. 2017, pp. 732–741.
- [Zha+20] Peng Zhao, Yu-Jie Zhang, Lijun Zhang, and Zhi-Hua Zhou. “Dynamic Regret of Convex and Smooth Functions”. In: *Advances in Neural Information Processing Systems 33: Annual Conference on Neural Information Processing Systems 2020*. 2020.

- [Zha+22] Mengxiao Zhang, Peng Zhao, Haipeng Luo, and Zhi-Hua Zhou. “No-Regret Learning in Time-Varying Zero-Sum Games”. In: *International Conference on Machine Learning, ICML 2022*. Vol. 162. Proceedings of Machine Learning Research. PMLR, 2022, pp. 26772–26808.
- [Zin03] Martin Zinkevich. “Online Convex Programming and Generalized Infinitesimal Gradient Ascent”. In: *International Conference on Machine Learning (ICML)*. Washington, DC, USA, 2003, pp. 928–936.

A Omitted Proofs

In this section, we provide the proofs from Section 3.

A.1 Proofs from Section 3.1

First, we start with the proof of Lemma 3.1 from Section 3.1. Before we proceed, it will be useful to express the update rule (OGD) in the following equivalent form:

$$\begin{aligned} \mathbf{x}^{(t)} &:= \arg \max_{\mathbf{x} \in \mathcal{X}} \left\{ \Psi_x^{(t)}(\mathbf{x}) := \langle \mathbf{x}, \mathbf{m}_x^{(t)} \rangle - \frac{1}{\eta} \mathcal{B}_{\phi_x}(\mathbf{x} \parallel \hat{\mathbf{x}}^{(t)}) \right\}, \\ \hat{\mathbf{x}}^{(t+1)} &:= \arg \max_{\hat{\mathbf{x}} \in \mathcal{X}} \left\{ \hat{\Psi}_x^{(t)}(\hat{\mathbf{x}}) := \langle \hat{\mathbf{x}}, \mathbf{u}_x^{(t)} \rangle - \frac{1}{\eta} \mathcal{B}_{\phi_x}(\hat{\mathbf{x}} \parallel \hat{\mathbf{x}}^{(t)}) \right\}. \end{aligned} \quad (13)$$

Here, $\mathcal{B}_{\phi_x}(\cdot \parallel \cdot)$ denotes the *Bregman divergence* induced by the (squared) Euclidean regularizer $\phi_x : \mathbf{x} \mapsto \frac{1}{2} \|\mathbf{x}\|_2^2$; namely, $\mathcal{B}_{\phi_x}(\mathbf{x} \parallel \mathbf{x}') := \phi(\mathbf{x}) - \phi(\mathbf{x}') - \langle \nabla \phi(\mathbf{x}'), \mathbf{x} - \mathbf{x}' \rangle = \frac{1}{2} \|\mathbf{x} - \mathbf{x}'\|_2^2$, for $\mathbf{x}, \mathbf{x}' \in \mathcal{X}$. The update rule (13) for general Bregman divergences will be referred to as *optimistic mirror descent (OMD)*.

A.1.1 Dynamic RVU Bounds

We now show Lemma 3.1, the statement of which is recalled below for the convenience of the reader. Then, in Lemma A.1 we provide an extension of Lemma 3.1 to a broader class of regularizers.

Lemma 3.1 (RVU bound for dynamic regret). Consider any sequence of utilities $(\mathbf{u}_x^{(1)}, \dots, \mathbf{u}_x^{(T)})$ up to time $T \in \mathbb{N}$. The dynamic regret (1) of OGD with respect to any sequence of comparators $(\mathbf{x}^{(1,\star)}, \dots, \mathbf{x}^{(T,\star)}) \in \mathcal{X}^T$ can be bounded by

$$\frac{D_{\mathcal{X}}^2}{2\eta} + \frac{D_{\mathcal{X}}}{\eta} \sum_{t=1}^{T-1} \|\mathbf{x}^{(t+1,\star)} - \mathbf{x}^{(t,\star)}\|_2 + \eta \sum_{t=1}^T \|\mathbf{u}_x^{(t)} - \mathbf{m}_x^{(t)}\|_2^2 - \frac{1}{2\eta} \sum_{t=1}^T \left(\|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t)}\|_2^2 + \|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t+1)}\|_2^2 \right).$$

Proof. First, by $(1/\eta)$ -strong convexity of the function $\Psi_x^{(t)}$ (defined in (13)) for any time $t \in \llbracket T \rrbracket$, we have that

$$\langle \mathbf{x}^{(t)}, \mathbf{m}_x^{(t)} \rangle - \frac{1}{2\eta} \|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t)}\|_2^2 - \langle \hat{\mathbf{x}}^{(t+1)}, \mathbf{m}_x^{(t)} \rangle + \frac{1}{2\eta} \|\hat{\mathbf{x}}^{(t+1)} - \hat{\mathbf{x}}^{(t)}\|_2^2 \geq \frac{1}{2\eta} \|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t+1)}\|_2^2, \quad (14)$$

where we used [Sha12, Lemma 2.8, p. 135]. Similarly, by $(1/\eta)$ -strong convexity of the function $\hat{\Psi}_x^{(t)}$ (defined in (13)) for any time $t \in \llbracket T \rrbracket$, we have that for any comparator $\mathbf{x}^{(t,\star)} \in \mathcal{X}$,

$$\langle \hat{\mathbf{x}}^{(t+1)}, \mathbf{u}_x^{(t)} \rangle - \frac{1}{2\eta} \|\hat{\mathbf{x}}^{(t+1)} - \hat{\mathbf{x}}^{(t)}\|_2^2 - \langle \mathbf{x}^{(t,\star)}, \mathbf{u}_x^{(t)} \rangle + \frac{1}{2\eta} \|\mathbf{x}^{(t,\star)} - \hat{\mathbf{x}}^{(t)}\|_2^2 \geq \frac{1}{2\eta} \|\hat{\mathbf{x}}^{(t+1)} - \mathbf{x}^{(t,\star)}\|_2^2. \quad (15)$$

Thus, adding (14) and (15),

$$\begin{aligned} \langle \mathbf{x}^{(t,\star)} - \hat{\mathbf{x}}^{(t+1)}, \mathbf{u}_x^{(t)} \rangle + \langle \hat{\mathbf{x}}^{(t+1)} - \mathbf{x}^{(t)}, \mathbf{m}_x^{(t)} \rangle &\leq \frac{1}{2\eta} \left(\|\hat{\mathbf{x}}^{(t)} - \mathbf{x}^{(t,\star)}\|_2^2 - \|\hat{\mathbf{x}}^{(t+1)} - \mathbf{x}^{(t,\star)}\|_2^2 \right) \\ &\quad - \frac{1}{2\eta} \left(\|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t)}\|_2^2 + \|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t+1)}\|_2^2 \right). \end{aligned} \quad (16)$$

We further see that

$$\langle \mathbf{x}^{(t,\star)} - \mathbf{x}^{(t)}, \mathbf{u}_x^{(t)} \rangle = \langle \mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t+1)}, \mathbf{m}_x^{(t)} - \mathbf{u}_x^{(t)} \rangle + \langle \mathbf{x}^{(t,\star)} - \hat{\mathbf{x}}^{(t+1)}, \mathbf{u}_x^{(t)} \rangle + \langle \hat{\mathbf{x}}^{(t+1)} - \mathbf{x}^{(t)}, \mathbf{m}_x^{(t)} \rangle. \quad (17)$$

Now the first term on the right-hand side can be upper bounded using the fact that, by (14) and (15),

$$\langle \mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t+1)}, \mathbf{m}_x^{(t)} - \mathbf{u}_x^{(t)} \rangle \geq \frac{1}{\eta} \|\hat{\mathbf{x}}^{(t+1)} - \mathbf{x}^{(t)}\|_2^2 \implies \|\hat{\mathbf{x}}^{(t+1)} - \mathbf{x}^{(t)}\|_2 \leq \eta \|\mathbf{m}_x^{(t)} - \mathbf{u}_x^{(t)}\|_2,$$

by Cauchy-Schwarz, in turn implying that $\langle \mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t+1)}, \mathbf{m}_x^{(t)} - \mathbf{u}_x^{(t)} \rangle \leq \eta \|\mathbf{m}_x^{(t)} - \mathbf{u}_x^{(t)}\|_2^2$. Thus, the proof follows by combining this bound with (16) and (17), along with the fact that

$$\begin{aligned} \sum_{t=1}^T \left(\|\hat{\mathbf{x}}^{(t)} - \mathbf{x}^{(t,\star)}\|_2^2 - \|\hat{\mathbf{x}}^{(t+1)} - \mathbf{x}^{(t,\star)}\|_2^2 \right) &\leq \|\hat{\mathbf{x}}^{(1)} - \mathbf{x}^{(1,\star)}\|_2^2 \\ &+ \sum_{t=1}^{T-1} \left(\|\hat{\mathbf{x}}^{(t+1)} - \mathbf{x}^{(t+1,\star)}\|_2^2 - \|\hat{\mathbf{x}}^{(t+1)} - \mathbf{x}^{(t,\star)}\|_2^2 \right) \\ &\leq D_{\mathcal{X}}^2 + 2D_{\mathcal{X}} \sum_{t=1}^{T-1} \|\mathbf{x}^{(t+1,\star)} - \mathbf{x}^{(t,\star)}\|_2, \end{aligned}$$

where the last bound follows since

$$\begin{aligned} \|\hat{\mathbf{x}}^{(t+1)} - \mathbf{x}^{(t+1,\star)}\|_2^2 - \|\hat{\mathbf{x}}^{(t+1)} - \mathbf{x}^{(t,\star)}\|_2^2 &\leq 2D_{\mathcal{X}} \left| \|\hat{\mathbf{x}}^{(t+1)} - \mathbf{x}^{(t+1,\star)}\|_2 - \|\hat{\mathbf{x}}^{(t+1)} - \mathbf{x}^{(t,\star)}\|_2 \right| \\ &\leq 2D_{\mathcal{X}} \|\mathbf{x}^{(t+1,\star)} - \mathbf{x}^{(t,\star)}\|_2, \end{aligned}$$

where we recall that $D_{\mathcal{X}}$ denotes the ℓ_2 -diameter of \mathcal{X} . □

As an aside, we remark that assuming that $\mathbf{m}_x^{(t)} := \mathbf{0}$ and $\|\mathbf{u}_x^{(t)}\|_2 \leq 1$ for any $t \in \llbracket T \rrbracket$, Lemma 3.1 implies that dynamic regret can be upper bounded by $O\left(\sqrt{(1 + \sum_{t=1}^{T-1} \|\mathbf{x}^{(t+1,\star)} - \mathbf{x}^{(t,\star)}\|_2)T}\right)$, for *any* (bounded)—potentially adversarially selected—sequence of utilities $(\mathbf{u}_x^{(1)}, \dots, \mathbf{u}_x^{(T)})$, for $\eta := \sqrt{\frac{D_{\mathcal{X}}^2}{2T} + \frac{D_{\mathcal{X}} \sum_{t=1}^{T-1} \|\mathbf{x}^{(t+1,\star)} - \mathbf{x}^{(t,\star)}\|_2}{T}}$, which is a well-known result in online optimization [Zin03]; while that requires setting the learning rate based on the first-order variation of the (optimal) comparators, there are standard techniques that would allow bypassing that assumption.

Next, we provide an extension of Lemma 3.1 to the more general OMD algorithm under a broad class of regularizers.

Lemma A.1 (Extension of Lemma 3.1 beyond Euclidean regularization). *Consider a 1-strongly convex continuously differentiable regularizer ϕ with respect to a norm $\|\cdot\|$ such that (i) $\|\nabla\phi(\mathbf{x})\|_* \leq G$ for any \mathbf{x} , and (ii) $\mathcal{B}_{\phi_x}(\mathbf{x} \parallel \mathbf{x}') \leq L\|\mathbf{x} - \mathbf{x}'\|$ for any \mathbf{x}, \mathbf{x}' . Then, for any sequence of utilities $(\mathbf{u}_x^{(1)}, \dots, \mathbf{u}_x^{(T)})$ up to time $T \in \mathbb{N}$ the dynamic regret (1) of OMD with respect to any sequence of comparators $(\mathbf{x}^{(1,\star)}, \dots, \mathbf{x}^{(T,\star)}) \in \mathcal{X}^T$ can be bounded as*

$$\begin{aligned} \frac{\mathcal{B}_{\phi_x}(\mathbf{x}^{(1,\star)} \parallel \hat{\mathbf{x}}^{(1)})}{\eta} + \frac{L + 2G}{\eta} \sum_{t=1}^{T-1} \|\mathbf{x}^{(t+1,\star)} - \mathbf{x}^{(t,\star)}\| + \eta \sum_{t=1}^T \|\mathbf{u}_x^{(t)} - \mathbf{m}_x^{(t)}\|_*^2 \\ - \frac{1}{2\eta} \sum_{t=1}^T \left(\|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t)}\|^2 + \|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t+1)}\|^2 \right). \end{aligned}$$

The proof is analogous to that of Lemma 3.1, and relies on the well-known three-point identity for the Bregman divergence:

$$\mathcal{B}_{\phi_x}(\mathbf{x} \parallel \mathbf{x}') = \mathcal{B}_{\phi_x}(\mathbf{x} \parallel \mathbf{x}'') + \mathcal{B}_{\phi_x}(\mathbf{x}'' \parallel \mathbf{x}') - \langle \mathbf{x} - \mathbf{x}'', \nabla\phi(\mathbf{x}') - \nabla\phi(\mathbf{x}'') \rangle. \quad (18)$$

In particular, along with the assumptions of Lemma A.1 imposed on the regularizer ϕ_x , (18) implies that the term $\sum_{t=1}^{T-1} (\mathcal{B}_{\phi_x}(\mathbf{x}^{(t+1,\star)} \parallel \hat{\mathbf{x}}^{(t+1)}) - \mathcal{B}_{\phi_x}(\mathbf{x}^{(t,\star)} \parallel \hat{\mathbf{x}}^{(t+1)}))$ is equal to

$$\begin{aligned} & \sum_{t=1}^{T-1} \left(\mathcal{B}_{\phi_x}(\mathbf{x}^{(t+1,\star)} \parallel \mathbf{x}^{(t,\star)}) - \langle \mathbf{x}^{(t+1,\star)} - \mathbf{x}^{(t,\star)}, \nabla\phi(\hat{\mathbf{x}}^{(t+1)}) - \nabla\phi(\mathbf{x}^{(t,\star)}) \rangle \right) \\ & \leq (L + 2G) \sum_{t=1}^{T-1} \|\mathbf{x}^{(t+1,\star)} - \mathbf{x}^{(t,\star)}\|, \end{aligned}$$

since $\mathcal{B}_{\phi_x}(\mathbf{x}^{(t+1,\star)} \parallel \mathbf{x}^{(t,\star)}) \leq L\|\mathbf{x}^{(t+1,\star)} - \mathbf{x}^{(t,\star)}\|$ (by assumption) and

$$\langle \mathbf{x}^{(t+1,\star)} - \mathbf{x}^{(t,\star)}, \nabla\phi(\hat{\mathbf{x}}^{(t+1)}) - \nabla\phi(\mathbf{x}^{(t,\star)}) \rangle \leq \|\mathbf{x}^{(t+1,\star)} - \mathbf{x}^{(t,\star)}\| \|\nabla\phi(\hat{\mathbf{x}}^{(t+1)}) - \nabla\phi(\mathbf{x}^{(t,\star)})\|_* \quad (19)$$

$$\leq \left(\|\nabla\phi(\hat{\mathbf{x}}^{(t+1)})\|_* + \|\nabla\phi(\mathbf{x}^{(t,\star)})\|_* \right) \|\mathbf{x}^{(t+1,\star)} - \mathbf{x}^{(t,\star)}\| \quad (20)$$

$$\leq 2G\|\mathbf{x}^{(t+1,\star)} - \mathbf{x}^{(t,\star)}\|, \quad (21)$$

where (19) follows from the Cauchy-Schwarz inequality; (20) uses the triangle inequality for the dual norm $\|\cdot\|_*$; and (21) follows from the assumption of Lemma A.1 that $\|\nabla\phi(\cdot)\|_* \leq G$. The rest of the proof of Lemma A.1 is analogous to Lemma 3.1, and it is therefore omitted. An important question is whether Lemma A.1 can be extended under any regularizer; as we explain in Section 3.4, this is the main obstacle to improving Theorem 3.16.

A.1.2 Nonnegativity of Dynamic Regret

We next proceed with the proof of Property 3.2. To provide additional intuition, we first prove the following special case; the proof of Property 3.2 is then analogous.

Property A.2 (Special case of Property 3.2). *Suppose that $\mathcal{Z} \ni \mathbf{z}^{(t,\star)} = (\mathbf{x}^{(t,\star)}, \mathbf{y}^{(t,\star)})$ is a Nash equilibrium of the t -th game, for any time $t \in \llbracket T \rrbracket$. Then, for $s_x^{(T)} = (\mathbf{x}^{(t,\star)})_{1 \leq t \leq T}$ and $s_y^{(T)} = (\mathbf{y}^{(t,\star)})_{1 \leq t \leq T}$,*

$$\text{DReg}_x^{(T)}(s_x^{(T)}) + \text{DReg}_y^{(T)}(s_y^{(T)}) \geq 0.$$

Proof. Let $v^{(t)} := \langle \mathbf{x}^{(t,\star)}, \mathbf{A}^{(t)} \mathbf{y}^{(t,\star)} \rangle$ be the value of the t -th game, for some $t \in \llbracket T \rrbracket$. Then, we have that $v^{(t)} = \langle \mathbf{x}^{(t,\star)}, \mathbf{A}^{(t)} \mathbf{y}^{(t,\star)} \rangle \leq \langle \mathbf{x}, \mathbf{A}^{(t)} \mathbf{y}^{(t,\star)} \rangle$ for any $\mathbf{x} \in \mathcal{X}$, since $\mathbf{x}^{(t,\star)}$ is a best response to $\mathbf{y}^{(t,\star)}$; similarly, $v^{(t)} = \langle \mathbf{x}^{(t,\star)}, \mathbf{A}^{(t)} \mathbf{y}^{(t,\star)} \rangle \geq \langle \mathbf{x}^{(t,\star)}, \mathbf{A}^{(t)} \mathbf{y} \rangle$ for any $\mathbf{y} \in \mathcal{Y}$. Hence, $\langle \mathbf{x}^{(t)}, \mathbf{A}^{(t)} \mathbf{y}^{(t,\star)} \rangle - \langle \mathbf{x}^{(t,\star)}, \mathbf{A}^{(t)} \mathbf{y}^{(t)} \rangle \geq 0$, or equivalently, $\langle \mathbf{x}^{(t,\star)}, \mathbf{u}_x^{(t)} \rangle + \langle \mathbf{y}^{(t,\star)}, \mathbf{u}_y^{(t)} \rangle \geq 0$. But given that the game is zero-sum, it holds that $\langle \mathbf{x}^{(t)}, \mathbf{u}_x^{(t)} \rangle + \langle \mathbf{y}^{(t)}, \mathbf{u}_y^{(t)} \rangle = 0$, so the last inequality can be in turn cast as

$$\langle \mathbf{x}^{(t,\star)}, \mathbf{u}_x^{(t)} \rangle - \langle \mathbf{x}^{(t)}, \mathbf{u}_x^{(t)} \rangle + \langle \mathbf{y}^{(t,\star)}, \mathbf{u}_y^{(t)} \rangle - \langle \mathbf{y}^{(t)}, \mathbf{u}_y^{(t)} \rangle \geq 0,$$

for any $t \in \llbracket T \rrbracket$. As a result, summing over all $t \in \llbracket T \rrbracket$ we have shown that

$$\begin{aligned} & \text{DReg}_x^{(T)}(\mathbf{x}^{(1,\star)}, \dots, \mathbf{x}^{(T,\star)}) + \text{DReg}_y^{(T)}(\mathbf{y}^{(1,\star)}, \dots, \mathbf{y}^{(T,\star)}) \\ &= \sum_{t=1}^T \langle \mathbf{x}^{(t,\star)}, \mathbf{u}_x^{(t)} \rangle - \langle \mathbf{x}^{(t)}, \mathbf{u}_x^{(t)} \rangle + \langle \mathbf{y}^{(t,\star)}, \mathbf{u}_y^{(t)} \rangle - \langle \mathbf{y}^{(t)}, \mathbf{u}_y^{(t)} \rangle \geq 0. \end{aligned}$$

□

Property 3.2. *Suppose that $\mathcal{Z} \ni \mathbf{z}^{(t,\star)} = (\mathbf{x}^{(t,\star)}, \mathbf{y}^{(t,\star)})$ is an $\epsilon^{(t)}$ -approximate Nash equilibrium of the t -th game. Then, for $s_x^{(T)} = (\mathbf{x}^{(t,\star)})_{1 \leq t \leq T}$ and $s_y^{(T)} = (\mathbf{y}^{(t,\star)})_{1 \leq t \leq T}$,*

$$\text{DReg}_x^{(T)}(s_x^{(T)}) + \text{DReg}_y^{(T)}(s_y^{(T)}) \geq -2 \sum_{t=1}^T \epsilon^{(t)}.$$

Proof. Given that $(\mathbf{x}^{(t,\star)}, \mathbf{y}^{(t,\star)}) \in \mathcal{Z}$ is an $\epsilon^{(t)}$ -approximate Nash equilibrium of the t -th game, it follows that $\langle \mathbf{x}^{(t,\star)}, \mathbf{A}^{(t)} \mathbf{y}^{(t,\star)} \rangle \leq \langle \mathbf{x}^{(t)}, \mathbf{A}^{(t)} \mathbf{y}^{(t,\star)} \rangle + \epsilon_x^{(t)}$ and $\langle \mathbf{x}^{(t,\star)}, \mathbf{A}^{(t)} \mathbf{y}^{(t,\star)} \rangle \geq \langle \mathbf{x}^{(t,\star)}, \mathbf{A}^{(t)} \mathbf{y}^{(t)} \rangle - \epsilon_y^{(t)}$, for some $\epsilon_x^{(t)}, \epsilon_y^{(t)} \leq \epsilon^{(t)}$. Thus, we have that $\langle \mathbf{x}^{(t)}, \mathbf{A}^{(t)} \mathbf{y}^{(t,\star)} \rangle \geq \langle \mathbf{x}^{(t,\star)}, \mathbf{A}^{(t)} \mathbf{y}^{(t)} \rangle - \epsilon_x^{(t)} - \epsilon_y^{(t)}$, or equivalently, $\langle \mathbf{x}^{(t,\star)}, \mathbf{u}_x^{(t)} \rangle + \langle \mathbf{y}^{(t,\star)}, \mathbf{u}_y^{(t)} \rangle \geq -\epsilon_x^{(t)} - \epsilon_y^{(t)} \geq -2\epsilon^{(t)}$. As a result,

$$\langle \mathbf{x}^{(t,\star)}, \mathbf{u}_x^{(t)} \rangle - \langle \mathbf{x}^{(t)}, \mathbf{u}_x^{(t)} \rangle + \langle \mathbf{y}^{(t,\star)}, \mathbf{u}_y^{(t)} \rangle - \langle \mathbf{y}^{(t)}, \mathbf{u}_y^{(t)} \rangle \geq -2\epsilon^{(t)}, \quad (22)$$

for any $t \in \llbracket T \rrbracket$, and the statement follows by summing (22) over all $t \in \llbracket T \rrbracket$. □

In fact, as we show below (in Property A.3), Property A.2 is a more general consequence of the minimax theorem. In particular, for a nonlinear online learning problem, we define dynamic regret with respect to a sequence of comparators $(\mathbf{x}^{(1,\star)}, \dots, \mathbf{x}^{(T,\star)}) \in \mathcal{X}^T$ as follows:

$$\text{DReg}_x^{(T)}(\mathbf{x}^{(1,\star)}, \dots, \mathbf{x}^{(T,\star)}) := \sum_{t=1}^T \left(u_x^{(t)}(\mathbf{x}^{(t,\star)}) - u_x^{(t)}(\mathbf{x}^{(t)}) \right), \quad (23)$$

where $u_x^{(1)}, \dots, u_x^{(T)} : \mathbf{x} \mapsto \mathbb{R}$ are the continuous utility functions observed by the learner, which could be in general nonconcave, and $(\mathbf{x}^{(t)})_{1 \leq t \leq T}$ is the sequence of strategies produced by the learner; (23) generalizes the notion of dynamic regret (1) in online linear optimization, that is, when $u_x^{(t)} : \mathbf{x} \mapsto \langle \mathbf{x}, \mathbf{u}_x^{(t)} \rangle$, where $\mathbf{u}_x^{(t)} \in \mathbb{R}^{d_x}$, for any time $t \in \llbracket T \rrbracket$.

Property A.3. *Suppose that $f^{(t)} : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}$ is a continuous function such that for any $t \in \llbracket T \rrbracket$,*

$$\min_{\mathbf{x} \in \mathcal{X}} \max_{\mathbf{y} \in \mathcal{Y}} f^{(t)}(\mathbf{x}, \mathbf{y}) = \max_{\mathbf{y} \in \mathcal{Y}} \min_{\mathbf{x} \in \mathcal{X}} f^{(t)}(\mathbf{x}, \mathbf{y}).$$

Let also $\mathbf{x}^{(t,\star)} \in \arg \min_{\mathbf{x} \in \mathcal{X}} \max_{\mathbf{y} \in \mathcal{Y}} f^{(t)}(\mathbf{x}, \mathbf{y})$ and $\mathbf{y}^{(t,\star)} \in \arg \max_{\mathbf{y} \in \mathcal{Y}} \min_{\mathbf{x} \in \mathcal{X}} f^{(t)}(\mathbf{x}, \mathbf{y})$, for any $t \in \llbracket T \rrbracket$. Then, for $s_x^{(T)} = (\mathbf{x}^{(t,\star)})_{1 \leq t \leq T}$ and $s_y^{(T)} = (\mathbf{y}^{(t,\star)})_{1 \leq t \leq T}$,

$$\text{DReg}_x^{(T)}(s_x^{(T)}) + \text{DReg}_y^{(T)}(s_y^{(T)}) \geq 0.$$

Proof. By definition of dynamic regret (23), it suffices to show that $f^{(t)}(\mathbf{x}^{(t)}, \mathbf{y}^{(t,*)}) \geq f^{(t)}(\mathbf{x}^{(t,*)}, \mathbf{y}^{(t)})$, for any time $t \in \llbracket T \rrbracket$. Indeed,

$$f^{(t)}(\mathbf{x}^{(t)}, \mathbf{y}^{(t,*)}) \geq \min_{\mathbf{x} \in \mathcal{X}} f^{(t)}(\mathbf{x}, \mathbf{y}^{(t,*)}) \quad (24)$$

$$= \max_{\mathbf{y} \in \mathcal{Y}} \min_{\mathbf{x} \in \mathcal{X}} f^{(t)}(\mathbf{x}, \mathbf{y}) \quad (25)$$

$$= \min_{\mathbf{x} \in \mathcal{X}} \max_{\mathbf{y} \in \mathcal{Y}} f^{(t)}(\mathbf{x}, \mathbf{y}) \quad (26)$$

$$= \max_{\mathbf{y} \in \mathcal{Y}} f^{(t)}(\mathbf{x}^{(t,*)}, \mathbf{y}) \quad (27)$$

$$\geq f^{(t)}(\mathbf{x}^{(t,*)}, \mathbf{y}^{(t)}), \quad (28)$$

where (24) and (28) are obvious; (25) and (27) follow from the definition of $\mathbf{y}^{(t,*)} \in \mathcal{Y}$ and $\mathbf{x}^{(t,*)} \in \mathcal{X}$, respectively; and (26) holds by assumption. This concludes the proof. \square

Remark A.4 (MVI property). Property (3) can also be generalized beyond time-varying bilinear saddle-point problems to more general time-varying *variational inequality* (VI) problems as follows. Let $F^{(t)} : \mathcal{Z} \rightarrow \mathcal{Z}$ be the (single-valued) operator of the VI problem at time t . $F^{(t)}$ is said to satisfy the *MVI property* if there exists a point $\mathbf{z}^{(t,*)} \in \mathcal{Z}$ such that $\langle \mathbf{z} - \mathbf{z}^{(t,*)}, F^{(t)}(\mathbf{z}) \rangle \geq 0$ for any $\mathbf{z} \in \mathcal{Z}$. For example, in the special case of a bilinear saddle-point problem, we have that $F : \mathbf{z} := (\mathbf{x}, \mathbf{y}) \mapsto (\mathbf{A}\mathbf{y}, -\mathbf{A}^\top \mathbf{x})$, and the MVI property is satisfied by virtue of Von Neumann’s minimax theorem. It is direct to see that Property A.2 applies to any time-varying VI with respect to the sequence $(\mathbf{z}^{(t,*)})_{1 \leq t \leq T}$ as long as every operator in the sequence $(F^{(1)}, \dots, F^{(T)})$ satisfies the MVI property. (Even more broadly, it suffices if almost all operators in the sequence satisfy the MVI property—in that their fraction converges to 1 as $T \rightarrow +\infty$.) This observation enables extending Theorem 3.5 beyond time-varying bilinear saddle-point problems.

A.1.3 Variation of the Nash Equilibria

In our next example, we point out that an arbitrarily small change in the entries of the payoff matrix can lead to a substantial deviation in the Nash equilibrium.

Example A.5. Consider a 2×2 (two-player) zero-sum game, where $\mathcal{X} := \Delta^2, \mathcal{Y} := \Delta^2$, described by the payoff matrix

$$\mathbf{A} := \begin{bmatrix} 2\delta & 0 \\ 0 & \delta \end{bmatrix}, \quad (29)$$

for some $\delta > 0$. Then, it is easy to see that the unique Nash equilibrium of this game is such that $\mathbf{x}^*, \mathbf{y}^* := (\frac{1}{3}, \frac{2}{3}) \in \Delta^2$. Suppose now that the original payoff matrix (29) is perturbed to a new matrix

$$\mathbf{A}' := \begin{bmatrix} \delta & 0 \\ 0 & 2\delta \end{bmatrix}. \quad (30)$$

The new (unique) Nash equilibrium now reads $\mathbf{x}^*, \mathbf{y}^* := (\frac{2}{3}, \frac{1}{3}) \in \Delta^2$. We conclude that an arbitrarily small deviation in the entries of the payoff matrix can lead to a non-trivial change in the Nash equilibrium.

Next, we leverage the simple observation of the example above to establish Proposition 3.3, the statement of which is recalled below.

Proposition 3.3. *For any $T \geq 4$, there is a sequence of T games such that $\mathcal{V}_{NE}^{(T)} \geq \frac{T}{2}$ while $\mathcal{V}_{\epsilon-NE}^{(T)} \leq \delta$, for any $\delta > 0$.*

Proof. We consider a sequence of T games such that $\mathcal{X}, \mathcal{Y} := \Delta^2$, and

$$\mathbf{A}^{(t)} = \begin{cases} \mathbf{A} & \text{if } t \bmod 2 = 1, \\ \mathbf{A}' & \text{if } t \bmod 2 = 0, \end{cases}$$

where \mathbf{A}, \mathbf{A}' are the payoff matrices defined in (29) and (30), and are parameterized by $\delta > 0$ (Example A.5). Then, the exact Nash equilibria read

$$\mathbf{x}^{(t,*)}, \mathbf{y}^{(t,*)} = \begin{cases} \left(\frac{1}{3}, \frac{2}{3}\right) & \text{if } t \bmod 2 = 1, \\ \left(\frac{2}{3}, \frac{1}{3}\right) & \text{if } t \bmod 2 = 0. \end{cases}$$

As a result, it follows that $\mathcal{V}_{NE}^{(T)} := \sum_{t=1}^{T-1} \|\mathbf{z}^{(t+1,*)} - \mathbf{z}^{(t,*)}\|_2 = \frac{2}{3}(T-1) \geq \frac{T}{2}$, for $T \geq 4$. In contrast, it is clear that $\mathcal{V}_{\epsilon-NE}^{(T)} \leq C\delta T$, which follows by simply considering the sequence of strategies wherein both players are always selecting actions uniformly at random; we recall that $C > 0$ here is the value that parameterizes $\mathcal{V}_{\epsilon-NE}^{(T)}$. Thus, taking $\delta := \frac{\delta'}{CT}$, for some arbitrarily small $\delta' > 0$, concludes the proof. \square

A.1.4 Main Result

Next, we proceed with the proof of our main result, Theorem 3.5. The key ingredient is Theorem 3.4, which bounds the second-order path length of OGD in terms of the considered variation measures. We first give the precise statement of Theorem 3.4, and we then proceed with its proof.

Theorem A.6 (Detailed version of Theorem 3.4). *Suppose that both players employ OGD with learning rate $\eta \leq \frac{1}{4L}$ in a time-varying bilinear saddle-point problem, where $L := \max_{t \in [T]} \|\mathbf{A}^{(t)}\|_2$. Then, for any time horizon $T \in \mathbb{N}$,*

$$\sum_{t=1}^T \left(\|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t)}\|_2^2 + \|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t+1)}\|_2^2 \right) \leq 2D_Z^2 + 4\eta^2 L^2 \|\mathcal{Z}\|_2^2 + 4D_Z \mathcal{V}_{\epsilon-NE}^{(T)} + 8\eta^2 \|\mathcal{Z}\|_2^2 \mathcal{V}_{\mathbf{A}}^{(T)}.$$

Proof of Theorem 3.4. First, for any $t \geq 2$ we have that $\|\mathbf{u}_x^{(t)} - \mathbf{m}_x^{(t)}\|_2^2$ is equal to

$$\|\mathbf{A}^{(t)}\mathbf{y}^{(t)} - \mathbf{A}^{(t-1)}\mathbf{y}^{(t-1)}\|_2^2 \leq 2\|\mathbf{A}^{(t)}(\mathbf{y}^{(t)} - \mathbf{y}^{(t-1)})\|_2^2 + 2\|(\mathbf{A}^{(t)} - \mathbf{A}^{(t-1)})\mathbf{y}^{(t-1)}\|_2^2 \quad (31)$$

$$\leq 2\|\mathbf{A}^{(t)}\|_2^2 \|\mathbf{y}^{(t)} - \mathbf{y}^{(t-1)}\|_2^2 + 2\|\mathbf{A}^{(t)} - \mathbf{A}^{(t-1)}\|_2^2 \|\mathbf{y}^{(t-1)}\|_2^2 \quad (32)$$

$$\leq 2L^2 \|\mathbf{y}^{(t)} - \mathbf{y}^{(t-1)}\|_2^2 + 2\|\mathcal{Y}\|_2^2 \|\mathbf{A}^{(t)} - \mathbf{A}^{(t-1)}\|_2^2, \quad (33)$$

where (31) uses the triangle inequality for the norm $\|\cdot\|_2$ along with the inequality $2ab \leq a^2 + b^2$ for any $a, b \in \mathbb{R}$; (32) follows from the definition of the operator norm; and (33) uses the assumption that $\|\mathbf{A}^{(t)}\|_2 \leq L$ and $\|\mathbf{y}\|_2 \leq \|\mathcal{Y}\|_2$ for any $\mathbf{y} \in \mathcal{Y}$. A similar derivation shows that for $t \geq 2$,

$$\|\mathbf{u}_y^{(t)} - \mathbf{m}_y^{(t)}\|_2^2 \leq 2L^2 \|\mathbf{x}^{(t)} - \mathbf{x}^{(t-1)}\|_2^2 + 2\|\mathcal{X}\|_2^2 \|\mathbf{A}^{(t)} - \mathbf{A}^{(t-1)}\|_2^2. \quad (34)$$

Further, for $t = 1$ we have that $\|\mathbf{u}_x^{(1)} - \mathbf{m}_x^{(1)}\|_2 = \|\mathbf{u}_x^{(1)}\|_2 = \|\mathbf{A}^{(1)}\mathbf{y}^{(1)}\|_2 \leq L\|\mathcal{Y}\|_2$, and $\|\mathbf{u}_y^{(1)} - \mathbf{m}_y^{(1)}\|_2 = \|\mathbf{u}_y^{(1)}\|_2 = \|(\mathbf{A}^{(1)})^\top \mathbf{x}^{(1)}\|_2 \leq L\|\mathcal{X}\|_2$. Next, we will use the following simple corollary, which follows similarly to Lemma 3.1.

Corollary A.7. For any sequence $s_z^{(T)} := (\mathbf{z}^{(t,\star)})_{1 \leq t \leq T}$, the dynamic regret $\text{DReg}_z^{(T)}(s_z^{(T)}) := \text{DReg}_x^{(T)}(s_x^{(T)}) + \text{DReg}_y^{(T)}(s_y^{(T)})$ can be bounded by

$$\frac{D_{\mathcal{Z}}^2}{2\eta} + \frac{D_{\mathcal{Z}}}{\eta} \sum_{t=1}^{T-1} \|\mathbf{z}^{(t+1,\star)} - \mathbf{z}^{(t,\star)}\|_2 + \eta \sum_{t=1}^T \|\mathbf{u}_z^{(t)} - \mathbf{m}_z^{(t)}\|_2^2 - \frac{1}{2\eta} \sum_{t=1}^T \left(\|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t)}\|_2^2 + \|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t+1)}\|_2^2 \right),$$

where $\mathbf{m}_z^{(t)} := (\mathbf{m}_x^{(t)}, \mathbf{m}_y^{(t)})$ and $\mathbf{u}_z^{(t)} := (\mathbf{u}_x^{(t)}, \mathbf{u}_y^{(t)})$ for any $t \in \llbracket T \rrbracket$.

As a result, combining (34) and (33) with Corollary A.7 applied for the dynamic regret of both players with respect to the sequence of comparators $((\mathbf{x}^{(t,\star)}, \mathbf{y}^{(t,\star)}))_{1 \leq t \leq T}$ yields that $\text{DReg}_x^{(T)}(\mathbf{x}^{(1,\star)}, \dots, \mathbf{x}^{(T,\star)}) + \text{DReg}_y^{(T)}(\mathbf{y}^{(1,\star)}, \dots, \mathbf{y}^{(T,\star)})$ is upper bounded by

$$\begin{aligned} \frac{D_{\mathcal{Z}}^2}{2\eta} + \eta L^2 \|\mathcal{Z}\|_2^2 + \frac{D_{\mathcal{Z}}}{\eta} \sum_{t=1}^{T-1} \|\mathbf{z}^{(t+1,\star)} - \mathbf{z}^{(t,\star)}\|_2 + 2\eta \|\mathcal{Z}\|_2^2 \mathcal{V}_{\mathbf{A}}^{(T)} \\ - \frac{1}{4\eta} \sum_{t=1}^T \left(\|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t)}\|_2^2 + \|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t+1)}\|_2^2 \right), \end{aligned}$$

where we used the fact that

$$\begin{aligned} 2\eta L^2 \sum_{t=2}^T \|\mathbf{z}^{(t)} - \mathbf{z}^{(t-1)}\|_2^2 - \frac{1}{4\eta} \sum_{t=1}^T \left(\|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t)}\|_2^2 + \|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t+1)}\|_2^2 \right) \\ \leq \left(2\eta L^2 - \frac{1}{8\eta} \right) \sum_{t=2}^T \|\mathbf{z}^{(t)} - \mathbf{z}^{(t-1)}\|_2^2 \leq 0, \end{aligned}$$

for $\eta \leq \frac{1}{4L}$. Finally, using the fact that $\text{DReg}_x^{(T)}(\mathbf{x}^{(1,\star)}, \dots, \mathbf{x}^{(T,\star)}) + \text{DReg}_y^{(T)}(\mathbf{y}^{(1,\star)}, \dots, \mathbf{y}^{(T,\star)}) \geq -2 \sum_{t=1}^T \epsilon^{(t)}$ for a suitable sequence of $\epsilon^{(t)}$ -approximate Nash equilibria (Property 3.2)—one that attains the variation measure $\mathcal{V}_{\epsilon\text{-NE}}^{(T)}$ —yields that

$$0 \leq \frac{D_{\mathcal{Z}}^2}{2\eta} + \eta L^2 \|\mathcal{Z}\|_2^2 + \frac{D_{\mathcal{Z}}}{\eta} \mathcal{V}_{\epsilon\text{-NE}}^{(T)} + 2\eta \|\mathcal{Z}\|_2^2 \mathcal{V}_{\mathbf{A}}^{(T)} - \frac{1}{4\eta} \sum_{t=1}^T \left(\|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t)}\|_2^2 + \|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t+1)}\|_2^2 \right),$$

where it suffices if the parameter C of $\mathcal{V}_{\epsilon\text{-NE}}^{(T)}$ is such that $2 \leq \frac{D_{\mathcal{Z}}}{\eta} C$. Thus, rearranging the last displayed inequality concludes the proof. \square

Next, we refine this theorem in time-varying games in which the deviation of the payoff matrices is bounded by the deviation of the players' strategies, in the following formal sense.

Corollary A.8. Suppose that both players employ OGD with learning rate $\eta \leq \min \left\{ \frac{1}{4L}, \frac{1}{8W\|\mathcal{Z}\|} \right\}$ in a time-varying bilinear saddle-point problem, where $L := \max_{t \in \llbracket T \rrbracket} \|\mathbf{A}^{(t)}\|_2$ and $\mathcal{V}_{\mathbf{A}}^{(T)} \leq W^2 \sum_{t=1}^{T-1} \|\mathbf{z}^{(t+1)} - \mathbf{z}^{(t)}\|_2^2$, for some parameter $W \in \mathbb{R}_{>0}$. Then, for any time horizon $T \in \mathbb{N}$,

$$\sum_{t=1}^T \left(\|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t)}\|_2^2 + \|\hat{\mathbf{z}}^{(t)} - \hat{\mathbf{z}}^{(t+1)}\|_2^2 \right) \leq 4D_{\mathcal{Z}}^2 + 8\eta^2 L^2 \|\mathcal{Z}\|_2^2 + 8D_{\mathcal{Z}} \mathcal{V}_{NE}^{(T)},$$

where $\mathcal{V}_{NE}^{(T)}$ is defined in (4).

Proof. Following the proof of Theorem 3.4, we have that for any $\eta \leq \frac{1}{4L}$,

$$0 \leq \frac{D_{\mathcal{Z}}^2}{2\eta} + \eta L^2 \|\mathcal{Z}\|_2^2 + \frac{D_{\mathcal{Z}}}{\eta} \mathcal{V}_{\text{NE}}^{(T)} + 2\eta \|\mathcal{Z}\|_2^2 \mathcal{V}_{\mathbf{A}}^{(T)} - \frac{1}{4\eta} \sum_{t=1}^T \left(\|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t)}\|_2^2 + \|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t+1)}\|_2^2 \right).$$

Further, for $\eta \leq \frac{1}{8W\|\mathcal{Z}\|_2}$,

$$\begin{aligned} 2\eta \|\mathcal{Z}\|_2^2 \mathcal{V}_{\mathbf{A}}^{(T)} - \frac{1}{8\eta} \sum_{t=1}^T \left(\|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t)}\|_2^2 + \|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t+1)}\|_2^2 \right) \\ \leq \left(2\eta \|\mathcal{Z}\|_2^2 W^2 - \frac{1}{16\eta} \right) \sum_{t=1}^{T-1} \|\mathbf{z}^{(t+1)} - \mathbf{z}^{(t)}\|_2^2 \leq 0. \end{aligned}$$

Thus, we have shown that

$$0 \leq \frac{D_{\mathcal{Z}}^2}{2\eta} + \eta L^2 \|\mathcal{Z}\|_2^2 + \frac{D_{\mathcal{Z}}}{\eta} \mathcal{V}_{\text{NE}}^{(T)} - \frac{1}{8\eta} \sum_{t=1}^T \left(\|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t)}\|_2^2 + \|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t+1)}\|_2^2 \right),$$

and rearranging concludes the proof. \square

Thus, in such time-varying games it is the first-order variation term, $\mathcal{V}_{\text{NE}}^{(T)}$, that will drive our convergence bounds.

Now before proving Theorem 3.5, we state the connection between the equilibrium gap and the deviation of the players' strategies ($\|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t)}\|_2 + \|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t+1)}\|_2$). In particular, the following claim can be extracted by [Ana+22b, Claim A.14]. (We caution that we use a slightly different indexing for the secondary sequence $(\hat{\mathbf{x}}_i^{(t)})$ in the definition of OMD (13) compared to [Ana+22b].)

Claim A.9. *Suppose that the sequences $(\mathbf{x}_i^{(t)})_{1 \leq t \leq T}$ and $(\hat{\mathbf{x}}_i^{(t)})_{1 \leq t \leq T+1}$ are produced by OMD under a G -smooth regularizer 1-strongly convex with respect to a norm $\|\cdot\|$. Then, for any time $t \in \llbracket T \rrbracket$ and any $\mathbf{x}_i \in \mathcal{X}_i$,*

$$\langle \mathbf{x}_i^{(t)}, \mathbf{u}_i^{(t)} \rangle \geq \langle \mathbf{x}_i, \mathbf{u}_i^{(t)} \rangle - \frac{G}{\eta} \|\hat{\mathbf{x}}_i^{(t+1)} - \hat{\mathbf{x}}_i^{(t)}\| - \|\mathbf{u}_i^{(t)}\|_* \|\mathbf{x}_i^{(t)} - \hat{\mathbf{x}}_i^{(t+1)}\|.$$

We are now ready to prove Theorem 3.5, the precise version of which is stated below.

Theorem A.10 (Detailed version of Theorem 3.5). *Suppose that both players employ OGD with learning rate $\eta = \frac{1}{4L}$ in a time-varying bilinear saddle-point problem, where $L := \max_{t \in \llbracket T \rrbracket} \|\mathbf{A}^{(t)}\|_2$. Then,*

$$\sum_{t=1}^T \left(\text{EQGAP}^{(t)}(\mathbf{z}^{(t)}) \right)^2 \leq 2L^2 (4 + \|\mathcal{Z}\|_2)^2 \left(2D_{\mathcal{Z}}^2 + 4\eta^2 L^2 \|\mathcal{Z}\|_2^2 + 4D_{\mathcal{Z}} \mathcal{V}_{\epsilon\text{-NE}}^{(T)} + 8\eta^2 \|\mathcal{Z}\|_2^2 \mathcal{V}_{\mathbf{A}}^{(T)} \right),$$

where $(\mathbf{z}^{(t)})_{1 \leq t \leq T}$ is the sequence of joint strategy profiles produced by OGD.

Proof. Let us first fix a time $t \in \llbracket T \rrbracket$. For convenience, we denote by $\text{BR}_x^{(t)}(\mathbf{x}^{(t)}) := \max_{\mathbf{x} \in \mathcal{X}} \langle \mathbf{x}, \mathbf{u}_x^{(t)} \rangle - \langle \mathbf{x}^{(t)}, \mathbf{u}_x^{(t)} \rangle$, the best response gap of Player's x strategy $\mathbf{x}^{(t)} \in \mathcal{X}$, and similarly for $\text{BR}_y^{(t)}(\mathbf{y}^{(t)})$. By definition, it holds that $\text{EQGAP}^{(t)} := \max\{\text{BR}_x^{(t)}(\mathbf{x}^{(t)}), \text{BR}_y^{(t)}(\mathbf{y}^{(t)})\}$. By Claim A.9, we have that

$$\text{BR}_x^{(t)}(\mathbf{x}^{(t)}) \leq \frac{1}{\eta} \|\hat{\mathbf{x}}^{(t+1)} - \hat{\mathbf{x}}^{(t)}\|_2 + \|\mathbf{u}_x^{(t)}\|_2 \|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t+1)}\|_2 \quad (35)$$

$$\leq 4L \|\hat{\mathbf{x}}^{(t+1)} - \hat{\mathbf{x}}^{(t)}\|_2 + L \|\mathcal{Y}\|_2 \|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t+1)}\|_2 \quad (36)$$

$$\leq L(4 + \|\mathcal{Z}\|_2) \left(\|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t)}\|_2 + \|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t+1)}\|_2 \right), \quad (37)$$

where (35) follows from Claim A.9 for $G = 1$ (since the squared Euclidean regularizer $\phi_x : \mathbf{x} \mapsto \frac{1}{2} \|\mathbf{x}\|_2^2$ is 1-smooth; (36) uses the fact that $\eta := \frac{1}{4L}$ and $\|\mathbf{u}_x^{(t)}\|_2 = \|\mathbf{A}^{(t)} \mathbf{y}^{(t)}\|_2 \leq L \|\mathcal{Y}\|$; and (37) follows from the triangle inequality. A similar derivation shows that

$$\text{BR}_y^{(t)}(\mathbf{y}^{(t)}) \leq L(4 + \|\mathcal{Z}\|) \left(\|\mathbf{y}^{(t)} - \hat{\mathbf{y}}^{(t)}\|_2 + \|\mathbf{y}^{(t)} - \hat{\mathbf{y}}^{(t+1)}\|_2 \right). \quad (38)$$

Thus,

$$\begin{aligned} \sum_{t=1}^T \left(\text{EQGAP}^{(t)}(\mathbf{z}^{(t)}) \right)^2 &= \sum_{t=1}^T \left(\max\{\text{BR}_x^{(t)}(\mathbf{x}^{(t)}), \text{BR}_y^{(t)}(\mathbf{y}^{(t)})\} \right)^2 \\ &\leq \sum_{t=1}^T \left(\left(\text{BR}_x^{(t)}(\mathbf{x}^{(t)}) \right)^2 + \left(\text{BR}_y^{(t)}(\mathbf{y}^{(t)}) \right)^2 \right) \\ &\leq 2L^2(4 + \|\mathcal{Z}\|_2)^2 \sum_{t=1}^T \left(\|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t)}\|_2^2 + \|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t+1)}\|_2^2 \right), \end{aligned} \quad (39)$$

where the last bound uses (37) and (38). Combining (39) with Theorem A.6 concludes the proof. \square

A.1.5 Variation-Dependent Regret Bounds

Here we state an important implication of Theorem 3.5 for deriving variation-dependent regret bounds in time-varying bilinear saddle-point problems; cf. [Zha+22].

Corollary A.11 (Detailed version of Corollary 3.7). *In the setup of Theorem 3.4, it holds that*

$$\text{Reg}_x^{(T)} \leq \frac{D_{\mathcal{X}}^2}{\eta} + 8\eta L^2 D_{\mathcal{Z}}^2 + \eta L^2 \|\mathcal{Y}\|_2^2 + 16\eta^3 L^4 \|\mathcal{Z}\|_2^2 + 16\eta L^2 D_{\mathcal{Z}} \mathcal{V}_{NE}^{(T)} + (2\eta \|\mathcal{Y}\|_2^2 + 32\eta^3 L^2 \|\mathcal{Z}\|_2^2) \mathcal{V}_{\mathbf{A}}^{(T)},$$

and

$$\text{Reg}_y^{(T)} \leq \frac{D_{\mathcal{Y}}^2}{\eta} + 8\eta L^2 D_{\mathcal{Z}}^2 + \eta L^2 \|\mathcal{X}\|_2^2 + 16\eta^3 L^4 \|\mathcal{Z}\|_2^2 + 16\eta L^2 D_{\mathcal{Z}} \mathcal{V}_{NE}^{(T)} + (2\eta \|\mathcal{X}\|_2^2 + 32\eta^3 L^2 \|\mathcal{Z}\|_2^2) \mathcal{V}_{\mathbf{A}}^{(T)}.$$

Proof. First, applying Lemma 3.1 under $\mathbf{x}^{(1,\star)} = \dots = \mathbf{x}^{(T,\star)}$, we have

$$\text{Reg}_x^{(T)} \leq \frac{D_{\mathcal{X}}^2}{\eta} + \eta L^2 \|\mathcal{Y}\|_2^2 + 2\eta L^2 \sum_{t=2}^T \|\mathbf{y}^{(t)} - \mathbf{y}^{(t-1)}\|_2^2 + 2\eta \|\mathcal{Y}\|_2^2 \sum_{t=2}^T \|\mathbf{A}^{(t)} - \mathbf{A}^{(t-1)}\|_2^2, \quad (40)$$

and similarly,

$$\text{Reg}_y^{(T)} \leq \frac{D_y^2}{\eta} + \eta L^2 \|\mathcal{X}\|_2^2 + 2\eta L^2 \sum_{t=2}^T \|\mathbf{x}^{(t)} - \mathbf{x}^{(t-1)}\|_2^2 + 2\eta \|\mathcal{X}\|_2^2 \sum_{t=2}^T \|\mathbf{A}^{(t)} - \mathbf{A}^{(t-1)}\|_2^2. \quad (41)$$

Now, by Theorem 3.4 we have

$$\sum_{t=1}^T \left(\|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t)}\|_2^2 + \|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t+1)}\|_2^2 \right) \leq 2D_Z^2 + 4\eta^2 L^2 \|\mathcal{Z}\|_2^2 + 4D_Z \mathcal{V}_{\text{NE}}^{(T)} + 8\eta^2 \|\mathcal{Z}\|_2^2 \mathcal{V}_{\mathbf{A}}^{(T)}. \quad (42)$$

Further,

$$\begin{aligned} \sum_{t=1}^T \left(\|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t)}\|_2^2 + \|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t+1)}\|_2^2 \right) &\geq \sum_{t=1}^T \left(\|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t)}\|_2^2 + \|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t+1)}\|_2^2 \right) \\ &\geq \frac{1}{2} \sum_{t=2}^T \|\mathbf{x}^{(t)} - \mathbf{x}^{(t-1)}\|_2^2. \end{aligned}$$

Combining this bound with (42) and (41) gives the claimed regret bound on $\text{Reg}_y^{(T)}$, and a similar derivation also gives the claimed bound on $\text{Reg}_x^{(T)}$. \square

A.1.6 Meta-Learning

We next provide the implication of Theorem 3.5 in the meta-learning setting. We first make a remark regarding the effect of the prediction of OGD to Theorem 3.5, and how that relates to an assumption present in [Har+22].

Remark A.12 (Improved predictions). Throughout Section 3.1, we have considered the standard prediction $\mathbf{m}_x^{(t)} := \mathbf{u}_x^{(t-1)} = -\mathbf{A}^{(t-1)} \mathbf{y}^{(t-1)}$ for $t \geq 2$, and similarly for Player y . It is easy to see that using the predictions

$$\mathbf{m}_x^{(t)} := -\mathbf{A}^{(t)} \mathbf{y}^{(t-1)} \text{ and } \mathbf{m}_y^{(t)} := (\mathbf{A}^{(t)})^\top \mathbf{x}^{(t-1)} \quad (43)$$

for $t \geq 1$ (where $\mathbf{z}^{(0)} := \hat{\mathbf{z}}^{(1)}$) entirely removes the dependency on $\mathcal{V}_{\mathbf{A}}^{(T)}$ on all our convergence bounds. While such a prediction cannot be implemented in the standard online learning model, there are settings in which we might know the sequence of matrices in advance; the meta-learning setting offers such examples, and indeed, Harris et al. [Har+22] use the improved prediction of (43).

Proposition A.13 (Meta-learning). *Suppose that both players employ OGD with learning rate $\eta = \frac{1}{4L}$, where $L := \max_{h \in [H]} \|\mathbf{A}^{(h)}\|_2$, and the prediction of (43) in a meta-learning bilinear saddle-point problem with $H \in \mathbb{N}$ games, each repeated for $m \in \mathbb{N}$ consecutive iterations. Then, for an average game,*

$$\left[\frac{P}{H\epsilon^2} + \frac{P' \mathcal{V}_{\text{NE}}^{(H)}}{H\epsilon^2} \right] \quad (44)$$

iterations suffice to reach an ϵ -approximate Nash equilibrium, where $P := 4L^2(4 + \|\mathcal{Z}\|_2)^2 D_Z^2$, $P' := 8L^2(4 + \|\mathcal{Z}\|_2)^2 D_Z$, and

$$\mathcal{V}_{\text{NE}}^{(H)} := \inf_{\mathbf{z}^{(h,*)} \in \mathcal{Z}^{(h,*)}, \forall h \in [H]} \sum_{h=1}^{H-1} \|\mathbf{z}^{(h+1,*)} - \mathbf{z}^{(h,*)}\|_2.$$

The proof is a direct application of Theorem A.10, where we remark that the term depending on $\mathcal{V}_{\mathbf{A}}^{(T)}$ and the term $4\eta^2 L^2 \|\mathcal{Z}\|_2^2$ from Theorem A.10 are eliminated because of the improved prediction of Remark A.12. The first term in the iteration complexity bound (44) vanishes in the meta-learning regime—as the number of games increases $H \gg 1$ —while the second term is proportional to $\frac{\mathcal{V}_{\text{NE}}^{(H)}}{H}$, a natural similarity measure; (44) always recovers the $m^{-1/2}$ rate, but offers significant gains if the games are similar, in the sense that $\frac{\mathcal{V}_{\text{NE}}^{(H)}}{H} \ll 1$. It is worth noting that, unlike the similarity measure derived in [Har+22], $\frac{\mathcal{V}_{\text{NE}}^{(H)}}{H}$ depends on the order of the games. We further remark that Proposition A.13 can be readily extended even if each game in the meta-learning sequence is not repeated for the same number of iterations.

A.1.7 General Variational Inequalities

Although our main focus in this paper is on the convergence of learning algorithms in time-varying games, our techniques could also be of interest for solving (static) general variational inequality (VI) problems.

In particular, let $F : \mathcal{Z} \rightarrow \mathcal{Z}$ be a single-valued operator. Solving general VIs is well-known to be computationally intractable, and so instead focus has been on identifying broad subclasses that elude those intractability barriers (see our overview in Section 1.2). Our framework in Section 3.1 motivates introducing the following measure of complexity for a VI problem:

$$\mathfrak{C}(F) := \inf_{\mathbf{z}^{(1,\star)}, \dots, \mathbf{z}^{(T,\star)} \in \mathcal{Z}} \sum_{t=1}^{T-1} \|\mathbf{z}^{(t+1,\star)} - \mathbf{z}^{(t,\star)}\|_2, \quad (45)$$

subject to

$$\text{DReg}_z^{(T)}(\mathbf{z}^{(1,\star)}, \dots, \mathbf{z}^{(T,\star)}) \geq 0 \iff \sum_{t=1}^T \langle \mathbf{z}^{(t)} - \mathbf{z}^{(t,\star)}, F(\mathbf{z}^{(t)}) \rangle \geq 0. \quad (46)$$

In words, (45) expresses the infimum first-order variation that a sequence of comparators must have in order to guarantee nonnegative dynamic regret (46); it is evident that (46) always admits a feasible sequence, namely $\mathbf{s}_z^{(T)} := (\mathbf{z}^{(t)})_{1 \leq t \leq T}$. We note that, in accordance to our results in Section 3.1, one can also consider an approximate version of the complexity measure (45), which could behave much more favorably (recall Proposition 3.3).

Now in a (static) bilinear saddle-point problem, it holds that $\mathfrak{C}(F) = 0$ given that there exists a static comparator that guarantees nonnegativity of the dynamic regret. More broadly, our techniques imply $O(\text{poly}(1/\epsilon))$ iteration-complexity bounds for any VI problem such that $\mathfrak{C}(F) \leq CT^{1-\omega}$, for a time-independent parameter $C > 0$ and $\omega \in (0, 1]$:

Proposition A.14. *Consider a variational inequality problem described with the operator $F : \mathcal{Z} \rightarrow \mathcal{Z}$ such that F is L -Lipschitz continuous, in the sense that $\|F(\mathbf{z}) - F(\mathbf{z}')\|_2 \leq L\|\mathbf{z} - \mathbf{z}'\|_2$, and $\mathfrak{C}(F) \leq CT^{1-\omega}$ for $C > 0$ and $\omega \in (0, 1]$. Then, OGD with learning rate $\eta = \frac{1}{4L}$ reaches an ϵ -strong solution $\mathbf{z}^* \in \mathcal{Z}$ in $O(\epsilon^{-2/\omega})$ iterations; that is, $\langle \mathbf{z} - \mathbf{z}^*, F(\mathbf{z}^*) \rangle \geq -\epsilon$ for any $\mathbf{z} \in \mathcal{Z}$.*

It is worth comparing (45) with another natural complexity measure, namely $\inf_{\mathbf{z}^* \in \mathcal{Z}} \sum_{t=1}^T \langle \mathbf{z}^{(t)} - \mathbf{z}^*, F(\mathbf{z}^{(t)}) \rangle$; the latter measures how negative (external) regret can be, and has already proven useful in certain settings that go bilinear saddle-point problems [YM22], although unlike (45), it

does not appear to be useful in characterizing time-varying bilinear saddle-point problems. In this context, $O(\text{poly}(1/\epsilon))$ iteration-complexity bounds can also be established whenever

- $\inf_{\mathbf{z}^* \in \mathcal{Z}} \sum_{t=1}^T \langle \mathbf{z}^{(t)} - \mathbf{z}^*, F(\mathbf{z}^{(t)}) \rangle \geq -CT^{1-\omega}$ for a time-invariant $C > 0$, or
- $\inf_{\mathbf{z}^* \in \mathcal{Z}} \sum_{t=1}^T \langle \mathbf{z}^{(t)} - \mathbf{z}^*, F(\mathbf{z}^{(t)}) \rangle \geq -C \sum_{t=1}^{T-1} \|\mathbf{z}^{(t+1)} - \mathbf{z}^{(t)}\|_2^2$, for a sufficiently small $C > 0$.

Following [YM22], identifying VIs that satisfy those relaxed conditions but not the MVI property is an interesting direction. In particular, it is important to understand if those relaxations can shed light into the convergence properties of OGD in Shapley's two-player zero-sum *stochastic games*.

A.2 Proofs from Section 3.2

In this subsection, we provide the proofs from Section 3.2, leading to our main result in Theorem 3.8.

Let us first introduce some additional notation. We let $f^{(t)} : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}$ be a continuously differentiable function for any $t \in \llbracket T \rrbracket$. We recall that in Section 3.2 it is assumed that the objective function changes after $m \in \mathbb{N}$ (consecutive) repetitions, which is akin to the meta-learning setting. Analogously to our setup for bilinear saddle-point problems (Section 3.1), it is assumed that Player x is endeavoring to minimizing the objective function, while Player y is trying to maximize it. We will denote by $\text{Reg}_{\mathcal{L},x}^{(T)}(\mathbf{x}^*) := \sum_{t=1}^T \langle \mathbf{x}^{(t)} - \mathbf{x}^*, -\mathbf{u}_x^{(t)} \rangle$ and $\text{Reg}_{\mathcal{L},y}^{(T)}(\mathbf{y}^*) := \sum_{t=1}^T \langle \mathbf{y}^* - \mathbf{y}^{(t)}, \mathbf{u}_y^{(t)} \rangle$, where $\mathbf{u}_x^{(t)} := -\nabla_{\mathbf{x}} f(\mathbf{x}^{(t)}, \mathbf{y}^{(t)})$ and $\mathbf{u}_y^{(t)} := \nabla_{\mathbf{y}} f(\mathbf{x}^{(t)}, \mathbf{y}^{(t)})$ for any $t \in \llbracket T \rrbracket$; similar notation is used for $\text{DReg}_{\mathcal{L},x}^{(T)}$, $\text{DReg}_{\mathcal{L},y}^{(T)}$.

Furthermore, we let $s_z^{(T)} = ((\mathbf{x}^{(t,\star)}, \mathbf{y}^{(t,\star)}))_{1 \leq t \leq T}$, so that $\mathbf{x}^{(t,\star)} = \mathbf{x}^{(h,\star)}$ and $\mathbf{y}^{(t,\star)} = \mathbf{y}^{(h,\star)}$ for any $t \in \llbracket T \rrbracket$ such that $\lfloor (t-1)/m \rfloor = h \in \llbracket H \rrbracket$. The first important step in our analysis is that, following the proof of Lemma 3.1,

$$\begin{aligned} \text{DReg}_{\mathcal{L},x}^{(T)}(s_x^{(T)}) &\leq \frac{1}{2\eta} \sum_{h=1}^H \left(\|\hat{\mathbf{x}}^{(h,1)} - \mathbf{x}^{(h,\star)}\|_2^2 - \|\hat{\mathbf{x}}^{(h,m+1)} - \mathbf{x}^{(h,\star)}\|_2^2 \right) + \eta \sum_{t=1}^T \|\mathbf{u}_x^{(t)} - \mathbf{m}_x^{(t)}\|_2^2 \\ &\quad - \frac{1}{2\eta} \sum_{t=1}^T \left(\|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t)}\|_2^2 + \|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t+1)}\|_2^2 \right), \end{aligned} \quad (47)$$

where $\hat{\mathbf{x}}^{(h,k)} := \hat{\mathbf{x}}^{((h-1) \times m + k)}$ for any $(h, k) \in \llbracket H \rrbracket \times \llbracket m \rrbracket$, $\hat{\mathbf{x}}^{(h,m+1)} := \hat{\mathbf{x}}^{(h+1,1)}$ for $h \in \llbracket H-1 \rrbracket$, and $\hat{\mathbf{x}}^{(H,m+1)} := \hat{\mathbf{x}}^{(T+1)}$. Similarly,

$$\begin{aligned} \text{DReg}_{\mathcal{L},y}^{(T)}(s_y^{(T)}) &\leq \frac{1}{2\eta} \sum_{h=1}^H \left(\|\hat{\mathbf{y}}^{(h,1)} - \mathbf{y}^{(h,\star)}\|_2^2 - \|\hat{\mathbf{y}}^{(h,m+1)} - \mathbf{y}^{(h,\star)}\|_2^2 \right) + \eta \sum_{t=1}^T \|\mathbf{u}_y^{(t)} - \mathbf{m}_y^{(t)}\|_2^2 \\ &\quad - \frac{1}{2\eta} \sum_{t=1}^T \left(\|\mathbf{y}^{(t)} - \hat{\mathbf{y}}^{(t)}\|_2^2 + \|\mathbf{y}^{(t)} - \hat{\mathbf{y}}^{(t+1)}\|_2^2 \right). \end{aligned} \quad (48)$$

Next, we will use the following key observation, which lower bounds the sum of the players' (external) regrets under strong convexity-concavity.

Lemma A.15. *Suppose that $f : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}$ is a μ -strongly convex-concave function with respect to $\|\cdot\|_2$. Then, for any Nash equilibrium $\mathbf{z}^* = (\mathbf{x}^*, \mathbf{y}^*) \in \mathcal{Z}$,*

$$\text{Reg}_{\mathcal{L},x}^{(m)}(\mathbf{x}^*) + \text{Reg}_{\mathcal{L},y}^{(m)}(\mathbf{y}^*) \geq \frac{\mu}{2} \sum_{t=1}^m \|\mathbf{z}^{(t)} - \mathbf{z}^*\|_2^2.$$

Proof. First, by μ -strong convexity of $f(\mathbf{x}, \cdot)$, we have that for any time $t \in \llbracket m \rrbracket$,

$$\langle \mathbf{x}^{(t)} - \mathbf{x}^*, \nabla_{\mathbf{x}} f(\mathbf{x}^{(t)}, \mathbf{y}^{(t)}) \rangle \geq f(\mathbf{x}^{(t)}, \mathbf{y}^{(t)}) - f(\mathbf{x}^*, \mathbf{y}^{(t)}) + \frac{\mu}{2} \|\mathbf{x}^{(t)} - \mathbf{x}^*\|_2^2. \quad (49)$$

Similarly, by μ -strong concavity of $f(\cdot, \mathbf{y})$, we have that for any time $t \in \llbracket m \rrbracket$,

$$\langle \mathbf{y}^* - \mathbf{y}^{(t)}, \nabla_{\mathbf{y}} f(\mathbf{x}^{(t)}, \mathbf{y}^{(t)}) \rangle \geq f(\mathbf{x}^{(t)}, \mathbf{y}^*) - f(\mathbf{x}^{(t)}, \mathbf{y}^{(t)}) + \frac{\mu}{2} \|\mathbf{y}^{(t)} - \mathbf{y}^*\|_2^2. \quad (50)$$

Further, for any Nash equilibrium $(\mathbf{x}^*, \mathbf{y}^*) \in \mathcal{Z}$ it holds that $f(\mathbf{x}^{(t)}, \mathbf{y}^*) \geq f(\mathbf{x}^{(t)}, \mathbf{y}^{(t)}) \geq f(\mathbf{x}^*, \mathbf{y}^{(t)})$. Combining this fact with (49) and (50) and summing over all $t \in \llbracket m \rrbracket$ gives the statement. \square

In turn, this readily implies the following lower bound for the dynamic regret.

Lemma A.16. *Suppose that $f^{(h)} : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}$ is a μ -strongly convex-concave function with respect to $\|\cdot\|_2$, for any $h \in \llbracket H \rrbracket$. Consider a sequence $s_z^{(T)} = ((\mathbf{x}^{(t,\star)}, \mathbf{y}^{(t,\star)}))_{1 \leq t \leq T}$, so that $\mathbf{x}^{(t,\star)} = \mathbf{x}^{(h,\star)}$ and $\mathbf{y}^{(t,\star)} = \mathbf{y}^{(h,\star)}$ for any $t \in \llbracket T \rrbracket$ such that $\lfloor (t-1)/m \rfloor = h \in \llbracket H \rrbracket$. If $(\mathbf{x}^{(h,\star)}, \mathbf{y}^{(h,\star)}) \in \mathcal{Z}$ is a Nash equilibrium of $f^{(h)}$,*

$$\text{DReg}_{\mathcal{L},x}^{(T)}(s_x^{(T)}) + \text{DReg}_{\mathcal{L},y}^{(T)}(s_y^{(T)}) \geq \frac{\mu}{2} \sum_{h=1}^H \sum_{k=1}^m \|\mathbf{z}^{(h,k)} - \mathbf{z}^{(h,\star)}\|_2^2,$$

where $\mathbf{z}^{(h,k)} := \mathbf{z}^{((h-1) \times m + k)}$ for any $(h, k) \in \llbracket H \rrbracket \times \llbracket m \rrbracket$.

We next combine this with the following monotonicity property of OGD: If \mathbf{z}^* is a Nash equilibrium, $\|\hat{\mathbf{z}}^{(t)} - \mathbf{z}^*\|_2$ is a decreasing function in t [Har+22, Proposition C.10]. This leads to the following refinement of Lemma A.16.

Lemma A.17. *Under the assumptions of Lemma A.16, if $\eta \leq \frac{1}{2\mu}$,*

$$\text{DReg}_{\mathcal{L},x}^{(T)}(s_x^{(T)}) + \text{DReg}_{\mathcal{L},y}^{(T)}(s_y^{(T)}) + \frac{1}{4\eta} \sum_{t=1}^T \|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t+1)}\|_2^2 \geq \frac{\mu m}{4} \sum_{h=1}^H \|\hat{\mathbf{z}}^{(h,m+1)} - \mathbf{z}^{(h,\star)}\|_2^2.$$

Proof. By Lemma A.16,

$$\begin{aligned} \text{DReg}_{\mathcal{L},x}^{(T)}(s_x^{(T)}) + \text{DReg}_{\mathcal{L},y}^{(T)}(s_y^{(T)}) + \frac{1}{4\eta} \sum_{t=1}^T \|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t+1)}\|_2^2 &\geq \frac{\mu}{2} \sum_{h=1}^H \sum_{k=1}^m \|\mathbf{z}^{(h,k)} - \mathbf{z}^{(h,\star)}\|_2^2 \\ &\quad + \frac{1}{4\eta} \sum_{t=1}^T \|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t+1)}\|_2^2 \\ &\geq \frac{\mu}{4} \sum_{h=1}^H \sum_{k=1}^m \|\hat{\mathbf{z}}^{(h,k+1)} - \mathbf{z}^{(h,\star)}\|_2^2 \end{aligned} \quad (51)$$

$$\geq \frac{\mu m}{4} \sum_{h=1}^H \|\hat{\mathbf{z}}^{(h,m+1)} - \mathbf{z}^{(h,\star)}\|_2^2, \quad (52)$$

where (51) uses that $\frac{1}{4\eta} \geq \frac{\mu}{2}$ along with Young's inequality and triangle inequality, and (52) follows from [Har+22, Proposition C.10]. \square

Armed with this important lemma, we are ready to establish our main result (Theorem 3.8), the detailed version of which is given below. We first point out that a function $f : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}$ is said to be L -smooth if $\|F(\mathbf{z}) - F(\mathbf{z}')\|_2 \leq L\|\mathbf{z} - \mathbf{z}'\|_2$, where $F(\mathbf{z}) := (\nabla_{\mathbf{x}}f(\mathbf{x}, \mathbf{y}), -\nabla_{\mathbf{y}}f(\mathbf{x}, \mathbf{y}))$.

Theorem A.18 (Detailed version of Theorem 3.8). *Let $f^{(h)} : \mathcal{X} \times \mathcal{Y}$ be a μ -strongly convex-concave and L -smooth function, for $h \in \llbracket H \rrbracket$. Suppose that both players employ OGD with learning rate $\eta = \min\left\{\frac{1}{8L}, \frac{1}{2\mu}\right\}$ for T repetitions, where $T = m \times H$ and $m \geq \frac{2}{\eta\mu}$. Then,*

$$\sum_{t=1}^T \left(\|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t)}\|_2^2 + \|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t+1)}\|_2^2 \right) \leq 4D_{\mathcal{Z}}^2 + 8\eta^2\|F(\mathbf{z}^{(1)})\|_2^2 + 8\mathcal{S}_{NE}^{(H)} + 16\eta^2\mathcal{V}_{\nabla f}^{(H)},$$

where $\mathcal{S}_{NE}^{(H)}$ and $\mathcal{V}_{\nabla f}^{(H)}$ are defined in (9) and (11). Thus, $\sum_{t=1}^T \left(\text{EQGAP}^{(t)}(\mathbf{z}^{(t)}) \right)^2 = O(1 + \mathcal{S}_{NE}^{(H)} + \mathcal{V}_{\nabla f}^{(H)})$.

Proof. Combining Lemma A.17 with (47) and (48),

$$\begin{aligned} 0 &\leq \frac{1}{2\eta} \sum_{h=1}^H \left(\|\hat{\mathbf{z}}^{(h,1)} - \mathbf{z}^{(h,\star)}\|_2^2 - \frac{2 + \eta\mu m}{2} \|\hat{\mathbf{z}}^{(h,m+1)} - \mathbf{z}^{(h,\star)}\|_2^2 \right) \\ &\quad + \eta \sum_{t=1}^T \|\mathbf{u}_z^{(t)} - \mathbf{m}_z^{(t)}\|_2^2 - \frac{1}{4\eta} \sum_{t=1}^T \left(\|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t)}\|_2^2 + \|\mathbf{z}^{(t)} - \hat{\mathbf{z}}^{(t+1)}\|_2^2 \right), \end{aligned}$$

for a sequence of Nash equilibria $(\mathbf{z}^{(h,\star)})_{1 \leq h \leq H}$, where we used the notation $\mathbf{u}_z^{(t)} := (\mathbf{u}_x^{(t)}, \mathbf{u}_y^{(t)})$ and $\mathbf{m}_z^{(t)} := (\mathbf{m}_x^{(t)}, \mathbf{m}_y^{(t)})$. Now we bound the first term of the right-hand side above as

$$\begin{aligned} &\frac{1}{2\eta} \sum_{h=1}^H \left(\|\hat{\mathbf{z}}^{(h,1)} - \mathbf{z}^{(h,\star)}\|_2^2 - 2\|\hat{\mathbf{z}}^{(h,m+1)} - \mathbf{z}^{(h,\star)}\|_2^2 \right) \leq \\ &\quad \frac{1}{2\eta} \|\hat{\mathbf{z}}^{(1,1)} - \mathbf{z}^{(1,\star)}\|_2^2 + \frac{1}{2\eta} \sum_{h=1}^{H-1} \left(\|\hat{\mathbf{z}}^{(h+1,1)} - \mathbf{z}^{(h+1,\star)}\|_2^2 - 2\|\hat{\mathbf{z}}^{(h+1,1)} - \mathbf{z}^{(h,\star)}\|_2^2 \right), \end{aligned}$$

where we used the fact that $m \geq \frac{2}{\eta\mu}$ and $\hat{\mathbf{z}}^{(h,m+1)} = \hat{\mathbf{z}}^{(h+1,1)}$, for $h \in \llbracket H-1 \rrbracket$. Hence, continuing from above,

$$\frac{1}{2\eta} \sum_{h=1}^H \left(\|\hat{\mathbf{z}}^{(h,1)} - \mathbf{z}^{(h,\star)}\|_2^2 - 2\|\hat{\mathbf{z}}^{(h,m+1)} - \mathbf{z}^{(h,\star)}\|_2^2 \right) \leq \frac{1}{2\eta} \|\hat{\mathbf{z}}^{(1,1)} - \mathbf{z}^{(1,\star)}\|_2^2 + \frac{1}{\eta} \sum_{h=1}^{H-1} \|\mathbf{z}^{(h+1,\star)} - \mathbf{z}^{(h,\star)}\|_2^2,$$

since $\|\hat{\mathbf{z}}^{(h+1,1)} - \mathbf{z}^{(h+1,\star)}\|_2^2 \leq 2\|\hat{\mathbf{z}}^{(h+1,1)} - \mathbf{z}^{(h,\star)}\|_2^2 + 2\|\mathbf{z}^{(h,\star)} - \mathbf{z}^{(h+1,\star)}\|_2^2$, by the triangle inequality and Young's inequality. Moreover, for $t \geq 2$,

$$\|\mathbf{u}_z^{(t)} - \mathbf{u}_z^{(t-1)}\|_2^2 = \|F^{(t)}(\mathbf{z}^{(t)}) - F^{(t-1)}(\mathbf{z}^{(t-1)})\|_2^2 \leq 2L^2\|\mathbf{z}^{(t)} - \mathbf{z}^{(t-1)}\|_2^2 + 2\|F^{(t)}(\mathbf{z}^{(t-1)}) - F^{(t-1)}(\mathbf{z}^{(t-1)})\|_2^2,$$

by L -smoothness. As a result,

$$\sum_{t=1}^{T-1} \|\mathbf{u}_z^{(t+1)} - \mathbf{u}_z^{(t)}\|_2^2 \leq 2L^2 \sum_{t=1}^{T-1} \|\mathbf{z}^{(t)} - \mathbf{z}^{(t-1)}\|_2^2 + 2\mathcal{V}_{\nabla f}^{(T)},$$

and the claimed bound on the second-order path length follows. Finally, the second claim of the theorem follows from Claim A.9 using convexity-concavity, analogously to Theorem 3.5. \square

We point out that the improved prediction mechanism described in Remark A.12 can also be used in this setting as well, resulting in the elimination of the variation measure (11) from Theorem A.18. We conclude this subsection by pointing out an improved variation-dependent regret bound, which follows directly from Theorem A.18 (cf. Corollary A.11).

Corollary A.19. *In the setup of Theorem A.18,*

$$\text{Reg}_{\mathcal{L},x}^{(T)}, \text{Reg}_{\mathcal{L},y}^{(T)} \leq \frac{D_{\mathcal{Z}}^2}{\eta} + 4\eta L^2 D_{\mathcal{Z}}^2 + 32\eta^3 L^2 \|F(\mathbf{z}^{(1)})\|_2^2 + 32\eta L^2 \mathcal{S}_{NE}^{(H)} + (64\eta^3 L^2 + 2\eta) \mathcal{V}_{\nabla f}^{(H)}.$$

Thus, setting the learning rate optimally implies that $\text{Reg}_{\mathcal{L},x}^{(T)}, \text{Reg}_{\mathcal{L},y}^{(T)} = O\left(\sqrt{\mathcal{S}_{NE}^{(H)} + \mathcal{V}_{\nabla f}^{(H)}}\right)$.

A.3 Proofs from Section 3.3

In this subsection, we provide the proofs from Section 3.3.

A.3.1 Potential Games

We first characterize the behavior of GD in time-varying potential games. Below we give the formal definition of an unweighted potential game, represented in normal form.

Definition A.20 (Potential game). A game admits a potential if there exists a function $\Phi : \times_{i=1}^n \mathcal{X}_i \rightarrow \mathbb{R}$ such that for any Player $i \in \llbracket n \rrbracket$, any joint strategy profile $\mathbf{x}_{-i} \in \times_{i' \neq i} \mathcal{X}_{i'}$, and any pair of strategies $\mathbf{x}_i, \mathbf{x}'_i \in \mathcal{X}_i$,

$$\Phi(\mathbf{x}_i, \mathbf{x}_{-i}) - \Phi(\mathbf{x}'_i, \mathbf{x}_{-i}) = u_i(\mathbf{x}_i, \mathbf{x}_{-i}) - u_i(\mathbf{x}'_i, \mathbf{x}_{-i}).$$

We also recall that GD is equivalent to OGD under the prediction $\mathbf{m}_x^{(t)} = \mathbf{0}$ for all t . The key ingredient in the proof of Theorem 3.10 is the following key bound on the second-order path length of the dynamics.

Proposition A.21. *Suppose that each player employs GD with a sufficiently small learning rate $\eta > 0$ and initialization $(\mathbf{x}_1^{(1)}, \dots, \mathbf{x}_n^{(1)}) \in \times_{i=1}^n \mathcal{X}_i$. Then,*

$$\frac{1}{2\eta} \sum_{t=1}^T \sum_{i=1}^n \|\mathbf{x}_i^{(t+1)} - \mathbf{x}_i^{(t)}\|_2^2 \leq \sum_{t=1}^T \left(\Phi^{(t)}(\mathbf{x}_1^{(t+1)}, \dots, \mathbf{x}_n^{(t+1)}) - \Phi^{(t)}(\mathbf{x}_1^{(t)}, \dots, \mathbf{x}_n^{(t)}) \right). \quad (53)$$

This bound can be derived from [Ana+22b, Theorem 4.3]. We note that if $\Phi^{(1)} = \Phi^{(2)} = \dots = \Phi^{(T)}$, the right-hand side of (53) telescops, thereby implying that the second-order path-length is bounded. More generally, the right-hand side of (53) can be upper bounded by

$$2\Phi_{\max} + \sum_{t=1}^{T-1} \left(\Phi^{(t)}(\mathbf{x}_1^{(t+1)}, \dots, \mathbf{x}_n^{(t+1)}) - \Phi^{(t+1)}(\mathbf{x}_1^{(t+1)}, \dots, \mathbf{x}_n^{(t+1)}) \right) \leq 2\Phi_{\max} + \mathcal{V}_{\Phi}^{(T)}, \quad (54)$$

where Φ_{\max} is an upper bound on $|\Phi^{(t)}(\cdot)|$ for any $t \in \llbracket T \rrbracket$, and $\mathcal{V}_{\Phi}^{(T)}$ is the variation measure of the potential functions we introduced in Section 3.3. Furthermore, we know that the Nash equilibrium gap in the t -th potential game can be bounded in terms of $\sum_{i=1}^n \|\mathbf{x}_i^{(t+1)} - \mathbf{x}_i^{(t)}\|_2$ (Claim A.9). As a result, combining this property with Proposition A.21 and (54) establishes Theorem 3.10.

A.3.2 General-Sum Games

We next turn our attention to general-sum multi-player games using the bilinear formulation presented in Section 3.3. To establish Property 3.11, let us first define the regret of any Player $i \in \llbracket n \rrbracket$ as

$$\text{Reg}_i^{(T)}(\bar{\mathbf{x}}_i^*) := \sum_{t=1}^T \langle \bar{\mathbf{x}}_i^* - \bar{\mathbf{x}}_i^{(t)}, (\mathbf{A}_i^{(t)})^\top \boldsymbol{\mu}^{(t)} \rangle,$$

where $\bar{\mathbf{x}}_i^* \in \bar{\mathcal{X}}_i$, so that $\sum_{i=1}^n \text{Reg}_i^{(T)}$ is easily seen to be equal to the regret of the maximizing player in (12). Further, the dynamic regret of the mediator—the minimizing player in (12)—can be expressed as

$$\text{DReg}_\mu^{(T)}(\boldsymbol{\mu}^{(1,\star)}, \dots, \boldsymbol{\mu}^{(T,\star)}) := \sum_{t=1}^T \langle \boldsymbol{\mu}^{(t)} - \boldsymbol{\mu}^{(t,\star)}, \sum_{i=1}^n \mathbf{A}_i^{(t)} \bar{\mathbf{x}}_i^{(t)} \rangle.$$

Property 3.11. *Suppose that $\Xi \ni \boldsymbol{\mu}^{(t,\star)}$ is a correlated equilibrium of the game at any time $t \in \llbracket T \rrbracket$. Then,*

$$\text{DReg}_\mu^{(T)}(\boldsymbol{\mu}^{(1,\star)}, \dots, \boldsymbol{\mu}^{(T,\star)}) + \sum_{i=1}^n \text{Reg}_i^{(T)} \geq 0.$$

Proof. We have that

$$\text{DReg}_\mu^{(T)} + \sum_{i=1}^n \text{Reg}_i^{(T)}(\bar{\mathbf{x}}_i^*) = \sum_{i=1}^n \sum_{t=1}^T \langle \bar{\mathbf{x}}_i^*, (\mathbf{A}_i^{(t)})^\top \boldsymbol{\mu}^{(t)} \rangle - \sum_{t=1}^T \langle \boldsymbol{\mu}^{(t,\star)}, \sum_{i=1}^n \mathbf{A}_i^{(t)} \bar{\mathbf{x}}_i^{(t)} \rangle.$$

Now for any correlated equilibrium $\boldsymbol{\mu}^{(t,\star)}$ of the t -th game we have that $\langle \boldsymbol{\mu}^{(t,\star)}, \mathbf{A}_i^{(t)} \bar{\mathbf{x}}_i^{(t)} \rangle \leq 0$ for any Player $i \in \llbracket n \rrbracket$, $\bar{\mathbf{x}}_i \in \bar{\mathcal{X}}_i$, and time $t \in \llbracket T \rrbracket$, which in turn implies that $-\sum_{t=1}^T \langle \boldsymbol{\mu}^{(t,\star)}, \sum_{i=1}^n \mathbf{A}_i^{(t)} \bar{\mathbf{x}}_i^{(t)} \rangle \geq 0$. Moreover, $\sum_{i=1}^n \max_{\bar{\mathbf{x}}_i^* \in \bar{\mathcal{X}}_i} \sum_{t=1}^T \langle \bar{\mathbf{x}}_i^*, (\mathbf{A}_i^{(t)})^\top \boldsymbol{\mu}^{(t)} \rangle \geq 0$ given that, by definition, $\mathbf{0} \in \bar{\mathcal{X}}_i$. This concludes the proof. \square

Next, we provide the main implication of Theorem 3.12 in the meta-learning setting, which is similar to the meta-learning guarantee of Proposition A.13 we established earlier in two-player zero-sum games. Below, we denote by $\Xi^{(h,\star)}$ the set of correlated equilibria of the h -th game in the meta-learning sequence.

Corollary A.22 (Meta-learning general-sum). *Suppose that each player employ OGD in (12) with a suitable learning rate $\eta > 0$ and the prediction of (43) in a meta-learning general-sum problem with $H \in \mathbb{N}$ games, each repeated for $m \in \mathbb{N}$ consecutive iterations. Then, for an average game,*

$$O\left(\frac{1}{\epsilon^2 H} + \frac{\mathcal{V}_{CE}^{(H)}}{\epsilon^2 H}\right) \tag{55}$$

iterations suffice to reach an ϵ -approximate correlated equilibrium, where

$$\mathcal{V}_{CE}^{(H)} := \inf_{\boldsymbol{\mu}^{(h,\star)} \in \Xi^{(h,\star)}} \|\boldsymbol{\mu}^{(h+1,\star)} - \boldsymbol{\mu}^{(h,\star)}\|_2.$$

In particular, in the meta-learning regime, $H \gg 1$, the iteration-complexity bound (55) is dominated by the (algorithm-independent) similarity metric of the correlated equilibria $\frac{\mathcal{V}_{\text{CE}}^{(H)}}{H}$. Corollary A.22 establishes significant gains when $\frac{\mathcal{V}_{\text{CE}}^{(H)}}{H} \ll 1$.

Finally, we conclude this subsection by providing a variation-dependent regret bound in general-sum multi-player games. To do so, we combine Corollary 3.7 with Theorem 3.12, leading to the following guarantee.

Corollary A.23 (Regret in general-sum games). *In the setup of Theorem 3.12,*

$$\text{Reg}_\mu^{(T)}, \text{Reg}_i^{(T)} = O\left(\frac{1}{\eta} + \eta\left(\mathcal{V}_{\text{CE}}^{(T)} + \mathcal{V}_{\mathbf{A}}^{(T)}\right)\right),$$

for any Player $i \in \llbracket n \rrbracket$.

In particular, if one selects optimally the learning rate, Corollary A.23 implies that the individual regret of each player is bounded by $O\left(\sqrt{\mathcal{V}_{\text{CE}}^{(T)} + \mathcal{V}_{\mathbf{A}}^{(T)}}\right)$. We note again that there are techniques that would allow (nearly) recovering such regret guarantees without having to know the variation measures in advance [Zha+22].

A.4 Proofs from Section 3.4

Finally, in this subsection we present the proofs omitted from Section 3.4. We begin with Proposition 3.13, the statement of which is recalled below. We first recall that a regularizer ϕ_x , 1-strongly convex with respect to a norm $\|\cdot\|$, is said to be G -smooth if $\|\nabla\phi_x(\mathbf{x}) - \nabla\phi_x(\mathbf{x}')\|_* \leq G\|\mathbf{x} - \mathbf{x}'\|$, for all \mathbf{x}, \mathbf{x}' .

Proposition 3.13. *Suppose that both players in a (static) two-player zero-sum game employ OMD with a smooth regularizer. Then, $\text{DReg}_x^{(T)}, \text{DReg}_y^{(T)} = O(\sqrt{T})$.*

Proof. First, using Claim A.9, it follows that the dynamic regret $\text{DReg}_x^{(T)}$ of Player x up to time T can be bounded as

$$\begin{aligned} & \sum_{t=1}^T \left(\max_{\mathbf{x}^{(t,*)} \in \mathcal{X}} \left\{ \langle \mathbf{x}^{(t,*)}, \mathbf{u}_x^{(t)} \rangle \right\} - \langle \mathbf{x}^{(t)}, \mathbf{u}_x^{(t)} \rangle \right) \\ & \leq \sum_{t=1}^T \left(\left(\frac{G}{\eta} + \|\mathbf{u}_x^{(t)}\|_* \right) \|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t+1)}\| + \frac{G}{\eta} \|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t)}\| \right), \end{aligned} \quad (56)$$

where $G > 0$ is the smoothness parameter of the regularizer, and $\eta > 0$ is the learning rate. We further know that $\sum_{t=1}^T (\|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t)}\|^2 + \|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t+1)}\|^2) = O(1)$ for any instance of OMD in a two-player zero-sum game [Ana+22b], which in turn implies that $\sum_{t=1}^T (\|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t)}\| + \|\mathbf{x}^{(t)} - \hat{\mathbf{x}}^{(t+1)}\|) = O(\sqrt{T})$ by Cauchy-Schwarz. Thus, combining with (56) we have shown that $\text{DReg}_x^{(T)} = O(\sqrt{T})$. Similar reasoning yields that $\text{DReg}_y^{(T)} = O(\sqrt{T})$, concluding the proof. \square

In contrast, we next show that such a result is precluded in general-sum games. In particular, we note that the following computational-hardness result holds beyond the online learning setting.

It should be stressed that without imposing computational or memory restrictions there are trivial online algorithms that guarantee even $O(1)$ dynamic regret by first exploring the payoff matrices and then computing a Nash equilibrium; we suspect that under the memory limitations imposed in our work, as in [DDK11], there could be unconditional information-theoretic lower bounds, but that is left for future work.

Proposition 3.15. *Unless $\text{PPAD} \subseteq \text{P}$, any polynomial-time algorithm incurs $\sum_{i=1}^n \text{DReg}_i^{(T)} = \Omega(T)$, even if $n = 2$, where $\Omega(\cdot)$ here hides polynomial factors.*

Proof. We will use the fact that computing a Nash equilibrium in two-player (normal-form) games to a sufficiently small accuracy $\epsilon := 1/\text{poly}$ is PPAD-hard [CDT09]. Indeed, suppose that there exist polynomial-time algorithms that always guarantee that $\sum_{i=1}^n \text{DReg}_i^{(T)} \leq \epsilon T$, where $n := 2$. Then, this implies that there exists a time $t \in \llbracket T \rrbracket$ such that

$$\max_{\mathbf{x}_1^{(t,\star)} \in \mathcal{X}_1} \langle \mathbf{x}_1^{(t,\star)}, \mathbf{u}_1^{(t)} \rangle - \langle \mathbf{x}_1^{(t)}, \mathbf{u}_1^{(t)} \rangle + \max_{\mathbf{x}_2^{(t,\star)} \in \mathcal{X}_2} \langle \mathbf{x}_2^{(t,\star)}, \mathbf{u}_2^{(t)} \rangle - \langle \mathbf{x}_2^{(t)}, \mathbf{u}_2^{(t)} \rangle \leq \epsilon,$$

which in turn implies that $(\mathbf{x}_1^{(t)}, \mathbf{x}_2^{(t)})$ is an ϵ -approximate Nash equilibrium. Further, such a time $t \in \llbracket T \rrbracket$ can be identified in polynomial time. But this would imply that $\text{PPAD} \subseteq \text{P}$, concluding the proof. \square

Finally, we provide the proof of Theorem 3.16, the detailed version of which is provided below.

Theorem A.24 (Detailed version of Theorem 3.16). *Consider an n -player game such that $\|\nabla_{\mathbf{x}_i} u_i(\mathbf{z}) - \nabla_{\mathbf{x}_i} u_i(\mathbf{z}')\|_2 \leq L\|\mathbf{z} - \mathbf{z}'\|_2$, where $\mathbf{z}, \mathbf{z}' \in \times_{i=1}^n \mathcal{X}_i$, for any Player $i \in \llbracket n \rrbracket$. Then, if all players employ OGD with learning rate $\eta > 0$ it holds that*

1. $\sum_{i=1}^n K\text{-DReg}_i^{(T)} = O(K\sqrt{n}L)$ for $\eta = \Theta\left(\frac{1}{L\sqrt{n}}\right)$;
2. $K\text{-DReg}_i^{(T)} = O(K^{3/4}T^{1/4}n^{1/4}\sqrt{L})$, for any Player $i \in \llbracket n \rrbracket$, for $\eta = \Theta\left(\frac{K^{1/4}}{n^{1/4}L^{1/2}}\right)$.

Proof. First, applying Lemma 3.1 subject to the constraint that $\sum_{t=1}^{T-1} \mathbb{1}\{\mathbf{x}^{(t+1,\star)} \neq \mathbf{x}^{(t,\star)}\} \leq K - 1$ gives that for any Player $i \in \llbracket n \rrbracket$,

$$K\text{-DReg}_i^{(T)} \leq \frac{D_{\mathcal{X}_i}^2}{2\eta} (2K - 1) + \eta \|\mathbf{u}_i^{(1)}\|_2^2 + \eta \sum_{t=1}^{T-1} \|\mathbf{u}_i^{(t+1)} - \mathbf{u}_i^{(t)}\|_2^2 - \frac{1}{4\eta} \sum_{t=1}^{T-1} \|\mathbf{x}_i^{(t+1)} - \mathbf{x}_i^{(t)}\|_2^2. \quad (57)$$

Further, by L -smoothness we have that

$$\|\mathbf{u}_i^{(t+1)} - \mathbf{u}_i^{(t)}\|_2^2 = \|\nabla_{\mathbf{x}_i} u_i(\mathbf{z}^{(t+1)}) - \nabla_{\mathbf{x}_i} u_i(\mathbf{z}^{(t)})\|_2^2 \leq L^2 \sum_{i=1}^n \|\mathbf{x}_i^{(t+1)} - \mathbf{x}_i^{(t)}\|_2^2,$$

for any $t \in \llbracket T - 1 \rrbracket$, where $(\mathbf{x}_1^{(t)}, \dots, \mathbf{x}_n^{(t)}) = \mathbf{z}^{(t)} \in \times_{i=1}^n \mathcal{X}_i$ is the joint strategy profile at time t . Thus, summing (57) over all $i \in \llbracket n \rrbracket$ and taking $\eta \leq \frac{1}{2L\sqrt{n}}$ implies that $\sum_{i=1}^n K\text{-DReg}_i^{(T)} \leq \frac{2K-1}{2\eta} \sum_{i=1}^n D_{\mathcal{X}_i}^2 + \eta \sum_{i=1}^n \|\mathbf{u}_i^{(1)}\|_2^2$, yielding the first part of the statement. The second part follows directly from (57) using the stability property of OGD: $\|\mathbf{x}_i^{(t+1)} - \mathbf{x}_i^{(t)}\|_2 = O(\eta)$, for any time $t \in \llbracket T - 1 \rrbracket$. \square

B Experimental Examples

Finally, although the focus of this paper is theoretical, in this section we provide some illustrative experimental examples. In particular, Appendix B.1 contains experiments on time-varying potential games, while Appendix B.2 focuses on time-varying (two-player) zero-sum games. For simplicity, we will be assuming that each game is represented in normal form.

B.1 Time-Varying Potential Games

Here we consider time-varying 2-player identical-interest games. We point out that such games are potential games (recall Definition A.20), and as such they are indeed amenable to our theory in Section 3.3.

In our first experiment, we first sampled two matrices $\mathbf{A}, \mathbf{P} \in \mathbb{R}^{d_x \times d_y}$, where $d_x = d_y = 1000$. Then, we defined each payoff matrix as $\mathbf{A}^{(t)} := \mathbf{A}^{(t-1)} + \mathbf{P}t^{-\alpha}$ for $t \geq 1$, where $\mathbf{A}^{(0)} := \mathbf{A}$. Here, $\alpha > 0$ is a parameter that controls the variation of the payoff matrices. In this time-varying setup, we let each player employ (online) GD with learning rate $\eta := 0.1$. The results obtained under different random initializations of matrices \mathbf{A} and \mathbf{P} are illustrated in Figure 1.

Next, we operate in the same time-varying setup but each player is now employing multiplicative weights update (MWU), instead of gradient descent, with $\eta := 0.1$. As shown in Figure 2, while the cumulative equilibrium gap is much larger compared to using GD (Figure 1), the dynamics still appear to be approaching equilibria, although our theory does not cover MWU. We suspect that theoretical results such as Theorem 3.10 should hold for MWU as well, but that has been left for future work.

In our third experiment for identical-interest games, we again first sampled two matrices $\mathbf{A}, \mathbf{P} \in \mathbb{R}^{d_x \times d_y}$, where $d_x = d_y = 1000$. Then, we defined $\mathbf{A}^{(t)} := \mathbf{A}^{(t-1)} + \epsilon \mathbf{P}$ for $t \geq 1$, where $\mathbf{A}^{(0)} := \mathbf{A}$. Here, $\epsilon > 0$ is the parameter intended to capture the variation of the payoff matrices. The results obtained under different random initializations of \mathbf{A} and \mathbf{P} are illustrated in Figure 3. As an aside, it is worth pointing out that this particular setting can be thought of as a game in which the variation in the payoff matrices is controlled by another learning agent. In particular, our theoretical results are helpful for characterizing the convergence properties of *two-timescale* learning algorithms, in which the deviation of the game is controlled by a player constrained to be updating its strategies with a much smaller learning rate.

B.2 Time-Varying Zero-Sum Games

We next conduct experiments on time-varying bilinear saddle-point problems when players are employing OGD. Such problems have been studied extensively in Section 3.1 from a theoretical standpoint.

First, we sampled two matrices $\mathbf{A}, \mathbf{P} \in \mathbb{R}^{d_x \times d_y}$, where $d_x = d_y = 10$; here we consider lower-dimensional payoff matrices compared to the experiments in Appendix B.1 for convenience in the graphical illustrations. Then, we defined each payoff matrix as $\mathbf{A}^{(t)} := \mathbf{A}^{(t-1)} + \mathbf{P}t^{-\alpha}$ for $t \geq 1$, where $\mathbf{A}^{(1)} := \mathbf{A}$. The results obtained under different random initializations are illustrated in Figure 4.




Figure 1: The equilibrium gap and the players' regrets in 2-player time-varying identical-interest games when both players are employing (online) GD with learning rate $\eta := 0.1$ for $T := 200$ iterations. Each row corresponds to a different random initialization of the matrices $\mathbf{A}, \mathbf{P} \in \mathbb{R}^{d_x \times d_y}$, which in turn induces a different time-varying game. Further, each figure contains trajectories corresponding to three different values of $\alpha \in \{0.1, 0.2, 0.5\}$, but under the same initialization of \mathbf{A} and \mathbf{P} . As expected, smaller values of α generally increase the equilibrium gap since the variation of the games is more significant. Nevertheless, for all games we observe that the players are gradually approaching equilibria.




Figure 2: The equilibrium gap and the players’ regrets in 2-player time-varying identical-interest games when both players are employing (online) GD with learning rate $\eta := 0.1$ for $T := 200$ iterations. Each row corresponds to a different random initialization of the matrices $\mathbf{A}, \mathbf{P} \in \mathbb{R}^{d_x \times d_y}$, which in turn induces a different time-varying game. Further, each figure contains trajectories corresponding to three different values of $\alpha \in \{0.1, 0.2, 0.5\}$, but under the same initialization of \mathbf{A} and \mathbf{P} . The MWU dynamics still appear to be approaching equilibria, although the cumulative gap is much larger compared to GD (Figure 1).




Figure 3: The equilibrium gap and the players’ regrets in 2-player time-varying identical-interest games when both players are employing (online) GD with learning rate $\eta := 0.1$ for $T := 500$ iterations. Each row corresponds to a different random initialization of the matrices $\mathbf{A}, \mathbf{P} \in \mathbb{R}^{d_x \times d_y}$, which in turn induces a different time-varying game. Further, each figure contains trajectories from three different values of $\epsilon \in \{0.1, 0.01, 0.001\}$, but under the same initialization of \mathbf{A} and \mathbf{P} . As expected, larger values of ϵ generally increase the equilibrium gap since the variation of the games is more significant. Yet, even for the larger value $\epsilon = 0.1$, the dynamics are still appear to be approaching Nash equilibria.




Figure 4: The equilibrium gap and the players’ regrets in 2-player time-varying zero-sum games when both players are employing OGD with learning rate $\eta := 0.01$ and $T := 1000$ iterations. Each row corresponds to a different random initialization of the matrices $\mathbf{A}, \mathbf{P} \in \mathbb{R}^{d_x \times d_y}$, which in turn induces a different time-varying game. Further, each figure contains trajectories from three different values of $\alpha \in \{0.7, 1, 2\}$, but under the same initialization of \mathbf{A} and \mathbf{P} . The OGD dynamics appear to be approaching equilibria, albeit with a much slower rate compared to the ones observed earlier for potential games (Figure 1).