

# ANOTHER LOOK AT PARTIALLY OBSERVED OPTIMAL STOCHASTIC CONTROL: EXISTENCE, ERGODICITY, AND APPROXIMATIONS WITHOUT BELIEF-REDUCTION

SERDAR YÜKSEL \*

**Abstract.** We present an alternative view for the study of optimal control of partially observed Markov Decision Processes (POMDPs). We first revisit the traditional (and by now standard) separated-design method of reducing the problem to fully observed MDPs (belief-MDPs), and present conditions for the existence of optimal policies. Then, rather than working with this standard method, we define a Markov chain taking values in an infinite dimensional product space with control actions and the state process causally conditionally independent given the measurement/information process. We provide new sufficient conditions for the existence of optimal control policies. In particular, while in the belief-MDP reduction of POMDPs, weak Feller condition requirement imposes total variation continuity on either the system kernel or the measurement kernel, with the approach of this paper only weak continuity of both the transition kernel and the measurement kernel is needed (and total variation continuity is not) together with regularity conditions related to filter stability. For the average cost setup, we provide a general approach on how to initialize the randomness which we show to establish convergence to optimal cost. For the discounted cost setup, we establish near optimality of finite window policies via a direct argument involving near optimality of quantized approximations for MDPs under weak Feller continuity, where finite truncations of memory can be viewed as quantizations of infinite memory with a uniform diameter in each finite window restriction under the product metric. In the control-free case, our analysis leads to new and weak conditions for the existence and uniqueness of invariant probability measures for non-linear filter processes, where we show that unique ergodicity of the measurement process and a measurability condition related to filter stability leads to unique ergodicity.

**AMS subject classifications:** 60J20,60J05,93E11

**1. Introduction, Literature Review, and Main Results.** Partially observed Markov Decision processes (POMDPs) present challenging mathematical problems with significant applied relevance. It is known that any POMDP can be reduced to a (completely observable) MDP [69], [51], whose states are the posterior state distributions or *beliefs* of the observer; in particular, the belief-MDP is a (fully observed) Markov decision process.

However, the use of belief-MDPs requires one to establish several regularity, continuity and stability (such as filter stability or unique ergodicity) results for arriving at existence, finite memory approximations, robustness, as well as learning theoretic results (see e.g. [65, 39, 40, 48]). While significant progress on the regularity properties of belief-MDPs has been made in the literature, in this paper we will see that further refinements are possible if one does not restrict the analysis to a belief-MDP based formulation. In this paper, we present an alternative view for the study of infinite horizon average-cost or discounted cost optimal control of POMDPs. Our approach will be on a design which is not based on separation / or belief-MDP reduction. We define a Markov chain taking values in an infinite dimensional product space with control actions and the state process causally conditionally independent given the measurement process.

For the controlled case, we provide new sufficient conditions for the existence of optimal control policies for average and discounted cost criteria, and under the latter criterion our analysis will establish a general result on near optimality of finite memory policies. In the control-free case, our analysis leads to sufficient conditions

---

\*Department of Mathematics and Statistics, Queen's University, Kingston, Ontario, Canada, K7L 3N6. Email: yuksel@mast.queensu.ca. This research was partially supported by the Natural Sciences and Engineering Research Council of Canada (NSERC).

for the existence and uniqueness of an invariant probability measure for non-linear filters.

We now present the problem. Consider a stochastic process  $\{X_k, k \in \mathbb{Z}_+\}$ , where each element  $X_k$  takes values in some standard Borel space  $\mathbb{X}$ , with dynamics described by

$$X_{k+1} = F(X_k, U_k, W_k) \quad (1.1)$$

$$Y_k = G(X_k, V_k) \quad (1.2)$$

where  $Y_k$  is an  $\mathbb{Y}$ -valued measurement sequence; we take  $\mathbb{Y}$  also to be some standard Borel space. Suppose further that  $X_0 \sim \mu$ . Here,  $W_k, V_k$  are mutually independent i.i.d. noise processes. This system is subjected to a control/decision process where the control/decision at time  $n$ ,  $U_n$ , incurs a cost  $c(X_n, U_n)$ . The decision maker only has access to the measurement process  $Y_n$  and  $U_n$  causally: An *admissible policy*  $\gamma$  is a sequence of control/decision functions  $\{\gamma_t, t \in \mathbb{Z}_+\}$  such that  $\gamma_t$  is measurable with respect to the  $\sigma$ -algebra generated by the information variables

$$I_t = \{Y_{[0,t]}, U_{[0,t-1]}\}, \quad t \in \mathbb{N}, \quad I_0 = \{Y_0\}.$$

so that

$$U_t = \gamma_t(I_t), \quad t \in \mathbb{Z}_+ \quad (1.3)$$

are the  $\mathbb{U}$ -valued control/decision actions and we use the notation

$$Y_{[0,t]} = \{Y_s, 0 \leq s \leq t\}, \quad U_{[0,t-1]} = \{U_s, 0 \leq s \leq t-1\}.$$

We define  $\Gamma$  to be the set of all such (strong-sense) admissible policies. We emphasize the implicit assumption here that the control policy can also depend on the prior probability measure  $\mu$ .

We assume that all of the random variables are defined on a common probability space  $(\Omega, \mathcal{F}, P)$ . We note that (1.1)-(1.2) can also, equivalently (via stochastic realization results [26, Lemma 1.2] [9, Lemma 3.1]), be represented with transition kernels: the state transition kernel is denoted with  $T$  so that for Borel  $B \subset \mathbb{X}$

$$T(B|x, u) := P(X_1 \in B | X_0 = x, U_0 = u), \quad .$$

We will denote the measurement kernel with  $Q$  so that for Borel  $B \subset \mathbb{Y}$ :

$$Q(B|x) := P(Y_0 \in B | X_0 = x).$$

For (1.1)-(1.2), we are interested in minimizing either the average-cost optimization criterion

$$J(\mu, \gamma) := \limsup_{N \rightarrow \infty} \frac{1}{N} E_\mu^\gamma \left[ \sum_{k=0}^{N-1} c(X_k, U_k) \right] \quad (1.4)$$

or the discounted cost criterion (for some  $\beta \in (0, 1)$ )

$$J(\mu, \gamma) := E_\mu^\gamma \left[ \sum_{k=0}^{\infty} \beta^k c(X_k, U_k) \right] \quad (1.5)$$

over all admissible control policies  $\gamma = \{\gamma_0, \gamma_1, \dots\} \in \Gamma$  with  $X_0 \sim \mu$ . With  $\mathcal{P}(\mathbb{U})$  denoting the set of probability measures on  $\mathbb{U}$  endowed with the weak convergence topology, we will also, when needed, allow for independent randomizations so that  $\gamma_n(I_n)$  is  $\mathcal{P}(\mathbb{U})$ -valued for each realization of  $I_n$ . Here  $c : \mathbb{X} \times \mathbb{U} \rightarrow \mathbb{R}_+$  is the cost function.

We will also consider the control-free case where the system equation (1.1) does not have control dependence; in this case only a decision is to be made at every time stage;  $U$  is present only in the cost expression in (1.4). This important special case has been studied extensively in the theory of non-linear filtering.

**1.1. Literature review and preliminaries.** It is well-known that any POMDP can be reduced to a (completely observable) MDP [69], [51], whose states are the posterior state probabilities, or beliefs, of the observer; that is, the state at time  $t$  is

$$\pi_t(\cdot) := P\{X_t \in \cdot | Y_0, \dots, Y_t, U_0, \dots, U_{t-1}\} \in \mathcal{P}(\mathbb{X}).$$

We call this equivalent MDP the belief-MDP. The belief-MDP has state space  $\mathcal{P}(\mathbb{X})$  and action space  $\mathbb{U}$ . Here,  $\mathcal{P}(\mathbb{X})$  is equipped with the Borel  $\sigma$ -algebra generated by the topology of weak convergence [6]. Since  $\mathbb{X}$  is a Borel space,  $\mathcal{P}(\mathbb{X})$  is metrizable with the Prokhorov metric which makes  $\mathcal{P}(\mathbb{X})$  into a Borel space [49]. The transition probability  $\eta$  of the belief-MDP can be constructed as follows (see also [29]). If we define the measurable function

$$F(\pi, a, y) := Pr\{X_{t+1} \in \cdot | \pi_t = \pi, U_t = a, Y_{t+1} = y\}$$

from  $\mathcal{P}(\mathbb{X}) \times \mathbb{U} \times \mathbb{Y}$  to  $\mathcal{P}(\mathbb{X})$  and the stochastic kernel  $H(\cdot | \pi, u) := Pr\{Y_{t+1} \in \cdot | \pi_t = \pi, U_t = u\}$  on  $\mathbb{Y}$  given  $\mathcal{P}(\mathbb{X}) \times \mathbb{U}$ , then  $\eta$  can be written as

$$\eta(\cdot | \pi, u) = \int_{\mathbb{Y}} \mathbf{1}_{\{F(\pi, u, y) \in \cdot\}} H(dy | \pi, u). \quad (1.6)$$

The one-stage cost function  $c$  of the belief-MDP is given by

$$\tilde{c}(\pi, u) := \int_{\mathbb{X}} c(x, u) \pi(dx). \quad (1.7)$$

In particular, the belief-MDP is a (fully observed) Markov decision process with the components  $(\mathcal{P}(\mathbb{X}), \mathbb{U}, \eta, \tilde{c})$ .

For finite horizon problems and a large class of infinite horizon discounted cost problems it is a standard result that an optimal control policy will use the belief  $\pi_t$  as a sufficient statistic for optimal policies (see [69, 51, 8]).

For the special case without control, the belief process is known as the (non-linear) filter process, and by the discussion above, this itself is a Markov process. The stability properties of such processes has been studied, where the existence of an invariant probability measure for the belief process, as well as the uniqueness of such a measure (i.e., the unique ergodicity property) has been investigated under various conditions, see. e.g. [15]. For the control-free case, [16] provides a comprehensive discussion on both the ergodicity of the filter process as well as filter stability, when the state space is finite but the measurement space is not necessarily so, under the further assumption that the unobserved state process is stationary and ergodic. [46, Theorem 2] and [61, Prop 2.1] assume that the hidden state process is ergodic and the filter is stable (almost surely or in expectation under total variation); these papers

crucially embed the stationary state in the joint process  $(x_k, \pi_k)$  and note that when  $x_k$  is stationary, the Markov chain defined by this process admits an invariant probability measure. The unique ergodicity argument builds on the fact that any two invariant probability measures would have to have the same marginal invariant measure on the state process  $x_k$ , and this leads to a direct argument in [61, Lemma B.1] to relate unique ergodicity to filter stability. In the context of finite state and measurement spaces, another line of work, adopted in [33],[24] and [16], studies the ergodicity and reachability properties of random matrix products, see [34] for a countable space setup. In [24] a reachability condition (across all initial priors) through the approximability of a rank-one matrix of unnormalized product of transition matrices, and in [33] a more restrictive subrectangularity condition, is utilized to establish unique ergodicity. A related argument appears in [59], in a general context. Finally, [16] established that the conditions of [24] are tight; with a different sufficiency proof related to filter stability and convex ordering of measures [58].

For the controlled-setup, [50, 23, 52, 32] study the average-cost control problem under the assumption that the state space is finite; they provide reachability type conditions for the belief kernels. References [10] considers the finite model setup and [60] considers the case with finite-dimensional real-valued state spaces under some restrictive assumptions on the controlled state process leading to strong ergodicity conditions. In a related discussion, [21] studies the existence problem for the average-cost control problem under weak continuity conditions for the controlled kernels. In all of the aforementioned studies, the vanishing discount method is considered.

**Regularity of Belief-MDPs.** As noted earlier, if one wishes to follow the traditional method of reducing a POMDP to a belief-MDP for studying the existence and structure of the optimal policies, it would be important to obtain continuity properties of the belief-MDP so that the standard measurable selection theorems, e.g. [30, Chapter 3] can be invoked to establish the existence of optimal control policies. Building on [35] and [22], we briefly review the weak Feller property of the kernel defined in (1.6) under two different sets of assumptions.

**Assumption 1.**

- (i) *The transition probability  $T(\cdot|x, u)$  is weakly continuous in  $(x, u)$ , i.e., for any  $(x_n, u_n) \rightarrow (x, u)$ ,  $T(\cdot|x_n, u_n) \rightarrow T(\cdot|x, u)$  weakly.*
- (ii) *The observation channel  $Q(\cdot|x)$  is continuous in total variation, i.e., for any  $x_n \rightarrow x$ ,  $Q(\cdot|x_n) \rightarrow Q(\cdot|x)$  in total variation.*

**Assumption 2.** *The transition probability  $T(\cdot|x, u)$  is continuous in total variation in  $(x, u)$ , i.e., for any  $(x_n, u_n) \rightarrow (x, u)$ ,  $T(\cdot|x_n, u_n) \rightarrow T(\cdot|x, u)$  in total variation.*

**THEOREM 1.1.**

- (i) [22] *Under Assumption 1, the transition kernel  $\eta(F(\pi, u, Y_1) \in \cdot | \pi, u)$  of the filter process is weakly continuous in  $(\pi, u)$ .*
- (ii) [35] *Under Assumption 2, the transition kernel  $\eta(F(\pi, u, Y_1) \in \cdot | \pi, u)$  of the filter process is weakly continuous in  $(\pi, u)$ .*

If the cost function  $c(x, u)$  is continuous and bounded, an application of the dominated convergence theorem implies that  $\tilde{c}(z, u) = E^z[c(X, u)]$  is also continuous and bounded.

In the uncontrolled setting, [5] and [15] have established similar weak continuity conditions (i.e., the weak-Feller property) of the non-linear filter process (i.e., the belief process) in continuous time and discrete time, respectively, where Assumption 1 is present for an additive noise measurement model.

The convex analytic approach [45, 11] is a powerful approach to the optimization of infinite-horizon problems. It is particularly effective in proving results on the optimality of stationary (and possibly randomized stationary) policies, through an infinite-dimensional linear program for constrained optimization problems and infinite horizon average cost optimization problems. For the average cost criterion, recall that via (1.6)-(1.7) under the belief-MDP reduction, we are interested in the minimization:

$$\inf_{\gamma \in \Gamma} \limsup_{N \rightarrow \infty} \frac{1}{N} E_{\mu}^{\gamma} \left[ \sum_{t=0}^{N-1} \tilde{c}(\pi_t, u_t) \right], \quad (1.8)$$

where  $E_{x_0}^{\gamma}[\cdot]$  denotes the expectation over all sample paths with initial state given by  $x_0$  under the admissible policy  $\gamma$ .

**Assumption 3.**

- (i) The  $\mathbb{R}_+$ -valued one-stage cost function  $c$  is bounded and continuous.
- (ii) Assumption 1 or 2 holds (this leads to the belief-MDP  $\eta$  to be weakly continuous by Theorem 1.1).
- (iii)  $\mathbb{X}$  and  $\mathbb{U}$  are compact.

**THEOREM 1.2.** *Under Assumption 3, there exists an optimal invariant measure and an associated optimal control policy, for either average cost (1.4) or for the discounted cost (1.5) criteria.*

**Proof.** For the average cost criterion, the proof follows from the convex analytic method, studied by Borkar in the weakly continuous case [11]; for completeness, and its use later in the paper, a proof is presented in Section A.1. For the discounted cost criterion, the result follows from a standard verification and measurable selection analysis on the discounted cost optimality equation [30].  $\square$

While this result presents sufficient findings on the existence of optimal control policies, the approach has a number of limitations.

- (i) [Fully observed MDPs viewed as a special case of POMDPs] Consider the case with the state transition kernel  $T$  being weakly continuous and where the measurements satisfy  $y_t = x_t$ , that is, with full state information, in which case the measurement kernel is  $Q(dy|x) = \delta_x(dy)$ , and thus the system does not satisfy Assumption 1 or 2 (as the channel is only weakly continuous but not total variation continuous). On the other hand, this model, which is a fully observed setup, has been studied well and leads to well-known existence (and further, structural and approximation) results. Therefore, we observe that one should be able to relax the conditions further, which we will indeed find to be the case.
- (ii) For the average cost criterion, Assumption 3 provides weaker conditions when compared with the efforts in the literature that typically have utilized the vanishing discount approach: [10] considers the finite model setup and [60] consider restrictive assumptions on the controlled state process, and [21] which in the POMDP case would require yet to be established rates of convergence conditions to invariant measures (needed for uniform boundedness properties of relative discounted cost value functions as the discount parameter approaches unity). We will see that further relaxations can be provided.
- (iii) There is a subtle question of how to select the initial measure, that is the question of achieving the optimal cost under a given initial belief measure and whether the support of an invariant probability measure attracts the belief process from a given initial condition. For an interesting counterexample,

see [37, Example 2.3]. For belief MDPs, there are few results on unique ergodicity properties, therefore the impact of the initial priors is an open problem under the belief-separated-approach. In our case, the infinite dimensional formulation to be presented leads to a flexibility on how to select the initial prior which we show to lead to global optimality under a mild and testable absolute continuity condition (this approach does not seem to easily carry over to the belief-separated-approach).

- (iv) Related to (iii), the unique ergodicity problem for the controlled or uncontrolled non-linear filtering process entails open problems. An alternative approach to what currently exists in the literature is needed, especially for the controlled setup. For the control-free case, building on [46, Theorem 2] and [61, Prop 2.1], it can be shown that almost sure filter stability in the total variation sense (or weak merging sense, with the arguments tailored to this case) and the uniqueness of an invariant probability measure on the hidden state process leads to unique ergodicity of the filter process. We will see that a complementary condition for the control-free case can be established (unique ergodicity of the measurement process and a measurability condition related to filter stability also leads to unique ergodicity).

**1.2. Statement of main results and contributions.** The paper makes the following contributions.

- (i) We present an approach of defining the state as an infinite-dimensional controlled Markov chain where the control policies satisfy conditional independence between the state and control actions given the information. For the controlled case, we provide new sufficient conditions for the existence of optimal control policies which turn out to be stationary in the history variables. In particular, while in the belief-MDP reduction of POMDPs, weak Feller condition requirement imposes total variation continuity on either the system kernel or the measurement kernel, with the approach of this paper only weak continuity of both the transition kernel and the measurement kernel is sufficient together with a stability condition (Theorem 3.3). For the average cost criterion, the paper provides a general approach on how to generate initial priors and beliefs so that optimal performance can be attained. The infinite dimensional formulation presents a natural flexibility in initialization, which can be used to arrive at optimal performance under a mild absolute continuity condition, see Theorem 3.5 (which does not seem to be lenient under the standard belief-MDP approach).
- (ii) For the discounted cost criterion, more relaxed existence conditions will be presented in Section 4. We first establish the existence of a stationary optimal policy. We then establish near optimality of finite window policies via a direct argument involving near optimality of quantized approximations for MDPs under weak Feller continuity (see Theorem 4.1), where finite truncations of memory can be viewed as (uniform) quantizations of infinite memory (which is the state definition adopted for the discounted cost criterion by Theorem 2.5) with a uniform diameter in each finite window restriction under the product metric. Building on recent results on near-optimality of quantized policies for weak Feller MDPs [55], near optimality of finite memory policies follows, complementing and generalizing recent works on the subject [39, 40]. This also facilitates reinforcement learning theoretic methods for POMDPs, which is discussed in further detail in the paper.

- (iii) Via our approach, further results on existence and uniqueness of invariant probability measures for control-free non-linear filter processes will be established in Section 5. In particular, we will see that unique ergodicity of the measurement process and a measurability condition related to filter stability leads to unique ergodicity (this complements, with an alternative argument, several results in the literature).

## 2. Another Look at POMDPs: Infinite Dimensional Markov Decision Process Formulation.

**2.1. Controlled-Markov construction with infinite-memory state space for the average cost criterion.** To facilitate the solution approach presented in this paper, we first assume that the measurements and the control actions have been taking place since  $-\infty$ . Later on we will motivate and justify this approach.

Let  $\mathbb{Y}^{\mathbb{Z}^-}$  be the one-sided product space consisting of elements of the form

$$y = \{\cdots, y_n, \cdots, y_{-2}, y_{-1}, y_0\}$$

with  $y_k \in \mathbb{Y}$ . We endow  $\mathbb{Y}^{\mathbb{Z}^-}$  with the product topology; this makes  $\mathbb{Y}^{\mathbb{Z}^-}$  a metric space, which is complete and separable. Likewise, we will view  $U_{(-\infty, -1]}$  also as a  $\mathbb{U}^{\mathbb{Z}^-}$ -valued random variable.

A related view, on using the infinite past as a Markov process, was presented in [66] to establish stochastic stability of control-free dynamical systems driven by noise processes which are not necessarily independent and identically distributed, but which are stationary. A further related approach was adopted in [1]; see also [28] (as well as [7] and [67] as discussed earlier).

THEOREM 2.1. *With*

$$Z_k = (Y_{(-\infty, k]}, U_{(-\infty, k-1]}, X_k),$$

the pair  $(Z_k, U_k)$  is a controlled Markov chain where  $Z_k$  is  $\mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U}^{\mathbb{Z}^-} \times \mathbb{X}$  valued.

**Proof.** Let  $A = \prod_{k=-\infty}^0 A_k^y$  be an open cylinder set in  $\mathbb{Y}^{\mathbb{Z}^-}$  and  $A^u = \prod_{k=-\infty}^0 A_k^u$  be an open cylinder set in  $\mathbb{U}^{\mathbb{Z}^-}$  and let  $B \in \mathbb{X}$  be a Borel set:

$$\begin{aligned} & P\left(Z_{t+1} \in (A^y \times A^u \times B) \mid Z_t = z, U_t = u, Z_{[0, t-1]} = z_{[0, t-1]}, U_{[0, t-1]} = u_{[0, t-1]}\right) \\ &= \int Q(y_{t+1} \in A_0^y \mid x_{t+1}) T(x_{t+1} \in B \mid x_t = x, u_t = u) \\ & \quad \times P(y_{(-\infty, t]} \in \prod_{k=-\infty}^0 A_k^y \mid y_{(-\infty, t]}) P(u_{(-\infty, t]} \in \prod_{k=-\infty}^0 A_k^u \mid u_{(-\infty, t]}) \\ &= \int Q(y_{t+1} \in A_0^y \mid x_{t+1}) T(x_{t+1} \in B \mid x_t = x, u_t = u) 1_{\{y_{(-\infty, t]} \in \prod_{k=-\infty}^0 A_k^y, u_{(-\infty, t]} \in \prod_{k=-\infty}^0 A_k^u\}} \\ & \tag{2.1} \end{aligned}$$

$$=: \mathbb{P}(A^y \times A^u \times B \mid Z_t = z, U_t = u) \tag{2.2}$$

where for a Borel  $C$ ,  $Q(y \in C \mid x) := P(G(X_0, V_0) \in C \mid X_0 = x)$  by (1.2) and  $T$  is the transition kernel defined by (1.1).  $\square$

In the above, we define  $\mathbb{P}$  to be the transition kernel for this Markov chain.

DEFINITION 2.2 (Weak-Feller Condition for  $\mathbb{P}$ ).  $\mathbb{P}(dz_{t+1} \mid z_t, u_t)$  is weak-Feller if for every continuous and bounded  $f \in C_b(\mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U}^{\mathbb{Z}^-} \times \mathbb{X})$ :

$$\int f(z_{t+1}) \mathbb{P}(dz_{t+1} \mid z_t = z, u_t = u)$$

is continuous in  $(z, u)$ .

The following will be invoked often.

**Assumption 4.**  $Q(\cdot|x)$  is weakly continuous in  $x$ , and  $T(\cdot|x, u)$  is weakly continuous in  $x, u$ .

LEMMA 2.3. Under Assumption 4,  $\mathbb{P}$  is weak-Feller.

**Proof.** First note that by (2.1) we have

$$\mathbb{P}(dz_{t+1}|z_t, u_t) = Q(dy_{t+1}|x_{t+1})T(dx_{t+1}|x_t, u_t)P(dy_{(-\infty, t]}|y_{(-\infty, t]})P(du_{(-\infty, t]}|u_{(-\infty, t]})$$

where  $P(y_{(-\infty, t]} \in \cdot | y_{(-\infty, t]}) = \delta_{y_{(-\infty, t]}}(\cdot)$  is the Dirac measure (as in (2.1)), with the same holding for  $P(du_{(-\infty, t]}|u_{(-\infty, t]})$ .

We wish to show that the following is continuous in  $x_t, y_{(-\infty, t]}, u_{(-\infty, t-1]}, u_t$ , for continuous and bounded  $f$ :

$$\begin{aligned} & \int f\left(x_{t+1}, y_{(-\infty, t+1]}, u_{(-\infty, t]}\right) \mathbb{P}(dx_{t+1}, dy_{(-\infty, t]}+1, du_{(-\infty, t-1]}|x_t, y_{(-\infty, t]}, u_{(-\infty, t-1]}, u_t) \\ &= \int f\left(x_{t+1}, y_{(-\infty, t+1]}, u_{(-\infty, t]}\right) Q(dy_{t+1}|x_{t+1})T(dx_{t+1}|x_t, u_t) \\ & \quad \times P(dy_{(-\infty, t]}|y_{(-\infty, t]})P(du_{(-\infty, t]}|u_{(-\infty, t]}) \\ &= \int \int_{x_{t+1}} \left( \left( \int_{y_{t+1}} f(x_{t+1}, y_{(-\infty, t+1]}, u_{(-\infty, t]}) Q(dy_{t+1}|x_{t+1}) \right) T(dx_{t+1}|x_t, u_t) \right) \\ & \quad \times P(dy_{(-\infty, t]}|y_{(-\infty, t]})P(du_{(-\infty, t]}|u_{(-\infty, t]}) \end{aligned}$$

The expression

$$g(x_{t+1}, y_{(-\infty, t+1]}, u_{(-\infty, t]}) := \left( \int_{y_{t+1}} f(x_{t+1}, y_{(-\infty, t+1]}, u_{(-\infty, t]}) Q(dy_{t+1}|x_{t+1}) \right)$$

is continuous by a generalized dominated convergence theorem for varying measures under *continuous-convergence* [44, Thm. 3.5] (see also [57, Thm. 3.3] for a more restricted version). Since  $g$  is continuous in  $x_{t+1}$ ,

$$h(x_t, u_t, y_{(-\infty, t]}, u_{(-\infty, t]}) := \int_{x_{t+1}} \left( g(x_{t+1}, y_{(-\infty, t]}, u_{(-\infty, t]}) T(dx_{t+1}|x_t, u_t) \right)$$

is also continuous in its parameters. Finally, again by [44, Thm. 3.5],

$$\int h(x_t, u_t, y_{(-\infty, t]}, u_{(-\infty, t]}) P(dy_{(-\infty, t]}|y_{(-\infty, t]})P(du_{(-\infty, t]}|u_{(-\infty, t]})$$

is continuous in  $x_t, y_{(-\infty, t]}, u_{(-\infty, t-1]}, u_t$ .  $\square$

The above completes the description of the controlled Markov model, that we will utilize for the average cost problem.

**2.2. Key technical assumptions on continuity and stability.** We will occasionally invoke a subset of the following assumptions on stationarity, stability, and continuity. We will see, in Appendix B, that these are related to filter stability [17].

**Assumption 5.** [A Continuity Condition] The stochastic kernel  $P(dx_k|y_{(-\infty, k]}, u_{(-\infty, k-1]})$  is weakly continuous, that is, for every continuous and bounded  $g : \mathbb{X} \rightarrow \mathbb{R}$ ,

$$\int g(x) \mathbb{P}(X_k \in dx | y_{(-\infty, k]}, u_{(-\infty, k-1]})$$



is continuous in  $(y_{(-\infty, k]}, u_{(-\infty, k-1]})$  (under the product topology on  $\mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U}^{\mathbb{Z}^-}$ ).

It will be shown in the appendix that a *uniform filter stability condition*, such as [27, Corollary 4.2] (via a Hilbert metric approach) as well as [47, Corollary 3.7] (via Dobrushin's coefficient method), implies Assumption 5 for the case where the measurement and action spaces are finite. Further discussion is available in Appendix B. Our next assumption is the following.

**Assumption 6.** [A Stationarity / Stability Condition] For every Borel  $B$ , and with  $y = (\dots, y_{-2}, y_{-1}, y_0)$ ,  $u = (\dots, u_{-2}, u_{-1}, u_0)$ , under any control policy, for almost every  $(y, u)$ , we have that

$$P(X_t \in B | Y_{(-\infty, t]} = y, U_{(-\infty, t-1]} = u) = P(X_{t+1} \in B | Y_{(-\infty, t+1]} = y, U_{(-\infty, t]} = u)$$

This assumption is also implied by an almost sure filter stability condition. In fact, [27, Corollary 4.2] also implies the next, and final, assumption:  $P\left(X_t \in \cdot \mid \bigcap_{n=1}^{\infty} \sigma(Y_{(-\infty, t]}, U_{(-\infty, t-1]}) \vee \sigma(\pi_{(-\infty, -n]})\right)$  is  $\sigma(I_t)$ -measurable in the sense that  $P$  a.s.

$$\begin{aligned} P\left(X_t \in \cdot \mid \bigcap_{n=1}^{\infty} \sigma(Y_{(-\infty, t]}, U_{(-\infty, t-1]}) \vee \sigma(\pi_{(-\infty, -n]})\right) \\ = P(X_t \in \cdot | \sigma(Y_{(-\infty, t]}, U_{(-\infty, t-1]})) \end{aligned} \quad (2.3)$$

Let  $\pi_t$  be the conditional probability on the state, given a (distant) prior at time  $-n \in \mathbb{Z}_-$ , and the information since then (or, equivalently, the information even prior to  $-n$ ): For all Borel  $A \subset \mathbb{X}$

$$\pi_t(A) = E[1_{X_t \in A} | \sigma(Y_{(-\infty, t]}, U_{(-\infty, t-1]}) \vee \sigma(\pi_{(-\infty, -n]})]$$

In view of the above, we state the following assumption.

**Assumption 7.** [A Filter Stability Condition]  $\pi_t$  is  $\sigma(I_t)$  measurable where  $I_t = (Y_{(-\infty, t]}, U_{(-\infty, t-1]})$ . In particular, there exists a measurable  $g$  such that  $g(y_{(-\infty, t]}, u_{(-\infty, t-1]})(A) = \pi_t(A)$  for every Borel  $A$ .

LEMMA 2.4. Condition (2.3) implies Assumption 7.

**Proof.** Observe that  $\pi_t(A) = E[1_{X_t \in A} | \sigma(Y_{(-\infty, t]}, U_{(-\infty, t-1]}) \vee \sigma(\pi_{(-\infty, -n]})]$  for every  $n > 0$  and thus

$$\pi_t(A) = \lim_{n \rightarrow \infty} E[1_{X_t \in A} | \sigma(Y_{(-\infty, t]}, U_{(-\infty, t-1]}) \vee \sigma(\pi_{(-\infty, -n]})].$$

Now,  $E[1_{X_t \in A} | \sigma(Y_{(-\infty, t]}, U_{(-\infty, t-1]}) \vee \sigma(\pi_{(-\infty, -n]})]$  is a bounded backwards martingale sequence with respect to the decreasing filtration  $\sigma(Y_{(-\infty, t]}, U_{(-\infty, t-1]}) \vee \sigma(\pi_{(-\infty, -n]})$ ,  $n \in \mathbb{N}$  and by the backwards martingale theorem [19], the above limit will converge to

$$P\left(X_t \in A \mid \bigcap_{n=1}^{\infty} \sigma(Y_{(-\infty, t]}, U_{(-\infty, t-1]}) \vee \sigma(\pi_{(-\infty, -n]})\right)$$

This then implies, via (2.3),  $\pi_t(A)$  is  $\sigma(I_t)$ -measurable for any given  $A$ .

Recall the following which builds on Theorem 2.1 of Dubins and Freedman [18] and Proposition 7.25 in Bertsekas and Shreve [4]: Let  $\mathbb{S}$  be a Polish space,  $\mathcal{P}(\mathbb{S})$  be the set of

probability measures under the weak convergence topology and  $(M, \mathcal{M})$  be a measurable space. A function  $F : (M, \mathcal{M}) \rightarrow \mathcal{P}(\mathbb{S})$  is measurable on  $\mathcal{M}$  if for all  $B \in \mathcal{B}(\mathbb{S})$   $(F(\cdot))(B) : \mathcal{M} \rightarrow \mathbb{R}$  is measurable on  $\mathcal{M}$ , that is for every  $B \in \mathcal{B}(\mathbb{S})$ ,  $(F(\pi))(B)$  is a measurable function when viewed as a function from  $M$  to  $\mathbb{R}$ . See also [4, Proposition 7.26]. By this result, it follows that  $\pi_t$  is measurable on  $\bigcap_{n=1}^{\infty} \sigma(Y_{(-\infty, t]}, U_{(-\infty, t-1]}) \vee \sigma(\pi_{(-\infty, -n]})$ . But by condition (2.3), it is also  $\sigma(I_t)$  measurable, and as the considered spaces are standard Borel, a functional representation via the measurable function  $g$  follows.  $\square$

Further analysis and sufficient conditions for Assumptions 5, 6 and 7 are presented in Appendix B. We note that (2.3) is essentially a filter stability condition: Indeed, the assumption above, in the control-free setup, is related to the statement in [16, Theorem 3.1(2)], building on [43].

**2.3. Controlled-Markov construction with infinite-memory state space for the discounted cost criterion.** Given the construction and stability properties, for the discounted cost criterion, towards arriving at approximate optimality results using finite window policies, we will present an alternative controlled Markov model. The proof follows from identical arguments presented in Section 2.1, but here we apriori impose Assumption 5 and Assumption 7.

THEOREM 2.5.

(i) With

$$S_k = (Y_{(-\infty, k]}, U_{(-\infty, k-1]}),$$

the pair  $(S_k, U_k)$  is a controlled Markov chain where  $S_k$  is  $\mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U}^{\mathbb{Z}^-}$  valued.  
(ii) Under Assumption 7, we have that for all  $u \in \mathbb{U}$

$$E \left[ c(X_k, u) \middle| (Y_{(-\infty, k]}, U_{(-\infty, k-1]}) = s \right] = \bar{c}(s, u),$$

for some measurable  $\bar{c} : \mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U}^{\mathbb{Z}^-} \times \mathbb{U} \rightarrow \mathbb{R}$ .

(iii) Under Assumption 5,  $\bar{c}$  is continuous and bounded, when  $c$  is bounded continuous.

(iv) The kernel  $P(z_{t+1} \in \cdot | Z_t = z, U_t = u)$  is weak Feller under Assumptions 4 and 5.

The final result, (iv), in the above follows by considering  $P(Y_t \in dy | Z_t = z, U_t = u) = \int Q(dy|x)P(X_t \in dx | Z_t = z, U_t = u)$ , and the weak continuity of each of the kernels in the integration and generalized weak convergence [44, Thm. 3.5] as used earlier in the proof of Lemma 2.3; (iii) follows from Assumption 5 by definition. (i) follows by construction, and (ii) directly by Assumption 7.

As a result, under Assumptions 7 and 5, we have an equivalent controlled Markov model with cost  $\bar{c} : (\mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U}^{\mathbb{Z}^-}) \times \mathbb{U} \rightarrow \mathbb{R}_+$ ,  $\mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U}^{\mathbb{Z}^-}$ -valued state  $s_k$ , and transition kernel  $P(Z_{t+1} \in \cdot | Z_t = z, U_t = u)$ .

**2.4. Conditionally state independent control policies.** Once we have established controlled Markov models, we now discuss the classes of control policies considered under the infinite dimensional controlled Markov model presented. We recall that in the theory of stochastic control, to facilitate stochastic analysis for arriving at existence, structural or approximation results (e.g. via continuity-compactness properties), the set of control policies may be enlarged. This is often referred to as a *relaxation* of control policies. Relaxations have been very effective in optimal control,

with a very prominent example being Young measures in deterministic optimal control [64]. These allow one to use topologies on the sets of probability measures to study existence, optimality, and structural results (especially that of weak convergence), rather than working with the space of measurable functions only (whose compactness conditions under an appropriate metric would be too restrictive). A key aspect of such relaxations is that any relaxation should not allow for optimal expected cost values to be improved; they should only be means to facilitate stochastic analysis. We can classify some policies and various relaxations as follows:

(i) *Wide sense admissible policies* introduced by Fleming and Pardoux [25] and prominently used to establish the existence of optimal solutions for partially observed stochastic control problems. Borkar [10, 13, 12] (see also Borkar and Budhiraja [14]) have utilized these policies for a coupling/simulation method to arrive at optimality results for average cost partially observed stochastic control problems. The approach here is to first apply a Girsanov type (see Borkar [10, 13] for discrete-time models and Witsenhausen [62] for decentralized stochastic control where the change of measure argument leads to *static reduction*) transformation to decouple the measurements from the system via an absolute continuity condition of measurement variables conditioned on the state, with respect to some reference measure; and then use independence properties: In the discrete-time case,  $\{Y_n\}$  is i.i.d. and independent of  $X_0$  and the system noise  $\{W_n\}$ , and  $\{U_0, \dots, U_n, Y_0, \dots, Y_n\}$  is independent of  $\{W_n\}$ ,  $X_0$ , and  $\{Y_m, m > n\}$ , for all  $n$ .

(ii) Policies defined by conditional independence (as in the dynamic programming formulation [67] for decentralized stochastic control). In the context of our setup here, the actions are those that satisfy  $U_n$  being conditionally independent of the past  $(X_0, W_{[0,n]}, V_{[0,n]})$  given the local information  $\{Y_{[0,n]}, U_{[0,n-1]}\}$ :

$$U_n \leftrightarrow \{Y_{[0,n]}, U_{[0,n-1]}\} \leftrightarrow \{X_0, W_{[0,\dots]}, V_{[0,\dots]}\} \quad (2.4)$$

We note that, here an absolute continuity condition required for the static reduction or Girsanov-type measure transformation is not necessary a priori. We will call such policies *Conditionally-Exogenous Variable-Independent Policies* since the conditional independence holds between action, information, and the exogenous random variables in the system. In our paper, we will consider a further refinement (to be called *Conditionally-State-Independent Policies*) which satisfies instead of (2.4)

$$U_n \leftrightarrow \{Y_{[0,n]}, U_{[0,n-1]}\} \leftrightarrow X_n \quad (2.5)$$

to facilitate the infinite horizon analysis.

If we have independent static reduction, with measurements  $\tilde{Y}_n$ , by considering the method for static measurements with expanding information structure (this is the information structure that one would obtain for a POMDP under the absolute continuity conditions) in [68, Theorem 5.6] or [54, Theorem 4.7] by expressing all the cost-relevant uncertainty in terms of  $\tilde{Y}_n$ , we have the following version of (2.4)

$$U_n \leftrightarrow \{\tilde{Y}_{[0,n]}, U_{[0,n-1]}\} \leftrightarrow \{\tilde{Y}_{[n+1,\dots]}\} \quad (2.6)$$

If the process continues on since the indefinite past, we write (2.5) with

$$U_n \leftrightarrow \{Y_{(-\infty,n]}, U_{(-\infty,n-1]}\} \leftrightarrow X_n \quad (2.7)$$

### 3. Existence Results for Optimal Policies: Average Cost Criterion.

**3.1. On the existence of an optimal stationary policy.** We first present a supporting result in the following, which will be crucial in our analysis to follow.

LEMMA 3.1. a) Let  $Y_n^1, Y_n^2, U_n^2$  be a sequence of random variables that satisfies for every  $n$ , the conditional independence property:

$$Y_n^1 \leftrightarrow Y_n^2 \leftrightarrow U_n^2,$$

with joint measure  $P_n$ , where the marginal on  $(Y_n^1, Y_n^2)$  is fixed throughout the sequence. In this case, if  $P_n \rightarrow P$  weakly, the limit measure  $P$  also satisfies

$$Y^1 \leftrightarrow Y^2 \leftrightarrow U^2$$

b) Let  $Y_n^1, Y_n^2, U_n^2$  be a sequence of random variables that satisfies for every  $n$ :

$$Y_n^1 \leftrightarrow Y_n^2 \leftrightarrow U_n^2$$

with measure  $P_n$  where the conditional probability  $P_n(Y_n^1 \in dy^1 | Y_n^2 = y^2) = \kappa(dy^1 | y^2)$  is fixed throughout the sequence and that  $\kappa(dy^1 | y^2)$  is a weakly continuous kernel (i.e.  $y^2 \mapsto \int f(y^1) \kappa(dy^1 | y^2)$  is continuous and bounded for every bounded continuous  $f$ ). In this case, if  $P_n \rightarrow P$  weakly, then, the limit measure also satisfies

$$Y^1 \leftrightarrow Y^2 \leftrightarrow U^2$$

**Proof.** See Section A.2.  $\square$

We now present a key intermediate result.

LEMMA 3.2. Let  $\mathbb{X}, \mathbb{Y}, \mathbb{U}$  be compact. Under Assumptions 4, 6 and 5, the set of invariant occupation measures

$$\begin{aligned} \mathcal{G} = \{v \in \mathcal{P}(\mathbb{X} \times \mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U}^{\mathbb{Z}^-}) : \\ v(B \times \mathbb{U}) = \int_{z,u} \mathbb{P}(z_{t+1} \in B | z, u) v(dz, du), \quad B \in \mathcal{B}(\mathbb{X} \times \mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U}^{\mathbb{Z}^-})\} \end{aligned}$$

that simultaneously satisfy the Markov chain

$$X_0 \leftrightarrow (U_{(-\infty, -1]}, Y_{(-\infty, 0]}) \leftrightarrow U_0,$$

or, equivalently, that belong to

$$\begin{aligned} \mathcal{H} = \{v \in \mathcal{P}(\mathbb{X} \times \mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U}^{\mathbb{Z}^-}) : \\ v(dx_0, dy_{(-\infty, 0]}, du_{(-\infty, -1]}, du_0) \\ = v(dx_0, dy_{(-\infty, 0]}, du_{(-\infty, -1]}) P(du_0 | dy_{(-\infty, 0]}, du_{(-\infty, -1]})\}, \quad (3.1) \end{aligned}$$

is a weakly compact set.

**Proof.** We note that under weak continuity of the transition kernel for  $\mathbb{P}$ ,  $\mathcal{G}$  is a closed set under weak convergence. By Lemma 3.1(b), as the kernel

$$P(dx_t | y_{(-\infty, t]}, u_{(-\infty, t-1]})$$

is weakly continuous (by Assumption 5) and time-invariant (by Assumption 6),  $\mathcal{H}$  is also closed. Since the state space is compact, the set of probability measures on  $(\mathbb{X} \times \mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U})$  is tight. The result follows since a closed subset of a tight set is weakly compact.  $\square$

Building on the above, we are able to state the following:

**THEOREM 3.3.** *Let  $\mathbb{X}, \mathbb{Y}, \mathbb{U}$  be compact and Assumptions 4, 6 and 5 hold. Then an optimal control policy exists. This policy is stationary and can be written as:*

$$U_t = \gamma(Y_{(-\infty, t]}, U_{(-\infty, t-1]}, R_t),$$

where  $R_t$  is an i.i.d. sequence.

**Proof.** See Section A.3.  $\square$

**Remark 3.1.** *The above is a relaxation on the known results where one needs to reduce the POMDP to a belief-MDP which is weak Feller. As noted earlier, the weak Feller conditions require total variation continuity on either the system kernel or the measurement kernel. Only weak continuity of both the transition kernel and the measurement kernel is needed for the result above, though filter stability is also imposed.*

**3.2. On the initial state distribution realization for an optimal stationary policy.** In the above, there is an important operational question: The actual system is a one-sided process with  $X_0 \sim \mu$  being the starting variable. To address this issue, in the following we explain that we can randomly generate the past and use the past realizations to apply optimal control. Under some conditions, the average cost will converge to the optimal cost.

Our strategy on the initialization is as follows: (i) We will first show that an optimal invariant measure can be taken to be ergodic (i.e., an invariant measure which cannot be expressed as a convex combination of multiple invariant measures). (ii) Then we will construct an initialization which is in the attractor set of this optimal invariant measure, under a mild absolute continuity condition. We will generate the measurement and control actions  $Y_{(-\infty, -1]}, U_{(-\infty, -1]}$  according to a stationary measure, and then  $X_0 \sim \mu, Y_0$  accordingly, and continue with the process according to the optimal policy under Theorem 3.3.

**LEMMA 3.4.** *Without any loss of optimality, an optimal invariant occupation measure can be assumed to be ergodic.*

**Proof.** Let  $\nu$  be an optimal occupation measure, which leads to a policy  $\gamma$ . Under  $\gamma$ , the *state* process

$$(Y_{(-\infty, k]}, U_{(-\infty, k-1]}, X_k),$$

is a Markov chain with the marginal of the invariant measure  $\nu$  on this state being invariant. By an ergodic decomposition theorem, every stationary measure can be expressed as a convex combination of ergodic invariant measures with disjoint supports: let  $\nu$  be expressed as a convex combination of measures  $\nu_\beta$ , parametrized by  $\beta$  (see Theorem C.1 for some related discussion), where each of them are also *invariant* and *ergodic*. Now, a question is whether each of the  $\nu_\beta$  measures satisfies (3.1) and (3.1). The first holds by the invariance of each of these measures. The latter holds by definition: conditioned on the support of each  $\nu_\beta$ ,  $\nu$  satisfies (3.1) (note that the control policy is fixed) and hence this holds also for  $\nu_\beta$ .

Since the cost in (A.14):

$$\inf_{v \in \mathcal{G} \cap \mathcal{H}} \int v(dx, dy_{(-\infty, 0]}, du_{(-\infty, -1]}, du) c(x, u) \quad (3.2)$$

is linear in the space of (signed) measures, we conclude that without any loss we can take  $\nu$  to be from the set of ergodic invariant measures.  $\square$

The above then completes our realization result:

**THEOREM 3.5.** *Let Assumptions 5 and 6 hold. Let  $U_t = \gamma(Y_{(-\infty,t]}, U_{(-\infty,t-1]}, R_t)$  be an optimal stationary policy with  $R_t$  being an i.i.d. noise process. Let*

$$g(y_{(-\infty,k]}, u_{(-\infty,k-1]}, x_k) := E[c(x_k, \gamma(y_{(-\infty,k]}, u_{(-\infty,k-1]}, R_k))]$$

where the expectation is over  $R_k$ . Let  $\bar{P}$  denote an (invariant) process measure on

$$(Y_{(-\infty,k-1]}, U_{(-\infty,k-1]}, Y_k, X_k),$$

and let  $P_0$  be the marginal of the invariant measure on  $\{Y_{(-\infty,-1]}, U_{(-\infty,-1]}\}$  and  $\mu$  be the prior measure on  $X_0$  as imposed (i.e., given in the problem statement) on the controller/decision maker. If  $P_0 \times \mu \ll \bar{P}$  (note that the distribution on  $X_0$  specifies that on  $Y_0$ ); that is, absolute continuity holds when the (past history) initialization is independent of the state process initialization, then

$$\begin{aligned} & \lim_{T \rightarrow \infty} \frac{1}{T} E_{P_0 \times \mu} \left[ \sum_{k=0}^{T-1} c(X_k, \gamma(Y_{(-\infty,k]}, U_{(-\infty,k-1]}, R_k)) \right] \\ &= \int g(y_{(-\infty,0]}, u_{(-\infty,-1]}, x_0) \bar{Q}(dy_{(-\infty,0]}, du_{(-\infty,-1]}, x_0), \end{aligned} \quad (3.3)$$

for some invariant  $\bar{Q}$ , which would necessarily be  $\bar{P}$  under the assumed ergodicity of  $\bar{P}$  via Lemma 3.4, in which case the optimal control/decision cost will be attained.

**Remark 3.2.** *Note that, in the above, with  $\bar{P}$  and  $P_0$  both stationary, we have that  $\bar{P}$  decomposes as  $P_0(dy_{(-\infty,-1]}, du_{(-\infty,-1]})\bar{P}(dx_0|y_{(-\infty,-1]}, u_{(-\infty,-1]})$ . A sufficient condition for the aforementioned absolute continuity condition then is,  $\bar{P}$  a.e.,*

$$\mu(dx_0) \ll \bar{P}(dx_0|dy_{(-\infty,-1]}, du_{(-\infty,-1]})$$

**Remark 3.3.** *Observe that in the analysis above we constructed  $y_{(-\infty,-1]}, u_{(-\infty,-1]}$  and not  $y_{(-\infty,0]},$  since  $\mu$  on  $X_0$  induces directly a measure on  $Y_0$ . Note that with  $P_0 \times \mu$  viewed as an initial measure, when the process evolves, the evolution is consistent with the actual realizations under the true measure. That is, the initialization does not affect the evolution of the measure. The reason is that the process  $X_0$  determines the probability of the future events, independent of the history initialization via:*

$$\mathbb{P}(dz_{t+1}|z_t, u_t) = Q(dy_{t+1}|x_{t+1})P(dx_{t+1}|x_t, u_t)P(dy_{(-\infty,t]}|y_{(-\infty,t]})P(du_{(-\infty,t]}|u_{(-\infty,t]}).$$

**4. Discounted Cost Criterion: Refined Existence Results and Near Optimality of Finite Window/Memory Policies.** In this section, we study the discounted cost setup. We will see a particular utility in this criterion: near optimality of finite window policies.

For the discounted criterion, we will restrict our control policies to those that are strict-sense admissible. We could also consider the relaxed framework, however the analysis for the discounted criterion setup will be seen to be more direct. Compared with the average cost criterion, we are able to utilize a verification theorem directly following the discounted cost optimality equation (DCOE).

Let, as in Theorem 2.5,  $s_k = (y_{(-\infty, k]}, \tilde{u}_{(-\infty, k-1]}) \in \mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U}^{\mathbb{Z}^-}$ . Now, under strict-sense policies, with the assumptions on weak-Feller continuity and bounded continuous cost function, if there exists a solution to

$$J_\beta(s) = \min_{u_k \in \mathbb{U}} E \left[ c(x_k, u_k) + \beta J_{k+1}(S_{k+1}) | S_k = s, U_k = u \right],$$

then we can declare this policy to be optimal using a standard verification argument [30]. As before, we assume that  $\mathbb{X}, \mathbb{Y}, \mathbb{U}$  are compact.

**THEOREM 4.1.** *An optimal solution exists under Assumptions 4 and 5 and Assumption 7.*

**Proof.** Note first that

$$E[(c(X_k, U_k)) | S_k = s, U_k = u]$$

will be a measurable function of  $(s, u)$  under Assumption 7, call this  $\bar{c}(s, u)$ . We then have

$$J_\beta(s) = \min_{u \in \mathbb{U}} \left( E[\bar{c}(S_k, U_k) + \beta J_{k+1}(S_{k+1}) | S_k = s, U_k = u] \right)$$

Then, if we have weak-Feller continuity of the controlled kernel (by Assumption 4) and if we have the continuity of  $f$  (by Assumption 5), by measurable selection theorems [30], there exists a solution to the discounted cost optimality equation.  $\square$

As a consequential application, we note the following. We say that, for some  $N \in \mathbb{N}$ , a control policy is an  $N$ -memory policy if for  $t > N$ ,

$$U_t = \gamma_t(Y_{[t-N+1, t]}, U_{[t-N+1, t-1]}),$$

and for  $0 \leq t \leq N$ ,

$$U_t = \gamma_t(Y_{[0, t]}, U_{[0, t-1]}).$$

**THEOREM 4.2.** *Under the conditions of Theorem 4.1, the set of  $N$ -memory policies are asymptotically optimal; i.e., for every  $\epsilon > 0$ , there exists  $N$  so that  $N$  window policies are  $\epsilon$ -optimal.*

Before the prove the theorem, let us note that the near-optimality is operationally justified in two scenarios. If the costs start becoming active from time  $N$  onwards, or the policies in the first  $N$  stages are designed via a policy-iteration argument; see [39, p. 16].

**Proof.** Finite window truncation of  $z$  can be viewed as a uniform binning/quantization (even though the number of bins may not be countable) operation applied to the infinite-dimensional state vector  $z$  under the product topology. Indeed, define (the binning/quantization map)

$$\rho : \mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U}^{\mathbb{Z}^-} \rightarrow \mathbb{Y}^N \times \mathbb{U}^{N-1},$$

with

$$\rho(y_{(-\infty, 0]}, \tilde{u}_{(-\infty, -1]}) = (y_{[-N, 0]}, u_{[-N, -1]})$$

In this case, for any  $(\tilde{y}_{(-\infty, 0]}, \tilde{u}_{(-\infty, -1]})$  and  $(\tilde{y}_{(-\infty, 0]}, \tilde{u}_{(-\infty, -1]})$  with

$$\rho(\tilde{y}_{(-\infty, 0]}, \tilde{u}_{(-\infty, -1]}) = \rho(\tilde{y}_{(-\infty, 0]}, \tilde{u}_{(-\infty, -1]}),$$

and  $\bar{d}$  being the product metric defined by

$$\begin{aligned} & \bar{d}(\bar{y}_{(-\infty,0]}, \bar{u}_{(-\infty,-1]}, \tilde{y}_{(-\infty,0]}, \tilde{u}_{(-\infty,-1]}) \\ & := \sum_{k=0}^{\infty} 2^{-k} \frac{d_{\mathbb{Y}}(\bar{y}_k, \tilde{y}_k)}{1 + d_{\mathbb{Y}}(\bar{y}_k, \tilde{y}_k)} + \sum_{m=0}^{\infty} 2^{-m} \frac{d_{\mathbb{U}}(\bar{u}_m, \tilde{u}_m)}{1 + d_{\mathbb{U}}(\bar{u}_m, \tilde{u}_m)}, \end{aligned} \quad (4.1)$$

where  $d_{\mathbb{Y}}$  and  $d_{\mathbb{U}}$  are the metrics on  $\mathbb{Y}$  and  $\mathbb{U}$ , respectively; we have that

$$\bar{d}(\bar{y}_{(-\infty,0]}, \bar{u}_{(-\infty,-1]}, \tilde{y}_{(-\infty,0]}, \tilde{u}_{(-\infty,-1]}) \leq 2^{-N+1}.$$

Since the constructed kernel  $\mathbb{P}$  is weakly continuous, the cost function is bounded continuous, and the state space is compact, the proof is then a corollary of [53, Theorem 4.27] (see also [41, Theorem 11]) by viewing the finite window truncation as a quantization (with a uniformly bounded radius for each bin) of the state space under the product topology.

In particular, one can construct an approximate MDP model with state space  $\mathbb{Y}^N \times \mathbb{U}^{N-1}$  and action space  $\mathbb{U}$  and transition kernel  $\mathbb{P}^N$  whose solution can be extended, via the quantization rule  $\rho$ , to the whole  $\mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U}^{\mathbb{Z}^-}$  and which will be near optimal for the original problem [53, Theorem 4.27].  $\square$

**Algorithmic and Numerical Implications.** Note that a finite memory truncation leads to a uniform quantization error, therefore the mathematical analysis adopted in this paper is particularly suitable for such a problem in view of recent near optimality results for quantized approximations to weakly continuous kernels. We note that Theorem 4.2 has a rather direct relation with [39] and [40], where near optimality of finite window policies were established via alternative and more tedious methods, and more restrictive conditions (though with rates of convergence properties, which we do not present). In our current paper, the filter stability condition manifests itself via Assumption 7 with no additional assumptions other than the weak Feller property. The result above thus presents further sufficient conditions on the applicability of re-inforcement learning methods presented in [40] for finite memory near-optimal control. In particular, if one applies an independently randomized exploration control policy for  $u_t$  (for learning) in the  $Q$ -learning algorithm presented in [40, Section 4], one arrives at a control-free system under which the unique ergodicity of the measurement process is satisfied under mild conditions so that the conditions in [40, Theorem 4.1] (and in this particular fully observed interpretation, [36, Corollary 3.3]) is applicable for arriving at near-optimal control policies which use only a finite window of recent measurement and past actions.

We also note here that another positive attribute for the discounted criterion is that under weak Feller continuity, both the value functions and optimal policies are robust to model approximations [38, Theorem 4.4].

**5. The Control-Free Case: Implications on Unique Ergodicity of Non-Linear Filters and Existence of an Optimal Stationary Policy.** In this section, we interpret some of the results presented in the controlled-case in the context of the control-free case. This serves both as an application and validation of the approach, but also presents new results on the non-linear filtering problem.

Even though the control-free case can be viewed as an instance of the controlled case, here some of the assumptions can be relaxed. Accordingly, we will state two results. One is on the stationarity properties on the optimal decision policies, and the other is on how to realize it by selecting the initial prior appropriately.



We have the following assumption for some of the results to follow. This will be a complementary condition.

**Assumption 8.**  $\{X_t\}$  is stationary.

Note that if  $X_t$  is stationary, so is the pair process  $(X_t, Y_t)$ . By a standard argument (e.g. Chapter 7 in [19]), we can embed the one-sided stationary process  $\{X_k, k \in \mathbb{Z}_+\}$  into a bilateral (double-sided) stationary process  $\{X_k, k \in \mathbb{Z}\}$ .

**THEOREM 5.1.** *Let  $\mathbb{X}, \mathbb{Y}, \mathbb{U}$  be compact and Assumption 4 hold. Under either Assumption 8 or Assumptions 5 and 6 an optimal decision/control policy exists. This policy is stationary.*

**Proof.** See Section A.5.  $\square$

**COROLLARY 5.2.** *Let either (i) Assumption 8, or, (ii) under the hypotheses in Theorem 5.1, Assumption 7, hold. Then, the filter process admits an invariant probability measure.*

As noted above, in the uncontrolled setting, [5] and [15] have established weak continuity conditions (i.e., the weak-Feller property) of the non-linear filter process (i.e., the belief process) in continuous time and discrete time, respectively; where the total variation continuity of the measurement channel was imposed. These results were used to establish the existence of an invariant probability measure for the belief process. The above shows that for the existence of an invariant probability measure, one may not need to invoke the continuity conditions.

A corollary (to Theorem 5.1 and Lemma 2.4) is the following.

**COROLLARY 5.3.** *If  $Y_t$  is uniquely ergodic and if Assumption 7 holds, then the filter is uniquely ergodic.*

**Proof.** We have seen that there exists at least one invariant probability measure. Suppose that there were two distinct invariant probability measures,  $\eta_1$  and  $\eta_2$ . Since a countable collection of continuous and bounded functions can be used to distinguish probability measures, we will consider  $\int \eta_i(d\pi)f(\pi)$  for such a countable collection of functions  $f$ , for  $i = 1, 2$ . Consider the joint process  $(\pi_t, Y_{(-\infty, t]})$  with invariant measure  $\kappa_i$  and with a marginal invariant measure on  $\pi_t$  as  $\eta_i$ . Since the marginal on  $Y_{(-\infty, t]}$  is uniquely ergodic, for every invariant measure  $\kappa_i$  on the joint process, the marginal will be a constant measure, call  $\psi$ . Therefore:

$$\int \eta_i(d\pi)f(\pi) = \int \kappa_i(d\pi, dy_{(-\infty, t]})f(\pi) = E_{\kappa_i}[f(\pi)] = E_{\kappa_i}[E_{\kappa_i}[f(\pi)|Y_{(-\infty, t]}]]$$

However,  $E_{\kappa_i}[f(\pi)|Y_{(-\infty, t]}] = E_{\kappa_i}[g(Y_{(-\infty, t]})] = E_{\psi}[g(Y_{(-\infty, t]})]$  for some measurable  $g$ , by Assumption 7. Therefore, as this argument applies for any  $f$  from the distinguishing family, we must have that  $\eta_1 = \eta_2$ .  $\square$

Note that for  $Y_t$  to be uniquely ergodic, we don't require  $X_t$  to be uniquely ergodic. On the other hand, if  $X_t$  is uniquely ergodic,  $Y_t$  must be uniquely ergodic as well. For sufficient conditions on unique ergodicity of infinite-memory processes such as  $Y_t$ , see [66, Theorem 3.8]; in particular the existence of an accessible element and a continuity condition (instead of weak continuity, where setwise continuity is imposed in Assumption 5; e.g., [27, Corollary 4.2] implies this condition as well).

Notably, building on [46, Theorem 2] and [61, Prop 2.1], it can be shown, via a functional analytic argument, that almost sure filter stability in the total variation sense (which can be relaxed to weak merging) and the uniqueness of an invariant probability measure on the state process leads to unique ergodicity of the filter process. It thus turns out that a complementary condition, and with an alternative argument (e.g. to [46, Theorem 2]), can also be established: unique ergodicity of the measure-

ment process and a measurability condition related to filter stability leads to unique ergodicity.

**Remark 5.1.** *For the problem at hand, for the control-free setup, it is evident that the decision maker can always apply  $\arg \min E[c(X_k, U_k)|Y_{[0,k]}]$  at any  $k \in \mathbb{N}$ . This policy will lower bound the expected cost under any policy, including the aforementioned policy. The conclusion is that, asymptotically, they are equivalent. The main novelty here is the optimality of a stationary policy for an average cost problem.*

**On the realization problem for an optimal stationary policy**

Assumption 8 will not be applicable if the initialization of the process is to be arbitrary. Let  $\bar{P}$  be the (unique) invariant probability measure on  $\{Y_{(-\infty,t]}\}$ . If the initial measure  $P_0$  is such that  $P_0 \ll \bar{P}$ , then a result due on the ergodic theory of Markov chains (see Theorem C.1(ii)) shows that for every measurable bounded  $g$ :

$$\lim_{T \rightarrow \infty} \frac{1}{T} E_{P_0} \left[ \sum_{k=0}^{T-1} g(Y_{(-\infty,k]}) \right] = \int g(y_{(-\infty,0]}) \bar{P}(dy_{(-\infty,0]})$$

We state the following, building on Theorem C.1(ii), and the proof method of Theorem 3.5.

**THEOREM 5.4.** *Let Assumptions 6 and 5 hold. Let  $u_t = \gamma(y_{(-\infty,t]}, R_t)$  be an optimal stationary policy (by Theorem 5.1) with  $R_t$  being an i.i.d. noise process. Let  $g(y_{(-\infty,k]}, x_k) := E[c(X_k, \gamma(Y_{(-\infty,k]}, R_k))]$  where the expectation is over  $R_k$ . Let  $\bar{P}$  denote an invariant process measure on  $(Y_{(-\infty,k-1]}, X_k)$  and let  $P_0$  be the invariant measure on  $\{Y_{(-\infty,-1]}\}$  and  $\mu$  be the prior measure on  $X_0$  as imposed by the controller/decision maker. If  $P_0 \times \mu \ll \bar{P}$ ; that is, if the (filter) initialization is independent of the state process initialization and under the absolute continuity condition,*

$$\lim_{T \rightarrow \infty} \frac{1}{T} E_{P_0 \times \mu} \left[ \sum_{k=0}^{T-1} g(Y_{(-\infty,k]}, X_k) \right] = \int g(y_{(-\infty,0]}, x_0) \bar{Q}(dy_{(-\infty,0]}, x_0). \quad (5.1)$$

*The optimal control/decision cost will be attained.*

Observe again, as in the controlled setup, that in the analysis above we constructed  $y_{(-\infty,-1]}$  and not  $y_{(-\infty,0]}$ , since  $\mu$  on  $X_0$  induces directly a measure on  $Y_0$ . Note that with  $P_0 \times \mu$  viewed as an initial measure, when the process evolves, the evolution is the correct one consistent with the actual realizations under the true measure and thus the initialization does not affect the evolution of the measure:  $X_0$  determines the probability of the future events: the transition kernel:

$$\mathbb{P}(dz_{t+1}|z_t) = Q(dy_{t+1}|x_{t+1})P(dx_{t+1}|x_t)\delta_{y_{(-\infty,t]}}(dy_{(-\infty,t]})$$

is such that  $Y_t$  is generated according to both the true process and the actual process.

Note that if  $\bar{P}$  and  $P_0$  are both stationary, we have that  $\bar{P}$  decomposes as  $P_0(dy)Q(dx_0|y_{(-\infty,-1]})$ . Thus, as in the controlled-case, what we need is that  $\mu(dx_0) \ll Q(dx_0|y_{(-\infty,-1]}) P_0$  a.e., so that convergence in the sense of (5.1) holds.

**Appendix A. Proofs.**

**A.1. Proof of Theorem 1.2.** Following [2, 11], we study the limit distribution of the following occupation measures, under any policy  $\gamma$  in  $\Gamma$ . Let for  $T \geq 1$

$$v_T(D) = \frac{1}{T} \sum_{t=0}^{T-1} 1_{\{(\pi_t, u_t) \in D\}}, \quad D \in \mathcal{B}(\mathcal{P}(\mathbb{X}) \times \mathbb{U}).$$

Consider any policy  $\gamma$  in  $\Gamma$ ,  $\pi_0 \sim \mu$ , and let for  $T \geq 1$ ,

$$\mu_T(D) = E_\mu^\gamma[v_T(D)] = E_\mu^\gamma \frac{1}{T} \left[ \sum_{t=0}^{T-1} 1_{\{\pi_t, u_t\} \in D} \right], \quad D \in \mathcal{B}(\mathcal{P}(\mathbb{X}) \times \mathbb{U})$$

Let for  $\mu \in \mathcal{P}(\mathbb{X})$ ,  $\mu P(A) := \int \mu(d\pi, du) P(\pi_{t+1} \in A | \pi_t = \pi, u_t = u)$ . Then through what is often referred to as a *Krylov-Bogoliubov-type* argument, for every Borel  $A \subset \mathcal{P}(\mathbb{X})$

$$|\mu_T(A \times \mathbb{U}) - \mu_T P(A)| = E_{\mu_0}^\gamma \frac{1}{T} \left[ \sum_{t=0}^{T-1} 1_{\{\pi_t, u_t\} \in (A \times \mathbb{U})} - \sum_{t=0}^{T-1} 1_{\{\pi_{t+1}, u_{t+1}\} \in (A \times \mathbb{U})} \right] \leq \frac{1}{T} \rightarrow 0,$$

as  $T \rightarrow \infty$ . Notice that the above applies for any policy  $\gamma \in \Gamma$ . Now, if we can ensure that, for some subsequence  $\mu_{t_k}$ ,  $\mu_{t_k} \rightarrow \mu$  weakly for some probability measure  $\mu$  (which holds by the assumption that the state and action spaces are compact, which makes the set of probability measures on the state weakly compact), it would follow that  $\mu_{t_k} \eta(A \times \mathbb{U}) \rightarrow \mu \eta(A \times \mathbb{U})$ . It would follow that  $\mu_{t_k} P \rightarrow \nu P$  also, by the weak Feller condition, and hence  $\nu = \nu P$  and  $\nu$  would be stationary.

Define, with  $\Gamma_S$  denoting the set of stationary control policies mapping the belief state to actions,

$$\mathcal{G} = \{v \in \mathcal{P}(\mathcal{P}(\mathbb{X}) \times \mathbb{U}) : \exists \gamma \in \Gamma_S, v(A) = \int_{\pi, u} P^\gamma((\pi_{t+1}, u_{t+1}) \in A | x) v(dx, du), \\ A \in \mathcal{B}(\mathcal{P}(\mathbb{X}) \times \mathbb{U})\} \quad (\text{A.1})$$

Thus, every weakly converging sequence  $\mu_t$  will satisfy the above equation, and therefore, under any admissible policy, every converging occupation measure sequence converges to the set  $\mathcal{G}$ . Let us define

$$\gamma^* = \inf_{v \in \mathcal{G}} \int v(d\pi, du) \tilde{c}(\pi, u)$$

Since the expression  $\int v(dx, du) c(x, u)$  is lower semi-continuous in  $v$ , a compactness condition on  $\mathcal{G}$  will ensure the existence of an optimal occupation measure which is in  $\mathcal{G}$ : we now show that there exists an optimal occupation measure in  $\mathcal{G}$  if the transition kernel is weak-Feller and  $\mathcal{G}$  is weakly compact: The problem has now reduced to

$$\inf_{\mu \in \mathcal{G}} \int \mu(dx, du) c(x, u),$$

The set  $\mathcal{G}$  is closed, since if  $\nu_n \rightarrow \nu$  and  $\nu_n \in \mathcal{G}$ , then for continuous and bounded  $f \in C_b(\mathcal{P}(\mathbb{X}))$ ,  $\langle \nu_n, f \rangle \rightarrow \langle \nu, f \rangle$ . By weak-Feller continuity of the kernel  $\int f(\pi') \eta(d\pi' | \pi, u)$  is also continuous and thus,  $\langle \nu_n, \eta f \rangle \rightarrow \langle \nu, \eta f \rangle = \langle \nu \eta, f \rangle$ . Thus,  $\nu(f) = \nu \eta(f)$  and  $\nu \in \mathcal{G}$ . Therefore,  $\mathcal{G}$  is weakly sequentially compact. Since the integral  $\int \mu(dx, du) c(x, u)$  is lower semi-continuous on the set of measures under weak convergence, and the existence result follows from Weierstrass' Theorem. As a result, there exists an optimal occupation measure, say  $v(d\pi, du)$ . This defines a stationary control policy by the Radon-Nikodym derivative:  $\mu(u \in \cdot | \pi) = \frac{dv(d\pi, \cdot)}{d \int_u v(d\pi, \cdot)}(\pi)$ , for  $v$  a.e.  $\pi$ .  $\square$

**A.2. Proof of Lemma 3.1.** a) Let us recall the  $w$ - $s$  topology [3, 56]: Let  $\mathbb{A}, \mathbb{B}$  be complete, separable, metric spaces. The  $w$ - $s$  topology on the set of probability measures  $\mathcal{P}(\mathbb{A} \times \mathbb{B})$  is the coarsest topology under which  $\int f(a, b) \nu(da, db) : \mathcal{P}(\mathbb{A} \times \mathbb{B}) \rightarrow \mathbb{R}$  is continuous for every measurable and bounded  $f$  which is continuous in  $b \in \mathbb{B}$  for every  $a \in \mathbb{A}$  (but unlike weak topology,  $f$  does not need to be continuous in  $a$ ).

For every  $n$ , we have that for every function  $g$  that is continuous and bounded:

$$\begin{aligned} \int P_n(dy^1, dy^2, du^2) g(y^1, y^2, u^2) &= \int P_n(dy^1 | y^2) P_n(du^2, y^2) g(y^1, y^2, u^2) \\ &= \int P(dy^1 | y^2) P_n(du^2, y^2) g(y^1, y^2, u^2) \end{aligned} \quad (\text{A.2})$$

Testing the equality above on continuous and bounded functions implies this property for any measurable and bounded function (that is, continuous and bounded functions form a *separating class*, see e.g. p. 13 in [6] or Theorem 3.4.5 in [20]) for weak convergence of probability measures. Since the marginals on  $y^1, y^2$  is fixed, [56, Theorem 3.10] (see also [3, Theorem 2.5]) establishes that the sequence  $\{P_n\}$  is relatively compact under the  $w$ - $s$  topology under the stated tightness condition;  $\{P_n\}$  is tight by Prohorov's theorem.

Now, taking the limit of both sides in (A.2), we have that the left hand side converges to  $\int P(dy^1, dy^2, du^2) g(y^1, y^2, u^2)$ . The right-hand side, on the other hand can be written as:

$$\int \left( \int P(dy^1 | dy^2) g(y^1, y^2, u^2) \right) P_n(du^2, y^2) \quad (\text{A.3})$$

The expression  $\left( \int P(dy^1 | dy^2) g(y^1, y^2, u^2) \right)$  is measurable in  $y^2$  and continuous in  $u^2$  by an application of the dominated convergence theorem. As a result, by the  $w$ - $s$  convergence, in the limit we have

$$\int \left( \int P(dy^1 | dy^2) g(y^1, y^2, u^2) \right) P(du^2, y^2) \quad (\text{A.4})$$

and thus the conditional independence property is satisfied.

b) For every  $n$ , we again have that for every function  $g$  continuous and bounded:

$$\begin{aligned} \int P_n(dy^1, dy^2, du^2) g(y^1, y^2, u^2) &= \int P_n(dy^1 | y^2) P_n(du^2, y^2) g(y^1, y^2, u^2) \\ &= \int \kappa(dy^1 | y^2) P_n(du^2, y^2) g(y^1, y^2, u^2) \end{aligned} \quad (\text{A.5})$$

As in (i), testing the equality above on continuous and bounded functions implies this property for any measurable and bounded function [6, p. 13] or [20, Theorem 3.4.5]) for weak convergence of probability measures. Since the marginals on  $y^1, y^2$  is fixed, [56, Theorem 3.10] (see also [3, Theorem 2.5]) establishes that the sequence  $\{P_n\}$  is relatively compact under the  $w$ - $s$  topology under the stated tightness condition;  $\{P_n\}$  is tight by Prohorov's theorem. Now, taking the limit of both sides in (A.5), we have that the left hand side converges to  $\int P(dy^1, dy^2, du^2) g(y^1, y^2, u^2)$ . The right-hand side can be written as:

$$\int \left( \int \kappa(dy^1 | dy^2) g(y^1, y^2, u^2) \right) P_n(du^2, y^2) \quad (\text{A.6})$$

The expression  $\left(\int \kappa(dy^1|dy^2)g(y^1, y^2, u^2)\right)$  is measurable in  $y^2$  and continuous in  $u^2$  by an application of dominated convergence theorem. Furthermore, by the condition that  $\kappa$  is a continuous kernel, we have that  $\left(\int \kappa(dy^1|dy^2)g(y^1, y^2, u^2)\right)$  is continuous in  $y^2$ , by an application of a generalized dominated convergence theorem [57, Theorem 3.5] or [44, Theorem 3.5]. As a result, in the limit we have

$$\int \left(\int P(dy^1|dy^2)g(y^1, y^2, u^2)\right)P(du^2, y^2) \quad (\text{A.7})$$

and thus the conditional independence property is satisfied.  $\square$

**A.3. Proof of Theorem 3.3.** Recall (A.1) and let

$$\mu_T(D) = E[v_T(D)] = E_{v_0} \frac{1}{T} \left[ \sum_{t=1}^T 1_{\{z_t, u_t \in D\}} \right], \quad D \in \mathcal{B}(\mathbb{X} \times \mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U}^{\mathbb{Z}^-} \times \mathbb{U}).$$

This can also be written as  $\frac{1}{T} \left[ \sum_{t=1}^T P(\{(z_t, u_t) \in D\}) \right]$ . Observe that for any  $t$ ,  $x_t \leftrightarrow y_{(-\infty, t]} \leftrightarrow u_t$  holds. Let  $(x_t, y_{(-\infty, t]}, u_t) \sim P_t$ . As in the proof of Theorem A.1, through a *Krylov-Bogoliubov-type* argument, for every Borel  $A \in \mathcal{B}(\mathbb{X} \times \mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U}^{\mathbb{Z}^-}$

$$\begin{aligned} & |\mu_N(A \times \mathbb{U}) - \mu_N \mathbb{P}(A)| \\ &= \left| \frac{1}{N} \left( (\mu_0(A \times \mathbb{U}) + \dots + \mu \mathbb{P}^{(N-1)}(A)) - (m u_0 \mathbb{P}(A) + \dots + v \mathbb{P}^N(A)) \right) \right| \\ &\leq \frac{1}{N} |\mu_0(A \times \mathbb{U}) - \mu_0 \mathbb{P}^N(A)| \rightarrow 0. \end{aligned} \quad (\text{A.8})$$

As earlier, in the proof of Theorem A.1, if we can ensure that for some subsequence,  $\mu_{t_k} \rightarrow \mu$  for some probability measure  $\mu$ , it would follow that  $\mu_{t_k} P \rightarrow \mu$  also and hence  $\mu = \mu P$  and  $\mu$  would be invariant and  $\mu \in \mathcal{G}$ , where

$$\begin{aligned} \mathcal{G} &= \{v \in \mathcal{P}(\mathbb{X} \times \mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U}^{\mathbb{Z}^-}) : \\ & \quad v(B \times \mathbb{U}) = \int_{x, u} \mathbb{P}(z_{t+1} \in B | z, u) v(dz, du), B \in \mathcal{B}(\mathbb{X} \times \mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U}^{\mathbb{Z}^-})\} \end{aligned}$$

Convergence to  $\mathcal{G}$  is by (A.8) is a consequence of Lemma 2.3 under Assumption 4.

We will show that, if convergence occurs, it must also be that  $\mu \in \mathcal{H}$ ;

$$\begin{aligned} \mathcal{H} &= \{v \in \mathcal{P}(\mathbb{X} \times \mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U}^{\mathbb{Z}^-}) : \\ & \quad v(dx_0, dy_{(-\infty, 0]}, du_{(-\infty, -1]}, du_0) \\ & \quad = v(dx_0, dy_{(-\infty, 0]}, du_{(-\infty, -1]}) v(du_0 | dy_{(-\infty, 0]}, du_{(-\infty, -1]}), \end{aligned} \quad (\text{A.9})$$

that is the control action variables are conditionally independent from the state variables given the information variable  $y_{(-\infty, 0]}, u_{(-\infty, -1]}$ , thus satisfying (2.5).

We show now that  $\mu_t$ , possibly along a subsequence, will also converge to  $\mathcal{H}$ . Observe that

$$\int \mu_T(dx, dy_{(-\infty, 0]}, du_{(-\infty, -1]}, du) f(x, y_{(-\infty, 0]}, u_{(-\infty, -1]}, u)$$

$$\begin{aligned}
&= \int \frac{1}{T} \sum_{t=1}^T P_t(dx, dy_{(-\infty, 0]}, du_{(-\infty, -1]}, du_0) f(x, y_{(-\infty, 0]}, u_{(-\infty, -1]}, u_0) \\
&= \int \frac{1}{T} \sum_{t=1}^T P_t(dx|y_{(-\infty, 0]}, du_{(-\infty, -1]}) P_t(dy_{(-\infty, 0]}, du_{(-\infty, -1]}, du_0) f(x, y_{(-\infty, 0]}, u_{(-\infty, -1]}, u_0) \\
&= \int \frac{1}{T} \sum_{t=1}^T \left( P_t(dx|y_{(-\infty, 0]}, du_{(-\infty, -1]}) f(x, y_{(-\infty, 0]}, u_{(-\infty, -1]}, u_0) \right) P_t(dy_{(-\infty, 0]}, du_{(-\infty, -1]}, du_0) \\
&= \int \frac{1}{T} \sum_{t=1}^T \left( P(dx|y_{(-\infty, 0]}, du_{(-\infty, -1]}) f(x, y_{(-\infty, 0]}, u_{(-\infty, -1]}, u_0) \right) P_t(dy_{(-\infty, 0]}, du_{(-\infty, -1]}, du_0)
\end{aligned} \tag{A.10}$$

$$\begin{aligned}
&= \int \frac{1}{T} \sum_{t=1}^T \left( g(y_{(-\infty, 0]}, u_{(-\infty, -1]}, u_0) \right) P_t(dy_{(-\infty, t]}, du_{(-\infty, t-1]}, du_t)
\end{aligned} \tag{A.11}$$

$$\begin{aligned}
&= \int \frac{1}{T} \sum_{t=1}^T \left( g(y_{(-\infty, t]}, u_{(-\infty, t-1]}, u_t) \right) P_t(dy_{(-\infty, t]}, du_{(-\infty, t-1]}, du_t) \\
&\rightarrow \int \left( g(y_{(-\infty, 0]}, u_{(-\infty, -1]}, u_0) \right) \mu(dy_{(-\infty, 0]}, du_{(-\infty, -1]}, du_0)
\end{aligned} \tag{A.12}$$

$$\begin{aligned}
&= \int \left( P(dx|y_{(-\infty, 0]}, u_{(-\infty, -1]}) f(x, y_{(-\infty, 0]}, u_{(-\infty, -1]}, u_0) \right) \mu(dy_{(-\infty, 0]}, du_{(-\infty, -1]}, du_0) \\
&= \int \left( P(dx|y_{(-\infty, 0]}, u_{(-\infty, -1]}) \mu(dy_{(-\infty, 0]}, du_{(-\infty, -1]}, du_0) \right) f(x, y_{(-\infty, 0]}, u_{(-\infty, -1]}, u_0)
\end{aligned} \tag{A.13}$$

Here (A.10) is due to Assumption 6<sup>1</sup> and in (A.11) we define

$$g(y_{(-\infty, 0]}, u_{(-\infty, -1]}, u_0) = \left( P(dx|y_{(-\infty, 0]}, u_{(-\infty, -1]}) f(x, y_{(-\infty, 0]}, u_{(-\infty, -1]}, u_0) \right).$$

Here, (A.12) follows from weak continuity of the kernel under Assumption 5. Thus,  $\mu_t(\cdot)$  converges weakly to  $\mu(\cdot)$ ,  $\mu$  satisfies the conditional independence property and is thus in  $\mathcal{H}$ .

In view of this, the existence problem reduces to

$$\inf_{v \in \mathcal{G} \cap \mathcal{H}} \int v(dx, dy_{(-\infty, 0]}, du_{(-\infty, -1]}, du) c(x, u) \tag{A.14}$$

By Lemma 3.2 we have that  $\mathcal{G} \cap \mathcal{H}$  is a compact set under weak topology, and since  $c$  is continuous, there exists an optimal measure  $v \in \mathcal{G} \cap \mathcal{H}$ .

□

**A.4. Proof of Theorem 3.5.** Let  $\bar{P}$  instead denote an invariant process measure on  $(Y_{(-\infty, k]}, U_{(-\infty, k-1]}, U_k, X_k)$ . Let  $P_0$  be the projection of  $\bar{P}$  on  $dy_{(-\infty, -1]}, du_{(-\infty, -1]}$ . If  $P_0(dy_{(-\infty, -1]}, du_{(-\infty, -1]}) \times \pi_0(dx_0, dy_0, du_u) \ll \bar{P}$ ; that is, with the initial state distribution selected independently of the *past* process; then with  $u_t = \gamma(y_{(-\infty, t]}, u_{(-\infty, t-1]}, r_t)$

<sup>1</sup>In the absence of Assumption 6 (A.10) would be incorrect; as a counterexample, consider  $y_k$  giving no information (e.g.  $y_k$  being constant); in this case, we clearly require  $X_t$  to be stationary for this to be correct since there is no information:  $P(X_t|Y_{-\infty, t]) = P(X_{t+1}|Y_{-\infty, t+1])$ .

and

$$g(y_{(-\infty,k]}, u_{(-\infty,k-1]}, x_k) = E[c(x_k, u_k) | x_k, y_{(-\infty,k]}, u_{(-\infty,k-1]}],$$

$$\lim_{T \rightarrow \infty} \frac{1}{T} E_{P_0 \times \pi_0} \sum_{k=0}^{T-1} g(y_{(-\infty,k]}, u_{(-\infty,k-1]}, x_k) = \int g(y_{(-\infty,k]}, u_{(-\infty,k-1]}, x_k) \bar{Q}(y_{(-\infty,0]}, u_{(-\infty,-1]})$$

for some  $\bar{Q}$  which is invariant. Note here that  $\bar{Q}$  does not need to be equal to  $\bar{P}$ .

□

**A.5. Proof of Theorem 5.1.** The proof closely follows that of Theorem 3.3. Recall (A.1) and let

$$\mu_T(D) = E[v_T(D)] = E_{v_0} \frac{1}{T} \left[ \sum_{t=1}^T 1_{\{z_t, u_t \in D\}} \right], \quad D \in \mathcal{B}(\mathbb{X} \times \mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U}^{\mathbb{Z}^-} \times \mathbb{U}).$$

This can also be written as

$$\frac{1}{T} \left[ \sum_{t=1}^T P(\{(z_t, u_t) \in D\}) \right]$$

Observe that for any  $t$ ,  $x_t \leftrightarrow y_{(-\infty,t]} \leftrightarrow u_t$  holds. Let  $(x_t, y_{(-\infty,t]}, u_t) \sim P_t$ . Now, again through the *Krylov-Bogoliubov-type* argument, for every Borel  $A$

$$\begin{aligned} & |\mu_N(A \times \mathbb{U}) - \mu_N \mathbb{P}(A)| \\ &= \left| \frac{1}{N} \left( (\mu_0(A \times \mathbb{U}) + \dots + \mu \mathbb{P}^{(N-1)}(A)) - (v_0 \mathbb{P}(A) + \dots + v \mathbb{P}^N(A)) \right) \right| \\ &\leq \frac{1}{N} |\mu_0(A \times \mathbb{U}) - \mu_0 \mathbb{P}^N(A)| \rightarrow 0. \end{aligned} \tag{A.15}$$

Now, if we can ensure that for some subsequence,  $\mu_{t_k} \rightarrow \mu$  for some probability measure  $\mu$ , it would follow that  $\mu_{t_k} P(B) \rightarrow \mu(B)$  also and hence  $\mu(B) = \mu P(B)$  and  $\mu$  would be invariant.

Thus, if  $\mu_t$  converges, it must be that  $\mu_t \rightarrow \mu$  for some  $\mu \in \mathcal{G} \cap \mathcal{H}$ , where

$$\begin{aligned} \mathcal{G} &= \{v \in \mathcal{P}(\mathbb{X} \times \mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U}) : \\ & v(B \times \mathbb{U}) = \int_{x,u} P(X_1, Y_{(-\infty,1]} \in B | x_0, y_{(-\infty,0]}, u) v(dx_0, dy_{(-\infty,0]}, du), \\ & \quad B \in \mathcal{B}(\mathbb{X} \times \mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U}) \} \end{aligned} \tag{A.16}$$

and

$$\begin{aligned} \mathcal{H} &= \{v \in \mathcal{P}(\mathbb{X} \times \mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U}) : \\ & v(dx, dy_{(-\infty,0]}, du_0) = v(dx, dy_{(-\infty,0]}) P(du_0 | dy_{(-\infty,0]}) \}, \end{aligned} \tag{A.17}$$

that is the control action variables are conditionally independent from the state variables given the information variable  $y_{(-\infty,0]}, u_{(-\infty,-1]}$ .

Convergence to  $\mathcal{G}$  is by (A.15) as a consequence of Lemma 2.3 under Assumption

4.

We show now that  $\mu_t$  will also converge to  $\mathcal{H}$ .

**Case 1. Under Assumption 8.** For a continuous and bounded  $f$ , if  $\mu_t \rightarrow \mu$  for some  $\mu$ , we have that

$$\begin{aligned}
& \int \mu_t(dx, dy_{(-\infty, 0]}, du) f(x, y_{(-\infty, 0]}, u) \\
&= \int \frac{1}{T} \sum_{t=1}^T P_t(dx, dy_{(-\infty, 0]}, du_0) f(x, y_{(-\infty, 0]}, u_0) \\
&= \int \frac{1}{T} \sum_{t=1}^T P_t(dx|y_{(-\infty, 0]}) P_t(dy_{(-\infty, 0]}, du_0) f(x, y_{(-\infty, 0]}, u_0) \\
&= \int \frac{1}{T} \sum_{t=1}^T \left( P_t(dx|y_{(-\infty, 0]}) f(x, y_{(-\infty, 0]}, u_0) \right) P_t(dy_{(-\infty, 0]}, du_0) \\
&= \int \frac{1}{T} \sum_{t=1}^T \left( P(dx|y_{(-\infty, 0]}) f(x, y_{(-\infty, 0]}, u_0) \right) P_t(dy_{(-\infty, 0]}, du_0) \quad (\text{A.18})
\end{aligned}$$

$$= \int \frac{1}{T} \sum_{t=1}^T \left( g(y_{(-\infty, 0]}, u_0) \right) P_t(dy_{(-\infty, t]}, du_t) \quad (\text{A.19})$$

$$\begin{aligned}
&= \int \frac{1}{T} \sum_{t=1}^T \left( g(y_{(-\infty, t]}, u_t) \right) P_t(dy_{(-\infty, t]}, du_t) \\
&\rightarrow \int \left( g(y_{(-\infty, 0]}, u_0) \right) \mu(dy_{(-\infty, 0]}, du_0) \\
&= \int \left( P(dx|y_{(-\infty, 0]}) f(x, y_{(-\infty, 0]}, u_0) \right) \mu(dy_{(-\infty, 0]}, du_0) \\
&= \int \left( P(dx|y_{(-\infty, 0]}) \mu(dy_{(-\infty, 0]}, du_0) \right) f(x, y_{(-\infty, 0]}, u_0) \quad (\text{A.20})
\end{aligned}$$

where (A.18) uses stationarity and in (A.19) we define

$$g(y_{(-\infty, 0]}, u_0) = \left( P(dx|y_{(-\infty, 0]}) f(x, y_{(-\infty, 0]}, u_0) \right).$$

This expression will converge as soon as  $\mu_t(dy_{(-\infty, 0]}, du_0)$  converges weakly to  $\mu$ , where  $\mu$  satisfies the conditional independence property. Here,  $\mu_t$  converges to  $\mu$  in the  $w-s$  sense. Thus, even in the absence of Assumption 5, convergence holds in this case.<sup>2</sup>

**Case 2. Under Assumptions 5 and 6.** If we don't assume stationarity, we can modify the above as follows, but then we need the weak continuity condition stated in Assumption 5:

$$\int \mu_t(dx, dy_{(-\infty, 0]}, du) f(x, y_{(-\infty, 0]}, u)$$

---

<sup>2</sup>Here, we crucially assume that the marginal on  $x_k, y_{(-\infty, k]}$  is fixed, and the realizations are generated according to the stationary measure. This is crucial for the argument in (A.18). In the above, we don't need weak continuity.



$$\begin{aligned}
&= \int \frac{1}{T} \sum_{t=1}^T P_t(dx, dy_{(-\infty, 0]}, du_0) f(x, y_{(-\infty, 0]}, u_0) \\
&= \int \frac{1}{T} \sum_{t=1}^T P_t(dx|y_{(-\infty, 0]}) P_t(dy_{(-\infty, 0]}, du_0) f(x, y_{(-\infty, 0]}, u_0) \\
&= \int \frac{1}{T} \sum_{t=1}^T \left( P_t(dx|y_{(-\infty, 0]}) f(x, y_{(-\infty, 0]}, u_0) \right) P_t(dy_{(-\infty, 0]}, du_0) \\
&= \int \frac{1}{T} \sum_{t=1}^T \left( P(dx|y_{(-\infty, 0]}) f(x, y_{(-\infty, 0]}, u_0) \right) P_t(dy_{(-\infty, 0]}, du_0) \quad (\text{A.21})
\end{aligned}$$

$$= \int \frac{1}{T} \sum_{t=1}^T \left( g(y_{(-\infty, 0]}, u_0) \right) P_t(dy_{(-\infty, t]}, du_t) \quad (\text{A.22})$$

$$\begin{aligned}
&= \int \frac{1}{T} \sum_{t=1}^T \left( g(y_{(-\infty, t]}, u_t) \right) P_t(dy_{(-\infty, t]}, du_t) \\
&\rightarrow \int \left( g(y_{(-\infty, 0]}, u_0) \right) \mu(dy_{(-\infty, 0]}, du_0) \quad (\text{A.23})
\end{aligned}$$

$$\begin{aligned}
&= \int \left( P(dx|y_{(-\infty, 0]}) f(x, y_{(-\infty, 0]}, u_0) \right) \mu(dy_{(-\infty, 0]}, du_0) \\
&= \int \left( P(dx|y_{(-\infty, 0]}) \mu(dy_{(-\infty, 0]}, du_0) \right) f(x, y_{(-\infty, 0]}, u_0) \quad (\text{A.24})
\end{aligned}$$

<sup>3</sup> Here (A.21) is due to Assumption 6 and in (A.22) we define

$$g(y_{(-\infty, 0]}, u_0) = \left( P(dx|y_{(-\infty, 0]}) f(x, y_{(-\infty, 0]}, u_0) \right).$$

This expression will converge as soon as  $\mu_t(dy_{(-\infty, 0]}, du_0)$  converges weakly to  $\mu$ , where  $\mu$  satisfies the conditional independence property. Here, (A.23) follows from weak continuity of the kernel under Assumption 5.

In view of this, the existence problem reduces to

$$\inf_{v \in \mathcal{G} \cap \mathcal{H}} \int v(dx, dy_{(-\infty, 0]}, du) c(x, u)$$

where

We note that under weak continuity of the transition kernel for  $\mathbb{P}$ ,  $\mathcal{G}$  is a closed set under weak convergence. By Lemma 3.1,  $\mathcal{H}$  is also closed. Since the state space is compact, the set of probability measures on  $(\mathbb{X} \times \mathbb{Y}^{\mathbb{Z}^-} \times \mathbb{U})$  is tight. Thus  $\mathcal{G} \cap \mathcal{H}$  is a compact set under weak topology and since  $c$  is continuous, there exists an optimal measure  $v \in \mathcal{G} \cap \mathcal{H}$ .  $\square$

## Appendix B. Discussion on Assumptions 5, 7 and 6 .

---

<sup>3</sup>In the absence of Assumption 6 (A.21) would be incorrect. As a counterexample, consider  $y_k =$  no information; in this case, we clearly require  $X_t$  to be stationary for this to be correct since there is no information:  $P(X_t|Y_{-\infty, t]) = P(X_{t+1}|Y_{-\infty, t+1])$ .

**B.1. Sufficient Conditions for Assumption 5.** For the case with finite measurement and actions, this condition is satisfied by almost sure filter stability, though with a uniformity condition over priors, in the following sense

$$\sup_{\nu} \|\pi_n^{\mu} - \pi_n^{\nu}\|_{BL} \rightarrow 0 \text{ } P^{\mu} \text{ a.s.}, \quad (\text{B.1})$$

where  $BL$  denotes the bounded Lipschitz norm (or any other weak convergence inducing metric can be used). This condition (including uniformity) holds under a contraction analysis via the *Hilbert metric*, as shown in [27, Corollary 4.2]. Furthermore, by modifying the proof of [47, Lemma 3.5] to place  $\sup_{\nu}$  inside the expectation, [47, Corollary 3.7] also leads to this condition in view of Borel-Cantelli Lemma as noted in [47, Remark 3.10].

The sufficiency of (B.1) builds on the following: Observe that

$$\begin{aligned} & E[f(X_0)|y_{(-\infty,0]}^n, u_{(-\infty,-1]}^n] - E[f(X_0)|y_{(-\infty,0]}, u_{(-\infty,-1]}] \\ &= E[f(X_0)|y_{(-\infty,0]}^n, u_{(-\infty,-1]}^n] - E[f(X_0)|(y, u)_{(-\infty,-N]}^n, y_{(-N+1,0]}, u_{(-N+1,-1]}] \end{aligned} \quad (\text{B.2})$$

$$+ E[f(X_0)|(y, u)_{(-\infty,-N]}^n, y_{(-N+1,0]}, u_{(-N+1,-1]}] - E[f(X_0)|y_{(-\infty,0]}, u_{(-\infty,-1]}] \quad (\text{B.3})$$

The uniform filter stability condition (B.1) will allow us to truncate the past finite window: For every  $\epsilon > 0$  and  $\pi = P(X_n \in \cdot | y_{(\infty,n-)}, u_{(\infty,n-1)})$  select  $N$  so that, the effect of the history is uniformly, over all priors  $P(X_{-N} \in \cdot | (y, u)_{(-\infty,-N]}^n)$ , is less than  $\frac{\epsilon}{2}$ , so that (B.3) is bounded uniformly over all sequences prior to  $-N$ .

We then apply a continuity argument for the first term, (B.2):

For this first term, in case the the measurements and actions are finitely valued, the desired result follows since for sufficiently large  $n$ , the first  $N$  coordinates of measurement and actions  $y_{(-N+1,0]}, u_{(-N+1,-1]}$ , will need to match to satisfy proximity under the product metric so that for all sufficiently large  $n$ :

$$y_{(-N+1,0]}^n, u_{(-N+1,-1]}^n = y_{(-N+1,0]}, u_{(-N+1,-1]},$$

making (B.2) zero.

For the case with continuous measurements and actions, conditions in [48, Lemma 4.6] suffices, together with the uniform filter stability condition presented above above.

**B.2. Sufficient Conditions for Assumptions 6 and 7.** In the control-free case; Assumption 6 is implied by stationarity or both Assumptions 7 and 6 hold under almost sure filter stability, see e.g. [17, 16]. On Assumption 7, as noted earlier a further related sufficient condition, obtained via the Hilbert metric, is [27, Corollary 4.2]. Complementing this, for both controlled and control-free setups, conditions in [47] (via [42, Theorem 2, Part 2] leading also to almost sure stability under total variation) based on Dobrushin's coefficients of the measurement channel and the controlled transition kernel leads to almost sure filter stability and accordingly Assumption 6.

### Appendix C. An ergodic theorems for Markov chains.

Suppose that  $\{X_t\}_{t \geq 0}$  denote a discrete-time Markov chain with state space  $\mathbb{X}$ , a Polish space.

**THEOREM C.1.** [31] [63] *Let  $\bar{P}$  be an invariant probability measure for a Markov process.*

- (i) [Ergodic decomposition and weak convergence] For  $x, \bar{P}$  a.s.,  $\frac{1}{N}E_x[\sum_{t=0}^{N-1} 1_{\{x_n \in \cdot\}}] \rightarrow P_x(\cdot)$  weakly and  $\bar{P}$  is invariant for  $P_x(\cdot)$  in the sense that

$$\bar{P}(B) = \int P_x(B)\bar{P}(dx)$$

- (ii) [Convergence in total variation] For all  $\mu \in \mathcal{P}(\mathbb{X}^{\mathbb{N}})$  which satisfies that  $\mu \ll \bar{P}$  (that is,  $\mu$  is absolutely continuous with respect to  $\bar{P}$ ), there exists  $v^*$  such that

$$\|E_\mu[\frac{1}{N} \sum_{t=0}^{N-1} 1_{\{T^n X \in \cdot\}}] - v^*(\cdot)\|_{TV} \rightarrow 0.$$

#### REFERENCES

- [1] F. Albertini, P. Dai Pra, and C. Prior. Small parameter limit for ergodic, discrete-time, partially observed, risk-sensitive control problems. *Mathematics of Control, Signals and Systems*, 14(1):1–28, 2001.
- [2] A. Arapostathis, V. S. Borkar, E. Fernandez-Gaucherand, M. K. Ghosh, and S. I. Marcus. Discrete-time controlled Markov processes with average cost criterion: A survey. *SIAM J. Control and Optimization*, 31:282–344, 1993.
- [3] E. J. Balder. On ws-convergence of product measures. *Mathematics of Operations Research*, 26(3):494–518, 2001.
- [4] D. P. Bertsekas and S. Shreve. *Stochastic Optimal Control: The Discrete Time Case*. Academic Press, New York, 1978.
- [5] A. G. Bhatt, A. Budhiraja, and R. L. Karandikar. Markov property and ergodicity of the nonlinear filter. *SIAM Journal on Control and Optimization*, 39(3):928–949, 2000.
- [6] P. Billingsley. *Convergence of probability measures*. New York: Wiley, 2nd edition, 1999.
- [7] J.-M. Bismut. Partially observed diffusions and their control. *SIAM Journal on Control and Optimization*, 20(2):302–309, 1982.
- [8] D. Blackwell. Memoryless strategies in finite-stage dynamic programming. *Annals of Mathematical Statistics*, 35:863–865, 1964.
- [9] V. S. Borkar. White-noise representations in stochastic realization theory. *SIAM J. on Control and Optimization*, 31:1093–1102, 1993.
- [10] V. S. Borkar. Average cost dynamic programming equations for controlled Markov chains with partial observations. *SIAM J. Control Optim.*, 39(3):673–681, 2000.
- [11] V. S. Borkar. Convex analytic methods in Markov decision processes. In *Handbook of Markov Decision Processes*, E. A. Feinberg, A. Shwartz (Eds.), pages 347–375. Kluwer, Boston, MA, 2001.
- [12] V. S. Borkar. Dynamic programming for ergodic control with partial observations. *Stochastic Processes and their Applications*, 103:293–310, 2003.
- [13] V. S. Borkar. Dynamic programming for ergodic control of Markov chains under partial observations: A correction. *SIAM J. Control Optim.*, 45(6):2299–2304, 2007.
- [14] V. S. Borkar and A. Budhiraja. A further remark on dynamic programming for partially observed Markov processes. *Stochastic Processes and their Applications*, 112:79–93, 2004.
- [15] A. Budhiraja. On invariant measures of discrete time filters in the correlated signal-noise case. *The Annals of Applied Probability*, 12(3):1096–1113, 2002.
- [16] P. Chigansky and R. Van Handel. A complete solution to Blackwell’s unique ergodicity problem for hidden Markov chains. *The Annals of Applied Probability*, 20(6):2318–2345, 2010.
- [17] P. Chigansky, R. Liptser, and R. van Handel. Intrinsic methods in filter stability. *Handbook of Nonlinear Filtering*, 2009.
- [18] L. Dubins and D. Freedman. Measurable sets of measures. *Pacific J. Math.*, 14:1211–1222, 1964.
- [19] R. Durrett. *Probability: theory and examples*, volume 3. Cambridge university press, 2010.
- [20] S. N. Ethier and T. G. Kurtz. *Markov processes: characterization and convergence*, volume 282. John Wiley & Sons, 2009.
- [21] E. A. Feinberg, P. O. Kasyanov, and N. V. Zadoianchuk. Average cost markov decision processes with weakly continuous transition probabilities. *Mathematics of Operations Research*, 37(4):591–607, 2012.

- [22] E.A. Feinberg, P.O. Kasyanov, and M.Z. Zgurovsky. Partially observable total-cost Markov decision process with weakly continuous transition probabilities. *Mathematics of Operations Research*, 41(2):656–681, 2016.
- [23] E. Fernández-Gaucherand, A. Arapostathis, and S. I. Marcus. Remarks on the existence of solutions to the average cost optimality equation in markov decision processes. *Systems & control letters*, 15(5):425–432, 1990.
- [24] F.Kochman and J. Reeds. A simple proof of kajiser’s unique ergodicity result for hidden markov  $\alpha$ -chains. *The Annals of Applied Probability*, 16(4):1805–1815, 2006.
- [25] W.H. Fleming and E. Pardoux. Optimal control for partially observed diffusions. *SIAM J. Control Optim.*, 20(2):261–285, 1982.
- [26] I. I. Gihman and A. V. Skorohod. *Controlled stochastic processes*. Springer Science & Business Media, 2012.
- [27] F. Le Gland and N. Oudjane. Stability and uniform approximation of nonlinear filters using the Hilbert metric and application to particle filters. *The Annals of Applied Probability*, 14(1):144–187, 2004.
- [28] M. Hairer. Ergodic properties of a class of non-Markovian processes. In *Trends in stochastic analysis, London Math. Soc. Lecture Note Ser.*, pages 65–98. Cambridge Univ. Press, 2009.
- [29] O. Hernández-Lerma. *Adaptive Markov Control Processes*. Springer-Verlag, 1989.
- [30] O. Hernández-Lerma and J. B. Lasserre. *Discrete-Time Markov Control Processes: Basic Optimality Criteria*. Springer, 1996.
- [31] O. Hernández-Lerma and J. B. Lasserre. *Markov Chains and Invariant Probabilities*. Birkhäuser, Basel, 2003.
- [32] S.-P. Hsu, D.-M Chuang, and A. Arapostathis. On the existence of stationary optimal policies for partially observed mdps under the long-run average cost criterion. *Systems & control letters*, 55(2):165–173, 2006.
- [33] T. Kaijser. A limit theorem for partially observed Markov chains. *Annals of Probability*, 4(677), 1975.
- [34] T. Kaijser. On markov chains induced by partitioned transition probability matrices. *Acta Mathematica Sinica, English Series*, 27(3):441–476, 2011.
- [35] A.D Kara, N. Saldi, and S. Yüksel. Weak Feller property of non-linear filters. *Systems & Control Letters*, 134:104–512, 2019.
- [36] A.D Kara, N. Saldi, and S. Yüksel. Q-learning for mdps with general spaces: Convergence and near optimality via quantization under weak continuity. *arXiv:2111.06781*, 2021.
- [37] A.D Kara and S. Yüksel. Robustness to incorrect priors in partially observed stochastic control. *SIAM Journal on Control and Optimization*, 57(3):1929–1964, 2019.
- [38] A.D Kara and S. Yüksel. Robustness to incorrect system models in stochastic control. *SIAM Journal on Control and Optimization*, 58(2):1144–1182, 2020.
- [39] A.D Kara and S. Yüksel. Near optimality of finite memory feedback policies in partially observed markov decision processes. *Journal of Machine Learning Research*, 23(11):1–46, 2022.
- [40] A.D Kara and S. Yüksel. Convergence of finite memory Q-learning for POMDPs and near optimality of learned policies under filter stability. *Mathematics of Operations Research (also arXiv:2103.12158)*, 2023.
- [41] Ali D. Kara and S. Yüksel. Robustness to approximations and model learning in mdps and pomdps. In A. B. Piunovskiy and Y. Zhang, editors, *Modern Trends in Controlled Stochastic Processes: Theory and Applications, Volume III*. Luniver Press, 2021.
- [42] ML Kleptsyna and A Yu Veretennikov. On discrete time ergodic filters with wrong initial data. *Probability Theory and Related Fields*, 141(3-4):411–444, 2008.
- [43] H. Kunita. Asymptotic behavior of the nonlinear filtering errors of markov processes. *Journal of Multivariate Analysis*, 1(4):365–393, 1971.
- [44] H.J. Langen. Convergence of dynamic programming models. *Math. Oper. Res.*, 6(4):493–512, Nov. 1981.
- [45] A. S. Manne. Linear programming and sequential decision. *Management Science*, 6:259–267, April 1960.
- [46] G. Di Masi and L. Stettner. Ergodicity of hidden markov models. *Mathematics of Control, Signals and Systems*, 17(4):269–296, 2005.
- [47] C. McDonald and S. Yüksel. Exponential filter stability via Dobrushin’s coefficient. *Electronic Communications in Probability*, 25, 2020.
- [48] C. McDonald and S. Yüksel. Robustness to incorrect priors and controlled filter stability in partially observed stochastic control. *SIAM Journal on Control and Optimization*, 60(2):842–870, 2022.
- [49] K.R. Parthasarathy. *Probability Measures on Metric Spaces*. AMS Bookstore, 1967.

- [50] L. K. Platzman. Optimal infinite-horizon undiscounted control of finite probabilistic systems. *SIAM Journal on Control and Optimization*, 18(4):362–380, 1980.
- [51] D. Rhenius. Incomplete information in Markovian decision models. *Ann. Statist.*, 2:1327–1334, 1974.
- [52] Wolfgang J Runggaldier and Lukasz Stettner. *Approximations of discrete time partially observed control problems*. Giardini Pisa, 1994.
- [53] N. Saldi, T. Linder, and S. Yüksel. *Finite Approximations in Discrete-Time Stochastic Control: Quantized Models and Asymptotic Optimality*. Springer, Cham, 2018.
- [54] N. Saldi and S. Yüksel. Geometry of information structures, strategic measures and associated control topologies. *Probability Surveys*, 19:450–532, 2022.
- [55] N. Saldi, S. Yüksel, and T. Linder. On the asymptotic optimality of finite approximations to markov decision processes with borel spaces. *Mathematics of Operations Research*, 42(4):945–978, 2017.
- [56] M. Schäl. Conditions for optimality in dynamic programming and for the limit of n-stage optimal policies to be optimal. *Z. Wahrscheinlichkeitsth.*, 32:179–296, 1975.
- [57] R. Serfozo. Convergence of lebesgue integrals with varying measures. *Sankhyā: The Indian Journal of Statistics, Series A*, pages 380–402, 1982.
- [58] V. Strassen. The existence of probability measures with given marginals. *The Annals of Mathematical Statistics*, 36(2):423–439, 1965.
- [59] T. Szarek. Feller processes on nonlocally compact spaces. *Annals of Probability*, pages 1849–1863, 2006.
- [60] A. Budhiraja V. S. Borkar. A further remark on dynamic programming for partially observed markov processes. *Stochastic processes and their applications*, 112(1):79–93, 2004.
- [61] R. van Handel. Uniform time average consistency of Monte Carlo particle filters. *Stochastic Processes and their Applications*, 119(11):3835–3861, 2009.
- [62] H. S. Witsenhausen. Equivalent stochastic control problems. *Math. Control, Signals and Systems*, 1:3–11, 1988.
- [63] D. T. H. Worm and S. C. Hille. Ergodic decompositions associated with regular markov operators on polish spaces. *Ergodic Theory and Dynamical Systems*, 31(2):571 – 597, 2010.
- [64] L.C. Young. Generalized curves and the existence of an attained absolute minimum in the calculus of variations. *Comptes Rendus de la Societe des Sci. et des Lettres de Varsovie*, 30:212–234, 1937.
- [65] H. Yu and D. P. Bertsekas. On near optimality of the set of finite-state controllers for average cost pomdp. *Mathematics of Operations Research*, 33(1):1–11, 2008.
- [66] S. Yüksel. On stochastic stability of a class of non-Markovian processes and applications in quantization. *SIAM J. on Control and Optimization*, 55:1241–1260, 2017.
- [67] S. Yüksel. A universal dynamic program and refined existence results for decentralized stochastic control. *SIAM Journal on Control and Optimization*, 58(5):2711–2739, 2020.
- [68] S. Yüksel and N. Saldi. Convex analysis in decentralized stochastic control, strategic measures and optimal solutions. *SIAM J. on Control and Optimization*, 55:1–28, 2017.
- [69] A.A. Yushkevich. Reduction of a controlled Markov model with incomplete data to a problem with complete information in the case of Borel state and control spaces. *Theory Prob. Appl.*, 21:153–158, 1976.