

# DENSITIES FOR ELLIPTIC CURVES OVER GLOBAL FUNCTION FIELDS

ANDREW YAO

ABSTRACT. Let  $K$  be a global function field. Using Haar measures, we compute the densities of the Kodaira types and Tamagawa numbers of elliptic curves over a completion of  $K$ . Also, we prove results about the number of iterations of Tate's algorithm that are completed when the algorithm is used on elliptic curves over a completion of  $K$ .

## 1. INTRODUCTION

Let  $p$  be a prime and  $q = p^n$  for a positive integer  $n$ . Let  $K$  be a finite extension of  $\mathbb{F}_q(t)$ . Define  $M_K$  to be the set of places of  $K$ . Suppose  $P \in M_K$ . Let  $K_P$  be the completion of  $K$  at  $P$  and  $R_P$  be the valuation ring of  $K_P$ . Suppose  $E$  is an elliptic curve over  $K$  with equation

$$E : y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6$$

such that  $a_1, a_2, a_3, a_4,$  and  $a_6$  are elements of  $K$ .  $E$  has a long Weierstrass form, and if  $a_1 = a_2 = a_3 = 0$ ,  $E$  has a short Weierstrass form. We study densities for elliptic curves over  $K$  that have a long Weierstrass form.

As an elliptic curve over  $K_P$ ,  $E$  has a Kodaira type, which describes its geometry. Particularly,  $E$  has a Tamagawa number  $c_P = [E(K_P) : E_0(K_P)]$  over  $K_P$ . A method to determine the Kodaira type and Tamagawa number of an elliptic curve over  $K_P$  is Tate's algorithm ([6], [7]). The description of Tate's algorithm in [6] is used in this paper to compute local densities. Often, steps from this description of Tate's algorithm are referred to.

The papers [2] and [3] discuss densities of Kodaira types and Tamagawa products for elliptic curves over  $\mathbb{Q}$ . In these papers, the densities at the nonarchimedean places of  $\mathbb{Q}$  are considered. In [2] and [3], the densities are for elliptic curves in long and short Weierstrass form, respectively. Moreover, [1] discusses densities of Kodaira types and Tamagawa products for elliptic curves over number fields in short Weierstrass form. Note that some of the methods for computing local densities with Tate's algorithm used in Section 4, Section 5, and Section 6 of this paper are similar to methods used in [1], [2], and [3].

Local densities over  $K_P$  can be obtained using the Haar measure. Let  $N$  be a positive integer. Note that  $K_P^N$  as an additive group is locally compact, and because of this, Haar's theorem can be used on  $K_P^N$ . Particularly, suppose  $\mu_P$  is the Haar measure on  $K_P^N$  such that  $\mu_P(R_P^N) = 1$ .

Let  $G_P$  be the set of curves  $y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6$  over  $K_P$  such that  $a_1, a_2, a_3, a_4, a_6 \in R_P$ . Because the discriminant of an elliptic curve must be nonzero, not

all elements of  $G_P$  are elliptic curves. Also, note that  $G_P$  can be considered to be  $R_P^5$ . The local densities for  $G_P$  are obtained from the Haar measure on  $R_P^5$ .

**Definition 1.1.** For an elliptic curve  $E \in G_P$ , let  $N_P(E)$  be the number of iterations of Tate's algorithm that are completed when the algorithm is used on  $E$ .

Suppose  $T$  is the set of Kodaira types. Let  $\mathfrak{r}$  be an element of  $T$  and  $n$  be a positive integer. Define  $\delta_K(\mathfrak{r}, n; P)$  to be the Haar measure of the set of elliptic curves  $E$  over  $K_P$  with coefficients in  $R_P$  such that  $E$  has Kodaira type  $\mathfrak{r}$  and the Tamagawa number of  $E$  is  $n$ . For  $k \geq 0$ , define  $\delta_K(\mathfrak{r}, n, k; P)$  to be the Haar measure of the set of elliptic curves  $E$  over  $K_P$  with coefficients in  $R_P$  such that  $E$  has Kodaira type  $\mathfrak{r}$ , the Tamagawa number of  $E$  is  $n$ , and  $N_P(E) = k$ .

In this paper, we often consider the number of iterations that Tate's algorithm completes when the algorithm is used on an elliptic curve over  $K_P$ . Note that in order to study this topic, Proposition 2.4 is useful. Next, we give an important result of the paper.

**Theorem 1.2.** For a Kodaira type  $\mathfrak{r}$ , positive integer  $n$ , and nonnegative integer  $k$ ,

$$\delta_K(\mathfrak{r}, n, k; P) = \frac{1}{Q_P^{10k}} \delta_K(\mathfrak{r}, n, 0; P).$$

We prove Theorem 1.2 by considering the cases  $p \geq 5$ ,  $p = 3$ , and  $p = 2$ . Note that the general method used to prove the theorem is to use translations. The proof of this result is given in Section 7.1.

**Organization.** The paper is organized as follows. In Section 2, we introduce elliptic curves and Tate's algorithm. Next, in Section 3, for a nonempty finite subset  $S$  of  $M_K$  and a positive integer  $N$ , we discuss how to obtain global densities for  $\mathcal{O}_{K,S}^N$ . Afterwards, in Section 4, Section 5, and Section 6, we compute the local densities if the characteristic  $p$  of  $K$  is at least 5, equal to 2, and equal to 3, respectively. Finally, in Section 7, we prove additional results about local and global densities.

**Notation.** Suppose  $P$  is a place of  $K$ . Let the degree of  $P$  be  $[R_P/\pi_P R_P : \mathbb{F}_q]$ . Also, let  $Q_P = |R_P/\pi_P R_P|$ . Moreover, let  $\pi_P$  be a uniformizer of  $P$  in  $K$ . Denote  $v_P$  to be the valuation  $v_{\pi_P}$  over  $K_P$ ; note that  $v_P$  is also a valuation over  $K$  because  $K \subset K_P$ . Additionally, for a nonnegative integer  $k$ , let  $L_{P,k}$  be a set of representatives of the cosets of  $R_P/\pi_P^k R_P$  such that  $0 \in L_{P,k}$ .

Suppose  $S$  is a finite nonempty subset of  $M_K$ . We let  $\mathcal{O}_{K,S}$  be the set of  $x \in K$  such that if  $P \in S^C = M_K \setminus S$ ,  $v_P(x) \geq 0$ . Also, let  $W_S$  be the set of curves  $y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6$  such that  $a_1, a_2, a_3, a_4, a_6 \in \mathcal{O}_{K,S}$ .

For  $d \geq 1$ , let  $T_d$  be the number of places of  $P$  with degree  $d$ . The zeta function of  $K$  is

$$\zeta_K(s) = \prod_{d=1}^{\infty} \left(1 - \frac{1}{q^{ds}}\right)^{-T_d}.$$

Suppose  $D$  is a divisor of  $K$ . Define  $L(D)$  as the set of  $x \in K$  such that  $x = 0$  or  $x \neq 0$  and  $(x) + D \geq 0$ .

**Acknowledgements.** This research was done in MIT SPUR. The author would like to thank Hao Peng for providing useful guidance. Also, the author would like to thank Zhiyu Zhang for suggesting the problem. Additionally, the author would like to thank David Jerison and Ankur Moitra for giving advice about the project.

## 2. ELLIPTIC CURVES AND GLOBAL DENSITIES

Suppose  $P$  is a place of  $K$ . Let  $E$  be an elliptic curve over  $K_P$ . There exist  $a_1, a_2, a_3, a_4, a_6 \in K_P$  such that  $E$  has equation

$$E : y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6.$$

Suppose  $a_1, a_2, a_3, a_4, a_6 \in K_P$  satisfy this condition. Additionally, define

$$\begin{aligned} b_2(E) &= a_1^2 + 4a_2, & b_4(E) &= a_1a_3 + 2a_4, & b_6(E) &= a_3^2 + 4a_6, \\ b_8(E) &= a_1^2a_6 + 4a_2a_6 - a_1a_3a_4 + a_2a_3^2 - a_4^2. \end{aligned}$$

Also, the discriminant of  $E$  is

$$\Delta(E) = -b_2(E)^2b_8(E) - 8b_4(E)^3 - 27b_6(E)^2 + 9b_2(E)b_4(E)b_6(E).$$

**Definition 2.1** ([7]). Elliptic curves  $E$  and  $F$  over  $K_P$  are *equivalent* if there exists  $l, m, n, u \in K_P$  such that  $u \neq 0$  and the equation for  $F$  can be obtained from the equation for  $E$  by first replacing  $x$  with  $u^2x + n$  and  $y$  with  $u^3y + lu^2x + m$  and then dividing by  $u^6$ .

**Definition 2.2** ([7]). An elliptic curve  $E$  over  $K_P$  is *minimal* if the equation for  $E$  has coefficients in  $R_P$  and if there does not exist an elliptic curve  $F$  over  $K_P$  such that the equation for  $F$  has coefficients in  $R_P$ ,  $F$  is equivalent to  $E$ , and  $v_P(\Delta(F)) < v_P(\Delta(E))$ .

The following proposition generalizes Theorem 3.2 of [7] to nonminimal equivalent elliptic curves. Note that this proposition is used later in the paper to compute local densities.

**Proposition 2.3.** Let  $E$  and  $F$  be elliptic curves over  $K_P$  that have equations with coefficients in  $R_P$ . Assume that  $E$  and  $F$  are equivalent and satisfy  $v_P(\Delta(E)) = v_P(\Delta(F))$ . Then, there exists  $l, m, n, u \in R_P$  such that  $v_P(u) = 0$  and the equation of  $F$  can be obtained from the equation of  $E$  by first replacing  $x$  with  $u^2x + n$  and  $y$  with  $u^3y + lu^2x + m$  and then dividing by  $u^6$ .

*Proof.* The proof of Theorem 3.2 of [7] can be used to prove this proposition. ■

**Proposition 2.4.** Let  $k$  be a nonnegative integer. Suppose  $E$  is an elliptic curve over  $K_P$  with equation

$$E : y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6$$

and assume that  $a_1, a_2, a_3, a_4, a_6 \in R_P$ . For  $l, m, n \in K_P$ , let  $E'(l, m, n)$  be the elliptic curve that is  $E$  with  $x$  replaced by  $x + n$  and  $y$  replaced by  $y + lx + m$ .  $N_P(E) \geq k$  if and only if there exists  $l, m, n \in R_P$  such that if  $E'(l, m, n)$  has equation

$$E'(l, m, n) : y^2 + a'_1xy + a'_3y = x^3 + a'_2x^2 + a'_4x + a'_6,$$

$a'_i \in \pi_P^{ki}R_P$  for  $i \in \{1, 2, 3, 4, 6\}$ .

*Proof.* Suppose  $l, m, n$  exist. Let  $l, m, n$  satisfy the condition. From Tate's algorithm, we have that  $N_P(E) = N_P(E'(l, m, n)) \geq k$ .

Next, we prove that if  $N_P(E) \geq k$ ,  $l, m$ , and  $n$  exist using induction on  $k$ . The base case  $k = 0$  is clear. Let  $a$  be a nonnegative integer and assume the result is true for  $k = a$ . We prove the result is true for  $k = a + 1$ . Assume  $N_P(E) \geq a + 1$ . Because  $N_P(E) \geq a$ ,

$l, m, n \in R_P$  exist such that if  $x$  is replaced with  $x + n$  and  $y$  is replaced with  $y + lx + m$ , the resulting curve  $E'(l, m, n) : y^2 + a'_1xy + a'_3y = x^3 + a'_2x^2 + a'_4x + a'_6$  has  $a'_i \equiv 0 \pmod{\pi_P^{ia}}$  for  $i \in \{1, 2, 3, 4, 6\}$ . Suppose  $l, m, n \in R_P$  satisfy this condition. Suppose that the curve that is obtained after Tate's algorithm is used for  $a$  iterations on  $E'(l, m, n)$  is

$$F : y^2 + \frac{a'_1}{\pi_P^a}xy + \frac{a'_3}{\pi_P^{3a}}y = x^3 + \frac{a'_2}{\pi_P^{2a}}x^2 + \frac{a'_4}{\pi_P^{4a}}x + \frac{a'_6}{\pi_P^{6a}}.$$

We have that  $F$  is  $E$  with  $x$  replaced with  $\pi_P^{2a}x + n$  and  $y$  replaced with  $\pi_P^{3a}y + l\pi_P^{2a}x + m$  divided by  $\pi_P^{6a}$ .

Because  $N_P(E'(l, m, n)) = N_P(E) \geq a + 1$ ,  $F$  will complete at least one more iteration. During this iteration, suppose  $x$  is replaced with  $x + n'$  and  $y$  is replaced with  $y + l'x + m'$ . We have that the resulting elliptic curve

$$F' : y^2 + a''_1xy + a''_3y = x^3 + a''_2x^2 + a''_4x + a''_6$$

has  $a''_i \equiv 0 \pmod{\pi_P^i}$  for  $i \in \{1, 2, 3, 4, 6\}$ . Moreover,  $F'$  is  $E$  with  $x$  replaced with

$$\pi_P^{2a}x + n + n'\pi_P^{2a}$$

and  $y$  replaced with

$$\pi_P^{3a}y + (l + l'\pi_P^a)\pi_P^{2a}x + m + m'\pi_P^{3a} + ln'\pi_P^{2a}$$

divided by  $\pi_P^{6a}$ . The equation of

$$E'(l + l'\pi_P^a, m + m'\pi_P^{3a} + ln'\pi_P^{2a}, n + n'\pi_P^{2a})$$

is

$$y^2 + \pi_P^a a''_1 xy + \pi_P^{3a} a''_3 y = x^3 + \pi_P^{2a} a''_2 x^2 + \pi_P^{4a} a''_4 x + \pi_P^{6a} a''_6,$$

and  $\pi_P^{ai} a''_i \in \pi_P^{(a+1)i} R_P$  for  $i \in \{1, 2, 3, 4, 6\}$ . This completes the induction. We are done.  $\blacksquare$

Note that Tate's algorithm cannot be used on a curve in  $G_P$  with discriminant 0. However, this is not considered in the calculations of local densities later in the paper. Suppose  $\mathfrak{r} \in T$ ,  $n$  is a positive integer, and  $k$  is a nonnegative integer. The set  $U$  of elliptic curves  $E \in G_P$  with Kodaira type  $\mathfrak{r}$ , Tamagawa number  $n$ , and  $M(E) = k$  is an open subset of  $G_P$ , because if  $E \in U$ , if multiples of  $\pi_P^M$  are added to the coefficients of  $E$  for sufficiently positive large integers  $M$ , the resulting curve will be an element of  $U$ . Particularly, the set of elliptic curves is an open subset of  $G_P$ . In the next proposition, we prove that the Haar measure of this set is 1; note that it follows that the Haar measure of the set of curves in  $G_P$  with discriminant 0 is 0.

**Proposition 2.5.** The Haar measure of the set of elliptic curves is 1.

*Proof.* Let  $M$  be a positive integer. For  $E : y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6$ , we see that the number of solutions for  $a_i, i \in \{1, 2, 3, 4, 6\}$  modulo  $\pi_P^M$  to  $\Delta(E) \equiv 0 \pmod{\pi_P^M}$  is  $O(Q_P^{4M})$ . Therefore, the Haar measure of the set of elliptic curves with discriminant equal to 0 is at most  $\frac{O(Q_P^{4M})}{Q_P^{5M}} = O(\frac{1}{Q_P^M})$ . The result follows from taking  $M \rightarrow \infty$ .  $\blacksquare$

## 3. GLOBAL DENSITIES

Next, global densities are established. Definitions and theorems from [4] are used in this section.

Let  $S$  be a finite nonempty subset of  $M_K$ . Also, suppose  $N$  is a positive integer. Let  $\text{Div}(S)$  be the set of divisors

$$\sum_{P \in S} n_P P$$

such that for  $P \in S$ ,  $n_P$  is a nonnegative integer, and there exists  $P \in S$  such that  $n_P > 0$ . Suppose  $U \subset \mathcal{O}_{K,S}^N$ . The upper density of  $U$  at  $S$  is

$$\bar{d}_S(U) = \limsup_{D \in \text{Div}(S)} \frac{|U \cap L(D)^N|}{|L(D)|^N},$$

and the lower density of  $U$  at  $S$  is

$$\underline{d}_S(U) = \liminf_{D \in \text{Div}(S)} \frac{|U \cap L(D)^N|}{|L(D)|^N}.$$

If  $\bar{d}_S(U) = \underline{d}_S(U)$ , the density  $d_S(U)$  of  $U$  at  $S$  exists, and equals  $\bar{d}_S(U) = \underline{d}_S(U)$ .

**Theorem 3.1** ([4], Theorem 2.1). For  $P \in S^C$ , let  $U_P \subset K_P^N$  be a measurable set such that  $\mu_P(\partial U_P) = 0$ . For a positive integer  $M$ , let  $V_M$  be the set of  $x \in \mathcal{O}_{K,S}^N$  such that  $x \in U_P$  for some  $P \in S^C$  with degree at least  $M$ . Suppose  $\lim_{M \rightarrow \infty} \bar{d}_S(V_M) = 0$ . Let  $\mathcal{P} : \mathcal{O}_{K,S}^N \rightarrow 2^{S^C}$ ,  $\mathcal{P}(a) = \{P \in S^C : a \in U_P\}$ . Then:

- (1)  $\sum_{P \in S^C} \mu_P(U_P)$  is convergent.
- (2) For  $T \subset 2^{S^C}$ ,  $\nu(T) := d_S(\mathcal{P}^{-1}(T))$  exists. Also,  $\nu$  defines a measure on  $2^{S^C}$ .
- (3)  $\nu$  is concentrated at finite subsets of  $S^C$ , and for a finite set  $T$  of places in  $S^C$ ,

$$\nu(T) = \prod_{P \in T} \mu_P(U_P) \prod_{P \in S^C \setminus T} (1 - \mu_P(U_P)).$$

**Theorem 3.2** ([4], Theorem 2.2). Let  $f$  and  $g$  be polynomials in  $\mathcal{O}_{K,S}[x_1, \dots, x_d]$  that are relatively prime. For  $M \geq 1$ , let  $V_M$  be the set of  $x \in \mathcal{O}_{K,S}^N$  such that  $f(x) \equiv g(x) \equiv 0 \pmod{\pi_P}$  for some  $P \in S^C$  with degree at least  $M$ . Then,  $\lim_{M \rightarrow \infty} \bar{d}_S(V_M) = 0$ .

In this paper, we consider global densities for elliptic curves over  $K$  with coefficients in  $\mathcal{O}_{K,S}$  in long Weierstrass form. We see that  $W_S$  can be considered to be  $\mathcal{O}_{K,S}^5$ , and particularly, the global density definitions from above for  $\mathcal{O}_{K,S}^5$  can be used on  $W_S$ . Similar methods are used in [2] for elliptic curves over  $\mathbb{Q}$  with coefficients in  $\mathbb{Z}$ . Note that an elliptic curve must have a nonzero discriminant, meaning that not all curves in  $W_S$  are elliptic curves. However, for  $D \in \text{Div}(S)$ , the number of curves in  $W_S$  with discriminant 0 that are elements of  $L(D)^5$ , where  $W_S$  is considered to be  $\mathcal{O}_{K,S}^5$ , is  $O(|L(D)|^4)$ . Particularly, if proportions over elliptic curves in  $W_S$  is considered rather than the proportions over  $W_S$ , the density is not changed.

Proposition 3.3 is about the global density of nonminimal elliptic curves. Note that the lemma is used to prove Theorem 7.2.

**Proposition 3.3.** For a positive integer  $M$ , let  $V_M$  be the set of elliptic curves  $E \in W_S$  such that there exists  $P \in S^C$  with degree at least  $M$  such that  $N_P(E) \geq 1$ . Then,  $\lim_{M \rightarrow \infty} \overline{d}_S(V_M) = 0$ .

*Proof.* We prove this with casework on the characteristic  $p$  of  $K$ . Suppose that  $E$  is an elliptic curve in  $G_P$  with equation  $E : y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6$  for  $a_1, a_2, a_3, a_4, a_6 \in R_P$  such that  $N_P(E) \geq 1$ .

Assume  $p \geq 5$ . We have that  $E$  can be translated to the curve

$$y^2 = x^3 + \left( -\frac{b_2(E)^2}{48} + \frac{b_4(E)}{2} \right) x - \frac{b_2(E)^3}{864} - \frac{b_2(E)b_4(E)}{24} + \frac{b_6(E)}{4}.$$

Because  $N_P(E) \geq 1$ , using Proposition 2.4,  $-\frac{b_2(E)^2}{48} + \frac{b_4(E)}{2} \equiv 0 \pmod{\pi_P}$  and  $-\frac{b_2(E)^3}{864} - \frac{b_2(E)b_4(E)}{24} + \frac{b_6(E)}{4} \equiv 0 \pmod{\pi_P}$ . Then, Theorem 3.2 with

$$f(x_1, x_2, x_3, x_4, x_6) = -\frac{(x_1^2 + 4x_2)^2}{48} + \frac{x_1x_3 + 2x_4}{2}$$

and

$$g(x_1, x_2, x_3, x_4, x_6) = -\frac{(x_1^2 + 4x_2)^3}{864} - \frac{(x_1^2 + 4x_2)(x_1x_3 + 2x_4)}{24} + \frac{x_3^2 + 4x_6}{4}$$

proves this proposition for  $p \geq 5$ .

Next, assume  $p = 3$ . We have that  $E$  can be translated to the curve

$$y^2 = x^3 + \frac{b_2(E)}{4}x^2 + \frac{b_4(E)}{2}x + \frac{b_6(E)}{4}$$

Using Proposition 2.4,  $\frac{b_2(E)}{4} \equiv 0 \pmod{\pi_P}$  from the coefficient of  $x^2$ . Additionally,  $\Delta(E) \equiv 0 \pmod{\pi_P}$ . Next, Theorem 3.2 with

$$f(x_1, x_2, x_3, x_4, x_6) = -(x_1^2 + x_2)^2(x_1^2x_6 + x_2x_6 - x_1x_3x_4 + x_2x_3^2 - x_4^2) + (x_1x_3 + 2x_4)^3$$

and

$$g(x_1, x_2, x_3, x_4, x_6) = x_1^2 + x_2$$

proves this proposition for  $p = 3$ .

Suppose  $p = 2$ . Using Proposition 2.4,  $a_1 \equiv 0 \pmod{\pi_P}$  from the coefficient of  $xy$ . Also,  $\Delta(E) \equiv 0 \pmod{\pi_P}$ . Therefore, Theorem 3.2 with

$$f(x_1, x_2, x_3, x_4, x_6) = x_1^4(x_1^2x_6 + x_1x_3x_4 + x_2x_3^2 + x_4^2) + x_3^4 + x_1^3x_3^3$$

and

$$g(x_1, x_2, x_3, x_4, x_6) = x_1$$

proves this proposition for  $p = 2$ . ■

#### 4. LOCAL DENSITIES FOR $p \geq 5$

**4.1. Setup.** Suppose that the characteristic of  $K$  is  $p \geq 5$ . Let  $P$  be a place of  $K$ . We compute the local densities over  $K_P$  of Kodaira types  $\mathfrak{r}$  and Tamagawa numbers  $n$  for elliptic curves in  $G_P$ . Let  $G_P^{(1)}$  be the set of curves

$$y^2 = x^3 + a_4x + a_6$$

over  $K_P$  such that  $a_4, a_6 \in R_P$ . Note that  $G_P^{(1)}$  can be considered to be  $R_P^2$ . Define  $\varphi : G_P \rightarrow G_P^{(1)}$  as the function such that if  $E$  is a curve in  $G_P$ ,  $\varphi(E)$  is the curve in  $G_P^{(1)}$  with equation

$$\varphi(E) : y^2 = x^3 + \left( -\frac{b_2(E)^2}{48} + \frac{b_4(E)}{2} \right) x - \frac{b_2(E)^3}{864} - \frac{b_2(E)b_4(E)}{24} + \frac{b_6(E)}{4}.$$

If  $E$  is an elliptic curve,  $\varphi(E)$  is an elliptic curve equivalent to  $E$ .

**Lemma 4.1.** If  $U$  is an open subset of  $G_P^{(1)}$ ,  $\mu_P(\varphi^{-1}(U)) = \mu_P(U)$ .

*Proof.* Let  $V$  be the set of  $y^2 = x^3 + a'_4x + a'_6$  with  $a'_4 \in r_4 + \pi_P^{n_4}R_P$  and  $a'_6 \in r_6 + \pi_P^{n_6}R_P$ . It suffices to prove that  $\mu_P(\varphi^{-1}(V)) = \mu_P(V) = \frac{1}{Q^{n_4+n_6}}$  because all open subsets of  $G_P^{(1)}$  can be written as a disjoint countable union of sets with the form of  $V$ . Suppose  $E : y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6 \in G_P$ .  $\varphi(E) \in V$  if and only if

$$-\frac{b_2(E)^2}{48} + \frac{b_4(E)}{2} \in r_4 + \pi_P^{n_4}R_P$$

and

$$-\frac{b_2(E)^3}{864} - \frac{b_2(E)b_4(E)}{24} + \frac{b_6(E)}{4} \in r_6 + \pi_P^{n_6}R_P.$$

Assume that  $\varphi(E) \in V$ . Let  $M = \max(n_4, n_6)$ . First, select  $a_1, a_2$ , and  $a_3$  modulo  $\pi_P^M$ . Each has  $Q_P^M$  possible residues. Afterwards,  $a_4$  will have  $Q_P^{M-n_4}$  residues modulo  $\pi_P^M$ ; select the residue for  $a_4$ . Finally,  $a_6$  has  $Q_P^{M-n_6}$  residues modulo  $\pi_P^M$ ; select the residue for  $a_6$ . We see that if each of  $a_1, a_2, a_3, a_4, a_6$  are taken modulo  $\pi_P^M$ , the number of combinations of residues is  $Q_P^{5M-n_4-n_6}$ . Also, because  $a_i$  is modulo  $\pi_P^M$  for  $i \in \{1, 2, 3, 4, 6\}$ , each combination of residues has a Haar measure of  $\frac{1}{Q_P^{5M}}$ . We are done.  $\blacksquare$

**4.2. Multiple Iterations.** Let  $k$  be a nonnegative integer. Suppose  $S_k$  is the set of elliptic curves  $E \in G_P^{(1)}$  such that  $N_P(E) \geq k$ .

Suppose  $E$  is an elliptic curve in  $G_P^{(1)}$  with equation  $E : y^2 = x^3 + a_4x + a_6$ . Assume  $E \in S_k$ . Then, using Proposition 2.4,  $l, m, n \in R_P$  exist such that

$$\left( y + \frac{l}{\pi_P^k}x + \frac{m}{\pi_P^{3k}} \right)^2 - \left( x + \frac{n}{\pi_P^{2k}} \right)^3 - \frac{a_4}{\pi_P^{4k}} \left( x + \frac{n}{\pi_P^{2k}} \right) - \frac{a_6}{\pi_P^{6k}} \in R_P[x, y].$$

The coefficient of  $xy$  is  $\frac{2l}{\pi_P^k}$ , giving that  $v_P(l) \geq k$ , and the coefficient of  $y$  is  $\frac{2m}{\pi_P^{3k}}$ , giving that  $v_P(m) \geq 3k$ . Also, the coefficient of  $x^2$  is  $\frac{3n-l^2}{\pi_P^{2k}}$ , giving that  $v_P(n) \geq 2k$ . From this, we have that  $v_P(a_4) \geq 4k$  and  $v_P(a_6) \geq 6k$ .

Define the function  $\phi_k : S_k \rightarrow S_0$ ,  $y^2 = x^3 + a_4x + a_6 \mapsto y^2 = x^3 + \frac{a_4}{\pi_P^{4k}}x + \frac{a_6}{\pi_P^{6k}}$ . Note that  $S_k \subset S_0 \subset G_P^{(1)}$ . From Proposition 2.5 and Lemma 4.1,  $\mu_P(S_0) = 1$ . Next, we show how we can use  $\phi_k$  to compute densities for  $S_k$ .

**Lemma 4.2.** If  $U$  is an open subset of  $G_P^{(1)}$ ,  $\mu_P(\phi_k^{-1}(U)) = \frac{1}{Q_P^{10k}}\mu_P(U)$ .

*Proof.* Suppose  $r_4, r_6 \in R_P$ . Also, suppose  $n_4$  and  $n_6$  are nonnegative integers. Let  $V$  be the set of elliptic curves  $y^2 = x^3 + a'_4x + a'_6$  with  $a'_4 \in r_4 + \pi_P^{n_4}R_P$  and  $a'_6 \in r_6 + \pi_P^{n_6}R_P$ . Because  $\mu_P(S_0) = 1$ ,  $\mu_P(V) = \frac{1}{Q_P^{n_4+n_6}}$ . To prove the lemma, it suffices to prove that

$$\mu_P(\phi_k^{-1}(V)) = \frac{1}{Q_P^{10k}}\mu_P(V) = \frac{1}{Q_P^{n_4+n_6+10k}}.$$

Suppose  $E : y^2 = x^3 + a_4x + a_6 \in G_P^{(1)}$  is an elliptic curve. We prove that  $E \in S_k$  and  $\phi_k(E) \in V$  if and only if  $\frac{a_4}{\pi_P^{4k}} \in r_4 + \pi_P^{n_4}R_P$  and  $\frac{a_6}{\pi_P^{6k}} \in r_6 + \pi_P^{n_6}R_P$ . If  $\phi_k(E) \in V$ , then  $\frac{a_4}{\pi_P^{4k}} \in r_4 + \pi_P^{n_4}R_P$  and  $\frac{a_6}{\pi_P^{6k}} \in r_6 + \pi_P^{n_6}R_P$ . Assume that  $\frac{a_4}{\pi_P^{4k}} \in r_4 + \pi_P^{n_4}R_P$  and  $\frac{a_6}{\pi_P^{6k}} \in r_6 + \pi_P^{n_6}R_P$ . From Tate's algorithm, we have that  $E \in S_k$ . Then, it is true that  $\phi_k(E) \in V$ .

Assume that  $E \in S_k$  and  $\phi_k(E) \in V$ . This is true if and only if  $a_4 \in \pi_P^{4k}r_4 + \pi_P^{n_4+4k}R$  and  $a_6 \in \pi_P^{6k}r_6 + \pi_P^{n_6+6k}R$ . Moreover, because  $\mu_P(S_0) = 1$ , the density of curves  $y^2 = x^3 + a_4x + a_6$  with discriminant 0 such that  $a_4 \in \pi_P^{4k}r_4 + \pi_P^{n_4+4k}R$  and  $a_6 \in \pi_P^{6k}r_6 + \pi_P^{n_6+6k}R$  is 0. Because of this,  $\mu_P(\phi_k^{-1}(V)) = \frac{1}{Q_P^{n_4+n_6+10k}}$ , completing the proof.  $\blacksquare$

**4.3. Density Calculations.** Note that the density of a set of curves in  $G_P^{(1)}$  is the Haar measure of the set. In this subsection, we compute the density of the set of minimal elliptic curves with a given Kodaira type and Tamagawa number over  $G_P^{(1)}$ . This can be extended to nonminimal elliptic curves using Theorem 1.2. Moreover, in this subsection, we use that the set of curves in  $G_P^{(1)}$  that have a discriminant equal to 0 has a Haar measure of 0.

Suppose the discriminant is not divisible by  $\pi_P$ . We compute the density for this set by considering  $a_4$  and  $a_6$  modulo  $\pi_P$ . Suppose  $a_4 \in r_4 + \pi_P R_P$  and  $a_6 \in r_6 + \pi_P R_P$ . We find the number of pairs  $(r_4, r_6)$  in  $L_{P,1}^2$  such that  $\left(\frac{r_4}{3}\right)^3 + \left(\frac{r_6}{2}\right)^2 \equiv 0 \pmod{\pi_P}$ . If  $r_4 = 0$ ,  $r_6$  has 1 choice, and if  $-\frac{r_4}{3}$  is a square modulo  $\pi_P$ ,  $r_6$  has 2 choices. Otherwise,  $r_6$  has 0 choices. We see that the number of pairs  $(r_4, r_6)$  is  $Q_P$ . Therefore, where each pair  $(r_4, r_6)$  has a density of  $\frac{1}{Q_P^2}$ , the density of the discriminant not being divisible by  $\pi_P$  is  $\frac{Q_P-1}{Q_P}$ . For this case, Tate's algorithm ends in step 1 and we get that  $\delta_K(I_0, 1, 0; P) = \frac{Q_P-1}{Q_P}$ .

Next, assume that the discriminant is divisible by  $\pi_P$ . Furthermore, assume that  $a_4, a_6 \not\equiv 0 \pmod{\pi_P}$ . Because there are  $Q_P - 1$  pairs  $(r_4, r_6)$  in  $L_{P,1}^2$  for this case, the total density is  $\frac{Q_P-1}{Q_P^2}$ . Let  $\alpha$  be the element of  $L_{P,1}$  such that  $a_4 \equiv -3\alpha^2 \pmod{\pi_P}$  and  $a_6 \equiv 2\alpha^3 \pmod{\pi_P}$ . The singular point is  $(\alpha, 0)$  and in step 2,  $x$  is replaced with  $x + n$  where  $n = \alpha$ . Because  $\alpha \not\equiv 0 \pmod{\pi_P}$ , Tate's algorithm ends in step 2. The quadratic considered in step 2 is  $T^2 - 3\alpha$ . We see that for  $\frac{Q_P-1}{2}$  values of  $\alpha$ , this quadratic has roots in  $R_P/\pi_P R_P$  and  $c = v_P(\Delta(E))$ . Otherwise,  $c = 1$  if  $v_P(\Delta(E))$  is odd and  $c = 2$  if  $v_P(\Delta(E))$  is even.

Let  $N$  be a positive integer. Suppose  $a_4 \in r_4 + \pi_P^N R_P$  and  $a_6 \in r_6 + \pi_P^N R_P$ . We find the number of pairs  $(r_4, r_6)$  in  $L_{P,1}^2$  such that  $\left(\frac{r_4}{3}\right)^3 + \left(\frac{r_6}{2}\right)^2 \equiv 0 \pmod{\pi_P^N}$  and  $r_4, r_6 \neq 0$ . Because there are  $\frac{Q_P^N - Q_P^{N-1}}{2}$  nonzero residues that are squares modulo  $\pi_P^M$ , we have that

the number of pairs  $(r_4, r_6)$  is  $Q_P^N - Q_P^{N-1}$ . Therefore, the density of  $v_P(\Delta(E)) \geq N$  for  $a_4, a_6 \not\equiv 0 \pmod{\pi_P}$  is  $\frac{Q_P-1}{Q_P^{N+1}}$ .

Suppose  $N$  is a positive integer. The density of  $v_P(\Delta(E)) = N$  is  $\frac{Q_P-1}{Q_P^{N+1}} - \frac{Q_P-1}{Q_P^{N+2}} = \frac{(Q_P-1)^2}{Q_P^{N+2}}$ . We therefore have that  $\delta_K(I_1, 1, 0; P) = \frac{(Q_P-1)^2}{Q_P^3}$ ,  $\delta_K(I_2, 2, 0; P) = \frac{(Q_P-1)^2}{Q_P^4}$ , and

$$\delta_K(I_N, N, 0; P) = \delta_K\left(I_N, 2 \left\lfloor \frac{N}{2} \right\rfloor - N + 2, 0; P\right) = \frac{(Q_P - 1)^2}{2Q_P^{N+2}}$$

for  $N \geq 3$ .

If  $v_P(a_4), v_P(a_6) \geq 1$ , the singular point modulo  $\pi_P$  from step 2 of Tate's algorithm is  $(0, 0)$ . The total density for this case is  $\frac{1}{Q_P^2}$ . If  $v_P(a_6) = 1$ , the algorithm ends in step 3.

For this, we get  $\delta_K(II, 1, 0; P) = \frac{Q_P-1}{Q_P^3}$ .

Assume that  $v_P(a_6) \geq 2$ . The total density for this case is  $\frac{1}{Q_P^3}$ . If  $v_P(a_4) = 1$ , the algorithm ends in step 4, and we get that  $\delta_K(III, 2, 0; P) = \frac{Q_P-1}{Q_P^4}$ .

Next, suppose  $v_P(a_4) \geq 2$ . The total density for this case is  $\frac{1}{Q_P^4}$ . If  $v_P(a_6) = 2$ , the algorithm ends in step 5. From this, we have that  $\delta_K(IV, 1, 0; P) = \delta_K(IV, 3, 0; P) = \frac{Q_P-1}{2Q_P^5}$ .

Suppose  $v_P(a_6) \geq 3$ . The total density for this case is  $\frac{1}{Q_P^5}$ . In step 6, the polynomial  $P(T) \in (R_P/\pi_P R_P)[T]$  has coefficient of  $T^2$  equal to 0. From adding multiples of  $\pi_P^2$  to  $a_4$ , the choices for the coefficient of  $T$  are  $L_{P,1}$ . Also, from adding multiples of  $\pi_P^3$  to  $a_6$ , the choices for the constant term are  $L_{P,1}$ . Then, we have that each polynomial  $P(T) \in (R_P/\pi_P R_P)[T]$  with coefficient of  $T^2$  equal to 0 corresponds to a density of  $\frac{1}{Q_P^7}$  in  $G_P^{(1)}$ .

Assume  $P(T)$  has distinct roots in  $\overline{R_P/\pi_P R_P}$ . The total number of  $P(T)$  for this case is  $Q_P^2 - Q_P$ ; therefore, the total density for this case is  $\frac{Q_P-1}{Q_P^6}$ . We have that Tate's algorithm ends in step 6 here. The number of  $P(T)$  with 0, 1, and 3 roots in  $R_P/\pi_P R_P$  is  $\frac{Q_P^2-1}{3}$ ,  $\frac{Q_P^2-Q_P}{2}$ , and  $\frac{Q_P^2-3Q_P+2}{6}$ , respectively. With this,  $\delta_K(I_0^*, 1, 0; P) = \frac{Q_P^2-1}{3Q_P^7}$ ,  $\delta_K(I_0^*, 2, 0; P) = \frac{Q_P-1}{2Q_P^6}$ , and  $\delta_K(I_0^*, 4, 0; P) = \frac{Q_P^2-3Q_P+2}{6Q_P^7}$ .

Next, assume that  $P(T)$  has a double root and a simple root in  $\overline{R_P/\pi_P R_P}$ . Then, Tate's algorithm enters the subprocedure in step 7. For this case, the total number of  $P(T)$  is  $Q_P - 1$  and the total density is therefore  $\frac{Q_P-1}{Q_P^7}$ . In Section 4.4, we compute that  $\delta_K(I_N^*, 2, 0; P) = \delta_K(I_N^*, 4, 0; P) = \frac{(Q_P-1)^2}{2Q_P^{N+7}}$  for all positive integers  $N$ .

Assume  $P(T)$  has a triple root in  $\overline{R_P/\pi_P R_P}$ . For this case, the total number of  $P(T)$  is 1 and the total density is therefore  $\frac{1}{Q_P^8}$ . Because the coefficient of  $T^2$  in  $P(T)$  is 0, the triple root is 0. If  $v_P(a_6) = 4$ , the algorithm ends in step 8. For this case,  $\delta_K(IV^*, 1, 0; P) = \delta_K(IV^*, 3, 0; P) = \frac{Q_P-1}{2Q_P^9}$ .

Next, assume that  $v_P(a_6) \geq 5$ . The total density for this case is  $\frac{1}{Q_P^9}$ . If  $v_P(a_4) = 3$ , the algorithm ends in step 9. We then have that  $\delta_K(III^*, 2, 0; P) = \frac{Q_P-1}{Q_P^9}$ .

Suppose  $v_P(a_4) \geq 4$ . The total density for this case is  $\frac{1}{Q_P^9}$ . If  $v_P(a_6) = 5$ , the algorithm ends in step 10. Therefore,  $\delta_K(II^*, 1, 0; P) = \frac{Q_P - 1}{Q_P^{10}}$ .

With density  $\frac{1}{Q_P^{10}}$ , we have that  $v_P(a_4) \geq 4$  and  $v_P(a_6) \geq 6$ , meaning that the curve is not minimal. That is, the curve will complete iteration 1 and continue iteration 2. Note that the density of nonminimal curves calculated from the algorithm matches Lemma 4.2.

**4.4. Subprocedure Density Calculations.** Next, we study the densities for the subprocedure in step 7 of Tate's algorithm. We compute the subprocedure densities by studying the translation of  $x$  in Tate's algorithm. In the step 7 subprocedure, because the coefficient of  $y$  is initially 0, there will be no translations of  $y$ .

Let  $X$  be the set of elliptic curves  $E \in G_P^{(1)}$  such that  $N_P(E) = 0$  and Tate's algorithm enters the step 7 subprocedure when used on  $E$ . For  $E \in X$ , let  $L(E)$  be the number of iterations of the step 7 subprocedure that are completed when Tate's algorithm is used on  $E$ . For a nonnegative integer  $N$ , let  $X_N$  be the set of  $E \in X$  such that  $L(E) \geq N$ .

Suppose  $N$  is an even nonnegative integer. Iteration  $N$  of the step 7 subprocedure is completed if and only if  $n \in R_P$  exists such that  $v_P(n) = 1$ ,  $v_P(a_4 + 3n^2) \geq \frac{N+6}{2}$ , and  $v_P(n^3 + 3na_4 + a_6) \geq N + 4$ . Suppose  $n = n_1$  satisfies this condition. Suppose  $n = n_2$  also satisfies this condition. We then have that  $n_1^2 \equiv n_2^2 \pmod{\pi_P^{\frac{N+6}{2}}}$ . This gives that  $n_1$  is equivalent to  $n_2$  or  $-n_2$  modulo  $\pi_P^{\frac{N+4}{2}}$ . However, because  $n_1^3 + n_1a_4 \equiv n_2^3 + n_2a_4 \pmod{\pi_P^{N+4}}$ , we have that  $v_P(n_1 - n_2) \geq \frac{N+4}{2}$ . Moreover, if  $v_P(n_1 - n_2) \geq \frac{N+4}{2}$ ,  $n = n_2$  also satisfies the condition.

Next, suppose  $N$  is an odd nonnegative integer. Iteration  $N$  of the subprocedure is completed if and only if  $n \in R_P$  exists such that  $v_P(n) = 1$ ,  $v_P(a_4 + 3n^2) \geq \frac{N+5}{2}$ , and  $v_P(n^3 + na_4 + a_6) \geq N + 4$ . Similarly, we have that if  $n = n_1$  satisfies the condition,  $n = n_2$  satisfies the condition if and only if  $v_P(n_1 - n_2) \geq \frac{N+3}{2}$ .

Suppose  $N$  is a nonnegative integer. Suppose  $n$  is an element of  $L_{P, \lfloor \frac{N+4}{2} \rfloor}$  such that  $v_P(n) = 1$ . Let  $Y_{n,N}$  be the set of curves  $x^3 + 3nx^2 + a'_4x + a'_6$  such that  $v_P(a'_4) \geq \lfloor \frac{N+6}{2} \rfloor$  and  $v_P(a'_6) \geq N + 4$ . Note that  $Y_{n,N}$  can be considered to be an open subset of  $R_P^2$ .

For  $E \in X_N$ , let  $n_N(E)$  be the unique value of  $n \in L_{P, \lfloor \frac{N+4}{2} \rfloor}$  such that  $v_P(n) = 1$ ,  $v_P(a_4 + 3n^2) \geq \lfloor \frac{N+6}{2} \rfloor$ , and  $v_P(n^3 + na_4 + a_6) \geq N + 4$ . Let  $\theta_N$  be the function such that if  $E : y^2 = x^3 + a_4x + a_6$  is an element of  $X_N$ ,

$$\begin{aligned} \theta_N(E) : y^2 &= (x + n_N(E))^3 + a_4(x + n_N(E)) + a_6 \\ &= x^3 + 3n_N(E)x^2 + (a_4 + 3n_N(E)^2)x + n_N(E)a_4 + a_6 + n_N(E)^3. \end{aligned}$$

**Lemma 4.3.** If  $U$  is an open subset of  $Y_{n,N}$ ,  $\mu_P(\theta_N^{-1}(U)) = \mu_P(U)$ .

*Proof.* Suppose  $r_4, r_6 \in R_P$ . Also, suppose  $n_4$  and  $n_6$  are nonnegative integers. Assume that  $v_P(r_4), n_4 \geq \lfloor \frac{N+4}{2} \rfloor$  and  $v_P(r_6), n_6 \geq N + 4$ . Let  $V \subset Y_{n,N}$  be the set of  $E' : y^2 = x^3 + 3nx^2 + a'_4x + a'_6$  such that  $a'_4 \in r_4 + \pi_P^{n_4}R_P$  and  $a'_6 \in r_6 + \pi_P^{n_6}R_P$ . It suffices to prove that  $\mu_P(\theta_N^{-1}(V)) = \mu_P(V)$ . Suppose  $E : y^2 = x^3 + a_4x + a_6$  is an elliptic curve.

We prove that that  $E \in X_N$  and  $\theta_N(E) \in V$  if and only if

$$a_4 + 3n^2 \in r_4 + \pi_P^{n_4}R_P, na_4 + a_6 + n^3 \in r_6 + \pi_P^{n_6}R_P.$$

Assume that  $E \in X_N$  and  $\theta_N(E) \in V$ . Because  $\theta_N(E) \in V$ , we have that  $n_N(E) = n$ . Therefore,  $a_4 + 3n^2 \in r_4 + \pi_P^{n_4} R_P$  and  $na_4 + a_6 + n^3 \in r_6 + \pi_P^{n_6} R_P$ . Next, assume that  $a_4 + 3n^2 \in r_4 + \pi_P^{n_4}$  and  $na_4 + a_6 + n^3 \in r_6 + \pi_P^{n_6} R_P$ . Because  $v_P(a_4 + 3n^2) \geq \lfloor \frac{N+6}{2} \rfloor$  and  $v_P(na_4 + a_6 + n^3) \geq N + 4$ ,  $E \in X_N$ . We then have that  $\theta_N(E) \in V$ .

Let  $M = \max(n_4, n_6)$ . Modulo  $\pi_P^M$ , there are  $Q_P^{M-n_4}$  choices for the residue of  $a_4$ . After choosing  $a_4$  modulo  $\pi_P^M$ , there are  $Q_P^{M-n_6}$  choices for the residue of  $a_6$  modulo  $\pi_P^M$ . Each of these combinations of residues modulo  $\pi_P^M$  for  $a_4$  and  $a_6$  has a density of  $\frac{1}{Q_P^{2M}}$  in  $G_P^{(1)}$ . The Haar measure of the  $Q_P^{2M-n_4-n_6}$  combinations is  $\frac{1}{Q_P^{n_4+n_6}}$ . Because the set of curves in  $G_P^{(1)}$  with discriminant 0 has a Haar measure of 0,

$$\mu_P(\theta_N^{-1}(V)) = \frac{1}{Q_P^{n_4+n_6}} = \mu_P(V).$$

This finishes the proof.  $\blacksquare$

Let  $N$  be a positive integer. We compute the density of  $I_N^*$ . Let  $n$  be an element of  $L_P, \lfloor \frac{N+3}{2} \rfloor$  such that  $v_P(n) = 1$ . We have that the Haar measure of the set of  $E \in Y_{n, N-1}$  that do not complete iteration  $N$  is  $\frac{Q_P-1}{Q_P^{\lfloor \frac{N+3}{2} \rfloor + N+4}}$ . With Lemma 4.3, because there are  $(Q_P-1)Q_P^{\lfloor \frac{N-1}{2} \rfloor}$  values of  $n$ , the density of  $I_N^*$  is  $\frac{(Q_P-1)^2}{Q_P^{N+7}}$ . From adding multiples of  $\pi_P^{N+4}$  to  $a_6$ ,  $c = 2$  and  $c = 4$  have equal density. Therefore,

$$\delta_K(I_N^*, 2, 0; P) = \delta_K(I_N^*, 4, 0; P) = \frac{(Q_P-1)^2}{2Q_P^{N+7}}.$$

## 5. LOCAL DENSITIES FOR $p = 3$

**5.1. Setup.** Suppose that the characteristic of  $K$  is  $p = 3$ . Let  $P$  be a place of  $K$  and  $G_P^{(2)}$  be the set of curves

$$y^2 = x^3 + a_2x^2 + a_4x + a_6$$

over  $K_P$  such that  $a_2, a_4, a_6 \in R_P$ . Note that  $G_P^{(2)}$  can be considered to be  $R_P^3$ . Define  $\varphi : G_P \rightarrow G_P^{(2)}$  as the function such that if  $E$  is a curve in  $G_P$ ,  $\varphi(E)$  is the curve in  $G_P^{(2)}$  with equation

$$y^2 = x^3 + \frac{b_2(E)}{4}x^2 + \frac{b_4(E)}{2}x + \frac{b_6(E)}{4}.$$

Note that if  $E$  is an elliptic curve,  $E$  and  $\varphi(E)$  are equivalent.

**Lemma 5.1.** If  $U$  is an open subset of  $G_P^{(2)}$ ,  $\mu_P(\varphi^{-1}(U)) = \mu_P(U)$ .

*Proof.* This can be proved using a method similar to the proof of Lemma 4.1.  $\blacksquare$

**5.2. Multiple Iterations.** Let  $k$  be a nonnegative integer. Suppose  $S_k$  is the set of elliptic curves  $E \in G_P^{(2)}$  such that  $N_P(E) \geq k$ .

Suppose  $E \in S_k$  has equation  $E : y^2 = x^3 + a_2x^2 + a_4x + a_6$ . From Proposition 2.4,  $l, m, n \in R_P$  exist such that

$$\left(y + \frac{l}{\pi_P^k}x + \frac{m}{\pi_P^{3k}}\right)^2 = \left(x + \frac{n}{\pi_P^{2k}}\right)^3 + \frac{a_2}{\pi_P^{2k}}\left(x + \frac{n}{\pi_P^{2k}}\right)^2 + \frac{a_4}{\pi_P^{4k}}\left(x + \frac{n}{\pi_P^{2k}}\right) + \frac{a_6}{\pi_P^{6k}}$$

has coefficients in  $R_P$ . From the coefficient of  $xy$ ,  $v_P(l) \geq k$ , and from the coefficient of  $y$ ,  $v_P(m) \geq 3k$ . Therefore, we have that

$$y^2 = \left(x + \frac{n}{\pi_P^{2k}}\right)^3 + \frac{a_2}{\pi_P^{2k}} \left(x + \frac{n}{\pi_P^{2k}}\right)^2 + \frac{a_4}{\pi_P^{4k}} \left(x + \frac{n}{\pi_P^{2k}}\right) + \frac{a_6}{\pi_P^{6k}}$$

has coefficients in  $R_P$ . Note that  $v_P(a_2) \geq 2k$  also.

For an elliptic curve  $E \in G_P^{(2)}$  with equation  $E : y^2 = x^3 + a_2x^2 + a_4x + a_6$ , let  $A_k(E)$  be the set of  $n \in R_P$  such that

$$y^2 = x^3 + \frac{a_2}{\pi_P^{2k}}x^2 + \frac{2na_2 + a_4}{\pi_P^{4k}}x + \frac{n^2a_2 + na_4 + a_6 + n^3}{\pi_P^{6k}}$$

has coefficients in  $R_P$ . The next proposition is useful for computing local densities for multiple iterations.

**Proposition 5.2.** Let  $E$  be an elliptic curve in  $G_P^{(2)}$ .  $E \in S_k$  if and only if a unique element  $n \in L_{P,k}$  exists such that  $n \in A_k(E)$ .

*Proof.* Assume a unique element  $n \in L_{P,k}$  exists such that  $n \in A_k(E)$ . Then,  $A_k(E)$  is nonempty, and using Proposition 2.4,  $E \in S_k$ .

Next, assume  $E \in S_k$ . From Proposition 2.4, we have that  $A_k(E)$  is nonempty. Let the equation of  $E$  be  $E : y^2 = x^3 + a_2x^2 + a_4x + a_6$  for  $a_2, a_4, a_6 \in R_P$ .

Suppose  $n \in A_k(E)$ . From replacing  $x$  with  $x+n'$  for  $n' \in R_P$ , we have that  $n+n'\pi_P^{2k} \in A_k(E)$ . Therefore,  $n \in L_{P,k}$  exists such that  $n \in A_k(E)$ .

Next, we prove uniqueness. Assume  $n_1, n_2 \in A_k(E) \cap L_{P,k}$ . Let

$$F : y^2 = x^3 + \frac{a_2}{\pi_P^{2k}}x^2 + \frac{a_4}{\pi_P^{4k}}x + \frac{a_6}{\pi_P^{6k}}.$$

For  $1 \leq i \leq 2$ , let  $F_i$  be  $F$  with  $x$  replaced by  $x + \frac{n_i}{\pi_P^{2k}}$ . Note that  $F_1, F_2 \in G_P^{(2)}$ .

From the coefficients of  $x$  in  $F_1$  and  $F_2$ ,

$$2n_1a_2 + a_4 \equiv 2n_2a_2 + a_4 \equiv 0 \pmod{\pi_P^{4k}}.$$

Also, from the constant terms of  $F_1$  and  $F_2$ ,

$$n_1^2a_2 + n_1a_4 + n_1^3 \equiv n_2^2a_2 + n_2a_4 + n_2^3 \pmod{\pi_P^{6k}}.$$

For the sake of contradiction, assume that  $v_P(n_1 - n_2) < 2k$ . Let  $a = v_P(n_1 - n_2)$ . Note that

$$v_P(n_1^3 - n_2^3) = v_P((n_1 - n_2)^3) = 3a.$$

We have that

$$n_1^2a_2 + n_1a_4 - n_2^2a_2 - n_2a_4 = (n_1 - n_2)(n_1a_2 + n_2a_2 + a_4).$$

Because  $a_4 \equiv n_1a_2 \equiv n_2a_2 \pmod{\pi_P^{4k}}$ ,

$$n_1a_2 + n_2a_2 + a_4 \equiv 3a_4 \equiv 0 \pmod{\pi_P^{4k}}.$$

From this,

$$\begin{aligned} v_P(n_1^2a_2 + n_1a_4 - n_2^2a_2 - n_2a_4) &= v_P((n_1 - n_2)(n_1a_2 + n_2a_2 + a_4)) \\ &\geq a + 4k \\ &> 3a. \end{aligned}$$

Since  $v_P(n_1^3 - n_2^3) = 3a$ ,

$$v_P(n_1^2 a_2 + n_1 a_4 + n_1^3 - n_2^2 a_2 - n_2 a_4 - n_2^3) = 3a < 6k,$$

which is a contradiction. Therefore,  $v_P(n_1 - n_2) \geq 2k$  and  $n_1 = n_2$ .  $\blacksquare$

Using Proposition 5.2, for  $E \in S_k$ , let  $n(E)$  be the unique  $n \in L_{P,2k}$  such that the  $n \in A_k(E)$ . Define  $\phi_k : S_k \rightarrow S_0$  to be the function such that if  $E \in S_k$  has equation  $E : y^2 = x^3 + a_2 x^2 + a_4 x + a_6$ ,  $\phi_k(E) \in S_0$  has equation

$$\phi_k(E) : y^2 = x^3 + \frac{a_2}{\pi_P^{2k}} x^2 + \frac{2n(E)a_2 + a_4}{\pi_P^{4k}} x + \frac{n(E)^2 a_2 + n(E)a_4 + a_6 + n(E)^3}{\pi_P^{6k}}.$$

Note that  $S_k \subset S_0 \subset G_P^{(2)}$ . Also, using Proposition 2.5 and Lemma 5.1,  $\mu_P(S_0) = 1$ . For  $n \in L_{P,2k}$ , suppose  $S_{k,n}$  is the set of  $E \in S_k$  such that  $n(E) = n$ , and let  $\phi_{k,n}$  be  $\phi_k$  restricted to  $S_{k,n}$ .

**Lemma 5.3.** Suppose  $n \in L_{P,k}$ . If  $U$  is an open subset of  $G_P^{(2)}$ ,  $\mu_P(\phi_{k,n}^{-1}(U)) = \frac{1}{Q_P^{12k}} \mu_P(U)$ .

*Proof.* Suppose  $r_2, r_4, r_6 \in R_P$ . Also, suppose  $n_2, n_4$ , and  $n_6$  are nonnegative integers. Let  $V$  be the set of  $y^2 = x^3 + a'_2 x^2 + a'_4 x + a'_6$  such that  $a'_2 \in r_2 + \pi_P^{n_2} R_P$ ,  $a'_4 \in r_4 + \pi_P^{n_4} R_P$ , and  $a'_6 \in r_6 + \pi_P^{n_6} R_P$ . Suppose  $E : y^2 = x^3 + a_2 x^2 + a_4 x + a_6 \in G_P^{(2)}$ .  $E \in S_{k,n}$  and  $\phi_{k,n}(E) \in V$  if and only if

$$\frac{a_2}{\pi_P^{2k}} \in r_2 + \pi_P^{n_2} R_P, \frac{2na_2 + a_4}{\pi_P^{4k}} \in r_4 + \pi_P^{n_4} R_P, \frac{n^2 a_2 + na_4 + a_6 + n^3}{\pi_P^{6k}} \in r_6 + \pi_P^{n_6} R_P.$$

Assume that  $E \in S_{k,n}$  and  $\phi_{k,n}(E) \in V$ . Let  $M = \max(n_2 + 2k, n_4 + 4k, n_6 + 6k)$ . There are  $Q_P^{M-n_2-2k}$  ways to pick  $a_2$  modulo  $\pi_P^M$ . Afterwards,  $a_4$  will have  $Q_P^{M-n_4-4k}$  choices for the residue modulo  $\pi_P^M$ ; pick  $a_4$  modulo  $\pi_P^M$ . Next,  $a_6$  has  $Q_P^{M-n_6-6k}$  choices for the residue modulo  $\pi_P^M$ . Select the residue for  $a_6$ . The number of combinations of residues is  $Q_P^{3M-n_2-n_4-n_6-12k}$  and each combination of residues has a Haar measure of  $Q_P^{-3M}$ . Also, because  $\mu_P(S_0) = 1$ , the set of curves with discriminant 0 counted in these combinations of residues has a Haar measure 0. Therefore,  $\mu_P(\phi_{k,n}^{-1}(V)) = \frac{1}{Q_P^{n_2+n_4+n_6+12k}}$ . With this,  $\mu_P(\phi_{k,n}^{-1}(U)) = \frac{1}{Q_P^{12k}} \mu_P(U)$  for all open subsets  $U$  of  $G_P^{(2)}$ .  $\blacksquare$

**Lemma 5.4.** If  $U$  is an open subset of  $G_P^{(2)}$ ,  $\mu_P(\phi_k^{-1}(U)) = \frac{1}{Q_P^{10k}} \mu_P(U)$ .

*Proof.* Let  $U$  be an open subset of  $G_P^{(2)}$ . We have that  $\phi_k^{-1}(U) = \bigsqcup_{n \in L_{P,2k}} \phi_{k,n}^{-1}(U)$ . Using Lemma 5.3,

$$\mu_P(\phi_k^{-1}(U)) = \sum_{n \in L_{P,2k}} \mu_P(\phi_{k,n}^{-1}(U)) = \sum_{n \in L_{P,2k}} \frac{1}{Q_P^{12k}} \mu_P(U) = \frac{1}{Q_P^{10k}} \mu_P(U),$$

completing the proof.  $\blacksquare$

**5.3. Density Calculations for  $v_P(a_2) = 0$ .** Suppose  $v_P(a_2) = 0$ . The density for this case over  $G_P^{(2)}$  is  $\frac{Q_P-1}{Q_P}$ . The discriminant is  $-a_2^3a_6 + a_2^2a_4^2 - a_4^3$ .

From adding multiples of  $\pi_P$  to  $a_6$ , the set of curves with discriminant not divisible by  $\pi_P$  has density  $\frac{(Q_P-1)^2}{Q_P^2}$ . Then, we add  $\frac{(Q_P-1)^2}{Q_P^2}$  to  $\delta_K(I_0, 1, 0; P)$ .

Assume the discriminant is divisible by  $\pi_P$ . The algorithm ends in step 2. Because  $v_P(a_2) = 0$ , the coefficient of  $a_6$  in the discriminant is not divisible by  $\pi_P$ . Then, we see that for  $N \geq 0$ , the density over  $G_P^{(2)}$  of curves such that  $v_P(a_2) = 0$  and  $v_P(\Delta(E)) = N$  is  $\frac{(Q_P-1)^2}{Q_P^{N+2}}$ . If  $a_2 \equiv r_2 \pmod{\pi_P}$  for  $r_2 \in L_{P,1}$  such that  $r_2 \neq 0$ ,  $T^2 + a_2$  is irreducible over  $R_P/\pi_P R_P$  for  $\frac{Q_P-1}{2}$  values of  $r_2$ . Using step 2 of Tate's algorithm, we have that  $\delta_K(I_1, 1, 0; P) = \frac{(Q_P-1)^2}{Q_P^3}$ ,  $\delta_K(I_2, 2, 0; P) = \frac{(Q_P-1)^2}{Q_P^4}$ , and

$$\delta_K(I_N, N, 0; P) = \delta_K\left(I_N, 2 \left\lfloor \frac{N}{2} \right\rfloor - N + 2, 0; P\right) = \frac{(Q_P - 1)^2}{2Q_P^{N+2}}$$

for  $N \geq 3$ .

**5.4. Density Calculations for  $v_P(a_2) \geq 1$ .** Next, suppose  $v_P(a_2) \geq 1$ . The density for this is  $\frac{1}{Q_P}$  and modulo  $\pi_P$ , the discriminant is  $-a_4^3$ .

Assume the discriminant is not divisible by  $\pi_P$ . This occurs if and only if  $a_4$  is not divisible by  $\pi_P$ , and the density of this case is  $\frac{Q_P-1}{Q_P^2}$ . Adding this density to  $\delta_K(I_0, 1, 0; P)$  gives that  $\delta_K(I_0, 1, 0; P) = \frac{Q_P-1}{Q_P}$ .

Next, assume the discriminant is divisible by  $\pi_P$ . The total density for these cases will be  $\frac{1}{Q_P^2}$ . Suppose  $\alpha_1$  is an element of  $L_{P,1}$  such that  $a_6 + \alpha_1^3 \equiv 0 \pmod{\pi_P}$ . A singular point is  $(\alpha_1, 0)$ . We have that  $x$  is replaced with  $x + n$  where  $n = \alpha_1$ . The resulting curve has equation

$$y^2 = (x + n)^3 + a_2(x + n)^2 + a_4(x + n) + a_6.$$

We have that  $n^2a_2 + na_4 + a_6 + n^3$  is not divisible by  $\pi_P^2$  with density  $\frac{Q_P-1}{Q_P^3}$  by adding multiples of  $\pi_P$  to  $a_6$ . For this case,  $\delta_K(II, 1, 0; P) = \frac{Q_P-1}{Q_P^3}$ .

Assume  $n^2a_2 + na_4 + a_6 + n^3$  is divisible by  $\pi_P^2$ . The total density for this case is  $\frac{1}{Q_P^3}$ . The density of  $v_P(2na_2 + a_4) = 1$  is  $\frac{Q_P-1}{Q_P^4}$  from replacing  $a_4$  with  $a_4 + \pi_P d$  and  $a_6$  with  $a_6 - \alpha_1 \pi_P d$  for  $d \in L_{P,1}$ . If  $v_P(2na_2 + a_4) = 1$ , the algorithm ends in step 4. We then have that  $\delta_K(III, 2, 0; P) = \frac{Q_P-1}{Q_P^4}$ .

Assume  $2na_2 + a_4$  is divisible by  $\pi_P^2$ . The total density for this case is  $\frac{1}{Q_P^4}$ . We have that  $v_P(n^2a_2 + na_4 + a_6 + n^3) = 2$  with density  $\frac{Q_P-1}{Q_P^5}$  from adding multiples of  $\pi_P^2$  to  $a_6$ . If this is true, the algorithm ends in step 5. Afterwards, we have that  $\delta_K(IV, 1, 0; P) = \delta_K(IV, 3, 0; P) = \frac{Q_P-1}{2Q_P^5}$ .

Suppose  $v_P(n^2a_2 + na_4 + a_6 + n^3) \geq 3$ . The total density for this case is  $\frac{1}{Q_P^5}$ . In step 6, there is no translation. Suppose  $a_2$  is replaced by  $a_2 + d_1 \pi_P$ ,  $a_4$  is replaced with  $a_4 - 2\alpha_1 d_1 \pi_P$ , and  $a_6$  is replaced with  $a_6 + \alpha_1^2 d_1 \pi_P$  for  $d_1 \in L_{P,1}$ . Note that the previous parts of the algorithm will not be changed. However, this changes the coefficient of  $x^2$  from  $a_2$  to  $a_2 + d_1 \pi_P$ , which changes the coefficient of  $T^2$  of  $P(T)$  in step 6. Next, replace

$a_4$  with  $a_4 + d_2\pi_P^2$  and  $a_6$  with  $a_6 - \alpha_1 d_2\pi_P^2$  for  $d_2 \in \pi_P$ . Similarly, this does not change the previous parts of the algorithm. However,  $d_2\pi_P^2$  will be added to the coefficient of  $x$ , which adds  $d_2$  to the coefficient of  $T$  of  $P(T)$ . Afterwards, replace  $a_6$  with  $a_6 + d_3\pi_P^3$  for  $d_3 \in L_{P,1}$ . This adds  $d_3$  to the constant term  $P(T)$ . With this, the choices for  $P(T)$  are the monic polynomials with degree 3 in  $(R_P/\pi_P R_P)[T]$ ; each choice for  $P(T)$  corresponds to a density of  $\frac{1}{Q_P^8}$ . Moreover, the number of  $P(T)$  with a double root and triple root are  $Q_P(Q_P - 1)$  and  $Q_P$ , respectively.

Assume  $P(T)$  has distinct roots. We have that the algorithm ends in step 6, with  $\delta_K(I_0^*, 1, 0; P) = \frac{Q_P^2 - 1}{3Q_P^7}$ ,  $\delta_K(I_0^*, 2, 0; P) = \frac{Q_P - 1}{2Q_P^6}$ , and  $\delta_K(I_0^*, 4, 0; P) = \frac{Q_P^2 - 3Q_P + 2}{6Q_P^7}$ .

Assume  $P(T)$  has a double root. For this case, Tate's algorithm ends in step 7 and the total density is  $\frac{Q_P - 1}{Q_P^7}$ . In Section 5.5, we compute that  $\delta_K(I_N^*, 2, 0; P) = \delta_K(I_N^*, 4, 0; P) = \frac{(Q_P - 1)^2}{2Q_P^{N+7}}$  for all positive integers  $N$ .

Next, assume  $P(T)$  has a triple root. The density for this case is  $\frac{1}{Q_P^7}$ . Let  $\alpha_2$  be the element of  $L_{P,1}$  such that

$$n^2 a_2 + n a_4 + a_6 + n^3 \equiv -\pi_P^3 \alpha_2^3 \pmod{\pi_P^4}.$$

Then, for the translation in step 8, we let  $n = \alpha_1 + \alpha_2 \pi_P$ . Suppose  $v_P(n^2 a_2 + n a_4 + a_6 + n^3) = 4$ . This occurs with density  $\frac{Q_P - 1}{Q_P^8}$  by adding multiples of  $\pi_P^4$  to  $a_6$ . In this case, Tate's algorithm ends in step 8, and  $\delta_K(IV^*, 1, 0; P) = \delta_K(IV^*, 3, 0; P) = \frac{Q_P - 1}{2Q_P^8}$ .

Assume  $v_P(n^2 a_2 + n a_4 + a_6 + n^3) \geq 5$ . The total density for this case is  $\frac{1}{Q_P^8}$ . Consider replacing  $a_4$  with  $a_4 + d\pi_P^3$  and  $a_6$  with  $a_6 - (\alpha_1 + \alpha_2 \pi_P)d\pi_P^3$  for  $d \in L_{P,1}$ . This does not change previous parts of the algorithm but adds  $d\pi_P^3$  to the coefficient of  $x$ . Therefore,  $v_P(2na_2 + a_4) = 3$  with density  $\frac{Q_P - 1}{Q_P^9}$ . For this, we have that Tate's algorithm ends in step 9 and  $\delta_K(III^*, 2, 0; P) = \frac{Q_P - 1}{Q_P^9}$ .

Suppose  $v_P(2na_2 + a_4) \geq 4$ . The total density of this case is  $\frac{1}{Q_P^9}$ . From adding multiples of  $\pi_P^6$  to  $a_6$ ,  $v_P(n^3 + a_2 n^2 + a_4 n + a_6) = 5$  with density  $\frac{Q_P - 1}{Q_P^{10}}$ . Also, if  $v_P(n^3 + a_2 n^2 + a_4 n + a_6) = 5$ , the algorithm ends in step 10. This gives that  $\delta_K(II^*, 1, 0; P) = \frac{Q_P - 1}{Q_P^{10}}$ .

**5.5. Subprocedure Density Calculations.** Let  $X$  be the set of elliptic curves  $E \in G_P^{(2)}$  such that  $N_P(E) = 0$  and Tate's algorithm enters the step 7 subprocedure when used on  $E$ . For  $E \in X$ , let  $L(E)$  be the number of iterations of the step 7 subprocedure that are completed when Tate's algorithm is used on  $E$ . For a nonnegative integer  $N$ , let  $X_N$  be the set of  $E \in X$  such that  $L(E) \geq N$ .

Suppose  $N$  is an even nonnegative integer. Iteration  $N$  of the step 7 subprocedure is completed if and only if  $n \in R_P$  exists such that  $v_P(a_2) = 1$ ,  $v_P(2na_2 + a_4) \geq \frac{N+6}{2}$ , and  $v_P(n^3 + n^2 a_2 + n a_4 + a_6) \geq N + 4$ . Assume  $n = n_1$  satisfies the condition. Suppose  $n = n_2$  satisfies the condition also. Because  $v_P(a_2) = 1$ ,  $v_P(n_1 - n_2) \geq \frac{N+4}{2}$ . Next, assume  $v_P(n_1 - n_2) \geq \frac{N+4}{2}$ . We show that  $n = n_2$  also satisfies the condition. Clearly,  $v_P(2n_2 a_2 + a_4) \geq \frac{N+6}{2}$ . Moreover, we have that

$$n_2^2 a_2 + n_2 a_4 = n_1^2 a_2 + n_1 a_4 + \frac{1}{2}(n_2 - n_1)((2n_1 a_2 + a_4) + (2n_2 a_2 + a_4)).$$

Therefore,  $v_P(n_2^3 + n_2^2 a_2 + n_2 a_4 + a_6) \geq N + 4$ . We have that  $n = n_2$  satisfies the condition if and only if  $v_P(n_1 - n_2) \geq \frac{N+4}{2}$ .

Next, suppose  $N$  is an odd nonnegative integer. Iteration  $N$  of the step 7 subprocedure is completed if and only if  $n \in R_P$  exists such that  $v_P(n^2 a_2 + n a_4 + a_6 + n^3) \geq N + 4$  and  $v_P(2n a_2 + a_4) \geq \frac{N+5}{2}$ . Assume  $n = n_1$  satisfies the condition. Similarly to when  $N$  is even, we have that  $n = n_2$  also satisfies the condition if and only if  $v_P(n_1 - n_2) \geq \frac{N+3}{2}$ .

Suppose  $N$  is a nonnegative integer. Let  $Y_N$  be the set of curves  $y^2 = x^3 + a_2' x^2 + a_4' x + a_6'$  with  $v_P(a_2') = 1$ ,  $v_P(a_4') \geq \lfloor \frac{N+6}{2} \rfloor$ , and  $v_P(a_6') \geq N + 4$ . For  $E \in X_N$ , let  $n_N(E)$  be the unique value of  $n$  in  $L_P, \lfloor \frac{N+4}{2} \rfloor$  from above. Suppose  $\theta_N(E)$ , with  $\theta_N : X_N \rightarrow Y_N$ , is the curve

$$\begin{aligned} \theta_N(E) : y^2 &= (x + n_N(E))^3 + a_2(x + n_N(E))^2 + a_4(x + n_N(E)) + a_6 \\ &= x^3 + a_2 x^2 + (2n_N(E)a_2 + a_4)x + n_N(E)^2 a_2 + n_N(E)a_4 + a_6. \end{aligned}$$

**Lemma 5.5.** If  $U$  is an open subset of  $Y_N$ ,  $\mu_P(\theta_N^{-1}(U)) = Q_P^{\lfloor \frac{N+4}{2} \rfloor} \mu_P(U)$ .

*Proof.* Suppose  $n \in L_P, \lfloor \frac{N+4}{2} \rfloor$ . Let  $X_{N,n}$  be the set of  $E \in X_N$  with  $n_N(E) = n$  and  $\theta_{N,n}$  be  $\theta_N$  restricted to  $X_{N,n}$ . Suppose  $U$  is an open subset of  $Y_N$ . Using a method similar to the proof of Lemma 4.3, we have that

$$\mu_P(\theta_{N,n}^{-1}(U)) = \mu_P(U).$$

Because there are  $Q_P^{\lfloor \frac{N+4}{2} \rfloor}$  values of  $n$ , the result follows.  $\blacksquare$

Suppose  $N$  is a positive integer. Using Lemma 5.5, we can compute the density of the curves  $E$  with  $N_P(E) = 0$  that have type  $I_N^*$  and Tamagawa number 2 or 4. The Haar measure of the curves in  $Y_{N-1}$  that end in iteration  $N$  is  $\frac{(Q_P-1)^2}{Q_P^{N+6+\lfloor \frac{N+5}{2} \rfloor}}$ . With Lemma 4.1,

we have that  $\delta_K(I_N^*, 2, 0; P) = \delta_K(I_N^*, 4, 0; P) = \frac{(Q_P-1)^2}{2Q_P^{N+7}}$ .

## 6. LOCAL DENSITIES FOR $p = 2$

**6.1. Setup.** Assume that the characteristic of  $K$  is  $p = 2$ . Let  $P$  be a place of  $K$  and  $G_P^{(3)}$  be the set of curves

$$y^2 + a_1 xy + a_3 y = x^3 + a_4 x + a_6$$

over  $K_P$  such that  $a_1, a_3, a_4, a_6 \in R_P$ . Note that  $G_P^{(3)}$  can be considered to be  $R_P^4$ . Define  $\varphi : G_P \rightarrow G_P^{(3)}$  as the function such that if  $E$  is the curve in  $G_P$  with equation  $E : y^2 + a_1 xy + a_3 y = x^3 + a_2 x^2 + a_4 x + a_6$ ,  $\varphi(E)$  is the curve in  $G_P^{(3)}$  with equation

$$\varphi(E) : y^2 + a_1 xy + \left(a_3 - \frac{a_1 a_2}{3}\right) y = x^3 + \left(a_4 - \frac{a_2^2}{3}\right) x + \frac{2a_2^3}{27} - \frac{a_2 a_4}{3} + a_6.$$

Note that if  $E$  is an elliptic curve,  $E$  and  $\varphi(E)$  are equivalent.

**Lemma 6.1.** If  $U$  is an open subset of  $G_P^{(3)}$ ,  $\mu_P(\varphi^{-1}(U)) = \mu_P(U)$ .

*Proof.* This can be proved using a method similar to the proof of Lemma 4.1.  $\blacksquare$

**6.2. Multiple Iterations.** Let  $k$  be a nonnegative integer. Suppose  $S_k$  is the set of elliptic curves  $E \in G_P^{(3)}$  such that  $N_P(E) \geq k$ .

For an elliptic curve  $E \in G_P^{(3)}$  with equation  $E : y^2 + a_1xy + a_3y = x^3 + a_4x + a_6$ , let  $A_k(E)$  be the set of  $(l, m, n) \in R_P^3$  such that if  $X = x + \frac{n}{\pi_P^{2k}}$  and  $Y = y + \frac{l}{\pi_P^k}x + \frac{m}{\pi_P^{3k}}$ ,

$$\begin{aligned} & \left( y + \frac{l}{\pi_P^k}x + \frac{m}{\pi_P^{3k}} \right)^2 + \frac{a_1}{\pi_P^k} \left( x + \frac{l^2 + a_1l}{\pi_P^{2k}} \right) \left( y + \frac{l}{\pi_P^k}x + \frac{m}{\pi_P^{3k}} \right) + \frac{a_3}{\pi_P^{3k}} \left( y + \frac{l}{\pi_P^k}x + \frac{m}{\pi_P^{3k}} \right) \\ & - \left( x + \frac{l^2 + a_1l}{\pi_P^{2k}} \right)^3 - \frac{a_4}{\pi_P^{4k}} \left( x + \frac{l^2 + a_1l}{\pi_P^{2k}} \right) - \frac{a_6}{\pi_P^{6k}} \in R_P[x, y]. \end{aligned}$$

**Proposition 6.2.** Let  $E$  be an elliptic curve in  $G_P^{(3)}$ .  $E \in S_k$  if and only if a unique pair  $(l, m) \in L_{P,k} \times L_{P,3k}$  exists such that  $(l, m, l^2 + a_1l) \in A_k(E)$ .

*Proof.* Suppose a unique pair  $(l, m)$  satisfying the conditions exists. Because  $A_k(E)$  is nonempty,  $E \in S_k$  from Proposition 2.4.

Assume  $E \in S_k$ . Then, using Proposition 2.4,  $A_k(E)$  is nonempty. Let the equation of  $E$  be  $E : y^2 + a_1xy + a_3y = x^3 + a_4x + a_6$  for  $a_1, a_3, a_4, a_6 \in R_P$ .

From replacing  $y$  with  $y + l'x$  for  $l' \in R_P$ , if  $(l, m, n) \in A_k(E)$ ,  $(l + l'\pi_P^k, m, n) \in A_k(E)$ . Therefore, there exist  $l \in L_{P,k}$  and  $m, n \in R_P$  such that  $(l, m, n) \in A_k(E)$ . Moreover, if  $(l, m, n) \in A_k(E)$ ,  $l^2 + a_1l + n \equiv 0 \pmod{\pi_P^{2k}}$ . With this, from replacing  $x$  with  $x + \frac{l^2 + a_1l + n}{\pi_P^{2k}}$ , if  $(l, m, n) \in A_k(E)$ ,  $(l, m + l(l^2 + a_1l + n), l^2 + a_1l) \in A_k(E)$ . Therefore, there exist  $l \in L_{P,k}$  and  $m \in R_P$  such that  $(l, m, l^2 + a_1l) \in A_k(E)$ . Next, from replacing  $y$  with  $y + m'$  for  $m' \in R_P$ , there exists  $l \in L_{P,k}$  and  $m \in L_{P,3k}$  such that  $(l, m, l^2 + a_1l) \in A_k(E)$ .

Next, we prove that  $(l, m)$  is unique. Assume that  $(l_1, m_1), (l_2, m_2) \in L_{P,k} \times L_{P,3k}$  and  $(l_1, m_1, l_1^2 + a_1l_1), (l_2, m_2, l_2^2 + a_1l_2) \in A_k(E)$ . We prove that  $(l_1, m_1) = (l_2, m_2)$ .

Let  $F$  be the curve

$$F : y^2 + \frac{a_1}{\pi_P^k}xy + \frac{a_3}{\pi_P^{3k}}y = x^3 + \frac{a_4}{\pi_P^k}x + \frac{a_6}{\pi_P^{6k}}.$$

For  $1 \leq i \leq 2$ , let  $F_i$  be  $F$  with  $x$  replaced by  $x + \frac{l_i^2 + a_1l_i}{\pi_P^{2k}}$  and  $y$  replaced by  $y + \frac{l_i}{\pi_P^k}x + \frac{m_i}{\pi_P^{3k}}$ . Note that  $F_i \in G_P^{(3)}$  because  $(l_i, m_i, l_i^2 + a_1l_i) \in A_k(E)$  for  $1 \leq i \leq 2$ . From this,  $a_1 \equiv 0 \pmod{\pi_P^k}$ .

Suppose  $a_1 \neq 0$ . We have that  $F_1$  and  $F_2$  are equivalent and  $v_P(\Delta(F_1)) = v_P(\Delta(F_2))$ . Then, using Proposition 2.3, let  $\tau$  be a translation from the equation of  $F_1$  to the equation of  $F_2$  that replaces  $x$  with  $u^2x + n'$  and  $y$  with  $u^3y + l'u^2x + m'$ , where  $u, l', m', n' \in R_P$  and  $v_P(u) = 0$ .

The coefficient of  $xy$  after  $\tau$  is applied to the equation of  $F_1$  is  $\frac{a_1}{u\pi_P^k}$ . However, the coefficient of  $xy$  in  $F_2$  is  $\frac{a_1}{\pi_P^k}$ . Therefore,  $u = 1$  and  $a_1 \equiv 0 \pmod{\pi_P^k}$ .

Next, the coefficient of  $y$  after  $\tau$  is applied to the equation of  $F_1$

$$\frac{a_1l_1^2 + a_1^2l_1 + a_3 + \pi_P^{2k}a_1n'}{\pi_P^{3k}}.$$

However, the coefficient of  $y$  in  $F_2$  is

$$\frac{a_1l_2^2 + a_1^2l_2 + a_3}{\pi_P^{3k}}.$$

Therefore,

$$l_1^2 + a_1 l_1 + \pi_P^{2k} n' = l_2^2 + a_1 l_2.$$

Because  $a_1 \equiv 0 \pmod{\pi_P^k}$ , we have that  $l_1 \equiv l_2 \pmod{\pi_P^k}$ . Therefore,  $l_1 = l_2$ . From this,  $n' = 0$ .

The coefficient of  $x^2$  after  $\tau$  is applied to the equation of  $F_1$  is

$$n' + (l')^2 + \frac{a_1 l'}{\pi_P^k}.$$

This equals the coefficient of  $x^2$  in  $F_2$ , which is 0. Because  $n' = 0$ , we have that  $l' = 0$  or  $l' = \frac{a_1}{\pi_P^k}$ .

From setting the coefficient of  $x$  after  $\tau$  is applied to the equation of  $F_1$  equal to the coefficient of  $x$  in  $F_2$ ,

$$\frac{a_1}{\pi_P^k} \cdot \left( \frac{m_1}{\pi_P^{3k}} + m' \right) + \frac{a_1(l_1^2 + a_1 l_1) + a_3}{\pi_P^{3k}} \cdot l' = \frac{a_1}{\pi_P^k} \cdot \frac{m_2}{\pi_P^{3k}}.$$

Suppose  $l' = 0$ . Then  $\frac{m_1}{\pi_P^{3k}} + m' = \frac{m_2}{\pi_P^{3k}}$ . It follows that  $m_1 \equiv m_2 \pmod{\pi_P^{3k}}$  and  $m_1 = m_2$ . Suppose  $l' = \frac{a_1}{\pi_P^k}$ . We have that

$$\frac{m_1}{\pi_P^{3k}} + m' + \frac{a_1(l_1^2 + a_1 l_1) + a_3}{\pi_P^{3k}} = \frac{m_2}{\pi_P^{3k}}.$$

However, using that the coefficient of  $y$  in  $F_2$  is an element of  $R_P$ ,

$$a_1(l_1^2 + a_1 l_1) + a_3 \equiv a_1(l_2^2 + a_1 l_2) + a_3 \equiv 0 \pmod{\pi_P^{3k}}.$$

Therefore,  $m_1 \equiv m_2 \pmod{\pi_P^{3k}}$  and  $m_1 = m_2$ .

Assume  $a_1 = 0$ . From the coefficient of  $y$  in  $F_2$ , we have that  $a_3 \equiv 0 \pmod{\pi_P^{3k}}$ . Also, from the coefficients of  $x$  in  $F_1$  and  $F_2$ ,  $l_1^4 + a_3 l_1 \equiv l_2^4 + a_3 l_2 \pmod{\pi_P^{4k}}$ . This gives that  $l_1 = l_2$ . Afterwards, from the constant terms of  $F_1$  and  $F_2$ ,  $m_1^2 + a_3 m_1 \equiv m_2^2 + a_3 m_2 \pmod{\pi_P^6}$ . From this, we obtain that  $m_1 = m_2$ .  $\blacksquare$

Using Proposition 6.2, for  $E \in S_k$ , let the unique pair  $(l, m) \in L_{P,k} \times L_{P,3k}$  such that  $(l, m, l^2 + a_1 l) \in A_k(E)$  be  $(l(E), m(E))$ . Define  $\phi_k : S_k \rightarrow S_0$  to be the function such that if  $E \in S_k$  has equation  $E : y^2 + a_1 xy + a_3 y = x^3 + a_4 x + a_6$ ,  $\phi_k(E)$  has equation

$$\begin{aligned} \phi_k(E) : y^2 + \frac{a_1}{\pi_P^k} xy + \frac{a_1(l(E)^2 + a_1 l(E)) + a_3}{\pi_P^{3k}} &= x^3 + \\ \frac{l(E)^2(l(E)^2 + a_1 l(E)) + a_1 m(E) + a_3 l(E) + a_4}{\pi_P^{4k}} x + & \\ \frac{(a_1 m(E) + a_4 + a_1^2 l(E)^2 + l(E)^4)(l(E)^2 + a_1 l(E)) + a_3 m(E) + a_6 + m(E)^2}{\pi_P^{6k}} &. \end{aligned}$$

The equation for  $\phi_k(E)$  is equivalent to

$$\begin{aligned} & \left( y + \frac{l(E)}{\pi_P^k} x + \frac{m(E)}{\pi_P^{3k}} \right)^2 + \frac{a_1}{\pi_P^k} \left( x + \frac{l(E)^2 + a_1 l(E)}{\pi_P^{2k}} \right) \left( y + \frac{l(E)}{\pi_P^k} x + \frac{m(E)}{\pi_P^{3k}} \right) + \\ & \frac{a_3}{\pi_P^{3k}} \left( y + \frac{l(E)}{\pi_P^k} x + \frac{m(E)}{\pi_P^{3k}} \right) = \\ & \left( x + \frac{l(E)^2 + a_1 l(E)}{\pi_P^{2k}} \right)^3 + \frac{a_4}{\pi_P^{4k}} \left( x + \frac{l(E)^2 + a_1 l(E)}{\pi_P^{2k}} \right) + \frac{a_6}{\pi_P^{6k}}. \end{aligned}$$

Note that  $S_0 \subset G_P^{(3)}$ , and from Proposition 2.5 and Lemma 6.1,  $\mu_P(S_0) = 1$ . For  $l \in L_{P,k}$  and  $m \in L_{P,3k}$ , let  $S_{k,l,m}$  be the set of  $E \in S_k$  such that  $l(E) = l$  and  $m(E) = m$ . Assume that  $\phi_{k,l,m}$  is  $\phi_k$  restricted to  $S_{k,l,m}$ .

**Lemma 6.3.** Suppose  $l \in L_{P,k}$  and  $m \in L_{P,3k}$ . If  $U$  is an open subset of  $G_P^{(3)}$ ,  $\mu_P(\phi_{k,l,m}^{-1}(U)) = \frac{1}{Q_P^{14k}} \mu_P(U)$ .

*Proof.* This can be proved with a method that is similar to the proof of Lemma 5.3.  $\blacksquare$

**Lemma 6.4.** If  $U$  is an open subset of  $G_P^{(3)}$ ,  $\mu_P(\phi_k^{-1}(U)) = \frac{1}{Q_P^{10k}} \mu_P(U)$ .

*Proof.* Let  $U$  be an open subset of  $G_P^{(3)}$ . We have that  $\phi_k^{-1}(U) = \bigsqcup_{l \in L_{P,k}, m \in L_{P,3k}} \phi_{k,l,m}^{-1}(U)$ . Using Lemma 6.3,

$$\mu_P(\phi_k^{-1}(U)) = \sum_{l \in L_{P,k}} \sum_{m \in L_{P,3k}} \mu_P(\phi_{k,l,m}^{-1}(U)) = \sum_{l \in L_{P,k}} \sum_{m \in L_{P,3k}} \frac{1}{Q_P^{14k}} \mu_P(U) = \frac{1}{Q_P^{10k}} \mu_P(U),$$

completing the proof.  $\blacksquare$

**6.3. Density Calculations for  $v_P(a_1) = 0$ .** Suppose that  $v_P(a_1) = 0$ . This case has density  $\frac{Q_P-1}{Q_P}$ . The discriminant is

$$a_1^4(a_1^2 a_6 + a_1 a_3 a_4 + a_4^2) + a_3^4 + a_1^3 a_3^3.$$

Note that by considering  $a_6$  modulo  $\pi_P$ , the discriminant is not divisible by  $\pi_P$  with density  $\frac{(Q_P-1)^2}{Q_P^2}$ . For this case, the algorithm ends in step 1. Then, we add  $\frac{(Q_P-1)^2}{Q_P^2}$  to  $\delta_K(I_0, 1, 0; P)$ .

Assume the discriminant is divisible by  $\pi_P$ . Let  $(\alpha_1, \alpha_2)$  be the singular point modulo  $\pi_P$ ; it can be proven that  $\alpha_1, \alpha_2 \in R_P$ . Also,  $\alpha_1 \equiv -\frac{a_3}{a_1} \pmod{\pi_P}$ . In step 2, replace  $x$  by  $x+n$  and  $y$  by  $y+m$  with  $n = \alpha_1$  and  $m = \alpha_2$ . Afterwards, the coefficient of  $xy$  is  $a_1$ , which is not divisible by  $\pi_P$ . The algorithm then ends in step 2.

We see that the discriminant is linear in  $a_6$ . Therefore, we have that  $v_P(a_1) = 0$  and  $v_P(\Delta(E)) = N$  with density  $\frac{(Q_P-1)^2}{Q_P^{N+2}}$  for  $N \geq 0$ . Note that the polynomial considered in step 2 is  $T^2 + a_1 T + \alpha_1$ . Suppose  $a_1 \equiv r_1 \pmod{\pi_P}$  and  $a_3 \equiv r_3 \pmod{\pi_P}$  for  $r_1, r_3 \in L_{P,1}$  such that  $r_1 \neq 0$ . Given  $r_1$ ,  $T^2 + a_1 T + \alpha_1$  is irreducible over  $R_P/\pi_P R_P$  for  $\frac{Q_P}{2}$  values of  $r_3$ . Afterwards, using step 2 of Tate's algorithm, we get that in this case,

$\delta_K(I_1, 1, 0; P) = \frac{(Q_P-1)^2}{Q_P^3}$ ,  $\delta_K(I_2, 2, 0; P) = \frac{(Q_P-1)^2}{Q_P^4}$ , and

$$\delta_K(I_N, N, 0; P) = \delta_K\left(I_N, 2 \left\lfloor \frac{N}{2} \right\rfloor - N + 2, 0; P\right) = \frac{(Q_P-1)^2}{2Q_P^{N+2}}$$

for  $N \geq 3$ .

**6.4. Density Calculations for  $v_P(a_1) \geq 1$ .** In this subsection, we assume that  $v_P(a_1) \geq 1$ . The density for this is  $\frac{1}{Q_P}$ , and the discriminant modulo  $\pi_P$  is  $a_3^4$ .

Suppose  $v_P(a_3) = 0$ . The density for this case is  $\frac{Q_P-1}{Q_P^2}$ . Here, the discriminant is not divisible by  $\pi_P$ . Tate's algorithm then ends in step 1, and we add  $\frac{Q_P-1}{Q_P^2}$  to  $\delta_K(I_0, 1, 0; P)$ . Following this, we obtain that  $\delta_K(I_0, 1, 0; P) = \frac{Q_P-1}{Q_P}$ .

Next, assume  $v_P(a_3) \geq 1$ . The total density for this is  $\frac{1}{Q_P}$ . The singular point modulo  $\pi_P$  is  $(x, y) = (\alpha_1, \alpha_2)$  for  $\alpha_1, \alpha_2 \in L_{P,1}$  such that  $a_4 \equiv \alpha_1^2 \pmod{\pi_P}$  and  $a_6 \equiv \alpha_2^2 \pmod{\pi_P}$ . We replace  $x$  with  $x + n$  and  $y$  with  $y + m$ , where  $n = \alpha_1$  and  $m = \alpha_2$ . The curve is

$$(y + m)^2 + a_1(x + n)(y + m) + a_3(y + m) = (x + n)^3 + a_4(x + n) + a_6.$$

If  $\pi_P^2$  does not divide  $mna_1 + ma_3 + na_4 + a_6 + m^2 + n^3$ , the algorithm ends in step 3. By adding multiples of  $\pi_P$  to  $a_6$ , this occurs with density  $\frac{Q_P-1}{Q_P^3}$ . We have that  $\delta_K(II, 1, 0; P) = \frac{Q_P-1}{Q_P^3}$ .

Assume  $\pi_P^2$  divides  $mna_1 + ma_3 + na_4 + a_6 + m^2 + n^3$ . The total density for this case is  $\frac{1}{Q_P}$ . We have that

$$b_8 = n(na_1 + a_3)^2 + (ma_1 + a_4 + n^2)^2.$$

If  $b_8$  is not divisible by  $\pi_P^3$ , the algorithm ends in step 4. By adding multiples of  $\pi_P$  to  $a_4$ , we have that  $\delta_K(III, 2, 0; P) = \frac{Q_P-1}{Q_P^4}$ .

Assume  $b_8$  is divisible by  $\pi_P^3$ . The total density for this case is  $\frac{1}{Q_P}$ . If  $v_P(na_1 + a_3) = 1$ , the algorithm ends in step 5. Assume  $a_4 \equiv 0 \pmod{\pi_P}$ . Then, replace  $a_3$  with  $a_3 + d\pi_P$  and  $a_4$  with  $a_4 + \beta d\pi_P$  for  $\beta, d \in L_{P,1}$  such that  $\beta^2 \equiv \alpha_1 \pmod{\pi_P}$ . This will not affect previous parts of the algorithm; particularly, this will not change  $b_8$  modulo  $\pi_P^3$ . However,  $na_1 + a_3$  will be increased by  $d\pi_P$ . Therefore, we have that  $v_P(na_1 + a_3) = 1$  with density  $\frac{Q_P-1}{Q_P^5}$ . From this,  $\delta_K(IV, 1, 0; P) = \delta_K(IV, 3, 0; P) = \frac{Q_P-1}{2Q_P^5}$ .

Assume  $v_P(na_1 + a_3) \geq 2$ . The total density for this case is  $\frac{1}{Q_P}$ . Let  $\alpha_3$  be the element of  $L_{P,1}$  such that  $n \equiv \alpha_3^2 \pmod{\pi_P}$ . Also, let  $\alpha_4$  be the element of  $L_{P,1}$  such that  $mna_1 + ma_3 + na_4 + a_6 + m^2 + n^3 \equiv \alpha_4^2 \pi_P^2 \pmod{\pi_P^3}$ . After the transformation in step 6, the equation of the curve is

$$\begin{aligned} & (y + lx + m)^2 + a_1(x + n)(y + lx + m) + a_3(y + lx + m) \\ & = (x + n)^3 + a_4(x + n) + a_6, \end{aligned}$$

where  $n = \alpha_1$ ,  $l = \alpha_3$ , and  $m = \alpha_2 + \alpha_4\pi_P$ . Suppose that in step 6, the polynomial  $P(T) \in (R_P/\pi_P R_P)[T]$  is  $P(T) = T^3 + w_2T^2 + w_1T + w_0$ .

Suppose  $a_4 \equiv 0 \pmod{\pi_P}$ . Because  $0 \in L_{P,1}$ , we have that  $n = l = 0$ . This means that  $w_2 = 0$ . Then, we can replace  $a_4$  with  $a_4 + d_1\pi_P^2$  for  $d_1 \in L_{P,1}$ , and the previous

parts of the algorithm will not be changed. With this, the choices for  $w_1$  modulo  $\pi_P$  are the elements of  $L_{P,1}$ . Following this, from replacing  $a_6$  with  $a_6 + d_2\pi_P^3$  for  $d_2 \in L_{P,1}$ , the choices for  $w_0$  modulo  $\pi_P$  are the elements of  $L_{P,1}$ . We have that the number of  $P(T)$  with a double root and no roots in  $\overline{R_P/\pi_P R_P}$  are  $Q_P - 1$  and  $1$ , respectively. Moreover, we have that the number of  $P(T)$  with 3 distinct roots in  $\overline{R_P/\pi_P R_P}$  and 0 roots, 1 root, and 3 roots in  $R_P/\pi_P R_P$  are  $\frac{Q_P^2-1}{3}$ ,  $\frac{Q_P^2-Q_P}{2}$ , and  $\frac{Q_P^2-3Q_P+2}{6}$ , respectively.

Suppose  $a_4 \not\equiv 0 \pmod{\pi_P}$ . Consider the translation of replacing  $a_1$  with  $a_1 + d_1\pi_P$ ,  $a_3$  with  $a_3 + \alpha_1 d_1\pi_P$ ,  $a_4$  with  $a_4 + (\alpha_2 + \alpha_4\pi_P)d_1\pi_P$ , and  $a_6$  with  $a_6 + \alpha_1(\alpha_2 + \alpha_4\pi_P)d_1\pi_P$  for  $d_1 \in L_{P,1}$ . After this, the parts of the algorithm before step 6 do not change. In step 6,  $w_0$  and  $w_1$  do not change. However,  $w_2$  increases by  $\alpha_3 d_1$ . Because  $\alpha_3 \neq 0$ , the choices for  $w_2$  are the elements of  $L_{P,1}$ . Next, replace  $a_6$  with  $a_6 + d_2\pi_P^3$  for  $d_2 \in L_{P,1}$ . With this, the choices for  $w_0$  are also the elements of  $L_{P,1}$ . The number of  $P(T)$  with a double root and no roots in  $\overline{R_P/\pi_P R_P}$  are the same as above. Also, the number of  $P(T)$  with 3 distinct roots in  $\overline{R_P/\pi_P R_P}$  and 0 roots, 1 root, and 3 roots in  $R_P/\pi_P R_P$  are the same as above.

Suppose  $P(T)$  has distinct roots. For this case, the total density is  $\frac{Q_P-1}{Q_P^6}$  and Tate's algorithm ends in step 6. We see that  $\delta_K(I_0^*, 1, 0; P) = \frac{Q_P^2-1}{3Q_P^7}$ ,  $\delta_K(I_0^*, 2, 0; P) = \frac{Q_P-1}{2Q_P^6}$ , and  $\delta_K(I_0^*, 4, 0; P) = \frac{Q_P^2-3Q_P+2}{6Q_P^7}$ .

Assume that  $P(T)$  has a double root and a simple root. For this case, the total density is  $\frac{Q_P-1}{Q_P^7}$  and Tate's algorithm ends in step 7. In Section 6.5, we compute that  $\delta_K(I_N^*, 2, 0; P) = \delta_K(I_N^*, 4, 0; P) = \frac{(Q_P-1)^2}{2Q_P^{N+7}}$  for all positive integers  $N$ .

Next, suppose  $P(T)$  has a triple root. For this case, the density is  $\frac{1}{Q_P^7}$ , and the root of  $P(T)$  is  $\sqrt{w_1}$  modulo  $\pi_P$ . If  $a_4 \equiv 0 \pmod{\pi_P}$ , the triple root is 0 modulo  $\pi_P$ . Let  $\alpha_5$  be an element of  $L_{P,1}$  such that

$$(m + ln)a_1 + la_3 + a_4 + n^2 \equiv \alpha_5^2 \pi_P^2 \pmod{\pi_P^3}.$$

Then, the translation in step 8 sets  $n$  to be  $n = \alpha_1 + \alpha_5\pi_P$ .

Suppose  $a_4 \equiv 0 \pmod{\pi_P}$ . Replace  $a_3$  with  $a_3 + d\pi_P^2$  and  $a_6$  with  $a_6 + (\alpha_2 + \alpha_4\pi_P)d\pi_P^2$  for some  $d \in L_{P,1}$ . Then, note that the previous parts of the algorithm, including  $P(T)$ , are unchanged. However, the coefficient of  $y$  increases by  $d\pi_P^2$ . We have that for one value of  $d$ , the coefficient of  $y$  is divisible by  $\pi_P^3$ . Next, suppose  $a_4 \not\equiv 0 \pmod{\pi_P}$ . Replace  $a_1$  with  $a_1 + d\pi_P^2$  and  $a_4$  with  $a_4 + (\alpha_2 + \alpha_4\pi_P)d\pi_P^2$  for some  $d \in L_{P,1}$ . The previous parts of the algorithm, including  $P(T)$ , are unchanged. However, the coefficient of  $y$  increases by  $(\alpha_1 + \alpha_5\pi_P)d\pi_P^2$ . Similarly, we have that for one value of  $d$ , the coefficient of  $y$  is divisible by  $\pi_P^3$ . From this, we get that the coefficient of  $y$  is not divisible by  $\pi_P^3$  and the algorithm ends in step 8 with density  $\frac{Q_P-1}{Q_P^8}$ . We then have that  $\delta_K(IV^*, 1, 0; P) = \delta_K(IV^*, 3, 0; P) = \frac{Q_P-1}{2Q_P^8}$ .

Assume the coefficient of  $y$  is divisible by  $\pi_P^3$ . The total density of this case is  $\frac{1}{Q_P^8}$ . Let  $\alpha_6$  be the element of  $L_{P,1}$  such that

$$mna_1 + ma_3 + na_4 + a_6 + m^2 + n^3 \equiv \alpha_6^2 \pi_P^4 \pmod{\pi_P^5}.$$

Then,  $m$  is set to  $m = \alpha_2 + \alpha_4\pi_P + \alpha_6\pi_P^2$  in step 9. If  $\pi_P^4$  does not divide the  $x$  coefficient of this curve, the algorithm ends in step 9. Consider the translation of replacing  $a_4$  with

$a_4 + d\pi_P^3$  and  $a_6$  with  $a_6 + (\alpha_1 + \alpha_5\pi_P)d\pi_P^3$  for  $d \in L_{P,1}$ . The previous parts of the algorithm do not change, but the coefficient of  $x$  is increased by  $d\pi_P^3$ . Therefore,  $\pi_P^4$  does not divide the  $x$  coefficient with density  $\frac{Q_P-1}{Q_P^9}$ . We have that  $\delta_K(III^*, 2, 0; P) = \frac{Q_P-1}{Q_P^9}$

Assume  $\pi_P^4$  divides the coefficient of  $x$  of the curve. The total density for this case is  $\frac{1}{Q_P^9}$ . If  $\pi_P^6$  does not divide  $mna_1 + ma_3 + na_4 + a_6 + m^2 + n^3$ , Tate's algorithm ends in step 10. This occurs with density  $\frac{Q_P-1}{Q_P^{10}}$  from adding multiples of  $\pi_P^6$  to  $a_6$ . We then have that  $\delta_K(II^*, 1, 0; P) = \frac{Q_P-1}{Q_P^{10}}$ .

**6.5. Subprocedure Density Calculations.** We calculate the density of Kodaira types  $\mathfrak{r} = I_N^*$  for  $N \geq 1$  and Tamagawa numbers  $n = 2, 4$ . Note that previously, the curve was reduced by removing  $a_2$  with a translation on  $x$  to obtain  $G_P^{(3)}$ . However, here the density is calculated in  $G_P$  without the reduction. That is, the density is calculated for curves in long Weierstrass form.

Let  $X$  be the set of elliptic curves  $E \in G_P$  such that  $N_P(E) = 0$  and Tate's algorithm enters the step 7 subprocedure when used on  $E$ . For  $E \in X$ , let  $L(E)$  be the number of iterations of the step 7 subprocedure that are completed when Tate's algorithm is used on  $E$ . For a nonnegative integer  $N$ , let  $X_N$  be the set of  $E \in X$  such that  $L(E) \geq N$ .

Suppose  $N$  is an even nonnegative integer. Assume that  $N = 0$ . In iteration  $N = 0$ , there is a translation. Note that the double root of  $P(T)$  is the squareroot of  $w_1$ . Because of this, in step 7, we add  $\gamma_0\pi_P$  to  $n$  and  $l\gamma_0\pi_P$  to  $m$  for some  $\gamma_0 \in L_{P,1}$  such that

$$(m + ln)a_1 + la_3 + a_4 + n^2 \equiv \gamma_0^2\pi_P^2 \pmod{\pi_P^3}$$

Next, assume  $N \geq 2$ . Suppose iteration  $N$  of the step 7 subprocedure is reached and the quadratic has a double root. Then,

$$v_P((m + ln)a_1 + la_3 + a_4 + n^2) \geq \frac{N + 6}{2}.$$

Also, we add  $\gamma_N\pi_P^{\frac{N+2}{2}}$  to  $n$  and  $l\gamma_N\pi_P^{\frac{N+2}{2}}$  to  $m$  for some  $\gamma_N \in L_{P,1}$  such that

$$mna_1 + ma_3 + na_4 + a_6 + m^2 + n^3 \equiv (la_1 + a_2 + n + l^2)\gamma_N^2\pi_P^{N+2} \pmod{\pi_P^{N+4}}.$$

Note that  $v_P(la_1 + a_2 + n + l^2) = 1$ .

Suppose  $N$  is an odd nonnegative integer. Suppose iteration  $N$  of the step 7 subprocedure is reached and the quadratic has a double root. Then,  $v_P(na_1 + a_3) \geq \frac{N+5}{2}$ . Also,  $\gamma_N\pi_P^{\frac{N+3}{2}}$  is added to  $m$  for some  $\gamma_N \in L_{P,1}$  such that

$$mna_1 + ma_3 + na_4 + a_6 + m^2 + n^3 \equiv \gamma_N^2\pi_P^{N+3} \pmod{\pi_P^{N+4}}$$

Let  $N$  be a nonnegative integer. Let  $Y_N$  be the set of curves  $y^2 + a'_1xy + a'_3y = x^3 + a'_2x^2 + a'_4x + a'_6$  with  $v_P(a'_1) \geq 1$ ,  $v_P(a'_2) = 1$ ,  $v_P(a'_3) \geq \lfloor \frac{N+5}{2} \rfloor$ ,  $v_P(a'_4) \geq \lfloor \frac{N+6}{2} \rfloor$ , and  $v_P(a'_6) \geq N + 4$ .

Suppose  $E \in X_N$  and that the translations of Tate's algorithm when it is used on  $E$  are  $\alpha_1, \alpha_2, \alpha_3, \alpha_4, \gamma_0, \gamma_1, \dots, \gamma_N$ . Let  $T_N(E) = (\alpha_1, \alpha_2, \alpha_3, \alpha_4, \gamma_0, \gamma_1, \dots, \gamma_N)$ . Note that because the characteristic of  $K$  is  $p = 2$ ,  $T_N(E)$  is well defined. Also, let  $\theta_N(E) : X_N \rightarrow Y_N$

be  $E$  with  $x$  replaced by  $x + n$  and  $y$  replaced by  $y + lx + m$ , where

$$n = \alpha_1 + \sum_{i=0}^{\lfloor \frac{N}{2} \rfloor} \gamma_{2i} \pi_P^{i+1}, l = \alpha_3, m = \alpha_2 + \alpha_4 \pi_P + \alpha_3 \sum_{i=0}^{\lfloor \frac{N}{2} \rfloor} \gamma_{2i} \pi_P^{i+1} + \sum_{i=0}^{\lfloor \frac{N-1}{2} \rfloor} \gamma_{2i+1} \pi_P^{i+2}.$$

**Lemma 6.5.** If  $U$  is an open subset of  $Y_N$ ,  $\mu_P(\theta_N^{-1}(U)) = Q_P^{N+5} \mu_P(U)$ .

*Proof.* Let  $a = (\alpha_1, \alpha_2, \alpha_3, \alpha_4, \gamma_0, \gamma_1, \dots, \gamma_N)_{0 \leq i \leq N}$  be an element of  $L_{P,1}^{N+5}$ . Suppose that  $X_{N,a}$  is the set of  $E \in X_N$  such that  $T_N(E) = a$ . Suppose that  $\theta_{N,a}$  is  $\theta_N$  restricted to  $X_{N,a}$ . Let  $U$  be an open subset of  $Y_N$ . Using a method similar to the proof of Lemma 4.3, we have that

$$\mu_P(\theta_{N,a}^{-1}(U)) = \mu_P(U).$$

Because there are  $Q_P^{N+5}$  choices of  $a$ , the result follows.  $\blacksquare$

Suppose  $N$  is a positive integer. With Lemma 6.5, we can compute the density for curves that enter step 7 in the first iteration and have type  $I_N^*$ . We have that  $\mu_P(Y_{N-1}) = \frac{Q_P-1}{Q_P^{2N+10}}$ , and the Haar measure in  $G_P^{(3)}$  of curves that have type  $I_N^*$  is then  $\frac{(Q_P-1)^2}{Q_P^{N+7}}$ . Particularly,  $\delta_K(I_N^*, 2, 0; P) = \delta_K(I_N^*, 4, 0; P) = \frac{(Q_P-1)^2}{2Q_P^{N+7}}$ .

## 7. LOCAL AND GLOBAL DENSITY RESULTS

In Section 4, Section 5, and Section 6, we compute the local densities of Kodaira types and Tamagawa numbers for  $p \geq 5$ ,  $p = 3$ , and  $p = 2$ , respectively. The methods we use involved first removing some terms from the equations of elliptic curves with translations, and then using translations to compute the local densities. Let  $\mathfrak{r}$  be a Kodaira type and  $n$  be a positive integer. Note that  $\delta_K(\mathfrak{r}, n; P)$  only depends on  $Q_P$ . Additionally, in [3], the local densities of  $\mathfrak{r}$  and the Tamagawa number  $n$  for elliptic curves in short Weierstrass form over  $\mathbb{Q}_r$  for primes  $r \geq 5$  have the same form as  $\delta_K(\mathfrak{r}, n; P)$  for global function fields  $K$  and  $P \in M_K$ . In [1], the local densities of  $\mathfrak{r}$  and the Tamagawa number  $n$  for elliptic curves in short Weierstrass form over completions of number fields at places that lie above primes  $r \geq 5$  also have the same form as  $\delta_K(\mathfrak{r}, n; P)$  for global function fields  $K$  and  $P \in M_K$ .

Next, we discuss some results about local and global densities, including a proof of Theorem 1.2. Particularly, we compute the density of completing at most  $k \geq 0$  iterations of Tate's algorithm.

**7.1. Proof of Theorem 1.2.** Let  $U$  and  $V$  be the sets of elliptic curves  $E \in G_P$  with Kodaira type  $\mathfrak{r}$  and Tamagawa number  $n$  such that  $N_P(E) = 0$  and  $N_P(E) = k$ , respectively. Note that  $\varphi(U)$  and  $\varphi(V)$  are the sets of curves  $E \in S_0$  with Kodaira type  $\mathfrak{r}$  and Tamagawa number  $n$  such that  $N_P(E) = 0$  and  $N_P(E) = k$ , respectively.

Suppose  $E \in G_P$  and  $\varphi(E) \in \varphi(U)$ . Then,  $E$  has Kodaira type  $\mathfrak{r}$ , Tamagawa number  $n$ , and  $N_P(E) = 0$ . This means that  $E \in U$ . From this,  $\varphi^{-1}(\varphi(U)) \subset U$ . Moreover,  $U \subset \varphi^{-1}(\varphi(U))$ . It follows that  $\varphi^{-1}(\varphi(U)) = U$ . Similarly,  $\varphi^{-1}(\varphi(V)) = V$ .

We have that  $U$  and  $V$  are open sets. Moreover,  $\varphi(U)$  and  $\varphi(V)$  are open sets. With this, we have that  $\mu_P(U) = \mu_P(\varphi(U))$  and  $\mu_P(V) = \mu_P(\varphi(V))$  for all characteristics  $p$

from Lemma 4.1, Lemma 5.1, and Lemma 6.1. Therefore, it suffices to prove that

$$\mu_P(\varphi(V)) = \frac{1}{Q_P^{10k}} \mu_P(\varphi(U)).$$

Suppose  $E \in \varphi(V)$ . We have that  $\phi_k(E)$  has Kodaira type  $\mathfrak{r}$ , Tamagawa number  $n$ , and  $N_P(\phi_k(E)) = 0$ . Therefore,  $\phi_k(E) \subset \varphi(U)$ . It follows that  $\varphi(V) \subset \phi_k^{-1}(\varphi(U))$ . Next, suppose  $E \in S_k$  and  $\phi_k(E) \in \varphi(U)$ . Then, the Kodaira type of  $E$  is  $\mathfrak{r}$  and the Tamagawa number of  $E$  is  $n$ . Moreover, because  $N_P(\phi_k(E)) = 0$ ,  $N_P(E) = k$ . It follows that  $E \in \varphi(V)$ . Therefore,  $\phi_k^{-1}(\varphi(U)) \subset \varphi(V)$ . From this,  $\phi_k^{-1}(\varphi(U)) = \varphi(V)$ . The result then follows from Lemma 4.2, Lemma 5.4, and Lemma 6.4.

**7.2. Density for Multiple Iterations.** Let  $k$  be a nonnegative integer. For  $P \in M_K$ , let  $U_P^k$  denote the set of elliptic curves  $E$  in  $G_P$  such that  $N_P(E) \geq k + 1$ . The following proposition is important for the proof of Theorem 7.2.

**Proposition 7.1.** For  $P \in M_K$ ,  $\mu_P(U_P^k) = \frac{1}{Q_P^{10(k+1)}}$ .

*Proof.* Suppose  $P \in M_k$ . From Lemma 4.2, Lemma 5.4, and Lemma 6.4 with  $k + 1$  as  $k$  and  $G_P$  as  $U$ , we have that

$$\mu_P(U_P^k) = \frac{1}{Q_P^{10(k+1)}} \cdot \mu_P(G_P) = \frac{1}{Q_P^{10(k+1)}}.$$

This finishes the proof. ■

**Theorem 7.2.** Let  $S$  be a finite nonempty subset of  $M_K$ . Suppose  $U$  is the set of elliptic curves in  $W_S$  such that  $N_P(E) \leq k$  for all  $P \in S^C$ . Then,

$$d_S(U) = \frac{1}{\zeta_K(10(k+1))} \cdot \prod_{P \in S} \left( \frac{Q_P^{10(k+1)}}{Q_P^{10(k+1)} - 1} \right).$$

*Proof.* For a positive integer  $M$ , let  $V_M$  be the set of elliptic curves  $E \in W_S$  such that there exists  $P \in S^C$  with degree at least  $M$  such that  $E \in U_P^k$ . From Proposition 3.3, we have that  $\lim_{M \rightarrow \infty} \bar{d}_S(V_M) = 0$ . Therefore, we can use Theorem 3.1 with  $U_P^k$  as  $U_P$  for  $P \in S^C$  and  $T = \{\}$ . The result follows from Proposition 7.1. ■

**Example 7.3.** We give an example of Theorem 7.2. Let  $K = \mathbb{F}_q(t)$ . Suppose  $P_\infty$  is the infinite place of  $\mathbb{F}_q(t)$  and let  $S = \{P_\infty\}$ . Let  $k$  be a nonnegative integer and  $U$  be the set of elliptic curves in  $W_S$  such that  $N_P(E) \leq k$  for all  $P \in S^C$ . From Theorem 5.9 of [5], because the genus of  $K$  is 0, we have that  $\zeta_K(10(k+1)) = \frac{q^{20k+19}}{(q^{10k+9}-1)(q^{10k+10}-1)}$ . Because  $P_\infty$  has degree 1, from Theorem 7.2,  $d_S(U) = 1 - \frac{1}{q^{10k+9}}$ .

## REFERENCES

- [1] Yunseo Choi, Sean Li, Apoorva Panidapu, and Casia Siegel, *Tamagawa products for elliptic curves over number fields*, arXiv, 2021.
- [2] John Cremona and Mohammad Sadek, *Local and global densities for Weierstrass models of elliptic curves*, arXiv, 2020.
- [3] Michael Griffin, Ken Ono, and Wei-Lun Tsai, *Tamagawa products of elliptic curves over  $\mathbb{Q}$* , The Quarterly Journal of Mathematics **72** (2021), no. 4, 1517–1543.
- [4] Giacomo Micheli, *A local to global principle for densities over function fields*, arXiv, 2017.
- [5] Michael Rosen, *Number theory in function fields*, 1st ed., Springer, 2002.
- [6] Joseph H. Silverman, *Advanced topics in the arithmetic of elliptic curves*, 1st ed., Springer, 1994.
- [7] John Tate, *Algorithm for determining the type of a singular fiber in an elliptic pencil*, Modular Functions of One Variable IV, 1975, pp. 33–52.