

Dispersed graph labellings

William J. Martin^{*1} and Douglas R. Stinson^{†2,3}

¹Department of Mathematical Sciences, Worcester Polytechnic Institute,
Worcester MA, 01609, USA

²David R. Cheriton School of Computer Science, University of Waterloo,
Waterloo ON, N2L 3G1, Canada

³School of Mathematics and Statistics, Carleton University, Ottawa, Ontario,
K1S 5B6, Canada

January 30, 2023

Abstract

A *k-dispersed labelling* of a graph G on n vertices is a labelling of the vertices of G by the integers $1, \dots, n$ such that $d(i, i+1) \geq k$ for $1 \leq i \leq n-1$. $\text{DL}(G)$ denotes the maximum value of k such that G has a k -dispersed labelling. In this paper, we study upper and lower bounds on $\text{DL}(G)$. Computing $\text{DL}(G)$ is NP-hard. However, we determine the exact values of $\text{DL}(G)$ for cycles, paths, grids, hypercubes and complete binary trees. We also give a product construction and we prove a degree-based bound.

1 Introduction

Many graph labelling problems have been studied over the years, starting with the graceful labellings introduced by Rosa. Gallian's dynamic survey [3] is an excellent starting point for this area of research. We assume standard graph-theoretic terminology throughout this paper, e.g., as defined in [1].

Let G be a graph having vertex set V , where $|V| = n$. Let $d(u, v)$ denote the distance between any two vertices u and v in G . It is easy to observe that G has a hamiltonian path if and only if there is a labelling of the vertices with the integers $1, \dots, n$ such that $d(i, i+1) = 1$ for $1 \leq i \leq n-1$. Here we consider a labelling problem motivated by the requirement that consecutively labelled vertices should be far apart. Thus we define a *k-dispersed labelling* to be a labelling of the vertices of G by the integers $1, \dots, n$ such that $d(i, i+1) \geq k$ for $1 \leq i \leq n-1$. Equivalently,

^{*}W.J. Martin's research is supported by NSF DMS Award #1808376.

[†]D.R. Stinson's research is supported by NSERC discovery grant RGPIN-03882.

for a graph $G = (V, E)$ with $|V| = n$, we could define a k -dispersed labelling to be a bijection $\phi : \{1, \dots, n\} \rightarrow V$ such that $d(\phi(i), \phi(i+1)) \geq k$ for $1 \leq i \leq n-1$.

Although it is not a main topic of this paper, we could also consider a “circular” variant of the above definition. We define a k -circular-dispersed labelling to be a k -dispersed labelling that satisfies the additional property that $d(n, 1) \geq k$. Equivalently, a k -circular-dispersed labelling could be defined to be a bijection $\phi : \mathbb{Z}_n \rightarrow V$ such that $d(\phi(i), \phi(i+1)) \geq k$ for $0 \leq i \leq n-1$ (in this definition, for convenience, the vertices are labelled with the elements of \mathbb{Z}_n).

We note that most of the labelling problems discussed in [3] do not involve distances between the vertices in a graph. One exception is the problem of *radio labellings* [3, §7.4].

Let $\text{DL}(G)$ denote the maximum value of k such that G has a k -dispersed labelling and let $\text{DL}^\circ(G)$ denote the maximum value of k such that G has a k -circular-dispersed labelling. Here are two easy preliminary lemmas that we state without proof.

Lemma 1.1. *If H is a spanning subgraph of G (obtained by removing edges but no vertices), then $\text{DL}(H) \geq \text{DL}(G)$ and $\text{DL}^\circ(H) \geq \text{DL}^\circ(G)$.*

Lemma 1.2. $\text{DL}^\circ(G) \leq \text{DL}(G)$.

Our first real result relates the values $\text{DL}(G)$ to the distance k graph of G . The *distance k graph* of G , denoted G_k , is the graph in which two vertices x and y are joined by an edge if $d(x, y) = k$. Clearly $G_1 = G$. Denote $G_{k-1}^* = G_1 \cup \dots \cup G_{k-1}$ and let $H_k = (G_{k-1}^*)^c$. So two vertices of H_k are adjacent if the distance between them (in G) is at least k .

Theorem 1.3. *For a graph G , $\text{DL}(G) \geq k$ if and only if H_k contains a hamiltonian path. Further, $\text{DL}^\circ(G) \geq k$ if and only if H_k contains a hamiltonian cycle.*

Proof. Suppose that G is a graph on vertex set V , where $|V| = n$. Suppose G has a k -dispersed labelling, say α , and define $P = (1, \dots, n)$. For any i , $1 \leq i \leq n-1$, it holds that $d(i, i+1) \geq k$ (because α is a k -dispersed labelling) and hence $\{i, i+1\}$ is an edge of H_k . It follows that P is a hamiltonian path in H_k .

The proof of the converse result is similar, as is the proof of the corresponding result for $\text{DL}^\circ(G)$. \square

Corollary 1.4. *Computing $\text{DL}(G)$ is NP-hard.*

Proof. Suppose that \mathcal{O} is an oracle that computes $\text{DL}(G)$ in polynomial time. We can use \mathcal{O} to solve the NP-complete HAMILTONIAN PATH problem as follows. Given a graph G , run \mathcal{O} on G^c . Observe that the $H_2(G^c) = G$. So $\mathcal{O}(G^c) \geq 2$ if and only if G has a hamiltonian path. \square

The rest of the paper is organized as follows. In Section 2, we determine the exact value of $\text{DL}(G)$ for cycles, paths, grids, hypercubes and complete binary trees. For these classes of graphs, $\text{DL}(G) = r(G)$ or $r(G) - 1$, where $r(G)$ is the radius

of the graph G . In Section 3, we prove a product construction: we show that $\text{DL}^\circ(G \square H) \geq \text{DL}^\circ(G) + \text{DL}^\circ(H)$, provided that the number of vertices in G and H is relatively prime. In Section 4, we prove a degree-based lower bound on $\text{DL}(G)$. Finally, in Section 5, we list a few interesting open questions.

2 Computing $\text{DL}(G)$ for some classes of graphs

We first consider cycles, for which the values $\text{DL}(G)$ can easily be determined.

Theorem 2.1. *Let C_n denote a cycle of length n . Then $\text{DL}(C_n) = (n - 1)/2$ if n is odd, and $\text{DL}(C_n) = (n - 2)/2$ if n is even.*

Proof. The maximum distance between two points of C_n is $n/2$ if n is even and $(n - 1)/2$ if n is odd.

First, suppose that n is odd and let $k = (n - 1)/2$. The graph H_k is a single (hamiltonian) cycle of length n , so $\text{DL}(C_n) \geq k$ follows from Theorem 1.3. Also, H_{k+1} is the empty graph, so $\text{DL}(C_n) \leq k$.

Next, suppose that n is even and let $k = n/2$. H_k consists of k disjoint edges, so H_k is not hamiltonian and therefore $\text{DL}(C_n) \leq k - 1$. We now study the structure of the graph H_{k-1} , which is a cubic graph. We consider two subcases.

First, suppose $n \equiv 0 \pmod{4}$. Here, the edges in H_{k-1} that are not in H_k form a hamiltonian cycle, so we are done. If $n \equiv 2 \pmod{4}$, then H_{k-1} is a prism; the edges in H_{k-1} that are not in H_k form two disjoint cycles of length $n/2$. It is an easy exercise to verify that the prism contains a hamiltonian path. Thus $\text{DL}(C_n) \geq k - 1$ when n is even, and the proof is complete. \square

The *eccentricity* of a vertex v is the quantity $\epsilon(v) = \max\{d(v, u) : u \in V\}$. The *radius* of a graph G , denoted $r(G)$, is the minimum eccentricity of any vertex, i.e., $r(G) = \min\{\epsilon(v) : v \in V\}$.

Theorem 2.2. *For a graph G , $\text{DL}(G) \leq r(G)$.*

Proof. Suppose that G has a k -dispersed labelling. Let i be a vertex such that $\epsilon(i) = r(G)$. If $i < n$, then let $j = i + 1$; if $i = n$, then let $j = i - 1$. We must have $d(i, j) \geq k$ since the labelling is k -dispersed. However, $d(i, j) \leq \epsilon(i) = r(G)$. Therefore, since $k \geq \text{DL}(G)$, it follows that $\text{DL}(G) \leq r(G)$. \square

We computed $\text{DL}(C_n)$ for all cycles C_n in Theorem 2.1. It is easy to verify that $r(C_n) = n/2$ if n is even and $r(C_n) = (n - 1)/2$ if n is odd. Hence, $\text{DL}(C_n) = r(C_n)$ if n is odd and $\text{DL}(C_n) = r(C_n) - 1$ if n is even.

We will require some additional related definitions for later use. A vertex is $v \in V$ is *uniquely eccentric* if there is a unique vertex u such that $d(u, v) = \epsilon(v)$. A vertex $v \in V$ is a *central vertex* if $\epsilon(v) = r(G)$.




Figure 1: The graph $L_{4,6}$

2.1 Paths

A simple class of graphs to consider are the paths. Let P_m denote the path having m edges and $m + 1$ vertices. It is easy to check that the radius of a path is given by the following formula:

$$r(P_m) = \begin{cases} \frac{m}{2} & \text{if } m \text{ is even} \\ \frac{m+1}{2} & \text{if } m \text{ is odd.} \end{cases}$$

Theorem 2.3. $\text{DL}(P_m) = r(P_m)$ for any path P_m .

Proof. First, suppose m is even. An $\frac{m}{2}$ -dispersed labelling is as follows:

$$2 \ 4 \ \cdots \ m \ 1 \ 3 \ \cdots \ m + 1.$$

For odd m , an $\frac{m+1}{2}$ -dispersed labelling is as follows:

$$2 \ 4 \ \cdots \ m + 1 \ 1 \ 3 \ \cdots \ m. \quad \square$$

2.2 Grids

As another, more complicated class of graphs, we consider the $m \times n$ grid graphs (or lattice graphs), which we denote by $L_{m,n}$. The graph $L_{4,6}$ is depicted in Figure 1. The following lemma will be useful.

Lemma 2.4. *Suppose a graph G contains three central vertices, each of which is uniquely eccentric. Then $\text{DL}(G) \leq r(G) - 1$.*

Proof. Suppose that G contains n vertices that are labelled $1, \dots, n$. Further, suppose that $\text{DL}(G) = r(G)$. At least one of the three hypothesized central vertices must receive a label i , where $2 \leq i \leq n - 1$. Consider the vertices labelled $i - 1$ and $i + 1$. Since vertex i is uniquely eccentric, either $d(i - 1, i) > r(G)$ or $d(i, i + 1) > r(G)$. This contradicts the assumption that $\text{DL}(G) = r(G)$. \square




Figure 2: A 4-dispersed labelling of $L_{2,7}$

The graph $L_{4,6}$, has radius $r(L_{4,6}) = 5$ and there are four central vertices, namely vertices 9, 10, 15 and 16. Each of these four central vertices is uniquely eccentric: $d(9, 24) = 5$, $d(10, 19) = 5$, $d(15, 6) = 5$ and $d(16, 11) = 5$. (To illustrate, vertex 24 is the only vertex that is distance five from vertex 9. A path of length five from 9 to 24 is indicated by the blackened vertices in Figure 1.) Therefore, from Lemma 2.4, it follows that $\text{DL}(L_{4,6}) \leq 4$. We will prove a bit later that $\text{DL}(L_{4,6}) = 4$.

More generally, we have the following upper bound.

Theorem 2.5. *Suppose m and n are both even. Then $\text{DL}(L_{m,n}) \leq (m+n)/2 - 1$.*

Proof. When m and n are even, it is easy to see that $r(L_{m,n}) = (m+n)/2$ and this graph has four central vertices, each of which is uniquely eccentric. Apply Lemma 2.4. \square

For other cases of m and n , we note the following.

Lemma 2.6.

1. *Suppose m and n are both odd. Then $r(L_{m,n}) = (m+n)/2 - 1$ and $L_{m,n}$ has one central vertex, which is not uniquely eccentric.*
2. *Suppose $m+n$ is odd. Then $r(L_{m,n}) = (m+n-1)/2$ and $L_{m,n}$ has two central vertices, neither of which is uniquely eccentric.*

Thus Lemma 2.4 cannot be applied in these cases, so we cannot rule out the possibility that $\text{DL}(L_{m,n}) = r(L_{m,n})$ if at least one of m and n is odd. In fact, we will prove in this section that $\text{DL}(L_{m,n}) = r(L_{m,n})$ if at least one of m and n is odd; and $\text{DL}(L_{m,n}) = r(L_{m,n}) - 1$ if m and n are both even.

First, we solve the case of $2 \times n$ grids.

Theorem 2.7. $\text{DL}(L_{2,n}) = r(L_{2,n}) = (n+1)/2$ for all odd $n \geq 3$.

Proof. The first row of $L_{2,n}$ is labelled $2, 4, \dots, 2n$ and the second row is labelled $n+2, n+4, \dots, 2n-1, 1, 3, \dots, n$. \square

Theorem 2.8. $\text{DL}(L_{2,n}) = r(L_{2,n}) - 1 = n/2$ for all even $n \geq 4$.

Proof. The first row of $L_{2,n}$ is labelled $2, 4, \dots, 2n$ and the second row is labelled $n+3, n+5, \dots, 2n-1, 1, 3, \dots, n+1$. \square




Figure 3: A 4-dispersed labelling of $L_{2,8}$

See Figures 2 and 3 for illustrations of the constructions in Theorems 2.7 and 2.8, respectively.

Suppose $m \geq 4$ is even. We can construct optimal labellings of $m \times n$ grids from optimal labellings of $2 \times n$ grids recursively.

Theorem 2.9. *Suppose $m \geq 2$ is even. Then*

$$DL(L_{m,n}) = \begin{cases} \frac{m+n-2}{2} & \text{if } n \text{ is even} \\ \frac{m+n-1}{2} & \text{if } n \text{ is odd.} \end{cases}$$

Proof. For odd n , we start with the $\frac{n+1}{2}$ -dispersed labelling of $L_{2,n}$ constructed in Theorem 2.7; for even n , we start with the $\frac{n}{2}$ -dispersed labelling of $L_{2,n}$ constructed in Theorem 2.8. Let the first row of one of these optimal labellings be denoted A and let the second row be denoted B . Then construct the labelling of $L_{m,n}$ having the following m rows:

A
$A + 2n$
\vdots
$A + (m - 2)n$
B
$B + 2n$
\vdots
$B + (m - 2)n$

See Figure 4 for an illustration of the construction when $m = 4$ and $n = 7$.

It is not hard to prove that the result is an $\frac{m+n-2}{2}$ -dispersed labelling of $L_{m,n}$ when n is even, and an $\frac{m+n-1}{2}$ -dispersed labelling of $L_{m,n}$ when n is odd. Basically, we have interleaved $m/2$ isomorphic copies of the labellings of $L_{2,n}$. The interleaving increases the distances between consecutively labelled vertices within a particular $L_{2,n}$ by $m/2 - 1$ (this is because, for any two consecutively labelled vertices, one is in A and the other is in B). In the case of even n , the minimum distance of $n/2$ is increased to $n/2 + m/2 - 1 = (m + n - 2)/2$, as desired. For odd n , the minimum distance of $(n + 1)/2$ is increased to $(n + 1)/2 + m/2 - 1 = (m + n - 1)/2$, as desired.

It is also necessary to consider the distance between the “last vertex” in one copy of $L_{2,n}$ and the “first” vertex in the next copy. The first vertex in a copy of $L_{2,n}$ is the middle element (when n is odd) or the leftmost of the two middle elements




Figure 4: A 4-dispersed labelling of $L_{4,7}$

(when n is even) of row B , while the last element is the rightmost vertex in row A . The distance between these two vertices is $(n + 1)/2$ when n is odd, and $(n + 2)/2$ when n is even. The interleaving increases the distance by $m/2$, which is more than what is required in the resulting labelling of $L_{m,n}$. \square

We now study $L_{m,n}$ for odd m . We can also assume that n is odd, because the case of odd m and even n is equivalent to the case where m is even and m is odd, and this is covered by Theorem 2.9. We begin with $m = 3$.

Theorem 2.10. $\text{DL}(L_{3,n}) = r(L_{3,n}) = (n + 1)/2$ for all odd $n \geq 3$.

Proof. It suffices to present an $\frac{n+1}{2}$ -dispersed labelling of $L_{3,n}$ when n is odd. The first row of $L_{2,n}$ is labelled

$$6, 9, \dots, 3n, 3;$$

the second row is labelled

$$\frac{3n+7}{2}, \frac{3n+13}{2}, \dots, 3n-1, 2, 5, \dots, \frac{3n+1}{2};$$

and the third row is labelled

$$1, 4, \dots, 3n-2$$

To get from vertex j to vertex $j + 1$, we proceed $(n - 1)/2$ vertices to the right and one row up, wrapping around if necessary. So the distance is at least $(n - 1)/2 + 1 = (n + 1)/2$ (because wrapping around cannot decrease the distance between the two vertices). \square

For future use, we observe that the vertex labelled 1 is in the bottom left corner of the grid. The vertex labelled 2 is the central vertex of the grid). The last vertex (i.e., the one labelled $3m$) is immediately to the left of the vertex in the top right corner of the grid.

Example 2.1. We construct a 3-dispersed labelling of $L_{3,5}$ using Theorem 2.10; see Figure 5. Therefore, $\text{DL}(L_{3,5}) = 3$.




Figure 5: A 3-dispersed labelling of $L_{3,5}$

Theorem 2.11. $DL(L_{m,n}) = r(L_{m,n}) = (n + m)/2 - 1$ for all odd $m, n \geq 3$.

Proof. In view of Theorem 2.10, we can assume that $m \geq 5$. Denote $m = 3 + 2t$; then $t = (m - 3)/2$ and $t \geq 1$.

Let the $(n + 1)/2$ -dispersed labelling of $L_{2,n}$ have rows A and B , as in the proof of Theorem 2.9. Also, let the $(n + 1)/2$ -dispersed labelling of $L_{3,n}$ constructed in Theorem 2.10 have rows C, D and E . Then construct the labelling of $L_{m,n}$ having the following m rows:

C
$A + 3n$
$A + 5n$
\vdots
$A + (2t + 1)n$
D
$B + 3n$
$B + 5n$
\vdots
$B + (2t + 1)n$
E

See Figure 6 for an illustration of the construction when $m = 5$ and $n = 5$.

We are interleaving the $\frac{n+1}{2}$ -dispersed labelling of $L_{3,n}$ with t copies of the $\frac{n+1}{2}$ -dispersed labelling of $L_{2,n}$. We need to prove that the distance between consecutively labelled vertices (say j and $j + 1$) is at least $(m + n - 2)/2$. The proof naturally divides into a number of cases:

case 1 Vertices j and $j + 1$ are both in the same copy of $L_{2,n}$.

case 2 Vertex j is the last vertex in one copy of $L_{2,n}$ and vertex $j + 1$ is the first vertex in the next copy of $L_{2,n}$.

case 3 Vertices j and $j + 1$ are both in the $L_{3,n}$.

case 4 Vertex j is the last vertex in the $L_{3,n}$ and vertex $j + 1$ is the first vertex in the first copy of $L_{2,n}$ (i.e., $j = 3n$).




Figure 6: A 4-dispersed labelling of $L_{5,5}$

Case 1 is similar to the analogous case in the proof of Theorem 2.9. The distance between vertices j and $j + 1$ in $L_{m,n}$ is t larger than the distance in $L_{3,n}$. So the distance is at least

$$\frac{n+1}{2} + t = \frac{n+1+m-3}{2} = \frac{m+n-2}{2}.$$

Case 2 is also similar to the analogous case in the proof of Theorem 2.9.

For case 3, the distance between vertices j and $j + 1$ in $L_{m,n}$ is t larger than the distance in $L_{3,n}$. As in case 1, the distance is at least $(m+n-2)/2$.

For case 4, we use the fact that vertex $3n$ is in row 1 and column $n-1$, and vertex $3n+1$ is in row $t+3$ and column $(n+1)/2$. So the distance is at least

$$\frac{n-3}{2} + t + 2 = \frac{n-3+m-3}{2} + 2 = \frac{m+n-2}{2}.$$

This completes the proof. □

2.3 Hypercubes

The n -dimensional hypercube, denoted \mathcal{Q}_n , is a graph having vertex set $V = (\mathbb{Z}_2)^n$. Two vertices are adjacent in \mathcal{Q}_n if they differ in exactly one co-ordinate. For $\mathbf{v}, \mathbf{w} \in V$, where $\mathbf{v} = (v_1, \dots, v_n)$ and $\mathbf{w} = (w_1, \dots, w_n)$, it is easy to see that

$$d(\mathbf{v}, \mathbf{w}) = |\{i : v_i \neq w_i\}|.$$

Here are some other easily verified properties of \mathcal{Q}_n .

Lemma 2.12. *Let $\mathbf{v} \in (\mathbb{Z}_2)^n$ and suppose $0 \leq i \leq n$. Then there are precisely $\binom{n}{i}$ vertices \mathbf{w} such that $d(\mathbf{v}, \mathbf{w}) = i$ in the graph \mathcal{Q}_n .*

Corollary 2.13. *For any integer $n \geq 2$, $r(\mathcal{Q}_n) = n$. Further, every vertex is a uniquely eccentric central vertex.*

Lemma 2.4 and Corollary 2.13 immediately imply that $\text{DL}(\mathcal{Q}_n) \leq n - 1$ for all $n \geq 2$. We will prove that $\text{DL}(\mathcal{Q}_n) \geq n - 1$ by constructing a suitable labelling of the vertices of \mathcal{Q}_n . It suffices to find a permutation of the n -tuples in V , say $\mathbf{v}_1, \dots, \mathbf{v}_{2^n}$, such that $d(\mathbf{v}_j, \mathbf{v}_{j+1}) \geq n - 1$ for $1 \leq j \leq 2^n - 1$. In the resulting dispersed labelling, vertex \mathbf{v}_j is labelled j for $1 \leq j \leq 2^n$.

We construct the permutations recursively. In order for the recursive construction to work, we require some additional properties. Hence, for each $n \geq 2$, we will construct a particular permutation $\Pi_n = (\mathbf{v}_1, \dots, \mathbf{v}_{2^n})$ satisfying the following properties.

1. $d(\mathbf{v}_j, \mathbf{v}_{j+1}) \geq n - 1$ for $1 \leq j \leq 2^n - 1$,
2. $\mathbf{v}_1 = (0, \dots, 0)$,
3. $\mathbf{v}_{2^{n-1}} = (1, \dots, 1, 0)$,
4. $\mathbf{v}_{2^{n-1}+1} = (0, \dots, 0, 1)$,
5. $\mathbf{v}_{2^n} = (1, \dots, 1)$.

Construction 2.14. We describe how to construct the permutations Π_n . To begin, when $n = 2$, Π_2 is the following permutation:

$$(0, 0), (1, 0), (0, 1), (1, 1).$$

Now we construct Π_{n+1} from Π_n . For notational convenience, we denote $\Pi_n = (\mathbf{v}_1, \dots, \mathbf{v}_{2^n})$ and $\Pi_{n+1} = (\mathbf{w}_1, \dots, \mathbf{w}_{2^{n+1}})$.

- For $1 \leq j \leq 2^{n-1}$, define

$$\mathbf{w}_j = \begin{cases} (\mathbf{v}_j, 0) & \text{if } j \text{ is odd} \\ (\mathbf{v}_j, 1) & \text{if } j \text{ is even.} \end{cases}$$

- For $2^{n-1} + 1 \leq j \leq 2^n$, define

$$\mathbf{w}_j = \begin{cases} (\mathbf{v}_j, 1) & \text{if } j \text{ is odd} \\ (\mathbf{v}_j, 0) & \text{if } j \text{ is even.} \end{cases}$$

- For $2^n + 1 \leq j \leq 2^n + 2^{n-1}$, define

$$\mathbf{w}_j = \begin{cases} (\mathbf{v}_{j-2^n}, 1) & \text{if } j \text{ is odd} \\ (\mathbf{v}_{j-2^n}, 0) & \text{if } j \text{ is even.} \end{cases}$$

- For $2^n + 2^{n-1} + 1 \leq j \leq 2^{n+1}$, define

$$\mathbf{w}_j = \begin{cases} (\mathbf{v}_{j-2^n}, 0) & \text{if } j \text{ is odd} \\ (\mathbf{v}_{j-2^n}, 1) & \text{if } j \text{ is even.} \end{cases}$$

Example 2.2. Π_3 is the following permutation:

$$(0, 0, 0), (1, 0, 1), (0, 1, 1), (1, 1, 0), (0, 0, 1), (1, 0, 0), (0, 1, 0), (1, 1, 1).$$

Theorem 2.15. For all $n \geq 2$, $\text{DL}(\mathcal{Q}_n) = n - 1$.

Proof. We have already observed that $\text{DL}(\mathcal{Q}_n) \leq n - 1$, so it suffices to prove that $\text{DL}(\mathcal{Q}_n) \geq n - 1$. This follows by showing that each of the permutations Π_n obtained from Construction 2.14 satisfies the properties 1–5 enumerated above, which we prove by induction on n .

First, we observe that each Π_n is a permutation. This is because every n -tuple in Π_n is appended with both a 0 and a 1 in the construction of Π_{n+1} .

For $n = 2$, the stated properties can be verified easily. Now, suppose that properties 1–5 hold for a particular Π_n , where $n \geq 2$. We will show that properties 1–5 hold for Π_{n+1} . We use the notation from Construction 2.14.

When $j \neq 2^n$, \mathbf{w}_j and \mathbf{w}_{j+1} are obtained from two consecutive \mathbf{v}_j 's, which are assumed to have distance at least $n - 1$, by induction. In most cases, one of \mathbf{w}_j and \mathbf{w}_{j+1} is appended with a 1 and the other is appended with a 0. In these cases, $d(\mathbf{w}_j, \mathbf{w}_{j+1}) \geq n$.

The only cases where \mathbf{w}_j and \mathbf{w}_{j+1} are appended with the same symbol are when $j = 2^{n-1}, 2^n$ or $2^n + 2^{n-1}$. When $j = 2^{n-1}$ or $2^n + 2^{n-1}$, properties 3 and 4 ensure that \mathbf{w}_j and \mathbf{w}_{j+1} are obtained from two consecutive \mathbf{v}_j 's that have distance n . When $j = 2^n$, \mathbf{w}_j and \mathbf{w}_{j+1} are obtained from \mathbf{v}_{2^n} and \mathbf{v}_1 (resp.), which have distance n due to properties 2 and 5. Hence, property 1 is satisfied for Π_{n+1} .

It is easy to verify that properties 2–5 hold for Π_{n+1} , so the proof is complete. \square

We note that there is an alternative approach that gives an easier proof of Theorem 2.15 in the case where n is even. It is well-known that, for n even, the distance $n - 1$ graph of \mathcal{Q}_n is isomorphic to \mathcal{Q}_n (see, e.g., [2, p. 265, Remark (i)]). One such isomorphism is obtained by flipping the bits of all odd-weight n -tuples. It is also well-known that \mathcal{Q}_n has a hamiltonian cycle; the *binary reflected gray code* is one example (see, e.g., [4]). Hence it immediately follows that $\text{DL}^\circ(\mathcal{Q}_n) = \text{DL}(\mathcal{Q}_n) = n - 1$ if n is even.

The distance $n - 1$ graph of \mathcal{Q}_n is disconnected if n is odd, so this particular approach will not work.

2.4 Complete Binary Trees

The radius of the complete binary tree of depth n is n . We will show that the complete binary tree of depth n admits an n -dispersed labelling.




Figure 7: A 3-dispersed labelling of the complete binary tree of depth three.

Let T have vertex set $\{\varepsilon\} \cup \{0, 1\} \cup \dots \cup \{0, 1\}^n$ where ε denotes the empty string. We have an edge joining \mathbf{x} to \mathbf{y} if one can be obtained from the other by deleting the rightmost bit. Let $|\mathbf{x}|$ denote the length of the binary string \mathbf{x} . A 3-dispersed labelling of a complete binary tree of depth three is given in Figure 7.

Our labelling is as follows. The root node, ε , receives label 1. The “left” nodes—those vertices whose strings begin with zero—receive even labels, with the rule that labels $2, 4, 6, \dots, 2^n$ are used on leaf nodes and the remaining even labels occur on nodes of degree three. The “right” nodes—those vertices whose strings begin with one—receive odd labels, with the rule that labels $2^n + 1, 2^n + 3, \dots, 2^{n+1} - 1$ are used for leaf nodes and the remaining odd labels $\ell \geq 3$ occur on nodes of degree three.

A concrete rule for such a labelling is given as follows, where $N(\mathbf{x})$ denotes the integer whose binary representation is \mathbf{x} . We set

$$\ell(\mathbf{x}) = \begin{cases} 2N(\mathbf{x}) + 1 & \text{if } x_1 = 1; \\ 2N(\mathbf{x}) + 2 & \text{if } x_1 = 0 \text{ and } \mathbf{x} \text{ is a leaf node;} \\ 2N(\mathbf{x}) + 2^n + 2^{|\mathbf{x}|} & \text{if } x_1 = 0 \text{ and } \mathbf{x} \text{ is not a leaf node.} \end{cases}$$

One may verify by inspection that each label $0 < i < 2^{n+1}$ is used once. Among any consecutive pair of labels $i, i + 1$, one of these appears at a leaf node and the path from this node to the other one passes through the root of the tree and therefore has length at least n .

Therefore we have proven the following result.

Theorem 2.16. *Let T_n denote the complete binary tree of depth n . Then $\text{DL}(T_n) = n$.*

Proof. The labelling described above shows that $\text{DL}(T_n) \geq n$. Since T_n has radius n , it follows immediately that $\text{DL}(T_n) = n$. \square

3 A product construction

The *Cartesian product* of graphs $G = (V(G), E(G))$ and $H = (V(H), E(H))$, denoted $G \square H$, is the graph with vertex set $V(G) \times V(H)$ and edge set

$$E(G \square H) = \{ \{(u, v), (u', v')\} : u = u', \{v, v'\} \in E(H) \text{ or } \{u, u'\} \in E(G), v = v' \}.$$

Suppose G is a graph on m vertices having a k -circular-dispersed labelling and H is a graph on n vertices having a k' -circular-dispersed labelling. When $\gcd(m, n) = 1$, we will show how to combine these two labellings to give a $(k + k')$ -circular-dispersed labelling of the product graph $G \square H$.

Theorem 3.1. *Let G and H be graphs with $|V(G)| = m$, $|V(H)| = n$ and $\gcd(m, n) = 1$. Then $\text{DL}^\circ(G \square H) \geq \text{DL}^\circ(G) + \text{DL}^\circ(H)$.*

Proof. Denote $k = \text{DL}^\circ(G)$ and $k' = \text{DL}^\circ(H)$. Suppose $\ell_G : \mathbb{Z}_m \rightarrow V(G)$ is a k -circular-dispersed labelling of G and $\ell_H : \mathbb{Z}_n \rightarrow V(H)$ is a k' -circular-dispersed labelling of H . For $0 \leq i \leq mn - 1$, define

$$\ell(i) = (\ell_G(i \bmod m), \ell_H(i \bmod n)).$$

Since $\gcd(m, n) = 1$, ℓ is a bijection from \mathbb{Z}_{mn} to the vertices of $G \square H$. The distance between two vertices in $G \square H$ is just the sum of the distances between the corresponding projections (i.e., to the first and second coordinates) in G and H . Hence, for any i , where $0 \leq i \leq mn - 1$, The distance between $\ell(i)$ and $\ell(i + 1 \bmod mn)$ in $G \square H$ is the sum

$$d_G(i \bmod m, i + 1 \bmod m) + d_H(i \bmod n, i + 1 \bmod n) \geq k + k',$$

as desired. □

We illustrate the product construction by returning to grids. Our goal is not to compute all the values $\text{DL}^\circ(L_{m,n})$, but rather to illustrate how Theorem 3.1 can be applied to a specific family of graphs.

Recall that $L_{m,n}$ is an m by n grid and P_{m-1} is a path having m vertices. It is easy to see that $P_{m-1} \square P_{n-1}$ is isomorphic to $L_{m,n}$. We can obtain lower bounds on $\text{DL}^\circ(L_{m,n})$ if we have lower bounds on $\text{DL}^\circ(P_{m-1})$ and $\text{DL}^\circ(P_{n-1})$. Therefore, we first look at circular-dispersed labellings of paths.

Lemma 3.2. *Suppose a graph $G = (V, E)$ contains two central vertices, say u and v , both of which are uniquely eccentric. Suppose also that $\{u, v\} \in E$ and suppose $r(G) \geq 2$. Then $\text{DL}^\circ(G) \leq r(G) - 1$.*

Proof. Suppose that $|V| = n$ and $\text{DL}^\circ(G) = r(G)$. Vertices u and v must receive the labels 1 and n since they are uniquely eccentric. However, $d(u, v) = 1 \leq r(G) - 1$, so it follows that $\text{DL}^\circ(G) \leq r(G) - 1$. □

Lemma 3.3. *Let P_m be the path with m edges (and $m + 1$ vertices). If m is even, then $\text{DL}^\circ(P_m) = \text{DL}(P_m) = \frac{m}{2}$. If $m \geq 3$ is odd, then $\text{DL}^\circ(P_m) = \text{DL}(P_m) - 1 = \frac{m-1}{2}$.*

Proof. We already have shown that $\text{DL}(P_m) = r(P_m)$ in Theorem 2.3. For m even, the labelling

$$2 \ 4 \ \cdots \ m \ 1 \ 3 \ \cdots \ m+1$$

given in Theorem 2.3 has $d(1, m+1) = \frac{m}{2}$. Hence, $\text{DL}^\circ(P_m) = \text{DL}(P_m) = \frac{m}{2}$ when m is even.

However, when $m \geq 3$ is odd, P_m contains two uniquely eccentric central vertices that are adjacent, so Lemma 3.2 asserts that $\text{DL}^\circ(P_m) \leq r(P_m) - 1 = (m-1)/2$. To achieve a $\frac{m-1}{2}$ -circular dispersed labelling of P_m , we start with the above labelling of P_{m-1} (on m vertices) and attach a new vertex labelled $m+1$ to the left end of the path. \square

Theorem 3.4. *Suppose $\text{gcd}(m, n) = 1$, $m \geq 3$ and $n \geq 3$. Then*

$$\text{DL}^\circ(L_{m,n}) \geq \begin{cases} \frac{m+n}{2} - 1 & \text{if } m, n \text{ are both odd} \\ \frac{m+n-3}{2} & \text{if } m+n \text{ is odd.} \end{cases}$$

Proof. This is an immediate application of Theorem 3.2 and Lemma 3.3. \square

We do not consider the case where both m and n are even in Theorem 3.4. This is because $\text{gcd}(m, n) \geq 2$ when m and n are even and hence the hypotheses cannot be satisfied.

Corollary 3.5. *Suppose $\text{gcd}(m, n) = 1$, $m \geq 3$ is odd and $n \geq 3$ is odd. Then*

$$\text{DL}^\circ(L_{m,n}) = \frac{m+n}{2} - 1.$$

Proof. From Theorem 2.11, we have $\text{DL}(L_{m,n}) = \frac{m+n}{2} - 1$ when $n \geq 3$ and $m \geq 3$ are odd. Lemma 1.2 shows that $\text{DL}^\circ(L_{m,n}) \leq \text{DL}(L_{m,n})$. Finally, Theorem 3.4 shows that $\text{DL}^\circ(L_{m,n}) \geq \frac{m+n}{2} - 1$ if $n \geq 3$ and $m \geq 3$ are odd and $\text{gcd}(m, n) = 1$. \square

4 Degree-based bounds

We now consider a degree-based lower bound on $\text{DL}(G)$. Suppose G has n vertices. For $x \in V$ and $i \geq 0$, let

$$\eta_i(x) = |\{y \in V : d(x, y) = i\}|,$$

and

$$\kappa(x) = \max \left\{ j : \sum_{i=0}^j \eta_i(x) \leq n/2 \right\}.$$

For example, in \mathcal{Q}_m , it is easy to see that $\eta_i(x) = \binom{m}{i}$ for each vertex x and $\kappa(x) = \lfloor \frac{m-1}{2} \rfloor$.

Lemma 4.1. $\text{DL}(G) \geq \min\{1 + \kappa(x) : x \in V(G)\}$.

Proof. For any vertex x , there are at least $n/2$ vertices y such that $d(x, y) \geq 1 + \kappa(x)$. Therefore, by Dirac's Theorem (e.g., see [1, Theorem 18.4]), H_k is hamiltonian for $k = \min\{1 + \kappa(x) : x \in V(G)\}$. Now apply Theorem 1.3. \square

Theorem 4.2. *If G has n vertices and maximum degree $\Delta > 2$, then*

$$\text{DL}(G) \geq 1 + \log_{\Delta-1} \left[1 + \frac{(n-2)(\Delta-2)}{2\Delta} \right]. \quad (1)$$

Proof. For any $x \in V(G)$ and $i \geq 1$, $n_i(x) \leq \Delta(\Delta-1)^{i-1}$. In order to apply Lemma 4.1, we compute the largest value of ℓ such that

$$1 + \Delta + \Delta(\Delta-1) + \cdots + \Delta(\Delta-1)^\ell \leq n/2.$$

This inequality is equivalent to

$$1 + \frac{\Delta(1 - (\Delta-1)^{\ell+1})}{2 - \Delta} \leq \frac{n}{2}.$$

Elementary algebra tells us that this can be rewritten as

$$\ell + 1 \leq \log_{\Delta-1} \left[1 + \frac{(n-2)(\Delta-2)}{2\Delta} \right].$$

Since $\kappa(x) \geq \ell + 1$, it therefore follows from Lemma 4.1 that $\text{DL}(G) \geq \ell + 2$ and hence the inequality (1) holds. \square

Corollary 4.3. *If G has n vertices and maximum degree three, then*

$$\text{DL}(G) \geq \log_2 \left(\frac{n+4}{6} \right) + 1.$$

Proof. Apply Theorem 4.2 with $\Delta = 3$. \square

For example, if $n = 3 \times 2^t - 4$ and $\Delta = 3$, then Corollary 4.3 says that $\text{DL}(G) \geq t$.

5 Open questions

There are many interesting open questions. We list a few now.

1. Which graphs G have $\text{DL}(G) = r(G)$?
2. We note that $\text{DL}(G) \in \{r(G), r(G) - 1\}$ for all the graphs considered in Section 2. We ask if there is an infinite family of graphs G_1, G_2, \dots such that

$$\lim_{i \rightarrow \infty} (r(G_i) - \text{DL}(G_i)) = \infty.$$

Actually, at this point, we do not have any examples of graphs G with $\text{DL}(G) \leq r(G) - 2$.

3. Is there an efficient algorithm to find “good” dispersed labellings of graphs (i.e., labellings where $DL(G)$ is “close to” $r(G)$) for certain classes of graphs (e.g., trees)?
4. Is there a generalization of the product construction (Theorem 3.1) that handles cases where $\gcd(m, n) > 1$?

References

- [1] J.A. Bondy and U.S.R. Murty. *Graph Theory*, Springer, 2008.
- [2] A.E. Brouwer, A.M. Cohen and A. Neumaier. *Distance-Regular Graphs*, Springer, 1989.
- [3] J.A. Gallian. Graph labeling, version 25. *Electron. J. Combin.*, Article #DS6, Dec. 2, 2022.
- [4] C. Savage. A survey of combinatorial Gray codes. *SIAM Review* **39** (1997), 605–629.